

Eccentricity and algebraic connectivity of graphs

B. AFSHARI, M. AFSHARI*

*Department of Computer Science and Engineering, Michigan State University

Abstract

Let G be a graph on n nodes with algebraic connectivity λ_2 . The eccentricity of a node is defined as the length of a longest shortest path starting at that node. If s_ℓ denotes the number of nodes of eccentricity at most ℓ , then for $\ell \geq 2$,

$$\lambda_2 \geq \frac{4 s_\ell}{(\ell - 2 + \frac{4}{n}) n^2}.$$

As a corollary, if d denotes the diameter of G , then

$$\lambda_2 \geq \frac{4}{(d - 2 + \frac{4}{n}) n}.$$

It is also shown that

$$\lambda_2 \geq \frac{s_\ell}{1 + \ell (e(G^\ell) - m)},$$

where m and $e(G^\ell)$ denote the number of edges in G and in the ℓ -th power of G , respectively.

2010 AMS Classification: 05C50, 15A18

Keywords: Laplacian eigenvalues of graphs, eccentricity, algebraic connectivity

Let G be a simple graph on $V = \{v_1, v_2, \dots, v_n\}$ with m edges E . The *distance* between nodes v_i and v_j , denoted by $\text{dist}(v_i, v_j)$, is the number of edges in a shortest path joining them. If there is no such path, then we define this value to be ∞ . The *diameter* of G , denoted by d is the maximum distance between any pair of nodes of G . The ℓ -th *power* of G , denoted by G^ℓ , is the graph with the same node set as G such that two nodes are adjacent in G^ℓ if and only

if their distance is at most ℓ in G . The *eccentricity* of a node in G is defined as the length of a longest shortest path starting at that node. We denote the number of nodes of eccentricity at most ℓ by s_ℓ .

The Laplacian matrix of G is defined as $L(G) = D - A$, where A is the adjacency matrix of G and D is a diagonal degree matrix with $D(i, i) = \sum_j A(i, j)$. The smallest eigenvalue of $L(G)$ is 0, where the corresponding eigenvector is the all one vector. The second smallest eigenvalue of $L(G)$, denoted by λ_2 , is called the *algebraic connectivity* of G . It is easy to see that the number of nodes with degree $n - 1$ is a lower bound for this parameter, that is,

$$\lambda_2 \geq s_1.$$

Also it is known that [1]

$$\lambda_2 \geq \frac{s_2}{n}.$$

Here it is shown that for $\ell \geq 3$,

$$\lambda_2 \geq \frac{4 s_\ell}{(\ell - 2 + \frac{4}{n})n^2}. \quad (1)$$

This result can be viewed as a generalization, with a slight improvement, of the well-known bound [3]: $\lambda_2(G) \geq \frac{4}{dn}$, as setting $\ell = d$ in (1) leads to $s_d = n$, and yields

$$\lambda_2 \geq \frac{4}{(d - 2 + \frac{4}{n})n}.$$

It is also shown that

$$\lambda_2 \geq \frac{s_\ell}{1 + \ell(e(G^\ell) - m)}, \quad (2)$$

where $e(G^\ell)$ denotes the number of edges in G^ℓ . This also generalizes the known result [2]:

$$\lambda_2 \geq \frac{n}{1 + d e(\overline{G})}.$$

Proofs

Proof of (1). If G is not connected, then $s_\ell = 0$, and the statement holds trivially. Therefore, we assume that G is connected. Let $\gamma = (\ell - 2 + \frac{4}{n})\frac{n^2}{4}$. It is sufficient to show that $M =$

$\gamma L(G) - L(G^\ell)$ is positive semi-definite. Because if so, assuming that z is the unit eigenvector corresponding to λ_2 , we can deduce (1) as follows:

$$\lambda_2 = z^T L(G) z \geq \frac{1}{\gamma} z^T L(G^\ell) z \geq \frac{1}{\gamma} \lambda_2(G^\ell) \geq \frac{1}{\gamma} s_1(G^\ell) = \frac{1}{\gamma} s_\ell.$$

Assume, to the contrary, that the minimum eigenvalue of M , denoted by μ , is negative, and let $x = (x_1, x_2, \dots, x_n)^T$ be an eigenvector corresponding to μ . Since the all one vector is an eigenvector corresponding to eigenvalue 0 of M , we have $\sum_{i=1}^n x_i = 0$. We may assume that

$$x_n \leq x_{n-1} \leq \dots \leq x_1.$$

For $k \in \{1, 2, \dots, n-1\}$, we define E_k as the set of edges $v_i v_j$ in G where $i \leq k < j$. Let k be chosen such that

$$\sum_{v_i v_j \in E_k} |x_i - x_j|$$

is maximized, with the assumption that this sum equals 1 without loss of generality. So, in particular, every edge $v_i v_j$ in G satisfies $|x_i - x_j| \leq 1$.

Let $S = \{v_1, v_2, \dots, v_k\}$. For $r \geq 1$, we define A_r as the set of nodes in S whose shortest paths to $V \setminus S$ have length r . Similarly, B_r is the set of nodes in $V \setminus S$ whose shortest paths to S have length r .

If $\text{dist}(v_i, v_j) = q$ for some $i \leq k < j$, then $v_i \in A_r$ and $v_j \in B_{r'}$ for some $r, r' \geq 1$ where $r + r' = q + 1$. Therefore,

$$\sum_{v_i v_j \in E(G^\ell): i \leq k < j} |x_i - x_j| \leq \sum_{r, r' \geq 1: r+r' \leq \ell+1} \sum_{v_i \in A_r, v_j \in B_{r'}} (x_i - x_j).$$

Let $r, r' \geq 2$. Let s be the smallest number such that $v_s \in A_{r-1}$ and t be the largest number such that $v_t \in B_{r'-1}$. We have

$$\sum_{\substack{v_i \in A_r \\ v_j \in B_{r'}}} (x_i - x_j) \leq \sum_{\substack{v_i \in A_r \\ v_j \in B_{r'}}} (x_s - x_t) + \sum_{v_i \in A_r} \sum_{v_j \in B_{r'}: j > t} (x_t - x_j) + \sum_{v_j \in B_{r'}} \sum_{v_i \in A_r: i < s} (x_i - x_s).$$

There is a path P of length $r-1$ from v_s to a node in $V \setminus S$, and a path P' of length $r'-1$ from v_t to a node in S . Both paths P and P' have an edge in E_k . Therefore, we conclude:

$$x_s - x_t \leq \sum_{v_i v_j \in E(P) \cup E(P')} |x_i - x_j| \leq r + r' - 3.$$

Moreover, for each $v_i \in A_r$ with $i < s$, there is some edge $v_i v_j$ with $j \geq s$ in G . Consequently,

$$\sum_{v_i \in A_r: i < s} (x_i - x_s) \leq \sum_{v_i v_j \in E_{s-1}} |x_i - x_j| \leq 1.$$

Similarly,

$$\sum_{v_j \in B_{r'}: j > t} (x_t - x_j) \leq \sum_{v_i v_j \in E_t} |x_i - x_j| \leq 1.$$

In consequence,

$$\sum_{v_i \in A_r, v_j \in B_{r'}} (x_i - x_j) \leq |A_r| |B_{r'}| (r + r' - 3) + |A_r| + |B_{r'}|.$$

For each $v_j \in B_1$, let $v_{j'}$ be the node adjacent to v_j in a shortest path from v_s to v_j . Also, denote by $P_{s j'}$ a shortest path from v_s to $v_{j'}$. We then have:

$$\begin{aligned} \sum_{v_i \in A_r, v_j \in B_1} (x_i - x_j) &\leq \sum_{v_j \in B_1} \left(\sum_{v_i \in A_r} (x_s - x_j) + \sum_{v_i \in A_r: i < s} (x_i - x_s) \right) \\ &\leq |A_r| \sum_{v_j \in B_1} ((x_s - x_{j'}) + (x_{j'} - x_j)) + |B_1| \\ &\leq |A_r| \sum_{v_j \in B_1} \sum_{v_a v_b \in P_{s j'}} |x_a - x_b| + |A_r| \sum_{v_i v_j \in E_k} |x_i - x_j| + |B_1| \\ &\leq |A_r| |B_1| (r - 2) + |A_r| + |B_1|. \end{aligned}$$

Similarly,

$$\sum_{v_i \in A_1, v_j \in B_{r'}} (x_i - x_j) \leq |A_1| |B_{r'}| (r - 2) + |A_1| + |B_{r'}|.$$

Let $A_1 = \{v_{s_1}, v_{s_2}, \dots, v_{s_p}\}$ and $B_1 = \{v_{s_{p+1}}, v_{s_{p+2}}, \dots, v_{s_q}\}$ where $s_1 < s_2 < \dots < s_q$. Note that for $i = 1, 2, \dots, q - 1$, we have

$$|E_k \cap E_{s_i}| \geq \min\{i, q - i\},$$

as there are edges in E_k with end nodes $v_{s_1}, v_{s_2}, \dots, v_{s_i}$ if $i \leq p$, and $v_{s_{i+1}}, v_{s_{i+2}}, \dots, v_{s_q}$ if $i \geq p + 1$. Suppose the vector y is obtained from x by translation such that $y_{s_{\lceil q/2 \rceil}}$ becomes

zero. This lead to,

$$\begin{aligned}
\sum_{v_i \in A_1, v_j \in B_1} (x_i - x_j) &= \sum_{v_i \in A_1, v_j \in B_1} (y_i - y_j) \leq \sum_{i < j} |y_{s_i} - y_{s_j}| = \frac{1}{2} \sum_{i=1}^q \sum_{j=1}^q |y_{s_i} - y_{s_j}| \\
&\leq \frac{1}{2} \sum_{i=1}^q \sum_{j=1}^q (|y_{s_i}| + |y_{s_j}|) \\
&= q \sum_{i=1}^q |y_{s_i}| \\
&= q \left(\sum_{i \leq q/2} y_{s_i} - \sum_{i > q/2} y_{s_i} \right) \\
&= q \sum_{i=1}^{q-1} \min\{i, q-i\} (y_{s_i} - y_{s_{i+1}}) \\
&\leq q \sum_{i=1}^{q-1} |E_k \cap E_{s_i}| (y_{s_i} - y_{s_{i+1}}) \\
&= q \sum_{v_i v_j \in E_k} |y_i - y_j| \leq q.
\end{aligned}$$

Now, we can deduce that,

$$\sum_{v_i v_j \in E(G^\ell): i \leq k < j} |x_i - x_j| \leq |A_1| + |B_1| + \sum_{r, r' \geq 1: 3 \leq r+r' \leq \ell+1} (r+r'-3)|A_r||B_{r'}| + |A_r| + |B_{r'}|.$$

The last term is bounded by the maximum possible value of

$$a_1 + b_1 + \sum_{r, r' \geq 1: 3 \leq r+r' \leq \ell+1} (r+r'-3)a_r b_{r'} + a_r + b_{r'},$$

where $a_r, b_r \geq 0$ for each r , subject to the constraints $\sum_{r=1}^{\ell} a_r = k$ and $\sum_{r'=1}^{\ell} b_{r'} = n - k$. By fixing the $b_{r'}$ values, this term reaches its maximum when $a_t = k$ for some t and $a_r = 0$ for $r \neq t$. Subsequently, after fixing the values for a_r 's, the term is maximized when $b_{\ell-t+1} = n - k$ and $b_{r'} = 0$ for $r' \neq \ell - t + 1$. Therefore,

$$\sum_{v_i v_j \in E(G^\ell): i \leq k < j} |x_i - x_j| \leq (\ell - 2)k(n - k) + n \leq (\ell - 2)\frac{n^2}{4} + n = \gamma.$$

Now, if $e_i \in \mathbb{R}^n$ is the i -th standard basis vector, then

$$\begin{aligned}
0 &\leq \gamma \sum_{v_i v_j \in E_k} |x_i - x_j| - \sum_{v_i v_j \in E(G^\ell): i \leq k < j} |x_i - x_j| \\
&= \sum_{i \leq k} \left(\gamma \sum_{v_i v_j \in E: j > k} (x_i - x_j) - \sum_{v_i v_j \in E(G^\ell): j > k} (x_i - x_j) \right) \\
&= \sum_{i \leq k} \left(\gamma \sum_{v_i v_j \in E} (x_i - x_j) - \sum_{v_i v_j \in E(G^\ell)} (x_i - x_j) \right) \\
&= (e_1 + e_2 + \dots + e_k)^T M x \\
&= \mu \sum_{i=1}^k x_i \\
&< 0,
\end{aligned}$$

which is a contradiction. Therefore, the proof is now complete. \square

Proof of (2). We need to show that $(1 + \ell (e(G^\ell) - m)) L(G) - L(G^\ell)$ is positive semi-definite. Consider a vector $x \in \mathbb{R}^n$, and let $v_r v_s \in E(G^\ell)$ be such that $|x_r - x_s|$ is maximum. Denote by P a shortest path between v_r and v_s . We then have:

$$\begin{aligned}
x^T L(G^\ell) x &= \sum_{v_i v_j \in E(G^\ell)} (x_i - x_j)^2 \\
&\leq x^T L(G) x + (e(G^\ell) - m) (x_r - x_s)^2 \\
&\leq x^T L(G) x + \ell (e(G^\ell) - m) \sum_{v_i v_j \in P} (x_i - x_j)^2 \\
&\leq (1 + \ell (e(G^\ell) - m)) x^T L(G) x.
\end{aligned}$$

This completes the proof. \square

References

- [1] B. Afshari, Algebraic connectivity of the second power of a graph, J. Graph Theory. (2023), 1–7.

- [2] M. Lu, L.-Z. Zhang and F. Tian, Lower bounds of the Laplacian spectrum of graphs based on diameter, *Linear Algebra Appl.* 420 (2007), 400–406.
- [3] B. Mohar, Eigenvalues, diameter, and mean distance in graphs, *Graph Combinator.* 7 (1991), 53–64.