# Low-Rank Toeplitz Matrix Restoration: Descent Cone Analysis and Structured Random Matrix\*

Gao Huang<sup>†</sup>and Song Li<sup>‡</sup>

July 4, 2024

#### Abstract

This note demonstrates that we can stably recover rank-r Toeplitz matrix  $X \in \mathbb{R}^{n \times n}$  from a number of rank-one subgaussian measurements on the order of  $r \log^2 n$  with an exponentially decreasing failure probability by employing a nuclear norm minimization program. Our approach utilizes descent cone analysis through Mendelson's small ball method with the Toeplitz constraint. The key ingredient is to determine the spectral norm of the random matrix of the Topelitz structure, which may be of independent interest. This improves upon earlier analyses and resolves the conjecture in Chen et al. (IEEE Transactions on Information Theory, 61(7):4034–4059, 2015).

**Keywords:** Toeplitz Matrix; Rank-One Measurement; Small Ball Method; Structured Random Matrix.

### **1** Introduction

Toeplitz matrices enjoy an important number of applications, including signal processing, numerical analysis, stochastic processes, time series analysis, image processing and so on [4]. Recovery of the matrix that is simultaneously low-rank and Toeplitz, which arises when the random process is generated by a few spectral spikes, is crucial in many tasks in wireless communications and array signal processing; see, e.g., [6, 7].

Following some ideas in [7], we aim to reconstruct an unknown symmetric matrix  $X_0 \in \mathbb{R}^{n \times n}$  with a structure that is simultaneously low-rank and Toeplitz from a small number of rank-one measurements. In particular, we explore sampling methods of the form

$$b_k = \langle \boldsymbol{\xi}_k \boldsymbol{\xi}_k^{\mathsf{I}}, \boldsymbol{X}_0 \rangle + e_k, \quad k = 1, \cdots, m,$$

<sup>\*</sup>This work was supported in part by the NSFC under grant numbers U21A20426 and 12071426

<sup>&</sup>lt;sup>†</sup>School of Mathematical Science, Zhejiang University, Hangzhou 310027, P. R. China, E-mail address: hgmath@zju.edu.cn

<sup>&</sup>lt;sup>‡</sup>Corresponding Author, School of Mathematical Science, Zhejiang University, Hangzhou 310027, P. R. China, E-mail address: songli@zju.edu.cn

where  $b := \{b_k\}_{k=1}^m$  denotes the measurements,  $\xi_k \in \mathbb{R}^n$  represents the sensing vector,  $e := \{e_k\}_{k=1}^m$  stands for the noise term, and m is the number of measurements. The rankone sampling scheme is utilised in a wide spectrum of practical settings, such as covariance sketching for data streams, noncoherent energy measurements in communications, and phaseless measurements in optical imaging [2, 7]. Furthermore, it admits near-optimal estimation with tractable algorithms and offers computational and storage advantages over other types of measurements.

Let us formally describe our setup and model, in accordance with [7]. We focus exclusively on the random sampling model. Specifically, we assume that the sensing vectors are given as independent copies of a random vector  $\boldsymbol{\xi}$ , whose entries  $\xi$  are assumed to be i.i.d. with subgaussian norm  $K^1$  and satisfy the moment properties:

$$\mathbb{E}\xi = 0, \ \mathbb{E}\xi^2 = 1, \ \text{and} \ \mu := \mathbb{E}\xi^4 \ge 1.$$
 (1)

We assume that the noise term e is bounded in  $\ell_p$ -norm  $(p \ge 1)$ . We also define the linear operator  $\mathcal{A}(\mathbf{X}) : \mathbb{R}^{n \times n} \to \mathbb{R}^m$  that maps a matrix  $\mathbf{X} \in \mathbb{R}^{n \times n}$  to  $\{\langle \boldsymbol{\xi} \boldsymbol{\xi}^*, \mathbf{X} \rangle\}_{k=1}^m$ . Thus, we express the measurements as

$$\boldsymbol{b}=\mathcal{A}\left(\boldsymbol{X}_{0}
ight)+\boldsymbol{e}.$$

A natural heuristic method is rank minimization to promote low-rank structures. However, the rank minimization problem is generally considered to be NP-hard. An alternative scheme is to seek a nuclear norm minimizer over all matrices compatible with the measurements, as well as the Toeplitz constraint:

minimize 
$$\|X\|_{*}$$
  
subject to  $\|\mathcal{A}(X) - y\|_{\ell_{p}} \leq \eta$  (2)  
 $X$  is Toeplitz.

Here  $\eta > 0$  is an a priori bound for the noise level; that is, we assume  $\|e\|_{\ell_n} \leq \eta$ .

Before we elaborate on our theorem, let's first introduce the previous work [7]. To solve the program (2) having an excess of positive semi-definite cone constraints, Chen et al. in [7] used stochastic RIP- $\ell_2/\ell_2$  and stated that when the noise is bounded in  $\ell_2$ -norm, rank-rToeplitz matrix  $X_0$  could be recovered from a number of rank-one subgaussian measurements, inspired by the Gaussian processes argument involving chaining techniques in [11]. Their method could be applicable to a wider range of settings; for instance, recovery of low-rank matrices with random subsampling using Fourier-type basis, and simultaneously low-rank and Toeplitz matrices with i.i.d. Gaussian measurement system.

Nevertheless, as they pointed out in [7] that their method, has certain constraints when dealing with the Toeplitz matrix situation, hence giving rise to some unresolved conjectures. First, their method has limited applicability for rank-one measurements. As  $\mu = \mathbb{E}\xi^4 \ge$ 

<sup>&</sup>lt;sup>1</sup>The subgaussian norm of a random variable X is defined as  $||X||_{\psi_2} := \inf\{t > 0 : \mathbb{E}\exp\left(|X|^2/t^2\right) \le 2\}.$ 

 $(\mathbb{E}\xi^2)^2 = 1$ , they are only able to provide theoretical guarantees for  $1 < \mu \le 3$ ; otherwise, the tails of measurement's distributions are typically not heavier than the Gaussian measure (e.g.,  $\mu = 3$  for Gaussian distribution), but it cannot be the Bernoulli measure (e.g.,  $\mu = 1$  for Bernoulli distribution). The main obstacle is that the rank-one measurement does not meet the stochastic RIP- $\ell_2/\ell_2$  condition (see, e.g., [3]) and requires the construction of equivalent sampling operators to satisfy the assumptions in [7, Theorem 5]. Secondly, unlike other low-rank matrix recovery problems [5, 10], they do not ensure recovery with an exponentially decreasing failure probability but with a polynomial decreasing failure probability. This is also due to the use of stochastic RIP- $\ell_2/\ell_2$ . For these two specific aspects, they conjecture in [7] that the results can be improved through other methods <sup>2</sup>.

**Conjecture.** Suppose sample matrices are  $\{\boldsymbol{\xi}_k \boldsymbol{\xi}_k^{\mathsf{T}}\}_{k=1}^m$ , where  $\boldsymbol{\xi}_k$  are given as independent copies of a random vector  $\boldsymbol{\xi}$ , whose entries  $\boldsymbol{\xi}$  are assumed to be i.i.d. subgaussian with moment condition (1). Then for all  $\mu = \mathbb{E}\boldsymbol{\xi}^4 > 1$  and with probability exceeding  $1 - e^{-cm}$ , we can recovery low-rank Toeplitz matrices through program (2) with near-optimal sample number, where *c* is a numerical absolute constant.

To resolve the aforementioned conjecture, we adopt descent cone analysis via Mendelson's small ball method, but confine it to the Toeplitz matrix space, abandoning the stochastic RIP- $\ell_2/\ell_2$  method. In the past decade, a number of works have studied low-rank matrix recovery [5], phase retrieval [10] and blind deconvolution [9] via the descent cone analysis. Descent cone analysis via Gordon's "escape through a mesh" theorem is suitable for i.i.d. Gaussian measurements [5], but it suffers from pitfalls when dealing with rank-one measurements. However, Mendelson's small ball method [8] is a powerful strategy for establishing a lower bound on a nonnegative empirical process that can address rank-one subgaussian measurements [10].

Two other approaches capable of handling rank-one measures are rank RIP- $\ell_2/\ell_1$  [2, 7] and dual certificate analysis [3]. However, the bottleneck of the former lies in establishing a precise upper bound on the covering number of all Toeplitz matrices with rank r, see e.g., [7]; while the latter resides in constructing dual elements, but the number of measurements is typically not optimal with respect to r, see e.g., [6]. The small ball method confine to the Toeplitz matrix space will be more effective than these two strategies. The following theorem resolves the conjecture above.

**Theorem 1.** Let  $p \ge 1$  and  $\|\boldsymbol{e}\|_{\ell_p} \le \eta$ . Suppose the sample matrices  $\{\boldsymbol{\xi}_k \boldsymbol{\xi}_k^{\mathsf{T}}\}_{k=1}^m$  satisfy  $\boldsymbol{\xi}_k$  are given as independent copies of a random vector  $\boldsymbol{\xi}$ , whose entries  $\boldsymbol{\xi}$  are assumed to be i.i.d. with subgaussian norm K and satisfy moment condition (1). For all  $\mu \ge 1$  with probability exceeding  $1 - e^{-cm}$ , the solution  $\hat{\boldsymbol{X}}$  to (2) satisfies

$$\left\|\widehat{\boldsymbol{X}} - \boldsymbol{X}_0\right\|_F \le C \frac{\eta}{m^{1/p}},\tag{3}$$

<sup>&</sup>lt;sup>2</sup>In [7, Remark 2], it states that "We conjecture that these two aspects can be improved via other proof techniques."

holds simultaneously for all symmetric Toeplitz matrices  $X_0$  of rank at most r, provided that  $m \ge Lr \log^2 n$ . Here, c, C and L are some universal constants dependent on K and  $\mu$ .

**Remark 1.** [7, Theorem 2] is applicable to models that include positive semi-definite cone constraints, as mentioned in [7], our result is also suitable for this scenario; our theorem is applicable to broader range of  $\ell_p$ -norm bounded noise.

Our theorem improves the analyses in [7] and gives positive answers to aforementioned conjecture. Firstly, our result applies to all subgaussian measurements that  $\mu \ge 1$ , not just for  $1 < \mu \le 3$ . Note that Bernoulli measurements are also applicable, which is different from the fact that in phase retrieval (when  $X_0$  is rank-1), it is impossible to distinguish between vector  $e_1$  and vector  $e_2$  from Bernoulli measurements. This is due to the fact that  $e_1e_1^T$  and  $e_2e_2^T$  correspond to the same Toeplitz matrix such that  $\mathcal{T}(e_1e_1^T) = \mathcal{T}(e_2e_2^T) = \frac{1}{n}I$ , where  $\mathcal{T}$  is orthogonal projection operator onto Toeplitz matrices. Aside from that, the result stabilizes with an exponentially decreasing failure probability. As  $m = \mathcal{O}(r \operatorname{poly}(n))$ , the established probability  $1 - e^{-cm}$  is about  $1 - \mathcal{O}(n^r)$ . Thus, exponentially decreasing failure probability is substantial. Besides, the sampling number in [7, Theorem 2] requires  $m = \mathcal{O}(r \log^{10} n)$ , which is optimal for rank constraint r but still far from the optimal sampling order. Our result improves the measurement number from  $m = \mathcal{O}(r \log^{10} n)$  to  $m = \mathcal{O}(r \log^2 n)$ . Furthermore, our result have the capability to address a broader range of  $\ell_p$ -norm bounded noise.

Our proof consists of three ingredients. The first ingredient is that the descent cone for the nuclear norm has clear characterization. The second is that the small ball function in the small ball method, when confined to the Toeplitz matrices, is bounded away from zero for rank-one subgaussian measurements. Finally, we provide an upper bound for the spectral norm of a random matrix with the Toeplitz matrix structure. Our approach relies on the feature of embedding Toeplitz matrices into circulant matrices, allowing for the explicit expression of the latter's eigenvalues, which diverges from the moment method [1] and the random process method [13] and may be of separate interest.

### 2 Proofs

We prove our results in this section. Before we begin, we introduce some notations. The symmetric Toeplitz matrix considered in this paper refers to the following form

$$\begin{pmatrix} z_0 & z_1 & \cdots & z_{n-1} \\ z_1 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & z_1 \\ z_{n-1} & \cdots & z_1 & z_0 \end{pmatrix}$$

We recall that we assume the sample vectors  $\{\xi_k\}_{k=1}^m$  in sample matrices  $\{\xi_k\xi_k^{\mathsf{T}}\}_{k=1}^m$  are composed of independent copies of random vectors  $\xi$ . Specifically, each entry of  $\xi$  is i.i.d.

drawn from a distribution with the following properties:

$$\mathbb{E}\xi = 0, \mathbb{E}\xi^2 = 1, \mathbb{E}\xi^4 = \mu \ge 1 \text{ and } \|\xi\|_{\psi_2} \le K$$

For a rank-*r* matrix X, we use  $||X||_*$  to denote its nuclear norm,  $||X||_F$  to denote its Frobenius norm and  $||X||_{op}$  to denote its spectral norm. We denote its corresponding singular value decomposition by  $X = U\Sigma V^*$ , where  $\Sigma \in \mathbb{R}^{r \times r}$  is a diagonal matrix with nonnegative entries and  $U, V \in \mathbb{R}^{n \times r}$  are unitary matrices. Then the tangent space of the manifold of rank-*r* matrices at X is defined by

$$T_{\boldsymbol{X}} := \left\{ \boldsymbol{U}\boldsymbol{A}^* + \boldsymbol{B}\boldsymbol{V}^* : \boldsymbol{A}, \boldsymbol{B} \in \mathbb{R}^{n \times r} \right\}.$$

We denote the orthogonal complement to  $T_X$  as  $T_X^{\perp}$ . In the following, we use  $Z_{T_X}$  to denote the orthogonal projection of Z onto  $T_{X_0}$  and  $Z_{T_{X_0}^{\perp}}$  to denote the orthogonal projection of Zonto  $T_{X_0}^{\perp}$ . In addition,  $\mathbb{S}_F$  denotes the Frobenius unit ball of  $\mathbb{R}^{n \times n}$ ,  $\mathcal{S}^n$  denotes the vector space of all symmetric matrices in  $\mathbb{R}^{n \times n}$ ,  $\mathcal{T}$  denotes the orthogonal projection operator onto Toeplitz matrices.

### 2.1 Preliminaries

We present the Hanson-Wright inequality, which gives a concentration bound for quadratic forms of random variables; for instance, see [13].

**Lemma 1.** Let  $\boldsymbol{\xi} = (\xi_1, \dots, \xi_n)$  be a random vector with independent components  $\xi$  which satisfy  $\mathbb{E}\xi = 0$ ,  $\mathbb{E}\xi^2 = 1$  and  $\|\xi\|_{\psi_2} \leq K$ . For any  $\boldsymbol{Z} \in S^n$ , there is a numerical constant c > 0 such that for all t > 0

$$\mathbb{P}\left(\left|\boldsymbol{\xi}^{\mathsf{T}}\boldsymbol{Z}\boldsymbol{\xi} - \mathbb{E}\left[\boldsymbol{\xi}^{\mathsf{T}}\boldsymbol{Z}\boldsymbol{\xi}\right]\right| > t\right) \le 2\exp\left(-c\min\left\{\frac{t^{2}}{K^{4}\left\|\boldsymbol{Z}\right\|_{F}^{2}}, \frac{t}{K^{2}\left\|\boldsymbol{Z}\right\|_{op}}\right\}\right).$$
(4)

We next present the Mendelson's small ball method, which can provide a lower bound for nonnegative empirical processes that can address a broad class of complexity measures and sampling matrices, especially for heavy tailed measurements. The proof can refer to [8, Theorem 2.1] or [12, Proposition 5.1].

**Lemma 2.** Fix  $\mathcal{K} \in \mathbb{R}^n$  and let  $\{\phi\}_{k=1}^m$  be independent copies of a random vector  $\phi$  in  $\mathbb{R}^n$ . Consider the small ball function

$$\mathcal{Q}_{\alpha}\left(\mathcal{K};\boldsymbol{\phi}\right) = \inf_{\boldsymbol{u}\in\mathcal{K}} \mathbb{P}\left(\left|\langle\boldsymbol{\phi},\boldsymbol{u}\rangle\right| \geq \alpha\right)$$
(5)

and suprema of empirical process

$$\mathcal{R}_{m}(\mathcal{K};\boldsymbol{\phi}) = \mathbb{E}\sup_{\boldsymbol{u}\in\mathcal{K}} \left| \frac{1}{m} \sum_{k=1}^{m} \varepsilon_{k} \langle \boldsymbol{\phi}, \boldsymbol{u} \rangle \right|,$$
(6)

where  $\{\varepsilon_k\}_{k=1}^m$  is a Rademacher sequence independent of everything else.

Then for any  $p \ge 1, \xi > 0$  and t > 0, with probability exceeding  $1 - \exp(-2t^2)$ ,

$$\inf_{\boldsymbol{u}\in\mathcal{K}}\left(\sum_{k=1}^{m}|\langle\boldsymbol{\phi},\boldsymbol{u}\rangle|^{p}\right)^{1/p}\geq m^{1/p}\left(\alpha\mathcal{Q}_{\alpha}\left(\mathcal{K};\boldsymbol{\phi}\right)-2\mathcal{R}_{m}\left(\mathcal{K};\boldsymbol{\phi}\right)-\frac{\alpha t}{\sqrt{m}}\right).$$
(7)

### 2.2 Descent Cone Analysis

In this section, we provide a framework for using the descent cone analysis method and a specific analysis of the small ball method confine to the Toeplitz matrix.

The descent cone of a norm at a point  $X_0 \in \mathbb{R}^{n \times n}$  is the set of all possible directions  $Z \in \mathbb{R}^{n \times n}$  such that the norm does not increase. For the nuclear norm, this leads to the following definition:

**Definition 1.** For any matrix  $X_0 \in \mathbb{R}^{n \times n}$  define its descent cone  $\mathcal{D}(X_0)$  by

$$\mathcal{D}\left(\boldsymbol{X}_{0}\right) := \left\{ \boldsymbol{Z} \in \mathbb{R}^{n \times n} : \left\| \boldsymbol{X}_{0} + t\boldsymbol{Z} \right\|_{*} \le \left\| \boldsymbol{X}_{0} \right\|_{*} \text{ for some } t > 0 \right\}.$$
(8)

The first crucial ingredient for Theorems 1 is the following proposition that characterizes the geometric property of the descent cone. The proof is inspired by [9, Lemma 4.1], and it can serves as a substitute for [10, Lemma 10].

**Proposition 1.** Let  $X_0 \in S^n$  be at most rank r. Then

$$\left\|\boldsymbol{Z}\right\|_{*} \leq \left(\sqrt{2} + 1\right) \sqrt{r} \left\|\boldsymbol{Z}\right\|_{F}, \text{ for all } \boldsymbol{Z} \in \overline{\mathcal{D}\left(\boldsymbol{X}_{0}\right)},$$
(9)

where  $\overline{\mathcal{D}(X_0)}$  denotes the topological closure of  $\mathcal{D}(X_0)$ .

*Proof.* We denote the SVD of  $X_0$  by  $X_0 = U\Sigma V^*$ . Then [9, Lemma 4.1] stated that

$$\overline{\mathcal{D}\left(\boldsymbol{X}_{0}\right)} = \left\{\boldsymbol{Z} \in \mathbb{R}^{n \times n} : -\langle \boldsymbol{U}\boldsymbol{V}^{*}, \boldsymbol{Z} \rangle \geq \left\|\boldsymbol{Z}_{T_{\boldsymbol{X}_{0}}^{\perp}}\right\|_{*}\right\}.$$
(10)

Thus, for any  $Z \in \overline{\mathcal{D}(X_0)}$ , we have

$$\begin{aligned} \|\boldsymbol{Z}\|_{*} &\leq \left\|\boldsymbol{Z}_{T_{\boldsymbol{X}_{0}}}\right\|_{*} + \left\|\boldsymbol{Z}_{T_{\boldsymbol{X}_{0}}}\right\|_{*} \\ &\leq \sqrt{2r} \left\|\boldsymbol{Z}_{T_{\boldsymbol{X}_{0}}}\right\|_{F} - \langle \boldsymbol{U}\boldsymbol{V}^{*}, \boldsymbol{Z} \rangle \\ &\leq \sqrt{2r} \left\|\boldsymbol{Z}_{T_{\boldsymbol{X}_{0}}}\right\|_{F} + \left\|\boldsymbol{U}\boldsymbol{V}^{*}\right\|_{F} \left\|\boldsymbol{Z}\right\|_{F} \\ &\leq \sqrt{2r} \left\|\boldsymbol{Z}\right\|_{F} + \left\|\boldsymbol{U}\right\|_{op} \left\|\boldsymbol{V}^{*}\right\|_{F} \left\|\boldsymbol{Z}\right\|_{F} \\ &\leq \left(\sqrt{2r} + \sqrt{r}\right) \left\|\boldsymbol{Z}\right\|_{F}. \end{aligned}$$

The second inequality deduced by  $Z_{T_{X_0}}$  is at most rank-2r and (10). The forth inequality follows from that for any U, V, we have  $\|UV^*\|_F \leq \|U\|_{op} \|V^*\|_F$ .

In the noiseless scenario, i.e.,  $\eta = 0$ , the matrix  $X_0$  is the unique minimizer to the semidefinite program (2) if and only if the null space of  $\mathcal{A}$  does not intersect the descent cone  $\mathcal{D}(X_0)$ . Then if the following quantity for a matrix  $X_0$  is far away 0, which is often referred to as minimum conic singular value, the intersection of  $\mathcal{A}$  and  $\mathcal{D}(X_0)$  is empty

$$\lambda_{\min}\left(\mathcal{A}; \mathcal{D}\left(\boldsymbol{X}_{0}
ight) \cap \mathbb{S}_{F}
ight) := \inf_{\boldsymbol{Z} \in \mathcal{D}\left(\boldsymbol{X}_{0}
ight) \setminus \left\{0
ight\} \cap \mathbb{S}_{F}} rac{\left\|\mathcal{A}\left(\boldsymbol{Z}
ight)
ight\|_{\ell_{p}}}{\left\|\boldsymbol{Z}
ight\|_{F}}.$$

For our specific situation, we only need to focus on the toeplitz matrices that fall within the descent cone  $\mathcal{D}(X_0)$ . Therefore, we only need to evaluate the minimum conic singular value within set  $\mathcal{D}(X_0) \cap \mathcal{T}$ . On the other hand, to ensure universal outcomes for all Toeplitz matrices with a rank at most r, the descent cone  $\mathcal{D}(X_0)$  can be extended to

$$\mathcal{K}_{\mathcal{T}}^{r} := \bigcup_{\boldsymbol{X}_{0}} \left( \mathcal{D}\left( \boldsymbol{X}_{0} \right) \cap \mathcal{T} \right), \tag{11}$$

where the union runs over all  $X_0 \in S^n \setminus \{0\}$  of rank at most r. Therefore, the minimum conic singular value this paper focus on is

$$\lambda_{\min}\left(\mathcal{A}; \mathcal{K}_{\mathcal{T}}^{r} \cap \mathbb{S}_{F}\right) := \inf_{\boldsymbol{Z} \in \mathcal{K}_{\mathcal{T}}^{r} \setminus \{0\} \cap \mathbb{S}_{F}} \frac{\left\|\mathcal{A}\left(\boldsymbol{Z}\right)\right\|_{\ell_{p}}}{\left\|\boldsymbol{Z}\right\|_{F}}.$$
(12)

We provide a concise description to prove Theorem 1 through  $\lambda_{\min} (\mathcal{A}; \mathcal{K}_{\mathcal{T}}^r \cap \mathbb{S}_F)$ . It follows by the triangle inequality that

$$\left\|\mathcal{A}\left(\boldsymbol{X}_{\star}-\boldsymbol{X}_{0}\right)\right\|_{\ell_{p}} \leq \left\|\mathcal{A}\left(\boldsymbol{X}_{\star}\right)-\boldsymbol{b}\right\|_{\ell_{p}}+\left\|\mathcal{A}\left(\boldsymbol{X}_{0}\right)-\boldsymbol{b}\right\|_{\ell_{p}} \leq 2\eta.$$

Thus, we have that

$$\left\|\boldsymbol{X}_{\star} - \boldsymbol{X}_{0}\right\|_{F} = \frac{\left\|\boldsymbol{\mathcal{A}}\left(\boldsymbol{X}_{\star} - \boldsymbol{X}_{0}\right)\right\|_{\ell_{p}}}{\left\|\boldsymbol{\mathcal{A}}\left(\boldsymbol{X}_{\star} - \boldsymbol{X}_{0}\right)\right\|_{\ell_{p}} / \left\|\boldsymbol{X}_{\star} - \boldsymbol{X}_{0}\right\|_{F}} \le \frac{2\eta}{\lambda_{\min}\left(\boldsymbol{\mathcal{A}}; \mathcal{K}_{\mathcal{T}}^{r} \cap \mathbb{S}_{F}\right)}.$$
(13)

The above (13) indicates that to prove Theorem 1 we need provide  $\lambda_{\min} (\mathcal{A}; \mathcal{K}_{\mathcal{T}}^r \cap \mathbb{S}_F)$  is bounded away from zero. Mendelson's small ball method can lead to the geometric analysis of  $\lambda_{\min} (\mathcal{A}; \mathcal{K}_{\mathcal{T}}^r \cap \mathbb{S}_F)$ . By Lemma 2, we have that

$$\lambda_{\min}\left(\mathcal{A};\mathcal{K}_{\mathcal{T}}^{r}\cap\mathbb{S}_{F}\right)\geq m^{1/p}\left(\alpha\mathcal{Q}_{\alpha}\left(\mathcal{K}_{\mathcal{T}}^{r};\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\right)-2\mathcal{R}_{m}\left(\mathcal{K}_{\mathcal{T}}^{r};\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\right)-\frac{\alpha t}{\sqrt{m}}\right).$$
(14)

Thus, we need a lower bound on small ball function  $\mathcal{Q}_{\alpha}\left(\mathcal{K}_{\mathcal{T}}^{r};\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\right)$  and an upper bound on suprema of empirical process  $\mathcal{R}_{m}\left(\mathcal{K}_{\mathcal{T}}^{r};\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\right)$ .

The second ingredient, is that when confined to the space of the Toeplitz matrix  $\mathcal{T}$  (it can be seen that  $\mathcal{K}^r_{\mathcal{T}} \subset \mathcal{T}$ ), the small ball function is bounded away from zero for al  $\mu \geq 1$ .

**Theorem 2.** Let  $\boldsymbol{\xi} = (\xi_1, \dots, \xi_n)$  be a random vector with i.i.d. entries  $\xi$  which satisfy  $\mathbb{E}\xi = 0$ ,  $\mathbb{E}\xi^2 = 1$ ,  $\mathbb{E}\xi^4 = \mu \ge 1$  and  $\|\xi\|_{\psi_2} \le K$ . Then it holds that

$$\mathcal{Q}_{\alpha}\left(\mathcal{T};\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\right) := \inf_{\boldsymbol{Z}\in\mathcal{T}\cap\mathbb{S}_{F}} \mathbb{P}\left(\left|\langle \boldsymbol{Z},\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\rangle\right| \geq \alpha\right)$$
$$\gtrsim (1-\alpha)^{2} \min\left\{\frac{4}{K^{8}}, \left(\frac{\mu}{1+K^{4}}\right)^{2}, 1\right\}.$$
(15)

*Proof.* It is useful to record two facts that shall be used. Fact 1: The first is that for any  $Z \in S^n$  it holds that

$$\mathbb{E} \left| \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} \right|^{2} = (\operatorname{Tr} \boldsymbol{Z})^{2} + \left( \mathbb{E} \boldsymbol{\xi}^{4} - 1 \right) \sum_{k=1}^{n} \boldsymbol{Z}_{i,i}^{2} + 2 \sum_{i \neq j}^{n} \boldsymbol{Z}_{i,j}^{2}.$$
(16)

By direct calculation,

$$\mathbb{E} \left| \boldsymbol{\xi}^* \boldsymbol{Z} \boldsymbol{\xi} \right|^2 = \mathbb{E} \left( \sum_{i,j} \xi_i \boldsymbol{Z}_{i,j} \xi_j \right)^2 = \mathbb{E} \sum_{i,\tilde{i},j,\tilde{i}} \xi_i \xi_{\tilde{i}} \xi_j \xi_{\tilde{j}} \boldsymbol{Z}_{i,j} \boldsymbol{Z}_{\tilde{i},\tilde{j}}$$
$$= \mathbb{E} \boldsymbol{\xi}^4 \cdot \sum_{i=1}^n \boldsymbol{Z}_{i,i}^2 + 2 \sum_{i,j} \boldsymbol{Z}_{i,j}^2 + \sum_{i,\tilde{i}} \boldsymbol{Z}_{i,i} \boldsymbol{Z}_{\tilde{i},\tilde{i}}$$
$$= (\operatorname{Tr} \boldsymbol{Z})^2 + (\mathbb{E} \boldsymbol{\xi}^4 - 1) \sum_{i=1}^n \boldsymbol{Z}_{i,i}^2 + 2 \sum_{i\neq j}^n \boldsymbol{Z}_{i,j}^2.$$

Fact 2: The second is that for any  $Z \in S^n$  it holds that

$$\mathbb{E}\left|\boldsymbol{\xi}^{\mathsf{T}}\boldsymbol{Z}\boldsymbol{\xi}\right|^{4} \lesssim (\operatorname{Tr}\boldsymbol{Z})^{4} + K^{8} \left\|\boldsymbol{Z}\right\|_{F}^{4}.$$
(17)

By Hanson-Wright inequality in Lemma 1, we have that

$$\begin{split} & \mathbb{E} \left| \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} - \mathbb{E} \left[ \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} \right] \right|^{4} \\ &= \int_{0}^{\infty} 4t^{3} \mathbb{P} \left( \left| \boldsymbol{\xi}^{*} \boldsymbol{Z} \boldsymbol{\xi} - \mathbb{E} \left[ \boldsymbol{\xi}^{*} \boldsymbol{Z} \boldsymbol{\xi} \right] \right| > t \right] dt \\ &\leq 8 \left( \int_{0}^{\infty} t^{3} \exp \left( -c \frac{t^{2}}{K^{4} \left\| \boldsymbol{Z} \right\|_{F}^{2}} \right) dt + \int_{0}^{\infty} t^{3} \exp \left( -c \frac{t}{K^{2} \left\| \boldsymbol{Z} \right\|_{op}} \right) dt \right) \\ &= 8 \left( K^{8} \left\| \boldsymbol{Z} \right\|_{F}^{4} \int_{0}^{\infty} x^{3} \exp \left( -cx^{2} \right) dx + K^{8} \left\| \boldsymbol{Z} \right\|_{op}^{4} \int_{0}^{\infty} x^{3} \exp \left( -cx \right) dx \right) \\ &\lesssim K^{8} \left\| \boldsymbol{Z} \right\|_{F}^{4}. \end{split}$$

Then the triangle inequality yields that

$$\begin{split} \mathbb{E} \left| \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} \right|^{4} &\lesssim \left| \mathbb{E} \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} \right|^{4} + \mathbb{E} \left| \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} - \mathbb{E} \left[ \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} \right] \right|^{4} \\ &= \left( \operatorname{Tr} \boldsymbol{Z} \right)^{4} + \mathbb{E} \left| \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} - \mathbb{E} \left[ \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} \right] \right|^{4} \\ &\lesssim \left( \operatorname{Tr} \boldsymbol{Z} \right)^{4} + K^{8} \left\| \boldsymbol{Z} \right\|_{F}^{4}. \end{split}$$

We now turn to proving (15). For  $Z \in \mathcal{T} \cap \mathbb{S}_F$ , let diagonal element of Z be  $z_0$ . Since  $\|Z\|_F = 1$ , we have  $0 \le z_0 \le \frac{1}{\sqrt{n}}$ . By (16), we have that

$$\mathbb{E} \left| \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} \right|^{2} = (\operatorname{Tr} \boldsymbol{Z})^{2} + \left( \mathbb{E} \boldsymbol{\xi}^{4} - 1 \right) \sum_{k=1}^{n} \boldsymbol{Z}_{i,i}^{2} + 2 \sum_{i \neq j}^{n} \boldsymbol{Z}_{i,j}^{2}$$
$$= (nz_{0})^{2} + (\mu - 1) nz_{0}^{2} + 2 \left( 1 - nz_{0}^{2} \right)$$
$$= \left[ n^{2} + (\mu - 3) n \right] z_{0}^{2} + 2.$$

Then it can be checked that  $\mathbb{E} \left| \boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi} \right|^{2} \geq 1 = \| \boldsymbol{Z} \|_{F}$ . Thus, by Paley-Zygmund inequality,

$$\mathbb{P}\left(\left|\boldsymbol{\xi}^{\mathsf{T}}\boldsymbol{Z}\boldsymbol{\xi}\right|^{2} \geq \alpha \left\|\boldsymbol{Z}\right\|_{F}^{2}\right) \geq \mathbb{P}\left(\left|\boldsymbol{\xi}^{\mathsf{T}}\boldsymbol{Z}\boldsymbol{\xi}\right|^{2} \geq \alpha \mathbb{E}\left|\boldsymbol{\xi}^{\mathsf{T}}\boldsymbol{Z}\boldsymbol{\xi}\right|^{2}\right) \geq (1-\alpha)^{2} \frac{\left(\mathbb{E}\left|\boldsymbol{\xi}^{\mathsf{T}}\boldsymbol{Z}\boldsymbol{\xi}\right|^{2}\right)^{2}}{\mathbb{E}\left|\boldsymbol{\xi}^{\mathsf{T}}\boldsymbol{Z}\boldsymbol{\xi}\right|^{4}}.$$

By (16) and (17), we have that

$$f(z_{0}) := \frac{\left(\mathbb{E} \left|\boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi}\right|^{2}\right)^{2}}{\mathbb{E} \left|\boldsymbol{\xi}^{\mathsf{T}} \boldsymbol{Z} \boldsymbol{\xi}\right|^{4}} \gtrsim \frac{\left[\left(\operatorname{Tr} \boldsymbol{Z}\right)^{2} + \left(\mathbb{E} \boldsymbol{\xi}^{4} - 1\right) \sum_{k=1}^{n} \boldsymbol{Z}_{i,i}^{2} + 2 \sum_{i \neq j}^{n} \boldsymbol{Z}_{i,j}^{2}\right]^{2}}{\left(\operatorname{Tr} \boldsymbol{Z}\right)^{4} + K^{8} \left\|\boldsymbol{Z}\right\|_{F}^{4}}$$
$$\geq \frac{\left[\left(n^{2} + (\mu - 3) n\right) z_{0}^{2} + 2\right]^{2}}{\left(nz_{0}\right)^{4} + K^{8}}$$
$$\geq 2 \left[\frac{\left(n^{2} + (\mu - 3) n\right) z_{0}^{2} + 2}{\left(nz_{0}\right)^{2} + K^{4}}\right]^{2} := g(z_{0}).$$

Since g is monotonic on interval  $\left[0, \frac{1}{\sqrt{n}}\right]$ , we have

$$\mathcal{Q}_{\alpha}\left(\mathcal{T};\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\right) \geq (1-\alpha)^{2}\min f\left(z_{0}\right) \gtrsim (1-\alpha)^{2}\min\left\{g\left(0\right), g\left(\frac{1}{\sqrt{n}}\right)\right\}$$
$$\geq 2\left(1-\alpha\right)^{2}\min\left\{\frac{4}{K^{8}}, \left(\frac{n+\mu-1}{n+K^{4}}\right)^{2}\right\}$$
$$\gtrsim (1-\alpha)^{2}\min\left\{\frac{4}{K^{8}}, \left(\frac{\mu}{1+K^{4}}\right)^{2}, 1\right\}.$$

#### 2.3 Structured Random Matrix

We investigate the spectral norm of the structured random matrix  $\sum_{k=1}^{m} \mathcal{T}(\boldsymbol{\xi}_k \boldsymbol{\xi}_k^*)$ , which constitutes our third ingredient. Our proof relies on embedding Toeplitz matrices into circulant matrices, which can be regarded as another method for characterizing the spectral method of random matrices. We provide the following theorem.

**Theorem 3.** Let  $\boldsymbol{\xi} = (\xi_1, \dots, \xi_n)$  be a random vector with i.i.d. entries  $\xi$  which satisfy  $\mathbb{E}\xi = 0$ ,  $\mathbb{E}\xi^2 = 1$  and  $\|\boldsymbol{\xi}\|_{\psi_2} \leq K$ . Assume that  $\{\boldsymbol{\xi}_k\}_{k=1}^m$  are i.i.d copies of  $\boldsymbol{\xi}$ , then we have

$$\mathbb{E}\left\|\sum_{k=1}^{m} \mathcal{T}\left(\boldsymbol{\xi}_{k} \boldsymbol{\xi}_{k}^{\mathsf{T}}\right) - m \mathbb{E} \mathcal{T}\left(\boldsymbol{\xi} \boldsymbol{\xi}^{\mathsf{T}}\right)\right\|_{op} \leq C K^{2} \left(\sqrt{m} \log n + \log^{3/2} n\right),$$
(18)

where C is a numerical constant.

*Proof.* We devide the proof into four steps. **Step 1: Expectation Calculation.** It can be seen that

$$\mathbb{E}\mathcal{T}\left(\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}
ight)=\mathcal{T}\left(\mathbb{E}\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}
ight)=\boldsymbol{I}_{n}\in\mathcal{T}.$$

We then set random matrix

$$\boldsymbol{Z}_{0} := \sum_{k=1}^{m} \mathcal{T}\left(\boldsymbol{\xi}_{k} \boldsymbol{\xi}_{k}^{\mathsf{T}}\right) - m \mathbb{E} \mathcal{T}\left(\boldsymbol{\xi} \boldsymbol{\xi}^{\mathsf{T}}\right) = \begin{pmatrix} z_{0} & z_{1} & \cdots & z_{n-1} \\ z_{1} & z_{0} & \cdots & z_{n-2} \\ \vdots & \vdots & \ddots & \vdots \\ z_{n-1} & z_{n-2} & \cdots & z_{0} \end{pmatrix},$$

and let  $z := (z_0, z_1, \dots, z_{n-1})$ . As the entry of each descending diagonal of a Toeplitz matrix is given by the average of the corresponding diagonal, we have that

$$\boldsymbol{z}_{\ell} = \begin{cases} \frac{1}{n} \sum_{k=1}^{m} \sum_{i=1}^{n} \left(\xi_{k,i}^{2} - 1\right), \ \ell = 0; \\ \frac{1}{n-\ell} \sum_{k=1}^{m} \sum_{i=1}^{n-\ell} \xi_{k,i} \xi_{k,i+\ell}, \ \ell = 1, 2, \cdots, n-1. \end{cases}$$

Apparently, one has  $\mathbb{E} \mathbf{z}_{\ell} = 0$  for all  $0 \leq \ell \leq n - 1$ .

Step 2: Toeplitz Matrix Embedding. The harmonic structure of the Toeplitz matrix motivates us to embed  $Z_0$  into a new circulant matrix  $C_{Z_0} \in \mathbb{R}^{(2n-1) \times (2n-1)}$  such that

$$\boldsymbol{C}_{\boldsymbol{Z}_{0}} = \begin{pmatrix} z_{0} & z_{1} & \cdots & z_{n-1} & z_{n-1} & \cdots & z_{1} \\ z_{1} & z_{0} & \cdots & z_{n-2} & z_{n-1} & \cdots & z_{2} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \cdots & \vdots \\ z_{n-1} & z_{n-2} & \cdots & z_{0} & z_{1} & \cdots & z_{n-1} \\ \hline z_{n-1} & z_{n-1} & \cdots & z_{1} & z_{0} & \cdots & z_{n-2} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ z_{1} & z_{2} & \cdots & z_{n-1} & z_{n-2} & \cdots & z_{0} \end{pmatrix}$$

Note that  $Z_0$  is a submatrix of  $C_{Z_0}$ , we only need to give an upper bound for  $\mathbb{E} \|C_{Z_0}\|_{op}$ .

Set  $\omega = e^{\frac{2\pi i}{2n-1}}$  to be the (2n-1)th root of unity. Then the corresponding eigenvalues of  $C_{Z_0}$  are given by

$$\lambda_{j} = z_{0} + z_{1}\omega^{j} + z_{2}\omega^{2j} + \dots + z_{n-1}\omega^{(n-1)j} + z_{n-1}\omega^{nj} + \dots + z_{1}\omega^{(2n-2)j}$$
  
=  $z_{0} + z_{1} \left(\omega^{j} + \omega^{(2n-2)j}\right) + \dots + z_{n-1} \left(\omega^{(n-1)j} + \omega^{nj}\right)$   
=  $z_{0} + 2\sum_{\ell=1}^{n-1} z_{\ell} \cos\left(\frac{2\pi j\ell}{2n-1}\right)$ , for all  $j = 0, 1, \dots, 2n-2$ .

It can be seen that  $\mathbb{E}\lambda_j = 0$  and

$$\|m{Z}_0\|_{op} \le \|m{C}_{m{Z}_0}\|_{op} = \max_{0 \le j \le 2n-2} |\lambda_j|$$

Step 3: Eigenvalue Estimation. Note that  $\lambda_j$  is a quadratic form in  $\{\xi_1, \dots, \xi_m\}$ . Let the symmetric coefficient matrix  $H^j$  be

$$\boldsymbol{H}_{\alpha,\beta}^{j} := \frac{1}{n - |\ell|} \cos\left(\frac{2\pi j |\ell|}{2n - 1}\right), \quad 1 \le \alpha, \beta \le n \text{ and } \alpha - \beta = \ell.$$
(19)

Thus,  $\lambda_i$  can be writeen as

$$\lambda_{j} = \begin{pmatrix} \boldsymbol{\xi}_{1}^{\mathsf{T}} & \boldsymbol{\xi}_{2}^{\mathsf{T}} & \cdots & \boldsymbol{\xi}_{m}^{\mathsf{T}} \end{pmatrix}_{1 \times mn} \begin{pmatrix} \boldsymbol{H}^{j} & & \\ & \boldsymbol{H}^{j} & \\ & & \ddots & \\ & & & \boldsymbol{H}^{j} \end{pmatrix}_{mn \times mn} \begin{pmatrix} \boldsymbol{\xi}_{1} \\ \boldsymbol{\xi}_{2} \\ \vdots \\ \boldsymbol{\xi}_{m} \end{pmatrix}_{mn \times 1} := \widetilde{\boldsymbol{\xi}}^{\mathsf{T}} \widetilde{\boldsymbol{H}^{j}} \widetilde{\boldsymbol{\xi}}.$$

Then by Hanson-Wright inequality in Lemma 1, we can get

$$\mathbb{P}\left(\left|\lambda_{j}\right| \geq t\right) \leq 2\exp\left(-c\min\left\{\frac{t^{2}}{K^{4}\left\|\widetilde{\boldsymbol{H}^{j}}\right\|_{F}^{2}}, \frac{t}{K^{2}\left\|\widetilde{\boldsymbol{H}^{j}}\right\|_{op}}\right\}\right)$$

It can be seen that  $\left\|\widetilde{H^{j}}\right\|_{op} = \left\|H^{j}\right\|_{op}$  and  $\left\|\widetilde{H^{j}}\right\|_{F} = \sqrt{m} \left\|H^{j}\right\|_{F}$ . We only estimate  $\left\|H^{j}\right\|_{F}$  as  $\left\|H^{j}\right\|_{op} \leq \left\|H^{j}\right\|_{F}$ . Since  $H^{j}$  is a symmetric Toeplitz matrix, we have

$$\|\boldsymbol{H}^{j}\|_{F}^{2} = \sum_{1 \le \alpha, \beta \le n} |\boldsymbol{H}_{\alpha,\beta}^{j}|^{2} \le \sum_{1 \le \alpha, \beta \le n; \alpha - \beta = |\ell|} \frac{1}{(n - |\ell|)^{2}}$$
$$= 2\sum_{\ell=1}^{n-1} \frac{1}{n - \ell} + \frac{1}{n} \le 2\log n + 2.$$

Therefore,

$$\left\|\widetilde{H}^{j}\right\|_{op} \leq \sqrt{2\log n + 2} \quad \text{and} \quad \left\|\widetilde{H}^{j}\right\|_{F} \leq \sqrt{m\left(2\log n + 2\right)}.$$
 (20)

Step 4: Uniform Argument. We define the event  $\Omega_{j,t}$  by  $|\lambda_j| \leq t$  and define  $\Omega_t = \bigcap_{0 \leq j \leq 2n-2} \Omega_{j,t}$ . Then

$$\mathbb{P}\left(\max_{0\leq j\leq 2n-2}|\lambda_{j}|\geq t\right) = \mathbb{P}\left(\Omega_{t}^{c}\right) \leq \sum_{0\leq j\leq 2n-2}\mathbb{P}\left(\Omega_{j,t}^{c}\right)$$
$$\leq \sum_{0\leq j\leq 2n-2}2\exp\left(-c\min\left\{\frac{t^{2}}{K^{4}\left\|\widetilde{\boldsymbol{H}^{j}}\right\|_{F}^{2}}, \frac{t}{K^{2}\left\|\widetilde{\boldsymbol{H}^{j}}\right\|_{op}}\right\}\right)$$
$$<4n\cdot\exp\left(-c\min\left\{\frac{t^{2}}{K^{4}m\left(2\log n+2\right)}, \frac{t}{K^{2}\sqrt{2\log n+2}}\right\}\right).$$

Finally, we have that

$$\mathbb{E} \| \mathbf{Z}_0 \|_{op} \leq \mathbb{E} \| \mathbf{C}_{\mathbf{Z}_0} \|_{op} = \mathbb{E} \max_{0 \leq j \leq 2n-2} |\lambda_j|$$

$$\leq \int_0^\infty \mathbb{P} \left( \max_{0 \leq j \leq 2n-2} |\lambda_j| \geq t \right) dt$$

$$\leq \int_0^{C_1 K^2 \sqrt{m} \log n} 1 dt + \int_{C_1 K^2 \sqrt{m} \log n}^\infty 4n \cdot \exp\left( -c \frac{t^2}{2K^4 m \log n} \right) dt \qquad (21)$$

$$+ \int_0^{C_2 K^2 \log^{3/2} n} 1 dt + \int_{C_2 K^2 \log^{3/2} n}^\infty 4n \cdot \exp\left( -c \frac{t}{K^2 \sqrt{2 \log n}} \right) dt$$

$$\leq \widetilde{C}_1 K^2 \sqrt{m} \log n + \widetilde{C}_2 K^2 \log^{3/2} n.$$

2.4 **Proof of Theorem 1** 

We now put all things together. By Theorem 2, the small ball function satisfies

$$\mathcal{Q}_{\alpha}\left(\mathcal{K}_{\mathcal{T}}^{r};\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\right) \geq \mathcal{Q}_{\alpha}\left(\mathcal{T};\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\right) \gtrsim (1-\alpha)^{2} \min\left\{\frac{4}{K^{8}}, \left(\frac{\mu}{1+K^{4}}\right)^{2}, 1\right\}.$$

Theorem 3 yields that

$$\mathbb{E} \left\| \frac{1}{m} \sum_{k=1}^{m} \varepsilon_{k} \mathcal{T}\left(\boldsymbol{\xi}_{k} \boldsymbol{\xi}_{k}^{\mathsf{T}}\right) \right\|_{op} \leq 2\mathbb{E} \left\| \frac{1}{m} \sum_{k=1}^{m} \mathcal{T}\left(\boldsymbol{\xi}_{k} \boldsymbol{\xi}_{k}^{\mathsf{T}}\right) - \mathbb{E} \mathcal{T}\left(\boldsymbol{\xi} \boldsymbol{\xi}^{\mathsf{T}}\right) \right\|_{op} \\ \leq CK^{2} \left( \frac{\log n}{\sqrt{m}} + \frac{\log^{3/2} n}{m} \right),$$

where in the first line we use the symmetrization principle [13, Lemma 6.4.2]. Thus, by Proposition 1, the suprema of empirical process

$$\mathcal{R}_{m}\left(\mathcal{K}_{\mathcal{T}}^{r};\boldsymbol{\xi}\boldsymbol{\xi}^{\mathsf{T}}\right) = \mathbb{E}\sup_{\boldsymbol{Z}\in\mathcal{K}_{\mathcal{T}}^{r}\cap\mathbb{S}_{F}} \left|\frac{1}{m}\sum_{k=1}^{m}\varepsilon_{k}\langle\boldsymbol{Z},\boldsymbol{\xi}_{k}\boldsymbol{\xi}_{k}^{\mathsf{T}}\rangle\right|$$
$$= \mathbb{E}\sup_{\boldsymbol{Z}\in\mathcal{K}_{\mathcal{T}}^{r}\cap\mathbb{S}_{F}} \left|\langle\boldsymbol{Z},\frac{1}{m}\sum_{k=1}^{m}\varepsilon_{k}\mathcal{T}\left(\boldsymbol{\xi}_{k}\boldsymbol{\xi}_{k}^{\mathsf{T}}\right)\rangle\right|$$
$$\leq \left(\sqrt{2}+1\right)\sqrt{r}\cdot\mathbb{E}\left\|\frac{1}{m}\sum_{k=1}^{m}\varepsilon_{k}\mathcal{T}\left(\boldsymbol{\xi}_{k}\boldsymbol{\xi}_{k}^{\mathsf{T}}\right)\right\|_{op}$$
$$\leq CK^{2}\left(\sqrt{2}+1\right)\sqrt{r}\cdot\left(\frac{\log n}{\sqrt{m}}+\frac{\log^{3/2} n}{m}\right)$$

Finally, we choose  $\alpha = c_1, t = c_2 \sqrt{m}$  in Lemma 2. By (14), provided  $m \ge Lr \log^2 n$  for sufficient large L, with probability exceeding  $1 - e^{-cm}$  we have that

$$\lambda_{\min}\left(\mathcal{A};\mathcal{K}_{\mathcal{T}}^{r}\cap\mathbb{S}_{F}\right)\gtrsim\tilde{c}m^{1/p}$$

Here  $c_1, c_2, c$  and  $\tilde{c}$  are numerical constants dependent on K and  $\mu$ . Thus by (13) we finish the proof.

## References

- [1] Afonso S Bandeira, March T Boedihardjo, and Ramon van Handel. Matrix concentration inequalities and free probability. *Inventiones mathematicae*, 234(1):419–487, 2023.
- [2] T Tony Cai and Anru Zhang. Rop: Matrix recovery via rank-one projections. *The Annals of Statistics*, 43(1):102–138, 2015.
- [3] Emmanuel J Candes, Thomas Strohmer, and Vladislav Voroninski. Phaselift: Exact and stable signal recovery from magnitude measurements via convex programming. *Communications on Pure and Applied Mathematics*, 66(8):1241–1274, 2013.
- [4] Raymond H Chan and Michael K Ng. Conjugate gradient methods for toeplitz systems. *SIAM Review*, 38(3):427–482, 1996.
- [5] Venkat Chandrasekaran, Benjamin Recht, Pablo A Parrilo, and Alan S Willsky. The convex geometry of linear inverse problems. *Foundations of Computational mathematics*, 12(6):805–849, 2012.
- [6] Yuxin Chen and Yuejie Chi. Robust spectral compressed sensing via structured matrix completion. *IEEE Transactions on Information Theory*, 60(10):6576–6601, 2014.

- [7] Yuxin Chen, Yuejie Chi, and Andrea J Goldsmith. Exact and stable covariance estimation from quadratic sampling via convex programming. *IEEE Transactions on Information Theory*, 61(7):4034–4059, 2015.
- [8] Vladimir Koltchinskii and Shahar Mendelson. Bounding the smallest singular value of a random matrix without concentration. *International Mathematics Research Notices*, 2015(23):12991–13008, 2015.
- [9] Felix Krahmer and Dominik Stöger. On the convex geometry of blind deconvolution and matrix completion. *Communications on Pure and Applied Mathematics*, 74(4):790–832, 2021.
- [10] Richard Kueng, Holger Rauhut, and Ulrich Terstiege. Low rank matrix recovery from rank one measurements. *Applied and Computational Harmonic Analysis*, 42(1):88– 116, 2017.
- [11] Mark Rudelson and Roman Vershynin. On sparse reconstruction from fourier and gaussian measurements. *Communications on Pure and Applied Mathematics*, 61(8):1025– 1045, 2008.
- [12] Joel A Tropp. Convex recovery of a structured signal from independent random linear measurements. *Sampling Theory, a Renaissance: Compressive Sensing and Other Developments*, pages 67–101, 2015.
- [13] Roman Vershynin. *High-Dimensional Probability: An Introduction with Applications in Data Science*, volume 47. Cambridge University Press, 2018.