

Entanglement Polygon Inequalities for A Class of Mixed States

Xian Shi^{1,*}

¹College of Information Science and Technology,
Beijing University of Chemical Technology, Beijing 100029, China

(Dated: July 8, 2024)

The study on the entanglement polygon inequality of multipartite systems has attracted much attention. However, most of the results are on pure states. Here we consider the property for a class of mixed states, which are the reduced density matrices of generalized W -class states in multipartite higher dimensional systems. First we show the class of mixed states satisfies the entanglement polygon inequalities in terms of Tsallis- q entanglement, then we propose a class of tighter inequalities for mixed states in terms of Tsallis- q entanglement. At last, we get an inequality for the mixed states, which can be regarded as a relation for bipartite entanglement.

PACS numbers: 03.65.Ud, 03.67.Mn

I. INTRODUCTION

Entanglement is an essential feature in quantum mechanics compared with the classical [1, 2]. It serves a key role in many information tasks, such as quantum teleportation [3, 4], quantum dense coding [5], and quantum cryptography [6, 7].

One of the most meaningful properties of entanglement in multipartite systems is that entanglement cannot be freely shared. Monogamy of entanglement (MoE) [8, 9] is one manifestation of the property. For a tripartite entangled state ρ_{ABC} , the mathematical representation of MoE in terms of an entanglement measure E is that

$$E_{A|B} + E_{A|C} \leq E_{A|BC}. \quad (1)$$

The above inequality means that the sum of entanglement in AB and AC is bounded by entanglement between subsystem A and the remaining BC . This relation is valid in terms of concurrence [8], the squashed entanglement measure [10], and Tsallis- q entanglement [11, 12] for n -qubit systems. Except (1), other representations of MoE have been proposed [13–17].

Recently, the other entanglement distribution property, entanglement polygon inequality (EPI), was proposed for pure states in multiqubit systems in terms of some entanglement measure E [18],

$$E_{j|\bar{j}} \leq \sum_{k \neq j} E_{k|\bar{k}},$$

here j denotes a subsystem \mathcal{H}_j , \bar{j} is the remaining subsystems. Recently, EPI was proved valid for pure states in multiqubit systems in terms of some entanglement measures [19, 20]. Based on the property, a method to build the genuine entanglement measure for multipartite systems was proposed [21]. However, there are few results on EPI for mixed states of multipartite systems in terms of some entanglement measure.

The generalized W class (GW) states of multipartite higher dimensional systems own fine properties on their entanglement distribution. In 2008, Kim and Sanders showed that the GW states in multipartite higher dimensional systems satisfy the monogamy relations in terms of the squared concurrence [22]. Moreover, there the authors showed that inequality (1) is saturated for the GW states in terms of the squared concurrence. In recent years, there have been many results on monogamy of entanglement for the GW states in terms of convex roof extended negativity [23], Tsallis- q entanglement [24], Renyi- α entanglement [25], and the unified- (q, s) entanglement [26].

This article is organized as follows. In section II, we present the preliminary knowledge of this article. In section III, we present our main results. First, we present a class of tripartite mixed states satisfy the EPI in terms of Tsallis- q entanglement, which are generated by the partial trace map over a generalized W -class state in terms of Tsallis- q entanglement, and then we show a tighter inequality for the mixed states. At last, we present another type of inequalities for the mixed states. In section IV, we end with a summary.

II. PRELIMINARY KNOWLEDGE

In this section, we present the preliminary knowledge needed here. First, we present the definitions and properties of entanglement measures for bipartite systems, and then we recall the generalized W -class states in multipartite higher dimensional systems.

A. Entanglement measures

Given a bipartite pure state $|\psi\rangle_{AB} = \sum_i \sqrt{\lambda_i} |ii\rangle$, the concurrence is defined as

$$C(|\psi\rangle_{AB}) = \sqrt{2(1 - \text{tr}\rho_A^2)}, \quad (2)$$

*shixian01@gmail.com

where $\rho_A = \text{tr}_B \rho_{AB}$. When ρ_{AB} is a mixed state, its concurrence is defined as

$$C(\rho_{AB}) = \min_{\{p_i, |\psi_i\rangle_{AB}\}} \sum_i p_i C(|\psi_i\rangle_{AB}), \quad (3)$$

where the minimum takes over all the decompositions of $\rho_{AB} = \sum_i p_i |\psi_i\rangle_{AB} \langle \psi_i|$.

For a quantum state ρ , the Tsallis- q entropy is

$$T_q(\rho) = \frac{1}{q-1} [1 - \text{tr} \rho^q],$$

for $q \geq 1$. When $q \rightarrow 1$, $T_q(\cdot)$ converges to the von Neumann entropy $S(\cdot)$

$$\lim_{q \rightarrow 1} T_q(\rho) = -\text{tr} \rho \log \rho = S(\rho). \quad (4)$$

For a bipartite pure state $|\psi\rangle_{AB}$, when $q \geq 1$, its Tsallis- q entanglement is

$$\mathbf{T}_q(|\psi\rangle_{AB}) = T_q(\rho_A),$$

where $\rho_A = \text{tr}_B |\psi\rangle_{AB} \langle \psi|$. For a mixed state ρ_{AB} , its Tsallis- q entanglement is defined as

$$\mathbf{T}_q(\rho_{AB}) = \min \sum_i p_i \mathbf{T}_q(|\psi_i\rangle_{AB}),$$

where the minimum takes over all possible pure state decompositions of $\rho_{AB} = \sum_i p_i |\psi_i\rangle_{AB} \langle \psi_i|$.

By combing (4), we have

$$\lim_{q \rightarrow 1} \mathbf{T}_q(\rho_{AB}) = E(\rho_{AB}),$$

where $E(\rho_{AB})$ is entanglement of formation for the mixed state ρ_{AB} [27, 28].

Next assume $|\phi\rangle$ is a pure state in $\mathcal{H}_2 \otimes \mathcal{H}_d$ with its Schmidt decomposition $|\phi\rangle_{AB} = \sqrt{\lambda}|00\rangle + \sqrt{1-\lambda}|11\rangle$, its Tsallis- q entanglement is

$$\mathbf{T}_q(|\phi\rangle_{AB}) = \frac{1}{(1-q)} [\lambda^q + (1-\lambda)^q - 1].$$

Combing the definition of concurrence for pure states, we have

$$\mathbf{T}_q(|\phi\rangle_{AB}) = f_q(C^2(|\phi_{AB}\rangle)),$$

where $f_q(x)$ is defined in $x \in [0, 1]$,

$$f_q(x) = \frac{((1 + \sqrt{1-x})^q + (1 - \sqrt{1-x})^q) - 2^q}{(1-q)2^q}.$$

Next we recall several lemmas on the properties of $f_q(x)$.

Lemma 1 [11] *The function $f_q(x)$ is a monotonously increasing and concave function when $q \in [\frac{5-\sqrt{13}}{2}, 2] \cup [3, \frac{5+\sqrt{13}}{2}]$.*

Lemma 2 [12] *The function $f_q(x^2)$ is a monotonically increasing function of the variable x for $q \in (0, \infty)$ and $x \in (0, 1)$, it is a convex function of x when $q \in [\frac{5-\sqrt{13}}{2}, \frac{5+\sqrt{13}}{2}]$.*

B. Generalized W-class State in Multipartite Higher Dimensional Systems

Here we first recall the generalized W-class (GW) state $|W_n^d\rangle_{A_1 A_2 \dots A_n}$ in multi-qudit systems [22],

$$\begin{aligned} & |W_n^d\rangle_{A_1 A_2 \dots A_n} \\ &= \sum_{j=1}^{d-1} (a_{1j}|j0 \dots 0\rangle + a_{2j}|0j \dots 0\rangle + \dots + a_{nj}|00 \dots 0j\rangle), \end{aligned} \quad (5)$$

here $\{|j\rangle_{A_i}\}_{j=0}^{d-1}$ is an orthonormal basis of qudit subsystems A_i with $i = 1, 2, \dots, n$ and $\sum_{i=1}^n \sum_{j=1}^{d-1} |a_{ij}|^2 = 1$.

Lemma 3 [22] *For any n -qudit generalized W-class state $|\psi\rangle_{AB_1 \dots B_{n-1}}$ in (5) and a partition $P = \{P_1, \dots, P_m\}$ for the subset $S = \{A, B_1, \dots, B_{l-1}\}$, $m \leq l \leq n$,*

$$C_{P_s | \overline{P_s}}^2 = \sum_{k \neq s} C_{P_s | P_k}^2 = \sum_{k \neq s} (C_{P_s | P_k}^a)^2, \quad (6)$$

for all $k \neq s$, here we denote $\{P_1, \dots, \overline{P_s}, \dots, P_m\} = \{P_1, \dots, P_s, \dots, P_m\} - \{P_s\}$

Based on Lemma 1 and the relations between $\mathbf{T}_q(\cdot)$ and $C(\cdot)$, the authors in [24] showed that

Lemma 4 *Assume $\rho_{A_{j_1} A_{j_2} \dots A_{j_m}}$ is a reduced density matrix of an n -qudit GW state in (5), here $m \leq n$, then we have*

$$\mathbf{T}_q(\rho_{A_{j_1} | A_{j_2} \dots A_{j_m}}) = f_q(C^2(\rho_{A_{j_1} | A_{j_2} \dots A_{j_m}})),$$

for $q \in [\frac{5-\sqrt{13}}{2}, \frac{5+\sqrt{13}}{2}]$.

III. MAIN RESULTS

This section mainly considers classes of entanglement polygon inequalities in terms of the Tsallis- q entanglement for a class of mixed states, which are the reduced density matrices of a GW state. First, we show that the EPI is satisfied by the class of mixed states. Then a tighter EPI is presented for the class of mixed states. At last, we present a generalized EPI for the mixed states, this inequality can be regarded as a relationship of a bipartite entanglement.

Theorem 1 *Assume $\rho_{A_{j_1} A_{j_2} A_{j_3}}$ is a tripartite reduced density matrix of an n -qudit GW state in $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2} \otimes \dots \mathcal{H}_{A_n}$, then*

$$\mathbf{T}_q(\rho_{A_{j_1} | A_{j_2} A_{j_3}}) \leq \mathbf{T}_q(\rho_{A_{j_2} | A_{j_1} A_{j_3}}) + \mathbf{T}_q(\rho_{A_{j_3} | A_{j_1} A_{j_2}}),$$

when $q \in [\frac{5-\sqrt{13}}{2}, 2] \cup [3, \frac{5+\sqrt{13}}{2}]$.




FIG. 1: Characterization of Corollary 1. In this figure, the length of segment $i - j$ indicates the value of $\mathbf{T}_q(\rho_{A_k|A_i A_j})$, here k is another party different from parties i and j .

Proof. As $\rho_{A_{j_1} A_{j_2} A_{j_3}}$ is a tripartite reduced density matrix of a GW state, then

$$\begin{aligned} & \mathbf{T}_q(\rho_{A_{j_2}|A_{j_1} A_{j_3}}) + \mathbf{T}_q(\rho_{A_{j_3}|A_{j_1} A_{j_2}}) \\ &= f_q(C_{A_{j_2}|A_{j_1} A_{j_3}}^2) + f_q(C_{A_{j_3}|A_{j_2} A_{j_1}}^2) \\ &= f_q(C_{A_{j_2} A_{j_1}}^2 + C_{A_{j_2} A_{j_3}}^2) + f_q(C_{A_{j_3} A_{j_2}}^2 + C_{A_{j_3} A_{j_1}}^2) \\ &\geq f_q(C_{A_{j_1} A_{j_2}}^2 + C_{A_{j_1} A_{j_3}}^2) \\ &= f_q(C_{A_{j_1}|A_{j_2} A_{j_3}}^2) \\ &= \mathbf{T}_q(\rho_{A_{j_1}|A_{j_2} A_{j_3}}), \end{aligned}$$

here the first equality is due to the Lemma 4, the second equality is due to Lemma 3, the first inequality is due to the Lemma 1, and the last equality is due to Lemma 3. \square

Due to the proof of Theorem 1, we can prove the following similarly,

$$\mathbf{T}_q(\rho_{A_{j_2}|A_{j_1} A_{j_3}}) \leq \mathbf{T}_q(\rho_{A_{j_1}|A_{j_2} A_{j_3}}) + \mathbf{T}(\rho_{A_{j_3}|A_{j_1} A_{j_2}}),$$

that is,

Corollary 1 Assume $\rho_{A_{j_1} A_{j_2} A_{j_3}}$ is a tripartite reduced density matrix of an n -qudit GW state in $\mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2} \otimes \cdots \mathcal{H}_{A_n}$, then

$$\begin{aligned} & |\mathbf{T}_q(\rho_{A_{j_2}|A_{j_1} A_{j_3}}) - \mathbf{T}_q(\rho_{A_{j_3}|A_{j_1} A_{j_2}})| \leq \\ & \mathbf{T}_q(\rho_{A_{j_1}|A_{j_2} A_{j_3}}) \leq \mathbf{T}_q(\rho_{A_{j_2}|A_{j_1} A_{j_3}}) + \mathbf{T}(\rho_{A_{j_3}|A_{j_1} A_{j_2}}), \end{aligned}$$

when $q \in [\frac{5-\sqrt{13}}{2}, 2] \cup [3, \frac{5+\sqrt{13}}{2}]$.

In Fig. 1, we plot a triangle to present the meaning of Corollary 1, the length of segment $i - j$ denotes the entanglement of $\mathbf{T}_q(\rho_{A_k|A_i A_j})$, here k is another party.

Corollary 2 Assume $\rho_{A_{j_1} A_{j_2} \cdots A_{j_m}}$ is a multipartite reduced density matrix of an n -qudit GW state in $\mathcal{H}_{A_1} \otimes$

$\mathcal{H}_{A_2} \otimes \cdots \otimes \mathcal{H}_{A_n}$, and $\{P_1, P_2, \cdots, P_h\}$ is a partition of the set $S = \{A_{j_1}, A_{j_2}, \cdots, A_{j_m}\}$, that is, $\cup_{g=1}^h P_g = S$,

$$\mathbf{T}_q(\rho_{P_{j_k}|\overline{P_{j_k}}}) \leq \sum_{l \neq k} \mathbf{T}_q(\rho_{P_{j_l}|\overline{P_{j_l}}}),$$

here $\overline{P_{j_i}} = P_{j_0} P_{j_1} \cdots P_{j_{i-1}} P_{j_{i+1}} \cdots P_{j_m}$, $i = 0, 1, \cdots, m$, and $m \leq n$.

Assume $j \in \mathbb{N}^+$, then j can be written as

$$j = \sum_{i=0}^{n-1} j_i 2^i, \quad (7)$$

here we assume $\log_2 j \leq n$, $j_i \in \{0, 1\}$. According to the equality (7), we have the following bijection:

$$\begin{aligned} j &\rightarrow \vec{j} \\ j &\rightarrow (j_0, j_1, \cdots, j_{n-1}), \end{aligned}$$

and we denote its Hamming distance $w_H(\vec{j})$ as the number 1 of the set $\{j_0, j_1, \cdots, j_{n-1}\}$. Next we present a tighter inequality of the GW states in terms of the Tsallis- q entanglement.

Theorem 2 Let $\beta \in [0, 1]$, assume $\rho_{A_{i_1} A_{i_2} \cdots A_{i_m}}$ is a reduced density matrix of a GW state $|\psi\rangle_{A_0 A_1 \cdots A_{n-1}}$. Let $P = \{P_{j_0}, P_{j_1}, \cdots, P_{j_{m-1}}, P_{j_m}\}$ be a partition of the set $\{A_{i_1}, A_{i_2}, \cdots, A_{i_m}\}$, when $q \in [\frac{5-\sqrt{13}}{2}, 2] \cup [3, \frac{5+\sqrt{13}}{2}]$, there exists an appropriate order of $P_{j_0}, P_{j_1}, \cdots, P_{j_{m-1}}, P_{j_m}$ such that

$$[\mathbf{T}_q(\rho_{P_{j_0}|\overline{P_{j_0}}})]^\beta \leq \sum_{i=1}^m (2^\beta - 1)^{w_H(\vec{j}_i)} [\mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}})]^\beta, \quad (8)$$

here $\overline{P_{j_i}} = P_{j_0} P_{j_1} \cdots P_{j_{i-1}} P_{j_{i+1}} \cdots P_{j_m}$, $i = 0, 1, \cdots, m$.

Here we place the proof of Theorem 2 in Sec. VI.

At last, we present the other class of entanglement polygon inequalities for bipartite entanglement. Assume $|\phi\rangle_{\mathbf{A}\mathbf{B}\mathbf{C}}$ is a GW state in $\mathcal{H}_{\mathbf{A}} \otimes \mathcal{H}_{\mathbf{B}} \otimes \mathcal{H}_{\mathbf{C}}$, $\rho_{\mathbf{A}\mathbf{B}} = \text{tr}_{\mathbf{C}} |\phi\rangle_{\mathbf{A}\mathbf{B}\mathbf{C}} \langle \phi|$ is a mixed state of the Hilbert space $\mathcal{H}_{\mathbf{A}\mathbf{B}}$. Here we denote $\mathbf{A} = A_1 A_2 \cdots A_m$ and $\mathbf{B} = B_1 B_2 \cdots B_m$.

Theorem 3 Assume $|\phi\rangle_{\mathbf{A}\mathbf{B}\mathbf{C}}$ is a GW state in $\mathcal{H}_{\mathbf{A}\mathbf{B}\mathbf{C}}$, $\rho_{\mathbf{A}\mathbf{B}} = \text{tr}_{\mathbf{C}} |\phi\rangle_{\mathbf{A}\mathbf{B}\mathbf{C}} \langle \phi|$, here $\mathbf{A} = A_1 A_2 \cdots A_m$ and $\mathbf{B} = B_1 B_2 \cdots B_m$, then

$$\mathbf{T}_q^{\mathbf{A}|\mathbf{B}} \leq \sum_{i,j=1}^m \mathbf{T}_q^{A_i|B_j}, \quad (9)$$

when $q \in [\frac{5-\sqrt{13}}{2}, 2] \cup [3, \frac{5+\sqrt{13}}{2}]$.

Proof. As $\rho_{\mathbf{AB}}$ is reduced density matrix of a GW state, then

$$\begin{aligned} \mathbf{T}_q(\rho_{\mathbf{AB}}) &= f_q(C_{\mathbf{A}|\mathbf{B}}^2) \\ &= f_q\left(\sum_{i,j=1}^m C_{A_i B_j}^2\right) \\ &\leq \sum_{i,j=1}^m f_q(C_{A_i B_j}^2) \\ &= \sum_{i,j=1}^m T_q(\rho_{A_i B_j}). \end{aligned}$$

□

IV. SUMMARY

In this manuscript, we have mainly considered the distribution property of multipartite entanglement for a

class of mixed states in terms of the Tsallis- q entanglement. First, we have proved the EPI in terms of Tsallis- q entanglement for the reduced density matrices of the GW states. With Hamming distance, we have presented a tighter EPI in terms of Tsallis- q entanglement for the mixed states. At last, we have shown inequalities for the mixed states in terms of Tsallis- q entanglement. We hope our results can provide a reference for future work on the study of multipartite quantum entanglement.

V. ACKNOWLEDGEMENT

This work was supported by the National Natural Science Foundation of China (Grant No. 12301580).

-
- [1] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, “Quantum entanglement,” *Reviews of modern physics*, vol. 81, no. 2, pp. 865–942, 2009.
- [2] M. B. Plenio and S. S. Virmani, “An introduction to entanglement theory,” in *Quantum information and coherence*. Springer, 2014, pp. 173–209.
- [3] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, “Teleporting an unknown quantum state via dual classical and einstein-podolsky-rosen channels,” *Physical review letters*, vol. 70, no. 13, p. 1895, 1993.
- [4] D. Bouwmeester, J.-W. Pan, K. Mattle, M. Eibl, H. Weinfurter, and A. Zeilinger, “Experimental quantum teleportation,” *Nature*, vol. 390, no. 6660, pp. 575–579, 1997.
- [5] A. Harrow, P. Hayden, and D. Leung, “Superdense coding of quantum states,” *Physical review letters*, vol. 92, no. 18, p. 187901, 2004.
- [6] A. K. Ekert, “Quantum cryptography based on bell’s theorem,” *Physical review letters*, vol. 67, no. 6, p. 661, 1991.
- [7] M. Mirhosseini, O. S. Magana-Loaiza, M. N. O Sullivan, B. Rodenburg, M. Malik, M. P. Lavery, M. J. Padgett, D. J. Gauthier, and R. W. Boyd, “High-dimensional quantum cryptography with twisted light,” *New Journal of Physics*, vol. 17, no. 3, p. 033033, 2015.
- [8] V. Coffman, J. Kundu, and W. K. Wootters, “Distributed entanglement,” *Physical Review A*, vol. 61, no. 5, p. 052306, 2000.
- [9] X.-L. Zong, H.-H. Yin, W. Song, and Z.-L. Cao, “Monogamy of quantum entanglement,” *Frontiers in Physics*, vol. 10, p. 880560, 2022.
- [10] M. Christandl and A. Winter, ““squashed entanglement”: an additive entanglement measure,” *Journal of mathematical physics*, vol. 45, no. 3, pp. 829–840, 2004.
- [11] J. S. Kim, “Tsallis entropy and entanglement constraints in multiqubit systems,” *Physical Review A*, vol. 81, no. 6, p. 062328, 2010.
- [12] Y. Luo, T. Tian, L.-H. Shao, and Y. Li, “General monogamy of tsallis q -entropy entanglement in multi-qubit systems,” *Physical Review A*, vol. 93, no. 6, p. 062340, 2016.
- [13] G. Gour and Y. Guo, “Monogamy of entanglement without inequalities,” *Quantum*, vol. 2, p. 81, 2018.
- [14] Y. Guo and L. Zhang, “Multipartite entanglement measure and complete monogamy relation,” *Physical Review A*, vol. 101, no. 3, p. 032301, 2020.
- [15] X. Shi, L. Chen, and M. Hu, “Multilinear monogamy relations for multiqubit states,” *Physical Review A*, vol. 104, no. 1, p. 012426, 2021.
- [16] Z.-X. Jin, S.-M. Fei, X. Li-Jost, and C.-F. Qiao, “A new parameterized monogamy relation between entanglement and equality,” *Advanced Quantum Technologies*, vol. 5, no. 6, p. 2100148, 2022.
- [17] Y. Guo and L. Huang, “Complete monogamy of multipartite quantum mutual information,” *Physical Review A*, vol. 107, no. 4, p. 042409, 2023.
- [18] X.-F. Qian, M. A. Alonso, and J. H. Eberly, “Entanglement polygon inequality in qubit systems,” *New Journal of Physics*, vol. 20, no. 6, p. 063012, 2018.
- [19] X. Yang, Y.-H. Yang, and M.-X. Luo, “Entanglement polygon inequality in qudit systems,” *Physical Review A*, vol. 105, no. 6, p. 062402, 2022.
- [20] X. Shi, “Entanglement polygon inequalities for pure states in qudit systems,” *The European Physical Journal Plus*, vol. 138, no. 8, pp. 1–10, 2023.
- [21] S. Xie and J. H. Eberly, “Triangle measure of tripartite entanglement,” *Physical Review Letters*, vol. 127, p. 040403, Jul 2021.
- [22] J. San Kim and B. C. Sanders, “Generalized w -class state and its monogamy relation,” *Journal of Physics A: Mathematical and Theoretical*, vol. 41, no. 49, p. 495301, 2008.
- [23] J. H. Choi and J. S. Kim, “Negativity and strong monogamy of multipartite quantum entanglement beyond qubits,” *Phys. Rev. A*, vol. 92, p. 042307, Oct 2015.
- [24] X. Shi and L. Chen, “Monogamy relations for the gener-

alized w-class states beyond qubits,” *Physical Review A*, vol. 101, no. 3, p. 032344, 2020.

- [25] L.-M. Lai, S.-M. Fei, and Z.-X. Wang, “Tighter monogamy and polygamy relations for a superposition of the generalized w-class state and vacuum,” *Journal of Physics A: Mathematical and Theoretical*, vol. 54, no. 42, p. 425301, 2021.
- [26] B. Li, B. Xie, Z. Zhang, and H. Fan, “Monogamy and polygamy for the generalized w-class states using unified-(q, s) entropy,” *Science China Physics, Mechanics & Astronomy*, vol. 67, no. 1, p. 210312, 2024.
- [27] C. H. Bennett, H. J. Bernstein, S. Popescu, and B. Schumacher, “Concentrating partial entanglement by local operations,” *Phys. Rev. A*, vol. 53, pp. 2046–2052, Apr 1996.
- [28] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, “Mixed-state entanglement and quantum error correction,” *Phys. Rev. A*, vol. 54, pp. 3824–3851, Nov 1996.

VI. APPENDIX

Lemma 5 *Let $\beta \in [0, 1], x \in (0, 1]$, then we have*

$$(1+x)^\beta \leq 1 + (2^\beta - 1)x^\beta. \quad (10)$$

Proof. Let $t = \frac{1}{x}$, then the lemma is equivalent to get the maximum of $f(t)$ when $t \in [1, \infty)$,

$$f(t) = (1+t)^\beta - t^\beta. \quad (11)$$

As $t \in [1, \infty)$, and $f'(t) \leq 0$, that is, when $t = 1$, $f(t)$ get the maximum $2^\beta - 1$. At last, When we replace t with $\frac{1}{x}$, we finish the proof. \square

Theorem 2: *Let $\beta \in [0, 1]$, assume $\rho_{A_{i_1} A_{i_2} \dots A_{i_m}}$ is a reduced density matrix of a GW state $|\psi\rangle_{A_0 A_1 \dots A_{n-1}}$, let $P = \{P_{j_0}, P_{j_1}, \dots, P_{j_{m-1}}, P_{j_m}\}$ be a partition of the set $\{A_{i_1}, A_{i_2}, \dots, A_{i_m}\}$, then there exists an appropriate order of $P_{j_0}, P_{j_1}, \dots, P_{j_{m-1}}, P_{j_m}$ such that*

$$[\mathbf{T}_q(\rho_{P_{j_0}|\overline{P_{j_0}}})]^\beta \leq \sum_{i=1}^m (2^\beta - 1)^{w_H(\vec{j}_i)} [\mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}})]^\beta, \quad (12)$$

here $\overline{P_{j_i}} = P_{j_0} P_{j_1} \dots P_{j_{i-1}} P_{j_{i+1}} \dots P_{j_m}$, $i = 0, 1, \dots, m$.

Proof. In the process of the proof, we always can order the partite $P_{j_0}, P_{j_1}, \dots, P_{j_{m-1}}$ such that

$$\mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}}) \geq \mathbf{T}_q(\rho_{P_{j_{i+1}}|\overline{P_{j_{i+1}}}}), i = 0, 1, \dots, m-1. \quad (13)$$

Through computation, we have that all the reduced density matrices of a GW state is can be written as

$$\rho_{P_{j_0} \dots P_{j_{m-1}}} = \gamma_{P_{j_0} \dots P_{j_{k-1}}} \otimes |0_{m-k}\rangle \langle 0_{m-k}|$$

Here we will use the mathematical induction method to prove the theorem. When $\rho_{P_{j_0} P_{j_1} P_{j_2}}$ is a tripartite

reduced density matrix of a GW state $|\psi\rangle_{AA_1 \dots A_{n-1}}$,

$$\begin{aligned} & [\mathbf{T}_q(\rho_{P_{j_0}|P_{j_1} P_{j_2}})]^\beta \\ & \leq [\mathbf{T}_q(\rho_{P_{j_1}|P_{j_0} P_{j_2}}) + \mathbf{T}_q(\rho_{P_{j_2}|P_{j_0} P_{j_1}})]^\beta \\ & = [\mathbf{T}_q(\rho_{P_{j_1}|P_{j_0} P_{j_2}})]^\beta \left[1 + \left[\frac{\mathbf{T}_q(\rho_{P_{j_2}|P_{j_0} P_{j_1}})}{\mathbf{T}_q(\rho_{P_{j_1}|P_{j_0} P_{j_2}})} \right] \right]^\beta \\ & \leq (\mathbf{T}_q(\rho_{P_{j_1}|P_{j_0} P_{j_2}}))^\beta + (2^\beta - 1)(\mathbf{T}_q(\rho_{P_{j_2}|P_{j_0} P_{j_1}}))^\beta, \quad (14) \end{aligned}$$

here the first inequality is due to the Theorem 1, and when $a > c > 0, b > 0, a^b > c^b$, and the second inequality is due to the Lemma 5.

Next assume $m < 2^n$, the theorem is correct. Then when $m = 2^n$, from the inequality (14), we have

$$\begin{aligned} & [\mathbf{T}_q(\rho_{P_{j_0}|\overline{P_{j_0}}})]^\beta \\ & \leq \left[\sum_{i=1}^{m/2-1} \mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}}) \right]^\beta \left[1 + \frac{\sum_{i=m/2}^{m-1} \mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}})}{\sum_{i=0}^{m/2-1} \mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}})} \right]^\beta \\ & \leq \left[\sum_{i=1}^{m/2-1} \mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}}) \right]^\beta + (2^\beta - 1) \left[\sum_{i=m/2}^{m-1} \mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}}) \right]^\beta \\ & \leq \sum_{i=1}^{m/2-1} (2^\beta - 1)^{w_H(\vec{j}_i)} (\mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}}))^\beta \\ & \quad + \sum_{i=m/2}^{m-1} (2^\beta - 1) \times (2^\beta - 1)^{w_H(\vec{j}_i)-1} [\mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}})]^\beta \\ & \leq \sum_{i=1}^{m-1} (2^\beta - 1)^{w_H(\vec{j}_i)} [\mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}})]^\beta. \quad (15) \end{aligned}$$

When m is an arbitrary number, we always can choose an $n \in \mathbb{N}^+$ such that $2^{n-1} \leq m \leq 2^n$. Let

$$\gamma_{P_{j_0} \dots P_{j_{2^n-1}}} = \rho_{P_{j_0} P_{j_1} \dots P_{j_{m-1}}} \otimes \delta_{P_{j_m} P_{j_{m+1}} \dots P_{j_{2^n-1}}},$$

here

$$\delta_{P_{j_m} P_{j_{m+1}} \dots P_{j_{2^n-1}}} = |00 \dots 0\rangle_{P_{j_m} P_{j_{m+1}} \dots P_{j_{2^n-1}}} \langle 00 \dots 0|.$$

Then due to the inequality (15), we have

$$\left[\sum_{i=0}^{2^n-1} \mathbf{T}_q(\gamma_{P_{j_i}|\overline{P_{j_i}}}) \right]^\beta \leq \sum_{i=0}^{2^n-1} (2^\beta - 1)^{w_H(\vec{j}_i)} [\mathbf{T}_q(\gamma_{P_{j_i}|\overline{P_{j_i}}})]^\beta. \quad (16)$$

From the definition of the state $\gamma_{P_{j_0} \dots P_{j_{2^n-1}}}$, we have

$$\mathbf{T}_q(\gamma_{P_{j_i}|\overline{P_{j_i}}}) = \mathbf{T}_q(\rho_{P_{j_i}|\overline{P_{j_i}}}), \quad i = 1, \dots, m-1, \quad (17)$$

$$\mathbf{T}_q(\gamma_{P_{j_i}|\overline{P_{j_i}}}) = 0, \quad i = m, m+1, \dots, 2^n, \quad (18)$$

hence,

$$\begin{aligned}
& [\mathbf{T}_q(\gamma_{P_{j_i}}|\overline{P_{j_i}})]^\beta \\
&= [\mathbf{T}_q(\rho_{P_{j_i}}|\overline{P_{j_i}})]^\beta \\
&\leq \sum_{i=0}^{2^n-1} (2^\beta - 1)^{w_H(\vec{j}_i)} [\mathbf{T}_q(\gamma_{P_{j_i}}|\overline{P_{j_i}})]^\beta
\end{aligned}$$

$$= \sum_{i=0}^{m-1} (2^\beta - 1)^{w_H(\vec{j}_i)} [\mathbf{T}_q(\rho_{P_{j_i}}|\overline{P_{j_i}})]^\beta, \quad (19)$$

here the first equality is due to (17), the second equality is due to (18) \square