

On the Compatibility of Constructive Predicative Mathematics with Weyl’s Classical Predicativity

Michele Contente

Czech Academy of Sciences, Prague, Czechia

Maria Emilia Maietti

Department of Mathematics, University of Padova, Italy

July 8, 2024

Abstract

It is well known that most foundations for Bishop’s constructive mathematics are incompatible with a classical predicative development of analysis as put forward by Weyl in his *Das Kontinuum*.

Here, we first recall how this incompatibility arises from the possibility, present in most constructive foundations, to define sets by quantifying over (the exponentiation of) functional relations. This possibility is not allowed in modern formulations of Weyl’s logical system.

Then, we argue how a possible way out is offered by foundations, such as the Minimalist Foundation, where exponentiation is limited to a primitive notion of function defined by λ -terms as in dependent type theory.

The price to pay is to renounce the so-called rule of unique choice identifying functional relations with λ -terms, and to number-theoretic choice principles, characteristic of foundations aimed to formalize Bishop’s constructive analysis. This restriction calls for a *point-free* constructive development of topology as advocated by P. Martin-Löf and G. Sambin with the introduction of Formal Topology.

Hence, we conclude that the Minimalist Foundation promises to be a natural crossroads between Bishop’s constructivism and Weyl’s classical predicativity provided that a point-free reformulation of classical analysis is viable.

Contents

1	Introduction	2
2	Weyl’s Classical Predicativity	3
3	Constructive Predicative Theories	6
3.1	Aczel’s Constructive Zermelo-Fraenkel set theory	6
3.2	Martin-Löf’s type theory	7
3.3	Homotopy Type Theory	9
4	The incompatibility of $\mathbf{HA}^\omega + \mathbf{iRC!}_{\mathbb{N},\mathbb{N}}$ with Weyl’s predicativity	10
5	The incompatibility of MLTT, HoTT and CZF with Weyl’s predicativity	12

6	A possible way out: the Minimalist Foundation	14
7	Conclusion	18

1 Introduction

The book *Das Kontinuum* [55, 56] is a milestone in the foundations of mathematics and, in particular, of classical predicative mathematics. Weyl’s monograph is mainly devoted to the predicative reformulation of real analysis.

Another major reformulation of real analysis within a foundation weaker than the usual classical one was given by Bishop in [6]. While Bishop followed Brouwer in adopting intuitionistic logic, he refrained in his approach from including Brouwer’s continuity principles that are inconsistent with classical mathematics. Bishop’s aim was to develop *constructive* mathematics not in opposition to classical mathematics but as a generalization of the latter.

After the publication of Bishop’s book, there was a flourishing of foundations for constructive mathematics aimed at formally capturing Bishop’s approach. A relevant example in axiomatic set theory is Aczel’s constructive set theory **CZF** [1], and in type theory Martin-Löf Type Theory **MLTT** [38, 39] and the more recent Homotopy Type Theory **HoTT** in [53] inspired by Voevodsky’s Univalent Foundations. In particular, **HoTT** gained a lot of attention for its application to the formalization of synthetic homotopy theory and higher category theory. However, these systems are sometimes not even mutually compatible and this consideration led the second author in joint work with G. Sambin to conceive the Minimalist Foundation (for short **MF**) in [33], later fully worked out as a two-level system in [24]. The two-level structure of **MF** in [24], which includes an intensional level, an extensional level, and an interpretation of the latter into the first, was introduced to better accomplish the compatibility of **MF** with constructive and classical foundations including the mentioned ones (see [24, 9]) and internal theories of categorical universes such as that of elementary topoi in [22, 23].

It is also well known that, among all the foundations of constructive and predicative mathematics, at least the mentioned ones, **CZF**, **MLTT** and **HoTT**, are incompatible with a classical predicative development of mathematics as advocated by Weyl in *Das Kontinuum*, since they become impredicative when extended with the law of excluded middle.

Here we first discuss the source of this incompatibility. The reason lies in their possibility of defining sets by quantifying over (the exponentiation of) functional relations. Such a possibility is not allowed in modern formulations of Weyl’s system [5, 4, 3, 16].

Then, we argue that it is not necessary to sacrifice exponentiation altogether to reconcile Bishop’s mathematics with Weyl’s approach if we adopt a foundation like the Minimalist Foundation and this for a number of reasons.

First of all, the intuitionistic version of the subsystem of second-order arithmetic known as **ACA** [48] can be interpreted in the extensional level of **MF** by preserving the meaning of all the logical operators, as well as the classical **ACA** can be analogously interpreted in the classical extension of **MF**. Therefore, Weyl’s reconstruction of analysis can be performed in the classical version of **MF**, since it is known from results of reverse mathematics in [16, 17, 48] that the formalization of Weyl’s analysis can be performed at least in **ACA**, whilst the proof-theoretic strength of Weyl’s original logical system exceeds it, see [5, 4, 3, 16].

Second, the classical version of **MF** is equi-consistent, as shown in [27, 44], with the original

constructive **MF**, whose proof-theoretic strength is strictly predicative a’ la Feferman being interpretable in first-order Martin-Löf’s type theory [24] and directly in Feferman’s \widehat{ID}_1 (see [20]).

Third, in both levels of **MF**, whilst the exponentiation of functional relations does not necessarily form a set, it does so the exponentiation of a primitive notion of function defined by λ -terms. This is possible since, in both levels of **MF**, the logic is many-sorted with sorts given by dependent types equipped with elements defined by λ -terms as in Martin-Löf’s type theory.

The price to be paid in **MF** is to renounce the so-called *rule of unique choice* identifying functional relations with λ -terms. In this way exponentiation of type-theoretic functions does not entail exponentiation of functional relations as it happens instead in Martin-Löf’s type theory and in Homotopy Type Theory, whose logics are also sorted on a dependent type theory. Accordingly, dropping the rule of unique choice entails dropping all the number-theoretic choice principles characteristic of Bishop’s mathematics as well as the fact that real numbers a’ la Dedekind or Cauchy form a set.

As a consequence, these limitations call for a *point-free* constructive development of topology as advocated by Martin-Löf and Sambin with the introduction of Formal Topology [45] to avoid the use of Brouwer’s Fan theorem as explained in [34]. Evidence that such an approach can be successfully developed is provided by the point-free results in [7, 41, 42, 7, 8, 21, 47].

Given that the point-free topology of Dedekind real numbers and that of Cantor space are definable in **MF**, if a point-free reformulation of classical analysis is fully viable, then **MF** would become a natural crossroads between Bishop’s constructivism and Weyl’s classical predicativity.

In the future, besides determining the exact proof-theoretic strength of **MF**, which is currently an open problem, we intend to investigate whether the extensions of **MF** with inductive and coinductive definitions in [28, 29, 32] is still predicative, or even still equi-consistent with their classical counterparts as established for **MF** in [27, 44]. A positive answer to one of these questions would provide a foundation more expressive than **MF** and capable of formalizing a whole development of point-free topology, including for example the results in [36, 15, 10, 47], while still maintaining compatibility with both Bishop’s constructive mathematics and Weyl’s classical predicative mathematics.

The structure of the paper is the following: first, we give a short overview of Weyl’s classical predicative conception in *Das Kontinuum* and highlight the points of convergence with some ideas underlying the Minimalist Foundation. In Section 3 we present some well-known examples of constructive predicative foundations. Then, in Section 4 we show that already Heyting Arithmetic in all finite types is incompatible with classical predicativity if it is extended with classical logic and the (number-theoretic) internal rule of unique choice. In Section 5 we extend this argument to other systems and analyze the incompatibility of **MLTT**, **HoTT** and **CZF** with classical predicativity. Finally, in Section 6 we introduce the Minimalist Foundation and we argue that it might represent a natural solution to the issues discussed in the previous sections.

2 Weyl’s Classical Predicativity

Das Kontinuum [55, 56] can be regarded as the first systematic attempt to develop *predicative* mathematics. The main focus of the book is the foundation of analysis and, moreover, the reconstruction of many fundamental results in this area without resorting to impredicative principles. Examples of theorems shown by Weyl in his framework include the Fundamental Theorem of the Calculus, Uniform continuity for continuous functions over a closed interval and existence of a maximum and a minimum for them as well as *sequential* versions of the Heine-Borel Theorem and the Least Upper Bound principle. Therefore, the main merit of *Das Kontinuum* was to show that predicative

mathematics was a viable project. Feferman has shown that the above mentioned results can be formalized in the subsystem of second-order arithmetic known as \mathbf{ACA}_0 and further evidence in support of this claim has been provided by the Reverse Mathematics program [48]. In this section, we review some basic tenets of Weyl’s approach in relation to the Minimalist Foundation, which will be introduced later in Section 6.

Weyl acknowledged - in agreement with Poincaré- the absolutely primitive character of the natural numbers and the associated notion of iteration. According to Weyl, every attempt to understand natural numbers in terms of more fundamental notions – as it happens in Zermelo’s set theory or in the logicist program of developing arithmetic on the basis of purely logical notions– was doomed to failure because it necessarily presupposes this very notion:

‘And I became firmly convinced (in agreement with Poincaré, whose philosophical position I share in so few other respects) that the *idea of iteration*, i.e., *of the sequence of natural numbers*, is an ultimate foundation of mathematical thought - in spite of Dedekind’s “theory of chains” which seeks to give a logical foundation for definition and inference by complete induction without employing our intuition of the natural numbers. For if it is true that the basic concepts of set theory can be grasped only through this “pure” intuition, it is unnecessary and deceptive to turn around them and offer a set-theoretic foundation for the concept of “natural number”.’ (*ibid.*, p.48)

Weyl’s conception of set bears some resemblance to the way sets are understood in Martin-Löf’s type theory (\mathbf{MLTT}) and also in both levels of the Minimalist Foundation (\mathbf{MF}), which will be presented in the next sections. In such theories, a set is determined by rules for forming its elements equipped with an induction principle to prove properties on it and a recursive definition to define a function by recursion from it to another set, see [39].

Furthermore, Weyl considered the notion of set (and function) underlying the classical foundation of analysis to be irremediably circular. Weyl noted that many basic definitions of analysis (e.g. that of least upper bound) were actually circular, and decided to reformulate analysis by avoiding such viciously circular definitions in favour of a predicative development. In particular, such a circularity clearly shows up in Dedekind’s construction of the real numbers: real numbers defined in this way do not form an *extensionally determinate collection*¹. Weyl elaborates on this point in a letter to Hölder, published as appendix in [56]:

‘If A is a property of properties, then one may form the property P_A which belongs to an object x if and only if there is a property constructed by means of these principles which belongs to x and itself possesses the property A . That would be a blatant *circulus vitiosus*; yet our current version of analysis commits this error and I consider it ground for censure.’ ([56], p.113)

In particular, it follows from this quotation that the power of a set is not a set according to Weyl. Thus we cannot define a new set by quantifying over the powerset to which it should belong. The subtle point is that this does not mean that the power of a set does not exist, but simply that it is not a set but a collection that is not inductively generated. In the Minimalist Foundation, this distinction between sets and collections is explicitly assumed. Accordingly, in Weyl’s work collections exceed the power-collection of a set, as shown in [4, 3, 5], but they are not allowed to enter into the definition of new sets.

¹See [5, 14] for a detailed analysis of this concept.

A major distinction between sets and collections in the Minimalist Foundation is that, whilst both entities are generated by rules forming their canonical elements as sets in Martin-Löf’s type theory, sets distinguish from collections for being equipped with an induction principle and a recursive definition of functions from them toward any other entity. Collections instead do not enjoy any induction principle or recursive definition from them, since inductive or recursive definitions acting on them would not be *invariant* under extensions of the theory with new sets. This approach aligns closely with the idea that predicative sets should be invariant, which traces back to Poincaré and has been carefully analyzed in [13].

Both in *Das Kontinuum* and in the Minimalist Foundation real numbers provide an example of a collection which is not a set. In particular, Weyl introduced real numbers as Dedekind cuts of rationals and are formalized as quadruples of natural numbers. The fact that the resulting collection \mathbb{R} of real numbers is not *extensionally determinate* in Weyl’s terminology, means that *we cannot define new sets by quantification over \mathbb{R}* , in sharp contrast with the case of natural numbers. This implies, for instance, that the least upper bound principle for sets of real numbers is not predicatively provable, while the principle is provable for sequences of real numbers, since the latter do not involve quantification over \mathbb{R} .

Finally, we want to point out that Weyl in *Das Kontinuum* places a significant emphasis on his notion of function and contrasts it with the set-theoretical definition. To this purpose, Avron writes in [5]:

‘Weyl totally rejected the modern view of a function as any set of pairs that satisfies the functionality condition. He insists that (like sets) functions can be determined only by a *rule*’.

With reference to this, it might be intriguing to attribute to Weyl a kind of computational understanding of what a function is. Indeed, Weyl works in a classical framework and assumes that functions are extensional objects just as sets in his system. However, Weyl’s quotation is in line with the fact that in the Minimalist Foundation we distinguish between functions as functional relations and functions as primitive operations defined by λ -terms. The distinct character of these two notions of functions is a peculiarity of **MF** among foundations for constructive predicative mathematics. While it might be argued that Weyl foreshadowed this distinction, nevertheless his technical treatment of the notion of function is not exactly the same. Therefore, we can only highlight the significance of the insight of viewing functions as rule-based objects. Moreover, this resembles the notion of Brouwer lawlike sequence, which we will identify with a λ -term in the Minimalist foundation (see [46]).

There have been various attempts in the literature to formalize Weyl’s system in a modern fashion, since Weyl’s original presentation presents some ambiguities. An influential reconstruction of Weyl’s system has been offered by Feferman in several works [16, 17, 18]. Feferman has extensively argued that Weyl’s system should correspond to the theory **ACA**₀ or, if the full induction scheme is accepted, to **ACA**, which are well-known subsystems of second-order arithmetic [48]. However, Feferman’s proposal has been criticized by Adams and Luo in [4, 3] and more recently by Avron in [5]. They observed that in certain sections of *Das Kontinuum*, Weyl introduces non-arithmetical sets and makes use of induction for non-arithmetical formulas. Therefore, they concluded that Weyl’s system should be stronger than what thought by Feferman, even if **ACA**₀ should be sufficient to reconstruct predicative analysis as developed in *Das Kontinuum*.

The systems introduced in these papers are many-sorted logics axiomatized over a dependent type theory, in a way similar to both levels of the Minimalist Foundation. But they differ between

them and with **MF** for what regards exponentiation of functions. Indeed, Adams and Luo’s system in [4, 3] includes some exponentiation of λ -functions in the form of function types, which are treated as *large types* in the sense that they are not contained in the universe of small types, that in turn can be thought of as the universe of sets. Then, Avron in [5] restricts the possibility to form exponentials by assuming that only certain types might serve as codomains. However, in his system we cannot define a set by quantifying over functions. Instead, in **MF** there is no exponentiation of functions between collections but only sets exponentiating λ -functions between sets.

3 Constructive Predicative Theories

Among the most popular frameworks introduced as foundations for Bishop’s constructive mathematics there are Constructive Zermelo-Fraenkel set theory **CZF** [1] and Martin-Löf Type Theory **MLTT** [39]. More recently, an extension of the latter called Homotopy Type Theory **HoTT** [53] has gained much attention, especially for its applications to synthetic homotopy theory and higher categories. All these systems are examples of constructive and predicative theories. However, the predicativity of these theories is closely tied to the adoption of a constructive logic. Indeed, whenever they are extended with the law of excluded middle or equivalent classical principles, they become impredicative and thus they are not compatible with classical predicativity à la Weyl. Briefly, the issue is that such theories are closed under the *exponentiation of functional relations* and the two-element set that, when classical logic is added, are strong enough to allow for impredicative constructions. Therefore, we cannot establish the predicativity of their classical counterparts, as it happens instead in the case of Heyting arithmetic or of the Minimalist Foundation, which we will introduce in detail in Section 6.

3.1 Aczel’s Constructive Zermelo-Fraenkel set theory

CZF [1] is a theory formulated in the same language of classical **ZF** and based on intuitionistic logic. The theory includes the following axioms: *extensionality, pairing, union, restricted separation, infinity, \in -induction, strong collection, and subset collection*.

The predicative nature of the theory is reflected in the restriction of *separation to Δ_0 -formulas*, which means that the defining formula ϕ might quantify only over sets that have already been introduced, and in the rejection of the full powerset axiom. The powerset axiom is replaced by the weaker axiom of subset collection, which in turn implies the exponentiation axiom for functions. Functions are defined as functional relations in the standard set-theoretic manner.

We recall that the Axiom of Choice in the form

$$(\mathbf{AC}) \quad \forall_{(x \in A)} \exists_{(y \in B)} R(x, y) \rightarrow \exists_{(f \in A \rightarrow B)} \forall_{(x \in A)} R(x, f(x))$$

cannot be added to **CZF** while preserving its constructive character:

Theorem 3.1. **CZF** + **AC** derives the law of excluded middle for Δ_0 -formulas.

Proof. See [2]. □

Remark 3.2. In [2], it is shown that this proof can be carried out in a weaker subsystem of **CZF**. On the other hand, the weaker principles of *countable choice* and *dependent choice* can be added to **CZF** without leading to undesirable results.

Moreover, in **CZF** real numbers form a set:

Theorem 3.3. *Real numbers both as Dedekind cuts or as Cauchy's sequences form a set in **CZF**.*

Proof. See [2]. □

3.2 Martin-Löf's type theory

Martin-Löf Type Theory (**MLTT**) [39] is a foundation for constructive mathematics, which was explicitly introduced to formalize Bishop's mathematics [37]. The theory is not presented in the usual language of first-order logic, but it is formulated in the language of dependent type theory.

Types of **MLTT** are introduced by means of four rules: formation, introduction, elimination and computation. In particular, the constructors of a given type are specified by the introduction rules, while the elimination rule states simultaneously a principle of induction at the level of types and of definition by recursion at the level of terms with the aid of the computation rules. The predicative nature of the theory is related to the fact that all the types are inductively generated by the relevant rules. Two fundamental aspects of **MLTT** are the presence of a *primitive notion of function*, called here λ -*function* to disambiguate it from the notion of functions as *functional relation*, and the *identification between propositions and types*. λ -functions are represented as λ -terms and this differentiates this foundational theory from other theories such as **CZF**. The identification between propositions and types makes the constructive logic of **MLTT** stronger than usual intuitionistic logic. This additional strength is visible in the case of the existential quantifier that is identified with the dependent sums type Σ . The elimination rule for this type is enough strong to allow for the definition of projection operators ². These projections play a fundamental role in establishing the following fact:

Theorem 3.4. *The type-theoretic Axiom of Choice*

$$(\mathbf{AC}) \quad \Pi_{(x:A)} \Sigma_{(y:B)} R(x, y) \rightarrow \Sigma_{(f:A \rightarrow B)} \Pi_{(x:A)} R(x, f(x))$$

*is derivable in **MLTT**.*

We remark that the rules for Σ -type are enough strong to derive also the corresponding *choice rule*. Furthermore, the Axiom of Unique Choice **AC!**, which identifies *functional relations* with λ -*functions*,

$$(\mathbf{AC}!) \quad \Pi_{(x:A)} \Sigma!_{(y:B)} R(x, y) \rightarrow \Sigma_{(f:A \rightarrow B)} \Pi_{(x:A)} R(x, f(x))$$

(where $\Sigma!_{(y:B)} R(x, y) \equiv \Sigma_{(y:B)} R(x, y) \times \Pi_{(y_1, y_2:B)} (R(x, y_1) \times R(x, y_2)) \rightarrow \text{Id}(B, y_1, y_2)$) also follows from the validity of **AC**.

Henceforth, we can refer in **MLTT** with the name *function* to either a *functional relation* or a λ -*function* without ambiguity. As anticipated in section 2, the two notions of function are instead kept distinct in the Minimalist Foundation as well as in some modern presentations of Weyl's system as those given in [3, 5].

Due to the absence of quotient sets in **MLTT**, real numbers, both à la Dedekind or à la Cauchy, are formally represented there using *setoids* whose elements are then indexed by a set. A setoid [40], called *set* by Bishop in ([6], p. 13), is a pair $(A, =_A)$, consisting of a set called *support* equipped

²We recall that the canonical elements of Σ -types given by the introduction rule are pairs.

with a suitable equivalence relation $a =_A a' [a, a' : A]$ used to equate the elements in the same equivalence class.

For example, Bishop reals in [6] can be represented as the setoid whose support is given by the set of Cauchy regular sequences from the set of natural numbers to that of rational numbers equipped with the equivalence relation $=_{\mathcal{R}_C}$ as in [6], pp.15-16. Bishop defined them by using the notion of *operation* which we can identify in **MLTT** with the notion of λ -function, i.e. with a λ -term of **MLTT**. Moreover, for what we have said about the notion of function in **MLTT**, *Cauchy reals* defined as Cauchy functional relations can be identified in **MLTT** with Bishop reals. For unifying the various concepts, we name them *Cauchy sequences a' la Bishop*.

Instead, in **MLTT** Dedekind reals can be defined only relatively to a universe of sets U_k (indexed by a natural number k) and are represented by a setoid whose support is given by the set of Dedekind cuts on the rationals relative to U_k as follows:

Definition 3.5. A Dedekind cut on the rationals is a pair of propositional functions $L, U : \mathbb{Q} \rightarrow U_k$ for some universe U_k that satisfy the usual conditions of being inhabited, disjoint, open, monotone, and located, all defined as in [35].

Definition 3.6. We denote with \mathbb{R}_{D_k} the support of Dedekind cuts made of propositional functions with values in the universe U_k for any natural number k .

Again thanks to the validity in **MLTT** of choice principles on natural numbers and thanks to the exponentiation of λ -terms by means of the dependent product type $\Pi_{(x:A)} B(x)$ of any family of types $B(x)(x : A)$, in **MLTT** each real number à la Dedekind in the form of a Dedekind cut corresponds bijectively to a Bishop real and hence to a Dedekind cut in \mathbb{R}_{D_0} .

Proposition 3.7. *In MLTT, Dedekind reals \mathbb{R}_{D_0} are in bijective correspondence with the Cauchy reals, defined as Cauchy sequences à la Bishop.*

Proof. See Prop. 17 in [7] and section 4.3 of [34]. □

As a consequence, we deduce that:

Proposition 3.8. *In MLTT, Dedekind reals \mathbb{R}_{D_0} are in bijective correspondence with \mathbb{R}_{D_k} for any natural number k .*

Proof. Given a Dedekind cut as in def. 3.5 in \mathbb{R}_{D_k} for any natural number k , by using the axiom of choice we can define a Bishop real given by an operation $\lambda n. x_n$, usually written $(x_n)_n$, as in Prop. 17 in [7] or section 4.3 in [34], whose cut (L_0, D_0) in \mathbb{R}_{D_0} is defined as

$$L_0 \equiv \{p : \mathbb{Q} \mid \exists n : \mathbb{N} \ p \leq x_n - 2/n\} \quad U_0 \equiv \{q : \mathbb{Q} \mid \exists n : \mathbb{N} \ q \geq x_n + 2/n\}$$

and is propositionally equivalent to the starting cut (L, U) in \mathbb{R}_{D_k} thanks to the correspondence between Dedekind cuts, formal points of point-free topology of real numbers and Bishop reals in section 4.3 of [34]. □

Remark 3.9. The previous proposition shows that some instances of resizing can be derived from choice principles constructively. In particular, this can be regarded as the fact that special cases of Russell's Reducibility Axiom can be derived in the presence of choice principles provided that the logic is constructive.

3.3 Homotopy Type Theory

Homotopy Type Theory (**HoTT**) [53] is an extension of **MLTT** and is an instance of Voevodsky's Univalent Foundations. It extends **MLTT** with the *Univalence Axiom* and *Higher Inductive types*, examples of which include quotients of homotopy sets and propositional truncation. The Univalence Axiom is formulated as follows:

the map $\text{idtoeqv} : (A =_{\mathcal{U}} B) \rightarrow (A \simeq B)$ is an equivalence

where $(A \simeq B)$ is the type of equivalences between A and B . The map idtoeqv and the notion of equivalence \simeq are defined as in Section 2 of [53]. Among the consequences of this axiom, there are certain extensional principles (e.g. function extensionality) that are not available in intensional **MLTT**.

A significant feature of **HoTT** is allowing for a representation of logical notions alternative to the propositions-as-types paradigm. The key notion here is that of *h-proposition*, where a type P is a h-proposition whenever for all $x, y : P$, $x =_P y$ holds, and it is analogous to the notion of *mono type* in the internal theory of topos in [23]. Further, a type A is a *h-set* if the associated identity type $=_A$ is a h-proposition.

The logic of h-propositions behaves differently from the propositions-as-types logic and is similar to the internal logic of elementary topoi or of regular categories as presented in [23]. In particular, the existential quantifier is not identified with the Σ -type any longer. Rather, it is identified with the propositional truncation of the Σ :

$$\exists_{(x:A)} P(x) \quad \equiv \quad \parallel \Sigma_{(x:A)} P(x) \parallel$$

Observe that the propositional truncation of a type is always a h-proposition. Indeed the constructors for the propositional truncation of a type A are the following: if $a : A$, then $| a | : \parallel A \parallel$ and $x =_{\parallel A \parallel} y$ holds for all $x, y : \parallel A \parallel$. Moreover, it holds that if P is a h-proposition, then it is equivalent to its truncation.

Observe also that in **HoTT**, while it is obviously possible to derive the type-theoretic **AC** under the identification propositions-as-types, it is not instead possible to derive the type-theoretic **AC** using the identification of propositions as h-propositions in the version:

$$\Pi_{(x:A)} \exists_{(y:B)} R(x, y) \rightarrow \exists_{(f:A \rightarrow B)} \Pi_{(x:A)} R(x, f(x))$$

with A, B h-sets and $R(x, y)$ h-proposition for $x : A, y : B$, as shown in [53]. Indeed, in **HoTT** it is only possible to derive the axiom of unique choice **AC!** formulated as follows

$$(\mathbf{AC!}) \Pi_{(x:A)} \exists!_{(y:B)} R(x, y) \rightarrow \Sigma_{(f:A \rightarrow B)} \Pi_{(x:A)} R(x, f(x))$$

with A, B types and $R(x, y)$ h-proposition and where unique existence $\exists!_{(y:B)} R(x, y)$ amounts to the following condition:

$$\exists_{(y:B)} R(x, y) \times \Pi_{(y_1, y_2:A)} R(x, y_1) \wedge R(x, y_2) \rightarrow y_1 =_B y_2.$$

Proposition 3.10. **AC!** can be derived in **HoTT**.

Proof. See [43]. □

Remark 3.11. Unique existence can be formulated in an equivalent way in **HoTT** by employing the notion of contractible type. A type A is said to be contractible if the following type is inhabited:

$$\Sigma_{(x:A)} \Pi_{(y:A)} x =_A y.$$

Therefore, we can rewrite unique choice by requiring the contractibility of $\Sigma_{(y:B)} R(x, y)$.

Remark 3.12. The proof of **AC!** in [43] is a special case of a result that holds in the internal dependent type theory \mathcal{T}_{reg} of any regular category [23]. Following [23], observe that the propositional truncation of a Σ -type $\Sigma_{(x:A)} P(x)$ can be equivalently expressed in \mathcal{T}_{reg} as a quotient, namely the quotient $\Sigma_{(x:A)} P(x)/\mathbf{1}$ over the unit type identifying all the proof-terms. Then, the proof in [43] corresponds to the argument in [23] relying on the fact that unique existence can be written as $\Sigma_{(y:B)} R(x, y)$ together with the usual uniqueness condition. In this case, $\Sigma_{(y:B)} R(x, y)$ is a mono type, according to the terminology introduced in [23], and is equivalent to its quotient over the unit type, and hence we can use Σ -projections to prove **AC!**.

A crucial feature of **HoTT** is the presence of quotient sets as higher inductive types. This means that Cauchy real numbers à la Bishop can be formalized as a quotient of the type of Cauchy sequences under the equivalence relation $=_{\mathbb{R}_C}$ without resorting to setoids. In particular, they form a set in **HoTT**.

The case of Dedekind reals is more delicate. The notion of cut can be formalized as a pair of propositional functions $(L, U) : \mathbb{Q} \rightarrow \mathbf{Prop}_{U_k}$ relative to some univalent universe of set U_k , that satisfy the conditions of Def.3.5. So it is possible to show that the Dedekind cuts up to a given universe form a set, but we cannot collect into a set all the Dedekind cuts varying in any universe unless impredicativity in the form of *propositional resizing* is assumed. Indeed, for the absence of dependent choice Prop.3.8 fails in **HoTT**, and hence there is no reduction of Dedekind reals of level k to those of the lowest level.

This fact can be contrasted with what happens in the Minimalist Foundation, and in particular in its extensional level where the presence of the power *collection* of a set allows us to collect all Dedekind cuts together inside the theory.

In any case, it must be noticed that in **HoTT** real numbers à la Cauchy are defined as a higher inductive type \mathbb{R}_C ([53], see Sec. 11.3). Hence, they form a set. However, the construction is different from the usual type-theoretic definition of Cauchy reals or Bishop reals. In particular, the definition of \mathbb{R}_C is intended to formalize the idea that Cauchy reals can be regarded as *the free complete metric space generated by the rationals*.

4 The incompatibility of $\mathbf{HA}^\omega + \mathbf{iRC!}_{\mathbb{N}, \mathbb{N}}$ with Weyl's predicativity

In this section, we want to show that already the weaker system of Heyting Arithmetic in all finite types (\mathbf{HA}^ω), when extended with the internal rule of unique choice and the law of excluded middle, becomes impredicative. Since the most common constructive and predicative foundations interpret this system, then the result can be easily extended to these more expressive theories.

For the presentation of \mathbf{HA}^ω , we refer to the standard reference [52].

The finite type structure \mathbf{T} is defined as follows: \mathbf{N} is a finite type and if σ, τ are finite types, then so are $\sigma \rightarrow \tau$ and $\sigma \times \tau$.

The language of \mathbf{HA}^ω includes countably infinite variables $x^\sigma, y^\sigma, z^\sigma, \dots$ for each type symbol $\sigma \in \mathbf{T}$. A binary predicate for equality $=_\sigma$ at each type $\sigma \in \mathbf{T}$ and an application operator $\mathbf{Ap}_{\sigma,\tau}$ for all $\sigma, \tau \in \mathbf{T}$. Furthermore, the language comprehends constants for zero and successor, recursor, pairing and projections and the combinators \mathbf{k} and \mathbf{s} .

The axioms of \mathbf{HA}^ω include the axioms of many-sorted intuitionistic predicate logic with equality, congruence rules for equality, the usual axioms for zero and successor with the induction scheme, and the defining conditions for the constants.

We recall two fundamental results about \mathbf{HA}^ω . It is immediate to realize that Heyting Arithmetic (\mathbf{HA}) is a subsystem of \mathbf{HA}^ω . Furthermore, if we extend \mathbf{HA}^ω with the Axiom of Choice for all finite types

$$(\mathbf{AC}_{\sigma,\tau}) \quad \forall x^\sigma \exists y^\tau \phi(x, y) \supset \exists f^{\sigma \rightarrow \tau} \forall x^\sigma \phi(x, fx)$$

the resulting theory is conservative over \mathbf{HA} . This result is known as Goodman's Theorem [19]. Another significant fact is that \mathbf{HA}^ω is equi-consistent with its classical version by a double negation translation. This result can be found in [51]. However, this result does not apply to the case of $\mathbf{HA}^\omega + \mathbf{AC}_{\sigma,\tau}$. Indeed, it is possible to show that $\mathbf{HA}^\omega + \mathbf{AC}_{\sigma,\tau} + \mathbf{LEM}$ is enough strong to derive the second-order comprehension axiom and thus to interpret full second-order classical arithmetic \mathbf{PA}^2 , which is an impredicative theory. The first to observe that \mathbf{HA}^ω extended with the Axiom of Choice and classical logic allows to derive the second-order comprehension principle was Spector in [49]. We remark that already the *internal rule of unique choice* ($\mathbf{iRC!}_{\sigma,\tau}$) is sufficient to carry out the proof.

Definition 4.1. The *internal rule of unique choice* for \mathbf{HA}^ω is formulated as follows:

$$\text{if } \mathbf{HA}^\omega \vdash \forall x^\sigma \exists! y^\tau \phi(x, y) \quad \text{then } \mathbf{HA}^\omega \vdash \exists f^{\sigma \rightarrow \tau} \forall x^\sigma \phi(x, fx)$$

where $\exists! y^\tau \phi(x, y) \equiv \exists y^\tau \phi(x, y) \wedge \forall y_1^\tau \forall y_2^\tau (\phi(x, y_1) \wedge \phi(x, y_2) \supset y_1 =_\tau y_2)$.

Remark 4.2. We call the rule displayed above *internal rule of unique choice* to contrast it with the following *rule of unique choice*: if $\mathbf{HA}^\omega \vdash \forall x^\sigma \exists! y^\tau \phi(x, y)$, then there exists a function $f^{\sigma \rightarrow \tau}$ such that $\mathbf{HA}^\omega \vdash \forall x^\sigma \phi(x, fx)$, where the existence of the choice function is stated externally. Admittedly, in \mathbf{HA}^ω the two rules are equivalent by the results in [51]. In general, this equivalence holds for all the systems that satisfy the existence property, but it cannot be stated in general. For instance, it does not hold for classical systems.

The theory \mathbf{PA}^2 is usually formulated as an extension of the language of \mathbf{PA} by a second sort of variables $X, Y, Z \dots$ standing for subsets of natural numbers and a membership symbol \in , where $t \in X$ is a new atomic formula for t first-order term. The axioms include the second-order induction axiom and the full scheme of comprehension for sets of natural numbers. However, the theory can be equally formulated in a language with typed primitive function symbols. Then first-order variables are interpreted as variables of type \mathbf{N} , second-order variables are interpreted by their characteristic functions with type $\mathbf{N} \rightarrow \mathbf{N}$ and the atomic formulas $t \in X$ are interpreted by $X(t) = 1$. In particular, the comprehension axiom is formulated as follows:

$$(\mathbf{CA}) \quad \exists f^{\mathbf{N} \rightarrow \mathbf{N}} \forall x^{\mathbf{N}} (fx = 1 \Leftrightarrow \phi(x))$$

where $\phi(x)$ is an arbitrary formula of the language provided that f does not occur free in ϕ .

Now, we show that in $\mathbf{HA}^\omega + \mathbf{iRC!}_{\mathbf{N},\mathbf{N}} + \mathbf{LEM}$ this comprehension principle is derivable.

Proposition 4.3. *The theory $\mathbf{HA}^\omega + \mathbf{iRC!}_{\mathbf{N},\mathbf{N}} + \mathbf{LEM}$ derives the second-order comprehension principle \mathbf{CA} .*

Proof. Let us consider a formula $\phi(x)$ with x of type \mathbf{N} . Then, we can define the following relation

$$\chi_\phi(x, y) := (\phi(x) \wedge y^{\mathbf{N}} = 1) \vee (\neg\phi(x) \wedge y^{\mathbf{N}} = 0)$$

By the validity of \mathbf{LEM} , we can argue by cases:

- if $\phi(x)$ holds, then $\chi_\phi(x, 1)$ holds.
- if $\neg\phi(x)$ holds, then $\chi_\phi(x, 0)$ holds.

We recall that $0 \neq 1$ follows from the axioms of \mathbf{HA}^ω . Thus we obtain $\forall x^{\mathbf{N}} \exists! y^{\mathbf{N}} \chi_\phi(x, y)$, that is a functional relation.

Therefore, we can apply $\mathbf{iRC!}_{\sigma, \tau}$ (for $\sigma, \tau = \mathbf{N}$) and we obtain that

$$\exists f^{\mathbf{N} \rightarrow \mathbf{N}} \forall x^{\mathbf{N}} \chi_\phi(x, fx)$$

which, unfolding the definition of χ_ϕ , amounts to

$$\exists f^{\mathbf{N} \rightarrow \mathbf{N}} \forall x^{\mathbf{N}} ((\phi(x) \wedge fx = 1) \vee (\neg\phi(x) \wedge fx = 0))$$

From which \mathbf{CA} follows. □

Corollary 4.4. The theory \mathbf{PA}^2 can be interpreted within $\mathbf{HA}^\omega + \mathbf{iRC!}_{\mathbf{N},\mathbf{N}} + \mathbf{LEM}$, which is then impredicative and incompatible with Weyl's predicativity.

5 The incompatibility of MLTT, HoTT and CZF with Weyl's predicativity

In this section, we show that the classical extensions of the systems presented in Section 3 are not compatible with classical predicativity.

By *compatibility of a theory T_1 with another theory T_2* we mean that *there exists a translation from T_1 to T_2 preserving the meaning of logical and set-theoretical constructors*. In this manner, a proof of a theorem in T_1 can be transported into T_2 with the same meaning.

It is easy to see that \mathbf{HA}^ω extended with $\mathbf{iRC!}_{\mathbf{N},\mathbf{N}}$ is compatible both with \mathbf{MLTT} and \mathbf{HoTT} since these are essentially extensions of \mathbf{HA}^ω validating $\mathbf{iRC!}_{\mathbf{N},\mathbf{N}}$.

Definition 5.1. A compatible translation from $\mathbf{HA}^\omega + \mathbf{iRC!}_{\mathbf{N},\mathbf{N}}$ to \mathbf{MLTT} can be defined as follows:

- $(\mathbf{N})^* := \mathbb{N}$, while $(\sigma \rightarrow \tau)^* := \sigma^* \rightarrow \tau^*$ and $(\sigma \times \tau)^* := \sigma^* \times \tau^*$.
- variables $x : \sigma$ are translated as variables $x : \sigma^*$ and application terms as application terms in \mathbf{MLTT} .
- constants of \mathbf{HA}^ω are translated as the corresponding terms in \mathbf{MLTT} : e.g., $(\mathbf{k})^* := \lambda x. \lambda y. x : \sigma^* \rightarrow (\tau^* \rightarrow \sigma^*)$.

- the formulas of many-sorted intuitionistic logic are sent to the corresponding types according to the propositions-as-types principle: e.g., $(\exists x^\sigma \phi(x))^* \equiv \Sigma_{(x:\sigma^*)} \phi^*(x)$.

Proposition 5.2. *The translation $(-)^*$ is a compatible translation of $\mathbf{HA}^\omega + \mathbf{iRC!}_{\mathbf{N},\mathbf{N}}$ in \mathbf{MLTT} .*

Proof. The proof is by induction over the structure of derivations. In particular, the interpretation of the number-theoretic internal rule of unique choice follows from the validity of the more general rule of choice in \mathbf{MLTT} . \square

Therefore, when \mathbf{MLTT} is extended with the law of excluded middle, we can reproduce the argument showing the incompatibility of \mathbf{HA}^ω with Weyl's predicativity:

Corollary 5.3. \mathbf{MLTT} with the addition of the law of excluded middle becomes impredicative.

Proof. It follows from Prop. 5.2 and Cor. 4.4. \square

Remark 5.4. In general, in \mathbf{MLTT} with classical logic we can use the type of booleans $\mathbf{2}$ as a classifier, however we do not get a (strong) power object. For instance, the type $\mathbf{N} \rightarrow \mathbf{2}$ behaves more like a *weak power object*, because \mathbf{MLTT} lacks of the relevant extensionality principles.

The translation of \mathbf{HA}^ω into \mathbf{MLTT} defined above can be refined in the case of \mathbf{HoTT} . A finite type σ is translated as a pair $\sigma^* \equiv (\sigma^*, p_{\sigma^*})$, where the second component is a proof that σ^* is a h-set in \mathbf{HoTT} . Similarly, formulas of \mathbf{HA}^ω can be translated as h-propositions in \mathbf{HoTT} , while the rest of the translation is defined as in the case of \mathbf{MLTT} . Therefore, the following fact holds:

Proposition 5.5. *The translation $(-)^*$ is a compatible translation of $\mathbf{HA}^\omega + \mathbf{iRC!}_{\mathbf{N},\mathbf{N}}$ in \mathbf{HoTT} .*

As in the case of \mathbf{MLTT} , we can reproduce the same proof of Prop.4.3 but with a significant difference. The law of excluded middle cannot be formulated according to the propositions-as-types identifications for *all* types in \mathbf{HoTT} , because this would yield a contradiction in presence of Univalence (see Ch.3 [53]). However, it is still possible to formulate **LEM** for all those types that are h-propositions. This formulation serves our purposes and is consistent with Univalence.

Corollary 5.6. \mathbf{HoTT} with the addition of the law of excluded middle for h-propositions becomes impredicative.

Proof. It follows from Prop. 5.5 and Cor. 4.4. \square

Remark 5.7. When \mathbf{HoTT} is extended with **LEM** for h-propositions, it is possible to show that the type of h-propositions within any universe \mathcal{U} , that is $\mathbf{Prop}_{\mathcal{U}}$, is equivalent (and hence equal by Univalence) to the type of Booleans $\mathbf{2}$ in \mathcal{U} . Therefore, $\mathbf{2}$ behaves like a *subobject classifier* and we can prove that the category of h-sets, as defined in [53], is an elementary topos.

Definition 5.8. A compatible translation from $\mathbf{HA}^\omega + \mathbf{iRC!}_{\mathbf{N},\mathbf{N}}$ to \mathbf{CZF} can be defined as follows:

- $(\mathbf{N})^* \equiv \omega$, while $(\sigma \rightarrow \tau)^* \equiv \sigma^* \rightarrow \tau^*$, which is the set of functional relations from σ^* to τ^* , and $(\sigma \times \tau)^* \equiv \sigma^* \times \tau^*$, which is the usual cartesian product of sets.
- terms of \mathbf{HA}^ω are interpreted by the corresponding definable terms of \mathbf{CZF} .
- the formulas of many-sorted intuitionistic logic are interpreted by the corresponding formulas of intuitionistic logic, with sorts interpreted as predicates: e.g., $(\forall x^\sigma \phi(x))^* \equiv \forall x (x \in \sigma^* \rightarrow \phi^*(x))$.

Proposition 5.9. *The translation $(-)^*$ is a compatible translation of $\mathbf{HA}^\omega + \mathbf{iRC}!_{\mathbf{N},\mathbf{N}}$ in \mathbf{CZF} .*

Corollary 5.10. \mathbf{CZF} with the addition of the law of excluded middle becomes impredicative.

Proof. It follows from Prop. 5.9 and Cor. 4.4. □

It is worth noting that in the case of \mathbf{CZF} , the incompatibility with classical predicativity can be shown by a very direct and simple argument proved by Aczel in [1]:

Proposition 5.11. *The axiom of exponentiation and the law of excluded middle for Δ_0 -formulas imply the powerset axiom.*

Proof. It is enough to show that $\mathcal{P}(1)$ is a set. This follows from an application of **LEM**. The rest follows from the fact that the powerset axiom is equivalent to exponentiation plus the existence of the powerset of the singleton set. □

Moreover, since the powerset axiom implies subset collection, which in turn implies exponentiation, it follows:

Proposition 5.12. \mathbf{CZF} with classical logic and \mathbf{ZF} prove the same set of theorems.

Proof. See [1]. □

6 A possible way out: the Minimalist Foundation


In this section, we want to show that there exists a constructive predicative system that does not present the same incompatibility with classical predicativity as the other systems previously considered and, at the same time, is enough expressive to allow for the development of non-trivial mathematics by keeping a form of *exponentiation* for a primitive notion of functions defined by λ -terms. This system is the *Minimalist Foundation* **MF**.

MF is a two-level foundation for constructive mathematics, that was first conceived in [33] and then fully developed in [24]. It consists of an intensional level, called **mTT**, and an extensional one, called **emTT**, together with an interpretation of the latter into the first, as defined in [24].

The two-level structure facilitates the compatibility of **MF** with the most relevant constructive and classical foundations as shown in [24]: the intensional level **mTT** is compatible with **MLTT** and Coquand-Paulin's Calculus of Inductive Constructions [11], while the extensional level is compatible with **CZF**, **HoTT** and the internal theory of elementary topoi in [22, 23]. It is a remarkable property that *both* levels of **MF** are compatible with **HoTT** as shown in [9].

Moreover, both levels of **MF** extend a version of Martin-Löf's type theory with a primitive notion of proposition: **mTT** extends the intensional type theory in [39], while **emTT** extends the extensional version presented in [38].

In **MF** there are four primitive distinct sorts: small propositions, propositions, sets and collections. The following diagram shows the relations between these primitive sorts



The basic forms of judgement in **MF** include:

$$A \text{ set } [\Gamma] \quad B \text{ coll } [\Gamma] \quad \phi \text{ prop } [\Gamma] \quad \psi \text{ prop}_s [\Gamma]$$

to which we add the meta-judgement

$$A \text{ type } [\Gamma]$$

where ‘*type*’ is to be interpreted as a meta-variable ranging over the four basic sorts.

The set-constructors of **mTT** and **emTT** include those of first order Martin-Löf’s type theory, respectively as presented in [39] and [38]. Hence we have the following set constructors: the empty set N_0 , the singleton set N_1 , the set of lists $List(A)$ over a set A , hence the set of natural numbers \mathbb{N} , the indexed sum and the dependent product of the family of sets $B(x)$ *set* $[x \in A]$ denoted respectively as $\Sigma_{x \in A} B(x)$ and $\Pi_{x \in A} B(x)$, the disjoint sum $A + B$ of the set A with the set B . In the case of the extensional level **emTT** we have that sets are closed under effective quotients A/R over a set A , provided that R is a small equivalence relation $R(x, y) \text{ prop}_s [x \in A, y \in A]$. This set constructors is not included in the intensional level **mTT**. Furthermore, both the sets of **mTT** and those of **emTT** include also their small propositions $\phi \text{ prop}_s$ regarded as sets of their proofs. Small propositions are those propositions closed under intuitionistic connectives and under quantifiers and equalities restricted to sets.

Collections of **mTT** and **emTT** include their sets and the indexed sum $\Sigma_{x \in A} B(x)$ of the family of collections $B(x)$ *coll* $[x \in A]$ indexed over a collection A . Moreover, they also include propositions $\psi \text{ prop}$ regarded as collections of their proofs. A significant distinction between **mTT**-collections and **emTT**-collections is the following: the former includes the proper collection of small propositions prop_s and the collection of small propositional functions $A \rightarrow \text{prop}_s$ over a set A (which are definitely not sets predicatively when A is not empty), while the latter includes the power-collection of the singleton $\mathcal{P}(1)$, which is the quotient of the collection of small propositions under the relation of equiprovability, and the power-collection $A \rightarrow \mathcal{P}(1)$ of a set A , that can be written simply as $\mathcal{P}(A)$. In **mTT** we identify *predicates over the set* A with *propositional functions* defined as primitive λ -terms. Hence, in **emTT** a subset of a set A is identified with the equivalence class of propositional function up to equiprovability. So this use of λ -terms corresponding to predicates is a major difference with axiomatic set theory where subsets of a set A are simply defined as sets in a certain relation with A and hence are identified with functional relations into the boolean set $\{0, 1\}$ when the logic is classical.

Both propositions of **mTT** and **emTT** include standard connectives and quantifiers. However, they differ in the way the rules for propositional equality are given. Indeed, **mTT**-propositions include an *intensional* propositional equality defined as in [39], except for the fact that the elimination rule is *restricted* to propositions. While **emTT**-propositions include an *extensional* propositional equality with an elimination rule formulated like that in [38].

Finally, we recall that **emTT**-propositions are *proof-irrelevant* and have one canonical element denoted as **true**.

It is relevant for our discussion to remark that elimination rules for propositions act only towards propositions and not towards collections. The same applies to small propositions. In this way, **mTT** and **emTT** do not generally validate choice principles. In particular, even unique choice cannot be derived in **MF**, thanks to the results in [30, 31, 25] and similarly to what happens in Coquand’s Calculus of Constructions **CoC**, as first proved in [50]. This is in contrast with what happens in **MLTT** and in **HoTT**, as we have seen in Section 3. The fact that choice principles are not validated in **MF** is important in connection with the notion of compatibility introduced in Section 5. In this

way, **MF** turns out to be also compatible with theories where choice does not generally hold, such as **CoC**.

Now, we discuss the status of the internal rule of unique choice in **MF**, in particular in **mTT**.

Definition 6.1. In **mTT** the *internal rule of unique choice* holds if for any derivable small proposition

$$R(x, y) \text{ prop}_s [x \in A, y \in B]$$

and for any derivable judgement of the form

$$p(x) \in \exists!_{y \in B} R(x, y) [x \in A]$$

there exists a proof-term q such that the following judgement is derivable

$$q \in \exists_{f \in A \rightarrow B} \forall_{x \in A} R(x, \mathbf{Ap}(f, x))$$

We call *number-theoretic internal rule of unique choice* the instance of the internal rule of unique choice when A and B coincide with the set \mathbb{N} of natural numbers.

Instead, we reserve the name *rule of unique choice* for the version of the above rule requiring the external existence of the choice function term f , i.e. if $\exists!_{y \in B} R(x, y) [x \in A]$ holds then we can derive a term $f \in A \rightarrow B$ such that $R(x, \mathbf{Ap}(f, x)) [x \in A]$ holds.

From [25] we know that the rule of unique choice is not valid in **mTT**:

Proposition 6.2. *The rule of unique choice is not valid in mTT.*

Proof. See proof of Th.1 of [25]. □

The existence property of **mTT**, following from normalization results, would allow to deduce the equivalence of the rule of unique choice with its internal version and hence the non-validity of the internal rule of unique choice in **mTT**, too.

Nevertheless, it is worth pointing out that without relying on the existence property, we can show the not validity of the internal rule of unique choice following the same argument in [25] since this can be exported also to extensions of **mTT**, not validating the existence property, such as its classical version:

Proposition 6.3. *If mTT satisfies the internal rule of unique choice, then it satisfies the axiom of unique choice.*

Proof. The proof of Prop.3 in [25] can be easily adapted to the case of the internal rule of unique choice. □

Now, thanks to the model in [50], we deduce from Prop. 6.3 that

Corollary 6.4. **mTT** does not validate the internal rule of unique choice.

It is important to remark that already the intensional level **mTT** can be regarded as an extension of **HA**^ω. Indeed, it is possible to show that there is a compatible translation of this theory into **mTT** following a similar translation of Section 5 for **MLTT** by interpreting **HA**^ω-propositions as **mTT**-propositions. What is most interesting is that **MF** extends **HA**^ω while preserving the fundamental property of equi-consistency with its classical counterpart. Indeed, recently it has been shown that *both levels of MF are equi-consistent with their classical counterparts* (collapsed to the extension of the extensional level **emTT** with the law of excluded middle), unlike theories such as **MLTT**, **HoTT** and **CZF**:

Proposition 6.5. *Both levels of **MF** are equi-consistent with **emTT** with the addition of the law of excluded middle.*

Proof. For a proof, we refer to [27, 44]. □

Therefore, the extension of **MF** with the law of excluded middle is still predicative. From this, knowing that **HA**^ω can be embedding in **mTT** and **emTT** in a compatible way, we can deduce the non-derivability of unique choice on natural numbers in their classical version:

Corollary 6.6. The classical version of **emTT** and of **mTT** do not validate the number-theoretic internal rule of choice and hence the number-theoretic axiom of unique choice:

$$(\mathbf{AC}^!_{\mathbb{N},\mathbb{N}}) \forall_{x \in \mathbb{N}} \exists!_{y \in \mathbb{N}} R(x, y) \rightarrow \exists_{f \in \mathbb{N} \rightarrow \mathbb{N}} \forall_{x \in \mathbb{N}} R(x, f(x))$$

Proof. First, observe that **AC**[!]_{ℕ,ℕ} implies the number-theoretic internal rule of unique choice. Then, the claim follows from Prop. 6.5 and Cor. 4.4. □

Since the validity of the axiom of unique choice in a theory can be transferred to its extensions, contrary to the rule of unique choice, from Cor. 6.6 we can immediately deduce the non-validity of **AC**[!]_{ℕ,ℕ} in **mTT** and **emTT** themselves:

Corollary 6.7. Both **emTT** and of **mTT** do not validate the number-theoretic axiom of unique choice.

It is worth noting that in the extensional level **emTT** of **MF** we can also interpret the intuitionistic version of **ACA** in a way that preserves the meaning of all logical operators:

Proposition 6.8. *The intuitionistic version of **ACA** is interpretable in the extensional level **emTT** of **MF**. Hence, **ACA** can be interpreted in its classical version.*

Proof. It is sufficient to define a translation from **ACA** to **emTT** as follows. First-order variables of **ACA** are translated as variables of type \mathbb{N} in **emTT**. The constants for zero and the successor are interpreted by the corresponding constructors of type \mathbb{N} in **emTT**. Second-order variables for sets of natural numbers are interpreted as propositional functions $X \in \mathbb{N} \rightarrow \mathcal{P}(1)$ in **emTT** and the atomic formulas of the form $t \in X$ are translated as applications of these functions $X(t)$.

In general, first-order formulas of **ACA** are interpreted as small propositions of **emTT** with quantifiers and equalities restricted to \mathbb{N} . While second-order formulas are interpreted by propositions of **emTT** quantifying over collections.

Peano axioms can be derived from the rules for natural numbers in **emTT** and the presence of power-collections, see [24]. The induction scheme of **ACA** follows from the elimination rule for \mathbb{N} which acts towards all types including all propositions. Finally, the arithmetical comprehension axiom is validated as follows: given an arithmetical formula $\phi(x)$ in **ACA**, we know that it is interpreted as a small proposition in **emTT**. Then we can introduce $[\phi](x) \in [x \in \mathbb{N}]$ in $\mathcal{P}(1)$ and we get $X := \lambda x. \phi(x) \in \mathbb{N} \rightarrow \mathcal{P}(1)$. Therefore, it follows that there exists X such that for all $n \in \mathbb{N}$, $X(x) \leftrightarrow \phi$. □

Hence, from prop. 6.8 it follows that the classical version of **MF** can formalize Weyl's classical predicative reformulation of analysis in *Das Kontinuum* following Feferman's analysis in [17, 18].

Furthermore, from [24, 20] we know a proof-theoretic upper bound for **MF**:

Proposition 6.9. *The proof-theoretic strength of both levels of \mathbf{MF} are bounded by Feferman's theory \widehat{ID}_1 of non-iterative fixpoints.*

Corollary 6.10. *The proof-theoretic strength of the classical extension of \mathbf{MF} is bounded by Feferman's theory \widehat{ID}_1 of non-iterative fixpoints.*

Proof. It follows from Prop.6.5 and Prop.6.9. □

Another significant feature of \mathbf{MF} which makes it compatible with Weyl's classical predicativity, is that *real numbers do not form a set* in its extensional level \mathbf{emTT} (and this holds also for their intensional representation in the quotient model over \mathbf{mTT}). This reflects Weyl's idea that the collection of real numbers is not an extensionally determinate collection.

Proposition 6.11. *In the extensional level \mathbf{emTT} of \mathbf{MF} real numbers à la Dedekind or Cauchy do not form a set as well as in their classical version.*

Proof. A direct proof of this was given [26] but it follows also from the predicativity of the classical version of \mathbf{MF} . If real numbers form a set in \mathbf{MF} then this is so in the classical version of \mathbf{MF} and every interval of real numbers form a set too. Since classically, the power-collection of natural numbers can be put in bijection with an interval of real numbers, then it would follow that impredicative definitions can be employed in the classical version of \mathbf{MF} contrary to its predicativity established thanks in prop.6.10. □

Despite the absence of choice principles and quantification over all real numbers to form new sets, a development of mathematics, and in particular of topology and analysis, within \mathbf{emTT} is possible if point-free methods are adopted as advocated by Martin-Löf and Sambin with the introduction of Formal Topology in [45]. Indeed, in \mathbf{emTT} the following point-free topologies are definable:

Proposition 6.12. *In the extensional level \mathbf{emTT} of \mathbf{MF} the point-free topology of Dedekind real numbers and Cantor space are definable.*

Proof. The formalization of the point-free topology of the Cantor space can be carried on in \mathbf{emTT} as in section 2.1 of [54]. Then, observe that the point-free topology of real numbers can be defined as in Def.3 of [7] in terms of another cover generated by a finite set of axioms that can be represented in \mathbf{emTT} with a completely analogous proof to that just mentioned for the point-free topology of Cantor space. □

Therefore, one can carry on in \mathbf{MF} the development of analysis initiated by Palmgren in [7, 41, 42] (see also [21]) and that of Positive Topology in [47] and *loc.cit.*.

7 Conclusion

We have discussed the incompatibility with Weyl's classical predicativity of some foundations for Bishop mathematics like Aczel's constructive set theory \mathbf{CZF} , Martin-Löf Type Theory \mathbf{MLTT} and Homotopy Type Theory \mathbf{HoTT} . We have observed that this incompatibility reduces to the fact that these theories interpret Heyting arithmetic in all finite types \mathbf{HA}^ω , which is a many-sorted theory where sorts include finite types, with the addition of the internal rule of number-theoretic unique choice $\mathbf{iRC!}_{\mathbf{N},\mathbf{N}}$, identifying functional relations over natural numbers with a primitive notion of

function as λ -terms. Indeed, $\mathbf{HA}^\omega + \mathbf{iRC!}_{\mathbb{N},\mathbb{N}}$ with the further addition of the law of excluded middle becomes impredicative since it interprets second-order Peano arithmetic.

Whilst Weyl’s foundational system does not contain quantification over exponentiation of functions as observed by [5], we have argued that it is not necessary to sacrifice exponentiation altogether to reconcile Bishop’s mathematics with Weyl’s approach if we adopt a foundation like the Minimalist Foundation.

Given that the Minimalist Foundation interprets \mathbf{HA}^ω , the price to pay is to renounce the rule of unique choice as well as all the number-theoretic choice principles characteristic of Bishop’s mathematics and furthermore that real numbers à la Dedekind or Cauchy form a set. All these facts together call for a point-free development of analysis by adopting the topological methods advocated by Martin-Löf and Sambin with the introduction of Formal Topology [45].

Hence, provided that a point-free reformulation of classical analysis is viable, as hinted in [7, 41, 42, 21], \mathbf{MF} promises to be a natural crossroads between Bishop’s constructivism and Weyl’s classical predicativity.

In the future, besides establishing the exact proof-theoretic strength of \mathbf{MF} , which is currently an open problem, we will explore whether the extension of \mathbf{MF} with inductive and coinductive definitions in [28, 29, 32] is still predicative, or even is still equiconsistent with its classical counterpart, as established for \mathbf{MF} in [27, 44]. A positive answer to either of these questions would provide a foundation capable of formalizing general results of point-free mathematics as those presented in [12, 36, 15, 47], and still compatible with both Bishop’s constructivism and Weyl’s classical predicativity, as in the case of \mathbf{MF} .

References

- [1] P. Aczel. The type theoretic interpretation of constructive set theory. In *Studies in Logic and the Foundations of Mathematics*, volume 96, pages 55–66. Elsevier, 1978.
- [2] P. Aczel and M. Rathjen. \mathbf{CST} Book Draft. unpublished manuscript, 2010.
- [3] R. Adams and Z. Luo. Classical predicative logic-enriched type theories. *Ann. Pure Appl. Log.*, 161(11):1315–1345, 2010.
- [4] R. Adams and Z. Luo. Weyl’s predicative classical mathematics as a logic-enriched type theory. *ACM Transactions on Computational Logic (TOCL)*, 11(2):1–29, 2010.
- [5] A. Avron. Weyl Reexamined: “Das Kontinuum” 100 Years Later. *Bull. Symb. Log.*, 26(1):26–79, 2020.
- [6] E. Bishop. *Foundations of constructive analysis*, volume 60. McGraw-Hill, 1967.
- [7] J. Cederquist and S. Negri. A Constructive Proof of the Heine-Borel Covering Theorem for Formal reals. In S. Berardi and M. Coppo, editors, *Types for Proofs and Programs, International Workshop TYPES’95, Torino, Italy, June 5-8, 1995, Selected Papers*, volume 1158 of *Lecture Notes in Computer Science*, pages 62–75. Springer, 1995.
- [8] F. Ciraulo and G. Sambin. Reducibility, a constructive dual of spatiality. *Journal of Logic and Analysis*, 11, 2019.

- [9] M. Contente and M. E. Maietti. The compatibility of the minimalist foundation with homotopy type theory. *Theoretical Computer Science*, 991:114421, 2024.
- [10] T. Coquand, E. Palmgren, and B. Spitters. Metric complements of overt closed sets. *Math. Log. Q.*, 57(4):373–378, 2011.
- [11] T. Coquand and C. Paulin. Inductively defined types. In P. Martin-Löf and G. Mints, editors, *COLOG-88, International Conference on Computer Logic, Tallinn, USSR, December 1988, Proceedings*, volume 417 of *Lecture Notes in Computer Science*, pages 50–66. Springer, 1988.
- [12] T. Coquand, G. Sambin, J. Smith, and S. Valentini. Inductively generated formal topologies. *Ann. Pure Appl. Log.*, 124(1-3):71–106, 2003.
- [13] L. Crosilla. Predicativity and Constructive Mathematics. In G. Oliveri, C. Ternullo, and S. Boscolo, editors, *Objects, Structures and Logics*. Springer Cham, 2022.
- [14] L. Crosilla and Ø. Linnebo. Weyl and two kinds of potential domains. *Noûs*, 2023.
- [15] G. Curi. Abstract Inductive and Co-Inductive Definitions. *J. Symb. Log.*, 83(2):598–616, 2018.
- [16] S. Feferman. Weyl Vindicated: Das Kontinuum, 70 Years Later. In *Proceedings of the Cesena Conference on Logic and Philosophy of Science*, 1988.
- [17] S. Feferman. *In the Light of Logic*. Oxford University Press, 1998.
- [18] S. Feferman. The significance of Weyl’s Das Kontinuum. In *Proof Theory- History and Philosophical Significance*, pages 179–194. Springer, 2000.
- [19] N. D. Goodman. Relativized realizability in intuitionistic arithmetic of all finite types. *The Journal of Symbolic Logic*, 43(1):23–44, 1978.
- [20] H. Ishihara, M. E. Maietti, S. Maschio, and T. Streicher. Consistency of the intensional level of the Minimalist Foundation with Church’s Thesis and Axiom of Choice. *Archive for Mathematical Logic*, 57(7-8):873–888, 2018.
- [21] T. Kawai. Bishop Metric Spaces in Formal Topology. In M. R. Douglas Bridges, Hajime Ishihara and H. Schwichtenberg, editors, *Handbook of Constructive Mathematics*, pages 395–425. Cambridge University Press, 2023.
- [22] J. Lambek and P. J. Scott. *Introduction to Higher Order Categorical Logic*. Cambridge University Press, 1986.
- [23] M. E. Maietti. Modular correspondence between dependent type theories and categories including pretopoi and topoi. *Math. Struct. Comput. Sci.*, 15(6):1089–1149, 2005.
- [24] M. E. Maietti. A minimalist two-level foundation for constructive mathematics. *Annals of Pure and Applied Logic*, 160(3):319–354, 2009.
- [25] M. E. Maietti. On choice rules in dependent type theory. In *TAMC*, volume 10185 of *Lecture Notes in Computer Science*, pages 12–23, 2017.

- [26] M. E. Maietti. On minimality of the Minimalist Foundation. In *Mathematisches Forschungsinstitut Oberwolfach Report No. 34/2020, Mathematical Logic: Proof Theory, Constructive Mathematics (hybrid meeting)*. MFO, 2020.
- [27] M. E. Maietti. Equiconsistency of the Minimalist Foundation with its classical version. In *Mathematisches Forschungsinstitut Oberwolfach Report n.50/2023- Mathematical Logic: Proof Theory, Constructive Mathematics*. MFO, 2024.
- [28] M. E. Maietti, S. Maschio, and M. Rathjen. A realizability semantics for inductive formal topologies, Church’s Thesis and Axiom of Choice. *Logical Methods in Computer Science*, 17(2), 2021.
- [29] M. E. Maietti, S. Maschio, and M. Rathjen. Inductive and Coinductive Topological Generation with Church’s thesis and the Axiom of Choice. *Log. Methods Comput. Sci.*, 18(4), 2022.
- [30] M. E. Maietti and G. Rosolini. Quotient completion for the foundation of constructive mathematics. *Logica Universalis*, 7(3):371–402, 2013.
- [31] M. E. Maietti and G. Rosolini. Relating quotient completions via categorical logic. In D. Probst and P. Schuster, editors, *Concepts of Proof in Mathematics, Philosophy, and Computer Science*, pages 229–250, 2016.
- [32] M. E. Maietti and P. Sabelli. A topological counterpart of well-founded trees in dependent type theory. In M. Kerjean and P. B. Levy, editors, *Proceedings of the 39th Conference on the Mathematical Foundations of Programming Semantics, MFPS XXXIX, Indiana University, Bloomington, IN, USA, June 21-23, 2023*, volume 3 of *EPTICS*. EpiSciences, 2023.
- [33] M. E. Maietti and G. Sambin. Toward a Minimalist Foundation for Constructive Mathematics. In *From sets and types to topology and analysis*, volume 48 of *Oxford logic guides*. Oxford University Press, 2005.
- [34] M. E. Maietti and G. Sambin. Why Topology in the Minimalist Foundation Must Be Pointfree. *Logic and Logical Philosophy*, 22(2):167–199, 2013.
- [35] M. E. Maietti and G. Sambin. The Minimalist Foundation and Bishop’s Constructive Mathematics. In M. R. Douglas Bridges, Hajime Ishihara and H. Schwichtenberg, editors, *Handbook of Constructive Mathematics*, pages 525–563. Cambridge University Press, 2023.
- [36] M. E. Maietti and S. Valentini. A structural investigation on formal topology: coreflection of formal covers and exponentiability. *Journal of Symbolic Logic*, 69:967–1005, 2004.
- [37] P. Martin-Löf. An intuitionistic theory of types: Predicative part. In *Studies in Logic and the Foundations of Mathematics*, volume 80, pages 73–118. Elsevier, 1975.
- [38] P. Martin-Löf. *Intuitionistic type theory*, volume 1 of *Studies in Proof Theory*. Bibliopolis, 1984.
- [39] B. Nordström, K. Petersson, and J. M. Smith. *Programming in Martin-Löf’s type theory*, volume 200. Oxford University Press Oxford, 1990.
- [40] E. Palmgren. Bishop’s set theory, 2005. Slides for lecture at the TYPES summer school.

- [41] E. Palmgren. Continuity on the real line and in formal spaces. In L. Crosilla and P. M. Schuster, editors, *From Sets and Types to Topology and Analysis*, volume 48 of *Oxford Logic Guides*. Oxford University Press, 2005.
- [42] E. Palmgren. A constructive and functorial embedding of locally compact metric spaces into locales. *Topology and its Applications*, 154(9):1854–1880, 2007.
- [43] E. Rijke and B. Spitters. Sets in homotopy type theory. *Mathematical Structures in Computer Science*, 25(5):1172–1202, 2015.
- [44] P. Sabelli. *Around the Minimalist Foundation: (co)-induction and equiconsistency*. PhD thesis, Department of Mathematics, University of Padova, 2024. to appear.
- [45] G. Sambin. Intuitionistic formal spaces - a first communication. In D. G. Skordev, editor, *Mathematical logic and its Applications*, pages 187–204. Springer, 1987.
- [46] G. Sambin. Two applications of dynamic constructivism: Brouwer’s continuity principle and choice sequences in formal topology. In M. B. Mark Van Atten, Pascal Boldini and G. Heinzmann, editors, *One Hundred Years of Intuitionism (1907–2007) The Cerisy Conference*, pages 301–315. Springer, 2008.
- [47] G. Sambin. *Positive Topology: A New Practice in Constructive Mathematics*. Oxford University Press, 2024.
- [48] S. G. Simpson. *Subsystems of Second Order Arithmetic*, volume 1. Cambridge University Press, 2009.
- [49] C. Spector. Provably recursive functionals of analysis: a consistency proof of analysis by an extension of principles formulated in current intuitionistic mathematics. In F. Dekker, editor, *Recursive function theory: Proceedings of Symposia in Pure Mathematics*, volume 5, pages 1–27. American Mathematical Society, 1962.
- [50] T. Streicher. Independence of the induction principle and the axiom of choice in the pure calculus of constructions. *Theoretical computer science*, 103(2):395–408, 1992.
- [51] A. S. Troelstra. *Metamathematical investigation of intuitionistic arithmetic and analysis*. Springer, 1973.
- [52] A. S. Troelstra and D. van Dalen. *Constructivism in Mathematics : An Introduction*. Studies in logic and the foundations of mathematics 121/123. North Holland, Amsterdam, 1988.
- [53] Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. <https://homotopytypetheory.org/book>, Institute for Advanced Study, 2013.
- [54] S. Valentini. Constructive characterizations of bar subsets. *Ann. Pure Appl. Logic*, 145(3):368–378, 2007.
- [55] H. Weyl. *Das Kontinuum. Kritische Untersuchungen über die Grundlagen der Analysis*. De Gruyter, Berlin, Boston, 1918.
- [56] H. Weyl. *The Continuum: a critical examination of the foundation of analysis*. Courier Corporation, 1994.