

Small families of partially shattering permutations

António Girão[†] Lukas Michel[†] Youri Tamitegama[†]

9 July 2024

Abstract

We say that a family of permutations t -shatters a set if it induces at least t distinct permutations on that set. What is the minimum number $f_k(n, t)$ of permutations of $\{1, \dots, n\}$ that t -shatter all subsets of size k ? For $t \leq 2$, $f_k(n, t) = \Theta(1)$. Spencer showed that $f_k(n, t) = \Theta(\log \log n)$ for $3 \leq t \leq k$ and $f_k(n, k!) = \Theta(\log n)$. In 1996, Füredi asked whether partial shattering with permutations must always fall into one of these three regimes. Johnson and Wickes recently settled the case $k = 3$ affirmatively and proved that $f_k(n, t) = \Theta(\log n)$ for $t > 2(k - 1)!$.

We give a surprising negative answer to the question of Füredi by showing that a fourth regime exists for $k \geq 4$. We establish that $f_k(n, t) = \Theta(\sqrt{\log n})$ for certain values of t and prove that this is the only other regime when $k = 4$. We also show that $f_k(n, t) = \Theta(\log n)$ for $t > 2^{k-1}$. This greatly narrows the range of t for which the asymptotic behaviour of $f_k(n, t)$ is unknown.

1 Introduction

A family \mathcal{P} of permutations of $[n] = \{1, \dots, n\}$ *shatters* a set $X \subseteq [n]$ if the permutations of \mathcal{P} induce every possible permutation on the elements of X . Shattering families of permutations were first studied by Spencer [12] who asked the following question.

What is the smallest family of permutations of $[n]$ that shatters all subsets of a fixed size k ?

Spencer [12] showed that such families have size $\Theta(\log n)$, with subsequent work improving the constant of the lower bound [6, 5, 9].

A natural refinement of this problem is to consider partial shattering. For $t \geq 1$, we say that a family \mathcal{P} *t -shatters* a set X if \mathcal{P} induces at least t distinct permutations on X . Let $f_k(n, t)$ be the minimum number of permutations of $[n]$ that t -shatter all subsets of size k . From above we know that $f_k(n, k!) = \Theta(\log n)$, and monotone permutations can be used to prove that $f_k(n, t) = t$ for $t \leq 2$. Moreover, an argument of Hajnal

[†]Mathematical Institute, University of Oxford, United Kingdom ([{girao,michel,tamitegama}@maths.ox.ac.uk](mailto:{girao,michel,tamitegama}@maths.ox.ac.uk})).

and Spencer [12] shows that $f_k(n, t) = \Theta(\log \log n)$ for $3 \leq t \leq k$.¹ Therefore, the asymptotic behaviour of $f_k(n, t)$ falls into at least three distinct regimes.

In 1996, Füredi [5] asked whether these might be the only possible regimes, even in a much more general version of partial shattering. Let \mathcal{S} be a family of sets of permutations of $[k]$. Then, a family \mathcal{P} of permutations of $[n]$ is *\mathcal{S} -mixing* if for every subset $X \subseteq [n]$ of size k , the set of permutations that \mathcal{P} induces on X is a member of \mathcal{S} . Moreover, \mathcal{S} is *monotone* if for all $S \in \mathcal{S}$ and $S \subseteq T$ we have $T \in \mathcal{S}$. If \mathcal{S} is the family of sets with at least t permutations of $[k]$, then \mathcal{S} is monotone and \mathcal{S} -mixing families are exactly those families that t -shatter all subsets of size k .

Even in this more general \mathcal{S} -mixing framework, the minimum size of an \mathcal{S} -mixing family in all previously known cases was in one of the three regimes $\Theta(1)$, $\Theta(\log \log n)$, and $\Theta(\log n)$. This prompted Füredi [5] to ask the following question.

Question 1.1 (Füredi, 1996). *If \mathcal{S} is a monotone family of sets of permutations of $[k]$, is the minimum size of an \mathcal{S} -mixing family either $\Theta(1)$, $\Theta(\log \log n)$, or $\Theta(\log n)$?*

Johnson and Wickes [7] recently made progress on this question for $f_k(n, t)$. They showed that $f_k(n, t) = \Theta(\log n)$ for $t > 2(k-1)!$. Together with the previously known asymptotics, this yields the following partial classification.

Theorem 1.2 (Johnson, Wickes, 2023). *Let $k \geq 3$ be an integer. Then,*

$$f_k(n, t) = \begin{cases} t & \text{for } t \leq 2, \\ \Theta(\log \log n) & \text{for } 3 \leq t \leq k, \\ \Theta(\log n) & \text{for } 2(k-1)! < t \leq k!. \end{cases}$$

Moreover, Johnson and Wickes [7] settled the case $k = 3$ completely by additionally proving that $f_3(n, 4) = \Theta(\log \log n)$. Given these results, they reiterated Füredi's question and asked specifically whether $f_k(n, t)$ must always fall into one of the three regimes $\Theta(1)$, $\Theta(\log \log n)$, and $\Theta(\log n)$ [1, 7].

We answer the questions of Füredi and of Johnson and Wickes negatively. For $k \geq 4$, we show that a fourth regime exists with $f_k(n, t) = \Theta(\sqrt{\log n})$. More generally, we improve the partial classification of the asymptotic behaviour of $f_k(n, t)$ as follows.

Theorem 1.3. *Let $k \geq 4$ be an integer. Then,*

$$f_k(n, t) = \begin{cases} t & \text{for } t \leq 2, \\ \Theta(\log \log n) & \text{for } 3 \leq t \leq 2^{\lceil \log_2 k \rceil}, \\ \Theta(\sqrt{\log n}) & \text{for } 2^{\lceil \log_2 k \rceil} < t \leq \min\{2k, 2^{\lceil \log_2 k \rceil} + 4\}, \\ \Theta(\log n) & \text{for } 2^{k-1} < t \leq k!. \end{cases}$$

For $k = 4$, this settles the asymptotic behaviour of $f_4(n, t)$ completely as all values of t are covered. However, if k is large, there is still an exponential gap between the regime $\Theta(\sqrt{\log n})$ and $\Theta(\log n)$.

¹Spencer [12] considered the slightly different problem of determining the minimum number of permutations such that for every subset $X \subseteq [n]$ of size k and every element $x \in X$, there is a permutation where x is the largest element of X . Johnson and Wickes [7] observed that this transfers to $f_k(n, t)$.

Theorem 1.3 is based on a series of new lower and upper bounds on $f_k(n, t)$. First, we show that the lower bound $f_k(n, t) = \Omega(\log n)$ holds for a wider range of values of t .

Theorem 1.4. *Let $k \geq 3$ and $t > 2^{k-1}$. Then, $f_k(n, t) = \Omega(\log n)$.*

The main observation for this result is that for any small family \mathcal{P} of permutations of $[n]$, we can construct two large subsets $A, B \subseteq [n]$ that are ordered in the following sense: for each permutation of \mathcal{P} either all elements of A are smaller than all elements of B , or all elements of B are smaller than all elements of A . By recursively constructing ordered sets in A , we find a subset of size k that is only 2^{k-1} -shattered by \mathcal{P} .

For smaller values of t , we provide a new lower bound of the form $\Omega(\sqrt{\log n})$.

Theorem 1.5. *Let $k \geq 3$ and $t > 2^{\lceil \log_2 k \rceil}$. Then, $f_k(n, t) = \Omega(\sqrt{\log n})$.*

The proof of this result is inspired by the proof of **Theorem 1.4**. However, instead of only constructing ordered sets in A , we recursively construct ordered sets both in A and in B . We then use a Ramsey-theoretic argument about vertex-coloured binary trees to find a subset of size k that is only $2^{\lceil \log k \rceil}$ -shattered.

We note that the lower bound on t in **Theorem 1.5** cannot be replaced by anything smaller. Indeed, a careful analysis of the construction of Hajnal and Spencer [12] shows that $f_k(n, t) = \Theta(\log \log n)$ for $3 \leq t \leq 2^{\lceil \log_2 k \rceil}$. We provide an equivalent construction which proves this and which serves as a motivating example for what follows.

Theorem 1.6. *Let $k \geq 3$ and $t \leq 2^{\lceil \log_2 k \rceil}$. Then, $f_k(n, t) = \mathcal{O}(\log \log n)$.*

Our construction identifies $[n]$ with $[2]^d$ for $d = \log_2 n$. Then, we consider lexicographic permutations of $[2]^d$ which are permutations where the order of $x, y \in [2]^d$ only depends on the values x_i and y_i for the first position i with $x_i \neq y_i$. We show that an appropriate choice of these permutations ensures that all subsets of size k are $2^{\lceil \log_2 k \rceil}$ -shattered.

Finally, for $k \geq 4$, we show that there exist $t > 2^{\lceil \log_2 k \rceil}$ with $f_k(n, t) = \mathcal{O}(\sqrt{\log n})$. Together with **Theorem 1.5**, this establishes the existence of a fourth regime for $f_k(n, t)$.

Theorem 1.7. *Let $k \geq 4$ and $t \leq \min\{2k, 2^{\lceil \log_2 k \rceil} + 4\}$. Then, $f_k(n, t) = \mathcal{O}(\sqrt{\log n})$.*

This result is proved similar to **Theorem 1.6**, but we identify $[n]$ with $[2^d]^d$ for $d = \sqrt{\log_2 n}$. Then, most subsets of size k are $2k$ -shattered by lexicographic permutations, and the remaining subsets have a very specific structure. We exploit this structure and add a few more permutations which ensure that all subsets of size k are t -shattered.

We remark that partial shattering with permutations is quite different to partial shattering with sets. A family \mathcal{F} of subsets of $[n]$ *shatters* a set $X \subseteq [n]$ if for every subset $Y \subseteq X$ there exists $F \in \mathcal{F}$ with $F \cap X = Y$, and \mathcal{F} *t -shatters* X if for at least t distinct subsets $Y \subseteq X$ there exists $F \in \mathcal{F}$ with $F \cap X = Y$. The study of shattering families of sets dates back to the seminal works of Sauer [10], Shelah [11], and Vapnik and Chervonenkis [13].

As for permutations, Kleitman and Spencer [8] showed that the minimum number of subsets of $[n]$ that shatter all subsets of size k is $\Theta(\log n)$. However, in the case of partial shattering, the family $\{\emptyset, [n]\}$ 2-shatters all subsets of size k , and every family that

3-shatters all subsets of size k already needs $\Omega(\log n)$ sets.² Therefore, partial shattering with sets only has the two regimes $\Theta(1)$ and $\Theta(\log n)$.

The rest of the paper is structured as follows. In [Section 2](#) we prove the lower bounds of [Theorems 1.4](#) and [1.5](#). Afterwards, in [Section 3](#), we prove the upper bounds of [Theorems 1.6](#) and [1.7](#). We finish with some open problems in [Section 4](#).

Notation. Throughout the paper, a permutation ρ of a set X is a total order of the elements of X . We denote this order by $<_\rho$. Note that if $Y \subseteq X$, then ρ induces a permutation on Y . If ρ is a permutation of $A \cup B$, we write $A <_\rho B$ if for all $a \in A$ and $b \in B$ we have $a <_\rho b$.

2 Lower bounds

To prove lower bounds for $f_k(n, t)$, we first need to define the concept of ordered sets. Let \mathcal{P} be a family of permutations of a set X . Then, we say that a pair of disjoint subsets $A, B \subseteq X$ is *\mathcal{P} -ordered* if for each permutation $\rho \in \mathcal{P}$ either $A <_\rho B$ or $B <_\rho A$. The following result shows that any set contains large \mathcal{P} -ordered subsets.

Lemma 2.1. *Let X be a set and let \mathcal{P} be a family of m permutations of X . Then, there exists a \mathcal{P} -ordered pair $A, B \subseteq X$ with $\min\{|A|, |B|\} \geq \lfloor |X|/2^{m+1} \rfloor$.*

Proof. Let $\mathcal{P} = \{\rho_1, \rho_2, \dots, \rho_m\}$ and define $\mathcal{P}_i = \{\rho_1, \rho_2, \dots, \rho_i\}$. We inductively construct a \mathcal{P}_i -ordered pair $A_i, B_i \subseteq X$ with $|A_i| = |B_i| \geq \lfloor |X|/2^{i+1} \rfloor$. For $i = 0$, we choose $A_0, B_0 \subseteq X$ as two disjoint subsets of size $\lfloor |X|/2 \rfloor$.

Now suppose we have constructed A_i and B_i . Let $\ell = |A_i| = |B_i| \geq \lfloor |X|/2^{i+1} \rfloor$ and consider the subset L of the ℓ largest elements of $A_i \cup B_i$ under $<_{\rho_{i+1}}$. Note that either $|L \cap A_i| \geq \ell/2$ or $|L \cap B_i| \geq \ell/2$. Without loss of generality, assume that $|L \cap A_i| \geq \ell/2$.

Set $A_{i+1} = L \cap A_i$ and $B_{i+1} = B_i \setminus L$. Since $b <_{\rho_{i+1}} a$ for all $a \in A_{i+1}$ and $b \in B_{i+1}$, the pair (A_{i+1}, B_{i+1}) is \mathcal{P}_{i+1} -ordered. We also have $|A_{i+1}| \geq \ell/2 \geq \lfloor |X|/2^{i+2} \rfloor$ and $|B_{i+1}| = \ell - |L \cap B_i| = \ell - (\ell - |L \cap A_i|) = |A_{i+1}|$, as required.

For $i = m$, the pair (A_m, B_m) is \mathcal{P} -ordered with $\min\{|A_m|, |B_m|\} \geq \lfloor |X|/2^{m+1} \rfloor$. \square

Using \mathcal{P} -ordered pairs, we can prove [Theorem 1.4](#). For this, we construct a sequence of nested \mathcal{P} -ordered pairs $A_i, B_i \subseteq A_{i-1}$ for $i \in [k]$, where $A_0 = [n]$. Then, if we pick an element $x_i \in B_i$ for each $i \in [k]$, the fact that all pairs are \mathcal{P} -ordered implies that $\{x_1, \dots, x_k\}$ is only 2^{k-1} -shattered by \mathcal{P} , as required.

Proof of Theorem 1.4. Let $t > 2^{k-1}$. We claim that $f_k(n, t) > (\log_2 n)/k - 1$. Indeed, suppose that \mathcal{P} is a family of m permutations of $[n]$ with $m \leq (\log_2 n)/k - 1$, and so $n \geq 2^{k(m+1)}$. We show that there is a subset of size k that is not t -shattered by \mathcal{P} .

²If $n \geq k \cdot 2^{|\mathcal{F}|}$, one of the intersections $\bigcap_{F \in \mathcal{F}} A_F$ with $A_F \in \{F, [n] \setminus F\}$ must have size at least k , and any subset $X \subseteq \bigcap_{F \in \mathcal{F}} A_F$ of size k satisfies $F \cap X \in \{\emptyset, X\}$ for all $F \in \mathcal{F}$.

To do so, we inductively construct a sequence of \mathcal{P} -ordered pairs as follows. Start with $A_0 = [n]$. Then, for $i = 1, \dots, k$, apply [Lemma 2.1](#) to A_{i-1} to obtain a \mathcal{P} -ordered pair $A_i, B_i \subseteq A_{i-1}$ with $\min\{|A_i|, |B_i|\} \geq \lfloor |A_{i-1}|/2^{m+1} \rfloor$. Note that this implies that $\min\{|A_i|, |B_i|\} \geq \lfloor n/2^{i(m+1)} \rfloor \geq \lfloor n/2^{k(m+1)} \rfloor \geq 1$, and so A_i and B_i are non-empty.

For each $i \in [k]$, pick $x_i \in B_i$. We claim that \mathcal{P} does not t -shatter the set $\{x_1, \dots, x_k\}$. Indeed, let $i \in [k-1]$. Note that $x_j \in A_i$ for all $j > i$. Since (A_i, B_i) is \mathcal{P} -ordered, it follows that for each permutation $\rho \in \mathcal{P}$ we either have $x_i <_\rho \{x_{i+1}, \dots, x_k\}$ or $\{x_{i+1}, \dots, x_k\} <_\rho x_i$. This provides two choices for the position of x_i relative to $\{x_{i+1}, \dots, x_k\}$. Moreover, given such a choice for each $i \in [k-1]$, this uniquely determines the permutation that ρ induces on $\{x_1, \dots, x_k\}$. Hence, \mathcal{P} induces at most 2^{k-1} permutations on $\{x_1, \dots, x_k\}$, and therefore does not t -shatter $\{x_1, \dots, x_k\}$. \square

To prove [Theorem 1.5](#), we will construct a complete binary tree of nested \mathcal{P} -ordered pairs. Namely, for each \mathcal{P} -ordered pair (A, B) that we obtain, we recursively construct new \mathcal{P} -ordered pairs in each of A and B .

This is useful for the following reason. Suppose that $k = 4$, (A, B) is a \mathcal{P} -ordered pair, and two \mathcal{P} -ordered pairs $A', B' \subseteq A$ and $A'', B'' \subseteq B$ in the tree are *synchronised* in the sense that for all $\rho \in \mathcal{P}$ we have $A' <_\rho B'$ if and only if $A'' <_\rho B''$. Pick $x_1 \in A', x_2 \in B', x_3 \in A'',$ and $x_4 \in B''$. Then, since (A, B) is \mathcal{P} -ordered, we have $\{x_1, x_2\} <_\rho \{x_3, x_4\}$ or $\{x_3, x_4\} <_\rho \{x_1, x_2\}$ for all $\rho \in \mathcal{P}$, and since (A', B') and (A'', B'') are synchronised, we have $x_1 <_\rho x_2$ if and only if $x_3 <_\rho x_4$. This implies that \mathcal{P} induces at most 4 permutations on $\{x_1, x_2, x_3, x_4\}$ which is less than $2^{k-1} = 8$.

To find synchronised \mathcal{P} -ordered pairs, we use a Ramsey-theoretic argument about vertex-coloured binary trees. For that, we need to introduce some terminology. A *binary tree* T rooted at a vertex r is a tree where r has degree 0 or 2 and every other vertex has degree 1 or 3. A leaf of T is a vertex of degree at most 1. The *layer* of T at height h is the set $N_h(r) \subseteq V(T)$ of those vertices at graph distance exactly h from r . For a vertex $v \in N_h(r)$, its children are its neighbours in $N_{h+1}(r)$, and if $v \neq r$ then its parent is its neighbour in $N_{h-1}(r)$. Clearly, $|N_h(r)| \leq 2^h$. If there exists some h such that $|N_h(r)| = 2^h$ and $|N_{h+1}(r)| = 0$, then T is a *complete binary tree* of height h .

Let S be a second binary tree with root q . A *subdivision* of S in T is an injective map $\varphi: V(S) \rightarrow V(T)$ such that for all vertices $v \in V(S) \setminus \{q\}$ with parent p , $\varphi(v)$ is contained in the subtree of T rooted at $\varphi(p)$. If the vertices of T are coloured (not necessarily properly), we say that the layers of the subdivision are *monochromatic* if $\varphi(N_h(q))$ is monochromatic for all h . We often identify $V(S)$ with $\varphi(V(S))$.

In our proof, T will be a complete binary tree whose vertices correspond to nested \mathcal{P} -ordered pairs, and we colour the vertices of T in such a way that monochromatic layers correspond to collections of synchronised \mathcal{P} -ordered pairs. Then, if we find a subdivision of a complete binary tree of height $\lceil \log_2 k \rceil$ in T with monochromatic layers, we can pick one element from each of its leaves to obtain a set of size k that is only $2^{\lceil \log_2 k \rceil}$ -shattered by \mathcal{P} , as required.

To do this, we need to find such subdivisions in large vertex-coloured complete binary trees. For all integers $c \geq 1$ and $h \geq 0$, let $g(c, h)$ be the smallest integer such that

every complete binary tree of height $g(c, h)$ whose vertices are coloured with c colours contains a subdivision of a complete binary tree of height h with monochromatic layers. We prove the following upper bound on $g(c, h)$.

Lemma 2.2. *For all integers $c \geq 1$ and $h \geq 0$ we have $g(c, h) \leq h \lceil \log_2(c^{h-1} + 1) \rceil$.*

Proof. We proceed by induction on h . If $h = 0$, a complete binary tree of height h is a single vertex, and so $g(c, 0) = 0$.

If $h \geq 1$, let $d = \lceil \log_2(c^{h-1} + 1) \rceil$ and let T be a complete binary tree with root r and height $d + g(c, h - 1)$ that is coloured with c colours. For each vertex $v \in N_d(r)$, the subtree of T rooted at v has height $g(c, h - 1)$ and must therefore contain a subdivision S_v of a complete binary tree of height $h - 1$ with monochromatic layers. Since such subdivisions admit at most c^{h-1} valid colourings and $|N_d(r)| = 2^d \geq c^{h-1} + 1$, there must be two subdivisions S_u and S_v whose colourings coincide. Then, the subdivision formed by S_u , S_v , and the lowest common ancestor of u and v is a subdivision of a complete binary tree of height h with monochromatic layers.

Therefore, we get that

$$g(c, h) \leq d + g(c, h - 1) \leq d + (h - 1) \lceil \log_2(c^{h-2} + 1) \rceil \leq h \lceil \log_2(c^{h-1} + 1) \rceil. \quad \square$$

Using the strategy outlined above, we now prove our lower bound for $t > 2^{\lceil \log_2 k \rceil}$.

Proof of Theorem 1.5. Let $h = \lceil \log_2 k \rceil$ and $t > 2^h$. We claim $f_k(n, t) > \sqrt{\log_2 n}/h - 1$. Indeed, suppose that \mathcal{P} is a family of m permutations of $[n]$ with $m \leq \sqrt{\log_2 n}/h - 1$, and so using Lemma 2.2 we get $\log_2 n \geq h^2(m + 1)^2 \geq g(2^m, h)(m + 1)$. We show that there is a subset of size k that is not t -shattered by \mathcal{P} .

Consider a complete binary tree T of height $g(2^m, h)$. We associate to each vertex v of T a set $X_v \subseteq [n]$ and a \mathcal{P} -ordered pair $A_v, B_v \subseteq X_v$ as follows. For the root r of T , let $X_r = [n]$ and let $A_r, B_r \subseteq X_r$ be a \mathcal{P} -ordered pair given by Lemma 2.1. Then, for a vertex v of T with \mathcal{P} -ordered pair (A_v, B_v) and children v_1 and v_2 , let $X_{v_1} = A_v$ and $X_{v_2} = B_v$, and let $A_{v_1}, B_{v_1} \subseteq X_{v_1}$ and $A_{v_2}, B_{v_2} \subseteq X_{v_2}$ be \mathcal{P} -ordered pairs given by Lemma 2.1. This implies that $|X_w| \geq \lfloor n/2^{g(2^m, h)(m+1)} \rfloor \geq 1$ for every $w \in V(T)$, and so X_w is non-empty.

Colour each vertex $v \in V(T)$ with a colour c_v as follows. For $\rho \in \mathcal{P}$, let

$$c_v(\rho) = \begin{cases} 1 & \text{if } A_v <_\rho B_v, \\ 0 & \text{if } B_v <_\rho A_v. \end{cases}$$

Since (A_v, B_v) is \mathcal{P} -ordered, this is well-defined. This colours each vertex of T with one of 2^m colours. By Lemma 2.2, T admits a subdivision of a complete binary tree S of height h with monochromatic layers. Let q be the root of S .

For each leaf ℓ of S , pick $y_\ell \in X_\ell$, and let $Y = \{y_\ell : \ell \in N_h^S(q)\}$ be the set of all of these elements. Note that $|Y| = 2^h \geq k$. We claim that Y is not t -shattered. Indeed, consider the colouring that a fixed permutation $\rho \in \mathcal{P}$ induces on the first h layers $N_{<h}^S(q)$ of S

by assigning $c_v(\rho)$ to each $v \in N_{<h}^S(q)$. Since each layer of S is monochromatic, this colours $N_{<h}^S(q)$ with one of at most 2^h distinct colourings (across all $\rho \in \mathcal{P}$).

Now, for any two distinct leaves $\ell, \ell' \in N_h^S(q)$, the lowest common ancestor $v \in N_{<h}^S(q)$ of ℓ and ℓ' satisfies $y_\ell \in A_v$ and $y_{\ell'} \in B_v$ (or vice versa). Then, if $c_v(\rho) = 1$, we have $A_v <_\rho B_v$ and so $y_\ell <_\rho y_{\ell'}$, and if $c_v(\rho) = 0$ we have $B_v <_\rho A_v$ and so $y_{\ell'} <_\rho y_\ell$. Thus, the colours $c_v(\rho)$ for the vertices $v \in N_{<h}^S(q)$ uniquely determine the permutation that ρ induces on Y . Since there are at most 2^h distinct colourings of $N_{<h}^S(q)$, it follows that \mathcal{P} induces at most 2^h permutations on Y , and therefore does not t -shatter Y . \square

3 Upper bounds

To prove our upper bounds on $f_k(n, t)$, we use lexicographic permutations. Identify $[n]$ with a subset of $[b]^d$ where $b \geq 2$ and $d \geq \lceil \log_b n \rceil$ and define a *lex-permutation* ρ of $[b]^d$ to be a permutation that is constructed as follows. For each $i \in [d]$, pick a permutation ρ_i of $[b]$. Then, for distinct $x, y \in [b]^d$, set $x <_\rho y$ if and only if $x_i <_{\rho_i} y_i$ where $i = i(x, y) = \min\{i \in [d] : x_i \neq y_i\}$. That is, the relative order of x and y in ρ is entirely determined by the first position where x and y differ. It is easy to check that $<_\rho$ is a total order, and so ρ is a permutation.

We say that a family of permutations \mathcal{P} of $[b]^d$ is *k-lex-shattering* if it is a family of lex-permutations such that the following holds. Let $\mathcal{I} \subseteq [d]$ be a subset of size at most k , and for each $i \in \mathcal{I}$ fix a permutation σ_i of a subset $Y_i \subseteq [b]$ of size at most k . Then, there exists a permutation $\rho \in \mathcal{P}$ such that ρ_i induces σ_i on Y_i for all $i \in \mathcal{I}$. We will later show that k -lex-shattering families $2^{\lceil \log_2 k \rceil}$ -shatter every subset of size k . However, we first have to show that small k -lex-shattering families exist.

Lemma 3.1. *For $k \geq 1$, there exists a k -lex-shattering family of $[b]^d$ with size $\mathcal{O}(\log(bd))$.*

Proof. By [Theorem 1.2](#), there is a family \mathcal{Q} of permutations of $[bd]$ with size $\mathcal{O}(\log(bd))$ that shatters every subset of size k^2 . For each $\tau \in \mathcal{Q}$, construct a lex-permutation ρ^τ such that ρ_i^τ is the permutation that τ induces on $(i-1)b + [b] = \{(i-1)b + 1, \dots, ib\}$. We claim that $\mathcal{P} = \{\rho^\tau : \tau \in \mathcal{Q}\}$ is k -lex-shattering.

Indeed, let $\mathcal{I} \subseteq [d]$ have size at most k , and for every $i \in \mathcal{I}$ let σ_i be a permutation of a subset $Y_i \subseteq [b]$ of size at most k . Since \mathcal{Q} shatters the set $\bigcup_{i \in \mathcal{I}} ((i-1)b + Y_i)$, there exists a permutation $\tau \in \mathcal{Q}$ which induces σ_i on $(i-1)b + Y_i$ for all $i \in \mathcal{I}$. This implies that ρ_i^τ induces σ_i on Y_i for all $i \in \mathcal{I}$. \square

To determine the permutations that a k -lex-shattering family \mathcal{P} induces on a subset X , consider the lexicographic structure of X . Let $i(X) = \inf\{i(x, y) : x, y \in X\}$ be the first position where two elements of X differ, and let $S(X) = \{x_{i(X)} : x \in X\}$ denote the *slice* of X at that position if $i(X) < \infty$. This partitions X into the subsets $X_s = \{x \in X : x_{i(X)} = s\}$ for $s \in S(X)$. Note that the sets X_s are pairwise \mathcal{P} -ordered and that their relative order in a permutation $\rho \in \mathcal{P}$ is determined by the permutation that $\rho_{i(X)}$ induces on $S(X)$. We will denote this permutation by $\rho(X)$.

If we partition a set X_s in the same way, this yields a partition of X_s into \mathcal{P} -ordered sets whose relative order is determined by $\rho(X_s)$. Continuing recursively like this, it follows that the order that ρ induces on X is determined by the permutations $\rho(Y)$ for $Y \subseteq X$. This is captured by the following lemma.

Lemma 3.2. *Let ρ be a lex-permutation of $[b]^d$ and let $X \subseteq [b]^d$. Then, the permutation that ρ induces on X is uniquely determined by the set of permutations $\rho(Y)$ for $Y \subseteq X$.*

Proof. Let $x, y \in X$ and $i = i(x, y)$. Then, $Y = \{x, y\} \subseteq X$ and $\rho(Y)$ is the permutation that ρ_i induces on $S(Y) = \{x_i, y_i\}$. This implies that $x <_\rho y$ if and only if $x_i <_{\rho_i} y_i$ which in turn is equivalent to $x_i <_{\rho(Y)} y_i$. So, the permutation that ρ induces on X is determined by the set of permutations $\rho(Y)$ for $Y \subseteq X$.

Conversely, suppose that ρ and τ are lex-permutations with $\rho(Y) \neq \tau(Y)$ for some $Y \subseteq X$. Let $i = i(Y)$. Since $\rho(Y) \neq \tau(Y)$, there must exist $x, y \in Y$ with $i(x, y) = i$ such that $x_i <_{\rho(Y)} y_i$ and $y_i <_{\tau(Y)} x_i$ (or vice versa). As above, this is equivalent to $x <_\rho y$ and $y <_\tau x$, and so ρ and τ induce different permutations on X . \square

Next, we deduce a lower bound on the number of permutations that \mathcal{P} induces on X . Let $\mathcal{I}(X) = \{i(Y) : Y \subseteq X \text{ and } i(Y) < \infty\}$. We rely on the following observation.

Lemma 3.3. *Let \mathcal{P} be a k -lex-shattering family of permutations of $[b]^d$ and let $X \subseteq [b]^d$ be a subset of size k . For each $i \in \mathcal{I}(X)$, let $Y_i \subseteq X$ be such that $i(Y_i) = i$. Then, X is $(\prod_{i \in \mathcal{I}(X)} |S(Y_i)|!)$ -shattered by \mathcal{P} .*

Proof. We claim that $|\mathcal{I}(X)| < k$. Indeed, write $<$ for the standard lexicographic order on $[b]^d$, so $x < y$ if and only if $x_i < y_i$ where $i = i(x, y)$. Note that if $x, y, z \in [b]^d$ satisfy $x < y < z$, then $i(x, z) = \min\{i(x, y), i(y, z)\}$. Therefore, if we write $X = \{x_1, \dots, x_k\}$ with $x_1 < \dots < x_k$, this implies that $\mathcal{I}(X) = \{i(x_\ell, x_{\ell+1}) : \ell \in [k-1]\}$, and so $|\mathcal{I}(X)| \leq k-1$ as claimed.

Since \mathcal{P} is k -lex-shattering, it follows that if σ_i is a permutation of $S(Y_i)$ for every $i \in \mathcal{I}(X)$, then there is a permutation $\rho \in \mathcal{P}$ with $\rho(Y_i) = \sigma_i$ for all $i \in \mathcal{I}(X)$. By [Lemma 3.2](#), each choice of the permutations σ_i induces a different permutation on X , and so X is $(\prod_{i \in \mathcal{I}(X)} |S(Y_i)|!)$ -shattered by \mathcal{P} . \square

To maximise this product, consider again the decomposition of X into its lexicographic structure from above. Suppose that we only pick the largest set X_s whenever we recursively continue the decomposition, and let the sets obtained during this process be the sets Y_i . Then, we will show that $\prod_{i \in \mathcal{I}} |S(Y_i)|! \geq 2^{\lceil \log_2 k \rceil}$, and so every set of size k is $2^{\lceil \log_2 k \rceil}$ -shattered by \mathcal{P} . This suffices to prove [Theorem 1.6](#).

For [Theorem 1.7](#), we will additionally infer some structural information about X whenever X is not $2k$ -shattered by \mathcal{P} . Later, this information will allow us to add more permutations to \mathcal{P} which ensure that X is t -shattered for some $t > 2^{\lceil \log_2 k \rceil}$. Overall, we obtain the following lemma.

Lemma 3.4. *Let \mathcal{P} be a k -lex-shattering family of permutations of $[b]^d$, and let $h = \lceil \log_2 k \rceil$. Then, \mathcal{P} is a family that 2^h -shatters all subsets of $[b]^d$ of size k . Moreover, if $X \subseteq [b]^d$ is a subset*

of size k that \mathcal{P} does not $2k$ -shatter, then the set $\mathcal{I}(X)$ has size h , all $Y \subseteq X$ with $i(Y) < \infty$ satisfy $|S(Y)| = 2$, and all $Y, Z \subseteq X$ with $i(Y) = i(Z) < \infty$ satisfy $S(Y) = S(Z)$.

Proof. Let $X \subseteq [b]^d$ be a subset of size k and let $\mathcal{I} = \mathcal{I}(X)$.

Claim 3.5. *There exist sets $Y_i \subseteq X$ with $i(Y_i) = i$ for each $i \in \mathcal{I}$ such that $\prod_{i \in \mathcal{I}} |S(Y_i)| \geq k$.*

Proof. We prove by induction on j that for all $0 \leq j \leq d$ there exist sets $Y_i \subseteq X$ with $i(Y_i) = i$ for each $i \in \mathcal{I} \cap [j]$ and a subset $Z \subseteq X$ which satisfy $i(Z) > j$ and $|Z| \cdot \prod_{i \in \mathcal{I} \cap [j]} |S(Y_i)| \geq k$. Applying this with $j = d$ proves the claim.

For $j = 0$, let $Z = X$. Now suppose we have constructed such sets for $j - 1$. If $i(Z) > j$, the same sets work for j , where we choose any $Y_j \subseteq X$ with $i(Y_j) = j$ if $j \in \mathcal{I}$.

Otherwise, $i(Z) = j$. Let $Y_j = Z$, and for every $s \in S(Z)$ let $Z_s = \{z \in Z : z_j = s\}$. Pick $s \in S(Z)$ such that $|Z_s| \geq |Z|/|S(Z)|$. Note that $i(Z_s) > j$ and $|Z_s| \cdot \prod_{i \in \mathcal{I} \cap [j]} |S(Y_i)| \geq (|Z|/|S(Z)|) \cdot \prod_{i \in \mathcal{I} \cap [j-1]} |S(Y_i)| \cdot |S(Z)| \geq k$. So we have constructed the sets for j . ■

For all $i \in \mathcal{I}$, let $Y_i \subseteq X$ be a set with $i(Y_i) = i$ that maximises $|S(Y_i)|$. By **Claim 3.5**, $\prod_{i \in \mathcal{I}} |S(Y_i)| \geq k$, and by **Lemma 3.3** \mathcal{P} induces at least $\prod_{i \in \mathcal{I}} |S(Y_i)|!$ permutations on X . If $|S(Y_i)| \geq 3$ for some $i \in \mathcal{I}$, then $\prod_{i \in \mathcal{I}} |S(Y_i)|! \geq 2 \prod_{i \in \mathcal{I}} |S(Y_i)| \geq 2k \geq 2^h$. Otherwise, $|S(Y_i)| = 2$ for all $i \in \mathcal{I}$. Then, since $2^{|\mathcal{I}|} = \prod_{i \in \mathcal{I}} |S(Y_i)| \geq k$, we have $|\mathcal{I}| \geq h$ and so $\prod_{i \in \mathcal{I}} |S(Y_i)|! = 2^{|\mathcal{I}|} \geq 2^h$. This shows that X is 2^h -shattered.

Moreover, if X is not $2k$ -shattered, these arguments imply that $|S(Y_i)| = 2$ for all $i \in \mathcal{I}$ and $|\mathcal{I}| = h$. Suppose that for some $j \in \mathcal{I}$ there exists a set $Z \subseteq X$ with $i(Z) = j$ and $S(Y_j) \neq S(Z)$. Then, by the maximality of Y_j , $|S(Z)| = 2$ and so $|S(Y_j) \cap S(Z)| \leq 1$. In particular, for any pair of permutations σ of $S(Y_j)$ and τ of $S(Z)$, there is a permutation σ_j of $S(Y_j) \cup S(Z)$ that induces σ on $S(Y_j)$ and τ on $S(Z)$. By the same argument as in the proof of **Lemma 3.3**, it follows that \mathcal{P} induces at least $|S(Z)|! \cdot \prod_{i \in \mathcal{I}} |S(Y_i)|! \geq 2k$ permutations on X . This contradicts the fact that X is not $2k$ -shattered. □

With $b = 2$ and $d = \lceil \log_2 n \rceil$, **Theorem 1.6** is an immediate consequence of **Lemmas 3.1** and **3.4**. We remark that while the description of our construction differs significantly from that of Spencer [12], these constructions are essentially equivalent.

To prove **Theorem 1.7**, start with a k -lex-shattering family \mathcal{P} . By **Lemma 3.4**, most subsets of size k are $2k$ -shattered by \mathcal{P} , and even if a subset X is not $2k$ -shattered, it is nevertheless $2^{\lceil \log_2 k \rceil}$ -shattered and it has a very specific structure. We exploit this structure by adding new permutations to \mathcal{P} which induce at least four additional permutation on X and thereby ensure that X is $(2^{\lceil \log_2 k \rceil} + 4)$ -shattered.

To do so, we add a constant number of new permutations for each position $i \in [d]$. Let $<$ and $>$ denote the standard and reverse permutations of $[b]$.³ Then, for all $\sigma, \tau \in \{<, >\}$, let $\pi_{i,\sigma,\tau}$ be the permutation of $[b]^d$ with $x <_{\pi_{i,\sigma,\tau}} y$ if and only if either $x_i <_{\sigma} y_i$, or

³That is, for $x, y \in [b]$ set $x << y$ if and only if $x < y$, and $x <> y$ if and only if $x > y$.

$x_i = y_i$ and $x_{i(x,y)} <_{\tau} y_{i(x,y)}$. That is, $\pi_{i,\sigma,\tau}$ first sorts according to position i and only afterwards behaves like a lex-permutation. Define

$$\mathcal{Q}_i = \{\pi_{i,\sigma,\tau} : \sigma, \tau \in \{<, >\}\}.$$

If X is not $2k$ -shattered, we show that for an appropriate $i \in [d]$ the permutations of \mathcal{Q}_i induce four additional permutations on X , as required.

Proof of Theorem 1.7. Let $b = 2^d$, $d = \lceil \sqrt{\log_2 n} \rceil$, and $h = \lceil \log_2 k \rceil$. By Lemma 3.1, there exists a k -lex-shattering family \mathcal{P} of $[b]^d$ with size $\mathcal{O}(\sqrt{\log n})$. Define $\mathcal{R} = \mathcal{P} \cup \bigcup_{i \in [d]} \mathcal{Q}_i$ and note that \mathcal{R} has size $\mathcal{O}(\sqrt{\log n})$.

We claim that every subset $X \subseteq [b]^d$ of size k is $\min\{2k, 2^h + 4\}$ -shattered by \mathcal{R} . Indeed, suppose that X is not $2k$ -shattered by \mathcal{P} . Then, by Lemma 3.4, we know that the set $\mathcal{I}(X)$ has size h , all $Y \subseteq X$ with $i(Y) < \infty$ satisfy $|S(Y)| = 2$, and all $Y, Z \subseteq X$ with $i(Y) = i(Z) < \infty$ satisfy $S(Y) = S(Z)$.

Write $X = \{x_1, \dots, x_k\}$ with $x_1 < \dots < x_k$. Note that $i(x_\ell, x_{\ell+1}) \in \mathcal{I}(X)$ for all $\ell \in [k-1]$. Since $k \geq 4$, one can check that $k-1 > h = |\mathcal{I}(X)|$. So there must exist $1 \leq \ell < m < k$ with $i(x_\ell, x_{\ell+1}) = i(x_m, x_{m+1})$. Let $i = i(x_\ell, x_{\ell+1})$ and $j = i(x_{\ell+1}, x_m)$.

If $j \geq i$, then $i(x_{\ell+1}, x_{m+1}) = i$ and $i(x_\ell, x_{m+1}) = i$. So, $Y = \{x_\ell, x_{\ell+1}, x_{m+1}\} \subseteq X$ satisfies $|S(Y)| = 3$. This contradicts the fact that all $Y \subseteq X$ with $i(Y) < \infty$ satisfy $|S(Y)| = 2$. Therefore, $j < i$. In particular, $(x_\ell)_j = (x_{\ell+1})_j < (x_m)_j = (x_{m+1})_j$, and so $(\{x_\ell, x_{\ell+1}\}, \{x_m, x_{m+1}\})$ is \mathcal{P} -ordered.

Note that $i(\{x_\ell, x_{\ell+1}\}) = i = i(\{x_m, x_{m+1}\})$. Since all $Y, Z \subseteq X$ with $i(Y) = i(Z) < \infty$ satisfy $S(Y) = S(Z)$, we must therefore have $(x_\ell)_i = (x_m)_i < (x_{\ell+1})_i = (x_{m+1})_i$, and so $(\{x_\ell, x_m\}, \{x_{\ell+1}, x_{m+1}\})$ is \mathcal{Q}_i -ordered.

In particular, \mathcal{P} and \mathcal{Q}_i induce different permutations on X . Moreover, note that $\{x_\ell, x_{\ell+1}, x_m, x_{m+1}\}$ is 4-shattered by \mathcal{Q}_i . Since Lemma 3.4 implies that X is 2^h -shattered by \mathcal{P} , it follows that X is $(2^h + 4)$ -shattered by \mathcal{R} . \square

4 Open Problems

In this paper, we have shown that at least four regimes exist for the asymptotic behaviour of $f_k(n, t)$ when $k \geq 4$, and we narrowed the range of values of t for which the asymptotic behaviour is unknown. The main open problem is to determine the asymptotic behaviour of $f_k(n, t)$ for $k \geq 5$ and $\min\{2k, 2^{\lceil \log_2 k \rceil} + 4\} < t \leq 2^{k-1}$. The following remains possible.

Question 4.1. For all integers k and t , is the asymptotic behaviour of $f_k(n, t)$ either $\Theta(1)$, $\Theta(\log \log n)$, $\Theta(\sqrt{\log n})$, or $\Theta(\log n)$?

We conjecture that Theorem 1.4 determines the entire range of values of t that satisfy $f_k(n, t) = \Theta(\log n)$.

Conjecture 4.2. Let $k \geq 3$ and $t \leq 2^{k-1}$. Then, $f_k(n, t) = o(\log n)$.

Since we did not try to optimise the upper bound on t in [Theorem 1.7](#), we believe that constructions similar to ours could prove that $f_k(n, t) = o(\log n)$ for a larger range of values of t . However, they seem to be far away from reaching $t = 2^{k-1}$.

In this context it seems important to mention the following inspiration for our approach that comes from the Erdős-Gyárfás problem [4]. If \mathcal{P} is a family of permutations of $[n]$, assign a colour $c_{x,y}$ to every pair $x, y \in [n]$ as follows. For $\rho \in \mathcal{P}$, let

$$c_{x,y}(\rho) = \begin{cases} 1 & \text{if } x <_{\rho} y, \\ 0 & \text{otherwise.} \end{cases}$$

This colouring uses $2^{|\mathcal{P}|}$ colours, and if a subset $X \subseteq [n]$ of size k spans at most ℓ colours, then \mathcal{P} induces at most 2^{ℓ} permutations on X . We implicitly used this approach to prove [Theorems 1.4](#) and [1.5](#). Unfortunately, the converse does not hold: X can simultaneously span at least ℓ colours and fail to be 2^{ℓ} -shattered by \mathcal{P} .

Nevertheless, Eichhorn and Mubayi [3] gave a colouring of K_n with $2^{\Theta(\sqrt{\log n})}$ colours such that all subsets of size k span at least $\ell = 2 \lceil \log_2 k \rceil - 2$ colours. This inspired our construction for [Theorem 1.7](#), even if our construction does not 2^{ℓ} -shatter all subsets of size k . Note that Conlon, Fox, Lee, and Sudakov [2] gave a colouring of K_n with $2^{o(\log n)}$ colours such that all subsets of size k span at least $k - 1$ colours. We hope that these colourings could inspire further progress towards better upper bounds on $f_k(n, t)$.

References

- [1] R. Baber, N. Behague, A. Calbet, D. Ellis, J. Erde, R. Gray, M.-R. Ivan, B. Janzer, R. Johnson, L. Milićević et al. *A collection of open problems in celebration of Imre Leader's 60th birthday*. arXiv preprint arXiv:2310.18163, 2023.
- [2] D. Conlon, J. Fox, C. Lee and B. Sudakov. *The Erdős–Gyárfás problem on generalized Ramsey numbers*. Proceedings of the London Mathematical Society, **110**(1):1–18, 2015.
- [3] D. Eichhorn and D. Mubayi. *Note-edge-coloring cliques with many colors on subcliques*. Combinatorica, **20**(3):441–444, 2000.
- [4] P. Erdős and A. Gyárfás. *A variant of the classical Ramsey problem*. Combinatorica, **17**:459–467, 1997.
- [5] Z. Füredi. *Scrambling permutations and entropy of hypergraphs*. Random Structures & Algorithms, **8**(2):97–104, 1996.
- [6] Y. Ishigami. *An extremal problem of d permutations containing every permutation of every t elements*. Discrete Mathematics, **159**(1-3):279–283, 1996.
- [7] J. R. Johnson and B. Wickes. *Shattering k -sets with permutations*. Order, pages 1–18, 2023.
- [8] D. J. Kleitman and J. Spencer. *Families of k -independent sets*. Discrete Mathematics, **6**(3):255–262, 1973.

- [9] J. Radhakrishnan. *A note on scrambling permutations*. *Random Structures & Algorithms*, **22**(4):435–439, 2003.
- [10] N. Sauer. *On the density of families of sets*. *Journal of Combinatorial Theory, Series A*, **13**(1):145–147, 1972.
- [11] S. Shelah. *A combinatorial problem; stability and order for models and theories in infinitary languages*. *Pacific Journal of Mathematics*, **41**(1):247–261, 1972.
- [12] J. H. Spencer. *Minimal scrambling sets of simple orders*. *Acta Mathematica Academiae Scientiarum Hungaricae*, **22**:349–353, 1972.
- [13] V. N. Vapnik and A. Y. Chervonenkis. *On the uniform convergence of relative frequencies of events to their probabilities*. In *Measures of complexity: festschrift for Alexey Chervonenkis*, pages 11–30. Springer, 2015.