

DOMAIN THEORY IN UNIVALENT FOUNDATIONS II: CONTINUOUS AND ALGEBRAIC DOMAINS

TOM DE JONG AND MARTÍN HÖTZEL ESCARDÓ

ABSTRACT. We develop the theory of continuous and algebraic domains in constructive and predicative univalent foundations, building upon our earlier work on basic domain theory in this setting. That we work predicatively means that we do not assume Voevodsky’s propositional resizing axioms. Our work is constructive in the sense that we do not rely on excluded middle or the axiom of (countable) choice. To deal with size issues and give a predicatively suitable definition of continuity of a dcpo, we follow Johnstone and Joyal’s work on continuous categories. Adhering to the univalent perspective, we explicitly distinguish between data and property. To ensure that being continuous is a property of a dcpo, we turn to the propositional truncation, although we explain that some care is needed to avoid needing the axiom of choice. We also adapt the notion of a domain-theoretic basis to the predicative setting by imposing suitable smallness conditions, analogous to the categorical concept of an accessible category. All our running examples of continuous dcpos are then actually examples of dcpos with small bases which we show to be well behaved predicatively. In particular, such dcpos are exactly those presented by small ideals. As an application of the theory, we show that Scott’s D_∞ model of the untyped λ -calculus is an example of an algebraic dcpo with a small basis. Our work is formalised in the AGDA proof assistant and its ability to infer universe levels has been invaluable for our purposes.

1. INTRODUCTION

In previous work [dJon21b; dJon24a] we developed domain theory in constructive and predicative univalent foundations and considered basic applications in the semantics of programming languages, such as the Scott model of PCF [Plo77; Sco93]. However, we did not discuss a rich and deep topic in domain theory: *algebraic* and *continuous* dcpos [Gie+03]. We present a treatment of the basic theory and examples in our constructive and predicative approach, where we deal with size issues by taking direct inspiration from category theory and the work of Johnstone and Joyal on continuous categories [JJ82] in particular.

Classically, a dcpo D is said to be *continuous* if for every element x of D the set of elements *way below* it is directed and has supremum x . The problem with this definition in our foundational setup is that the type of elements way below x is not necessarily small. Although this does not stop us from asking it to be directed and having supremum x , this still poses a problem: for example, there would be no guarantee that its supremum is preserved by a Scott continuous function, as it is only required to preserve suprema of directed families indexed by small types.

Our solution is to use the ind-completion to give a predicatively suitable definition of continuity of a dcpo, following the category theoretic work by Johnstone and Joyal [JJ82]. Some care is needed to ensure that the resulting definition expresses a property of a dcpo, rather than an equipment with additional structure. This is of course where the propositional truncation comes in useful, but there are two

Key words and phrases. constructivity, predicativity, univalent foundations, homotopy type theory, HoTT/UF, propositional resizing, domain theory, continuous domain, algebraic domain.

natural ways of using the truncation. We show that one of them yields a well behaved notion that serves as our definition of continuity, while the other, which we call *pseudocontinuity*, is problematic in a constructive context. In a classical setting where the axiom of choice is assumed, the two notions (continuity and pseudocontinuity) are equivalent.

Another approach is to turn to the notion of a *basis* [AJ94, Section 2.2.2], but to include smallness conditions. While we cannot expect the type of elements way below an element x to be small, in many examples it is the case that the type of basis elements way below x is small. We show that if a dcpo has a small basis, then it is continuous. In fact, all our running examples of continuous dcpos are actually examples of dcpos with small bases. Moreover, dcpos with small bases are better behaved. For example, they are locally small and so are their exponentials. Furthermore, we show that having a small basis is equivalent to being presented by ideals. For algebraic dcpos, bases work especially well constructively, at least in the presence of set quotients and univalence, as explained in Section 5.3. In particular, we show that Scott's D_∞ , as originally conceived in [Sco72], and ported to predicative univalent foundations in [dJon24a], is algebraic and that it has a small compact basis.

Finally, we pause to mention two other domain-theoretic aspects in this work that require particular attention when working constructively. Firstly, it is well-known that there are several inequivalent notions of a finite subset in constructive mathematics and to characterise the compact elements of a powerset we need to use the *Kuratowski* finite subsets. Secondly, *single step function* are classically defined by a case distinction (using excluded middle) on the ordering of elements. Constructively, we cannot, in general, make this case distinction, so we use subsingleton suprema to define single step functions instead.

1.1. Prerequisites. This is the second part of two papers on domain theory in univalent foundations. Part I [dJon24a] develops the concepts of dcpos, lifting, Scott continuous maps and exponentials, among others. We also rely on some examples, such as the dcpo of propositions, the powerset as a dcpo, and Scott's D_∞ , all of which we show to be algebraic dcpos in this paper. Part I further develops the material and notation for univalent foundations needed here. In particular, we write \equiv for definitional equality, reserving $=$ for identity types. We will give explicit references to appropriate sections, definitions, examples and theorems in Part I when we invoke them here, so it is advisable to have Part I available for reference.

1.2. Related work. For a general discussion of related work on (constructive and predicative) domain theory, we refer to the related work section of Part I.

Our treatment of continuous (and algebraic) dcpos is based on the work of Johnstone and Joyal [JJ82] which is situated in category theory where attention must be paid to size issues even in an impredicative setting. In the categorical context, a smallness criterion similar to our notion of having a small basis appears in [JJ82, Proposition 2.16]. In contrast to Johnstone and Joyal [JJ82], we use the propositional truncation to ensure that the type of continuous dcpos is a subtype of the type of dcpos. This, together with the related notion of pseudocontinuity, is discussed in Section 4.2. The particular case of a dcpo with a small compact basis is analogous to the notion of an accessible category [MP89].

In constructive set theory, our approach corresponds to working with partially ordered classes as opposed to sets [Acz06]. Our notion of a small basis for a dcpo (Section 5) is similar, but different from Aczel's notion of a set-generated dcpo [Acz06, Section 6.4]. While Aczel requires the set $\{b \in B \mid b \sqsubseteq x\}$ to be directed, we instead

require the set of elements in B that are way-below x to be directed in line with the usual definition of a basis [AJ94, Section 2.2.6].

Finally, abstract bases were introduced by Smyth as “R-structures” [Smy77]. Our treatment of them and the rounded ideal completion is closer to that of Abramsky and Jung in the aforementioned [AJ94, Section 2.2.6], although ours is based on families and avoids impredicative constructions.

1.3. Relation to our previous work. Along with Part I, this paper presents a revised and expanded treatment of the results in our conference paper [dJE21]. With the notable exception of Example 5.21 and Section 5.3, the results of this paper can also be found in the first author’s PhD thesis [dJon22]. Compared to the thesis, we also mention two terminological changes:

- Instead of writing “ α is cofinal in β ”, we now say that “ β exceeds α ” (see Definition 3.1). The issue with saying “cofinal” was that this word is ordinarily used for two subsets where one is already contained in the other. In particular, two cofinal subsets have the same least upper bound (if it exists) and this was not the case with our usage of the word “cofinal”.
- We have swapped “continuity structure” for “continuity data” when discussing continuous dcpos to reflect that the morphisms do not preserve the data (see also Remark 4.3). Accordingly, we no longer say that a dcpo is “structurally continuous”; instead writing that a dcpo is “equipped with continuity data”. Similar terminological changes apply to the algebraic case.

The present treatment of continuous and algebraic dcpos and small (compact) bases is significantly different from that of our earlier work [dJE21]. There, the definition of continuous dcpo was an amalgamation of pseudocontinuity and having a small basis, although it did not imply local smallness. In this work we have disentangled the two notions and based our definition of continuity on Johnstone and Joyal’s notion of a continuous category [JJ82] without making any reference to a basis. The current notion of a small basis is simpler and slightly stronger than that of our conference paper [dJE21], which allows us to prove that having a small basis is equivalent to being presented by ideals.

1.4. Formalisation. All of our results are formalised in AGDA, building on Escardó’s TYPE TOPOLOGY development [Esc+]. The reference [dJon24b] precisely links each numbered environment (including definitions, examples and remarks) in this paper to its implementation. The HTML rendering has clickable links and so is particularly suitable for exploring the development: <https://www.cs.bham.ac.uk/~mhe/TypeTopology/DomainTheory.Part-II.html>.

1.5. Organisation. The paper is organised as follows:

- Section 2: Definition and examples of the way-below relation and compact elements.
- Section 3: The ind-completion of a preorder: a tool used to discuss continuity and pseudocontinuity of dcpos.
- Section 4: Definitions of continuous and algebraic dcpos accompanied by a discussion on pseudocontinuity and issues concerning the axiom of choice.
- Section 5: The notion of a small (compact) basis: strengthening continuity (resp. algebraicity) by imposing smallness conditions.
- Section 6: The (rounded) ideal completion of an abstract basis as a continuous dcpo with a small basis.
- Section 7.1: Bilimits and exponentials of structurally continuous (or algebraic) dcpos (with small bases), including a proof that Scott’s D_∞ is algebraic with a small compact basis.

2. THE WAY-BELOW RELATION AND COMPACTNESS

The way-below relation is the fundamental ingredient in the development of continuous dcpos. Following Scott [Sco70], a computational intuition of $x \ll y$ says that every computation of y has to print x , or something better than x , at some stage. We recall the notions of a directed family, its supremum (denoted by \sqcup), and a \mathcal{V} -dcpo from Part I (Definitions 3.4, 3.6 and 3.7, respectively).

Definition 2.1 (Way-below relation, $x \ll y$). An element x of a \mathcal{V} -dcpo D is *way below* an element y of D if whenever we have a directed family $\alpha : I \rightarrow D$ indexed by $I : \mathcal{V}$ such that $y \sqsubseteq \sqcup \alpha$, then there exists $i : I$ such that $x \sqsubseteq \alpha_i$ already. We denote this situation by $x \ll y$.

Lemma 2.2. *The way-below relation enjoys the following properties:*

- (i) *it is proposition-valued;*
- (ii) *if $x \ll y$, then $x \sqsubseteq y$;*
- (iii) *if $x \sqsubseteq y \ll v \sqsubseteq w$, then $x \ll w$;*
- (iv) *it is antisymmetric;*
- (v) *it is transitive.*

Proof. (i) Using that a dependent product of propositions (over an arbitrary type) is again a proposition together with the fact that we propositionally truncated the existence of $i : I$ in the definition. (ii) Simply take $\alpha : \mathbf{1}_{\mathcal{V}} \rightarrow D$ to be $u \mapsto y$. (iii) Suppose that $\alpha : I \rightarrow D$ is directed with $w \sqsubseteq \sqcup \alpha$. Then $v \sqsubseteq \sqcup \alpha$, so by assumption that $y \ll v$ there exists $i : I$ with $y \sqsubseteq \alpha_i$ already. But then $x \sqsubseteq \alpha_i$. (iv) Follows from (ii). (v) Follows from (ii) and (iii). \square

In general, the way below relation is not reflexive. The elements for which it is have a special status and are called compact. We illustrate this notion by a series of examples.

Definition 2.3 (Compactness). An element of a dcpo is *compact* if it is way below itself.

Example 2.4. The least element of a pointed dcpo is always compact.

We recall the \mathcal{V} -dcpo of propositions $\Omega_{\mathcal{V}}$ with designated elements $\mathbf{0}_{\mathcal{V}}$ (the empty type) and $\mathbf{1}_{\mathcal{V}}$ (the unit type) from Part I (Example 3.14).

Example 2.5 (Compact elements in $\Omega_{\mathcal{V}}$). The compact elements of $\Omega_{\mathcal{V}}$ are exactly $\mathbf{0}_{\mathcal{V}}$ and $\mathbf{1}_{\mathcal{V}}$. In other words, the compact elements of $\Omega_{\mathcal{V}}$ are precisely the decidable propositions.

Proof. By Example 2.4 we know that $\mathbf{0}_{\mathcal{V}}$ must be compact. For $\mathbf{1}_{\mathcal{V}}$, suppose that we have $Q_{(-)} : I \rightarrow \Omega_{\mathcal{V}}$ directed such that $\mathbf{1}_{\mathcal{V}} \sqsubseteq \exists_{i:I} Q_i$. Then there exists $i : I$ such that Q_i holds, and hence, $\mathbf{1}_{\mathcal{V}} \sqsubseteq Q_i$. Now suppose that $P : \Omega_{\mathcal{V}}$ is compact. We show that P is decidable. The family $\alpha : (P + \mathbf{1}_{\mathcal{V}}) \rightarrow \Omega_{\mathcal{V}}$ given by $\text{inl}(p) \mapsto \mathbf{1}_{\mathcal{V}}$ and $\text{inr}(\star) \mapsto \mathbf{0}_{\mathcal{V}}$ is directed and $P \sqsubseteq \sqcup \alpha$. Hence, by compactness, there exists $i : P + \mathbf{1}_{\mathcal{V}}$ such that $P \sqsubseteq \alpha_i$ already. Since being decidable is a property of a proposition, we actually get such an i and by case distinction on it we get decidability of P . \square

We recall the lifting $\mathcal{L}_{\mathcal{V}}(X)$ of a set X as a \mathcal{V} -dcpo from Part I (Section 5), which built on [EK17; Kna18].

Example 2.6 (Compact elements in the lifting). An element (P, φ) of the lifting $\mathcal{L}_{\mathcal{V}}(X)$ of a set $X : \mathcal{V}$ is compact if and only if P is decidable. Hence, the compact elements of $\mathcal{L}_{\mathcal{V}}(X)$ are exactly \perp and $\eta(x)$ for $x : X$.

Proof. To see that compactness implies decidability of the domain of the partial element, we proceed as in the proof of Example 2.5, but for a partial element (P, φ) , we consider the family $\alpha : (P + \mathbf{1}_{\mathcal{V}}) \rightarrow \mathcal{L}_{\mathcal{V}}(X)$ given by $\text{inl}(p) \mapsto \eta(\varphi(p))$ and $\text{inr}(\star) \mapsto \perp$. Conversely, if we have a partial element (P, φ) with P decidable, then either P is false in which case $(P, \varphi) = \perp$ which is compact by Example 2.4, or P holds. So suppose that P holds and let $\alpha : I \rightarrow \mathcal{L}_{\mathcal{V}}(X)$ be directed with $P \sqsubseteq \bigsqcup \alpha$. Since P holds, the element $\bigsqcup \alpha$ must be defined, which means that there exists $i : I$ such that α_i is defined. But for this $i : I$ we also have $\bigsqcup \alpha = \alpha_i$ by construction of the supremum, and hence, $P \sqsubseteq \alpha_i$, proving compactness of (P, φ) . \square

For characterising the compact elements of the powerset, we introduce a lemma, as well as the notion of Kuratowski finiteness and the induction principle for Kuratowski finite subsets.

Lemma 2.7. *The compact elements of a dcpo are closed under (existing) binary joins.*

Proof. Suppose that x and y are compact elements of a \mathcal{V} -dcpo D with z as their least upper bound and suppose that we have $\alpha : I \rightarrow D$ directed with $z \sqsubseteq \bigsqcup \alpha$. Then $x \sqsubseteq \bigsqcup \alpha$ and $y \sqsubseteq \bigsqcup \alpha$, so by compactness there exist $i, j : I$ such that $x \sqsubseteq \alpha_i$ and $y \sqsubseteq \alpha_j$. By semidirectedness of α , there exists $k : I$ with $\alpha_i, \alpha_j \sqsubseteq \alpha_k$, so that $x, y \sqsubseteq \alpha_k$. But z is the join of x and y , so $z \sqsubseteq \alpha_k$, as desired. \square

We recall the type $\mathcal{P}_{\mathcal{T}}(X)$ of \mathcal{T} -valued subsets of X and the set-theoretic notation \in and \subseteq from Part I (Definitions 2.3 and 2.4).

Definition 2.8 (Total space of a subset, \mathbb{T}). The *total space* of a \mathcal{T} -valued subset S of a type X is defined as $\mathbb{T}(S) \equiv \Sigma_{x:X}(x \in S)$.

Definition 2.9 (Kuratowski finiteness).

- (i) A type X is *Kuratowski finite* if there exists some natural number $n : \mathbb{N}$ and a surjection $e : \text{Fin}(n) \twoheadrightarrow X$, where $\text{Fin}(n)$ is the standard finite type with exactly n elements [Rij22, Section 7.3].
- (ii) A subset is *Kuratowski finite* if its total space is a Kuratowski finite type.

Thus, a type X is *Kuratowski finite* if its elements can be finitely enumerated, possibly with repetitions, although the repetitions can be removed when X has decidable equality.

Lemma 2.10. *The Kuratowski finite subsets of a set are closed under finite unions and contain all singletons.*

Proof. The empty set and any singleton are clearly Kuratowski finite. Moreover, if A and B are Kuratowski finite subsets, then we may assume to have natural numbers n and m and surjections $\sigma : \text{Fin}(n) \twoheadrightarrow \mathbb{T}(A)$ and $\tau : \text{Fin}(m) \twoheadrightarrow \mathbb{T}(B)$. We can then patch these together to obtain a surjection $\text{Fin}(n+m) \twoheadrightarrow \mathbb{T}(A \cup B)$, as desired. \square

Lemma 2.11 (Induction for Kuratowski finite subsets). *A property of subsets of a type X holds for all Kuratowski finite subsets of X as soon as*

- (i) *it holds for the empty set,*
- (ii) *it holds for any singleton subset, and*
- (iii) *it holds for $A \cup B$, whenever it holds for A and B .*

Proof. Let Q be a such a property and let A be an arbitrary Kuratowski finite subset of X . Since Q is proposition-valued, we may assume to have a natural number n and a surjection $\sigma : \text{Fin}(n) \twoheadrightarrow \mathbb{T}(A)$. Then the subset A must be equal to the finite join of singletons $\{\sigma_0\} \cup \{\sigma_1\} \cup \dots \cup \{\sigma_{n-1}\}$, which can be shown to satisfy Q by induction on n , and hence, so must A . \square

Definition 2.12 (β). For a set $X : \mathcal{U}$, we write $\beta : \text{List}(X) \rightarrow \mathcal{P}_{\mathcal{U}}(X)$ for the canonical map which takes a list to its set of elements. (For the inductive definition of the type $\text{List}(X)$, recall e.g. [Uni13, Section 5.1].)

Lemma 2.13. *A subset $A : \mathcal{P}_{\mathcal{U}}(X)$ of a set $X : \mathcal{U}$ is Kuratowski finite if and only if it is in the image of β .*

Proof. The left to right direction follows from Lemma 2.10, while the converse follows easily from the induction principle for Kuratowski finite subsets where we use list concatenation in case (iii). \square

Example 2.14 (Compact elements in $\mathcal{P}_{\mathcal{U}}(X)$). The compact elements of $\mathcal{P}_{\mathcal{U}}(X)$ for a set $X : \mathcal{U}$ are exactly the Kuratowski finite subsets of X .

Proof. Suppose first that $A : \mathcal{P}_{\mathcal{U}}(X)$ is a compact element. The family

$$\left(\sum_{l:\text{List}(X)} \beta(l) \subseteq A\right) \xrightarrow{\beta \circ \text{pr}_1} \mathcal{P}_{\mathcal{U}}(X)$$

is directed, as it contains \emptyset and we can concatenate lists to establish semidirectedness. Moreover, $\left(\sum_{l:\text{List}(X)} \beta(l) \subseteq A\right)$ lives in \mathcal{U} and we clearly have $A \subseteq \bigsqcup \beta \circ \text{pr}_1$. So by compactness, there exists $l : \text{List}(X)$ with $\beta(l) \subseteq A$ such that $A \subseteq \beta(l)$ already. But this says exactly that A is Kuratowski finite by Lemma 2.13.

For the converse we use the induction principle for Kuratowski finite subsets: the empty set is compact by Example 2.4, singletons are easily shown to be compact, and binary unions are compact by Lemma 2.7. \square

We end this section by presenting a few lemmas connecting the way-below relation and compactness to Scott continuous sections, which we recall from Part I (Definition 4.13).

Lemma 2.15. *If we have a Scott continuous retraction $D \xrightleftharpoons[r]{s} E$, then $y \ll s(x)$ implies $r(y) \ll x$ for every $x : D$ and $y : E$.*

Proof. Suppose that $y \ll s(x)$ and that $x \sqsubseteq \bigsqcup \alpha$ for a directed family $\alpha : I \rightarrow D$. Then $s(x) \sqsubseteq s(\bigsqcup \alpha) = \bigsqcup (s \circ \alpha)$ by Scott continuity of s , so there exists $i : I$ such that $y \sqsubseteq s(\alpha_i)$ already. Now monotonicity of r implies $r(y) \sqsubseteq r(s(\alpha_i)) = \alpha_i$ which completes the proof that $r(y) \ll x$. \square

We also recall embedding-projection pairs from Part I (Definition 7.2).

Lemma 2.16. *The embedding in an embedding-projection pair $D \xrightleftharpoons[\pi]{\varepsilon} E$ preserves and reflects the way-below relation, i.e. $x \ll y \iff \varepsilon(x) \ll \varepsilon(y)$. In particular, an element x is compact if and only if $\varepsilon(x)$ is.*

Proof. Suppose that $x \ll y$ in D and let $\alpha : I \rightarrow E$ be directed with $\varepsilon(y) \sqsubseteq \bigsqcup \alpha$. Then $y = \pi(\varepsilon(y)) \sqsubseteq \bigsqcup \pi \circ \alpha$ by Scott continuity of π . Hence, there exists $i : I$ such that $x \sqsubseteq \pi(\alpha_i)$. But then $\varepsilon(x) \sqsubseteq \varepsilon(\pi(\alpha_i)) \sqsubseteq \alpha_i$ by monotonicity of ε and the fact that $\varepsilon \circ \pi$ is a deflation. This proves that $x \ll y$. Conversely, if $\varepsilon(x) \ll \varepsilon(y)$, then $x = \pi(\varepsilon(x)) \ll y$ by Lemma 2.15. \square

3. THE IND-COMPLETION

The ind-completion will be a useful tool for phrasing and proving results about directed complete posets and is itself a directed complete preorder, cf. Lemma 3.3. It was introduced by Grothendieck and Verdier in [GV72, Section 8] in the context of category theory, but its role in order theory is discussed in [JJ82, Section 1]. We will also use it in the context of order theory, but our treatment will involve a careful consideration of the universes involved, very similar to the original treatment in [GV72].

Definition 3.1 (\mathcal{V} -ind-completion $\mathcal{V}\text{-Ind}(X)$, exceed, \lesssim). The \mathcal{V} -ind-completion $\mathcal{V}\text{-Ind}(X)$ of a preorder X is the type of directed families in X indexed by types in the universe \mathcal{V} . Such a family $\beta : J \rightarrow X$ *exceeds* another family $\alpha : I \rightarrow X$ if for every $i : I$, there exists $j : J$ such that $\alpha_i \sqsubseteq \beta_j$, and we denote this relation by $\alpha \lesssim \beta$.

The following is straightforward.

Lemma 3.2. *The relation \lesssim defines a preorder on the ind-completion.*

Lemma 3.3. *The \mathcal{V} -ind-completion $\mathcal{V}\text{-Ind}(X)$ of a preorder X is \mathcal{V} -directed complete.*

Proof. Suppose that we have a directed family $\alpha : I \rightarrow \mathcal{V}\text{-Ind}(X)$ with $I : \mathcal{V}$. Then each α_i is a directed family in X indexed by a type $J_i : \mathcal{V}$. We define the family $\hat{\alpha} : (\Sigma_{i:I} J_i) \rightarrow X$ by $(i, j) \mapsto \alpha_i(j)$. It is clear that $\hat{\alpha}$ exceeds each α_i , and that β exceeds $\hat{\alpha}$ if β exceeds every α_i . So it remains to show that $\hat{\alpha}$ is in fact an element of $\mathcal{V}\text{-Ind}(X)$, i.e. that it is directed. Because α and each α_i are directed, I and each J_i are inhabited. Hence, so is the domain of $\hat{\alpha}$. It remains to show that $\hat{\alpha}$ is semidirected. Suppose we have $(i_1, j_1), (i_2, j_2)$ in the domain of $\hat{\alpha}$. By directedness of α , there exists $i : I$ such that α_i exceeds both α_{i_1} and α_{i_2} . Hence, there exist $j'_1, j'_2 : J_i$ with $\alpha_{i_1}(j_1) \sqsubseteq \alpha_i(j'_1)$ and $\alpha_{i_2}(j_2) \sqsubseteq \alpha_i(j'_2)$. Because the family α_i is directed in X , there exists $j : J_i$ such that $\alpha_i(j'_1), \alpha_i(j'_2) \sqsubseteq \alpha_i(j)$. Hence, we conclude that $\hat{\alpha}(i_1, j_1) \equiv \alpha_{i_1}(j_1) \sqsubseteq \alpha_i(j'_1) \sqsubseteq \alpha_i(j) \equiv \hat{\alpha}(i, j)$, and similarly for (i_2, j_2) , which proves semidirectedness of $\hat{\alpha}$. \square

Lemma 3.4. *Taking directed suprema defines a monotone map from a \mathcal{V} -dcpo to its \mathcal{V} -ind-completion, denoted by $\bigsqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$.*

Proof. We have to show that $\bigsqcup \alpha \sqsubseteq \bigsqcup \beta$ for directed families α and β such that β exceeds α . Note that the inequality holds as soon as $\alpha_i \sqsubseteq \bigsqcup \beta$ for every i in the domain of α . For this, it suffices that for every such i , there exists a j in the domain of β such that $\alpha_i \sqsubseteq \beta_j$. But the latter says exactly that β exceeds α . \square

Johnstone and Joyal [JJ82] generalise the notion of continuity from posets to categories and do so by phrasing it in terms of $\bigsqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$ having a left adjoint. We follow their approach and now work towards this. It turns out to be convenient to use the following two notions, which are in fact equivalent by Lemma 3.8:

Definition 3.5 (Approximate, left adjunct). For an element x of a dcpo D and a directed family $\alpha : I \rightarrow D$, we say that

- (i) α *approximates* x if the supremum of α is x and each α_i is way below x , and
- (ii) α is *left adjunct* to x if $\alpha \lesssim \beta \iff x \sqsubseteq \bigsqcup \beta$ for every directed family β .

Remark 3.6. For a \mathcal{V} -dcpo D , a function $L : D \rightarrow \mathcal{V}\text{-Ind}(D)$ is a left adjunct to $\bigsqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$ precisely when $L(x)$ is left adjunct to x for every $x : D$. Of course, we need to know that L is monotone and this is shown in the next lemma.

Lemma 3.7. *Every function $L : D \rightarrow \mathcal{V}\text{-Ind}(D)$ satisfying that $L(x)$ is left adjunct to x for every $x : D$ is monotone and order-reflecting.*

Proof. Suppose we are given elements $x, y : D$. By assumption, we know that $L(x) \lesssim L(y) \iff x \sqsubseteq \bigsqcup L(y)$, but $L(y)$ approximates y , so $\bigsqcup L(y) = y$ and hence $L(x) \lesssim L(y) \iff x \sqsubseteq y$, so L preserves and reflects the order. \square

Lemma 3.8. *A directed family α approximates an element x if and only if it is left adjunct to it.*

Proof. Suppose first that α approximates x . If $\alpha \lesssim \beta$, then $x = \bigsqcup \alpha \sqsubseteq \bigsqcup \beta$, by Lemma 3.4. Conversely, if $x \sqsubseteq \bigsqcup \beta$, then β exceeds α : for if i is in the domain of α , then $\alpha_i \ll x$, so there exists j such that $\alpha_i \sqsubseteq \beta_j$ already.

In the other direction, suppose that α is left adjunct to x . We show that each α_i is way below x . If β is a directed family with $x \sqsubseteq \bigsqcup \beta$, then β exceeds α as α is assumed to be left adjunct to x . Hence, for every i , there exists j with $\alpha_i \sqsubseteq \beta_j$, proving that $\alpha_i \ll x$. Since α exceeds itself, we get $x \sqsubseteq \bigsqcup \alpha$ by assumption. For the other inequality, we note that $x \sqsubseteq \bigsqcup \hat{x}$, where $\hat{x} : \mathbf{1} \rightarrow D$ is the directed family that maps to x . Hence, as α is left adjunct to x , we must have that \hat{x} exceeds α , which means that each α_i is below x . Thus, $\bigsqcup \alpha \sqsubseteq x$ and $x = \bigsqcup \alpha$ hold, as desired. \square

Proposition 3.9. *For a \mathcal{V} -dcpo D , a function $L : D \rightarrow \mathcal{V}\text{-Ind}(D)$ is a left adjoint to $\bigsqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$ if and only if $L(x)$ approximates x for every $x : D$.*

Proof. Immediate from Lemma 3.8 and Remark 3.6. \square

4. CONTINUOUS AND ALGEBRAIC DCPOS

Using the ind-completion from the previous section, we turn to defining continuous and algebraic dcpos, paying special attention to size and constructivity issues regarding the axiom of choice. This second issue is discussed in Section 4.2 from two perspectives: type-theoretically, via a discussion on the placement of the propositional truncation, and categorically, via left adjoints.

4.1. Continuous dcpos. We define what it means for a \mathcal{V} -dcpo to be continuous and prove the fundamental interpolation properties for the way-below relation. Examples are postponed (see Sections 5.2 and 6.2) until we have developed the theory further.

Definition 4.1 (Continuity data, I_x, α_x). *Continuity data* for a \mathcal{V} -dcpo D assigns to every $x : D$ a type $I_x : \mathcal{V}$ and a directed *approximating family* $\alpha_x : I \rightarrow D$ such that α_x has supremum x and each element $\alpha_x(i)$ is way below x .

The notion of continuity data can be understood categorically as follows.

Proposition 4.2. *The type of continuity data for a \mathcal{V} -dcpo D is equivalent to the type of left adjoints to $\bigsqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$.*

Proof. Given continuity data for D , we define a function $D \rightarrow \mathcal{V}\text{-Ind}(D)$ by sending $x : D$ to the directed family α_x . This is indeed a left adjoint to $\bigsqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$ because of Proposition 3.9. Conversely, given a left adjoint $L : D \rightarrow \mathcal{V}\text{-Ind}(D)$, the assignment $x \mapsto L(x)$ is continuity data for D , again by Proposition 3.9. That these two mappings make up a type equivalence can be checked directly, using that the type expressing that a map is a left adjoint to $\bigsqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$ is a proposition. \square

Remark 4.3. It should be noted that having continuity data is *not* a property of a dcpo, i.e., the type of continuity data for a dcpo is not a subsingleton, because an element $x : D$ can have several different approximating families, e.g. if $\alpha : I \rightarrow D$ approximates x , then so does $[\alpha, \alpha] : (I + I) \rightarrow D$. In other words, the left adjoint to $\bigsqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$ is not unique, although it is unique up to isomorphism (of the preorder $\mathcal{V}\text{-Ind}(D)$). In category theory this is typically sufficient (and often the best one can do). Johnstone and Joyal follow this philosophy in [JJ82], but we want the type of continuous \mathcal{V} -dcpos to be a subtype of the \mathcal{V} -dcpos. One reason that a property is preferred is that it is considered good mathematical practice to only consider morphisms that preserve imposed structure. In the case of continuity data, this would imply preservation of the way-below relation, but this excludes many Scott continuous maps, e.g. if the value of a constant map is not compact, then it does not preserve the way-below relation.

It is natural to ask whether the univalence axiom can be used to identify these isomorphic objects. However, this is not the case because the ind-completion $\mathcal{V}\text{-Ind}(D)$ is *not* a univalent category in the sense of [AKS15], because it is a preorder and not a poset. One way to obtain a subtype is to propositionally truncate the notion of continuity data and this is indeed the approach that we will take. However, another choice that would yield a property is to identify isomorphic elements of $\mathcal{V}\text{-Ind}(D)$. This approach is discussed at length in Section 4.2 and in particular it is explained to be inadequate in a constructive setting.

Definition 4.4 (Continuity of a dcpo). A \mathcal{V} -dcpo is *continuous* if it has unspecified continuity data, i.e. if the type of continuity data is inhabited.

Thus, a dcpo is continuous if we have an *unspecified* function assigning an approximating family to every element of the dcpo.

Proposition 4.5. *Continuity of a \mathcal{V} -dcpo D is equivalent to having an unspecified left adjoint to $\sqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$.*

Proof. By Proposition 4.2 and functoriality of the propositional truncation. \square

Lemma 4.6. *For elements x and y of a dcpo with continuity data, the following are equivalent:*

- (i) $x \sqsubseteq y$;
- (ii) $\alpha_x(i) \sqsubseteq y$ for every $i : I_x$;
- (iii) $\alpha_x(i) \ll y$ for every $i : I_x$.

Proof. Note that (iii) implies (ii) and (ii) implies (i), because if $\alpha_x(i) \sqsubseteq y$ for every $i : I_x$, then $x = \sqcup \alpha_x \sqsubseteq y$, as desired. So it remains to prove that (i) implies (iii), but this holds, because $\alpha_x(i) \ll x$ for every $i : I_x$. \square

Lemma 4.7. *For elements x and y of a dcpo with continuity data, x is way below y if and only if there exists $i : I_y$ such that $x \sqsubseteq \alpha_y(i)$.*

Proof. The left-to-right implication holds, because α_y is a directed family with supremum y , while the converse holds because $\alpha_y(i) \ll y$ for every $i : I_y$. \square

We now prove three interpolation lemmas for dcpos with continuity data. Because the conclusions of the lemmas are propositions, the results will follow for continuous dcpos immediately.

Lemma 4.8 (Nullary interpolation for the way-below relation). *For every $x : D$ of a continuous dcpo D , there exists $y : D$ such that $y \ll x$.*

Proof. The approximating family α_x is directed, so there exists $i : I_x$ and hence we can take $y := \alpha_x(i)$ since $\alpha_x(i) \ll x$. \square

Our proof of the following lemma is inspired by [JJ82, Proposition 2.12].

Lemma 4.9 (Unary interpolation for the way-below relation). *If $x \ll y$ in a continuous dcpo D , then there exists an interpolant $d : D$ such that $x \ll d \ll y$.*

Proof. Since we are proving a proposition, we may assume to be given continuity data for D . Thus, we can approximate every approximant $\alpha_y(i)$ of y by an approximating family $\beta_i : J_i \rightarrow D$. This defines a map $\hat{\beta}$ from I_y to $\mathcal{V}\text{-Ind}(D)$, the ind-completion of the \mathcal{V} -dcpo D , by sending $i : I_y$ to the directed family β_i . We claim that $\hat{\beta}$ is directed in $\mathcal{V}\text{-Ind}(D)$. Since α_y is directed, I_y is inhabited, so it remains to prove that $\hat{\beta}$ is semidirected. So suppose we have $i_1, i_2 : I_y$. Because α_y is semidirected, there exists $i : I_y$ such that $\alpha_y(i_1), \alpha_y(i_2) \sqsubseteq \alpha_y(i)$. We claim that β_i exceeds β_{i_1} and β_{i_2} , which would prove semidirectedness of $\hat{\beta}$. We give the argument for i_1 only

as the case for i_2 is completely analogous. We have to show that for every $j : J_{i_1}$, there exists $j' : J_i$ such that $\beta_{i_1}(j) \sqsubseteq \beta_i(j')$. But this holds because $\beta_{i_1}(j) \ll \bigsqcup \beta_i$ for every such j , as we have $\beta_{i_1}(j) \ll \alpha_y(i_1) \sqsubseteq \alpha_y(i) \sqsubseteq \bigsqcup \beta_i$.

Thus, $\hat{\beta}$ is directed in $\mathcal{V}\text{-Ind}(D)$ and hence we can calculate its supremum in $\mathcal{V}\text{-Ind}(D)$ to obtain the *directed* family $\gamma : (\Sigma_{i:I} J_i) \rightarrow D$ given by $(i, j) \mapsto \beta_i(j)$.

We now show that y is below the supremum of γ . By Lemma 4.6, it suffices to prove that $\alpha_y(i) \sqsubseteq \bigsqcup \gamma$ for every $i : I_y$, and, in turn, to prove this for an $i : I_y$ it suffices to prove that $\beta_i(j) \sqsubseteq \bigsqcup \gamma$ for every $j : J_i$. But this is immediate from the definition of γ . Thus, $y \sqsubseteq \bigsqcup \gamma$. Because $x \ll y$, there exists $(i, j) : \Sigma_{i:I} J_i$ such that $x \sqsubseteq \gamma(i, j) \equiv \beta_i(j)$.

Finally, for our interpolant, we take $d \equiv \alpha_y(i)$. Then, indeed, $x \ll d \ll y$, because $x \sqsubseteq \beta_i(j) \ll \alpha_y(i) \equiv d$ and $d \equiv \alpha_y(i) \ll y$, completing the proof. \square

Lemma 4.10 (Binary interpolation for the way-below relation). *If $x \ll z$ and $y \ll z$ in a continuous dcpo D , then there exists an interpolant $d : D$ such that $x, y \ll d$ and $d \ll z$.*

The proof is a straightforward application of unary interpolation.

Proof. Using that $x \ll z$ and $y \ll z$, there exist interpolants $d_1, d_2 : D$ such that $x \ll d_1 \ll z$ and $y \ll d_2 \ll z$. Hence, there exist $i_1, i_2 : I_z$ such that $d_1 \sqsubseteq \alpha_z(i_1)$ and $d_2 \sqsubseteq \alpha_z(i_2)$. By semidirectedness of α_z , there then exists $i : I_z$ for which $d_1, d_2 \sqsubseteq \alpha_z(i)$. Our final interpolant is defined as $d \equiv \alpha_z(i)$, which works because $x \ll d_1 \sqsubseteq d$, $y \ll d_2 \sqsubseteq d$ and $d \equiv \alpha_z(i) \ll z$. \square

Continuous dcpos are closed under Scott continuous retracts. Keeping track of universes, it holds in the following generality, where we recall from Part I (Definition 3.9) that we write $\mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$ for the type of \mathcal{V} -dcpos with carriers in \mathcal{U} and order relations taking values in \mathcal{T} .

Theorem 4.11. *If we have dcpos $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$ and $E : \mathcal{V}\text{-DCPO}_{\mathcal{U}', \mathcal{T}'}$ such that D is a Scott continuous retract of E , then D is continuous if E is. Moreover, we can give continuity data for D if we have such data for E .*

Proof. We prove the result in case we are given continuity data for E , as the other will follow from that and the fact that the propositional truncation is functorial. So suppose that we have a Scott continuous section $s : D \rightarrow E$ and retraction $r : E \rightarrow D$. We claim that for every $x : D$, the family $r \circ \alpha_{s(x)}$ approximates x . Firstly, it is directed, because $\alpha_{s(x)}$ is and r is Scott continuous. Secondly,

$$\begin{aligned} \bigsqcup r \circ \alpha_{s(x)} &= r(\bigsqcup \alpha_{s(x)}) && \text{(by Scott continuity of } r\text{)} \\ &= r(s(x)) && \text{(as } \alpha_{s(x)} \text{ is the approximating family of } s(x)\text{)} \\ &= x && \text{(because } s \text{ is a section of } r\text{),} \end{aligned}$$

so the supremum of $r \circ \alpha_{s(x)}$ is x . Finally, we must prove that $r(\alpha_{s(x)}(i)) \ll x$ for every $i : I_x$. By Lemma 2.15, this is implied by $\alpha_{s(x)}(i) \ll s(x)$, which holds as $\alpha_{s(x)}$ is the approximating family of $s(x)$. \square

Recall from Part I (Definition 3.12) that a \mathcal{V} -dcpo is locally small if $x \sqsubseteq y$ is equivalent to a type in \mathcal{V} for all elements x and y .

Proposition 4.12. *A continuous dcpo is locally small if and only if its way-below relation has small values.*

Proof. By Lemmas 4.6 and 4.7, we have

$$x \sqsubseteq y \iff \forall i : I_x (\alpha_x(i) \ll y) \quad \text{and} \quad x \ll y \iff \exists i : I_y (x \sqsubseteq \alpha_y(i)),$$

for every two elements x and y of a dcpo with continuity data. But the types I_x and I_y are small, finishing the proof. The result also holds for continuous dcpos, because what we are proving is a proposition. \square

Proposition 4.12 is significant because the definition of the way-below relation for a \mathcal{V} -dcpo D quantifies over all families into D indexed by types in \mathcal{V} .

4.2. Pseudocontinuity. In light of Proposition 4.2, we see that a \mathcal{V} -dcpo D can have continuity data in more than one way: the map $\sqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$ can have two left adjoints L_1, L_2 such that for some $x : D$, the directed families $L_1(x)$ and $L_2(x)$ exceed each other, yet are unequal. In order for the left adjoint to be truly unique (and not just up to isomorphism), the preorder $\mathcal{V}\text{-Ind}(D)$ would have to identify families that exceed each other. Of course, we could enforce this identification by passing to the poset reflection $\mathcal{V}\text{-Ind}(D)/\approx$ of $\mathcal{V}\text{-Ind}(D)$ and this section studies exactly that.

Another perspective on the situation is the following: The type-theoretic definition of having continuity data for a \mathcal{V} -dcpo D is of the form $\prod_{x:D} \sum_{I:\mathcal{V}} \sum_{\alpha:I \rightarrow D} \dots$, while continuity is defined as its propositional truncation $\|\prod_{x:D} \sum_{I:\mathcal{V}} \sum_{\alpha:I \rightarrow D} \dots\|$. Yet another way to obtain a property is by putting the propositional truncation on the *inside* instead: $\prod_{x:D} \|\sum_{I:\mathcal{V}} \sum_{\alpha:I \rightarrow D} \dots\|$. We study what this amounts to and how it relates to continuity and the poset reflection. Our results are summarised in Table 1 below.

Definition 4.13 (Pseudocontinuity). A \mathcal{V} -dcpo D is *pseudocontinuous* if for every $x : D$ there exists an unspecified directed family that approximates x .

Notice that continuity data \Rightarrow continuity \Rightarrow pseudocontinuity, but reversing the first implication is an instance of global choice [Esc19, Section 3.35.6], while reversing the second amounts to an instance of the axiom of choice that we do not expect to be provable in our constructive setting. We further discuss this point in Remark 4.15.

For a \mathcal{V} -dcpo D , the map $\sqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$ is monotone, so it induces a unique monotone map $\sqcup_{\approx} : \mathcal{V}\text{-Ind}(D)/\approx \rightarrow D$ such that the diagram

$$\begin{array}{ccc}
 \mathcal{V}\text{-Ind}(D)/\approx & \xrightarrow{\sqcup_{\approx}} & D \\
 \searrow [-] & & \nearrow \sqcup \\
 & \mathcal{V}\text{-Ind}(D) &
 \end{array} \tag{1}$$

commutes.

Proposition 4.14. A \mathcal{V} -dcpo D is pseudocontinuous if and only if the map of posets $\sqcup_{\approx} : \mathcal{V}\text{-Ind}(D)/\approx \rightarrow D$ has a necessarily unique left adjoint.

Observe that the type of left adjoints to $\sqcup_{\approx} : \mathcal{V}\text{-Ind}(D)/\approx \rightarrow D$ is a proposition, precisely because $\mathcal{V}\text{-Ind}(D)/\approx$ is a poset, cf. [AKS15, Lemma 5.2].

Proof. Suppose that $\sqcup_{\approx} : \mathcal{V}\text{-Ind}(D)/\approx \rightarrow D$ has a left adjoint L and let $x : D$ be arbitrary. We have to prove that there exists a directed family $\alpha : I \rightarrow D$ that approximates x . By surjectivity of the universal map $[-]$, there exists a directed family $\alpha : I \rightarrow D$ such that $L(x) = [\alpha]$. Moreover, α approximates x by virtue of Lemma 3.8, since for every $\beta : \mathcal{V}\text{-Ind}(D)$, we have

$$\begin{aligned}
 \alpha \lesssim \beta &\iff L(x) \leq [\beta] && \text{(since } L(x) = [\alpha]\text{)} \\
 &\iff x \sqsubseteq \sqcup_{\approx} [\beta] && \text{(since } L \text{ is a left adjoint to } \sqcup_{\approx}\text{)} \\
 &\iff x \sqsubseteq \sqcup \beta && \text{(by Equation (1)).}
 \end{aligned}$$

The converse is more involved and we apply [Kra+17, Theorem 5.4], which says that every weakly constant map with values in a set factors through the propositional truncation of its domain. Assume that D is pseudocontinuous. We start by constructing the left adjoint, so let $x : D$ be arbitrary. Writing \mathcal{A}_x for the type of directed families that approximate x , we have an obvious map $\varphi_x : \mathcal{A}_x \rightarrow \mathcal{V}\text{-Ind}(D)$ that forgets that the directed family approximates x .

We claim that all elements in the image of φ_x exceed each other. For if α and β are directed families both approximating x , then for every i in the domain of α we know that $\alpha_i \ll x = \bigsqcup \beta$, so that there exists j with $\alpha_i \sqsubseteq \beta_j$. Hence, passing to the poset reflection, the composite $[-] \circ \varphi_x$ is constant. Thus, by [Kra+17, Theorem 5.4] we have a (necessarily unique) map ψ_x making the diagram

$$\begin{array}{ccc} \mathcal{A}_x & \xrightarrow{[-] \circ \varphi_x} & \mathcal{V}\text{-Ind}(D)/\approx \\ & \searrow |-\!| & \nearrow \psi_x \\ & & \|\mathcal{A}_x\| \end{array}$$

commute. Since D is assumed to be pseudocontinuous, we have that $\|\mathcal{A}_x\|$ holds for every $x : D$, so together with ψ_x this defines a map $L : D \rightarrow \mathcal{V}\text{-Ind}(D)/\approx$ by $L(x) \equiv \psi_x(p)$, where p witnesses pseudocontinuity at x .

Lastly, we prove that L is indeed a left adjoint to \bigsqcup_{\approx} . So let $x : D$ be arbitrary. Since we're proving a property, we can use pseudocontinuity at x to specify a directed family α that approximates x . We now have to prove $[\alpha] \leq \beta' \iff x \sqsubseteq \bigsqcup_{\approx} \beta'$ for every $\beta' : \mathcal{V}\text{-Ind}(D)/\approx$. This is a proposition, so using quotient induction, it suffices to prove $[\alpha] \leq [\beta] \iff x \sqsubseteq \bigsqcup_{\approx} [\beta]$ for every $\beta : \mathcal{V}\text{-Ind}(D)$. Indeed, for such β we have

$$\begin{aligned} [\alpha] \leq [\beta] &\iff \alpha \lesssim \beta \\ &\iff x \sqsubseteq \bigsqcup \beta && \text{(by Lemma 3.8 and the fact that } \alpha \text{ approximates } x) \\ &\iff x \sqsubseteq \bigsqcup_{\approx} [\beta] && \text{(by Equation (1)),} \end{aligned}$$

finishing the proof. \square

Thus, the explicit type-theoretic formulation and the formulation using left adjoints in each row of Table 1 (which summarises our findings) are equivalent.

	Type-theoretic formulation	Formulation with adjoints	Property
Continuity data	$\prod_{x:D} \Sigma_{I:\mathcal{V}} \Sigma_{\alpha:I \rightarrow D} \delta(\alpha, x)$	Specified left adjoint to $\bigsqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$	\times
Continuity	$\ \prod_{x:D} \Sigma_{I:\mathcal{V}} \Sigma_{\alpha:I \rightarrow D} \delta(\alpha, x)\ $	Unspecified left adjoint to $\bigsqcup : \mathcal{V}\text{-Ind}(D) \rightarrow D$	\checkmark
Pseudocont.	$\prod_{x:D} \ \Sigma_{I:\mathcal{V}} \Sigma_{\alpha:I \rightarrow D} \delta(\alpha, x)\ $	Specified left adjoint to $\bigsqcup_{\approx} : \mathcal{V}\text{-Ind}(D)/\approx \rightarrow D$	\checkmark

TABLE 1. Continuity (data) and pseudocontinuity of a dcpo D , where $\delta(\alpha, x)$ abbreviates that α is directed and approximates x .

Remark 4.15. The issue with pseudocontinuity is that taking the quotient introduces a dependence on instances of the axiom of choice when it comes to proving properties of pseudocontinuous dcpos. An illustrative example is the proof of unary interpolation (Lemma 4.9), where we used the continuity data to first approximate an element y by α_y and then, in turn, approximate every approximant $\alpha_y(i)$. With pseudocontinuity this argument would require *choosing* an approximating family for

every i . Another example is that while the preorder $\mathcal{V}\text{-Ind}(D)$ is \mathcal{V} -directed complete, a direct lifting of the proof of this fact to the poset reflection $\mathcal{V}\text{-Ind}(D)/\approx$ requires the axiom of choice. The same issues concerning the axiom of choice occur in [JJ82, pp. 260–261] and the notion of continuity data follows their solution precisely. We then truncate this to get a property of dcpos (recall Remark 4.3), resulting in our definition of a continuity.

4.3. Algebraic dcpos. Many of our examples of dcpos are not just continuous, but satisfy the stronger condition of being algebraic, meaning their elements can be approximated by compact elements only.

Definition 4.16 (Algebraicity data, I_x, κ_x). *Algebraicity data* for a \mathcal{V} -dcpo D assigns to every $x : D$ a type $I_x : \mathcal{V}$ and a directed family $\kappa_x : I \rightarrow D$ of *compact* elements such that κ_x has supremum x .

Definition 4.17 (Algebraicity). A \mathcal{V} -dcpo is *algebraic* if it has some unspecified algebraicity data, i.e. if the type of algebraicity data is inhabited.

Lemma 4.18. *Every algebraic dcpo is continuous.*

Proof. We prove that algebraicity data for a dcpo yields continuity data. The claim for algebraic and continuous then follows by functoriality of the propositional truncation. It suffices to prove that $\kappa_x(i) \ll x$ for every $i : I_x$. By assumption, $\kappa_x(i)$ is compact and has supremum x . Hence, $\kappa_x(i) \ll \kappa_x(i) \sqsubseteq \bigsqcup \kappa_x = x$, so $\kappa_x(i) \ll x$. \square

5. SMALL BASES

Recall that the traditional, set-theoretic definition of a dcpo D being continuous says that for every element $x \in D$, the subset $\{y \in D \mid y \ll x\}$ is directed with supremum x . As explained in the introduction, the problem with this definition in a predicative context is that the subset $\{y \in D \mid y \ll x\}$ is not small in general. But, as is well-known in domain theory, it is sufficient (and in fact equivalent) to instead ask that D has a subset B , known as a *basis*, such that the subset $\{b \in B \mid b \ll x\} \subseteq B$ is directed with supremum x , see [AJ94, Section 2.2.2] and [Gie+03, Definition III-4.1]. The idea developed in this section is that in many examples we can find a *small* basis giving us a predicative handle on the situation.

If a dcpo has a small basis, then it is continuous. In fact, all our running examples of continuous dcpos are actually examples of dcpos with small bases. Moreover, dcpos with small bases are better behaved. For example, they are all locally small and so are their exponentials, which also have small bases (Section 7.2). In Section 6.3 we also show that having a small basis is equivalent to being presented by ideals.

Following the flow of Section 4, we first consider small bases for continuous dcpos, before turning to small compact bases for algebraic dcpos (Section 5.1). After presenting examples of dcpos with small compact bases in Section 5.2, we describe the canonical small compact basis for an algebraic dcpo and the role that the univalence axiom and a set replacement principle play in Section 5.3.

Definition 5.1 (Small basis). For a \mathcal{V} -dcpo D , a map $\beta : B \rightarrow D$ with $B : \mathcal{V}$ is a *small basis* for D if the following conditions hold:

- (i) for every $x : D$, the family $(\Sigma_{b:B}(\beta(b) \ll x)) \xrightarrow{\beta \text{opr}_1} D$ is directed and has supremum x ;
- (ii) for every $x : D$ and $b : B$, the proposition $\beta(b) \ll x$ is \mathcal{V} -small.

We will write $\downarrow_\beta x$ for the type $\Sigma_{b:B}(\beta(b) \ll x)$ and conflate this type with the canonical map $\downarrow_\beta x \xrightarrow{\beta \text{opr}_1} D$.

Item (ii) ensures not only that the type $\Sigma_{b:B}(\beta(b) \ll x)$ is \mathcal{V} -small, but also that a dcpo with a small basis is locally small (Proposition 5.5).

Remark 5.2. If $\beta : B \rightarrow D$ is a small basis for a \mathcal{V} -dcpo D , then the type $\downarrow_\beta x$ is small. Hence, we have a type $I : \mathcal{V}$ and an equivalence $\varphi : I \simeq \downarrow_\beta x$ and we see that the family $I \xrightarrow{\varphi} \downarrow_\beta x \xrightarrow{\beta \circ \text{pr}_1} D$ is directed and has the same supremum as $\downarrow_\beta x \rightarrow D$. We will use this tacitly and write as if the type $\downarrow_\beta x$ is actually a type in \mathcal{V} .

Lemma 5.3. *If a dcpo comes equipped with a small basis, then it can be equipped with continuity data. Hence, if a dcpo has an unspecified small basis, then it is continuous.*

Proof. For every element x of a dcpo D , the family $\downarrow_\beta x \rightarrow D$ approximates x , so the assignment $x \mapsto \downarrow_\beta x$ is continuity data for D . \square

Lemma 5.4. *In a dcpo D with a small basis $\beta : B \rightarrow D$, we have $x \sqsubseteq y$ if and only if $\forall_{b:B}(\beta(b) \ll x \rightarrow \beta(b) \ll y)$ for every $x, y : D$.*

Proof. If $x \sqsubseteq y$ and $\beta(b) \ll x$, then $\beta(b) \ll y$, so the left-to-right implication is clear. For the converse, suppose that the condition of the lemma holds. Because $x = \bigsqcup \downarrow_\beta x$, the inequality $x \sqsubseteq y$ holds as soon as $\beta(b) \sqsubseteq y$ for every $b : B$ with $\beta(b) \ll x$, but this is implied by the condition. \square

Proposition 5.5. *A dcpo with a small basis is locally small. Moreover, the way-below relation on all of the dcpo has small values.*

Proof. The first claim follows from Lemma 5.4 and the second follows from the first and Proposition 4.12. \square

A notable feature of dcpos with a small basis is that interpolants for the way-below relation, cf. Lemmas 4.8 to 4.10, can be found in the basis, as we show now.

Lemma 5.6 (Nullary interpolation in the basis for the way-below relation). *In a dcpo D with a small basis $\beta : B \rightarrow D$, there exists $b : B$ with $\beta(b) \ll x$ for every $x : D$.*

Proof. For every $x : D$, the approximating family $\downarrow x$ is directed, so there exists $b : B$ with $\beta(b) \ll x$. \square

Lemma 5.7 (Unary interpolation in the basis for the way-below relation). *If $x \ll y$ in a dcpo D with a small basis $\beta : B \rightarrow D$, then there exists an interpolant $b : B$ such that $x \ll \beta(b) \ll y$.*

Proof. The small basis ensures that we have continuity data for D by Lemma 5.3. Hence, if $x \ll y$, then there exists an interpolant $d : D$ with $x \ll d \ll y$. Now $d \ll y \sqsubseteq \bigsqcup \downarrow_\beta y$, so there exists $b : B$ such that $d \sqsubseteq \beta(b) \ll y$. Moreover, $x \ll d \sqsubseteq \beta(b)$, completing the proof. \square

Lemma 5.8 (Binary interpolation in the basis for the way-below relation). *If $x \ll z$ and $y \ll z$ in a dcpo D with a small basis $\beta : B \rightarrow D$, then there exists an interpolant $b : B$ such that $x, y \ll \beta(b)$ and $\beta(b) \ll z$.*

Proof. Again, the small basis ensures that we have continuity data for D by Lemma 5.3. Hence, if $x \ll y$ and $y \ll z$, then there exists an interpolant $d : D$ with $x, y \ll d \ll z$. Now $d \ll z \sqsubseteq \bigsqcup \downarrow_\beta z$, so there exists $b : B$ such that $d \sqsubseteq \beta(b) \ll z$. Moreover, $x, y \ll d \sqsubseteq \beta(b)$, completing the proof. \square

Before proving the analogue of Theorem 4.11 (closure under Scott continuous retracts) for small bases, we need a type-theoretic analogue of [AJ94, Proposition 2.2.4] and [Gie+03, Proposition III-4.2], which essentially says that it is sufficient for a “subset” of $\downarrow_\beta x$ (given by σ in the lemma) to be directed and have supremum x .

Lemma 5.9. *Suppose that we have an element x of a \mathcal{V} -dcpo D together with two maps $\beta : B \rightarrow D$ and $\sigma : I \rightarrow \Sigma_{b:B}(\beta(b) \ll x)$ with $I : \mathcal{V}$. If $\downarrow_\beta x \circ \sigma$ is directed and has supremum x , then $\downarrow_\beta x$ is directed with supremum x too.*

Proof. Suppose that $\downarrow_\beta x \circ \sigma$ is directed and has supremum x . Obviously, x is an upper bound for $\downarrow_\beta x$, so we are to prove that it is the least. If y is an upper bound for $\downarrow_\beta x$, then it is also an upper bound for $\downarrow_\beta x \circ \sigma$ which has supremum x , so that $x \sqsubseteq y$ follows. So the point is directedness of $\downarrow_\beta x$. Its domain is inhabited, because σ is directed. Now suppose that we have $b_1, b_2 : B$ with $\beta(b_1), \beta(b_2) \ll x$. Since $x = \bigsqcup(\downarrow_\beta x \circ \sigma)$, there exist $i_1, i_2 : I$ such that $\beta(b_1) \sqsubseteq \beta(\text{pr}_1(\sigma(i_1)))$ and $\beta(b_2) \sqsubseteq \beta(\text{pr}_1(\sigma(i_2)))$. Since $\downarrow_\beta x \circ \sigma$ is directed, there exists $i : I$ with $\beta(\text{pr}_1(\sigma(i_1))), \beta(\text{pr}_1(\sigma(i_2))) \sqsubseteq \beta(\text{pr}_1(\sigma(i)))$. Hence, writing $b \equiv \text{pr}_1(\sigma(i))$, we have $\beta(b) \ll x$ and $\beta(b_1), \beta(b_2) \sqsubseteq \beta(b)$. Thus, $\downarrow_\beta x$ is directed, as desired. \square

Theorem 5.10. *If we have a Scott continuous retract $D \xrightleftharpoons[r]{s} E$ and $\beta : B \rightarrow E$ is a small basis for E , then $r \circ \beta$ is a small basis for D .*

Proof. First of all, note that E is locally small by Proposition 5.5. But being locally small is closed under Scott continuous retracts by Lemma 4.14 of Part I, so D is locally small too. Moreover, we have continuity data for D by virtue of Theorem 4.11 and Lemma 5.3. Hence, the way-below relation is small-valued by Proposition 4.12. In particular, the proposition $r(\beta(b)) \ll x$ is small for every $b : B$ and $x : D$.

We are going to use Lemma 5.9 to show that $\downarrow_{r \circ \beta} x$ is directed and has supremum x for every $x : D$. By Lemma 2.15, the identity on B induces a well-defined map $\sigma : (\Sigma_{b:B}(\beta(b) \ll s(x))) \rightarrow (\Sigma_{b:B}(r(\beta(b)) \ll x))$. Now Lemma 5.9 tells us that it suffices to prove that $r \circ \downarrow_\beta s(x)$ is directed with supremum x . But $\downarrow_\beta s(x)$ is directed with supremum x , so by Scott continuity of r , the family $r \circ \downarrow_\beta s(x)$ is directed with supremum $r(s(x)) = x$, as desired. \square

Finally, a nice use of dcpos with small bases is that they yield locally small exponentials, as we can restrict the quantification in the pointwise order to elements of the small basis.

Proposition 5.11. *If D is a dcpo with an unspecified small basis and E is a locally small dcpo, then the exponential E^D is locally small too.*

Proof. Being locally small is a proposition, so in proving the result we may assume that D comes equipped with a small basis $\beta : B \rightarrow D$. For arbitrary Scott continuous functions $f, g : D \rightarrow E$, we claim that $f \sqsubseteq g$ precisely when $\forall_{b:B}(f(\beta(b)) \sqsubseteq g(\beta(b)))$, which is a small type using that E is locally small. The left-to-right implication is obvious, so suppose that $f(\beta(b)) \sqsubseteq g(\beta(b))$ for every $b : B$ and let $x : D$ be arbitrary. We are to show that $f(x) \sqsubseteq g(x)$. Since $x = \bigsqcup \downarrow_\beta x$, it suffices to prove $f(\bigsqcup \downarrow_\beta x) \sqsubseteq g(\bigsqcup \downarrow_\beta x)$ and in turn, that $f(\beta(b)) \sqsubseteq g(\bigsqcup \downarrow_\beta x)$ for every $b : B$. But is easily seen to hold, because $f(\beta(b)) \sqsubseteq g(\beta(b))$ for every $b : B$ by assumption. \square

5.1. Small compact bases. Similarly to the progression from continuous dcpos (Section 4.1) to algebraic ones (Section 4.3), we now turn to small *compact* bases.

Definition 5.12 (Small compact basis). For a \mathcal{V} -dcpo D , a map $\beta : B \rightarrow D$ with $B : \mathcal{V}$ is a *small compact basis* for D if the following conditions hold:

- (i) for every $b : B$, the element $\beta(b)$ is compact in D ;
- (ii) for every $x : D$, the family $(\Sigma_{b:B}(\beta(b) \sqsubseteq x)) \xrightarrow{\beta \circ \text{pr}_1} D$ is directed and has supremum x ;
- (iii) for every $x : D$ and $b : B$, the proposition $\beta(b) \sqsubseteq x$ is \mathcal{V} -small.

We will write $\downarrow_\beta x$ for the type $\Sigma_{b:B}(\beta(b) \sqsubseteq x)$ and conflate this type with the canonical map $\downarrow_\beta x \xrightarrow{\beta \circ \text{pr}_1} D$.

Remark 5.13. If $\beta : B \rightarrow D$ is a small compact basis for a \mathcal{V} -dcpo D , then the type $\downarrow_\beta x$ is small. Similarly to Remark 5.2, we will use this tacitly and write as if the type $\downarrow_\beta x$ is actually a type in \mathcal{V} .

Lemma 5.14. *If a dcpo comes equipped with a small compact basis, then it can be equipped with algebraicity data. Hence, if a dcpo has an unspecified small compact basis, then it is algebraic.*

Proof. For every element x of a dcpo D with a small compact basis $\beta : B \rightarrow D$, the family $\downarrow_\beta x \rightarrow D$ consists of compact elements and approximates x , so the assignment $x \mapsto \downarrow_\beta x$ is algebraicity data for D . \square

Actually, with suitable assumptions, we can get canonical algebraicity data from an unspecified small compact basis, as discussed in detail in Section 5.3. This observation relies on Proposition 5.16 below and we prefer to present examples of dcpos with small compact bases first (Section 5.2).

Lemma 5.15. *A map $\beta : B \rightarrow D$ is a small compact basis for a dcpo D if and only if β is a small basis for D and $\beta(b)$ is compact for every $b : B$.*

Proof. If $\beta(b)$ is compact for every $b : B$, then $\beta(b) \sqsubseteq x$ if and only if $\beta(b) \ll x$ for every $b : B$ and $x : D$, so that $\downarrow_\beta x \simeq \downarrow_\beta x$ for every $x : D$. In particular, $\downarrow_\beta x$ approximates x if and only if $\downarrow_\beta x$ does, which completes the proof. \square

Proposition 5.16. *A small compact basis contains every compact element. That is, if $\beta : B \rightarrow D$ is a small compact basis for a dcpo D and $x : D$ is compact, then there exists $b : B$ such that $\beta(b) = x$.*

Proof. Suppose we have a compact element $x : D$. By compactness of x and the fact that $x = \downarrow_\beta x$, there exists $b : B$ with $\beta(b) \ll x$ such that $x \sqsubseteq \beta(b)$. But then $\beta(b) = x$ by antisymmetry. \square

5.2. Examples of dcpos with small compact bases. Armed with the theory of small bases we turn to examples illustrating small bases in practice. Our examples will involve small *compact* bases and an example of a dcpo with a small basis that is not compact will have to wait until Section 6.2 when we have developed the ideal completion.

Example 5.17. The map $\beta : \mathbf{2} \rightarrow \Omega_{\mathcal{U}}$ defined by $0 \mapsto \mathbf{0}_{\mathcal{U}}$ and $1 \mapsto \mathbf{1}_{\mathcal{U}}$ is a small compact basis for $\Omega_{\mathcal{U}}$. In particular, $\Omega_{\mathcal{U}}$ is algebraic.

The basis $\beta : \mathbf{2} \rightarrow \Omega_{\mathcal{U}}$ defined above has the special property that it is *dense* in the sense of [Esc+, TypeTopology.Density]: its image has empty complement, i.e. the type $\Sigma_{P:\Omega_{\mathcal{U}}} \neg(\Sigma_{b:\mathbf{2}} \beta(b) = P)$ is empty.

Proof of Example 5.17. By Example 2.5, every element in the image of β is compact. Moreover, $\Omega_{\mathcal{U}}$ is locally small, so we only need to prove that for every $P : \Omega_{\mathcal{U}}$ the family $\downarrow_\beta P$ is directed with supremum P . The domain of the family is inhabited, because $\beta(0)$ is the least element. Semidirectedness also follows easily, since $\mathbf{2}$ has only two elements for which we have $\beta(0) \sqsubseteq \beta(1)$. Finally, the supremum of $\downarrow_\beta P$ is obviously below P . Conversely, if P holds, then $\bigsqcup \downarrow_\beta P = \mathbf{1} = P$. The final claim follows from Lemma 5.14. \square

Example 5.18. For a set $X : \mathcal{U}$, the map $\beta : (\mathbf{1} + X) \rightarrow \mathcal{L}_{\mathcal{U}}(X)$ given by $\text{inl}(\star) \mapsto \perp$ and $\text{inr}(x) \mapsto \eta(x)$ is a small compact basis for $\mathcal{L}_{\mathcal{U}}(X)$. In particular, $\mathcal{L}_{\mathcal{U}}(X)$ is algebraic.

Similar to Example 5.17, the basis $\beta : (\mathbf{1} + X) \rightarrow \mathcal{L}_{\mathcal{U}}(X)$ defined above is also dense.

Proof of Example 5.18. By Example 2.6, every element in the image of β is compact. Moreover, the lifting is locally small, so we only need to prove that for every partial element l , the family $\downarrow_{\beta} l$ is directed with supremum l . The domain of the family is inhabited, because $\beta(\text{inl}(\star))$ is the least element. Semidirectedness also follows easily: First of all, $\beta(\text{inl}(\star))$ is the least element. Secondly, if we have $x, x' : X$ such that $\beta(\text{inr}(x)), \beta(\text{inr}(x')) \sqsubseteq l$, then because $\beta(\text{inr}(x)) \equiv \eta(x)$ is defined, we must have $\beta(\text{inr}(x)) = l = \beta(\text{inr}(x'))$ by definition of the order. Finally, the supremum of $\downarrow_{\beta} l$ is obviously a partial element below l . Conversely, if l is defined, then $l = \eta(x)$ for some $x : X$, and hence, $l = \eta(x) \sqsubseteq \bigsqcup \downarrow_{\beta} l$. The final claim follows from Lemma 5.14. \square

Example 5.19. For a set $X : \mathcal{U}$, the map $\beta : \text{List}(X) \rightarrow \mathcal{P}_{\mathcal{U}}(X)$ from Definition 2.12 (whose image is the type of Kuratowski finite subsets of X) is a small compact basis for $\mathcal{P}_{\mathcal{U}}(X)$. In particular, $\mathcal{P}_{\mathcal{U}}(X)$ is algebraic.

Proof of Example 5.19. By Lemma 2.13 and Example 2.14, all elements in the image of β are compact. Moreover, $\mathcal{P}_{\mathcal{U}}(X)$ is locally small, so we only need to prove that for every $A : \mathcal{P}(X)$ the family $\downarrow_{\beta} A$ is directed with supremum A , but this was also proven in Example 2.14. The final claim follows from Lemma 5.14. \square

At this point the reader may ask whether we have any examples of dcpos which can be equipped with algebraicity data but that do not have a small compact basis. The following example shows that this can happen in our predicative setting:

Example 5.20. The lifting $\mathcal{L}_{\mathcal{V}}(P)$ of a proposition $P : \mathcal{U}$ can be given algebraicity data, but has a small compact basis if and only if P is \mathcal{V} -small.

Thus, requiring that $\mathcal{L}_{\mathcal{V}}(P)$ has a small basis for every proposition $P : \mathcal{U}$ is equivalent to the propositional resizing principle that every proposition in \mathcal{U} is equivalent to one in \mathcal{V} .

Proof of Example 5.20. Note that $\mathcal{L}_{\mathcal{V}}(P)$ is simply the type of propositions in \mathcal{V} that imply P . It has algebraicity data, because given such a proposition Q , the family

$$\begin{aligned} Q + \mathbf{1}_{\mathcal{V}} &\rightarrow \mathcal{L}_{\mathcal{V}}(P) \\ \text{inl}(q) &\mapsto \mathbf{1}_{\mathcal{V}} \\ \text{inr}(\star) &\mapsto \mathbf{0}_{\mathcal{V}} \end{aligned}$$

is directed, has supremum Q and consists of compact elements. But if $\mathcal{L}_{\mathcal{V}}(P)$ had a small compact basis $\beta : B \rightarrow \mathcal{L}_{\mathcal{V}}(P)$, then we would have $P \simeq \exists_{b:B} (\beta(b) \simeq \mathbf{1}_{\mathcal{V}})$ and the latter is \mathcal{V} -small. Conversely, if P is equivalent to $P_0 : \mathcal{V}$, then $\mathcal{L}_{\mathcal{V}}(P)$ is isomorphic to $\mathcal{L}_{\mathcal{V}}(P_0)$, which has a small compact basis by Example 5.18. \square

Example 5.21. The ordinals in univalent foundations, as introduced in [Uni13, Section 10.3] and further developed by the second author in [Esc+18], give an example of a large (even impredicatively) algebraic sup-lattice without a small basis. In [dJE23, Theorem 5.8] we showed that the large poset of small ordinals has small suprema, so indeed the ordinals form a sup-lattice¹. Moreover, they cannot

¹This result needs small set quotients, or equivalently, set replacement; see Section 5.3 for a discussion of set replacement.

have a small basis, as otherwise we could take the supremum of all ordinals in the basis which would yield a greatest ordinal which does not exist (as a consequence of [Uni13, Lemma 10.3.21]).

It remains to show that the sup-lattice of ordinals is algebraic. This follows from the following two facts.

- (i) Every successor ordinal, i.e. one of the form $\alpha + \mathbf{1}$, is compact.
- (ii) Every ordinal α is the supremum of the family $x : \alpha \mapsto \alpha \downarrow x + \mathbf{1}$, where $\alpha \downarrow x$ denotes the ordinal of elements of α that are strictly less than x .

While the family in (ii) is not necessarily directed, this does not pose a problem, since we can take its directification (see Definition 7.10 later) by considering finite joins of elements in the family which are necessarily compact again by Lemma 2.7.

For proving these facts, we recall from [Esc+18, Ordinals.OrdinalOfOrdinals] that the order \preceq on ordinals can be characterised as follows:

$$\alpha \preceq \beta \iff \forall x:\alpha \exists y:\beta \alpha \downarrow x = \beta \downarrow y.$$

We now prove (i): Suppose that $\alpha + \mathbf{1} \preceq \bigsqcup_{i:I} \beta_i$. Since $\alpha = (\alpha + \mathbf{1}) \downarrow \text{inr}(\star)$, there exists $s : \bigsqcup_{i:I} \beta_i$ with $\alpha = \bigsqcup_{i:I} \beta_i \downarrow s$. By [dJon+23, Lemma 15] there exist $j : I$ and $b : \beta_j$ with $\bigsqcup_{i:I} \beta_i \downarrow s = \beta_j \downarrow b$. Hence, $\alpha = \beta_j \downarrow b$, but then it follows that $\alpha + \mathbf{1} \preceq \beta_j$.

To see that (ii) is true, we first notice that $\alpha \downarrow x + \mathbf{1} \preceq \alpha$ holds (using the characterisation of the partial order) so that α is an upper bound for the family. Now suppose that β is another upper bound. We need to show that $\alpha \preceq \beta$. So let $x : \alpha$ be arbitrary. Since β is an upper bound of the family, there is $b : \beta$ with $\alpha \downarrow x = (\alpha \downarrow x + \mathbf{1}) \downarrow \text{inr} \star = \beta \downarrow b$, so we are done.

5.3. The basis of compact elements. So far, our development of algebraic dcpos (with small compact bases) has resulted from specialising the treatment of continuous dcpos with small bases. In this section we take a closer look at the algebraic case.

Classically, the subset K of compact elements of an algebraic dcpo D forms a basis for D . In our predicative context, we only consider small bases, and a priori there is no reason for K to be a small type. However, if D comes equipped with a small compact basis, then set replacement implies that K is in fact small.

We recall the *set replacement* principle from [dJE23, Definition 3.27]: it asserts that the image of a map $f : X \rightarrow Y$ is $\mathcal{U} \sqcup \mathcal{V}$ -small if X is \mathcal{U} -small and Y is a locally \mathcal{V} -small set, where local smallness refers to smallness of the identity types. We also recall [dJE23, Theorem 3.29] that set replacement is logically equivalent to the existence of small set quotients.

If we additionally assume univalence, then the relevant smallness condition is a property [dJE23, Section 2.3], which means that having an unspecified small compact basis is sufficient.

In particular, with set replacement and univalence, we can show:

$$\begin{aligned} & \|D \text{ has a specified small compact basis}\| \rightarrow \\ & D \text{ has a specified small compact basis} \end{aligned}$$

Using the terminology of [Kra+17, Definition 3.9], we may thus say that the type expressing that D has a specified small compact basis has split support. This observation is due to Ayberk Tosun (private communication) who also formalised the result for spectrality in the context of locale theory in predicative univalent foundations [AET24, Theorem 4.17].

Lemma 5.22. *Every element x of an algebraic dcpo is the directed supremum of all compact elements below x .*

Proof. Writing D for the algebraic dcpo, and letting $x : D$ be arbitrary, we have to show that the inclusion family

$$\iota_x : (\Sigma_{c:D}(c \text{ is compact}) \times (c \sqsubseteq x)) \rightarrow D$$

is directed with supremum x . Since this is a proposition, we may assume to be given algebraicity data for D . Thus, we have a directed family $\kappa_x : I_x \rightarrow D$ of compact elements with supremum x . By directedness, I_x is inhabited, so we see that the domain of ι_x is inhabited too. For semidirectedness, assume we have compact elements $c_1, c_2 : D$ below x . Since x is the directed supremum of κ_x , there exist elements $i_1, i_2 : I_x$ with $c_1 \sqsubseteq \kappa_x(i_1)$ and $c_2 \sqsubseteq \kappa_x(i_2)$. By directedness of κ_x , there then exists $i : I_x$ such that c_1 and c_2 are both below $\kappa_x(i)$. But $\kappa_x(i)$ is a compact element below x so we are done. Finally, we show that ι_x has supremum x . Clearly, x is an upper bound for ι_x . Now suppose that y is any other upper bound. It then suffices to show that $\kappa_x(i) \sqsubseteq y$ for all $i : I_x$. But each $\kappa_x(i)$ is a compact element below x , so this holds since y is an upper bound for ι_x . \square

Lemma 5.23. *Assuming set replacement, if a \mathcal{V} -dcpo is equipped with a small compact basis, then the subtype of compact elements is \mathcal{V} -small.*

If we additionally assume univalence, then having an unspecified small compact basis suffices.

Proof. Let $\beta : B \rightarrow D$ be the small compact basis of the \mathcal{V} -dcpo D . Notice that β factors through the subtype K of compact elements of D . Moreover, by Proposition 5.16 the map $\beta : B \rightarrow K$ is surjective. Hence, K is equivalent to the image of $\beta : B \rightarrow D$. Now an application of set replacement finishes the proof, since B is small and K is locally small because it is a subtype of D which is locally small by antisymmetry, Proposition 5.5 and Lemma 5.15.

Assuming univalence, being small is a property [dJE23, Proposition 2.8], so that the result follows from the above and the universal property of the truncation. \square

Proposition 5.24. *Assuming univalence and set replacement, the types expressing that a dcpo has a specified, resp. unspecified, small compact basis are logically equivalent.*

Proof. In one direction, we simply apply the propositional truncation. In the other direction, we apply Lemmas 5.22 and 5.23 to see that

$$K_s \simeq K \hookrightarrow D$$

is a small compact basis for the \mathcal{V} -dcpo D , where K denotes the subtype of compact elements and K_s is the \mathcal{V} -small copy of K . \square

We note that the above cannot be promoted to an equivalence of types, because the type of specified small compact bases is not a proposition. This may seem puzzling to a domain theorist because there is a unique basis—as a subset—which consists of compact elements. If we had asked for the map $\beta : B \rightarrow D$ in the definition of a small compact basis to be an embedding, then (ignoring size issues for a moment) the resulting type is a proposition: it has a unique element in case the dcpo is algebraic, given by the *subset* of compact elements. (This is true because any basis must contain all compact elements.)

We illustrate why we do not impose this requirement by revisiting Example 5.19. This example showed that the canonical map from lists into the powerset of a set X is a small compact basis for the algebraic dcpo $\mathcal{P}(X)$. This map is not an embedding, as any permutation of a list will give rise to the same subset. If we insisted on having an embedding, we would instead have to use the inclusion of the Kuratowski finite subsets $\mathcal{K}(X)$ into $\mathcal{P}(X)$. However, $\mathcal{K}(X)$ is not a small type

without additional assumptions, such as HITs or more specifically, set replacement (as $\mathcal{K}(X)$ is precisely the image of the inclusion of lists).

Returning to the main line of thought, we conclude that, in the presence of set replacement and univalence, if there is some unspecified small compact basis, then the subset of compact elements is small.

6. THE ROUNDED IDEAL COMPLETION

We have seen that in continuous dcpos, the basis essentially “generates” the whole dcpo, because the basis suffices to approximate any of its elements. It is natural to ask whether one can start from a more abstract notion of basis and “complete” it to a continuous dcpo. Following Abramsky and Jung [AJ94, Section 2.2.6], but keeping track of size, this is exactly what we do here using the notion of an *abstract basis* and the *rounded ideal completion*.

Definition 6.1 (Abstract basis). An *abstract \mathcal{V} -basis* is a type $B : \mathcal{V}$ with a binary relation $\prec : B \rightarrow B \rightarrow \mathcal{V}$ that is proposition-valued, transitive and satisfies *nullary* and *binary interpolation*:

- (i) for every $a : B$, there exists $b : B$ with $b \prec a$, and
- (ii) for every $a_1, a_2 \prec b$, there exists $a : B$ with $a_1, a_2 \prec a \prec b$.

Definition 6.2 (Ideal, (rounded) ideal completion, $\mathcal{V}\text{-Idl}(B, \prec)$).

- (i) A subset $I : \mathcal{P}_{\mathcal{V}}(B)$ of an abstract \mathcal{V} -basis (B, \prec) is a *\mathcal{V} -ideal* if it is a directed lower set with respect to \prec . That it is a lower set means: if $b \in I$ and $a \prec b$, then $a \in I$ too.
- (ii) We write $\mathcal{V}\text{-Idl}(B, \prec)$ for the type of \mathcal{V} -ideals of an abstract \mathcal{V} -basis (B, \prec) and call $\mathcal{V}\text{-Idl}(B, \prec)$ the *(rounded) ideal completion* of (B, \prec) .

For the remainder of this section, we will fix an abstract \mathcal{V} -basis (B, \prec) and consider its \mathcal{V} -ideals. The name rounded ideal completion is justified by Lemmas 6.5 and 6.6 below.

Definition 6.3 (Union of ideals, $\bigcup \mathcal{I}$). Given a family $\mathcal{I} : S \rightarrow \mathcal{V}\text{-Idl}(B, \prec)$ of ideals, indexed by $S : \mathcal{V}$, we write

$$\bigcup \mathcal{I} := \{b \in B \mid \exists_{s:S} (b \in \mathcal{I}_s)\}.$$

Lemma 6.4. *If $\mathcal{I} : S \rightarrow \mathcal{V}\text{-Idl}(B, \prec)$ is directed, then $\bigcup \mathcal{I}$ is an ideal.*

Proof. The subset $\bigcup \mathcal{I}$ is easily seen to be a lower set, for if $a \prec b \in \bigcup \mathcal{I}$, then there exists $s : S$ such that $a \prec b \in \mathcal{I}_s$, so $a \in \mathcal{I}_s$ as \mathcal{I}_s is a lower set, but then $a \in \bigcup \mathcal{I}$. Moreover, $\bigcup \mathcal{I}$ is inhabited: Since \mathcal{I} is directed, there exists $s : S$, but \mathcal{I}_s is an ideal and therefore inhabited, so there exists $b \in \mathcal{I}_s$ which implies $b \in \bigcup \mathcal{I}$. Finally, suppose we have $b_1, b_2 \in \bigcup \mathcal{I}$. By definition, there exist $s_1, s_2 : S$ such that $b_1 \in \mathcal{I}_{s_1}$ and $b_2 \in \mathcal{I}_{s_2}$. By directedness of S , there exists $s : S$ such that $\mathcal{I}_{s_1}, \mathcal{I}_{s_2} \subseteq \mathcal{I}_s$. Hence, $b_1, b_2 \in \mathcal{I}_s$, which is an ideal, so there exists $b \in \mathcal{I}_s$ with $b_1, b_2 \prec b$. But then also $b \in \bigcup \mathcal{I}$, which proves that $\bigcup \mathcal{I}$ is directed and hence an ideal, completing the proof. \square

Lemma 6.5. *The rounded ideal completion is a \mathcal{V} -dcpo when ordered by subset inclusion.*

Proof. Since taking unions yields the least upper bound in the powerset, we only have to prove that the union of ideals is again an ideal, but this is Lemma 6.4. \square

Paying attention to the universe levels, the ideals form a large but locally small \mathcal{V} -dcpo because $\mathcal{V}\text{-Idl}(B, \prec) : \mathcal{V}\text{-DCPO}_{\mathcal{V}^+, \mathcal{V}}$.

Lemma 6.6 (Roundedness). *The ideals of an abstract basis are rounded: for every element a of an ideal I , there exists $b \in I$ such that $a \prec b$.*

Proof. Because ideals are semidirected. \square

Roundedness makes up for the fact that we have not required an abstract basis to be reflexive. If it is, then (Section 6.1) the ideal completion is algebraic.

Definition 6.7 (Principal ideal, $\downarrow b$). The *principal ideal* of an element $b : B$ is defined as the subset $\downarrow b \equiv \{a \in B \mid a \prec b\}$. Observe that the principal ideal is indeed an ideal: it is a lower set by transitivity of \prec , and inhabited and semidirected precisely by nullary and binary interpolation, respectively.

Lemma 6.8. *The assignment $b \mapsto \downarrow b$ is monotone, i.e. if $a \prec b$, then $\downarrow a \subseteq \downarrow b$.*

Proof. By transitivity of \prec . \square

Lemma 6.9. *Every ideal is the directed supremum of its principal ideals. That is, for an ideal I , the family $(\Sigma_{b:B}(b \in I)) \xrightarrow{b \mapsto \downarrow b} \mathcal{V}\text{-Idl}(B, \prec)$ is directed and has supremum I .*

Proof. Since ideals are lower sets, we have $\downarrow b \subseteq I$ for every $b \in I$. Hence, the union $\bigcup_{b \in I} \downarrow b$ is a subset of I . Conversely, if $a \in I$, then by roundedness of I there exists $a' \in I$ with $a \prec a'$, so that $a \in \bigcup_{b \in I} \downarrow b$. So it remains to show that the family is directed. Notice that it is inhabited, because I is an ideal. Now suppose that $b_1, b_2 \in I$. Since I is directed, there exists $b \in I$ such that $b_1, b_2 \prec b$. But this implies $\downarrow b_1, \downarrow b_2 \subseteq \downarrow b$ by Lemma 6.8, so the family is semidirected, as desired. \square

Lemma 6.10. *The following are equivalent for every two ideals I and J :*

- (i) $I \ll J$;
- (ii) there exists $b \in J$ such that $I \subseteq \downarrow b$;
- (iii) there exist $a \prec b$ such that $I \subseteq \downarrow a \subseteq \downarrow b \subseteq J$.

In particular, if b is an element of an ideal I , then $\downarrow b \ll I$.

Proof. We show that (i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (i). So suppose that $I \ll J$. Then J is the directed supremum of its principal ideals by Lemma 6.9. Hence, there exists $b \in J$ such that $I \subseteq \downarrow b$ already, which is exactly (ii). Now suppose that we have $a \in J$ with $I \subseteq \downarrow a$. By roundedness of J , there exists $b \in J$ with $a \prec b$. But then $I \subseteq \downarrow a \subseteq \downarrow b \subseteq J$ by Lemma 6.8 and the fact that J is a lower set, establishing (iii). Now suppose that condition (iii) holds and that J is a subset of some directed join of ideals \mathcal{J} indexed by a type $S : \mathcal{V}$. Since $a \in \downarrow b \subseteq J$, there exists $s : S$ such that $a \in \mathcal{J}_s$. In particular, $\downarrow a \subseteq \mathcal{J}_s$ because ideals are lower sets. Hence, if $a' \in I \subseteq \downarrow a$, then $a' \in \mathcal{J}_s$, so $I \subseteq \mathcal{J}_s$, which proves that $I \ll J$.

Finally, if b is an element of an ideal I , then $\downarrow b \ll I$, because (ii) implies (i) and $\downarrow b \subseteq \downarrow b$ obviously holds. \square

Theorem 6.11. *The principal ideals $\downarrow(-) : B \rightarrow \mathcal{V}\text{-Idl}(B, \prec)$ yield a small basis for $\mathcal{V}\text{-Idl}(B, \prec)$. In particular, $\mathcal{V}\text{-Idl}(B, \prec)$ is continuous.*

Proof. First of all, note that the way-below relation on $\mathcal{V}\text{-Idl}(B, \prec)$ is small-valued because of Lemma 6.10. So it remains to show that for every ideal I , the family $(\Sigma_{b:B}(\downarrow b \ll I)) \xrightarrow{b \mapsto \downarrow b} \mathcal{V}\text{-Idl}(B, \prec)$ is directed with supremum I . That the domain of this family is inhabited follows from Lemma 6.10 and the fact that I is inhabited. For semidirectedness, suppose we have $b_1, b_2 : B$ with $\downarrow b_1, \downarrow b_2 \ll I$. By Lemma 6.10 there exist $c_1, c_2 \in I$ such that $\downarrow b_1 \subseteq \downarrow c_1$ and $\downarrow b_2 \subseteq \downarrow c_2$. Since I is directed, there exists $b \in I$ with $c_1, c_2 \prec b$. But now $\downarrow b_1 \subseteq \downarrow c_1 \subseteq \downarrow b \ll I$ by Lemmas 6.8 and 6.10 and similarly, $\downarrow b_2 \subseteq \downarrow b \ll I$. Hence, the family is semidirected, as we wished to

show. Finally, we show that I is the supremum of the family. If $b \in I$, then, since I is rounded, there exists $c \in I$ with $b \prec c$. Moreover, $\downarrow c \ll I$ by Lemma 6.10. Hence, b is included in the join of the family. Conversely, if we have $b : B$ with $\downarrow b \ll I$, then $\downarrow b \subseteq I$, so I is also an upper bound for the family. \square

6.1. The rounded ideal completion of a reflexive abstract basis. If the relation of an abstract basis is reflexive, then we obtain an algebraic dcpo, as we show now.

Lemma 6.12. *If $\prec : B \rightarrow B \rightarrow \mathcal{V}$ is proposition-valued, transitive and reflexive, then (B, \prec) is an abstract basis.*

Proof. The interpolation properties for \prec are easily proved when it is reflexive. \square

Lemma 6.13. *If an element $b : B$ is reflexive, i.e. $b \prec b$ holds, then $b \in I$ if and only if $\downarrow b \subseteq I$ for every ideal I .*

Proof. The left-to-right implication holds because I is a lower set and the converse holds because $b \in \downarrow b$ as b is assumed to be reflexive. \square

Lemma 6.14. *If $b : B$ is reflexive, then its principal ideal $\downarrow b$ is compact.*

Proof. Suppose that we have $b : B$ such that $b \prec b$ holds and that $\downarrow b \subseteq \bigcup \mathcal{I}$ for some directed family \mathcal{I} of ideals. By Lemma 6.13, we have $b \in \bigcup \mathcal{I}$, which means that there exists s in the domain of \mathcal{I} such that $b \in \mathcal{I}_s$. Using Lemma 6.13 once more, we see that $\downarrow b \subseteq \mathcal{I}_s$, proving that $\downarrow b$ is compact. \square

Theorem 6.15. *If \prec is reflexive, then the principal ideals $\downarrow(-) : B \rightarrow \mathcal{V}\text{-Idl}(B, \prec)$ yield a small compact basis for $\mathcal{V}\text{-Idl}(B, \prec)$. In particular, $\mathcal{V}\text{-Idl}(B, \prec)$ is algebraic.*

Proof. This follows from Theorem 6.11 and Lemmas 5.15 and 6.14. \square

Theorem 6.16. *If $f : B \rightarrow D$ is a monotone map to a \mathcal{V} -dcpo D , then the map $\bar{f} : \mathcal{V}\text{-Idl}(B, \prec) \rightarrow D$ defined by taking an ideal I to the supremum of the directed family $f \circ \text{pr}_1 : (\Sigma_{b:B}(b \in I)) \rightarrow D$ is Scott continuous. Moreover, if \prec is reflexive, then \bar{f} is the unique Scott continuous map making the diagram*

$$\begin{array}{ccc} B & \xrightarrow{f} & D \\ \downarrow(-) \searrow & & \nearrow \bar{f} \\ & \mathcal{V}\text{-Idl}(B, \prec) & \end{array}$$

commute.

Proof. Note that $f \circ \text{pr}_1 : (\Sigma_{b:B}(b \in I)) \rightarrow D$ is indeed a directed family, because I is a directed subset of B and f is monotone. For Scott continuity of \bar{f} , assume that we have a directed family \mathcal{I} of ideals indexed by $S : \mathcal{V}$. We first show that $\bar{f}(\bigcup \mathcal{I})$ is an upper bound of $\bar{f} \circ \mathcal{I}$. So let $s : S$ be arbitrary and note that $\bar{f}(\bigcup \mathcal{I}) \supseteq \bar{f}(\mathcal{I}_s)$ as soon as $\bar{f}(\bigcup \mathcal{I}) \supseteq f(b)$ for every $b \in \mathcal{I}_s$. But for such b we have $b \in \bigcup \mathcal{I}$, so this holds. Now suppose that y is an upper bound of $\bar{f} \circ \mathcal{I}$. To show that $\bar{f}(\bigcup \mathcal{I}) \sqsubseteq y$, it is enough to prove that $f(b) \sqsubseteq y$ for every $b \in \mathcal{I}$. But for such b , there exists $s : S$ such that $b \in \mathcal{I}_s$ and hence, $f(b) \sqsubseteq \bar{f}(\mathcal{I}_s) \sqsubseteq y$.

Finally, if \prec is reflexive, then we prove that $\bar{f}(\downarrow b) = f(b)$ for every $b : B$ by antisymmetry. Since \prec is assumed to be reflexive, we have $b \in \downarrow b$ and therefore, $f(b) \sqsubseteq \bar{f}(\downarrow b)$. Conversely, for every $c \prec b$ we have $f(c) \sqsubseteq f(b)$ by monotonicity of f and hence, $\bar{f}(\downarrow b) \sqsubseteq f(b)$, as desired. Uniqueness is proved easily, because if $g : \mathcal{V}\text{-Idl}(B, \prec) \rightarrow D$ is Scott continuous with $g(\downarrow b) = f(b)$, then for an arbitrary ideal I we have $g(I) = g(\bigcup_{b \in I} \downarrow b) = \bigcup_{b \in I} g(\downarrow b) = \bigcup_{b \in I} f(b) \equiv \bar{f}(I)$ by Lemma 6.9 and Scott continuity of g . \square

6.2. Example: the ideal completion of dyadics. We describe an example of a continuous dcpo, built using the ideal completion, that is not algebraic. In fact, this dcpo has no compact elements at all.

We inductively define a type and an order representing dyadic rationals $m/2^n$ in the interval $(-1, 1)$ for integers m, n . This type is isomorphic to the lower Dedekind reals extended with ∞ . We could work with lower Dedekind reals but we prefer the type of dyadics for its simplicity.

The intuition for the upcoming definitions is the following: Start with the point 0 in the middle of the interval (represented by *middle* below). Then consider the two functions (respectively represented by *left* and *right* below)

$$\begin{aligned} l, r &: (-1, 1) \rightarrow (-1, 1) \\ l(x) &:\equiv (x - 1)/2 \\ r(x) &:\equiv (x + 1)/2 \end{aligned}$$

that generate the dyadic rationals. Observe that $l(x) < 0 < r(x)$ for every $x : (-1, 1)$. Accordingly, we inductively define the following types:

Definition 6.17 (Dyadics, \mathbb{D} , \prec). The type of *dyadics* $\mathbb{D} : \mathcal{U}_0$ is the inductive type with these three constructors

$$\text{middle} : \mathbb{D} \quad \text{left} : \mathbb{D} \rightarrow \mathbb{D} \quad \text{right} : \mathbb{D} \rightarrow \mathbb{D}.$$

We also inductively define $\prec : \mathbb{D} \rightarrow \mathbb{D} \rightarrow \mathcal{U}_0$ as

$$\begin{aligned} \text{middle} \prec \text{middle} &:\equiv \mathbf{0} & \text{left}(x) \prec \text{middle} &:\equiv \mathbf{1} & \text{right}(x) \prec \text{middle} &:\equiv \mathbf{0} \\ \text{middle} \prec \text{left}(y) &:\equiv \mathbf{0} & \text{left}(x) \prec \text{left}(y) &:\equiv x \prec y & \text{right}(x) \prec \text{left}(y) &:\equiv \mathbf{0} \\ \text{middle} \prec \text{right}(y) &:\equiv \mathbf{1} & \text{left}(x) \prec \text{right}(y) &:\equiv \mathbf{1} & \text{right}(x) \prec \text{right}(y) &:\equiv x \prec y. \end{aligned}$$

Lemma 6.18. *The type of dyadics is a set with decidable equality.*

Proof. Sethood follows from having decidable equality by Hedberg's Theorem. To see that \mathbb{D} has decidable equality, one can use a standard inductive proof. \square

Definition 6.19 (Trichotomy, density, having no endpoints). We say that a binary relation $<$ on a type X is

- *trichotomous* if exactly one of $x < y$, $x = y$ or $y < x$ holds.
- *dense* if for every $x, y : X$, there exists some $z : X$ such that $x < z < y$.
- *without endpoints* if for every $x : X$, there exist some $y, z : X$ with $y < x < z$.

Lemma 6.20. *The relation \prec on the dyadics is proposition-valued, transitive, irreflexive, trichotomous, dense and without endpoints.*

Proof. That \prec is proposition-valued, transitive, irreflexive and trichotomous is all proven by a straightforward induction on the definition on \mathbb{D} . That it has no endpoints is witnessed by the fact that for every $x : \mathbb{D}$, we have

$$\text{left } x \prec x \prec \text{right } x \tag{\dagger}$$

which is proven by induction on \mathbb{D} as well. We spell out the inductive proof that it is dense, making use of (\dagger) . Suppose that $x \prec y$. Looking at the definition of the order, we see that we need to consider five cases.

- If $x = \text{middle}$ and $y = \text{right } y'$, then we have $x \prec \text{right}(\text{left}(y')) \prec y$.
- If $x = \text{left}(x')$ and $y = \text{middle}$, then we have $x \prec \text{left}(\text{right}(x')) \prec y$.
- If $x = \text{left}(x')$ and $y = \text{right } y'$, then we have $x \prec \text{middle} \prec y$.
- If $x = \text{right}(x')$ and $y = \text{right } y'$, then we have $x' \prec y'$ and therefore, by induction hypothesis, there exists $z' : \mathbb{D}$ such that $x' \prec z' \prec y'$. Hence, $x \prec \text{right}(z') \prec y$.

- If $x = \text{left}(x')$ and $y = \text{left}(y')$, then $x' \prec y'$ and so, by induction hypothesis, there exists $z' : \mathbb{D}$ such that $x' \prec z' \prec y'$. Thus, $x \prec \text{left}(z') \prec y$. \square

Proposition 6.21. *The pair (\mathbb{D}, \prec) is an abstract \mathcal{U}_0 -basis.*

Proof. By Lemma 6.20 the relation \prec is proposition-valued and transitive. Moreover, that it has no endpoints implies unary interpolation. For binary interpolation, suppose that we have $x \prec z$ and $y \prec z$. Then by trichotomy there are three cases.

- If $x = y$, then using density and our assumption that $x \prec z$, there exists $d : \mathbb{D}$ with $y = x \prec d \prec z$, as desired.
- If $x \prec y$, then using density and our assumption that $y \prec z$, there exists $d : \mathbb{D}$ with $y \prec d \prec z$, but then also $x \prec d$ since $x \prec y$, so we are done.
- If $x \prec y$, then the proof is similar to that of the second case. \square

Proposition 6.22. *The ideal completion $\mathcal{U}_0\text{-Idl}(\mathbb{D}, \prec) : \mathcal{U}_0\text{-DCPO}_{\mathcal{U}_1, \mathcal{U}_0}$ is continuous with small basis $\downarrow(-) : \mathbb{D} \rightarrow \mathcal{U}_0\text{-Idl}(\mathbb{D}, \prec)$. Moreover, it cannot be algebraic, because none of its elements are compact.*

Proof. The first claim follows from Theorem 6.11. Now suppose for a contradiction that we have a compact ideal I . By Lemma 6.10, there exists $x \in I$ with $I \subseteq \downarrow x$. But this implies $x \prec x$, which is impossible as \prec is irreflexive. \square

6.3. Ideal completions of small bases. Given a \mathcal{V} -dcpo D with a small basis $\beta : B \rightarrow D$, we show that there are two natural ways of turning B into an abstract basis. Either define $b \prec c$ by $\beta(b) \ll \beta(c)$, or by $\beta(b) \sqsubseteq \beta(c)$. Taking their \mathcal{V} -ideal completions we show that the former yields a continuous dcpo isomorphic to D , while the latter yields an algebraic dcpo (with a small compact basis) in which D can be embedded. The latter fact will find application in Section 7.2, while the former gives us a presentation theorem: every dcpo with a small basis is isomorphic to a dcpo of ideals. In particular, if $D : \mathcal{V}\text{-DCPO}_{\mathcal{U}, \mathcal{T}}$ has a small basis, then it is isomorphic to a dcpo with simpler universe parameters, namely $\mathcal{V}\text{-Idl}(B, \ll_\beta) : \mathcal{V}\text{-DCPO}_{\mathcal{V}^+, \mathcal{V}}$. Of course a similar result holds for dcpos with a small compact basis. In studying these variations, it is helpful to first develop some machinery that all of them have in common.

Fix a \mathcal{V} -dcpo D with a small basis $\beta : B \rightarrow D$. In what follows we conflate the family $\downarrow_\beta x : (\sum_{b:B} (\beta(b) \ll x)) \xrightarrow{\beta_{\text{opr1}}} D$ with its associated subset $\{b \in B \mid \beta(b) \ll x\}$, formally given by the map $B \rightarrow \Omega_{\mathcal{V}}$ defined as $b \mapsto \exists_{b:B} (\beta(b) \ll x)$.

Lemma 6.23. *The assignment $x : D \mapsto \downarrow_\beta x : \mathcal{P}(B)$ is Scott continuous.*

Proof. Note that $\downarrow_\beta(-)$ is monotone: if $x \sqsubseteq y$ and $b : B$ is such that $\beta(b) \ll x$, then also $\beta(b) \ll y$. So it suffices to prove that $\downarrow_\beta(\bigsqcup \alpha) \subseteq \bigcup_{i:I} \downarrow_\beta \alpha_i$. So suppose that $b : B$ is such that $\beta(b) \ll \bigsqcup \alpha$. By Lemma 5.7, there exists $c : B$ with $\beta(b) \ll \beta(c) \ll \bigsqcup \alpha$. Hence, there exists $i : I$ such that $\beta(b) \ll \beta(c) \sqsubseteq \alpha_i$ already, and therefore, $b \in \bigcup_{j:J} \downarrow_\beta \alpha_j$, as desired. \square

By virtue of the fact that β is a small basis for D , we know that taking the directed supremum of $\downarrow_\beta x$ equals x for every $x : D$. In other words, $\downarrow_\beta(-)$ is a section of $\bigsqcup(-)$. The following lemma gives conditions for the other composite to be an inflation or a deflation.

Lemma 6.24. *Let $I : \mathcal{P}_{\mathcal{V}}(B)$ be a subset of B such that its associated family $\bar{I} : (\sum_{b:B} (b \in I)) \xrightarrow{\beta_{\text{opr1}}} D$ is directed.*

- (i) *If the conjunction of $\beta(b) \sqsubseteq \beta(c)$ and $c \in I$ implies $b \in I$, then $\downarrow_\beta \bigsqcup \bar{I} \subseteq I$.*
- (ii) *If for every $b \in I$ there exists $c \in I$ such that $\beta(b) \ll \beta(c)$, then $I \subseteq \downarrow_\beta \bigsqcup \bar{I}$.*

In particular, if both conditions hold, then $I = \downarrow_{\beta} \bigsqcup \bar{I}$.

Note that the first condition says that I is a lower set with respect to the order of D , while the second says that I is rounded with respect to the way-below relation.

Proof. (i) Suppose that I is a lower set and let $b : B$ be such that $\beta(b) \ll \bigsqcup \bar{I}$. Then there exists $c \in I$ with $\beta(b) \sqsubseteq \beta(c)$, which implies $b \in I$ as desired, because I is assumed to be a lower set. (ii) Assume that I is rounded and let $b \in I$ be arbitrary. By roundedness of I , there exists $c \in I$ such that $\beta(b) \ll \beta(c)$. But then $\beta(b) \ll \beta(c) \sqsubseteq \bigsqcup \bar{I}$, so that $b \in \downarrow_{\beta} \bigsqcup \bar{I}$, as we wished to show. \square

Lemma 6.25. *Suppose that we have $\prec : B \rightarrow B \rightarrow \mathcal{V}$ and let $x : D$ be arbitrary.*

- (i) *If $b \prec c$ implies $\beta(b) \sqsubseteq \beta(c)$ for every $b, c : B$, then $\downarrow_{\beta} x$ is a lower set w.r.t. \prec .*
- (ii) *If $\beta(b) \ll \beta(c)$ implies $b \prec c$ for every $b, c : B$, then $\downarrow_{\beta} x$ is semidirected w.r.t. \prec .*

Proof. (i) This is immediate, because $\downarrow_{\beta} x$ is a lower set with respect to the order relation on D . (ii) Suppose that the condition holds and that we have $b_1, b_2 : B$ such that $\beta(b_1), \beta(b_2) \ll x$. Using binary interpolation in the basis, there exist $c_1, c_2 : B$ with $\beta(b_1) \ll \beta(c_1) \ll x$ and $\beta(b_2) \ll \beta(c_2) \ll x$. Hence, $c_1, c_2 \in \downarrow_{\beta} x$ and moreover, by assumption we have $b_1 \prec c_1$ and $b_2 \prec c_2$, as desired. \square

6.3.1. *Ideal completion with respect to the way-below relation.*

Lemma 6.26. *If $\beta : B \rightarrow D$ is a small basis for a \mathcal{V} -dcpo D , then (B, \ll_{β}) is an abstract \mathcal{V} -basis where $b \ll_{\beta} c$ is defined as $\beta(b) \ll \beta(c)$.*

Remark 6.27. The definition of an abstract \mathcal{V} -basis requires the relation on it to be \mathcal{V} -valued. Hence, for the lemma to make sense we appeal to the fact that β is a small basis which tells us that we can substitute $\beta(b) \ll \beta(c)$ by an equivalent type in \mathcal{V} .

Proof of Lemma 6.26. The way-below relation is proposition-valued and transitive. Moreover, \ll_{β} satisfies nullary and binary interpolation precisely because we have nullary and binary interpolation in the basis for the way-below relation by Lemmas 5.6 and 5.8. \square

The following theorem is a presentation result for dcpos with a small basis: every such dcpo can be presented as the rounded ideal completion of its small basis.

Theorem 6.28. *The map $\downarrow_{\beta}(-) : D \rightarrow \mathcal{V}\text{-Idl}(B, \ll_{\beta})$ is an isomorphism of \mathcal{V} -dcpos.*

Proof. First of all, we should check that the map is well-defined, i.e. that $\downarrow_{\beta} x$ is an (B, \ll_{β}) -ideal. It is an inhabited subset by nullary interpolation in the basis and a semidirected lower set because the criteria of Lemma 6.25 are satisfied when taking \prec to be \ll_{β} . Secondly, the map $\downarrow_{\beta}(-)$ is Scott continuous by Lemma 6.23.

Now notice that the map $\beta : (B, \ll_{\beta}) \rightarrow D$ is monotone and that the Scott continuous map it induces by Theorem 6.16 is exactly the map $\bigsqcup : \mathcal{V}\text{-Idl}(B, \ll_{\beta}) \rightarrow D$ that takes an ideal I to the supremum of its associated directed family $\beta \circ \text{pr}_1 : (\Sigma_{b:B} (b \in I)) \rightarrow D$.

Since β is a basis for D , we know that $\bigsqcup \downarrow_{\beta} x = x$ for every $x : D$. So it only remains to show that $\downarrow_{\beta} \circ \bigsqcup$ is the identity on $\mathcal{V}\text{-Idl}(B, \ll_{\beta})$, for which we will use Lemma 6.24. So suppose that $I : \mathcal{V}\text{-Idl}(B, \ll_{\beta})$ is arbitrary. Then we only need to prove that

- (i) the conjunction of $\beta(b) \sqsubseteq \beta(c)$ and $c \in I$ implies $b \in I$ for every $b, c : B$;
- (ii) for every $b \in I$, there exists $c \in I$ such that $\beta(b) \ll \beta(c)$.

Note that (ii) is just saying that I is a rounded ideal w.r.t. \ll_β , so this holds. For (i), suppose that $\beta(b) \sqsubseteq \beta(c)$ and $c \in I$. By roundedness of I , there exists $c' \in I$ such that $c \ll_\beta c'$. But then $\beta(b) \sqsubseteq \beta(c) \ll \beta(c')$, so that $b \ll_\beta c'$ which implies that $b \in I$, because ideals are lower sets. \square

6.3.2. *Ideal completion with respect to the order relation.*

Lemma 6.29. *If $\beta : B \rightarrow D$ is a small basis for a \mathcal{V} -dcpo D , then (B, \sqsubseteq_β) is an abstract \mathcal{V} -basis where $b \sqsubseteq_\beta c$ is defined as $\beta(b) \sqsubseteq \beta(c)$.*

Proof. The relation \sqsubseteq_β is reflexive, so this follows from Lemma 6.12. \square

Remark 6.30. The definition of an abstract \mathcal{V} -basis requires the relation on it to be \mathcal{V} -valued. Hence, for the lemma to make sense we appeal to Proposition 5.5 to know that D is locally small which tells us that we can substitute $\beta(b) \sqsubseteq \beta(c)$ by an equivalent type in \mathcal{V} .

Theorem 6.31. *The map $\downarrow_\beta(-) : D \rightarrow \mathcal{V}\text{-Idl}(B, \sqsubseteq_\beta)$ is the embedding in an embedding-projection pair. In particular, D is a Scott continuous retract of the algebraic dcpo $\mathcal{V}\text{-Idl}(B, \sqsubseteq_\beta)$ that has a small compact basis. Moreover, if β is a small compact basis, then the map is an isomorphism.*

Proof. First of all, we should check that the map is well-defined, i.e. that $\downarrow_\beta x$ is an (B, \sqsubseteq_β) -ideal. It is an inhabited subset by nullary interpolation in the basis and a semidirected lower set because the criteria of Lemma 6.25 are satisfied when taking \prec to be \sqsubseteq_β . Secondly, the map $\downarrow_\beta(-)$ is Scott continuous by Lemma 6.23.

Now notice that the map $\beta : (B, \sqsubseteq_\beta) \rightarrow D$ is monotone and that the continuous map it induces by Theorem 6.16 is exactly the map $\bigsqcup : \mathcal{V}\text{-Idl}(B, \sqsubseteq_\beta) \rightarrow D$ that takes an ideal I to the supremum of the directed family $\beta \circ \text{pr}_1 : (\Sigma_{b:B}(b \in I)) \rightarrow D$.

Since β is a basis for D , we know that $\bigsqcup \downarrow_\beta x = x$ for every $x : D$. So it only remains to show that $\downarrow_\beta \circ \bigsqcup$ is a deflation, for which we will use Lemma 6.24. So suppose that $I : \mathcal{V}\text{-Idl}(B, \sqsubseteq_\beta)$ is arbitrary. Then we only need to prove that the conjunction of $\beta(b) \sqsubseteq \beta(c)$ and $c \in I$ implies $b \in I$, but this holds, because I is a lower set with respect to \sqsubseteq_β .

Finally, assume that β is a small compact basis. We show that $\downarrow_\beta \circ \bigsqcup$ is also inflationary in this case. So let I be an arbitrary ideal. By Lemma 6.24 it is enough to show that for every $b \in I$, there exists $c \in I$ such that $\beta(b) \ll \beta(c)$. But by assumption, $\beta(b)$ is compact, so we can simply take c to be b . \square

Combining Theorem 5.10, Theorems 6.11 and 6.28, and Theorems 6.15 and 6.31, we obtain the following result:

Corollary 6.32.

- (i) *A \mathcal{V} -dcpo has a small basis if and only if it is isomorphic to $\mathcal{V}\text{-Idl}(B, \prec)$ for an abstract basis (B, \prec) .*
- (ii) *A \mathcal{V} -dcpo has a small compact basis if and only if it is isomorphic to $\mathcal{V}\text{-Idl}(B, \prec)$ for an abstract basis (B, \prec) where \prec is reflexive.*
- (iii) *A \mathcal{V} -dcpo has a small basis if and only if it is a Scott continuous retract of a \mathcal{V} -dcpo with a small compact basis.*

In particular, every \mathcal{V} -dcpo with a small basis is isomorphic to one whose order takes values in \mathcal{V} and whose carrier lives in \mathcal{V}^+ .

7. BILIMITS AND EXPONENTIALS

7.1. Structurally continuous and algebraic bilimits. We show that bilimits are closed under equipment with continuity/algebraicity data. For the remainder of this section, fix a directed diagram of \mathcal{V} -dcpo's $(D_i)_{i:I}$ with embedding-projection

pairs $(\varepsilon_{i,j}, \pi_{i,j})_{i \sqsubseteq j \text{ in } I}$ between them, as in Section 7 of Part I. We stress that, throughout this section, the word “embedding” is only used in the domain-theoretic sense, i.e. it is reserved for one half of an embedding-projection pair, rather than in the homotopy type theory sense of having subsingleton fibers.

Now suppose that for every $i : I$, we have $\alpha_i : J_i \rightarrow D_i$ with each $J_i : \mathcal{V}$. Then we define $J_\infty \equiv \Sigma_{i:I} J_i$ and $\alpha_\infty : J_\infty \rightarrow D_\infty$ by $(i, j) \mapsto \varepsilon_{i,\infty}(\alpha_i(j))$, where $\varepsilon_{i,\infty}$ is as in Definition 7.12 of Part I.

Lemma 7.1. *If every α_i is directed and we have $\sigma : D_\infty$ such that α_i approximates σ_i , then α_∞ is directed and approximates σ .*

Proof. Observe that α_∞ is equal to the supremum, if it exists, of the directed families $(\varepsilon_{i,\infty} \circ \alpha_i)_{i:I}$ in the ind-completion of D_∞ , cf. the proof of Lemma 3.3. Hence, for directedness of α_∞ , it suffices to prove that the family $i \mapsto \varepsilon_{i,\infty} \circ \alpha_i$ is directed with respect to the exceeds-relation. The index type I is inhabited, because we are working with a directed diagram of dpos. For semidirectedness, we will first prove that if $i \sqsubseteq i'$, then $\varepsilon_{i',\infty} \circ \alpha_{i'}$ exceeds $\varepsilon_{i,\infty} \circ \alpha_i$.

So suppose that $i \sqsubseteq i'$ and $j : J_i$. As α_i approximates σ_i , we have $\alpha_i(j) \ll \sigma_i$. Because $\varepsilon_{i,i'}$ is an embedding, it preserves the way-below relation (Lemma 2.16), so that we get $\varepsilon_{i,i'}(\alpha_i(j)) \ll \varepsilon_{i,i'}(\sigma_i) \sqsubseteq \sigma_{i'} = \bigsqcup \alpha_{i'}$. Hence, there exists $j' : J_{i'}$ with $\varepsilon_{i,i'}(\alpha_i(j)) \sqsubseteq \alpha_{i'}(j')$ which yields $\varepsilon_{i,\infty}(\alpha_i(j)) = \varepsilon_{i',\infty}(\varepsilon_{i,i'}(\alpha_i(j))) \sqsubseteq \varepsilon_{i',\infty}(\alpha_{i'}(j'))$, completing the proof that $\varepsilon_{i',\infty} \circ \alpha_{i'}$ exceeds $\varepsilon_{i,\infty} \circ \alpha_i$.

Now to prove that the family $i \mapsto \varepsilon_{i,\infty} \circ \alpha_i$ is semidirected with respect to the exceeds-relation, suppose we have $i_1, i_2 : I$. Since I is a directed preorder, there exists $i : I$ such that $i_1, i_2 \sqsubseteq i$. But then $\varepsilon_{i,\infty} \circ \alpha_i$ exceeds both $\varepsilon_{i_1,\infty} \circ \alpha_{i_1}$ and $\varepsilon_{i_2,\infty} \circ \alpha_{i_2}$ by the above.

Thus, α_∞ is directed. To see that its supremum is σ , observe that

$$\begin{aligned} \sigma &= \bigsqcup_{i:I} \varepsilon_{i,\infty}(\sigma_i) && \text{(by Lemma 7.19 of Part I)} \\ &= \bigsqcup_{i:I} \varepsilon_{i,\infty}(\bigsqcup \alpha_i) && \text{(since } \alpha_i \text{ approximates } \sigma_i) \\ &= \bigsqcup_{i:I} \bigsqcup \varepsilon_{i,\infty} \circ \alpha_i && \text{(by Scott continuity of } \varepsilon_{i,\infty}) \\ &= \bigsqcup_{(i,j):J_\infty} \alpha_\infty(i, j), \end{aligned}$$

as desired.

Finally, we wish to show that $\alpha_\infty(i, j) \ll \sigma$ for every $(i, j) : J_\infty$. But $\varepsilon_{i,\infty}$ is an embedding and therefore preserves the way-below relation while $\alpha_i(j)$ approximates σ_i , so we get $\alpha_\infty(i, j) \equiv \varepsilon_{i,\infty}(\alpha_i(j)) \ll \varepsilon_{i,\infty}(\sigma_i) \sqsubseteq \sigma$ where the final inequality holds because $\varepsilon_{i,\infty} \circ \pi_{i,\infty}$ is a deflation. \square

Lemma 7.2. *If $\alpha_i(j)$ is compact for every $i : I$ and $j : J_i$, then all the values of α_∞ are compact too.*

Proof. Let $(i, j) : J_\infty$ be arbitrary. Since $\varepsilon_{i,\infty}$ is an embedding it preserves compact elements, so $\alpha_\infty(i, j) \equiv \varepsilon_{i,\infty}(\alpha_i(j))$ is compact. \square

Theorem 7.3. *If each D_i comes equipped with continuity (resp. algebraicity) data, then we can give continuity (resp. algebraicity) data for D_∞ .*

Proof. Let $\sigma : D_\infty$ be arbitrary. By assumption on each D_i , we have a directed family $\alpha_i : J_i \rightarrow D_i$ approximating σ_i . Hence, by Lemma 7.1, the family α_∞ is directed and approximates σ , giving continuity data for D_∞ . For the algebraic case, we apply the above and Lemma 7.2. \square

Note that we do not expect to be able to prove that D_∞ is continuous if each D_i is, because it would require an instance of the axiom of choice to get continuity data on each D_i , and without those we have nothing to operate on.

Theorem 7.4. *If each D_i has a small basis $\beta_i : B_i \rightarrow D_i$, then the map*

$$\begin{aligned} \beta_\infty : (B_\infty \equiv \Sigma_{i:I} B_i) &\rightarrow D_\infty \\ (i, b) &\mapsto \varepsilon_{i,\infty}(\beta_i(b)) \end{aligned}$$

is a small basis for D_∞ . Furthermore, if each β_i is a small compact basis, then β_∞ is a small compact basis too.

Proof. First of all, we must show that the proposition $\beta_\infty(i, b) \ll \sigma$ is small for every $i : I$, $b : B_i$ and $\sigma : D_\infty$. We claim that this is the case as the way-below relation on D_∞ has small values. By Proposition 4.12 and Theorem 7.3, it suffices to prove that D_∞ is locally small. But this holds by Proposition 7.22 of Part I as each D_i is locally small by Proposition 5.5.

It remains to prove that, for an arbitrary element $\sigma : D_\infty$, the family $\downarrow_{\beta_\infty} \sigma$ given by $(\Sigma_{(i,b):B_\infty} \beta_\infty(i, b) \ll \sigma) \xrightarrow{\beta_\infty \circ \text{pr}_1} D_\infty$ is directed with supremum σ . Note that for every $i : I$ and $b : B_i$, we have that $\beta_i(b) \ll \sigma_i$ implies

$$\beta_\infty(i, b) \equiv \varepsilon_{i,\infty}(\beta_i(b)) \ll \varepsilon_{i,\infty}(\sigma_i) \sqsubseteq \sigma,$$

since Lemma 2.16 tells us that the embedding $\varepsilon_{i,\infty}$ preserves the way-below relation. Hence, the identity map induces a well-defined map

$$\iota : (\Sigma_{i:I} \Sigma_{b:B_i} \beta_i(b) \ll \sigma_i) \rightarrow (\Sigma_{(i,b):B_\infty} \beta_\infty(i, b) \ll \sigma).$$

Lemma 5.9 now tells us that we only need to show that $\downarrow_{\beta_\infty} \sigma \circ \iota$ is directed and has supremum σ . But if we write $\alpha_i : (\Sigma_{b:B_i} \beta_i(b) \ll \sigma_i) \rightarrow D_i$ for the map $b \mapsto \beta_i(b)$, then we see that $\downarrow_{\beta_\infty} \sigma \circ \iota$ is given by α_∞ defined from α_i as in the start of this section. But then α_∞ is indeed seen to be directed with supremum σ by virtue of Lemma 7.1 and the fact that α_i approximates σ_i .

Finally, if every β_i is a small compact basis, then β_∞ is also a small compact basis because by Lemma 5.15 all we need to know is that $\beta_\infty(i, b) \equiv \varepsilon_{i,\infty}(\beta_i(b))$ is compact for every $i : I$ and $b : B_i$. But this follows from the fact that embeddings preserve compactness and that each $\beta_i(b)$ is compact. \square

7.2. Exponentials with small (compact) bases. Just as in the classical, impredicative setting, the exponential of two continuous dcpos need not be continuous [Jun89]. However, with some work, we are able to show that E^D has a small basis provided that both D and E do and that E has all (not necessarily directed) \mathcal{V} -suprema, that is, E is a continuous lattice. We first establish this for small compact bases using *step functions* and then derive the result for compact bases using Theorem 6.31.

7.2.1. Single step functions. Suppose that we have a dcpo D and a pointed dcpo E . Classically [Gie+03, Exercise II-2.31], the single step function given by $d : D$ and $e : E$ is defined as

$$\begin{aligned} \langle d \Rightarrow e \rangle : D &\rightarrow E \\ x &\mapsto \begin{cases} e & \text{if } d \sqsubseteq x; \\ \perp & \text{otherwise.} \end{cases} \end{aligned}$$

Constructively, we can't expect to make this case distinction in general, so we define single step functions using subsingleton suprema instead.

Definition 7.5 (Single step function, $\langle d \Rightarrow e \rangle$). The *single step function* given by two elements $d : D$ and $e : E$, where D is a locally small \mathcal{V} -dcpo and E is a pointed \mathcal{V} -dcpo, is the function $\langle d \Rightarrow e \rangle : D \rightarrow E$ that maps a given $x : D$ to the supremum of the family indexed by the subsingleton $d \sqsubseteq x$ that is constantly e .

Note that we need the domain D to be locally small, because we need the type $d \sqsubseteq x$ to be a subsingleton in \mathcal{V} to use the \mathcal{V} -directed-completeness of E . For the definition of $\langle d \Rightarrow e \rangle$ to make sense, we need to know that the supremum of the constant family exists. This is the case by Lemma 4.9 of Part I, which says that the supremum of a subsingleton-indexed family $\alpha : P \rightarrow E$ is given by the supremum of the directed family $\mathbf{1} + P \rightarrow E$ defined by $\text{inl}(\star) \mapsto \perp$ and $\text{inr}(p) \mapsto \alpha(p)$.

Lemma 7.6. *If $d : D$ is compact, then $\langle d \Rightarrow e \rangle$ is Scott continuous for every $e : E$.*

Proof. Suppose that $d : D$ is compact and that $\alpha : I \rightarrow D$ is a directed family. We first show that $\langle d \Rightarrow e \rangle(\bigsqcup \alpha)$ is an upper bound of $\langle d \Rightarrow e \rangle \circ \alpha$. So let $i : I$ be arbitrary. Then we have to prove $\bigsqcup_{d \sqsubseteq \alpha_i} e \sqsubseteq \langle d \Rightarrow e \rangle \circ \alpha$. Since the supremum gives a lower bound of the upper bounds, it suffices to prove that $e \sqsubseteq \langle d \Rightarrow e \rangle \circ \alpha$ whenever $d \sqsubseteq \alpha_i$. But in this case we have $e = \langle d \Rightarrow e \rangle(\alpha_i) \sqsubseteq \langle d \Rightarrow e \rangle \circ \alpha$, so we are done.

To see that $\langle d \Rightarrow e \rangle(\bigsqcup \alpha)$ is a lower bound of the upper bounds, suppose that we are given $y : E$ such that y is an upper bound of $\langle d \Rightarrow e \rangle \circ \alpha$. We are to prove that $(\bigsqcup_{d \sqsubseteq \alpha} e) \sqsubseteq y$. Note that it suffices for $d \sqsubseteq \bigsqcup \alpha$ to imply $e \sqsubseteq y$. So assume that $d \sqsubseteq \bigsqcup \alpha$. By compactness of d there exists $i : I$ such that $d \sqsubseteq \alpha_i$ already. But then $e = \langle d \Rightarrow e \rangle(\alpha_i) \sqsubseteq y$, as desired. \square

Lemma 7.7. *A Scott continuous function $f : D \rightarrow E$ is above the single step function $\langle d \Rightarrow e \rangle$ with $d : D$ compact if and only if $e \sqsubseteq f(d)$.*

Proof. Suppose that $\langle d \Rightarrow e \rangle \sqsubseteq f$. Then $\langle d \Rightarrow e \rangle(d) = e \sqsubseteq f(d)$, proving one implication. Now assume that $e \sqsubseteq f(d)$ and let $x : D$ be arbitrary. To prove that $\langle d \Rightarrow e \rangle(x) \sqsubseteq f(x)$, it suffices that $e \sqsubseteq f(x)$ whenever $d \sqsubseteq x$. But if $d \sqsubseteq x$, then $e \sqsubseteq f(d) \sqsubseteq f(x)$ by monotonicity of f . \square

Lemma 7.8. *If d and e are compact, then so is $\langle d \Rightarrow e \rangle$ in the exponential E^D .*

Proof. Suppose that we have a directed family $\alpha : I \rightarrow E^D$ such that $\langle d \Rightarrow e \rangle \sqsubseteq \bigsqcup \alpha$. Then we consider the directed family $\alpha^d : I \rightarrow E$ given by $i \mapsto \alpha_i(d)$. We claim that $e \sqsubseteq \bigsqcup \alpha^d$. Indeed, by Lemma 7.7 and our assumption that $\langle d \Rightarrow e \rangle \sqsubseteq \bigsqcup \alpha$ we get $e \sqsubseteq (\bigsqcup \alpha)(d) = \bigsqcup \alpha^d$. Now by compactness of e , there exists $i : I$ such that $e \sqsubseteq \alpha^d(i) \equiv \alpha_i(d)$ already. But this implies $\langle d \Rightarrow e \rangle \sqsubseteq \alpha_i$ by Lemma 7.7 again, finishing the proof. \square

7.2.2. Exponentials with small compact bases. Fix \mathcal{V} -dcpo's D and E with small compact bases $\beta_D : B_D \rightarrow D$ and $\beta_E : B_E \rightarrow E$ and moreover assume that E has suprema for all (not necessarily directed) families indexed by types in \mathcal{V} . We are going to construct a small compact basis on the exponential E^D .

Lemma 7.9. *If E is sup-complete, then every Scott continuous function $f : D \rightarrow E$ is the supremum of the collection of single step functions $(\langle \beta_D(b) \Rightarrow \beta_E(c) \rangle)_{b : B_D, c : B_E}$ that are below f .*

Proof. Note that f is an upper bound by definition, so it remains to prove that it is the least. Therefore suppose we are given an upper bound $g : D \rightarrow E$. We have to prove that $f(x) \sqsubseteq g(x)$ for every $x : D$, so let $x : D$ be arbitrary. Now $x = \bigsqcup \downarrow_{\beta_D} x$, because β_D is a small compact basis for D , so by Scott continuity of f and g , it suffices to prove that $f(\beta_D(b)) \sqsubseteq g(\beta_D(b))$ for every $b : B_D$. So let $b : B_D$ be arbitrary. Since β_E is a small compact basis for E , we have $f(\beta_D(b)) = \bigsqcup \downarrow_{\beta_E} f(\beta_D(b))$. So to prove $f(\beta_D(b)) \sqsubseteq g(\beta_D(b))$ it is enough to know that $\beta_E(c) \sqsubseteq g(\beta_D(b))$ for every $c : B_E$ with $\beta_E(c) \sqsubseteq f(\beta_D(b))$. But for such $c : B_E$ we have $\langle \beta_D(b) \Rightarrow \beta_E(c) \rangle \sqsubseteq f$ and therefore $\langle \beta_D(b) \Rightarrow \beta_E(c) \rangle \sqsubseteq g$ because f is an upper bound of such single step functions, and hence $\beta_E(c) \sqsubseteq g(\beta_D(b))$ by Lemma 7.7, as desired. \square

Definition 7.10 (Directification). In a poset P with finite joins, the *directification* of a family $\alpha : I \rightarrow P$ is the family $\bar{\alpha} : \text{List}(I) \rightarrow P$ that maps a finite list to the join of its elements. It is clear that $\bar{\alpha}$ has the same supremum as α , and by concatenating lists, one sees that the directification yields a directed family, hence the name.

Lemma 7.11. *If each element of a family into a sup-complete dcpo is compact, then so are all elements of its directification.*

Proof. The supremum of the empty list is \perp by Example 2.4, and hence the join of finitely many compact elements is compact by Lemma 2.7. \square

Let us write $\sigma : B_D \times B_E \rightarrow E^D$ for the map that takes (b, c) to the single step function $(\beta_D(b) \Rightarrow \beta_E(c))$ and

$$\beta : B := \text{List}(B_D \times B_E) \rightarrow E^D$$

for its directification, which exists because E^D is \mathcal{V} -sup-complete as E is and suprema are calculated pointwise.

Theorem 7.12. *The map β is a small compact basis for the exponential E^D , where E is assumed to be sup-complete.*

Proof. Firstly, every element in the image of β is compact by Lemmas 7.8 and 7.11. Secondly, for every $b : B$ and Scott continuous map $f : D \rightarrow E$, the type $\beta(b) \sqsubseteq f$ is small, because E^D is locally small by Proposition 5.11. Thirdly, for every such f , the family

$$(\Sigma_{b:B}(\beta(b) \sqsubseteq f)) \xrightarrow{\beta \circ \text{pr}_1} E^D$$

is directed because β is the directification of σ . Finally, this family has supremum f by Lemma 7.9. \square

7.2.3. *Exponentials with small bases.* We now present a variation of Theorem 7.12 but for (sup-complete) dcpos with small bases. In fact, we will prove it using Theorem 7.12 and the theory of Scott continuous retracts (Theorem 6.31 in particular).

Definition 7.13 (Closure under finite joins). A small basis $\beta : B \rightarrow D$ for a sup-complete poset is *closed under finite joins* if we have $b_\perp : B$ with $\beta(b_\perp) = \perp$ and a map $\vee : B \rightarrow B \rightarrow B$ such that $\beta(b \vee c) = \beta(b) \vee \beta(c)$ for every $b, c : B$.

Lemma 7.14. *If D is a sup-complete dcpo with a small basis $\beta : B \rightarrow D$, then the directification of β is also a small basis for D . Moreover, by construction, it is closed under finite joins.*

Proof. Since β is a small basis for D , it follows by Proposition 5.5 that the way-below relation on D is small-valued. Hence, writing $\bar{\beta}$ for the directification of β , it remains to prove that $\downarrow_{\bar{\beta}} x$ is directed with supremum x for every $x : D$. But this follows easily from Lemma 5.9, because $\downarrow_\beta x$ is directed with supremum x and this family is equal to the composite

$$(\Sigma_{b:B}(\beta(b) \ll x)) \xrightarrow{b \mapsto [b]} (\Sigma_{l:\text{List}(B)}(\bar{\beta}(l) \ll x)) \xrightarrow{\bar{\beta} \circ \text{pr}_1} D,$$

where $[b]$ denotes the singleton list. \square

Lemma 7.15. *If D is a \mathcal{V} -sup-complete poset with a small basis $\beta : B \rightarrow D$ closed under finite joins, then the ideal-completion $\mathcal{V}\text{-Idl}(B, \sqsubseteq)$ is \mathcal{V} -sup-complete too.*

Proof. Since the \mathcal{V} -ideal completion is \mathcal{V} -directed complete, it suffices to show that $\mathcal{V}\text{-Idl}(B, \sqsubseteq)$ has finite joins, because then we can turn an arbitrary small family into a directed one via Definition 7.10. As $\beta : B \rightarrow D$ is closed under finite joins, we have $b_\perp : B$ with $\beta(b_\perp) = \perp$ and we easily see that $\{b \in B \mid \beta(b) \sqsubseteq \beta(b_\perp)\}$ is the

least element of $\mathcal{V}\text{-Idl}(B, \sqsubseteq)$. Now suppose that we have two ideals $I, J : \mathcal{V}\text{-Idl}(B, \sqsubseteq)$ and consider the subset K defined as

$$K := \{b \in B \mid \exists_{c \in I} \exists_{d \in J} (\beta(b) \sqsubseteq \beta(c \vee d))\}.$$

Notice that K is clearly a lower set. Also note that ideals are closed under finite joins as they are directed lower sets. Hence, $b_\perp \in I$ and $b_\perp \in J$, so that $b_\perp \in K$ and K is inhabited. For semidirectedness, assume $b_1, b_2 \in K$ so that there exist $c_1, c_2 \in I$ and $d_1, d_2 \in J$ with $\beta(b_1) \sqsubseteq \beta(c_1 \vee d_1)$ and $\beta(b_2) \sqsubseteq \beta(c_2 \vee d_2)$. Then $b_1 \vee b_2 \in K$, because $c_1 \vee c_2 \in I$ and $d_1 \vee d_2 \in J$ and $\beta(b_1 \vee b_2) \sqsubseteq \beta((c_1 \vee c_2) \vee (d_1 \vee d_2))$. Hence, K is a directed lower set. We claim that K is the join of I and J . First of all, I and J are both subsets of K , so it remains to prove that K is the least upper bound. To this end, suppose that we have an ideal L that includes I and J , and let $b \in K$ be arbitrary. Then there exist $c \in I$ and $d \in J$ with $\beta(b) \sqsubseteq \beta(c \vee d)$. But $I, J \subseteq L$ and ideals are closed under finite joins, so $c \vee d \in L$, which implies that $b \in L$ since L is a lower set. Therefore, K is the least upper bound of I and J , completing the proof. \square

Theorem 7.16. *The exponential E^D of dcpos has a specified small basis if D and E have specified small bases and E is sup-complete.*

Proof. Suppose that $\beta_D : B_D \rightarrow D$ and $\beta_E : B_E \rightarrow E$ are small bases and that E is sup-complete. By Lemma 7.14 we can assume that $\beta_E : B_E \rightarrow E$ is closed under finite joins. We will write \overline{D} and \overline{E} for the respective ideal completions $\mathcal{V}\text{-Idl}(B_D, \sqsubseteq)$ and $\mathcal{V}\text{-Idl}(B_E, \sqsubseteq)$. Then Theorem 6.31 tells us that we have Scott continuous retracts

$$D \begin{array}{c} \xrightarrow{s_D} \\ \xleftarrow{r_D} \end{array} \overline{D} \text{ and } E \begin{array}{c} \xrightarrow{s_E} \\ \xleftarrow{r_E} \end{array} \overline{E}.$$

Composition yields a Scott continuous retract

$$E^D \begin{array}{c} \xrightarrow{s} \\ \xleftarrow{r} \end{array} \overline{E}^{\overline{D}}$$

where $s(f) := s_E \circ f \circ r_D$ and $r(g) := r_E \circ g \circ s_D$. Now \overline{D} and \overline{E} have small compact basis by Theorem 6.15 and \overline{E} is sup-complete by Lemma 7.15. Therefore, the exponential of \overline{D} and \overline{E} has a small basis by Theorem 7.12. Finally, Theorem 5.10 tells us that the retraction r yields a small basis on E^D , as desired. \square

Note how, unlike Theorem 7.12, the above theorem does not give a nice description of the small basis for the exponential when we unfold the definitions. It may be possible to do so using function graphs, as is done in the classical setting of effective domain theory in [Smy77, Section 4.1].

An application of the closure results of Theorems 7.4 and 7.12 is that Scott's D_∞ model of the untyped λ -calculus [Sco72] has a small compact basis. We recall that Scott's construction of iterating function spaces was ported to predicative univalent foundation in Part I.

Theorem 7.17. *Scott's D_∞ has a small compact basis and in particular is algebraic.*

Proof. By Example 5.18 the \mathcal{U}_0 -dcpo D_0 has a small compact basis. Moreover, it is not just a \mathcal{U}_0 -dcpo as it has suprema for all (not necessarily directed) families indexed by types in \mathcal{U}_0 , as D_0 is isomorphic to $\Omega_{\mathcal{U}_0}$. Hence, by induction it follows that each D_n is \mathcal{U}_0 -sup-complete. Therefore, by induction and Theorem 7.12 we get a small compact basis for each D_n . Thus, by Theorem 7.4, the bilimit D_∞ has a small basis too. \square

8. CONCLUDING REMARKS

Taking inspiration from work in category theory by Johnstone and Joyal [JJ82], we gave predicatively adequate definitions of continuous and algebraic dcpos, and discussed issues related to the absence of the axiom of choice. We also presented predicative adaptations of the notions of a basis and the rounded ideal completion. The theory was accompanied by several examples: we described canonical small compact bases for the lifting and the powerset, and considered the rounded ideal completion of the dyadics. We also showed that Scott’s D_∞ has a small compact basis and is thus algebraic in particular.

To prove that D_∞ had a small compact basis, we used that each D_n is a \mathcal{U}_0 -sup-lattice, so that we could apply the results of Section 7.2. Example 5.18 tells us that $\mathcal{L}_{\mathcal{U}_0}(\mathbb{N})$ has a small compact basis too, but to prove that the \mathcal{U}_0 -dcpos in the Scott model of PCF (see [dJon21b]) have small compact bases using the techniques of Section 7.2, we would need $\mathcal{L}_{\mathcal{U}_0}(\mathbb{N})$ to be a \mathcal{U}_0 -sup-lattice, but it isn’t. However, it is complete for *bounded* families indexed by types in \mathcal{U}_0 and we believe that is possible to generalise the results of Section 7.2 from sup-lattices to bounded complete posets. Classically, this is fairly straightforward, but from preliminary considerations it appears that constructively one needs to impose certain decidability criteria on the bases of the dcpos. For instance that the partial order is decidable when restricted to basis elements. We also studied such decidability conditions in our paper [dJon21a]. These conditions should be satisfied by the bases of the dcpos in the Scott model of PCF, but we leave a full treatment of bounded complete dcpos with bases satisfying such conditions for future investigations.

9. ACKNOWLEDGEMENTS

We are grateful to Andrej Bauer and Vincent Rahli for valuable feedback, and to Ayberk Tosun for discussions on algebraic dcpos in univalent foundations and for the results in Section 5.3 in particular. We also thank Jon Sterling for feedback on a previous version of this paper.

The first author was supported by Cambridge Quantum Computing and Ilyas Khan [Dissertation Fellowship in Homotopy Type Theory]; and The Royal Society [grant reference URF\R1\191055].

REFERENCES

- [Acz06] Peter Aczel. “Aspects of general topology in constructive set theory”. In: *Annals of Pure and Applied Logic* 137.1–3 (2006): *Papers presented at the 2nd Workshop on Formal Topology (2WFTop 2002)*. Ed. by Bernhard Banaschewski, Thierry Coquand, and Giovanni Sambin, pp. 3–29. DOI: [10.1016/j.apal.2005.05.016](https://doi.org/10.1016/j.apal.2005.05.016).
- [AET24] Igor Arrieta, Martín Hötzel Escardó, and Ayberk Tosun. *The Patch Topology in Univalent Foundations*. 2024. arXiv: [2402.03134](https://arxiv.org/abs/2402.03134) [cs.LG].
- [AJ94] Samson Abramsky and Achim Jung. “Domain theory”. In: *Handbook of Logic in Computer Science*. Ed. by S. Abramsky, Dov M. Gabray, and T. S. E. Maibaum. Vol. 3. Clarendon Press, 1994, pp. 1–168. Updated online version available at <https://www.cs.bham.ac.uk/~axj/pub/papers/handy1.pdf>.
- [AKS15] Benedikt Ahrens, Krzysztof Kapulkin, and Michael Shulman. “Univalent categories and the Rezk completion”. In: *Mathematical Structures in Computer Science* 25.5 (2015): *From type theory and homotopy theory to Univalent Foundations of Mathematics*, pp. 1010–1039. DOI: [10.1017/S0960129514000486](https://doi.org/10.1017/S0960129514000486).

- [dJE21] Tom de Jong and Martín Hötzel Escardó. “Domain Theory in Constructive and Predicative Univalent Foundations”. In: *29th EACSL Annual Conference on Computer Science Logic (CSL 2021)*. Ed. by Christel Baier and Jean Goubault-Larrecq. Vol. 183. Leibniz International Proceedings in Informatics (LIPIcs). Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2021, 28:1–28:18. DOI: [10.4230/LIPIcs.CSL.2021.28](https://doi.org/10.4230/LIPIcs.CSL.2021.28). Expanded version with full proofs available on arXiv: [2008.01422 \[math.LO\]](https://arxiv.org/abs/2008.01422).
- [dJE23] Tom de Jong and Martín Hötzel Escardó. “On Small Types in Univalent Foundations”. In: *Logical Methods in Computer Science* 19.2 (2023), 8:1–8:33. DOI: [10.46298/lmcs-19\(2:8\)2023](https://doi.org/10.46298/lmcs-19(2:8)2023).
- [dJon+23] Tom de Jong et al. “Set-Theoretic and Type-Theoretic Ordinals Coincide”. In: *2023 38th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*. IEEE, 2023. DOI: [10.1109/lics56636.2023.10175762](https://doi.org/10.1109/lics56636.2023.10175762). Publicly available as arXiv: [2301.10696](https://arxiv.org/abs/2301.10696).
- [dJon21a] Tom de Jong. “Sharp Elements and Apartness in Domains”. In: *37th Conference on the Mathematical Foundations of Programming Semantics (MFPS 2021)*. Ed. by Ana Sokolova. Vol. 351. Electronic Proceedings in Theoretical Computer Science. Open Publishing Association, 2021, pp. 134–151. DOI: [10.4204/EPTCS.351.9](https://doi.org/10.4204/EPTCS.351.9).
- [dJon21b] Tom de Jong. “The Scott model of PCF in univalent type theory”. In: *Mathematical Structures in Computer Science* 31.10 (2021): *Homotopy Type Theory 2019*, pp. 1270–1300. DOI: [10.1017/S0960129521000153](https://doi.org/10.1017/S0960129521000153).
- [dJon22] Tom de Jong. “Domain Theory in Constructive and Predicative Univalent Foundations”. PhD thesis. University of Birmingham, 2022. arXiv: [2301.12405 \[cs.LO\]](https://arxiv.org/abs/2301.12405). URL: <https://etheses.bham.ac.uk/id/eprint/13401/>.
- [dJon24a] Tom de Jong. “Domain theory in univalent foundations I: Directed complete posets and Scott’s D_∞ ”. 2024. arXiv: [2407.06952 \[cs.LO\]](https://arxiv.org/abs/2407.06952).
- [dJon24b] Tom de Jong. *Index for “Domain Theory in Univalent Foundations II: Continuous and algebraic domains”*. <https://www.cs.bham.ac.uk/~mhe/TypeTopology/DomainTheory.Part-II.html>. 2024. <https://github.com/martinescardo/TypeTopology/tree/master/source/DomainTheory/Part-II.lagda>, commit [1d83144](https://github.com/martinescardo/TypeTopology/commit/1d83144).
- [EK17] Martín H. Escardó and Cory M. Knapp. “Partial Elements and Recursion via Dominances in Univalent Type Theory”. In: *26th EACSL Annual Conference on Computer Science Logic (CSL 2017)*. Ed. by Valentin Goranko and Mads Dam. Vol. 82. Leibniz International Proceedings in Informatics (LIPIcs). Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2017, 21:1–21:16. DOI: [10.4230/LIPIcs.CSL.2017.21](https://doi.org/10.4230/LIPIcs.CSL.2017.21).
- [Esc+] Martín Hötzel Escardó et al. *TypeTopology*. <https://www.cs.bham.ac.uk/~mhe/TypeTopology/index.html>. AGDA development. <https://github.com/martinescardo/TypeTopology>, commit [1d83144](https://github.com/martinescardo/TypeTopology/commit/1d83144).
- [Esc+18] Martín Hötzel Escardó et al. *Ordinals in univalent type theory in Agda notation*. <https://www.cs.bham.ac.uk/~mhe/TypeTopology/Ordinals.index.html>. AGDAdevelopment. Since 2018. <https://github.com/martinescardo/TypeTopology/tree/master/source/Ordinals/index.lagda>, commit [1d83144](https://github.com/martinescardo/TypeTopology/commit/1d83144).
- [Esc19] Martín Hötzel Escardó. “Introduction to Univalent Foundations of Mathematics with Agda”. Nov. 2019. arXiv: [1911.00580 \[cs.LO\]](https://arxiv.org/abs/1911.00580).

- [Gie+03] G. Gierz et al. *Continuous Lattices and Domains*. Vol. 93. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2003. DOI: [10.1017/CBO9780511542725](https://doi.org/10.1017/CBO9780511542725).
- [GV72] A. Grothendieck and J. L. Verdier. “Prefaisceaux”. In: *Théorie des Topos et Cohomologie Etale des Schémas*. Tome 1 (Séminaire de Géométrie Algébrique du Bois Marie 1963/1964, SGA4). Ed. by A. Dold. Heidelberg and B. Eckmann. Vol. 269. Lecture Notes in Mathematics. Springer, 1972, pp. 1–217. DOI: [10.1007/BFB0081552](https://doi.org/10.1007/BFB0081552).
- [JJ82] Peter Johnstone and André Joyal. “Continuous categories and exponentiable toposes”. In: *Journal of Pure and Applied Algebra* 25.3 (1982), pp. 255–296. DOI: [10.1016/0022-4049\(82\)90083-4](https://doi.org/10.1016/0022-4049(82)90083-4).
- [Jun89] Achim Jung. *Cartesian Closed Categories of Domains*. Vol. 66. CWI Tracts. Centrum voor Wiskunde en Informatica (Centre for Mathematics and Computer Science), 1989. URL: <https://ir.cwi.nl/pub/13191>.
- [Kna18] Cory Knapp. “Partial Functions and Recursion in Univalent Type Theory”. PhD thesis. School of Computer Science, University of Birmingham, June 2018. URL: <https://etheses.bham.ac.uk/id/eprint/8448/>.
- [Kra+17] Nicolai Kraus et al. “Notions of Anonymous Existence in Martin-Löf Type Theory”. In: *Logical Methods in Computer Science* 13.1 (2017). DOI: [10.23638/LMCS-13\(1:15\)2017](https://doi.org/10.23638/LMCS-13(1:15)2017).
- [MP89] Michael Makkai and Robert Paré. *Accessible Categories: The Foundations of Categorical Model Theory*. Vol. 104. Contemporary Mathematics. American Mathematical Society, 1989. DOI: [10.1090/conm/104](https://doi.org/10.1090/conm/104).
- [Plo77] G.D. Plotkin. “LCF considered as a programming language”. In: *Theoretical Computer Science* 5.3 (1977), pp. 223–255. DOI: [10.1016/0304-3975\(77\)90044-5](https://doi.org/10.1016/0304-3975(77)90044-5).
- [Rij22] Egbert Rijke. “Introduction to Homotopy Type Theory”. Pre-publication of forthcoming textbook. 2022. arXiv: [2212.11082](https://arxiv.org/abs/2212.11082) [math.LO].
- [Sco70] Dana Scott. *Outline of a Mathematical Theory of Computation*. Tech. rep. PRG02. Oxford University Computing Laboratory, Nov. 1970. URL: <https://www.cs.ox.ac.uk/publications/publication3720-abstract.html>.
- [Sco72] Dana S. Scott. “Continuous lattices”. In: *Toposes, Algebraic Geometry and Logic. Dalhousie University, Halifax, January 16–19, 1971*. Ed. by F. W. Lawvere. Vol. 274. Lecture Notes in Mathematics. Springer, 1972, pp. 97–136. DOI: [10.1007/BFB0073967](https://doi.org/10.1007/BFB0073967).
- [Sco93] Dana S. Scott. “A type-theoretical alternative to ISWIM, CUCH, OWHY”. In: *Theoretical Computer Science* 121.1–2 (1993), pp. 411–440. DOI: [10.1016/0304-3975\(93\)90095-B](https://doi.org/10.1016/0304-3975(93)90095-B).
- [Smy77] M. B. Smyth. “Effectively given domains”. In: *Theoretical Computer Science* 5.3 (1977), pp. 257–274. DOI: [10.1016/0304-3975\(77\)90045-7](https://doi.org/10.1016/0304-3975(77)90045-7).
- [Uni13] The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. Institute for Advanced Study: <https://homotopytypetheory.org/book>, 2013.