Directed Transit Functions

Arun Anil¹, Manoj Changat¹, Lekshmi Kamal K-Sheela¹, Ameera Vaheeda Shanavas¹, John J. Chavara^{2,3}, Prasanth G. Narasimha-Shenoi^{4,5}, Bruno J. Schmidt^{6,7}, Peter F. Stadler^{7,8,6,9,10,11}
¹Department of Futures Studies, University of Kerala, Karyavattom Campus 695 581 Thiruvananthapuram, India.
²Department of Mathematics, University of Kerala, Karyavattom Campus 695 581 Thiruvananthapuram, India.
³Department of Mathematics, St. Berchmans College, Changanassery, Kottayam, 686101, India.

⁴Department of Mathematics, Government College, Chittur, Palakkad, 678 104, India.

⁵Department of Collegiate Education, Government of Kerala, Thiruvananthapuram, India.

⁶Max Planck Institute for Mathematics in the Sciences, Inselstraße 22, D-04103 Leipzig, Germany.

 $^7\mathrm{Bioinformatics}$ Group, Department of Computer Science &

Interdisciplinary Center for Bioinformatics, Leipzig University, Härtelstraße 16–18, D-04107 Leipzig, Germany

⁹Department of Theoretical Chemistry, University of Vienna, Währingerstraße 17, A-1090 Wien, Austria.
¹⁰Facultad de Ciencias, Universidad National de Colombia, Bogotá, Colombia.

¹¹Santa Fe Institute, 1399 Hyde Park Rd., Santa Fe, NM 87501, USA.

Contributing authors: arunanil93@gmail.com; mchangat@keralauniversity.ac.in; lekshmisanthoshgr@gmail.com; ameerasv@gmail.com; jjchavara@gmail.com; prasanthgns@gmail.com; bruno@bioinf.uni-leipzig.de; studla@bioinf.uni-leipzig.de;

Abstract

Transit functions were introduced as models of betweenness on undirected structures. Here we introduce directed transit function as the directed analogue on directed structures such as posets and directed graphs. We first show that betweenness in posets can be expressed by means of a simple set of first order axioms. Similar characterizations can be obtained for graphs with natural partial orders, in particular, forests, trees, and mangroves. Relaxing the acyclicity conditions leads to a generalization of the well-known geometric transit function to the directed structures. Moreover, we discuss some properties of the directed analogues of prominent transit functions, including the all-paths, induced paths, and shortest paths (or interval) transit functions. Finally we point out some open questions and directions for future work.

Keywords: Transit function; Directed Graph; Poset; Reachability; Interval Function; All-Path Function

1 Introduction

Transit functions $R: V \times V \to 2^V$ on a non-empty finite set V were introduced as an abstract model of betweenness, see [1]. In the most general form, R satisfies the three basic axioms (t1) $\{u, v\} \in R(u, v)$, (t2) R(u, v) = R(v, u), and (t3) $R(u, u) = \{u\}$. Important examples of transit functions are obtained by defining R(u, v) as the set of all vertices that lie on a certain class of paths and walks connecting u and v. This includes shortest paths [2, 3, 4], induced paths [5, 6, 7], all-paths [8], toll walks [9, 10], and weak toll walks [11] in graphs. In a very different context, transit functions also appear as descriptions of so-called clustering systems, in which each cluster C is the intersection of all clusters that contain a pair of suitably chosen points $x, y \in C$ [12, 13, 14, 15, 16, 17].

Recently, axioms (t1) and (t3) were relaxed in [18] to study the interval function of not necessarily connected undirected graphs, using the axioms (t1^{*}) $R(u, v) \neq \emptyset$ implies $u \in R(u, v)$, and (compt) $R(u, v) \neq \emptyset$ and $R(v, w) \neq \emptyset$ then $R(u, w) \neq \emptyset$. This generalization does not seem to have drawn much attention so far beyond the work reported in [18].

Interval functions satisfying (t1), (t2), and (t3) have been constructed for directed graphs (digraphs) in [19] as the union of all shortest directed path from u to v and v to u, setting $R(u, v) = \{u, v\}$ if no such path exists. This notion has been explored in particular for oriented (antisymmetric) graphs in relation to convexities derived from these intervals. Here, however, we will pursue a different avenue, considering also the intervals as directed structures.

Measures of betweenness, in particular betweenness centrality, i.e., the average fraction of shortest s-t-paths passing through a vertex v, have been studied extensively also for directed networks [20, 21]. Surprisingly, the theory of transit functions, so far has not been extended to directed structures.

Notions of paths connecting two points are not restricted to graphs. Directed hypergraphs have received increasing attention of constructive dynamical systems including

chemical reaction networks [22, 23]. As in graphs, hyperpaths are defined as alternating sequences of vertices and edges $(x_1, E_1, x_2, \ldots, x_{n-1}, E_n, x_n)$ subject to a wide variety of additional conditions, see e.g. [24, 25, 26].

A directed form of betweenness also arises from bubble structures in directed graphs, defined as sets of vertices reachable from an entrance s and from which an exit t can be reached. Such structures appear in conjunction with additional isolation properties e.g. in the context of genome assembly [27, 28, 29] and as a model of alternative splicing [30] in computational biology.

In this contribution we introduce transit functions for directed structures. As a very natural starting point we will first consider partial orders in Sect. 3. While the standard notion of betwenness for partial orders using symmetric transit functions, cannot be characterized by a single first-order sentence [31], its directed counterpart turns out to have a simple and very natural characterization. We explore the directed transit functions on posets further by describing general directed acyclic graphs (DAG) as well as some special subclasses such as rooted trees in more detail. In Sect. 4 we relax our assumptions and consider directed betweenness in general reachability relations. This leads us in particular to a natural generalization of Nebeský's geometric transit functions on graphs that have been studied in detail in the undirected case, namely the all-paths, the induced-path, and shortest-path transit functions. As we shall see, there are non-trivial differences between undirected and directed transit functions.

2 Notation

We consider finite, loopless, simple directed graphs (digraphs) G = (V, E) and write V(G) and E(G) to refer explicitly to the vertex and edge set of G. A directed edge from u to v in G will be denoted by $(u, v) \in E(G)$. A walk in G is an alternating sequence $x_0, e_1, x_1, \ldots, e_n, x_n$ of vertices $x_i \in V(G)$ and edges $e_i = (x_{i-1}, x_i) \in E(G)$. Since the edges are implied by the vertices, we will simply write (x_0, x_1, \ldots, x_n) for the walk starting from vertex x_0 and ending in the vertex x_n . The number n of edges is the length of a walk. A path is a walk such that $x_i \neq x_j$ for all $i \neq j$. A directed cycle is composed of a path (x_1, \ldots, x_n) and the edge (x_n, x_1) .

A digraph is symmetric if $(u, v) \in E(G)$ implies $(v, u) \in E(G)$. Symmetric digraphs can also be interpreted as undirected graphs. A digraph G is strongly connected if for any two vertices $u, v \in V(G)$ there exists a path from u to v and from v to u. It is unilaterally connected if for any two vertices $u, v \in V(G)$ there exists a path from u to v or from v to u. Finally, G is weakly connected if for any two vertices $u, v \in V(G)$ there is a sequence $(u = x_0, x_1, \ldots, x_n = v)$ such that for $1 \le i \le n$ we have $(x_{i-1}, x_i) \in E(G)$ or $(x_i, x_{i-1}) \in E(G)$.

A vertex $x \in V(G)$ is a source if it has no incoming edges $(u, x) \in E(G)$, and a sink if it has no outgoing edges $(x, v) \in E(G)$. If G is a DAG, a vertex $x \in V(G)$ is called a hybrid vertex if it has more than one incoming edge. A source vertex $w \in V(G)$ is called root vertex, or simply root, if there is a path from w to every other vertex in V(G). A DAG is rooted (at w) if it contains a root vertex w. A graph containing a

root vertex is always weakly connected by definition. If $(x, y) \in E(G)$ in a DAG G, then y is called a successor of x, and x a predecessor of y.

H = (W, F) is a sub(di)graph of G = (V, E) if H is a digraph, $W \subseteq V$ and $F \subseteq E$. The subgraph H = G[W] is induced by W if for all $u, v \in W$ we have $(u, v) \in E$ if and only if $(u, v) \in F$. A strongly (or weakly) connected component of G is a maximal subset $W \subseteq V(G)$ such that the subgraph G[W] induced by W is strongly (or weakly) connected. A rooted (directed) tree is a rooted DAG without hybrid vertices. Thus any path between two vertices is unique. A (directed) forest \vec{F} is the disjoint union of rooted trees.

A partial order is a reflexive, antisymmetric, transitive relation on a non-empty set V, i.e., (i) $x \succeq x$ for all $x \in V$, (ii) $x \succeq y$ and $y \succeq x$ implies x = y for all $x, y \in V$, and (iii) $x \succeq z$ and $z \succeq y$ implies $x \succeq y$ for all $x, y, z \in V$.

3 Directed Transit Functions on Partially Ordered Sets

3.1 Motivation

Partially ordered sets (posets) are probably the simplest "directed" structures, comprising a set V endowed with a reflexive, transitive, and antisymmetric relation. Posets are associated with a straightforward notion of betweenness, which can be captured as follows:

Definition 1. Let (V, \succeq) be a non-empty finite partial order. Then the function R_{\succeq} : $V \times V \to 2^V$ define by

$$R_{\succeq}(u,v) \coloneqq \{x \in V | u \succeq x \succeq v\}$$

$$\tag{1}$$

is the directed transit function $of \succeq$.

For incomparable elements u and v, i.e., for u and v such that $u \not\geq v$ and $v \not\geq u$, we use $u \parallel v$. Note that $R_{\succeq}(u, v) = \emptyset$ if and only if $u \not\geq v$. Moreover, we have $u \parallel v$ if and only if $R_{\succeq}(u, v) = R_{\succeq}(v, u) = \emptyset$.

We will usually simply write R instead of R_{\succeq} when there is no danger of confusion. We note that *symmetric* transit functions also have been associated with posets [1]. Similar to eqn.(1), if u and v are comparable, $T(u, v) = \{x \in V | u \succeq x \succeq v\}$ if $u \succeq v$ and $T(v, u) = \{x \in V | v \succeq x \succeq u\}$ if $v \succeq u$. In contrast to our definition, however, $T(u, v) = \{u, v\}$ is used if u and v are incomparable. We use the empty set here in the same manner as in the generalized transit functions on disconnected graphs in [18]. **Lemma 1.** Let R be the transit function of a partial order. Then R satisfies the following statements:

- (t1) If $R(u, v) \neq \emptyset$ then $\{u, v\} \subseteq R(u, v)$ for all $u, v \in V$.
- (t2a) If $R(u,v) \neq \emptyset$ and $u \neq v$ then $R(v,u) = \emptyset$ for all $u, v \in V$.
- (t3) $R(u, u) = \{u\} \text{ for all } u \in V.$
- (tr1) If $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$ then $R(u, w) \cup R(w, v) \subseteq R(u, v)$ for all $u, v \in V$ with $u \neq v$.
- (tr2) If $R(u, w) = \emptyset$ or $R(w, v) = \emptyset$ then $w \notin R(u, v)$ for all $u, v \in V$ with $u \neq v$.

Proof. (t1) follows directly from the definition, and $u \succeq u$ yields $R(u, u) = \{u\}$ for all $u \in V$, i.e., (t3). Since $R(u, v) \neq \emptyset$ implies $u \succeq v$ and $u \neq v$, we have $v \not\succeq u$ and thus $R(u, v) = \emptyset$, and consequently (t2a) is satisfied. Similarly, $R(u, v) \neq \emptyset$ and $R(v, w) \neq \emptyset$ implies $u \succeq v$ and $v \succeq w$, which by transitivity of the partial order yields $u \succeq x \succeq v \succeq w$ and thus equ.(1) implies $R(u, v) \subseteq R(u, w)$. Analogously, we obtain $R(v, w) \subseteq R(u, w)$, and hence (tr1) is satisfied. Finally, $u \not\succeq x$ or $x \not\succeq v$ implies immediately that $x \notin R(u, v)$, and thus (tr2) holds.

We note that in the presence of (t2a), the statements of (tr1) and (tr2) trivially extends to the excluded cases u = v as an immediate consequence of (t3). We explicitly exclude u = v in the statement of the axioms (tr1) and (tr2) because this will be relevant later in the context of reachability relations.

Before we proceed we consider two simple consequences of (tr1).

Observation 2. Let $R: V \times V \to 2^V$ be a function satisfying (t3), and (tr1). Then R satisfies

(tr0) If $u \neq v$, $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$ then $w \in R(u, v)$.

(t0) If $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$ then $R(u, v) \neq \emptyset$.

Proof. If $u \neq v$, and $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$ then $w \in R(u, w) \cup R(w, v) \subseteq R(u, v) \neq \emptyset$, and thus (tr0) holds and the implication of (t0) is true. If u = v, then (t3) yields $u \in R(u, v) \neq \emptyset$, and hence (t0) holds for all $u, v \in V$.

We note that under the assumptions of Observation 2, (tr0) implies (t0), while the converse is not true. In [18], (t0) (there denoted by "compt"), (t1), and (t3) appear in this form for undirected, but not necessarily connected, graphs. The analogy of the basic properties of R in Lemma 1 on properties with symmetric transit functions is to be motivated by the following

Definition 2. Let V be a finite set. A directed transit function is a function $R : V \times V \rightarrow 2^V$ satisfying (t0), (t1) and (t3).

3.2 Characterization of the Directed Transit Functions of Posets

Given a directed transit function, we define a relation \succeq_R by $u \succeq_R v$ if and only if $R(u, v) \neq \emptyset$. Next, we show that the construction of R_{\succeq} preserves the partial order in the following sense:

Lemma 3. Let \succeq be a partial order and R_{\succeq} its directed transit function. Then $\succeq_{R_{\succ}} = \succeq$.

Proof. If $u \succeq v$ then $R_{\succeq}(u, v) \neq \emptyset$ and thus $u \succeq_{R_{\succeq}} v$. If $u \not\succeq v$ then $R_{\succeq}(u, v) = \emptyset$ and thus $u \not\succeq_{R_{\succ}} v$. Hence $\succeq_{R_{\succ}} = \succeq$.

Definition 3. A directed transit function is a poset function if it is antisymmetric, (t2a), and satisfies (tr1) and (tr2).

The main result of this section is that poset functions faithfully describe posets. **Theorem 4.** Let R be a poset function. Then the relation \succeq_R defined by $u \succeq_R v$ if and only if $R(u, v) \neq \emptyset$ is a partial order. Moreover $R_{\succeq_R} = R$.

Proof. First we note that by (t3) we have $u \succeq_R u$ for all $u \in V$. Hence \succeq_R is reflexive. Suppose $u \neq v$. Then $u \succeq_R v$ implies $R(u, v) \neq \emptyset$ and thus $R(v, u) = \emptyset$ by (t2a), i.e., $v \not\succeq u$, i.e., \succeq_R is antisymmetric. Finally assume that $u \succeq_R v$ and $v \succeq_R w$. Thus $R(u, v) \neq \emptyset$ and $R(v, w) \neq \emptyset$, whence (tr1) implies $R(u, w) \neq \emptyset$, and thus $u \succeq_R w$, i.e., \succeq_R is transitive. Thus \succeq_R is a partial order.

By definition, $x \in R_{\succeq_R}(u, v)$ if and only if $u \succeq_R x$ and $x \succeq_R v$, which is the case if and only if $R(u, x) \neq \emptyset$ and $R(x, v) \neq \emptyset$; thus (tr1), which implies $x \in R(u, v)$. If $x \notin R_{\succeq_R}(u, v)$ then $u \not\succeq_R x$ or $x \not\succeq_R v$, i.e., $R(u, x) = \emptyset$ or $R(x, v) = \emptyset$, which by (tr2) implies $x \notin R(u, v)$. Hence $R \succeq_R = R$.

Before we proceed we note that (tr1) and (tr2) can be expressed in the following, more compact form.

Lemma 5. Let R be a directed transit function. Then R satisfies (tr1) and (tr2) if and only R satisfies, for all $u, v, w \in V$ with $u \neq v$:

(tr) $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$ if and only if $w \in R(u, v)$

We remark that in the presence of (t3) and (t2a), the equivalence also remains valid for u = v.

Proof. Assume (tr1), (tr2). If $w \in R(u, v)$ and $R(u, w) = \emptyset$ then $w \notin R(u, v)$ by (tr2), a contradiction. The same argument can be repeated for $R(w, v) = \emptyset$, hence the if-part of (tr) hold. For the only-if part, assume $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$ and $w \notin R(u, v)$. By (tr1), we have $R(u, w) \cup R(w, v) \subseteq R(u, v)$ and since $w \in R(u, w), R(w, v)$ by (t1) we have $w \in R(u, v)$, a contradiction. Thus (tr1) and (tr2) imply (tr).

For the converse, assume (tr). For (tr1), assume $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$ and $R(u, w) \cup R(w, v) \not\subseteq R(u, v)$. By (tr), we know $w \in R(u, v)$ and by (t1), that $u, v \in R(u, v)$, hence there must be an $x \in R(u, w)$ or $x \in R(w, v)$ such that $x \notin R(u, v)$. If $x \in R(u, w)$ then $R(u, x) \neq \emptyset$, $R(x, w) \neq \emptyset$, and $R(w, v) \neq \emptyset$; Now $x \in R(x, v)$ and (tr) implies $x \in R(u, v)$, a contradiction. We argue analogously for $x \in R(w, u)$. (tr2) is the contrapositive of the only-if direction in (tr).

Since a poset function satisfies (tr1) and (tr2) we have

Remark 1. A poset function R always satisfies (tr)

The following property of poset functions will be convenient for the proofs in the following sections.

Lemma 6. Let R be a poset function. Then R satisfies, for all $u, v, w \in V$

(q) $R(u, w) \cap R(w, v) \in \{w, \emptyset\}$

Proof. (q). Suppose there is $x \in R(u, w) \cap R(w, v)$ with $x \neq w$. Since $x \in R(u, w)$ and $x \in R(w, v)$, the contrapositive of (tr2) implies $R(x, w) \neq \emptyset$ and $R(w, x) \neq \emptyset$, contradicting (t2a). Thus, if $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$ then $R(u, w) \cap R(w, v) = \{w\}$. Clearly, $R(u, w) \cap R(w, v) = \emptyset$ if (and only if) $R(u, w) = \emptyset$ and $R(w, v) = \emptyset$. \Box

3.3 Directed Acyclic Graphs

A graph G = (V, E) is acyclic if it does not contain a directed cycle. The directed paths in directed acyclic graph (DAG) define a partial order \succeq on the vertex set V

such that $u \succeq v$ if and only if there is a directed path from u to v. Thus the directed all-path transit function R_G of G coincides with R_{\succeq_G} . This begs the questions whether the DAG G can be identified from R_G .

A key concept linking transit functions with graphs is the *underlying graph* [1]. The idea directly carries over to the directed case:

Definition 4. Let R be a directed transit function on V. The underderlying digraph of R is $G_R = (X, E_R)$, where $(x, y) \in E_R$ if and only if $R(x, y) = \{x, y\}$ and $x \neq y$.

The underlying graph G_R is symmetric if R is symmetric (i.e., if R satisfies (t2) R(u,v) = R(v,u). In this case G_R corresponds to the undirected underlying graph defined for symmetric transit functions.

Lemma 7. Let R be a poset function. If $R(u,v) \neq \emptyset$ and $u \neq v$ then there are $x, y \in R(u,v)$ such that $R(u,x) = \{u,x\}$ and $R(y,v) = \{y,v\}$.

Proof. If $R(u, v) = \{u, v\}$ there is nothing to show. Otherwise, there is $w \in R(u, v) \setminus \{u, v\}$, (q) implies $R(u, w) \cap R(w, v) = \{w\}$, and thus $u \notin R(w, v)$ and $v \notin R(u, w)$. On the other hand, (tr1) ensures $R(u, w) \subseteq R(u, v)$ and $R(w, v) \subseteq R(u, v)$, and thus |R(u, w)|, |R(w, v)| < |R(u, v)|. Applying the argument separately to R(u, w) and R(w, v) yields, elements $x_2 \in R(u, w), y_2 \in R(w, v)$, with $R(u, x_2) \subsetneq R(u, w)$ and, $R(y_2, v) \subseteq R(w, v)$ and, thus a sequence of transit sets having elements $x_{k-1}, x_k, x_{k+1}, \ldots$, and $y_{\ell-1}, y_{\ell}, y_{\ell+1}, \ldots$, with $x_k \in R(u, x_{k-1}), x_{k+1} \in R(u, x_k)$ and $y_{\ell} \in R(y_{\ell-1}, v), y_{\ell+1} \in R(y_{\ell}, v)$, respectively, such that $R(u, x_{k+1}) \subsetneq R(u, x_k)$ and $R(y_{\ell+1}, v) \subsetneq R(y_{\ell}, v)$. By finiteness, these sequences, eventually terminate with $R(u, x) = \{u, x\}$ and $R(y, v) = \{y, v\}$.

Corollary 8. Let R be a poset function. If $w \in R(u, v) \neq \emptyset$ then there is a directed path in G_R from u to v that passes through w.

Proof. Let $w \in R(u, v)$. Then Lemma 7 ensures that there is $x_1 \in R(u, w)$ and $y_1 \in R(w, v)$ such that $(u, x_1) \in E(G_R)$, (y_1, v) in $E(G_R)$. Repeating the argument separately on $R(x_1, w)$ and $R(w, y_1)$ yields edges $(x_1, x_2) \in E(R_G)$ and $(y_1, y_2) \in E(R_G)$ and residual sets $R(x_2, w) \subseteq R(x_1, w)$ and $R(w, y_1) \subseteq R(w, y_2)$, which eventually yields a path $u, x_1, x_2 \dots, x_k, w, y_h, \dots, y_2, y_1, v$ in G_R .

Lemma 9. Let R be a poset function. If $R(u, v) = \emptyset$ then G_R contains no directed path from u to v.

Proof. Let $R(u, v) = \emptyset$ and assume for contradiction that there is a directed path $P = (u, z_1, \ldots, z_k, v)$. Then $R(u, z_1) = \{u, z_1\}$ and $R(z_1, z_2) = \{z_1, z_2\}$. Since R satisfies (tr) by Remark 1, this yield $z_1 \in R(u, z_2)$ and thus $\{u, z_1, z_2\} \subseteq R(u, z_2)$. We proceed by induction. Assume $\{u, z_1, \ldots, z_i\} \subseteq R(u, z_i)$. Using $R(z_i, z_{i+1}) = \{z_i, z_{i+1}\}$ yields $z_i \in R(u, z_{i+1})$ and by (tr1) $R(u, z_i) \cup \{z_i, z_{i+1}\} \subseteq R(u, z_{i+1})$. Thus, in particular R(u, v) contains all vertices of the path P, and thus Let $R(u, v) = \emptyset$ and assume, for contradiction, that there is a directed path $P = (u, z_1, \ldots, z_k, v)$. Then $R(u, z_1) = \{u, z_1\}$ and $R(z_1, z_2) = \{z_1, z_2\}$. Since R satisfies (tr), this yield $z_1 \in R(u, z_2)$ and thus $\{u, z_1, z_2\} \subseteq R(u, z_2)$. We proceed by induction. Assume $\{u, z_1, \ldots, z_i\} \subseteq R(u, z_i)$. Using $R(z_i, z_{i+1}) = \{z_i, z_{i+1}\}$ yields $z_i \in R(u, z_{i+1})$ and by (tr1) $R(u, z_i) \cup \{z_i, z_{i+1}\} \subseteq R(u, z_{i+1})$. Thus, in particular R(u, v) contains all vertices of the path P, and thus $R(u, z_i) \cup \{z_i, z_{i+1}\} \subseteq R(u, z_i)$. We proceed by induction. Assume $\{u, z_1, \ldots, z_i\} \subseteq R(u, z_i)$. Using $R(z_i, z_{i+1}) = \{z_i, z_{i+1}\}$ yields $z_i \in R(u, z_{i+1})$ and by (tr1) $R(u, z_i) \cup \{z_i, z_{i+1}\} \subseteq R(u, z_{i+1})$. Thus, in particular R(u, v) contains all vertices of the path P, and thus $R(u, v) \neq \emptyset$, a contradiction. □

Corollary 10. Let R be a poset function. Then the following statements are true:

(i) There is a directed path connecting u and v in G_R if and only if $R(u, v) \neq \emptyset$.

(ii) If $w \notin R(u, v)$ then there is no directed path from u to v through w in G_R .

(iii) The graph G_R is a DAG.

Proof. Statement (i) is an immediate consequence of Corollary 8 and Lemma 9. (ii) If $w \notin R(u, v)$ then the contrapositive of (tr) implies that $R(u, w) = \emptyset$ or $R(w, v) = \emptyset$. By Lemma 9, then there is no directed path from u to w or no directed path from w to v in G_R . (iii) following immediately from the fact that R defines a partial order by virtue of being a poset function.

Analogous to the symmetric case of the all-paths function of a undirected graph, the all paths function of a directed graph is defined as:

Definition 5. Let G be a directed graph with vertex set V. The all-paths transit function $A_G: V \times V \to 2^V$ of G is defined by

$$A_G(u,v) \coloneqq \{x \in V \mid x \text{ lies on a directed path from } u \text{ to } v\}$$

$$(2)$$

We are now in the position to state the main result of this section:

Theorem 11. Let R be a poset function, G_R its underlying graph, and A_{G_R} the all-paths transit function of G_R . Then G_R is a DAG and $R = A_{G_R}$.

Proof. It follows from Corollary 10 that G_R is a DAG. We have $w \in A_{G_R}(u, v)$ if and only if there is a directed path from u to v through w in G_R , which by Cor.8 and Cor. 10(ii) is the case if and only $w \in R(u, v)$. Thus $R(u, v) = A_{G_R}(u, v)$ for all $u, v \in V$.

We shall return to the function A_G in a more general setting in Section 4.3.

3.4 Shortcut free DAGs

The directed paths of a directed acyclic graph(DAG) G define a partial order \succeq_G and thus a poset function R_G . It is tempting to assume that G equals the underlying graph G_{R_G} of the directed transit function R_G . This is not true in full generality, however, and the obstructions are *shortcuts*, i.e., transitive edges in G [33]. More formally, if $P = (x_1, x_2, \ldots, x_k)$ is a path of length $k \ge 2$ in G, we call an edge (x_1, x_k) a *shortcut*. Obviously, shortcuts do not affect the set of vertices lying on paths from u to v, and $R_G = R_{G'}$ if G and G' differ only by the presence or absence of shortcut edges. In the following we denote the set of directed graphs without shortcuts by \mathfrak{G}° .

Lemma 12. Let G be a DAG. Then $G = G_{R_G}$ if and only if $G \in \mathfrak{G}^\circ$.

Proof. Assume that $G = G_{R_G}$. If $G \notin \mathfrak{G}^\circ$, then G contains a path $P = (x_1, x_2, x_3, \ldots, x_n)$ and the edge (x_1, x_n) . Then $P \subseteq R_G(x_1, x_n) \neq \{x_1, x_n\}$, and thus (x_1, x_n) is not an edge in G_{R_G} . Conversely, assume that $G \neq G_{R_G}$. Note that (x_1, x_2) is an arc in G_{R_G} if and only if $R_G(x_1, x_2) = \{x_1, x_2\}$. That is, (x_1, x_2) is an arc in G. Therefore, $G \neq G_{R_G}$ implies that there is an edge (x_1, x_2) in G that is not in G_{R_G} . Thus there is $x_3 \in R_G(x_1, x_2)$ such that $x_3 \notin \{x_1, x_2\}$. Hence, there is a directed path from x_1 to x_2 other than the edge (x_1, x_2) , and thus (x_1, x_2) is shortcut in G.

The graph G_{R_G} is obtained from G by removing all shortcuts. For DAG, it coincides with the unique *transitive reduction* of G [34] and the unique minimum equivalent graph [35].

Observation 13. For every directed acyclic graph G holds $G_{R_G} \in \mathfrak{G}^{\circ}$.

3.5 Connection to the undirected poset function

A (symmetric) transit function P has been associated with a poset (X, \leq) by defining the order intervals $[u, v] := \{x \in V | u \leq x \leq v \text{ or } v \leq x \leq u\}$ and setting P(u, v) := [u, v] if $u \leq v$ or $v \leq u$, and if both $u \not\leq v$ and $v \not\leq u$ then $P(u, v) = \{u, v\}$. The function P has a long history and was part of folklore in the study of order intervals and order convexity; for further references we refer to [36, 37, 38]. It satisfies a rich portfolio of properties that admit immediate generalizations to the non-symmetric case. In Lemma 14, we show that this is true in particular for the betweenness axioms most commonly studied for symmetric transit functions [39].

Lemma 14. Let R be a poset function. Then R satisfies the following properties

- (m) $x, y \in R(u, v)$ implies $R(x, y) \subseteq R(u, v)$
- (b1₁) $x \in R(u, v)$ and $x \neq v$ implies $v \notin R(u, x)$
- (b1₂) $x \in R(u, v)$ and $x \neq u$ implies $u \notin R(x, v)$
- (b2₁) $x \in R(u, v)$ implies $R(u, x) \subseteq R(u, v)$
- (b2₂) $x \in R(u, v)$ implies $R(x, v) \subseteq R(u, v)$
- (b3₁) $x \in R(u, v)$ and $y \in R(u, x)$ implies $x \in R(y, v)$
- (b3₂) $x \in R(u, v)$ and $y \in R(x, v)$ implies $x \in R(u, y)$
- (b4) If $x \in R(u, v)$ then $R(u, x) \cap R(x, v) = \{x\}$
- (Ch) $x \in R(u, v)$ and $y \in R(x, w)$ implies $y \in R(u, w)$ or $y \in R(v, w)$ or $y \in R(u, v)$.
- (j0) $x \in R(u, y)$ and $y \in R(x, v)$ implies $x \in R(u, v)$.

In the following we write $(b1_{1,2})$, $(b2_{1,2})$, or $(b2_2)$ if R satisfies both $(b1_1)$ and $(b1_2)$, both $(b2_1)$ and $(b2_2)$, or both $(b3_1)$ and $(b3_2)$, respectively.

Proof. If $R(x, y) = \emptyset$, there is nothing to show. Thus we assume $R(x, y) \neq \emptyset$. By (tr2), $x, y \in R(u, v)$ implies $R(u, x) \neq \emptyset$, $R(x, v) \neq \emptyset$, $R(u, y) \neq \emptyset$, and $R(y, v) \neq \emptyset$. By (tr1), $R(u, x) \neq \emptyset$ and $R(x, y) \neq \emptyset$ implies $R(x, y) \subseteq R(u, y) \neq \emptyset$, which together with $R(y, v) \neq \emptyset$ implies $R(u, y) \subseteq R(u, v)$, and thus also $R(x, y) \subseteq R(u, v)$.

 $(b1_{1,2})$ is a direct consequence of (q). The contrapositive of (tr2) and (tr1) yield $w \in R(u, v) \implies R(u, w) \cup R(w, v) \subseteq R(u, v)$. Together with (q) implies $(b2_{1,2})$.

To see (b3₁), we assume $x \notin R(y, v)$, whence by (tr), we have $R(y, x) = \emptyset$ or $R(x, v) = \emptyset$. The latter is impossible since $x \in R(u, v)$ and thus $R(y, x) = \emptyset$. However, we have $y \in R(u, x)$ and thus both $R(u, y) \neq \emptyset$ and $R(y, x) \neq \emptyset$, a contradiction. (b3₂) follows analogously.

(b4) is an immediate consequence of (q).

 $x \in R(u, v)$ and $y \in R(x, w)$ implies $R(u, x) \neq \emptyset$, $R(x, v) \neq \emptyset$, $R(x, y) \neq \emptyset$ and $R(y, w) \neq \emptyset$. Hence $y \in R(u, x) \cup R(x, y) \cup R(y, w) \subseteq R(u, w)$, and thus (Ch) holds.

Since $x \in R(u, y)$ and $y \in R(x, v)$ implies $R(u, x) \neq \emptyset$, $R(x, y) \neq \emptyset$ and $R(y, v) \neq \emptyset$, we obtain $x \in R(u, y) \cup R(y, v) \subseteq R(u, v)$ and thus (j0) is satisfied. Note that the analogs of axioms (b1), (b2), and (b3) come in pairs in the directed case. We shall return to the implications of the betweenness axioms in a more general context in Sect. 4.2. Axiom (Ch) first appeared in the context of convex geometries in [40]. It may be worth noting that one of the three alternatives appearing in this condition is always true. Axiom (j0) was initially introduced in [7] and also plays a role in the study of convex geometries. It remains an open question whether (Ch) and (j0) also play a major role in the directed setting, given that convex geometries are inherently undirected structures.

Lemma 15. Let R be a directed transit function. Then (q) implies (t2a)

Proof. Let $R(u, v) \neq \emptyset$ and $u \neq v$. Suppose that $R(v, u) \neq \emptyset$. Then $\{u, v\} \subseteq R(u, v) \cap R(v, u)$. Now (q) implies that u = v, a contradiction. Thus $R(v, u) = \emptyset$ and hence R satisfies (t2a).

3.6 Rooted Directed Forests and Trees

Rooted directed forests \vec{F} and rooted directed trees \vec{T} are an interesting special case of posets. We note that the various notions of path transit functions, which we will discuss in more detail in the following section, coincide in \vec{F} and \vec{T} , because any two vertices u and v are connected by directed path if and only if u lies on the unique directed path from the root of a tree \vec{T} (which may be a component of a forest \vec{F}) to a vertex v, i.e., if and only if u is an ancestor of v. The uniqueness of paths on \vec{F} incurs additional conditions that restrict further the directed transit function. The main purpose of this section is to show that directed rooted forests and trees can be characterized by suitable axioms on directed transit functions.

Clearly the directed path transit function of \vec{F} , and thus also of a tree \vec{T} is a poset function. Since all paths between vertices in \vec{F} and \vec{T} are unique, it is easy to see that all transit functions defined by means of various types of paths in \vec{F} and \vec{T} are equal. Therefore we use the ambiguous term "path function" in this section instead of referring to a specific type, such as the all-paths function.

Lemma 16. The R be the path function of a rooted forest \vec{F} . Then R is a poset function satisfying

(p) $w \in R(u, v)$ implies $R(u, w) \cup R(w, v) = R(u, v)$.

(hy) If $R(x,v) \neq \emptyset$ and $R(y,v) \neq \emptyset$ then $x \in R(y,v)$ or $y \in R(x,v)$.

Proof. (p). If $w \in R(u, v)$, then there is a unique directed path from u to v in \vec{F} passing through w. If w = u or w = v, or u = v = w there is nothing to show, hence assume $w \neq u \neq v$. In particular then, there is a unique path from u to v, which is the concatenation of the unique paths from u to w and from w to v, and thus $R(u, v) = R(u, w) \cup R(w, v)$. Hence R satisfies (p).

(hy). Let $R(x, v) \neq \emptyset$ and $R(y, v) \neq \emptyset$, then clearly x, y, v are in the same connected component in \overrightarrow{F} . Assume $x \notin R(y, v)$ and $y \notin R(x, v)$ then we have a path in \overrightarrow{F} from x to v that does not include y, and from y to v that does not include x. Clearly, now either v does not have a unique predecessor, or there is a third vertex $w \in R(x, v) \cap R(y, v)$ that does not have a unique predecessor, contradicting the assumption that \overrightarrow{F} is a forest.

Observation 17. Since the path transit function R of a rooted tree \vec{T} in particular derives from a poset, we have also

(a5*) $R(u, x) \neq \emptyset$ and $R(x, v) \neq \emptyset$ implies $R(u, x) \cup R(x, v) = R(u, v)$. (a5) $R(u, x) \cap R(x, v) = \{x\}$ implies $R(u, x) \cup R(x, v) = R(u, v)$

Proof. To see this, note that $R(u, x) \neq \emptyset$ and $R(x, v) \neq \emptyset$ by (tr1) implies $x \in R(x, v) \cap R(x, v)$ and hence (p) and (tr2) yields (a5^{*}); moreover, (a5^{*}) implies (a5).

Axiom (a5) appears, e.g., in [8]. We shall see below that axiom (p) yields unique paths for the underlying graph of a poset function.

Lemma 18. Let R be a poset function satisfying (p). If $R(u, v) \neq \emptyset$ and $u \neq v$ then there are unique $x, y \in R(u, v)$ such that $R(u, x) = \{u, x\}$ and $R(y, v) = \{y, v\}$.

Proof. The existence of x and y has been established in Lemma 7 for all poset functions. It remains to show uniqueness. To this end, consider $x, x' \in R(u, v)$ such that $R(u, x) = \{u, x\}$ and $R(u, x') = \{u, x'\}$. Assume that $x' \neq x$. Then (**p**) implies $x' \in R(x, v) = R(x, x') \cup R(x', v)$. Since we also have $R(u, v) = R(u, x') \cup R(x', v)$ by (**p**) and $R(u, x') \cap R(x', v) = \{x'\}$ by (**q**)(by Lemma 6), we conclude that $x' \in R(u, x)$, a contradiction. Thus x = x'. An analogous argument shows that $y, y' \in R(u, v)$ with $R(y, v) = \{y, v\}$ and $R(y', v) = \{y', v\}$ implies y = y'.

Consider $x, x' \in R(u, v)$ such that $R(u, x) = \{u, x\}$ and $R(u, x') = \{u, x'\}$. Assume that $x' \neq x$. Then (**p**) implies $x' \in R(x, v) = R(x, x') \cup R(x', v)$. Moreover, $x \in R(u, v) \implies R(u, x) \cup R(x, v) = R(u, v)$ and $x' \in R(x, v) \implies R(x, x') \cup R(x'v) = R(x, v)$. By similar argument we can show that $x \in R(x', v) \implies R(x', x) \cup R(x, v) = R(x'v)$. Now either $R(x, x') = \emptyset$ or $R(x', x) = \emptyset$. W.l.o.g. let $R(x, x') = \emptyset$ and so $R(x', x) \neq \emptyset \implies x, x' \in R(x', x)$. This in turn implies $R(x', x) \cap R(x, v) = \{x, x'\}$, a contradiction to the axiom (**q**). Hence x = x'. In a similar manner it can be shown that y = y'.

An immediate consequence of Lemma 18 is the existence of unique paths in the underlying graph G_R of a poset function R that satisfies (p):

Corollary 19. Let R be a poset function satisfying (p). Then for any two $u, v \in V$ for which $R(u, v) \neq \emptyset$, R(u, v) consists of precisely the vertices that lie on the unique path from u to v in G_R .

A mangrove or strongly unambigious graphs [41], also termed multitree [42] is a DAG G with at most one directed path between any two vertices. Equivalently, the reachability relation \succeq of such a DAG G is diamond-free poset, i.e., there are no four distinct elements $a, b, c, d \in V$ such that $a \succeq b \succeq d$, $a \succeq c \succeq d$, and $b \parallel c$ [42]. As an immediate consequence of Theorem 11 and Cor. 19 we obtain:

Corollary 20. Let R be a poset function that satisfies (p), G_R its underlying graph, and R_{G_R} the path transit function of G_R . Then G_R is a mangrove and $R = R_{G_R}$.

The intuition behind condition (hy) is to forbid "hybrid vertices", i.e., the confluence of multiple paths in a vertex v.

Lemma 21. Let R be a poset function satisfying (p) and (hy). Then all vertices in G_R have in-degree at most 1.

Proof. Let R be a poset function that satisfies (p) and (hy). Assume G_R contains a vertex v with in-degree larger than one. Then we have two vertices x, y such that $R(x, v) = \{x, v\}$ and $R(y, v) = \{y, v\}$ but, by (hy), either $x \in R(y, v)$ or $y \in R(x, v)$, a contradiction.

As an immediate consequence of Corollary 19 and Lemma 21 we obtain

Corollary 22. Let R be a transit function that satisfies (p) and (hy). Then every weakly connected component of G_R contains a unique source vertex.

The unique source vertex is the root of the trees that form the weakly connected components of \vec{F} .

Theorem 23. Let R be a poset function that satisfies (p) and (hy), G_R its underlying graph, and R_{G_R} the path transit function of G_R . Then G_R is a directed rooted forest and $R_{G_R} = R$.

Proof. By Theorem 11, G_R is a DAG, by Corollary 19 we know that any path between two vertices in G_R is unique, and by Corollary 22 we know that every weakly connected component in G_R contains a unique root. Since R is a poset function, Theorem 11 implies $R_{G_R} = R$.

Let us now turn to trees. Here, we have to establish a unique root and connectedness.

Lemma 24. Let R be the path function of a rooted tree \vec{T} . Then R is a poset function satisfying (p) and

(r) $R(u,v) = R(v,u) = \emptyset$ implies that there is $w \in V$ such that $R(w,u) \neq \emptyset$ and $R(w,v) \neq \emptyset$.

Proof. Since every rooted tree is in particular a rooted forest, Lemma 16 implies that (p) holds. Suppose $R(u, v) = R(v, u) = \emptyset$ with $u, v \in V(\vec{T})$, i.e., there is neither a path from u to v or from v to u in \vec{T} . Since \vec{T} is a rooted tree there is a root vertex $w \in V(\vec{T})$ such that there is a path from w to u and v. Consequently $R(w, u) \neq \emptyset$ and $R(w, v) \neq \emptyset$ and (r) holds.

Intriguingly, a poset function R that satisfies (p) together with (r) yields a graph G_R that does not contain any hybrid vertices.

Lemma 25. Let R be a poset function that satisfies (p) and (r). Then R satisfies (hy).

Proof. Let $x, y, v \in V$ such that $R(x, v) \neq \emptyset$ and $R(y, v) \neq \emptyset$. Without loss of generality assume $R(x, y) \neq \emptyset$, then (tr) implies $y \in R(x, v)$.

Now consider the case that both $R(x, y) = \emptyset$ and $R(y, x) = \emptyset$ and hence $x \neq y$. By (r), there is $w \in V$ such that $R(w, x) \neq \emptyset$ and $R(w, y) \neq \emptyset$, and hence (tr) implies $x, y \in R(w, v)$. By (p) we have $R(w, v) = R(w, x) \cup R(x, v) = R(w, y) \cup R(y, v)$. If $x \in R(y, v)$ or $y \in R(x, v)$ there is nothing to show. Hence assume $y \in R(w, x)$ and $x \in R(w, y)$. Using (p) again we obtain $R(w, x) = R(w, y) \cup R(y, x)$. Moreover, by (q) $R(w, x) = R(w, y) \cap R(y, x) = \{y\}$ and thus $x \notin R(w, y)$, a contradiction.

Lemma 26. Let R be a poset function that satisfies (r), then G_R contains a unique source vertex.

Proof. Let *R* be a poset function on *V* that satisfies (**r**). If for every *u*, *v* we have u = v, or $R(u, v) \neq \emptyset$, or $R(v, u) \neq \emptyset$, then \succeq_R is a total order, and thus there is a unique $w \in V$ such that $R(w, u) \neq \emptyset$ for all $u \in V$, i.e., there is unique source vertex. Now suppose, for contradiction, that no such vertex exists. Then \succeq_R cannot be a total order, and thus there is a pair of vertices $u, v \in V$ such that $R(u, v) = R(v, u) = \emptyset$. Let $U_1 = \{u, v\}$. By (**r**), there is $u_1 \in V$ with $R(u_1, u) \neq \emptyset$ and $R(u_1, v) \neq \emptyset$, and thus $u_1 \notin U_1$. By assumption, there is $v_1 \in V$ such that $R(u_1, v_1) = \emptyset$ and $R(v_1, u_1) = \emptyset$, which in particular implies $v_1 \notin U_1$. By (**r**), there is $u_2 \in V$ such that $R(u_2, u_1) \neq \emptyset$ and $R(u_2, v_1) \neq \emptyset$, thus by (**t**0) also $R(u_2, x) \neq \emptyset$ for all $x \in U_1 \cup \{u_1, v_1\} =: U_2$. Repeating this argument yields a sequence of vertices $u_i, v_i \in V \setminus U_{i-1}$ such that $R(u_i, v_i) = R(v_i, u_i) = \emptyset$ and $R(u_i, x) \neq \emptyset$ for all $x \in U_{i-1}$ and $U_i = U_{i-1} \cup \{u_{i-1}, v_{i-1}\}$. In particular the set U_i grows in each step, which is impossible since *V* is finite. Hence, there must be a source vertex.

The existence of a unique source vertex in a graph in particular implies that this graph is (at least) weakly connected.

Lemma 27. Let R be a poset function that satisfies (r) and (p). Then G_R is weakly connected.

Proof. Let $u, v \in V$ such that $R(u, v) = R(v, u) = \emptyset$ then we know by (r) that there is a $w \in V$ such that $R(w, u) \neq \emptyset$ and $R(w, v) \neq \emptyset$. By Corollary 19, there is a path from w to u in G_R and from w to v in G_R . Hence, w, u, and v are contained within the same (weakly) connected component in G_R .

If $R(u, v) \neq \emptyset$ and $R(v, u) = \emptyset$, we use Corollary 19 again to show that there is a path from u to v in G_R (analogously for $R(v, u) \neq \emptyset$ and $R(u, v) = \emptyset$). Since trivially (t2a) prohibits symmetric edges, the case that $R(u, v) \neq \emptyset$ and $R(v, u) \neq \emptyset$ cannot occur. Summarizing, for any two vertices in G_R , there is either a path between them, or there is a third vertex w such that there is a path from w to u and from w to v. Consequently, G_R is weakly connected.

So far, we have established that the underlying graph G_R of a poset function R that satisfies (p) and (r) is weakly connected, contains a unique source vertex, and there is at most one directed path between any two vertices of G_R . This puts us into position to state the main result of this subsection.

Theorem 28. Let R be a poset function that satisfies (r) and (p), G_R its underlying graph, and R_{G_R} the all-path transit function of G_R . Then G_R is a directed rooted tree and $R = R_{G_R}$.

Proof. By Theorem 11, G_R is a DAG and by Lemma 27 G_R is weakly connected. We know by Corollary 19 that every path from one vertex to another in G_R is unique. The existence of a root in G_R (by Lemma 26), from which there exists a unique path to every other vertex in V together with the fact that G_R is a DAG implies that the root is unique. Consequently, G_R is a rooted tree. Since R is a poset function, Theorem 11 implies $R_{G_R} = R$.



Fig. 1: The reachability function of this digraph satisfies $R_{\rightarrow}(x_1, x_2) = \{u, v, w\}$ for any two distinct vertices $x_1, x_2 \in \{u, v, w\}$. Thus $G_{R_{\rightarrow}}$ is edge-less and hence $G \not\cong G_{R_{\rightarrow}}$.

4 Directed transit functions of general graphs

4.1 Reachability

Reachability in its most general form can be understood as a reflexive, transitive relation \rightarrow on a set V. Naturally, we set

$$R_{\rightarrow}(u,v) = \begin{cases} \{x \in V | u \to x \text{ and } x \to v\} & \text{if } u \neq v \\ \{u\} & \text{if } u = v \end{cases}$$
(3)

Note that $R_{\rightarrow}(u, v) = \emptyset$ if and only if $u \neq v$, i.e., if v is not reachable from u. One easily checks that R_{\rightarrow} satisfies (t1), (t3), (tr1), and (tr2), using the same arguments as in Lemma 1.

The statements of (tr1) and (tr2) do not extend to the case u = v in the absence of the anti-symmetry axiom (t2a), which is not assumed here. Therefore, (q) no longer holds in this more general setting.

Let G = (V, E) be a connected graph. Then for every u and every $v \neq u$, we have $u \to x$ and $x \to v$, and thus $x \in R_{\to}(u, v)$. That is, $R_{\to}(u, v) = V$ for $u \neq v$ and $R_{\to}(u, u) = \{u\}$. The example in Fig. 1 show that in general the underlying graph of the reachability relation does not recover the original graph.

Reachability is closely related to notions of connectedness. We see that V is strongly connected if $R(u, v) \neq \emptyset$ for all $u, v \in V$ and unilaterally connected if $R(u, v) \neq \emptyset$ or $R(v, u) \neq \emptyset$ holds for all $u, v \in V$.

Lemma 29. Let R be a directed transit function on V. Then the binary relation ~ on V defined by $u \sim v$ if $R(u, v) \neq \emptyset$ and $R(v, u) \neq \emptyset$ is an equivalence relation.

Proof. Symmetry is inherent in the definition and by (t3), \sim is reflexive. Suppose $u \sim w$ and $w \sim v$. Then $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$, which by (t0) implies $R(u, v) \neq \emptyset$; analogously, $R(v, w) \neq \emptyset$ and $R(w, u) \neq \emptyset$ by (t0) implies $R(v, u) \neq \emptyset$, and thus $v \sim u$, i.e., \sim is an equivalence relation.

The equivalence classes of \sim can be viewed as strongly connected components w.r.t. the directed transit function R. The set V is partitioned into a set C of strongly connected components if and and only if $R(u, v) \neq \emptyset$ implies $R(v, u) \neq \emptyset$ for all $u, v \in C$ with $C \in C$. Moreover, V is strongly connected w.r.t. R if and only if $R(u, v) \neq \emptyset$ for all $u, v \in V$. **Lemma 30.** Let R be a directed transit function. Then $R(u, v) = \emptyset$ implies that there is no path from u to v in the underlying graph G_R .

Proof. Let $C_+(u) := \{w \in V | R(u, w) \neq \emptyset\}$ and $C_-(v) := \{w \in V | R(w, v) \neq \emptyset\}$. By (t0), $R(u, v) = \emptyset$ implies $C_+(u) \cap C_-(v) = \emptyset$. Suppose there is an edge $(p, q) \in E(G_R)$ with $p \in C_+(u)$ and $q \in C_-(u)$. Then $R(p,q) = \{p,q\} \neq \emptyset$ and thus $R(u,p) \neq \emptyset$ and $R(p,q) \neq \emptyset$ implies $R(u,q) \neq \emptyset$, i.e., $q \in C_+(u)$, a contradiction. Thus G_R cannot contain an edge connecting u with any vertex $q \notin C_+(u)$, and thus there cannot be a path in G for u to any $q \notin C_+(u)$.

Lemma 30 extends Lemma 9 to all directed transit functions. The converse is not true in general for directed transit functions since R need not have any transit sets of size 2.

4.2 Geometric directed transit functions

It is interesting to note that key implications between properties of poset functions remain true when the condition of the anti-symmetry axiom (t2a) is lifted. In particular, it is well-known for the case of symmetric transit function that (b3) implies (b4) implies (b1). This statement generalizes to the directed transit functions:

Lemma 31. Let R be a directed transit function satisfying (tr2). If R satisfies $(b3_1)$ and $(b3_2)$, then it also satisfies (b4), $(b1_1)$, and $(b1_2)$.

Proof. Suppose $x \in R(u, v)$. By (tr2), $R(u, x) \neq \emptyset$ and $R(x, v) \neq \emptyset$ and hence $x \in R(u, x) \cap R(x, v)$. Let $y \in R(u, x)$ and $y \in R(x, v)$. Then by (b3₁) $x \in R(y, v)$ and by (b3₂) $x \in R(u, y)$. Applying (b3₁) to $x \in R(u, y)$ and $y \in R(u, x)$ with v = y implies $x \in R(y, y) = \{y\}$ and thus x = y, and thus (b4) holds. Now consider $x \in R(u, v) \neq \{u, v\}$. Then $R(u, x) \cap R(x, v) = \{x\}$ by (b4) and thus $u \in R(u, x)$ implies $u \notin R(x, v)$, and $v \in R(v, x)$ implies $v \notin R(u, x)$, i.e., (b1₁) and (b1₂) hold.

For symmetric transit functions without empty transit sets, the notion of a geometric (transit) functions was introduced in [32] as those that satisfy axioms (b2) and (b3). Since then, the term has been in common use; see also [43, 38]. In the following we combine $(b2_1)$ and $(b2_2)$ defined in Lemma 14 into a single axiom

(b2) $w \in R(u, v)$ implies $R(u, w) \cup R(w, v) \subseteq R(u, v)$

Recalling (tr0) from Observation 2, it is easy to check

Observation 32. A directed transit function R satisfies (tr1) if and only it satisfies (tr0) and (b2).

Axiom (tr0) is clearly true for the reachability transit function, and for the directed all-paths transit function in the setting of partial orders. In general it fails for pathbased transit functions since the concatenation of a u, w-path and a w, v-path is, in general, only guaranteed to contain a u, v-path. However, this u, v-path need not pass through the concatenation point w. In [18], therefore the weaker condition (t0) appears to ensure consistency of connected components in undirected graphs. Axiom (t0) is an immediate consequence of (tr1). Notably, even in conjunction with (tr2) and (b3_{1,2}), (t0) does not imply (tr1).

A natural generalization of Nebeský's notion of a geometric transit function for (not necessarily connected) structures is the following:

Definition 6. A directed transit function R is geometric if it satisfies (tr2), (b2) and $(b3_{1,2})$. Moreover, we say that directed transit function R is weakly geometric if it satisfies (tr2), (b2) and $(b1_{1,2})$.

It follows from Lemma 31 that geometric directed transit function is always weakly geometric. The converse is not true in general, as shown by examples 1 and 2. **Example 1.** $(b1_{1,2})$, (b2), but not $(b3_1)$.

 $V = \{a, b, c, d\}$ and $R(a, d) = \{a, b, c, d\}$, $R(a, c) = \{a, b, c\}$, $R(a, b) = \{a, b\}$, $R(b, c) = \{b, c\}$, $R(c, d) = \{c, d\}$, $R(b, d) = \{b, d\}$, $R(x, x) = \{x\}$ for all $x \in V$, and $R(u, v) = \emptyset$ otherwise.

Example 2. $(b1_{1,2})$, (b2), but not $(b3_2)$.

 $V = \{a, b, c, d\} \text{ and } R(a, d) = \{a, b, c, d\}, R(b, d) = \{b, c, d\}, R(a, b) = \{a, b\}, R(b, c) = \{b, c\}, R(c, d) = \{c, d\}, R(a, c) = \{a, c\}, R(x, x) = \{x\} \text{ for all } x \in V, \text{ and } R(u, v) = \emptyset \text{ otherwise.}$

The following examples of functions $R: V \times V \to 2^V$ satisfying (t1) and (t3) show that the axioms (t0), (b1₁), (b1₂), (b2), and (tr2) are independent of each other, i.e., weakly geometric transit functions are not defined by any subset of this collection of axioms.

Example 3. (b1₁), (b1₂), (b2), and (t0) but not (tr2).

 $V = \{a, b, c\}$ and $R(a, b) = \{a, c, b\}$, $R(a, c) = \{a, c\}$, $R(b, c) = \{b, c\}$, $R(x, x) = \{x\}$ for all $x \in V$, and $R(u, v) = \emptyset$, otherwise.

Example 4. (b1₁), (b1₂), (b2), and (tr2) but not (t0).

 $V = \{a, b, c\}$ and $R(a, b) = \{a, c, b\}$, $R(b, a) = \emptyset$, $R(x, x) = \{x\}$ for all $x \in V$, and $R(u, v) = \{u, v\}$ otherwise.

Example 5. $(b1_2)$, (b2), (t0), and (tr2) but not $(b1_1)$.

 $V = \{a, b, c, d\} \text{ and } R(a, b) = \{a, c, d, b\}, R(a, c) = \{a, b, d, c\}, R(a, d) = \{a, d\}, R(b, c) = \{b, c\}, R(c, b) = \{c, b\}, R(d, b) = \{d, b\}, R(d, c) = \{d, c\}, R(x, x) = \{x\} \text{ for all } x \in V, \text{ and } R(u, v) = \emptyset \text{ otherwise.}$

Example 6. $(b1_1)$, (b2), (t0), and (tr2) but not $(b1_2)$.

Let $V = \{a, b, c, d\}$ and $R(a, b) = \{a, b, c\} = R(c, b)$, $R(a, c) = \{a, c\}$, $R(a, d) = \{a, d\}$, $R(b, c) = \{b, c\}$, $R(b, d) = \{b, d\}$, $R(c, d) = \{c, d\}$, $R(x, x) = \{x\}$ and $R(u, v) = \emptyset$ otherwise.

Example 7. $(b1_1)$, $(b1_2)$, (t0), and (tr2) but not (b2).

 $V = \{a, b, c, d\}$ and $R(a, b) = \{a, c, b\}$, $R(a, c) = \{a, d, c\}$, $R(a, d) = \{a, d\}$, $R(c, b) = \{c, b\}$, $R(d, b) = \{d, b\}$, $R(d, c) = \{d, c\}$, $R(x, x) = \{x\}$ for all $x \in V$ and $R(u, v) = \emptyset$ otherwise.

Similarly, the axioms in the definition of geometric directed transit functions are independent:

Example 8. (t0), $(b3_{1,2})$, (b2) but not (tr2).

 $V = \{a, b, c, d\}$ and $R(a, d) = \{a, b, c, d\}$, $R(a, c) = \{a, b, c\}$, $R(a, b) = \{a, b\}$, $R(b, c) = \{b, c\}$, $R(c, d) = \{c, d\}$, $R(b, d) = \{b, c, d\}$, $R(d, c) = \{d, c\}$, $R(d, b) = \{d, c, b\}$, $R(x, x) = \{x\}$ for all $x \in V$, and $R(u, v) = \emptyset$ otherwise. **Example 9.** (tr2), (b3_{1.2}), (b2) but not (t0).

 $V = \{a, b, c, d\}$ and $R(a, d) = \{a, b, c, d\}, R(a, c) = \{a, b, c\}, R(a, b) = \{a, b\},$

 $\begin{aligned} R(b,c) &= \{b,c\}, \ R(c,d) = \{c,d\}, \ R(b,d) = \{b,c,d\}, \ R(d,c) = \{d,c\}, \ R(c,b) = \{c,b\}, \\ R(x,x) &= \{x\} \ for \ all \ x \in V, \ and \ R(u,v) = \emptyset \ otherwise. \end{aligned}$ Example 10. (tr2), (b3_{1,2}), (t0), (b2₂) but not (b2₁). $V &= \{a,b,c,d\} \ and \ R(a,d) = \{a,b,c,d\}, \ R(a,c) = \{a,b,c\}, \ R(a,b) = \{a,b\}, \ R(b,c) = \{b,c\}, \ R(c,d) = \{c,d\}, \ R(b,d) = \{b,c,d\}, \ R(d,b) = \{d,c,b\}, \ R(d,c) = \{a,d,c\}, \\ R(x,x) &= \{x\} \ for \ all \ x \in V, \ and \ R(u,v) = \emptyset \ otherwise. \end{aligned}$ Example 11. (tr2), (b3_{1,2}), (t0), (b2₁) but not (b2₂). $V &= \{a,b,c,d\} \ and \ R(a,d) = \{a,b,c,d\}, \ R(a,c) = \{a,b,c\}, \ R(a,b) = \{a,b\}, \ R(b,c) = \{b,c\}, \ R(c,d) = \{c,d\}, \ R(b,d) = \{b,c,d\}, \ R(d,b) = \{a,c,b\}, \ R(b,c) = \{b,c\}, \ R(c,d) = \{c,d\}, \ R(b,d) = \{b,c,d\}, \ R(d,b) = \{d,c,b\}, \ R(c,b) = \{a,c,b\}, \\ R(x,x) &= \{x\} \ for \ all \ x \in V, \ and \ R(u,v) = \emptyset \ otherwise. \end{aligned}$ Example 12. (t0), (b3₂), (b2), (tr2) \ but \ not (b3₁) $X &= \{a,b,c,d\} \ and \ R(a,b) = V, \ R(a,c) = \{a,c,d\}, \ R(a,d) = \{a,c,d\}, \ R(b,c) = \{a,c,d\},$

Example 13. (t0), (b3₁), (b2), (tr2) but not (b3₂).

 $\begin{array}{l} X = \{a,b,c,d\} \ and \ R(a,b) = V, \ R(c,b) = \{b,c,d\}, \ R(a,d) = \{a,d\}, \ R(b,c) = \{b,c\}, \\ R(c,d) = \{c,d\}, \ R(d,c) = \{d,c\}, \ R(c,b) = \{c,b\}, \ R(x,x) = \{x\} \ for \ all \ x \in V, \ and \\ R(u,v) = \emptyset \ otherwise \end{array}$

The reachability and poset functions discussed above are special cases of geometric transit functions:

Observation 33. Let R be a geometric transit function. Then:

(i) R is a reachability transit function if and only if it satisfies (tr0).

(ii) R is a poset function if and only if it satisfies (tr0) and (t2a).

Theorem 34. Let R be a weakly geometric directed transit function. Then, $R(u, v) \neq \emptyset$ if and only if there exists a directed path from u to v in G_R that is contained in R(u, v). If R in addition satisfies (b4), then for every $w \in R(u, v)$ such a path can be found that runs through w.

Proof. If $R(u, v) = \emptyset$, then Lemma 30 rules out the existence of path from u to vin G_R . In the following we assume $R(u, v) \neq \emptyset$. If u = v then $R(u, u) = \{u\}$ and there is nothing to show. Thus we assume $u \neq v$ and proceed by induction over |R(u, v)| that there is a directed path from u to v in G_R . The base case |R(u, v)| = 2is trivial since we have $R(u, v) = \{u, v\}$ and thus $(u, v) \in E(G_R)$. For |R(u, v)| = 3let $R(u, v) = \{u, v, x\}$. We have $R(u, x) \subseteq R(u, v)$ and $R(x, v) \subseteq R(u, v)$ by (b2) and $R(u, x) \subsetneq R(u, v)$ and $R(x, v) \subsetneq R(u, v)$ by (b1₁) and (b1₂), respectively, while (tr2) implies that both $R(u, x) \neq \emptyset$ and $R(x, v) \neq \emptyset$. Therefore $R(u, x) = \{u, x\}$ and $R(x, v) = \{x, v\}$, and thus (u, x, v) is path in G_R .

Now, assume that assertion holds for all u' to v' such that $2 \leq |R(u',v')| < k$ and assume |R(u,v)| = k. Thus there is $w \in R(u,v) \setminus \{u,v\}$ and hence $R(u,w) \neq \emptyset$ and $R(w,v) \neq \emptyset$. From (b2) and (b1₁) we obtain $R(u,w) \subsetneq R(u,v)$ since $v \notin R(u,w)$, and thus |R(u,w)| < k. Analogously, (b2) and (b1₂) yields $R(w,v) \subsetneq R(u,v)$ and |R(w,v)| < k. By induction hypothesis, therefore, there is a directed path from u to w contained in R(u,w) and a directed path from w to v contained in R(w,v). The concatenation of these two directed paths contains a path from u to v that, by (b2), is contained in R(u,v).

If R satisfies (b4), then $R(u, w) \cap R(w, v) = \{w\}$ and thus the concatenation of any directed path from u to w in R(u, w) with any directed path from w to v in R(w, v) is again a directed path. Thus there is directed path from u to v in R(u, v) that runs through w.

Theorem. 34 extends Cor.8 from poset functions to geometric transit functions. The assumption of axiom (b4) in the second part of Theorem 34 is essential:

Example 14. Let $V = \{u, x, w, v\}$ and $R(u, v) = \{u, x, w, v\}$, $R(u, w) = \{u, x, w\}$, $R(w, v) = \{w, x, v\}$, $R(u, x) = \{u, x\}$, $R(x, w) = \{x, w\}$, $R(w, x) = \{w, x\}$, $R(x, v) = \{x, v\}$, $R(y, y) = \{y\}$ for all $y \in V$, and $R(y, z) = \emptyset$ otherwise. The directed transit function R satisfies (b1₁), (b1₂), (b2). However (b4), and hence (b3₁) and (b3₂) do not hold. We have $w \in R(u, v)$, there is a path from u to v in R(u, v), namely (u, x, v), but there is no path from u to v passing through w.

4.3 Directed all-path transit function

Here we continue with more properties of the all-paths functions A_G on an arbitrary graph G.

Lemma 35. Let G be a directed graph. Then $R = A_G$ is a directed transit function satisfying (tr2) as well as

(b5) $R(u,w) \cap R(w,v) = \{w\}$ implies $R(u,w) \cup R(w,v) \subseteq R(u,v)$. (Mod) $R(u,w) \neq \emptyset$ and $R(w,v) \neq \emptyset$ implies $R(u,v) \cap R(u,w) \cap R(w,v) \neq \emptyset$.

Proof. Clearly A_G satisfies (t1) and (t3). If $A_G(u, w) \neq \emptyset$ and $A_G(w, v) \neq \emptyset$ there is a u, w-path P_1 and a w, v-path P_2 . Since $w \in V(P_1) \cap V(P_2) \neq \emptyset$, the first vertex xalong P_1 also contained in P_2 is well-defined. Then the concatenation Q of the u, xsubpath of P_1 and the x, v-subpath of P_2 is a u, v-path and thus $A_G(u, v) \neq \emptyset$, i.e., (t0) is satisfied. In particular, therefore, A_G is a directed transit function.

Moreover, we have $x \in V(P_1) \subseteq A_G(u, w)$, $x \in V(P_2) \subseteq A_G(u, w)$, and $x \in V(Q) \subseteq A_G(u, v)$. Thus $x \in A_G(u, w) \cap A_G(w, v) \cap A_G(u, v) \neq \emptyset$, and thus (Mod) holds.

If $x \in A_G(u, v)$ then in particular there exist a path from u to v through x, which contains a sub-path from u to x and a sub-path from x to v. Thus $A_G(u, x) \neq \emptyset$ and $A_G(x, v) \neq \emptyset$, i.e., (tr2) is satisfied.

Let $R(u, w) \cap R(w, v) = \{w\}$ and assume $R(u, w) \cup R(w, v) \not\subseteq R(u, v)$. Then there is an $x \in R(u, w) \cup R(w, v)$ such that $x \notin R(u, v)$. Assume $x \in R(u, w)$ then there is a path from u over x to w. Since $x \notin A_G(u, v)$ but there is at least one path from w to v, the uxw-path and the wv-path must overlap. But we have $A_G(u, w) \cap A_G(w, v) = \{w\}$, a contradiction. An analogous argument can be made for the case that $x \in A_G(w, v)$. Thus A_G satisfies (b5).

If $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$ then there are paths P_1 from u to w and P_2 from w to v. Let x be the first vertex along P_1 that is also contained in P_2 . Denote by P'_1 the subpath of P_1 from u to x and by P'_2 the subpath of P_2 from x to v. Then $P'_1 \cup P'_2$ is a path from u to v.

Lemma 35 generalizes the results in [44], where it is shown that the all-paths function A_G of any connected undirected graph satisfies (b5) and (Mod). Connectedness in this setting implies that the precondition in (Mod) is always satisfied.



Fig. 2: Example of a directed graph that does not satisfy many of the axioms that are essential to prove the existence of paths for DAGs, or their posets. For example we do not have $R(u, w) \neq \emptyset$ and $R(w, v) \neq \emptyset$ implies that $w \in R(u, v)$, hence (tr0) is not satisfied. Similarly, we have that $R(u, w) \cup R(w, v) \not\subseteq R(u, v)$ and (tr1) does not hold. Since $R(u, w) \cap R(w, v) = \{w, x\}$ also (q) is not satisfied.



Fig. 3: The all-path transit function A_G of the graphs violates one of the betweenness axioms. (a) (b1₁) is violated since $x \in R(u, v) = \{u, x, v\}$ but $v \in R(u, x) = \{u, v, x\}$. (b) (b1₂) is violated by $x \in R(u, v) = \{u, x, v\}$ and $u \in R(x, v) = \{u, x, v\}$. (c) (b2₁) does not hold since $x \in R(u, v) = \{u, x, v\}$ and $w \in R(u, x) = \{u, v, w, x\}$ but $w \notin R(u, v)$ and hence $R(u, x) \not\subseteq R(u, v)$. (d) Analogously, (b2₂) is violated by $x \in R(u, v)$ but $w \notin R(u, v)$ and $w \in R(x, v)$ but $w \notin R(u, v)$.

While the all-path transit function has many appealing properties in the symmetric case, this is no longer the case for directed graphs. Fig. 3 shows that A_G in general satisfies neither (b1_{1,2}) nor (b2), and hence A_G is not weakly geometric but only satisfies the much weaker axiom (b5). It is worth noting that directed transit functions satisfying (b4) and (b5) also satisfy (b2). Moreover, (b4) immediately implies (the contrapositive of) (tr2). Thus a directed transit function satisfying (b4) and (b5) is weakly geometric.

Lemma 36. A directed transit function R satisfying axiom (b2) and (Mod) also satisfies (b5).



Fig. 4: A graph G for which the transit function J_G does not satisfy (b5): we have $J_G(u, w) = \{u, ..., w\}, J_G(u, w) = \{w, ..., v\},$ but $J_G(u, v) = \{u, v\}.$

Proof. Suppose $R(u, x) \cap R(x, v) = \{x\}$. By (Mod), we have $R(u, v) \cap R(u, x) \cap R(x, v) = R(u, v) \cap \{x\} \neq \emptyset$ and thus $x \in R(u, v)$. Now (b2) immediately yield $R(u, x) \cup R(x, v) \subseteq R(u, v)$, and thus R satisfies (b5).

Note that a geometric directed transit function does not necessarily satisfy (b5). As an example, consider $V = \{u, v, w\}$, $R(u, v) = \{u, v\}$, $R(u, w) = \{u, w\}$, $R(w, v) = \{w, v\}$, $R(v, u) = R(w, u) = R(v, w) = \emptyset$, and $R(x, x) = \{x\}$ for all $x \in V$. It is not difficult to check that R is a geometric directed transit function. However, $R(u, w) \cap R(w, v) = \{w\}$ but $R(u, v) = \{u, v\}$, violating (b5).

4.4 Induced path transit function

A path $P = (u = x_0, x_1, ..., x_k = v)$ in a (directed) graph G = (V, E) is *induced* if there is no edge $(i, j) \in E$ with $j \neq i-1$ for $1 \leq j \leq n$. That is, the subgraph G[P] of Ginduced by the vertex set P comprises only the directed edges (x_{i-1}, x_i) for $1 \leq i \leq k$.

A path P is shortcut-free if there is no edge $(i, j) \in E$ with $j \neq i - 1$ for $1 \leq j \leq n$ and i < j. We caution the reader that what we call here shortcut-free paths is sometimes called "induced directed paths" [45]. For symmetric digraphs (and thus undirected graphs) shortcut-free and induced paths coincide. In the directed case, every induced path is shortcut-free, but the converse is not true.

The transit functions deriving from induced paths in undirected graphs have been considered e.g. in [46, 6, 7]. Nebeský's proof that it is impossible to characterize the induced path function of connected graphs using a set of first order axioms [47] suggests that it will be difficult to obtain general results on this class of transit functions. Research in the undirected cases therefore focuses on the characterization of induced path transit functions for specific graph classes. In the directed case, it seems quite natural to consider the shortcut-free paths.

Definition 7. For a digraph G, let $x \in J_G(u, v)$ if x lies along a shortcut-free path from u to v.

Lemma 37. The function J_G is a directed transit function for every digraph G.

Proof. It follows immediately from the definition that J_G satisfies (t1) and (t3). Moreover, every paths P in G contains a shortcut-free subpath, obtained by removing, iteratively, from P all vertices that are "bridged" by a shortcut. Thus $J_G(u, v) = \emptyset$ if and only if $A_G(u, v) = \emptyset$ for all $u, v \in V$. It follows immediately that J_G satisfies (t0) since A_G satisfies (t0).

The example in Figure 4 shows that J_G in general does not satisfy (b5). The following two properties are satisfied by The undirected induced path transit function

satisfies the following two properties, see e.g. [7]. Below, we show that this remains true in the directed case:

Lemma 38. The directed transit function J_G of any digraph G satisfies

- (j2) $R(u,x) = \{u,x\}, R(x,v) = \{x,v\}, R(u,v) \neq \{u,v\}$ implies $x \in R(u,v)$.
- (j2') $x \in R(u, y)$ and $y \in R(x, v)$, $R(u, x) = \{u, x\}$, $R(x, y) = \{x, y\}$, $R(y, v) = \{y, v\}$, and $R(u, v) \neq \{u, v\}$ implies $x \in R(u, v)$.

Proof. Let G be a directed graph. If u = v then $J_G(u, v) = \{u, v\}$, and thus the precondition of both (j2) and (j2') is not met. Hence we may assume $u \neq v$. Setting x = y in (j2') and removing the preconditions that become trivially true, i.e., $x \in R(u, x)$, $x \in R(x, v)$, and $R(x, x) = \{x, x\}$, shows that (j2') reduces to (j2) for x = y. For x = u or x = v the precondition in (j2) is never met, and hence the axiom holds trivially. If u, v, x are pairwise distinct, then $(u, x) \in E(G)$ and $(x, v) \in E(G)$, $u \neq v$ and $(u, v) \notin E(G)$. Hence (u, x, v) is a shortcut-free path of length 2 in G. By definition, we therefore have $\{u, x, v\} \subseteq J_G(u, v)$ and thus in particular $x \in J_G(u, v)$, i.e., J_G satisfies (j2).

It thus remains to consider (j2') for $x \neq y$. Moreover, the precondition is void if u = v. For x = u and y = v or x = v and y = u, the implication is trivially true. Thus it remains to consider the case that u, v, x, y are pairwise distinct. The precondition of (j2') implies that $(u, x) \in E(G)$, $(x, y) \in E(G)$, $(y, v) \in E(G)$ and thus (u, x, y, v) is a directed path of length 3. Moreover, the condition $x \in R(u, y)$ requires that there is a shortcut-free path from u to y that runs through x, and thus in particular $(u, y) \notin E(G)$. Analogously, $y \in R(x, v)$ implies $(x, v) \notin E(G)$. Finally, $R(u, v) \neq \{u, v\}$ implies $(u, v) \notin E(G)$, and thus (u, x, y, v) is a shortcut-free directed path from u to v and thus $\{u, x, y, v\} \subseteq J_G(u, v)$.

The induced path function $J_G(u, v)$ is not (weakly) geometric as shown by the following examples Fig. 5.



Fig. 5: Examples of graphs for which J_G is not geometric/weakly geometric.

4.5 Directed interval functions

A finite quasimetric space (X, d) consists of a finite set V and a distance $d: V \times V \rightarrow \mathbb{R}_+ \cup \{\infty\}$ satisfies, for all $x, y, z \in V$, (i) d(x, y) = 0 iff x = y and (ii) the triangle inequality $d(x, z) + d(z, y) \ge d(x, y)$.

Definition 8. Let (X, d) be a quasimetric space. Then the function $I_d : V \times V \to 2^V$ given by

$$I_d(u,v) \coloneqq \begin{cases} \{w \mid d(u,w) + d(w,v) = d(u,v)\} & \text{if } d(u,v) < \infty \\ \emptyset & \text{if } d(u,v) = \infty \end{cases}$$
(4)

is called the interval function of (X, d).

This definition is the obvious analog of the undirected interval function defined in [1, 8]. For strongly directed graphs, the related set system $\{I_d(u, v)|u, v \in V(G)\}$ was used in [48] to define a convexity for digraph G.

The next result shows that the directed version shares the basic of properties of the undirected interval function:

Lemma 39. The interval function I_d of a finite quasimetric space (X, d) is a geometric directed transit function.

Proof. (t1) follows from d(u, v) = d(u, u) + d(u, v) = d(u, v) + d(v, v) since d(u, u) = d(v, v) = 0 by (i), and (t3) follow from d(u, u) = 0 = d(u, w) + d(w, u) only if w = u. If $I_d(u, w) \neq \emptyset$ and $I_d(w, v) \neq \emptyset$ then $d(u, w) < \infty$ and $d(w, v) < \infty$. The triangle inequality thus implies $d(u, v) \leq d(u, w) + d(w, v) < \infty$ and thus $I_d(u, v) \neq \emptyset$, i.e., (t0) is satisfied. Thus I_d is a directed transit function.

Now suppose $d(u, x) = \infty$ or $d(x, v) = \infty$. In the first case, we have $d(u, v) = \infty$ and thus $I_d(u, y) = \emptyset$, which in turn implies $x \notin I_d(u, v)$. Otherwise, $d(u, v) < \infty$, in which case $d(u, x) + d(x, v) \neq d(u, v)$ and thus $x \notin I_d(u, v)$. Hence I_d satisfies (tr2).

Next consider the betweenness axioms. $x \in I_d(u, v)$ and $y \in I_d(u, x)$ implies d(u, v) = d(u, x) + d(x, v) and d(u, x) = d(u, y) + d(y, x), and thus d(u, v) = d(u, y) + d(y, x) + d(x, v). Now suppose $d(y, x) + d(x, v) \neq d(y, v)$. Then the triangle inquality implies d(y, x) < d(y, x) + d(x, v) and thus d(u, v) > d(u, y) + d(y, v), violating the triangle inequality. Thus d(y, x) + d(x, v) = d(y, v), i.e., $x \in I_d(y, v)$, i.e., $(b3)_1$) holds. Moreover, we have d(u, y) + d(y, v) = d(u, v), i.e., $y \in I_d(u, v)$ for all $y \in I_d(u, x)$, and thus $I_d(u, x) \subseteq I_d(u, v)$. An analogous argument yield $(b2_2)$ and $(b2_2)$, and hence (b2) is satisfied.

It is not difficult to check that Lemma 18 remains true for geometric directed transit function. Recall that we have shown in Theorem. 34 above that for a geometric directed transit function, the underlying graph G_R contains paths contained entirely in R(u, v)and running through any chosen $w \in R(u, v)$ provided $R(u, v) \neq \emptyset$. In particular, therefore, there is $x, y \in R(u, v)$ such that $R(u, x) = \{u, x\}$ and $R(y, v) = \{y, v\}$. To show that x and y are unique if (p) is assumed, it suffices to to appeal (b4) instead of (q) in the proof Lemma 18. It follows immediately that the entire path from u to v is uniquely determined:

Observation 40. If R is a geometric directed transit function that in addition satisfies (p), then $R(u, v) \neq \emptyset$ implies that there is a unique path from u to v in G_R .

It is natural to ask under which conditions a geometric directed transit function R derives from a graph G such that $R = I_d$ and $G_R = G$, where I_d is the interval function of the quasi-metric defined by the shortest paths in G. For the symmetric case, i.e., for undirected graphs and their metrics, several characterizations for such

"graphic" transit functions have been given [3, 49]. In the directed case, however, this problem is more difficult and will be tackled in the forthcoming work.

5 Concluding Remarks

We have introduced here a notion of transit functions for directed and not necessarily connected structures. It is a proper generalization of the well-studied undirected transit functions for connected structures that also encompasses the extension this framework to disconnected graphs in [18]. Directed transit functions also capture the idea of directed betweenness that is common in network science [20]. As a consequence, we obtain a simple characterization of directed transit functions that describe posets. This is in contrast to the difficulties of describing posets by means of connected, undirected transit functions [36, 37, 38].

We observed that the most widely used concepts and in particular the wellestablished betweenness axioms naturally generalize to the directed case. For example, there is natural notion of weakly geometric transit functions that ensures that $R(u, v) \neq \emptyset$ implies that there is a directed path from u to v in the underlying digraph G_R , generalizing a basic fact from the theory of undirected transit functions. Similarly, it is possible to define many of the walk- and path-based constructions of transit function on graphs to the directed case. We briefly consider the directed analogs of interval (geodetic) functions, as well as all-paths, and induced path transit functions. For directed acyclic graphs, we have obtained characterizations closely matching results for undirected transit functions on cycle-free graphs and trees. On the other hand, there are also some quite drastic differences. While interval functions of digraphs turn out to be a special case of geometric directed transit functions in complete analogy with the undirected case, it does not seem possible to extend the graphic transit functions [3, 49] to the directed case. For further work on this topic we refer to forthcoming work.

Nebeský [5] showed that no first order characterization of the induced-path function exists for undirected graphs. Since the symmetric case is obtained from the definition of J_G by simply adding the symmetry axiom (t2s), such a characterization also may not exists for the the general case. The all-paths transit function of undirected graphs has a well-known characterization in terms of first order axioms [8]. These conditions do not appear to generalize to directed graphs, however, since it does not satisfy (b2) for directed graphs. It remains an open problem, therefore, whether a simple characterization for the all-paths directed transit functions can be obtained.

Even if a simple characterization for transit functions deriving from graphs is not possible, it remains a rich and interesting topic to identify transit functions of particular graphs classes. In the undirected case, for instance, there is a rich literature on distance hereditary graphs, see e.g. [50, 51]. The concept is extended to directed graph in [45]: A directed graph G = (V, E) is distance hereditary if for all induced sub-digraphs G' and all $u, v \in V(G')$ we have $d_{G'}(u, v) = \infty$ or $d_{G'}(u, v) = d_G(u, v)$. As shown in Lemma 1 of [45], G is a distance hereditary if and only if every shortcutfree directed path from u to v is a shortest directed path. In [52] it it show that the interval function of undirected distance hereditary graph has a simple characterization

in terms of first order axioms. Similar results have been obtained for various graph classes for interval function, for example see [53] and for induced path see[7, 39, 54]. It will be interesting to explore this type of results also of directed transit functions and digraphs. Related questions arising from the present concern e.g. the characterization of directed graphs G in which the all-paths function A_G or the induced path function J_G is (geometric/weakly geometric). In recent years also the betweenness relations defined other undirected path systems, such a triangle paths or toll walks have been considered [10, 11]. We expect that some of these notions of betweenness will have interesting generalizations to directed paths.

Transit functions also feature interesting connections to other set systems and relations. Among symmetric transit functions, the geometric transit functions can be characterized in terms of their "base relations" \leq_b defined as

$$u \leq_b v \iff R(b,u) \subseteq R(b,v) \quad \text{for all } u, v, b \in V$$

$$\tag{5}$$

which must be a partial order satisfying $R(u, v) = \{x \mid u \leq_b x \leq_b v\}$, see [38, Prop. 5.2] and [43]. In the general case, however, equ.(5) yields $u \leq_b v$ and $v \leq_b u$ whenever $R(b, u) = R(b, v) = \emptyset$, i.e., \leq_b is not a total order on V. It will be interesting to explore whether this construction can be adjusted to yield a similar characterization for geometric directed transit functions.

Taken together, the directed transit functions introduced here appear to be a natural and useful generalization of the well-studied transit functions.

Declarations

Competing interests

The authors declare that they have no competing interests, or other interests that might be perceived to influence the results and/or discussion reported in this paper.

Funding

This work was supported in part by the DST, Govt. of India (Grant No. DST/INT/DAAD/P-03/ 2023 (G)), the DAAD, Germany (Grant No. 57683501), and the CSIR-HRDG for the Senior Research Fellowship (09/0102(12336)/2021-EMR-I).

Authors' contributions

MC, PGNS, and PFS designed the study. All authors contributed the mathematical results and the drafting of the manuscript.

References

 Mulder HM (2008) Transit functions on graphs (and posets). In: Changat M, Klavžar S, Mulder HM, Vijayakumar A (eds) Convexity in Discrete Structures, RMS Lecture Notes Series, pp 117–130

- [2] Mulder HM (1980) The interval function of a graph. Tech. rep., Centrum Voor Wiskunde en Informatica
- [3] Nebeský L (1994) A characterization of the interval function of a connected graph. Czechoslovak Mathematical Journal 44(1):173–178, DOI 10.21136/CMJ. 1994.128449
- [4] Mulder HM, Nebeský L (2009) Axiomatic characterization of the interval function of a graph. European Journal of Combinatorics 30(5):1172–1185, DOI 10.1016/j. ejc.2008.09.007
- [5] Nebeský L (2002) The induced paths in a connected graph and a ternary relation determined by them. Mathematica Bohemica 127(3):397–408, DOI 10.21136/MB. 2002.134072
- [6] Changat M, Mathew J (2004) Induced path transit function, monotone and Peano axiom. Discrete Mathematics 286(3):185–194, DOI 10.1016/j.disc.2004.02.017
- [7] Changat M, Mathew J, Mulder HM (2010) The induced path function, monotonicity and betweenness. Discrete Applied Mathematics 158(5):426–433, DOI 10. 1016/j.dam.2009.10.004
- [8] Changat M, Klavžar S, Mulder HM (2001) The all-paths transit function of a graph. Czechoslovak Math J 51:439–448, DOI 10.1023/A:1013715518448
- [9] Alcón L, Brešar B, Gologranc T, Gutierrez M, Sumenjak TK, Peterin I, Tepeh A (2015) Toll convexity. European Journal of Combinatorics 46:161–175, DOI 10. 1016/j.ejc.2015.01.002
- [10] Sheela LKK, Changat M, Peterin I (2023) Axiomatic characterization of the toll walk function of some graph classes. In: Bagchi A, Muthu R (eds) Conference on Algorithms and Discrete Applied Mathematics (CALDAM 2023), Springer, Cham, Lect. Notes Comp. Sci., vol 13947, pp 427–446, DOI 10.1007/ 978-3-031-25211-2.33
- [11] Sheela LKK, Changat M, Jacob J (2024) The weak-toll function of a graph: Axiomatic characterizations and first-order non-definability. In: Conference on Algorithms and Discrete Applied Mathematics (CALDAM 2024), Springer, Cham, Lect. Notes Computer Sci., vol 14508, pp 286–301, DOI 10.1007/ 978-3-031-52213-0.20
- [12] Bandelt HJ, Dress AWM (1989) Weak hierarchies associated with similarity measures — an additive clustering technique. Bull Math Biol 51:133–166, DOI 10.1007/BF02458841
- [13] Barthélemy JP, Brucker F (2008) Binary clustering. Discr Appl Math 156:1237– 1250, DOI 10.1016/j.dam.2007.05.024
- Bertrand P, Diatta J (2024) An interval convexity-based framework for multilevel clustering with applications to single-linkage clustering. Discr Appl Math 342:38– 63, DOI 10.1016/j.dam.2023.08.003
- [15] Changat M, Nezhad FH, Stadler PF (2018) Axiomatic characterization of transit functions of hierarchies. Ars Math Contemp 14:117–128, DOI 10.26493/ 1855-3974.831.e12
- [16] Changat M, Shanavas AV, Stadler PF (2024) Transit functions and pyramidlike binary clustering systems. Discr Appl Math In press; arXiv preprint doi: 10.48550/arXiv.2212.08721

- [17] Shanavas AV, Changat M, Hellmuth M, Stadler PF (2024) Unique least common ancestors and clusters in directed acyclic graphs. In: Kalyanasundaram S, Maheshwari A (eds) Algorithms and Discrete Applied Mathematics. CAL-DAM 2024, Springer, Cham, Lect. Notes Comp. Sci., vol 14508, pp 148–161, DOI 10.1007/978-3-031-52213-0_11, in press
- [18] Changat M, Nezhad FH, Mulder HM, Narayanan N (2018) A note on the interval function of a disconnected graph. Discussiones Math: Graph Th 28:39–38, DOI 10. 7151/dmgt.1990
- [19] Chartrand G, Zhang P (2000) The geodetic number of an oriented graph. European J Combin 21(2):181–189, DOI 10.1006/eujc.1999.0301
- [20] White DR, Borgatti SP (1994) Betweenness centrality measures for directed graphs. Social Networks 16(4):335–346, DOI 10.1016/0378-8733(94)90015-9
- [21] Brandes U (2001) A faster algorithm for betweenness centrality. J Math Sociology 25(2):163–177, DOI 10.1080/0022250X.2001.9990249
- [22] Estrada E (2011) The Structure of Complex Networks: Theory and Applications. Oxford Academic, Oxford, UK, DOI 10.1093/acprof:oso/
- [23] Fagerberg R, Flamm C, Kianian R, Merkle D, Stadler PF (2018) Finding the k best synthesis plans. J Cheminformatics 10:19, DOI 10.1186/s13321-018-0273-z
- [24] Thakur M, Tripathi R (2009) Linear connectivity problems in directed hypergraphs. Theor Comp Sci 410:2592–2618, DOI 10.1016/j.tcs.2009.02.038
- [25] Ausiello G, Italiano GF, Laura L, Nanni U, Sarracco F (2012) Structure theorems for optimum hyperpaths in directed hypergraphs. In: Mahjoub AR, Markakis V, Milis I, Paschos VT (eds) Combinatorial Optimization. ISCO 2012., Springer, Berlin, Heidelberg, Lect. Notes Comp. Sci., vol 7422, DOI 10.1007/ 978-3-642-32147-4_1
- [26] Dharmarajan R, Kannan K (2015) Hyper path and hyper cycles. Intl J Pure Appl Math 98(3):309–312, DOI 10.12732/ijpam.v98i3.2
- [27] Onodera T, Sadakane K, Shibuya T (2013) Detecting superbubbles in assembly graphs. In: Darling A, Stoye J (eds) International Workshop on Algorithms in Bioinformatics, Springer Verlag, Berlin, Heidelberg, vol 8126, pp 338–348, DOI 10. 1007/978-3-642-40453-5_26
- [28] Paten B, Eizenga JM, Rosen YM, Novak AM, Garrison E, Hickey G (2018) Superbubbles, ultrabubbles, and cacti. J Comp Biol 25:649–663, DOI 10.1089/cmb. 2017.0251
- [29] Gärtner F, Müller L, Stadler PF (2018) Superbubbles revisited. Alg Mol Biol 13:16, DOI 10.1186/s13015-018-0134-3
- [30] Sammeth M (2009) Complete alternative splicing events are bubbles in splicing graphs. J Comput Biol 16:1117–1140, DOI 10.1089/cmb.2009.0108
- [31] Courcelle B (2020) Betweenness of partial orders. RAIRO Theor Inf Appl 54:7, DOI 10.1051/ita/2020007
- [32] Nebeský L (2001) A characterization of the interval function of a (finite or infinite) connected graph. Czechoslovak Mathematical Journal 51(3):635–642, DOI doi. org/10.1023/A:101374432480x
- [33] Hellmuth M, Schaller D, Stadler PF (2023) Clustering systems of phylogenetic networks. Th Biosci 142:301–358, DOI 10.1007/s12064-023-00398-w

- [34] Aho AV, Garey MR, Ullman JD (1972) The transitive reduction of a directed graph. SIAM Journal on Computing 1:131–137, DOI 10.1137/0201008
- [35] Moyles DM, Thompson GL (1969) An algorithm for finding a minimum equivalent graph of a digraph. J ACM 16(3):455–460, DOI 10.1145/321526.321534
- [36] Brešar B, Changat M, Klavžar S, Mathews J, Mathews A, Prasanth NS (2009) Characterizing posets for which their natural transit functions coincide. Ars mathematica contemporanea 2(1)
- [37] Mathews A, Mathews J (2008) Transit functions on posets and lattices. In: Changat M, Klavžar S, Mulder HM, Vijayakumar A (eds) Convexity in Discrete Structures, RMS Lecture Notes Series, pp 105–116
- [38] Van de Vel MLJ (1993) Theory of Convex Sturctures, North Holland Mathematical Library, vol 50. North Holland, Amsterdam, DOI 10.1016/s0924-6509(09) x7015-7
- [39] Changat M, Narasimha-Shenoi PG, Seethakuttyamma G (2019) Betweenness in graphs: A short survey on shortest and induced path betweenness. AKCE Intl J Graphs Combinatorics 16:96–109, DOI 10.1016/j.akcej.2018.06.007
- [40] Chvátal V, Rautenbach D, Schäfer PM (2011) Finite Sholander trees, trees, and their betweenness. Discrete mathematics 311(20):2143–2147, DOI 10.1016/j.disc. 2011.06.011
- [41] Allender E, Lange KJ (1998) RUSPACE($\log n$) \subseteq DSPACE($\log 2n / \log \log n$). Theory of Computing Systems 31:539–550, DOI 10.1007/s002240000102
- [42] Furnas GW, Zacks J (1994) Multitrees: Enriching and reusing hierarchical structure. In: Proceedings of the SIGCHI Conference on Human Factors in Computing Systems, ser. CHI '94, ACM, New York, NY, pp 330–336, DOI 10.1145/191666. 191778
- [43] Bandelt HJ, Chepoi V (1996) A Helly theorem in weakly modular space. Discr Math 160:25–39, DOI 10.1016/0012-365X(95)00217-K
- [44] Nezhad FH (2016) Axiomatic characterizations of transit function on a hierarchy and standard path transit functions of special graph classes. PhD thesis, University of Kerala, Thiruvanathapuram, IND, unpublished PhD thesis
- [45] Lätsch M, Schrader R (2010) Distance-hereditary digraphs. J Discrete Algorithms 8(2):231–240, DOI 10.1016/j.jda.2009.11.002
- [46] Changat M, Mathew J (1999) Interval monotone graphs: minimal path convexity. In: Balakrishnan R, Mulder HM, Vijayakumar A (eds) Proceedings of the Conference on Graph Connections, Allied Publishers, New Delhi, IND, pp 87–90
- [47] Nebeský L (2002) The induced paths in a connected graph and a ternary relation determined by them. Mathematica Bohemica 127(3):397–408, DOI 10.21136/MB. 2002.134072
- [48] Sur N, Cataranciuc S (2008) About directed d-convex simple graphs. Comp Sci J Moldova 16(3):323–346
- [49] Nebeský L (1998) Characterizing the interval function of a connected graph. Mathematica Bohemica 123(2):137–144, DOI 10.21136/MB.1998.126307
- [50] Howorka E (1977) A characterization of distance-hereditary graphs. Quarterly J Math, 2nd Series 28(112):417–420, DOI 10.1093/qmath/28.4.417

- [51] Bandelt HJ, Mulder HM (1986) Distance-hereditary graphs. J Comb Th Ser B 41(2):182–208, DOI 10.1016/0095-8956(86)90043-2
- [52] Changat M, Kamal Kamalolbhavan-Sheela L, Narasimha-Shenoi PG (2024) The axiomatic characterization of the interval function of distance hereditary graphs. Discr Appl Math 350:62–70
- [53] Chalopin J, Changat M, Chepoi V, Jacob J (2024) First-order logic axiomatization of metric graph theory. Theoretical Computer Science 993:114,460, DOI 10.1016/j.tcs.2024.114460
- [54] Changat M, Sheela LKK, Narasimha-Shenoi PG (2023) Axiomatic characterizations of Ptolemaic and chordal graphs. Opuscula Math 43(3):293–407, DOI 10. 7494/OpMath.2023.43.3.393