On Tree Automata, Generating Functions, and Differential Equations

Rida Ait El Manssour CNRS, IRIF, France Vincent Cheval University of Oxford, UK Mahsa Shirmohammadi CNRS, IRIF, France

James Worrell University of Oxford, UK

Abstract

In this paper we introduce holonomic tree automata: a common extension of weighted tree automata and holonomic recurrences. We show that the generating function of the tree series represented by such an automaton is differentially algebraic. Conversely, we give an algorithm that inputs a differentially algebraic power series, represented as a solution of a rational dynamical system, and outputs an automaton whose generating function is the given series. Such an automaton yields a recurrence that can be used to compute the terms of the power series. We use the algorithm to obtain automaton representations of exponential generating functions of families of combinatorial objects given as combinatorial species. Using techniques from differential algebra, we show that it is decidable both whether two automata represent the same formal tree series and whether they have the same generating function.

1 Introduction

1.1 Overview

Weighted automata are a means of representing and effectively manipulating formal power series. Weighted word automata determine power series on non-commuting variables [BR10, Chapter 1], while weighted tree automata represent formal tree series: see [DKV09, DPV05]. (A formal tree series is a mapping that assigns numbers to trees, e.g., the function sending an arithmetic expression to its value is realisable by a weighted tree automaton [BR82, Example 6.2].) Each such automaton also has a *census generating function*, which is a univariate power series whose *n*-th coefficient denotes the total weight of all words or trees, respectively, of size n.

The generating function of a weighted word automaton is a rational power series while the generating function of a weighted tree automaton in an algebraic power series. The correspondence between weighted automata and their generating functions extends to the class of differentially finite power series (power series f(x) that satisfy linear differential equations $P_0(x)f(x) + P_1(x)f'(x) + \cdots + P_k(x)f^{(k)}(x) = 0$ with polynomial coefficients) which subsumes the class of algebraic power series. It is well-known that a power series $f(x) = \sum_{n=0}^{\infty} a_n x^n$ is differentially finite if and only if the sequence $(a_n)_{n=0}^{\infty}$ of coefficients is holonomic (also called *P*-recursive), i.e., the sequence satisfies a linear recurrence with polynomial coefficients [Kau23]. An automata formulation of such sequences is given in [Reu12].

In this paper we introduce a common generalisation of weighted tree automata and holonomic recurrences, called holonomic tree automata, that compute formal tree series. In a holonomic tree automaton the weight of each transition is a rational function in the size of the trees involved in the transition. Our main results concern decidability and expressiveness of the new model and focus on the link between automata and their generating functions. The corresponding class of generating functions goes beyond differentially finite series to include a general class of differentially algebraic power series.

Recall that a formal power series $f \in \mathbb{Q}[\![x]\!]$ is differentially algebraic if it satisfies a differential equation $P(f, f', \ldots, f^{(k)}) = 0$ for some polynomial $P(y_0, \ldots, y_k)$ with coefficients in $\mathbb{Q}[x]$. Every such series can be expressed as (a component) of a solution of a system of firstorder differential equations $y'_1 = P_1(y_1, \ldots, y_k), \ldots, y'_k = P_k(y_1, \ldots, y_k)$ for rational functions $P_1, \ldots, P_k \in \mathbb{Q}(x_1, \ldots, x_k)$. The latter is called a rational dynamical system [OPV22].¹ Our main results are as follows:

- We show that the class of generating functions of holonomic tree automata is closed under sum, Cauchy product, inverse, derivative, integration, forward and backward shift (Theorem 7 in Appendix A). We moreover show that every such series is differentially algebraic (Theorem 2).
- Conversely, we give an algorithm that inputs a differentially algebraic power series f, expressed as a (component of a) solution of a rational dynamical system

$$y'_1 = Q_1(y_1, \dots, y_k) \quad \dots \quad y'_k = Q_k(y_1, \dots, y_k),$$
 (1)

in which the rational functions Q_1, \ldots, Q_k are defined at $(y_1(0), \ldots, y_k(0))$, and outputs an automaton whose generating function equals f (Theorem 6). To illustrate the utility of this procedure, we give in Section 4.1 a collection of rational dynamical systems that characterise the exponential generation functions arising from standard constructions of combinatorial

¹Whether there exists a single rational dynamical system that has the same set of solutions as the equation $P(f, f', \ldots, f^{(k)}) = 0$ is called the *rational realisation* problem [FOR92, NS09] and is equivalent to the variety defined by P admitting a rational parameterisation.

species [BLL98]. This allows us to go from a specification of a combinatorial species to an automaton that represents its exponential generating function.

• We give a procedure to determine whether two automata represent the same formal tree series (Corollary 1). More specifically we give an effective bound on the size of the smallest tree on which two non-equivalent automata differ. This result is obtained as a corollary of a procedure for determining whether the generating function of a given holonomic tree automaton is identically zero (Theorem 5). The latter relies on techniques from differential algebra—specifically an effective version of the differential Nullstellensatz.

1.2 Weighted and Holonomic Tree Automata

Let Σ be a ranked alphabet, that is, Σ is a finite set of function symbols each having a non-negative integer arity. For $k \in \mathbb{N}$ we write Σ_k for the set of symbols in Σ of arity k. Let T_{Σ} be the set of terms over this alphabet, i.e., T_{Σ} is the smallest set that contains all symbols of arity 0 and such that if σ has arity k and $t_1, \ldots, t_k \in T_{\Sigma}$ then $\sigma(t_1, \ldots, t_k) \in T_{\Sigma}$. We consider elements of T_{Σ} as ordered trees where the vertices of $t \in T_{\Sigma}$ are its sub-terms and, for a k-ary symbol $\sigma \in \Sigma$, the tree $\sigma(t_1, \ldots, t_k)$ has list of children t_1, \ldots, t_k . The size ||t|| of a term $t \in T_{\Sigma}$ is the number of internal nodes and is defined inductively by $||\sigma|| := 0$ for σ a nullary symbol and $||\sigma(t_1, \ldots, t_k)|| := 1 + ||t_1|| + \cdots + ||t_k||$ for σ having arity $k \ge 1$.

Weighted Tree Automata As a prelude to the central definition of the paper we introduce the notion of weighted tree automaton (also called a multilinear representation) [BR82]. To this end, recall that the Kronecker product of two row vectors is defined by

$$(u_1 \quad u_2 \quad \cdots \quad u_m) \otimes (v_1 \quad v_2 \quad \cdots \quad v_n) := (u_1v_1 \quad u_1v_2 \quad \cdots \quad u_mv_{n-1} \quad u_mv_n)$$

The operation is associative and the nullary Kronecker product is the 1×1 identity matrix.

A weighted tree automaton over alphabet Σ is a pair $\mathcal{A} = (d, \mu)$, where $d \in \mathbb{N}$ is the dimension and μ is a function with domain Σ such that $\mu(\sigma) \in \mathbb{Q}^{d^k \times d}$ is a matrix of rational numbers for $\sigma \in \Sigma_k$. The map μ induces a function $\tilde{\mu} : T_{\Sigma} \to \mathbb{Q}^{1 \times d}$ that is defined inductively by specifying that

$$\widetilde{\mu}(\sigma(t_1,\ldots,t_k)) := (\widetilde{\mu}(t_1) \otimes \cdots \otimes \widetilde{\mu}(t_k)) \cdot \mu(\sigma)$$

for a k-ary symbol σ . In other words, $\tilde{\mu}(t)$ is given by an iterated matrix product determined by parsing the tree t from the leaves to the root. The formal tree series defined by \mathcal{A} is the map $\mathcal{A}[\![\cdot]\!]: T_{\Sigma} \to \mathbb{Q}$ by $\mathcal{A}[\![t]\!]:= \tilde{\mu}(t)_1$.

Two weighted tree automata over the same alphabet are said to be *equivalent* if they induce the same tree series. Equivalence of weighted tree automata was shown to be decidable in randomised polynomial time in [Sei90] and later to be logspace interreducible with Polynomial Identity Testing in [MW15].

We also associate with \mathcal{A} its generating function

$$f_{\mathcal{A}}(x) = \sum_{n=0}^{\infty} \left(\sum_{t: \|t\| = n} \mathcal{A}\llbracket t \rrbracket \right) x^{n}, \qquad (2)$$

in which the coefficient of x^n is the total value of all trees of size n. The generating function of a weighted tree automaton is an algebraic power series [BR82, Proposition 7.2]. Note that two

automata that represent different tree series can have the same generating function, since the latter groups together all trees of the same size. Equivalence of generating functions of weighted tree automata lies in the counting hierarchy [BCN⁺23].

Holonomic Tree Automata Let $\mathbb{Q}_u(x_0, \ldots, x_k)$ be the subring of $\mathbb{Q}(x_0, \ldots, x_k)$ consisting of rational functions of the form

$$\frac{P(x_0,\ldots,x_k)}{Q_0(x_0)\cdots Q_k(x_k)}$$

where $P \in \mathbb{Q}[x_0, \ldots, x_k]$ is a multivariate polynomial and $Q_i \in \mathbb{Q}[x_i]$ for $i \in \{0, \ldots, k\}$ are univariate polynomials such that Q_0 has no positive integer root and Q_1, \ldots, Q_k have no nonnegative integer root. Since the set of allowed denominators is closed under products $\mathbb{Q}_u(x_0, \ldots, x_k)$ is indeed a ring. The restriction of the codomain of μ to matrices with entries in the subring \mathbb{Q}_u will be used in the proof that the generating function of a holonomic tree automaton is differentially algebraic.

The notion of holonomic tree automata generalises that of weighted tree automata. A holonomic tree automaton over alphabet Σ is a pair $\mathcal{A} = (d, \mu)$, where $d \in \mathbb{N}$ is the dimension and μ is a map with domain Σ such that $\mu(\sigma) \in \mathbb{Q}^{1 \times d}$ for $\sigma \in \Sigma_0$ and, for $k \geq 1$ and $\sigma \in \Sigma_k$, $\mu(\sigma) \in \mathbb{Q}_u(x_0, \ldots, x_k)^{d^k \times d}$ is a matrix of rational functions.

The map μ induces a function $\widetilde{\mu}: T_{\Sigma} \to \mathbb{Q}^{1 \times d}$, inductively defined by specifying that $\widetilde{\mu}(\sigma) = \mu(\sigma)$ for $\sigma \in \Sigma_0$ and for $k \ge 1$ and $\sigma \in \Sigma_k$ and $t_0 = \sigma(t_1, \ldots, t_k)$,

$$\widetilde{\mu}(t_0) := (\widetilde{\mu}(t_1) \otimes \cdots \otimes \widetilde{\mu}(t_k)) \cdot \mu(\sigma)(\|t_0\|, \dots, \|t_k\|).$$

We define the formal tree series represented by \mathcal{A} to be $\mathcal{A}[\![\cdot]\!]: T_{\Sigma} \to \mathbb{Q}$ by $\mathcal{A}[\![t]\!]:=\widetilde{\mu}(t)_1$.

The definition of the generating function $f_{\mathcal{A}}$ of a holonomic tree automaton \mathcal{A} is exactly as for a weighted tree automaton, namely via Equation (2). In general $f_{\mathcal{A}}$ need not be algebraic, unlike for weighted tree automata, but we will show that it is differentially algebraic.

Examples. In Theorem 6 we give a procedure that inputs a power series, represented as a rational dynamical system and outputs a holonomic automata whose generating function equals the given power series. Here, we give some specific examples of holonomic automata that arise from differential equations that have a combinatorial flavour. In all the following examples the signature is $\Sigma = \{\sigma_0, \sigma_1, \sigma_2\}$ where σ_k has arity k.

Example 1 (Bell numbers). Recall that the sequence $(b_n)_{n=0}^{\infty} = (1, 1, 2, 5, 15, 52, 203, ...)$ of Bell numbers has general term the number of partitions of the set $\{1, \ldots, n\}$. As noted in [BR90], the exponential generating function $f_B(x) := \sum_{n=0}^{\infty} \frac{b_n}{n!} x^n$ is (the *f*-component of) a solution of the system of differential equations f'(x) = f(x)g(x) and g'(x) = g(x).

We specify an automaton $\mathcal{A} = (2, \mu)$ over alphabet Σ such that $f_{\mathcal{A}} = f_B$. The weight function is defined as follows:

$$\mu(\sigma_0) := \begin{bmatrix} 1 & 1 \end{bmatrix} \qquad \mu(\sigma_1) := \begin{bmatrix} 0 & 0 \\ 0 & \frac{1}{x_0} \end{bmatrix} \qquad \mu(\sigma_2) := \begin{bmatrix} \mathbf{0}_{2\times 1} & \mathbf{0}_{2\times 1} \\ \frac{1}{x_0} & 0 \\ 0 & 0 \end{bmatrix}$$

We outline a combinatorial interpretation of the equality $f_{\mathcal{A}} = f_B$. The set $\{t \in T_{\Sigma} : [\mathcal{A}]](t) \neq 0\}$ comprises those trees $t \in T_{\Sigma}$ such that all internal nodes in t reachable from the root by zero or more steps of the left-child relation have arity two, while all other internal nodes have arity one (see Figure 1.2). For such a tree t of size n, deleting all edges between a node and its left-child one



Figure 1: Encoding partitions as ranked binary trees.

obtains a forest of chains. If the chains have respective sizes k_1, \ldots, k_s , reading from left to right in t, we have

$$\llbracket \mathcal{A} \rrbracket(t) = \frac{1}{k_1! \cdots k_s!} \frac{k_1}{k_1 + k_2} \cdots \frac{k_{s-1}}{k_1 + \cdots + k_s}.$$
(3)

Note that every labelling of the internal nodes of t in the numbers $\{1, \ldots, n\}$ naturally yields a partition of $\{1, \ldots, n\}$. For any fixed partition π of $\{1, \ldots, n\}$, summing the expression for $[\![\mathcal{A}]\!](t)$ in (3) over all trees $t \in T_{\Sigma}$ and labellings of the nodes of t that yield π , we obtain n!.

Example 2 (Labelled trees). Let $(t_n)_{n=0}^{\infty} = (0, 1, 2, 9, 64, 625, ...)$ be the sequence whose general term counts the number of labelled rooted trees with n nodes. It is classical that $t_n = n^{n-1}$ for all $n \ge 1$. From this closed form (or from the characterisation of labelled trees as a combinatorial species) it can be seen that the exponential generating function $f_T(x) = \sum_{n=0}^{\infty} \frac{t_n}{n!} x^n$ is the f-component of the solution of the system

$$f(x) = xg(x), \quad g'(x) = f'(x)g(x).$$

From the above system of equations we obtain an automaton $\mathcal{A} = (2, \mu)$ over alphabet Σ whose generating function coincides with f_T . The definition of μ is as follows:

$$\mu(\sigma_0) = \begin{bmatrix} 0 & 1 \end{bmatrix} \qquad \mu(\sigma_1) = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \qquad \mu(\sigma_2) = \begin{bmatrix} \mathbf{0}_{3\times 1} & \mathbf{0}_{3\times 1} \\ 0 & \frac{x_2+1}{x_0} \end{bmatrix}$$

The equality $f_{\mathcal{A}} = f_T$ again admits a combinatorial interpretation. The set $\{t \in T_{\Sigma} : [\![\mathcal{A}]\!](t) \neq 0\}$ comprises those trees $t \in T_{\Sigma}$ in which the root has arity one and all other internal nodes have arity two. Each such tree t determines in a natural way a rooted unordered tree τ (in which there is no bound on the number of children of a node): the idea is that a left-child in t encodes the sibling relation in τ (see Figure 2). Now every labelling of t yields a labelling of τ and, similarly to the previous example, summing $[\![\mathcal{A}]\!](t)$ over all trees t and labellings of the nodes of t that give the same labelled rooted unordered tree gives n!.

Example 3. Consider the solution $y_1(x)$ of the differential equation $(y'_1)^3 + y_1^3 = 1$ such that $y_1(0) = 0$ and $y'_1(0) = 1$. Differentiating the equation we deduce that $y_1(x)$ also satisfies $3(y'_1)^2 y''_1 + y''_1 = 1$.



Figure 2: Encoding rooted non-plane trees as ranked binary trees.

 $3(y_1)^2 y'_1 = 0$, which leads to the following rational dynamical system:

$$\begin{cases} y_1' = y_2 \\ y_2' = -\frac{y_1^2}{y_2} \end{cases}$$

As $y_2(0) = y'_1(0) \neq 0$, the proof of Theorem 6 constructs an automaton $\mathcal{A} = (4, \mu)$ such that $f_{\mathcal{A}}(x) = y_1(x)$ (see Example 5 for more details). The weight function is defined by

$$\mu(\sigma_0) = \begin{bmatrix} 0 & 1 & 0 & 0 \end{bmatrix} \qquad \mu(\sigma_1)(x_0, x_1) = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & \frac{1}{x_0 + 1} & 0 & 0 \\ 0 & 0 & \frac{1}{x_0 + 1} & 0 \end{bmatrix}$$

and

$$\mu(\sigma_2)(x_0, x_1, x_2) = \begin{bmatrix} \mathbf{0}_{6\times 1} & \mathbf{0}_{6\times 1} & \mathbf{0}_{6\times 1} & \mathbf{0}_{6\times 1} \\ 0 & 0 & 0 & -1 \\ \mathbf{0}_{3\times 1} & \mathbf{0}_{3\times 1} & \mathbf{0}_{3\times 1} & \mathbf{0}_{3\times 1} \\ 0 & 0 & -\frac{x_2+1}{x_0+1} & 0 \\ \mathbf{0}_{5\times 1} & \mathbf{0}_{5\times 1} & \mathbf{0}_{5\times 1} & \mathbf{0}_{5\times 1} \end{bmatrix}$$

Running the automaton, we compute its generating function $f_{\mathcal{A}}$ as:

$$y_1(x) = x - \frac{2}{4!}x^4 - \frac{20}{7!}x^7 - \frac{3320}{10!}x^{10} - \frac{1598960}{13!}x^{13} - \frac{1757280800}{16!}x^{16} - \dots$$

1.3 Organisation of the Paper

Our main contributions concern the expressiveness and effectiveness of holonomic tree automata, both in terms of their generating functions and the formal tree series that they define.

In Section 2 we show that the generating function $f_{\mathcal{A}}$ of a holonomic tree automaton \mathcal{A} is (effectively) differentially algebraic. To show this we extract from the definition of \mathcal{A} a system of differential polynomials that have a unique common zero in power series, a component of which is $f_{\mathcal{A}}$. Now a differential-algebraic version of Artin's approximation theorem [DL84, Theorem 2.1] states that any power-series zero of a finite collection of differential polynomials can be approximated to arbitrary order by a zero that is a differentially algebraic power-series. We conclude that $f_{\mathcal{A}}$, being (a component of) a *unique* solution of such a system, is itself differentially algebraic. In Section 3 we give a procedure for determining whether the generating function $f_{\mathcal{A}}$ of a holonomic automaton \mathcal{A} is identically zero. Here we again exploit the fact that $f_{\mathcal{A}}$ is a component of the unique zero in power series of a system of differential polynomials. This allows us to reduce the problem of determining zeroness of $f_{\mathcal{A}}$ to that of checking whether a certain collection of differential polynomials has a common zero in power series. Here we use an effective version of the differential Nullstellensatz [OPV22]. This gives an explicit bound B such that our collection of differential polynomials has no zero in power series if and only if the polynomial ideal generated by the first B derivatives has no common zero. From this we deduce that $f_{\mathcal{A}}$ is identically zero if and only if it is zero modulo x^{B+1} and decidability follows immediately.

As a corollary of the ability to check zeroness of the generating function $f_{\mathcal{A}}$ we can decide whether the generating functions of two given holonomic tree automata \mathcal{A} and \mathcal{B} are equal and also whether their formal tree series are equal, that is, whether $[\![\mathcal{A}]\!](t) = [\![\mathcal{B}]\!](t)$ for all trees t. In the latter case we obtain an explicit upper bound B such that if $[\![\mathcal{A}]\!] \neq [\![\mathcal{B}]\!]$ then the two functions differ already on a tree of size B.

In Section 4 we give an effective procedure that inputs a power series f, presented as the solution of a rational dynamical system (subject to a condition on the poles of the rational function, as specified above), and outputs a holonomic tree automaton \mathcal{A} whose generating function equals f. We illustrate this procedure with several examples. Furthermore, in Section 4.1 we give a translation of a standard collection of specifications on combinatorial species to rational dynamical systems that define their exponential generating functions. Combining this translation with the abovementioned procedure, we obtain a general method that given the specification of a combinatorial species outputs an automaton \mathcal{A} such that $f_{\mathcal{A}}$ is the exponential generating function of the species.

Finally, in Appendix A we provide effective constructions to show that the class of generating functions of holonomic tree automata is closed under sum, Cauchy product, inverse, derivative, integration, forward and backward shift. We also show that the class of formal tree series recognizable by a holonomic tree automata is closed under sum and Hadamard product.

1.4 Related Work

A power series that satisfies the system (1) in the special case that P_1, \ldots, P_k are polynomials is called *constructibly differentially algebraic (CDA)*. Combinatorial interpretations of CDA series were given in [BR90] (via an automaton-like notion called tree labelling tables) and in [LV86] (via combinatorial species). However the class of CDA power series does not include all differentially algebraic series. Indeed, as observed in [BR90], there are already *D*-finite power series that are not CDA.

Reutnauer [Reu12] introduces and studies a matrix representation of holonomic sequences. These correspond to the special case of holonomic tree automata over words, that is, over an alphabet with one nullary and one unary symbol. Among other things, [Reu12] shows the existence of a minimal representation of a given sequence that is unique up to change of base.

A Parikh automaton is a non-deterministic finite automaton in which the acceptance condition is a formula of linear arithmetic that is evaluated on the Parikh image of the run (i.e., the count of each edge in the run). It is shown in [BCKN20, CM17] that the multivariate generating series of the language accepted by an unambiguous Parikh automaton is *D*-finite.

The paper [SMK15] introduces a class of tree automata with registers that have polynomial updates and proves decidability of equivalence of the formal trees series they induce. As a corollary, the paper shows decidability of equivalence of deterministic top-down tree-to-word transducers, resolving a longstanding open problem. The decision procedure is, however, non-primitive recursive—the authors show that both equivalence and non-equivalence of formal tree series are semi-decidable. The semi-decidability of non-equivalence is obvious, while semi-decidability of equivalence follows from the fact that if two such automata are equivalent then there exists an algebraic inductive invariant that certifies equivalence and such an invariant can be found by exhaustive search. We believe that the same invariant technique could be applied to show decidability of equivalence of holonomic tree automata (although not necessarily of the associated generating functions).

2 From Automata to Differentially Algebraic Functions

In this section, we show that the generating functions of all holonomic tree automata are differentially algebraic. We start by introducing some notation that will be used throughout this section. We assume the existence of an automaton $\mathcal{A} = (d, \mu)$ over the alphabet Σ . We associate d formal power series $f_1(x), \ldots, f_d(x)$ defined for $i \in \{1, \ldots, d\}$ by $f_i(x) := \sum_{n=0}^{\infty} (\boldsymbol{a}_n)_i x^n$ where for all $n \in \mathbb{N}$, $\boldsymbol{a}_n = \sum_{\substack{t \in T_{\Sigma} \\ |t|=n}} \tilde{\mu}(t)$. Moreover, we denote by $\boldsymbol{f}(x)$ the vector of power series $[f_1(x) \ldots f_d(x)]$. In other word, we have $\boldsymbol{f}(x) = \sum_{n=0}^{\infty} \boldsymbol{a}_n x^n$ and $\boldsymbol{f}_1(x) = f_{\mathcal{A}}(x)$.

Definition 1. Let $f(x) \in \mathbb{Q}[\![x]\!]$ be a power series, we say that f(x) is differentially algebraic, or *D*-algebraic for short if f(x) satisfies the differential equation $P(y, y', \ldots, y^{(k)}) = 0$, where $P(y_0, \ldots, y_k)$ is a polynomial with coefficients in $\mathbb{Q}(x)$.

Remark 1. We can without loss of generality assume that the coefficients of the polynomial P in definition 1 are in \mathbb{Q} . Indeed, as shown in [vDH19, Proposition 2], for example, f(x) satisfies a polynomial differential equation with coefficients in \mathbb{Q} if and only if the transcendence degree $\operatorname{trdeg}(\mathbb{Q}(\{f^{(i)}(x) : i \in \mathbb{N}\})/\mathbb{Q})$ is finite. However, as $\operatorname{trdeg}(\mathbb{Q}(x)(\{f^{(i)}(x) : i \in \mathbb{N}\})/\mathbb{Q}(x))$ being finite implies $\operatorname{trdeg}(\mathbb{Q}(\{f^{(i)}(x) : i \in \mathbb{N}\})/\mathbb{Q})$ is also finite, we deduce that f(x) is D-algebraic if and only if f(x) satisfies a polynomial differential equation with coefficients in $\mathbb{Q}(x)$.

The argument to prove that the generating functions of automata are differentially algebraic is organised as follows: (i) we start by stating some useful properties of f, its derivatives and the coefficient vectors a_n ; (ii) we show that f(x) is a component of a solution of a system of differential equations; (iii) we show that this system of equations admits a unique vector of power series solution; (iv) finally we show that the unique solution is differentially algebraic.

Useful properties. By definition, we have:

$$f(x) = \sum_{n=0}^{\infty} a_n x^n$$
 with $a_n = \sum_{\substack{t \in T_{\Sigma} \\ ||t||=n}} \widetilde{\mu}(t) \quad \forall n \in \mathbb{N}$

Notice that the only trees of size 0 are the nullary function symbols, hence $a_0 = \sum_{a \in \Sigma_0} \mu(a)$. For all n > 0, splitting the sum into the possible shapes that trees can take, we obtain:

$$\boldsymbol{a}_n = \sum_{\substack{t \in T_{\Sigma} \\ \|t\| = n}} \widetilde{\mu}(t) = \sum_{\substack{k > 0 \\ g \in \Sigma_k}} \sum_{\substack{n_1, \dots, n_k \in \mathbb{N} \\ n_1 + \dots + n_k = n-1}} \sum_{\substack{t_1 \in T_{\Sigma} \\ \|t\| = n_1}} \dots \sum_{\substack{t_k \in T_{\Sigma} \\ \|t\| = n_k}} (\widetilde{\mu}(t_1) \otimes \dots \otimes \widetilde{\mu}(t_k)) \cdot \mu(g)(n, n_1, \dots, n_k)$$

Using the bilinearity of the Kronecker product, we obtain:

$$\boldsymbol{a}_{n} = \sum_{\substack{k>0\\g\in\Sigma_{k}}} \sum_{\substack{n_{1},\dots,n_{k}\in\mathbb{N}\\n_{1}+\dots+n_{k}=n-1}} \left(\sum_{\substack{t_{1}\in T_{\Sigma}\\\|t\|=n_{1}}} \widetilde{\mu}(t_{1})\otimes\dots\otimes\sum_{\substack{t_{k}\in T_{\Sigma}\\\|t\|=n_{k}}} \widetilde{\mu}(t_{k}) \right) \cdot \mu(g)(n,n_{1},\dots,n_{k})$$

$$=\sum_{\substack{k>0\\g\in\Sigma_k}}\sum_{\substack{n_1,\dots,n_k\in\mathbb{N}\\n_1+\dots+n_k=n-1}} (a_{n_1}\otimes\dots\otimes a_{n_k})\cdot\mu(g)(n,n_1,\dots,n_k)$$
(4)

The property in Equation (4) is very useful as it expresses the coefficient vectors a_n in terms of the previous coefficient vectors a_m , m < n. We will reuse this property in most sections of the paper. For instance, by applying this equation to the definition of f, we obtain:

$$f(x) - a_0 = \sum_{n=1}^{\infty} a_n x^n$$

=
$$\sum_{n=1}^{\infty} \sum_{\substack{k>0\\g\in\Sigma_k}} \sum_{\substack{n_1,\dots,n_k\in\mathbb{N}\\n_1+\dots+n_k=n-1}} (a_{n_1}\otimes\dots\otimes a_{n_k}) \cdot \mu(g)(n,n_1,\dots,n_k) x^n$$

=
$$\sum_{\substack{k>0\\g\in\Sigma_k}} \sum_{n=0}^{\infty} \sum_{\substack{n_1,\dots,n_k\in\mathbb{N}\\n_1+\dots+n_k=n}} (a_{n_1}\otimes\dots\otimes a_{n_k}) \cdot \mu(g)(n+1,n_1,\dots,n_k) x^{n+1}$$

Observe that for any expression $fun(n, n_1, \ldots, n_k)$, we have:

$$\sum_{n=0}^{\infty} \sum_{\substack{n_1,\dots,n_k \in \mathbb{N} \\ n_1+\dots+n_k=n}} fun(n,n_1,\dots,n_k) = \sum_{n_1=0}^{\infty} \dots \sum_{n_k=0}^{\infty} fun(\sum_{i=1}^k n_i,n_1,\dots,n_k)$$
(5)

Hence:

$$\boldsymbol{f}(x) - \boldsymbol{a}_0 = x \sum_{\substack{k>0\\g\in\Sigma_k}} \sum_{n_1=0}^{\infty} \dots \sum_{n_k=0}^{\infty} (\boldsymbol{a}_{n_1} x^{n_1} \otimes \dots \otimes \boldsymbol{a}_{n_k} x^{n_k}) \mu(g) (1 + \sum_{i=1}^k n_i, n_1, \dots, n_k)$$
(6)

Equation (6) is the starting point for building the system of differential equations satisfied by $f_{\mathcal{A}}(x)$. Indeed, notice that the sums $\sum_{n_1=0}^{\infty} a_{n_1} x^{n_1}, \ldots$ are exactly f. However, this is not sufficient as the term $\mu(g)(1 + \sum_{i=1}^{k} n_i, n_1, \ldots, n_k)$ contains $n_1, n_2 \ldots$ which need to be removed. To do so, we will rely on the derivatives of f.

Let us denote by $\mathbf{f}^{(i)}(x)$ the *i*-th derivative of the vector of power series $\mathbf{f}(x)$, that is $\mathbf{f}^{(i)}(x) = \begin{bmatrix} f_1^{(i)}(x) & \dots & f_d^{(i)}(x) \end{bmatrix}$. We also introduce the operator $\Theta \mathbf{f}(x) := x \mathbf{f}'(x)$. We write $\Theta^{(0)} \mathbf{f}(x) = \mathbf{f}(x)$ and for all $i \in \mathbb{N}$, $\Theta^{(i)} \mathbf{f}(x) = \Theta(\Theta^{(i-1)} \mathbf{f})(x)$.

Proposition 1. For all vectors of power series $f(x) = \sum_{n=0}^{\infty} a_n x^n$, writing $0^0 = 1$, we have for all $i \in \mathbb{N}, \ \Theta^{(i)} f(x) = \sum_{n=0}^{\infty} n^i a_n x^n$.

Proof. We prove this result by induction on i. The base case (i = 0) is trivial. In the inductive step (i > 0), $\Theta^{(i-1)} f(x) = \sum_{n=0}^{\infty} n^{i-1} a_n x^n$ by inductive hypothesis, hence

$$\Theta(\Theta^{(i-1)}\boldsymbol{f})(x) = x(\Theta^{(i-1)}\boldsymbol{f})'(x) = x\sum_{n=1}^{\infty} n^{i}\boldsymbol{a}_{n}x^{n-1} = \sum_{n=1}^{\infty} n^{i}\boldsymbol{a}_{n}x^{n-1}$$

As i > 0, we have $0^i = 0$ and so $\Theta^{(i)} \boldsymbol{f}(x) = \sum_{n=0}^{\infty} n^i \boldsymbol{a}_n x^n$.

Building the system of differential equations. By definition, when $g \in \Sigma_k$, the coefficients of the matrix $\mu(g)$ are in $\mathbb{Q}_u(x_0, \ldots, x_k)$, i.e., they are the rational functions of the form

$$\frac{P(x_0,\ldots,x_k)}{Q_0(x_0)\cdots Q_k(x_k)}$$

where $P \in \mathbb{Q}[x_0, \ldots, x_k]$ and $Q_i \in \mathbb{Q}[x_i]$ for $i \in \{1, \ldots, k\}$ are polynomials with no nonnegative integer root and $Q_0 \in \mathbb{Q}[x_0]$ is a polynomial with no positive integer root. Without loss of generality we can always assume that the coefficients in the matrix $\mu(g)$ have a common denominator. We cannot make this assumption throughout all the matrices of the weight function as two symbols may have different arity, and so their denominators are polynomials in different numbers of variables. However, the matrices of the weight function associated to all non-nullary symbols share at least the variable x_0 , meaning that we can assume that the univariate polynomials associated to x_0 in all denominators are the same. In other words, we can assume without loss of generality that there exist $Q \in \mathbb{Q}[x]$, and $r \in \mathbb{N}$, such that for all $k \geq 1$, for all $g \in \Sigma_k$, there exist some matrices $\mu_{g,(i_1,\ldots,i_k)} \in \mathbb{Q}^{d^k \times d}$ for $i_1, \ldots, i_k \in \{0, \ldots, r\}$ and polynomials $Q_{g,1}, \ldots, Q_{g,k} \in \mathbb{Q}[x]$ such that:

$$\mu(g)(x_0, x_1, \dots, x_k) = \frac{1}{Q(x_0)} \sum_{i_1, \dots, i_k \in \{0, \dots, r\}} \frac{x_1^{i_1} \dots x_k^{i_k}}{Q_{g,1}(x_1) \dots Q_{g,k}(x_k)} \mu_{g,(i_1, \dots, i_k)}$$
(7)

Once the weight function is in the shape described in Equation (7), we can build the system of differential equations satisfied by f(x) as described in the following proposition.

Proposition 2. Assume that $Q(x) = \sum_{i=0}^{u} c_i x^i$ for some $c_0, \ldots, c_u \in \mathbb{Q}$ and $u \in \mathbb{N}$. Moreover, for all $k \geq 1$, for all $g \in \Sigma_k$, for all $i \in \{1, \ldots, k\}$,

- denote by $p_{g,i}(x)$ the vector of power series $\sum_{n=0}^{\infty} a_n \frac{x^n}{Q_{g,i}(n)}$; and
- assume that $Q_{g,i}(x) = \sum_{j=0}^{u} b_{g,i,j} x^j$ for some $b_{g,i,0}, \ldots, b_{g,i,u} \in \mathbb{Q}$.

The vectors of power series f(x) and $p_{g,i}(x)$ for all $k \ge 1, g \in \Sigma_k, i \in \{1, \ldots, k\}$ satisfy the following system of $d(1 + \sum_{k\ge 1} |\Sigma_k| k)$ differential equations:

$$\sum_{i=0}^{u} c_{i} \Theta^{(i)} \boldsymbol{f} - c_{0} \boldsymbol{a}_{0} = x \sum_{\substack{k>0\\g\in\Sigma_{k}}} \sum_{i_{1},\dots,i_{k}\in\{0,\dots,r\}} (\Theta^{(i_{1})} \boldsymbol{p}_{g,1} \otimes \dots \otimes \Theta^{(i_{k})} \boldsymbol{p}_{g,k}) \mu_{g,(i_{1},\dots,i_{k})} \tag{8}$$

$$\left\{ \boldsymbol{f} = \sum_{j=0}^{u} b_{g,i,j} \Theta^{(j)} \boldsymbol{p}_{g,i} \qquad \forall k \ge 1, \forall g \in \Sigma_k, \forall i \in \{1, \dots, k\} \right\}$$
(9)

Proof. By definition, $\mathbf{p}_{g,i}(x) = \sum_{n=0}^{\infty} \mathbf{a}_n \frac{x^n}{Q_{g,i}(n)}$. Hence, by Proposition 1, for all $j \in \mathbb{N}$, $\Theta^{(j)} \mathbf{p}_{g,i}(x) = \sum_{n=0}^{\infty} \mathbf{a}_n \frac{n^j x^n}{Q_{g,i}(n)}$. Moreover, as $Q(x) = \sum_{i=0}^{u} c_i x^i$, we combine all these into the following sum:

$$\sum_{i=0}^{u} c_i \Theta^{(i)} \boldsymbol{f}(x) - c_0 \boldsymbol{a}_0 = c_0 \sum_{n=0}^{\infty} \boldsymbol{a}_n x^n - c_0 \boldsymbol{a}_0 + \sum_{i=1}^{u} c_i \sum_{n=0}^{\infty} n^i \boldsymbol{a}_n x^n$$
$$= \sum_{i=0}^{u} c_i \sum_{n=0}^{\infty} (n+1)^i \boldsymbol{a}_{n+1} x^{n+1}$$

$$= \sum_{n=0}^{\infty} \sum_{i=0}^{u} c_i (n+1)^i a_{n+1} x^{n+1}$$

$$= \sum_{n=0}^{\infty} Q(n+1) \sum_{\substack{k>0\\g\in\Sigma_k}} \sum_{\substack{n_1,\dots,n_k\in\mathbb{N}\\n_1+\dots+n_k=n}} (a_{n_1}\otimes\dots\otimes a_{n_k}) \cdot \mu(g)(n+1,n_1,\dots,n_k) x^{n+1}$$

$$= \sum_{n=0}^{\infty} Q(n+1) \sum_{\substack{k>0\\g\in\Sigma_k}} \sum_{\substack{n_1,\dots,n_k\in\mathbb{N}\\n_1+\dots+n_k=n}} (a_{n_1}\otimes\dots\otimes a_{n_k}) \cdot \frac{1}{Q(n+1)} \sum_{\substack{i_1,\dots,i_k\in\{0,\dots,r\}\\i_1,\dots,i_k\in\{0,\dots,r\}}} \frac{n_1^{i_1}\dots n_k^{i_k}}{Q_{g,1}(n_1)\dots Q_{g,k}(n_k)} \mu_{g,(i_1,\dots,i_k)} x^{n+1}$$

$$= \sum_{\substack{k>0\\g\in\Sigma_k}} \sum_{\substack{i_1,\dots,i_k\in\{0,\dots,r\}\\n_1+\dots+n_k=n}} \sum_{\substack{n_1,\dots,n_k\in\mathbb{N}\\n_1+\dots+n_k=n}} (a_{n_1}\otimes\dots\otimes a_{n_k}) \cdot \frac{n_1^{i_1}\dots n_k^{i_k}}{Q_{g,1}(n_1)\dots Q_{g,k}(n_k)} \mu_{g,(i_1,\dots,i_k)} x^{n+1}$$

Once again, applying Equation (5) and the bilinearity of Kronecker product, we establish the first part of our system of differential equations, that is Equation (8):

$$\sum_{i=0}^{u} c_{i} \Theta^{(i)} \boldsymbol{f}(x) - c_{0} \boldsymbol{a}_{0} = x \sum_{\substack{k>0\\g\in\Sigma_{k}}} \sum_{i_{1},\dots,i_{k}\in\{0,\dots,r\}} \sum_{n_{1}=0}^{\infty} \dots \sum_{n_{k}=0}^{\infty} \left(\frac{n_{1}^{i_{1}} \boldsymbol{a}_{n_{1}} x^{n_{1}}}{Q_{g,1}(n_{1})} \otimes \dots \otimes \frac{n_{k}^{i_{k}} \boldsymbol{a}_{n_{k}} x^{n_{k}}}{Q_{g,k}(n_{k})}\right) \cdot x^{n_{1}}$$
$$= x \sum_{\substack{k>0\\g\in\Sigma_{k}}} \sum_{i_{1},\dots,i_{k}\in\{0,\dots,r\}} (\Theta^{(i_{1})} \boldsymbol{p}_{g,1}(x) \otimes \dots \otimes \Theta^{(i_{k})} \boldsymbol{p}_{g,k}(x)) \mu_{g,(i_{1},\dots,i_{k})}$$

We complete the verification of the system of differential equations by showing, for all k > 0, all $g \in \Sigma_k$ and all $i \in \{1, \ldots, k\}$, that f(x) and $p_{g,i}(x)$ satisfy Equation (9). Recall that $Q_{g,i}(x) = \sum_{i=0}^{u} b_{g,i,j} x^j$. This leads to the following:

$$\sum_{j=0}^{u} b_{g,i,j} \Theta^{(j)} \boldsymbol{p}_{g,i}(x) = \sum_{j=0}^{u} b_{g,i,j} \sum_{n=0}^{\infty} n^{j} \boldsymbol{a}_{n} \frac{x^{n}}{Q_{g,i}(n)}$$
$$= \sum_{n=0}^{\infty} \left(\sum_{j=0}^{u} b_{g,i,j} n^{j} \right) \boldsymbol{a}_{n} \frac{x^{n}}{Q_{g,i}(n)}$$
$$= \sum_{n=0}^{\infty} Q_{g,i}(n) \boldsymbol{a}_{n} \frac{x^{n}}{Q_{g,i}(n)}$$
$$= \boldsymbol{f}(x)$$

Uniqueness of solutions. The system of differential equations given in Proposition 2 is fully defined by the automaton \mathcal{A} and its weight function μ . From Proposition 2, we showed that the vector of power series f(x) defined at the beginning of the section in conjunction with the vectors of power series $p_{g,i}(x)$ are solution of this system. We now show that these vectors were the only possible solution.

Proposition 3. For a given $\overline{a}_0 \in \mathbb{Q}^d$, there exists a unique power series solution $(\overline{f}(x), \{\overline{p}_{g,1}(x), \ldots, \overline{p}_{g,k}(x)\}_{k>0,g\in\Sigma_k})$ to the system of differential equations 8 and 9 from Proposition 2 where $\overline{f}(x) = \sum_{i=0}^{\infty} \overline{a}_i x^i$. In other words, if $\overline{a}_0 = a_0$ then $\overline{f}(x) = f(x)$, where f(x) and a_0 are those coming from the automaton.

Proof. Note that the equations of 9 give unique relations between the n^{th} coefficient of $\overline{f}(x)$ and the n^{th} coefficient of $\overline{p}_{g,i}(x)$ for every i, n. In fact, for all k > 0, for all $g \in \Sigma_k$, for all $i \in \{1, \ldots, k\}$, denoting by $d_{g,i,n}$ the n^{th} coefficient of $\overline{p}_{g,i}(x)$ (note that $d_{g,i,n}$ is a vector), we have for all $n \in \mathbb{N}$,

$$\sum_{j=0}^{u} b_{g,i,j} n^{j} \boldsymbol{d}_{g,i,n} = Q_{g,i}(n) \boldsymbol{d}_{g,i,n} = \overline{\boldsymbol{a}}_{n}.$$
(10)

Therefore, as $Q_{g,i}(n)$ does not vanish for all $n \in \mathbb{N}$, we deduce from 10 that $d_{g,i,n}$ is uniquely determined by \overline{a}_n .

We show by a simple inductive proof on n that every \overline{a}_n are uniquely defined. The base case is trivial since \overline{a}_0 is given. In the inductive step (n > 0), we know from our inductive hypothesis that all $\overline{a}_0, \ldots, \overline{a}_{n-1}$ are uniquely defined. Thus, $\overline{d}_{g,i,0}, \overline{d}_{g,i,1}, \ldots, \overline{d}_{g,i,n-1}$ are also uniquely defined for all k > 0, for all $g \in \Sigma_k$, for all $i \in \{1, \ldots, k\}$.

Note that the n^{th} coefficient of $\sum_{i=0}^{u} c_i \Theta^{(i)} \overline{f}(x) - c_0 \overline{a}_0$ is $\sum_{i=0}^{u} c_i n^i \overline{a}_n = Q(n) \overline{a}_n$ (since n > 0). On the other hand, the n^{th} coefficient of the right hand side of the Equation (8) is the $(n-1)^{th}$ coefficient of

$$\sum_{\substack{k>0\\g\in\Sigma_k}} \sum_{i_1,\ldots,i_k\in\{0,\ldots,r\}} (\Theta^{(i_1)}\overline{p}_{g,1}(x)\otimes\ldots\otimes\Theta^{(i_k)}\overline{p}_{g,k}(x))\mu_{g,(i_1,\ldots,i_k)}$$

which is determined uniquely by $\overline{d}_{g,i,0}, \ldots, \overline{d}_{g,i,m}$ with $k > 0, g \in \Sigma_k$, and $i \in \{1, \ldots, k\}$. Therefore, since Q(n) is never zero for all $n \in \mathbb{N}$ by hypothesis, \overline{a}_n is uniquely determined.

Existence of a differentially algebraic solution. We will use the uniqueness of the power series solution to prove that every entry of this solution is D-algebraic. This is based on a differential version of the Artin approximation theorem given in [DL84].

Theorem 1. [DL84, Theorem 2.1] Let S be a system of differential polynomials in the differential variables y_1, \ldots, y_k with coefficients in a field K. Let $(y_1(x), \ldots, y_k(x)) \in K[\![x]\!]^k$ be a solution of S and $m \in \mathbb{N}$, then there exists another solution $(\bar{y}_1(x), \ldots, \bar{y}_k(x)) \in K[\![x]\!]^k$ such that all $\bar{y}_i(x)$'s are D-algebraic and $(y_1(x), \ldots, y_k(x)) \equiv (\bar{y}_1(x), \ldots, \bar{y}_k(x)) \pmod{x^m}$.

This theorem states that any solution (not necessarily D-algebraic) of a system of a differential equations can be approximated by a D-algebraic solution that coincides on the first m coefficients. Since we showed uniqueness of the power series solutions, we can state our first main result.

Theorem 2. Let $\mathcal{A} = (d, \mu)$ be a holonomic tree automaton over an alphabet Σ . The generating function $f_{\mathcal{A}}(x)$ is differentially algebraic.

Proof. From Proposition 2, we know that f(x) and the vectors of power series $p_{g,i}(x)$ for k > 1, $g \in \Sigma_k$, and $i \in \{1, \ldots, k\}$, are a solution of the system of differential equations 8 and 9. From Theorem 1, we know that there exists another power series solution $\overline{f}(x)$, $\overline{p}_{g,i}(x)$ for k > 1, $g \in \Sigma_k$, and $i \in \{1, \ldots, k\}$ such that all the power series are D-algebraic and they coincide in particular with the first coefficients of f(x), i.e. $\overline{a}_0 = a_0$. From Proposition 3, we deduce that $\overline{f}(x) = f(x)$ and in particular $f_{\mathcal{A}}(x) = \overline{f}_1(x)$. Since $\overline{f}_1(x)$ is D-algebraic the proof is complete.

3 Equivalence of Generating Functions and Formal Tree Series

We are interested in deciding the following two problems:

- (Equivalence of Formal Tree Series) Given two automata $\mathcal{A}_1, \mathcal{A}_2$ over the same alphabet Σ , does $\mathcal{A}_1[\![t]\!] = \mathcal{A}_2[\![t]\!]$ for all $t \in T_{\Sigma}$?
- (Equivalence of Generating Functions) Given two automata $\mathcal{A}_1, \mathcal{A}_2$, does $f_{\mathcal{A}_1}(x) = f_{\mathcal{A}_2}(x)$?

As previously mentioned, the class of holonomic tree automata is closed under subtraction, both for formal tree series and generating functions (see Appendix A for more details). Therefore, we can reduce the respective equivalence problems to the corresponding zeroness problems:

- (Zeroness of Formal Tree Series) Given an automaton \mathcal{A} , does $\mathcal{A}[\![t]\!] = 0$ for all $t \in T_{\Sigma}$?
- (Zeroness of Generating Functions) Given an automaton \mathcal{A} , does $f_{\mathcal{A}}(x) = 0$?

We start by focusing on the zeroness of generating functions. Given a holonomic tree automaton $\mathcal{A} = (d, \mu)$, we use the defining differential equations of $f_{\mathcal{A}}$ given in Proposition 2 to decide whether $f_{\mathcal{A}}(x) = 0$. Since the system of differential equations has a unique power series solution by Proposition 3, the zeroness problem can be reduced to checking whether the system in Proposition 2 remains satisfiable after adding the equation $f_1 = 0$.

Satisfiability of a system of differential equations. To check the satisfiability of a system of differential algebraic equations, we rely on a version of the Hilbert Nullstelensatz theorem for differential ideals [Lan52]. First, we provide some background on differential algebra.

Definition 2. A differential ring (R, ') is a commutative ring with a derivation ': $R \to R$, that is, a map such that, for all $a, b \in R$, (a + b)' = a' + b' and (ab)' = a'b + ab'. A differential field is a differential ring that is a field. For i > 0, $a^{(i)}$ denotes the *i*-th order derivative of $a \in R$.

For a variable y, write $y^{(\infty)}$ for (y, y', y'', \ldots) and for the vector $\mathbf{y} = (y_1, \ldots, y_d)$ of variables write $\mathbf{y}^{(\infty)}$ for $(y_1^{\infty}, \ldots, y_d^{\infty})$. Likewise define $y^{(\leq h)} := (y, y', \ldots, y^{(h)})$ and $\mathbf{y}^{(\leq h)} := (y_1^{(\leq h)}, \ldots, y_d^{(\leq h)})$.

Definition 3 (Differential polynomials). Let R be a differential ring and $\mathbf{y} = (y_1, \ldots, y_d)$ a vector of indeterminates. A differential polynomial is a polynomial with coefficients in R and variables drawn from $\mathbf{y}^{(\infty)}$. We denote such a polynomial by $P(y_1, \ldots, y_d)$. The collection of all such differential polynomials forms a ring $R[\mathbf{y}^{(\infty)}]$. The derivation map extends from R to $R[\mathbf{y}^{(\infty)}]$ by defining $(y_i^{(j)})' = y_i^{(j+1)}$ and lifting $(\cdot)'$ to all of $R[\mathbf{y}^{(\infty)}]$ by linearity and the product rule. Thus $R[\mathbf{y}^{(\infty)}]$ is a differential ring. The order of a differential polynomial is the highest order derivative occurring among its variables.

Definition 4 (Differential ideals). Let $R := K[\mathbf{y}^{(\infty)}]$ be a ring of differential polynomials over a differential field K. An ideal $I \subset R$ is called a *differential ideal* if $a' \in I$ for every $a \in I$. Given a differential polynomial P, write $P^{(\infty)}$ for the set $\{P, P', P'', \ldots\}$ of all derivatives of P. One can verify that, for every $P_1, \ldots, P_s \in R$, the ideal $\langle P_1^{(\infty)}, \ldots, P_s^{(\infty)} \rangle$ is a differential ideal, which we denote $\langle P_1, \ldots, P_s \rangle^{(\infty)}$.

In our setting, we are mostly interested in the case that K is $\mathbb{Q}(x)$ or $\overline{\mathbb{Q}}(x)$ and the derivation is the usual derivative of rational functions. Here $\overline{\mathbb{Q}}$ denotes the algebraic closure of \mathbb{Q} . We can now recall a version of Hilbert Nullstelensatz theorem for differential ideals. **Theorem 3.** [Lan52] Let P_1, \ldots, P_n be differential polynomials in $\mathbb{Q}(x)[\mathbf{y}^{(\infty)}]$, then there exists a tuple of power series y(x) with coefficients in $\overline{\mathbb{Q}}$ such that $P_1(y(x)) = \cdots = P_n(y(x)) = 0$ if and only if $1 \notin \langle P_1, \ldots, P_n \rangle^{(\infty)}$.

Notice here that the coefficients of the power series in y(x) lie in $\overline{\mathbb{Q}}$. We will show that this is not a problem as Proposition 3 on the uniqueness of solutions does not rely on whether the coefficients are in \mathbb{Q} or $\overline{\mathbb{Q}}$, as long as the initial values are the same as the ones of the power series $\boldsymbol{f}, \, \boldsymbol{p}_{q,i}, \, \ldots$ of the automaton.

Ådditionally, instead of checking satisfiability over an ideal generated by infinitely many polynomials there is a bound B for which one can check satisfiability only over the ideal $(P_1, \ldots, P_n)^{(\leq B)} \coloneqq$ $\langle P_1^{(\leq B)}, \ldots, P_n^{(\leq B)} \rangle$. This bound was given explicitly in [OPV22].

Theorem 4. [OPV22, Theorem 1] Let $I = \langle P_1, \ldots, P_n \rangle^{(\infty)}$ be a differential ideal in $\mathbb{Q}(x)[\mathbf{y}^{(\infty)}]$, then (ac) $(\langle R \rangle$

$$\langle P_1, \ldots, P_n \rangle^{(\infty)} \cap \mathbb{Q}(x) = \langle 0 \rangle$$
 if and only if $\langle P_1, \ldots, P_n \rangle^{(\leq B)} \cap \mathbb{Q}(x) = \langle 0 \rangle$

- $B = \max(D^{(\bar{m}+1)2^{m+1}}, m+1)$
- $D = \max_{1 \le i \le n} \deg(P_i),$
- m, \bar{m} are respectively the dimension and the codimension of $\langle P_1, \ldots, P_n \rangle$ in any suitable polynomial ring.

The bound described in this theorem can be computed from the automaton given as input, which leads to the following decision procedure.

Theorem 5 (Zeroness of generating functions). Let $\mathcal{A} = (d, \mu)$ be a holonomic tree automaton over alphabet Σ . Assume that the weight function μ has the form described in Equation (7) and let s be the maximum of r and the largest degree of the polynomials Q and $Q_{q,i}$ for all k > 0, all $g \in \Sigma_k$, all $i \in \{1, ..., k\}$. Finally, let $M := d(s+2)(1+\sum_{k>1} |\Sigma_k|k) + 1$, and $D := \max\{k \in \mathbb{N} \mid \Sigma_k \neq \emptyset\}$, and $B := D^{M2^M}$. Then we have:

$$f_{\mathcal{A}}(x) = 0$$
 if and only if $f_{\mathcal{A}}(x) \equiv 0 \pmod{x^{B+1}}$

Proof. By assumption, the weight function μ has the form described in Equation (7). Let us rewrite the system of differential equations (8) and (9) as follows:

$$\begin{cases} P_{1}(\boldsymbol{f}, \{\boldsymbol{p}_{g,1}, \dots, \boldsymbol{p}_{g,k}\}_{k>0, g \in \Sigma_{k}}) &= 0 \\ \vdots \\ P_{N}(\boldsymbol{f}, \{\boldsymbol{p}_{g,1}, \dots, \boldsymbol{p}_{g,k}\}_{k>0, g \in \Sigma_{k}}) &= 0 \end{cases}$$
(11)

where $N = d \cdot (1 + \sum_{k>0} |\Sigma_k| k)$. This system is satisfied by the vector of power series $\boldsymbol{f} := \sum_{n=0}^{\infty} \boldsymbol{a}_n x^n$ associated to \mathcal{A} , where $\boldsymbol{a}_n := \sum_{t:||t||=n} \tilde{\mu}(t)$. In particular, $\boldsymbol{a}_0 = \sum_{g \in \Sigma_0} \mu(g)$. We define the polynomials V_1, \ldots, V_d by $V_i := f_i - (\boldsymbol{a}_0)_i - x h_i$ where $\boldsymbol{h} = \begin{bmatrix} h_1 & \ldots & h_d \end{bmatrix}$ are fresh variables. We now define the following ideal $I \subseteq \mathbb{Q}(x)[\boldsymbol{f}^{(\leq u)}, \boldsymbol{h}, \{\boldsymbol{p}_{g,1}^{(\leq s)}, \ldots, \boldsymbol{p}_{g,k}^{(\leq s)}\}_{k>0, g \in \Sigma_k}]$:

 $I = \langle P_1, \ldots, P_n, f_1, V_1, \ldots, V_d \rangle$

where $s = \max(u, r)$ (see Equation (7)). By including f_1 as a generator of I we ensure that the power series solution $\overline{f}_1, \overline{f}_2, \ldots$ obtained by applying Theorem 3 satisfies $\overline{f}_1(x) = 0$. Similarly, by including V_1, \ldots, V_d as generators of I, we ensure that the solution has the same initial values as f, that is: $\overline{f}_1(0) = (a_0)_1, \ldots, \overline{f}_d(0) = (a_0)_d$.

The number of variables that appear in the generators of I is $d(u+2)+d\sum_{k\geq 1} |\Sigma_k| k(s+1)$, which is bounded by M. Therefore the dimension and the codimension of I (m and \bar{m} in Theorem 4) are bounded by M. Moreover, from Equation (8), we deduce that the maximal degree of the polynomials P_1, \ldots, P_N are bounded by the largest arity in Σ , that is $\max\{k \in \mathbb{N} \mid \Sigma_k \neq \emptyset\} = D$.

We can therefore apply Theorem 4 and obtain that:

$$I^{(\infty)} \cap \mathbb{Q}(x) = \langle 0 \rangle$$
 if and only if $I^{(\leq B)} \cap \mathbb{Q}(x) = \langle 0 \rangle$.

Observe that $I^{(\infty)} \cap \mathbb{Q}(x) = \langle 0 \rangle$ is equivalent to $1 \notin I^{(\infty)}$. Applying Theorem 3, we deduce that $I^{(\leq B)} \cap \mathbb{Q}(x) = \langle 0 \rangle$ if and only if there exist power series $\overline{f}(x), \overline{h}(x), \{\overline{p}_{g,1}(x), \dots, \overline{p}_{g,k}(x)\}_{k>0,g\in\Sigma_k}$ with coefficients in $\overline{\mathbb{Q}}$ that are roots of the differential polynomials $P_1, \dots, P_N, f_1, V_1, \dots, V_d$.

Note that by definition of V_1, \ldots, V_d , we know that if $\overline{f}(x), \overline{h}(x), \{\overline{p}_{g,1}(x), \ldots, \overline{p}_{g,k}(x)\}_{k>0,g\in\Sigma_k}$ is a solution of V_1, \ldots, V_n then $\overline{f}_i(x) = (a_0)_i + x \overline{h}_i(x)$ for all $i \in \{1, \ldots, d\}$. In other words, $\overline{f}(0) = a_0$. However, as $\overline{f}(x), \{\overline{p}_{g,1}(x), \ldots, \overline{p}_{g,k}(x)\}_{k>0,g\in\Sigma_k}$ are also solutions of P_1, \ldots, P_N , we deduce from Proposition 3 that $\overline{f}(x) = f(x)$. As $\overline{f}(x)$ is also a solution of the polynomial f_1 , we obtain that $f_1(x) = f_A(x) = 0$. On the other hand, we know from Proposition 2 that $f(x), \{p_{g,1}(x), \ldots, p_{g,k}(x)\}_{k>0,g\in\Sigma_k}$ is a solution of P_1, \ldots, P_N and we can easily build $h(x) = f(x) - a_0$ such that f(x), h(x) is a solution of the differential polynomials V_1, \ldots, V_d . Thus, if $f_A(x) = 0$ then $f(x), h(x), \{p_{g,1}(x), \ldots, p_{g,k}(x)\}_{k>0,g\in\Sigma_k}$ is a solution of $P_1, \ldots, P_N, f_1, V_1, \ldots, V_d$. This allows us to obtain the following statement:

$$f_{\mathcal{A}}(x) = 0$$
 if and only if $I^{(\infty)} \cap \mathbb{Q}(x) = \langle 0 \rangle$

We conclude the proof by showing that $I^{(\leq B)} \cap \mathbb{Q}(x) = \langle 0 \rangle$ is equivalent to $f_{\mathcal{A}}(x) \equiv 0 \pmod{x^{B+1}}$.

On the one hand, assume that $I^{(\leq B)} \cap \mathbb{Q}(x) = \langle 0 \rangle$. We have already showed that this implies $f_{\mathcal{A}}(x) = 0$, which clearly implies that $f_{\mathcal{A}}(x) \equiv 0 \pmod{x^{B+1}}$.

On the other hand, assume that $f_{\mathcal{A}}(x) \equiv 0 \pmod{x^{B+1}}$. Towards a contradiction, assume that $f_{\mathcal{A}}(x) \neq 0$. Defining $\mathbf{h}(x) = \frac{\mathbf{f}(x) - \mathbf{a}_0}{x}$, we deduce that $\mathbf{f}(x), \mathbf{h}(x), \{\mathbf{p}_{g,1}(x), \dots, \mathbf{p}_{g,k}(x)\}_{k>0,g\in\Sigma_k}$ is not a solution of $P_1, \dots, P_N, f_1, V_1, \dots, V_d$ and so $1 \in I^{(\leq B)}$. However, by construction and by Proposition 2, $\mathbf{f}(x), \mathbf{h}(x), \{\mathbf{p}_{g,1}(x), \dots, \mathbf{p}_{g,k}(x)\}_{k>0,g\in\Sigma_k}$ is a solution of the differential polynomials $P_1, \dots, P_N, V_1, \dots, V_d$. Moreover, $I^{(\leq B)} = \langle f_1 \rangle^{(\leq B)} + \langle P_1, \dots, P_N, V_1, \dots, V_d \rangle^{(\leq B)}$. Therefore, as $1 \in \langle f_1 \rangle^{(\leq B)} + \langle P_1, \dots, P_N, V_1, \dots, V_d \rangle^{(\leq B)}$, we can write:

$$1 = \sum_{0 \le i \le B} R_i \cdot f_1^{(i)} + \sum_{0 \le i \le B} \sum_{1 \le j \le N} T_i \cdot P_j^{(i)} + \sum_{0 \le i \le B} \sum_{1 \le j \le d} S_i \cdot V_j^{(i)}$$
(12)

By substituting $(\boldsymbol{f}(0), \boldsymbol{h}(0), \{\boldsymbol{p}_{g,1}(0), \dots, \boldsymbol{p}_{g,k}(0)\}_{k>0, g\in\Sigma_k})$ in (12), we get $1 = \sum_{i=1}^B \overline{R}_i \cdot f_1^{(i)}(0)$ where \overline{R}_i is R_i after the substitution at $(\boldsymbol{f}(0), \boldsymbol{h}(0), \{\boldsymbol{p}_{g,1}(0), \dots, \boldsymbol{p}_{g,k}(0)\}_{k>0, g\in\Sigma_k})$. Thus there exists $1 \leq i \leq B$ such that $f_1^{(i)}(0) \neq 0$ and so $f_{\mathcal{A}}^{(i)}(0) \neq 0$. This is in contradiction with our hypothesis $f_{\mathcal{A}}(x) \equiv 0 \pmod{x^{B+1}}$. Therefore, our contradiction assumption does not hold, meaning that $f_{\mathcal{A}}(x) = 0$.

With Theorem 5, we obtain a decision procedure for the zeroness of automata generating functions, consisting of checking whether the B initial values of the generating function are 0. The decidability of zeroness of generating functions also gives the dedidability of zeroness of formal tree series, as shown below. **Proposition 4** (Zeroness of formal tree series). The zeroness problem of formal tree series is decidable.

Proof. Let \mathcal{A} be a holonomic tree automaton over Σ . Since the formal tree series recognisable by holonomic tree automata are closed under product (Proposition 11 in Appendix A), we deduce that there exists a holonomic tree automaton \mathcal{A}' over Σ such that $\mathcal{A}'[t] = \mathcal{A}[t]^2 \geq 0$ for all $t \in T_{\Sigma}$. Since $f_{\mathcal{A}'}(x) = \sum_{n=0}^{\infty} \left(\sum_{t:||t||=n} \mathcal{A}'[t] \right) x^n$ and $\mathcal{A}'[t]$ is always nonnegative, we deduce that

 $f_{\mathcal{A}'}(x) = 0$ if and only if $\forall t \in T_{\Sigma}, \mathcal{A}'[t] = 0$ if and only if $\forall t \in T_{\Sigma}, \mathcal{A}[t] = 0$

As the zeroness problem of generating series is decidable by Theorem 5, the proof is complete. \Box

Corollary 1 (Equivalence). The equivalence problem of formal tree series and the equivalence problem of generating functions are decidable.

Proof. Direct from Propositions 2 and 4 and theorem 5.

4 From Differential Algebraic Functions to Automata

We have shown in Section 2 that the generating functions of holonomic tree automata are differentially algebraic. Conversely, we are interested in characterising which differential algebraic power series arise as the generating function of an automaton.

The main result of this section focuses on the power series that are solutions of first-order rational systems of differential equations, as defined below.

Definition 5. A Rational Dynamical System is a system of differential equations

$$y'_1 = Q_1(y_1, \dots, y_k) \quad \dots \quad y'_k = Q_k(y_1, \dots, y_k),$$

where Q_1, \ldots, Q_k are rational functions.

We say that a power series y(x) is solution of a rational dynamical system when there exists a vector $(y_1(x), \ldots, y_k(x))$ of power series that satisfy a rational dynamical system $y'_1 = Q_1(y_1, \ldots, y_k), \ldots, y'_k = Q_k(y_1, \ldots, y_k)$ such that $y_1(x) = y(x)$. In addition, we say that the power series y(x) is *Rationally Dynamically Algebraic* (RDA) if Q_1, \ldots, Q_k are defined at $(y_1(0), \ldots, y_k(0))$.

The class of power series solution of an RDS is very large as we prove that it includes all differentially algebraic power series.

Proposition 5. All differentially algebraic power series arise as solutions of rational dynamical systems.

Proof. Let y(x) be a differentially algebraic power series. Let $F \in \mathbb{Q}[y^{(\infty)}]$ be a minimal-order differential polynomial such that F(y(x)) = 0. Let *n* be the order of *F*. Observe that $F' = y^{(n+1)}S + R$ where $S, R \in \mathbb{Q}[y^{(\leq n)}]$. Moreover we can assume that $S(y(x), y'(x), \ldots, y^{(n)}(x)) \neq 0$ as otherwise the resultant of *F* and *S* would give a differential polynomial of order strictly smaller than *n* with root y(x). Let us introduce the following system of differential equations

We directly have that $(y(x), y'(x), \dots, y^{(n)}(x))$ is solution of the system (13) which allows us to conclude.

As previously mentioned, the main result of this section shows that every RDA power series is the generating function of a holonomic tree automaton. However, as the transformation from rational dynamical systems to automata is quite complex, we start by exhibiting a transformation for D-finite power series, followed by one for CDA power series, before finally treating the whole class of RDA power series.

Proposition 6 (D-finite power series). For all D-finite power series f(x), there exists a holonomic tree automaton \mathcal{A} such that $f_{\mathcal{A}}(x) = f(x)$.

Proof. By definition, if $f(x) = \sum_{n=0}^{\infty} a_n x^n$ then its series $(a_n)_{n=0}^{\infty}$ of coefficients satisfies a polynomial recurrence

$$Q_0(n)a_n + Q_1(n)a_{n-1} + \ldots + Q_k(n)a_{n-k} = 0$$
 $(n \ge k)$

such that for all $n \geq k$, $Q_0(n) \neq 0$. We build the automaton $\mathcal{A} = (d, \mu)$ over an alphabet $\Sigma = \{\sigma_0, \sigma_1\}$ such that for all $n \in \mathbb{N}$,

$$\sum_{\substack{t \in T_{\Sigma} \\ \|t\| = n}} \widetilde{\mu}(t) = \begin{bmatrix} a_n & \dots & a_{n+k-1} \end{bmatrix}$$

To do so, we consider d = k with the following weight function:

$$\mu(\sigma_0) = \begin{bmatrix} a_0 & \dots & a_{k-1} \end{bmatrix} \quad \text{and} \quad \mu(\sigma_1)(x_0, x_1) = \begin{bmatrix} \mathbf{0}_{1 \times d-1} & \frac{Q_k(x_1+k-1)}{Q_0(x_1+k-1)} \\ & \frac{Q_{k-1}(x_1+k-1)}{Q_0(x_1+k-1)} \end{bmatrix}$$
$$I_{d-1} \quad \vdots \\ & \frac{Q_1(x_1+k-1)}{Q_0(x_1+k-1)} \end{bmatrix}$$

Let us show the desired property by induction on n. The base case (n = 0) being trivial, we focus on the induction step n > 0. In such a case, denoting $\mathbf{a}_n = \sum_{\substack{t \in T_{\Sigma} \\ \|t\| = n}} \widetilde{\mu}(t)$ for all $n \in \mathbb{N}$, we obtain from Equation (4) that $\mathbf{a}_n = \mathbf{a}_{n-1}\mu(\sigma)(n+1,n)$. By the induction hypothesis, we deduce that $\mathbf{a}_{n-1} = \begin{bmatrix} a_{n-1} & \dots & a_{n+k-2} \end{bmatrix}$ and so:

$$\begin{aligned} \boldsymbol{a}_{n} &= \begin{bmatrix} a_{n-1} & \dots & a_{n+k-2} \end{bmatrix} \mu(\sigma)(n+1,n) \\ &= \begin{bmatrix} a_{n} & \dots & a_{n+k-2} & \frac{1}{Q_{0}(n+k-1)} \sum_{i=0}^{k-1} Q_{k-i}(n+k-1)a_{n-1+i} \end{bmatrix} \\ &= \begin{bmatrix} a_{n} & \dots & a_{n+k-2} & a_{n+k-1} \end{bmatrix} \end{aligned}$$

For D-finite power series, the construction of the automaton consists of storing consecutive values of the recurrence in states of our automaton, i.e. $\sum_{\substack{t \in T_{\Sigma} \\ \|t\| = n}} \tilde{\mu}(t) = \begin{bmatrix} a_n & \dots & a_{n+k-1} \end{bmatrix}$. For CDA power series, i.e., power series defined by a rational dynamical system in y_1, \dots, y_k with polynomials instead of rational functions, the intuition is similar. The values of the states of the automaton contain the current coefficients of the solution $y_1(x), \dots, y_k(x)$. In other words, we have $\sum_{\substack{t \in T_{\Sigma} \\ \|t\| = n}} \tilde{\mu}(t) = \begin{bmatrix} a_{n,1} & \dots & a_{n,k} \end{bmatrix}$ where $a_{n,i}$ is the n^{th} coefficient of $y_i(x)$. This similarity is explained by the fact that the rational dynamical system for which a D-finite power series f(x) is solution can intuitively be obtained by introducing new variables f_1, \dots, f_{k-1} for each derivative $f^{(1)}, \dots, f^{(k-1)}$ (and possibly additional variables). The current values of each f_1, \dots, f_{k-1} will thus correspond to coefficients $a_{n+1}, \dots, a_{n+k-1}$ of the recurrence relation.

Proposition 7 (CDA power series). For all CDA power series f(x), there exists a holonomic tree automaton \mathcal{A} such that $f_{\mathcal{A}}(x) = f(x)$.

Proof. By definition, there is a system of differential equations of the form

$$y'_1 = P_1(y_1, \dots, y_k) \quad \dots \quad y'_k = P_k(y_1, \dots, y_k)$$

and $(y_1(x), \ldots, y_k(x))$ power series solution of this system such that $y_1(x) = f(x)$ and for all $i \in \{1, \ldots, k\}, P_i \in \mathbb{Q}[y_1, \ldots, y_k]$.

Let us denote by r the maximal degree of all polynomials P_1, \ldots, P_k . As such, each P_j has the form:

$$P_j(y_1,\ldots,y_k) = \sum_{\ell=0}^{r} \sum_{i_1,\ldots,i_\ell \in \{1,\ldots,k\}} \alpha_{(i_1,\ldots,i_\ell),j} y_{i_1}\ldots y_{i_\ell}$$

where the all coefficients $\alpha_{(i_1,\ldots,i_\ell),j}$ are in \mathbb{Q} . Note that the given system of equations can be transformed into:

$$\Theta y_1 = x P_1(y_1, \dots, y_k) \quad \dots \quad \Theta y_k = x P_k(y_1, \dots, y_k)$$

Since $(y_1(x), \ldots, y_k(x))$ are power series, if we denote $y_i(x) = \sum_{n=0}^{\infty} a_{n,i} x^n$ for all $i \in \{1, \ldots, k\}$, then we deduce that the coefficients of the power series satisfy the following recurrence relations: For all $j \in \{1, \ldots, k\}$, for all $n \ge 1$,

$$n \, a_{n,j} = \sum_{\ell=0}^{\prime} \sum_{\substack{(i_1,\dots,i_\ell) \in \{1,\dots,k\}^*}} \sum_{\substack{n_1+\dots+n_\ell = n-1\\n_1,\dots,n_\ell \in \mathbb{N}}} \alpha_{(i_1,\dots,i_\ell),j} \, a_{n_1,i_1}\dots a_{n_\ell,i_\ell}$$

Let us build the automaton \mathcal{A} over the alphabet $\Sigma = \{\varepsilon, \sigma, g_1, \ldots, g_r\}$ with ε being a nullary symbol, σ being a unary symbol and for all $i \in \{1, \ldots, r\}$, g_i is a *i*-ary symbol. Intuitively, each g_i with $i \geq 1$ will represent all the monomials of degree i > 0. The symbol σ will represent the constants. We now build the weight function μ of the automaton \mathcal{A} such that for all $n \in \mathbb{N}$,

$$\sum_{\substack{t \in T_{\Sigma} \\ \|t\|=n}} \widetilde{\mu}(t) = \begin{bmatrix} a_{n,1} & \dots & a_{n,k} & 1 \end{bmatrix}$$

To do so, our automaton \mathcal{A} must have dimension k+1 and weight function defined by:

- $\mu(\varepsilon) = \begin{bmatrix} a_{0,1} & \dots & a_{0,k} & 1 \end{bmatrix}$
- for all $\forall \ell \in \{1, ..., r\}$, for all $i_1, ..., i_\ell, j \in \{1, ..., k+1\}$,

$$\mu(g_{\ell})_{(i_1,\ldots,i_{\ell}),j}(x_0,\ldots,x_{\ell}) = \begin{cases} \frac{\alpha_{(i_1,\ldots,i_{\ell}),j}}{x_0} & \text{if } \forall i_1,\ldots,i_{\ell}, j \in \{1,\ldots,k\}\\ 0 & \text{otherwise} \end{cases}$$

• for all $i, j \in \{1, \dots, k+1\},\$

$$\mu(\sigma)_{i,j}(x_0, x_1) = \begin{cases} \frac{\alpha_{(j,j)}}{x_0} & \text{if } i = k+1, j \le k\\ 1 & \text{if } i = j = k+1\\ 0 & \text{otherwise} \end{cases}$$

We prove the desired property by induction on n. The base case n = 0 being trivial, we focus on the induction step (n > 0). Denoting $a_n = \sum_{t \in T_{\Sigma}} \tilde{\mu}(t)$ for all $n \in \mathbb{N}$, we know from Equation (4) and from our induction hypothesis that for all $n \ge 1$,

$$\begin{aligned} \boldsymbol{a}_{n} &= \sum_{\substack{\ell > 0 \\ g \in \Sigma_{\ell}}} \sum_{\substack{n_{1}, \dots, n_{\ell} \in \mathbb{N} \\ n_{1} + \dots + n_{\ell} = n - 1}} (\boldsymbol{a}_{n_{1}} \otimes \dots \otimes \boldsymbol{a}_{n_{\ell}}) \cdot \mu(g)(n, n_{1}, \dots, n_{\ell}) \\ &= \sum_{\substack{\ell > 0 \\ g \in \Sigma_{\ell}}} \sum_{\substack{n_{1}, \dots, n_{\ell} \in \mathbb{N} \\ n_{1} + \dots + n_{\ell} = n - 1}} ([a_{n_{1}, 1} \dots a_{n_{1}, k} \quad 1] \otimes \dots \otimes [a_{n_{\ell}, 1} \dots a_{n_{\ell}, k} \quad 1]) \cdot \mu(g)(n, n_{1}, \dots, n_{\ell}) \end{aligned}$$

Therefore, by definition of μ , we have for $j \in \{1, \ldots, k\}$,

$$\begin{aligned} \boldsymbol{a}_{n,j} &= \boldsymbol{a}_{n-1,k+1} \frac{\alpha_{(),j}}{n} + \sum_{\ell=1}^{r} \sum_{\substack{n_1,\dots,n_\ell \in \mathbb{N} \\ n_1 + \dots + n_\ell = n-1}} \sum_{\substack{(i_1,\dots,i_\ell) \in \{1,\dots,k\}^*}} a_{n_1,i_1} \dots a_{n_\ell,i_\ell} \frac{\alpha_{(i_1,\dots,i_\ell),j}}{n} \\ &= \sum_{\ell=0}^{r} \sum_{\substack{n_1,\dots,n_\ell \in \mathbb{N} \\ n_1 + \dots + n_\ell = n-1}} \sum_{\substack{(i_1,\dots,i_\ell) \in \{1,\dots,k\}^*}} a_{n_1,i_1} \dots a_{n_\ell,i_\ell} \frac{\alpha_{(i_1,\dots,i_\ell),j}}{n} \\ &= a_{n,j} \end{aligned}$$

Furthermore, we also have that:

$$a_{n,k+1} = a_{n-1,k+1}\mu(\sigma)_{k+1,k+1}(n,n-1) = 1$$

This allows us to conclude the proof that for all $n \in \mathbb{N}$, $a_n = \begin{bmatrix} a_{n,1} & \dots & a_{n,k} & 1 \end{bmatrix}$, and so $f_{\mathcal{A}}(x) = y_1(x) = f(x)$.

Example 4. As an example, take the function $y_1(x) = \frac{1}{\cos(x)}$ and $y_2(x) = \tan(x)$. They satisfy the system of differential equations:

$$\begin{cases} y_1' = y_1 y_2 \\ y_2' = 1 + y_2^2 \end{cases}$$

It was shown that in [Sta80, Example 2.5] that $y_1(x)$ is not D-finite. Let us build the corresponding holonomic tree automaton based on the proof of Proposition 7. The maximal degree r of the polynomials is 2 but there is no monomial of degree 1. Therefore, we consider the alphabet $\Sigma = \{\varepsilon, \sigma, g_2\}$ (no need for the symbol g_1 representing the monomial of degree 1). Since we only have two variables, y_1, y_2 , the dimension of our automaton is 3. As $y_1(0) = 1$ and $y_2(0) = 0$, the weight function is defined as follows:

$$\mu(\varepsilon) = \begin{bmatrix} 1 & 0 & 1 \end{bmatrix} \qquad \mu(\sigma)(x_0, x_1) = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \frac{1}{x_0} & 1 \end{bmatrix} \qquad \mu(g_2)(x_0, x_1, x_2) = \begin{bmatrix} 0 & 0 & 0 \\ \frac{1}{x_0} & 0 & 0 \\ 0_{1\times 2} & \mathbf{0}_{1\times 2} & \mathbf{0}_{1\times 2} \\ 0 & \frac{1}{x_0} & 1 \\ \mathbf{0}_{1\times 4} & \mathbf{0}_{1\times 4} & \mathbf{0}_{1\times 4} \end{bmatrix}$$

The construction for RDA power series will take a similar approach, for instance by splitting the rational functions into their monomials. However, to handle the presence of polynomials in both denominators and numerators, we will first transform the system to reduce its maximal degree to be at most 2. This is done in particular by introducing new variables whose values will also appear in the states of the automaton, thus increasing the dimension of the automaton. Another change in the construction is that, instead of keeping the current values, the automaton keeps the forward-shifted values of each variable and only keep the current value of the target function. In other words, we seek to maintain the invariant

$$\sum_{\substack{t \in T_{\Sigma} \\ \|t\| = n}} \widetilde{\mu}(t) = \begin{bmatrix} a_{n,1} & a_{n+1,1} & a_{n+1,2} & \dots & a_{n+1,k} & b_{n+1,1} & \dots & b_{n+1,s} & 1 \end{bmatrix}$$

where the $a_{n,i}$ is the *n*-th coefficient of the power series $y_i(x)$ and the $b_{n,i}$ are auxiliary variables introduced during the transformation of the system.

Theorem 6 (RDA power series). For all RDA power series f(x), there exists a holonomic tree automaton \mathcal{A} such that $f_{\mathcal{A}}(x) = f(x)$.

Proof. By definition, there exists a rational dynamical system of the form

$$y'_1 = \frac{P_1(y_1, \dots, y_k)}{Q_1(y_1, \dots, y_k)} \quad \dots \quad y'_k = \frac{P_k(y_1, \dots, y_k)}{Q_k(y_1, \dots, y_k)}$$

and $(y_1(x), \ldots, y_k(x))$ power series solution of this system such that $f(x) = y_1(x)$ and for all $i \in \{1, \ldots, k\}, P_i, Q_i \in \mathbb{Q}[y_1, \ldots, y_k]$ and $Q_i(y_1(0), \ldots, y_k(0)) \neq 0$.

Let us denote by r the maximal degree of all polynomials $P_1, Q_1, \ldots, P_k, Q_k$. As such, each P_j and Q_j are of the form:

$$P_{j}(y_{1},\ldots,y_{k}) = \sum_{\ell=0}^{r} \sum_{(i_{1},\ldots,i_{\ell})\in\{1,\ldots,k\}^{*}} \alpha_{(i_{1},\ldots,i_{\ell}),j} y_{i_{1}}\ldots y_{i_{\ell}}$$
$$Q_{j}(y_{1},\ldots,y_{k}) = \sum_{\ell=0}^{r} \sum_{(i_{1},\ldots,i_{\ell})\in\{1,\ldots,k\}^{*}} \beta_{(i_{1},\ldots,i_{\ell}),j} y_{i_{1}}\ldots y_{i_{\ell}}$$

where all the coefficients $\alpha_{(i_1,...,i_\ell),j}$ and $\beta_{(i_1,...,i_\ell),j}$ are in \mathbb{Q} . Note that the given system of equations can be transformed into:

$$Q_1(y_1,\ldots,y_k)\Theta y_1 = xP_1(y_1,\ldots,y_k) \quad \dots \quad Q_k(y_1,\ldots,y_k)\Theta y_k = xP_k(y_1,\ldots,y_k) \tag{14}$$

We introduce several new variables in order to transform the system of equations 14 into a set of equations of degree 2. To start, we introduce variables $z_1, \ldots, z_k, w_1, \ldots, w_k$ and transform 14 to the equivalent system:

$$\begin{cases} z_1 \Theta y_1 = x \, w_1 & \dots & z_k \Theta y_k = x \, w_k \\ z_1 = Q_1(y_1, \dots, y_k) & \dots & z_k = Q_k(y_1, \dots, y_k) \\ w_1 = P_1(y_1, \dots, y_k) & \dots & w_k = P_k(y_1, \dots, y_k) \end{cases}$$
(15)

Then, for each monomial $y_{i_1} \ldots y_{i_\ell}$ in $Q_j(y_1, \ldots, y_k)$ and $P_j(y_1, \ldots, y_k)$ with $\ell > 2$, we introduce fresh variables t_1, \ldots, t_ℓ to split the equations $z_j = Q_j(y_1, \ldots, y_k)$ and $w_j = P_j(y_1, \ldots, y_k)$ into an equivalent system of equations of degree at most 2 as follows:

$$t_1 = y_{i_1} t_2$$
 $t_2 = y_{i_2} t_3$... $t_{\ell-1} = y_{\ell-1} t_\ell$ $t_\ell = y_\ell$

By applying this transformation to every monomial in every polynomial $Q_j(y_1, \ldots, y_k)$ and $P_j(y_1, \ldots, y_k)$ and after renaming the newly introduced variables, we deduce that the set of solutions of system (15) in variables y_1, \ldots, y_k equals the set of solutions in y_1, \ldots, y_k of the following system:

$$\begin{cases} z_1 \Theta y_1 = x \, w_1 \quad \dots \quad z_k \Theta y_k = x \, w_k \\ z_1 = R_1(y_1, \dots, y_k, t_1, \dots, t_s) \quad \dots \quad z_k = R_k(y_1, \dots, y_k, t_1, \dots, t_s) \\ w_1 = S_1(y_1, \dots, y_k, t_1, \dots, t_s) \quad \dots \quad w_k = S_k(y_1, \dots, y_k, t_1, \dots, t_s) \\ t_1 = T_1(y_1, \dots, y_k, t_2, \dots, t_s) \quad t_2 = T_2(y_1, \dots, y_k, t_3, \dots, t_s) \quad \dots \quad t_s = T_s(y_1, \dots, y_k) \end{cases}$$
(16)

where $R_1, S_1, \ldots, R_k, S_k$ and T_1, \ldots, T_s are polynomials of degree at most 2. Moreover in T_1, \ldots, T_s , all monomials of degree 2 have one variable from $\{y_1, \ldots, y_k\}$ and one variable from $\{t_1, \ldots, t_s\}$.

Since the power series $(y_1(x), \ldots, y_k(x))$ are solutions of the initial system (14), we also deduce the existence of power series $z_1(x), \ldots, z_k(x), w_1(x), \ldots, w_k(x), t_1(x), \ldots, t_s(x)$ that, together with $(y_1(x), \ldots, y_k(x))$, form a solution of the system system (16). Let us denote by $y_{n,i}$ (resp. $z_{n,i}, w_{n,i}, t_{n,i}), n \in \mathbb{N}$ the coefficients of the power series $y_i(x)$ (resp. $z_i(x), w_i(x), t_i(x)$). From (16) we deduce that these coefficients satisfy the following relations:

• For all $n \ge 1$, for all $j \in \{1, \ldots, k\}$,

$$\sum_{\ell=0}^{n+1} z_{\ell,j}(n+1-\ell)y_{n+1-\ell,j} = w_{n,j}$$

$$\iff z_{0,j}(n+1)y_{n+1,j} + \sum_{\ell=1}^{n} z_{\ell,j}(n+1-\ell)y_{n+1-\ell,j} = w_{n,j}$$

$$\iff y_{n+1,j} = \frac{1}{z_{0,j}(n+1)} \left(w_{n,j} - \sum_{\ell=1}^{n} z_{\ell,j}(n+1-\ell)y_{n+1-\ell,j} \right)$$

$$\iff y_{n+1,j} = \frac{1}{z_{0,j}(n+1)} \left(w_{n,j} - \sum_{\ell=0}^{n-1} z_{\ell+1,j}(n-\ell)y_{n-\ell,j} \right)$$

By the assumption that $Q_i(y_1(0), \ldots, y_k(0)) \neq 0$ for all $i \in \{1, \ldots, k\}$ we deduce from (15) that $z_{0,j} \neq 0$ for all $j \in \{1, \ldots, k\}$. Introducing new variables $\overline{z}_{n,j} = z_{n+1,j}$, $\overline{y}_{n,j} = y_{n+1,j}$ and $\overline{w}_{n,j} = w_{n+1,j}$, for all $j \in \{1, \ldots, k\}$ and $n \geq 1$, we obtain :

$$\overline{y}_{n,j} = \frac{1}{z_{0,j}(n+1)} \left(\overline{w}_{n-1,j} - \sum_{\ell=0}^{n-1} \overline{z}_{\ell,j}(n-\ell) \overline{y}_{n-1-\ell,j} \right)$$
(17)

Notice that the shape of this equation is reminiscent of Equation (4) and they will intuitively correspond to the application of several symbols of the alphabet Σ : the term $\overline{w}_{n-1,j}$ will correspond to the application of a unary symbol whereas the term $\sum_{\ell=0}^{n-1} \overline{z}_{\ell,j}(n-\ell)\overline{y}_{n-1-\ell,j}$ will correspond to the application of a binary symbol.

• Let us denote $\Gamma = \{w_i, y_i, z_i\}_{i=1}^k \cup \{t_1, \dots, t_s\}$. We show by induction on s - j that for all $j \in \{1, \dots, s\}$, for all $n \ge 1$,

$$\overline{t}_{n,j} = A(n) + \sum_{h \in \Gamma} B_h(n, n-1)\overline{h}_{n-1} + \sum_{h,g \in \Gamma} \sum_{\ell=0}^{n-1} C_{h,g}(n, \ell, n-1-\ell) \overline{h}_\ell \overline{g}_{n-1-\ell}$$
(18)

where $A \in \mathbb{Q}_u(x_0), B_h \in \mathbb{Q}_u(x_0, x_1)$ and $C_{h,g} \in \mathbb{Q}_u(x_0, x_1, x_2)$

In the base case (s = j), we know that $t_s = T_s(y_1, \ldots, y_k)$. Moreover, we also know that T_s is of degree at most 2 and all monomials of degree 2 must have a variable from $\{t_1, \ldots, t_s\}$. In other words, T_s is of degree at most 1. Therefore, for all $n \ge 1$,

$$\overline{t}_{n,s} = \alpha + \sum_{j=1}^{k} \beta_j \overline{y}_{n,j}$$
$$= \alpha + \sum_{j=1}^{k} \frac{\beta_j}{z_{0,j}(n+1)} \left(\overline{w}_{n-1,j} - \sum_{\ell=0}^{n-1} \overline{z}_{\ell,j}(n-\ell) \overline{y}_{n-1-\ell,j} \right)$$

$$= \alpha + \sum_{j=1}^{k} \frac{\beta_j \overline{w}_{n-1,j}}{z_{0,j}(n+1)} - \sum_{j=1}^{k} \sum_{\ell=0}^{n-1} \frac{(n-\ell)\beta_j}{z_{0,j}(n+1)} \overline{z}_{\ell,j} \overline{y}_{n-1-\ell,j}$$

Hence the result holds.

In the inductive step (s > j), we know that $t_j = T_j(y_1, \ldots, y_k, t_{j+1}, \ldots, t_s)$ with T_j being of degree at most 2 and that all monomials of degree 2 must have a variable from $\{t_1, \ldots, t_s\}$ and one from $\{y_1, \ldots, y_k\}$. For a monomial of degree 1, i.e. αu , as $u \in \{y_1, \ldots, y_k, t_{j+1}, \ldots, t_s\}$ and by Equation (17) and our inductive hypothesis (18) on each t_{j+1}, \ldots, t_s , we have the equation $(\alpha u)_{n+1} = \alpha \overline{u}_n$, which already has the desired form.

We now focus on the monomials of degree 2. Assume that αuv is a monomial of T_j with $u, v \in \{y_1, \ldots, y_k, t_{j+1}, \ldots, t_s\}$. Then we have for all $n \ge 1$,

$$(\alpha u v)_{n+1} = \alpha \sum_{\ell=0}^{n+1} u_{\ell} v_{n+1-\ell}$$

$$= \alpha u_0 v_{n+1} + \alpha u_{n+1} v_0 + \alpha \sum_{\ell=1}^n u_{\ell} v_{n+1-\ell}$$

$$= \alpha u_0 \overline{v}_n + \alpha \overline{u}_n v_0 + \alpha \sum_{\ell=0}^{n-1} \overline{u}_{\ell} \overline{v}_{n-1-\ell}$$

$$(19)$$

By Equation (17) and by our inductive hypothesis (18) on each t_{j+1}, \ldots, t_s , we deduce that:

$$\overline{u}_n = A(n) + \sum_{h \in \Gamma} B_h(n, n-1)\overline{h}_{n-1} + \sum_{h,g \in \Gamma} \sum_{\ell=0}^{n-1} C_{h,g}(n, \ell, n-1-\ell) \overline{h}_\ell \overline{g}_{n-1-\ell}$$
$$\overline{v}_n = D(n) + \sum_{h \in \Gamma} E_h(n, n-1)\overline{h}_{n-1} + \sum_{h,g \in \Gamma} \sum_{\ell=0}^{n-1} F_{h,g}(n, \ell, n-1-\ell) \overline{h}_\ell \overline{g}_{n-1-\ell}$$

for some $A, D \in \mathbb{Q}_u(x_0)$, $B_h, E_h \in \mathbb{Q}_u(x_0, x_1)$ and $C_{h,g}, F_{h,g} \in \mathbb{Q}_u(x_0, x_1, x_2)$. Therefore, by combining these two equations with Equation (20), we obtain that the monomial $(\alpha u v)_{n+1}$ has the desired form.

• For all $j \in \{1, \ldots, k\}$, for all $n \ge 1$, since $z_j = R_j(y_1, \ldots, y_k, t_1, \ldots, t_s)$ and $w_j = S_j(y_1, \ldots, y_k, t_1, \ldots, t_s)$, where $R_1, S_1, \ldots, R_k, S_k$ are also polynomials of degree at most 2, we can apply the same proof as in the previous case (the $\overline{t}_{n,j}$ s) to show that both $\overline{w}_{n,j}$ and $\overline{z}_{n,j}$ are of the form: n-1

$$A(n) + \sum_{h \in \Gamma} B_h(n, n-1)\overline{h}_{n-1} + \sum_{h,g \in \Gamma} \sum_{\ell=0}^{n-1} C_{h,g}(n, \ell, n-1-\ell) \overline{h}_\ell \overline{g}_{n-1-\ell}$$

with $A \in \mathbb{Q}_u(x_0)$, $B_h \in \mathbb{Q}_u(x_0, x_1)$ and $C_{h,g} \in \mathbb{Q}_u(x_0, x_1, x_2)$.

To summarize, we have proved that for all $n \ge 1$, the coefficients $\overline{y}_{n,i}$, $\overline{z}_{n,i}$, $\overline{w}_{n,i}$ and $\overline{t}_{n,i}$ are all of the form

$$A(n) + \sum_{h \in \Gamma} B_h(n, n-1)\overline{h}_{n-1} + \sum_{h,g \in \Gamma} \sum_{\ell=0}^{n-1} C_{h,g}(n, \ell, n-1-\ell) \overline{h}_\ell \overline{g}_{n-1-\ell}$$

with $\Gamma = \{w_i, y_i, z_i\}_{i=1}^k \cup \{t_1, \dots, t_s\}, A \in \mathbb{Q}_u(x_0), B_h \in \mathbb{Q}_u(x_0, x_1) \text{ and } C_{h,g} \in \mathbb{Q}_u(x_0, x_1, x_2).$

From this point the construction of our automaton is straightforward. First, we will consider an automaton $\mathcal{A} = (d, \mu)$ of dimension d = 2 + 3k + s over the alphabet $\Sigma = \{\varepsilon, \sigma_1, \sigma_2\}$ where ε is nullary, σ_1 is unary and σ_2 is binary. The weight function μ is defined such that for all $n \in \mathbb{N}$:

$$\sum_{\substack{t \in T_{\Sigma} \\ \|t\|=n}} \widetilde{\mu}(t) = \begin{bmatrix} y_{n,1} & \overline{y}_{n,1} & \dots & \overline{y}_{n,k} & \overline{z}_{n,1} & \dots & \overline{z}_{n,k} & \overline{w}_{n,1} & \dots & \overline{w}_{n,k} & \overline{t}_{n,1} & \dots & \overline{t}_{n,s} & 1 \end{bmatrix}$$

To show how we build such weight function, let us write $a_n = \sum_{\substack{t \in T_{\Sigma} \\ \|t\|=n}} \widetilde{\mu}(t)$. Thus, we obtain from Equation (4) that for all $n \ge 1$,

$$a_n = a_{n-1} \mu(\sigma_1)(n, n-1) + \sum_{\ell=0}^{n-1} (a_\ell \otimes a_{n-1-\ell}) \mu(\sigma_2)(n, \ell, n-1-\ell).$$

The matrices $\mu(\sigma_1)$ and $\mu(\sigma_2)$ are defined as follows:

- By definition, $\overline{y}_{n-1,1} = y_{n,1}$. Thus, we can set $\mu(\sigma_1)_{2,1} = 1$ and $\mu(\sigma_1)_{i,1} = 0$ for all $i \neq 2$. Moreover we also set $\mu(\sigma_2)_{(i,j),1} = 0$ for all $i, j \in \{1, \ldots, d\}$.
- Consider $j \in \{2, \ldots, d-1\}$. We know that $(a_n)_j$ is of the form:

$$A(n) + \sum_{h \in \Gamma} B_h(n, n-1)\overline{h}_{n-1} + \sum_{h, g \in \Gamma} \sum_{\ell=0}^{n-1} C_{h,g}(n, \ell, n-1-\ell) \overline{h}_\ell \overline{g}_{n-1-\ell}$$

with $\Gamma = \{w_i, y_i, z_i\}_{i=1}^k \cup \{t_1, \dots, t_s\}, A \in \mathbb{Q}_u(x_0), B_h \in \mathbb{Q}_u(x_0, x_1) \text{ and } C_{h,g} \in \mathbb{Q}_u(x_0, x_1, x_2).$ We now define:

- $-\mu(\sigma_1)_{0,j} := 0$
- $\mu(\sigma_1)_{d,j}(x_0, x_1) := A(x_0)$
- for all $h \in \Sigma$, if *i* is the index corresponding to *h* then $\mu(\sigma_1)_{i,j}(x_0, x_1) := B_h(x_0, x_1)$.
- for all $h, g \in \Gamma$, if i_1 and i_2 are respectively the indices corresponding to h and g then $\mu(\sigma_2)_{(i_1,i_2),j}(x_0, x_1, x_2) := C_{h,g}(x_0, x_1, x_2)$
- $-\mu(\sigma_2)_{(i_1,i_2),j} = 0$ when either $i_1 = 1$ or $i_1 = d$ or $i_2 = 1$ of $i_2 = d$.
- Finally, $\mu(\sigma_1)_{d,d} := 1$ and $\mu(\sigma_1)_{i,d} = 0$ for $i \neq d$.

Provided that

$$\mu(\varepsilon) = \begin{bmatrix} y_{0,1} & \overline{y}_{0,1} & \dots & \overline{y}_{0,k} & \overline{z}_{0,1} & \dots & \overline{z}_{0,k} & \overline{w}_{0,1} & \dots & \overline{w}_{0,k} & \overline{t}_{0,1} & \dots & \overline{t}_{0,s} & 1 \end{bmatrix}$$

an easy proof by induction on n (unfolding the definition of $\mu(\sigma_1)$ and $\mu(\sigma_2)$) allows us to show that for all $n \in \mathbb{N}$,

$$\boldsymbol{a}_n = \begin{bmatrix} y_{n,1} & \overline{y}_{n,1} & \dots & \overline{y}_{n,k} & \overline{z}_{n,1} & \dots & \overline{z}_{n,k} & \overline{w}_{n,1} & \dots & \overline{w}_{n,k} & \overline{t}_{n,1} & \dots & \overline{t}_{n,s} & 1 \end{bmatrix}$$

We can therefore conclude that $f_{\mathcal{A}}(x) = y_1(x)$.

Example 5. Consider the solution $y_1(x)$ of the differential equation $(y'_1)^3 + y_1^3 = 1$ such that $y_1(0) = 0$ and $y'_1(0) = 1$. This equation cannot be put directly into a rational dynamical system. However, by deriving the equation (following Proposition 5), we deduce that $y_1(x)$ also satisfies $3(y'_1)^2y''_1 + 3(y_1)^2y'_1 = 0$ which leads to the following system:

$$\begin{cases} y_1' = y_2 \\ y_2' = -\frac{y_1^2}{y_2} \end{cases}$$

Notice that as $y_2(0) = y'_1(0) \neq 0$, we know from Theorem 6 that there exists an automaton \mathcal{A} such that $f_{\mathcal{A}}(x) = y_1(x)$. To build this automaton, we follow the construction in the proof of Theorem 6. Introducing only the non-trivial variables, we obtain :

$$\begin{cases} \Theta y_1 = x \, y_2 \quad y_2 \Theta y_2 = x \, w_2 \\ w_2 = -y_1^2 \end{cases}$$

In term of relations between coefficients, we thus obtain:

$$\overline{y}_{n,1} = \frac{1}{n+1}\overline{y}_{n-1,2} \qquad \overline{y}_{n,2} = \frac{1}{n+1}\left(\overline{w}_{n-1,2} - \sum_{\ell=0}^{n-1}\overline{y}_{\ell,2}(n-\ell)\overline{y}_{n-1-\ell,2}\right)$$

Moreover, we have

$$\overline{w}_{n,2} = -2y_{0,1}\overline{y}_{n,1} - \sum_{\ell=0}^{n-1} \overline{y}_{\ell,1}\overline{y}_{n-1-\ell,1} = -\sum_{\ell=0}^{n-1} \overline{y}_{\ell,1}\overline{y}_{n-1-\ell,1}$$

Since none of the coefficient has a constant term, we build an automaton \mathcal{A} such that $a_n = \begin{bmatrix} y_{n,1} & \overline{y}_{n,2} & \overline{w}_{n,2} \end{bmatrix}$. Therefore, we take

$$\mu(\varepsilon) = \begin{bmatrix} 0 & 1 & 0 & 0 \end{bmatrix} \qquad \mu(\sigma_1)(x_0, x_1) = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & \frac{1}{x_0 + 1} & 0 & 0 \\ 0 & 0 & \frac{1}{x_0 + 1} & 0 \end{bmatrix}$$

and

$$\mu(\sigma_2)(x_0, x_1, x_2) = \begin{bmatrix} \mathbf{0}_{6\times 1} & \mathbf{0}_{6\times 1} & \mathbf{0}_{6\times 1} & \mathbf{0}_{6\times 1} \\ 0 & 0 & 0 & -1 \\ \mathbf{0}_{3\times 1} & \mathbf{0}_{3\times 1} & \mathbf{0}_{3\times 1} & \mathbf{0}_{3\times 1} \\ 0 & 0 & -\frac{x_2+1}{x_0+1} & 0 \\ \mathbf{0}_{5\times 1} & \mathbf{0}_{5\times 1} & \mathbf{0}_{5\times 1} & \mathbf{0}_{5\times 1} \end{bmatrix}$$

Running the automaton leads to the power series:

$$y_1(x) = x - \frac{2}{4!}x^4 - \frac{20}{7!}x^7 - \frac{3320}{10!}x^{10} - \frac{1598960}{13!}x^{13} - \frac{1757280800}{16!}x^{16} - \dots$$

4.1 Application to counting combinatorial structures

Combinatorial species are a formalism for defining classes of combinatorial structures [FZC94]. A species A is a mapping (technically, a functor) that takes a finite set of labels U and returns a set, denoted A[U], of combinatorial structures labelled by U. For example, if B is the species of plane rooted binary trees then $B[\{1,4,5\}]$ is the set of trees shown below:



Note that all nodes do not have to be labelled. If C is the species of plane rooted binary trees with labels only on external nodes then $C[\{1,2\}]$ contains only two trees:



To define species [FZC94] relies on a collection of constructions: (i) The initial object 1, which represents the empty structure without label. (ii) The variable X, which represents a single labelled node. (iii) The addition A + B, which represents the disjoints union of structures from A and B. (iv) The product $A \cdot B$, which represents all the pairs of elements from A and B over the input set of labels. (v) The sequence sequence(A), which generates the sequences of elements of A. (vi) The set set(A), which generates the sets of elements of A. (vii) The cycle cycle(A), which generates the cycles of elements of A.

The constructions set, sequence and cycle also allow constraints on the cardinality. For example, set($A, card \geq 3$) represents the sets of at least 3 elements of A. A specification of species is then a (set of) equations that use these constructions. In this setting, [FZC94] are interested in computing the exponential generating series of a species A, corresponding to the power series $\sum_{n=0}^{\infty} \frac{\|A[\{1,...,n\}]\|}{n!} x^n$. In other words, the *n*-th coefficient of the power corresponds to the numbers of combinatorial structures in the species A labelled by $\{1,...,n\}$.

For instance, [FZC94] provides the following examples of specifications:

Specifications	Objects	
$A = X \cdot set(A)$	Non-plane trees	
$B = X + B \cdot B$	Plane binary trees with only external labels	
$C = X \cdot sequence(C)$	Plane general trees	
D = set(cycle(X))	Permutations	
E = set(cycle(A))	Functional graph	
$F = set(set(X, card \geq 1))$	Set partitions	
$G = X + X \cdot set(G, card = 3)$	Non-plane ternary trees	
$H = X + set(H, card \ge 2)$	Hierarchies	
$K = set(cycle(X \cdot set(G, card = 2)))$	3-constrained functional graphs	
$L = set(set(set(X, card \geq 1), card \geq 1))$	3-balanced hierarchies	
$M = sequence(set(X, card \geq 1))$	Surjections	

These specifications represents standard objects in the literature. Here, non-plane trees are trees in which the children of a node are unordered, whereas in plane trees siblings are ordered. Functional graphs are directed graphs with every node having outdegree 1 and 3-functional graphs are function graphs where each node has indegree being 0 or 3.

Fortunately, [FZC94] also provides a way to translate every construct of the specification into a system of differential equations on the associated exponential generating functions. For instance, (i) B = set(A) is translated to the equation $y'_B = y_B \cdot y'_A$; (ii)ZB = sequence(A) is translated

System	RDS	Initial values	Objects
$\begin{cases} y_a = x z \\ z' = z y'_a \end{cases}$	$\begin{cases} y'_a = z + \frac{xz^2}{1-xz}\\ z' = \frac{z^2}{1-xz} \end{cases}$	_	Non plane trees
$y_b = x + y_b^2$	$y'_b = rac{1}{1-2y_b}$	$y_b(0) = 0$	Plane binary trees with external labels
$y_c = \frac{x}{1 - y_c}$	$y'_c = \frac{(1-y_c)^3}{(1-y_c)^2 - x}$	$y_c(0) = 0$	Plane general trees
$\begin{cases} y'_d = y_d z' \\ z' = \frac{1}{1-x} \end{cases}$	$y'_d = \frac{y_d}{1-x}$		Permutations
$\begin{cases} y'_e = y_e z'_2 \\ z'_2 = \frac{y'_a}{1 - y_a} \\ y_a = x z \\ z' = z y'_a \end{cases}$	$\begin{cases} y'_e = y_e \frac{z}{1-y_a} + \frac{x z^2}{(1-x z)(1-y_a)} \\ y'_A = z + \frac{x z^2}{1-x z} \\ z' = \frac{z^2}{1-x z} \end{cases}$	$y_a(0) = 0$	Functional graph
$\left\{\begin{array}{l} y'_f = y_f z\\ z' = z\end{array}\right.$	$\begin{cases} y'_f = y_f z \\ z' = z \end{cases}$	_	Set partitions (Bell Number)
$\begin{cases} y_g = x + x z_2 \\ z'_2 = z_1 y'_g \\ z'_1 = y_g y'_g \end{cases}$	$\begin{cases} y'_g = \frac{1+z_2}{1-xz_1} \\ z'_2 = \frac{y_1(1+z_2)}{1-xz_1} \\ z'_1 = y_g \frac{1+z_2}{1-xz_1} \end{cases}$		Non plane ternary trees
$\begin{cases} y_h = x + z_2 \\ z'_2 = z_1 y'_h \\ z'_1 = z_0 y'_h \\ z'_0 = z_0 y'_h \end{cases}$	$\begin{cases} y'_h = 1 + \frac{z_1}{1 - z_1} \\ z'_2 = \frac{z_1}{1 - z_1} \\ z'_1 = z_0 + \frac{z_0 z_1}{1 - z_1} \\ z'_0 = z_0 + \frac{z_0 z_1}{1 - z_1} \end{cases}$	$y_h(0) = 0$ $z_1(0) = 0$	Hierarchies
$\begin{cases} y'_k = y_k z'_1 \\ z'_1 = \frac{z'_2}{1 - z_2} \\ z_2 = x z_3 \\ z'_3 = y_g y'_g \\ y_g = x + x z_4 \\ z'_4 = z_5 y'_g \\ z'_5 = y_g y'_g \end{cases}$	$\begin{cases} y'_k = y_k \frac{z_3}{1-x z_3} + y_k \frac{x y_g(1+z_4)}{(1-x z_5)(1-x z_3)} \\ z'_1 = \frac{z_3}{1-x z_3} + \frac{x y_g(1+z_4)}{(1-x z_5)(1-x z_3)} \\ z'_2 = z_3 + x \frac{y_g(1+z_4)}{1-x z_5} \\ z'_3 = \frac{y_g(1+z_4)}{1-x z_5} \\ y'_g = \frac{1+z_4}{1-x z_5} \\ y'_g = \frac{1+z_4}{1-x z_5} \\ z'_4 = \frac{z_5(1+z_4)}{1-x z_5} \\ z'_5 = y_g \frac{1+z_4}{1-x z_5} \end{cases}$		3-constrained functional graphs
$\begin{cases} y'_{\ell} = y_{\ell} z_1 z'_2 \\ z'_1 = z_1 z'_2 \\ z'_2 = z_3 \\ z'_3 = y_3 \end{cases}$	$\begin{cases} y'_{\ell} = y_{\ell} z_1 z_3 \\ z'_1 = z_1 z_3 \\ z'_3 = z_3 \end{cases}$		3-balanced hierarchies
$\begin{cases} y_m = \frac{1}{1 - z_1} \\ z'_1 = z_0 \\ z'_0 = z_0 \end{cases}$	$\begin{cases} y'_m = \frac{z_0}{(1-z_1)^2} \\ z'_1 = z_0 \\ z'_0 = z_0 \end{cases}$	$y_m(0) = 1$	Surjections

Figure 3: Example of Species with RDA exponential generating series. Unspecified initial values indicate that all power series solutions are Rationally Dynamically Algebraic, no matter the initial values.

into $y_B = \frac{1}{1-y_A}$; (iii) B = cycle(A) is translated into $y'_B = \frac{y'_A}{1-y_A}$. We refer the reader to [FZC94, Thereom 2] for a more detailed description of this translation.

We show in Figure 3 that each of the specifications presented above translates into a system of differential equations and in addition we show that their exponential generating series are all RDA power series.

Once the system of differential equations is transformed into a rational dynamical system, checking that the exponential generating series is RDA only requires checking that the rational functions are defined on the initial values. In some cases, all initial values yield RDA power series: for example for the systems of non-plane trees, permutations, 3-balanced hierarchies. In the other cases, only specific initial values yield RDA power series (e.g. Functional graphs). Thus so we present in Figure 3 the admissible initial value of the exponential generating series of the species.

We highlight the cases of the Hierarchies and Surjections. For Hierarchies, the initial value $y_h(0) = 0$ is given by the specification. However, understanding the initial value of $z_1(x)$ is less evident. For this, we rely on the expression satisfied by $y_h(x)$ given in [FZC94]: $y_h(x) = x + e^{y_h(x)} - 1 - y_h(x)$. From this we have $y'_h(x) = \frac{1}{2 - e^{y_h(x)}}$ and $z'_2(x) = y'_h(x) - 1$. Since $y_h(0) = 0$, we deduce that $y'_h(0) = 1$ and $z'_2(0) = 0$. Hence, $z'_2(x) = z_1(x)y'_h(x)$ yields $y_1(0) = 0$.

For Surjections, the initial value $y_m(0) = 1$ is given by the specification. Therefore, as $(1 - z_1(x))y_m(x) = 1$, we deduce that $(1 - z_1(0)) \neq 0$.

5 Future Work

We have given algorithms for determining equivalence of holonomic tree automata and their generating functions. For both holonomic recurrences and weighted tree automata there are canonical notions of minimality and associated minimisation algorithms [PZ13, KMW17]. Minimisation of recurrences prove useful in practical applications [Kau23, Chapter 2.3] in which one wants execute and manipulate automata and recurrences. However, when minimal realisations are unique up to isomorphism, the problem of minimisation transcends equality testing—two automata are equivalent if their minimal forms are isomorphic. In future work we plan to investigate whether holonomic tree automata admit canonical minimal realisations and, if so, to apply this to obtaining optimal recurrences for combinatorial classes.

Our exposition has emphasised the generating functions of holonomic tree automata over their formal tree series. It remains to explore the link between such formal tree series and combinatorial objects. Such a relationship was hinted at in Examples 1 and 2, and it is well understood that for families of combinatorial objects with rational generating functions there is typically a natural bijection between the objects and the words of a regular language. Likewise for families of combinatorial objects with algebraic generating functions, there is often a bijection between the objects and trees of of a regular tree language (or the words of an unambiguous context-free language); see [BM08, FS09] for a comprehensive exposition of this phenomenon.

Acknowledgement

We would like to thank Gleb Pogudin for pointing out the paper [DL84]. Mahsa Shirmohammadi and Ait El Manssour are supported by International Emerging Actions grant (IEA'22) and by ANR grant VeSyAM (ANR-22- CE48-0005). James Worrell was supported by UKRI Frontier Research Grant EP/X033813/1.

References

- [BCKN20] Alin Bostan, Arnaud Carayol, Florent Koechlin, and Cyril Nicaud. Weaklyunambiguous parikh automata and their link to holonomic series. In 47th International Colloquium on Automata, Languages, and Programming, ICALP, volume 168 of LIPIcs, pages 114:1–114:16. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- [BCN⁺23] Nikhil Balaji, Lorenzo Clemente, Klara Nosan, Mahsa Shirmohammadi, and James Worrell. Multiplicity problems on algebraic series and context-free grammars. In 38th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2023, Boston, MA, USA, June 26-29, 2023, pages 1–12. IEEE, 2023.
- [BLL98] François Bergeron, Gilbert Labelle, and Pierre Leroux. Combinatorial species and treelike structures. Number 67. Cambridge University Press, 1998.
- [BM08] Mireille Bousquet-Mélou. Rational and algebraic series in combinatorial enumeration. arXiv preprint arXiv:0805.0588, 2008.
- [BR82] Jean Berstel and Christophe Reutenauer. Recognizable formal power series on trees. *Theor. Comput. Sci.*, 18:115–148, 1982.
- [BR90] F. Bergeron and Christophe Reutenauer. Combinatorial resolution of systems of differential equations iii: a special class of differentially algebraic series. *Eur. J. Comb.*, 11:501–512, 1990.
- [BR10] Jean Berstel and Christophe Reutenauer. Noncommutative rational series with applications. 2010.
- [CM17] Giusi Castiglione and Paolo Massazza. On a class of languages with holonomic generating functions. *Theoretical Computer Science*, 658:74–84, 2017.
- [DKV09] Manfred Droste, Werner Kuich, and Heiko Vogler. *Handbook of weighted automata*. Springer Science & Business Media, 2009.
- [DL84] Jan Denef and Leonard Lipshitz. Power series solutions of algebraic differential equations. *Mathematische annalen*, pages 213–238, 1984.
- [DPV05] Manfred Droste, Christian Pech, and Heiko Vogler. A kleene theorem for weighted tree automata. *Theory Comput. Syst.*, 38(1):1–38, 2005.
- [FOR92] On rational state space realizations. IFAC Proceedings Volumes, 25(13):341–346, 1992. 2nd IFAC Symposium on Nonlinear Control Systems Design 1992, Bordeaux, France, 24-26 June.
- [FS09] Philippe Flajolet and Robert Sedgewick. *Analytic combinatorics*. cambridge University press, 2009.
- [FZC94] Philippe Flajolet, Paul Zimmermann, and Bernard Van Cutsem. A calculus for the random generation of labelled combinatorial structures. *Theor. Comput. Sci.*, 132(2):1– 35, 1994.
- [Kau23] Manuel Kauers. *D-finite Functions*. Springer, 2023.

- [KMW17] Stefan Kiefer, Ines Marusic, and James Worrell. Minimisation of multiplicity tree automata. Log. Methods Comput. Sci., 13(1), 2017.
- [Lan52] Serge Lang. Hilbert's nullstellensatz in infinite-dimensional space. Proceedings of the American Mathematical Society, pages 407–410, 1952.
- [LV86] Pierre Leroux and Gérard Viennot. Combinatorial resolution of systems of differential equations, i. ordinary differential equations. 1986.
- [MW15] Ines Marusic and James Worrell. Complexity of equivalence and learning for multiplicity tree automata. J. Mach. Learn. Res., 16:2465–2500, 2015.
- [NS09] Jana Nemcová and Jan Schuppen. Realization theory for rational systems: The existence of rational realizations. *SIAM J. Control and Optimization*, 48:2840–2856, 01 2009.
- [OPV22] Alexey Ovchinnikov, Gleb Pogudin, and Thieu N Vo. Bounds for elimination of unknowns in systems of differential-algebraic equations. International Mathematics Research Notices, pages 12342–12377, 2022.
- [PZ13] Marko Petkovšek and Helena Zakrajšek. Solving linear recurrence equations with polynomial coefficients. In Computer Algebra in Quantum Field Theory: Integration, Summation and Special Functions, pages 259–284. Springer, 2013.
- [Reu12] Christophe Reutenauer. On a matrix representation for polynomially recursive sequences. The Electronic Journal of Combinatorics [electronic only], 19, 09 2012.
- [Sei90] Helmut Seidl. Deciding equivalence of finite tree automata. SIAM J. Comput., 19(3):424–437, 1990.
- [SMK15] Helmut Seidl, Sebastian Maneth, and Gregor Kemper. Equivalence of deterministic topdown tree-to-string transducers is decidable. In Venkatesan Guruswami, editor, IEEE 56th Annual Symposium on Foundations of Computer Science, FOCS, pages 943–962. IEEE Computer Society, 2015.
- [Sta80] Richard P. Stanley. Differentiably finite power series. *Eur. J. Comb.*, 1(2):175–188, 1980.
- [vDH19] Joris van Der Hoeven. Computing with d-algebraic power series. Applicable Algebra in Engineering, Communication and Computing, pages 17–49, 2019.

A Algebraic operations on holonomic tree automata

This appendix is dedicated to the proof of the following theorem.

Theorem 7 (Closure properties). The class of formal tree series recognisable by holonomic tree automata is closed under addition, scalar product (in \mathbb{Q}) and product.

The class of generating functions of holonomic tree automata is closed under addition, scalar product (in \mathbb{Q}), product, derivative, integral, inverse, forward shift and backward shift.

For the class of formal tree series recognisable by holonomic tree automata, the closure properties are given by the following propositions:

• addition: Proposition 9

- scalar product: Proposition 10
- product: Proposition 11

For the class of generating functions of holonomic tree automata, the closure properties are given by the following propositions:

- addition and scalar product: Corollary 2
- product: Corollary 4
- derivation and integral: Corollary 3
- inverse: Proposition 18
- backward shift: Proposition 16
- forward shift: Proposition 17

A.1 The class of formal tree series recognisable by holonomic tree automata

Proposition 8. Let $\mathcal{A} = (d, \mu)$ be a holonomic tree automaton over Σ . Let $\mathcal{B}(x) \in \mathbb{Q}(x)^{d \times 1}$ defined on all non-negative integers. There exists an automaton \mathcal{A}' over Σ such that for all $t \in T_{\Sigma}$, $\mathcal{A}'[tt] = \tilde{\mu}(t)\mathcal{B}(||t||)$.

Proof. We build the holonomic tree automaton by incrementing the dimension of \mathcal{A} by 1 and by building the weight function μ' such that for all $t \in T_{\Sigma}$, $\tilde{\mu}'(t) = [\tilde{\mu}(t)\beta(||t||) \quad \tilde{\mu}(t)]$. Formally, for all $a \in \Sigma_0$, we define $\mu'(a) = [\mu(a)\beta(0) \quad \mu(a)]$; and for all $g \in \Sigma_k$ with k > 0, $\mu'(g) \in \mathbb{Q}[x_0, \ldots, x_k]^{(d+1)^k \times (d+1)}$ and, denoting $M(x_0, \ldots, x_n) = \mu(g)(x_0, \ldots, x_n)\beta(x_0)$, we have:

• for all $j \in \{2, \dots, d+1\}$, for all $i_1, \dots, i_k \in \{1, \dots, d+1\}$,

$$\mu'(g)_{(i_1,\dots,i_k),j} = \begin{cases} \mu(g)_{(i_1-1,\dots,i_k-1),j-1} & \text{if } \forall \ell.i_\ell \in \{2,\dots,d+1\} \\ 0 & \text{otherwise} \end{cases}$$

• for all $i_1, \ldots, i_k \in \{2, \ldots, d+1\},\$

$$\mu'(g)_{(i_1,\dots,i_k),1} = \begin{cases} M_{(i_1-1,\dots,i_k-1),1} & \text{if } \forall \ell.i_\ell \in \{2,\dots,d+1\} \\ 0 & \text{otherwise} \end{cases}$$

Take $\mathcal{A}' = (d', \mu')$ with d' = d + 1, we prove by induction on the structure of trees that for all $t \in T_{\Sigma}$, $\tilde{\mu}'(t) = \begin{bmatrix} \tilde{\mu}(t)\mathcal{B}(||t||) & \tilde{\mu}(t) \end{bmatrix}$. In the base case, t is a leaf a with $a \in \Sigma_0$, meaning that $\tilde{\mu}'(a) = \mu'(a) = \begin{bmatrix} \mu(a)\beta(0) & \mu(a) \end{bmatrix} = \begin{bmatrix} \tilde{\mu}(a)\beta(0) & \tilde{\mu}(a) \end{bmatrix}$.

In the inductive step, $t = g(t_1, \ldots, t_k)$ for some $g \in \Sigma_k$ and $t_1, \ldots, t_k \in T_{\Sigma}$. Let us denote $t_0 = t$. Applying our inductive hypothesis on t_1, \ldots, t_k , we have for all $i \in \{1, \ldots, k\}$, $\tilde{\mu}'(t_i) = [\tilde{\mu}(t_i)\beta(||t_i||) \quad \tilde{\mu}(t_i)]$. Hence,

$$\widetilde{\mu}'(g(t_1,\ldots,t_k)) = (\begin{bmatrix} \widetilde{\mu}(t_1)\beta(\|t_1\|) & \widetilde{\mu}(t_1) \end{bmatrix} \otimes \ldots \otimes \begin{bmatrix} \widetilde{\mu}(t_k)\beta(\|t_k\|) & \widetilde{\mu}(t_k) \end{bmatrix})\mu'(g)(\|t_0\|,\ldots,\|t_k\|)$$

By construction, for all $j \in \{1, ..., d+1\}$, for all $(i_1, ..., i_k) \in \{1, ..., d+1\}^k$, if $i_{\ell} = 1$ for some ℓ then $\mu'(g)_{(i_1,...,i_k),j}(x_0, ..., x_k) = 0$. Thus, for all $j \in \{2, ..., d+1\}$,

$$\widetilde{\mu}'(g(t_1,\ldots,t_k))_j = \sum_{(i_1,\ldots,i_k)\in\{2,\ldots,d+1\}} \left(\prod_{\ell=1}^k \widetilde{\mu}'(t_\ell)_{i_\ell}\right) \mu'(g)_{(i_1,\ldots,i_k),j}(\|t_0\|,\ldots,\|t_k\|)$$

$$= \sum_{(i_1,\dots,i_k)\in\{2,\dots,d+1\}} \left(\prod_{\ell=1}^k \widetilde{\mu}(t_\ell)_{i_\ell-1} \right) \mu(g)_{(i_1-1,\dots,i_k-1),j-1}(\|t_0\|,\dots,\|t_k\|) \\ = \sum_{(i_1,\dots,i_k)\in\{1,\dots,d\}} \left(\prod_{\ell=1}^k \widetilde{\mu}(t_\ell)_{i_\ell} \right) \mu(g)_{(i_1,\dots,i_k),j-1}(\|t_0\|,\dots,\|t_k\|) \\ = ((\widetilde{\mu}(t_1)\otimes\dots\otimes\widetilde{\mu}(t_k))\mu(g)(\|t_0\|,\dots,\|t_k\|))_{j-1} \\ = \widetilde{\mu}(g(t_1,\dots,t_k))_{j-1}$$

Similarly, we also have:

$$\widetilde{\mu}'(g(t_1,\ldots,t_k))_1 = \sum_{(i_1,\ldots,i_k)\in\{2,\ldots,d+1\}} \left(\prod_{\ell=1}^k \widetilde{\mu}(t_\ell)_{i_\ell-1}\right) M_{(i_1-1,\ldots,i_k-1),1}(\|t_0\|,\ldots,\|t_k\|)$$

$$= \sum_{(i_1,\ldots,i_k)\in\{1,\ldots,d\}} \left(\prod_{\ell=1}^k \widetilde{\mu}(t_\ell)_{i_\ell}\right) M_{(i_1,\ldots,i_k),1}(\|t_0\|,\ldots,\|t_k\|)$$

$$= (\widetilde{\mu}(t_1)\otimes\ldots\otimes\widetilde{\mu}(t_k))\mu(g)(\|t_0\|,\ldots,\|t_k\|)\boldsymbol{\beta}(\|t_0\|)$$

$$= \widetilde{\mu}'(g(t_1,\ldots,t_k))\boldsymbol{\beta}(\|g(t_1,\ldots,t_k)\|)$$

This concludes the proof of $\widetilde{\mu}'(t) = \begin{bmatrix} \widetilde{\mu}(t)\beta(||t||) & \widetilde{\mu}(t) \end{bmatrix}$ for all $t \in T_{\Sigma}$. As $\mathcal{A}'[t] = \widetilde{\mu}'(t)_1$, we conclude that $\mathcal{A}'[t] = \widetilde{\mu}(t)\beta(||t||) = \mathcal{A}[t]$ for all $t \in T_{\Sigma}$.

Proposition 9. Let \mathcal{A}_1 and \mathcal{A}_2 be two holonomic tree automata over Σ . There exists a holonomic tree automaton \mathcal{A} over Σ such that for all $t \in T_{\Sigma}$, $\mathcal{A}[t] = \mathcal{A}_1[t] + \mathcal{A}_2[t]$.

Proof. Assume that $A_1 = (d_1, \mu_1)$ and $A_2 = (d_2, \mu_2)$. We build the automaton $A' = (d, \mu)$ of dimension $d = d_1 + d_2$ such that

for all
$$t \in T_{\Sigma}, \widetilde{\mu}(t) = \begin{bmatrix} \widetilde{\mu}_1(t) & \widetilde{\mu}_2(t) \end{bmatrix}$$

and we will then conclude by applying Proposition 8 with the automaton \mathcal{A}' and the vector $\boldsymbol{\beta}(x)$ defined as follows to obtain the desired automaton \mathcal{A} .

$$\boldsymbol{\beta}(x) = \begin{bmatrix} 1\\ \mathbf{0}_{(d_1-1)\times 1}\\ 1\\ \mathbf{0}_{(d_2-1)\times 1} \end{bmatrix}$$

The definition of $\tilde{\mu}$ is fairly straightforward: For all $a \in \Sigma_0$, $\mu(a) = \begin{bmatrix} \mu_1(a) & \mu_2(a) \end{bmatrix}$. Moreover, for all $g \in \Sigma_k$ with k > 0, the matrix $\mu(g)$ is in $\mathbb{Q}[x_0, \ldots, x_k]^{d^k \times d}$ such that for all $i_1, \ldots, i_k \in \{1, \ldots, d\}$,

• for all $j \in \{1, ..., d_1\}$,

$$\mu(g)_{(i_1,\dots,i_k),j} = \begin{cases} \mu_1(g)_{(i_1,\dots,i_k),j} & \text{if } \forall \ell. i_\ell \in \{1,\dots,d_1\} \\ 0 & \text{otherwise} \end{cases}$$

• for all $j \in \{d_1 + 1, \dots d\},\$

$$\mu(g)_{(i_1,\dots,i_k),j} = \begin{cases} \mu_2(g)_{(i_1-d_1,\dots,i_k-d_1),j-d_1} & \text{if } \forall \ell.i_\ell \in \{d_1+1,\dots,d\} \\ 0 & \text{otherwise} \end{cases}$$

The proof of $\tilde{\mu}(t) = \begin{bmatrix} \tilde{\mu}_1(t) & \tilde{\mu}_2(t) \end{bmatrix}$ for all $t \in T_{\Sigma}$ is done by induction on the structure of the tree tand follows by construction of μ . The base case, that is t = a with $a \in \Sigma_0$, is trivial by construction of $\mu(a)$. In the inductive step, $t = g(t_1, \ldots, t_k)$ for some $g \in \Sigma_k$ and $t_1, \ldots, t_k \in T_{\Sigma}$. Let us denote $t_0 = t$. By construction, for all $j \in \{1, \ldots, d_1\}$, for all $i_1, \ldots, i_k \in \{1, \ldots, d\}$, $\mu(g)_{(i_1, \ldots, i_k), j} = 0$ when there exists ℓ such that $i_{\ell} \notin \{1, \ldots, d_1\}$. Thus, by applying our inductive hypothesis on t_1, \ldots, t_k , we obtain that for all $j \in \{1, \ldots, d_1\}$,

$$\widetilde{\mu}(t)_{j} = \sum_{i_{1},\dots,i_{k}\in\{1,\dots,d\}} \left(\prod_{\ell=1}^{k} \widetilde{\mu}(t_{\ell})_{i_{\ell}} \right) \mu(g)_{(i_{1},\dots,i_{k}),j}(\|t_{0}\|,\dots,\|t_{k}\|) \\ = \sum_{i_{1},\dots,i_{k}\in\{1,\dots,d_{1}\}} \left(\prod_{\ell=1}^{k} \widetilde{\mu}(t_{\ell})_{i_{\ell}} \right) \mu(g)_{(i_{1},\dots,i_{k}),j}(\|t_{0}\|,\dots,\|t_{k}\|) \\ = \sum_{i_{1},\dots,i_{k}\in\{1,\dots,d_{1}\}} \left(\prod_{\ell=1}^{k} \widetilde{\mu}_{1}(t_{\ell})_{i_{\ell}} \right) \mu_{1}(g)_{(i_{1},\dots,i_{k}),j}(\|t_{0}\|,\dots,\|t_{k}\|) \\ = \widetilde{\mu}_{1}(t)_{j}$$

Similarly, we have by construction that for all $j \in \{d_1 + 1, \ldots, d\}$, for all $i_1, \ldots, i_k \in \{1, \ldots, d\}$, $\mu(g)_{(i_1,\ldots,i_k),j} = 0$ when there exists ℓ such that $i_\ell \notin \{d_1 + 1, \ldots, d\}$. Hence, by applying our inductive hypothesis on t_1, \ldots, t_k , we obtain that for all $j \in \{d_1 + 1, \ldots, d\}$,

$$\widetilde{\mu}(t)_{j} = \sum_{i_{1},\dots,i_{k}\in\{1,\dots,d\}} \left(\prod_{\ell=1}^{k} \widetilde{\mu}(t_{\ell})_{i_{\ell}} \right) \mu(g)_{(i_{1},\dots,i_{k}),j}(\|t_{0}\|,\dots,\|t_{k}\|) \\ = \sum_{i_{1},\dots,i_{k}\in\{d_{1}+1,\dots,d\}} \left(\prod_{\ell=1}^{k} \widetilde{\mu}_{2}(t_{\ell})_{i_{\ell}-d_{1}} \right) \mu_{2}(g)_{(i_{1}-d_{1},\dots,i_{k}-d_{1}),j-d_{1}}(\|t_{0}\|,\dots,\|t_{k}\|) \\ = \sum_{i_{1},\dots,i_{k}\in\{1,\dots,d_{2}\}} \left(\prod_{\ell=1}^{k} \widetilde{\mu}_{2}(t_{\ell})_{i_{\ell}} \right) \mu_{2}(g)_{(i_{1},\dots,i_{k}),j-n}(\|t_{0}\|,\dots,\|t_{k}\|) \\ = \widetilde{\mu}_{2}(t)_{j-d_{1}}$$

Proposition 10. Let \mathcal{A} be a holonomic tree automaton over Σ . Let $\alpha \in \mathbb{Q}$. There exists a holonomic tree automaton \mathcal{A}' over Σ such that for all $t \in T_{\Sigma}$, $\mathcal{A}'[t]] = \alpha \mathcal{A}[t]$.

Proof. It suffices to apply Proposition 8 on \mathcal{A} with the vector $\boldsymbol{\beta}(x) = \alpha e_1$, where e_1 is the canonical vector.

Proposition 11. Let \mathcal{A}_1 and \mathcal{A}_2 be two holonomic tree automata over Σ . There exists a holonomic tree automaton \mathcal{A} over Σ such that for all $t \in T_{\Sigma}$, $\mathcal{A}[t] = \mathcal{A}_1[t] \cdot \mathcal{A}_2[t]$.

Proof. We take $\mathcal{A}_1 = (d_1, \mu_1)$ and $\mathcal{A}_2 = (d_2, \mu_2)$. We build the automaton $\mathcal{A} = (d, \mu)$ of dimension $d = d_1 d_2$ such that for all $t \in T_{\Sigma}$,

$$\widetilde{\mu}(t) = \begin{bmatrix} \widetilde{\mu}_1(t)_1 \widetilde{\mu}_2(t)_1 & \dots & \widetilde{\mu}_1(t)_1 \widetilde{\mu}_2(t)_{d_2} & \widetilde{\mu}_1(t)_2 \widetilde{\mu}_2(t)_1 & \dots & \widetilde{\mu}_1(t)_{d_1} \widetilde{\mu}_2(t)_{d_2} \end{bmatrix}$$

For that purpose, we construct the function μ as follows. For all $a \in \Sigma_0$, for all $i \in \{1, \ldots, d_1\}$, for all $j \in \{1, \ldots, d_2\}$, $\mu(a)_{(i,j)} = \mu_1(a)_i \mu_2(a)_j$. For all $g \in \Sigma_k$ with k > 0, for all $i, i_1, \ldots, i_k \in \{1, \ldots, d_1\}$, for all $j, j_1, \ldots, j_k \in \{1, \ldots, d_2\}$,

$$\mu(g)_{((i_1,j_1),\dots,(i_k,j_k)),(i,j)} = \mu_1(g)_{(i_1,\dots,i_k),i} \cdot \mu_2(g)_{(j_1,\dots,j_k),j}$$

By definition, $\widetilde{\mu}(g(t_1,\ldots,t_k)) = (\widetilde{\mu}(t_1)\otimes\ldots\otimes\widetilde{\mu}(t_k))\mu(g)(||t_0||,\ldots,||t_k||)$. Therefore, denoting $t_0 = t$ and applying our inductive hypothesis on each t_1, \ldots, t_k , we obtain that for all $i \in \{1, \ldots, d_1\}$, for all $j \in \{1, \ldots, d_2\}$,

$$\begin{split} \widetilde{\mu}(t)_{(i,j)} &= \sum_{\substack{i_1, \dots, i_k \in \{1, \dots, d_1\} \\ j_1, \dots, j_k \in \{1, \dots, d_2\}}} \left(\prod_{\ell=1}^k \widetilde{\mu}(t_\ell)_{(i_\ell, j_\ell)} \right) \cdot \mu(g)(\|t_0\|, \dots, \|t_k\|)_{((i_1, j_1), \dots, (i_k, j_k)), (i, j)} \\ &= \sum_{\substack{i_1, \dots, i_k \in \{1, \dots, d_1\} \\ j_1, \dots, j_k \in \{1, \dots, d_2\}}} \left(\prod_{\ell=1}^k \widetilde{\mu}_1(t_\ell)_{i_\ell} \widetilde{\mu}_2(t_\ell)_{j_\ell} \right) \cdot \mu_1(g)(\|t_0\|, \dots, \|t_k\|)_{(i_1, \dots, i_k), i} \cdot \mu_2(g)(\|t_0\|, \dots, \|t_k\|)_{(j_1, \dots, j_k), j} \right) \\ &= \sum_{\substack{i_1, \dots, i_k \in \{1, \dots, d_1\} \\ j_1, \dots, j_k \in \{1, \dots, d_2\}}} \left(\left(\prod_{\ell=1}^k \widetilde{\mu}_1(t_\ell)_{i_\ell} \right) \mu_1(g)(\|t_0\|, \dots, \|t_k\|)_{(i_1, \dots, i_k), i} \right) \cdot \left(\left(\prod_{\ell=1}^k \widetilde{\mu}_2(t_\ell)_{j_\ell} \right) \mu_2(g)(\|t_0\|, \dots, \|t_k\|)_{(j_1, \dots, j_k), j} \right) \\ &= \left(\sum_{\substack{i_1, \dots, i_k \in \{1, \dots, d_2\} \\ (i_1, \dots, i_k \in \{1, \dots, d_2\}}} \left(\prod_{\ell=1}^k \widetilde{\mu}_2(t_\ell)_{j_\ell} \right) \mu_2(g)(\|t_1\|, \dots, \|t_k\|)_{(j_1, \dots, j_k), j} \right) \\ &= \left((\widetilde{\mu}_1(t_1) \otimes \dots \otimes \widetilde{\mu}_1(t_k)) \mu_1(g)(\|t_1\|, \dots, \|t_k\|))_i \cdot ((\widetilde{\mu}_2(t_1) \otimes \dots \otimes \widetilde{\mu}_2(t_k)) \mu_2(g)(\|t_0\|, \dots, \|t_k\|))_j \right) \\ &= \widetilde{\mu}_1(t_i \cdot \widetilde{\mu}_2(t_j)_j \end{split}$$

We conclude by noticing that $\mathcal{A}\llbracket t \rrbracket = \widetilde{\mu}(t)_1 = \widetilde{\mu}_1(t)_1 \widetilde{\mu}_2(t)_1 = \mathcal{A}_1\llbracket t \rrbracket \cdot \mathcal{A}_2\llbracket t \rrbracket$.

Operations on the generating functions A.2

We start by showing that when considering generating functions, we can always restrict ourselves to alphabets Σ where two different function symbols must have different arity. In other words, there can only be one nullary function symbol, one unary function symbol, one binary function symbol, etc. We say in this case that Σ is arity distinct.

Proposition 12. Let \mathcal{A} be a holonomic tree automaton over Σ . There exists a holonomic tree automaton \mathcal{A}' over arity distinct Σ' such that $f_{\mathcal{A}}(x) = f_{\mathcal{A}'}(x)$.

Proof. Let $\mathcal{A} = (d, \mu)$. Let r be the largest arity in Σ . For all $k \in \{0, \ldots, r\}$, we define a new function symbol h_k of arity k and we build $\Sigma' = \{h_k\}_{k=0}^r$. Additionally, we build the automaton

function symbol h_k of arity κ and we build $\Delta = \lim_{g \in \Sigma_k} \mu(g)$. $\mathcal{A}' = (d, \mu')$ such that for all $k \in \{0, \dots, r\}, \ \mu'(h_k) = \sum_{g \in \Sigma_k} \mu(g)$. Consider now the vector of power series $\boldsymbol{f}(x) = \sum_{n=0}^{\infty} \boldsymbol{a}_n x^n$ where $\boldsymbol{a}_n = \sum_{\substack{t \in T_{\Sigma} \\ \|t\|=n}} \widetilde{\mu}(t)$. Similarly, consider the vector of power series $\boldsymbol{g}(x) = \sum_{n=0}^{\infty} \boldsymbol{b}_n x^n$ where $\boldsymbol{b}_n = \sum_{\substack{t \in T_{\Sigma'} \\ \|t\|=n}} \widetilde{\mu}'(t)$. We show by $\|t\|=n$

induction on n that for all $n \in \mathbb{N}$, $\boldsymbol{a}_n = \boldsymbol{b}_n$. In the base case (n = 0), $\boldsymbol{a}_0 = \sum_{a \in \Sigma_0} \widetilde{\mu}(a) = \sum_{a/0 \in \Sigma} \mu(a) = \mu'(h_0) = \widetilde{\mu}'(h_0) = \boldsymbol{b}_0$. In the

inductive step (n > 0), from Equation (4), we deduce that

$$\begin{aligned} \mathbf{a}_{n} &= \sum_{k=1}^{r} \sum_{g \in \Sigma_{k}} \sum_{\substack{n_{1}, \dots, n_{k} \in \mathbb{N} \\ n_{1} + \dots + n_{k} = n - 1}} (\mathbf{a}_{n_{1}} \otimes \dots \otimes \mathbf{a}_{n_{k}}) \mu(g)(n, n_{1}, \dots, n_{k}) \\ &= \sum_{k=1}^{r} \sum_{\substack{n, n_{1}, \dots, n_{k} \in \mathbb{N} \\ n_{1} + \dots + n_{k} = n - 1}} (\mathbf{a}_{n_{1}} \otimes \dots \otimes \mathbf{a}_{n_{k}}) \sum_{g \in \Sigma_{k}} \mu(g)(n, n_{1}, \dots, n_{k}) \\ &= \sum_{k=1}^{r} \sum_{\substack{n, n_{1}, \dots, n_{k} \in \mathbb{N} \\ n_{1} + \dots + n_{k} = n - 1}} (\mathbf{a}_{n_{1}} \otimes \dots \otimes \mathbf{a}_{n_{k}}) \mu'(h_{k})(n, n_{1}, \dots, n_{k}) \\ &= \sum_{k=1}^{r} \sum_{\substack{n, n_{1}, \dots, n_{k} \in \mathbb{N} \\ n_{1} + \dots + n_{k} = n - 1}} (\mathbf{b}_{n_{1}} \otimes \dots \otimes \mathbf{b}_{n_{k}}) \mu'(h_{k})(n, n_{1}, \dots, n_{k}) \\ &= \mathbf{b}_{n} \end{aligned}$$

As $f_{\mathcal{A}}(x) = \boldsymbol{f}(x)_1$ and $f_{\mathcal{A}'}(x) = \boldsymbol{g}(x)_1$, we conclude.

Proposition 13. Let \mathcal{A}_1 and \mathcal{A}_2 be two holonomic tree automata over Σ_1 and Σ_2 respectively. Therefore exist an alphabet Σ and two holonomic tree automata \mathcal{A}'_1 and \mathcal{A}'_2 over Σ such that $f_{\mathcal{A}_1}(x) = f_{\mathcal{A}'_1}(x)$ and $f_{\mathcal{A}_2}(x) = f_{\mathcal{A}'_2}(x)$.

Proof. Take $\mathcal{A}_1 = (d_1, \mu_1)$ and $\mathcal{A}_2 = (d_2, \mu_2)$. By Proposition 12, we can also assume that Σ_1 and Σ_2 are arity distinct.

Let r_1 and r_2 the maximum arity of symbols in Σ_1 and Σ_2 . Let $\{k_1, \ldots, k_\ell\}$ be the set of arity in $\{0, \ldots, r_2\}$ such that for all $i \in \{1, \ldots, \ell\}$, Σ_2 contains a symbol of arity k_i but not Σ_1 . We define Σ'_1 by extending Σ_1 with the fresh symbols $h_{k_1}, \ldots, h_{k_\ell}$ of arity k_1, \ldots, k_ℓ respectively and we extend \mathcal{A}_1 into $\mathcal{A}'_1 = (d_1, \mu'_1)$ such that $\mu'_1(h_{k_i}) = \mathbf{0}_{d_1^{k_i} \times d_1}$ for all $i \in \{1, \ldots, \ell\}$. We trivially have $f_{\mathcal{A}_1}(x) = f_{\mathcal{A}'_1}(x)$.

We extend in a similar fashion Σ_2 and \mathcal{A}_2 by computing $\{k'_1, \ldots, k'_{\ell'}\}$ to be the set of arities in $\{0, \ldots, r_1\}$ such that for all $i \in \{1, \ldots, \ell'\}$, Σ_1 contains a symbol of arity k'_i but not Σ_2 . This yields an automaton \mathcal{A}'_2 over Σ'_2 such that $f_{\mathcal{A}'_2}(x) = f_{\mathcal{A}_2}(x)$. As Σ_1 and Σ_2 are arity distinct, we have by construction that Σ'_1 is a renaming of Σ'_2 . Therefore, we can fully rename the automaton \mathcal{A}'_1 to be over Σ'_2 with $f_{\mathcal{A}_1}(x) = f_{\mathcal{A}'_1}(x)$.

Corollary 2. Let \mathcal{A}_1 and \mathcal{A}_2 be two holonomic tree automata. Let $\alpha \in \mathbb{Q}$.

- there exists a holonomic tree automaton such that $f_{\mathcal{A}}(x) = f_{\mathcal{A}_1}(x) + f_{\mathcal{A}_2}(x)$.
- there exists a holonomic tree automaton such that $f_{\mathcal{A}}(x) = \alpha f_{\mathcal{A}_1}(x)$.

Proof. Direct from Propositions 9, 10 and 13.

Proposition 14. Let \mathcal{A} be a holonomic tree automaton. There exists a holonomic tree automaton \mathcal{A}' such that $f_{\mathcal{A}'}(x) = \Theta f_{\mathcal{A}}(x) = x f'_{\mathcal{A}}(x)$.

Proof. Let $\mathcal{A} = (d, \mu)$ over the alphabet Σ . Let us consider the vector $\boldsymbol{\beta}(x) = \begin{bmatrix} x \\ \mathbf{0}_{(d-1)\times 1} \end{bmatrix}$. By Proposition 8, there exists a holonomic tree automaton \mathcal{A}' such that $\mathcal{A}'[t] = \tilde{\mu}(t)\boldsymbol{\beta}(||t||) = ||t||\tilde{\mu}(t)_1 = ||t||\mathcal{A}[t]$ for all $t \in T_{\Sigma}$. Hence, we directly obtain that $f_{\mathcal{A}'}(x) = \sum_{n=0}^{\infty} n \sum_{\substack{t \in T_{\Sigma} \\ \|t\| = n}} \mathcal{A}[t] x^n$. By Proposition 1, we conclude that $f_{\mathcal{A}'}(x) = \Theta f_{\mathcal{A}}(x)$.

Proposition 15. Let \mathcal{A} be a holonomic tree automaton. There exists a holonomic tree automaton \mathcal{A}' such that $f_{\mathcal{A}'}(x) = \frac{1}{x} \int_0^x f_{\mathcal{A}}(x) dx$.

Proof. Let $\mathcal{A} = (d, \mu)$ over an alphabet Σ . Let us consider the vectore $\boldsymbol{\beta}(x) = \begin{bmatrix} \frac{1}{1+x} \\ \mathbf{0}_{(d-1)\times 1} \end{bmatrix}$. By Proposition 8, there exists a holonomic tree automaton \mathcal{A}' such that $\mathcal{A}'[t] = \tilde{\mu}(t)\boldsymbol{\beta}(||t||) = \frac{1}{1+||t||}\tilde{\mu}(t)_1 = \frac{1}{1+||t||}\mathcal{A}[t]$. Hence, we directly obtain that:

$$f_{\mathcal{A}'}(x) = \sum_{n=0} \frac{1}{1+n} \sum_{\substack{t \in T_{\Sigma} \\ \|t\|=n}} \mathcal{A}\llbracket t \rrbracket x^n = \frac{1}{x} \int_0^x f_{\mathcal{A}}(x) dx \qquad \Box$$

Proposition 16. Let \mathcal{A}_1 and \mathcal{A}_2 be two holonomic tree automata.

- There exists a holonomic tree automaton \mathcal{A} such that $f_{\mathcal{A}}(x) = x \cdot f_{\mathcal{A}_1}(x)$.
- There exists a holonomic tree automaton \mathcal{A} such that $f_{\mathcal{A}}(x) = x \cdot f_{\mathcal{A}_1}(x) \cdot f_{\mathcal{A}_2}(x)$.

Proof. Without loss of generality, let us assume that \mathcal{A}_1 and \mathcal{A}_2 are automata over distinct alphabet Σ^1 and Σ^2 (we can always rename the function symbols otherwise). Let $\mathcal{A}_1 = (d_1, \mu_1)$ and $\mathcal{A}_2 = (d_2, \mu_2)$.

We start by building \mathcal{A} such that $f_{\mathcal{A}}(x) = x \cdot f_{\mathcal{A}_1}(x)$. Let us take a new unary symbol u/1 not already in Σ^1 . We build the automaton \mathcal{A} to have dimension $d_1 + 1$ over $\Sigma = \{u\} \cup \Sigma^1$ with the weight function μ such that for all $a \in \Sigma_0^1$, $\mu(a) = \begin{bmatrix} 0 & \mu_1(a) \end{bmatrix}$; and for all k > 0, for all $g \in \Sigma_k^1$, for all $i_1, \ldots, i_k, j \in \{1, \ldots, d_1 + 1\}$,

$$\mu(g)_{(i_1,\dots,i_k),j} = \begin{cases} \mu_1(g)_{(i_1-1,\dots,i_k-1),j-1} & \text{if } \forall \ell.i_\ell, j \in \{2,\dots,d_1+1\} \\ 0 & \text{otherwise} \end{cases}$$

and

$$\mu(u)(x_0, x_1) = \begin{bmatrix} 0 & \mathbf{0}_{1 \times d_1} \\ 1 & \mathbf{0}_{1 \times d_1} \\ \mathbf{0}_{(d_1 - 1) \times 1} & \mathbf{0}_{(d_1 - 1) \times d_1} \end{bmatrix}$$

With such a construction, we can show the following property: for all $t \in T_{\Sigma}$,

- if $t \in T_{\Sigma^1}$ then $\widetilde{\mu}(t) = \begin{bmatrix} 0 & \widetilde{\mu}_1(t) \end{bmatrix}$;
- if t = u(t') with $t' \in T_{\Sigma^1}$ then $\widetilde{\mu}(u(t')) = \begin{bmatrix} \widetilde{\mu}_1(t')_1 & \mathbf{0}_{1 \times d_1} \end{bmatrix}$
- and $\widetilde{\mu}(t) = \mathbf{0}_{1 \times (d_1+1)}$ otherwise

We prove these properties by induction on the structure of t. In the base case, t is necessarily a nullary symbol $a \in \Sigma_0^1$. By definition $\mu(a) = \begin{bmatrix} 0 & \mu_1(a) \end{bmatrix}$ hence the result holds. In the inductive step, we do a case analysis on t:

• Case $t = g(t_1, \ldots, t_k)$ with $g \in \Sigma_k^1$ and $t_1, \ldots, t_k \in T_{\Sigma^1}$: By inductive hypothesis on t_1, \ldots, t_k , we have that for all $i \in \{1, \ldots, d_1 + 1\}$,

$$\widetilde{\mu}(t)_i = (\begin{bmatrix} 0 & \widetilde{\mu}_1(t_1) \end{bmatrix} \otimes \ldots \otimes \begin{bmatrix} 0 & \widetilde{\mu}_1(t_k) \end{bmatrix}) \mu(g)(\|t\|, \|t_1\|, \ldots, \|t_k\|)_i$$

Thus, by definition of $\mu(g)$, we have $\widetilde{\mu}(t)_1 = 0$ and for all $i \in \{2, \ldots, d_1 + 1\}$, $\widetilde{\mu}(t)_i = (\widetilde{\mu}_1(t_1) \otimes \ldots \otimes \widetilde{\mu}_1(t_k)) \mu_1(g)(||t||, ||t_1||, \ldots, ||t_k||)_{i-1}$. We thus obtain:

$$\widetilde{\mu}(g(t_1,\ldots,t_k)) = \begin{bmatrix} 0 & \widetilde{\mu}_1(g(t_1,\ldots,t_k)) \end{bmatrix}$$

• Case $t = g(t_1, \ldots, t_k)$ with $g \in \Sigma_k^1$ and there exists $\ell \in \{1, \ldots, k\}$ such that $t_\ell \notin T_{\Sigma^1}$: By inductive hypothesis on t_ℓ , we know that for all $i \in \{2, \ldots, d_1 + 1\}$, $\tilde{\mu}(t_\ell)_i = 0$. Thus, we deduce from the definition of $\mu(g)$ that:

$$(\widetilde{\mu}(t_1)\otimes\ldots\otimes\widetilde{\mu}(t_k))\mu(g)(||t||,||t_1||,\ldots,||t_k||)=\mathbf{0}_{1\times(d_1+1)}$$

• Case t = u(t') with $t' \in T_{\Sigma^1}$: By inductive hypothesis on t', we know that $\tilde{\mu}(t') = \begin{bmatrix} 0 & \tilde{\mu}_1(t') \end{bmatrix}$. Thus, following the definition of $\mu(u)$, we directly have that:

$$\widetilde{\mu}(t) = \widetilde{\mu}(t')\mu(u)(\|t\|, \|t'\|) = \begin{bmatrix} \widetilde{\mu}_1(t)_1 & \mathbf{0}_{1 \times d_1} \end{bmatrix}$$

• Case t = u(t') with $t' \notin T_{\Sigma^1}$: By inductive hypothesis on t', we know that for all $i \in \{2, \ldots, d_1 + 1\}$, $\tilde{\mu}(t')_i = 0$. Hence, by definition of $\mu(u)$, we obtain:

$$\widetilde{\mu}(t) = \widetilde{\mu}(t')\mu(u)(||t||, ||t'||) = \mathbf{0}_{1 \times (d_1+1)}$$

Let us now conclude the proof by computing the generating series. First notice that for all $t \in T_{\Sigma}$, $\tilde{\mu}(t)_1 \neq 0$ implies t = u(t') with $t' \in T_{\Sigma^1}$. Hence,

$$f_{\mathcal{A}}(x) = \sum_{n=0}^{\infty} \sum_{\substack{t \in T_{\Sigma} \\ \|t\| = n}} \widetilde{\mu}(t)_{1} x^{n} = \sum_{n=1}^{\infty} \sum_{\substack{t' \in T_{\Sigma^{1}} \\ \|t'\| = n-1}} \widetilde{\mu}(u(t'))_{1} x^{n} = x \sum_{n=1}^{\infty} \sum_{\substack{t' \in T_{\Sigma^{1}} \\ \|t'\| = n-1}} \widetilde{\mu}_{1}(t')_{1} x^{n-1} = x \cdot f_{\mathcal{A}_{1}}(x)$$

We now build a holonomic tree automaton \mathcal{A} such that $f_{\mathcal{A}}(x) = x \cdot f_{\mathcal{A}_1}(x) \cdot f_{\mathcal{A}_2}(x)$. The proof is in fact very similar to the above proof. Instead of considering a new unary symbol, we consider a new binary symbol u (hence arity 2) not already in Σ^1 and Σ^2 . We build the automaton \mathcal{A} over $\Sigma = \{u\} \cup \Sigma^1 \cup \Sigma^2$ with dimension $d = d_1 + d_2 + 1$, and the weight function μ such that:

- for all $a \in \Sigma_0^1$, $\mu(a) = \begin{bmatrix} 0 & \mu_1(a) & \mathbf{0}_{1 \times d_2} \end{bmatrix}$
- for all $a \in \Sigma_0^2$, $\mu(a) = \begin{bmatrix} 0 & \mathbf{0}_{1 \times d_1} & \mu_2(a) \end{bmatrix}$
- for all k > 0, for all $g \in \Sigma_k^1$, for all $i_1, \ldots, i_k, j \in \{1, \ldots, d\}$,

$$\mu(g)_{(i_1,\dots,i_k),j} = \begin{cases} \mu_1(g)_{(i_1-1,\dots,i_k-1),j-1} & \text{if } \forall \ell.i_\ell, j \in \{2,\dots,d_1+1\} \\ 0 & \text{otherwise} \end{cases}$$

• for all k > 0, for all $g \in \Sigma_k^2$, for all $i_1, \ldots, i_k, j \in \{1, \ldots, d\}$, denoting $n = d_1 + 1$,

$$\mu(g)_{(i_1,\dots,i_k),j} = \begin{cases} \mu_1(g)_{(i_1-n,\dots,i_k-n),j-n} & \text{if } \forall \ell. i_\ell, j \in \{d_1+2,\dots,d\} \\ 0 & \text{otherwise} \end{cases}$$

• $\mu(u)_{(1,1),0} = 1$ and otherwise $\mu(u)_{(i_1,i_2),j} = 0$. In other words:

$$\mu(u) = \begin{bmatrix} \mathbf{0}_{(d_1+1)\times(d_1+d_2)} & \mathbf{0}_{(d_1+1)\times(d_1+d_2)} \\ 1 & \mathbf{0}_{1\times(d_1+d_2)} \\ \mathbf{0}_{(d^2-d_1-2)\times1} & \mathbf{0}_{(d^2-d_1-2)\times(d_1+d_2)} \end{bmatrix}$$

This construction naturally entails a similar property: for all $t \in T_{\Sigma}$,

- if $t \in T_{\Sigma^1}$ then $\widetilde{\mu}(t) = \begin{bmatrix} 0 & \widetilde{\mu}_1(t) & \mathbf{0}_{1 \times d_2} \end{bmatrix}$;
- if $t \in T_{\Sigma^2}$ then $\widetilde{\mu}(t) = \begin{bmatrix} 0 & \mathbf{0}_{1 \times d_1} & \widetilde{\mu}_2(t) \end{bmatrix}$;
- if $t = u(t_1, t_2)$ with $t_1 \in T_{\Sigma^1}$ and $t_2 \in T_{\Sigma^2}$ then

$$\widetilde{\mu}(u(t_1, t_2)) = \begin{bmatrix} \widetilde{\mu}_1(t_1)_1 \cdot \widetilde{\mu}_2(t_2)_1 & \mathbf{0}_{1 \times (d_1 + d_2)} \end{bmatrix}$$

• and $\widetilde{\mu}(t) = \mathbf{0}_{1 \times (d_1 + d_2 + 1)}$ otherwise

The proof of this property is done once again by induction on the structure of t and is very similar to the above unary case, hence we omit the details. We now conclude the main proof by computing the generating series: Once again, notice that for all $t \in T_{\Sigma}$, $\tilde{\mu}(t)_1 \neq 0$ implies $t = u(t_1, t_2)$. Therefore, we have:

$$\begin{split} f_{\mathcal{A}}(x) &= \sum_{n=0}^{\infty} \sum_{\substack{t \in T_{\Sigma} \\ \|t\| = n}} \widetilde{\mu}(t)_{1} x^{n} \\ &= \sum_{n=1}^{\infty} \sum_{k=0}^{n-1} \sum_{\substack{t_{1} \in T_{\Sigma^{1}} \\ \|t_{1}\| = k}} \sum_{\substack{t_{2} \in T_{\Sigma^{2}} \\ \|t_{2}\| = n-1-k}} \widetilde{\mu}_{1}(t_{1})_{1} \cdot \widetilde{\mu}_{2}(t_{2})_{1} x^{n} \\ &= x \sum_{n=1}^{\infty} \sum_{k=0}^{n-1} \left(\sum_{\substack{t_{1} \in T_{\Sigma^{1}} \\ \|t_{1}\| = k}} \widetilde{\mu}_{1}(t_{1})_{1} x^{k} \right) \cdot \left(\sum_{\substack{t_{2} \in T_{\Sigma^{2}} \\ \|t_{2}\| = n-1-k}} \widetilde{\mu}_{2}(t_{2})_{1} x^{n-1-k} \right) \\ &= x \cdot \left(\sum_{n=0}^{\infty} \sum_{\substack{t_{1} \in T_{\Sigma^{1}} \\ \|t_{1}\| = n}} \widetilde{\mu}_{1}(t_{1})_{1} x^{n} \right) \cdot \left(\sum_{\substack{n=0 \\ \|t_{2} \in T_{\Sigma^{2}} \\ \|t_{2}\| = n}} \widetilde{\mu}_{2}(t_{2})_{1} x^{n} \right) \\ &= x \cdot f_{\mathcal{A}_{1}}(x) \cdot f_{\mathcal{A}_{2}}(x) \end{split}$$

Proposition 17. Let \mathcal{A} be a holonomic tree automaton. Let $f_{\mathcal{A}} = \sum_{n=0}^{\infty} a_n x^n$. There exists a holonomic tree automaton \mathcal{A}' such that $f_{\mathcal{A}'}(x) = \frac{f_{\mathcal{A}}(x) - a_0}{x} = \sum_{n=0}^{\infty} a_{n+1} x^n$.

Proof. Let $\mathcal{A} = (d, \mu)$ over Σ . Thanks for Proposition 12, we assume that Σ is arity distinct. In other words, we can assume that $\Sigma = \{g_k\}_{k=0}^r$ for some r with g_0, \ldots, g_r of arity $0, \ldots, r$ respectively. The difficulty of this proof is that we need to *decrease* the size of the trees of \mathcal{A} by 1. We thus consider a new alphabet $\Sigma' = \Sigma \cup \bigcup_{k=1}^r \bigcup_{i=0}^k \{h_{k,i}\}$ where each $h_{k,i}$ has arity i. Intuitively a term $h_{k,i}(t_1, \ldots, t_{i-1}, u)$ will have the same value in \mathcal{A}' as the term $g_k(t_1, \ldots, t_{i-1}, t_i, g_0, \ldots, g_0)$ in \mathcal{A} with t_i having the same value as u but with the size decreased by 1. Formally, we build an injective transformation Γ from trees t of T_{Σ} of size ||t|| > 0 to $T_{\Sigma'}$ as follows: For all $t \in T_{\Sigma}$, if $t = g_k(t_1, \ldots, t_k)$ with k > 0 and $i = \min(\{k\} \cup \{j-1 \mid \forall k \ge \ell \ge j.t_\ell = g_0\})$ then

$$\Gamma(t) = h_{k,i}(t_1, \dots, t_{i-1}, \Gamma(t_i))$$

Notice a simple inductive proof allows us to show that for all $t \in T_{\Sigma}$, $||t|| = ||\Gamma(t)|| + 1$.

We will build the automata \mathcal{A}' to have dimension d' = 2d over Σ' with the weight function μ' such that for all $t \in T_{\Sigma'}$,

• if $t \in T_{\Sigma}$ then $\widetilde{\mu}'(t) = \begin{bmatrix} \mathbf{0}_{d \times 1} & \widetilde{\mu}(t) \end{bmatrix}$

- if there exists $t' \in T_{\Sigma}$ such that $\Gamma(t') = t$ then $\widetilde{\mu}'(t) = \begin{bmatrix} \widetilde{\mu}(t') & \mathbf{0}_{d \times 1} \end{bmatrix}$
- $\widetilde{\mu}'(t) = \mathbf{0}_{2d \times 1}$ otherwise.

To achieve this property, we build μ' as follows:

- $\mu'(g_0) = \begin{bmatrix} \mathbf{0}_{d \times 1} & \mu(g_0) \end{bmatrix}$
- for all $k \in \{1, ..., r\}$, for all $i_1, ..., i_k, j \in \{1, ..., 2d\}$,

$$\mu'(g_k)_{(i_1,\dots,i_k),j} = \begin{cases} \mu(g)_{(i_1-d,\dots,i_k-d),j-d} & \text{if } \forall \ell.i_\ell, j \in \{d+1,\dots,2d\} \\ 0 & \text{otherwise} \end{cases}$$

- for all $k \in \{1, ..., r\}, \mu'(h_{k,0}) = \begin{bmatrix} \widetilde{\mu}(g_k(g_0, ..., g_0)) & \mathbf{0}_{d \times 1} \end{bmatrix}$.
- for all $k \in \{1, \ldots, r\}$, for all $\ell \in \{1, \ldots, k\}$, denoting

$$M^{k,\ell}(x_0, x_1, \dots, x_\ell) = (I_{d^\ell} \otimes \bigotimes_{j=\ell+1}^k \mu(g_0))\mu(g_k)(x_0+1, x_1, \dots, x_{\ell-1}, x_\ell+1, 0, \dots, 0)$$

we define for all for all $i_1, \ldots, i_\ell, j \in \{1, \ldots, 2d\}$,

$$\mu'(h_{k,\ell})_{(i_1,\dots,i_\ell),j} = \begin{cases} M^{k,\ell}_{(i_1-d,\dots,i_{\ell-1}-d,i_\ell),j} & \text{if } i_1,\dots,i_{\ell-1} \in \{d+1,\dots,2d\}, i_\ell, j \in \{1,\dots,d\}\\ 0 & \text{otherwise} \end{cases}$$

We now prove the desired property by induction on the structure of t. In the base case, t is a nullary symbol. Hence, either $t = g_0$ or $t = h_{k,0}$ for some $k \in \{1, \ldots, r\}$. In the former case, we have by definition that $\mu'(g_0) = \begin{bmatrix} \mathbf{0}_{d\times 1} & \mu(g_0) \end{bmatrix}$ hence the result directly holds. In the latter case, by definition of $h_{k,0}$, we know that $\Gamma(g_k(g_0, \ldots, g_0)) = h_{k,0}$. Moreover, by definition, we have $\mu'(h_{k,0}) = \begin{bmatrix} \widetilde{\mu}(g_k(g_0, \ldots, g_0)) & \mathbf{0}_{d\times 1} \end{bmatrix}$.

In the inductive step, we have that either $t = g_k(t_1, \ldots, t_k)$ or $t = h_{k,i}(t_1, \ldots, t_i)$ for some k > 0and i > 0.

• Case $t = g_k(t_1, \ldots, t_k)$ when $t_1, \ldots, t_k \in T_{\Sigma}$: By inductive hypothesis on each t_ℓ s, we have that for all $j \in \{1, \ldots, 2d\}$,

$$\widetilde{\mu}'(t)_j = (\begin{bmatrix} \mathbf{0}_{1 \times d} & \widetilde{\mu}(t_1) \end{bmatrix} \otimes \ldots \otimes \begin{bmatrix} \mathbf{0}_{1 \times d} & \widetilde{\mu}(t_k) \end{bmatrix}) \mu'(g_k) (\|t\|, \|t_1\|, \dots, \|t_k\|)_j$$

Thus, by definition of $\mu'(g_k)$, we have $\widetilde{\mu}'(t)_j = 0$ for $j \in \{1, \ldots, d\}$ and $\widetilde{\mu}'(t)_j = (\widetilde{\mu}(t_1) \otimes \ldots \otimes \widetilde{\mu}(t_k))\mu(g_k)(||t||, ||t_1||, \ldots, ||t_k||)_{j-d}$ when $j \in \{d+1, \ldots, 2d\}$. We deduce that $\widetilde{\mu}'(t) = [\mathbf{0}_{d\times 1} \quad \widetilde{\mu}(t)]$.

• Case $t = g_k(t_1, \ldots, t_k)$ and there exists $\ell \in \{1, \ldots, k\}$ such that $t_\ell \in T_{\Sigma}$: By inductive hypothesis on t_ℓ , we know that for all $j \in \{d + 1, \ldots, 2d\}$, $\tilde{\mu}(t_\ell)_j = 0$. Thus we deduce from the definition of $\mu'(g_k)$ that

$$(\widetilde{\mu}'(t_1)\otimes\ldots\otimes\widetilde{\mu}'(t_k))\mu(g_k)(\|t\|,\|t_1\|,\ldots,\|t_k\|)=\mathbf{0}_{1\times 2d}$$

• Case $t = h_{k,i}(t_1, \ldots, t_i)$ and $t_1, \ldots, t_{i-1} \in T_{\Sigma}$ and there exists $t'_i \in T_{\Sigma}$ such that $\Gamma(t'_i) = t_i$: By inductive hypothesis on each t_ℓ s, we know that for all $j \in \{1, \ldots, 2d\}$,

$$\widetilde{\mu}'(t)_j = (\begin{bmatrix} \mathbf{0}_{1 \times d} & \widetilde{\mu}(t_1) \end{bmatrix} \otimes \ldots \otimes \begin{bmatrix} \mathbf{0}_{1 \times d} & \widetilde{\mu}(t_{i-1}) \end{bmatrix} \otimes \begin{bmatrix} \widetilde{\mu}(t'_i) & \mathbf{0}_{1 \times d} \end{bmatrix}) \mu'(h_{k,i}) (\|t\|, \|t_1\|, \dots, \|t_i\|)_j$$

Recall that as $\Gamma(t'_i) = t_i$, we have $||t'_i|| = ||t_i|| + 1$. Denoting $t' = g_k(t_1, \ldots, t_{i-1}, t'_i, g_0, \ldots, g_0)$, we thus obtain ||t'|| = ||t|| + 1. Hence unfolding the definition of $\mu'(h_{k,i})$ gives us for all $j \in \{d+1, \ldots, 2d\}, \widetilde{\mu}'(t)_j = 0$ and for all $j \in \{1, \ldots, d\}$,

$$\widetilde{\mu}'(t)_j = (\widetilde{\mu}(t_1) \otimes \ldots \otimes \widetilde{\mu}(t_{i-1}) \otimes \widetilde{\mu}(t')) \cdot M_j^{k,i}(||t||, ||t_1||, \ldots, ||t_i||)$$
$$= (\widetilde{\mu}(t_1) \otimes \ldots \otimes \widetilde{\mu}(t_{i-1}) \otimes \widetilde{\mu}(t'_i)) \cdot$$
$$(I_{d^i} \otimes \otimes_{\ell=i+1}^k \mu(g_0)) \cdot \mu(g_k)(||t'||, ||t_1||, \ldots, ||t_{i-1}||, ||t'_i||, 0, \ldots, 0)_j$$

Notice that $A = (\tilde{\mu}(t_1) \otimes \ldots \otimes \tilde{\mu}(t_{i-1}) \otimes \tilde{\mu}(t'_i)) \in \mathbb{Q}^{1 \times d^i}$ and $B = \bigotimes_{\ell=i+1}^k \mu(g_0) \in \mathbb{Q}^{1 \times d^{k-1}}$. Hence, by the mixed-product property of the Kronecker product, we obtain that $A(I_{d^i} \otimes B) = (AI_{d^i}) \otimes (I_1B) = A \otimes B$. Therefore:

$$\widetilde{\mu}'(t)_j = (\widetilde{\mu}(t_1) \otimes \ldots \otimes \widetilde{\mu}(t_{i-1}) \otimes \widetilde{\mu}(t'_i) \otimes \bigotimes_{\ell=i+1}^k \mu(g_0)) \cdot \mu(g_k)(||t'||, ||t_1||, \ldots, ||t_{i-1}||, ||t'_i||, 0, \ldots, 0)_j$$
$$= \widetilde{\mu}(g_k(t_1, \ldots, t_{i-1}, t'_i, g_0, \ldots, g_0))_j$$

Notice that by definition of Γ , we have $\Gamma(t') = t$ which allows us to conclude that $t' \in T_{\Sigma}$, and $t = \Gamma(t')$, and $\tilde{\mu}'(t) = \begin{bmatrix} \tilde{\mu}(t') & \mathbf{0}_{d \times 1} \end{bmatrix}$.

• Case $t = h_{k,i}(t_1, \ldots, t_i)$ and either $t_{\ell} \notin T_{\Sigma}$ with $\ell \in \{1, \ldots, i-1\}$ or for all $t'_i \in T_{\Sigma}$, $\Gamma(t'_i) \neq t_i$: In the former case, by inductive hypothesis on t_{ℓ} , we deduce that for all $j \in \{d+1, \ldots, 2d\}$, $\tilde{\mu}'(t_{\ell})_j = 0$. In the latter case, applying our inductive hypothesis on t_i gives us that for all $j \in \{1, \ldots, d\}$, $\tilde{\mu}'(t_i)_j = 0$. In both cases, we deduce from the definition of $\mu'(h_{k,i})$ that

$$(\widetilde{\mu}'(t_1)\otimes\ldots\otimes\widetilde{\mu}'(t_i))\mu'(h_{k,i})(||t||,||t_1||,\ldots,||t_i||)=\mathbf{0}_{1\times 2d}$$

This conclude the proof of the desired property. Let us now conclude the main result by computing the power series. Thanks to our desired property, notice that for all $t \in T_{\Sigma'}$, $\mathcal{A}'[\![t]\!] \neq 0$ implies that there exists $t' \in T_{\Sigma}$ such that $\Gamma(t') = t$ and $\mathcal{A}'[\![t]\!] = \mathcal{A}[\![t']\!]$. As Γ is injective, we deduce that for all $n \in \mathbb{N}$,

$$\sum_{\substack{t \in T_{\Sigma'} \\ |t\|=n}} \mathcal{A}'\llbracket t \rrbracket = \sum_{\substack{t \in T_{\Sigma'}, \|t\|=n \\ \exists t' \in T_{\Sigma}. \Gamma(t')=t}} \mathcal{A}'\llbracket t \rrbracket = \sum_{\substack{t' \in T_{\Sigma} \\ \|t'\|=n+1}} \mathcal{A}\llbracket t'\rrbracket$$

which allows us to conclude.

Corollary 3. Let \mathcal{A} be a holonomic tree automata. Let $f_{\mathcal{A}}(x) = \sum_{n=0}^{\infty} a_n x^n$.

- There exists a holonomic tree automaton \mathcal{A}' such that $f_{\mathcal{A}'}(x) = f'_{\mathcal{A}}(x)$.
- There exists a holonomic tree automaton \mathcal{A}' such that $f_{\mathcal{A}'}(x) = \int_0^x f_{\mathcal{A}}(x) dx$.

Proof. From Proposition 14, we have a $f_{\mathcal{A}''}(x) = x f'_{\mathcal{A}}(x)$ for some \mathcal{A}'' and $f_{\mathcal{A}''}(x)$ has no constant term. Hence by Proposition 17, we have $f_{\mathcal{A}'}(x) = \frac{f_{\mathcal{A}''}(x)}{x} = f'_{\mathcal{A}}(x)$. From Propositions 15 and 16, we directly obtain that $f_{\mathcal{A}'}(x) = \int_0^x f_{\mathcal{A}}(x) dx$ for some \mathcal{A}' .

Corollary 4. Let \mathcal{A}_1 , \mathcal{A}_2 be two holonomic tree automata. There exists a holonomic tree automatom \mathcal{A} such that $f_{\mathcal{A}}(x) = f_{\mathcal{A}_1}(x) \cdot f_{\mathcal{A}_2}(x)$.

Proof. Direct from Propositions 11 and 17.

Proposition 18. Let \mathcal{A} be a holonomic tree automaton. Let $f_{\mathcal{A}}(x) = \sum_{n=0}^{\infty} a_n x^n$. If $a_0 \neq 0$ then there exists a holonomic tree automaton \mathcal{A}' such that $f_{\mathcal{A}'}(x) = \frac{1}{f_{\mathcal{A}}(x)}$.

Proof. It is well known that when $a_0 \neq 0$, $f_A(x)$ admits an multiplicative inverse power series $f_A^{-1}(x) = \sum_{n=0}^{\infty} b_n x^n$ where the coefficients of $f_A^{-1}(x)$ satisfy the following property:

$$\begin{cases} b_0 = \frac{1}{a_0} \\ b_n = -\frac{1}{a_0} \sum_{i=0}^{n-1} a_{i+1} b_{n-1-i} \quad \forall n > 0 \end{cases}$$

Using Proposition 17, we first build the automaton \mathcal{A}'' such that $f_{\mathcal{A}''}(x) = \sum_{n=0}^{\infty} a_{n+1}x^n$. By Proposition 12, we can assume that $\mathcal{A}'' = (d, \mu)$ over some alphabet Σ that is arity distinct. Let us denote $\mathbf{f}(x) = \sum_{n=0} \mathbf{c}_n x^n$ the vector of power series corresponding to \mathcal{A}'' , that is $\mathbf{c}_n = \sum_{\substack{t \in T_{\Sigma} \\ \|t\| = n}} \mu(t)$. Notice that $\mathbf{c}_{n,1} = a_{n+1}$ for $n \in \mathbb{N}$.

We build the automaton $\mathcal{A}' = (d', \mu')$ over $\Sigma' = \Sigma \cup \{u\}$ with d' = d + 1, and u a fresh binary function symbol and the weight function μ' such that for all $n \ge 0$, $\mathbf{a}'_n = \sum_{\substack{t \in T_{\Sigma'} \\ \|t\| = n}} \mu'(t) = \begin{bmatrix} b_n & \mathbf{c}_n \end{bmatrix}$.

For that purpose, let us define μ' such that

- $\mu'(a) = \begin{bmatrix} \frac{1}{a_0} & \mu(a) \end{bmatrix}$ where *a* is the unique nullary function symbol in Σ (recall that Σ is arity distinct)
- for all k > 0, for all $g \in \Sigma_k$, for all $i_1, \ldots, i_k, j \in \{1, \ldots, d+1\}$,

$$\mu'(g)_{(i_1,\dots,i_k),j} = \begin{cases} \mu(g)_{(i_1-1,\dots,i_k-1),j-1} & \text{if } \forall \ell.i_\ell, j \in \{2,\dots,d+1\} \\ 0 & \text{otherwise} \end{cases}$$

•

$$\mu'(u) = \begin{bmatrix} 0 & \mathbf{0}_{1 \times d} \\ -\frac{1}{a_0} & \mathbf{0}_{1 \times d} \\ \mathbf{0}_{(d'^2 - 2) \times 1} & \mathbf{0}_{(d'^2 - 2) \times d} \end{bmatrix}$$

In the inductive step n > 0, from Equation (6) and from our inductive hypothesis, we have

$$\begin{aligned} \mathbf{a}_{n}' &= \sum_{\substack{k>0\\g\in\Sigma_{k}'}} \sum_{\substack{n_{1},\dots,n_{k}\in\mathbb{N}\\n_{1}+\dots,n_{k}=n-1}} (\mathbf{a}_{n_{1}}'\otimes\dots\otimes\mathbf{a}_{n_{k}}')\mu'(g)(n,n_{1},\dots,n_{k}) \\ &= \sum_{\substack{k>0\\g\in\Sigma_{k}'}} \sum_{\substack{n_{1},\dots,n_{k}\in\mathbb{N}\\n_{1}+\dots,n_{k}=n-1}} ([b_{n_{1}} \ \mathbf{c}_{n_{1}}]\otimes\dots\otimes[b_{n_{k}} \ \mathbf{c}_{n_{k}}])\mu'(g)(n,n_{1},\dots,n_{k}) \\ &= \sum_{\substack{k>0\\g\in\Sigma_{k}}} \sum_{\substack{n_{1},\dots,n_{k}\in\mathbb{N}\\n_{1}+\dots,n_{k}=n-1}} ([b_{n_{1}} \ \mathbf{c}_{n_{1}}]\otimes\dots\otimes[b_{n_{k}} \ \mathbf{c}_{n_{k}}])\mu'(g)(n,n_{1},\dots,n_{k}) \\ &+ \sum_{i=0}^{n-1} ([b_{i} \ \mathbf{c}_{i}]\otimes[b_{n-1-i} \ \mathbf{c}_{n-1-i}])\mu'(u)(n,i,n-1-i) \end{aligned}$$

Following the definition of $\mu'(u)(x_0, x_1, x_2)$, we deduce that

$$(\begin{bmatrix} b_i & c_i \end{bmatrix} \otimes \begin{bmatrix} b_{n-1-i} & c_{n-1-i} \end{bmatrix}) \mu'(u)(n,i,n-1-i) = \begin{bmatrix} -\frac{1}{a_0} c_{i,1} & b_{n-1-i} & \mathbf{0}_{1\times d} \end{bmatrix} = \begin{bmatrix} -\frac{a_{n+1}b_{n-1-i}}{a_0} & \mathbf{0}_{1\times d} \end{bmatrix}$$

Moreover, by definition of $\mu'(g)(x_0, \ldots, x_k)$ for all k > 0, for all $g \in \Sigma_k$, we deduce that:

$$(\begin{bmatrix} b_{n_1} & \boldsymbol{c}_{n_1} \end{bmatrix} \otimes \ldots \otimes \begin{bmatrix} b_{n_k} & \boldsymbol{c}_{n_k} \end{bmatrix}) \mu'(g)(n, n_1, \dots, n_k) = \begin{bmatrix} 0 & (\boldsymbol{c}_1 \otimes \ldots \otimes \boldsymbol{c}_k) \mu(g)(n, n_1, \dots, n_k) \end{bmatrix}$$

We therefore obtain:

$$\begin{aligned} \boldsymbol{a}_{n}^{\prime} &= \sum_{\substack{k>0\\g\in\Sigma_{k}}} \sum_{\substack{n_{1},\dots,n_{k}\in\mathbb{N}\\n_{1}+\dots+n_{k}=n-1}} \begin{bmatrix} 0 & (\boldsymbol{c}_{1}\otimes\dots\otimes\boldsymbol{c}_{k})\mu(g)(n,n_{1},\dots,n_{k}) \end{bmatrix} + \sum_{i=0}^{n-1} \begin{bmatrix} -\frac{a_{n+1}b_{n-1-i}}{a_{0}} & \boldsymbol{0}_{1\times d} \end{bmatrix} \\ &= \begin{bmatrix} 0 & \boldsymbol{c}_{n} \end{bmatrix} + \begin{bmatrix} b_{n} & \boldsymbol{0}_{1\times d} \end{bmatrix} \\ &= \begin{bmatrix} b_{n} & \boldsymbol{c}_{n} \end{bmatrix} \end{aligned}$$

As for all $n \ge 0$, $a'_{n,1} = b_n$, we conclude that $f_{\mathcal{A}'}(x) = f_{\mathcal{A}}^{-1}(x)$.