

# The Memory of Primordial Gravitational Waves

Paolo Creminelli<sup>a,b</sup><sup>1</sup> and Filippo Vernizzi<sup>c</sup><sup>2</sup>

<sup>a</sup> *ICTP, International Centre for Theoretical Physics  
Strada Costiera 11, 34151, Trieste, Italy*

<sup>b</sup> *IFPU - Institute for Fundamental Physics of the Universe,  
Via Beirut 2, 34014, Trieste, Italy*

<sup>c</sup> *Institut de Physique Théorique, Université Paris Saclay,  
CEA, CNRS, 91191 Gif-sur-Yvette, France*

## Abstract

Primordial gravitational waves, after they enter the horizon and decay away, leave a residual displacement in test particles: a memory, in analogy with gravitational waves generated by astrophysical sources. The displacement in the distance between test particles  $\xi^i$  does not depend on the cosmological evolution, but only on the primordial value  $\bar{h}_j^i$  of the gravitational wave:  $\Delta\xi^i = -\frac{1}{2}\bar{h}_j^i\xi^j$ . The memory is thus related to the adiabatic tensor mode that maps the unperturbed FLRW geometries at early and late times; this is analogous to the relation between memory in Minkowski spacetime and the BMS group. The primordial memory is also connected to the consistency relations of cosmological correlators, as the flat-space memory is related to the soft theorems for gravitational wave emission. We comment on the signature of the effect on the CMB  $B$ -modes and on the large-scale structure. There is also a primordial memory effect that is subleading in the spatial gradients of the wave: it is encoded in the rotation of free-falling gyroscopes.

*To the memory of Valery A. Rubakov.*

*Contribution to the special issue of the International Journal of Modern Physics A.*

---

<sup>1</sup>Email: [creminel@ictp.it](mailto:creminel@ictp.it)

<sup>2</sup>Email: [filippo.vernizzi@ipht.fr](mailto:filippo.vernizzi@ipht.fr)

The memory effect of gravitational waves (GWs) is the residual displacement of test particles after the passage of the wave. GWs emitted by astrophysical sources give a memory effect and this phenomenon has been the subject of intense studies starting from [1] (for a review, see [2]). The memory can be measured by present or planned GW experiments, see for instance [3, 4, 5]. Theoretically, gravitational memory is a crucial feature of the infrared (IR) limit of gravity [6] and, as such, it is deeply connected with the asymptotic symmetries of Minkowski space and soft theorems for the emission of low-energy gravitons (for a review of the subject see Strominger's lectures [7]).

Here we point out that the same effect exists for cosmological GWs generated in the early Universe. During inflation, GWs are produced from quantum vacuum fluctuations. The homogeneous expanding Universe is described by a Friedmann-Lemaître-Robertson-Walker (FLRW) metric,  $ds^2 = a^2(\eta) (-d\eta^2 + d\vec{x}^2)$ , where  $\eta$  is the conformal time and  $x^i$  are comoving spatial coordinates. The cosmic expansion rate is set by the Hubble parameter  $H \equiv \frac{1}{a^2} \frac{da}{d\eta}$ . Consider a Fourier mode of physical wavenumber  $k_{\text{phys}}(\eta) = k/a(\eta)$ , where  $k$  is a constant *comoving* wavenumber. When  $k_{\text{phys}}$  is stretched out of the horizon by the accelerated expansion, i.e., for  $k_{\text{phys}} \ll H$ , GWs freeze to a constant value. In this regime they are unobservable because they locally correspond to an anisotropic rescaling of the coordinates. Later on, during the decelerated expansion of the Universe, they come back into the horizon,  $k_{\text{phys}} \gtrsim H$ , and oscillate. In this regime they affect cosmological observables, most notably the  $B$ -mode polarisation of the Cosmic Microwave Background (CMB) anisotropies [8, 9], which is the target of a massive observational program; see e.g. [10, 11, 12]. As the Universe expands, the amplitude of their oscillations decays and they disappear at late times. One can thus ask the question: is there a residual displacement of test particles that were initially at rest before the GWs came back into the horizon? Is there a memory of the primordial GWs?

The calculation to the answer is very simple using the usual transverse-traceless (TT) gauge for GWs in a FLRW background. The metric in this case reads

$$ds^2 = a^2(\eta) [-d\eta^2 + (\delta_{ij} + h_{ij}(\eta, \vec{x})) dx^i dx^j] , \quad (1)$$

where the tensor perturbation  $h_{ij}$  is transverse,  $\partial_i h_{ij} = 0$ , and traceless,  $h^i_i = 0$ . At linear order in  $h_{ij}(\eta, \vec{x})$ , each of its Fourier modes are independent and we can focus on a single one, evolving as  $\tilde{h}_{ij}(\eta, \vec{k}) = \bar{h}_{ij}(\vec{k}) F(\eta, k)$ , where  $F(\eta, k)$  is the transfer function and  $\bar{h}_{ij}(\vec{k})$  the initial condition at early times, i.e. for  $k \ll aH$  or, equivalently,  $k\eta \ll 1$ . The time evolution of a mode depends on the cosmological evolution  $a(\eta)$  during the decelerated expansion; however, in general one has  $F \rightarrow 1$  at early times and  $F \rightarrow 0$  at late times (see Fig. 1).

Notice that only the spatial part of the metric is perturbed. This implies, in the same way as in the treatment of GWs around Minkowski spacetime, that in this gauge test particles that are initially at rest remain at constant spatial coordinates. Indeed, the geodesic equation reads

$$\frac{d^2 x^i}{d\tau^2} = -\Gamma_{\mu\nu}^i \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} = -\Gamma_{00}^i \frac{d\eta}{d\tau} \frac{d\eta}{d\tau} = 0 , \quad (2)$$

where the second equality holds since there is no initial velocity and the third because  $\Gamma_{00}^i = 0$  at linear order in  $h_{ij}$ . Therefore, test particles keep the same spatial coordinates in this gauge.




Figure 1: Time evolution of a Fourier mode of cosmological GWs. Here we assume that the Universe is matter dominated, in which case  $F(\eta, k) = 3[\sin(k\eta)/(k\eta) - \cos(k\eta)]/(k\eta)^2$ , with  $k = |\vec{k}|$ ; however, the evolution is qualitatively the same for other decelerating cosmologies. The offset between early and late times is responsible of the memory effect. Notice how this is much larger than the late-time oscillations.

If we now compare around a given point the *physical* distance  $\vec{\xi}$  between test masses at early times, when the GW is frozen outside the horizon, and much later, when it has decayed to zero, one finds (see Fig. 2)

$$\Delta\xi^i = \xi_{\text{late}}^i - \xi_{\text{early}}^i = -\frac{1}{2}\bar{h}_{ij}\xi^j. \quad (3)$$

This is the memory of the primordial GWs. To derive this equation we have assumed that the distance between the test particles is much smaller than the GW wavelength. Notice also that even if the GW time-dependence  $F(\eta, k)$ , and therefore its effect on test particles and cosmological observables, depends on the cosmological history  $a(\eta)$ , the residual displacement does not: *the memory of primordial GWs does not depend on the cosmological evolution*.<sup>3</sup> As evident from figure 1, the memory of primordial GWs is large compared to the late-time oscillations. This is at variance with the memory of astrophysical binary sources, which is tiny compared to the oscillations in the waveform, since it is only a non-linear effect [15].

Of course, it is impossible to go back in the early Universe and measure the position of a given test mass to compare it with its final one today. But if we look at many test masses, we can compare their initial and final positions *statistically*, assuming that they were homogeneously and isotropically distributed at early time. For instance, one can observe the effect of the memory of primordial GWs in the correlation function of the dark matter

---

<sup>3</sup>The memory of primordial GWs, which is the topic of interest here, should not be confused with the modification of the usual Minkowski memory when the source and the detectors are in a FLRW universe (see [13, 14] and references therein).




Figure 2: At early times (left figure), the GW is much longer than the comoving Hubble horizon  $(aH)^{-1}_{\text{early}}$ , i.e.  $k \ll (aH)_{\text{early}}$ , and test particles are distributed in a spherical shell. At late times (right), the GW is much shorter than the horizon,  $k \gg (aH)_{\text{late}}$ , and has decayed away. The shell of particles is left permanently deformed to an ellipsoid.




Figure 3: Cosmological Infrared Triangle: Triangle of relations in the cosmological setting (with corners Primordial Memory, Consistency Relation, Adiabatic Mode), analogue of the Infrared Triangle (with corners Memory Effect, Soft Theorem, Asymptotic Symmetry) discussed in flat space [7].

distribution. The effect of a primordial GW on short-scale density perturbations was studied in [16, 17] (see also [18]), including the displacement of test particles as mapping between Lagrangian and Eulerian coordinates, although these references did not interpret the final displacement as a memory effect. We will discuss observational signatures below.

In flat spacetime, the memory of GWs is connected to two other aspects of gravity in the IR

limit: the asymptotic Bondi-Metzner-Sachs (BMS) symmetries [19, 20] and the soft theorems for GW emission [21]. These three concepts and their relative connection can be generalized to other massless theories and they have been the object of a fervent research activity, with the hope of obtaining a deeper understanding of gauge and gravitational theories, see e.g. [22] for a recent review. In the gravitational case, BMS symmetries, soft theorems and memory effects form a “triangle”, as discussed in [7]. Here we point out that, similarly, the memory of primordial GWs is part of an analogous triangle, shown in Fig. 3, as explained below (see also [23]).

Both at early times, when the GW is frozen outside the horizon, and at late times, after it has decayed away, the wave has no measurable effect and the metric is the same as an unperturbed FLRW one. However, the two unperturbed metrics are written in a different way with a diffeomorphism relating the two,

$$x_{\text{late}}^i = \left( \delta_j^i + \frac{1}{2} \bar{h}_j^i \right) x_{\text{early}}^j . \quad (4)$$

The memory effect is a measure of this “mismatch” of the coordinates. This is in complete analogy with what happens in flat space. The flat space before and after the passage of the GW are related by a diffeomorphism, a BMS transformation, and the memory effect is a measure of this mapping. Cosmological perturbations that reduce to a pure diffeomorphism in the long-wavelength limit are called adiabatic [24]. Adiabaticity has important implications: for instance it implies that  $h_{ij}$  is constant in time, independently of the evolution of the Universe. Notice that the memory effect does not depend on the production mechanism (classical or quantum) of GWs during inflation. What matters is that GWs get frozen out of the cosmological horizon during the accelerated expansion of the Universe and become adiabatic.<sup>4</sup>

Adiabatic modes are related to the third corner of the triangle: the consistency relations of cosmological perturbations. These originate from the adiabaticity of the modes *during* inflation. Let us imagine we want to calculate the correlation between a long GW (with polarisation  $s$  and polarisation tensor  $\epsilon_{ij}^s$ ) and short scalar perturbations described by the variable  $\zeta$ . The long tensor mode is already frozen outside the horizon during inflation when the short scalar modes are generated. Therefore there is no physical effect of the GW apart from the diffeomorphism of eq. (4). This logic gives a universal relation in this limit of the form [26]

$$\langle h_{\vec{k}_1}^s \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \rangle' \simeq - \langle h_{\vec{k}_1}^s h_{-\vec{k}_1}^s \rangle' \epsilon_{ij}^s k_2^i k_2^j \frac{\partial}{\partial k_2^2} \langle \zeta_{\vec{k}_3} \zeta_{\vec{k}_2} \rangle' , \quad (5)$$

where the prime denotes that we have stripped the momentum conserving delta function off the statistical averages  $\langle \dots \rangle$ . This relation shows the effect in Fourier space of the anisotropic rescaling of the coordinates.<sup>5</sup>

<sup>4</sup>A memory effect associated with the quantum production of the GWs *during* inflation has been studied in [25], although it is not an observable quantity.

<sup>5</sup>Consistency relations and adiabatic modes exist also for scalar perturbations. However, since a scalar perturbation does not decay away when entering the horizon, one cannot define a memory effect.

Not only do consistency relations apply during inflation, but they can be extended to observable quantities by including the late-time effect of the long mode on the short ones, see for instance [27, 28, 29, 30]. To show the link between consistency relations and memory, let us consider for example the baryon acoustic oscillations (BAOs), i.e. the fluctuations in the density of matter due to acoustic waves in the primordial plasma. The BAO peak can be crudely represented as a thin spherical shell of radius  $r_{\text{BAO}}$  around a given galaxy, where it is more likely to find another galaxy. (Notice the analogy with the spherical shell of inertial detectors of Fig. 2.) What happens to the BAO shell in the presence of a GW with wavelength much longer than  $r_{\text{BAO}}$ ? Since the scalar modes giving rise to the BAOs are much shorter than the GW, they are generated during inflation when the GW is already frozen outside the horizon. The BAO is thus a sphere with constant physical radius when the GW is outside the horizon. When the GW comes back and fade away, the BAO shell will be distorted to an ellipsoid due to the memory effect. This represents a physical effect associated with the consistency relation eq. (5).

The effect of anisotropic distortion due to the GW on the late-time dark matter density contrast  $\delta$  can be computed rescaling the coordinates using eq. (4) and reads [17]

$$\delta(\eta, \vec{x})|_h = \left(1 + \frac{1}{2}\bar{h}_{ij}x^i\partial^j\right)\delta(\eta, \vec{x}). \quad (6)$$

The same formula applies to the galaxy number over-density or to the overdensity of other cosmological tracers. The GW modifies the small-scale correlation between dark matter particles or galaxies. This effect can be compared to that of the gravitational tidal field caused by dark matter itself, responsible for the deformation of the BAO sphere. It is obtained by replacing  $\bar{h}_{ij}$  in the above equation by  $\partial_i\partial_j\nabla^{-2}\delta$ : this effect is therefore much larger than the one of the memory of primordial GWs.

Let us comment on the possibility of measuring the memory of primordial GWs in galaxy surveys by observing its effect on the alignment of the shapes of the observed galaxies: the so-called intrinsic alignment.<sup>6</sup> Although small, this effect can be measured in the *B*-mode of the weak lensing shear. The memory of the primordial GWs contributes to the intrinsic alignment, since the matter that will give rise to a certain galaxy is displaced following eq. (6). Going beyond the test particle limit and taking into account the gravitational interaction of matter, GWs change the growth on short scales: this tidal interaction is an effect of the same order as the memory, but with a different scale dependence and with an amplitude that depends on the cosmological evolution [16, 17, 33]. It would be nice to study the separability of the two effects on the intrinsic alignment or to explore the possibility of detecting the primordial memory effect using other cosmological probes.

Notably, the polarization of the CMB is affected by the primordial memory. Indeed, as we discussed, the most direct way of detecting primordial GWs is by observing *B*-modes of the CMB. Their spectrum can be read from the book of Gorbunov and Rubakov [34] and is

---

<sup>6</sup>The second source of alignment is the small deflection of light due to foreground dark matter distribution: the weak lensing. Light is also deflected by the presence of GWs, although this is a very tiny effect [31, 32].

given by [see eq. (10.83) of that reference]

$$C_\ell^{BB} = \frac{4\pi}{25} \Delta\eta_r^2 \int_0^\infty \frac{dk}{k} \mathcal{P}_T(k) |F'(k, \eta)|_{\eta=\eta_r}^2 \left[ \frac{\ell+2}{2\ell+1} j_{\ell-1}(k\eta_0) - \frac{\ell-1}{2\ell+1} j_{\ell+1}(k\eta_0) \right]^2, \quad (7)$$

where  $\Delta\eta_r$  is the thickness of recombination surface,  $\mathcal{P}_T(k)$  is the power spectrum of the primordial tensor modes  $\bar{h}_{ij}$  and  $F' \equiv \partial F / \partial \eta$ . This observable is sensitive to the derivative of  $F$  with respect to  $\eta$  evaluated at recombination,  $\eta = \eta_r$ . Therefore, it is not directly sensitive to the memory of primordial gravitational waves. However, with sufficient precision this observation can still be used to reconstruct the variation of  $F$  from early to late time and measure the memory of the waves. Indeed, while  $C_\ell^{BB}$  depends on  $F'$  at  $\eta_r$ , its  $\eta$ -dependence can be reconstructed from its  $k$ -dependence. For instance, in matter domination  $F$  is a function of the combination  $k\eta$ .<sup>7</sup> As a matter of fact, since the first peak of the  $B$ -mode power spectrum is much larger than the others, one can see “by eye” that the wave is not oscillating around zero. Notice that the detection of the primordial memory requires to be able to compare moments when the GW is outside and inside the horizon. This is clearly not possible for interferometers, even in inflationary models with a detectable GW spectrum.

A new kind of gravitational memory in Minkowski spacetime, dubbed “Spin Memory”, has been recently proposed [35], consisting in the relative time delay acquired by a Sagnac interferometer after the passage of a GW. In analogy to the standard gravitational memory, this effect is deeply connected to extensions of the asymptotic symmetries of Minkowski spacetime [36] and to soft-graviton theorems at *subleading* order in the soft expansion [37].

Also the primordial memory can be extended to subleading order. Imagine that the test particles previously considered are replaced by gyroscopes. These gyroscopes are initially aligned when the GW is well outside the Hubble radius. We want to show that the gyroscopes are not aligned anymore after the GW has decayed away and that this memory is sensitive to the derivative of the initial value of the wave,  $\partial_k \bar{h}_{ij}$ . The gyroscopes are described by space-like 4-vectors,  $S^\nu$ , which are parallel-transported along time-like geodesics with tangent vector  $V^\mu$ . This gives

$$0 = V^\mu \nabla_\mu S^i = \partial_0 S^i + \Gamma_{0\alpha}^i S^\alpha = \dot{S}^i + \frac{1}{2} \dot{h}_{ij} S^j. \quad (8)$$

In the second equality we specialized to the geodesics that are initially at rest: in the coordinates we are using their tangent vector is simply  $V^\mu = (1, \vec{0})$ . In the third equality we used that the 4-vectors  $V^\mu$  and  $S^\mu$  are orthogonal, since they are orthogonal at the beginning and then both parallelly transported. We have also evaluated the Christoffel’s symbol in our gauge, eq. (1).

This equation does not describe any rotation of the gyroscopes, but simply a rescaling, which is necessary to keep the norm of the vector constant. For simplicity, let us assume that  $\bar{h}_{ij}$  vanishes at the position of the experiment. The equation above then says that the spatial

---

<sup>7</sup>Since the power spectrum is sensitive to  $|F'|^2$ , and not directly to  $F'$ , there is a sign “ambiguity” in reconstructing the derivative.

derivatives  $\partial_k \bar{h}_{ij}$  do not induce a change of  $S^i$ : the vectors  $S^i$  remain constant in this gauge, like the coordinates of the geodesics in the previous discussion. However this does not mean there is no physical rotation of the gyroscopes. At early times, in order to start with parallel gyroscopes, one has to perform a parallel transport on a surface of constant time<sup>8</sup>. To move in a spatial direction one has to parallel transport the vector  $S^i$  along a spatial vector  $W^j$ , i.e. solve  $W^j(\partial_j S^i + \Gamma_{jk}^i S^k) = 0$ . This gives the expression of parallel gyroscopes around a central one  $S_{(0)}^i$

$$S^i = S_{(0)}^i + x^j(-\Gamma_{jk}^i)S_{(0)}^k , \quad (9)$$

at first order in  $x^i$ . When the GW has decayed away, the spatial metric is just  $\delta_{ij}$  and the gyroscopes are clearly not aligned. Writing the Christoffel's symbols in terms of  $\bar{h}_{ij}$ , one gets

$$\Delta S^i = -\frac{1}{2}(\partial_j \bar{h}_{ik} + \partial_k \bar{h}_{ij} - \partial_i \bar{h}_{jk})x^j S^k . \quad (10)$$

The memory of the direction of the gyroscopes is encoded in the spatial derivative of the GW.

Adiabatic modes and consistency relations were studied at subleading order in the soft limit, see for instance [38, 39]: therefore the whole triangle exists also at subleading order, in analogy with the Minkowski case.

**Acknowledgements:** We thank Laura Donnay, Marc Favata, Luca Santoni, Fabian Schmidt and Andy Strominger for useful discussions. We thank Fabian Schmidt for comments on the draft.

## References

- [1] Y. B. Zel'dovich and A. G. Polnarev, “Radiation of gravitational waves by a cluster of superdense stars,” *Sov. Astron.* **18** (1974) 17.
- [2] M. Favata, “The gravitational-wave memory effect,” *Class. Quant. Grav.* **27** (2010) 084036, 1003.3486.
- [3] P. D. Lasky, E. Thrane, Y. Levin, J. Blackman, and Y. Chen, “Detecting gravitational-wave memory with LIGO: implications of GW150914,” *Phys. Rev. Lett.* **117** (2016), no. 6 061102, 1605.01415.
- [4] D. A. Nichols, “Spin memory effect for compact binaries in the post-Newtonian approximation,” *Phys. Rev. D* **95** (2017), no. 8 084048, 1702.03300.
- [5] H. Inchauspé, S. Gasparotto, D. Blas, L. Heisenberg, J. Zosso, and S. Tiwari, “Measuring gravitational wave memory with LISA,” 2406.09228.
- [6] A. Strominger and A. Zhiboedov, “Gravitational Memory, BMS Supertranslations and Soft Theorems,” *JHEP* **01** (2016) 086, 1411.5745.
- [7] A. Strominger, “Lectures on the Infrared Structure of Gravity and Gauge Theory,” 1703.05448.
- [8] U. Seljak and M. Zaldarriaga, “Signature of gravity waves in polarization of the microwave background,” *Phys. Rev. Lett.* **78** (1997) 2054–2057, astro-ph/9609169.

---

<sup>8</sup>Since we are only considering first derivatives of the metric, the space is flat and the parallel transport does not depend on the path.

- [9] M. Kamionkowski, A. Kosowsky, and A. Stebbins, “A Probe of primordial gravity waves and vorticity,” *Phys. Rev. Lett.* **78** (1997) 2058–2061, [astro-ph/9609132](#).
- [10] **BICEP, Keck** Collaboration, P. A. R. Ade *et. al.*, “Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations through the 2018 Observing Season,” *Phys. Rev. Lett.* **127** (2021), no. 15 151301, [2110.00483](#).
- [11] **CMB-S4** Collaboration, K. Abazajian *et. al.*, “Snowmass 2021 CMB-S4 White Paper,” [2203.08024](#).
- [12] **LiteBIRD** Collaboration, M. Hazumi *et. al.*, “LiteBIRD: JAXA’s new strategic L-class mission for all-sky surveys of cosmic microwave background polarization,” *Proc. SPIE Int. Soc. Opt. Eng.* **11443** (2020) 114432F, [2101.12449](#).
- [13] L. Bieri, D. Garfinkle, and S.-T. Yau, “Gravitational wave memory in de Sitter spacetime,” *Phys. Rev. D* **94** (2016), no. 6 064040, [1509.01296](#).
- [14] A. Tolish and R. M. Wald, “Cosmological memory effect,” *Phys. Rev. D* **94** (2016), no. 4 044009, [1606.04894](#).
- [15] D. Christodoulou, “Nonlinear nature of gravitation and gravitational wave experiments,” *Phys. Rev. Lett.* **67** (1991) 1486–1489.
- [16] L. Dai, D. Jeong, and M. Kamionkowski, “Anisotropic imprint of long-wavelength tensor perturbations on cosmic structure,” *Phys. Rev. D* **88** (2013), no. 4 043507, [1306.3985](#).
- [17] F. Schmidt, E. Pajer, and M. Zaldarriaga, “Large-Scale Structure and Gravitational Waves III: Tidal Effects,” *Phys. Rev. D* **89** (2014), no. 8 083507, [1312.5616](#).
- [18] K. W. Masui and U.-L. Pen, “Primordial gravity wave fossils and their use in testing inflation,” *Phys. Rev. Lett.* **105** (2010) 161302, [1006.4181](#).
- [19] H. Bondi, M. G. J. van der Burg, and A. W. K. Metzner, “Gravitational waves in general relativity. 7. Waves from axisymmetric isolated systems,” *Proc. Roy. Soc. Lond. A* **269** (1962) 21–52.
- [20] R. K. Sachs, “Gravitational waves in general relativity. 8. Waves in asymptotically flat space-times,” *Proc. Roy. Soc. Lond. A* **270** (1962) 103–126.
- [21] S. Weinberg, “Infrared photons and gravitons,” *Phys. Rev.* **140** (1965) B516–B524.
- [22] T. McLoughlin, A. Puhm, and A.-M. Raclariu, “The SAGEX Review on Scattering Amplitudes, Chapter 11: Soft Theorems and Celestial Amplitudes,” [2203.13022](#).
- [23] M. Mirbabayi and M. Simonović, “Weinberg Soft Theorems from Weinberg Adiabatic Modes,” [1602.05196](#).
- [24] S. Weinberg, “Adiabatic modes in cosmology,” *Phys. Rev. D* **67** (2003) 123504, [astro-ph/0302326](#).
- [25] R. Z. Ferreira, M. Sandora, and M. S. Sloth, “Asymptotic Symmetries in de Sitter and Inflationary Spacetimes,” *JCAP* **04** (2017) 033, [1609.06318](#).
- [26] J. M. Maldacena, “Non-Gaussian features of primordial fluctuations in single field inflationary models,” *JHEP* **0305** (2003) 013, [astro-ph/0210603](#).
- [27] P. Creminelli, C. Pitrou, and F. Vernizzi, “The CMB bispectrum in the squeezed limit,” *JCAP* **11** (2011) 025, [1109.1822](#).
- [28] Z. Huang and F. Vernizzi, “Cosmic Microwave Background Bispectrum from Recombination,” *Phys. Rev. Lett.* **110** (2013), no. 10 101303, [1212.3573](#).
- [29] M. Peloso and M. Pietroni, “Galilean invariance and the consistency relation for the nonlinear squeezed bispectrum of large scale structure,” *JCAP* **05** (2013) 031, [1302.0223](#).

- [30] P. Creminelli, J. Noreña, M. Simonović, and F. Vernizzi, “Single-Field Consistency Relations of Large Scale Structure,” *JCAP* **12** (2013) 025, 1309.3557.
- [31] S. Dodelson, E. Rozo, and A. Stebbins, “Primordial gravity waves and weak lensing,” *Phys. Rev. Lett.* **91** (2003) 021301, astro-ph/0301177.
- [32] F. Schmidt and D. Jeong, “Large-Scale Structure with Gravitational Waves II: Shear,” *Phys. Rev. D* **86** (2012) 083513, 1205.1514.
- [33] K. Akitsu, Y. Li, and T. Okumura, “Gravitational wave fossils in nonlinear regime: Halo tidal bias and intrinsic alignments from gravitational wave separate universe simulations,” *Phys. Rev. D* **107** (2023), no. 6 063531, 2209.06226.
- [34] D. S. Gorbunov and V. A. Rubakov, *Introduction to the theory of the early universe: Cosmological perturbations and inflationary theory*. 2011.
- [35] S. Pasterski, A. Strominger, and A. Zhiboedov, “New Gravitational Memories,” *JHEP* **12** (2016) 053, 1502.06120.
- [36] G. Barnich and C. Troessaert, “Symmetries of asymptotically flat 4 dimensional spacetimes at null infinity revisited,” *Phys. Rev. Lett.* **105** (2010) 111103, 0909.2617.
- [37] F. Cachazo and A. Strominger, “Evidence for a New Soft Graviton Theorem,” 1404.4091.
- [38] P. Creminelli, J. Noreña, and M. Simonović, “Conformal consistency relations for single-field inflation,” *JCAP* **07** (2012) 052, 1203.4595.
- [39] K. Hinterbichler, L. Hui, and J. Khoury, “An Infinite Set of Ward Identities for Adiabatic Modes in Cosmology,” *JCAP* **01** (2014) 039, 1304.5527.