

Conformal BK equation at QCD Wilson-Fisher point

I. Balitsky^{a,b} and G.A. Chirilli^{c,d}

^a *Physics Department, Old Dominion University, Norfolk, VA 23529, USA.*

^b *Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA.*

^c *Department of Mathematics and Physics, University of Salento, Via Arnesano, Lecce, I-73100, Italy.*

^d *Sezione di Lecce, Istituto Nazionale di Fisica Nucleare, Via Arnesano, Lecce, I-73100, Italy.*

E-mail: ^{a,b} balitsky@jlab.org, ^{c,d} chirilli@le.infn.it

ABSTRACT: High-energy scattering in pQCD in the Regge limit is described by the evolution of Wilson lines governed by the BK equation [1, 2]. In the leading order, the BK equation is conformally invariant and the eigenfunctions of the linearized BFKL equation are powers. It is a common belief that at $d \neq 4$ the BFKL equation is useless since unlike $d = 4$ case it cannot be solved by usual methods. However, we demonstrate that at critical Wilson-Fisher point of QCD the relevant part of NLO BK restores the conformal invariance so the solutions are again powers. As a check of our approach to high-energy amplitudes at the Wilson-Fisher point, we calculate the anomalous dimensions of twist-2 light-ray operators in the Regge limit $j \rightarrow 1$.

Contents

1	Introduction	1
2	BK equation at the WF point	2
3	Pomeron intercept and anomalous dimensions of the twist-2 light-ray operators	5
3.1	“Regge+LC” limit of the CF (3.4)	6
3.2	“LC+Regge” limit of the CF (3.4)	8
3.3	Anomalous dimensions at $j \rightarrow 1$ from comparison of the two limits	11
4	Conclusions	12
5	Appendix: NLO BFKL eigenvalues	13
6	Appendix: gluon light-ray operators	14

1 Introduction

Despite the fact that QCD is far from being a conformal theory, some aspects of perturbative QCD are very close to those of a conformal theory. First and foremost, since bare QCD Lagrangian is conformally invariant, the leading-log (LL) evolutions in QCD have this property with respect of some subgroup of full $SO(4, 2)$. As an example, one may recall $SL(2, R)$ invariance of the leading-order evolution of light-ray QCD operators [3] and $SL(2, C)$ invariance of the BFKL equation [4]. Moreover, the common belief is that at higher orders of perturbation theory, the result in $\mathcal{N} = 4$ SYM theory gives the most complicated part of the corresponding result in QCD. This so-called maximal transcendentality (MT) feature [5] was checked in several examples. For instance, the MT part of three-loop DGLAP kernels is exactly the three-loop result in $\mathcal{N} = 4$ SYM theory [6]. The same property is correct for NLO pomeron intercept [5] and one should expect it to be true for the NNLO intercept which unfortunately is not obtained in QCD yet, although the $\mathcal{N} = 4$ intercept is available [7–9].

Another example of a conformal theory with perturbation theory “close” to that of QCD, is QCD at $d \neq 4$ at the so-called Wilson-Fisher (WF) point [10] $d = 4 - 2\varepsilon_*$ where β -function vanishes. An example of practical use of this closeness is the restoration of three-loop result for the GPD evolution in QCD from that of PDF evolution by a two-loop calculation of Ward identity at the WF point [11–13]. The goal of this paper is to lay a groundwork for similar analysis for the BFKL/BK equation describing high-energy (small- x) limit of QCD.

The common belief is that in $d \neq 4$ the BFKL equation is useless since unlike $d = 4$ it is not conformally invariant, and as a result, cannot be solved by power-like eigenfunctions. However, we will demonstrate below that at the WF point the conformal invariance of the BFKL equation is restored, at least in the leading order, so the solutions are powers again. This opens up the whole machinery of conformal field theories (CFTs) like general Conformal Regge theory discussed in Refs. [14–16] and theory of conformal light-ray operators [17–20].

As an application, we use the conformal BK equation at WF point to find anomalous dimensions of twist-2 light-ray operators of spin j in the “BFKL limit” $j \rightarrow 1$. Usually the extraction of $j \rightarrow 1$ limit of anomalous dimensions is done using the BFKL kernel in the momentum space [21–23], but in a conformal theory there is an alternative way. Since BFKL equation gives the amplitude in the Regge limit and anomalous dimensions are relevant for the DGLAP evolution of LR operators, one can consider a correlation function (CF) of 4 scalar currents in the double “Regge+light-cone” limit. On one side, one can first take Regge limit of that CF which is given by conformal Regge formula [24] (see below) and take a light-cone (LC) limit of that formula afterwards. On the other hand, we can start with an expansion in light-ray (LR) operators. Since LR operators are an analytic continuation of local operators (see the discussion in Ref. [17]) one can use an extension of Polyakov three-point formula and get the light-cone expansion of the same CF with explicit dependence on anomalous dimensions. After that we can take the small- x $j \rightarrow 1$ limit and compare this “LC+Regge” limit to “Regge+LC” representation discussed above. As was mentioned, the former representation involves anomalous dimensions of the leading-twist LR operators while the latter is governed by the BFKL pomeron. The comparison gives us an opportunity to relate anomalous dimension of LR operator to BFKL intercept. The result is as expected: the pomeron intercept at $d = 4 - 2\varepsilon_*$ is different from intercept at $d = 4$, but the anomalous dimensions (in the \overline{MS} scheme) are the same.

The paper is organized as follows: in Sect. 2 we derive the conformal LO BK equation at the WF point using NLO BK in $d = 4$. In Sect. 3.1 we derive the “Regge+LC” representation discussed above while Sect. 3.2 is devoted to “LC+Regge” one. In Sect. 3.3 we compare the two representations and get the anomalous dimensions of LR operators in the $j \rightarrow 1$ limit while Sect. 4 is a “conclusions+outlook” section. The necessary $d=4$ NLO BFKL kernel and eigenvalues are listed in the Appendix 5 and the formal definition of gluon LR operators is presented in Appendix 6.

2 BK equation at the WF point

We consider QCD at non-integer $d = 4 - 2\varepsilon_*$ such that the β -function vanishes

$$\frac{1}{g}\beta(g) = -\varepsilon - \bar{\beta}(a) = -\varepsilon - (b_0 a + b_1 a^2 + \dots) = 0, \quad a \equiv \frac{\alpha_s}{4\pi} \quad (2.1)$$

To be on the safe side, one should consider QCD at small ε near the Banks-Zaks point [25] where n_f is such that $\bar{\beta}(a) = 0$. At small ε we are still in the conformal window. Since we are interested in the first few orders in perturbative expansion, after calculation we can analytically continue our formulas to n_f outside of the conformal window.

Thus, we consider QCD at the critical Wilson-Fisher point ε_* such that

$$\varepsilon_* = -ab_0 - a^2b_1, \quad b_0 = \frac{11}{3}N_c - \frac{2}{3}n_f, \quad b_1 = \frac{2}{3}[17N_c^2 - 5n_fN_c - 3c_Fn_f] \quad (2.2)$$

With out NLO accuracy we need only the b_0 part of Eq. (2.2), the b_1 term will be important for NNLO calculations.

The main observation of the paper is the following. The LO BK equation for the rapidity evolution of color dipole $\text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}$ at arbitrary dimension of the transverse space $d_\perp = d - 2$ has the form:

$$\frac{d}{d\eta} \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\} = \frac{\alpha_s}{2\pi^{d_\perp}} \frac{\Gamma^2(\frac{d_\perp}{2})}{\mu_{\overline{MS}}^{d_\perp-2}} \int d^{d_\perp} z_3 K_{\text{LO}}^{d_\perp}(z_i) \left[\text{tr}\{U_{z_1} U_{z_3}^\dagger\} \text{tr}\{U_{z_3} U_{z_2}^\dagger\} - N_c \text{tr}\{U_{z_1} U_{z_2}^\dagger\} \right] \quad (2.3)$$

where $\mu_{\overline{MS}}^2 = \mu^2 \frac{e^{\gamma_E}}{4\pi}$ ¹ and

$$K_{\text{LO}}^{d_\perp}(z_i) = \frac{4\pi^{d_\perp}}{\Gamma^2(\frac{d_\perp}{2})} \left[(z_1 | \frac{p_i}{p^2} | z_3)(z_3 | \frac{p_i}{p^2} | z_1) + (z_2 | \frac{p_i}{p^2} | z_3)(z_3 | \frac{p_i}{p^2} | z_2) - 2(z_1 | \frac{p_i}{p^2} | z_3)(z_3 | \frac{p_i}{p^2} | z_2) \right] \quad (2.4)$$

Here we use Schwinger's notation for propagators $(x | \frac{1}{p^2} | y) \equiv \int \frac{d^n p}{(2\pi)^n} e^{-ip(x-y)}$ and the rapidity cutoff η is defined as the logarithm of maximal p_+ of the gluons forming Wilson lines

$$U(x_i) = \text{Pexp} \left\{ ig \int dx_+ A_-(x_+, 0, x_{i\perp}) \right\} \quad (2.5)$$

The formula (2.3) is the same as the LO BK kernel in Ref. [1], with the only difference that the free propagators (before and after interaction with the shock wave) are in $d_\perp = 2 - 2\varepsilon$. Now, at $d_\perp = 2$ we get the familiar conformal dipole kernel [26, 27]

$$4\pi^2 \left[(z_1 | \frac{p_i}{p_\perp^2} | z_3)(z_3 | \frac{p_i}{p_\perp^2} | z_1) + (z_2 | \frac{p_i}{p_\perp^2} | z_3)(z_3 | \frac{p_i}{p_\perp^2} | z_2) - 2(z_1 | \frac{p_i}{p_\perp^2} | z_3)(z_3 | \frac{p_i}{p_\perp^2} | z_2) \right] = \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \quad (2.6)$$

while at $d_\perp = 2 - 2\varepsilon$ one obtains the kernel

$$K_{\text{LO}}^{d_\perp}(z_i) = \frac{1}{(z_{13}^2)^{d_\perp-1}} + \frac{1}{(z_{23}^2)^{d_\perp-1}} - \frac{2z_{13} \cdot z_{23}}{(z_{13}^2 z_{23}^2)^{\frac{d_\perp}{2}}} \quad (2.7)$$

which is not conformal so the corresponding LO BFKL equation cannot be solved by powers.

However, at the critical point ε_* one should expand in ε and α_s simultaneously. The expansion of the LO kernel $K_{\text{LO}}^{d_\perp}(z_i)$ can be rewritten as

$$K_{\text{LO}}^{2-2\varepsilon_*}(z_i) = \left(\frac{z_{12}^2}{z_{13}^2 z_{23}^2} \right)^{1-\varepsilon_*} [1 + \varepsilon_* \ln z_{12}^2] + \left(\frac{1}{z_{13}^2} - \frac{1}{z_{23}^2} \right) \ln \frac{z_{13}^2}{z_{23}^2} + O(\varepsilon^2) \quad (2.8)$$

The NLO BK equation in d=4 QCD reads [28, 29]

$$\begin{aligned} \frac{d}{d\eta} [\text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}] &\stackrel{d=4}{=} \frac{\alpha_s}{2\pi^2} \int d^2 z_3 \frac{z_{12}^2}{z_{13}^2 z_{23}^2} \left\{ 1 + \frac{\alpha_s}{4\pi} b_0 \left[\ln \frac{z_{12}^2 \mu^2}{4} + 2\gamma_E - \frac{z_{13}^2 - z_{23}^2}{z_{12}^2} \ln \frac{z_{13}^2}{z_{23}^2} \right] \right. \\ &+ \left. \frac{\alpha_s N_c}{4\pi} \left(\frac{67}{9} - \frac{\pi^2}{3} - \frac{10n_f}{9N_c} \right) \right\} [\text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta}\} \text{tr}\{\hat{U}_{z_3}^\eta \hat{U}_{z_2}^{\dagger\eta}\} - N_c \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}] + \frac{\alpha_s^2 N_c}{16\pi^2} K_{\text{conf}} \end{aligned} \quad (2.9)$$

¹Throughout the paper, μ is the normalization point in the \overline{MS} scheme and $\alpha_s \equiv \alpha_s(\mu)$

where K_{conf} is the conformal part presented in the Appendix 5. Consequently, the NLO BK equation at $d = 4 - 2\varepsilon$ can be written as

$$\begin{aligned} \frac{d}{d\eta} \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\} &= \frac{\alpha_s \Gamma(1-\varepsilon)}{2\pi^{2-\varepsilon}} \int d^{2-2\varepsilon} z_3 \left(\frac{z_{12}^2}{z_{13}^2 z_{23}^2} \right)^{1-\varepsilon} \left\{ 1 + \left(\varepsilon + \frac{\alpha_s}{4\pi} b_0 \right) \left[\ln \frac{z_{12}^2 \mu^2}{4} \right. \right. \\ &+ 2\gamma_E - \frac{z_{13}^2 - z_{23}^2}{z_{12}^2} \ln \frac{z_{13}^2}{z_{23}^2} \left. \left. \right] + \frac{\alpha_s N_c}{4\pi} \left(\frac{67}{9} - \frac{\pi^2}{3} - \frac{10n_f}{9N_c} \right) \right\} [\text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta}\} \text{tr}\{\hat{U}_{z_3}^\eta \hat{U}_{z_2}^{\dagger\eta}\} \\ &- N_c \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}] + \frac{\alpha_s^2 N_c}{16\pi^2} K_{\text{conf}} + O(\alpha_s^3, \alpha_s^2 \varepsilon, \alpha_s \varepsilon^2) \end{aligned} \quad (2.10)$$

which turns to conformal equation

$$\begin{aligned} \frac{d}{d\eta} \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\} &= \frac{\alpha_s \Gamma(1-\varepsilon_*)}{2\pi^{2-\varepsilon_*}} \int d^{2-2\varepsilon_*} z_3 \left(\frac{z_{12}^2}{z_{13}^2 z_{23}^2} \right)^{1-\varepsilon_*} \left[1 + \frac{\alpha_s N_c}{4\pi} \left(\frac{67}{9} - \frac{\pi^2}{3} - \frac{10n_f}{9N_c} \right) \right] \\ &\times [\text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta}\} \text{tr}\{\hat{U}_{z_3}^\eta \hat{U}_{z_2}^{\dagger\eta}\} - N_c \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_2}^{\dagger\eta}\}] + \frac{\alpha_s^2 N_c}{16\pi^2} K_{\text{conf}} + O(\alpha_s^3, \alpha_s^2 \varepsilon_*, \alpha_s \varepsilon_*^2) \end{aligned} \quad (2.11)$$

at the critical point $d = 4 - 2\varepsilon_*$.

Thus, the BK equation at $d = 4 - 2\varepsilon_*$ is conformally (Möbius) invariant up to NLO order.² It should be noted that this property, albeit expected at the LO, is not a direct consequence of the invariance of the WF QCD since rapidity cutoff of Wilson lines violates conformal invariance (even in the conformal $\mathcal{N} = 4$ SYM). We hope that at the next order in α_s the conformal invariance will survive after appropriate replacement of the dipole $U_x U_y^\dagger$ by a suitable ‘‘conformal dipole’’ similar to the $\mathcal{N} = 4$ SYM case at $d = 4$ [29].

The conformal invariance of BK equation at the QCD critical point opens up the whole machinery of CFT for the analysis of high-energy amplitudes. For example, applying the logic of this Section in reverse, one could have predicted the b_0 part of NLO BK (2.9) coming from the running of the QCD coupling constant. It is worth noting that for now, only this b_0 part is used in the numerical estimates of the solution of BK equation with running coupling.

As an another example of using CFT methods for high-energy amplitudes in QCD, we will obtain the anomalous dimensions of twist-two gluon operators $F^{-i} \nabla_-^{j-2} F_i^-$ at the ‘‘BFKL point’’ $j \rightarrow 1$. We will see that while the pomeron intercept at $d = 4 - 2\varepsilon_*$ is different from that of $d = 4$, the anomalous dimensions of these operators in \overline{MS} scheme are the same as discussed in Refs. [11–13].

Let us start with calculation of the pomeron intercept at $d = 4 - 2\varepsilon_*$. The intercept is the rightmost eigenvalue of the BFKL equation which is a linearization of Eq. (2.11). To find the intercept, it is sufficient to consider forward matrix elements of the dipole

$$\mathcal{U}(z_1, z_2) \equiv 1 - \frac{1}{N_c} \text{tr}\{\hat{U}(z_{1\perp}) \hat{U}^\dagger(z_{2\perp})\}, \quad \langle \mathcal{U}(z_1, z_2) \rangle = \mathcal{U}(z_{12})$$

²For now, we see only the invariance under the inversion in the transverse plane. At $d = 4$, it is demonstrated in Ref. [29] that the LO BK kernel satisfies equations following from standard commutation relations of leading-order $\text{SL}(2, \mathbb{C})$ operators. These relations are not satisfied at $d = 4 - 2\varepsilon_*$ which means that the corresponding leading-order $\text{SL}(2, \mathbb{C})$ operators get $O(\alpha_s, \varepsilon)$ corrections similarly to $\text{SL}(2, \mathbb{R})$ operators discussed in Refs. [11, 30]. The authors hope to return to this matter in future publications.

The forward linearized equation (2.11) takes the form

$$\begin{aligned} \frac{d}{d\eta}\mathcal{U}(z) &= \frac{\alpha_s N_c \Gamma(1 - \varepsilon_*)}{2\pi^{2-\varepsilon_*}} \int d^{2-2\varepsilon_*} z' \left[1 + \frac{\alpha_s N_c}{4\pi} \left(\frac{67}{9} - \frac{\pi^2}{3} - \frac{10n_f}{9N_c} \right) \right] \left(\frac{z^2}{(z-z')^2 z'^2} \right)^{1-\varepsilon_*} \\ &\times [2\mathcal{U}(z') - \mathcal{U}(z)] + \frac{\alpha_s^2 N_c^2}{4\pi^3} \int dz' \mathcal{K}_{\text{conf}}(z, z') \mathcal{U}(z') + O(\alpha_s^3, \alpha_s^2 \varepsilon_*, \alpha_s \varepsilon_*^2) \end{aligned} \quad (2.12)$$

where the linearized forward kernel $\mathcal{K}_{\text{conf}}(z, z')$ is presented in the Appendix 5. Since the pomeron have quantum numbers of the vacuum, it should be related to the angle-independent eigenfunctions of this equation, namely powers $(z'^2/z^2)^\xi$. Using the integral

$$\frac{\Gamma(\frac{d_\perp}{2})}{2\pi^{\frac{d_\perp}{2}}} \int d^{d_\perp} z' \left(\frac{z^2}{z'^2(z-z')^2} \right)^{\frac{d_\perp}{2}} \left[2\left(\frac{z'^2}{z^2}\right)^\xi - 1 \right] = \left[\psi\left(\frac{d_\perp}{2}\right) - \psi(\xi) - \psi\left(\frac{d_\perp}{2} - \xi\right) - \gamma_E \right] \quad (2.13)$$

we obtain the eigenvalues in the form

$$\begin{aligned} \bar{\mathfrak{N}}(\xi) &= \frac{\alpha_s N_c}{\pi} \left[\bar{\chi}(\xi) + \frac{\alpha_s N_c}{4\pi} \bar{\delta}(\xi) + O(\alpha_s^2, \alpha_s \varepsilon_*, \varepsilon_*^2) \right] \\ \bar{\chi}(\xi) &= \psi(1 - \varepsilon_*) - \gamma_E - \psi(\xi) - \psi(1 - \varepsilon_* - \xi) \end{aligned} \quad (2.14)$$

where $\psi(x)$ is the logarithmic derivative of the gamma-function ($\psi(1) = -\gamma_E$). With our accuracy, we need the function $\bar{\delta}(\xi)$ at $d = 4$. It was calculated in Ref. [22] and we represent it in the Appendix 5.³

It is easy to see that the pomeron intercept is the eigenvalue corresponding to eigenfunction with power $\xi = \frac{1}{2} - \frac{\varepsilon_*}{2} - i\nu$ (and no angular dependence). Indeed, writing the identity

$$\mathcal{U}(|z|) = \frac{\Gamma(\frac{d_\perp}{2})}{\pi^{\frac{d_\perp}{2}}} \int d^{d_\perp} z' \int \frac{d\nu}{2\pi} (z'^2)^{-\frac{d_\perp}{2} - \frac{d_\perp}{4} + i\nu} (z^2)^{\frac{d_\perp}{2} - \frac{d_\perp}{4} - i\nu} \mathcal{U}(|z'|), \quad (2.15)$$

taking the derivative with respect to rapidity according to Eq. (2.12) and exponentiating, one obtains

$$\mathcal{U}(|z|) = \frac{\Gamma(\frac{d_\perp}{2})}{\pi^{\frac{d_\perp}{2}}} \int d^{d_\perp} z' \int \frac{d\nu}{2\pi} (z'^2)^{-\frac{d_\perp}{2} - \frac{d_\perp}{4} + i\nu} (z^2)^{\frac{d_\perp}{2} - \frac{d_\perp}{4} - i\nu} e^{\eta \bar{\mathfrak{N}}(\frac{d_\perp}{4} - i\nu)} \mathcal{U}(|z'|). \quad (2.16)$$

So the pomeron intercept (at $\nu = 0$) is $\frac{\alpha_s N_c}{\pi} \chi\left(\frac{1}{2} - \frac{\varepsilon_*}{2}\right) = \frac{\alpha_s N_c}{\pi} \left[4 \ln 2 - \frac{\pi^2}{3} \varepsilon_* + O(\varepsilon_*^2) \right]$.

3 Pomeron intercept and anomalous dimensions of the twist-2 light-ray operators

It is well known that the analytic continuation of anomalous dimensions of twist-2 local operators $F^{-i} \nabla_-^{j-2} F_i^-$ to $j \simeq 1$ can be obtained from BFKL pomeron intercept. Since $j = 1$ is the unphysical point for local gluon operators, one should consider the generalization of local twist-2 operators to gluon light-ray operators discussed in Refs. [3, 31]. A gluon LR

³To avoid confusion with the notations in Ref. [22] we denote our (slightly different) functions $\bar{\chi}$ and $\bar{\delta}$ with bars.

operator is an analytic continuation in spin of a local operator $F^{-i}\nabla_-^{j-2}F_i^-$, see Appendix 6 for detailed definitions. For simplicity, we consider pure gluodynamics where gluon light-ray operators are multiplicatively renormalizable.

As mentioned in the Introduction, usually the relation between Pomeron intercept and anomalous dimensions of twist-2 operators is obtained from the BFKL kernel in the momentum space, but in a conformal theory it can be done directly in the coordinate space. One way is to consider the correlation function of two ‘‘Wilson frames’’ as was done in Ref. [31] for $\mathcal{N} = 4$ SYM. However, this method requires calculating many nontrivial integrals at $d_\perp \neq 2$. An easier way, which does not require any computation, is described in Ref. [32]. One considers the CF of four scalar operators, for example in $d = 4 - 2\varepsilon_*$ QCD and takes $\mathcal{O}(x) = F_{\mu\nu}^a(x)F^{a\mu\nu}(x)$, and get

$$A(R, r) = [x_{12}^2 x_{34}^2]^{4-2\varepsilon_* + \gamma_{\mathcal{O}}} (\mu^2)^{2\gamma_{\mathcal{O}}} \langle \mathcal{O}(x_1) \mathcal{O}(x_2) \mathcal{O}(x_3) \mathcal{O}(x_4) \rangle \quad (3.1)$$

where $\gamma_{\mathcal{O}}$ is the anomalous dimension of the operator \mathcal{O} . With Regge limit in view, it is convenient to choose two conformal ratios as follows:

$$R = \frac{x_{13}^2 x_{24}^2}{x_{12}^2 x_{34}^2}, \quad \mathfrak{r} = R^{-1} \left[1 - \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2} + \frac{1}{R} \right]^{-2} \quad (3.2)$$

Now, we consider the CF (3.1) in the double Regge and light-cone limit. The Regge limit can be specified as

$$x_{1+} \rightarrow \rho x_{1+}, x_{2+} \rightarrow \rho x_{2+}, \quad x_{1-} = x_{2-} = 0, \quad x_{3-} \rightarrow \rho' x_{3-}, x_{4-} \rightarrow \rho' x_{4-}, \quad x_{3+} = x_{4+} = 0 \quad (3.3)$$

with $\rho\rho' \rightarrow \infty$ and the light-cone limit corresponds to $x_{12\perp}^2 \rightarrow 0$ (we use metric $x^2 = 2x_+x_- - x_\perp^2$). It is convenient to consider ‘‘forward’’ correlation function ⁴

$$A(L_+, L_-; x_{1\perp}, x_{2\perp}, x_{3\perp}, x_{4\perp}) = (x_{12\perp}^2 x_{34\perp}^2)^{4-2\varepsilon_* + \gamma_{\mathcal{O}}} (\mu^2)^{2\gamma_{\mathcal{O}}} \times \int dx_{2+} dx_{3-} \langle \mathcal{O}(L_+ + x_{2+}, x_{1\perp}) \mathcal{O}(x_{2+}, x_{2\perp}) \mathcal{O}(L_- + x_{4-}, x_{3\perp}) \mathcal{O}(x_{4-}, x_{4\perp}) \rangle \quad (3.4)$$

in the double limit: Regge ($L_+L_- \rightarrow \infty$) plus light-cone ($x_{12}^2 \rightarrow 0$). By comparison of two orders: ‘‘Regge+LC’’ and ‘‘LC+Regge’’ we will be able to relate anomalous dimensions of gluon LR operators in the limit

$$j - 1 = \omega \rightarrow 0, \quad \frac{\alpha_s}{\omega} \sim 1 \quad (3.5)$$

to the BFKL pomeron intercept.

3.1 ‘‘Regge+LC’’ limit of the CF (3.4)

The 4-point CF (3.1) is a function of two conformal ratios (3.2) which reduce to

$$R = \frac{x_{13}^2 x_{24}^2}{x_{12}^2 x_{34}^2} \rightarrow \frac{4x_{1+}x_{2+}x_{3-}x_{4-}}{x_{12\perp}^2 x_{34\perp}^2}, \quad \mathfrak{r} = R^{-1} \left[1 - \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2} + \frac{1}{R} \right]^{-2} \rightarrow \frac{x_{12\perp}^2 x_{34\perp}^2 x_{1+}x_{2+}x_{3-}x_{4-}}{[x_{24\perp}^2 x_{1+}x_{3-} + x_{2+}x_{4-} x_{13\perp}^2 - x_{1+}x_{4-} x_{23\perp}^2 - x_{2+}x_{3-} x_{14\perp}^2]^2} \quad (3.6)$$

⁴In our formulas throughout the paper L_+ and L_- are positive.

in the Regge limit (3.3). It is easy to see that under the rescaling (3.3) R increases with “energy” ($\sim \sqrt{\rho\rho'}$) while \mathfrak{r} is energy-independent. As demonstrated in Ref. [14, 24, 33]), the general formula for the Regge limit of a 4-point CF in a conformal theory has the form

$$A(R, \mathfrak{r}) \stackrel{s \sim \rho\rho' \rightarrow \infty}{=} \frac{i}{2} \int d\nu f_+(\aleph(\alpha_s, \nu)) F(\alpha_s, \nu) \Omega(\mathfrak{r}, \nu) R^{\aleph(\alpha_s, \nu)/2} \quad (3.7)$$

where $f_+(\omega) = \frac{e^{i\pi\omega} - 1}{\sin \pi\omega}$ is a signature factor and

$$\Omega(\mathfrak{r}, \nu) = \frac{2\nu \sinh 2\pi\nu \Gamma(2 - \frac{d}{2} + 2i\nu) \Gamma(\frac{d}{2} - 1 - 2i\nu) \Gamma(d - 2)}{2^{d-1} \pi^{\frac{d+1}{2}} \Gamma(\frac{d}{2} - \frac{1}{2})} C_{-\frac{d}{2} + 1 + 2i\nu}^{\frac{d}{2} - 1} \left(\frac{1}{2\sqrt{\mathfrak{r}}} \right) \quad (3.8)$$

is a solution of the Laplace equation for H_{d-1} hyperboloid ($\partial_{H_{d-1}}^2 + \nu^2 + 1$) $\Omega(\mathfrak{r}, \nu) = 0$. The dynamics is described by the pomeron intercept $\aleph(\alpha_s, \nu)$ and the “pomeron residue” $F(\alpha_s, \nu)$. The formula (3.7) was proved in Ref. [24] (see also Refs. [14, 33]) by considering the leading Regge pole in a conformal theory. Also, it was demonstrated up to the NLO level that the structure (3.7) is reproduced by the high-energy OPE in Wilson lines [34].

Now let us take the light-cone limit $x_{12}^2 = -x_{12_\perp}^2 \rightarrow 0$ on top of the Regge limit (3.7). In this limit

$$R = \frac{x_{13}^2 x_{24}^2}{x_{12}^2 x_{34}^2} \rightarrow \frac{4x_{1+} x_{2+} x_{3-} x_{4-}}{x_{12_\perp}^2 x_{34_\perp}^2}, \quad \mathfrak{r} \rightarrow \frac{x_{1+} x_{2+} x_{3-} x_{4-} x_{12_\perp}^2 x_{34_\perp}^2}{x_{12_\perp}^2 (x_{3-} x_{14_\perp}^2 - x_{4-} x_{13_\perp}^2)^2} \quad (3.9)$$

and

$$\Omega(\mathfrak{r}, \nu) \stackrel{\mathfrak{r} \rightarrow \infty}{=} \frac{2^{\varepsilon_* - 2} \nu \sinh 2\pi\nu \Gamma(2 - 2\varepsilon_*)}{\pi^{\frac{5}{2} - \varepsilon_*} \Gamma(\frac{3}{2} - \varepsilon_*) \Gamma(1 - \varepsilon_*)} \left[\Gamma(-2i\nu) \Gamma(1 - \varepsilon_* + 2i\nu) \mathfrak{r}^{\frac{1}{2} - \frac{\varepsilon_*}{2} + i\nu} + \text{c.c.} \right] \quad (3.10)$$

so Eq. (3.4) reduces to

$$\begin{aligned} A(L_+, L_-; x_{i_\perp}) &= i \int d\nu f_+(\aleph(\alpha_s, \nu)) \tilde{F}(\alpha_s, \nu) \int dx_{2+} dx_{4-} \theta(L_+ + x_{2+}) \theta(-x_{2+}) \\ &\times \theta(L_- + x_{4-}) \theta(-x_{4-}) \left(\frac{(L_+ + x_{2+}) x_{2+} (L_- + x_{4-}) x_{4-}}{x_{12_\perp}^2 x_{34_\perp}^2} \right)^{\frac{\aleph(\nu, \alpha_s)}{2}} \\ &\times \left(\frac{L_+^2 [(L_- + x_{4-}) x_{14_\perp}^2 - x_{4-} x_{13_\perp}^2]^2}{(L_+ + x_{2+}) x_{2+} (L_- + x_{4-}) x_{4-} x_{12_\perp}^2 x_{34_\perp}^2} \right)^{-\frac{1}{2} + \frac{\varepsilon_*}{2} - i\nu} \end{aligned} \quad (3.11)$$

where the prefactors in Eq. (3.10) are absorbed in the function $\tilde{F}(\alpha_s, \nu)$.

The theta-function factors in Eq. (3.11) require some explanation. Qualitatively, one should put them in since Regge limit corresponds to $x_{1+}, x_{3-} > 0$ and $x_{2+}, x_{4-} < 0$. Quantitatively, it follows from the derivation of Eq. (3.7) by high-energy factorization [34]. In that approach, the amplitude (3.4) is a product of two “impact factors” - coefficients of high-energy OPE in color dipoles, and the amplitude of dipole-dipole scattering. The “top” impact factors is the CF of two currents $\mathcal{O}(L_+ + x_{2+}, x_{1_\perp}) \mathcal{O}(x_{2+}, x_{2_\perp})$ in the background of the shock wave positioned at $x_+ = 0$, so the region $L_+ + x_{2+} > 0 > x_{2+}$ contributes with the weight one (there is an intersection with the shock wave) while regions $L_+ + x_{2+}, x_{2+} > 0$

and $L_+ + x_{2+}, x_{2+} < 0$ give zero contribution since there are no interactions with the shock wave in that regions. Similarly, from the consideration of the “bottom” impact factors one sees that only the region $L_- + x_{4-} > 0 > x_{4-}$ gives nonzero contribution. ⁵

Now, introducing variables $v = \frac{|x_{2+}|}{L_+}$, $u = \frac{|x_{4-}|}{L_-}$ we get

$$A(L_+, L_-; x_{1\perp}, x_{2\perp}, x_{3\perp}, x_{4\perp}) = iL_+L_- \times \int_0^1 du dv \int d\nu F(\nu, \alpha_s) \left(\frac{x_{12\perp}^2 x_{34\perp}^2 \bar{u} u \bar{v} v}{[x_{13\perp}^2 u + x_{14\perp}^2 \bar{u}]^2} \right)^{\frac{1}{2} - \frac{\varepsilon_*}{2} + i\nu} \left(\frac{L_+^2 L_-^2 \bar{u} u \bar{v} v}{x_{12\perp}^2 x_{34\perp}^2} \right)^{\aleph(\nu, \alpha_s)/2} f_+(\aleph(\nu, \alpha_s))$$

Performing integral over v one obtains the representation of the CF (3.4) in the double “Regge+LC” limit in the form

$$A(L_+, L_-; x_{1\perp}, x_{2\perp}, x_{3\perp}, x_{4\perp}) = \int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty} \frac{d\xi}{2\pi} f_+(\bar{\aleph}(\xi)) F(\xi, \alpha_s) (L_+ L_-)^{1+\aleph(\xi)} \times \int_0^1 du \frac{(\bar{u}u)^{1-\varepsilon_*-\xi+\frac{\aleph(\xi)}{2}} (x_{12\perp}^2 x_{34\perp}^2)^{1-\xi-\varepsilon_*-\frac{\aleph}{2}}}{[x_{13\perp}^2 u + x_{14\perp}^2 \bar{u}]^{2-2\xi-2\varepsilon_*}} \quad (3.12)$$

where $\xi = \frac{1}{2} - i\nu - \frac{\varepsilon_*}{2}$, $\bar{\aleph}(\xi, \alpha_s) \equiv \aleph\left(-i\left(\frac{1}{2} - \xi - \frac{\varepsilon_*}{2}\right), \alpha_s\right)$, and

$$F(\xi, \alpha_s) = \frac{\Gamma^2(2 - \varepsilon_* - \xi + \frac{\aleph(\xi)}{2})}{\Gamma(4 - 2\varepsilon_* - 2\xi - \aleph(\xi))} \tilde{F}\left(\alpha_s, -i\left(\frac{1}{2} - \xi - \frac{\varepsilon_*}{2}\right)\right) \quad (3.13)$$

3.2 “LC+Regge” limit of the CF (3.4)

This time we start with the light-cone limit $x_{12\perp}^2 \rightarrow 0$ and perform the light-cone expansion of two “top” operators $\mathcal{O}(x_1)\mathcal{O}(x_2)$ in light-ray gluon operators. We will demonstrate that

$$(x_{12\perp}^2)^{3-\varepsilon_*} (\mu^2 x_{12\perp}^2)^{\gamma_{\mathcal{O}}} \int dx_{2+} \mathcal{O}(L_+ + x_{2+}, x_{1\perp}) \mathcal{O}(x_{2+}, x_{2\perp}) = \int_{\frac{3}{2}-i\infty}^{\frac{3}{2}+i\infty} \frac{dj}{2\pi} d(j, \alpha_s) L_+^j (\mu^2 x_{12\perp}^2)^{\frac{\gamma_j}{2}} \mathcal{F}_j^{(+)}(x_{1\perp}) + \mathcal{O}(x_{12\perp}^2) \quad (3.14)$$

where $\mathcal{F}_j^{(+)}(x_{\perp})$ is a “forward” light-ray operator with spin j defined in Eq. (6.2) in the Appendix 6. After that, we will write down three-point CF of light-ray operator $\mathcal{F}_j^{(+)}(x_{\perp})$ with two “bottom” operators $\mathcal{O}(x_3)\mathcal{O}(x_4)$ and obtain the representation of 4-point CF (3.4) as an integral over spin j . The Regge limit would correspond to the behavior of the integrand around $j = 1$.

It is convenient to start with the three-point CF of the LR operator $\mathcal{F}_j^{(+)}(x_{\perp})$ and two local operators. A textbook formula for CF of three primary local operators - one with spin n and two scalars - in a conformal theory reads

$$\langle \mathcal{O}_{\mu_1 \dots \mu_n}(z_1) \mathcal{O}(z_2) \mathcal{O}(z_3) \rangle \sim \frac{[1 + (-1)^n] \left(\frac{z_{12}}{z_{12}^2} - \frac{z_{13}}{z_{13}^2} \right)_{\mu_1} \dots \left(\frac{z_{12}}{z_{12}^2} - \frac{z_{13}}{z_{13}^2} \right)_{\mu_n}}{(z_{12}^2)^{\frac{\Delta_1 + \Delta_2 - \Delta_3}{2}} (z_{13}^2)^{\frac{\Delta_1 + \Delta_3 - \Delta_2}{2}} (z_{23}^2)^{\frac{\Delta_2 + \Delta_3 - \Delta_1}{2}}} \quad (3.15)$$

⁵For similar but more detailed discussion, see Ref. [31]

In our case, the conformal gluon operators with spin n are made from Gegenbauer polynomials $C_n^{\frac{5}{2}}(\overleftrightarrow{\nabla} / \overleftrightarrow{\partial})$ (at the one-loop level). However, for our purposes we need formula (3.15) integrated over z_{1+} so the only surviving term in $\mathcal{O}_G^{(n)}$ will be $F_{-i}^a \nabla_-^{n-2} F_-^{ai}$.

As explained in Ref. [17], to be on the safe side, the analytic continuation in spin of 3-point CF (3.15) should be done for Wightman CFs rather than for time-ordered ones so first we need to split this time-ordered CF into Wightman ones. Anticipating the next integral over total translation in x_- , consider the case $x_{3-} > x_{4-}$. First, we write down

$$\begin{aligned} & \langle \mathcal{O}(x_{3-}, x_{3\perp}) \mathcal{O}_G^{(n)}(x_{0+}, x_{0\perp}) \mathcal{O}(x_{4-}, x_{4\perp}) \rangle \\ &= \theta(x_{3-} > 0 > x_{4-}) \langle \mathcal{O}(x_{3-}, x_{3\perp}) \mathcal{O}_G^{(n)}(x_{0+}, x_{0\perp}) \mathcal{O}(x_{4-}, x_{4\perp}) \rangle^W \\ &+ \theta(x_{3-} > x_{4-} > 0) \langle \mathcal{O}(x_{3-}, x_{3\perp}) \mathcal{O}(x_{4-}, x_{4\perp}) \mathcal{O}_G^{(n)}(x_{0+}, x_{0\perp}) \rangle^W \\ &+ \theta(0 > x_{3-} > x_{4-}) \langle \mathcal{O}_G^{(n)}(x_{0+}, x_{0\perp}) \mathcal{O}(x_{3-}, x_{3\perp}) \mathcal{O}(x_{4-}, x_{4\perp}) \rangle^W \end{aligned} \quad (3.16)$$

where $\langle \dots \rangle^W$ denotes a vacuum average of operators as they stand (without time ordering). First, let us consider the first term in the above equation. After integration over x_{0+} we get

$$\begin{aligned} & (\mu^2)^{\gamma\mathcal{O} + \frac{\gamma n}{2}} \int dx_{0+} \langle \mathcal{O}(x_{3-}, x_{3\perp}) F_{-i}^a \nabla_-^{n-2} F_-^{ai}(x_{0+}, x_{0\perp}) \mathcal{O}(x_{4-}, x_{4\perp}) \rangle^W \stackrel{x_{3-} > 0 > x_{4-}}{=} \quad (3.17) \\ &= \int dx_{0+} \frac{c_n [1 + (-1)^n]}{(x_{34\perp}^2)^{3-\varepsilon_* + \gamma\mathcal{O} - \frac{\gamma n}{2}}} \left(\frac{x_{3-}}{2x_{0+}x_{3-} + x_{03\perp}^2 + i\epsilon x_{3-}} - \frac{x_{4-}}{2x_{0+}x_{4-} + x_{04\perp}^2 - i\epsilon x_{4-}} \right)^n \\ &\times \frac{1}{[(2x_{0+}x_{3-} + x_{03\perp}^2 + i\epsilon x_{3-})(2x_{0+}x_{4-} + x_{04\perp}^2 - i\epsilon x_{4-})]^{1-\varepsilon_* + \frac{\gamma n}{2}}} \\ &= -\pi i \frac{c_n [1 + (-1)^n]}{(x_{34\perp}^2)^{4-2\varepsilon_* + \gamma\mathcal{O}}} \frac{\Gamma(1 + 2n + \gamma_n - 2\varepsilon_*)}{\Gamma^2(1 + n + \frac{1}{2}\gamma_n - \varepsilon_*)} \left(\frac{x_{34\perp}^2}{x_{3-}|x_{4-}|} \right)^{1 + \frac{\gamma n}{2} - \varepsilon_*} \left(\frac{x_{03\perp}^2}{x_{3-}} - \frac{x_{04\perp}^2}{x_{4-}} \right)^{2\varepsilon_* - 1 - n - \gamma_n} \end{aligned}$$

Now we can continue analytically in j using Eqs. (6.2)

$$\begin{aligned} & -i\pi (\mu^2)^{\gamma\mathcal{O} + \frac{\gamma j}{2}} \langle \mathcal{O}(x_{3-}, x_{3\perp}) \mathcal{F}_j^{(+)}(x_{0\perp}) \mathcal{O}(x_{4-}, x_{4\perp}) \rangle^W \stackrel{x_{3-} > 0 > x_{4-}}{=} \quad (3.18) \\ &= -i\pi \frac{c(j, \alpha_s) [1 + e^{i\pi j}]}{(x_{34\perp}^2)^{4-2\varepsilon_* + \gamma\mathcal{O}}} \frac{\Gamma(1 + 2j + \gamma_j - 2\varepsilon_*)}{\Gamma^2(1 + j + \frac{1}{2}\gamma_n - \varepsilon_*)} \left(\frac{x_{34\perp}^2}{x_{3-}|x_{4-}|} \right)^{1 + \frac{\gamma j}{2} - \varepsilon_*} \left(\frac{x_{03\perp}^2}{x_{3-}} + \frac{x_{04\perp}^2}{|x_{4-}|} \right)^{2\varepsilon_* - 1 - j - \gamma_j} \end{aligned}$$

For now, we have proved this equation only for Wightman CF. However, it is easy to see that two other orderings in Eq. (3.16) give zero result after integration over x_{0+} . Indeed, at $x_{3-} > x_{4-} > 0$ all singularities in the denominators in Eq. (3.17) are of the form $(x_{0+} + i\epsilon)$ which gives zero contribution to the integral. Similarly, at $0 > x_{3-} > x_{4-}$ the singularities in the denominators are $\sim (x_{0+} - i\epsilon)$ and the result vanishes. This is in accordance with the general statement in Ref. [17] that the CFs

$$\langle \mathcal{F}_j^{(+)}(x_{0\perp}) \mathcal{O}(x_{4-}, x_{4\perp}) \mathcal{O}(x_{3-}, x_{3\perp}) \rangle^W = \langle \mathcal{O}(x_{4-}, x_{4\perp}) \mathcal{O}(x_{3-}, x_{3\perp}) \mathcal{F}_j^{(+)}(x_{0\perp}) \rangle^W = 0$$

because a (light-ray) operator with non-integer spin annihilates the vacuum. Thus, we get for the T-product of operators

$$\begin{aligned}
& -i\pi(\mu^2)^{\gamma_{\mathcal{O}}+\frac{\gamma_j}{2}} \langle \mathcal{O}(x_{3_{\perp}}, x_{3_{\perp}}) \mathcal{F}_j^{(+)}(x_{0_{\perp}}) \mathcal{O}(x_{4_{\perp}}, x_{4_{\perp}}) \rangle \stackrel{x_{3_{\perp}} > x_{4_{\perp}}}{=} -i\pi \frac{c(j, \alpha_s)[1 + e^{i\pi j}]}{(x_{3_{4_{\perp}}}^2)^{4-2\varepsilon_*+\gamma_{\mathcal{O}}}} \\
& \times \frac{\Gamma(1+2j+\gamma_j-2\varepsilon_*)}{\Gamma^2(1+j+\frac{1}{2}\gamma_n-\varepsilon_*)} \left(\frac{x_{3_{4_{\perp}}}^2}{-x_{3_{\perp}}x_{4_{\perp}}} \right)^{1+\frac{\gamma_j}{2}-\varepsilon_*} \left(\frac{x_{0_{3_{\perp}}}^2}{x_{3_{\perp}}} - \frac{x_{0_{4_{\perp}}}^2}{x_{4_{\perp}}} \right)^{2\varepsilon_*-1-j-\gamma_j} \theta(x_{3_{\perp}})\theta(-x_{4_{\perp}})
\end{aligned} \tag{3.19}$$

Finally, let us perform an integration over total translation in x_{\perp} direction:

$$\begin{aligned}
& (\mu^2)^{\gamma_{\mathcal{O}}+\frac{\gamma_j}{2}} (x_{3_{4_{\perp}}}^2)^{4-2\varepsilon_*+\gamma_{\mathcal{O}}} \int dx_{4_{\perp}} \langle \mathcal{O}(L_{\perp} + x_{4_{\perp}}, x_{3_{\perp}}) \mathcal{F}_j^{(+)}(x_{0_{\perp}}) \mathcal{O}(x_{4_{\perp}}, x_{4_{\perp}}) \rangle \\
& = -i\pi L_{\perp}^j c(j, \alpha_s)[1 + e^{i\pi j}] \frac{\Gamma(1+2j+\gamma_j-2\varepsilon_*)}{\Gamma^2(1+j+\frac{\gamma_j}{2}-\varepsilon_*)} \int_0^1 du \frac{(x_{3_{4_{\perp}}}^2)^{1+\frac{\gamma_j}{2}-\varepsilon_*} (\bar{u}u)^{j+\frac{\gamma_j}{2}-\varepsilon_*}}{(x_{0_{3_{\perp}}}^2 u + x_{0_{4_{\perp}}}^2 \bar{u})^{1+j+\gamma_j-2\varepsilon_*}}
\end{aligned} \tag{3.20}$$

where $u = \frac{|x_{4_{\perp}}|}{L_{\perp}}$ and $\bar{u} \equiv 1 - u$.

Now we are in a position to prove expansion in LR operators (3.14). To this end, first we take Eq. (3.20), make the top \leftrightarrow bottom replacement $x_{3_{\perp}} \rightarrow x_{1_{\perp}}, x_{4_{\perp}} \rightarrow x_{2_{\perp}}, x_{3_{\perp}} \rightarrow x_{1_{\perp}}, x_{4_{\perp}} \rightarrow x_{2_{\perp}}, x_{0_{+}} \rightarrow x_{0_{\perp}}$ and $\mathcal{F}_j^{(+)}(x_{0_{\perp}}) \rightarrow \mathcal{F}_j^{(-)}(x_{0_{\perp}})$ ⁶ and take the light-cone limit $x_{12_{\perp}} \rightarrow 0$. We get

$$\begin{aligned}
& (\mu^2)^{\gamma_{\mathcal{O}}+\frac{\gamma_j}{2}} (x_{12_{\perp}}^2)^{4-2\varepsilon_*+\gamma_{\mathcal{O}}} \int dx_{2_{\perp}} \langle \mathcal{O}(L_{\perp} + x_{2_{\perp}}, x_{1_{\perp}}) \mathcal{F}_j^{(-)}(x_{0_{\perp}}) \mathcal{O}(x_{2_{\perp}}, x_{2_{\perp}}) \rangle \\
& \stackrel{x_{12_{\perp}} \rightarrow 0}{=} -i\pi L_{\perp}^j \frac{c(j, \alpha_s)[1 + e^{i\pi j}]}{(1+2j+\gamma_j-2\varepsilon_*)} \frac{(x_{12_{\perp}}^2)^{1+\frac{\gamma_j}{2}-\varepsilon_*}}{(x_{0_{1_{\perp}}}^2)^{1+j+\gamma_j-2\varepsilon_*}}
\end{aligned} \tag{3.21}$$

Second, the CF of two LR operators has the form [31, 35]

$$\langle \mathcal{F}_{j=\frac{3}{2}+i\nu}^{(+)}(x_{\perp}) \mathcal{F}_{j'=\frac{3}{2}+i\nu'}^{(-)}(y_{\perp}) \rangle = \frac{2\pi\delta(\nu-\nu')a(j, \alpha_s)(1 + e^{i\pi j})}{[(x-y)_{\perp}^2]^{j+1-2\varepsilon_*+\gamma_j}(\mu^2)^{\gamma_j}} \tag{3.22}$$

Note that it is essential to have T-product of two LR operators in the l.h.s., for the Wightman product one gets zero as discussed in Ref. [17]⁷.

Now let us expand the l.h.s. of Eq. (3.21) at $x_{12_{\perp}} \rightarrow 0$ using Eq. (3.14):

$$\begin{aligned}
& (\mu^2)^{\frac{\gamma_j}{2}} (x_{12_{\perp}}^2)^{1-\varepsilon_*} \int_{\frac{3}{2}-i\infty}^{\frac{3}{2}+i\infty} \frac{dj'}{2\pi} d(j', \alpha_s) L_{\perp}^{j'} (\mu^2 x_{12_{\perp}}^2)^{\frac{\gamma_{j'}}{2}} \langle \mathcal{F}_{j'}^{(+)}(x_{1_{\perp}}) \mathcal{F}_j^{(-)}(x_{0_{\perp}}) \rangle \\
& = iL_{\perp}^j (1 + e^{i\pi j}) a(j, \alpha_s) d(j, \alpha_s) (x_{12_{\perp}}^2)^{1-\varepsilon_*+\frac{\gamma_j}{2}} (x_{0_{1_{\perp}}}^2)^{2\varepsilon_*-j-1-\gamma_j}
\end{aligned} \tag{3.23}$$

⁶The explicit form of $\mathcal{F}_j^{(-)}(x_{0_{\perp}})$ is written in Eq. (6.3) from Appendix 6.

⁷We have checked by explicit calculation in the free massless electrodynamics that non-zero contributions to CF in Eq. (3.22) come from operator orderings like $F_{+i}(L_{\perp} + x_{\perp}) \mathcal{F}_{\perp}^j(y_{\perp}) F_{+i}(x_{\perp})$ with one of the LR operators split into two pieces. One may wonder if there will be problems with split gauge links for QCD light-rays, but the analysis in Ref. [31] shows that there are none, at least in the Regge limit.

Comparing this to the r.h.s. of Eq. (3.21), we confirm Eq. (3.14) with

$$d(j, \alpha_s) = \frac{c(j, \alpha_s)}{a(j, \alpha_s)} \frac{-\pi}{1 + 2j + \gamma_j - 2\varepsilon_*} \quad (3.24)$$

Finally, we combine the light-cone expansion (3.14) with CF (3.20) and obtain

$$\begin{aligned} A(L_+, L_-; x_{1\perp}, x_{2\perp}, x_{3\perp}, x_{4\perp}) &= [x_{12}^2 x_{34}^2]^{4-2\varepsilon_* + \gamma_{\mathcal{O}}} (\mu^2)^{2\gamma_{\mathcal{O}}} \\ &\times \int dx_{2+} dx_{4-} \langle \mathcal{O}(L_+ + x_{2+}, x_{1\perp}) \mathcal{O}(x_{2+}, x_{2\perp}) \mathcal{O}(L_- + x_{4-}, x_{3\perp}) \mathcal{O}(x_{4-}, x_{4\perp}) \rangle \\ &= \int_{\frac{3}{2}-i\infty}^{\frac{3}{2}+i\infty} \frac{dj}{2\pi} (1 + e^{i\pi j}) (L_+ L_-)^j F(j, \alpha_s) \int_0^1 du \frac{(\bar{u}u)^{j+\frac{\gamma_j}{2}-\varepsilon_*} (x_{12\perp}^2 x_{34\perp}^2)^{1-\varepsilon_*+\frac{\gamma_j}{2}}}{(x_{03\perp}^2 \bar{u} + x_{04\perp}^2 u)^{1+j+\gamma_j-2\varepsilon_*}} \end{aligned} \quad (3.25)$$

where

$$F(j, \alpha_s) = -i\pi c(j, \alpha_s) d(j, \alpha_s) \frac{\Gamma(1 + 2j - \gamma_j + 2\varepsilon_*)}{\Gamma^2(1 + j + \frac{\gamma_j}{2} - \varepsilon_*)} \quad (3.26)$$

The ‘‘Regge limit’’ of Eq. (3.25) corresponds to the behavior of the integrand around $j = 1$ which should be compared to behavior of the integrand of Eq. (3.12) at small $\aleph(\xi)$.

3.3 Anomalous dimensions at $j \rightarrow 1$ from comparison of the two limits

It is easy to see that the integrals (3.12) and (3.25) are actually the same if one makes identifications

$$\omega \equiv j - 1 = \bar{\aleph}(\xi, \alpha_s), \quad \gamma_j(\alpha_s) = -2\xi - \bar{\aleph}(\xi) = -2\xi - \omega \quad (3.27)$$

To compare with classical result of Refs. [22, 23] one should rewrite these two equations (3.27) in terms of $\tilde{\gamma}_\omega = -\frac{\gamma_j}{2}$ and combine them as follows

$$\omega = \bar{\aleph}\left(\tilde{\gamma}_\omega - \frac{\omega}{2}\right) \quad (3.28)$$

Next, we need to solve this equation at small ω and $\tilde{\gamma} \simeq 0$. With the NLO accuracy one obtains

$$\omega \simeq \bar{\aleph}(\tilde{\gamma}_\omega) - \frac{\omega}{2} \bar{\aleph}'(\tilde{\gamma}_\omega) \simeq \bar{\aleph}(\tilde{\gamma}_\omega) - \frac{1}{2} \bar{\aleph}(\tilde{\gamma}_\omega) \bar{\aleph}'(\tilde{\gamma}_\omega) \quad (3.29)$$

which can be rewritten as

$$\omega = \frac{\alpha_s N_c}{\pi} \left\{ \bar{\chi}(\tilde{\gamma}_\omega) + \frac{\alpha_s N_c}{4\pi} \left[\bar{\delta}(\tilde{\gamma}_\omega) - 2\bar{\chi}(\tilde{\gamma}_\omega) \bar{\chi}'(\tilde{\gamma}_\omega) \right] \right\} \quad (3.30)$$

In the BFKL limit (3.5) the anomalous dimensions are represented as a sum of series’ in $\frac{\alpha_s}{\omega}$

$$\tilde{\gamma}_\omega = \sum a_n \left(\frac{\alpha_s}{\omega}\right)^n + \omega \sum b_n \left(\frac{\alpha_s}{\omega}\right)^n + \omega^2 \sum c_n \left(\frac{\alpha_s}{\omega}\right)^n + \dots \quad (3.31)$$

The NLO BFKL result (3.30) gives an opportunity to get the first two series in this expansion. Unfortunately, we can inverse Eq. (3.30) only perturbatively and get a few a_n ’s and b_n ’s. To this end, we expand the r.h.s. of Eq. (3.30) in powers of $\tilde{\gamma}_\omega$. To compare with result of Ref. [22] we need three terms of the expansion: $\tilde{\gamma}_\omega^{-2}$, $\tilde{\gamma}_\omega^{-1}$, and constant.

The expansion of $\chi(\gamma)$ defined in Eq. (2.14) has the form

$$\chi(\gamma) = \psi(1 - \varepsilon_*) - \gamma_E - \psi(\gamma) - \psi(1 - \varepsilon_* - \gamma) = \frac{1}{\gamma} + 2\varepsilon_*\zeta(3)\gamma + \dots = \frac{1}{\gamma} + O(\gamma) \quad (3.32)$$

because to compare with Ref. [22] we neglect terms $\sim \alpha_s^2\gamma$ in the r.h.s. of Eq. (3.30). The expansion of $\bar{\delta}(\gamma) - 2\bar{\chi}(\gamma)\bar{\chi}'(\gamma)$ is presented in Eq. (5.6) in the Appendix, and combining it with the above expansion one obtains the equation

$$\omega = \frac{\alpha_s N_c}{\pi} \left\{ \frac{1}{\tilde{\gamma}_\omega} + \frac{\alpha_s N_c}{4\pi} \left[-\frac{11}{3} \frac{1}{\tilde{\gamma}_\omega^2} + 2\zeta(3) - \frac{395}{27} + \frac{11}{18}\pi^2 + O(\tilde{\gamma}_\omega) \right] \right\} \quad (3.33)$$

which is literally equation (23) from Ref. [22] at $n_f = 0$. For completeness, let us present the corresponding anomalous dimension

$$\tilde{\gamma}_\omega = \frac{\alpha_s N_c}{\omega} + O\left(\frac{\alpha_s}{\omega}\right)^4 + \omega \left[-\frac{11}{12} \frac{\alpha_s N_c}{\pi\omega} - \left(\frac{\alpha_s N_c}{\pi\omega}\right)^3 \frac{1}{4} \left(\frac{395}{27} - \frac{11}{18}\pi^2 - 2\zeta(3) \right) + O\left(\frac{\alpha_s}{\omega}\right)^4 \right] \quad (3.34)$$

We see that at the critical point $d = 4 - 2\varepsilon_*$ we get the same anomalous dimensions as at $d = 4$ in accordance with analysis of Refs. [11–13]. Incidentally, both the corrections proportional to b_0 in QCD result and correction $\sim \varepsilon_*$ do not contribute to Eq. (3.34) so the only difference with QCD calculation of Ref. [22] is the origin of the subtraction $\bar{\delta}(\tilde{\gamma}_\omega) - 2\bar{\chi}(\tilde{\gamma}_\omega)\bar{\chi}'(\tilde{\gamma}_\omega)$ in Eq. (3.30). In classical momentum-space calculation of Ref. [22], this subtraction comes from the change of the energy scale from the symmetric scale QP to non-symmetric Q^2 . Instead, in the coordinate space it comes directly from the symmetric “energy scale” R , through the identification $\gamma_\omega = \xi + \frac{\bar{N}(\xi)}{2}$ in Eq. (3.27) coming from the comparison of two representations of 4-point CF (3.1) in the Regge limit.

4 Conclusions

The main result of the paper is that high-energy amplitudes at QCD Wilson-Fisher point (2.2) are determined by the Mobius invariant BFKL/BK equation, at least up to NLO level. It should be emphasized that this result is not a trivial consequence of the conformal invariance of WF QCD because BK equation describes evolution of color dipoles with respect to rapidity cutoff, and the rapidity cutoff violates conformal invariance.⁸ As we mentioned in the introduction, the fact that high-energy amplitudes of WF QCD are determined by (pertubatively calculable) BFKL pomeron opens up a whole machinery of CFTs at high energies. We considered one example: calculation of anomalous dimensions of twist-two LR operators in the BFKL limit (3.5). Another possible application worth exploring is the relation between high-energy amplitudes and non-global logs which was used in Refs. [8, 36] to get the NNLO BK kernel in $\mathcal{N} = 4$ SYM⁹.

Of course, the next question is how to relate results obtained at the WF point to real pQCD. In Sect. 2 we explained how to find terms proportional to b_0 in the NLO result by performing the LO calculations at the LO level. One may hope to find terms proportional

⁸It is safe to say that the rapidity cutoff respecting conformal invariance, if it exists, has not been discovered up to now.

⁹Simon Caron-Huot, private communication.

to b_0 in the NNLO BK kernel is a similar way: calculate $O(\epsilon)$ corrections to NLO BK integrals from Ref. [28] and say that $O(\alpha_s \epsilon_*)$ deviations from the conformal NLO kernel at the WF point are due to $O(\alpha_s^2 b_0)$ terms in NNLO BK kernel. The study is in progress.

The authors are indebted to V. Braun, P. Kravchuk, and A. Manashov for valuable discussions. One of us (I.B.) is grateful to CERN TH for kind hospitality while a part of this work was done. This work of I.B. is supported by DOE contract DE-AC05-06OR23177 and by the grant DE-FG02-97ER41028. The work of G.A.C. is supported by the grant PRIN 2022BP52A MUR "The Holographic Universe for all Lambdas" Lecce-Naples.

5 Appendix: NLO BFKL eigenvalues

The conformal part of NLO BK equation for QCD ‘‘composite dipole’’ of Ref. [29] has the form

$$\begin{aligned}
K_{\text{conf}} = & \frac{\alpha_s^2}{16\pi^4} \int d^2 z_3 d^2 z_4 \frac{z_{12}^2}{z_{13}^2 z_{34}^2 z_{24}^2} \left\{ 2 \ln \frac{z_{12}^2 z_{34}^2}{z_{14}^2 z_{23}^2} + \left[1 + \frac{z_{12}^2 z_{34}^2}{z_{13}^2 z_{24}^2 - z_{14}^2 z_{23}^2} \right] \ln \frac{z_{13}^2 z_{24}^2}{z_{14}^2 z_{23}^2} \right. \\
& \times [2 \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta}\} \text{tr}\{\hat{U}_{z_3}^\eta \hat{U}_{z_4}^{\dagger\eta}\} \text{tr}\{\hat{U}_{z_4}^\eta \hat{U}_{z_2}^{\dagger\eta}\} - \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta} \hat{U}_{z_4}^\eta U_{z_2}^{\dagger\eta} \hat{U}_{z_3}^\eta \hat{U}_{z_4}^{\dagger\eta}\} \\
& \quad - \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_4}^{\dagger\eta} \hat{U}_{z_3}^\eta U_{z_2}^{\dagger\eta} \hat{U}_{z_4}^\eta \hat{U}_{z_3}^{\dagger\eta}\} - (z_4 \rightarrow z_3)] \\
& + \frac{\alpha_s^2}{16\pi^4} \int \frac{d^2 z_3 d^2 z_4}{z_{34}^4} \left\{ -2 + \frac{z_{13}^2 z_{24}^2 + z_{23}^2 z_{14}^2 - 4z_{12}^2 z_{34}^2}{z_{13}^2 z_{24}^2 - z_{14}^2 z_{23}^2} \ln \frac{z_{13}^2 z_{24}^2}{z_{14}^2 z_{23}^2} + (z_3 \leftrightarrow z_4) \right\} \\
& \times [(\text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta}\} \text{tr}\{\hat{U}_{z_3}^\eta \hat{U}_{z_4}^{\dagger\eta}\} \text{tr}\{\hat{U}_{z_4}^\eta \hat{U}_{z_2}^{\dagger\eta}\} - \text{tr}\{\hat{U}_{z_1}^\eta \hat{U}_{z_3}^{\dagger\eta} \hat{U}_{z_4}^\eta \hat{U}_{z_2}^{\dagger\eta} \hat{U}_{z_3}^\eta \hat{U}_{z_4}^{\dagger\eta}\}) - (z_4 \rightarrow z_3)] \\
& + \frac{\alpha_s^2 n_f}{2\pi^4} \int \frac{d^2 z_3 d^2 z_4}{z_{34}^4} \left\{ 2 - \frac{z_{13}^2 z_{24}^2 + z_{23}^2 z_{14}^2 - z_{12}^2 z_{34}^2}{z_{13}^2 z_{24}^2 - z_{14}^2 z_{23}^2} \ln \frac{z_{13}^2 z_{24}^2}{z_{14}^2 z_{23}^2} \right\} \\
& \times \text{tr}\{t^a \hat{U}_{z_1}^\eta t^b \hat{U}_{z_2}^{\dagger\eta}\} \text{tr}\{t^a \hat{U}_{z_3}^\eta t^b (\hat{U}_{z_4}^{\dagger\eta} - \hat{U}_{z_3}^\eta)\} \tag{5.1}
\end{aligned}$$

We need also the linearized version of this kernel in the ‘‘forward’’ case when $\langle \mathcal{U}(x, y) \rangle = \mathcal{U}(x - y)$:

$$\mathcal{K}_{\text{conf}}(z, z') = \frac{z^2}{z'^2} \left[-\frac{1}{(z - z')^2} \ln^2 \frac{z^2}{z'^2} + \Phi(z, z') + 6\pi\zeta(3)\delta^{(2)}(z - z') + F(z, z') \right] \tag{5.2}$$

where $F(z, z')$ and $\Phi(z, z')$ are given by Eqs. (55) and (56) from Ref. [37].

$$\begin{aligned}
\Phi(z, z') = & \frac{z^2}{z'^2} \left\{ \frac{(z^2 - z z'^2)}{(z - z')^2 (z + z')^2} \left[\ln \frac{z^2}{z'^2} \ln \frac{z^2 z'^2 (z - z')^4}{(z^2 + z'^2)^4} + 2\text{Li}_2\left(-\frac{z'^2}{z^2}\right) \right. \right. \\
& \left. \left. - 2\text{Li}_2\left(-\frac{z^2}{z'^2}\right) \right] - \left(1 - \frac{(z^2 - z'^2)^2}{(z - z')^2 (z + z')^2} \right) \left[\int_0^1 - \int_1^\infty \right] \frac{du}{(z - z'u)^2} \ln \frac{u^2 z'^2}{z^2} \right\} \tag{5.3}
\end{aligned}$$

$$\begin{aligned}
F(z, z') = & \frac{z^2}{z'^2} \left\{ \left(1 + \frac{n_f}{N_c^3} \right) \frac{3(z, z')^2 - 2z^2 z'^2}{16z^2 z'^2} \left(\frac{2}{z^2} + \frac{2}{z'^2} + \frac{z^2 - z'^2}{z^2 z'^2} \right) \ln \frac{z^2}{z'^2} - \left[3 \right. \right. \\
& \left. \left. + \left(1 + \frac{n_f}{N_c^3} \right) \left(1 - \frac{(z^2 + z'^2)^2}{8z^2 z'^2} + \frac{3z^4 + 3z'^4 - 2z^2 z'^2}{16z^4 z'^4} (z, z')^2 \right) \right] \int_0^\infty dt \frac{1}{z^2 + t^2 z'^2} \ln \frac{1+t}{|1-t|} \right\} \tag{5.4}
\end{aligned}$$

It should be noted that the first term in Eq. (5.1) and the first three terms in Eq. (5.2) are actually the $\mathcal{N} = 4$ result which has maximal transcendentality in comparison to the

remainder of the QCD result. This is especially clear if one writes down the corresponding eigenvalues

$$\begin{aligned}
\frac{1}{\pi} \int d^2 z' \left(\frac{z'^2}{z^2}\right)^{\xi-1} \frac{\ln^2 \frac{z'^2}{z^2}}{(z'-z)^2} &= \chi''(\xi), \quad \chi''(\xi) + 2\chi(\xi)\chi'(\xi) = 8\zeta(3) + O(\xi) \\
\frac{1}{\pi} \int d^2 z \Phi(z, z') \left(\frac{z'^2}{z^2}\right)^\xi &= -4[S_{-2,1}(\xi-1) + S_{-2,1}(-\xi) + \frac{5}{4}\zeta(3)] + \frac{\pi^3}{\sin \pi \xi} \\
&= \frac{\pi^2}{3\xi} + 4\zeta(3) + O(\xi) \\
F(\xi) &= \frac{1}{\pi} \int d^2 z' F(z, z') \left(\frac{z'^2}{z^2}\right)^\xi = -\left[3 + \left(1 + \frac{n_f}{N_c^3}\right) \frac{2 + 3\xi(1-\xi)}{(3-2\xi)(1+2\xi)}\right] \frac{\pi^2 \cos \pi \xi}{(1-2\xi) \sin^2 \pi \xi}, \\
&= -\frac{1}{\xi^2} \left(\frac{11}{3} + \frac{2n_f}{3N_c^3}\right) - \frac{1}{\xi} \left(\frac{67}{9} + \frac{13n_f}{9N_c^3}\right) - \frac{395}{27} + \frac{11}{18}\pi^2 - \frac{71n_f}{27N_c^3} + \frac{\pi^2 n_f}{9N_c^3} + O(\xi) \quad (5.5)
\end{aligned}$$

The representation of the r.h.s. of the second Eq. (5.5) in terms of harmonic sums was obtained in Ref. [14].

Thus, $\bar{\delta}(\xi)$, defined by Eq. (2.14), has the form

$$\begin{aligned}
\bar{\delta}(\xi) - 2\bar{\chi}(\xi)\bar{\chi}'(\xi) &= \chi(\xi) \left(\frac{67}{9} - \frac{\pi^2}{3} - \frac{10n_f}{9N_c}\right) - \chi''(\xi) + F(\xi) + \Phi(\xi) + 6\zeta(3) - 2\chi(\xi)\chi'(\xi) \\
&= -\frac{1}{\xi^2} \left(\frac{11}{3} + \frac{2n_f}{3N_c^3}\right) - \frac{1}{\xi} \left(\frac{10n_f}{9N_c} + \frac{13n_f}{9N_c^3}\right) + 2\zeta(3) - \frac{395}{27} + \frac{11}{18}\pi^2 - \frac{71n_f}{27N_c^3} + \frac{\pi^2 n_f}{9N_c^3} \quad (5.6)
\end{aligned}$$

6 Appendix: gluon light-ray operators

First, we need to define a light-ray operator as an analytic continuation of a local operator with spin j . As explained in Ref. [17], a proper way is to continue in spin a non-local object rather than the local one. A suitable non-local object is a “light transform” of a local operator - integral of the local operator along the null line with some weight [17]. In our case, we define “forward” local operator as

$$\mathcal{O}_j^{(+)}(z_\perp) = \int dz_+ F_{-i}^a \nabla_-^{j-2} F_-^{ai}(z_+, 0, z_\perp) \quad (6.1)$$

which is a specific case of the light transform, namely the integral over null line with unit weight. The analytic continuation of Eq. (6.1) in spin j has the form

$$\begin{aligned}
\mathcal{F}_j^{(+)}(z_\perp) &= \int dz_+ \int dz'_+ \left[\frac{i\Gamma(j-1)}{2\pi(z'_+ + i\epsilon)^{j-1}} + \frac{i\Gamma(j-1)}{2\pi(-z'_+ + i\epsilon)^{j-1}} \right] \\
&\times F_{-i}^a(z_+ + z'_+, 0, z_\perp) [z_+ + z'_+ + z_\perp, z_+ + z_\perp]^{ab} F_-^{bi}(z_+, 0, z_\perp) \quad (6.2)
\end{aligned}$$

where

$$[x, y] \equiv \text{Pexp} \, ig \int_0^1 du (x-y)^\mu A_\mu(ux + (1-u)y)$$

is a standard notation for straight-line gauge link connection points x and y . It is easy to see that at even integer j the expression in square brackets in Eq. (6.2) is $\left(\frac{d}{dz'_+}\right)^{j-2} \delta(z'_+)$ so after $(j-2)$ integrations by parts we return to “forward” local operator (6.1).

We will need also the light-ray operator integrated over z_- direction

$$\begin{aligned} \mathcal{F}_j^{(-)}(z_\perp) &= \int dz_- \int dz'_- \left[\frac{i\Gamma(j-1)}{2\pi(z'_- + i\epsilon)^{j-1}} + \frac{i\Gamma(j-1)}{2\pi(-z'_- + i\epsilon)^{j-1}} \right] \\ &\times F_{+i}^a(z_- + z'_-, 0, z_\perp) [z_- + z'_- + z_\perp, z_- + z_\perp]^{ab} F_+^{bi}(z_-, 0, z_\perp) \end{aligned} \quad (6.3)$$

Let us now demonstrate that the anomalous dimension of the LR operator (6.2) is an analytic continuation of γ_j of local operator (6.1). First, let us rewrite the operator (6.2) as follows

$$\mathcal{F}_j^{(+)}(z_\perp) = \frac{i\Gamma(j-1)}{\pi} (1 - e^{-i\pi j}) \int_0^\infty dL_+ L_+^{1-j} F(L_+, z_\perp) \quad (6.4)$$

where

$$F(L_+, z_\perp) = \int dz_+ F_{-i}^a(L_+ + z_+, 0, z_\perp) [z_+ + z'_+ + z_\perp, z_+ + z_\perp]^{ab} F_-^{bi}(z_+, 0, z_\perp) \quad (6.5)$$

is the ‘‘forward’’ gluon LR operator defined in Ref. [3].

Evolution equation for the LR operator (6.5) has the form (see e.g. Ref. [3])

$$\mu \frac{d}{d\mu} F(L_+, x_\perp) = \int_0^1 du K_{gg}(u, \alpha_s) F(uL_+, x_\perp) \quad (6.6)$$

where $K_{gg}(u, \alpha_s)$ is the DGLAP kernel in gluodynamics. The twist-2 operators (6.1) are obtained by expansion of LR operator (6.5) in powers of L_+ , so their anomalous dimensions are given by

$$\gamma_j(\alpha_s) = - \int_0^1 du u^{j-2} K_{gg}(u, \alpha_s) \quad \mu \frac{d}{d\mu} \mathcal{O}_j^{(+)} = - \gamma_j(\alpha_s) \mathcal{O}_j^{(+)} \quad (6.7)$$

Now consider the anomalous dimensions of LR operator (6.4). Combining Eqs. (6.5) and (6.6) one obtains

$$\mu \frac{d}{d\mu} \mathcal{F}_j(z_\perp) = \int_0^1 du K_{gg}(u, \alpha_s) u^{j-2} \mathcal{F}_j(z_\perp) \quad (6.8)$$

Thus, we see that the above equation gives the explicit formula for analytic continuation of the anomalous dimensions to non-integer spins.

References

- [1] I. Balitsky. Operator expansion for high-energy scattering. *Nucl. Phys. B*, 463:99–160, 1996.
- [2] Yuri V. Kovchegov. Small x F(2) structure function of a nucleus including multiple pomeron exchanges. *Phys. Rev.*, D60:034008, 1999.
- [3] I. I. Balitsky and Vladimir M. Braun. Evolution Equations for QCD String Operators. *Nucl. Phys. B*, 311:541–584, 1989.
- [4] L. N. Lipatov. The Bare Pomeron in Quantum Chromodynamics. *Sov. Phys. JETP*, 63:904–912, 1986.

- [5] A. V. Kotikov and L. N. Lipatov. DGLAP and BFKL equations in the $N = 4$ supersymmetric gauge theory. *Nucl. Phys. B*, 661:19–61, 2003. [Erratum: Nucl.Phys.B 685, 405–407 (2004)].
- [6] A. V. Kotikov. The property of maximal transcendentality in the $N=4$ SYM. *Phys. Part. Nucl.*, 41:951–953, 2010.
- [7] Nikolay Gromov, Fedor Levkovich-Maslyuk, and Grigory Sizov. Pomeron Eigenvalue at Three Loops in $\mathcal{N} = 4$ Supersymmetric Yang-Mills Theory. *Phys. Rev. Lett.*, 115(25):251601, 2015.
- [8] Simon Caron-Huot and Matti Herranen. High-energy evolution to three loops. *JHEP*, 02:058, 2018.
- [9] V. N. Velizhanin. BFKL pomeron in the next-to-next-to-leading approximation in the planar $N=4$ SYM theory. 8 2015.
- [10] Kenneth G. Wilson and Michael E. Fisher. Critical exponents in 3.99 dimensions. *Phys. Rev. Lett.*, 28:240–243, 1972.
- [11] V. M. Braun, A. N. Manashov, S. Moch, and M. Strohmaier. Two-loop conformal generators for leading-twist operators in QCD. *JHEP*, 03:142, 2016.
- [12] V. M. Braun, A. N. Manashov, S. Moch, and M. Strohmaier. Three-loop evolution equation for flavor-nonsinglet operators in off-forward kinematics. *JHEP*, 06:037, 2017.
- [13] Yao Ji, Alexander Manashov, and Sven-Olaf Moch. Evolution kernels of twist-two operators. *Phys. Rev. D*, 108(5):054009, 2023.
- [14] Miguel S. Costa, Vasco Goncalves, and Joao Penedones. Conformal Regge theory. *JHEP*, 12:091, 2012.
- [15] Miguel S. Costa, James Drummond, Vasco Goncalves, and Joao Penedones. The role of leading twist operators in the Regge and Lorentzian OPE limits. *JHEP*, 04:094, 2014.
- [16] Miguel S. Costa, Vasco Goncalves, Aaditya Salgarkar, and Joao Vilas Boas. Conformal multi-Regge theory. *JHEP*, 09:155, 2023.
- [17] Petr Kravchuk and David Simmons-Duffin. Light-ray operators in conformal field theory. *JHEP*, 11:102, 2018.
- [18] Murat Kologlu, Petr Kravchuk, David Simmons-Duffin, and Alexander Zhiboedov. The light-ray OPE and conformal colliders. *JHEP*, 01:128, 2021.
- [19] Cyuan-Han Chang, Murat Kologlu, Petr Kravchuk, David Simmons-Duffin, and Alexander Zhiboedov. Transverse spin in the light-ray OPE. *JHEP*, 05:059, 2022.
- [20] Simon Caron-Huot, Murat Kologlu, Petr Kravchuk, David Meltzer, and David Simmons-Duffin. Detectors in weakly-coupled field theories. *JHEP*, 04:014, 2023.
- [21] T. Jaroszewicz. Gluonic Regge Singularities and Anomalous Dimensions in QCD. *Phys. Lett. B*, 116:291–294, 1982.
- [22] Victor S. Fadin and L. N. Lipatov. BFKL pomeron in the next-to-leading approximation. *Phys. Lett. B*, 429:127–134, 1998.
- [23] Marcello Ciafaloni and Gianni Camici. Energy scale(s) and next-to-leading BFKL equation. *Phys. Lett. B*, 430:349–354, 1998.

- [24] Lorenzo Cornalba. Eikonal methods in AdS/CFT: Regge theory and multi-reggeon exchange. 10 2007.
- [25] Tom Banks and A. Zaks. On the Phase Structure of Vector-Like Gauge Theories with Massless Fermions. *Nucl. Phys. B*, 196:189–204, 1982.
- [26] Alfred H. Mueller. Soft gluons in the infinite momentum wave function and the BFKL pomeron. *Nucl. Phys. B*, 415:373–385, 1994.
- [27] Alfred H. Mueller and Bimal Patel. Single and double BFKL pomeron exchange and a dipole picture of high-energy hard processes. *Nucl. Phys. B*, 425:471–488, 1994.
- [28] Ian Balitsky and Giovanni A. Chirilli. Next-to-leading order evolution of color dipoles. *Phys. Rev. D*, 77:014019, 2008.
- [29] Ian Balitsky and Giovanni A. Chirilli. NLO evolution of color dipoles in N=4 SYM. *Nucl. Phys. B*, 822:45–87, 2009.
- [30] V. M. Braun and A. N. Manashov. Evolution equations beyond one loop from conformal symmetry. *Eur. Phys. J. C*, 73:2544, 2013.
- [31] Ian Balitsky, Vladimir Kazakov, and Evgeny Sobko. Two-point correlator of twist-2 light-ray operators in N=4 SYM in BFKL approximation. *Nucl. Phys. B*, 993:116267, 2023.
- [32] Ian Balitsky. NLO BFKL and anomalous dimensions of light-ray operators. *Int. J. Mod. Phys. Conf. Ser.*, 25:1460024, 2014.
- [33] Lorenzo Cornalba, Miguel S. Costa, and Joao Penedones. Eikonal Methods in AdS/CFT: BFKL Pomeron at Weak Coupling. *JHEP*, 06:048, 2008.
- [34] Ian Balitsky and Giovanni A. Chirilli. High-energy amplitudes in N=4 SYM in the next-to-leading order. *Phys. Lett.*, B687:204–213, 2010.
- [35] Ian Balitsky. Structure constants of twist-two light-ray operators in the triple Regge limit. *JHEP*, 04:042, 2019.
- [36] Simon Caron-Huot. Resummation of non-global logarithms and the BFKL equation. *JHEP*, 03:036, 2018.
- [37] Ian Balitsky and Giovanni A. Chirilli. Photon impact factor and k_T -factorization for DIS in the next-to-leading order. *Phys. Rev. D*, 87(1):014013, 2013.