Excitations of the Higgs boson

Gregory A. Wright^{*}

Nokia Bell Laboratories, 600 Mountain Avenue, Murray Hill, NJ 07974

(Dated: July 12, 2024)

In a simplified version of the standard model with a single quark doublet, I derive a transcendental equation for the complex Higgs boson mass. The equation involves a divergent integral which is regularized and renormalized conventionally. Setting the Higgs mass to its observed value, the decay width is narrow and within experimental limits. The mass equation has additional roots on other sheets in the complex energy plane. The first two are at 186 and 219 GeV. The lightest excitations are well defined though wider than the 125 GeV Higgs because decays to W^+W^- and ZZ are allowed.

In this letter I describe a simplified version of the standard model in which electroweak symmetry is dynamically broken by a gravitationally bound condensate of fermions. The collective modes of this condensate are three phase excitations, corresponding to Goldstone bosons, and a massive amplitude excitation which is the Higgs boson. The Goldstone bosons are absorbed by the usual Higgs mechanism [1]. In contrast to the standard model, where the Higgs mass is a free parameter, here it is determined by a transcendental equation with a well isolated smallest root and an infinite number of additional solutions accumulating at $\sqrt{2}v$, where v is the Higgs vacuum expectation value.

This work extends the old idea that fermion mass generation occurs analogously as in superconductivity [2, 3]. The new idea here is that initially massless fermions are bound in a condensate by their mutual gravitational attraction. Because the gravitational force is both longrange and unscreened, even though it is weak, it can cause the formation of a correlated ground state (as long as cosmological considerations can be neglected). Also, every fermion is attracted to the condensate with a strength equal to its radiatively corrected gravitational coupling. This provides a natural explanation for why all fermions have masses. Radiative corrections allow the fermions to have different masses, even though the model is invariant under $SU(2)_W \times U(1)_Y$. The model requires no nonlinear gravitational effects, beyond the radiative corrections of the graviton-fermion vertex by vector gauge loops, to produce the effects described here, and in particular no third-order or higher gravitational interaction. A shortcoming of the model is that as yet it provides no explanation for fermion flavor.

The model is defined by a lagrangian for two massless fermions, t and b. In what follows, gauge fields play no role and are omitted. I will use the momentum representation from the outset, with the fermion fields in the Weyl representation. The integral of the lagrangian is assumed to be the action within a functional integral. The trace minus two Minkowski metric is used throughout and $\hbar = c = 1$. The right and left-handed Weyl matrices are $\sigma^{\mu} = (\mathbb{1}, \boldsymbol{\sigma}), \ \overline{\sigma}^{\mu} = (\mathbb{1}, -\boldsymbol{\sigma})$, respectively, where $\mathbb{1}$ is the 2 × 2 identity matrix and $\boldsymbol{\sigma}$ are the Pauli matrices. For brevity, the momentum measure is written $dp = d^4 p/(2\pi)^4$ and the delta function in momentum space is $\delta(p) = (2\pi)^4 \delta^{(4)}(p)$.

The kinetic part of the lagrangian is

$$\mathscr{L}_{\text{fermi}} = t_L^{\dagger}(\overline{\sigma} \cdot p)t_L + t_R^{\dagger}(\sigma \cdot p)t_R + b_L^{\dagger}(\overline{\sigma} \cdot p)b_L + b_R^{\dagger}(\sigma \cdot p)b_R, \qquad (1)$$

where from now on the two fermions will be referred to as the top and bottom quarks. The validity of this interpretation will be shown.

As in the Bardeen-Cooper-Schrieffer (BCS) model of superconductivity [4], an effective quartic interaction approximates the interaction between the fermions:

$$\mathcal{L}_{\text{quartic}} = -f_t^2 \left[\alpha \left(t_R^{\dagger} t_L t_L^{\dagger} t_R + b_L^{\dagger} t_R t_R^{\dagger} b_L \right) + \\ \beta \left(b_L^{\dagger} b_R t_L^{\dagger} t_R + t_R^{\dagger} t_L b_R^{\dagger} b_L \\ - b_R^{\dagger} t_L t_R^{\dagger} b_L - b_L^{\dagger} t_R t_L^{\dagger} b_R \right) + \\ \gamma \left(b_L^{\dagger} b_R b_R^{\dagger} b_L + b_R^{\dagger} t_L t_L^{\dagger} b_R \right) \right].$$

$$(2)$$

The three terms in parentheses are separately invariant under $SU(2)_W \times U(1)_Y$. Because the action is a lorentz scalar, pairs of fermi fields must have opposite handedness (e.g., $t_I^{\mathsf{T}} b_R$). For the moment, the parameters α, β and γ are arbitrary, dimensionless numbers. The overall coupling constant f_t^2 is the product of a dimensionless ratio of momenta (two factors of momentum in the numerator from the quark energy-momentum tensors and a momentum squared in the denominator from the graviton propagator) and the gravitational constant with dimension -2. Following BCS, the momentum dependence of the coupling will be ignored: f_t^2 is taken as constant up to some cutoff, above which it is zero. Taking the coupling to be momentum independent is equivalent to swave pairing of the condensate. Each pair of Weyl spinors has color indices contracted to form a color singlet. Color indices are suppressed throughout but the color factor is provided in the final result.

The interaction (2) implicitly assumes that a condensate forms. The gravitational interaction is diagonal in left and right-handed fields, so if the interaction were perturbative, in each term the t or b fields would appear with the same handedness. The terms multiplied by β violate this condition and require the condensate to act as a reservoir of the appropriate helicity states. When a condensate is present all terms can be generated, as shown schematically in Fig. 1.



FIG. 1. The gravitational interactions that give the effective lagrangian terms in equation (2). Double wiggly lines are gravitons, blobs are the radiatively corrected gravitonfermion vertices, and the shaded area at the bottom denotes the condensate. Panel (a) shows the interactions that, after a Fierz rearrangement give the first term of equation (2). Panel (b) illustrates the interactions that give the sixth and seventh terms of (2). Because the calculation is nonperturbative, the contribution of (b) is not necessarily less than that of (a).

For the quartic interaction (2) to be diagonalizable by a Hubbard-Stratonovich shift, the constants α , β and γ must obey $\gamma/\alpha = (\beta/\alpha)^2$. This allows factoring the quartic into products of quark and scalar fields whose couplings reproduce those of the standard model. The radiative corrections due to weak hypercharge, to lowest order, satisfy this condition. To see this, consider the hypercharge factor in the correction to the gravitational vertices of Fig. 1(a). This correction gives the factor α . The left vertex contributes a factor of $(4/3)^2$ and the right vertex $(1/3)^2$, their product gives $\alpha = 16/81$. (All radiative corrections have a common momentum factor which is absorbed into the overall coupling, as is the interaction with the condensate.) Substituting the corresponding fields from the last term of (2), the diagram Fig. 1(a) gives $\gamma = 4/81$. The correction factor β is from the diagram in Fig. 1(b). The product of hypercharges in that diagram is $(8/81)^2$. This factor is shared between two of the middle terms of (2), and I divide it equally between the terms, giving $\beta = 8/81$. These assignments of α , β and γ satisfy the required condition.

This argument is the least satisfactory part of what is presented here. It is rooted in perturbation theory, and the rest of the calculation is non-perturbative. However, it is very suggestive and the results below indicate that it contains some truth. I conjecture that the full radiative corrections of weak hypercharge follow the same pattern. In what follows, I extract a factor of α , absorbing it into the overall coupling f_t^2 , and set $\beta/\alpha = \eta$ and $\gamma/\alpha = \eta^2$, for some positive η . The constant η will be the ratio of the bottom quark mass to the top quark masses.

Diagonalizing this model starts by adding scalar fields

to the lagrangian,

$$\mathscr{L} \to \frac{\phi_0^{\mathsf{T}} \phi_0 + \phi_+ \phi_-}{f_t^2} + \mathscr{L}. \tag{3}$$

In principle, the new term could be multiplied by a constant $\xi > 0$ without changing any results. I will show elsewhere that if this model is parameterized so that it reproduces the lowest order results of the standard model, for example, $m_W = gv/2$, then $\xi = 1$. Shifting the scalar fields eliminates the quartic term:

$$\begin{aligned} \phi_0 &\to \phi_0 - f_t^2 \left(t_L^{\dagger} t_R + \eta \, b_R^{\dagger} b_L \right) , \\ \phi_0^{\dagger} &\to \phi_0^{\dagger} - f_t^2 \left(t_R^{\dagger} t_L + \eta \, b_L^{\dagger} b_R \right) , \\ \phi_+ &\to \phi_+ + f_t^2 \left(b_L^{\dagger} t_R - \eta \, b_R^{\dagger} t_L \right) , \\ \phi_- &\to \phi_- + f_t^2 \left(t_R^{\dagger} b_L - \eta \, t_L^{\dagger} b_R \right) . \end{aligned}$$

$$(4)$$

The shifted fields are chosen so that $\Phi = (\phi_+, \phi_0)^T$ transforms under $SU(2)_W \times U(1)_Y$ with the correct quantum numbers. Using cartesian components and splitting the Higgs field split into the sum of the observable field H and the vacuum expectation value v,

$$\phi_0 = \frac{H + v + iw_3}{\sqrt{2}}, \qquad \phi_+ = \frac{w_1 + iw_2}{\sqrt{2}}, \quad (5)$$

the lagrangian becomes

$$\begin{aligned} \mathscr{L} &= \mathscr{L}_{\text{fermi}} + \frac{H^2 + 2Hv + v^2 + w_i w_i}{2 f_t^2} - \\ &\frac{v}{\sqrt{2}} \left(t_R^{\dagger} t_L + t_L^{\dagger} t_R \right) - \frac{\eta v}{\sqrt{2}} \left(b_R^{\dagger} b_L + b_L^{\dagger} b_R \right) - \\ \left\{ \frac{H}{\sqrt{2}} \left[t_R^{\dagger} t_L + t_L^{\dagger} t_R + \eta \left(b_R^{\dagger} b_L + b_L^{\dagger} b_R \right) \right] + \\ &i \frac{w_3}{\sqrt{2}} \left[t_R^{\dagger} t_L - t_L^{\dagger} t_R + \eta \left(b_L^{\dagger} b_R - b_R^{\dagger} b_L \right) \right] - \\ &\frac{w_1}{\sqrt{2}} \left[t_R^{\dagger} b_L + b_L^{\dagger} t_R - \eta \left(t_L^{\dagger} b_R + b_R^{\dagger} t_L \right) \right] - \\ &i \frac{w_2}{\sqrt{2}} \left[t_R^{\dagger} b_L - b_L^{\dagger} t_R - \eta \left(t_L^{\dagger} b_R - b_R^{\dagger} t_L \right) \right] \right\}. \end{aligned}$$
(6)

It is apparent that at the semiclassical level the top and bottom quarks acquire masses of $v/\sqrt{2}$ and $\eta v/\sqrt{2}$, respectively, justifying the statement that η is ratio of the quark masses, and the interpretation of the t and b fields as the top and bottom quarks.

As yet, the scalar excitations of the condensate have no kinetic terms. Well below the transition temperature, I can integrate out the fermions to get an effective action for the scalars alone, which will have the expected kinetic terms.

Collecting the fermions into a Nambu spinor,

$$\Psi(p) = [t_L(p), t_R(p), b_L(p), b_R(p)]^T,$$
(7)

the action is

$$S = \int \widetilde{dp} \, \widetilde{dp'} \, \Psi^{\dagger}(p) M(p,p') \Psi(p') \,, \tag{8}$$

where

$$M(p, p') = G^{-1}(p) + u(p, p'),$$

$$G^{-1}(p) = \operatorname{diag}\left[\overline{\sigma} \cdot p, \sigma \cdot p, \overline{\sigma} \cdot p, \sigma \cdot p\right],$$
(9)

and u =

$$\begin{bmatrix} 0 & -\frac{H+v+iw_3}{\sqrt{2}} & 0 & \frac{\eta(w_1-iw_2)}{\sqrt{2}} \\ -\frac{H+v-iw_3}{\sqrt{2}} & 0 & -\frac{w_1-iw_2}{\sqrt{2}} & 0 \\ 0 & -\frac{w_1+iw_2}{\sqrt{2}} & 0 & -\frac{\eta(H+v-iw_3)}{\sqrt{2}} \\ \frac{\eta(w_1+iw_2)}{\sqrt{2}} & 0 & -\frac{\eta(H+v+iw_3)}{\sqrt{2}} & 0 \end{bmatrix} .$$
(10)

Integrating out the fermi field Ψ gives an action for the scalars:

$$S_{\text{eff}} = \int \widetilde{dp} \, \frac{1}{2f_t^2} (H^2 + 2Hv + v^2 + w_i w_i) -$$

$$i \, \text{Tr} \log(\mathbb{1} + Gu) \,, \qquad (11)$$

where a factor of $G^{-1}(p)$ has been used to make the argument of the logarithm dimensionless at the expense of a change in the normalization of the functional integral. The 'Tr' symbol denotes the combined operator and matrix trace, 'tr' will be the matrix trace alone.

It is possible to treat Gu in equation (11) as small and expand the logarithm. However, this would be an expansion about the symmetric vacuum, valid only for energies near the electroweak transition. To find the low energy effective action (and therefore the observable particle spectrum), an expansion about the true (condensed) vacuum is needed. The first step is to find the condensate density by extremizing the action with respect to v:

$$0 = \frac{\delta S_{\text{eff}}}{\delta v} = \int \widetilde{dp} \left\{ \frac{v}{f_t^2} - i N_c \langle p | \operatorname{tr} \left[\Delta \left. \frac{\delta u}{\delta v} \right|_0 \right] | p \rangle \right\},$$
(12)

where $\Delta = (\mathbb{1} + Gu_0)^{-1}G$, N_c is the number of quark colors and a zero subscript indicates evaluation at $H = w_i = 0$. Inserting a resolution of the identity as free particle momentum states between Δ and $\delta u/\delta v$ gives

$$0 = \frac{1}{2N_c f_t^2} + i \int \widetilde{dp} \left[\frac{1}{p^2 - \frac{v^2}{2}} + \frac{\eta^2}{p^2 - \frac{\eta^2 v^2}{2}} \right].$$
 (13)

This is the equation for v, analogous to the gap equation of superconductivity. The integral is divergent, but it is not necessary to evaluate it. Instead, in what follows, it is used to eliminate f_t , which is not accessible to measurement, in favor of v, which is. The kinetic term for the Higgs field is the piece quadratic in H in the expansion of the effective action,

$$\int \tilde{d}r \tilde{d}s \, \frac{1}{2} H(r) \frac{\delta^2 S_{\text{eff}}}{\delta H(r) \delta H(s)} H(s) \,, \tag{14}$$

which, in the parameterization of the standard model, is

$$\int \tilde{dr} \, \frac{Z(r^2)}{2} H(-r)(r^2 - M_H^2) H(r) \,. \tag{15}$$

Because the kinetic term is generated by quantum effects, a field renormalization factor, Z, must be anticipated.

The functional derivative is

$$\frac{\delta^2 S_{\text{eff}}}{\delta H(r) \delta H(s)} = \frac{1}{f_t^2} + i \int \widetilde{dp} \langle p | \operatorname{tr} \left[\Delta(p) \frac{\delta u}{\delta H(r)} \Delta(p-r) \frac{\delta u}{\delta H(s)} \right] | p \rangle .$$
(16)

The first term on the right hand side can be eliminated by using the gap equation and the integral reduced to Feynman integrals as the gap equation was, giving

$$\frac{\delta^2 S_{\text{eff}}}{\delta H(r)\delta H(s)} = -i\,\tilde{\delta}(r+s)\int_0^1 dx \int \widetilde{dp} \quad \frac{r^2 - 2v^2}{\left[p^2 - \frac{v^2}{2} + r^2x(1-x)\right]^2} + \frac{\eta^2(r^2 - 2\eta^2v^2)}{\left[p^2 - \frac{\eta^2v^2}{2} + r^2x(1-x)\right]^2}.$$
(17)

If $\eta = 0$, the bottom quark decouples and the above has a real root $r = \pm \sqrt{2}v$, twice the mass of the top quark. If $\eta \neq 0$, it has no roots on the physical sheet, but there are complex roots on other Riemann sheets. The roots come in groups of four, with equal magnitude: positive and negative energy, and growing and decaying modes (these quartets of roots were observed for the collective modes of BCS by Andrianov and Popov [5]). The complex roots are on unphysical sheets because unitarity requires that the action be real [6].

The integral in (17) is regularized using a Pauli-Villars parameter Λ . Introducing dimensionless parameters $\rho^2 = r^2/v^2$ and $m_H^2 = M_H^2/v^2$, and equating the standard model Higgs kinetic term (15) to that of the effective model (16), gives

$$Z(\rho^{2})(\rho^{2} - m_{H}^{2}) = (\rho^{2} - 2)I_{n}(\rho, 1) + \eta^{2}(\rho^{2} - 2\eta^{2})I_{n}(\rho, \eta) + (18)$$
$$C_{\Lambda},$$

where C_{Λ} is the regularized divergent term and the finite part of the integral is

$$I_n(\rho,\eta) = \log\left(\eta^2/2\right) - 2 + \frac{2}{\rho}\sqrt{2\eta^2 - \rho^2} \left[\arctan\left(\frac{\rho}{\sqrt{2\eta^2 - \rho^2}}\right) + n\pi\right].$$
 (19)



FIG. 2. Higgs partial width to $b\bar{b}$ as a function of the effective mass ratio $\eta = m_t/m_b$. The dashed horizontal line is the partial width of the standard model Higgs to $b\bar{b}$ taken from [9].

In this expression, n is the sheet index. Using on-shell renormalization, the equation whose root determines the Higgs mass is

$$0 = (\rho^2 - 2)I_n(\rho, 1) + \eta^2(\rho^2 - 2\eta^2)I_n(\rho, \eta) - \operatorname{Re}\left[(m_H^2 - 2)I_1(m_H, 1) + \eta^2(m_H^2 - 2\eta^2)I_1(m_H, \eta)\right].$$
(20)

The second line of the above is the counterterm that fixes the real part of the excitation on the first unphysical sheet to be the mass of the observed Higgs boson. (The imaginary part can not be specified since it is determined by unitarity.) Note that this expression is independent of the number of quark colors, N_c . The Higgs mass does depend on the number of colors, but this dependence is contained entirely in the vacuum expectation value v.

On the right hand side of equation (20), before the counterterm, are a term multiplied by $\eta = 1$ (the "top term") and another multiplied by $\eta \neq 1$ (the "bottom term"). Roots exist if only top term is off the physical sheet $(n \geq 1)$ or if both the top and bottom terms are off the physical sheet. The roots are close to the real-axis branch cuts for both the bottom and top terms, and the numerical results given below put both top and bottom terms on the same sheet. However, roots still exist if only the top term is on a non-physical sheet and the bottom term remains on the physical (n = 0) sheet. In that case, the real part of the root is only slightly different from when both terms are on the same sheet, but the magnitude of the imaginary part is significantly smaller.

For numerical evaluation, I take the ratio of the bottom to top mass to be $\eta = 0.015$, corresponding to the ratio of the running bottom mass measured at the energy scale of the Higgs boson, $m_b(m_H)$ [7] and the measured top quark pole mass [8]. The typical residual of the numerical root finder is a few parts in 10¹⁵.

The partial width to $b\bar{b}$ for $\eta = 0.015$ is 3.35 MeV, about fifty percent larger the standard model prediction for a pointlike Higgs [10, 11]. It is not surprising that the Higgs decay rate here is different than in the standard model, since here the Higgs is not a fundamental scalar particle. Its mass is about half of the condensate density v, an energy scale at which the constituents of the condensate may begin to be resolved. For other values of η the partial width to $b\bar{b}$ is shown in Fig. (2). The first few roots of (20) for n > 1 are given in Table I, along with their estimated decay widths, assuming a pointlike Higgs with standard model couplings to W^+W^- and ZZ. Numerical examination of the roots of (20) indicates that they are all simple, corresponding to isolated poles in the propagator.

The real parts of the first twenty five roots are plotted in Fig. (3). The top panel plots the masses and the bottom panel shows the "binding energy", the difference between the mass and the asymptotic mass for large sheet index ($\approx 2m_t$). That the binding energy of the Higgs excitations decreases as $n^{-2/3}$ is evidence against interpreting the Higgs as a kind of bound state of valence quarks. This model has no gauge fields and the gravitational interaction is treated in the random phase approximation [12], so there is no potential between fermionic field oscillators. A physical interpretation of the Higgs excitations might be itinerant "bubbles" or dislocations in the ordered ground state.

What I have done above parallels the work of Andri-



FIG. 3. Panel (a): Masses of the first twenty five Higgs excitations vs. sheet index. The point corresponding to n = 1 is the 125 GeV Higgs boson. The dashed horizontal line is at 347.88 GeV, the root of the Higgs mass equation for $n = 10^6$, $\eta = 0.015$. Panel (b): The "binding energy", i.e., difference in mass between a Higgs excitation and the asymptotic mass of the Higgs excitations, where the last has been approximated by the mass on the sheet with $n = 10^6$. Over the range $10 \le n \le 10^4$, the binding energy is approximately proportional to $n^{-2/3}$.

TABLE I. The masses and estimated partial widths to W^+W^- and ZZ of the first six Higgs excitations. For excitations beyond H^{5*} , the width is comparable to or larger than the spacing between excitations, so they are not distinct. The third column is the partial width to weak vector bosons assuming a pointlike Higgs boson and standard model HZZ and HWW coupling strengths.

Excitation	Mass (GeV)	Width (GeV)
H^*	186.34	0.88
H^{**}	218.82	2.22
H^{3*}	239.00	3.30
H^{4*}	252.89	4.20
H^{5*}	263.10	4.95
H^{6*}	270.97	5.59

anov and Popov [5] and Popov [13] in which they computed the collective excitations of the condensate in BCS superconductivity. They predicted a series of Higgs-like amplitude excitations in superconductors which are unfortunately hard to detect because there is no external probe that couples linearly to the Higgs modes of a superconductor [14]. In contrast, the Higgs excitations described here appear to have the same coupling as the standard model Higgs, and are at energies in reach of existing accelerators, allowing their existence to be checked directly.

I would like to thank R.W. Wilson for listening to these ideas at an early stage and the late R.W. Lucky for additional discussions. At Nokia Bell Labs, Wolfgang Templ suggested the problem that led to this work, and Timo Soirinsuo provided continued support. Their help is gratefully acknowledged.

* email: Gregory.Wright@nokia-bell-labs.com

- P. W. Higgs, Broken symmetries and the masses of gauge bosons, Phys. Rev. Lett. 13, 508 (1964).
- [2] Y. Nambu and G. Jona-Lasinio, Dynamical model of elementary particles based on an analogy with superconductivity. I, Phys. Rev. **122**, 345 (1961).
- [3] V. G. Vaks and A. I. Larkin, On the application of the methods of superconductivity theory to the problem of the masses of elementary particles, Sov. Phys. JETP 13, 192 (1961).
- [4] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Theory of superconductivity, Phys. Rev. 108, 1175 (1957).
- [5] V. A. Andrianov and V. N. Popov, Hydrodynamic action and Bose spectrum of superfluid Fermi systems, Theoret. and Math. Phys. 28, 829 (1976).
- [6] R. J. Eden, P. V. Landshoff, D. I. Olive, and J. C. Polkinghorne, *The Analytic S-Matrix* (Cambridge Univ. Press, Cambridge, 1966).
- [7] J. Aparisi, J. Fuster, A. Irles, G. Rodrigo, M. Vos, H. Yamamoto, A. Hoang, C. Lepenik, M. Spira, S. Tairafune, and R. Yonamine, m_b at m_H : The running bottom quark mass and the higgs boson, Phys. Rev. Lett. **128**, 122001 (2022), arXiv:2110.10202 [hep-ph].
- [8] S. Navas *et al.* (Particle Data Group), Phys. Rev. D **110**, 030001 (2024), to be published.
- [9] L. G. Almeida, S. J. Lee, S. Pokorski, and J. D. Wells, Study of the standard model Higgs boson partial widths and branching fractions, Phys. Rev. D 89, 033006 (2014), arXiv:1311.6721 [hep-ph].
- [10] G. Aad *et al.* (ATLAS Collaboration), Evidence of offshell Higgs boson production from ZZ leptonic decay channels and constraints on its total width with the ATLAS detector, Phys. Lett. B 846, 138223 (2023), arXiv:2304.01532 [hep-ex].
- [11] A. Tumasyan *et al.* (CMS Collaboration), Measurement of the Higgs boson width and evidence of its off-shell contributions to ZZ production, Nature Physics 18, 1329 (2022), arXiv:2202.06923 [hep-ex].
- [12] P. W. Anderson, Random-phase approximation in the theory of superconductivity, Phys. Rev. 112, 1900 (1958).
- [13] V. N. Popov, Functional Integrals and Collective Excitations (Cambridge University Press, 1987).
- [14] R. Shimano and N. Tsuji, Higgs mode in superconductors, Annual Review of Condensed Matter Physics 11, 103 (2020), arXiv:1906.09401 [cond-mat].