

Role of Complexity on the Minimal Deformation of Black Holes

Z. Yousaf,^{1,*} Kazuharu Bamba,^{2,†} Bander Almutairi,^{3,‡} S. Khan,^{1,§} and M. Z. Bhatti^{1,¶}

¹Department of Mathematics, University of the Punjab, Quaid-i-Azam Campus, Lahore-54590, Pakistan.

²Faculty of Symbiotic Systems Science, Fukushima University, Fukushima 960-1296, Japan.

³Department of Mathematics, College of Science,

King Saud University, P.O.Box 2455 Riyadh 11451, Saudi Arabia.

We investigate spherically symmetric classes of anisotropic solutions within the realm of a schematic gravitational decoupling scheme, primarily decoupling through minimal geometric deformation, applied to non-rotating, ultra-compact, self-gravitational fluid distributions. In this respect, we employ the minimal complexity factor scheme to generate physically realistic models for anisotropic matter distributions, using a well-behaved model. The zero-complexity factor condition enables us to determine the deformation function for solving the decoupled system. We explore all the structure-defining scalar variables, such as density inhomogeneity, strong energy condition, density homogeneity, and the complexity factor (an alloy of density inhomogeneity and pressure anisotropy) for the decoupling constant ranging between 0 and 1. We observe that the anisotropy vanishes when the coupling constant is set to unity. This finding holds significance as it implies that, in the context of a zero-complexity factor approach, an anisotropic matter distribution becomes perfect without requiring any isotropy requirements. This work effectively explored the impact of complexity on the composition of self-gravitational stellar distributions. This effective approach enables the development of new, physically realistic isotropic stellar models for anisotropic matter distributions. Additionally, our findings indicate that the complexity factor in static, spherically symmetric self-gravitational objects can significantly affect the nature of the matter distribution within these systems. It is concluded that the minimally deformed Durgapal-IV model features an increasing pressure profile, and the local anisotropy of pressure vanishes throughout the model under complexity-free conditions.

PACS numbers: 04.20.Dw; 04.40.Dg; 0 4.50.Kd; 52.40.Db.

I. INTRODUCTION

The Einstein gravitational-field equations (EFEs), which describe the effects of spacetime curvature, present significant challenges due to their complex and nonlinear nature when attempting to construct closed-form analytical solutions. In this regard, Schwarzschild made a groundbreaking attempt by constructing an interior solution to the EFEs, endowed with spherical symmetry and isotropic matter content [1]. He also solved the EFEs for the exterior vacuum case with a perfect fluid configuration. The recognition of Schwarzschild's interior solution as a three-sphere geometry is attributed to Weyl [2]. In subsequent decades, significant development occurred in the advancement of closed-form analytical solutions to EFEs. Understanding the structure and evolution of compact stars requires an equation of state (EoS) for dense nuclear matter. This EoS should describe both the crust and core of a neutron star using a single, consistent physical model. Tolman [3] made significant contributions by constructing the internal solutions of the EFEs within the background of perfect fluid spheres governed by polytropic EoS. Subsequently, the properties of polytropes in cosmological spacetimes were explored in detail by various authors (see [4, 5] for details). The fascinating characteristics of these spherically symmetric polytropes have been thoroughly investigated in subsequent studies [6–8]. Besides proposing an EoS to correlate the matter variables, we can employ the geometrical constraints on the functions. By utilizing this scheme, Tolman formulated a family of eight isotropic solutions. Tolman's closed-form solutions characterize the configuration of static, stellar structures with a perfect fluid EoS, where radial and tangential stresses are equal, denoted as $P_r = P_\theta$.

* Email: zeeshan.math@pu.edu.pk

† Email: bamba@sss.fukushima-u.ac.jp; Corresponding author

‡ Email: baalmutairi@ksu.edu.sa

§ Email: suraj.pu.edu.pk@gmail.com

¶ Email: mzaeem.math@pu.edu.pk

However, follow-up studies revealed that spherical symmetry does not necessarily ensure isotropic pressure. Recognizing this limitation, the introduction of anisotropic matter ($P_r \neq P_\theta$) became crucial for understanding highly dense objects. Lemaître [9] explored the interiors of stars by solving the equations of general relativity (GR) for matter distributions with anisotropic properties. This approach has opened new avenues of research, attracting scientists interested in understanding the structure and behavior of stellar matter. Ruderman’s work [10] highlights the possibility of anisotropic nuclear matter at ultra-high densities ($> 10^{15} \text{g/cm}^3$). This implies a non-isotropic EoS in these regimes. Additionally, Bowers and Liang [11] elaborated on the importance of anisotropic EoS for highly dense self-gravitational configurations of fluid spheres. Finding physically viable solutions to the EFEs in the presence of anisotropy remains a significant challenge [12]. Despite its considerable challenge, a wide range of works in this direction have been available in the literature [13–23]. These investigations have successfully addressed the impact of anisotropy for understanding the evolution and formation of highly dense stellar distributions. Harko pointed out that various mechanisms can induce anisotropy in ultra-compact stellar distributions, including the presence of a type 3A superfluid [24], the existence of a solid core [10], pion condensation [25], or different types of phase transitions [26].

The existence of anisotropy gives rise to several characteristics in the matter distribution of self-gravitational objects. For example, a positive anisotropic factor $\Delta \equiv P_r - P_\theta > 0$ introduces a repulsive force (attractive when the anionic factor Δ is negative) that counters the gravitational collapse. This enables us to construct highly dense structures with anisotropic fluids compared to isotropic fluids [15, 16]. Additionally, positive anisotropy enhances the stability and equilibrium of the self-gravitational distribution.

For decades, self-gravitational compact entities like stars and galaxies have been crucial tools in unraveling the mysteries of our vast universe. Large-scale surveys like the Sloan Digital Sky Survey and the Large Synoptic Survey Telescope provide invaluable data on these cosmic bodies. However, these objects are incredibly complex heavenly configurations. Even slight fluctuations within them can significantly alter their physical properties, potentially offering new insights into the universe’s origin and evolution. Within this domain, a notable challenge resides in precisely gauging the complex interplay among various physical phenomena transpiring within stars and analogous self-gravitational compact structures. This is where the notion of a “complexity factor” stems and assumes paramount importance. This factor seeks to enfold the collective effects of numerous internal mechanisms and their collective influence on the system’s global dynamics. Within stable self-gravitational distributions, complexity emerges from the interconnection of these diverse elements, often giving rise to emergent phenomena that cannot be easily foreseen based solely on the behavior of individual components. Presently, ongoing research across diverse scientific fields is centered on establishing precise definitions and characterizations of complexity. Within the domain of compact configurations, especially those of significant mass, the analysis of complexity frequently entails evaluating metrics related to information and entropy present within the configuration of the system. The investigation of ultra-compact stellar configurations with anisotropic matter content often involves the examination of the complexity of self-gravitational structures. In physics, isolated gases are regarded as intricate systems due to their disorder and high information content, while perfect crystals, with their periodic behavior and symmetric dispersion, are seen as possessing minimal complexity. “disequilibrium” as a measure of complexity in systems. This measure quantifies the deviation from the system’s most probable, equally likely state, essentially capturing the system’s “distance” from equilibrium. They argued that for ideal gases and perfect crystals, the concept of complexity vanishes when considering both disequilibrium and information content. Considering the limitations in existing complexity measures for self-gravitating systems, Herrera [27] proposed a novel approach based on fluid properties. This includes fundamental quantities like energy density and pressure, capturing the system’s overall content. The complexity is then constructed using a “complexity factor”, one of the structural scalars derived from the orthogonal decomposition of the intrinsic curvature.

Building upon their prior work, Herrera *et al.* [28] extended the complexity concept to dissipative fluids. They went beyond mere complexity analysis and established conditions for evolutionary paths with minimal complexity. Their investigation revealed the existence of multiple solutions, with the fluid exhibiting shearing and geodesic properties in the dissipative regime. Herrera *et al.* further explored the interplay between complexity and geometry [29]. Utilizing axially symmetric geometries, they identified three distinct complexity categories and demonstrated the relationship between complexity and symmetry. Notably, they obtained analytical solutions in this specific case. On the other hand [30], they investigated the evolution of non-static geometries endowed with spherical symmetry using the notion of complexity, considering both dissipative and non-dissipative scenarios. In this context, they

constructed anisotropic stellar-type models and explored the implications for understanding their evolution using the quasi-homologous condition. Many researchers have employed this approach to explore the evolution of stellar distributions through alternative gravitational models [31–34]. Herrera et al. [35] extended the complexity factor concept to geometries with hyperbolic symmetry. They incorporated both the Misner-Sharp mass and Tolman mass, along with various structural scalars obtained through the orthogonal decomposition of the intrinsic curvature. Interestingly, their analysis revealed that the Tolman mass exhibits a negative nature in this specific case. For more related references, see [36] (recent review) and references therein.

The minimal geometric deformation (MGD) method has gained attraction as a powerful and versatile tool for extending isotropic solutions to EFEs. Its primary advantage lies in its direct and analytical methodology, which necessitates minimal geometric alterations. This feature renders it a valuable substitute for current techniques. Within the context of GR, the gravitational decoupling (GD) facilitated by the minimal geometric deformation (MGD) approach offers a seamless means of introducing the necessary alterations to equations or solutions. The MGD technique presents a strong capability: providing a pre-existing solution to EFEs, it enables the discovery of innovative solutions that integrate an additional gravitational-field source. Primarily, proposed for the Randall-Sundrum braneworld [37–44], the MGD scheme has found significant applications beyond its initial domain. Significantly, it has played a crucial role in investigating novel categories of black hole solutions [37, 38]. This feat is based on the application of MGD decoupling to the conventional Schwarzschild metric, showcasing the method’s adaptability in tackling various gravitational challenges. The original applications of this scheme also focused on black hole acoustics [45] and GUP Hawking fermions [46]. After its development, the MGD scheme was employed within the framework of GR [47, 48]. This implementation extended the method’s capability beyond perfect fluids, with a key achievement being the generalization of solutions from isotropic fluids to encompass anisotropic domains. As we will discuss in the following section, this methodology comprises two primary components: (i) two independent gravitational sectors, $T_{\mu\nu}$ and $\mathbb{X}_{\mu\nu}$, which interact solely gravitationally, and (ii) an MGD scheme introduced in the radial metric potential g_{rr} . This deformation enables the decoupling of the system into two sets of equations, one for each source. In the MGD scheme, the source $T_{\mu\nu}$ often serves as the source for a well-established isotropic interior solution and $\mathbb{X}_{\mu\nu}$ introduces a local anisotropic behavior into the system. Consequently, GD is known for extending isotropic solutions to encompass anisotropic scenarios. Utilizing the MGD scheme for anisotropic fluids within GR poses a distinctive challenge. The geometric deformation and the components of the \mathbb{X} -gravitational field source $\mathbb{X}_{\mu\nu}$ remain undetermined. To tackle this issue, researchers have investigated diverse approaches to impose additional constraints that facilitate solving for these unknowns. For example (i) mimic constraints scheme [45, 49] (ii) some anisotropy mechanism [50] (iii) zero complexity factor condition [51, 52].

This work explores anisotropic Class I stellar-type configurations by constructing solutions to EFEs that adhere to the complexity-free condition. In this context, we employ one of the effective theoretical techniques for constructing EFEs solutions: the GD scheme, which admits the zero-complexity factor condition. The well-behaved and viable metric ansatz, combined with the anisotropic fluid sphere is utilized to solve the system of equations. Karmarkar (see [53] and references therein) introduced the embedding Class I condition, which serves as a versatile and straightforward auxiliary condition for solving the EFEs. This condition is typically applicable to any n -dimensional (pseudo)-Riemannian metric. The Durgapal-IV metric ansatz [54] is a relativistic model proposed by M.C. Durgapal. He observed that by considering a simple relation $e^{\phi_1} \propto (1 + r^4)^n$ (where n denotes a positive real constant), closed-form analytical solutions of the EFEs could be derived. Based on our findings, the pressure anisotropy increases as the radial variable rises, resulting in a more effective stabilization of the shell compared to the central core areas. Additionally, we observe that contributions from MGD decoupling, characterized by a complexity-free condition, appear to diminish the anisotropy of the stellar structures.

The structure of this paper is as follows: In Sec. II, we briefly review the formalism of GD primarily in the context of anisotropically structured spherical matter distributions. Section III covers the introduction of MGD decoupling along with the formulation of junction conditions. In Sec. IV, we discuss the zero-complexity factor constraint and the embedding Class I scheme for constructing the MGD-based, anisotropic Durgapal-IV self-gravitational compact stars. Section V presents a detailed physical analysis of the results, and the concluding remarks are provided in Sec. VI.

II. THE STANDARD PROTOCOL FOR DECOUPLING GRAVITATIONAL SOURCES

To explore the physical properties inherent to decoupled, self-gravitating compact stars, the modified version of standard gravitational action reads [47, 48]

$$S = \int d^4x \sqrt{|g|} \left(\frac{R}{16\pi} + L_m \right) + \beta(\text{corrections}), \quad (1)$$

where β corresponds to the new gravitational source. The above action can also be defined as

$$S^\circ = \int d^4x \sqrt{|g|} \left(\frac{R}{16\pi} + L_m + \beta L_X \right), \quad (2)$$

where $R = g^{\mu\nu} R_{\mu\nu}$ denotes the Ricci scalar, while $R_{\mu\nu}$ and $g^{\mu\nu}$ represent the Ricci tensor and metric tensor, respectively.

Furthermore, the Lagrangian for ordinary matter is L_m , whereas the contribution from the additional gravitational field source, originating from scalar, vector, or tensor fields (beyond the purview of GR), is denoted by L_Φ . Additionally, g is the trace of $g^{\mu\nu}$ and β is a free parameter that determines the strength of decoupling. Thus, the EFEs featuring the decoupled stellar fluid configurations can be obtained by varying the action S° with respect to $g_{\mu\nu}$, yielding

$$G_{\mu\nu} \equiv R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = 8\pi T_{\mu\nu}^*, \quad (3)$$

where the stress-energy tensor (SET) $T_{\mu\nu}^*$ is defined as

$$T_{\mu\nu}^* = T_{\mu\nu} + \alpha \mathbb{X}_{\mu\nu}. \quad (4)$$

Here, $\mathbb{X}_{\mu\nu}$ represents generic type of fluid distribution (\mathbb{X} -gravitational sector, hereafter), induced by GD. Furthermore, $T_{\mu\nu}^*$ denotes the interior matter content of the decoupled, self-gravitational spherical object, which is approximated with an anisotropic distribution of energy density (σ), tangential pressure (P_θ) and radial pressure P_r . Thus, the SET $T_{\mu\nu}^*$ in canonical form is defined as

$$[T^*]_\nu^\mu = (\sigma + P_\theta) U^\mu U_\nu - P_\theta \delta_\nu^\mu + (P_r - P_\theta) \mathcal{X}^\mu \mathcal{X}_\nu, \quad (5)$$

where U^μ is the four-velocity and \mathcal{X}^μ denotes the unit four vector along the radial radiation direction and satisfy

$$U^\mu = (e^{-\phi_1/2}, 0, 0, 0), \quad \mathcal{X}^\mu = (0, e^{-\phi_2/2}, 0, 0), \quad (6)$$

respectively, such that $U^\mu U_\mu = -\mathcal{X}^\mu \mathcal{X}_\mu = 1$. In standard coordinates, $x^\mu = x^{0,1,2,3} = \{t, r, \vartheta, \varphi\}$, the most general form of the metric with static, spherical symmetry can be expressed as

$$ds^2 = e^{\phi_1} dt^2 - e^{\phi_2} dr^2 - r^2 d\Omega^2, \quad (7)$$

where $d\Omega^2 = dr^2 - \sin^2 \vartheta d\varphi^2$ denotes the metric of the unit 2-sphere, while $\phi_1 = \phi_1(r)$ and $\phi_2 = \phi_2(r)$ are functions of the areal radius r only. Then, under the spherical symmetry endowed metric ansatz (7), the non-null components of $T_{\mu\nu}^*$ read

$$[T^*]_0^0 = \sigma^*, \quad [T^*]_1^1 = -P_r^*, \quad [T^*]_2^2 = [T^*]_3^3 = -P_\theta^*. \quad (8)$$

As a consequence, the tt , rr and $\vartheta\vartheta$ components of the decoupled EFEs turn out to be

$$G_0^0 = 8\pi [T^*]_0^0 : \frac{1}{r^2} - \left(\frac{1}{r^2} - \frac{\phi_2'}{r} \right) e^{\phi_2} = 8\pi (T_0^0 + \beta \mathbb{X}_0^0), \quad (9)$$

$$G_1^1 = 8\pi [T^*]_1^1 : -\frac{1}{r^2} + \left(\frac{1}{r^2} - \frac{\phi_2'}{r} \right) e^{\phi_2} = 8\pi (T_1^1 + \beta \mathbb{X}_1^1), \quad (10)$$

$$G_2^2 = 8\pi[T^*]_2^2 : \frac{e^{-\phi_2}}{4} \left(2\frac{\phi_2'}{r} - 2\frac{\phi_1'}{r} + \phi_2'\phi_1' - \phi'^2 - 2\phi_1'' \right) = 8\pi(T_2^2 + \beta\mathbb{X}_2^2), \quad (11)$$

where $f' \equiv \partial_r f$. Because G_ν^μ is divergence-free, the SET $[T^*]_{\mu\nu}$ follows the subsequent conservation expression

$$0 \equiv [T^*]^{\mu\nu}{}_{;\nu} = \frac{d[T^*]^{\mu\nu}}{dx^\nu} + \Gamma_{\nu\epsilon}^\mu [T^*]^{\epsilon\nu} + \Gamma_{\nu\epsilon}^\nu [T^*]^{\mu\epsilon}, \quad (12)$$

which, for $\mu = 1$, provides a decoupled hydrostatic equilibrium equation, defined as

$$\frac{d}{dr} ([T^*]_1^1) = -\frac{1}{2}g^{00}\frac{dg_{00}}{dr} ([T^*]_0^0 - [T^*]_1^1) - g^{22}\frac{dg_{22}}{dr} ([T^*]_1^1 - [T^*]_2^2), \quad (13)$$

where $[T^*]_2^2 = [T^*]_3^3$ due to spherical symmetry. Therefore, under the GD approach Eq. (13) reads

$$\frac{dP_r}{dr} = -\frac{\phi_1'}{2}(\sigma - P_r) - \frac{2}{r}(P_\theta - P_r) + \beta \left[(\mathbb{X}_1^1)' - \frac{\phi_1'}{2}(\mathbb{X}_0^0 - \mathbb{X}_1^1) - \frac{2}{r}(\mathbb{X}_2^2) \right], \quad (14)$$

which is a linear combination of the Eqs. (9)-(11). Additionally, the values of σ , P_r , and P_θ correspond to the effective energy density, effective radial pressure, and effective tangential pressure, respectively, which are determined by

$$\sigma^* = T_0^0 + \beta\mathbb{X}_0^0 \equiv \sigma + \sigma^{\mathbb{X}}, \quad (15)$$

$$P_r^* = -T_1^1 - \beta\mathbb{X}_1^1 \equiv p_r + P_r^{\mathbb{X}}, \quad (16)$$

$$P_\theta^* = -T_2^2 - \beta\mathbb{X}_2^2 \equiv p_\perp + P_\theta^{\mathbb{X}}, \quad (17)$$

which are associated with the SET $[T^*]_{\mu\nu}$. Then, by making use of the above-stated physical variables, the pressure anisotropy for the source $[T^*]_{\mu\nu}$ can be defined as

$$\Delta^* \equiv P_r^* - P_\theta^* = \Delta + \Delta^{\mathbb{X}}. \quad (18)$$

Here,

$$\Delta \equiv P_r - P_\theta, \quad \Delta^{\mathbb{X}} \equiv P_r^{\mathbb{X}} - P_\theta^{\mathbb{X}} = -\alpha(\mathbb{X}_1^1 - \mathbb{X}_2^2), \quad (19)$$

where Π and $\Pi^{\mathbb{X}}$ denote the pressure anisotropies corresponding to seed-sector and \mathbb{X} -gravitational sector respectively. It is interesting to observe that even if the seed sector is isotropic, the \mathbb{X} -gravitational sector will tend to produce anisotropy in the compact system because, in general, $\mathbb{X}_1^1 \neq \mathbb{X}_2^2$. Now, we will construct the gravitational field equations corresponding to the seed sector as well as the θ -gravitational sector by employing the MGD-decoupling approach.

III. FIELD EQUATIONS FOR $[T^*]_{\mu\nu}$ AND $\mathbb{X}_{\mu\nu}$ SECTORS VIA MGD-DECOUPLING

In this section, we will examine how the gravitational system can be decomposed into two independent gravitational sectors such that the equations of motion associated with the additional fluid $\mathbb{X}_{\mu\nu}$ correspond to the effective “quasi-Einstein system”. We will now apply the GD approach by assuming a solution for the system Eqs. (9)–(11), with $\beta = 0$, whose dynamics can be described as the metric ansatz

$$ds^2 = e^{\lambda(r)}dt^2 - e^{\xi(r)}dr^2 - r^2d\Omega^2, \quad (20)$$

with

$$e^{\xi(r)} \equiv 1 - \frac{8\pi}{r} \int_0^r x^2 T_0^0(x) dx = 1 - \frac{2m_s}{r}, \quad (21)$$

where m_s is the mass function associated with the seed gravitational sector. Ovalle [47] proposed that the influence of the \mathbb{X} -gravitational sector on the seed SET, $T_{\mu\nu}$ can be determined by introducing the following geometric transformations.

$$e^\lambda \mapsto e^{\phi_1} = e^{\lambda + \beta h}, \quad (22)$$

$$e^{-\xi} \mapsto e^{-\phi_2} = e^{-\xi} + \beta f, \quad (23)$$

where $f(r)$ and $h(r)$ are the corresponding deformation functions governed by the free parameter β . The MGD-decoupling requires that

$$h(r) \mapsto 0; \quad f(r) \mapsto F^\diamond(r). \quad (24)$$

This shows that the temporal metric potential remains the same, while the presence of the \mathbb{X} -gravitational source induces a deformation of the radial metric potential (23). Therefore, all the effects stemming from the \mathbb{X} -gravitational sector is encoded in the radial metric potential $g_{rr}(r) = e^{\phi_2(r)}$. Thus, under the MGD-decoupling the gravitational system (9)-(11) is divided as follows:

1. the standard EFEs associated with seed gravitational sector

$$G^\mu_\nu(\lambda, \xi) = 8\pi T^\mu_\nu, \quad \nabla_\mu T^\mu_\nu = 0, \quad (25)$$

where

$$G^0_0(\lambda, \xi) = \frac{1}{r^2} - e^{-\xi} \left(\frac{1}{r^2} - \frac{\xi'}{r} \right), \quad (26)$$

$$G^1_1(\lambda, \xi) = \frac{1}{r^2} - e^{-\xi} \left(\frac{1}{r^2} + \frac{\lambda'}{r} \right), \quad (27)$$

$$G^2_2(\lambda, \xi) = G^3_3(\sigma, \varphi) = \frac{e^{-\xi}}{4} \left(2\frac{\xi'}{r} - 2\frac{\lambda'}{r} + \lambda'\xi' - \lambda'^2 - 2\lambda'' \right), \quad (28)$$

2. the gravitational equations corresponding to the \mathbb{X} -gravitational sector

$$\beta \mathcal{G}^\mu_\nu(\lambda, \xi, F^\diamond) = 8\pi \mathbb{X}^\mu_\nu, \quad \nabla_\mu \mathbb{X}^\mu_\nu = 0, \quad (29)$$

where

$$\mathcal{G}^0_0 = - \left(\frac{F^\diamond}{r^2} + \frac{f^{*'}}{r} \right), \quad (30)$$

$$\mathcal{G}^1_1 = -F^\diamond \left(\frac{1}{r^2} + \frac{\lambda'}{r} \right), \quad (31)$$

$$\mathcal{G}^2_2 = \mathcal{G}^3_3 = -\frac{F^\diamond}{4} \left(2\lambda'' + \lambda'^2 + 2\frac{\lambda'}{r} \right) - \frac{F^\diamond}{4} \left(\lambda' + \frac{2}{r} \right). \quad (32)$$

The hydrostatic equilibrium equations corresponding to the seed SET and \mathbb{X} -gravitational sector read

$$\frac{dP_r}{dr} = -\frac{m_s + 4\pi r^3 P_r}{(r - 2m_s)} (\sigma + P_r) + \frac{2}{r} (P_\theta - P_r), \quad (33)$$

and

$$\frac{dP_r^\mathbb{X}}{dr} = -\frac{m_s + 4\pi r^3 P_r^*}{(r - 2m_s)} (\sigma^\mathbb{X} + P_r^\mathbb{X}) + \frac{2}{r} (P_\theta^\mathbb{X} - P_r^\mathbb{X}), \quad (34)$$

respectively. Next, the mass functions associated with both gravitational sectors are related through the expression

$$m = m_s + \beta m_{\mathbb{X}}, \quad \text{where} \quad m_{\mathbb{X}} = -\frac{r}{2} F^\diamond. \quad (35)$$

Here, the mass function induced by the \mathbb{X} -gravitational sector and the seed SET are denoted by $m_{\mathbb{X}}$ and m_s , respectively. Herrera [27] proposed that the mass function corresponding to the spherical self-gravitational distribution with anisotropic matter content can be expressed as a sum of two quantities: the homogeneous density and the corresponding alteration due to the non-homogeneous behavior of diversity, which reads

$$m = \frac{4\pi}{3} r^3 \sigma^* - \frac{4\pi}{3} \int_0^r [\sigma^*(x)]' x^3 dx. \quad (36)$$

Then, the corresponding expressions of the mass function for the seed sector and the \mathbb{X} -gravitational source take the form

$$m = \frac{4\pi}{3} r^3 \sigma - \frac{4\pi}{3} \int_0^r [\sigma(s)]' s^3 ds, \quad (37)$$

$$m_{\mathbb{X}} = \frac{4\pi}{3} r^3 \sigma^{\mathbb{X}} - \frac{4\pi}{3} \int_0^r [\sigma^{\mathbb{X}}(x)]' x^3 dx. \quad (38)$$

Furthermore, Tolman's groundbreaking research provided a foundation for understanding the total energy content associated with relativistic self-gravitational objects, which is formally defined as (for details see [55])

$$m_T = 4\pi \int_0^{r_\Sigma} x^2 e^{(\phi_1 + \phi_2)/2} \left([T^*]_0^0 - [T^*]_1^1 - 2[T^*]_2^2 \right) dx, \quad (39)$$

or, alternatively,

$$m_T = e^{(\phi_1 + \phi_2)/2} \left(m + 4\pi r^3 P_r^* \right). \quad (40)$$

The above expression can be cast into the form

$$m_T = [m_T(r)]_\Sigma \left(\frac{r}{r_\Sigma} \right)^3 - r^3 \int_r^{r_\Sigma} \left[\frac{4\pi}{x^4} \int_0^r [\sigma^*(x)]' x^3 ds - \frac{8\pi}{x} \Delta^* \right] e^{(\phi_1 + \phi_2)/2} dx, \quad (41)$$

which can be expressed in terms of Weyl scalar E as

$$m_T = [m_T(r)]_\Sigma \left(\frac{r}{r_\Sigma} \right)^3 + r^3 \int_r^{r_\Sigma} \frac{1}{x} e^{(\phi_1 + \phi_2)/2} [4\pi \Delta^* + E] dx. \quad (42)$$

Across the boundary of the compact star, the continuity of the first and second fundamental forms imposes a matching condition for the compact object. This is achieved by ensuring that the interior and exterior metrics, which describe the geometry inside and outside the sphere respectively, match at the boundary surface $r = r_\Sigma$. In this respect, the Schwarzschild exterior metric reads

$$ds^2 = \left(1 - \frac{2M}{r} \right) dt^2 - \frac{dr^2}{\left(1 - \frac{2M}{r} \right)} - r^2 d\Omega^2. \quad (43)$$

The necessary and sufficient conditions at the boundary can be obtained by matching two metrics and applying the first and second fundamental forms as

$$\left(1 - \frac{2M}{r} \right) \Big|_\Sigma = e^\nu \Big|_\Sigma, \quad (44)$$

$$\left(1 - \frac{2M}{r} \right) \Big|_\Sigma = e^{-\lambda} \Big|_\Sigma, \quad (45)$$

$$P_r^* \Big|_\Sigma = 0. \quad (46)$$

IV. COMPLEXITY FACTOR AND OTHER SCALARS UNDER MGD-DECOUPLING

To explore the inherent features of relativistic compact self-gravitational structures endowed with anisotropic matter configurations, Herrera [27] coined the term “complexity factor”. This factor is primarily a scalar function derived from the orthogonal decomposition of the Riemann-Christoffel tensor and is referred to as such because it plays a role in computing the complexity of anisotropic compact stars. Herrera and his coworkers [56] analyzed the growth and structure of spherical anisotropic fluid distributions by constructing a full set of equations in terms of five scalar functions that reduce to $\{X_T, X_{TF}, Y_T, Y_{TF}\}$ in the case of non-dissipative conditions. Additionally, they have demonstrated that all possible solutions to the EFEs corresponding to the time-independent spherical fluid spheres can be expressed explicitly in terms of these scalar quantities. These properties include pressure anisotropy, inhomogeneous density, Tolman mass, heat flux as well as the homogeneous distribution of density. These scalar functions appear as trace and trace-free parts of the specific tensor terms (see [56, 57] for details)

$$Y_{\mu\nu} = \left(S_\mu S_\nu - \frac{1}{3} h_{\mu\nu} \right) Y_{TF}^* + \frac{1}{3} h_{\mu\nu} Y_T^*, \quad (47)$$

$$X_{\mu\nu} = \left(S_\mu S_\nu - \frac{1}{3} h_{\mu\nu} \right) X_{TF}^* + \frac{1}{3} h_{\mu\nu} X_T^*. \quad (48)$$

The term $h_{\mu\nu}$ represents the projection tensor. This tensor allows us to project vectors and tensors onto specific surfaces within spacetime, facilitating the analysis and manipulation of geometric structures. The projection tensor enables us to decompose various quantities, such as velocity, momentum, or energy density, into components that are parallel and perpendicular to the hypersurfaces under consideration. The vector S_μ represents the worldline (path through spacetime) of a massless particle traveling along a null geodesic. The expressions of the structural scalars in terms of physical variables read

$$Y_{TF}^* = 4\pi\Delta^* + E = 8\pi\Delta^* - \frac{4\pi}{r^3} \int_0^r [\sigma^*(x)]' x^3 dx, \quad (49)$$

$$Y_T^* = 4\pi(\sigma^* + 3P_r^* - 2\Delta^*), \quad (50)$$

$$X_{TF}^* = 4\pi\Delta^* - E = \frac{4\pi}{r^3} \int_0^r [\sigma^*(x)]' x^3 dx, \quad (51)$$

$$X_T^* = 8\pi\sigma^*(r), \quad (52)$$

where the values of Δ^* and Δ are defined in Eqs. (18) and (19). Then, according to the MGD decoupling, the aforementioned scalars split as

$$Y_{TF}^* \equiv Y_{TF} + Y_{TF}^X; \quad Y_T^* \equiv Y_T + Y_T^X; \quad X_{TF}^* \equiv X_{TF} + X_{TF}^X; \quad X_T^* \equiv X_T + X_T^X, \quad (53)$$

where

$$Y_{TF} = 4\pi\Delta + E = 8\pi\Delta - \frac{4\pi}{r^3} \int_0^r [\sigma(x)]' x^3 dx, \quad (54)$$

$$Y_{TF}^X = 4\pi\Pi^X + E = 8\pi\Delta^X - \frac{4\pi}{r^3} \int_0^r [\sigma^X(x)]' x^3 dx, \quad (55)$$

$$Y_T = 4\pi(\sigma + 3P_r - 2\Delta); \quad Y_T^X = 4\pi(\sigma^X + 3P_r^X - 2\Delta^X), \quad (56)$$

$$X_{TF} = 4\pi\Delta - E = \frac{4\pi}{r^3} \int_0^r [\sigma(x)]' x^3 dx; \quad X_{TF}^X = 4\pi\Pi^X - E = \frac{4\pi}{r^3} \int_0^r [\sigma^X(x)]' x^3 dx, \quad (57)$$

$$X_T = 8\pi\sigma, \quad X_T^X = 8\pi\sigma^X. \quad (58)$$

In this context, the scalars X_T^* and X_{TF}^* denote the density homogeneity and density inhomogeneity of the anisotropic compact star, while the remaining scalars Y_T^* and Y_{TF}^* denote the strong energy condition and the complexity factor, respectively. Next, the local anisotropy in decoupled anisotropic stellar distribution is governed by the interaction between X_{TF}^* and Y_{TF}^* , which can be expressed as

$$Y_{TF}^* + X_{TF}^* \equiv [Y_{TF} + Y_{TF}^X] + [X_{TF} + X_{TF}^X] = 8\pi (\Delta + \Delta^X). \quad (59)$$

The definition of Tolman mass in terms of the complexity factor reads

$$m_T = \left(\frac{r}{R}\right)^3 M_T + r^3 \int_r^R \frac{e^{(\phi_1+\phi_1)/2}}{x} Y_{TF}^* dx. \quad (60)$$

or, alternatively,

$$m_T = \left(\frac{r}{R}\right)^3 M_T + r^3 \int_r^R \frac{e^{(\phi_1+\phi_1)/2}}{x} Y_{TF} dx + \beta r^3 \int_r^R \frac{e^{(\phi_1+\phi_1)/2}}{x} Y_{TF}^{\mathbb{X}} dx, \quad (61)$$

which describes the influence of the \mathbb{X} -gravitational sector regulated by decoupling parameter β . Furthermore, M_T represents the total value of the Tolman mass for the self-gravitational compact star with radius R . The value of m_T , using the strong energy condition Y_T^* , can be written as

$$m_T = \int_0^r x^2 e^{(\phi_1+\phi_2)/2} Y_T dx + \beta \int_0^r x^2 e^{(\phi_1+\phi_2)/2} Y_T^{\mathbb{X}} dx. \quad (62)$$

Now, for solving both the gravitational sectors, we will adopt different strategies:

1. Seed sector $T_{\mu\nu}$: We will use the Embedding Class I condition (Karmarkar condition) to solve this system. This approach utilizes known solutions to constrain the solution space.
2. \mathbb{X} -gravitational sector $\mathbb{X}_{\mu\nu}$: This gravitational will be solved by applying the constraint of zero complexity factor ($Y_{TF}^* = 0$).

A. Complexity-free Embedding Class I Solutions

If the spherical symmetry endowed metric associated with the seed SET $T_{\mu\nu}$ satisfies the Karmarkar condition [58], then we have

$$R_{1010}R_{2323} = R_{1212}R_{3030} + R_{1220}R_{1330}. \quad (63)$$

The above condition fulfills the Pandey and Sharma condition [59], i.e., $R_{2323} \neq 0$. Then, according to the seed metric ansatz with geometric variables $\{\lambda, \zeta\}$, the Karmarkar condition [58] provides the following differential equation

$$\frac{\zeta' \lambda'}{1 - e^{\zeta(r)}} = \zeta' \lambda' - 2\lambda'' - \lambda'^2. \quad (64)$$

Notably, if a metric ansatz follows this condition, it can be embedded within a five-dimensional pseudo-Euclidean space. This kind of metric ansatz is called an embedding Class I metric. Next, the integration of Eq. (64) provides a significant relationship between the geometric variables $\{\lambda, \zeta\}$.

$$e^{\lambda(r)} = \left(L_0 + M_0 \int \sqrt{e^{\zeta(r)} - 1}\right)^2 \quad \text{or} \quad e^{\zeta(r)} = \left(1 + \frac{\lambda'^2 e^{\lambda(r)}}{4M_0}\right), \quad (65)$$

where L_0 and M_0 are integration constants. Now, we will construct the complexity-free compact star models corresponding to both the gravitation sectors. Firstly, we solve the compact system $\{G_{\mu\nu}, T_{\mu\nu}\}$ for the seed sector involving the variables $\{\lambda, \zeta\}$. Specifying both the metric potentials unlocks the complete set of physical variables (σ, P_r, P_\perp) for the SET $T_{\mu\nu}$. For physically realistic solutions of EFEs, Lake's algorithms [60] constraint the source variable $\lambda(r)$. In particular, $\lambda(r)$ needs to have a regular minimum at $r = 0$ and increase monotonically. This guarantees a time-independent, spherically symmetric solution to the EFEs, with a regular center $r = 0$. Therefore, we assume the source function $e^{\lambda(r)}$ as the Durgapal-IV metric ansatz [54]

$$e^{\lambda(r)} = D(1 + \mathcal{A}r^2)^4, \quad (66)$$

where \mathcal{A} and D are the integration constants with units [$length^{-2}$]. The Durgapal-IV metric ansatz remains regular at the center, $r = 0$, and consistently increases positively throughout the interior of the self-gravitational compact star,

resulting in a realistic model of a stellar distribution. Now, using the Durgapal-IV metric potential, the embedding Class I condition gives [61] as

$$e^{\xi(r)} = 1 + \mathcal{A}Br^2(1 + \mathcal{A}r^2)^2, \quad (67)$$

where $B = 16AD/M_0$. The metric potential $e^{\xi(r)}$ may be represented as $e^{\xi(r)} = 1 + O(r^2)$, which characterizes a necessary condition for the construction of a regular stellar solution at the origin. There, the solution associated with the gravitational system $\{G_{\mu\nu}, T_{\mu\nu}\}$ may be expressed through the metric

$$ds^2 = D(1 + \mathcal{A}r^2)^4 dt^2 - [1 + \mathcal{A}Br^2(1 + \mathcal{A}r^2)^2] dr^2 - r^2 d\Omega^2. \quad (68)$$

We can now obtain the physical variables $\{\sigma, P_r, P_\theta\}$ by inserting the values of the metric potentials $e^{\lambda(r)}$ and $e^{\xi(r)}$ into Eqs. (26)-(28) as

$$8\pi\sigma = \frac{\mathcal{A}B(1 + \mathcal{A}r^2)[-1 + (B - 5)\mathcal{A}r^2 + 3B\mathcal{A}^2r^4 + 3B\mathcal{A}^3r^6 + B\mathcal{A}^4r^8]}{(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)^2}, \quad (69)$$

$$8\pi P_r = -\frac{\mathcal{A}[-8 + B(1 + \mathcal{A}r^2)^3]}{(1 + \mathcal{A}r^2)(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)}, \quad (70)$$

$$8\pi P_\theta = \frac{\mathcal{A}[8 + 16\mathcal{A}r^2 - B(1 + \mathcal{A}r^2)^2(-1 + 9\mathcal{A}^2r^4)]}{(1 + \mathcal{A}r^2)^2(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)}. \quad (71)$$

Now, we solve the gravitational system $\{\mathcal{G}_{\mu\nu}, \mathbb{X}_{\mu\nu}\}$ for the \mathcal{X} -gravitational sector involving the variables $\{\lambda, \xi, F^\diamond\}$ by employing the complexity-free condition, i.e., $Y_{\text{TF}}^* = 0$ with $Y_{\text{TF}} \neq 0$. Then, from Eq. (49), we have

$$Y_{\text{TF}}^* = 4\pi\Delta^* + E \equiv Y_{\text{TF}} + Y_{\text{TF}}^{\mathbb{X}} = 0, \quad (72)$$

where Y_{TF} and $Y_{\text{TF}}^{\mathbb{X}}$ are the specific complexity factors associated with the seed sector and the \mathbb{X} -gravitational sector, respectively. The above equation implies

$$Y_{\text{TF}}^* = 8\pi\Delta^* - \frac{4\pi}{r^3} \int_0^r [\sigma^*(x)]' x^3 dx = 0. \quad (73)$$

As density (σ), radial pressure (P_r), and tangential pressure (P_θ) are fully determined by Eqs. (69)-(71), the expression for Y_{TF} can be directly obtained as

$$Y_{\text{TF}} = \frac{4\mathcal{A}^2r^2[-2 + B(1 + \mathcal{A}r^2)^3]}{(1 + \mathcal{A}r^2)^2(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)^2}. \quad (74)$$

Building on Eq. (49), we need to satisfy the following condition

$$Y_{\text{TF}}^* = 8\pi\Delta^* + 8\pi\Delta^{\mathbb{X}} - \frac{4\pi}{r^3} \int_0^r [\sigma^{\mathbb{X}}(x)]' x^3 dx = 0, \quad (75)$$

with $\beta = 1$. Next, substituting Eqs. (30)-(32) into the aforementioned equation, we have

$$\lambda' F^{\diamond'} + \frac{1}{r} \left[2r\lambda'' + (r\lambda' - 2)\lambda' \right] F^\diamond - 4Y_{\text{TF}} = 0, \quad (76)$$

whose solution can be obtained by using the Durgapal-IV metric ansatz (66) along with Eq. (75) as

$$F^\diamond(r) = \frac{C_1}{(1 + \mathcal{A}r^2)^2} - \frac{1}{1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6}, \quad (77)$$

where C_1 is an arbitrary integration constant. The specific value of C_1 can be determined by using a physically relevant condition, which reads

$$e^{-\phi_2(0)} = e^{-\xi(0)} + \beta F^\diamond(0) = 1 \Rightarrow F^\diamond(0) = 1. \quad (78)$$

Therefore, the deformed radial metric potential takes the following form

$$e^{-\phi_2(r)} = \frac{1 + [1 + \mathcal{A}Br^2(1 + \mathcal{A}r^2)^2]\beta F^\diamond}{[1 + \mathcal{A}Br^2(1 + \mathcal{A}r^2)^2]}, \quad (79)$$

and the corresponding metric reads

$$ds^2 = D(1 + \mathcal{A}r^2)^4 dt^2 - \frac{[1 + \mathcal{A}Br^2(1 + \mathcal{A}r^2)^2]}{1 + [1 + \mathcal{A}Br^2(1 + \mathcal{A}r^2)^2]\beta F^\diamond} dr^2 - r^2 d\Omega^2. \quad (80)$$

The physical variables associated with the above metric are defined in Appendix A as Eqs. (81)–(88).

V. PHYSICAL ANALYSIS OF THE MINIMALLY DEFORMED DURGAPAL-IV METRIC WITH $Y_{\text{TF}}^* = 0$

In this section, the physical feasibility of the anisotropically structured Class I Durgapal-IV metric is explored within the framework of the radial metric deformation scheme (MDG decoupling), satisfying the zero-complexity factor condition. It is well-established that physically and mathematically sound modeling of the relativistic stellar distributions must satisfy certain physical constraints. In this work, we analyze the salient features associated with the minimally deformed Durgapal-IV relativistic model under a complexity-free background. The physical properties of these types of anisotropic stellar solutions offer valuable insights into the formation and evolution of self-gravitational compact stars. We discuss the structural features corresponding to the complexity-free Durgapal-IV metric ansatz filled with anisotropic matter content using the radial metric deformation approach in FIGs. 1-4. These profiles depict various properties plotted against the radial variable r with decoupling parameter $\beta \in [0, 1]$. The characteristics include:

- Physical variables: energy density (σ^*), radial pressure (P_r^*), tangential pressure (P_θ^*) and pressure anisotropy (Δ^*).
- Structure scalars: density inhomogeneity (X_{TF}^*), complexity factor (Y_{TF}^*), strong energy condition (Y_{T}^*) and density homogeneity (X_{T}^*).

A. Behavior of Physical Variables

The behavior of energy density and stress components for different β -values are displayed in FIGs. 1 and 2. Their profiles describe that σ , P_r , and P_θ are gradually decreasing functions towards the stellar's boundary and exhibit a maximum at the center of the object, as required. We observe that σ^* , P_r^* , and P_θ^* decrease monotonically towards the stellar's surface for $\beta = 0.0, 0.19, 0.42, 0.61, 0.83$ and 1 as shown in FIGs. 1 and 2. Similarly, the anisotropy Δ^* decreases as we move away from the center for $\beta = 0.0, 0.19, 0.42, 0.61$, and 0.83 as displayed in the left panel of FIG. 2. However, for $\beta = 1$, anisotropy vanishes entirely throughout the model. This disappearance of anisotropy signifies a Class I isotropic solution, i.e., $P_r^* = P_\theta^*$ for all $r \in [0, R]$. This is consistent with a matter distribution described by the stress-energy tensor $T_{\mu\nu}^*$. The vanishing of pressure anisotropy reveals an important physical insight regarding the deviation of the anisotropic stellar-type configuration into the isotropic regime. Because under a zero-complexity factor background, with no explicit constraints imposing isotropy, the anisotropic matter content (represented by the SET, $T_{\mu\nu}^*$) effectively transforms into an isotropic fluid ($P_r = P_\theta$) without any explicit constraints enforcing isotropy. This interconnection between vanishing complexity and MGD decoupling offers a novel approach to constructing Durgapal-IV isotropic stellar models by assuming an initially anisotropic matter content. This finding has strong physical implications. Because, generally, GD has been employed for extending isotropic solutions to anisotropic domains or exploring more complex anisotropic scenarios. This solution shows that the nature of the matter distribution can be strongly influenced by the complexity factor in time-independent spherical stellar configurations. This points to a possible the connection between the geometric complexity of the system and the physical characteristics of the filled matter. Further investigation into this interconnection could provide valuable insights into the composition of relativistic stellar fluids, shedding light on the intricate relationships between their geometric and matter variables.

B. Behavior of Complexity factor

The variation of Y_{TF}^* for the minimally deformed Durgapal-IV solution is displayed in FIG. 3 (left panel) for different β -values. It is observed that Y_{TF}^* increases initially for a certain distance, followed by a decrease towards the boundary of the compact distribution. The value of Y_{TF}^* decreases as the parameter β is increased, and it vanishes when $\beta = 1$. This suggests that the complexity factor has less of an impact on the gravitationally decoupled stellar distribution. Another important finding is that Y_{TF}^* influences the matter variables, such as density and pressure. It affects the surface density, central pressure, and central density because the decline of these matter variables is directly linked with Y_{TF}^* . Furthermore, we see that the complexity factor Y_{TF}^* increases as we move towards the boundary of the model as displayed in FIG. 3 (left panel). However, for higher values of the decoupling constant β , the overall complexity factor decreases.

C. Energy Condition and Energy Density: Homogeneous and Inhomogeneous Scenarios

The behavior of other structural scalars, including the density homogeneity, strong energy condition, and density inhomogeneity, is observed for the permissible values of the parameter β as displayed in FIGS. 3 and 4. The solution profile encompasses the entire parameter space of the decoupling constant β . This profile satisfies all relevant scalar constraints, ensuring its physical viability. These constraints include the complexity factor, density inhomogeneity, strong energy conditions, and homogeneous energy density distribution. On the other hand, it is observed that as the value of β increases from 0 to 1, the values of Y_{T}^* and X_{T}^* decrease as shown in FIG. 4. Furthermore, the profile of X_{TF}^* displays that its value increases as β increases from 0 to 1 as shown in FIG. 4 (right panel).




FIG. 1: Plot of the energy density $[\sigma^*(r, \beta) \times 10^4]$ (left panel) and radial pressure $[P_r^*(r, \beta) \times 10^4]$ (right panel) for the Durgapal-IV model against the radial variable r with different values of $\beta \in [0, 1]$.




FIG. 2: Plot of the tangential pressure [$P_\theta^*(r, \beta) \times 10^4$] (left panel) and anisotropic factor [$\Delta^*(r, \beta) \times 10^4$] (right panel) corresponding to the Durgapal-IV model versus r with different β -values.




FIG. 3: Plot of the complexity factor [$Y_{TF}^*(r, \beta) \times 10^4$] (left panel) and density inhomogeneity [$X_{TF}^*(r, \beta) \times 10^4$] (right panel) subject to the Durgapal-IV model versus r with different values of the coupling constant β .




FIG. 4: Plot of strong energy condition $[Y_{\text{T}}^*(r, \beta) \times 10^4]$ (left panel) and density homogeneity $[X_{\text{T}}^*(r, \beta) \times 10^4]$ (right panel) for the Durgapal-IV model versus r with $\beta \in [0, 1]$.

VI. CONCLUSIONS

Determining the presence of anisotropic matter distribution in compact stars through direct observation is currently a significant challenge. The internal structure of compact stars cannot be directly observed using current observational methods. However, observations of phenomena like gravitational waves or extreme matter behavior can provide indirect clues about the internal structure of compact stars. Regarding the transition from an anisotropic matter configuration into an isotropic domain within compact stars, theoretical possibilities exist, but the extent to which this process occurs in the majority of compact stars remains uncertain. Evolutionary models and simulations indicate that physical processes like accretion or nuclear reactions can cause changes in the matter distribution within compact stars over time. However, the prevalence of such transformations among compact stars is an active area of research and debate. The MGD decoupling approach offers a straightforward and efficient framework for generating well-behaved solutions for EFEs. Despite its apparent simplicity, this approach provides profound insights into self-gravitating anisotropic configurations. Its primary advantage lies in breaking down a complex gravitational system into two simpler, decoupled sets. The first set corresponds to the standard EFEs for the considered seed matter content, while the second, influenced by the additional gravitational source term $\mathbb{X}_{\mu\nu}$, governs the anisotropic sector. This second system is often termed the quasi-Einstein gravitational system, leading to a departure from the standard EFEs. In this context, we have considered the Durgapal-IV model for exploring the dynamics of complexity-free stellar distribution. Subsequently, the quasi-Einstein equations necessitate supplementary information, either through constraints such as an EoS or the physically acceptable condition, such as a zero-complexity factor, for the source term $\mathbb{X}_{\mu\nu}$ or by specifying an appropriate representation of the MGD-function $F^\circ(r)$, as implemented in this study. This work demonstrates that the presented complexity-free Durgapal-IV model of anisotropic fluid spheres satisfies all necessary criteria for a valid solution. In this work, we have merged the embedding Class I approach with gravitational decoupling via the MGD scheme to construct complexity-free anisotropic solutions for stellar interiors. To investigate the potentiality of finding simple, analytical solutions for non-rotating, spherically symmetric, and stationary stellar structures with zero-complexity factor ($Y_{\text{TF}}^* = 0$), we use the MGD approach combined with the Durgapal-IV metric model. In this direction, we utilize the combination of the Class I condition and the condition $Y_{\text{TF}}^* = 0$ as the basis for the MGD function to construct physically realistic compact star models.

We have analyzed the structural features of self-gravitating, anisotropic stars within the Durgapal-IV model using gravitational decoupling. We analyzed various matter variables $\{\sigma^*, P_r^*, P_\theta^*, \Delta^*\}$ and four scalar quantities $\{Y_{\text{TF}}^*, X_{\text{TF}}^*, T_{\text{T}}^*, X_{\text{T}}^*\}$ for $\beta \in [0, 1]$. We have noted the impact of the decoupling parameter β on all matter variables within the framework of the Durgapal-IV relativistic model. The scalar function Y_{TF}^* used for measuring complexity notably affects the inherent structural characteristics of stellar interiors and facilitates the transition from an anisotropic to an isotropic solution. Specifically, when $\beta = 1$, indicating a vanishing complexity factor, solutions

derived from the Durgapal-IV ansatz shift from the initially considered anisotropic regime to the isotropic domain. This emphasizes the relevance of the complexity factor on physical features associated with self-gravitational stellar distributions through the mechanism of GD. In particular, we probed the captivating structural properties corresponding to the Durgapal-IV model: the disappearance of the anisotropic factor as the complexity factor approaches zero ($\beta = 1$). This finding involves the conversion of an anisotropic matter content of stress tensor constitution $T_{\mu\nu}^*$ into a homogeneous matter configuration, where $P_r = P_\theta$. This transformation is encouraged by GD, coupled with the condition $Y_{\text{TF}}^* = 0$, without requiring any isotropic precondition. Furthermore, we have observed significant trends in stress components throughout the solution employing the Durgapal-IV metric potential. Specifically, it has been observed that as $Y_{\text{TF}}^* \rightarrow 0$, there is a noticeable rise in pressure throughout the interior of the star, a situation considered undesirable. Moreover, we have revealed that contributions from GD, enforced by the constraint $T_{\text{TF}}^* = 0$, act to reduce the anisotropic factor. Interestingly, incorporating the structure scalar T_{TF}^* can lead to required outcomes, exerting substantial influence on all matter variables and producing a feasible solution for the minimum values of β .

In summary, this research effectively examined the influence of complexity on self-gravitational, anisotropic stars employing the MGD-based GD approach. This potent method allows for the generation of novel, physically plausible isotropic models for anisotropic fluid configurations by assuming the Durgapal metric potential within the formalism of embedding Class I method. Our results illustrate the pivotal role of the complexity factor in altering the distribution of matter within static, self-gravitational stellar interiors. The influence of complexity on the structural features of anisotropic stellar objects can be explored through the mechanism of the complete geometric deformation approach. However, this study specifically focuses on investigating the complexity of minimally deformed compact systems. The examination of the complexity factor for completely deformed stellar structures could be considered for future investigation. We have provided insights into the geometric properties and behavior of spacetime through matter-geometric relationships.

This finding is significant because the disappearance of pressure anisotropy throughout the model suggests that an anisotropic matter distribution transforms into an isotropic matter content under the paradigm of a zero-complexity factor, without necessitating any isotropy requirement. This result illustrates how the coupling of MGD decoupling with the concept of a zero-complexity factor can be employed to model the isotropic stellar structures that are originally filled with anisotropic matter content. Until now, gravitational decoupling has been utilized to extend isotropic solutions into anisotropic domains or from anisotropic to more potent anisotropic domains. This work differs from previous studies in which the notion of GD is generally used to extend isotropic stellar solutions to the anisotropic domain. In the past, several attempts have been made to extend isotropic stellar configurations into the anisotropic regime within the GD approach, including the Durgapal-IV metric [62–65]. However, this work features a novel aspect of the GD approach: the coupling of GD with a zero-complexity factor may be used to transform the originally anisotropic matter distribution into an isotropic domain with any pre-assumed isotropic condition.

APPENDIX A

The physical variables associated with the \mathbb{X} -gravitational sector take the form

$$8\pi\sigma^{\mathbb{X}} = \frac{\beta\mathcal{A}[6 + 3Ar^2 + \mathcal{A}^2r^4 + \mathcal{A}B^2r^2(1 + \mathcal{A}r^2)^4(-1 + 3Ar^2) - B(1 + \mathcal{A}r^2)^2(3 + Ar^2 + 11\mathcal{A}^2r^4 + 5\mathcal{A}^3r^6)]}{(1 + \mathcal{A}r^2)^3(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)^2}, \quad (81)$$

$$8\pi P_r^{\mathbb{X}} = \frac{\beta\mathcal{A}(1 + 9Ar^2)[-2 - Ar^2 + B(1 + \mathcal{A}r^2)^2]}{(1 + \mathcal{A}r^2)^3(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)}, \quad (82)$$

$$8\pi P_\theta^{\mathbb{X}} = -\frac{\beta\mathcal{A}[2(-1 + 3Ar^2 + 20\mathcal{A}^2r^4 + 8\mathcal{A}^3r^6) + B(1 + \mathcal{A}r^2)^2(1 - 2Ar^2 + 12\mathcal{A}^2r^4 + 78\mathcal{A}^3r^6 + 31\mathcal{A}^4r^8)]}{(1 + \mathcal{A}r^2)^4(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)^2} + \frac{\beta\mathcal{A}[2\mathcal{A}^2B^2(5 + 13Ar^2)(r + \mathcal{A}r^3)^4]}{(1 + \mathcal{A}r^2)^4(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)^2}, \quad (83)$$

where $\sigma^{\mathbb{X}}$, $P_r^{\mathbb{X}}$, and $P_\theta^{\mathbb{X}}$ denote the physical variables associated with the decoupling fluid $\mathbb{X}_{\mu\nu}$. The anisotropic factor associated with the Durgapal-IV metric potential takes the form

$$\begin{aligned} \Delta^*(r, \beta) = & [\mathcal{A}(-4\beta + \mathcal{A}B^2r^2(1 + \mathcal{A}r^2)^4(-2 - 6\mathcal{A}r^2 + 14\mathcal{A}^3r^6 + 8\mathcal{A}^4r^8 + \mathcal{A}^2r^4(2 - 17\beta) + \beta) + 4(-4 + 3\beta) \\ & \times \mathcal{A}^2r^4 + \mathcal{A}^3r^6(-8 + 7\beta) - \mathcal{A}r^2(8 + 15\beta) + 2B(1 + \mathcal{A}r^2)^2(-1 - 3\mathcal{A}^2r^4 + 3\mathcal{A}r^2(\beta - 1) + \beta + 11\beta\mathcal{A}^4 \\ & \times r^8 + \mathcal{A}^3(-1 + 25\beta)r^6)] / [(1 + \mathcal{A}r^2)^4(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)^2]. \end{aligned} \quad (84)$$

The corresponding value of the complexity factor Y_{TF}^* turn out to be

$$Y_{\text{TF}}^*(r, \beta) = (1 - \beta) \frac{4\mathcal{A}^2r^2[-2 + B(1 + \mathcal{A}r^2)^3]}{[(1 + \mathcal{A}r^2)^2(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)^2]}. \quad (85)$$

This expression captures the effects of the \mathbb{X} -gravitational source on the complexity of self-gravitational compact star by interpolating between the original value of β , i.e., $\beta = 0$ and complexity-free condition ($\beta = 1$). The corresponding values of the other structure scalars read

$$\begin{aligned} X_{\text{TF}}^*(r, \beta) = & -\mathcal{A}^2r^2[B^2(1 + \mathcal{A}r^2)^4(1 + 3\mathcal{A}^2r^4 + \mathcal{A}^3r^6 - 3\mathcal{A}r^2(\beta - 1) - \beta + 2B(1 + \mathcal{A}r^2)^2 \\ & \times (\beta - 1 + \mathcal{A}^2r^4(2\beta - 1) + \mathcal{A}r^2(5\beta - 2))] / [(1 + \mathcal{A}r^2)^3(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)^2], \end{aligned} \quad (86)$$

$$\begin{aligned} Y_{\text{T}}^*(r, \beta) = & -[A(-20 + 70\mathcal{A}^8B^2r^{16} + 11\mathcal{A}^9B^2r^{18} + 2\mathcal{A}^7B^2r^{14}(95 - 16\beta) + 2(B - 2)\beta + \mathcal{A}r^2(-68 - 20B + B^2 \\ & + 12\beta) + 2\mathcal{A}^6Br^{12}(-10 + B(143 - 72\beta) + 19\beta) + 2\mathcal{A}^4Br^8(-100 + B(73 - 112\beta) + 131\beta) - 2\mathcal{A}^2r^4 \\ & \times (38 + B(50 - 9\beta) - 26\beta + B^2(-5 + 8\beta)) + 2\mathcal{A}^3r^6(B^2(25 - 48\beta) + 2(-7 + 5\beta) + 4B(-25 + 18\beta)) \\ & - 4\mathcal{A}^5Br^{10}(25 - 44\beta + B(-65 + 64\beta))] / [(1 + \mathcal{A}r^2)^3(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)^2], \end{aligned} \quad (87)$$

$$\begin{aligned} X_{\text{T}}^*(r, \beta) = & [\mathcal{A}(-4\beta + \mathcal{A}B^2r^2(1 + \mathcal{A}r^2)^4(-2 - 6\mathcal{A}r^2 + 14\mathcal{A}^3r^6 + 8\mathcal{A}^4r^8 + \mathcal{A}^2r^4(2 - 17\beta) + \beta) + (-4 + 3\beta) \\ & \times 4\mathcal{A}^2r^4 + \mathcal{A}^3r^6(-8 + 7\beta) - \mathcal{A}r^2(8 + 15\beta) + 2B(1 + \mathcal{A}r^2)^2(-1 - 3\mathcal{A}^2r^4 + 3\mathcal{A}r^2(\beta - 1) + \beta + 11\mathcal{A}^4 \\ & \times \beta r^8 + \mathcal{A}^3r^6(-1 + 25\beta))] / [(1 + \mathcal{A}r^2)^4(1 + \mathcal{A}Br^2 + 2\mathcal{A}^2Br^4 + \mathcal{A}^3Br^6)^2]. \end{aligned} \quad (88)$$

where X_{TF}^* , Y_{TF}^* , X_{T}^* , and Y_{T}^* denote density inhomogeneity, complexity factor, density homogeneity, and the strong energy condition for the compact system, respectively.

ACKNOWLEDGEMENT

The work of KB was partially supported by the JSPS KAKENHI Grant Number 21K03547 and 23KF0008. The work by BA was supported by Researchers Supporting Project number: RSPD2024R650, King Saud University, Riyadh, Saudi Arabia.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

This manuscript has no associated data or the data will not be deposited. [Authors comment: This manuscript contains no associated data.]

- [2] H. Weyl, *Phys. Z* **20**, 65 (1919).
- [3] R. C. Tolman, *Phys. Rev.* **55**, 364 (1939).
- [4] Z. Stuchlík, S. Hledík, and J. Novotný, *Phys. Rev. D* **94**, 103513 (2016).
- [5] J. Novotný, J. Hladík, and Z. Stuchlík, *Phys. Rev. D* **95**, 043009 (2017).
- [6] S. Hod, *Phys. Rev. D* **97**, 084018 (2018).
- [7] S. Hod, *Eur. Phys. J. C* **78**, 417 (2018).
- [8] Z. Stuchlík, J. Schee, B. Toshmatov, J. Hladík, and J. Novotný, *J. Cosmol. Astropart. Phys.* **2017**, 056 (2017).
- [9] G. Lemaître, *Ann. Soc. Sci. Bruxelles A* **53**, 51 (1933).
- [10] M. Ruderman, *Ann. Rev. Astron. Astrophys* **10**, 427 (1972).
- [11] R. L. Bowers and E. Liang, *Astrophys. J.* **188**, 657 (1974).
- [12] M. Delgaty and K. Lake, *Comput. Phys. Commun.* **115**, 395 (1998).
- [13] L. Herrera, *Phys. Lett. A* **165**, 206 (1992).
- [14] L. Herrera and N. O. Santos, *Phys. Rep.* **286**, 53 (1997).
- [15] M. Mak and T. Harko, *Proc. R. Soc. Lond. A* **459**, 393 (2003).
- [16] B. Ivanov, *Phys. Rev. D* **65**, 104001 (2002).
- [17] F. E. Schunck and E. W. Mielke, *Class. Quantum Grav.* **20**, R301 (2003).
- [18] V. V. Usov, *Phys. Rev. D* **70**, 067301 (2004).
- [19] D. Deb, M. Khlopov, F. Rahaman, S. Ray, and B. Guha, *Eur. Phys. J. C* **78**, 465 (2018).
- [20] F. Rahaman, S. Ray, A. K. Jafry, and K. Chakraborty, *Phys. Rev. D* **82**, 104055 (2010).
- [21] V. Varela, F. Rahaman, S. Ray, K. Chakraborty, and M. Kalam, *Phys. Rev. D* **82**, 044052 (2010).
- [22] S. Khan, A. Adeel, and Z. Yousaf, *Eur. Phys. J. C* **84**, 572 (2024).
- [23] Z. Yousaf, B. M. Z., and S. Khan, *Ann. Phys.* **534**, 2200252 (2022).
- [24] R. Kippenhahn, A. Weigert, and A. Weiss, *Stellar structure and evolution*, vol. 192 (Springer, 1990).
- [25] R. F. Sawyer, *Phys. Rev. Lett.* **29**, 382 (1972).
- [26] J. B. Hartle, R. F. Sawyer, and D. J. Scalapino, *Astrophys. J.* **199**, 471 (1975).
- [27] L. Herrera, *Phys. Rev. D* **97**, 044010 (2018).
- [28] L. Herrera, A. Di Prisco, and J. Ospino, *Phys. Rev. D* **98**, 104059 (2018).
- [29] L. Herrera, A. Di Prisco, and J. Ospino, *Phys. Rev. D* **99**, 044049 (2019).
- [30] L. Herrera, A. Di Prisco, and J. Ospino, *Eur. Phys. J. C* **80**, 631 (2020).
- [31] M. Z. Bhatti, M. Y. Khlopov, Z. Yousaf, and S. Khan, *Mon. Not. R. Astron. Soc.* **506**, 4543 (2021).
- [32] M. Z. Bhatti, Z. Yousaf, and S. Khan, *Eur. Phys. J. Plus* **136**, 975 (2021).
- [33] Z. Yousaf, M. Z. Bhatti, S. Khan, and P. K. Sahoo, *Phys. Dark Universe* **36**, 101015 (2022).
- [34] Z. Yousaf, S. Khan, N. B. Turki, and T. Suzuki, *Chinese J. Phys.* **89**, 1595 (2024).
- [35] L. Herrera, A. Di Prisco, and J. Ospino, *Phys. Rev. D* **103**, 024037 (2021).
- [36] Z. Yousaf, K. Bamba, M. Bhatti, and U. Farwa, arXiv preprint [arXiv:2311.10369](https://arxiv.org/abs/2311.10369) (2023).
- [37] R. Casadio, J. Ovalle, and R. Da Rocha, *Class. Quantum Grav.* **32**, 215020 (2015).
- [38] J. Ovalle, in *Int. J. Mod. Phys. Conf. Ser.* (World Scientific, 2016), vol. 41, p. 1660132.
- [39] J. Ovalle, *Int. J. Mod. Phys. D* **18**, 837 (2009).
- [40] J. Ovalle, *Mod. Phys. Lett. A* **25**, 3323 (2010).
- [41] R. Casadio and J. Ovalle, *Phys. Lett. B* **715**, 251 (2012).
- [42] J. Ovalle and F. Linares, *Phys. Rev. D* **88**, 104026 (2013).
- [43] R. Casadio, J. Ovalle, and R. Da Rocha, *Class. Quantum Grav.* **31**, 045016 (2014).
- [44] J. Ovalle, L. A. Gergely, and R. Casadio, *Class. Quantum Grav.* **32**, 045015 (2015).
- [45] R. da Rocha, *Eur. Phys. J. C* **77**, 355 (2017).
- [46] R. Casadio, P. Nicolini, and R. Da Rocha, *Class. Quantum Grav.* **35**, 185001 (2018).
- [47] J. Ovalle, *Phys. Rev. D* **95**, 104019 (2017).
- [48] J. Ovalle, R. Casadio, R. da Rocha, and A. Sotomayor, *Eur. Phys. J. C* **78**, 122 (2018).
- [49] S. Khan and Z. Yousaf, *Phys. Scr.* **99**, 055303 (2024).
- [50] G. Abellán, V. Torres-Sánchez, E. Fuenmayor, and E. Contreras, *Eur. Phys. J. C* **80**, 177 (2020).
- [51] A. M. Albalahi, M.Z. Bhatti, A. Ali, and S. Khan, *Eur. Phys. J. C* **84**, 293 (2024).
- [52] A. M. Albalahi, Z. Yousaf, A. Ali, and S. Khan, *Eur. Phys. J. C* **84**, 9 (2024).
- [53] F. Tello-Ortiz, S. Maurya, and Y. Gomez-Leyton, *Eur. Phys. J. C* **80**, 324 (2020).
- [54] M. Durgapal and R. Fuloria, *Gen. Relativ. Gravit.* **17**, 671 (1985).
- [55] R. C. Tolman, *Phys. Rev.* **35**, 875 (1930).
- [56] L. Herrera, J. Ospino, A. Di Prisco, E. Fuenmayor, and O. Troconis, *Phys. Rev. D* **79**, 064025 (2009).
- [57] L. Herrera, A. Di Prisco, and J. Ibanez, *Phys. Rev. D* **84**, 107501 (2011).
- [58] K. Karmarkar, in *Proc. Indian Acad. Sci. A* (Springer, 1948), vol. 27, p. 56.

- [59] S. Pandey and S. Sharma, *Gen. Relativ. Gravit.* **14**, 113 (1982).
- [60] K. Lake, *Phys. Rev. D* **67**, 104015 (2003).
- [61] P. Vaidya and R. Tikekar, *J Astrophys. Astron.* **3**, 325 (1982).
- [62] E. Contreras, E. Fuenmayor, and G. Abellán, *Eur. Phys. J. C* **82**, 187 (2022).
- [63] J. Andrade and E. Contreras, *Eur. Phys. J. C* **81**, 889 (2021).
- [64] B. Das, S. Dey, S. Das, and B. C. Paul, *Eur. Phys. J. C* **82**, 519 (2022).
- [65] S. Maurya, A. Errehymy, M. Jasim, S. Hansraj, N. Al-Harbi, and A.-H. Abdel-Aty, *Eur. Phys. J. C* **82**, 1173 (2022).