

# Superluminous supernovae

Takashi J. Moriya<sup>a,b,c</sup>

<sup>a</sup>National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

<sup>b</sup>Graduate Institute for Advanced Studies, SOKENDAI, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

<sup>c</sup>Monash University, School of Physics and Astronomy, Clayton, Victoria 3800, Australia

© 20xx Elsevier Ltd. All rights reserved.

Chapter Article tagline: update of previous edition,, reprint..

## Glossary

**Circumstellar matter** Materials surrounding supernova progenitors.

**Luminosity function** Peak luminosity distribution of supernovae.

**Magnetar** Strongly magnetized rapidly rotating neutron star.

**Type I SNe** Supernovae without spectroscopic features of hydrogen. Type Ib SNe show helium features, while Type Ic SNe show neither hydrogen nor helium features.

**Type II SNe** Supernovae with spectroscopic features of hydrogen.

**Type IIn/Ibn/Icn SNe** Supernovae with narrow emission features indicating the existence of dense circumstellar matter.

## Nomenclature

CCSN	Core-collapse Supernova
CSM	Circumstellar Matter
FBOT	Fast Blue Optical Transient
FRB	Fast Radio Burst
GRB	Gamma-Ray Burst
LSST	Legacy Survey of Space and Time
PS1	Pan-STARRS1
PTF	Palomar Transient Factory
SLSN	Superluminous Supernova
SN	Supernova
ZAMS	Zero-Age Main Sequence
ZTF	Zwicky Transient Facility

## Abstract

Superluminous supernovae (SLSNe) are a population of supernovae (SNe) whose peak luminosities are much larger than those of canonical SNe. Although SLSNe were simply defined by their peak luminosity at first, it is currently recognized that they show rich spectroscopic diversities including hydrogen-poor (Type I) and hydrogen-rich (Type II) subtypes. The exact mechanisms making SLSNe luminous are still not fully understood, but there are mainly four major suggested luminosity sources (radioactive decay of  $^{56}\text{Ni}$ , circumstellar interaction, magnetar spin-down, and fallback accretion). We provide an overview of observational properties of SLSNe and major theoretical models for them. Future transient surveys are expected to discover SLSNe at high redshifts which will provide a critical information in revealing their nature.

### Key Points

- SLSNe are a class of SNe that become more luminous than around  $-20$  mag in the optical. Broadly, SLSNe have two spectroscopic types: hydrogen-poor (Type I) and hydrogen-rich (Type II).
- Hydrogen-poor (Type I) SLSNe are characterized by O II absorption features around the peak luminosity. Their luminosity sources and progenitors are still debated.
- Most hydrogen-rich (Type II) SLSNe have narrow hydrogen emission features (Type IIn) indicating the existence of dense CSM. Thus, they are powered by the interaction between SN ejecta and dense, massive ( $\geq 5 M_{\odot}$ ) CSM. The origin of such a CSM is still not clear.

## 1 Introduction

Superluminous supernovae (SLSNe) are a class of SNe that become more luminous than other kind of SNe at their peak luminosity. The existence of a population of such a luminous SN was not recognized until the 2000s. The first glimpse was found in SN 1999as (Knop et al., 1999; Deng et al., 2001), but its nature remained unclear for a long time. SLSNe started to be discovered frequently when unbiased SN

## 2 Superluminous supernovae

surveys were started to be conducted in the 2000s. The first well-observed SLSNe include SN 2005ap (Quimby et al., 2007), SN 2006gy (e.g., Smith et al., 2008, 2010), and SCP06F6 (Barbary et al., 2009). Some mysterious SNe whose origin was unclear when they were discovered were later identified as SLSNe (e.g., Kostrzewa-Rutkowska et al., 2013; Cartier et al., 2022). Several hundred SLSNe have been identified so far (Nicholl, 2021). Several review papers on SLSNe are available for further reading (Gal-Yam, 2012, 2019b; Howell, 2017; Moriya et al., 2018b; Inserra, 2019; Nicholl, 2021).

### 2 Definition

SLSNe have intrinsically higher luminosity than other SNe. SLSNe are initially defined as SNe that have peak luminosity more luminous than  $-21$  mag in the optical (see Gal-Yam 2012 for an early review). This luminosity cut of  $-21$  mag is about 10 times larger than the peak luminosities of commonly observed SNe. However, as the number of SN discovery increases, it was recognized that there are many SNe with peak luminosity fainter than  $-21$  mag showing similar spectroscopic characteristics to SNe with peak luminosity brighter than  $-21$  mag. In other words, it was recognized that SNe that show characteristic spectroscopic features of SLSNe do not necessarily have a clear magnitude cut. For example, the peak luminosity of SNe with hydrogen-free (Type I) SLSN spectroscopic features can be as faint as around  $-20$  mag as discussed in Section 3.1.2. In addition, some rapidly-evolving SNe like so-called fast blue optical transients (FBOTs) exceed  $-21$  mag in a short time ( $\lesssim 3$  days) after explosion, but they have different spectroscopic features from canonical SLSNe and they are usually not referred to as SLSNe (e.g., Ho et al., 2023). Thus, at least for Type I SLSNe, SLSNe are defined as a population of luminous SNe with characteristic spectroscopic features rather than SNe that exceed a certain luminosity cut. The observed number of other SLSNe such as Type II SLSNe (Section 3.2) is still too limited to characterize them by their spectroscopic features. In such cases, SNe more luminous than around  $-20$  mag are naively called SLSNe following the luminosity range of Type I SLSNe.

### 3 Observational properties

As in the case of other SNe, SLSNe can be broadly classified into two spectroscopic classes based on the presence or absence of hydrogen features in their spectra. Type I SLSNe are SLSNe without hydrogen features and Type II SLSNe are SLSNe with hydrogen features. We summarize their observational properties in this section.

#### 3.1 Type I SLSNe

Type I SLSNe have been actively observed by many transient surveys. Summaries of Type I SLSN samples from major transient surveys so far can be found in De Cia et al. (2018, PTF), Quimby et al. (2018, PTF), Lunnan et al. (2018a, PS1), Angus et al. (2019, DES), and Chen et al. (2023a, ZTF). Gomez et al. (2024) also provides a summary of Type I SLSN properties.


##### 3.1.1 Spectroscopic properties

Type I SLSNe are characterized by their O II absorption features in the wavelength range between  $3000 \text{ \AA}$  and  $5000 \text{ \AA}$  observed around their luminosity peak (Quimby et al. 2011, Figure 1). No other prominent spectroscopic features appear in optical spectra around the luminosity peak. Thermal excitation is suggested to be sufficient to excite oxygen to form the O II features if the photospheric temperature is around  $14,000 - 16,000 \text{ K}$  (Saito et al., 2024). Thus, the O II features can present diverse temporal evolution and strength depending on the photospheric temperature evolution of Type I SLSNe (Könyves-Tóth, 2022). Most Type I SLSNe do not show helium features and thus they are Type Ic SNe. Only a few SLSNe are found to have helium (Type Ib SN) features so far and their peak luminosity is rather faint (around  $-20$  mag in the optical, Yan et al. 2020). Photospheric velocities estimated by Fe II lines exceeds  $10,000 \text{ km s}^{-1}$  at around the luminosity peak (Nicholl et al., 2015a; Quimby et al., 2018). The temporal evolution of photospheric velocities is overall found to be similar to those of Type Ic-BL (Liu et al., 2017).

The rest-frame ultraviolet spectra around the light-curve peak of Type I SLSNe are diverse (Vreeswijk et al., 2014; Nicholl et al., 2017b; Yan et al., 2017b, 2018). Ultraviolet spectra below  $3000 \text{ \AA}$  have attenuation due to metal absorption, but the amount of attenuation varies. Strong absorptions in the ultraviolet spectra are likely formed by combinations of the absorptions of several metal lines (Howell et al., 2013; Mazzali et al., 2016; Quimby et al., 2018; Gal-Yam, 2019a).

As the photospheric temperature declines, optical spectra starts to show line features of diverse elements as in other Type I SNe and they evolve to nebular phases. In some cases (e.g., SN 2007bi, Gal-Yam et al. 2009), strong Ca II emissions start to appear on top of the photospheric spectral features. The origin of these early strong Ca II emissions is still unclear. Around 10 Type I SLSNe are observed until the entire ejecta become transparent (so-called the “nebular” phase). In the nebular phases, Type I SLSNe are found to have similar spectral features to those of Type Ic-BL (Nicholl et al., 2019). They show strong, broad emission lines of O II and Ca II, for example. This fact indicates that the physical conditions at the central regions of Type I SLSNe and Type Ic-BL SNe are likely similar.

Another notable characteristic of Type I SLSN is the emergence of hydrogen emission at late phases observed in a few Type I SLSNe (Yan et al., 2017a). The late-phase hydrogen emission is likely to indicate the existence of detached hydrogen-rich dense circumstellar matter (CSM) surrounding the progenitors, although it may also originate from hydrogen stripped from the companion star of the progenitors (Moriya et al., 2015). The late-time emergence of hydrogen emission lines is also observed in a couple of Type Ibc SNe (Milisavljevic et al., 2015; Chen et al., 2018; Kuncarayakti et al., 2018). Even if no clear interaction signatures are found, the existence of relatively dense



**Fig. 1** Spectra of Type I SLSNe compared with other hydrogen-free SNe. The shaded regions indicate the characteristic spectroscopic feature of each SN type. Type I SLSNe are characterized by the O II absorption features between 3000 Å and 5000 Å. The characteristic features of the other SN types are H I (Type II), Si II (Type Ia), and He I (Type Ib). Reproduced with permission from Quimby et al. (2018).

CSM surrounding Type I SLSNe is sometimes imprinted in their spectra (e.g., Lunnan et al., 2018b).

### 3.1.2 Light-curve properties

The luminosity evolution of Type I SLSNe shows diversity. Their peak luminosity ranges from  $-20$  mag to  $-23$  mag in optical or  $3 \times 10^{43}$  erg  $s^{-1}$  to  $8 \times 10^{44}$  erg  $s^{-1}$  in bolometric (e.g., Chen et al. 2023a). The peak luminosity distribution of Type I SLSNe is consistent with an extrapolation from the lower luminosity Type I SNe including Type Ibc and Type Ic-BL SNe (Figure 2). In other words, Type I SLSNe do not likely make a separate population in hydrogen-free SNe but they are higher luminosity extension of hydrogen-free SNe. Intermediate SNe are briefly discussed in Section 3.1.10.

The mean rise time of Type I SLSNe is estimated to be  $41.9 \pm 17.8$  days in the  $g$  band with the  $1\sigma$  dispersion in the recent ZTF sample study by Chen et al. (2023a). The rise time is much shorter than the mean rise time of Type Ibc SNe ( $\sim 20$  days, e.g., Drout et al. 2011). However, there are Type I SLSNe with the rise time as short as about 10 days (Chen et al., 2023a). To the other extreme, some Type I SLSNe have a very long rise time exceeding 100 days (e.g., PS1-14bj, Lunnan et al. 2016; SN 2018ibb, Schulze et al. 2024). These extreme cases are, however, found to be rare. The rise time and the decline time of Type I SLSNe are positively correlated, i.e., slowly rising Type I SLSNe tend to decline slowly. There has been a suggestion that Type I SLSNe can be divided into two populations of slowly evolving Type I SLSNe and rapidly evolving Type I SLSNe (e.g., Inserra et al., 2018a). It has also been suggested that Type I SLSNe may have a relation between light-curve decline rate and peak luminosity as in Type Ia SNe and they could be a potential standardizable candle (Inserra and Smartt, 2014; Inserra et al., 2021; Khetan et al., 2023).

The luminosity evolution of Type I SLSNe is not merely characterized by a simple rise and fall. First, it is known that some Type I SLSNe show a precursor before the major luminosity increase. The precursor was first identified in SN 2006oz by Leloudas et al. (2012), which showed a precursor luminosity “bump” before the major luminosity increase. Subsequently, LSQ14bdq (Nicholl et al., 2015b) and DES14X3taz (Smith et al., 2016) are found to have a clear precursor bump. The precursor bump lasts for about 10 days. No spectrum during the bump has been obtained so far, but the color during the bump indicates that the bump should have a very hot spectrum (Nicholl and Smartt, 2016). Recent studies on a large number of Type I SLSNe indicate that such a precursor luminosity bump is not a ubiquitous feature of Type I SLSNe and only a fraction (40% or less) of Type I SLSNe show the precursor bump (Angus et al., 2019). In one case of




Fig. 2 Luminosity function of hydrogen-poor SNe. Reproduced with permission from De Cia et al. (2018).

SN 2018bsz, the precursor was found to have a gradual increase in luminosity without showing the temporal luminosity decline before the major luminosity increase (Anderson et al. 2018). In some cases, the early bumps may not be prominent and they may be observed as early flux excess (Vreeswijk et al., 2017).

After the luminosity peak, a significant fraction of Type I SLSNe show undulations in their light curves (Inserra et al., 2017; Chen et al., 2023b). Even in Type I SLSNe without clear undulations, there may often exist an underlying flux excess in their light curves (Chen et al., 2017a; Hosseinzadeh et al., 2022). In some clear cases, we can even observe a secondary luminosity peak in light curves (Hosseinzadeh et al., 2022), which are also sometimes observed in Type Ic SNe that have lower luminosity than Type I SLSNe (e.g., Gomez et al., 2021). Optical color and spectral line features do not change significantly during the undulations.


Only a couple of Type I SLSNe have light-curve information beyond 1000 days after explosion (e.g., Blanchard et al., 2021). The light-curve decline rates at the very late phase are found to be diverse even among these cases. The late-phase luminosity evolution might provide an important clue in their luminosity source as discussed in Section 5.

### 3.1.3 Polarimetric properties

Polarimetric properties of Type I SLSNe have been investigated to constrain their ejecta geometry (see Pursiainen et al. 2023 for a summary of Type I SLSNe with polarimetric observations). No significant polarization is observed in many cases, which indicates that ejecta are not far from spherical symmetry. However, polarization is indeed observed in some cases. For Type I SLSNe with significant polarization, it is often found that the polarization degree increases with time. This indicates that the asphericity of the ejecta in Type I SLSNe increases with time. During the period when the significant polarization degree is observed, some signatures of the CSM interaction is also identified in many cases (Pursiainen et al., 2023). While most polarimetric observations have been conducted to measure linear polarization, PS17bek is the only Type I SLSN with circular polarimetric observations (Cikota et al., 2018). However, no significant circular polarization is identified in PS17bek.

### 3.1.4 X-ray properties

Type I SLSNe have been extensively observed in X-ray, but they have been detected only in a couple of cases (e.g., Margutti et al., 2018). The first case is SCP06F6 (Levan et al., 2013). It was detected in 0.2–2.0 keV by XMM-Newton at around 150 days after the discovery, and its corresponding rest-frame luminosity is  $\approx 10^{45}$  erg s<sup>-1</sup>. This X-ray luminosity is three orders of magnitude higher than those observed in other SNe (Levan et al., 2013), making SCP06F6 the most luminous SN observed in X-ray. SCP06F6 was observed again 3 months after the



**Fig. 3** Summaries of the observations of Type I SLSNe in X-rays (left) and radio (right). Reproduced with permission from Margutti et al. (2018, left) and Margutti et al. (2023, right).

detection, but it was not detected ( $< 2.5 \times 10^{44}$  erg s<sup>-1</sup>, Levan et al. 2013). Many other Type I SLSNe were observed in similar epochs with sufficient depths to detect them if they are as bright as SCP06F6, but SCP06F6 remains to be the only detection with such a high luminosity among Type I SLSNe (Margutti et al. 2018, Figure 3).

PTF12dam is another case of the X-ray detection from a Type I SLSN (Margutti et al., 2018). It was detected by the Chandra X-ray Observatory at around the optical luminosity peak, and its X-ray luminosity was  $\sim 2 \times 10^{40}$  erg s<sup>-1</sup> in 0.3 – 10 keV. The X-ray luminosity is, however, consistent with the expected diffuse X-ray luminosity from the underlying star forming activities. Thus, it is possible that a significant fraction of the observed X-ray luminosity is not from PTF12dam itself.

Finally, a potential Type I SLSN ASASSN-15lh (Dong et al., 2016) was detected in X-ray (Margutti et al., 2017), but its nature as a Type I SLSN has been debated and questioned (Leloudas et al., 2016).

### 3.1.5 Mid-infrared properties

Some observations of Type I SLSNe in mid-infrared are currently available. SN 2018bsz, a nearby Type I SLSN, was observed by Spitzer and it was detected in 3.6  $\mu$ m and 4.5  $\mu$ m at around 400 days and 550 days after the *r* band luminosity peak in the rest frame (Chen et al., 2021). It was also detected by WISE in 3.4  $\mu$ m and 4.6  $\mu$ m at around 250 days and 400 days after the *r* band luminosity peak. Sun et al. (2022) investigated WISE data, and found some additional Type I SLSNe with mid-infrared detections. The mid-infrared detection indicates the existence of dust in Type I SLSNe created before and/or after their explosions.


### 3.1.6 Radio properties

Radio information of Type I SLSNe is suggested to play a key role in constraining their powering mechanism (Section 5.3), and radio follow-up observations of Type I SLSNe have been conducted actively (see Margutti et al. 2023 for a current summary; Figure 3). A couple of Type I SLSNe have been detected in radio. Type I SLSN PTF10hgi was the first case of the radio detection from Type I SLSNe (e.g., Eftekhari et al. 2019). It was detected in 1 – 20 GHz from around 7.5 years after the explosion and its radio luminosity was  $\simeq 10^{28}$  erg s<sup>-1</sup> Hz<sup>-1</sup> at 6 GHz.

SN 2017ens (Chen et al., 2018) is a SN that reached the peak luminosity of  $-21.1$  mag in the *g* band and its luminosity is consistent with being a Type I SLSN. Although it did not show the characteristic O II features of Type I SLSNe, they might have been missed because of infrequent spectroscopic observations. SN 2017ens first had broad features similar to those of Type Ic-BL at around the peak luminosity. However, it showed Type II<sub>n</sub> SN features after 160 days since the luminosity peak which indicate that the ejecta started to interact with a detached hydrogen-rich dense CSM. SN 2017ens was detected in 3 – 10 GHz from around 3.3 years after explosion, and its radio luminosity was  $\simeq 10^{28}$  erg s<sup>-1</sup> Hz<sup>-1</sup> at 6 GHz (Margutti et al., 2023). Given the presence of the late-phase CSM interaction signatures in optical spectra, the radio luminosity may originate from the CSM interaction.

### 3.1.7 Gamma-ray properties

Possible gamma-ray detection from SN 2017egm by the Fermi satellite is reported by Li et al. (2024). The gamma-ray (500 MeV – 500 GeV) was detected at 100 – 150 days after the discovery.



**Fig. 4** Stellar masses and star-formation rates (SFRs) of SN host galaxies in the PTF core-collapse SN sample. The host galaxies of Type I SLSNe tend to have high star-formation efficiencies. Reproduced with permission from Schulze et al. (2021).

### 3.1.8 Association with gamma-ray bursts

A potential Type I SLSN, SN 2011kl, was associated with ultra-long GRB 111209A (Greiner et al., 2015). The spectrum of SN 2011kl was not good enough to identify their spectral features, but it reached  $-20$  mag at the peak luminosity. Long GRB 140506A was associated with a possible luminous blue SN component which might indicate a potential association between normal long GRBs and SLSNe (Kann et al., 2024). However, no spectrum was obtained for the potential SN component. Association between GRBs and SLSNe has not been fully investigated and further observational studies are required.

### 3.1.9 Host environments

Type I SLSNe are observed to prefer low-metallicity environments (Neill et al., 2011). Most Type I SLSNe are observed below around  $0.5 Z_{\odot}$  (Perley et al., 2016; Chen et al., 2017c; Schulze et al., 2018), although some SLSNe such as SN 2017egm are exceptionally found in high metallicity environments that are similar to typical core-collapse SNe (Bose et al., 2018; Nicholl et al., 2017a; Chen et al., 2017b). The host galaxies of Type I SLSNe tend to be low stellar-mass galaxies (Wiseman et al., 2020). They also tend to have high star-formation efficiencies of around  $10^{-9} - 10^{-7} \text{ yr}^{-1}$  (e.g., Schulze et al. 2021; Figure 4). This might indicate that Type I SLSNe are explosions of very massive stars that occur immediately after the star formation, although simply associating high star-formation efficiencies to massive progenitors has been questioned (Cleland et al., 2023). Some Type I SLSNe are also found to be associated with dense molecular clouds with active star formation (Arabsalmani et al., 2019; Hatsukade et al., 2020). Some host galaxies of Type I SLSNe are interacting galaxies (Chen et al., 2017a; Cikota et al., 2017; Ørum et al., 2020) or compact dwarf irregular galaxies with extremely strong emission lines likely experiencing active star formations (Leloudas et al., 2015; Lunnan et al., 2015; Angus et al., 2016). Type I SLSNe are found to have a tendency to explode further away from the host galaxy center than any other types of SNe as well as long GRBs (Hsu et al., 2024).

The similarities between the host galaxies of Type I SLSNe and long GRBs have been explored in many studies (e.g., Lunnan et al., 2014). While some differences among them are pointed out (Leloudas et al., 2015; Hsu et al., 2024), their host galaxies are rather similar without much statistical differences (Japelj et al., 2016; Taggart and Perley, 2021). The host galaxies of fast radio bursts (FRBs) are often found to be different from those of Type I SLSNe, although the host galaxies of repeating FRBs might have some similarities (Bhandari et al., 2022).


### 3.1.10 Transitional SNe

Type I SNe with the light-curve peak magnitude range of  $\sim -20$  mag  $-\sim -19$  mag in optical show diverse spectroscopic features (Gomez et al., 2022). Some of them show hot spectra similar to Type I SLSNe while others show cool spectra with more absorption features similar to Type I SNe. Some Type I SNe in this luminosity range have relatively slow photospheric velocities than those of Type I SLSNe and Type I SNe that may indicate an existence of interesting intermediate population in this luminosity range (e.g., Blanchard et al., 2019).

## 3.2 Type IIa and Type II SLSNe

While more than 100 hydrogen-poor (Type I) SLSNe have been found so far, the number of observed hydrogen-rich SLSNe is still far less (of the order of 10). Therefore, the spectroscopic features that characterize hydrogen-rich SLSNe have not been carefully investigated, and hydrogen-rich SLSNe are still classified based on their peak luminosities. Following the luminosity range of Type I SLSNe, SNe with hydrogen features having the peak magnitude brighter than around  $-20$  mag in optical are often referred as hydrogen-rich (Type IIa or Type II) SLSNe.

Most hydrogen-rich SLSNe show narrow hydrogen lines in their spectra and they are called Type IIa SNe. The prototype and the most studied SN of this class is SN 2006gy (e.g., Smith et al., 2010). The luminosity and luminosity evolution of Type IIa SNe are diverse. The most luminous Type IIa SLSNe reaches to around  $-22.5$  mag in optical (e.g., SN 2008fz, Drake et al. 2010; SN 2016aps, Nicholl et



**Fig. 5** Summary of the estimated event rates of SLSNe. The fractions to core-collapse SNe (CCSNe) are based on the core-collapse SN rate estimated by Strolger et al. (2015).

al. 2020) and these bright Type II<sub>n</sub> SLSNe tend to have round light-curve shapes around the luminosity peak. As the peak luminosity of Type II<sub>n</sub> SLSNe becomes smaller, their light-curve evolution tends to decline with a power law (e.g., SN 2010jl, Fransson et al. 2014). However, some Type II<sub>n</sub> SLSNe are known to evolve quite fast (e.g., SN 2003ma, Rest et al. 2011), while others evolve very slowly (e.g., SN 2015da, Tartaglia et al. 2020) regardless of their high luminosity. It is not clear if there is a separate population of Type II<sub>n</sub> SLSNe or they consist of the most luminous end of a continuous Type II<sub>n</sub> SN population (Richardson et al., 2014). Some Type II<sub>n</sub> SLSNe emit more than  $5 \times 10^{51}$  erg and the explosion inside such events are clearly distinct from other Type II<sub>n</sub> SNe (Nicholl et al., 2020).

Some very luminous transients having spectra that are consistent with Type II<sub>n</sub> SNe appear near the center of AGNs (Drake et al., 2011; Kankare et al., 2017). Because spectra of some AGNs and Type II<sub>n</sub> SNe are known to have similar narrow emission features, it is sometimes difficult to distinguish if they are Type II<sub>n</sub> SLSNe originating from stellar explosions or certain activities of AGNs (e.g., Moriya et al., 2017b). Some tidal disruption events may also be confused with Type II<sub>n</sub> SLSNe (Blanchard et al., 2017; Petrushevska et al., 2023). We note that SN 2006gy appeared in an X-ray bright galaxy (NGC 1260, Wang et al. 2016) which might be an AGN and it may have been identified as a nuclear transient if it appeared at high redshifts.

Type II<sub>n</sub> SLSNe have been often observed in infrared wavelengths and they are found to have observational features of dusts (Sun et al., 2022). For example, SN 2006gy was bright in infrared for a long time (Miller et al., 2010). SN 2010jl is also well observed in infrared with dust signatures (Maeda et al., 2013).

Not all hydrogen-rich SLSNe are Type II<sub>n</sub> SLSNe. Several hydrogen-rich SLSNe without narrow hydrogen emission lines are known and they are referred as Type II SLSNe (Inserra et al., 2018b; Kangas et al., 2022). SN 2008es is the first SLSN that was identified as a Type II SLSN (Gezari et al., 2009; Miller et al., 2009). The number of observed Type II SLSNe is still small (Inserra et al., 2018b; Kangas et al., 2022) and the characteristic properties of Type II SLSNe that distinguish them from less luminous Type II SNe are still not fully understood.

Host environments of Type II<sub>n</sub> SLSNe are more diverse than those of Type I SLSNe. They can appear in higher metallicity environments than Type I SLSNe (Neill et al., 2011). Their host galaxies can have higher metallicities and larger stellar masses than Type I SLSNe as well. They might tend to originate from lower metallicity and lower luminosity galaxies than typical core-collapse SNe (Schulze et al., 2021). The host environments of Type II SLSNe are found to be not so different from those of Type II<sub>n</sub> SLSNe (Kangas et al., 2022), but the number of Type II SLSNe is still too small to make a proper comparison. Some studies suggest that Type II SLSN environments are similar to those of Type I SLSNe (Inserra et al., 2018b).

## 4 Event rates

The current event rate estimates for SLSNe are summarized in Figure 5. The event rates of SLSNe were first studied by Quimby et al. (2013). Based on SLSNe discovered by ROTSE-IIIb, they estimated the Type I SLSN rate at  $z \approx 0.17$  as  $32_{-26}^{+77} h_{71}^3 \text{ Gpc}^{-3} \text{ yr}^{-1}$  and the hydrogen-rich SLSN rate as  $151_{-82}^{+151} h_{71}^3 \text{ Gpc}^{-3} \text{ yr}^{-1}$  at  $z \approx 0.15$ . The hydrogen-rich SLSN rate from Quimby et al. (2013) includes both Type II<sub>n</sub> and Type II SLSNe discovered by ROTSE-IIIb. McCrum et al. (2015) estimated the Type I SLSN rate at  $0.3 \lesssim z \lesssim 1.4$  as  $(3 - 8) \times 10^{-3}\%$

## 8 Superluminous supernovae

of the core-collapse SN rate. Prajs et al. (2017) estimated the Type I SLSN rate at  $z \approx 1.13$  as  $91_{-36}^{+76} h_{70}^3 \text{ Gpc}^{-3} \text{ yr}^{-1}$  based on the Type I SLSN sample from Supernova Legacy Survey. Frohmaier et al. (2021) estimated Type I SLSN rate at  $z \lesssim 0.2$  as  $35_{-13}^{+25} h_{70}^3 \text{ Gpc}^{-3} \text{ yr}^{-1}$  based on the PTF SLSN sample. Using the public Type I SLSN data from PS1, Zhao et al. (2021) estimated the Type I SLSN event rate of  $40 h_{70}^3 \text{ Gpc}^{-3} \text{ yr}^{-1}$  at  $z \approx 0.89$  with an unknown error. No updated event rates for Type IIn and Type II SLSNe have been obtained after Quimby et al. (2013).

There are some estimates for the total event rates of SLSNe based on photometric samples of SLSNe. For example, based on the fact that no SLSNe were discovered during the Supernova Diversity and Rate Evolution (SUDARE) survey, the total SLSN rate was constrained to be less than  $900 h_{70}^3 \text{ Gpc}^{-3} \text{ yr}^{-1}$  at  $z \approx 0.5$  (Cappellaro et al., 2015). Cooke et al. (2012) estimated the SLSN rate at  $z \approx 2 - 4$  to be  $\sim 400 h_{71}^3 \text{ Gpc}^{-3} \text{ yr}^{-1}$  using two SLSNe at  $z = 2.05$  and  $3.90$  discovered by the photometric SLSN search in the CFHT archival data. The high-redshift SN survey with Subaru/Hyper Suprime-Cam (HSC) provided the SLSN event rates of  $\sim 900_{-500}^{+900} h_{70}^3 \text{ Gpc}^{-3} \text{ yr}^{-1}$  at  $z \approx 2$ ,  $\sim 400_{-300}^{+900} h_{70}^3 \text{ Gpc}^{-3} \text{ yr}^{-1}$  at  $z \approx 3$ , and  $\sim 500_{-400}^{+1200} h_{70}^3 \text{ Gpc}^{-3} \text{ yr}^{-1}$  at  $z \approx 4$  (Moriya et al., 2019b; Curtin et al., 2019).

## 5 Luminosity sources and progenitors

We provide a brief overview of possible luminosity sources that make SLSNe so bright. For more details of the major proposed luminosity sources, we refer to Moriya et al. (2018b). For each possible luminosity source, we also discuss possible progenitors that can realize the conditions required to have the luminosity source.

### 5.1 Radioactive decay

The energy released by the radioactive decay of  $^{56}\text{Ni}$  synthesized during the SN explosion is a standard powering mechanism of SNe (Arnett, 1996). Especially, the early luminosity peak of stripped-envelope SNe is mainly powered by the  $^{56}\text{Ni}$  decay (Bersten and Mazzali, 2017). A simple approach extending the standard SN powering mechanism to SLSNe is to have a larger amount of  $^{56}\text{Ni}$  from the explosive nucleosynthesis. The more  $^{56}\text{Ni}$  is synthesized, the more radioactive energy is available to make SNe brighter. The synthesized  $^{56}\text{Ni}$  decays to  $^{56}\text{Co}$  with a decay time of  $8.76 \pm 0.01$  days (da Cruz et al., 1992) and then  $^{56}\text{Co}$  decays to  $^{56}\text{Fe}$  with a decay time of  $111.42 \pm 0.04$  days (Funck et al., 1992). The total available energy from this radioactive decay is

$$L_{56\text{Ni-decay}} = \left[ 6.48 \exp\left(-\frac{t}{8.76 \text{ days}}\right) + 1.44 \exp\left(-\frac{t}{111.42 \text{ days}}\right) \right] \frac{M_{56\text{Ni}}}{M_{\odot}} 10^{43} \text{ erg s}^{-1}, \quad (1)$$

where  $M_{56\text{Ni}}$  is the mass of  $^{56}\text{Ni}$  synthesized at the explosion (Nadyozhin 1994 with the updated physical values from <http://www.nndc.bnl.gov/chart>). The energy from the nuclear decay is mostly released as gamma-rays having the energy of the order of MeV. Because these gamma-rays may not necessarily be absorbed by the SN ejecta, the actual available energy to power the SN luminosity is less than the total decay energy in Equation (1).


In order to have a rough estimate for the amount of  $^{56}\text{Ni}$  that is required to account for the luminosity of SLSNe, we can use the rise time and peak bolometric luminosity of SLSNe. Arnett (1982) analytically showed that the peak luminosity of a SN is the same as the luminosity input at the time of the peak luminosity (so-called "Arnett-law"). If we take the average rise time of 41.9 days for Type I SLSNe, for example, the  $^{56}\text{Ni}$  mass required to explain their peak luminosities above  $3 \times 10^{43} \text{ erg s}^{-1}$  is  $M_{56\text{Ni}} \gtrsim 3 M_{\odot}$  (Equation 1). This  $^{56}\text{Ni}$  mass is much higher than the  $^{56}\text{Ni}$  mass estimated for typical SNe ( $M_{56\text{Ni}} \lesssim 0.3 M_{\odot}$ , e.g., Anderson 2019). Even broad-lined Type Ic SNe that are among the most energetic core-collapse SNe are mostly estimated to have  $M_{56\text{Ni}} \lesssim 1 M_{\odot}$  (Taddia et al., 2019).

Synthesizing  $M_{56\text{Ni}} \gtrsim 3 M_{\odot}$  is a challenge in the standard core-collapse SN explosion models. The maximum amount of  $^{56}\text{Ni}$  that can be synthesized by core-collapse SNe is estimated to be around  $10 M_{\odot}$ , which requires the explosion energy of  $10^{53} \text{ erg}$  (Umeda and Nomoto, 2008). Some relatively low-luminosity SLSNe that require  $M_{56\text{Ni}} \lesssim 10 M_{\odot}$  could be consistent with such energetic core-collapse SNe (Moriya et al., 2019a; Mazzali et al., 2019). However, a significant fraction of SLSNe require  $M_{56\text{Ni}} \gtrsim 10 M_{\odot}$  to explain their peak luminosity. However, pair-instability SNe (PISNe, Barkat et al. 1967; Rakavy and Shaviv 1967) are predicted to synthesize up to around  $70 M_{\odot}$  of  $^{56}\text{Ni}$  (Heger and Woosley, 2002). PISNe are predicted explosions of very massive stars with the helium core mass between  $\sim 65 M_{\odot}$  and  $\sim 135 M_{\odot}$ , although the exact mass range depends on uncertainties in, e.g., nuclear reaction rates (Takahashi et al., 2018; Farmer et al., 2019, 2020). This helium core mass corresponds to the zero-age main sequence (ZAMS) mass between  $\sim 140 M_{\odot}$  and  $\sim 260 M_{\odot}$  when mass loss and rotation are ignored (Heger and Woosley, 2002). If the rotation is significant, the ZAMS mass of PISN progenitors can be as low as  $65 M_{\odot}$  through chemically homogeneous evolution (e.g., Chatzopoulos and Wheeler, 2012). Stellar mergers are also an important path to form massive stars ending up with PISNe (Vigna-Gómez et al., 2019).

Because it is required to sustain massive cores until the instability is triggered, PISNe are not expected to occur frequently above a certain metallicity because of strong mass loss at high metallicity (Langer et al., 2007). This is consistent with the fact that Type I SLSNe prefer low metallicity environments. However, the exact metallicity cut is uncertain (Sabharwal et al., 2023), and it is also possible to suppress mass loss in high metallicity environments (Georgy et al., 2017). The predicted PISN event rates in the local Universe are also lower than the event rates of Type I SLSNe (Briel et al., 2022).

Predicted PISN light curves evolve slowly (e.g., Kasen et al., 2011; Dessart et al., 2013) and they are often consistent with those of slowly evolving SLSNe. In many rapidly evolving SLSNe, the required amount of  $^{56}\text{Ni}$  to explain their luminosity becomes more than the estimated ejecta mass (Kasen, 2017). Therefore, rapidly evolving SLSNe are generally not likely powered by the radioactive decay of  $^{56}\text{Ni}$ , although mixing of synthesized  $^{56}\text{Ni}$  in the ejecta may make luminosity evolution of PISNe faster (Kozyreva and Blinnikov, 2015).





**Fig. 6** Predicted rise times and peak luminosities of interaction-powered SNe and their comparisons with FBOTs and Type IIIn SNe including Type IIIn SLSNe. Reproduced with permission from Suzuki et al. (2020).

However, no significant mixing is predicted in multi-dimensional simulations of PISNe (e.g., Joggerst and Whalen, 2011; Chen et al., 2014). The nuclear decay of  $^{56}\text{Ni}$  as a possible power source of SLSNe is mainly discussed for slowly evolving SLSNe whose light-curve decline rates are consistent with the decay rate of  $^{56}\text{Ni} \rightarrow ^{56}\text{Co} \rightarrow ^{56}\text{Fe}$ . Even if the light-curve evolution is consistent with PISNe, their spectral features are often found to be inconsistent with those predicted by PISNe (Dessart et al., 2012, 2013; Jerkstrand et al., 2016, 2017). Especially, a large amount of  $^{56}\text{Fe}$  is expected to exist in late phases from the nuclear decay of  $^{56}\text{Ni}$ , but no strong  $^{56}\text{Fe}$  absorption and emission are usually observed in SLSNe. Currently, only SN 2018ibb is found to match most of the predicted PISN properties, but their late-phase spectra have blue flux excess which might be caused by the additional CSM interaction (Schulze et al., 2024).

It is difficult to explain the precursors and light-curve bumps and undulations occurring in different timescales solely by the nuclear decay energy input that is govern by the nuclear decay timescale. Thus, additional luminosity source such as CSM interaction is required to explain the whole luminosity evolution of SLSNe.

## 5.2 Circumstellar interaction

The collision of SN ejecta into their CSM can efficiently convert the kinetic energy of the ejecta to radiation especially when the CSM has a comparable mass to the SN ejecta. When the CSM density is high enough, the collision forms a strong radiative shock. The emission is first mostly in X-rays but they can be immediately absorbed through free-free absorption when the CSM density is high. Then the post-shock region can be around  $10^4$  K to emit photons in optical (e.g., Moriya et al., 2013). The unshocked CSM can be optically thick and photons will diffuse out in the dense CSM. This diffusion process can make light curves of interaction-powered SNe broad. The diffusion time can vary depending on the CSM density and radius. Therefore, the CSM interaction model can explain SLSNe of various timescales.

Type IIIn SNe show clear signatures of the CSM interaction in their spectra. Thus, Type IIIn SLSNe are considered to be powered by the CSM interaction as in the case of the lower luminosity Type IIIn SNe (e.g., Moriya et al., 2013). Compared to low-luminosity Type IIIn SNe, Type IIIn SLSNe are estimated to have higher explosion energies or more massive CSM because of their higher luminosity. The total radiated energy of Type IIIn SLSNe is of the order of  $10^{51}$  erg, which is a typical SN explosion energy. Thus, if we can efficiently convert the kinetic energy of SN ejecta to radiation through the CSM interaction, it is possible to explain the huge luminosities observed in Type IIIn SLSNe. The total CSM mass around Type IIIn SLSNe are estimated to be  $\gtrsim 5 M_{\odot}$  (Chatzopoulos et al. 2013; Suzuki et al. 2020, Figure 6). We note that the most luminous Type IIIn SLSNe emit more than  $10^{52}$  erg in total and the inner explosion sometimes needs to be energetic (Nicholl et al., 2020; Suzuki et al., 2021).

Because observational signatures of Type IIIn SLSNe are dominated by the CSM interaction, it is difficult to identify the nature of the explosions inside. The progenitors of Type IIIn SNe are known to be massive ( $\gtrsim 50 M_{\odot}$ ) luminous blue variables (LBVs, Gal-Yam and Leonard 2009) and low-mass ( $\approx 10 M_{\odot}$ ) red supergiants (Prieto et al., 2008). Among them, some mass eruptions from LBVs such as the Great Eruption of  $\eta$  Carinae (Humphreys and Davidson, 1994) are known to form CSM as massive as  $10 M_{\odot}$  (Morris et al., 1999), although their mass-loss mechanisms are not well understood. If LBVs explode immediately after they experience such an eruptive mass loss, they could be observed as Type IIIn SLSNe. LBVs are considered to be massive stars with ZAMS masses above around  $40 M_{\odot}$  (Humphreys and

Davidson, 1994), but they may also originate from stellar mergers of less massive stars (Justham et al., 2014). Even if massive stars are not in the LBV phase, they may experience strong mass loss triggered by the strong convection at the innermost layers of massive stars (Quataert and Shiode 2012; Mcley and Soker 2014, but see also Fuller 2017). Phase transition from nuclear matter to the quark-gluon plasma at the center of massive stars after the core collapse is also suggested to result in Type IIn SLSNe (Fischer et al., 2018).

Another mechanism to form massive CSM is common-envelope mass ejection (Chevalier, 2012; Ercolino et al., 2024). Unstable mass transfer in binary systems with massive stars can lead to a common-envelope phase. Although the exact outcome of the common-envelope phase is uncertain, one possible consequence is ejection of the massive stellar envelope. If the common-envelope mass ejection occurs shortly before the explosion of massive stars, the SN ejecta can collide with the massive ejected envelope and can be observed as Type IIn SLSNe.

Extensive mass loss from massive stars can also be related to pulsational pair-instability (Woosley et al., 2007). Massive stars slightly below the mass range of PISNe (Section 5.1) can still become dynamically unstable to eject a part of their mass to form a massive CSM. Several mass ejection triggered by this instability can occur sequentially and ejected shells can collide to each other to make Type IIn SLSNe (Woosley et al., 2007). This kind of transients triggered by the pulsational pair-instability is called pulsational pair-instability SNe (PPISNe). It is also possible that a core-collapse SN is followed by the pulsational pair-instability. In this case, the SN ejecta can collide to the dense CSM formed by the pulsational pair-instability to become Type IIn SLSNe. As in the case of PISN progenitors, PPISN progenitors can originate from stellar mergers (Vigna-Gómez et al., 2019).

It has also been suggested that some Type IIn SLSNe are related to explosions of white dwarfs (Type Ia SNe). The late-phase spectrum of SN 2006gy was suggested to be similar to that of Type Ia SN and its luminosity evolution is also suggested to be consistent with the explosion of a Type Ia SN within the hydrogen-rich CSM having the mass of  $\approx 10 M_{\odot}$  (Jerkstrand et al., 2020). A major question for this scenario is how to form such a massive hydrogen-rich CSM around Type Ia SN progenitors. Some Type Ia SNe are known to show signatures of hydrogen-rich CSM (so-called Type Ia-CSM), but their CSM mass is estimated to be of the order of  $0.1 M_{\odot}$  (Sharma et al., 2023). A rare evolutionary path involving common-envelope mass ejection shortly before Type Ia SN explosions may be able to realize such a massive CSM around Type Ia SNe (Ablimit, 2021).

We have discussed Type IIn SLSNe so far. Although the other types of SLSNe do not show clear signatures of the CSM interaction in their spectra in early phases, the CSM interaction has been considered to be their possible luminosity source. For example, the light curves of Type I SLSNe can be reproduced by the interaction between SN ejecta and massive hydrogen-poor CSM (Sorokina et al., 2016). The precursor bump observed in Type I SLSNe may also be explained by the existence of a massive CSM (Moriya and Maeda, 2012). Such a massive hydrogen-poor CSM can be formed by the pulsational pair-instability of massive hydrogen-poor stars (Woosley, 2017) and such massive hydrogen-poor stars can be formed through the chemically homogeneous evolution, for example (Marchant et al., 2016; Mandel and de Mink, 2016). Several Type I SLSNe have been suggested to be hydrogen-poor PPISNe (e.g., Tolstov et al., 2017). If Type I SLSNe are mainly powered by the CSM interaction, a major remaining question is why we do not see clear interaction signatures in their spectra. There are hydrogen-poor SNe that show clear CSM interaction signatures in their spectra known as Type Ibn (e.g., Pastorello et al., 2007; Hosseinzadeh et al., 2017) and Type Icn SNe (e.g., Fraser et al., 2021; Gal-Yam et al., 2022; Perley et al., 2022). Thus, we may expect to have similar spectroscopic signatures if Type I SLSNe are mainly powered by the CSM interaction, although the required CSM mass and density for Type I SLSNe are expected to be higher. More theoretical investigations on the expected spectroscopic features of hydrogen-poor interaction-powered SNe are required.


Even if the major luminosity source of Type I and Type II SLSNe is not the CSM interaction, it is very likely that their properties are often partially affected by the CSM interaction. The luminosity undulations observed in Type I SLSNe can be explained as an additional effect caused by the CSM interaction. Similarly multiple luminosity bumps in Type I SLSNe could be related to the existence of the multiple dense CSM components (Lin et al., 2023). Because many Type I SLSNe show the CSM interaction signatures at late phases, it is possible that some effects of the CSM interaction starts to appear in earlier epochs after the major luminosity source ends providing their energy.

### 5.3 Spin down of strongly magnetized neutron stars (“magnetars”)

A neutron star may remain at the center of an exploding massive star after a core-collapse SN explosion. If the neutron star has rotation and poloidal magnetic fields (in other word, if the neutron star becomes a “pulsar”), the rotational energy should be converted to the electromagnetic field mostly as Poynting-flux dominated outflows (e.g., Goldreich and Julian, 1969). The total spin-down energy that can be released in this form can be expressed as

$$L_{\text{spin-down}} = \frac{(l-1)E_p}{t_p} \left(1 + \frac{t}{t_p}\right)^{-l}, \quad (2)$$

where  $E_p$  is the initial rotational energy of the neutron star and  $t_p$  is the spin-down timescale. The temporal index  $l$  is determined by the braking index and it is often assumed to be  $l = 2$ , i.e., the magnetar spin-down is dominated by dipole radiation. A fraction of this spin-down energy can be thermalized to power the SN luminosity. The idea that pulsars may be able to power SN explosions and SN luminosity appeared shortly after the discovery of pulsars (Ostriker and Gunn, 1971). However, the pulsar-powering mechanism for typical core-collapse SNe was not found to match their observations. Later, Maeda et al. (2007) proposed that the pulsar power can be an extra energy source to illuminate SNe to explain a peculiar light-curve behavior of SN 2005bf (Anupama et al., 2005; Folatelli et al., 2006). Kasen and Bildsten (2010); Woosley (2010) applied this idea of powering SNe with pulsars for SLSNe. They demonstrated that if the pulsar has an initial rotational period of  $\sim 1$  ms with a dipole magnetic field of  $\sim 10^{14}$  G, it can reproduce the light curves of Type I SLSNe. Such a strongly-magnetized, rapidly-rotating pulsar powering SLSNe are often referred to as “magnetars.”




**Fig. 7** Estimated parameters for Type I SLSNe in the magnetar-powered model. Reproduced with permission from Chen et al. (2023b).

This magnetar scenario to power SLSN luminosity is currently the most popular scenario to explain SLSNe without clear CSM interaction signatures. Many studies applied the magnetar model to fit the light-curve evolution of Type I SLSNe to constrain the properties of magnetars powering Type I SLSNe (e.g., Inserra et al., 2013; Metzger et al., 2015; Kashiyama et al., 2016). The magnetar model can explain early-phase light curves of both slowly-evolving and rapidly-evolving Type I SLSNe well. The systematic model fitting to the magnetar-powered model for the Type I SLSN sample from ZTF obtained the average initial spin period of  $2.65^{+2.58}_{-0.68}$  ms and the dipole magnetic field strength of  $0.98^{+0.98}_{-0.63} \times 10^{14}$  G with the  $1\sigma$  range (Chen et al., 2023b). This fitting provides the average ejecta mass of  $5.03^{+4.01}_{-2.39} M_{\odot}$  and the average kinetic energy of  $2.13^{+1.89}_{-0.96} \times 10^{51}$  erg with the  $1\sigma$  range (Chen et al., 2023b). The distributions of the estimated parameters and their correlation can be found in Figure 7. Under the assumption of the magnetar spin-down model, a possible correlation between SN ejecta mass and initial spin periods is found (Blanchard et al., 2020).

The observed complexity in the light curves of Type I SLSNe is also suggested to be explained by the magnetar scenario. The precursor bump can be explained by the additional shock breakout of a strong shock formed by the central energy injection at the center (Kasen et al., 2016; Liu et al., 2021). The late-phase light curves powered by magnetar spin down are strongly affected by the uncertain thermalization efficiencies of the spin-down energy (Kotera et al., 2013; Wang et al., 2015). In order to distinguish magnetar spin-down and other energy source such as the  $^{56}\text{Ni}$  radioactive decay, light curves need to be followed for around 1000 days (Moriya et al., 2017a), but SLSNe with such a long-term observation is still limited (Blanchard et al., 2017). The late-phase light curve undulations can be related to temporal changes in magnetar activities (e.g., Yu and Li, 2017; Moriya et al., 2022b).

Spectroscopic properties are also found to be consistent with the magnetar model (e.g., Jerkstrand et al., 2017; Dessart, 2019). Especially, the brightness in ultraviolet in early phases as well as slow evolution in the photospheric velocity is consistent with the predicted properties from the magnetar powered models. Aspherical nature of Type I SLSNe observed in some Type I SLSNe can also be explained by the magnetar scenario because the spin-down energy injection inevitably occur in an aspherical form. Jet emergence from magnetars is also expected (e.g., Soker, 2022).



**Fig. 8** Ejecta mass and total accreted mass required to account for Type I SLSN light curves in the fallback accretion model. The energy conversion efficiency of  $\epsilon_{\text{acc}} = 10^{-3}$  is assumed. Modified from Moriya et al. (2018a).

The magnetar energy injection at the central region of SN ejecta is predicted to result in several interesting characteristic features in SLSNe. For example, the extra strong energy input at the innermost layers of the SN ejecta can lead to strong mixing within the ejecta and make the ejecta density structure flatter than other SNe (e.g., Chen et al., 2016; Suzuki and Maeda, 2017; Desai et al., 2023). Such a flat density structure may affect spectral formation in SLSNe, although their consequences have not been studied in detail. Another characteristic prediction is that high energy and X-ray emissions can be observed at late phases after the central nebular regions formed by the spin down of the central magnetar become transparent (e.g., Metzger et al., 2014; Murase et al., 2015). In addition, magnetar-powered SLSNe can become bright in radio because of the pulsar wind nebula formed by the central magnetar. However, X-ray and radio emissions from Type I SLSNe are often not consistent with simple predictions (e.g., Margutti et al., 2018; Murase et al., 2021). The thermalization process in the magnetar spin down is not understood well and more investigations are required to predict their expected observational properties (Margalit et al., 2018; Vurm and Metzger, 2021). Possible gamma-ray emission from SN 2017egm is suggested to be consistent with the magnetar spin-down model (Li et al., 2024). If the central magnetar is massive enough to require rotational energy to sustain its mass, the magnetar would eventually collapse to a black hole at some moment and leave some observational consequences (Metzger et al., 2015; Moriya et al., 2016). Finally, FRBs might be associated with SLSNe if they are both powered by magnetars (Metzger et al., 2017), although host galaxy properties of SLSNe and FRBs are different (Section 3.1.9).

The progenitors of magnetar-powered SLSNe should have rapid rotation. Rapidly rotating Type I SLSN progenitors could be realized through chemically homogeneous evolution (Aguilera-Dena et al., 2018, 2020) possibly in binary systems. The strong magnetic field may exist at the time of core collapse or amplified during the core collapse. Stellar mergers may also be responsible for the magnetic field amplification (Ablimit et al., 2022). We note that some Type I SLSNe showing late-phase CSM interaction features require an additional CSM component ejected from the progenitor in addition to the rapid rotation and strong magnetic field. Some Type I SLSNe show late-phase hydrogen-rich emission that requires hydrogen-rich mass loss from the progenitors shortly before their explosions.

#### 5.4 Black hole accretion

A black hole, instead of a neutron star, can be formed after the terminal collapse of a massive star. Accretion towards the black hole can launch a jet or a disk wind outflow that can be a luminosity source of SNe. If a massive star collapses directly to the black hole with a free-fall timescale, the accretion timescale would be too short to power SLSNe. However, if the outer layers of the SN ejecta are first ejected and then fall back onto the black hole, a long-lasting mass accretion towards the central black hole can be realized. Such a fallback accretion is suggested to be a potential power source for SLSNe (Dexter and Kasen, 2013). In a simplified picture, the luminosity input from the accretion can be expressed as

$$L_{\text{accretion}} = \epsilon_{\text{acc}} \dot{M}_{\text{acc}} c^2, \quad (3)$$

where  $\epsilon_{\text{acc}}$  is the thermalization efficiency of the accretion and  $\dot{M}_{\text{acc}}$  is the accretion rate to the black hole. The thermalization efficiency is quite uncertain, but it can be reasonably assumed to be  $\epsilon_{\text{acc}} \sim 10^{-3}$  (Dexter and Kasen, 2013). When the fallback accretion is dominant,  $\dot{M}_{\text{acc}}$  becomes proportional to  $t^{-3/5}$ . Moriya et al. (2018a) systematically investigated the properties of the fallback accretion required to account for Type I SLSNe. Although their light curves can be fitted well by the fallback accretion scenario, the required mass to accrete

onto the black hole is often found to be very massive (Figure 8).

## 5.5 Other proposed mechanisms

It is still possible that some unrecognized mechanisms lead to at least some fraction of the observed SLSN populations. For example, axion-instability SNe might be related to SLSNe (Sakstein et al., 2022; Mori et al., 2023). A latent energy released by the phase transition from neutron stars to quark stars is suggested to power SLSNe (e.g., Ouyed et al., 2012). Other unconsidered energy sources may also play an important role.

## 6 Outlook

As we face the era of large-scale transient surveys in various wavelengths, the number of SLSN discovery is still expected to increase. For example, the Vera C. Rubin Observatory’s Legacy Survey of Space and Time (LSST) is expected to discover  $\sim 10^4$  SLSNe in a year, although only a limited fraction among them will have sufficient light-curve information to reconstruct their physical properties (Villar et al., 2018; du Buisson et al., 2020).

Unexplored frontier in SLSN discoveries are at high redshifts. Only a couple of SLSNe have been observed at  $z \gtrsim 2$  so far (Cooke et al., 2012; Pan et al., 2017; Smith et al., 2018; Moriya et al., 2019b; Curtin et al., 2019). SLSNe at high redshifts will allow us to explore SLSN properties in, e.g., low metallicity environments that will be essential information to uncover their progenitors. In addition, the event rates of PISNe are predicted to be higher at higher redshifts (Briel et al., 2022) and we expect to discover many PISNe if we search for them at high redshifts. Exploring high-redshift SLSNe require near-infrared transient surveys. Fortunately, several wide-field, sensitive near-infrared imaging instruments will be available in the coming years and they will allow us to explore SLSNe at high redshifts (Tanaka et al., 2012, 2013). For example, Euclid has started its operation with successful SN discoveries (Euclid Collaboration et al., 2024). Euclid is expected to discover dozens of SLSNe and PISNe up to  $z \sim 4$  (Inserra et al., 2018a; Moriya et al., 2022a). Nancy Grace Roman Space Telescope, which is currently planned to be launched in 2026, can realize time domain surveys that allow us to discover SLSNe beyond  $z \sim 6$  (Moriya et al., 2022c). While the field-of-view is small, James Webb Space Telescope may also potentially discover high-redshift SLSNe (Regős and Vinkó, 2019; Venditti et al., 2024). SLSNe at high redshifts may be used as distance measurement (e.g., Inserra et al., 2021; Wei et al., 2015; Khetan et al., 2023) as well as light sources to explore interstellar media in the distant galaxies that hosted SLSNe (e.g., Berger et al., 2012).

## Acknowledgments

TJM is supported by the Grants-in-Aid for Scientific Research of the Japan Society for the Promotion of Science (JP24K00682, JP24H01824, JP21H04997, JP24H00002, JP24H00027, JP24K00668) and the Australian Research Council (ARC) through the ARC’s Discovery Projects funding scheme (project DP240101786).

**See Also:** Gal-Yam (2012, 2019b); Howell (2017); Moriya et al. (2018b); Inserra (2019); Nicholl (2021)

## References

- Ablimit I (2021), Jul. The CO White Dwarf + Intermediate-mass/Massive Star Binary Evolution: Possible Merger Origins for Peculiar Type Ia and II Supernovae. *PASP* 133 (1025), 074201. doi:10.1088/1538-3873/ac025c. 2101.03670.
- Ablimit I, Podsiadlowski P, Hirai R and Wicker J (2022), Jul. Stellar core-merger-induced collapse: new formation pathways for black holes, Thorne-Żytkow objects, magnetars, and superluminous supernovae. *MNRAS* 513 (4): 4802–4813. doi:10.1093/mnras/stac631. 2108.08430.
- Aguilera-Dena DR, Langer N, Moriya TJ and Schootemeijer A (2018), May. Related Progenitor Models for Long-duration Gamma-Ray Bursts and Type Ic Superluminous Supernovae. *ApJ* 858 (2), 115. doi:10.3847/1538-4357/aabfc1. 1804.07317.
- Aguilera-Dena DR, Langer N, Antoniadis J and Müller B (2020), Oct. Precollapse Properties of Superluminous Supernovae and Long Gamma-Ray Burst Progenitor Models. *ApJ* 901 (2), 114. doi:10.3847/1538-4357/abb138. 2008.09132.
- Anderson JP (2019), Aug. A meta-analysis of core-collapse supernova  $^{56}\text{Ni}$  masses. *A&A* 628, A7. doi:10.1051/0004-6361/201935027. 1906.00761.
- Anderson JP, Pessi PJ, Dessart L, Inserra C, Hiramatsu D, Taggart K, Smartt SJ, Leloudas G, Chen TW, Möller A, Roy R, Schulze S, Perley D, Selsing J, Prentice SJ, Gal-Yam A, Angus CR, Arcavi I, Ashall C, Bulla M, Bray C, Burke J, Callis E, Cartier R, Chang SW, Chambers K, Clark P, Denneau L, Dennefeld M, Flewelling H, Fraser M, Galbany L, Gromadzki M, Gutiérrez CP, Heinze A, Hosseinzadeh G, Howell DA, Hsiao EY, Kankare E, Kostrzewa-Rutkowska Z, Magnier E, Maguire K, Mazzali P, McBrien O, McCully C, Morrell N, Lowe TB, Onken CA, Onori F, Phillips MM, Rest A, Ridden-Harper R, Ruiter AJ, Sand DJ, Smith KW, Smith M, Stalder B, Stritzinger MD, Sullivan M, Tonry JL, Tucker BE, Valenti S, Wainscoat R, Waters CZ, Wolf C and Young D (2018), Nov. A nearby super-luminous supernova with a long pre-maximum & “plateau” and strong C II features. *A&A* 620, A67. doi:10.1051/0004-6361/201833725. 1806.10609.
- Angus CR, Levan AJ, Perley DA, Tanvir NR, Lyman JD, Stanway ER and Fruchter AS (2016), May. A Hubble Space Telescope survey of the host galaxies of Superluminous Supernovae. *MNRAS* 458 (1): 84–104. doi:10.1093/mnras/stw063. 1601.01874.
- Angus CR, Smith M, Sullivan M, Inserra C, Wiseman P, D’Andrea CB, Thomas BP, Nichol RC, Galbany L, Childress M, Asorey J, Brown PJ, Casas R, Castander FJ, Curtin C, Frohmaier C, Glazebrook K, Gruen D, Gutierrez C, Kessler R, Kim AG, Lidman C, Macaulay E, Nugent P, Pursiainen M, Sako M, Soares-Santos M, Thomas RC, Abbott TMC, Avila S, Bertin E, Brooks D, Buckley-Geer E, Burke DL, Carnero Rosell

- A, Carretero J, da Costa LN, De Vicente J, Desai S, Diehl HT, Doel P, Eifler TF, Flaughner B, Fosalba P, Frieman J, García-Bellido J, Gruendl RA, Gschwend J, Hartley WG, Hollowood DL, Honscheid K, Hoyle B, James DJ, Kuehn K, Kuropatkin N, Lahav O, Lima M, Maia MAG, March M, Marshall JL, Menanteau F, Miller CJ, Miquel R, Ogando RLC, Plazas AA, Romer AK, Sanchez E, Schindler R, Schubnell M, Sobreira F, Suchyta E, Swanson MEC, Tarle G, Thomas D, Tucker DL and DES Collaboration (2019), Aug. Superluminous supernovae from the Dark Energy Survey. *MNRAS* 487 (2): 2215–2241. doi:10.1093/mnras/stz1321. 1812. 04071.
- Anupama GC, Sahu DK, Deng J, Nomoto K, Tominaga N, Tanaka M, Mazzali PA and Prabhu TP (2005), Oct. The Peculiar Type Ib Supernova SN 2005bf: Explosion of a Massive He Star with a Thin Hydrogen Envelope? *ApJL* 631 (2): L125–L128. doi:10.1086/497336. astro-ph/0509625.
- Arabsalmani M, Roychowdhury S, Renaud F, Cormier D, Le Floc'h E, Emsellem E, Perley DA, Zwaan MA, Bournaud F, Arumugam V and Møller P (2019), Sep. A Superluminous Supernova in High Surface Density Molecular Gas within the Bar of a Metal-rich Galaxy. *ApJ* 882 (1), 31. doi:10.3847/1538-4357/ab2897. 1906. 00972.
- Arnett WD (1982), Feb. Type I supernovae. I - Analytic solutions for the early part of the light curve. *ApJ* 253: 785–797. doi:10.1086/159681.
- Arnett D (1996). Supernovae and Nucleosynthesis: An Investigation of the History of Matter from the Big Bang to the Present.
- Barbary K, Dawson KS, Tokita K, Aldering G, Amanullah R, Connolly NV, Doi M, Faccioli L, Fadeyev V, Fruchter AS, Goldhaber G, Goobar A, Gude A, Huang X, Ihara Y, Konishi K, Kowalski M, Lidman C, Meyers J, Morokuma T, Nugent P, Perlmutter S, Rubin D, Schlegel D, Spadafora AL, Suzuki N, Swift HK, Takahashi N, Thomas RC and Yasuda N (2009), Jan. Discovery of an Unusual Optical Transient with the Hubble Space Telescope. *ApJ* 690 (2): 1358–1362. doi:10.1088/0004-637X/690/2/1358. 0809. 1648.
- Barkat Z, Rakavy G and Sack N (1967), Mar. Dynamics of Supernova Explosion Resulting from Pair Formation. *Phys. Rev. Lett.* 18 (10): 379–381. doi:10.1103/PhysRevLett.18.379.
- Berger E, Chornock R, Lunnan R, Foley R, Czekala I, Rest A, Leibler C, Soderberg AM, Roth K, Narayan G, Huber ME, Milisavljevic D, Sanders NE, Drout M, Margutti R, Kirshner RP, Marion GH, Challis PJ, Riess AG, Smartt SJ, Burgett WS, Hodapp KW, Heasley JN, Kaiser N, Kudritzki RP, Magnier EA, McCrum M, Price PA, Smith K, Tonry JL and Wainscoat RJ (2012), Aug. Ultraluminous Supernovae as a New Probe of the Interstellar Medium in Distant Galaxies. *ApJL* 755 (2), L29. doi:10.1088/2041-8205/755/2/L29. 1206. 4050.
- Bersten MC and Mazzali PA (2017), Light Curves of Type I Supernovae, Alsabti AW and Murdin P, (Eds.), *Handbook of Supernovae*, pp. 723.
- Bhandari S, Heintz KE, Aggarwal K, Marnoch L, Day CK, Sydnor J, Burke-Spolaor S, Law CJ, Xavier Prochaska J, Tejos N, Bannister KW, Butler B, Deller AT, Ekers RD, Flynn C, Fong Wf, James CW, Lazio TJW, Luo R, Mahony EK, Ryder SD, Sadler EM, Shannon RM, Han J, Lee K and Zhang B (2022), Feb. Characterizing the Fast Radio Burst Host Galaxy Population and its Connection to Transients in the Local and Extragalactic Universe. *AJ* 163 (2), 69. doi:10.3847/1538-3881/ac3aec. 2108. 01282.
- Blanchard PK, Nicholl M, Berger E, Guillochon J, Margutti R, Chornock R, Alexander KD, Leja J and Drout MR (2017), Jul. PS16dtm: A Tidal Disruption Event in a Narrow-line Seyfert 1 Galaxy. *ApJ* 843 (2), 106. doi:10.3847/1538-4357/aa77f7. 1703. 07816.
- Blanchard PK, Nicholl M, Berger E, Chornock R, Milisavljevic D, Margutti R and Gomez S (2019), Feb. A Hydrogen-poor Superluminous Supernova with Enhanced Iron-group Absorption: A New Link between SLSNe and Broad-lined Type Ic SNe. *ApJ* 872 (1), 90. doi:10.3847/1538-4357/aaf13. 1810. 11051.
- Blanchard PK, Berger E, Nicholl M and Villar VA (2020), Jul. The Pre-explosion Mass Distribution of Hydrogen-poor Superluminous Supernova Progenitors and New Evidence for a Mass-Spin Correlation. *ApJ* 897 (2), 114. doi:10.3847/1538-4357/ab9638. 2002. 09508.
- Blanchard PK, Berger E, Nicholl M, Chornock R, Gomez S and Hosseinzadeh G (2021), Nov. Late-time Hubble Space Telescope Observations of a Hydrogen-poor Superluminous Supernova Reveal the Power-law Decline of a Magnetar Central Engine. *ApJ* 921 (1), 64. doi:10.3847/1538-4357/ac1b27. 2105. 03475.
- Bose S, Dong S, Pastorello A, Filippenko AV, Kochanek CS, Mauerhan J, Romero-Cañizales C, Brink TG, Chen P, Prieto JL, Post R, Ashall C, Grupe D, Tomasella L, Benetti S, Shappee BJ, Stanek KZ, Cai Z, Falco E, Lundqvist P, Mattila S, Mutel R, Ochner P, Pooley D, Stritzinger MD, Villanueva S, J, Zheng W, Beswick RJ, Brown PJ, Cappellaro E, Davis S, Fraser M, de Jaeger T, Elias-Rosa N, Gall C, Gaudi BS, Herczeg GJ, Hestenes J, Holoien TWS, Hosseinzadeh G, Hsiao EY, Hu S, Jeejin S, Jeffers B, Koff RA, Kumar S, Kurtenkov A, Lau MW, Prentice S, Reynolds T, Rudy RJ, Shahbandeh M, Somero A, Stassun KG, Thompson TA, Valenti S, Woo JH and Yunus S (2018), Jan. Gaia17biu/SN 2017egm in NGC 3191: The Closest Hydrogen-poor Superluminous Supernova to Date Is in a “Normal,” Massive, Metal-rich Spiral Galaxy. *ApJ* 853 (1), 57. doi:10.3847/1538-4357/aaa298. 1708. 00864.
- Briel MM, Eldridge JJ, Stanway ER, Stevance HF and Chrimess AA (2022), Jul. Estimating transient rates from cosmological simulations and BPASS. *MNRAS* 514 (1): 1315–1334. doi:10.1093/mnras/stac1100. 2111. 08124.
- Cappellaro E, Botticella MT, Pignata G, Grado A, Greggio L, Limatola L, Vaccari M, Baruffolo A, Benetti S, Bufano F, Capaccioli M, Cascone E, Covone G, De Cicco D, Falocco S, Della Valle M, Jarvis M, Marchetti L, Napolitano NR, Paolillo M, Pastorello A, Radovich M, Schipani P, Spiro S, Tomasella L and Turatto M (2015), Dec. Supernova rates from the SUDARE VST-OmegaCAM search. I. Rates per unit volume. *A&A* 584, A62. doi:10.1051/0004-6361/201526712. 1509. 04496.
- Cartier R, Hamuy M, Contreras C, Anderson JP, Phillips MM, Morrell N, Stritzinger MD, Hueichapan ED, Clocchiatti A, Roth M, Thomas-Osip J and González LE (2022), Aug. A puzzle solved after two decades: SN 2002gh among the brightest of superluminous supernovae. *MNRAS* 514 (2): 2627–2649. doi:10.1093/mnras/stac1371. 2108. 09828.
- Chatzopoulos E and Wheeler JC (2012), Mar. Effects of Rotation on the Minimum Mass of Primordial Progenitors of Pair-instability Supernovae. *ApJ* 748 (1), 42. doi:10.1088/0004-637X/748/1/42. 1201. 1328.
- Chatzopoulos E, Wheeler JC, Vinko J, Horvath ZL and Nagy A (2013), Aug. Analytical Light Curve Models of Superluminous Supernovae:  $\chi^2$ -minimization of Parameter Fits. *ApJ* 773 (1), 76. doi:10.1088/0004-637X/773/1/76. 1306. 3447.
- Chen KJ, Heger A, Woosley S, Almgren A and Whalen DJ (2014), Sep. Pair Instability Supernovae of Very Massive Population III Stars. *ApJ* 792 (1), 44. doi:10.1088/0004-637X/792/1/44. 1402. 5960.
- Chen KJ, Woosley SE and Sukhbold T (2016), Nov. Magnetar-Powered Supernovae in Two Dimensions. I. Superluminous Supernovae. *ApJ* 832 (1), 73. doi:10.3847/0004-637X/832/1/73. 1604. 07989.
- Chen TW, Nicholl M, Smartt SJ, Mazzali PA, Yates RM, Moriya TJ, Inserra C, Langer N, Krühler T, Pan YC, Kotak R, Galbany L, Schady P, Wiseman P, Greiner J, Schulze S, Man AWS, Jerkstrand A, Smith KW, Dennefeld M, Baltay C, Bolmer J, Kankare E, Knust F, Maguire K, Rabinowitz D, Rostami S, Sullivan M and Young DR (2017a), Jun. The evolution of superluminous supernova LSQ14mo and its interacting host galaxy system. *A&A* 602, A9. doi:10.1051/0004-6361/201630163. 1611. 09910.
- Chen TW, Schady P, Xiao L, Eldridge JJ, Schweyer T, Lee CH, Yu PC, Smartt SJ and Inserra C (2017b), Nov. Spatially Resolved MaNGA Observations of the Host Galaxy of Superluminous Supernova 2017egm. *ApJL* 849 (1), L4. doi:10.3847/2041-8213/aa8f40. 1708. 04618.
- Chen TW, Smartt SJ, Yates RM, Nicholl M, Krühler T, Schady P, Dennefeld M and Inserra C (2017c), Sep. Superluminous supernova progenitors have a half-solar metallicity threshold. *MNRAS* 470 (3): 3566–3573. doi:10.1093/mnras/stx1428. 1605. 04925.
- Chen TW, Inserra C, Fraser M, Moriya TJ, Schady P, Schweyer T, Filippenko AV, Perley DA, Ruitter AJ, Seitzzahl I, Sollerman J, Taddia F, Anderson JP, Foley RJ, Jerkstrand A, Ngeow CC, Pan YC, Pastorello A, Points S, Smartt SJ, Smith KW, Taubenberger S, Wiseman P, Young DR, Benetti S, Berton M, Bufano F, Clark P, Della Valle M, Galbany L, Gal-Yam A, Gromadzki M, Gutiérrez CP, Heinze A, Kankare E, Kilpatrick CD, Kuncarayakti H, Leloudas G, Lin ZY, Maguire K, Mazzali P, McBrien O, Prentice SJ, Rau A, Rest A, Siebert MR, Stalder B, Tonry JL

- and Yu PC (2018), Nov. SN 2017ens: The Metamorphosis of a Luminous Broadlined Type Ic Supernova into an SN II<sub>n</sub>. *ApJL* 867 (2), L31. doi:10.3847/2041-8213/aab2e. 1808. 04382.
- Chen TW, Brennan SJ, Wesson R, Fraser M, Schweyer T, Inserra C, Schulze S, Nicholl M, Anderson JP, Hsiao EY, Jerkstrand A, Kankare E, Kool EC, Kravtsov T, Kuncarayakti H, Leloudas G, Li CJ, Matsuura M, Pursiainen M, Roy R, Ruiters AJ, Schady P, Seitzzahl I, Sollerman J, Tartaglia L, Wang L, Yates RM, Yang S, Baade D, Carini R, Gal-Yam A, Galbany L, Gonzalez-Gaitan S, Gromadzki M, Gutierrez CP, Kotak R, Maguire K, Mazzali PA, Mueller-Bravo TE, Paraskeva E, Pessi PJ, Pignata G, Rau A and Young DR (2021), Sep. SN 2018bsz: significant dust formation in a nearby superluminous supernova. *arXiv e-prints*, arXiv:2109.07942doi:10.48550/arXiv.2109.07942. 2109. 07942.
- Chen ZH, Yan L, Kangas T, Lunnan R, Schulze S, Sollerman J, Perley DA, Chen TW, Taggart K, Hinds KR, Gal-Yam A, Wang XF, Andreoni I, Bellm E, Bloom JS, Burdge K, Burgos A, Cook D, Dahiwal A, De K, Dekany R, Dugas A, Frederik S, Fremling C, Graham M, Hankins M, Ho A, Jencson J, Karambelkar V, Kasliwal M, Kulkarni S, Laher R, Rusholme B, Sharma Y, Taddia F, Tartaglia L, Thomas BP, Tzanidakis A, Van Roestel J, Walter R, Yang Y, Yao YH and Yaron O (2023a), Jan. The Hydrogen-poor Superluminous Supernovae from the Zwicky Transient Facility Phase I Survey. I. Light Curves and Measurements. *ApJ* 943 (1), 41. doi:10.3847/1538-4357/aca161. 2202. 02059.
- Chen ZH, Yan L, Kangas T, Lunnan R, Sollerman J, Schulze S, Perley DA, Chen TW, Taggart K, Hinds KR, Gal-Yam A, Wang XF, De K, Bellm E, Bloom JS, Dekany R, Graham M, Kasliwal M, Kulkarni S, Laher R, Neill D and Rusholme B (2023b), Jan. The Hydrogen-poor Superluminous Supernovae from the Zwicky Transient Facility Phase I Survey. II. Light-curve Modeling and Characterization of Undulations. *ApJ* 943 (1), 42. doi:10.3847/1538-4357/aca162. 2202. 02060.
- Chevalier RA (2012), Jun. Common Envelope Evolution Leading to Supernovae with Dense Interaction. *ApJL* 752 (1), L2. doi:10.1088/2041-8205/752/1/L2. 1204. 3300.
- Cikota A, De Cia A, Schulze S, Vreeswijk PM, Leloudas G, Gal-Yam A, Perley DA, Cikota S, Kim S, Patat F, Lunnan R, Quimby R, Yaron O, Yan L and Mazzali PA (2017), Aug. Spatially resolved analysis of superluminous supernovae PTF 11hrq and PTF 12dam host galaxies. *MNRAS* 469 (4): 4705–4717. doi:10.1093/mnras/stx1110. 1705. 01948.
- Cikota A, Leloudas G, Bulla M, Inserra C, Chen TW, Spyromilio J, Patat F, Cano Z, Cikota S, Coughlin MW, Kankare E, Lowe TB, Maund JR, Rest A, Smartt SJ, Smith KW, Wainscoat RJ and Young DR (2018), Oct. Testing the magnetar scenario for superluminous supernovae with circular polarimetry. *MNRAS* 479 (4): 4984–4990. doi:10.1093/mnras/sty1891. 1805. 00025.
- Cleland C, McGee SL and Nicholl M (2023), Sep. Metallicity beats sSFR: the connection between superluminous supernova host galaxy environments and the importance of metallicity for their production. *MNRAS* 524 (3): 3559–3567. doi:10.1093/mnras/stad2118. 2307. 06368.
- Cooke J, Sullivan M, Gal-Yam A, Barton EJ, Carlberg RG, Ryan-Weber EV, Horst C, Omori Y and Díaz CG (2012), Nov. Superluminous supernovae at redshifts of 2.05 and 3.90. *Nature* 491 (7423): 228–231. doi:10.1038/nature11521. 1211. 2003.
- Curtin C, Cooke J, Moriya TJ, Tanaka M, Quimby RM, Bernard SR, Galbany L, Jiang J, Lee CH, Maeda K, Morokuma T, Nomoto K, Pignata G, Pritchard T, Suzuki N, Takahashi I, Tanaka M, Tominaga N, Yamaguchi M and Yasuda N (2019), Apr. First Release of High-redshift Superluminous Supernovae from the Subaru High-Z SUpernova CAmpaign (SHIZUCA). II. Spectroscopic Properties. *ApJS* 241 (2), 17. doi:10.3847/1538-4365/ab07c8. 1801. 08241.
- da Cruz MT, Chan Y, Larimer RM, Lesko KT, Norman EB, Stokstad RG, Wietfeldt FE and Žilim I (1992), Sep. Half-life of <sup>56</sup>Ni. *Phys. Rev. C* 46 (3): 1132–1135. doi:10.1103/PhysRevC.46.1132.
- De Cia A, Gal-Yam A, Rubin A, Leloudas G, Vreeswijk P, Perley DA, Quimby R, Yan L, Sullivan M, Flörs A, Sollerman J, Bersier D, Cenko SB, Gal-Yam M, Maguire K, Ofek EO, Prentice S, Schulze S, Spyromilio J, Valenti S, Arcavi I, Corsi A, Howell DA, Mazzali P, Kasliwal MM, Taddia F and Yaron O (2018), Jun. Light Curves of Hydrogen-poor Superluminous Supernovae from the Palomar Transient Factory. *ApJ* 860 (2), 100. doi:10.3847/1538-4357/aab9b6. 1708. 01623.
- Deng JS, Hatano K, Nakamura T, Maeda K, Nomoto K, Nugent P, Aldering G and Branch D (2001), Jan., Luminous SN 1999as: Hypernova with 4M<sub>⊙</sub> <sup>56</sup>Ni Ejected?, Inoue H and Kunieda H, (Eds.), *New Century of X-ray Astronomy*, Astronomical Society of the Pacific Conference Series, 251, pp. 238.
- Desai DK, Siegel DM and Metzger BD (2023), Sep. Three-dimensional General-relativistic Simulations of Neutrino-driven Winds from Magnetized Proto-Neutron Stars. *ApJ* 954 (2), 192. doi:10.3847/1538-4357/acea83. 2306. 03973.
- Dessart L (2019), Jan. Simulations of light curves and spectra for superluminous Type Ic supernovae powered by magnetars. *A&A* 621, A141. doi:10.1051/0004-6361/201834535. 1812. 03749.
- Dessart L, Hillier DJ, Waldman R, Livne E and Blondin S (2012), Oct. Superluminous supernovae: <sup>56</sup>Ni power versus magnetar radiation. *MNRAS* 426 (1): L76–L80. doi:10.1111/j.1745-3933.2012.01329.x. 1208. 1214.
- Dessart L, Waldman R, Livne E, Hillier DJ and Blondin S (2013), Feb. Radiative properties of pair-instability supernova explosions. *MNRAS* 428 (4): 3227–3251. doi:10.1093/mnras/sts269. 1210. 6163.
- Dexter J and Kasen D (2013), Jul. Supernova Light Curves Powered by fallback Accretion. *ApJ* 772 (1), 30. doi:10.1088/0004-637X/772/1/30. 1210. 7240.
- Dong S, Shappee BJ, Prieto JL, Jha SW, Stanek KZ, Holoien TWS, Kochanek CS, Thompson TA, Morrell N, Thompson IB, Basu U, Beacom JF, Bersier D, Brimacombe J, Brown JS, Bufano F, Chen P, Conseil E, Danilec AB, Falco E, Grupe D, Kiyota S, Masi G, Nicholls B, Olivares E, F, Pignata G, Pojmanski G, Simonian GV, Szczygiel DM and Woźniak PR (2016), Jan. ASASSN-15lh: A highly super-luminous supernova. *Science* 351 (6270): 257–260. doi:10.1126/science.aac9613. 1507. 03010.
- Drake AJ, Djorgovski SG, Prieto JL, Mahabal A, Balam D, Williams R, Graham MJ, Catelan M, Beshore E and Larson S (2010), Aug. Discovery of the Extremely Energetic Supernova 2008fz. *ApJL* 718 (2): L127–L131. doi:10.1088/2041-8205/718/2/L127. 0908. 1990.
- Drake AJ, Djorgovski SG, Mahabal A, Anderson J, Roy R, Mohan V, Ravindranath S, Frail D, Gezari S, Neill JD, Ho LC, Prieto JL, Thompson D, Thorstensen J, Wagner M, Kowalski R, Chiang J, Grove JE, Schinzel FK, Wood DL, Carrasco L, Recillas E, Kewley L, Archana KN, Basu A, Wadadekar Y, Kumar B, Myers AD, Phinney ES, Williams R, Graham MJ, Catelan M, Beshore E, Larson S and Christensen E (2011), Jul. The Discovery and Nature of the Optical Transient CSS100217:102913+404220. *ApJ* 735 (2), 106. doi:10.1088/0004-637X/735/2/106. 1103. 5514.
- Drout MR, Soderberg AM, Gal-Yam A, Cenko SB, Fox DB, Leonard DC, Sand DJ, Moon DS, Arcavi I and Green Y (2011), Nov. The First Systematic Study of Type Ibc Supernova Multi-band Light Curves. *ApJ* 741 (2), 97. doi:10.1088/0004-637X/741/2/97. 1011. 4959.
- du Buisson L, Marchant P, Podsiadlowski P, Kobayashi C, Abdalla FB, Taylor P, Mandel I, de Mink SE, Moriya TJ and Langer N (2020), Dec. Cosmic rates of black hole mergers and pair-instability supernovae from chemically homogeneous binary evolution. *MNRAS* 499 (4): 5941–5959. doi:10.1093/mnras/staa3225. 2002. 11630.
- Eftekhari T, Berger E, Margalit B, Blanchard PK, Patton L, Demorest P, Williams PKG, Chatterjee S, Cordes JM, Lunnan R, Metzger BD and Nicholl M (2019), May. A Radio Source Coincident with the Superluminous Supernova PTF10hgi: Evidence for a Central Engine and an Analog of the Repeating FRB 121102? *ApJL* 876 (1), L10. doi:10.3847/2041-8213/ab18a5. 1901. 10479.
- Ercolino A, Jin H, Langer N and Dessart L (2024), May. Interacting supernovae from wide massive binary systems. *A&A* 685, A58. doi:10.1051/0004-6361/202347646. 2308. 01819.
- Euclid Collaboration, Mellier Y, Abdurro'uf, Acevedo Barroso JA, Achúcarro A, Adamek J, Adam R, Addison GE, Aghanim N, Aguena M, Ajani V, Akrami Y, Al-Bahlan A, Alavi A, Albuquerque IS, Alestas G, Alguero G, Allaoui A, Allen SW, Allevalo V, Alonso-Tetilla AV, Altieri B,

Alvarez-Candal A, Amara A, Amendola L, Amiaux J, Andika IT, Andreon S, Andrews A, Angora G, Angulo RE, Annibaldi F, Anselmi A, Anselmi S, Arcari S, Archidiacono M, Aricò G, Arnaud M, Arnouts S, Asgari M, Asorey J, Atayde L, Atek H, Atrio-Barandela F, Aubert M, Aubourg E, Auphan T, Auricchio N, Aussel B, Aussel H, Avelino PP, Avgoustidis A, Avila S, Awan S, Azzollini R, Baccigalupi C, Bachelet E, Bacon D, Baes M, Bagley MB, Bahr-Kalus B, Balaguera-Antolinez A, Balbinot E, Balcells M, Baldi M, Baldry I, Balestra A, Ballardini M, Ballester O, Balogh M, Bañados E, Barbier R, Bardelli S, Barreiro T, Barriere JC, Barros BJ, Barthelemy A, Bartolo N, Basset A, Battaglia P, Battisti AJ, Baugh CM, Baumont L, Bazzanini L, Beaulieu JP, Beckmann V, Belikov AN, Bel J, Bellagamba F, Bella M, Bellini E, Benabed K, Bender R, Benevento G, Bennett CL, Benson K, Bergamini P, Bermejo-Climent JR, Bernardeau F, Bertacca D, Berthe M, Berthier J, Bethermin M, Beutler F, Bevillon C, Bhargava S, Bhatawdekar R, Bisigello L, Biviano A, Blake RP, Blanchard A, Blazek J, Blot L, Bosco A, Bodendorf C, Boenke T, Böhringer H, Bolzonella M, Bonchi A, Bonici M, Bonino D, Bonino L, Bonvin C, Bon W, Booth JT, Borgani S, Borlaff AS, Borsato E, Bosco A, Bose B, Botticella MT, Boucaud A, Bouche F, Boucher JS, Boutigny D, Bouvard T, Bouy H, Bowler RAA, Bozza V, Bozzo E, Branchini E, Brau-Nogue S, Brekke P, Bremer MN, Brescia M, Breton MA, Brinchmann J, Brinckmann T, Brockley-Blatt C, Brodwin M, Brouard L, Brown ML, Bruton S, Bucko J, Buddelmeijer H, Buenadicha G, Buitrago F, Burger P, Burgina C, Busillo V, Busonero D, Cabanac R, Cabayol-Garcia L, Cagliari MS, Caillat A, Caillat L, Calabrese M, Calabro A, Calderone G, Calura F, Camacho Quevedo B, Camera S, Campos L, Canas-Herrera G, Candini GP, Cantiello M, Capobianco V, Cappellaro E, Cappelluti N, Cappi A, Caputi KI, Cara C, Carbone C, Cardone VF, Carella E, Carlberg RG, Carle M, Carminati L, Caro F, Carrasco JM, Carretero J, Carrilho P, Carron Duque J, Carry B, Carvalho A, Carvalho CS, Casas R, Casas S, Casenove P, Casey CM, Cassata P, Castander FJ, Castela D, Castellano M, Castiblanco L, Castignani G, Castro T, Cavet C, Cavuoti S, Chabaud PY, Chambers KC, Charles Y, Charlot S, Chartab N, Chary R, Chaumeil F, Cho H, Chon G, Ciancetta E, Ciliegi P, Cimatti A, Cimino M, Cioni MRL, Claydon R, Cleland C, Clément B, Clements DL, Clerc N, Clesse S, Codis S, Cogato F, Colbert J, Cole RE, Coles P, Collett TE, Collins RS, Colodro-Conde C, Colombo C, Combes F, Conforti V, Congedo G, Conseil S, Conscience CJ, Contarini S, Contini T, Conversi L, Cooray AR, Copin Y, Corasaniti PS, Corcho-Caballero P, Corcione L, Cordes O, Corpace O, Correnti M, Costanzi M, Costille A, Courbin F, Courcoul Mfsud L, Courtois HM, Courinou MC, Covone G, Cowell T, Cragg C, Cresci G, Cristiani S, Croce M, Cropper M, E Crouzet P, Csizi B, Cuby JG, Cucchetti E, Cucciati O, Cuillandre JC, Cunha PAC, Cuozzo V, Daddi E, D'Addona M, Dafonte C, Dagoneau N, Dalessandro E, Dalton GB, D'Amico G, Dannerbauer H, Danto P, Das I, Da Silva A, da Silva R, Daste G, Davies JE, Davini S, de Boer T, Decarli R, De Caro B, Degaudenzi H, Degni G, de Jong JTA, de la Bella LF, de la Torre S, Delhaise F, Delley D, Delucchi G, De Lucia G, Denniston J, De Paolis F, De Petris M, Derosa A, Desai S, Desjacques V, Despali G, Desprez G, De Vicente-Albendea J, Deville Y, Dias JDF, Díaz-Sánchez A, Diaz JJ, Di Domizio S, Diego JM, Di Ferdinando D, Di Giorgio AM, Dimauro P, Dinis J, Dolag K, Dolding C, Dole H, Domínguez Sánchez H, Doré O, Dournac F, Douspis M, Dreihahn H, Droge B, Dryer B, Dubath F, Duc PA, Ducret F, Duffy C, Dufresne F, Duncan CAJ, Dupac X, Duret V, Durrer R, Durret F, Dusini S, Ealet A, Eggemeier A, Eisenhardt PRM, Elbaz D, Elkhachab MY, Ellien A, Endicott J, Enia A, Erben T, Escartin Vigo JA, Escoffier S, Escudero Sanz I, Essert J, Etori S, Ezziati M, Fabbian G, Fabricius M, Fang Y, Farina A, Farina M, Farinelli R, Farrens S, Faustini F, Feltre A, Ferguson AMN, Ferrando P, Ferrari AG, Ferré-Mateu A, Ferreira PG, Ferreras I, Ferrero I, Ferriol S, Ferruit P, Filleul D, Finelli F, Finkelstein SL, Finoguenov A, Fiorini B, Flentge F, Focardi P, Fonseca J, Fontana A, Fontanot F, Fornari F, Fosalba P, Fossati M, Fotopoulou S, Fouchez D, Fourmanoit N, Fraills M, Fraix-Burnet D, Franceschi E, Franco A, Franzetti P, Freihofer J, Frittoli G, Frugier PA, Frusciantè N, Fumagalli A, Fumagalli M, Fumana M, Fu Y, Gabarra L, Galeotta S, Galluccio L, Ganga K, Gao H, García-Bellido J, Garcia K, Gardner JP, Garilli B, Gaspar-Venancio LM, Gasparetto T, Gautard V, Gavazzi R, Gaztanaga E, Genolet L, Genova Santos R, Gentile F, George K, Ghaffari Z, Giacomini F, Gianotti F, Gibb GPS, Gillard W, Gillis B, Ginolfi M, Giocoli C, Girardi M, Giri SK, Goh LWK, Gómez-Alvarez P, Gonzalez AH, Gonzalez EJ, Gonzalez JC, Gouyou Beauchamps S, Gozalias G, Gracia-Carpio J, Grandis S, Granett BR, Granvik M, Grazian A, Grazio A, Grenet C, Grillo C, Grupp F, Gruppioni C, Gruppuso A, Guerbuez C, Guernini S, Guidi M, Guillard P, Gutierrez CM, Guttridge P, Guzzo L, Gwyn S, Haapala J, Haase J, Haddow CR, Hailey M, Hall A, Hall D, Hamaus N, Haridasu BS, Harnois-Déraps J, Harper C, Hartley WG, Hasinger G, Hassani F, Hatch NA, Haugan SVH, Häußler B, Heavens A, Heisenberg L, Helmi A, Helou G, Hemmati S, Henares K, Herent O, Hernández-Monteagudo C, Heuberger T, Hewett PC, Heydenreich S, Hildebrandt H, Hirschmann M, Hjorth J, Hoar J, Hoekstra H, Holland AD, Holliman MS, Holmes W, Hook I, Horeau B, Hormuth F, Hornstrup A, Hosseini S, Hu D, Hudelot P, Hudson MJ, Huertas-Company M, Huff EM, Hughes ACN, Humphrey A, Hunt LK, Huynh DD, Iбата R, Ichikawa K, Iglesias-Groth S, Ilbert O, Ilić S, Ingoglia L, Iodice E, Israel H, Israelsson UE, Izzo L, Jablonka P, Jackson N, Jacobson J, Jafariyazani M, Jahnke K, Jansen H, Jarvis MJ, Jasche J, Jauzac M, Jeffrey N, Jhabvala M, Jimenez-Teja Y, Jimenez Muñoz A, Joachimi B, Johansson PH, Joudaki S, Jullo E, Kajava JJE, Kang Y, Kannawadi A, Kansal V, Karagiannis D, Kärcher M, Kashlinsky A, Kazandjian MV, Keck F, Keihänen E, Kerins E, Kermiche S, Khalil A, Kiessling A, Kiiveri K, Kilbinger M, Kim J, King R, Kirkpatrick CC, Kitching T, Kluge M, Knabenhans M, Knapen JH, Knebe A, Kneib JP, Kohley R, Koopmans LVE, Koskinen H, Koulouridis E, Kou R, Kovács A, Kovačević I, Kowalczyk A, Koyama K, Kraljic K, Krause O, Kruk S, Kubik B, Kuchner U, Kuijken K, Kümmel M, Kunz M, Kurki-Suonio H, Lacasa F, Lacey CG, La Franca F, Lagarde N, Lahav O, Laigle C, La Marca A, La Marle O, Lamine B, Lam MC, Lançon A, Landt H, Langer M, Lapi A, Larcheveque C, Larsen SS, Lattanzi M, Laudisoia P, Laugier D, Laureijs R, Lavaux G, Lawrenson A, Lazanau A, Lazzaro G, Le Boulc'h Q, Le Brun AMC, Le Brun V, Leclercq F, Lee S, Le Graet J, Legrand L, Leirvik KN, Le Jeune M, Lembo M, Le Mignant D, Lepinzan MD, Lepori F, Lesci GF, Lesgourgues J, Leuzzi L, Levi ME, Liaudat TI, Libet G, Liebing P, Ligorì S, Lilje PB, Lin CC, Linde D, Linder E, Lindholm V, Linke L, Li SS, Liu SJ, Lloro I, Lobo FSN, Lodieu N, Lombardi M, Lombriser L, Lonare P, Longo G, López-Cañiego M, Lopez Lopez X, Alvarez JL, Loureiro A, Loveday J, Lusso E, Macias-Perez J, Maciaszek T, Magliocchetti M, Magnard F, Magnier EA, Magro A, Mahler G, Mainetti G, Maino D, Maiorano E, Maiorano E, Malavasi N, Mamon GA, Mancini C, Mandelbaum R, Manera M, Manjón-García A, Mannucci F, Mansutti O, Manteiga Outeiro M, Maoli R, Maraston C, Marcin S, Marcos-Arenal P, Margalef-Bentabol B, Marggraf O, Marinucci D, Marinucci M, Markovic K, Marleau FR, Marpaud J, Martignac J, Martín-Fleitas J, Martín-Moruno P, Martin EL, Martinelli M, Martinet N, Martin H, Martins CJAP, Marulli F, Massari D, Massey R, Masters DC, Matarrese S, Matsuoka Y, Matthew S, Maughan BJ, Mauri N, Maurin L, Maurogordato S, McCarthy K, McConnachie AW, McCracken HJ, McDonald I, McEwen JD, McPartland CJR, Medinaceli E, Mehta V, Mei S, Melchior M, Melin JB, Ménard B, Mendes J, Mendez-Abreu J, Meneghetti M, Mercurio A, Merlin E, Metcalf RB, Meylan G, Migliaccio M, Mignoli M, Miller L, Miluzio M, Milvang-Jensen B, Mimoso JP, Miquel R, Miyatake H, Mobasher B, Mohr JJ, Monaco P, Monguió M, Montoro A, Mora A, Moradinezhad Dizgah A, Moresco M, Moretti C, Morgante G, Morisset N, Moriya TJ, Morris JW, Mortlock DJ, Moscardini L, Mota DF, Moustakas LA, Moutard R, Müller T, Munari E, Murphree G, Murray C, Murray N, Musi P, Nadathur S, Nagam BC, Nagao T, Naidoo K, Nakajima R, Nally C, Natoli P, Navarro-Alsina A, Navarro Girones D, Neissner C, Nersesian A, Nesseris S, Nguyen-Kim HN, Nicastro L, Nichol RC, Nielbock M, Niemi SM, Nieto S, Nilsson K, Noller J, Norberg P, Nourizono A, Ntelis P, Nucita AA, Nugent P, Nunes NJ, Nutma T, Ocampo I, Odier J, Oesch PA, Oguri M, Magalhaes Oliveira D, Onoue M, Oosterbroek T, Oppizzi F, Ordenovic C, Osato K, Pacaud F, Pace F, Padilla C, Paech K, Pagano L, Page MJ, Palazzi E, Paltani S, Pamuk S, Pandolfi S, Paoletti D, Paolillo M, Papaderos P, Pardede K, Paribelli G, Parmar A, Partmann C, Pasian F, Passalacqua F, Paterson K, Patrizii L, Pattison C, Paulino-Afonso A, Paviot R, Peacock JA, Pearce FR, Pedersen K, Peel A, Peletier RF, Pellejero Ibanez M, Pello R, Penny MT, Percival WJ, Perez-Garrido A, Perotto L, Pettorino V, Pezzotta A, Pezzuto S, Philippon A, Piersanti O, Pietroni M, Piga L, Pilo L, Pires S, Pisani A, Pizzella A, Pizzuti L, Plana C, Polenta G, Pollack JE, Poncet M, Pöntinen M, Pool P, Popa LA, Popa V, Popp J, Porciani C, Porth L, Potter D, Poulain M, Pourtsidou A, Pozzetti L, Prandoni I, Pratt GW, Prezelus S, Prieto E, Pugno A, Quai S, Quilley L, Racca GD, Raccanelli A, Rácz G, Radinović S, Radovich M, Ragagnin A, Ragnin U, Raison F, Ramos-Chernenko N, Ranc C, Raylet N, Rebolo R, Refregier A, Reimberg P, Reiprich TH, Renk F, Renzi A, Retre J, Revaz Y, Reylé C, Reynolds L, Rhodes J, Ricci F, Ricci M, Riccio



- G, Ricken SO, Rissanen S, Risso I, Rix HW, Robin AC, Rocca-Volmerange B, Rocci PF, Rodenhuis M, Rodighiero G, Rodriguez Monroy M, Rollins RP, Romanello M, Roman J, Romelli E, Romero-Gomez M, Roncarelli M, Rosati P, Rosset C, Rossetti E, Roster W, Rottgering HJA, Rozas-Fernández A, Ruane K, Rubino-Martin JA, Rudolph A, Ruppin F, Rusholme B, Sacquegna S, Sáez-Casares I, Saga S, Saglia R, Sahlén M, Saifollahi T, Sakr Z, Salvalaggio J, Salvaterra R, Salvati L, Salvato M, Salvignol JC, Sánchez AG, Sanchez E, Sanders DB, Sapone D, Saponara M, Sarpa E, Sarron F, Sartori S, Sassolas B, Saunier L, Sauvage M, Sawicki M, Scaramella R, Scarlata C, Scharré L, Schaye J, Schewtschenko JA, Schindler JT, Schinnerer E, Schirmer M, Schmidt F, Schmidt F, Schmidt M, Schneider A, Schneider M, Schneider P, Schöneberg N, Schrabback T, Schultheis M, Schulz S, Schwartz J, Sciotti D, Scodreggio M, Scognamiglio D, Scott D, Scottez V, Secroun A, Sefusatti E, Seidel G, Seiffert M, Sellentin E, Selwood M, Semboloni E, Sereno M, Serjeant S, Serrano S, Shankar F, Sharples RM, Short A, Shulevski A, Shuntov M, Sias M, Sikkema G, Silvestri A, Simon P, Sirignano C, Sirri G, Skottfelt J, Slezak E, Sluse D, Smith GP, Smith LC, Smith RE, Smit SJA, Soldano F, Solheim BGB, Sorce JG, Sorrenti F, Soubrie E, Spinoglio L, Spurio Mancini A, Stadel J, Stagnaro L, Stanco L, Stanford SA, Starck JL, Stassi P, Steinwagner J, Stern D, Stone C, Strada P, Strafella F, Stramaccioni D, Surace C, Sureau F, Suyu SH, Swindells I, Szafraniec M, Szapudi I, Taamoli S, Talia M, Tallada-Crespí P, Tanidis K, Tao C, Tarrío P, Tavagnacco D, Taylor AN, Taylor JE, Taylor PL, Teixeira EM, Tenti M, Teodoro Idiago P, Teplitz HI, Tereno I, Tessore N, Testa V, Testera G, Tewes M, Teyssier R, Theret N, Thizy C, Thomas PD, Toba Y, Toft S, Toledo-Moreo R, Tolstoy E, Tommasi E, Torbaniuk O, Torradeflot F, Tortora C, Tosi S, Tosti S, Trifoglio M, Troja A, Trombetti T, Tronconi A, Tsedrik M, Tsyganov A, Tucci M, Tutusaus I, Uhlemann C, Ulivi L, Urbano M, Vacher L, Vaillon L, Valdes I, Valentijn EA, Valenziano L, Valieri C, Valiviita J, Van den Broeck M, Vassallo T, Vavrek R, Venemans B, Venhola A, Ventura S, Verdoes Kleijn G, Vergani D, Verma A, Vernizzi F, Veropalumbo A, Verza G, Vescovi C, Vibert D, Viel M, Vielzeuf P, Viglione C, Viitanen A, Villaescusa-Navarro F, Vinciguerra S, Visticot F, Voggel K, von Wietersheim-Kramsta M, Vriend WJ, Wachter S, Walmsley M, Walth G, Walton DM, Walton NA, Wander M, Wang L, Wang Y, Weaver JR, Weller J, Whalen DJ, Wiesmann M, Wilde J, Williams OR, Winther HA, Wittje A, Wong JHW, Wright AH, Yankelevich V, Yeung HW, Youles S, Yung LYA, Zacchei A, Zalesky L, Zamorani G, Zamorano Vitorelli A, Zannoni Marc M, Zennaro M, Zerbi FM, Zinchenko IA, Zoubian J, Zucca E and Zumalacargui M (2024), May. Euclid. I. Overview of the Euclid mission. *arXiv e-prints*, arXiv:2405.13491doi:10.48550/arXiv.2405.13491. 2405. 13491.
- Farmer R, Renzo M, de Mink SE, Marchant P and Justham S (2019), Dec. Mind the Gap: The Location of the Lower Edge of the Pair-instability Supernova Black Hole Mass Gap. *ApJ* 887 (1), 53. doi:10.3847/1538-4357/ab518b. 1910. 12874.
- Farmer R, Renzo M, de Mink SE, Fishbach M and Justham S (2020), Oct. Constraints from Gravitational-wave Detections of Binary Black Hole Mergers on the  $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$  Rate. *ApJL* 902 (2), L36. doi:10.3847/2041-8213/abbadd. 2006. 06678.
- Fischer T, Bastian NUF, Wu MR, Baklanov P, Sorokina E, Blinnikov S, Typel S, Klähn T and Blaschke DB (2018), Oct. Quark deconfinement as a supernova explosion engine for massive blue supergiant stars. *Nature Astronomy* 2: 980–986. doi:10.1038/s41550-018-0583-0. 1712. 08788.
- Folatelli G, Contreras C, Phillips MM, Woosley SE, Blinnikov S, Morrell N, Suntzeff NB, Lee BL, Hamuy M, González S, Krzemiński W, Roth M, Li W, Filippenko AV, Foley RJ, Freedman WL, Madore BF, Persson SE, Murphy D, Boissier S, Galaz G, González L, McCarthy PJ, McWilliam A and Pych W (2006), Apr. SN 2005bf: A Possible Transition Event between Type Ib/c Supernovae and Gamma-Ray Bursts. *ApJ* 641 (2): 1039–1050. doi:10.1086/500531. astro-ph/0509731.
- Fransson C, Ergon M, Challis PJ, Chevalier RA, France K, Kirshner RP, Marion GH, Milisavljevic D, Smith N, Bufano F, Friedman AS, Kangas T, Larsson J, Mattila S, Benetti S, Chornock R, Czekala I, Soderberg A and Sollerman J (2014), Dec. High-density Circumstellar Interaction in the Luminous Type IIc SN 2010jl: The First 1100 Days. *ApJ* 797 (2), 118. doi:10.1088/0004-637X/797/2/118. 1312. 6617.
- Fraser M, Stritzinger MD, Brennan SJ, Pastorello A, Cai Y, Piro AL, Ashall C, Brown P, Burns CR, Elias-Rosa N, Kotak R, Filippenko AV, Galbany L, Hsiao EY, Jha SW, Reguitti A, Zhang J, Moran S, Morrell N, Shappee BJ, Tomasella L, Anderson JP, Barna T, Ochner P, Phillips MM, Tucker M, Wang X, Baron E, Benetti S, Bersten MC, Brink TG, Camacho-Neves Y, Davis S, Dettman KG, Folatelli G, Gutierrez CP, Hoflich P, Holoien TW, Kankare E, Kumar S, Lu J, Mazzali P, Taubenberger S, Tinianont S, Kuncarayakti H, Kwok L, Shahbandeh M, Suntzeff NB, Yan S, Yang Y and Zheng W (2021), Aug. SN 2021csp – the explosion of a stripped envelope star within a H and He-poor circumstellar medium. *arXiv e-prints*, arXiv:2108.07278doi:10.48550/arXiv.2108.07278. 2108. 07278.
- Frohmaier C, Angus CR, Vincenzi M, Sullivan M, Smith M, Nugent PE, Cenko SB, Gal-Yam A, Kulkarni SR, Law NM and Quimby RM (2021), Jan. From core collapse to superluminous: the rates of massive stellar explosions from the Palomar Transient Factory. *MNRAS* 500 (4): 5142–5158. doi:10.1093/mnras/staa3607. 2010. 15270.
- Fuller J (2017), Sep. Pre-supernova outbursts via wave heating in massive stars - I. Red supergiants. *MNRAS* 470 (2): 1642–1656. doi:10.1093/mnras/stx1314. 1704. 08696.
- Funck E, Schötzig U, Woods MJ, Sephton JP, Munster AS, Dean JCJ, Blanchis P and Chauvenet B (1992), Feb.  $^{56}\text{Co}$  standardization and half-life. *Nuclear Instruments and Methods in Physics Research A* 312 (1): 334–338. doi:10.1016/0168-9002(92)90177-6.
- Gal-Yam A (2012), Aug. Luminous Supernovae. *Science* 337 (6097): 927. doi:10.1126/science.1203601. 1208. 3217.
- Gal-Yam A (2019a), Sep. A Simple Analysis of Type I Superluminous Supernova Peak Spectra: Composition, Expansion Velocities, and Dynamics. *ApJ* 882 (2), 102. doi:10.3847/1538-4357/ab2f79. 1806. 08224.
- Gal-Yam A (2019b), Aug. The Most Luminous Supernovae. *ARA&A* 57: 305–333. doi:10.1146/annurev-astro-081817-051819. 1812. 01428.
- Gal-Yam A and Leonard DC (2009), Apr. A massive hypergiant star as the progenitor of the supernova SN 2005gl. *Nature* 458 (7240): 865–867. doi:10.1038/nature07934.
- Gal-Yam A, Mazzali P, Ofek EO, Nugent PE, Kulkarni SR, Kasliwal MM, Quimby RM, Filippenko AV, Cenko SB, Chornock R, Waldman R, Kasen D, Sullivan M, Beshore EC, Drake AJ, Thomas RC, Bloom JS, Poznanski D, Miller AA, Foley RJ, Silverman JM, Arcavi I, Ellis RS and Deng J (2009), Dec. Supernova 2007bi as a pair-instability explosion. *Nature* 462 (7273): 624–627. doi:10.1038/nature08579. 1001. 1156.
- Gal-Yam A, Bruch R, Schulze S, Yang Y, Perley DA, Irani I, Sollerman J, Kool EC, Soumagnac MT, Yaron O, Strotjohann NL, Zimmerman E, Barbarino C, Kulkarni SR, Kasliwal MM, De K, Yao Y, Fremling C, Yan L, Ofek EO, Fransson C, Filippenko AV, Zheng W, Brink TG, Copperwheat CM, Foley RJ, Brown J, Siebert M, Leloudas G, Cabrera-Lavers AL, Garcia-Alvarez D, Marante-Barreto A, Frederick S, Hung T, Wheeler JC, Vinkó J, Thomas BP, Graham MJ, Duev DA, Drake AJ, Dekany R, Bellm EC, Rusholme B, Shupe DL, Andreoni I, Sharma Y, Riddle R, van Roessel J and Knezevic N (2022), Jan. A WC/WO star exploding within an expanding carbon-oxygen-neon nebula. *Nature* 601 (7892): 201–204. doi:10.1038/s41586-021-04155-1. 2111. 12435.
- Georgy C, Meynet G, Ekström S, Wade GA, Petit V, Keszthelyi Z and Hirschi R (2017), Mar. Possible pair-instability supernovae at solar metallicity from magnetic stellar progenitors. *A&A* 599, L5. doi:10.1051/0004-6361/201730401. 1702. 02340.
- Gezari S, Halpern JP, Grupe D, Yuan F, Quimby R, McKay T, Chamarro D, Sisson MD, Akerlof C, Wheeler JC, Brown PJ, Cenko SB, Rau A, Djordjevic JO and Terndrup DM (2009), Jan. Discovery of the Ultra-Bright Type II-L Supernova 2008es. *ApJ* 690 (2): 1313–1321. doi:10.1088/0004-637X/690/2/1313. 0808. 2812.
- Goldreich P and Julian WH (1969), Aug. Pulsar Electrodynamics. *ApJ* 157: 869. doi:10.1086/150119.
- Gomez S, Berger E, Hosseinzadeh G, Blanchard PK, Nicholl M and Villar VA (2021), Jun. The Luminous and Double-peaked Type Ic Supernova 2019stc: Evidence for Multiple Energy Sources. *ApJ* 913 (2), 143. doi:10.3847/1538-4357/abf5e3. 2103. 02611.
- Gomez S, Berger E, Nicholl M, Blanchard PK and Hosseinzadeh G (2022), Dec. Luminous Supernovae: Unveiling a Population between Superluminous and Normal Core-collapse Supernovae. *ApJ* 941 (2), 107. doi:10.3847/1538-4357/ac9842. 2204. 08486.

- Gomez S, Nicholl M, Berger E, Blanchard PK, Villar VA, Rest S, Hosseinzadeh G, Aamer A, Ajay Y, Athukoralalage W, Coulter DC, Eftekhari T, Fiore A, Franz N, Fox O, Gagliano A, Hiramatsu D, Howell DA, Hsu B, Karmen M, Siebert MR, Könyves-Tóth R, Kumar H, McCully C, Pellegrino C, Pierel J, Rest A and Wang Q (2024), Jul. The Type I Superluminous Supernova Catalog I: Light Curve Properties, Models, and Catalog Description. *arXiv e-prints*, arXiv:2407.079462407. 07946.
- Greiner J, Mazzali PA, Kann DA, Krühler T, Pian E, Prentice S, Olivares E. F, Rossi A, Klose S, Taubenberger S, Knust F, Afonso PMJ, Ashall C, Bolmer J, Delvaux C, Diehl R, Elliott J, Filgas R, Fynbo JPU, Graham JF, Guelbenzu AN, Kobayashi S, Leloudas G, Savaglio S, Schady P, Schmidl S, Schweyer T, Sudilovsky V, Tanga M, Updike AC, van Eerten H and Varela K (2015), Jul. A very luminous magnetar-powered supernova associated with an ultra-long  $\gamma$ -ray burst. *Nature* 523 (7559): 189–192. doi:10.1038/nature14579. 1509.03279.
- Hatsukade B, Morokuma-Matsui K, Hayashi M, Tominaga N, Tamura Y, Niinuma K, Motogi K, Morokuma T and Matsuda Y (2020), Aug. Spatially resolved molecular gas properties of host galaxy of Type I superluminous supernova SN 2017egm. *PASJ* 72 (4), L6. doi:10.1093/pasj/psaa052. 2005.06656.
- Heger A and Woosley SE (2002), Mar. The Nucleosynthetic Signature of Population III. *ApJ* 567 (1): 532–543. doi:10.1086/338487. astro-ph/0107037.
- Ho AYQ, Perley DA, Gal-Yam A, Lunnan R, Sollerman J, Schulze S, Das KK, Dobie D, Yao Y, Fremling C, Adams S, Anand S, Andreoni I, Bellm EC, Bruch RJ, Burdge KB, Castro-Tirado AJ, Dahiwalie A, De K, Dekany R, Drake AJ, Duev DA, Graham MJ, Helou G, Kaplan DL, Karambelkar V, Kasliwal MM, Kool EC, Kulkarni SR, Mahabal AA, Medford MS, Miller AA, Nordin J, Ofek E, Petitpas G, Riddle R, Sharma Y, Smith R, Stewart AJ, Taggart K, Tartaglia L, Tzanidakis A and Winters JM (2023), Jun. A Search for Extragalactic Fast Blue Optical Transients in ZTF and the Rate of AT2018cow-like Transients. *ApJ* 949 (2), 120. doi:10.3847/1538-4357/acc533. 2105.08811.
- Hosseinzadeh G, Arcavi I, Valenti S, McCully C, Howell DA, Johansson J, Sollerman J, Pastorello A, Benetti S, Cao Y, Cenko SB, Clubb KI, Corsi A, Duggan G, Elias-Rosa N, Filippenko AV, Fox OD, Fremling C, Horesh A, Karamehmetoglu E, Kasliwal M, Marion GH, Ofek E, Sand D, Taddia F, Zheng W, Fraser M, Gal-Yam A, Inserra C, Laher R, Masci F, Rebbapragada U, Smartt S, Sullivan M, Surace J and Woźniak P (2017), Feb. Type Icn Supernovae Show Photometric Homogeneity and Spectral Diversity at Maximum Light. *ApJ* 836 (2), 158. doi:10.3847/1538-4357/836/2/158. 1608.01998.
- Hosseinzadeh G, Berger E, Metzger BD, Gomez S, Nicholl M and Blanchard P (2022), Jul. Bumpy Declining Light Curves Are Common in Hydrogen-poor Superluminous Supernovae. *ApJ* 933 (1), 14. doi:10.3847/1538-4357/ac67dd. 2109.09743.
- Howell DA (2017), Superluminous Supernovae, Alsabti AW and Murdin P, (Eds.), *Handbook of Supernovae*, pp. 431.
- Howell DA, Kasen D, Lidman C, Sullivan M, Conley A, Astier P, Ballard C, Carlberg RG, Fouchez D, Guy J, Hardin D, Pain R, Palanque-Delabrouille N, Perrett K, Pritchett CJ, Regnault N, Rich J and Ruhlmann-Kleider V (2013), Dec. Two Superluminous Supernovae from the Early Universe Discovered by the Supernova Legacy Survey. *ApJ* 779 (2), 98. doi:10.1088/0004-637X/779/2/98. 1310.0470.
- Hsu B, Blanchard PK, Berger E and Gomez S (2024), Feb. An Extensive Hubble Space Telescope Study of the Offset and Host Light Distributions of Type I Superluminous Supernovae. *ApJ* 961 (2), 169. doi:10.3847/1538-4357/ad12be. 2308.07271.
- Humphreys RM and Davidson K (1994), Oct. The Luminous Blue Variables: Astrophysical Geysers. *PASP* 106: 1025. doi:10.1086/133478.
- Inserra C (2019), Aug. Observational properties of extreme supernovae. *Nature Astronomy* 3: 697–705. doi:10.1038/s41550-019-0854-4. 1908.02314.
- Inserra C and Smartt SJ (2014), Dec. Superluminous Supernovae as Standardizable Candles and High-redshift Distance Probes. *ApJ* 796 (2), 87. doi:10.1088/0004-637X/796/2/87. 1409.4429.
- Inserra C, Smartt SJ, Jerkstrand A, Valenti S, Fraser M, Wright D, Smith K, Chen TW, Kotak R, Pastorello A, Nicholl M, Bresolin F, Kudritzki RP, Benetti S, Botticella MT, Burgett WS, Chambers KC, Ergon M, Flewelling H, Fynbo JPU, Geier S, Hodapp KW, Howell DA, Huber M, Kaiser N, Leloudas G, Magill L, Magnier EA, McCrum MG, Metcalfe N, Price PA, Rest A, Sollerman J, Sweeney W, Taddia F, Taubenberger S, Tonry JL, Wainscoat RJ, Waters C and Young D (2013), Jun. Super-luminous Type Icn Supernovae: Catching a Magnetar by the Tail. *ApJ* 770 (2), 128. doi:10.1088/0004-637X/770/2/128. 1304.3320.
- Inserra C, Nicholl M, Chen TW, Jerkstrand A, Smartt SJ, Krühler T, Anderson JP, Baltay C, Della Valle M, Fraser M, Gal-Yam A, Galbany L, Kankare E, Maguire K, Rabinowitz D, Smith K, Valenti S and Young DR (2017), Jul. Complexity in the light curves and spectra of slow-evolving superluminous supernovae. *MNRAS* 468 (4): 4642–4662. doi:10.1093/mnras/stx834. 1701.00941.
- Inserra C, Prajs S, Gutierrez CP, Angus C, Smith M and Sullivan M (2018a), Feb. A Statistical Approach to Identify Superluminous Supernovae and Probe Their Diversity. *ApJ* 854 (2), 175. doi:10.3847/1538-4357/aaaaaa. 1711.03787.
- Inserra C, Smartt SJ, Gall EEE, Leloudas G, Chen TW, Schulze S, Jerkstrand A, Nicholl M, Anderson JP, Arcavi I, Benetti S, Cartier RA, Childress M, Della Valle M, Flewelling H, Fraser M, Gal-Yam A, Gutiérrez CP, Hosseinzadeh G, Howell DA, Huber M, Kankare E, Krühler T, Magnier EA, Maguire K, McCully C, Prajs S, Primak N, Scalzo R, Schmidt BP, Smith M, Smith KW, Tucker BE, Valenti S, Wilman M, Young DR and Yuan F (2018b), Mar. On the nature of hydrogen-rich superluminous supernovae. *MNRAS* 475 (1): 1046–1072. doi:10.1093/mnras/stx3179. 1604.01226.
- Inserra C, Sullivan M, Angus CR, Macaulay E, Nichol RC, Smith M, Frohmaier C, Gutiérrez CP, Vicenzi M, Möller A, Brout D, Brown PJ, Davis TM, D'Andrea CB, Galbany L, Kessler R, Kim AG, Pan YC, Pursiainen M, Scolnic D, Thomas BP, Wiseman P, Abbott TMC, Annis J, Avila S, Bertin E, Brooks D, Burke DL, Carnero Rosell A, Carrasco Kind M, Carretero J, Castander FJ, Cawthon R, Desai S, Diehl HT, Eifler TF, Finley DA, Flaugher B, Fosalba P, Frieman J, Garcia-Bellido J, Gaztanaga E, Gerdes DW, Giannantonio T, Gruen D, Gruendl RA, Gschwend J, Gutierrez G, Hollowood DL, Honscheid K, James DJ, Krause E, Kuehn K, Kuropatkin N, Li TS, Lidman C, Lima M, Maia MAG, Marshall JL, Martini P, Menanteau F, Miquel R, Plazas Malagón AA, Romer AK, Roodman A, Sako M, Sanchez E, Scarpine V, Schubnell M, Serrano S, Sevilla-Noarbe I, Soares-Santos M, Sobreira F, Suchyta E, Swanson MEC, Tarle G, Thomas D, Tucker DL, Vikram V, Walker AR, Zhang Y, Asorey J, Calcino J, Carollo D, Glazebrook K, Hinton SR, Hoormann JK, Lewis GF, Sharp R, Swann E, Tucker BE and DES Collaboration (2021), Jun. The first Hubble diagram and cosmological constraints using superluminous supernovae. *MNRAS* 504 (2): 2535–2549. doi:10.1093/mnras/stab978. 2004.12218.
- Japelj J, Vergani SD, Salvaterra R, Hunt LK and Mannucci F (2016), Oct. Taking stock of superluminous supernovae and long gamma-ray burst host galaxy comparison using a complete sample of LGRBs. *A&A* 593, A115. doi:10.1051/0004-6361/201628603. 1607.01045.
- Jerkstrand A, Smartt SJ and Heger A (2016), Jan. Nebular spectra of pair-instability supernovae. *MNRAS* 455 (3): 3207–3229. doi:10.1093/mnras/stv2369. 1510.02698.
- Jerkstrand A, Smartt SJ, Inserra C, Nicholl M, Chen TW, Krühler T, Sollerman J, Taubenberger S, Gal-Yam A, Kankare E, Maguire K, Fraser M, Valenti S, Sullivan M, Cartier R and Young DR (2017), Jan. Long-duration Superluminous Supernovae at Late Times. *ApJ* 835 (1), 13. doi:10.3847/1538-4357/835/1/13. 1608.02994.
- Jerkstrand A, Maeda K and Kawabata KS (2020), Jan. A type Ia supernova at the heart of superluminous transient SN 2006gy. *Science* 367 (6476): 415–418. doi:10.1126/science.aaw1469. 2002.10768.
- Joggerst CC and Whalen DJ (2011), Feb. The Early Evolution of Primordial Pair-instability Supernovae. *ApJ* 728 (2), 129. doi:10.1088/0004-637X/728/2/129. 1010.4360.
- Justham S, Podsiadlowski P and Vink JS (2014), Dec. Luminous Blue Variables and Superluminous Supernovae from Binary Mergers. *ApJ* 796

- (2), 121. doi:10.1088/0004-637X/796/2/121. 1410.2426.
- Kangas T, Yan L, Schulze S, Fransson C, Sollerman J, Lunnan R, Omand CMB, Andreoni I, Burruss R, Chen TW, Drake AJ, Fremling C, Gal-Yam A, Graham MJ, Groom SL, Lezmy J, Mahabal AA, Masci FJ, Perley D, Riddle R, Tartaglia L and Yao Y (2022), Oct. The Zwicky Transient Facility phase I sample of hydrogen-rich superluminous supernovae without strong narrow emission lines. *MNRAS* 516 (1): 1193–1218. doi:10.1093/mnras/stac2218. 2207.12059.
- Kankare E, Kotak R, Mattila S, Lundqvist P, Ward MJ, Fraser M, Lawrence A, Smartt SJ, Meikle WPS, Bruce A, Harmanen J, Hutton SJ, Inserra C, Kangas T, Pastorello A, Reynolds T, Romero-Cañizales C, Smith KW, Valenti S, Chambers KC, Hodapp KW, Huber ME, Kaiser N, Kudritzki RP, Magnier EA, Tonry JL, Wainscoat RJ and Waters C (2017), Nov. A population of highly energetic transient events in the centres of active galaxies. *Nature Astronomy* 1: 865–871. doi:10.1038/s41550-017-0290-2. 1711.04577.
- Kann DA, Rossi A, Oates SR, Klose S, Blazek M, Agúí Fernández JF, de Ugarte Postigo A, Thöne CC and Schulze S (2024), Apr. Highly luminous supernovae associated with gamma-ray bursts. II. The luminous blue bump in the afterglow of GRB 140506A. *A&A* 684, A164. doi:10.1051/0004-6361/202142344. 2110.00110.
- Kasen D (2017), Unusual Supernovae and Alternative Power Sources, Alsabti AW and Murdin P, (Eds.), *Handbook of Supernovae*, pp. 939.
- Kasen D and Bildsten L (2010), Jul. Supernova Light Curves Powered by Young Magnetars. *ApJ* 717 (1): 245–249. doi:10.1088/0004-637X/717/1/245. 0911.0680.
- Kasen D, Woosley SE and Heger A (2011), Jun. Pair Instability Supernovae: Light Curves, Spectra, and Shock Breakout. *ApJ* 734 (2), 102. doi:10.1088/0004-637X/734/2/102. 1101.3336.
- Kasen D, Metzger BD and Bildsten L (2016), Apr. Magnetar-driven Shock Breakout and Double-peaked Supernova Light Curves. *ApJ* 821 (1), 36. doi:10.3847/0004-637X/821/1/36. 1507.03645.
- Kashiyama K, Murase K, Bartos I, Kiuchi K and Margutti R (2016), Feb. Multi-messenger Tests for Fast-spinning Newborn Pulsars Embedded in Stripped-envelope Supernovae. *ApJ* 818 (1), 94. doi:10.3847/0004-637X/818/1/94. 1508.04393.
- Khetan N, Cooke J and Branchesi M (2023), May. The rest-frame ultraviolet of superluminous supernovae - I. Potential as cosmological probes. *MNRAS* 521 (2): 2814–2832. doi:10.1093/mnras/stad661. 2302.14659.
- Knop R, Aldering G, Deustua S, Goldhaber G, Kim M, Nugent P, Helin E, Pravdo S, Rabinowitz D and Lawrence K (1999), Mar. Supernovae 1999as and 1999at in Anonymous Galaxies. *IAU Circ.* 7128: 1.
- Könyves-Tóth R (2022), Nov. Premaximum Spectroscopic Diversity of Hydrogen-poor Superluminous Supernovae. *ApJ* 940 (1), 69. doi:10.3847/1538-4357/ac9903. 2210.02153.
- Kostrzewa-Rutkowska Z, Kozłowski S, Wyrzykowski Ł, Djorgovski SG, Glikman E, Mahabal AA and Kopsosov S (2013), Dec. A Plausible (Overlooked) Super-luminous Supernova in the Sloan Digital Sky Survey Stripe 82 Data. *ApJ* 778 (2), 168. doi:10.1088/0004-637X/778/2/168. 1308.2065.
- Kotera K, Phinney ES and Olinto AV (2013), Jul. Signatures of pulsars in the light curves of newly formed supernova remnants. *MNRAS* 432 (4): 3228–3236. doi:10.1093/mnras/stt680. 1304.5326.
- Kozyreva A and Blinnikov S (2015), Dec. Can pair-instability supernova models match the observations of superluminous supernovae? *MNRAS* 454 (4): 4357–4365. doi:10.1093/mnras/stv2287. 1510.00439.
- Kuncarayakti H, Maeda K, Ashall CJ, Prentice SJ, Mattila S, Kankare E, Fransson C, Lundqvist P, Pastorello A, Leloudas G, Anderson JP, Benetti S, Bersten MC, Cappellaro E, Cartier R, Denneau L, Della Valle M, Elias-Rosa N, Folatelli G, Fraser M, Galbany L, Gall C, Gal-Yam A, Gutiérrez CP, Hamaonowicz A, Heinze A, Inserra C, Kangas T, Mazzali P, Melandri A, Pignata G, Rest A, Reynolds T, Roy R, Smartt SJ, Smith KW, Sollerman J, Somero A, Stalder B, Stritzinger M, Taddia F, Tomasella L, Tonry J, Weiland H and Young DR (2018), Feb. SN 2017dio: A Type-Ic Supernova Exploding in a Hydrogen-rich Circumstellar Medium. *ApJL* 854 (1), L14. doi:10.3847/2041-8213/aaa1a. 1712.00027.
- Langer N, Norman CA, de Koter A, Vink JS, Cantiello M and Yoon SC (2007), Nov. Pair creation supernovae at low and high redshift. *A&A* 475 (2): L19–L23. doi:10.1051/0004-6361:20078482. 0708.1970.
- Leloudas G, Chatzopoulos E, Dilday B, Gorosabel J, Vinko J, Gallazzi A, Wheeler JC, Bassett B, Fischer JA, Frieman JA, Fynbo JPU, Goobar A, Jelínek M, Malesani D, Nichol RC, Nordin J, Östman L, Sako M, Schneider DP, Smith M, Sollerman J, Stritzinger MD, Thöne CC and de Ugarte Postigo A (2012), May. SN 2006oz: rise of a super-luminous supernova observed by the SDSS-II SN Survey. *A&A* 541, A129. doi:10.1051/0004-6361/201118498. 1201.5393.
- Leloudas G, Schulze S, Krühler T, Gorosabel J, Christensen L, Mehner A, de Ugarte Postigo A, Amorín R, Thöne CC, Anderson JP, Bauer FE, Gallazzi A, Helminiak KG, Hjorth J, Ibar E, Malesani D, Morell N, Vinko J and Wheeler JC (2015), May. Spectroscopy of superluminous supernova host galaxies. A preference of hydrogen-poor events for extreme emission line galaxies. *MNRAS* 449 (1): 917–932. doi:10.1093/mnras/stv320. 1409.8331.
- Leloudas G, Fraser M, Stone NC, van Velzen S, Jonker PG, Arcavi I, Fremling C, Maund JR, Smartt SJ, Krühler T, Miller-Jones JCA, Vreeswijk PM, Gal-Yam A, Mazzali PA, De Cia A, Howell DA, Inserra C, Patat F, de Ugarte Postigo A, Yaron O, Ashall C, Bar I, Campbell H, Chen TW, Childress M, Elias-Rosa N, Harmanen J, Hosseinzadeh G, Johansson J, Kangas T, Kankare E, Kim S, Kuncarayakti H, Lyman J, Magee MR, Maguire K, Malesani D, Mattila S, McCully CV, Nicholl M, Prentice S, Romero-Cañizales C, Schulze S, Smith KW, Sollerman J, Sullivan M, Tucker BE, Valenti S, Wheeler JC and Young DR (2016), Dec. The superluminous transient ASASSN-15lh as a tidal disruption event from a Kerr black hole. *Nature Astronomy* 1, 0002. doi:10.1038/s41550-016-0002. 1609.02927.
- Levan AJ, Read AM, Metzger BD, Wheatley PJ and Tanvir NR (2013), Jul. Superluminous X-Rays from a Superluminous Supernova. *ApJ* 771 (2), 136. doi:10.1088/0004-637X/771/2/136. 1304.1173.
- Li S, Liang YF, Liao NH, Lei L and Fan YZ (2024), Jul. Fermi-LAT discovery of the GeV emission of the superluminous supernovae SN 2017egm. *arXiv e-prints*, arXiv:2407.05968doi:10.48550/arXiv.2407.05968. 2407.05968.
- Lin W, Wang X, Yan L, Gal-Yam A, Mo J, Brink TG, Filippenko AV, Xiang D, Lunnan R, Zheng W, Brown P, Kasliwal M, Fremling C, Blagorodnova N, Mirzaqulov D, Ehgamberdiev SA, Lin H, Zhang K, Zhang J, Yan S, Zhang J, Chen Z, Deng L, Wang K, Xiao L and Wang L (2023), Jul. A superluminous supernova lightened by collisions with pulsational pair-instability shells. *Nature Astronomy* 7: 779–789. doi:10.1038/s41550-023-01957-3. 2304.10416.
- Liu YQ, Modjaz M and Bianco FB (2017), Aug. Analyzing the Largest Spectroscopic Data Set of Hydrogen-poor Super-luminous Supernovae. *ApJ* 845 (1), 85. doi:10.3847/1538-4357/aa7174. 1612.07321.
- Liu LD, Gao H, Wang XF and Yang S (2021), Apr. Magnetar-driven Shock Breakout Revisited and Implications for Double-peaked Type I Superluminous Supernovae. *ApJ* 911 (2), 142. doi:10.3847/1538-4357/abf042. 2103.09971.
- Lunnan R, Chornock R, Berger E, Laskar T, Fong W, Rest A, Sanders NE, Challis PM, Drout MR, Foley RJ, Huber ME, Kirshner RP, Leibler C, Marion GH, McCrum M, Milisavljevic D, Narayan G, Scolnic D, Smartt SJ, Smith KW, Soderberg AM, Tonry JL, Burgett WS, Chambers KC, Flewelling H, Hodapp KW, Kaiser N, Magnier EA, Price PA and Wainscoat RJ (2014), Jun. Hydrogen-poor Superluminous Supernovae and Long-duration Gamma-Ray Bursts Have Similar Host Galaxies. *ApJ* 787 (2), 138. doi:10.1088/0004-637X/787/2/138. 1311.0026.
- Lunnan R, Chornock R, Berger E, Rest A, Fong W, Scolnic D, Jones DO, Soderberg AM, Challis PM, Drout MR, Foley RJ, Huber ME, Kirshner RP, Leibler C, Marion GH, McCrum M, Milisavljevic D, Narayan G, Sanders NE, Smartt SJ, Smith KW, Tonry JL, Burgett WS, Chambers KC,

- Flewelling H, Kudritzki RP, Wainscoat RJ and Waters C (2015), May. Zooming In on the Progenitors of Superluminous Supernovae With the HST. *ApJ* 804 (2), 90. doi:10.1088/0004-637X/804/2/90. 1411.1060.
- Lunnan R, Chornock R, Berger E, Milisavljevic D, Jones DO, Rest A, Fong W, Fransson C, Margutti R, Drout MR, Blanchard PK, Challis P, Cowperthwaite PS, Foley RJ, Kirshner RP, Morrell N, Riess AG, Roth KC, Scolnic D, Smartt SJ, Smith KW, Villar VA, Chambers KC, Draper PW, Huber ME, Kaiser N, Kudritzki RP, Magnier EA, Metcalfe N and Waters C (2016), Nov. PS1-14bj: A Hydrogen-poor Superluminous Supernova With a Long Rise and Slow Decay. *ApJ* 831 (2), 144. doi:10.3847/0004-637X/831/2/144. 1605.05235.
- Lunnan R, Chornock R, Berger E, Jones DO, Rest A, Czekala I, Dittmann J, Drout MR, Foley RJ, Fong W, Kirshner RP, Laskar T, Leibler CN, Margutti R, Milisavljevic D, Narayan G, Pan YC, Riess AG, Roth KC, Sanders NE, Scolnic D, Smartt SJ, Smith KW, Chambers KC, Draper PW, Flewelling H, Huber ME, Kaiser N, Kudritzki RP, Magnier EA, Metcalfe N, Wainscoat RJ, Waters C and Willman M (2018a), Jan. Hydrogen-poor Superluminous Supernovae from the Pan-STARRS1 Medium Deep Survey. *ApJ* 852 (2), 81. doi:10.3847/1538-4357/aa9f1a. 1708.01619.
- Lunnan R, Fransson C, Vreeswijk PM, Woosley SE, Leloudas G, Perley DA, Quimby RM, Yan L, Blagorodnova N, Bue BD, Cenko SB, De Cia A, Cook DO, Fremling CU, Gatkine P, Gal-Yam A, Kasliwal MM, Kulkarni SR, Masci FJ, Nugent PE, Nyholm A, Rubin A, Suzuki N and Wozniak P (2018b), Sep. A UV resonance line echo from a shell around a hydrogen-poor superluminous supernova. *Nature Astronomy* 2: 887–895. doi:10.1038/s41550-018-0568-z. 1808.04887.
- Maeda K, Tanaka M, Nomoto K, Tominaga N, Kawabata K, Mazzali PA, Umeda H, Suzuki T and Hattori T (2007), Sep. The Unique Type Ib Supernova 2005bf at Nebular Phases: A Possible Birth Event of a Strongly Magnetized Neutron Star. *ApJ* 666 (2): 1069–1082. doi:10.1086/520054. 0705.2713.
- Maeda K, Nozawa T, Sahu DK, Minowa Y, Motohara K, Ueno I, Folatelli G, Pyo TS, Kitagawa Y, Kawabata KS, Anupama GC, Kozasa T, Moriya TJ, Yamanaka M, Nomoto K, Bersten M, Quimby R and Iye M (2013), Oct. Properties of Newly Formed Dust Grains in the Luminous Type II<sub>n</sub> Supernova 2010jl. *ApJ* 776 (1), 5. doi:10.1088/0004-637X/776/1/5. 1308.0406.
- Mandel I and de Mink SE (2016), May. Merging binary black holes formed through chemically homogeneous evolution in short-period stellar binaries. *MNRAS* 458 (3): 2634–2647. doi:10.1093/mnras/stw379. 1601.00007.
- Marchant P, Langer N, Podsiadlowski P, Tauris TM and Moriya TJ (2016), Apr. A new route towards merging massive black holes. *A&A* 588, A50. doi:10.1051/0004-6361/201628133. 1601.03718.
- Margalit B, Metzger BD, Thompson TA, Nicholl M and Sukhbold T (2018), Apr. The GRB-SLSN connection: misaligned magnetars, weak jet emergence, and observational signatures. *MNRAS* 475 (2): 2659–2674. doi:10.1093/mnras/sty013. 1705.01103.
- Margutti R, Metzger BD, Chornock R, Milisavljevic D, Berger E, Blanchard PK, Guidorzi C, Migliori G, Kamble A, Lunnan R, Nicholl M, Coppejans DL, Dall'Osso S, Drout MR, Perna R and Sbarufatti B (2017), Feb. X-Rays from the Location of the Double-humped Transient ASASSN-15lh. *ApJ* 836 (1), 25. doi:10.3847/1538-4357/836/1/25. 1610.01632.
- Margutti R, Chornock R, Metzger BD, Coppejans DL, Guidorzi C, Migliori G, Milisavljevic D, Berger E, Nicholl M, Zauderer BA, Lunnan R, Kamble A, Drout M and Modjaz M (2018), Sep. Results from a Systematic Survey of X-Ray Emission from Hydrogen-poor Superluminous SNe. *ApJ* 864 (1), 45. doi:10.3847/1538-4357/aad2df. 1704.05865.
- Margutti R, Bright JS, Matthews DJ, Coppejans DL, Alexander KD, Berger E, Bietenholz M, Chornock R, DeMarchi L, Drout MR, Eftekhari T, Jacobson-Galán WV, Laskar T, Milisavljevic D, Murase K, Nicholl M, Ormand CMB, Strohm M, Terreran G and VanderLey BA (2023), Sep. Luminous Radio Emission from the Superluminous Supernova 2017ens at 3.3 yr after Explosion. *ApJL* 954 (2), L45. doi:10.3847/2041-8213/ac1fd. 2306.13730.
- Mazzali PA, Sullivan M, Pian E, Greiner J and Kann DA (2016), Jun. Spectrum formation in superluminous supernovae (Type I). *MNRAS* 458 (4): 3455–3465. doi:10.1093/mnras/stw512. 1603.00388.
- Mazzali PA, Moriya TJ, Tanaka M and Woosley SE (2019), Apr. The nature of PISN candidates: clues from nebular spectra. *MNRAS* 484 (3): 3451–3462. doi:10.1093/mnras/stz177. 1901.07629.
- McCrum M, Smartt SJ, Rest A, Smith K, Kotak R, Rodney SA, Young DR, Chornock R, Berger E, Foley RJ, Fraser M, Wright D, Scolnic D, Tonry JL, Urata Y, Huang K, Pastorello A, Botticella MT, Valenti S, Mattila S, Kankare E, Farrow DJ, Huber ME, Stubbs CW, Kirshner RP, Bresolin F, Burgett WS, Chambers KC, Draper PW, Flewelling H, Jedicke R, Kaiser N, Magnier EA, Metcalfe N, Morgan JS, Price PA, Sweeney W, Wainscoat RJ and Waters C (2015), Apr. Selecting superluminous supernovae in faint galaxies from the first year of the Pan-STARRS1 Medium Deep Survey. *MNRAS* 448 (2): 1206–1231. doi:10.1093/mnras/stv034. 1402.1631.
- McIcley L and Soker N (2014), Dec. Wave-driven stellar expansion and binary interaction in pre-supernova outbursts. *MNRAS* 445 (3): 2492–2499. doi:10.1093/mnras/stu1952. 1408.1564.
- Metzger BD, Vurm I, Hascoët R and Beloborodov AM (2014), Jan. Ionization break-out from millisecond pulsar wind nebulae: an X-ray probe of the origin of superluminous supernovae. *MNRAS* 437 (1): 703–720. doi:10.1093/mnras/stt1922. 1307.8115.
- Metzger BD, Margalit B, Kasen D and Quataert E (2015), Dec. The diversity of transients from magnetar birth in core collapse supernovae. *MNRAS* 454 (3): 3311–3316. doi:10.1093/mnras/stv2224. 1508.02712.
- Metzger BD, Berger E and Margalit B (2017), May. Millisecond Magnetar Birth Connects FRB 121102 to Superluminous Supernovae and Long-duration Gamma-Ray Bursts. *ApJ* 841 (1), 14. doi:10.3847/1538-4357/aa633d. 1701.02370.
- Milisavljevic D, Margutti R, Kamble A, Patnaude DJ, Raymond JC, Eldridge JJ, Fong W, Bietenholz M, Challis P, Chornock R, Drout MR, Fransson C, Fesen RA, Grindlay JE, Kirshner RP, Lunnan R, Mackey J, Miller GF, Parrent JT, Sanders NE, Soderberg AM and Zauderer BA (2015), Dec. Metamorphosis of SN 2014C: Delayed Interaction between a Hydrogen Poor Core-collapse Supernova and a Nearby Circumstellar Shell. *ApJ* 815 (2), 120. doi:10.1088/0004-637X/815/2/120. 1511.01907.
- Miller AA, Chornock R, Perley DA, Ganeshalingam M, Li W, Butler NR, Bloom JS, Smith N, Modjaz M, Poznanski D, Filippenko AV, Griffith CV, Shiode JH and Silverman JM (2009), Jan. The Exceptionally Luminous Type II-Linear Supernova 2008es. *ApJ* 690 (2): 1303–1312. doi:10.1088/0004-637X/690/2/1303. 0808.2193.
- Miller AA, Smith N, Li W, Bloom JS, Chornock R, Filippenko AV and Prochaska JX (2010), Jun. New Observations of the Very Luminous Supernova 2006gy: Evidence for Echoes. *AJ* 139 (6): 2218–2229. doi:10.1088/0004-6256/139/6/2218. 0906.2201.
- Mori K, Moriya TJ, Takiwaki T, Kotake K, Horiuchi S and Blinnikov SI (2023), Jan. Light Curves and Event Rates of Axion Instability Supernovae. *ApJ* 943 (1), 12. doi:10.3847/1538-4357/acaaff. 2209.03517.
- Moriya TJ and Maeda K (2012), Sep. A Dip after the Early Emission of Superluminous Supernovae: A Signature of Shock Breakout within Dense Circumstellar Media. *ApJL* 756 (1), L22. doi:10.1088/2041-8205/756/1/L22. 1203.1451.
- Moriya TJ, Blinnikov SI, Tominaga N, Yoshida N, Tanaka M, Maeda K and Nomoto K (2013), Jan. Light-curve modelling of superluminous supernova 2006gy: collision between supernova ejecta and a dense circumstellar medium. *MNRAS* 428 (2): 1020–1035. doi:10.1093/mnras/sts075. 1204.6109.
- Moriya TJ, Liu ZW, Mackey J, Chen TW and Langer N (2015), Dec. Revealing the binary origin of Type Ic superluminous supernovae through nebular hydrogen emission. *A&A* 584, L5. doi:10.1051/0004-6361/201527515. 1510.01621.
- Moriya TJ, Metzger BD and Blinnikov SI (2016), Dec. Supernovae Powered by Magnetars that Transform into Black Holes. *ApJ* 833 (1), 64.

- doi:10.3847/1538-4357/833/1/64. 1606.09316.
- Moriya TJ, Chen TW and Langer N (2017a), Feb. Properties of Magnetars Mimicking  $^{56}\text{Ni}$ -powered Light Curves in Type IC Superluminous Supernovae. *ApJ* 835 (2), 177. doi:10.3847/1538-4357/835/2/177. 1612.06917.
- Moriya TJ, Tanaka M, Morokuma T and Ohsuga K (2017b), Jul. Superluminous Transients at AGN Centers from Interaction between Black Hole Disk Winds and Broad-line Region Clouds. *ApJL* 843 (2), L19. doi:10.3847/2041-8213/aa7af3. 1706.06855.
- Moriya TJ, Nicholl M and Guillochon J (2018a), Nov. Systematic Investigation of the Fallback Accretion-powered Model for Hydrogen-poor Superluminous Supernovae. *ApJ* 867 (2), 113. doi:10.3847/1538-4357/aac53d. 1806.00090.
- Moriya TJ, Sorokina EI and Chevalier RA (2018b), Mar. Superluminous Supernovae. *Space Sci. Rev.* 214 (2), 59. doi:10.1007/s11214-018-0493-6. 1803.01875.
- Moriya TJ, Mazzali PA and Tanaka M (2019a), Apr. Synthetic spectra of energetic core-collapse supernovae and the early spectra of SN 2007bi and SN 1999as. *MNRAS* 484 (3): 3443–3450. doi:10.1093/mnras/stz262. 1901.07628.
- Moriya TJ, Tanaka M, Yasuda N, Jiang Ja, Lee CH, Maeda K, Morokuma T, Nomoto K, Quimby RM, Suzuki N, Takahashi I, Tanaka M, Tominaga N, Yamaguchi M, Bernard SR, Cooke J, Curtin C, Galbany L, González-Gaitán S, Pignata G, Pritchard T, Komiya Y and Lupton RH (2019b), Apr. First Release of High-Redshift Superluminous Supernovae from the Subaru High-Z SUpernova CAmpaign (SHIZUCA). I. Photometric Properties. *ApJS* 241 (2), 16. doi:10.3847/1538-4365/ab07c5. 1801.08240.
- Moriya TJ, Inserra C, Tanaka M, Cappellaro E, Della Valle M, Hook I, Kotak R, Longo G, Mannucci F, Mattila S, Tao C, Altieri B, Amara A, Auricchio N, Bonino D, Branchini E, Brescia M, Brinchmann J, Camera S, Capobianco V, Carbone C, Carretero J, Castellano M, Cavuoti S, Cimatti A, Ciedassou R, Congedo G, Conselice CJ, Conversi L, Copin Y, Corcione L, Courbin F, Cropper M, Da Silva A, Degaudenzi H, Douspis M, Dubath F, Duncan CAJ, Dupac X, Dusini S, Ealet A, Farrens S, Ferriol S, Frailis M, Franceschi E, Fumana M, Garilli B, Gillard W, Gillis B, Giocoli C, Grazian A, Grupp F, Haugan SVH, Holmes W, Hornuth F, Hornstrup A, Jahnke K, Kermiche S, Kiessling A, Kilbinger M, Kitching R, Kurki-Suonio H, Ligorì S, Lilje PB, Llori I, Maiorano A, Mansutti O, Marggraf O, Markovic K, Marulli F, Massey R, McCracken HJ, Melchior M, Meneghetti M, Meylan G, Moresco M, Moscardini L, Munari E, Niemi SM, Padilla C, Paltani S, Pasian F, Pedersen K, Pettorino V, Poncet M, Popa L, Raison F, Rhodes J, Riccio G, Rossetti E, Saglia R, Sartoris B, Schneider P, Secroun A, Seidel G, Sirignano C, Sirri G, Stanco L, Tallada-Crespí P, Taylor AN, Tereno I, Toledo-Moreo R, Torradeflot F, Wang Y, Zamorani G, Zoubian J, Andreon S, Scottez V and Morris PW (2022a), Oct. Euclid: Searching for pair-instability supernovae with the Deep Survey. *A&A* 666, A157. doi:10.1051/0004-6361/202243810. 2204.08727.
- Moriya TJ, Murase K, Kashiyama K and Blinnikov SI (2022b), Jul. Variable thermal energy injection from magnetar spin-down as a possible cause of stripped-envelope supernova light-curve bumps. *MNRAS* 513 (4): 6210–6218. doi:10.1093/mnras/stac1352. 2202.03082.
- Moriya TJ, Quimby RM and Robertson BE (2022c), Feb. Discovering Supernovae at the Epoch of Reionization with the Nancy Grace Roman Space Telescope. *ApJ* 925 (2), 211. doi:10.3847/1538-4357/ac415e. 2108.01801.
- Morris PW, Waters LBFM, Barlow MJ, Lim T, de Koter A, Voors RHM, Cox P, de Graauw T, Henning T, Hony S, Lamers HJGLM, Mutschke H and Trams NR (1999), Dec. Discovery of a massive equatorial torus in the  $\eta$  Carinae stellar system. *Nature* 402 (6761): 502–504. doi:10.1038/990048.
- Murase K, Kashiyama K, Kiuchi K and Bartos I (2015), May. Gammy-Ray and Hard X-Ray Emission from Pulsar-aided Supernovae as a Probe of Particle Acceleration in Embryonic Pulsar Wind Nebulae. *ApJ* 805 (1), 82. doi:10.1088/0004-637X/805/1/82. 1411.0619.
- Murase K, Omand CMB, Coppejans DL, Nagai H, Bower GC, Chornock R, Fox DB, Kashiyama K, Law C, Margutti R and Mészáros P (2021), Nov. ALMA and NOEMA constraints on synchrotron nebular emission from embryonic superluminous supernova remnants and radio-gamma-ray connection. *MNRAS* 508 (1): 44–51. doi:10.1093/mnras/stab2506. 2105.05239.
- Nadyozhin DK (1994), Jun. The Properties of Ni CO Fe Decay. *ApJS* 92: 527. doi:10.1086/192008.
- Neill JD, Sullivan M, Gal-Yam A, Quimby R, Ofek E, Wyder TK, Howell DA, Nugent P, Seibert M, Martin DC, Overzier R, Barlow TA, Foster K, Friedman PG, Morrissey P, Neff SG, Schiminovich D, Bianchi L, Donas J, Heckman TM, Lee YW, Madore BF, Milliard B, Rich RM and Szalay AS (2011), Jan. The Extreme Hosts of Extreme Supernovae. *ApJ* 727 (1), 15. doi:10.1088/0004-637X/727/1/15. 1011.3512.
- Nicholl M (2021), Oct. Superluminous supernovae: an explosive decade. *Astronomy and Geophysics* 62 (5): 5.34–5.42. doi:10.1093/astrogeo/atab092. 2109.08697.
- Nicholl M and Smartt SJ (2016), Mar. Seeing double: the frequency and detectability of double-peaked superluminous supernova light curves. *MNRAS* 457 (1): L79–L83. doi:10.1093/mnras/slv210. 1511.03740.
- Nicholl M, Smartt SJ, Jerkstrand A, Inserra C, Sim SA, Chen TW, Benetti S, Fraser M, Gal-Yam A, Kankare E, Maguire K, Smith K, Sullivan M, Valenti S, Young DR, Baltay C, Bauer FE, Baumont S, Bersier D, Botticella MT, Childress M, Dennefeld M, Della Valle M, Elias-Rosa N, Feindt U, Galbany L, Hadjijska E, Le Guillou L, Leloudas G, Mazzali P, McKinnon R, Polshaw J, Rabinowitz D, Rostami S, Scalzo R, Schmidt BP, Schulze S, Sollerman J, Taddia F and Yuan F (2015a), Oct. On the diversity of superluminous supernovae: ejected mass as the dominant factor. *MNRAS* 452 (4): 3869–3893. doi:10.1093/mnras/stv1522. 1503.03310.
- Nicholl M, Smartt SJ, Jerkstrand A, Sim SA, Inserra C, Anderson JP, Baltay C, Benetti S, Chambers K, Chen TW, Elias-Rosa N, Feindt U, Fwelling HA, Fraser M, Gal-Yam A, Galbany L, Huber ME, Kangas T, Kankare E, Kotak R, Krühler T, Maguire K, McKinnon R, Rabinowitz D, Rostami S, Schulze S, Smith KW, Sullivan M, Tonry JL, Valenti S and Young DR (2015b), Jul. LSQ14bdq: A Type Ic Super-luminous Supernova with a Double-peaked Light Curve. *ApJL* 807 (1), L18. doi:10.1088/2041-8205/807/1/L18. 1505.01078.
- Nicholl M, Berger E, Margutti R, Blanchard PK, Guillochon J, Leja J and Chornock R (2017a), Aug. The Superluminous Supernova SN 2017egm in the Nearby Galaxy NGC 3191: A Metal-rich Environment Can Support a Typical SLSN Evolution. *ApJL* 845 (1), L8. doi:10.3847/2041-8213/aa82b1. 1706.08517.
- Nicholl M, Berger E, Margutti R, Blanchard PK, Milisavljevic D, Challis P, Metzger BD and Chornock R (2017b), Jan. An Ultraviolet Excess in the Superluminous Supernova Gaia16apd Reveals a Powerful Central Engine. *ApJL* 835 (1), L8. doi:10.3847/2041-8213/aa56c5. 1611.06993.
- Nicholl M, Berger E, Blanchard PK, Gomez S and Chornock R (2019), Jan. Nebular-phase Spectra of Superluminous Supernovae: Physical Insights from Observational and Statistical Properties. *ApJ* 871 (1), 102. doi:10.3847/1538-4357/aaf470. 1808.00510.
- Nicholl M, Blanchard PK, Berger E, Chornock R, Margutti R, Gomez S, Lunnan R, Miller AA, Fong Wf, Terreran G, Vigna-Gómez A, Bhirombhakdi K, Bieryla A, Challis P, Laher RR, Masci FJ and Paterson K (2020), Apr. An extremely energetic supernova from a very massive star in a dense medium. *Nature Astronomy* 4: 893–899. doi:10.1038/s41550-020-1066-7. 2004.05840.
- Ørum SV, Ivens DL, Strandberg P, Leloudas G, Man AWS and Schulze S (2020), Nov. The interacting nature of dwarf galaxies hosting superluminous supernovae. *A&A* 643, A47. doi:10.1051/0004-6361/202038176. 2009.03925.
- Ostriker JP and Gunn JE (1971), Mar. Do Pulsars Make Supernovae? *ApJL* 164: L95. doi:10.1086/180699.
- Ouyed R, Kostka M, Koning N, Leahy DA and Steffen W (2012), Jun. Quark nova imprint in the extreme supernova explosion SN 2006gy. *MNRAS* 423 (2): 1652–1662. doi:10.1111/j.1365-2966.2012.20986.x. 1010.5530.
- Pan YC, Foley RJ, Smith M, Galbany L, D'Andrea CB, González-Gaitán S, Jarvis MJ, Kessler R, Kovacs E, Lidman C, Nichol RC, Papadopoulos A, Sako M, Sullivan M, Abbott TMC, Abdalla FB, Annis J, Bechtol K, Benoit-Lévy A, Brooks D, Buckley-Geer E, Burke DL, Carrero Rosell A, Carrasco Kind M, Carretero J, Castander FJ, Cunha CE, da Costa LN, Desai S, Diehl HT, Doel P, Eifler TF, Finley DA, Flaugher B, Frieman

- J, García-Bellido J, Goldstein DA, Gruen D, Gruendl RA, Gschwend J, Gutierrez G, James DJ, Kim AG, Krause E, Kuehn K, Kuropatkin N, Lahav O, Lima M, Maia MAG, March M, Marshall JL, Martini P, Miquel R, Nugent P, Plazas AA, Romer AK, Sanchez E, Scarpine V, Schubnell M, Sevilla-Noarbe I, Smith RC, Sobreira F, Suchyta E, Swanson MEC, Thomas RC, Walker AR and DES Collaboration (2017), Oct. DES15E2mlf: a spectroscopically confirmed superluminous supernova that exploded 3.5 Gyr after the big bang. *MNRAS* 470 (4): 4241–4250. doi:10.1093/mnras/stx1467. 1707.06649.
- Pastorello A, Smartt SJ, Mattila S, Eldridge JJ, Young D, Itagaki K, Yamaoka H, Navasardyan H, Valenti S, Patat F, Agnoletto I, Augusteijn T, Benetti S, Cappellaro E, Boles T, Bonnet-Bidaud JM, Botticella MT, Bufano F, Cao C, Deng J, Dennefeld M, Elias-Rosa N, Harutyunyan A, Keenan FP, Iijima T, Lorenzi V, Mazzali PA, Meng X, Nakano S, Nielsen TB, Smoker JV, Stanishev V, Turatto M, Xu D and Zampieri L (2007), Jun. A giant outburst two years before the core-collapse of a massive star. *Nature* 447 (7146): 829–832. doi:10.1038/nature05825. astro-ph/0703663.
- Perley DA, Quimby RM, Yan L, Vreeswijk PM, De Cia A, Lunnan R, Gal-Yam A, Yaron O, Filippenko AV, Graham ML, Laher R and Nugent PE (2016), Oct. Host-galaxy Properties of 32 Low-redshift Superluminous Supernovae from the Palomar Transient Factory. *ApJ* 830 (1), 13. doi:10.3847/0004-637X/830/1/13. 1604.08207.
- Perley DA, Sollerman J, Schulze S, Yao Y, Fremling C, Gal-Yam A, Ho AYQ, Yang Y, Kool EC, Irani I, Yan L, Andreoni I, Baade D, Bellm EC, Brink TG, Chen TW, Cikota A, Coughlin MW, Dahiwal A, Dekany R, Duev DA, Filippenko AV, Hoeflich P, Kasliwal MM, Kulkarni SR, Lunnan R, Masci FJ, Maund JR, Medford MS, Riddle R, Rosnet P, Shupe DL, Strotjohann NL, Tzanidakis A and Zheng W (2022), Mar. The Type Icn SN 2021csp: Implications for the Origins of the Fastest Supernovae and the Fates of Wolf-Rayet Stars. *ApJ* 927 (2), 180. doi:10.3847/1538-4357/ac478e. 2111.12110.
- Petrushevska T, Leloudas G, Ilić D, Bronikowski M, Charalampopoulos P, Jaisawal GK, Paraskeva E, Pursiainen M, Rakić N, Schulze S, Taggart K, Wedderkopp CK, Anderson JP, de Boer T, Chambers K, Chen TW, Damjanović G, Fraser M, Gao H, Gomboc A, Gromadzki M, Ihanec N, Maguire K, Marčun B, Müller-Bravo TE, Nicholl M, Onori F, Reynolds TM, Smartt SJ, Sollerman J, Smith KW, Wevers T and Wyrzykowski Ł (2023), Jan. The rise and fall of the iron-strong nuclear transient PS16dtm. *A&A* 669, A140. doi:10.1051/0004-6361/202244623. 2211.17097.
- Prajs S, Sullivan M, Smith M, Levan A, Karpenka NV, Edwards TDP, Walker CR, Wolf WM, Balland C, Carlberg R, Howell DA, Lidman C, Pain R, Pritchett C and Ruhlmann-Kleider V (2017), Jan. The volumetric rate of superluminous supernovae at  $z \sim 1$ . *MNRAS* 464 (3): 3568–3579. doi:10.1093/mnras/stw1942. 1605.05250.
- Prieto JL, Kistler MD, Thompson TA, Yüksel H, Kochanek CS, Stanek KZ, Beacom JF, Martini P, Pasquali A and Bechtold J (2008), Jul. Discovery of the Dust-Enshrouded Progenitor of SN 2008S with Spitzer. *ApJL* 681 (1): L9. doi:10.1086/589922. 0803.0324.
- Pursiainen M, Leloudas G, Cikota A, Bulla M, Inserra C, Patat F, Wheeler JC, Aamer A, Gal-Yam A, Maund J, Nicholl M, Schulze S, Sollerman J and Yang Y (2023), Jun. Polarimetry of hydrogen-poor superluminous supernovae. *A&A* 674, A81. doi:10.1051/0004-6361/202345945. 2301.08111.
- Quataert E and Shiode J (2012), Jun. Wave-driven mass loss in the last year of stellar evolution: setting the stage for the most luminous core-collapse supernovae. *MNRAS* 423 (1): L92–L96. doi:10.1111/j.1745-3933.2012.01264.x. 1202.5036.
- Quimby RM, Aldering B, Wheeler JC, Höflich P, Akerlof CW and Rykoff ES (2007), Oct. SN 2005ap: A Most Brilliant Explosion. *ApJL* 668 (2): L99–L102. doi:10.1086/522862. 0709.0302.
- Quimby RM, Kulkarni SR, Kasliwal MM, Gal-Yam A, Arcavi I, Sullivan M, Nugent P, Thomas R, Howell DA, Nakar E, Bildsten L, Theissen C, Law NM, Dekany R, Rahmer G, Hale D, Smith R, Ofek EO, Zolkower J, Velur V, Walters R, Henning J, Bui K, McKenna D, Poznanski D, Cenko SB and Levitan D (2011), Jun. Hydrogen-poor superluminous stellar explosions. *Nature* 474 (7352): 487–489. doi:10.1038/nature10095. 0910.0059.
- Quimby RM, Yuan F, Akerlof C and Wheeler JC (2013), May. Rates of superluminous supernovae at  $z \sim 0.2$ . *MNRAS* 431 (1): 912–922. doi:10.1093/mnras/stt213. 1302.0911.
- Quimby RM, De Cia A, Gal-Yam A, Leloudas G, Lunnan R, Perley DA, Vreeswijk PM, Yan L, Bloom JS, Cenko SB, Cooke J, Ellis R, Filippenko AV, Kasliwal MM, Kleiser IKW, Kulkarni SR, Matheson T, Nugent PE, Pan YC, Silverman JM, Sternberg A, Sullivan M and Yaron O (2018), Mar. Spectra of Hydrogen-poor Superluminous Supernovae from the Palomar Transient Factory. *ApJ* 855 (1), 2. doi:10.3847/1538-4357/aaac2f. 1802.07820.
- Rakavy G and Shaviv G (1967), Jun. Instabilities in Highly Evolved Stellar Models. *ApJ* 148: 803. doi:10.1086/149204.
- Regós E and Vinkó J (2019), Apr. Detection and Classification of Supernovae Beyond  $z \sim 2$  Redshift with the James Webb Space Telescope. *ApJ* 874 (2), 158. doi:10.3847/1538-4357/ab0a73. 1811.00891.
- Rest A, Foley RJ, Gezari S, Narayan G, Draine B, Olsen K, Huber ME, Matheson T, Garg A, Welch DL, Becker AC, Challis P, Clocchiatti A, Cook KH, Damke G, Meixner M, Miknaitis G, Minniti D, Morelli L, Nikolaev S, Pignata G, Prieto JL, Smith RC, Stubbs C, Suntzeff NB, Walker AR, Wood-Vasey WM, Zenteno A, Wyrzykowski L, Udalski A, Szymański MK, Kubiak M, Pietrzyński G, Soszyński I, Szewczyk O, Ulaczyk K and Poleski R (2011), Mar. Pushing the Boundaries of Conventional Core-collapse Supernovae: The Extremely Energetic Supernova SN 2003ma. *ApJ* 729 (2), 88. doi:10.1088/0004-637X/729/2/88. 0911.2002.
- Richardson D, Jenkins Robert L. I, Wright J and Maddox L (2014), May. Absolute-magnitude Distributions of Supernovae. *AJ* 147 (5), 118. doi:10.1088/0004-6256/147/5/118. 1403.5755.
- Sabhahit GN, Vink JS, Sander AAC and Higgins ER (2023), Sep. Very massive stars and pair-instability supernovae: mass-loss framework for low metallicity. *MNRAS* 524 (1): 1529–1546. doi:10.1093/mnras/stad1888. 2306.11785.
- Saito S, Tanaka M, Mazzali PA, Hachinger S and Hotokezaka K (2024), May. On the Formation of the W-shaped O II Lines in Spectra of Type I Superluminous Supernovae. *ApJ* 967 (1), 13. doi:10.3847/1538-4357/ad39e8. 2404.03112.
- Sakstein J, Croon D and McDermott SD (2022), May. Axion instability supernovae. *Phys. Rev. D* 105 (9), 095038. doi:10.1103/PhysRevD.105.095038. 2203.06160.
- Schulze S, Krühler T, Leloudas G, Gorosabel J, Mehner A, Buchner J, Kim S, Ibar E, Amorín R, Herrero-Illana R, Anderson JP, Bauer FE, Christensen L, de Pasquale M, de Ugarte Postigo A, Gallazzi A, Hjorth J, Morrell N, Malesani D, Sparre M, Stalder B, Stark AA, Thöne CC and Wheeler JC (2018), Jan. Cosmic evolution and metal aversion in superluminous supernova host galaxies. *MNRAS* 473 (1): 1258–1285. doi:10.1093/mnras/stx2352. 1612.05978.
- Schulze S, Yaron O, Sollerman J, Leloudas G, Gal A, Wright AH, Lunnan R, Gal-Yam A, Ofek EO, Perley DA, Filippenko AV, Kasliwal MM, Kulkarni SR, Neill JD, Nugent PE, Quimby RM, Sullivan M, Strotjohann NL, Arcavi I, Ben-Ami S, Bianco F, Bloom JS, De K, Fraser M, Fremling CU, Horesh A, Johansson J, Kelly PL, Knežević N, Knežević S, Maguire K, Nyholm A, Papadogiannakis S, Petrushevska T, Rubin A, Yan L, Yang Y, Adams SM, Bufano F, Clubb KI, Foley RJ, Green Y, Harmanen J, Ho AYQ, Hook IM, Hosseinzadeh G, Howell DA, Kong AKH, Kotak R, Matheson T, McCully C, Milisavljević D, Pan YC, Poznanski D, Shivvers I, van Velzen S and Verbeek KK (2021), Aug. The Palomar Transient Factory Core-collapse Supernova Host-galaxy Sample. I. Host-galaxy Distribution Functions and Environment Dependence of Core-collapse Supernovae. *ApJS* 255 (2), 29. doi:10.3847/1538-4365/abff5e. 2008.05988.
- Schulze S, Fransson C, Kozlyeva A, Chen TW, Yaron O, Jerkstrand A, Gal-Yam A, Sollerman J, Yan L, Kangas T, Leloudas G, Omand CMB, Smartt SJ, Yang Y, Nicholl M, Sarin N, Yao Y, Brink TG, Sharon A, Rossi A, Chen P, Chen Z, Cikota A, De K, Drake AJ, Filippenko AV,

- Fremling C, Fréour L, Fynbo JPU, Ho AYQ, Inserra C, Irani I, Kuncarayakti H, Lunnan R, Mazzali P, Ofek EO, Palazzi E, Perley DA, Pursiainen M, Rothberg B, Shingles LJ, Smith K, Taggart K, Tartaglia L, Zheng W, Anderson JP, Cassara L, Christensen E, George Djorgovski S, Galbany L, Gkini A, Graham MJ, Gromadzki M, Groom SL, Hiramatsu D, Andrew Howell D, Kasiwal MM, McCully C, Müller-Bravo TE, Paiano S, Paraskava E, Pessi PJ, Polishook D, Rau A, Rigault M and Rusholme B (2024), Mar. 1100 days in the life of the supernova 2018ibb. The best pair-instability supernova candidate, to date. *A&A* 683, A223. doi:10.1051/0004-6361/202346855. 2305.05796.
- Sharma Y, Sollerman J, Fremling C, Kulkarni SR, De K, Irani I, Schulze S, Strotjohann NL, Gal-Yam A, Maguire K, Perley DA, Bellm EC, Kool EC, Brink TG, Bruch R, Deckers M, Dekany R, Dugas A, Filippenko AV, Goldwasser S, Graham MJ, Graham ML, Groom SL, Hankins M, Jencson J, Johansson JP, Karambelkar V, Kasiwal MM, Masci FJ, Medford MS, Neill JD, Nir G, Riddle RL, Rigault M, Schweyer T, Terwel JH, Yan L, Yang Y and Yao Y (2023), May. A Systematic Study of Ia-CSM Supernovae from the ZTF Bright Transient Survey. *ApJ* 948 (1), 52. doi:10.3847/1538-4357/acbc16. 2301.04637.
- Smith N, Foley RJ, Bloom JS, Li W, Filippenko AV, Gavazzi R, Ghez A, Konopacky Q, Malkan MA, Marshall PJ, Pooley D, Treu T and Woo JH (2008), Oct. Late-Time Observations of SN 2006gy: Still Going Strong. *ApJ* 686 (1): 485–491. doi:10.1086/590141. 0802.1743.
- Smith N, Chornock R, Silverman JM, Filippenko AV and Foley RJ (2010), Feb. Spectral Evolution of the Extraordinary Type II<sub>n</sub> Supernova 2006gy. *ApJ* 709 (2): 856–883. doi:10.1088/0004-637X/709/2/856. 0906.2200.
- Smith M, Sullivan M, D'Andrea CB, Castander FJ, Casas R, Prajs S, Papadopoulos A, Nichol RC, Karpenka NV, Bernard SR, Brown P, Cartier R, Cooke J, Curtin C, Davis TM, Finley DA, Foley RJ, Gal-Yam A, Goldstein DA, González-Gaitán S, Gupta RR, Howell DA, Inserra C, Kessler R, Lidman C, Marriner J, Nugent P, Pritchard TA, Sako M, Smartt S, Smith RC, Spinka H, Thomas RC, Wolf RC, Zenteno A, Abbott TMC, Benoit-Lévy A, Bertin E, Brooks D, Buckley-Geer E, Carnero Rosell A, Carrasco Kind M, Carretero J, Croce M, Cunha CE, da Costa LN, Desai S, Diehl HT, Doel P, Estrada J, Evrard AE, Flaughner B, Fosalba P, Frieman J, Gerdes DW, Gruen D, Gruendl RA, James DJ, Kuehn K, Kuropatkin N, Lahav O, Li TS, Marshall JL, Martini P, Miller CJ, Miquel R, Nord B, Ogando R, Plazas AA, Reil K, Romer AK, Roodman A, Rykoff ES, Sanchez E, Scarpine V, Schubnell M, Sevilla-Noarbe I, Soares-Santos M, Sobreira F, Suchyta E, Swanson MEC, Tarle G, Walker AR, Wester W and DES Collaboration (2016), Feb. DES14X3taz: A Type I Superluminous Supernova Showing a Luminous, Rapidly Cooling Initial Pre-peak Bump. *ApJL* 818 (1), L8. doi:10.3847/2041-8205/818/1/L8. 1512.06043.
- Smith M, Sullivan M, Nichol RC, Galbany L, D'Andrea CB, Inserra C, Lidman C, Rest A, Schirmer M, Filippenko AV, Zheng W, Cenko SB, Angus CR, Brown PJ, Davis TM, Finley DA, Foley RJ, González-Gaitán S, Gutiérrez CP, Kessler R, Kuhlmann S, Marriner J, Möller A, Nugent PE, Prajs S, Thomas R, Wolf R, Zenteno A, Abbott TMC, Abdalla FB, Allam S, Annis J, Bechtol K, Benoit-Lévy A, Bertin E, Brooks D, Burke DL, Carnero Rosell A, Carrasco Kind M, Carretero J, Castander FJ, Croce M, Cunha CE, da Costa LN, Davis C, Desai S, Diehl HT, Doel P, Eifler TF, Flaughner B, Fosalba P, Frieman J, García-Bellido J, Gaztanaga E, Gerdes DW, Goldstein DA, Gruen D, Gruendl RA, Gschwend J, Gutierrez G, Honscheid K, James DJ, Johnson MWG, Kuehn K, Kuropatkin N, Li TS, Lima M, Maia MAG, Marshall JL, Martini P, Menanteau F, Miller CJ, Miquel R, Ogando RLC, Petravick D, Plazas AA, Romer AK, Rykoff ES, Sako M, Sanchez E, Scarpine V, Schindler R, Schubnell M, Sevilla-Noarbe I, Smith RC, Soares-Santos M, Sobreira F, Suchyta E, Swanson MEC, Tarle G, Walker AR and DES Collaboration (2018), Feb. Studying the Ultraviolet Spectrum of the First Spectroscopically Confirmed Supernova at Redshift Two. *ApJ* 854 (1), 37. doi:10.3847/1538-4357/aaa126. 1712.04535.
- Soker N (2022), Aug. Powering Luminous Core Collapse Supernovae with Jets. *ApJ* 935 (2), 108. doi:10.3847/1538-4357/ac822d. 2205.09560.
- Sorokina E, Blinnikov S, Nomoto K, Quimby R and Tolstov A (2016), Sep. Type I Superluminous Supernovae as Explosions inside Non-hydrogen Circumstellar Envelopes. *ApJ* 829 (1), 17. doi:10.3847/0004-637X/829/1/17. 1510.00834.
- Strolger LG, Dahlen T, Rodney SA, Graur O, Riess AG, McCully C, Ravindranath S, Mobasher B and Shahady AK (2015), Nov. The Rate of Core Collapse Supernovae to Redshift 2.5 from the CANDELS and CLASH Supernova Surveys. *ApJ* 813 (2), 93. doi:10.1088/0004-637X/813/2/93. 1509.06574.
- Sun L, Xiao L and Li G (2022), Jul. A mid-infrared study of superluminous supernovae. *MNRAS* 513 (3): 4057–4073. doi:10.1093/mnras/stac1121. 2204.09419.
- Suzuki A and Maeda K (2017), Apr. Supernova ejecta with a relativistic wind from a central compact object: a unified picture for extraordinary supernovae. *MNRAS* 466 (3): 2633–2657. doi:10.1093/mnras/stw3259. 1612.03911.
- Suzuki A, Moriya TJ and Takiwaki T (2020), Aug. A Systematic Study on the Rise Time-Peak Luminosity Relation for Bright Optical Transients Powered by Wind Shock Breakout. *ApJ* 899 (1), 56. doi:10.3847/1538-4357/aba0ba. 2006.16494.
- Suzuki A, Nicholl M, Moriya TJ and Takiwaki T (2021), Feb. Extremely Energetic Supernova Explosions Embedded in a Massive Circumstellar Medium: The Case of SN 2016aps. *ApJ* 908 (1), 99. doi:10.3847/1538-4357/abd6ce. 2012.13283.
- Taddia F, Sollerman J, Fremling C, Barbarino C, Karamehmetoglu E, Arcavi I, Cenko SB, Filippenko AV, Gal-Yam A, Hiramatsu D, Hosseinzadeh G, Howell DA, Kulkarni SR, Laher R, Lunnan R, Masci F, Nugent PE, Nyholm A, Perley DA, Quimby R and Silverman JM (2019), Jan. Analysis of broad-lined Type Ic supernovae from the (intermediate) Palomar Transient Factory. *A&A* 621, A71. doi:10.1051/0004-6361/201834429. 1811.09544.
- Taggart K and Perley DA (2021), May. Core-collapse, superluminous, and gamma-ray burst supernova host galaxy populations at low redshift: the importance of dwarf and starbursting galaxies. *MNRAS* 503 (3): 3931–3952. doi:10.1093/mnras/stab174. 1911.09112.
- Takahashi K, Yoshida T and Umeda H (2018), Apr. Stellar Yields of Rotating First Stars. II. Pair-instability Supernovae and Comparison with Observations. *ApJ* 857 (2), 111. doi:10.3847/1538-4357/aab95f. 1803.06630.
- Tanaka M, Moriya TJ, Yoshida N and Nomoto K (2012), May. Detectability of high-redshift superluminous supernovae with upcoming optical and near-infrared surveys. *MNRAS* 422 (3): 2675–2684. doi:10.1111/j.1365-2966.2012.20833.x. 1202.3610.
- Tanaka M, Moriya TJ and Yoshida N (2013), Nov. Detectability of high-redshift superluminous supernovae with upcoming optical and near-infrared surveys - II. Beyond  $z = 6$ . *MNRAS* 435 (3): 2483–2493. doi:10.1093/mnras/stt1469. 1306.3743.
- Tartaglia L, Pastorello A, Sollerman J, Fransson C, Mattila S, Fraser M, Taddia F, Tomasella L, Turatto M, Morales-Garoffolo A, Elias-Rosa N, Lundqvist P, Harmanen J, Reynolds T, Cappellaro E, Barbarino C, Nyholm A, Kool E, Ofek E, Gao X, Jin Z, Tan H, Sand DJ, Ciabattari F, Wang X, Zhang J, Huang F, Li W, Mo J, Rui L, Xiang D, Zhang T, Hosseinzadeh G, Howell DA, McCully C, Valenti S, Benetti S, Callis E, Carracedo AS, Fremling C, Kangas T, Rubin A, Somero A and Terreran G (2020), Mar. The long-lived Type II<sub>n</sub> SN 2015da: Infrared echoes and strong interaction within an extended massive shell. *A&A* 635, A39. doi:10.1051/0004-6361/201936553. 1908.08580.
- Tolstov A, Nomoto K, Blinnikov S, Sorokina E, Quimby R and Baklanov P (2017), Feb. Pulsational Pair-instability Model for Superluminous Supernova PTF12dam: Interaction and Radioactive Decay. *ApJ* 835 (2), 266. doi:10.3847/1538-4357/835/2/266. 1612.01634.
- Umeda H and Nomoto K (2008), Feb. How Much  $^{56}\text{Ni}$  Can Be Produced in Core-Collapse Supernovae? Evolution and Explosions of 30-100  $M_{\odot}$  Stars. *ApJ* 673 (2): 1014–1022. doi:10.1086/524767. 0707.2598.
- Venditti A, Bromm V, Finkelstein SL, Graziani L and Schneider R (2024), Jan. The first fireworks: A roadmap to Population III stars during the epoch of reionization through pair-instability supernovae. *MNRAS* 527 (3): 5102–5116. doi:10.1093/mnras/stad3513. 2306.06501.
- Vigna-Gómez A, Justham S, Mandel I, de Mink SE and Podsiadlowski P (2019), May. Massive Stellar Mergers as Precursors of Hydrogen-rich Pulsational Pair Instability Supernovae. *ApJL* 876 (2), L29. doi:10.3847/2041-8213/ab1bdf. 1903.02135.
- Villar VA, Nicholl M and Berger E (2018), Dec. Superluminous Supernovae in LSST: Rates, Detection Metrics, and Light-curve Modeling. *ApJ*

- 869 (2), 166. doi:10.3847/1538-4357/aace6a. 1809.07343.
- Vreeswijk PM, Savaglio S, Gal-Yam A, De Cia A, Quimby RM, Sullivan M, Cenko SB, Perley DA, Filippenko AV, Clubb KI, Taddia F, Sollerman J, Leloudas G, Arcavi I, Rubin A, Kasliwal MM, Cao Y, Yaron O, Tal D, Ofek EO, Capone J, Kutyrev AS, Toy V, Nugent PE, Laher R, Surace J and Kulkarni SR (2014), Dec. The Hydrogen-poor Superluminous Supernova iPTF 13ajg and its Host Galaxy in Absorption and Emission. *ApJ* 797 (1), 24. doi:10.1088/0004-637X/797/1/24. 1409.8287.
- Vreeswijk PM, Leloudas G, Gal-Yam A, De Cia A, Perley DA, Quimby RM, Waldman R, Sullivan M, Yan L, Ofek EO, Fremling C, Taddia F, Sollerman J, Valenti S, Arcavi I, Howell DA, Filippenko AV, Cenko SB, Yaron O, Kasliwal MM, Cao Y, Ben-Ami S, Horesh A, Rubin A, Lunnan R, Nugent PE, Laher R, Rebbapragada UD, Woźniak P and Kulkarni SR (2017), Jan. On the Early-time Excess Emission in Hydrogen-poor Superluminous Supernovae. *ApJ* 835 (1), 58. doi:10.3847/1538-4357/835/1/58. 1609.08145.
- Vurm I and Metzger BD (2021), Aug. Gamma-Ray Thermalization and Leakage from Millisecond Magnetar Nebulae: Toward a Self-consistent Model for Superluminous Supernovae. *ApJ* 917 (2), 77. doi:10.3847/1538-4357/ac0826. 2101.05299.
- Wang SQ, Wang LJ, Dai ZG and Wu XF (2015), Jan. Superluminous Supernovae Powered by Magnetars: Late-time Light Curves and Hard Emission Leakage. *ApJ* 799 (1), 107. doi:10.1088/0004-637X/799/1/107. 1411.5126.
- Wang S, Liu J, Qiu Y, Bai Y, Yang H, Guo J and Zhang P (2016), Jun. CHANDRA ACIS Survey of X-Ray Point Sources: The Source Catalog. *ApJS* 224 (2), 40. doi:10.3847/0067-0049/224/2/40. 1603.08353.
- Wei JJ, Wu XF and Melia F (2015), May. Testing Cosmological Models with Type Ic Super Luminous Supernovae. *AJ* 149 (5), 165. doi:10.1088/0004-6256/149/5/165. 1503.06378.
- Wiseman P, Smith M, Childress M, Kelsey L, Möller A, Gupta RR, Swann E, Angus CR, Brout D, Davis TM, Foley RJ, Frohmaier C, Galbany L, Gutiérrez CP, Inserra C, Kessler R, Lewis GF, Lidman C, Macaulay E, Nichol RC, Pursiainen M, Sako M, Scolnic D, Sommer NE, Sullivan M, Tucker BE, Abbott TMC, Agüena M, Allam S, Avila S, Bertin E, Brooks D, Buckley-Geer E, Burke DL, Carnero Rosell A, Carollo D, Carrasco Kind M, da Costa LN, De Vicente J, Desai S, Diehl HT, Doel P, Eifler TF, Everett S, Fosalba P, Frieman J, García-Bellido J, Gaztanaga E, Gerdes DW, Gill MSS, Glazebrook K, Gruendl RA, Gschwend J, Hartley WG, Hinton SR, Hollowood DL, Honscheid K, James DJ, Kuehn K, Kurupatkin N, Lima M, Maia MAG, March M, Martini P, Melchior P, Menanteau F, Miquel R, Ogando RLC, Paz-Chinchón F, Plazas AA, Romer AK, Roodman A, Sanchez E, Scarpine V, Serrano S, Suchyta E, Swanson MEC, Tarle G, Thomas D, Tucker DL, Varga TN, Walker AR, Wilkinson RD and (DES Collaboration) (2020), Jul. Supernova host galaxies in the dark energy survey: I. Deep coadds, photometry, and stellar masses. *MNRAS* 495 (4): 4040–4060. doi:10.1093/mnras/staa1302. 2001.02640.
- Woodsley SE (2010), Aug. Bright Supernovae from Magnetar Birth. *ApJL* 719 (2): L204–L207. doi:10.1088/2041-8205/719/2/L204. 0911.0698.
- Woodsley SE (2017), Feb. Pulsational Pair-instability Supernovae. *ApJ* 836 (2), 244. doi:10.3847/1538-4357/836/2/244. 1608.08939.
- Woodsley SE, Blinnikov S and Heger A (2007), Nov. Pulsational pair instability as an explanation for the most luminous supernovae. *Nature* 450 (7168): 390–392. doi:10.1038/nature06333. 0710.3314.
- Yan L, Lunnan R, Perley DA, Gal-Yam A, Yaron O, Roy R, Quimby R, Sollerman J, Fremling C, Leloudas G, Cenko SB, Vreeswijk P, Graham ML, Howell DA, De Cia A, Ofek EO, Nugent P, Kulkarni SR, Hosseinzadeh G, Masci F, McCully C, Rebbapragada UD and Woźniak P (2017a), Oct. Hydrogen-poor Superluminous Supernovae with Late-time  $H\alpha$  Emission: Three Events From the Intermediate Palomar Transient Factory. *ApJ* 848 (1), 6. doi:10.3847/1538-4357/aa8993. 1704.05061.
- Yan L, Quimby R, Gal-Yam A, Brown P, Blagorodnova N, Ofek EO, Lunnan R, Cooke J, Cenko SB, Jencson J and Kasliwal M (2017b), May. Far-ultraviolet to Near-infrared Spectroscopy of a Nearby Hydrogen-poor Superluminous Supernova Gaia16apd. *ApJ* 840 (1), 57. doi:10.3847/1538-4357/aa6b02. 1611.02782.
- Yan L, Perley DA, De Cia A, Quimby R, Lunnan R, Rubin KHR and Brown PJ (2018), May. Far-UV HST Spectroscopy of an Unusual Hydrogen-poor Superluminous Supernova: SN2017egm. *ApJ* 858 (2), 91. doi:10.3847/1538-4357/aabad5. 1711.01534.
- Yan L, Perley DA, Schulze S, Lunnan R, Sollerman J, De K, Chen ZH, Fremling C, Gal-Yam A, Taggart K, Chen TW, Andreoni I, Bellm EC, Cunningham V, Dekany R, Duev DA, Fransson C, Laher RR, Hankins M, Ho AYQ, Jencson JE, Kaye S, Kulkarni SR, Kasliwal MM, Golkhou VZ, Graham M, Masci FJ, Miller AA, Neill JD, Ofek E, Porter M, Mróz P, Reiley D, Riddle R, Rigault M, Rusholme B, Shupe DL, Soumagnac MT, Smith R, Tartaglia L, Yao Y and Yaron O (2020), Oct. Helium-rich Superluminous Supernovae from the Zwicky Transient Facility. *ApJL* 902 (1), L8. doi:10.3847/2041-8213/abb8c5. 2006.13758.
- Yu YW and Li SZ (2017), Sep. A possible relation between flare activity in super-luminous supernovae and gamma-ray bursts. *MNRAS* 470 (1): 197–201. doi:10.1093/mnras/stx1028. 1607.00626.
- Zhao WC, Xue XX and Cao XF (2021), Feb. On the event rate and luminosity function of superluminous supernovae. *New Astron.* 83, 101506. doi:10.1016/j.newast.2020.101506.