

Bounding elastic photon-photon scattering at $\sqrt{s} \approx 1$ MeV using a laser-plasma platform

R. Watt^a, B. Kettle^a, E. Gerstmayr^a, B. King^b, A. Alejo^c, S. Astbury^h, C. Bairdⁱ, S. Bohlen^d, M. Campbell^p, C. Colgan^a, D. Dannheim^p, C. Gregory^h, H. Harsh^{f,g}, P. Hatfield^l, J. Hinojosa^c, D. Hollatz^{f,g}, Y. Katzir^h, J. Morton^k, C.D. Murphy^l, A. Nurnberg^p, J. Osterhoff^l, G. Pérez-Callejo^m, K. Pöder^d, P.P. Rajeev^h, C. Roedel^f, F. Roeder^g, F.C. Salgado^{f,g}, G.M. Samarin^e, G. Sarri^e, A. Seidel^{f,g}, C. Spindloe^h, S. Steinkeⁿ, M.J.V. Streeter^{e,o}, A.G.R. Thomas^{c,o}, C. Underwood^l, W. Wu^a, M. Zepf^{f,g}, S.J. Rose^a, S.P.D. Mangles^a

^aThe John Adams Institute for Accelerator Science, Imperial College London, London, SW7 2AZ, UK,

^bCentre for Mathematical Sciences, University of Plymouth, PL4 8AA, UK,

^cCenter for Ultrafast Optical Science, University of Michigan, Ann Arbor, MI 48109-2099, USA,

^dDeutsches Elektronen-Synchrotron DESY, Notkestr. 85, 22607 Hamburg, Germany,

^eSchool of Mathematics and Physics, Queen's University of Belfast, BT7 1NN, Belfast, UK,

^fHelmholtz Institut Jena, Fröbelstieg 3, 07743 Jena, Germany,

^gInstitut für Optik und Quantenelektronik, Friedrich-Schiller-Universität, 07743 Jena, Germany,

^hCentral Laser Facility, STFC Rutherford Appleton Laboratory, Didcot OX11 0QX, UK,

ⁱDepartment of Physics, University of York, York, YO10 5DD, UK,

^jDepartment of Physics, University of Oxford, Oxford, OX1 3PU, UK,

^kAWE Aldermaston, Reading RG7 4PR, UK,

^lUniversité de Bordeaux-CNRS-CEA, Centre Lasers Intenses et Applications (CELIA), UMR 5107, F-33405 Talence, France,

^mDepartamento de Física Teórica Atómica y Óptica, Universidad de Valladolid, 47011 Valladolid, Spain,

ⁿLawrence Berkeley National Laboratory, Berkeley, CA 94720, US,

^oPhysics Department, Lancaster University, Lancaster LA1 4YB, UK,

^pCERN, Geneva, Switzerland,

Abstract

We report on a direct search for elastic photon-photon scattering using x-ray and γ photons from a laser-plasma based experiment. A gamma photon beam produced by a laser wakefield accelerator provided a broadband gamma spectrum extending to above $E_\gamma = 200$ MeV. These were collided with a dense x-ray field produced by the emission from a laser heated germanium foil at $E_x \approx 1.4$ keV, corresponding to an invariant mass of $\sqrt{s} = 1.22 \pm 0.22$ MeV. In these asymmetric collisions elastic scattering removes one x-ray and one high-energy γ photon and outputs two lower energy γ photons. No changes in the γ photon spectrum were observed as a result of the collisions allowing us to place a 95% upper bound on the cross section of 1.5×10^{15} μb . Although far from the QED prediction, this represents the lowest upper limit obtained so far for $\sqrt{s} \lesssim 1$ MeV.

1. Introduction

Photon-photon scattering is one of the most fundamental processes in quantum electrodynamics (QED) and is of elementary importance in astrophysics. It is used in models that calculate primordial abundances, affects the observed spectra from γ -ray bursts from the first million years of the universe [1, 2] and plays an important role in models of the evolution of strongly magnetised neutron stars [3]. However, these calculations all use the QED cross section which is currently poorly bounded by experiment.

Photon-photon scattering involving virtual photons has previously been observed in several forms (see the summary in Tab. 1): the 1-to-1 process of Delbrück scattering ($\gamma\gamma^* \rightarrow \gamma\gamma^*$), where a real photon, γ , scatters from a virtual photon, γ^* , in the Coulomb field of an ion [4, 5, 6]; the 1-to-2 process of photon splitting ($\gamma\gamma^* \rightarrow \gamma\gamma$) in atomic fields [4, 7], and the 0-to-2 process of real double photon emission from colliding the virtual photons from Coulomb fields in ultra-peripheral heavy-ion collisions at the ATLAS and CMS experiments ($\gamma^*\gamma^* \rightarrow \gamma\gamma$) [8, 9, 10]. Instead, in this paper, we will report on a search for

the 2-to-2 process of photon-photon scattering involving only real photons ($\gamma\gamma \rightarrow \gamma\gamma$).




Table 1: Comparison of different photon-photon scattering processes. The top row shows the Feynman diagram, the second row shows the number of real incoming and outgoing photons (N -to- M) and the third row shows name of the relevant inelastic contribution (i.e. the real electron-positron pair creation process described by the left-hand side of dotted line in the Feynman diagram). In this paper we report results on bounding the cross-section of 2-to-2 scattering.

A crucial parameter in photon-photon collisions is the invariant mass of the collision \sqrt{s} . For two photons, energy E_1 and E_2 colliding at an angle ϕ , we have $s = 2E_1E_2(1 - \cos \phi)$. Photon-photon scattering has been searched for indirectly in the signal of vacuum birefringence at small invariant mass $\sqrt{s} \ll m_e c^2$ in cavity experiments such as PVLAS [14] and BMV [15] involving photons traversing a quasi-constant magnetic field.

At invariant mass $\sqrt{s} \sim O(\text{eV})$ it has also been searched for by directly colliding two optical laser pulses [16] and three optical laser pulses [17, 18]. At $\sqrt{s} \sim O(\text{keV})$, the cross-section has been bounded by experiments employing x-ray free electron lasers [19, 20]. By comparison, for large invariant mass $\sqrt{s} \gg m_e c^2$, photon scattering with quasi-real photons has been measured by the ATLAS and CMS experiments (in which $\sqrt{s} \approx 5 - 20 \text{ GeV}$). Despite these results, photon-photon scattering has yet to be measured using manifestly real photons and this has sustained interest in the process. The upcoming HIBEF experiment plans to provide the first measurement involving manifestly real photons at $\sqrt{s} \sim O(10^2 \text{ eV})$ by colliding an x-ray free electron laser with a high power optical laser pulse [21, 22] and there have been many suggestions for how to measure this effect using only PW-class optical lasers [23, 24, 25, 26].

Apart from being a test of fundamental QED, searches for photon-photon scattering can also provide bounds on physics beyond the Standard Model (BSM), e.g. the ATLAS results enabled bounds on Born-Infeld electrodynamics [27] for energy scales $> 100 \text{ GeV}$. Suggestions for improving these bounds by measuring photon-photon scattering at future colliders have also recently appeared in the literature [28].

This paper reports on a search for elastic photon-photon scattering at $\sqrt{s} \approx 1 \text{ MeV}$. The search is motivated at this energy scale because it is relevant to astrophysics, it is where the cross-section takes its maximum value, and at this energy, the QED effect has only very weak experimental bounds. This scale also includes energies over the threshold for creation of real electron-positron pairs via the linear Breit-Wheeler process, which therefore is a sub-process of photon-photon scattering at these energies (this can be understood by the optical theorem [29]). Linear Breit-Wheeler is also being searched for at PW-class optical lasers [30, 31], and multi-photon Breit-Wheeler pair-creation forms part of the science goals for the LUXE experiment planned at DESY [32] and the E320 experiment at SLAC [33].

Our experiment uses $\sim 1 \text{ GeV}$ electrons from a laser wakefield accelerator [34] which collide with a fixed target to generate a broadband bremsstrahlung distribution of γ photons extending to $E_\gamma \approx 800 \text{ MeV}$. The γ photons are then collided with the dense x-ray photon field in the vicinity of a laser heated germanium foil. The x-ray radiation is dominated by M-L-band emission in the region of $E_x \approx 1.4 \text{ keV}$. The spectrum of the γ photons is determined using a caesium-iodide stack spectrometer [35, 36].

In this asymmetric collision, a scattering event typically removes one x-ray and one high energy γ photon from the beam and replaces it with two lower energy γ photons. If sufficient numbers of γ photons were to scatter in the x-ray field, we would therefore be able to detect the effect in the photon energy spectrum. This is illustrated in Fig. 1 for the idealised case of a narrow energy spread 500 MeV γ photon beam scattering from a dense x-ray field. If however, insufficient scattering events occur to produce a measurable change in the spectrum, this allows us to place a limit on the photon-photon scattering cross-section.




Figure 1: Illustration of the effect on the photon energy spectrum of copious photon-photon scattering of a 500 MeV γ photon beam in a dense 1.5 keV x-ray field.

As well as the effect on the photon spectrum, scattering also leads to an increase in the divergence of the photon beam. Due to the asymmetry in our photon energies very few photons will be scattered outside of the original γ beam profile and so, for our set-up, the effect on the spectrum is more pronounced. We therefore use the energy spectrum diagnostic only.

The QED prediction was calculated by extending the Geant4 simulation framework [37, 38] to include the cross-section for photon-photon scattering [39, 40, 41, 42, 43]. Simulation results were compared to an independent direct numerical evaluation of the process integrated over the x-ray and γ -distributions reported in experiment, and found to be in agreement. The extended Geant4 model simulates the leading order (in fine-structure constant α) photon-photon scattering contribution involving four photons¹. This complements other recently developed simulation frameworks [45, 46, 47, 48, 49] of photon-photon scattering at $\sqrt{s} \sim O(\text{eV}) - O(\text{keV})$ based on the weak-field Heisenberg-Euler Lagrangian.

2. Experimental set-up

The experiment took place at the Gemini laser facility in the UK. This is a dual beam, 300 TW Ti:Sa system, allowing us to generate and collide two high energy-density photon sources. The experimental setup was based on the scheme by Pike *et al.*, (2014) with asymmetrical photon sources [50]. The γ photon source was provided by bremsstrahlung emission produced by an electron beam in a high Z-material target. The electron beam was produced using laser wakefield acceleration [51]. The x-ray photon source was generated through direct laser heating of a thin metal foil. The two photon beams were temporally overlapped at the interaction point within 2 picoseconds using the drive laser beams (same optical path) and a fast-response photodiode. A schematic of the experimental setup can be found in figure 2. A detailed description of our laser-plasma platform for photon-photon physics can be found in ref [30]. The experiment can be separated into three parts: the x-ray photon source, the γ photon source, and the γ photon spectrometer

¹During preparation of the manuscript, another simulation framework was developed that includes the same process [44].




Figure 2: Schematic of the experimental set-up (not to scale). γ -rays are produced using a laser wakefield accelerator, they are collimated and pass through the x-ray field in the vicinity of a laser-heated plasma. The spectrum of γ -rays after passage through the x-ray field is characterised using an array of caesium-iodide crystals.

2.1. x-ray photon source

One of the Gemini laser pulses was used to generate a dense x-ray field by rapidly heating a 100 nm germanium (Ge) foil. As this solid Ge foil is heated, it turns into a plasma, leading to the emission of intense x-ray radiation predominantly due to M-L band transitions [52, 53]. The heating laser pulse had a duration of 40 ps (FWHM intensity) and a total energy of 10.7 ± 0.3 J. It was focused to an elliptical spot using a distributive phase plate, with major and minor axes of (217 ± 6) μm and (77 ± 6) μm respectively, which contained 72% of the total energy. The Ge targets were mounted on a Kapton ($\text{C}_{22}\text{H}_{10}\text{N}_2\text{O}_5$) tape with a lower average atomic number, limiting the mass of Ge close to the interaction which is a potential noise source. A motorised tape-drive was used to change targets between shots.

To diagnose the x-ray field, a pinhole imaging system and crystal spectrometer were used. The pinhole imaging system gave an on-shot measure of both the emission spot size and the target alignment. The spectrometer used a flat, thallium acid phthalate (TIAP) crystal, with a spectral window of ≈ 700 eV, centred at approximately 1.6 keV, signal above approximately 1.5 keV was suppressed due to an aluminium filter. This spectral window is around the M-L band transitions of Ge.

It was found that the total conversion efficiency from laser energy to 1.3–1.5 keV x-rays was $(2.4 \pm 0.3)\%$. This corresponds to $(3.7 \pm 0.4) \times 10^{10}$ photons $\text{eV}^{-1} \text{J}^{-1} \text{srad}^{-1}$ emitted normal to the front surface of the germanium target. Taking into account the absorption in the kapton layer on the rear side of the target, at the interaction region (1 mm from the tape) this corresponds to an x-ray photon density of $(1.4 \pm 0.5) \times 10^{12} \text{mm}^{-3}$, over an effective length of approximately 3 mm.

2.2. γ photon source

The γ photon source was generated through bremsstrahlung emission, which first requires a beam of high energy electrons. To produce these electrons, one of the Gemini laser beams was focused into a 17.5 mm gas cell filled with helium and a 2% nitrogen dopant. The duration of the laser pulse was 45 ± 5 fs (FWHM intensity) and the focal spot was $(44 \pm 2) \mu\text{m} \times (53 \pm 2) \mu\text{m}$ (FWHM intensity). The laser energy on target was 5.5 ± 0.6 J, corresponding to a normalised vector potential $a_0 = 1.1 \pm 0.2$. Through the laser wakefield acceleration mechanism [34], a beam of high energy electrons (energy up to ≈ 800 MeV, charge ≈ 50 pC) was emitted from the gas cell. These electrons then

passed through a 0.5 mm thick bismuth (Bi) foil, acting as a bremsstrahlung converter. This emits a beam of high energy γ photons with a similar duration to that of the driving laser pulse.

Bremsstrahlung converters also generate a large number of low energy, divergent γ photons, which are a potential noise source. If these divergent γ photons were to interact with the Ge foil, or another part of the experimental setup, they would produce background through the Compton scattering process. To prevent this a 100 mm block of tungsten (W) with a 2 mm diameter hole drilled through the centre was used to collimate the γ photon beam. This collimator effectively removes any γ photon travelling at an angle greater than 10 mrad to the beam propagation direction.

To reduce the background further, a 50 mm tungsten block was placed just off-axis, shadowing the Ge foil from the γ photon beam. Placing such a high Z-material close the γ photon beam axis will itself generate a large number of background Bethe–Heitler pairs which would pass through the x-ray field and create a background signal through Compton scattering. Therefore, a 30 cm dipole magnet with a field strength of $B = 1$ T, was used to remove these before the photon-photon interaction zone. A CsI crystal screen was placed before the γ spectrometer to detect the footprint of the γ beam, but is not used for this analysis.

2.3. γ photon spectrometer

The γ photon spectrometer consisted of a stack of 33×47 CsI crystals doped with thallium and imaged onto a 14-bit EMCCD camera. This was positioned inside the γ photon beam path, with higher energy γ photons penetrating deeper into the stack than lower energy γ photons. The γ -photon spectrum is determined using a trial function based forward model. Bayesian inference is used to determine the best fit parameters of the trial function and their uncertainty.

The detector was calibrated to remove systematic effects such as variations in light yield in each crystal due to crystal imperfections and misalignment or effects produced by the imaging system. We compared the measured energy deposited in each crystal averaged over a large number of shots with that predicted in Geant4 using the average electron energy spectrum (as measured on a series of shots without the bismuth foil intercepting the electron beam) and generated a correction factor which can then be applied to each column of crystals.

By running Geant4 simulations for a series of mono-energetic γ photon beams over a range of energies we can model response of each crystal as a function of photon energy, $\rho_i(E_\gamma)$, where i is the crystal index and E_γ is the photon energy. From this the signal measured by the detector for an arbitrary γ photon spectrum can be quickly calculated with the following integral

$$I_i[f] = C_i \int_0^\infty \rho_i(E) f(E) dE \quad (1)$$

where C_i is the correction factor.

To enable a Bayesian inference of the γ photon spectrum on each shot, we first find a low dimensional parameterisation of the spectrum, $f(E)$. We performed Geant4 simulations

of the bremsstrahlung converter, using electron energy spectra observed on the experiment, to get a data set of typical γ photon spectra. From this data set, the following function was found to provide a good approximation to the spectra measured on the detector.

$$f(E; \alpha, E_c) = \alpha \left(1 - 0.18^2 \frac{E^2}{E_c^2}\right) E^{-0.94}, \quad (2)$$

where α controls the amplitude of the spectrum and E_c is a characteristic energy that controls both the slope and the position of the cut off at high energy. Using this parameterisation, we can obtain the γ photon spectrum on each shot by applying Bayesian inference to estimate a distribution over $\mathbf{x} = (E_c, \alpha)$. This involves applying Bayes' theorem

$$p(\mathbf{x} | \mathbf{y}) = \frac{p(\mathbf{y} | \mathbf{x})p(\mathbf{x})}{\int p(\mathbf{y} | \mathbf{x})p(\mathbf{x})d\mathbf{x}}, \quad (3)$$

where \mathbf{y} is an observed data point, corresponding to a vector of the crystal responses. In this equation $p(\mathbf{y} | \mathbf{x})$ is the likelihood and $p(\mathbf{x}) = p(E_c)p(\alpha)$ is a prior which we must set. If we make the assumption that the crystals exhibit random Gaussian noise, σ , we can write the likelihood function as

$$p(\mathbf{y} | \mathbf{x}) = \prod_i \frac{1}{\sigma \sqrt{2\pi}} \exp\left(-\frac{(y_i - I_i[f(\mathbf{x})])^2}{\sigma^2}\right).$$

This introduces a new parameter, σ , which is treated in the same way as E_c and α . Given that we have little prior knowledge of E_c , α or σ , other than the fact that they cannot be negative, we set uniform priors on each with a lower bound of zero. We know the upper bound for E_c cannot be greater than the maximum energy of the electrons (≈ 800 MeV) so the prior used is $p(E_c) = \mathcal{U}(0, 800\text{MeV})$. Through appropriate normalisation of the data set, we can ensure that α is never greater than 10, allowing us to apply the prior $p(\alpha) = \mathcal{U}(0, 10)$. Finally, we set the prior on σ to be $p(\sigma) = \mathcal{U}(0, 1)$ as if the limit is greater than this, the data will be too noisy to make any inference.

With the likelihood and priors set, we can use equation 3 to calculate the posterior. Given that the numerator involves a three-dimensional integral, it is most efficiently solved using a Markov chain Monte Carlo (MCMC) method. In figure 3 we can see an example of this calculation performed on a randomly selected shot from the data set.

3. Results

3.1. γ photon spectrum

Having developed a robust method for extracting the γ photon spectrum from the crystal response, we can test if the presence of the x-ray field has an effect on the γ photon spectrum. To do this, we run the Bayesian spectral retrieval algorithm on each shot of the experiment and compare the distributions over E_c for null and collision shots. Null shots involve firing only the beam that generates the γ photon beam. Collision shots involve firing both beams at a relative delay that ensures the γ photons pass through the x-ray field. The Ge foil was properly aligned




Figure 3: a) Gamma photon spectrum (Geant4 simulations based on measured electron beam) b) Measured x-ray photon spectrum c) Measured signal on the CsI array from a randomly selected null shot (red circles). The CsI signal generated by the forward model of the detector for the average (black) and 100 randomly selected samples (grey) samples from the posterior distribution of E_c . The mean value corresponds to $E_c = 83.2 \pm 3.2$ MeV.




Figure 4: Distribution of inferred value for the γ photon spectrum E_c for 32 null shots (blue) and 22 collision shots (red)

for both null and collision shots to ensure that any contribution to the γ photon spectrum measurement due to interactions with the foil are fully accounted for. The data set consists of 32 null shots and 22 collision shots. These shots were all performed on a single shot day on the Gemini laser system.

We compare the distribution of E_c on collision and null shots in various ways. Figure 4 shows the histogram of the inferred value for the γ photon spectrum E_c . The relatively small number of shots in each distribution means it is not immediately clear if differences between them are significant. To investigate this, figure 5 shows the distribution of mean and standard deviation calculated from 100,000 bootstrap samples of the null and collision shot data. The fact that the distributions overlap illustrates further that there is no significant difference between the distribution of E_c on null and collision shots.

Figure 6 shows the empirical cumulative distribution function (ecdf) of E_c for the data. Also shown are 50 ecdfs for bootstrap samples of the data, these effectively represent the uncertainty in the ecdfs. The overlap between these ecdfs again illustrates that there is no significant difference between the distributions.

A third method to assess differences in the distribution of E_c




Figure 5: Bootstrap estimate of (a) the mean and (b) the standard deviation of E_c for null shots (blue) and collision shots (red)




Figure 6: The empirical cumulative distribution (ecdf) function of E_c for null shots (blue) and collision shots (red). The thick lines show the ecdf for the experimental data, the thin lines show ecdfs calculated from 50 bootstrap samples of the data.

is to use the two-sample Kolmogorov–Smirnov (KS) test. The null hypothesis of this test is that both null and full data sets have been sampled from the same distribution, i.e. that there is no measurable difference effect of the photon-photon collisions on the measured γ -ray spectrum. To perform the KS test, the two-sample KS-test statistic must be calculated:

$$D = \sup |F_N(E_c) - F_C(E_c)|, \quad (4)$$

where $F_N(E_c)$ and $F_C(E_c)$ are the cumulative distribution functions for the null and collision shots respectively. The null hypothesis is accepted at the 95% confidence level if $D < 0.378$ [54] for a data sets with 32 and 22 samples. The value obtained for our data set is $D = 0.216$, so we cannot reject the null hypothesis.

We can also calculate the two-sample Kolmogorov-Smirnov test statistic for a large number of bootstrap samples from the data to estimate the uncertainty in the KS test statistic. This is shown in figure 7. It shows that the bulk of the distribution ($\approx 90\%$) lies below the critical value, providing further strong evidence that we cannot reject the null hypothesis, i.e. we must assume that collisions between γ photons and the dense x-ray field did not produce a detectable difference in the energy spectrum of the γ photons.

3.2. Bounding the cross section

As the various analyses all show that there is no significant difference between the distribution over E_c on null and collision shots, we can conclude there was not a detectable level




Figure 7: Distribution of the two sample Kolmogorov-Smirnov test statistic comparing the distribution of E_c for null shots and collision shots from 100,000 bootstrap samples from the data. The red line shows the value above which the null hypothesis (that the null and collision shots are from the same distribution) can be rejected at the $\alpha = 0.05$ significance level.

of photon-photon scattering. To find how much larger than the standard QED value the cross-section would have to be to produce a detectable, we performed multiple simulations of the experiment with an increasing cross-section. The factor by which the cross section would have to increase for us to have observed a significant difference in the value of E_c on collision shots provides a bound on the cross-section. These simulations were performed with Geant4 [37] which we have adapted to include photon-photon collisions between the γ photons and a dense x-ray field, using the QED cross section [38].

The simulation modelled all major aspects of the experimental geometry, including the γ -photon source, collimator and magnet before the collision point, the x-ray photon source (both the tape target and surrounding x-ray field), and the magnetic transport system, shielding and detectors and after the collision point. The x-ray photon source was modelled as a static (i.e. non evolving) photon field using the measured x-ray spectrum and a spatial distribution of photon density $n_x(x, y, z)$ calculated from the experimentally measured photon numbers and assuming emission from a uniform disk of radius 100 nm. Details of the geometry can be found in [30], and details of the modifications made to Geant4 are described in [38].

The result of these simulations is shown in figure 8. Also shown is the mean and 95% confidence limit of E_c for the null shots. Increasing the bias on the cross-section up to 10^{13} has little effect on the E_c our detector would measure. Beyond this point, E_c starts to decrease. The copious amount of elastic photon-photon scattering that would occur if the cross section were $10^{14} - 10^{15}$ times higher than the QED prediction would significantly lower the average energy of the γ photons exiting the collision volume. The simulations show that this would result in a measurably lower value of E_c on collision shots than that measured on null shots. The fact that we do not measure a lower value of E_c on collision shots therefore allows us to place an upper bound on the elastic photon-photon scattering cross section at $\approx 10^{15} \sigma_{\text{QED}}$.

The broadband nature of the photon spectra in this exper-




Figure 8: Simulated γ photon spectrum critical energy, E_c , after interaction with the x-ray field, as a function of the cross section bias factor. Blue shaded region represents the critical energy measured on null shots (95% confidence interval).

iment means that this measurement is not at a single, specified value of \sqrt{s} , but the effective \sqrt{s} be found by weighting the cross section, $\sigma(E_1, E_2)$, with the measured photon spectra $N_x(E_1)$ and $N_y(E_2)$ and considering the range of collision angles. The effective \sqrt{s} for this experiment is 1.22 ± 0.22 MeV.

4. Conclusions

The cross-section limits that have been made by previous direct searches for elastic photon-photon scattering are shown in figure 9. The closest of these to the QED cross-section for real photon-photon scattering is that of Bernard et al. (2000) [18] using optical photons. However, this is a factor of 10^{18} times higher than the QED prediction. More recently, work using x-ray photons provided by a free electron laser bounded the cross section at $\sqrt{s} \sim 10^{-2}m_e$ a factor of 10^{19} times higher than the QED prediction [19, 20]. These high bounds are due to the fact that these previous direct searches operated in a regime where $\sqrt{s} \ll m_e$, where the cross-section is severely suppressed. The experiment reported here provides the first bound at $\sqrt{s} \sim m_e$, where elastic scattering is expected to play a role in various astrophysical situations [2]. This experiment also provides the lowest ratio of the upper bound to the QED prediction for 2-to-2 photon-photon scattering to date and the lowest bound in the range close to $\sqrt{s} \approx 1$ MeV.

While this current work provides an upper bound on the cross section, it is also useful to consider if laser-plasma interactions are a potential route to directly observing photon-photon collisions in the laboratory. To do this we consider how long an experiment would have to operate to observe a single scatter event. A simple estimate of the number of scatter events per shot is $N_{\text{scatter}} \approx N_\gamma \sigma n_x L_x$, where N_γ is the number of γ photons, σ is the cross section, n_x is the x-ray photon density and L_x is the length of the x-ray field. For the current configuration described here $N_\gamma \sim 10^7$, $\sigma \sim 10^{-30}$ cm², $n_x \sim 10^{15}$ cm⁻³, and $L_x \sim 0.1$ cm, resulting in $N_{\text{scatter}} \sim 10^{-9}$ per laser shot. At the repetition rate of this experiment (0.05 Hz) this would require




Figure 9: Comparison of this work with different measurements of the total cross-section of photon-photon scattering. The theory prediction for the cross-section of the 2-to-2 process is plotted in the solid blue line (calculated from [41, 42, 43]). For the 1-to-1 process (also measured in [6]), the horizontal co-ordinate is given by the incident photon energy in the lab. For the 0-to-2 process, the measurements were for a diphoton mass > 5 GeV [9] or > 6 GeV [8, 10], to represent this on the plot, $\sqrt{s} = 10$ GeV has been chosen. Notable exceptions to this plotting scheme are the cavity experiments such as PVLAS [14] and BMV [15], measuring a 1-to-1 process in a quasi-constant magnetic field and photon-splitting experiments [4, 7] measuring a 1-to-2 process.

over 600 years of continuous operation, but a 100 Hz laser with similar capabilities would require only 100 days.

Higher energy lasers such as EPAC [55] and ELI-NP[56] will be capable of producing greater than 10 times more γ photons per shot due to the higher charge, higher energy electron beams they will be capable of producing. If such lasers could be operated at 100 Hz, the required time drops to ~ 1 day.

The development of such high repetition rate, high power lasers has already been identified as a key future direction for laser wakefield accelerators [57] and is an area of active research (see e.g. [58]). Such facilities could open up the real possibility of observing and studying photon-photon scattering.

Data Availability

Data will be made available on request.

Acknowledgement

We wish to acknowledge the support of the staff at the Central Laser Facility. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant Agreement No. 682399) and STFC (Grant No. ST/P002021/1). JH and AT acknowledge support from the US National Science Foundation grant #1804463. GS would like to acknowledge support from EPSRC (Grant Nos. EP/N027175/1, EP/P010059/1). GPC was supported by Research Grant No.

References

- [1] J. Ellis, G. B. Gelmini, J. L. Lopez, D. V. Nanopoulos, S. Sarkar, Astrophysical constraints on massive unstable neutral relic particles, *Nuclear Physics B* 373 (2) (1992) 399–437.
- [2] R. Svensson, A. A. Zdziarski, Photon-photon scattering of gamma rays at cosmological distances, *Astrophys. J.* 349 (1990) 415–428. doi:10.1086/168325.
- [3] M. G. Baring, A. K. Harding, Photon splitting and pair creation in highly magnetized pulsars, *The Astrophysical Journal* 547 (2) (2001) 929.
- [4] G. Jarlskog, L. Joensuu, S. Pruenster, H. D. Schulz, H. J. Willutzki, G. G. Winter, Measurement of delbruck scattering and observation of photon splitting at high energies, *Phys. Rev. D* 8 (1973) 3813–3823. doi:10.1103/PhysRevD.8.3813.
- [5] M. Schumacher, I. Borchert, F. Smend, P. Rullhusen, Delbruck scattering of 2.75 mev photons by lead, *Physics Letters B* 59 (2) (1975) 134–136.
- [6] S. Z. Akhmadaliev, et al., Delbruck scattering at energies of 140-450 MeV, *Phys. Rev. C* 58 (1998) 2844–2850. doi:10.1103/PhysRevC.58.2844.
- [7] S. Z. Akhmadaliev, et al., Experimental investigation of high-energy photon splitting in atomic fields, *Phys. Rev. Lett.* 89 (2002) 061802. arXiv:hep-ex/0111084, doi:10.1103/PhysRevLett.89.061802.
- [8] M. Aaboud, et al., Evidence for light-by-light scattering in heavy-ion collisions with the ATLAS detector at the LHC, *Nature Phys.* 13 (9) (2017) 852–858. arXiv:1702.01625, doi:10.1038/nphys4208.
- [9] A. M. Sirunyan, et al., Evidence for light-by-light scattering and searches for axion-like particles in ultraperipheral PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Lett. B* 797 (2019) 134826. arXiv:1810.04602, doi:10.1016/j.physletb.2019.134826.
- [10] G. Aad, et al., Observation of light-by-light scattering in ultraperipheral Pb+Pb collisions with the ATLAS detector, *Phys. Rev. Lett.* 123 (5) (2019) 052001. arXiv:1904.03536, doi:10.1103/PhysRevLett.123.052001.
- [11] L. D. Landau, E. M. Lifschitz, On the Production of Electrons and Positrons by a Collision of Two Particles, *Phys. Z. Sowjetunion* 6 (1934) 244. doi:10.1016/B978-0-08-010586-4.50021-3.
- [12] H. Bethe, W. Heitler, On the Stopping of fast particles and on the creation of positive electrons, *Proc. Roy. Soc. Lond. A* 146 (1934) 83–112. doi:10.1098/rspa.1934.0140.
- [13] G. Breit, J. A. Wheeler, Collision of two light quanta, *Phys. Rev.* 46 (12) (1934) 1087–1091. doi:10.1103/PhysRev.46.1087.
- [14] A. Ejlli, F. Della Valle, U. Gastaldi, G. Messineo, R. Pengo, G. Russo, G. Zavattini, The PVLAS experiment: A 25 year effort to measure vacuum magnetic birefringence, *Phys. Rept.* 871 (2020) 1–74. arXiv:2005.12913, doi:10.1016/j.physrep.2020.06.001.
- [15] J. Agil, R. Battesti, C. Rizzo, Monte Carlo study of the BMV vacuum linear magnetic birefringence experiment, *Eur. Phys. J. D* 75 (3) (2021) 90. doi:10.1140/epjd/s10053-021-00100-z.
- [16] F. Moulin, D. Bernard, F. Amiranoff, Photon-photon elastic scattering in the visible domain, *Z. Phys. C* 72 (1996) 607–611. doi:10.1007/s002880050282.
- [17] F. Moulin, D. Bernard, Four-wave interaction in gas and vacuum. Definition of a third order nonlinear effective susceptibility in vacuum: $\chi_{vacuum}^{(3)}$, *Opt. Commun.* 164 (1999) 137–144. arXiv:physics/0203069, doi:10.1016/S0030-4018(99)00169-8.
- [18] D. Bernard, F. Moulin, F. Amiranoff, A. Braun, J. P. Chambaret, G. Darpentigny, G. Grillon, S. Ranc, F. Perrone, Search for Stimulated Photon-Photon Scattering in Vacuum, *Eur. Phys. J. D* 10 (2000) 141. arXiv:1007.0104, doi:10.1007/s100530050535.
- [19] T. Inada, et al., Search for Photon-Photon Elastic Scattering in the X-ray Region, *Phys. Lett. B* 732 (2014) 356–359. arXiv:1403.2547, doi:10.1016/j.physletb.2014.03.054.
- [20] T. Yamaji, et al., An experiment of X-ray photon-photon elastic scattering with a Laue-case beam collider, *Phys. Lett. B* 763 (2016) 454–457. arXiv:1608.06369, doi:10.1016/j.physletb.2016.11.003.
- [21] H.-P. Schlenvoigt, T. Heinzl, U. Schramm, T. E. Cowan, R. Sauerbrey, Detecting vacuum birefringence with x-ray free electron lasers and high power optical lasers: a feasibility study, *Phys. Scripta* 91 (2) (2016) 023010. doi:10.1088/0031-8949/91/2/023010.
- [22] F. Karbstein, D. Ullmann, E. A. Mosman, M. Zepf, Direct accessibility of the fundamental constants governing light-by-light scattering, *Phys. Rev. Lett.* 129 (2022) 061802. doi:10.1103/PhysRevLett.129.061802. URL <https://link.aps.org/doi/10.1103/PhysRevLett.129.061802>
- [23] M. Marklund, P. K. Shukla, Nonlinear collective effects in photon-photon and photon-plasma interactions, *Rev. Mod. Phys.* 78 (2006) 591–640. arXiv:hep-ph/0602123, doi:10.1103/RevModPhys.78.591.
- [24] A. Di Piazza, C. Muller, K. Z. Hatsagortsyan, C. H. Keitel, Extremely high-intensity laser interactions with fundamental quantum systems, *Rev. Mod. Phys.* 84 (2012) 1177. arXiv:1111.3886, doi:10.1103/RevModPhys.84.1177.
- [25] B. King, T. Heinzl, Measuring Vacuum Polarisation with High Power Lasers, *High Power Laser Sci. Eng.* 4 (2016). arXiv:1510.08456, doi:10.1017/hpl.2016.1.
- [26] A. Fedotov, A. Ilderton, F. Karbstein, B. King, D. Seipt, H. Taya, G. Torgrimsson, Advances in QED with intense background fields, *Phys. Rept.* 1010 (2023) 1–138. arXiv:2203.00019, doi:10.1016/j.physrep.2023.01.003.
- [27] J. Ellis, N. E. Mavromatos, T. You, Light-by-light scattering constraint on born-infeld theory, *Phys. Rev. Lett.* 118 (2017) 261802. doi:10.1103/PhysRevLett.118.261802. URL <https://link.aps.org/doi/10.1103/PhysRevLett.118.261802>
- [28] J. Ellis, N. E. Mavromatos, P. Roloff, T. You, Light-by-light scattering at future e^+e^- colliders, *Eur. Phys. J. C* 82 (7) (2022) 634. arXiv:2203.17111, doi:10.1140/epjc/s10052-022-10565-w.
- [29] M. D. Schwartz, *Quantum Field Theory and the Standard Model*, Cambridge University Press, 2013.
- [30] B. Kettle, et al., A laser-plasma platform for photon-photon physics: the two photon Breit-Wheeler process, *New J. Phys.* 23 (11) (2021) 115006. arXiv:2106.15170, doi:10.1088/1367-2630/ac3048.
- [31] X. Ribeyre, E. d’Humières, O. Jansen, S. Jequier, V. Tikhonchuk, M. Lobet, Pair creation in collision of γ -ray beams produced with high-intensity lasers, *Physical Review E* 93 (1) (2016) 013201.
- [32] H. Abramowicz, et al., Conceptual design report for the LUXE experiment, *Eur. Phys. J. ST* 230 (11) (2021) 2445–2560. arXiv:2102.02032, doi:10.1140/epjs/s11734-021-00249-z.
- [33] Z. Chen, S. Meuren, E. Gerstmayr, V. Yakimenko, P. H. Bucksbaum, D. A. R. and, Preparation of strong-field qed experiments at facet-ii, in: *Optica High-brightness Sources and Light-driven Interactions Congress 2022*, Optica Publishing Group, 2022, p. HF4B.6. doi:10.1364/HILAS.2022.HF4B.6. URL <https://opg.optica.org/abstract.cfm?URI=HILAS-2022-HF4B.6>
- [34] E. Esarey, C. B. Schroeder, W. P. Leemans, Physics of laser-driven plasma-based electron accelerators, *Reviews of modern physics* 81 (3) (2009) 1229.
- [35] K. Behm, J. Cole, A. Joglekar, E. Gerstmayr, J. Wood, C. Baird, T. Blackburn, M. Duff, C. Harvey, A. Ilderton, et al., A spectrometer for ultrashort gamma-ray pulses with photon energies greater than 10 mev, *Review of Scientific Instruments* 89 (11) (2018) 113303.
- [36] J. Cole, K. Behm, E. Gerstmayr, T. Blackburn, J. Wood, C. Baird, M. J. Duff, C. Harvey, A. Ilderton, A. Joglekar, et al., Experimental evidence of radiation reaction in the collision of a high-intensity laser pulse with a laser-wakefield accelerated electron beam, *Physical Review X* 8 (1) (2018) 011020.
- [37] S. Agostinelli, J. Allison, K. Amako, J. Apostolakis, H. Araujo, P. Arce, M. Asai, D. Axen, S. Banerjee, G. Barrand, F. Behner, L. Bellegamba, J. Boudreau, L. Broglia, A. Brunengo, H. Burkhardt, S. Chauvie, J. Chuma, R. Chytracsek, G. Cooperman, G. Cosmo, P. Degt'yarenko, A. Dell’Acqua, G. Depaola, D. Dietrich, R. Enami, A. Fellicello, C. Ferguson, H. Fesefeldt, G. Folger, F. Foppiano, A. Forti, S. Garelli, S. Giani, R. Giannitrapani, D. Gibin, J. Gómez Cadenas, I. González, G. Gracia Abril, G. Greeniaus, W. Greiner, V. Grichine, A. Grossheim, S. Guatelli, P. Gumplinger, R. Hamatsu, K. Hashimoto, H. Hasui, A. Heikkinen, A. Howard, V. Ivanchenko, A. Johnson, F. Jones, J. Kallenbach, N. Kanaya, M. Kawabata, Y. Kawabata, M. Kawaguti, S. Kelner, P. Kent, A. Kimura, T. Kodama, R. Kokoulin, M. Kossov,

- H. Kurashige, E. Lamanna, T. Lampén, V. Lara, V. Lefebure, F. Lei, M. Liendl, W. Lockman, F. Longo, S. Magni, M. Maire, E. Medernach, K. Minamimoto, P. Mora de Freitas, Y. Morita, K. Murakami, M. Nagamatu, R. Nartallo, P. Nieminen, T. Nishimura, K. Ohtsubo, M. Okamura, S. O’Neale, Y. Oohata, K. Paech, J. Perl, A. Pfeiffer, M. Pia, F. Ranjard, A. Rybin, S. Sadilov, E. Di Salvo, G. Santin, T. Sasaki, N. Savvas, Y. Sawada, S. Scherer, S. Sei, V. Sirotenko, D. Smith, N. Starkov, H. Stoecker, J. Sulkimo, M. Takahata, S. Tanaka, E. Tcherniaev, E. Safai Tehrani, M. Tropeano, P. Truscott, H. Uno, L. Urban, P. Urban, M. Verderi, A. Walkden, W. Wander, H. Weber, J. Wellisch, T. Wenaus, D. Williams, D. Wright, T. Yamada, H. Yoshida, D. Zschiesche, [Geant4—a simulation toolkit](#), Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 506 (3) (2003) 250–303. doi:[https://doi.org/10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8). URL <https://www.sciencedirect.com/science/article/pii/S0168900203013688>
- [38] R. Watt, S. Rose, B. Kettle, S. Mangles, Monte carlo modeling of the linear breitt-wheeler process within the geant 4 framework, Physical Review Accelerators and Beams 26 (5) (2023) 054601.
- [39] R. Karplus, M. Neuman, The scattering of light by light, Phys. Rev. 83 (1951) 776–784. doi:[10.1103/PhysRev.83.776](https://doi.org/10.1103/PhysRev.83.776).
- [40] R. Karplus, M. Neuman, Non-Linear Interactions between Electromagnetic Fields, Phys. Rev. 80 (1950) 380–385. doi:[10.1103/PhysRev.80.380](https://doi.org/10.1103/PhysRev.80.380).
- [41] B. De Tollis, Dispersive approach to photon-photon scattering, Nuovo Cim. 32 (3) (1964) 757–768. doi:[10.1007/BF02735895](https://doi.org/10.1007/BF02735895).
- [42] B. De Tollis, The scattering of photons by photons, Nuovo Cim. 35 (4) (1965) 1182–1193. doi:[10.1007/BF02735534](https://doi.org/10.1007/BF02735534).
- [43] V. Costantini, B. De Tollis, G. Pistoni, Nonlinear effects in quantum electrodynamics, Nuovo Cim. A 2 (3) (1971) 733–787. doi:[10.1007/BF02736745](https://doi.org/10.1007/BF02736745).
- [44] M. Sangal, C. H. Keitel, M. Tamburini, Observing light-by-light scattering in vacuum with an asymmetric photon collider, Phys. Rev. D 104 (11) (2021) L111101. arXiv:[2101.02671](https://arxiv.org/abs/2101.02671), doi:[10.1103/PhysRevD.104.L111101](https://doi.org/10.1103/PhysRevD.104.L111101).
- [45] B. King, P. Böhl, H. Ruhl, Interaction of photons traversing a slowly varying electromagnetic background, Phys. Rev. D 90 (6) (2014) 065018. arXiv:[1406.4139](https://arxiv.org/abs/1406.4139), doi:[10.1103/PhysRevD.90.065018](https://doi.org/10.1103/PhysRevD.90.065018).
- [46] P. Böhl, B. King, H. Ruhl, Vacuum high harmonic generation in the shock regime, Phys. Rev. A 92 (3) (2015) 032115. arXiv:[1503.05192](https://arxiv.org/abs/1503.05192), doi:[10.1103/PhysRevA.92.032115](https://doi.org/10.1103/PhysRevA.92.032115).
- [47] A. Blinne, H. Gies, F. Karbstein, C. Kohlfürst, M. Zepf, All-optical signatures of quantum vacuum nonlinearities in generic laser fields, Phys. Rev. D 99 (1) (2019) 016006. arXiv:[1811.08895](https://arxiv.org/abs/1811.08895), doi:[10.1103/PhysRevD.99.016006](https://doi.org/10.1103/PhysRevD.99.016006).
- [48] T. Grismayer, R. Torres, P. Carneiro, F. Cruz, R. Fonseca, L. O. Silva, Quantum Electrodynamics vacuum polarization solver, New J. Phys. 23 (9) (2021) 095005. arXiv:[1607.04224](https://arxiv.org/abs/1607.04224), doi:[10.1088/1367-2630/ac2004](https://doi.org/10.1088/1367-2630/ac2004).
- [49] A. Lindner, B. Ölmez, H. Ruhl, HEWES: Heisenberg-Euler Weak-Field Expansion Simulator (2 2022). arXiv:[2202.09680](https://arxiv.org/abs/2202.09680).
- [50] O. Pike, F. Mackenroth, E. Hill, S. Rose, A photon–photon collider in a vacuum hohlraum, Nature Photonics 8 (6) (2014) 434–436.
- [51] T. Tajima, J. M. Dawson, Laser electron accelerator, Physical review letters 43 (4) (1979) 267.
- [52] J. A. Bearden, A. Burr, Reevaluation of x-ray atomic energy levels, Reviews of Modern Physics 39 (1) (1967) 125.
- [53] B. Kettle, T. Dzelzainis, S. White, L. Li, A. Rigby, C. Spindloe, M. Notley, R. Heathcote, C. Lewis, D. Riley, Ml band x-rays (3–3.5 keV) from palladium coated targets for isochoric radiative heating of thin foil samples, Journal of Physics B: Atomic, Molecular and Optical Physics 48 (22) (2015) 224002.
- [54] D. E. Knuth, Seminumerical algorithms, vol. 2: The art of the computer programming, Addison-Wesley, 1981.
- [55] P. Mason, N. Stuart, J. Phillips, R. Heathcote, S. Buck, A. Wojtusiak, M. Galimberti, T. de Faria Pinto, S. Hawkes, S. Tomlinson, et al., Progress on laser development at the extreme photonics applications centre, in: The European Conference on Lasers and Electro-Optics, Optica Publishing Group, 2023, p. ca_8_2.
- [56] F. Lureau, G. Matras, O. Chalus, C. Derycke, T. Morbieu, C. Radier, O. Casagrande, S. Laux, S. Ricaud, G. Rey, et al., High-energy hybrid femtosecond laser system demonstrating 2×10 pw capability, High Power Laser Science and Engineering 8 (2020) e43. doi:[10.1017/hpl.2020.41](https://doi.org/10.1017/hpl.2020.41).
- [57] F. Albert, M. Couprie, A. Debus, M. C. Downer, J. Faure, A. Flacco, L. A. Gizzi, T. Grismayer, A. Huebl, C. Joshi, et al., 2020 roadmap on plasma accelerators, New Journal of Physics 23 (3) (2021) 031101.
- [58] M. De Vido, K. Ertel, A. Wojtusiak, N. O’Donoghue, S. Tomlinson, M. Divoky, M. Sawicka-Chyla, J. Pilar, P. Mason, J. Phillips, et al., Design of a 10 j, 100 hz diode-pumped solid state laser, in: Applications of Lasers for Sensing and Free Space Communications, Optica Publishing Group, 2019, pp. JTU3A–14.