

MULTIPLIER IDEALS OF NORMAL SURFACE SINGULARITIES

LÁSZLÓ KOLTAI, TAMÁS LÁSZLÓ, AND ANDRÁS NÉMETHI

ABSTRACT. We study the multiplier ideals and the corresponding jumping numbers and multiplicities $\{m(c)\}_{c \in \mathbb{R}}$ in the following context: (X, o) is a complex analytic normal surface singularity, $\mathfrak{a} \subset \mathcal{O}_{X, o}$ is an $\mathfrak{m}_{X, o}$ -primary ideal, $\phi : \tilde{X} \rightarrow X$ is a log resolution of \mathfrak{a} such that $\mathfrak{a}\mathcal{O}_{\tilde{X}} = \mathcal{O}_{\tilde{X}}(-F)$, for some nonzero effective divisor F supported on $\phi^{-1}(0)$.

We show that $\{m(c)\}_{c > 0}$ is combinatorially computable from F and the resolution graph Γ of ϕ , and we provide several formulae. We also extend Budur's result (valid for $(X, o) = (\mathbb{C}^2, 0)$), which makes an identification of $\sum_{c \in [0, 1]} m(c)t^c$ with a certain Hodge spectrum. In our general case we use Hodge spectrum with coefficients in a mixed Hodge module.

We show that $\{m(c)\}_{c \leq 0}$ usually depends on the analytic type of (X, o) . However, for some distinguished analytic types we determine it concretely. E.g., when (X, o) is weighted homogeneous (and F is associated with the central vertex), we recover $\sum_c m(c)t^c$ from the Poincaré series of (X, o) and when (X, o) is a splice quotient then we recover $\sum_c m(c)t^c$ from the multivariable topological Poincaré (zeta) function of Γ .

1. INTRODUCTION

Assume that (X, o) is a complex analytic normal surface singularity, let $\mathcal{O}_{X, o}$ be its local algebra with maximal ideal $\mathfrak{m}_{X, o}$. Let us also fix an $\mathfrak{m}_{X, o}$ -primary ideal $\mathfrak{a} \subset \mathcal{O}_{X, o}$. Then there exists a log resolution $\phi : \tilde{X} \rightarrow X$ and a nonzero integral cycle F supported on the exceptional curve $\phi^{-1}(0) = E$ such that $\mathfrak{a}\mathcal{O}_{\tilde{X}} = \mathcal{O}_{\tilde{X}}(-F)$. In fact, in our study we will start with some ϕ and such a cycle F .

We denote by Z_K the anticanonical cycle associated with ϕ , $-c_1(\Omega_{\tilde{X}}^2) = -c_1(\mathcal{O}_{\tilde{X}}(K_{\tilde{X}})) \in H^2(\tilde{X}, \mathbb{Z})$, identified with a rational cycle (via the adjunction relations). We consider for any $c \in \mathbb{R}$ the cycle $L(c) := \lfloor Z_K + cF \rfloor$. Then the tower of *multiplier ideals* is defined as $\mathfrak{J}(c, F) := \phi_* \mathcal{O}_{\tilde{X}}(-L(c)) \subset \mathcal{O}_{X, o}$. They measure the complexity of the singularity (X, o) compatibly with F (or with \mathfrak{a}). They were introduced in [12, 21] and since then they were intensively studied, see [5, 3, 2, 14, 13, 15, 16, 17, 22, 23, 40] with all their connections with different aspects of singularities and singular map-germs. In fact, the definition and study extends to any dimension, see e.g. [12, 21, 31] and the references therein.

Date: June 20, 2024.

2010 Mathematics Subject Classification. Primary: 14B05, 14Fxx, 32S05; Secondary: 32S10, 32S25.

Key words and phrases. normal surface singularities, plane curve singularities, multiplier ideals, jumping numbers, jumping multiplicities, Hodge spectrum, mixed Hodge modules, canonical covering, weighted homogeneous singularities, splice quotient singularities .

The authors are partially supported by NKFIH Grant ‘‘Élvonal (Frontier)’’ KKP 144148. The second author acknowledges the support of the ‘János Bolyai Research Scholarship’ of the Hungarian Academy of Sciences, and the support of the project ‘‘Singularities and Applications’’ - CF 132/31.07.2023 funded by the European Union - NextGenerationEU - through Romania’s National Recovery and Resilience Plan.

It turns out that for any $c \in \mathbb{R}$, the ideal $\mathfrak{J}(c, F) \subset \mathcal{O}_{X,o}$ has finite codimension, $\mathfrak{J}(c, F) \subset \mathfrak{J}(c', F)$ for $c' < c$, and $\mathfrak{J}(c, F) = \mathcal{O}_{X,o}$ for $c \ll 0$. The number c is called *jumping number* if $m(c) := \dim \mathfrak{J}(c - \varepsilon, F) / \mathfrak{J}(c, F) \neq 0$ for $0 < \varepsilon \ll 1$, and $m(c)$ is the corresponding *jumping multiplicity*. One verifies that in any compact interval there are only finitely many jumping numbers and they are all rational.

Compared with the earlier literature in the subject we make two major extensions:

(A) We do not assume any restriction regarding the surface singularity (X, o) , e.g. we do not assume that the link is a rational homology sphere (i.e. the dual graph might have loops and the irreducible exceptional divisors might have higher genera). Also, we do not restrict the analytic type of (X, o) : on the supporting topological type the analytic type might vary (which in the non-rational or non-minimally elliptic cases might have considerable effects). We emphasize that in general the classification of such analytic types (or the description of the corresponding moduli spaces) is still an open problem. Under this generality the general analytic tools (e.g. vanishing theorems) are more complex (and the zones what they do not cover are wider).

(B) In the earlier studies only the cases $c > 0$ were treated. Our study of the cases $c \leq 0$ opens a new perspectives. If (X, o) is smooth or log terminal then $m(c) = 0$ for $c \leq 0$, however, in the general case (even for rational (X, o)) the negative zone appears as a very challenging part.

In fact (and this is one of the main messages of the note), we show that the study of the Poincaré series of multiplier ideals $\sum_{c \in \mathbb{R}} m(c)t^c$ (introduced first in [14]) breaks into two major parts with rather different aspects.

(i) (the $c > 0$ zone) We also call this the *Riemann–Roch zone*, since in this case we determine $\sum_{c > 0} m(c)t^c$ completely topologically from the resolution graph Γ of ϕ and the cycle F (without any restriction regarding the link or the analytic type of (X, o)).

We provide several combinatorial formulae. The first one, Proposition 3.3.4, is a generalization of expressions obtained in [2, 3, 40] proved in those articles for rational singularities.

We also compute $m(c+1) - m(c)$ (see Corollary 3.3.5), thus we recover $\sum_{c > 0} m(c)t^c$ from $\sum_{c \in (0,1]} m(c)t^c$ and (D_c, F) , where $D_c = L(c) - L(c - \varepsilon)$ ($0 < \varepsilon \ll 1$) is the *jumping divisor* (introduced in [2]), computable combinatorially. (This fact was also shown for rational (X, o) in [3].)

Hence, the first interval and $\sum_{c \in (0,1]} m(c)t^c$ has a distinguished role.

Regarding this we stress out two facts. Firstly, Budur in [5] identified $\sum_{c \in (0,1]} m(c)t^c$ with the Hodge spectrum $\text{Sp}_{(0,1]}(f)$ of a function germ f , where in his case, $(X, o) = (\mathbb{C}^2, 0)$ is smooth, $f : (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}, 0)$ is a function germ with an isolated singularity, ϕ is a log embedded resolution and F is the divisor $\text{div}_{\phi, E}(f)$ of $\phi^*(f)$ along $\phi^{-1}(0) = E$. This was reproved in [3] as well. (For the theory of mixed Hodge structures and of the spectrum associated with isolated hypersurface singularities see the work of Steenbrink, e.g. [36, 37] or [35].)

In this note we succeed to generalize this statement. The first step is that we rewrite $J_{[0,1]}(t) = \sum_{c \in [0,1]} m(c)t^c$ in a different way, in terms of Γ and the multiplicities of F , for any (X, o) and any F (cf. Theorem 4.2.3). This is the second combinatorial formula of the jumping multiplicities.

This formula will resonate perfectly with Hodge spectrum computations of a generic element $f \in \mathfrak{a}$. Indeed, if (X, o) is numerically Gorenstein (that is, Z_K is an integral cycle), and f is an

analytic function germs $(X, o) \rightarrow (\mathbb{C}, 0)$ with isolated singularities, $\operatorname{div}_{\phi, E}(f) = F$, we prove the generalization of the statement of Budur: $J_{[0,1]}(t) = \operatorname{Sp}_{[0,1]}(t)$, cf. Corollary 5.1.3. However, when Γ is not numerically Gorenstein, $\{Z_K\} \neq 0$, then we need to use a more general spectrum computation established in [24]. In [24] the usual spectrum is replaced by the spectrum of f with coefficients in a mixed Hodge module \mathcal{M} of (X, o) , such that \mathcal{M} restricted to $X \setminus \{o\}$ is a variation of Hodge structure whose underlying representation is abelian. Indeed, under the assumption that the link is a rational homology sphere, from Z_K we construct a character of the first homology of the link, hence a rank one variation of Hodge structure \mathbb{V} on $X \setminus \{o\}$, and we prove the identity $J_{[0,1]}(t) = \operatorname{Sp}_{[0,1]}(t)(f, j_*\mathbb{V})$ (where $j: X \setminus \{o\} \subset X$ is the inclusion), see Theorem 5.3.2. The construction of \mathbb{V} can be done using the canonical cyclic covering of (X, o) (associated with Z_K , or in \mathbb{Q} -Gorenstein case associated with the canonical divisor $K_{\tilde{X}}$, as well). We emphasize again, though a priori both objects $J_{[0,1]}(t)$ and $\operatorname{Sp}_{[0,1]}(t)(f, j_*\mathbb{V})$ are analytical, we provide combinatorial formulae for them.

The second new aspect regarding $\sum_{c \in (0,1]} m(c)t^c$ is the following connection. Assume that there exists again a function $f: (X, o) \rightarrow (\mathbb{C}, 0)$ with an isolated singularity such that $F = \operatorname{div}_{\phi, E}(f)$. Set $(C, o) := (f^{-1}(0), o)$ and let $\delta(C, o)$ be its delta invariant. Then $\sum_{c \in (0,1]} m(c) = \delta(C, o) + (\{Z_K\}, F)$.

(ii) (the $c \leq 0$ zone) We show by examples that the multiplicity $m(0)$ cannot be determined topologically. However, we characterize it completely in terms of the analytic type of (X, o) . The analytic contribution is a term $h^0 \in \{0, 1\}$ characterized by an h^0 -sheaf cohomology.

The $c \leq 0$ zone can be called the ‘ h^0 -zone’, where the Riemann–Roch expressions or vanishing theorems cannot be applied. In this zone $m(c)$ usually cannot be computed from Γ and F . We exemplify this fact by concrete examples when we fix Γ and F and we consider several different analytic realizations, and we compute the $m(c)$ ’s. Interestingly enough, already the numerical invariant $\sum_{c \leq 0} m(c)$ measures the analytic type and it is connected to the most important analytic invariant of (X, o) , the geometric genus $p_g(X, o)$. More precisely, we prove that $\sum_{c \leq 0} m(c) = p_g(X, o) + \chi(\lfloor Z_K \rfloor)$. Already when (X, o) is rational (i.e. $p_g(X, o) = 0$), $\sum_{c \leq 0} m(c)$ can be arbitrarily large (see e.g. Example 3.2.2).

On the other hand, even for this zone, we provide concrete formulae valid for certain *distinguished analytic types*.

- In the case of a *weighted homogeneous germs* (X, o) with local graded algebra $\bigoplus_{\ell \geq 0} R_\ell$ (and F is associated with the central vertex), we show the correspondence $\dim(R_\ell) = m(c)$ with $c = (\ell - \tau)/\alpha$ (for notations and precise statement see subsection 3.4). For $\ell > \tau$ (corresponding to $c > 0$) $\dim(R_\ell)$ is topological computable from the Seifert invariants of the link, however in general it is not. But, if the link is a rational homology sphere, then by Dolgachev-Pinkham-Demazure formulae all the terms for any ℓ are topological. (For the theory of weighted homogeneous germs see e.g. [32].)

- If the link is a rational homology sphere and we fix a resolution graph Γ , and we assume that the analytic type of (X, o) is *splice quotient* with respect to Γ (a family introduced by Neumann and Wahl, cf. [29]). Then in Proposition 3.1.4 we provide a combinatorial formula for $\sum_{c \in \mathbb{R}} m(c)t^c$ in terms of $\{L(c)\}_c$ and the multivariable topological Poincaré (zeta) function $Z(\mathbf{t})$ of Γ . This includes e.g. all the cases when (X, o) is rational or minimally elliptic (with Γ minimal). (For more on $Z(\mathbf{t})$ see [6, 7, 27].)

But we emphasize again that in general, the computation of $\{m(c)\}_{c \leq 0}$ is of analytic nature, hence it runs in parallel with the classification of the analytic types of (X, o) supported of a fixed topological type identified by Γ . We will return back to this study in the forthcoming manuscript.

2. PRELIMINARIES

2.1. Notations regarding resolutions and resolution graphs.

2.1.1. Let us fix a normal surface singularity (X, o) and one of its log (good) resolutions $\phi : \tilde{X} \rightarrow X$. Let Γ be the dual resolution graph of ϕ , we denote its vertices by \mathcal{V} and its edges by \mathcal{E} . We write E for the irreducible exceptional curve $\phi^{-1}(o)$, and let $\{E_i\}_{i \in \mathcal{V}}$ be its irreducible components.

The lattice $L := H_2(\tilde{X}, \mathbb{Z})$ is a free \mathbb{Z} -module generated by the classes of $\{E_i\}_{i \in \mathcal{V}}$ and it is endowed with the natural negative definite intersection form $(,)$. The dual lattice is $L' = \text{Hom}_{\mathbb{Z}}(L, \mathbb{Z}) \simeq \{l' \in L \otimes \mathbb{Q} : (l', L) \in \mathbb{Z}\} \subset L \otimes \mathbb{Q}$. We use the same symbol $(,)$ for the extended intersection form to $L \otimes \mathbb{Q}$. The overlattice L' is generated over \mathbb{Z} by the (anti)dual classes $\{E_i^*\}_{i \in \mathcal{V}}$ defined by $(E_i^*, E_j) = -\delta_{ij}$ (where δ_{ij} stays for the Kronecker symbol). L' can also be identified with $H^2(\tilde{X}, \mathbb{Z})$ (the target of the first Chern class $c_1 : \text{Pic}(\tilde{X}) \rightarrow H^2(\tilde{X}, \mathbb{Z})$).

We denote by M the link of (X, o) . It is known that M is a rational homology sphere if and only if each E_i is rational and Γ is a tree. It is an integral homology sphere if additionally $\det(,) = \pm 1$. More generally, if g_i denotes the genus of E_i , $g := \sum_i g_i$ and h the first Betti number of the topological realization of Γ (the number of independent cycle in the graph), then the first Betti number of M is $2g + h$, and $\text{Tors} H_1(M, \mathbb{Z}) \simeq L'/L$.

There is a natural partial ordering of $L \otimes \mathbb{Q}$: we write $l'_1 \geq l'_2$ if $l'_1 - l'_2 = \sum_i a_i E_i$ with every $a_i \geq 0$. We set $L_{\geq 0} = \{l \in L : l \geq 0\}$ and $L_{> 0} = L_{\geq 0} \setminus \{0\}$.

We define the rational *Lipman cone* as $\mathcal{S}_{\mathbb{Q}} := \{l' \in L \otimes \mathbb{Q} : (l', E_i) \leq 0 \text{ for all } i\}$, and we also set $\mathcal{S}' := \mathcal{S}_{\mathbb{Q}} \cap L'$ and $\mathcal{S} := \mathcal{S}_{\mathbb{Q}} \cap L$. As a monoid \mathcal{S}' is generated over $\mathbb{Z}_{\geq 0}$ by $\{E_i^*\}_i$. If $s' \in \mathcal{S}_{\mathbb{Q}} \setminus \{0\}$, then all its E_i -coordinates are strictly positive (see [27, Corollary 2.1.19]).

Let $K_{\tilde{X}}$ be the *canonical divisor* of \tilde{X} (well-defined up to a linear equivalence), that is, $\mathcal{O}_{\tilde{X}}(K_{\tilde{X}}) \simeq \Omega_{\tilde{X}}^2$, the sheaf of holomorphic 2-forms. We denote its first Chern class $c_1(\mathcal{O}_{\tilde{X}}(K_{\tilde{X}})) \in H^2(\tilde{X}, \mathbb{Z}) \simeq L'$ by $-Z_K$; Z_K is called the *(anti)canonical cycle* of ϕ . The (rational) cycle $Z_K \in L'$ can also be determined combinatorially from $(,)$ by the *adjunction formulae* $(Z_K, E_i) = (E_i, E_i) + 2 - 2g_i$ for all $i \in \mathcal{V}$.

We call (X, o) *numerically Gorenstein* if $Z_K \in L$, i.e. if all its E_i -coefficients are integral (this condition is independent of the choice of ϕ). We call (X, o) *Gorenstein* if $Z_K \in L$ and one can choose $K_{\tilde{X}}$ as $-Z_K$, i.e. $\mathcal{O}_{\tilde{X}}(K_{\tilde{X}} + Z_K) \simeq \mathcal{O}_{\tilde{X}}$. In a different language, (X, o) is numerically Gorenstein (resp. Gorenstein) exactly when the line bundle $\Omega_{\tilde{X} \setminus \{o\}}^2$ is topologically (resp. analytically) trivial. (For a different characterization see also 5.4.1.)

Finally, we set for any $l' \in L'$ the Riemann-Roch expression $\chi(l') = (Z_K - l', l')/2$. If $l \in L_{> 0}$, then $\chi(l) = h^0(\mathcal{O}_l) - h^1(\mathcal{O}_l)$. E.g., $\chi(E) = 1 - g - h$. As usual, we also write $\chi(\mathcal{O}_l(D))$ for $h^0(\mathcal{O}_l(D)) - h^1(\mathcal{O}_l(D))$, which, by Riemann-Roch, equals $\chi(l) + (D, l)$ ($D \in L$).

For any \mathbb{R} -divisor $D = \sum_i d_i E_i$ of \tilde{X} ($d_i \in \mathbb{R}$) we define its integral part $\lfloor D \rfloor$ and fractional part $\{D\}$ by applying the corresponding operation to each coefficient d_i .

For more details see e.g. [25, 27].

2.1.2. Assume that (C, o) is a reduced Weil divisor in (X, o) (i.e. it is an isolated curve singularity in (X, o)). Then we can consider a log embedded resolutions of the pair $(C, o) \subset (X, o)$, that is, we require that in the log resolution the union of E with the strict transform \tilde{C} of (C, o) form a simple normal crossing divisor. Moreover, we can consider the corresponding embedded resolution graph too. Usually, the strict transform will be denoted by arrows on the graph. Their index set will be denoted by \mathcal{A} corresponding to the irreducible decomposition $\cup_{a \in \mathcal{A}} C_a$ of (C, o) .

Once an embedded resolution and the strict transform $\tilde{C} \subset \tilde{X}$ is fixed, the embedded topological type of the pair $(C, o) \subset (X, o)$ is basically coded in the information that how many components of \tilde{C} intersect each E_i , that is, in the intersection numbers $(\tilde{C}, E_i)_{\tilde{X}}$. Then we define the rational cycle $l'_C \in \mathcal{S}' \setminus \{0\} \subset L'_{>0}$, associated with C and ϕ , via the linear system of equations $(l'_C, E_i) + (\tilde{C}, E_i)_{\tilde{X}} = 0$ for every $i \in \mathcal{V}$. (Hence, $l'_C + \tilde{C}$, as a rational divisor of \tilde{X} , is numerically trivial. It is the total transform of C denoted by ϕ^*C .) If the arrow a is supported by the vertex $i(a)$, then $l'_C = \sum_{a \in \mathcal{A}} E_{i(a)}^*$.

If (C, o) is cut out by a germ $f : (X, o) \rightarrow (\mathbb{C}, 0)$ such that $(C, o) = (f^{-1}(0), o)$ is reduced (that is, f defines an isolated singularity), then we say that it is a Cartier divisor of (X, o) . In such a case $\phi^*C = \text{div}_{\tilde{X}}(\phi^*f)$. Hence l'_C is integral; it will also be denoted by $\text{div}_{\phi, E}(f)$.

2.1.3. Next, we fix a cycle $F \in \mathcal{S} \setminus \{0\}$. Some geometrical situations how such a cycle might appear is listed below.

- (A) Let $\mathfrak{m}_{X, o}$ denote the maximal ideal of the local ring $\mathcal{O}_{X, o}$ of (X, o) . Let \mathfrak{a} be an $\mathfrak{m}_{X, o}$ -primary ideal of $\mathcal{O}_{X, o}$. Then there exists a log resolution ϕ such that $\mathfrak{a} \cdot \mathcal{O}_{\tilde{X}} = \mathcal{O}_{\tilde{X}}(-F)$ for some $F \in \mathcal{S} \setminus \{0\}$.
- (B) Let $(C, o) \subset (X, o)$ be a reduced embedded curve germ, and ϕ an embedded good resolution. Then $l'_C \in \mathcal{S}' \setminus \{0\}$, hence a convenient multiple of it is in $\mathcal{S} \setminus \{0\}$. Cf. also with Remark 2.2.2.
- (C) Let $f : (X, o) \rightarrow (\mathbb{C}, 0)$ be a germ, then in any good embedded resolution $F := \text{div}_{\phi, E}(f) \in \mathcal{S} \setminus \{0\}$.

If \mathfrak{a} is as in (A), and we fix a resolution ϕ with $\mathfrak{a} \cdot \mathcal{O}_{\tilde{X}} = \mathcal{O}_{\tilde{X}}(-F)$, then for any generic element f of \mathfrak{a} one has $\text{div}_{\phi, E}(f) = F$.

- (D) Conversely, let us fix a semigroup element $F \in \mathcal{S} \setminus \{0\}$, a divisor of a resolution of some analytic germ (X, o) . If (X, o) is rational then there exists a germ f such that $\text{div}_{\phi, E}(f) = F$, but for an arbitrary singularity (X, o) this is not the case. However, if we fix the topological type of (X, o) , identified by M or by some resolution graph Γ , then there exists an analytic type, say (X_F, o) supported on M and a germ $f_F : (X_F, 0) \rightarrow (\mathbb{C}, 0)$ such that $\text{div}_{\phi_F, E_F}(f_F) = F$.

- (E) The correspondence $\mathfrak{a} \mapsto F$ given by $\mathfrak{a} \cdot \mathcal{O}_{\tilde{X}} = \mathcal{O}_{\tilde{X}}(-F)$ (via some good resolution) is not injective. E.g., if we start with such an \mathfrak{a} and F , then $\phi_* \mathcal{O}_{\tilde{X}}(-F)$ is the integral closure $\bar{\mathfrak{a}}$ of \mathfrak{a} in $\mathcal{O}_{X, o}$, and $\bar{\mathfrak{a}} \cdot \mathcal{O}_{\tilde{X}} = \mathfrak{a} \cdot \mathcal{O}_{\tilde{X}} = \mathcal{O}_{\tilde{X}}(-F)$ too. Moreover, for any other ideal \mathfrak{b} the identity $\bar{\mathfrak{a}} = \bar{\mathfrak{b}}$ is equivalent with $\mathfrak{b} \cdot \mathcal{O}_{\tilde{X}} = \mathfrak{a} \cdot \mathcal{O}_{\tilde{X}}$.

2.2. The multiplier ideals.

2.2.1. The definition of multiplier ideals, jumping numbers and jumping multiplicities. We fix a normal surface singularity (X, o) , a good resolution ϕ , and a cycle $F \in \mathcal{S} \setminus \{0\}$.

Following [12, 21] we define the *multiplier ideal* associated with F and a real number $c \in \mathbb{R}$ as

$$\mathfrak{J}(c, F) = \phi_* \mathcal{O}_{\tilde{X}}(-\lfloor Z_K + cF \rfloor) \subset \mathcal{O}_{X, o}.$$

For simplicity, we denote the divisor $\lfloor Z_K + cF \rfloor$ by $L(c)$. Though Z_K and F are combinatorial cycles associated with the graph Γ , the ideals $\mathfrak{J}(c, F)$ might depend essentially on the analytic structure of the germ (X, o) . Since $\phi : \tilde{X} \setminus E \rightarrow X \setminus o$ is an isomorphism and $L(c)$ is supported on E , we obtain that $\mathfrak{J}(c, F) = \mathcal{O}_{X, o}$ whenever $L(c) \leq 0$, and $\mathfrak{J}(c, F) \subset \mathcal{O}_{X, o}$ is an $\mathfrak{m}_{X, o}$ -primary ideal otherwise. Also, $\phi_* \mathcal{O}_{\tilde{X}}(-L(c)) = \phi_* \mathcal{O}_{\tilde{X}}(-\max\{0, L(c)\})$ (see e.g. Lemma 3.1.1). As a corollary we get that

$$\dim(\mathcal{O}_{X, o}/\mathfrak{J}(c, F)) < \infty$$

If $c \leq c'$, then $\mathfrak{J}(c, F) \supseteq \mathfrak{J}(c', F)$. Whenever we have strict inclusion

$$\mathfrak{J}(c - \varepsilon, F) \supsetneq \mathfrak{J}(c, F)$$

for arbitrarily small $\varepsilon > 0$, we say that c is a *jumping number*, and its *jumping multiplicity* is:

$$m(c) := \dim(\mathfrak{J}(c - \varepsilon, F)/\mathfrak{J}(c, F)).$$

Define the jumping divisor corresponding to $c \in \mathbb{R}$ (cf. [2, 3, 40]) as

$$D_c := L(c) - L(c - \varepsilon) \quad (1 \gg \varepsilon > 0).$$

Obviously, D_c is a reduced divisor and it is ‘periodical’: $D_{c+1} = D_c$. In fact, if $Z_K = \sum k_i E_i$ and $F = \sum m_i E_i$, then E_i is in the support of D_c precisely when $k_i + cm_i \in \mathbb{Z}$. This shows that in any bounded interval there are only finitely many values c with $D_c \neq 0$, hence with $m(c) > 0$. Note that $D_c \neq 0$ usually does not imply $m(c) > 0$. We also define

$$\Delta_c := \{Z_K + cF\}.$$

Then any E_i is either in the support of D_c or of Δ_c , but not in both.

Since $F \in \mathcal{S} \setminus \{0\}$, hence all the coefficient of F are strict positive, if $c \ll 0$ then $Z_K + cF \leq 0$, and $\mathfrak{J}(c, F) = \mathcal{O}_{X, o}$. In particular, $m(c) = 0$ for $c \ll 0$. Thus, in any interval of type $(-\infty, c_0]$ there are only finitely many jumping numbers c .

Remark 2.2.2. The assumption that F is integral is just a convenient choice. In all our arguments regarding the cycle F , we can equally start with any $F \in \mathcal{S}_{\mathbb{Q}} \setminus \{0\}$, or, we can replace the integral F by any rF , where $r \in \mathbb{Q}_{>0}$. Indeed, the effect of the replacement F by rF is $m_{rF}(c) = m_F(rc)$.

3. JUMPING MULTIPLICITIES AND THE HILBERT FUNCTION

3.1. Multiplier ideals and the Hilbert function. Let us start with the following (known) lemma.

Lemma 3.1.1. *Fix some $l \in L$ and write it as $l = P - N$, with $P, N \in L_{\geq 0}$ and such that they have no common E_i -component. Then*

- (a) $H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(-P)) \rightarrow H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(-l))$ is an isomorphism,
- (b) $h^1(\mathcal{O}_{\tilde{X}}(-P)) - h^1(\mathcal{O}_{\tilde{X}}(-l)) = \chi(N) - (N, l)$.

Proof. Consider the short exact sequence of sheaves $0 \rightarrow \mathcal{O}_{\tilde{X}}(-P) \rightarrow \mathcal{O}_{\tilde{X}}(-l) \rightarrow \mathcal{O}_N(-l) \rightarrow 0$.

Then $(c_1(\mathcal{O}_{\tilde{X}}(-P)), E_i) = (-P, E_i) \leq 0$ for all E_i in the support of N . Therefore by the h^0 -vanishing theorem (see [27, Theorem 6.4.2.]), the dual of the Grauert-Riemenschneider vanishing, we obtain $H^0(N, \mathcal{O}_N(-P + N)) = 0$. Then use the long cohomological exact sequence. \square

This shows that for any $l \in L$ we have $H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(-l)) = H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(-\max\{0, l\}))$, which is canonically a subspace of $H^0(\tilde{X}, \mathcal{O}_{\tilde{X}})$. The Hilbert function $\mathfrak{h}(l)$ (associated with the divisorial filtration of the resolution) is defined as

$$L \ni l \mapsto \mathfrak{h}(l) := \dim H^0(\tilde{X}, \mathcal{O}_{\tilde{X}})/H^0(\tilde{X}, \mathcal{O}_{\tilde{X}}(-\max\{0, l\})) = \dim \mathcal{O}_{X,o}/\phi_* \mathcal{O}_{\tilde{X}}(-l).$$

Note that $\mathfrak{h}(L(c)) = \dim \mathcal{O}_{X,o}/\mathfrak{J}(c, F)$, hence

$$(3.1.2) \quad m(c) = \mathfrak{h}(L(c)) - \mathfrak{h}(L(c - \varepsilon)).$$

3.1.3. Some general comments regarding the formula (3.1.2). Usually, under the assumption that we fix the combinatorial data Γ , the Hilbert function \mathfrak{h} might depend essentially on the choice of the analytic type of (X, o) . However, for any rational singularity, for any fixed resolution with graph Γ , $\mathfrak{h}(l)$ can be determined from Γ for any $l \in L$. In fact, this is true for any singularity whose link is a rational homology sphere and its analytic type is *splice quotient*. This family of normal surface singularities was introduced by Neumann and Wahl and it includes (among others) all rational, minimally elliptic and weighted homogeneous singularities (with $\mathbb{Q}HS^3$ link). For such germs, $\mathfrak{h}(l)$ follows combinatorially via the following two steps: by [27, (8.2.10)] $\mathfrak{h}(l)$ can be computed from the multivariable (analytic) Poincaré series $P(\mathbf{t})$ of (X, o) (this is true for any normal surface singularities), while by [27, Corollary 8.5.35] $P(\mathbf{t}) = Z(\mathbf{t})$, where $Z(\mathbf{t})$ is the topological multivariable Poincaré series (or zeta function) associated with Γ (here the splice quotient assumption is crucial). (For the definition and certain properties of analytic and topological multivariable Poincaré series see also [6, 7, 8].) In particular, in principle, for all such singularities (and for any F) all the jumping multiplicities $\{m(c)\}_{c \in \mathbb{R}}$ are computable from Γ and F . Here is the precise statement.

Proposition 3.1.4. *Assume that the link of (X, o) is a rational homology sphere and (X, o) is splice quotient (associated with its resolution graph Γ). Consider free variables $\mathbf{t} = \{t_i\}_{i \in \mathcal{V}}$ indexed by the set of vertices, and for any $l' = \sum l'_i E_i \in L'$ write $\mathbf{t}^{l'} = \prod_i t_i^{l'_i}$. Let $Z(\mathbf{t}) = \sum_{l' \in \mathcal{S}'} \mathfrak{p}(l') \mathbf{t}^{l'}$ be the Taylor expansion at the origin of the rational function*

$$\prod_{i \in \mathcal{V}} (1 - \mathbf{t}^{E_i^*})^{\kappa_i - 2},$$

where κ_i is the valency of the vertex i . Let $Z_0(\mathbf{t})$ be its integral part, namely $Z_0(\mathbf{t}) = \sum_{l \in \mathcal{S}} \mathfrak{p}(l) \mathbf{t}^l \in \mathbb{Z}[\mathcal{S}]$. Then for any $c \in \mathbb{R}$ one has

$$m(c) = \sum_{l \geq L(c) - D_c, l \not\geq L(c)} \mathfrak{p}(l).$$

However, we wish to emphasise, that in the non-rational splice quotient cases, if we keep the topological type (say Γ) and we modify the analytic structure of (X, o) (into a non-splice quotient one) then the Hilbert function and the jumping multiplicities might change.

But, in the rational case, independently of the analytic realizations of the topological type, the Hilbert function and $m(c)$ are combinatorial expressible from Γ and F (via the above recipe).

In the general splice quotient case the explicit computation via the above recipe can be rather technical and computational.

In this note, for $c \geq 0$ and for arbitrary (X, o) we run a completely different approach (inspired from mixed Hodge theory). In fact, for $c > 0$ we will be able to determine $m(c)$ *combinatorially* for any (X, o) and F , independently of the analytic structure of (X, o) , and we provide explicit formulae. The number $m(0)$ might depend on the analytic structure, but we determine it completely.

Now, in fact, we arrived to a main point of the discussion: the behaviour of $m(c)$ is very different for $c > 0$ and for $c \leq 0$. As we already said, $m(c)$ for $c > 0$ is topological, independent of the analytic structure of (X, o) . For $c \leq 0$, the knowledge of Γ and F is not enough, the analytic structure of (X, o) plays a crucial role. Hence, in this case, the study of the jumping numbers and multiplicities goes in parallel with the study of (yet open problem of classification of) analytic structures supported on a fixed topological type.

In subsection 3.2 we provide some concrete examples in the case $c < 0$ to show how the variation on the analytic structure enters in the picture. Moreover, for weighted homogeneous germs we determine $m(c)$ for any $c \in \mathbb{R}$ in terms of the (analytic) Poincaré series of (X, o) . But a more complete study regarding $\{m(c)\}_{c < 0}$ will be presented in forthcoming manuscript including several other analytic aspects and connections.

The difference between the negative and positive half-lines is already visible via the following formula (cf. the first part of the proof of Proposition 3.3.4).

$$(3.1.5) \quad \sum_{c' \leq c} m(c') = \dim \mathcal{O}_{X,o}/\mathcal{J}(c, F) = \mathfrak{h}(L(c)) = \chi(L(c)) + p_g - h^1(\mathcal{O}_{\tilde{X}}(-L(c))).$$

Here $p_g = p_g(X, o)$ is the geometric genus of (X, o) , defined as $\dim H^1(\tilde{X}, \mathcal{O}_{\tilde{X}})$. Usually it is not a topological invariant of (X, o) , however, it is a c -independent ‘universal’ analytic invariant. Thus, let us focus on the difference $\mathfrak{h}(L(c)) - p_g$. It has two terms, $\chi(L(c))$ is combinatorial, however, $h^1(\mathcal{O}_{\tilde{X}}(-L(c)))$ is not. The point is that for $c \geq 0$ this last term is zero (see the second part of the proof of Proposition 3.3.4) by the Generalized Grauert-Riemenschneider vanishing theorem, independently of the analytic structure of (X, o) . On the other hand, for $c < 0$ this term can be nonzero, and usually depends essentially on the choice of analytic structure (see also examples from subsection 3.2, and Remark 3.4.6). We can say that $c > 0$ is the ‘Riemann–Roch zone’, while $c \leq 0$ is the ‘ h^0 -zone’.

3.1.6. On the other hand, the smallest jumping number — independently of the fact that it is negative or not — can be deduced combinatorially.

Lemma 3.1.7. (a) *The value $\mathfrak{c} := \min\{c : m(c) > 0\}$ can be determined from Γ and F as follows.*

Write $Z_K = \sum_i k_i E_i$ and $F = \sum_i m_i E_i$. Then

$$(3.1.8) \quad \mathfrak{c} = \min_{i \in \mathcal{V}} \{(1 - k_i)/m_i\}.$$

Moreover, $m(\mathfrak{c}) = 1$.

(b) (‘weak/support periodicity’) *Assume that there exists a function $f : (X, o) \rightarrow (\mathbb{C}, 0)$ and $n \in \mathbb{Z}_{>0}$ such that $\operatorname{div}_{\phi, E}(f^n) = F$. Then $m(c + \frac{1}{n}) \geq m(c)$ for any $c \in \mathbb{R}$.*

In the case of $\mathfrak{c} > 0$, part (a) was already known (see eg. [2]), \mathfrak{c} is known as the log-canonical threshold of F .

Note that if (X, o) is rational and ϕ arbitrary, or (X, o) is minimally elliptic and ϕ is minimal, then any $F \in \mathcal{S} \setminus \{0\}$ is realized as $\text{div}_{\phi, E}(f)$ for some f [27, Corollary 7.1.13 and Theorem 7.2.31].

Proof. (a) Let c' be the right hand side of (3.1.8). Then $L(c') \leq E$, but at least one of its coefficients is one. Hence $\mathfrak{J}(c', F) = \mathfrak{m}_{X, o}$ and $\mathfrak{J}(c' - \varepsilon, F) = 0$.

(b) Note that $F/n \in L$ hence $L(c + \frac{1}{n}) = L(c) + F/n$ for every $c \in \mathbb{R}$. Then consider the commutative diagram (with simplification $\mathfrak{J}(c) = \mathfrak{J}(c, F)$):

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathfrak{J}(c) & \longrightarrow & \mathfrak{J}(c - \varepsilon) & \longrightarrow & \mathfrak{J}(c - \varepsilon)/\mathfrak{J}(c) \\ & & \downarrow \cdot f & & \downarrow \cdot f & & \downarrow \cdot \bar{f} \\ 0 & \longrightarrow & \mathfrak{J}(c + 1/n) & \longrightarrow & \mathfrak{J}(c - \varepsilon + 1/n) & \longrightarrow & \mathfrak{J}(c - \varepsilon + 1/n)/\mathfrak{J}(c + 1/n) \end{array}$$

Above, the vertical morphisms are induced by multiplication by $f \in \mathcal{O}_{X, o}$. The first two vertical morphisms are injective. Since the intersection $f\mathfrak{J}(c - \varepsilon) \cap \mathfrak{J}(c + \frac{1}{n})$, as subspaces of $\mathfrak{J}(c - \varepsilon + \frac{1}{n})$, is $f\mathfrak{J}(c)$, the injectivity of $\cdot \bar{f}$ follows too. \square


Example 3.1.9. Write $Z_K = \sum_i k_i E_i$. Following e.g. [19, page 56], (X, o) is *numerically log terminal*, *nlt* (respectively *numerically log canonical*, *nlc*) if $k_i < 1$ (respectively $k_i \leq 1$) for all $i \in \mathcal{V}$. (If this happens in the minimal good resolution then it happens in any good resolution.) The *nlt* singularities are exactly the quotient singularities (see also [27, Example 6.3.33]).

By (3.1.8), if (X, o) is *nlt* (respectively *nlc*) then $m(c) = 0$ for all $c \leq 0$ (respectively $c < 0$) and for any $F \in \mathcal{S} \setminus \{0\}$. In the *nlc* case, (X, o) is either (special) rational or $Z_K = E$ (cf. [19] or [27, Example 6.3.33]). This latter case happens if (X, o) is either cusp or simple elliptic. If $Z_K = E$ then $c = 0$ and $m(0) = 1$.

In the special case when (X, o) is smooth or it is of type ADE then $Z_K \leq 0$, hence $c > 0$ again.

3.2. Some examples regarding $m(c)$ with $c \leq 0$.

Example 3.2.1. Negative c can appear even for (X, o) rational. Take e.g. the following rational graph, and let F be determined by the arrow: if $i(a)$ is the vertex which supports the arrow, then $F = E_{i(a)}^*$.



Then, by a computation, $c = -1/4$ and $m(-1/4) = 1$.

In this case $D_0 \neq 0$, but $m(0) = 0$. Indeed, Z_K has some integral coefficients hence $D_0 \neq 0$. On the other hand, both $L(0)$ and $L(-\varepsilon)$ are elements of $L_{>0}$, but they are smaller than the Artin fundamental cycle. Hence both jumping ideals are $\mathfrak{m}_{X, o}$.

Example 3.2.2. (a) For any (X, o) , for any good ϕ and any F , the identity (3.1.5) (via the vanishing $h^1(\mathcal{O}_{\tilde{X}}(-[Z_K])) = 0$, cf. the proof of Proposition 3.3.4) in the case of $c = 0$ reads as


$$\sum_{c' \leq 0} m(c') = \chi([Z_K]) + p_g.$$

Note that this number is independent of the choice of F , however, the distribution of the jumping numbers might depend on F .

In rational case of Example 3.2.1, $p_g = 0$ and $\chi(\lfloor Z_K \rfloor) = 1$.

If (X, o) is numerically Gorenstein, then $\chi(\lfloor Z_K \rfloor) = \chi(Z_K) = 0$, hence $\sum_{c' \leq 0} m(c') = p_g$.

(b) Consider the next rational graph, where the number of (-2) curves in all the legs is $(2n - 1)$.



Set $F := E$, it is $\text{div}_{\phi, E}$ of the generic function. A computation shows that the multiplicity of Z_K at the nodes is n , by (3.1.8) $\mathfrak{c} = 1 - n$, and $\chi(\lfloor Z_K \rfloor) = n$. Hence, by part (a) $\sum_{c \leq 0} m(c) = n$ and by Lemma 3.1.7(b), necessarily $\sum_{c \leq 0} m(c)t^c = t^{1-n} + \dots + t^{-1} + t^0$.

Remark 3.2.3. In the next subsection (Proposition 3.3.4) we determine $m(0)$ for any (X, o) . It depends on the choice of the analytic structure of (X, o) but it does not depend on the choice of F .

Example 3.2.4. Consider the following resolution graph.



The graph is weakly elliptic. It turns out that depending on the analytic structure $1 \leq p_g \leq 3$ (see e.g. [27, 7.2.C and 7.2.D]). $Z_K = 3E_1 + 2E_2 + E_3$, hence $Z_K \in L$, and $\sum_{c \leq 0} m(c) = p_g$.

Taking E_1 (the elliptic exceptional divisor) as a central vertex, the link is a Seifert 3-manifold, and (X, o) can be chosen weighted homogeneous. The point is that we can choose three different families of weighted homogeneous analytic structures, see [32] (or [27, Example 5.1.30]). Certain representatives are the following.

- (i) (X, o) is the hypersurface singularity $z^2 = y^3 + x^{18}$ with $p_g = 3$,
- (ii) (X, o) is the hypersurface singularity $z^2 = x(y^4 + x^6)$ with $p_g = 2$,
- (iii) (X, o) is not Gorenstein, with $p_g = 1$.

Take $F = E_1 + E_2 + E_3$. Then there are three values $c \leq 0$ with non-trivial jumping divisors: $c \in \{-2, -1, 0\}$. For any analytic structure, $L(-2) = E_1$ and $L(-2 - \varepsilon) = 0$, hence $m(-2) = 1$. On the other hand, taking $c = 0$ we get $L(0) = Z_K$, $L(-\varepsilon) = 2E_1 + E_2$, and $\chi(L(0)) = \chi(L(-\varepsilon)) = 0$. Therefore, as we will prove in Proposition 3.3.4, $m(0) = \mathfrak{h}^0$, which equals 1 if (X, o) is Gorenstein (in the first two cases). In case (iii) it should be 0 since $\sum_{c \leq 0} m(c) = 1$, but we already know that $m(-2) = 1$. Hence $\sum_{c \leq 0} m(c)t^c$ is $t^{-2} + t^{-1} + t^0$ in case (i), $t^{-2} + t^0$ in case (ii), and t^{-2} in case (iii).

3.3. The computation of $m(c)$ for $c \geq 0$.

3.3.1. In this subsection we determine $m(c)$ for all $c \geq 0$, for all possible F 's, and for all analytic structures.

Though the cycles $L(c)$ are very special, still, in principle, the output of the substitution $c \mapsto \mathfrak{h}(L(c))$ might depend on the analytic type of (X, o) , usually it cannot be determined from Γ . This

fact will follow from Proposition 3.3.4 where we connect $\mathfrak{h}(L(c))$ with the geometric genus (X, o) , which usually is not a topological invariant of (X, o) .

However, for any $c > 0$ we will show that $\mathfrak{h}(L(c)) - p_g$ is combinatorial. Hence, via (3.1.2), $m(c)$ is also combinatorial, and we will provide several formulae for it.

For $c = 0$ an additional analytic contribution should also be considered:

$$\mathfrak{h}^0 := h^0(D_0, \mathcal{O}_{D_0}(K_{\bar{X}} + [Z_K])).$$

Recall that from the definition of D_0 , $D_0 = E$ exactly when (X, o) is numerically Gorenstein: $Z_K \in L$.

Lemma 3.3.2. $\mathfrak{h}^0 \in \{0, 1\}$. More precisely,

(a) $\mathfrak{h}^0 = 0$ exactly when either (i) $D_0 \neq E$, or (ii) $D_0 = E$ but there exists E_i such that $\mathcal{O}_{\bar{X}}(K_{\bar{X}} + [Z_K])|_{E_i}$ is not trivial in $\text{Pic}^0(E_i) \simeq \mathbb{C}^{g_i}$. (In this second case, necessarily $g_i > 0$.)

(b) $\mathfrak{h}^0 = 1$ exactly when $D_0 = E$ and $\mathcal{O}_{\bar{X}}(K_{\bar{X}} + [Z_K])|_{E_i} = 0$ in $\text{Pic}^0(E_i)$ for every $i \in \mathcal{V}$.

Example 3.3.3. (a) If (X, o) is Gorenstein, then $\mathcal{O}_{\bar{X}}(K_{\bar{X}} + [Z_K]) = \mathcal{O}_{\bar{X}}$, and $\mathfrak{h}^0 = h^0(\mathcal{O}_E) = 1$.

(b) If $g = \sum_i g_i = 0$ then $\mathfrak{h}^0 = 1$ in the numerically Gorenstein case, and it is zero otherwise.

(c) If (X, o) is rational, then $g = 0$ automatically, hence $\mathfrak{h}^0 = 1$ in the ADE (and smooth) cases (the only numerically Gorenstein rational cases) and $\mathfrak{h}^0 = 0$ otherwise.

(d) If the link is an integral homology sphere (hence $L = L'$) then $g = 0$ and $Z_K \in L$, hence $\mathfrak{h}^0 = 1$.

(e) In general \mathfrak{h}^0 is not topological: for the germs from Example 3.2.4 $\mathfrak{h}^0 = 1$ in the Goresntein cases and 0 otherwise. (Use the formula $m(0) = \mathfrak{h}^0$ from Corollary 3.3.5(a').)

Proof of Lemma 3.3.2. Write $\mathcal{L} := \mathcal{O}_{\bar{X}}(K_{\bar{X}} + [Z_K])$. Then $c_1 \mathcal{L} = -\{Z_K\}$.

(a) Assume that $Z_K \notin L$, hence $\Delta_0 = \{Z_K\} \neq 0$. Recall that D_0 is reduced, let D' be one of its connected components. Then $(D', c_1 \mathcal{L}) < 0$. In particular, there exists $E_{i'}$ in the support of D' such that $(E_{i'}, c_1 \mathcal{L}) < 0$. Then we claim that $h^0(\mathcal{O}_{D'} \otimes \mathcal{L}) = 0$.

Indeed, consider a sequence of reduced connected divisors $\{D'_n\}_{n=1}^N$ such that $D'_N = D'$, $D'_{m+1} = D'_m + E_{i(m)}$ with $(D'_m, E_{i(m)}) > 0$ and $D'_1 = E_{i'}$. Then the vanishing follows inductively from the exact sequences $0 \rightarrow \mathcal{O}_{E_{i(m)}}(-D'_m) \otimes \mathcal{L} \rightarrow \mathcal{O}_{D'_{m+1}} \otimes \mathcal{L} \rightarrow \mathcal{O}_{D'_m} \otimes \mathcal{L} \rightarrow 0$.

Assume next that $Z_K \in L$ (hence $H_0 = E$) but $\mathcal{L}|_{E_{i'}}$ is not trivial in $\text{Pic}^0(E_{i'})$ for some i' . This means that $h^0(E_{i'}, \mathcal{L}|_{E_{i'}}) = 0$. Indeed, since $\mathcal{L}|_{E_{i'}}$ is topologically trivial (its Chern class is zero), any nonzero element of $H^0(E_{i'}, \mathcal{L}|_{E_{i'}})$ would provide a trivialization of $\mathcal{L}|_{E_{i'}}$.

Then we can repeat the above proof with a similar sequence with $D'_N = E$ and $D'_1 = E_{i'}$.

(b) It follows by induction using a similar sequence of divisors with $D'_N = E$ and $D'_1 = E_{i'}$ arbitrary. Indeed, $h^0(\mathcal{O}_E \otimes \mathcal{L}) = h^0(\mathcal{O}_{D'_m} \otimes \mathcal{L}) = h^0(\mathcal{O}_{E_{i'}} \otimes \mathcal{L}) = h^0(\mathcal{O}_{E_{i'}}) = 1$. \square

Proposition 3.3.4. (a) If $c \in \mathbb{R}_{\geq 0}$ then $\mathfrak{h}(L(c)) = \chi(L(c)) + p_g$.

Thus, $\dim \mathcal{O}_{X,o}/\mathfrak{J}(c, F) - p_g = \mathfrak{h}(L(c)) - p_g = \chi(L(c))$ is combinatorial computable from Γ .

In particular, if (X, o) is rational, then $\dim \mathcal{O}_{X,o}/\mathfrak{J}(c, F)$ is completely combinatorial.

(b) $m(c) = \chi(L(c)) - \chi(L(c - \varepsilon))$ for any $c > 0$, hence it is always combinatorial.

$m(0) = \chi(L(0)) - \chi(L(-\varepsilon)) + \mathfrak{h}^0$, hence $m^*(0) := m(0) - \mathfrak{h}^0$ is combinatorial.

The case $c > 0$ in part (b) for rational (X, o) was proved in [3, Prop. 4.10].

Proof. (a) Write $L(c) = P - N$ as in Lemma 3.1.1. Then, by this Lemma, $\mathfrak{h}(L(c)) = \mathfrak{h}(P)$. The short exact sequence of sheaves $0 \rightarrow \mathcal{O}_{\tilde{X}}(-P) \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow \mathcal{O}_P \rightarrow 0$ provides the exact sequence

$$0 \rightarrow H^0(\mathcal{O}_{\tilde{X}})/H^0(\mathcal{O}_{\tilde{X}}(-P)) \rightarrow H^0(\mathcal{O}_P) \rightarrow H^1(\mathcal{O}_{\tilde{X}}(-P)) \rightarrow H^1(\mathcal{O}_{\tilde{X}}) \rightarrow H^1(\mathcal{O}_P) \rightarrow 0.$$

By this sequence and part (b) of Lemma 3.1.1

$$\begin{aligned} \mathfrak{h}(P) &= \chi(P) - h^1(\mathcal{O}_{\tilde{X}}(-P)) + p_g \\ &= \chi(P - N + N) - \chi(\mathcal{O}_N(-P + N)) - h^1(\mathcal{O}_{\tilde{X}}(-P + N)) + p_g \\ &= \chi(P - N) + \chi(N) - (N, P - N) - (N, -P + N) - \chi(N) - h^1(\mathcal{O}_{\tilde{X}}(-P + N)) + p_g \\ &= \chi(L(c)) - h^1(\mathcal{O}_{\tilde{X}}(-L(c))) + p_g. \end{aligned}$$

Finally, we show that $h^1(\mathcal{O}_{\tilde{X}}(-L(c))) = 0$. To do this, set $\mathcal{L}' = \mathcal{O}_{\tilde{X}}(-L(c))$. Then $c_1(\mathcal{L}') = -(Z_K + cF) + \Delta_c$. Hence $c_1(\mathcal{L}'(-K_{\tilde{X}})) = \Delta_c - cF$, where $F \in \mathcal{S}'$. Thus we can apply the Generalized Grauert-Riemenschneider Theorem (see [27, Theorem 6.4.3.]), which gives $h^1(\tilde{X}, \mathcal{L}') = 0$.

Part (b) for $c > 0$ follows from part (a) whenever $c > 0$. Indeed, in that case $c - \varepsilon \geq 0$ too, hence (a) applies for both c and $c - \varepsilon$.

Next, we assume that $c = 0$. Then $L(0) = [Z_K]$. Consider the exact sequence $0 \rightarrow \mathcal{O}_{\tilde{X}}(-[Z_K]) \rightarrow \mathcal{O}_{\tilde{X}}(-[Z_K] + D_0) \rightarrow \mathcal{O}_{D_0}(-[Z_K] + D_0) \rightarrow 0$. Since by the Generalized Grauert-Riemenschneider Theorem (see [27, Theorem 6.4.3.]) $h^1(\mathcal{O}_{\tilde{X}}(-[Z_K])) = 0$, we get

$$\begin{aligned} m(0) &= \dim H^0(\mathcal{O}_{\tilde{X}}(-[Z_K] + D_0))/H^0(\mathcal{O}_{\tilde{X}}(-[Z_K])) = h^0(\mathcal{O}_{D_0}(-[Z_K] + D_0)) \\ &= \chi(\mathcal{O}_{D_0}(-[Z_K] + D_0)) + h^1(\mathcal{O}_{D_0}(-[Z_K] + D_0)) \\ &= \chi(L(0)) - \chi(L(-\varepsilon)) + h^1(\mathcal{O}_{D_0}(-[Z_K] + D_0)). \end{aligned}$$

But $h^1(\mathcal{O}_{D_0}(-[Z_K] + D_0)) = h^0(\mathcal{O}_{D_0}(K_{\tilde{X}} + [Z_K]))$ by Serre duality. \square

Corollary 3.3.5. (a) $m(c) = (D_c, \Delta_c) - \chi(D_c) - c(D_c, F)$ for $c > 0$.

$$(a') \quad m(0) = (D_0, \Delta_0) - \chi(D_0) + \mathfrak{h}^0.$$

$$(b) \quad m(c+1) = m(c) - (D_c, F) \text{ for } c > 0.$$

$$(b') \quad m(1) = m(0) - \mathfrak{h}^0 - (D_0, F).$$

Above, in (a)–(b'), for any fixed $c \geq 0$, the term (D_c, F) equals $\sum(E_i, F)$, where the sum is over those vertices $i = i(a) \in \mathcal{V}$ with $k_{i(a)} + cm_{i(a)} \in \mathbb{Z}$.

(c) Write $m^*(c)$ for $m(c)$ if $c > 0$ and $m(0) - \mathfrak{h}^0$ if $c = 0$. Then $J_{[0, \infty)}(t) := \sum_{c \geq 0} m(c)t^c$ (where t is a formal variable) equals

$$J_{[0, \infty)}(t) = \mathfrak{h}^0 t^0 + \sum_{c \in [0, 1)} t^c \cdot (m^*(c)/(1-t) - (D_c, F)t/(1-t)^2).$$

Proof. We prove (a), the others follow similarly or from this one.

$$\begin{aligned} m(c) &= \chi(L(c - \varepsilon) + D_c) - \chi(L(c - \varepsilon)) = \chi(D_c) - (D_c, L(c - \varepsilon)) \\ &= \chi(D_c) - (D_c, Z_K + cF - \Delta_c - D_c) = \chi(D_c) - 2\chi(D_c) + (D_c, \Delta_c - cF). \end{aligned} \quad \square$$

Example 3.3.6. For the rational graph from Example 3.2.1 one gets

$$J_{[0, \infty)}(t) = \frac{t}{(1-t)^2} (1 + t^{1/4} + t^{2/4} + t^{3/4}) + \frac{2}{1-t} t^{3/4} = 2t^{3/4} + t + t^{5/4} + t^{3/2} + 3t^{7/4} + \dots$$

An additional interesting connection is realized by the following sum.

Proposition 3.3.7.

$$\sum_{c \in (0,1]} m(c) = \chi(-F) + (\{Z_K\}, F).$$

Proof. By (3.1.5), the left hand side is $\chi(L(1)) - \chi(L(0)) = \chi(\lfloor Z_K \rfloor + F) - \chi(\lfloor Z_K \rfloor)$. \square

Corollary 3.3.8. *Assume that there exists a reduced Cartier divisor $(C, o) \subset (X, o)$, cut out by the function $f : (X, o) \rightarrow (\mathbb{C}, 0)$, such that $\text{div}_{\phi, E}(f) = F$, cf. 2.1.2. Then*

$$\sum_{c \in (0,1]} m(c) = \delta(C, o) + (\{Z_K\}, F),$$

where $\delta(C, o)$ is the delta invariant of the abstract analytic curve germ (C, o) .

Proof. For the identity $\delta(C, o) = \chi(-F)$ see [27, Example 6.3.8]. \square

For a generalization of the formula $\chi(-F) = \delta(C, o)$ when (X, o) is rational and (C, o) not necessarily Cartier, see [9, 10] (when an additional correction term appears).

3.4. The weighted homogenous case.

Our goal is to determine $\sum_{c \in \mathbb{R}} m(c)t^c$ completely and to discuss some examples.

3.4.1. The affine algebraic (complex) variety $Y \subset \mathbb{C}^n$ is called weighted homogeneous with integral weights $\{w_i\}_{i=1}^n$ if it is stable with respect to the action of \mathbb{C}^* on \mathbb{C}^n , $t * (x_1, \dots, x_n) = (t^{w_1}x_1, \dots, t^{w_n}x_n)$. We assume that (i) each w_i is positive (that is, the action is *good*: the origin is contained in the closure of any orbit), and (ii) $\text{gcd}\{w_i\}_i = 1$ (that is, the action is *effective*: if $t * x = x$ for all $x \in Y$ then $t = 1$). For such a variety Y the affine coordinate ring is $\mathbb{Z}_{\geq 0}$ -graded, $R = \bigoplus_{\ell \geq 0} R_\ell$. (Usually the normality of R is not guaranteed.)

A local normal surface singularity (X, o) is called weighted homogeneous if there exists a normal affine surface Y with a good and effective \mathbb{C}^* action, and a singular point $y \in Y$ such that (X, o) and (Y, y) are analytically isomorphic as complex analytic germs. In this case, the local algebra of (X, o) is graded: we use the same notation $\mathcal{O}_{X, o} = \bigoplus_{\ell \geq 0} R_\ell$.

We fix such a singularity and let ϕ be its minimal good resolution. The graph is star-shaped with ν legs (strings). Let $-b_0$ and g denote the Euler decoration and the genus of the central vertex v_0 . Each leg for $1 \leq j \leq \nu$ has Euler decorations $-b_{j1}, \dots, -b_{js_j}, b_{ji} \geq 2$, determined by the Hirzebruch continued fraction $\alpha_j/\omega_j = [[b_{j1}, \dots, b_{js_j}]]$, where $\text{gcd}(\alpha_j, \omega_j) = 1, 0 < \omega_j < \alpha_j$. In the graph, for each j the central vertex v_0 is connected to v_{j1} by an edge. We assume that either $g > 0$ or $\nu \geq 3$. (Otherwise, the graph is a string with all $g_i = 0$, hence (X, o) is a quotient singularity, or a numerically log terminal singularity with all $k_i < 1$; this case is treated in Example 3.1.9 and in subsection 3.3.)

The link is a Seifert 3-manifold with Seifert invariants $(b_0, g; \{(\alpha_j, \omega_j)\}_j)$. It has several important numerical invariants:

- the orbifold Euler number $e := -b_0 + \sum_j \omega_j/\alpha_j$ ($e < 0$ by the negative definiteness of $(,)$)
- the orbifold Euler characteristic $\chi := 2 - 2g - \sum_j (\alpha_j - 1)/\alpha_j$,
- $\tau = \chi/e$, where $-\tau$ is called the log discrepancy of (X, o) ,
- the order of $H = L'/L = \text{Tors}(H_1(M, \mathbb{Z}))$, it equals $|H| = |e| \cdot \prod_j \alpha_j$,

- the order of $[E_{v_0}^*]$ (of the generic S^1 orbit) in L'/L , it equals $\mathfrak{o} = |e|\alpha$, where $\alpha := \text{lcm}_j\{\alpha_j\}$,
- k_0 , the E_{v_0} -coefficient of Z_K , it equals $k_0 = 1 + \tau$,
- the E_{v_0} -coefficient of $E_{v_0}^*$, it equals $(\prod_j \alpha_j)/|H|$.

Associated with ϕ we choose $F := \mathfrak{o}E_{v_0}^*$. The rational cycle $E_{v_0}^*$ is associated with one arrow attached to the central vertex, and \mathfrak{o} is the smallest integer such that $\mathfrak{o}E_{v_0}^*$ is integral.

Note that m_{v_0} , the E_{v_0} -multiplicity of F , is $\mathfrak{o} \cdot \prod_j \alpha_j / |H| = \alpha$.

For any cycle l we denote by l_0 its E_{v_0} -coefficient.

Theorem 3.4.2. *Let (X, \mathfrak{o}) and F be as before. Then*

(a) *If $\mathfrak{J}(c, F) \subsetneq \mathfrak{J}(c - \varepsilon, F)$ then $k_0 + cm_{v_0} \in \mathbb{Z}$, and $\mathfrak{J}(c - \varepsilon, F)/\mathfrak{J}(c, F) \simeq R_{L(c)_0 - 1}$,*

(b)

$$\sum_{c \in \mathbb{R}} m(c)t^c = \sum_{\ell \geq 0} \dim(R_\ell)t^{(\ell+1-k_0)/\alpha} = \sum_{\ell \geq 0} \dim(R_\ell)t^{(\ell-\tau)/\alpha}.$$

Thus, $\sum_c m(c)t^c$ follows by a simple way from the Poincaré series $P(t) := \sum_{\ell \geq 0} \dim(R_\ell)t^\ell$ of (X, \mathfrak{o}) .

Theorem 3.4.2 follows from the next technical result.

Proposition 3.4.3. *For any $c \in \mathbb{R}$ and $s \in \mathcal{S}$, $L(c) \leq s$ if and only if $L(c)_0 \leq s_0$.*

Proof of Theorem 3.4.2 via Proposition 3.4.3. We consider the filtration $\{\mathcal{F}(\ell)\}_{\ell \geq 0}$ of $\mathcal{O}_{X, \mathfrak{o}}$ associated with the divisor E_{v_0} , i.e. $\mathcal{F}(\ell) := \{f \in \mathcal{O}_{X, \mathfrak{o}} : \text{div}_{\phi, E}(f) \geq \ell E_{v_0}\}$.

We claim that $\mathfrak{J}(c, F) = \mathcal{F}(L(c)_0)$.

Indeed, if $f \in \mathfrak{J}(c, F) = \phi_* \mathcal{O}_{\tilde{X}}(-L(c))$ then $\text{div}_{\phi, E}(f) \geq L(c) \geq L(c)_0 E_{v_0}$. Also, if $f \in \mathcal{F}(L(c)_0)$ then $\text{div}_{\phi, E}(f) \in \mathcal{S}$ and $\text{div}_{\phi, E}(f) \geq L(c)_0 E_{v_0}$, hence $f \in \phi_* \mathcal{O}_{\tilde{X}}(-L(c))$ too by the Proposition.

But, for homogeneous singularities the $\{\mathcal{F}(\ell)\}_\ell$ filtration coincides with the filtration induced by the \mathbb{C}^* -action: $\mathcal{F}(\ell) = \bigoplus_{\nu \geq \ell} R_\nu$, see e.g. [27, Example 8.6.13]. Therefore, $\mathfrak{J}(c, F) = \bigoplus_{\ell \geq L(c)_0} R_\ell$. If $\mathfrak{J}(c, F) \subsetneq \mathfrak{J}(c - \varepsilon, F)$ then necessarily $k_0 + cm_{v_0} \in \mathbb{Z}$, and $\mathfrak{J}(c - \varepsilon, F) = \bigoplus_{\ell \geq L(c)_0 - 1} R_\ell$. \square

Proof of Proposition 3.4.3. Fix $s \in \mathcal{S}$ and assume that $s_0 \geq L(c)_0$. Then $(Z_K + cF - E, E_v) = 0$ for any vertex except for the node and the end vertices, and it equals 1 for any end vertex. Therefore, $(s - Z_K - cF + E, E_v) \leq 0$ for any $v \neq v_0$. On the other hand, $(s - Z_K - cF + E)_0 = s_0 - (Z_K + cF)_0 + 1 \geq r$, where $r := (\Delta_c)_0 + 1 > 0$.

Recall that $(E_{v_0}^*)_0 = \prod_j \alpha_j / |H|$, and set $r' := (r|H|) / \prod_j \alpha_j > 0$. Then $(r'E_{v_0}^*)_0 = r$. This shows that the cycle

$$C := s - Z_K - cF + E - r'E_{v_0}^*$$

is supported on the legs, and $(C, E_v) \leq 0$ for any vertex v of the legs. Thus, C is in the Lipman cone of the legs, hence $C \geq 0$. This reads as

$$s \geq Z_K + cF - E + r'E_{v_0}^* \geq L(c) - E + r'E_{v_0}^*.$$

Finally use the facts that all the coefficients of $E_{v_0}^*$ are strict positive and s is an integral cycle. \square

Example 3.4.4. (i) Assume that M is an integral homology sphere. Then it is the Seifert 3-manifold $\Sigma(\alpha_1, \dots, \alpha_\nu)$. Then, by [18, 28] (X, \mathfrak{o}) is a Brieskorn-Hamm complete intersection, the $\{\alpha_j\}_j$'s are pairwise relative prime, $\alpha = \prod_j \alpha_j$, and its Poincaré series is

$$P(t) = \frac{(1-t^\alpha)^{\nu-2}}{\prod_j (1-t^{\alpha/\alpha_j})}.$$

(ii) If M is a rational homology sphere (equivalently, $g = 0$) then again, the Poincaré series can be determined from the Seifert invariants by the Dolgachev-Pinkham-Demazure formula [32]

$$P(t) = \sum_{\ell \geq 0} \max\{0, 1 + \ell b_0 - \sum_j [\ell \omega_j / \alpha_j]\} \cdot t^\ell.$$

(iii) If (X, o) is a hyperurface singularity given by an isolated quasihomogeneous germ with weights (w_1, w_2, w_3) and degree d then

$$P(t) = \frac{1 - t^d}{\prod_i (1 - t^{w_i})}.$$

Example 3.4.5. (a) Consider the weighted homogeneous hypersurface germs (i) and (ii) from Example 3.2.4. In that Example we considered $F = E = E_3^*$, which does not apply to the discussion from this subsection 3.4. Hence, let us take at this time $F = E_{v_0}^* = Z_K$. Then by a very similar computation $\sum_{c \leq 0} m_{(i)}(c)t^c = t^{-2/3} + t^{-1/3} + t^0$ in case (i), and $\sum_{c \leq 0} m_{(ii)}(c)t^c = t^{-2/3} + t^0$ in case (ii). On the other hand, for $c > 0$ (by subsection 3.3) the values $m(c)$ should agree (since the germs share the same graph). Hence $\sum_c m_{(i)}(c)t^c - \sum_c m_{(ii)}(c)t^c = t^{-1/3}$.

Let us compare this fact with the statement of Theorem 3.4.2. In case (i), $z^2 = y^3 + x^{18}$ has weights $(1, 6, 9)$ and degree 18. In case (ii), $z^2 = x(y^4 + x^6)$, the weights are $(2, 3, 7)$ and the degree is 14. Hence the Poincaré series are

$$P_{(i)}(t) = \frac{1 - t^{18}}{(1 - t)(1 - t^6)(1 - t^9)} \quad \text{and} \quad P_{(ii)}(t) = \frac{1 - t^{14}}{(1 - t^2)(1 - t^3)(1 - t^7)}.$$

Since the ‘substitution’ is $t^\ell \mapsto t^{(\ell-2)/3}$, this is indeed in agreement with the above computation since (rather interestingly) $P_{(i)}(t) - P_{(ii)}(t) = t$ (and the above substitution realizes $t \mapsto t^{-1/3}$).

(b) The above computation, valid for $F := E_{v_0}^* = E_1^* = Z_K$ can be compared with the computation from Example 3.2.4. In fact, let us take $F' := 3E_3^* = 3E$. Then the multiplicities in the cycles $F = Z_K$ and $F' = 3E$ of the central vertex agree. Then, surprisingly, the computation from Example 3.2.4 and from above show that the expression $\sum_{c \leq 0} m(c)t^c$, computed individually for any choice (i)–(iii) of the analytic type, is the same for F and F' (though F' is not a rational multiple of $E_{v_0}^*$).

This might suggest that if we drop the assumption that F is a rational multiple of $E_{v_0}^*$, then in the computation of $\sum_c m(c)t^c$, associated with F , only the the central vertex multiplicity of F counts. Well, for $c > 0$ this is not the case. For $c > 0$, though $\sum_{c > 0} m(c)t^c$ is independent of the choice of the analytic type (i)–(iii), it turns out that it depends on the choice of F and F' :

$$\sum_{c > 0} m(c)t^c = \begin{cases} \sum_{n > 0} \left\lceil \frac{n}{3} \right\rceil t^{n/3} & \text{for } F, \\ \sum_{n > 0} n t^{n/3} & \text{for } F'. \end{cases}$$

Remark 3.4.6. (a) Let us consider again the formula from Theorem 3.4.2(b), which gives the identification $m((\ell - \tau)/\alpha) = \dim(R_\ell)$. Though each homogeneous component R_ℓ of the local algebra of (X, o) a priori is analytical, in subsection 3.3 proved that $m(c)$ for $c > 0$ is topological. Hence, via our formula $\dim(R_\ell)$ is topological for any $\ell > \tau$. This can be seen in the theory of weighted homogeneous singularities as well. In that case the h^1 of a certain line bundle is vanishing, whose h^0 is exactly R_ℓ (for details see e.g. [32] or [27, Remark 5.1.29]).

(b) On the other hand, the formula $(\dagger) \sum_{c \leq 0} m(c) = \chi([Z_K]) + p_g$ from Example 3.2.2 at this generality is new even from the point of view of the Dolgachev-Pinkham-Demazure theory of

weighted homogeneous singularities (reinterpreted via Theorem 3.4.2(b)). Indeed, in the Gorenstein case, when $\chi([Z_K]) = 0$, it is known that by duality $\sum_{\ell \leq \tau} \dim(R_\ell)t^\ell$ corresponds to a finite Poincaré series of $H^1(\bar{X}, \mathcal{O}_{\bar{X}})$ (graded by the \mathbb{C}^* -action), see e.g. [27, Example 6.3.30 (d)]. Hence, $\sum_{\ell \leq \tau} \dim R_\ell = p_g$. However, for non-Gorenstein germ, the ‘broken duality’ formula (†), even at this numerical level $\sum_{\ell \leq \tau} \dim R_\ell = p_g + \chi([Z_K])$ was unknown (at least for the authors).

4. JUMPING MULTIPLICITIES FROM THE MULTIPLICITIES OF F

4.1. The definition of the multiplicities $\{m_i\}_{i \in \mathcal{V}}$.

4.1.1. Let us fix the resolution graph Γ with vertices \mathcal{V} and the cycle $F \in \mathcal{S} \setminus \{0\}$ as above. We write $F = \sum_{i \in \mathcal{V}} m_i E_i$. We complete Γ and its decorations (self-intersection numbers $e_i = E_i^2$, genera g_i and multiplicities m_i , all indexed by \mathcal{V}) to a ‘virtual embedded resolution graph’. We complete the vertices of Γ with $-(E, F)$ arrowhead vertices, each vertex $i \in \mathcal{V}$ will support $-(E_i, F)$ arrows, and each of them will be decorated by multiplicity 1. Let \mathcal{A} denote the set of arrows, $\mathcal{W} := \mathcal{V} \cup \mathcal{A}$. Hence each element of \mathcal{W} has a multiplicity, $\{m_i\}_{i \in \mathcal{V}}$ is given by F , all others are $m_a = 1$ for all $a \in \mathcal{A}$. (If we fix the topological type of (X, o) by Γ , then for a convenient analytic type of (X, o) there exists a function $f : (X, o) \rightarrow (\mathbb{C}, 0)$ such that $\text{div}_{\phi, E}(f) = F$ and the embedded resolution graph associated with ϕ and f is this completed graph with arrows and multiplicities $\{m_v\}_{v \in \mathcal{W}}$.)

For any $v \in \mathcal{V}$ let \mathcal{W}_v be the set of all (usual and arrowhead) vertices adjacent with v , and $\mathcal{V}_v = \mathcal{W}_v \cap \mathcal{V}$, $\mathcal{A}_v = \mathcal{W}_v \cap \mathcal{A}$.

Finally, we consider certain rational numbers $\{\beta_v\}_{v \in \mathcal{V}}$, $\beta_v \in [0, 1)$. It is convenient to set also $\beta_a := 0$ for any $a \in \mathcal{A}$. (For a geometric interpretation see Theorem 4.2.3 and subsection 5.3.)

4.2. The definition of $TSp_{[0,1)}$.

4.2.1. Next we define a combinatorial invariant $TSp_{[0,1)} = TSp_{[0,1)}(\{\beta_v\}_{v \in \mathcal{V}})$ associated with the completed Γ , multiplicity system $\{m_v\}_{v \in \mathcal{W}}$ and rational numbers $\{\beta_v\}_{v \in \mathcal{V}}$ ($\beta_v \in [0, 1)$). Motivated by the connections from section 5, we will call it ‘topological spectrum’.

For any $v \in \mathcal{V}$ and $s \in \{0, 1, \dots, m_v - 1\}$ we set

$$R_v^s(\beta) = \sum_{w \in \mathcal{W}_v} \left\{ -\beta_w + \frac{m_w}{m_v}(s + \beta_v) \right\} \text{ and } \alpha_v^s = \frac{s + \beta_v}{m_v}.$$

Then we define

$$TSp_{[0,1),v}^\beta = \sum_{s=0}^{m_v-1} (R_v^s(\beta) + g_v - 1)t^{\alpha_v^s}.$$

Let \mathcal{E} be the set of edges of Γ . If $e \in \mathcal{E}$ connects $v, w \in \mathcal{V}$, then we set $m_e := \text{gcd}(m_v, m_w)$. Also, we consider the following system of equations in γ_e

$$(4.2.2) \quad \begin{cases} \beta_v = \{m_v \gamma_e / m_e\} \\ \beta_w = \{m_w \gamma_e / m_e\}. \end{cases}$$

Either it has a solution $\gamma_e \in [0, 1)$ or it does not. If it has a solution then that solution is unique in $[0, 1)$. We also introduce

$$\gamma_e^s = \begin{cases} (\gamma_e + s)/m_e & \text{if } \gamma_e \in [0, 1) \text{ exists.} \\ -\infty & \text{otherwise.} \end{cases}$$

and define $TSp_{[0,1),e}^\beta = \sum_{s=0}^{m_e-1} t^{\gamma_e^s}$, where $t^{-\infty} = 0$.

Finally let

$$TSp_{[0,1)}^\beta := \sum_{v \in \mathcal{V}} TSp_{[0,1),v}^\beta + \sum_{e \in \mathcal{E}} TSp_{[0,1),e}^\beta.$$

Theorem 4.2.3. *Fix the resolution with graph Γ and $F \in \mathcal{S} \setminus \{0\}$. Consider the complete graph with multiplicities $\{m_w\}_{w \in \mathcal{W}}$ and $\{\beta_w\}_{w \in \mathcal{W}}$ defined as $\beta_v := \{-k_v\}$ for $v \in \mathcal{V}$ and $\beta_a = 0$ for $a \in \mathcal{A}$.*

Then the jumping multiplicities associated with jumping numbers $c \in [0, 1)$ are determined as follows:

$$\sum_{c \in [0,1)} m^*(c)t^c = -\mathfrak{h}^0 t^0 + \sum_{c \in [0,1)} m(c)t^c = TSp_{[0,1)}^\beta.$$

In the proof we will need the following lemma.

Lemma 4.2.4. *For a fixed $c \in [0, 1)$ and an edge $e \in \mathcal{E}$ which connects v and w , the following two facts are equivalent.*

(1) *The system (4.2.2) admits a solution $\gamma_e \in [0, 1)$ and there exists $s_e \in \{0, 1, \dots, m_e - 1\}$ such that $(\gamma_e + s_e)/m_e = c$,*

(2) *there exist $s_v \in \{0, 1, \dots, m_v - 1\}$ and $s_w \in \{0, 1, \dots, m_w - 1\}$ such that $(\beta_v + s_v)/m_v = (\beta_w + s_w)/m_w = c$.*

The pairs (γ_e, s_e) and (s_v, s_w) determine each other in a unique way.

Proof. If (1) holds then $s_v := \lfloor (\gamma_e + s_e)m_v/m_e \rfloor$ and $s_w := \lfloor (\gamma_e + s_e)m_w/m_e \rfloor$ satisfy (2). On the other hand, if (2) is satisfied by (s_v, s_w) , then $\beta_v = \{cm_v\}$ and $\beta_w = \{cm_w\}$. Hence if we write $cm_e = \gamma_e + s_e$ (i.e. $\gamma_e := \{cm_e\}$) then $\gamma_e \in [0, 1)$ is a solution of the system and $(\gamma_e + s_e)/m_e = c$ too. \square

Proof of Theorem 4.2.3. Using the definitions of $\{\beta_w\}_{w \in \mathcal{W}}$ we have $\alpha_v^s = \frac{s + \{-k_v\}}{m_v}$, and $R_v^s(\beta) = \sum_{w \in \mathcal{W}_v} \{k_w + \frac{m_w}{m_v}(s + \{-k_v\})\}$. In the proof below we just write R_v^s for $R_v^s(\beta)$.

Let us rewrite $TSp_{[0,1)}^\beta$ as $\sum_{c \in [0,1)} \tau(c)t^c$. We want to show that $\tau(c) = m^*(c)$.

For a fixed $c \in [0, 1)$ we have

$$(4.2.5) \quad \tau(c) = \sum_{v \in \mathcal{V}, \exists s \alpha_v^s = c} (R_v^s + g_v - 1) + \sum_{e \in \mathcal{E}, \exists s \gamma_e^s = c} 1.$$

Then we apply Lemma 4.2.4 and the fact that $-\beta_v + cm_v \in \mathbb{Z}$ if and only if v is in the support of D_c (and we use the notation $\mathcal{V}(D_c)$ and $\mathcal{E}(D_c)$ for the vertices and edges of D_c). Thus,

$$(4.2.6) \quad \tau(c) = \sum_{v \in \mathcal{V}, \exists s \alpha_v^s = c} (R_v^s + g_v - 1) + \sum_{e \in \mathcal{E}, \exists s \gamma_e^s = c} 1$$

$$(4.2.7) \quad = \sum_{v \in \mathcal{V}(D_c)} (R_v^{cm_v - \{-k_v\}} + g_v - 1) + \#\{e \in \mathcal{E}(D_c)\}$$

$$(4.2.8) \quad = \sum_{v \in \mathcal{V}(D_c)} R_v^{cm_v - \{-k_v\}} + \sum_{v \in \mathcal{V}(D_c)} g_v + \#\{e \in \mathcal{E}(D_c)\} - \#\{v \in \mathcal{V}(D_c)\}$$

$$(4.2.9) \quad = \sum_{v \in \mathcal{V}(D_c)} R_v^{cm_v - \{-k_v\}} - \chi(D_c).$$

Furthermore, for any $v \in \mathcal{V}(D_c)$,

$$(4.2.10) \quad R_v^{cm_v - \{-k_v\}} = \sum_{w \in \mathcal{W}_v} \{k_w + \frac{m_w}{m_v}(cm_v - \{-k_v\} + \{-k_v\})\}$$

$$(4.2.11) \quad = \sum_{u \in \mathcal{V}_v} \{k_u + m_u c\} + \sum_{a \in \mathcal{A}_v} c$$

$$(4.2.12) \quad = (\{Z_K + cF\}, E_v) - (cF, E_v).$$

Hence, via Corollary 3.3.5(a)

$$(4.2.13) \quad \tau(c) = \left(\sum_{v \in \mathcal{V}(D_c)} (\Delta_c, E_v) - (cF, E_v) \right) - \chi(D_c)$$

$$(4.2.14) \quad = (\Delta_c, D_c) - (cF, D_c) - \chi(D_c) = m^*(c) \quad \square$$

5. JUMPING MULTIPLICITIES AND HODGE SPECTRUM

5.1. The Milnor package and Hodge spectrum of an analytic germ.

5.1.1. Let (X, o) be a normal surface singularity and $f : (X, o) \rightarrow (\mathbb{C}, 0)$ an analytic germ which defines an isolated singularity $(C, o) := (f^{-1}(0), o) \subset (X, o)$. Let F_o be the Milnor fiber of f . It is a 1-dimensional Stein space, hence its cohomology $H^n := H^n(F_o, \mathbb{C})$ can be nonzero only for $n = 0, 1$. Moreover, F_o is connected, hence $H^0 = \mathbb{C}$. Let T^* be the monodromy operator $T^* : H^* \rightarrow H^*$. In fact, $T^0 = \text{id}_{\mathbb{C}}$. The operator T^n defines a generalized eigenspace decomposition $H^n = \bigoplus_{\lambda} H_{\lambda}^n$. The corresponding dimensions are codified in the monodromy zeta function

$$\zeta(t) = \prod_{\lambda} (t - \lambda)^{\dim H_{\lambda}^1 - \dim H_{\lambda}^0} = \det(t \cdot \text{Id} - T^1) / \det(t \cdot \text{Id} - T^0).$$

If ϕ is a good embedded resolution of $(C, o) \subset (X, o)$ (and we use the notations of 4.1.1) then by A'Campo's Theorem [1]

$$\zeta(t) = \prod_{v \in \mathcal{V}} (1 - t^{m_v})^{2g_v - 2 + \#\mathcal{W}_v}.$$

In particular, $\dim H_1^0 = 1$ and $\dim H_1^1 = \#\mathcal{A} + 2g + 2h - 1$.

By [36, 33, 34] H^1 carries a mixed Hodge structure, set $H^{pq} := Gr_{p+q}^W Gr_F^p H^1$. The semisimple part of T^1 acts on each H^{pq} , let H_{λ}^{pq} be the eigenspace with dimension h_{λ}^{pq} . Then the spectrum of

H^1 is defined as $Sp(H^1) := \sum_{p,q,|\alpha|=p} h_{e^{2\pi i\alpha}}^{pq} t^\alpha$. One defines similarly the spectrum $Sp(H^0)$ of H^0 too, but in fact it is t^0 . Hence, the ‘total’ (Euler characterisitic type) spectrum of f , as lifting of ζ , is

$$Sp(f,t) := Sp(H^1) - Sp(H^0) = -t^0 + \sum_{p,q,|\alpha|=p} h_{e^{2\pi i\alpha}}^{pq} t^\alpha \in \mathbb{Z}[\mathbb{Q}].$$

We write $Sp(f,t) = \sum_{\alpha} \sigma(\alpha) t^\alpha$. It turns out that $\sigma(\alpha) = 0$ whenever $\alpha \notin [0,2)$, $\sigma(0) = g + h - 1$ and $\sigma(1) = \#\mathcal{A} + g + h - 1$. Moreover, for any $\alpha \in (0,1)$, one has the symmetry $\sigma(\alpha) = \sigma(2 - \alpha)$. Hence $Sp(f,t)$ is completely determined by $Sp_{[0,1]}(f,t) := \sum_{\alpha \in [0,1]} \sigma(\alpha) t^\alpha$.

In the next theorem we fix an embedded resolution and its decorated graph (as in 4.1.1).

Theorem 5.1.2. [35, 24]

$$Sp_{[0,1]}(f,t) = (g + h - 1)t^0 + (\#\mathcal{A} + g + h - 1)t^1 + \sum_{e \in \mathcal{E}} \sum_{0 < s < m_e} t^{\frac{s}{m_e}} + \sum_{v \in \mathcal{V}} \sum_{0 < s < m_v} (R_v^s + g_v - 1)t^{\frac{s}{m_v}},$$

where $R_v^s := \sum_{w \in \mathcal{W}_v} \{sm_w/m_v\}$.

Note that the right hand side is exactly $TSP_{[0,1]}^{\beta=0}$. Hence, Theorems 4.2.3 and 5.1.2 combined give

Corollary 5.1.3. *Assume that (X,o) is a numerically Gorenstein singularity, and $f : (X,o) \rightarrow (\mathbb{C},0)$ is an analytic germ with $(f^{-1}(0),o)$ isolated singularity. Fix an embedded good resolution ϕ and set $F := \text{div}_{\phi,E}(f) \in \mathcal{S} \setminus \{0\}$. Then*

$$(5.1.4) \quad \sum_{c \in [0,1]} m^*(c)t^c = -\mathfrak{h}^0 t^0 + \sum_{c \in [0,1]} m(c)t^c = TSP_{[0,1]}^{\beta=0} = Sp_{[0,1]}(f,t).$$

This identity was proved in case $(X,o) = (\mathbb{C}^2,0)$ by Budur [5].

Though both invariants $J_{[0,1]}(t)$ and $Sp_{[0,1]}(f,t)$ are a priori analytical, and they might depend on the analytic structure of (X,o) , it turns out that this is not the case, and both of them equal the combinatorial expression $TSP_{[0,1]}^{\beta=0}$.

Recall that if (X,o) is smooth then in the above formulae $m^*(0) = -1$, $\mathfrak{h}^0 = 1$, $m(0) = 0$, $m(1) = \#\mathcal{A} - 1$ and $g = h = 0$. The $-t^0$ term corresponds to $Sp(H^0)$.

5.1.5. If (X,o) is not numerically Gorenstein then the identity $\sum_{c \in [0,1]} m^*(c)t^c = Sp_{[0,1]}(f,t)$ is not true anymore. It fails already for rational germs (X,o) .

Example 5.1.6. Consider the rational graph Γ as in Example 3.2.1. Assume that f is a germ whose strict transform corresponds to the arrow of the graph, and $F = \text{div}_{\phi,E}(f)$. Then $J_{[0,1]}(t) = 2t^{3/4} + t$, cf. 3.3.6. But $Sp_{[0,1]}(f,t) = -t^0 + t^{1/4} + t^{2/4} + t^{3/4}$, hence they are different.

The above example shows that we have to modify the above spectrum expression (which equals $TSP_{[0,1]}^{\beta=0}$) if we wish to recover $TSP_{[0,1]}^{\beta}$, where β is determined by $\{Z_K\}$ as in Theorem 4.2.3. This is the subject of the next subsections. We have to replace the cohomology of the Milnor fiber $H^*(F_o, \mathbb{C})$ by a cohomology with coefficient in a mixed Hodge module. That is, we will replace the constant pure Hodge module with a different Hodge module of (X,o) . The needed mixed Hodge module will be determined by a variation of Hodge structures on $X \setminus \{0\}$, in fact, by a finite order character of $\pi_1(X \setminus \{0\})$. This can be identified in many different ways, one conceptual way is via a (regular) covering of $X \setminus \{0\}$. If the link is a rational homology sphere then we can rely on the topological canonical covering, if (X,o) is \mathbb{Q} -Gorenstein then we can rely on the (analytic) canonical covering (see the construction below).

5.2. Mixed Hodge modules on (X, o) .

5.2.1. Let $MHM(X, o)$ be the category of *mixed Hodge modules* on (X, o) [33]. If $f : (X, o) \rightarrow (\mathbb{C}, 0)$ defines an isolated singularity $(C, o) \subset (X, o)$ then there exists a nearby cycle functor $\psi_f : MHM(X, o) \rightarrow MHM(C, o)$ [33]. It has a functor automorphism T_s of finite order (the semisimple part of the monodromy action). The nearby cycle functor is compatible with the corresponding functor at the level of perverse sheaves [11] by the forgetful functor $rat : MHM(X, o) \rightarrow Perv(X, o)$.

A mixed Hodge module \mathcal{M} is called *smooth* if $rat(\mathcal{M})$ is a local system [33]. A module \mathcal{M} is called *pure of weight n* (or polarized Hodge module of weight n) if $Gr_i^W \mathcal{M} = 0$ for $i \neq n$. The category of smooth polarized Hodge modules is equivalent with the category of (admissible) variations of polarized Hodge structures.

The relation with the Milnor fibers from the previous subsection 5.1 is the following. Let $i_o : \{o\} \rightarrow (C, o)$ be the inclusion. Then $H^*(i_o^* \psi_f \mathbb{C}_{X, o}) = H^*(F_o, \mathbb{C})$.

The point is that if we replace the constant Hodge module by an arbitrary $\mathcal{M} \in MHM(X, o)$, then each $H^k(i_o^* \psi_f \mathcal{M})$ carries a mixed Hodge structure (together with a finite monodromy action), hence we can define its spectrum $Sp(H^k(i_o^* \psi_f \mathcal{M}))$ similarly as we defined $Sp(H^k)$ in the previous subsection. Moreover, we set $Sp_\psi(f, \mathcal{M}, t)$ as $\sum_k (-1)^k Sp(H^k(i_o^* \psi_f \mathcal{M}))$.

Here we will construct \mathcal{M} as $j_* \mathbb{V}$, where $j : X \setminus \{o\} \rightarrow X$ is the inclusion and \mathbb{V} is a polarized variation of Hodge structure on $X \setminus \{o\}$ with finite cyclic rank one underlying representation.

5.3. The construction of \mathcal{M} when the link of (X, o) is a rational homology sphere.

5.3.1. Assume that the link M of (X, o) is a rational homology sphere. This means that in any good resolution $H_1(M, \mathbb{Z}) = L'/L$. Let \widehat{H} be the Pontrjagin dual of H , the group of characters $Hom(H, S^1)$. It is known that there exists an isomorphism $\theta : H \rightarrow \widehat{H}$, $[l'] \mapsto e^{2\pi i(l', \cdot)}$, where $[l']$ denotes the class of $l' \in L'$ in L'/L . In particular, any $l' \pmod{L}$ gives a character of H , hence by the composition $\pi_1(M) \rightarrow H_1(M, \mathbb{Z}) \rightarrow S^1$ a finite order character of $\pi_1(M) = \pi_1(X \setminus \{o\})$. The corresponding flat bundle can be lifted to the category of polarized variation of Hodge structures on $X \setminus \{o\}$, this will be denoted by $\mathbb{V}_{[l']}$.

$\mathbb{V}_{[l']}$ can also be constructed as follows. Let N be the order of $[l']$ in H . Then the above representation $\rho_{[l']} : \pi_1(X \setminus \{o\}) \rightarrow H_1(M, \mathbb{Z}) \rightarrow \mathbb{Z}_N$ determines a \mathbb{Z}_N -Galois o -ramified covering $cov : (X_{[l]}, o) \rightarrow (X, o)$, where $(X_{[l]}, o)$ is a normal surface singularity, and the restriction $X_{[l]} \setminus \{o\} \rightarrow X \setminus \{o\}$ is the regular covering associated with $\rho_{[l']}$. This follows from Stein theorem [38]. Then $cov_* \mathbb{C}_{X_{[l]} \setminus \{o\}} = \bigoplus_{i=1}^N \mathbb{V}^i$, where each \mathbb{V}^i is a variation of Hodge structures whose underlying monodromy representation is $\rho_{[l']}^i$. Our variation we searching for is $\mathbb{V}_{[l]} := \mathbb{V}^1$.

Theorem 5.3.2. [24] *Assume that the link of (X, o) is a $\mathbb{Q}HS^3$. Let $f : (X, o) \rightarrow (\mathbb{C}, 0)$ be an isolated singularity. Consider the variation $\mathbb{V}_{[l]}$, and define the rational numbers $\beta_v \in [0, 1)$ by*

$$\rho_{[l]}(-E_v^*) = e^{2\pi i(l', -E_v^*)} = e^{2\pi i \beta_v} \text{ for any } v \in \mathcal{V}.$$

For any $a \in \mathcal{A}$ set $\beta_a = 0$. Define $\mathcal{M}_{[l]} := j_* \mathbb{V}_{[l]}$, where $j : X \setminus \{o\} \rightarrow X$ is the inclusion. Then

$$Sp_{[0,1]}(f, \mathcal{M}_{[l]}, t) = TSP_{[0,1]}^\beta.$$

Hence, Theorems 4.2.3 and 5.1.2 combined give

Corollary 5.3.3. *Consider the situation of Theorem 5.3.2. Let us fix a good embedded resolution ϕ and set $F := \text{div}_{\phi,E}(f)$. Furthermore, define $\beta_v = \{-k_v\}$ for $v \in \mathcal{V}$ and $\beta_a = 0$ for $a \in \mathcal{A}$. Then*

$$(5.3.4) \quad \sum_{c \in [0,1]} m^*(c)t^c = -\mathfrak{h}^0 t^0 + \sum_{c \in [0,1]} m(c)t^c = \text{TS}p_{[0,1]}^\beta = \text{Sp}_{[0,1]}(f, \mathcal{M}_{[-Z_K]}, t).$$

The covering cov considered above, associated with $-Z_K$, is called the *topological canonical covering* of (X, o) (cf. [27, 6.1.19]). (Note that the coverings associated with $[l']$ and $[-l']$ agree, cf. [27, Remark 4.1.36].)

5.4. The construction of \mathcal{M} when (X, o) is \mathbb{Q} -Gorenstein.

5.4.1. A Weil divisor of (X, o) is called \mathbb{Q} -Cartier if its class in $\text{Cl}(X, o)$ (the local divisor class group of (X, o)) has finite order. Its order is called its index. If D is a \mathbb{Q} -Cartier divisor of index N then it determines a o -ramified Galois \mathbb{Z}_N -covering $\text{cov} : (X_D, o) \rightarrow (X, o)$, here (X_D, o) is a normal surface singularity. The covering depends only on the class of D in $\text{Cl}(X, o)$. For details see [20, 2.49], [30, 39] or [27, Proposition 6.1.10].

Write $\Omega_X^2 = \mathcal{O}_{\bar{X}}(K_{\bar{X}})$ and set $K_X = \phi_*(K_{\bar{X}})$, well-defined in $\text{Cl}(X, o)$. We say that (X, o) is Gorenstein (respectively \mathbb{Q} -Gorenstein) if $K_X = 0$ (respectively K_X has finite order in $\text{Cl}(X, o)$). If (X, o) is \mathbb{Q} -Gorenstein then the associated cyclic covering is called the ‘(analytic) canonical covering’.

Assume that the link is a rational homology sphere and (X, o) is \mathbb{Q} -Gorenstein. Then the coverings $\text{cov} : (X_{[l]}, o) \rightarrow (X, o)$ and $\text{cov} : (X_{K_X}, o) \rightarrow (X, o)$ agree [27,]. Thus, if we replace the covering $\text{cov} : (X_{[l]}, o) \rightarrow (X, o)$ from the previous subsection with the covering $\text{cov} : (X_{K_X}, o) \rightarrow (X, o)$, in a totally similar way we can define (via $\text{cov}_* \mathbb{C}_{X_{K_X} \setminus \{o\}}$ and \mathbb{V}^1) a variation of Hodge structure \mathbb{V}_{K_X} on $X \setminus \{o\}$ (which agrees with $\mathbb{V}_{[Z_K]}$, and it has the same underlying monodromy representation and the same β -coefficients). Hence, if we take $\mathcal{M}_{K_X} := j_* \mathbb{V}_{K_X}$ then by [24] and Theorem 4.2.3

$$(5.4.2) \quad \sum_{c \in [0,1]} m^*(c)t^c = -\mathfrak{h}^0 t^0 + \sum_{c \in [0,1]} m(c)t^c = \text{TS}p_{[0,1]}^\beta = \text{Sp}_{[0,1]}(f, \mathcal{M}_{K_X}, t).$$

Remark 5.4.3. Note that the jumping multiplicity/spectral number correspondence of Corollary 5.3.3 is realized for a character (or β) associated with the cycle $-Z_K$. The general construction of $\text{TS}p^\beta$, valid for any β , and the general Theorem 5.3.2, valid for any character, suggest that there exists a more general correspondence, where one has to generalize the multiplier ideals as well (by replacing the cycle Z_K by a more general cycle). We will return back to such generalizations in a forthcoming manuscript.

REFERENCES

- [1] A’Campo, N.: La fonction zêta d’une monodromie, *Comment. Math. Helv.* 50 (1975), 233–248.
- [2] Alberich-Carraminana, M., Dachs-Cadefau, F. and Montaner, J.A.: Multiplier ideals in two-dimensional local rings with rational singularities, *Michigan Mathematical Journal* 65.2 (2016): 287–320.
- [3] Alberich-Carraminana, M., Dachs-Cadefau, F., González-Alonso, V. and Montaner, J.A.: Poincaré series of multiplier ideals in two-dimensional local rings with rational singularities, *Advances in Mathematics* 304 (2017): 769–792.
- [4] Baumers, H., Veys, W., Smith, K.E. and Tucker, K.F.: Contribution of jumping numbers by exceptional divisors. *Journal of Algebra* 496 (2018): 24–47.

- [5] Budur, N.: On Hodge spectrum and multiplier ideal, *Math. Ann.* 327 (2003), no. 2, 257–270.
- [6] Campillo, A., Delgado de la Mata, F. and Gusein-Zade, S. M.: Poincaré series of a rational surface singularity, *Invent. math.* **155** (1), 41–53 (2004).
- [7] Campillo, A., Delgado de la Mata, F. and Gusein-Zade, S. M.: Universal Abelian covers of rational surface singularities and multi-index filtrations, *Functional Analysis and Its Appl.* **42** (2), 83–88 (2008).
- [8] Campillo, A., Delgado de la Mata, F. and Gusein-Zade, S. M.: Poincaré series of curves on rational surface singularities, *Comment. Math. Helv.* 80, 95–102 (2005).
- [9] Cogolludo-Agustín, J.I., László, T., Martín-Morales, J. and Némethi, A.: Delta invariant of curves on rational surfaces I. An analytic approach, *Communications in Contemporary Mathematics* **24**, No. 07 (2022), 2150052.
- [10] Cogolludo-Agustín, J.I., László, T., Martín-Morales, J. and Némethi, A.: Local invariants of minimal generic curves on rational surfaces, *Proceedings of Escuela de Matemáticas Lluís Santaló 2019: p-adic Analysis, Arithmetic and Singularities*, Contemporary Math.
- [11] Deligne, P.: Le formalisme des cycles évanescents, *SGA VII², Exp XIII*, lecture Notes in Math. 340 (1973), 82–115.
- [12] Ein, L., Lazarsfeld, R., Smith, K.E. and Varolin, D.: Jumping coefficients of multiplier ideals, *Duke Math. J.* 123 (2004): 469-506.
- [13] Galindo, C., Hernando, F. and Monserrat F.: The log-canonical threshold of a plane curve, *Mathematical Proceedings of the Cambridge Philosophical Society*. Vol. 160. No. 3. Cambridge University Press, 2016.
- [14] Galindo, C. and Monserrat, F.: The Poincaré series of multiplier ideals of a simple complete ideal in a local ring of a smooth surface, *Advances in Mathematics* 225.2 (2010): 1046-1068.
- [15] Howald, J.: Multiplier ideals of monomial ideals, *Transactions of the American Mathematical Society* 353.7 (2001): 2665-2671.
- [16] Hyry, E. and Järvilehto T.: A formula for jumping numbers in a two-dimensional regular local ring, *Journal of Algebra* 516 (2018): 437-470.
- [17] Järvilehto, T.: Jumping numbers of a simple complete ideal in a two-dimensional regular local ring, Vol. 214. No. 1009. American Mathematical Society, 2011.
- [18] Eisenbud, D. and Neumann, W. D.: Three-Dimensional Link Theory and Invariants of Plane Curve Singularities, *Annals of Math. Studies* 110, Princeton University Press 1985.
- [19] Kollár, J.: Singularities of the Minimal Model Program, *Cambridge Tracts in Mathematics*, vol. 200, Cambridge University Press, Cambridge 2013.
- [20] Kollár, J. and Mori, S.: *Birational Geometry of Algebraic Varieties*, Cambridge Tracts in Mathematics 134, Cambridge University Press, Cambridge 1998.
- [21] Lazarsfeld, R.: Positivity in algebraic geometry. II. Positivity for vector bundles, and multiplier ideals, *Ergebnisse der Mathematik und ihrer Grenzgebiete* 3 (2003): 49.
- [22] Montaner, J.À. and Núñez-Betancourt, L.: Poincaré series of multiplier and test ideals, *Revista matemática iberoamericana* 38.6 (2022): 1993-2009.
- [23] Naie, D.: Jumping numbers of a unibranch curve on a smooth surface. *Manuscripta mathematica* 128 (2009): 33-49.
- [24] Némethi, A. and Steenbrink, J.H.M.: Spectral pairs, mixed Hodge modules and series of plane curve singularities, *New York Journal of Mathematics*, August 16, 1995.
- [25] Némethi, A.: On the Ozsváth–Szabó invariant of negative definite plumbed 3-manifolds, *Geometry & Topology* 9(2), 991–1042 (2005).
- [26] Némethi, A.: Poincaré series associated with surface singularities, in *Singularities I: Algebraic and Analytic Aspects*, Contemp. Math. 474, AMS 271–297 (2008).
- [27] Némethi, A.: *Normal surface singularities*, *Ergebnisse der Math. und ihrer Grenzgebiete*, Springer 2022.
- [28] Neumann, W. D.: Abelian covers of quasihomogeneous surface singularities, In: Orlik, P. (ed.) *Singularities*, Part 2, Proc. of Symposia in Pure Math. 40/2 (1983), 233–243.
- [29] Neumann, W.D. and Wahl, J.: Complete intersection singularities of splice type as universal abelian covers. *Geom. Topol.* 9 (2005), 699-755.

- [30] Okuma, T.: Universal abelian covers of rational surface singularities, *J. London Math. Soc.*, 70(2), 307–324 (2004).
- [31] Pande, S.: Multiplicities of jumping numbers, *Algebra & Number Theory* 17.1 (2023): 83–110.
- [32] Pinkham, H.: Normal surface singularities with \mathbb{C}^* -action, *Math. Ann.* 227(2), 183–193 (1977).
- [33] Saito, M.: Mixed Hodge modules, *Publ. RIMS Kyoto University* 26 (1990), 221–333.
- [34] Saito, M.: On Steenbrink’s Conjecture, *math. Annalen* 289 (1991), 703–716.
- [35] Schrauwen, R., Steenbrink, J. and Stevens, J.: Spectral Pairs and the Topology of Curve Singularities, *Proceedings of Symposia in Pure Mathematics* 53, (1991), 305–328.
- [36] Steenbrink, J.H.M.: Mixed Hodge structures on the vanishing cohomology, In P.Holm ed.: *Real and Complex singularities, Oslo 1976*, Sijthoff-Noordhoff, Alphen a/d Rijn 1977, 525–563.
- [37] Steenbrink, J.H.M.: The spectrum of hypersurface singularities, *Astérisque* 179–180 (1989), 163–184 (Actes du Colloque de Théorie de Hodge, Luminy, 1987).
- [38] Stein, K.: Analytische Zerlegungen komplexer Räume, *Math. Ann.* 132 (1), (1956), 63–93.
- [39] Tomari, M. and Watanabe, K.: Normal \mathbb{Z}_r -graded rings and normal cyclic covers, *Manuscripta Math.* 76, 325–340 (1992).
- [40] Tucker, K.F.: Jumping numbers on algebraic surfaces with rational singularities, *Transactions of the American Mathematical Society* 362.6 (2010): 3223–3241.

HUN-REN ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, REÁLTANODA UTCA 13-15, H-1053, BUDAPEST, HUNGARY

ELTE - FACULTY OF SCIENCE, DEPT. OF GEOMETRY, PÁZMÁNY PÉTER SÉTÁNY 1/A, 1117 BUDAPEST, HUNGARY

Email address: koltai.laszlo@renyi.hu

BABEŞ-BOLYAI UNIVERSITY, STR. MIHAIL KOGĂLNICEANU NR. 1, 400084 CLUJ-NAPOCA, ROMANIA
SIMION STOILOW INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, BUCHAREST, ROMANIA

Email address: tamas.laszlo@ubbcluj.ro

HUN-REN ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, REÁLTANODA UTCA 13-15, H-1053, BUDAPEST, HUNGARY

ELTE - FACULTY OF SCIENCE, DEPT. OF GEOMETRY, PÁZMÁNY PÉTER SÉTÁNY 1/A, 1117 BUDAPEST, HUNGARY

BBU - BABEŞ-BOLYAI UNIV., STR. M. KOGĂLNICEANU 1, 400084 CLUJ-NAPOCA, ROMANIA

BCAM - BASQUE CENTER FOR APPLIED MATH., MAZARREDO, 14 E48009 BILBAO, BASQUE COUNTRY – SPAIN

Email address: nemethi.andras@renyi.hu