THE MULTIPLICATIVE STRUCTURE OF THE K-THEORETICAL MCKAY CORRESPONDENCE FOR THE HILBERT SCHEME OF POINTS IN THE COMPLEX PLANE

JAKUB KONCKI AND MAGDALENA ZIELENKIEWICZ

ABSTRACT. We consider the K-theory of the Hilbert scheme of points in the complex plane, which under McKay correspondence is isomorphic to the space of symmetric functions Λ^n . We prove a formula conjectured by Boissière for the endomorphism of Λ^n induced by multiplication by the classes of the Adams powers of the tautological bundle. We describe the structure constants for the multiplication on Λ^n induced by the tensor product in K-theory.

1. INTRODUCTION

Hilbert schemes, introduced in the 60's, quickly became standard objects in algebraic geometry. In this paper, we focus on the Hilbert scheme of n points in the complex plane $\operatorname{Hilb}_n := \operatorname{Hilb}_n(\mathbb{A}^2_{\mathbb{C}})$. It is a smooth quasiprojective variety parametrizing zero-dimensional subschemes of $\mathbb{A}^2_{\mathbb{C}}$ of length n. The McKay correspondence forms a bridge between the geometry of Hilb_n and the combinatorics of symmetric functions of degree n.

The permutation group S_n of n elements acts on the vector space $(\mathbb{C}^2)^n$ by permuting the \mathbb{C}^2 factors. This action preserves the standard symplectic form on \mathbb{C}^{2n} . The quotient space is a symplectic singularity in the sense of [Kal09, Bea00] and the Hilbert scheme Hilb_n is its symplectic resolution. The McKay correspondence predicts a relation between the geometry of Hilb_n and the S_n -equivariant geometry of \mathbb{C}^{2n} . Various instances of the McKay correspondence are known. It was studied as an isomorphism of cohomology or K-theory groups e.g. [Kal02, IN00], as an equality of elliptic classes [BL05, Wae08] or as an equivalence of derived categories [BKR01]. The K-theoretical correspondence was used by Haiman to prove several combinatorial conjectures [Hai01, Hai02]. The study of the McKay correspondence (often for different groups and resolutions of singularities) is an active area of research, e.g. [Ohm08, Kru18, MW21, DBW18].

We focus on the K-theoretical correspondence for the Hilbert scheme $\operatorname{Hilb}_n(\mathbb{A}^2_{\mathbb{C}})$. It provides us with an isomorphism

$$\Theta: \Lambda^n \simeq \mathrm{K}_{\mathrm{S}_n}(\mathbb{C}^{2n}) \to \mathrm{K}(\mathrm{Hilb}_n(\mathbb{A}^2_{\mathbb{C}}))\,,$$

where Λ^n is the space of symmetric functions of degree n. The codomain of this isomorphism has a product structure given by the tensor product. It induces interesting operations on the space Λ^n . We are interested in two natural questions:

A) For a given element $F \in K(Hilb_n)$, find a compact formula for the map $\mathcal{E}_F : \Lambda^n \to \Lambda^n$ such that $\Theta(\mathcal{E}_F(x)) = \Theta(x) \otimes F$, i.e. the diagram below commutes

ż

B) Describe the product structure on Λ^n induced from K(Hilb_n).

The Hilbert scheme Hilb_n comes with a distinguished vector bundle B_n called the tautological vector bundle. Kirwan surjectivity type results imply that the cohomology $\operatorname{H}^*(\operatorname{Hilb}_n)$ is generated as a ring by the graded parts of the Chern character $\operatorname{ch}^k(B_n)$ [ES93] and the K-theory is generated by the Adams powers $\psi^k B_n$ [MN18]. Therefore, it is enough to solve problem A) for $F = \psi^k B_n$. Characteristic classes of the tautological bundles and their relation with the McKay correspondence can also be considered for varieties other than the affine plane, e.g. [BNW07, Sca09, Kru14].

A cohomological version of problem A) was solved in [Leh99]. It was used in [LS01] to describe the multiplicative structure on Λ^n induced from $\mathrm{H}^*(\mathrm{Hilb}_n)$. In K-theory, the formula for the map \mathcal{E}_{B_n} was given in [Boi06b]

$$\mathcal{E}_{B_n} = \operatorname{Coeff}\left(t^0; \sum_{r\geq 1} rp_r t^r \cdot \exp\left(\sum_{r\geq 1} \frac{\partial}{\partial p_r} t^{-r}\right)\right).$$

Moreover, the formula for $F = \psi^k B_n$ was conjectured.

Theorem ([Boi06b, Conjecture 8.1]).

$$\mathcal{E}_{\psi^k B_n} = k^{-1} \cdot \operatorname{Coeff}\left(t^0; \sum_{r \ge 1} r p_r t^r \cdot \exp\left(k \cdot \sum_{r \ge 1} \frac{\partial}{\partial p_r} t^{-r}\right)\right) \,.$$

Our main result is Theorem 5.14, where we prove an equivalent version of the above formula. Both the results of [Boi06b] and our proof rely on the study of the torus action.

The standard action of the two dimensional complex torus $\mathbb{T} = (\mathbb{C}^*)^2$ on the affine plane induces an action on the Hilbert scheme Hilb_n. This action was first used to compute the Betti numbers of the Hilbert scheme in [ES87] and its use is a standard method since then, e.g. [CE15, Smi20, KZ23, SV13]. The fixed point set Hilb_n^T is finite, which allows for a combinatorial approach to the equivariant cohomology and K-theory using localization theorems [Tho92, AB84]. Roughly speaking, any class in K-theory is determined by local contributions coming from the fixed point set. These contributions are governed by the combinatorics of the tangent weights. Therefore, many cohomological statements can be reduced to combinatorial problems. Unfortunately, they are usually complicated. This is the case for the computation of the map \mathcal{E}_F .

Our idea is to simplify the problem using a reduction to a well-chosen subtorus $\mathbb{T}_1 \subset \mathbb{T}$. For a general \mathbb{T}_1 , the fixed point $\operatorname{Hilb}_n^{\mathbb{T}_1}$ is finite, yet the description of the tangent weights for \mathbb{T}_1 is no easier than for \mathbb{T} . On the other hand, the combinatorics of the tangent weights visibly simplifies when the fixed point set $\operatorname{Hilb}_n^{\mathbb{T}_1}$ is bigger than $\operatorname{Hilb}_n^{\mathbb{T}}$. Yet, in such case one needs to consider the geometry of the fixed point set which may be complicated, see [Eva04]. We choose $\mathbb{T}_q \subset \mathbb{T}$ to be a coordinate subtorus. The fixed point set $\operatorname{Hilb}_n^{\mathbb{T}_q}$ is much bigger than $\operatorname{Hilb}_n^{\mathbb{T}}$, therefore the tangent weights become much easier. Moreover, $\operatorname{Hilb}_n^{\mathbb{T}_q}$ is a disjoint union of affine spaces, thus it has trivial geometry (from the cohomological point of view). As expected, after restriction to the subtorus \mathbb{T}_q , problem A) simplifies.

Our proof is purely combinatorial but it is based on the geometric intuition presented above. Let q and t be the coordinate characters of \mathbb{T} . The restriction to \mathbb{T}_q corresponds to the substitution t = 1 in the combinatorial formulas. After this restriction the adjoint operator $\mathcal{E}_{\psi_k B}^*$ satisfies the Leibniz rule, which allows for an effective calculation.

In the last section, we study problem B). We consider the multiplication $\odot : \Lambda^n \times \Lambda^n \to \Lambda^n$ induced by the tensor product in the K-theory ring K(Hilb_n). This product is determined by its structure constants $c^{\mu}_{\lambda_1,\lambda_2} \in \mathbb{Q}$ in the power sum basis, i.e.

$$p_{\lambda_1} \odot p_{\lambda_2} = \sum_{\mu \vdash n} c^{\mu}_{\lambda_1, \lambda_2} p_{\mu}.$$

In Theorem 7.3, we obtain a formula for the constants $c^{\mu}_{\lambda_1,\lambda_2}$ as the coefficients of a certain power series. Our method of proof is again the study of the action of the one-dimensional torus \mathbb{T}_q .

The structure of this paper is as follows. Sections 2–5 provide the necessary background. In Section 5 we state equivalent formulations of [Boi06b, Conjecture 8.1] in Theorem 5.14 and Proposition 5.15. Section 6 contains the proof of the main theorem, Theorem 5.14. In Section 7 we study the product \odot on Λ^n . In Appendix A we include a proof of the equivariant K-theoretic McKay correspondence. It is a well-known result, but we were unable to find a reference.

Acknowledgements: JK is supported by NCN grant SONATINA 2023/48/C/ST1/00002. Both authors are grateful to Joachim Jelisiejew, Bruno Stonek and Andrzej Weber for their helpful comments. JK wants to thank Jörg Schürmann for introducing him to the papers of Boissière.

2. K-THEORY

The standard reference for equivariant K-theory is [CG10]. Let \mathbb{T} be an algebraic torus and X a \mathbb{T} -variety. The equivariant K-theory $K_{\mathbb{T}}(X)$ is the K-group of the additive category of \mathbb{T} -equivariant vector bundles on X. If X is smooth, it is isomorphic to the K-group of the category of equivariant coherent sheaves on X. All our varieties are quasiprojective. Moreover, we consider only K-theory with rational coefficients, i.e.

$$\mathrm{K}(X) := \mathrm{K}(X, \mathbb{Q}) = \mathrm{K}(X, \mathbb{Z}) \otimes \mathbb{Q}, \qquad \mathrm{K}_{\mathbb{T}}(X) := \mathrm{K}_{\mathbb{T}}(X, \mathbb{Q}) = \mathrm{K}_{\mathbb{T}}(X, \mathbb{Z}) \otimes \mathbb{Q}.$$

For an algebraic torus \mathbb{T} , let $S_{\mathbb{T}} \subset K_{\mathbb{T}}(\mathrm{pt})$ be the multiplicative system of all nonzero elements. The localised K-theory of a point is $S_{\mathbb{T}}^{-1} K_{\mathbb{T}}(\mathrm{pt})$. For any \mathbb{T} -variety X, the ring $K_{\mathbb{T}}(X)$ is a $K_{\mathbb{T}}(\mathrm{pt})$ -algebra. We denote the localised K-theory of X by

$$S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(X) := \operatorname{K}_{\mathbb{T}}(X) \otimes_{\operatorname{K}_{\mathbb{T}}(\operatorname{pt})} S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(\operatorname{pt})$$

Analogously, for a finite group G and a $\mathbb{T} \times G$ -variety X we have

$$S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T} \times G}(X) := \operatorname{K}_{\mathbb{T} \times G}(X) \otimes_{\operatorname{K}_{\mathbb{T}}(\operatorname{pt})} S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(\operatorname{pt}).$$

Theorem 2.1 (Localisation Theorem [Tho92, Theorem 2.1]). Let \mathbb{T} be an algebraic torus and X a \mathbb{T} -variety. The inclusion $i: X^{\mathbb{T}} \hookrightarrow X$ induces an isomorphism of rings

$$i^*: S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(X) \simeq S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(X^{\mathbb{T}}).$$

Corollary 2.2. Let \mathbb{T} be an algebraic torus and X be a \mathbb{T} -variety.

- (1) Suppose that the fixed point set $X^{\mathbb{T}}$ is finite. Then the fundamental classes of the fixed points form a basis of the localised K-theory $S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(X)$ over $S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(\operatorname{pt})$.
- (2) More generally, suppose that the fixed point set is a disjoint union of affine spaces. Then the fundamental classes of the components of the fixed point set form a basis of the localised K-theory $S_{\mathbb{T}}^{-1} K_{\mathbb{T}}(X)$ over $S_{\mathbb{T}}^{-1} K_{\mathbb{T}}(\operatorname{pt})$.

For a T-variety X, the K-theory $K_{\mathbb{T}}(X)$ is a λ -ring with λ -operations induced by taking exterior powers, see [FL85] for λ -rings. The theory of λ -rings provides us with Adams operations

$$\psi^m: \mathrm{K}_{\mathbb{T}}(X) \to \mathrm{K}_{\mathbb{T}}(X) \,,$$

for every $m \in \mathbb{N}$. They were first introduced in the non-equivariant setting in [Ada62]. Let us recall some basic properties of the Adams operations.

Proposition 2.3. Let \mathbb{T} be an algebraic torus and X be a \mathbb{T} -variety.

- (1) The Adams operation $\psi^m : K_{\mathbb{T}}(X) \to K_{\mathbb{T}}(X)$ is a ring homomorphism.
- (2) For an equivariant line bundle $\mathcal{L} \in \operatorname{Pic}^{\mathbb{T}}(X)$ we have

$$\psi^m(\mathcal{L}) = \mathcal{L}^{\otimes m}$$

(3) The Adams operations are natural transformations with respect to pullbacks, i.e. for any \mathbb{T} -equivariant map $f: Y \to X$ we have

$$f^* \circ \psi^m = \psi^m \circ f^*$$

3. Combinatorics

3.1. **Partitions.** A partition is a non-increasing sequence of positive integers, i.e.

$$\lambda = (\lambda_1, \dots, \lambda_k), \qquad \lambda_1 \ge \lambda_2 \ge \dots \ge \lambda_k > 0.$$

We say that $s(\lambda) := \sum_{i=1}^{k} \lambda_i$ is the sum of the partition λ or that λ is a partition of $s(\lambda)$, and write $\lambda \vdash s(\lambda)$. We call $l(\lambda) := k$ the length of the partition λ . A partition can be presented in an alternative way $\lambda = (1^{a_1(\lambda)}, 2^{a_2(\lambda)}, \dots)$ where

$$a_j(\lambda) := \left| \{ i \in \mathbb{N} | \lambda_i = j \} \right|.$$

In further considerations we use the numbers

(1)
$$z_{\lambda} = \prod_{i=1}^{\infty} \left(a_i(\lambda)! \cdot i^{a_i(\lambda)} \right) = \prod_{i=1}^{\infty} a_i(\lambda)! \cdot \prod_{i=1}^{l(\lambda)} \lambda_i \,.$$

Definition 3.1. Let λ be a partition of n and μ a partition of m. The partition $\lambda \cdot \mu$ of m + n is defined by

$$a_i(\lambda \cdot \mu) = a_i(\lambda) + a_i(\mu)$$
.

Example 3.2. For $\lambda = (5, 2, 1)$ and $\mu = (3, 2, 1)$ we obtain $\lambda \cdot \mu = (5, 3, 2, 2, 1, 1)$.

3.2. Symmetric functions. We recall some properties of symmetric functions. For a full exposition and proofs see [Mac15]. Let S_n denote the permutation group of nelements. Let Λ^n be the Q-vector space of symmetric functions of degree n and Λ be the ring of symmetric functions, i.e.

$$\Lambda^{n} := \lim_{\leftarrow} \{ W \in \mathbb{Q}[x_{1}, \dots, x_{m}]^{\mathbb{S}_{m}} | W \text{ is homogeneous, } \deg W = n \}, \qquad \Lambda = \bigoplus_{n \in \mathbb{N}} \Lambda^{n}.$$

Denote by $\Lambda_a^n, \Lambda_{a,t}^n, \Lambda_q, \Lambda_{q,t}$ the spaces of symmetric functions with additional parameters

$$\Lambda_q^n = \Lambda^n \otimes_{\mathbb{Q}} \mathbb{Q}(q) \,, \qquad \Lambda_{q,t}^n = \Lambda^n \otimes_{\mathbb{Q}} \mathbb{Q}(q,t) \,, \qquad \Lambda_q = \Lambda \otimes_{\mathbb{Q}} \mathbb{Q}(q) \,, \qquad \Lambda_{q,t} = \Lambda \otimes_{\mathbb{Q}} \mathbb{Q}(q,t) \,.$$

The space Λ^n has several standard bases indexed by partitions of n, such as: the power sum functions p_{λ} , the full symmetric functions h_{λ} and the Schur functions s_{λ} .

Definition 3.3. • The involution $\omega : \Lambda^n \to \Lambda^n$ is defined by $\omega(p_\lambda) = (-1)^{l(\lambda)} p_\lambda$. We extend it $\mathbb{Q}(q,t)$ -linearly to an involution $\omega : \Lambda^n_{q,t} \to \Lambda^n_{q,t}$. • There is an inner product $\langle -, - \rangle$ on the space Λ^n defined by one of the equivalent conditions:

 $\langle p_{\lambda}, p_{\mu} \rangle = \delta_{\lambda,\mu} z_{\lambda} , \qquad \langle s_{\lambda}, s_{\mu} \rangle = \delta_{\lambda,\mu} .$ We extend it to a $\mathbb{Q}(q, t)$ -bilinear form on $\Lambda_{q,t}^n$.

Remark 3.4. In the literature there are two conventions concerning the involution ω . One either defines $\omega(p_{\lambda}) = (-1)^{l(\lambda)} p_{\lambda}$ (e.g. [Boi06b]), or $\omega(p_{\lambda}) = (-1)^{n+l(\lambda)} p_{\lambda}$ (e.g. [Mac15, Hai03]). As we prove a conjecture originating from [Boi06b], we use the first convention to be consistent with the notation therein. This convention has a nice interpretation in terms of plethystic substitutions, see Example 3.9 (1).

The ring of symmetric functions is a polynomial ring generated by the power sums $\{p_n\}_{n\in\mathbb{N}}$, i.e.

(2)
$$\Lambda = \mathbb{Q}[p_1, p_2, \dots].$$

This allows us to define derivations $\frac{\partial}{\partial p_n} : \Lambda \to \Lambda$ in the standard way.

Definition 3.5. For a partition $\mu = (\mu_1, \ldots, \mu_l)$, the map $\partial_\mu : \Lambda \to \Lambda$ is defined by

$$\partial_{\mu} = \frac{\partial}{\partial p_{\mu_1}} \circ \frac{\partial}{\partial p_{\mu_2}} \circ \cdots \circ \frac{\partial}{\partial p_{\mu_l}}.$$

This map may be extended $\mathbb{Q}(q,t)$ -linearly to a map $\partial_{\mu} : \Lambda_{q,t} \to \Lambda_{q,t}$.

The following widely known fact can be easily checked using the power sum basis.

Proposition 3.6. Let f and g be symmetric functions.

(1) Let m be a natural number. Multiplication by $\frac{p_m}{m}$ and differentiation $\frac{\partial}{\partial p_m}$ are adjoint, i.e.

$$\left\langle \frac{f \cdot p_m}{m}, g \right\rangle = \left\langle f, \frac{\partial}{\partial p_m} g \right\rangle.$$

(2) Let λ be a partition. Multiplication by $\frac{p_{\lambda}}{\prod_{i=1}^{l(\lambda)} \lambda_i}$ and differentiation ∂_{λ} are adjoint, *i.e.*

$$\left\langle \frac{f \cdot p_{\lambda}}{\prod_{i=1}^{l(\lambda)} \lambda_i}, g \right\rangle = \left\langle f, \partial_{\lambda} g \right\rangle \,.$$

3.3. Plethystic substitution. The plethystic substitution describes the λ -ring structure on the ring of symmetric functions, see e.g. [GH96] for a detailed description. Thanks to the presentation (2) of Λ , in order to define a Q-algebra map $\varphi : \Lambda^n \to \Lambda^n$ it is enough to specify the values $\varphi(p_n)$.

Definition 3.7. For an element $A \in \Lambda$, the plethystic substitution $f \to f[A]$ is a Q-algebra morphism $-[A] : \Lambda \to \Lambda$ such that

$$p_k[A] = A(x_1^k, x_2^k, \dots) \,.$$

For $A \in \Lambda_{q,t}$, the plethystic substitution is a $\mathbb{Q}(q,t)$ -algebra morphism $-[A] : \Lambda_{q,t} \to \Lambda_{q,t}$ such that $p_k[A] = A(x_1^k, x_2^k, \dots, q^k, t^k)$.

In plethystic notation we write X for the sum of all variables: $X = x_1 + x_2 + ...$

Definition 3.8. Let n be a natural number and λ a partition. We write

$$[n]_q := \frac{q^n - 1}{q - 1} \in \mathbb{Q}(q) , \qquad [\lambda]_q := \prod_{i=1}^{l(\lambda)} [\lambda_i]_q \in \mathbb{Q}(q) .$$

Example 3.9. Let $f, g \in \Lambda_q$ be symmetric functions.

- (1) We have f[X] = f and $f[-X] = \omega(f)$.
- (2) Let $A(q) \in \mathbb{Q}(q)$ be a non-zero rational function. The substitution $f \to f[A(q) \cdot X]$ is a $\mathbb{Q}(q)$ -linear isomorphism of Λ_q^n with inverse $f \to f[\frac{X}{A(q)}]$.
- (3) Let $A(q) \in \mathbb{Q}(q)$ be a rational function. The substitution $f \to f[A(q) \cdot X]$ is self-adjoint, i.e.

$$\langle f, g[A(q) \cdot X] \rangle = \langle f[A(q) \cdot X], g \rangle$$

(4) For any partition λ , we have

$$\langle p_{\lambda}, f[X(q-1)] \rangle = \langle p_{\lambda}[X(q-1)], f \rangle = (q-1)^{l(\lambda)}[\lambda]_q \cdot \langle p_{\lambda}, f \rangle.$$

3.4. Macdonald Polynomials. Macdonald polynomials form an important basis of the space $\Lambda_{q,t}$, see [Mac15, Chapter 6]. To a partition λ of n we associate a symmetric function $\tilde{H}_{\lambda}(q,t) \in \Lambda_{q,t}^{n}$ called the modified Macdonald polynomial, see [Hai01, Proposition 2.1.1] for a definition. The functions \tilde{H}_{λ} are in fact polynomials in q, t. It was proved in [Hai01, Theorem 2] that their coefficients in the Schur basis are polynomials with non-negative integer coefficients, i.e.

$$\tilde{H}_{\mu}(q,t) = \sum_{\lambda \vdash n} \tilde{K}_{\lambda\mu}(q,t) s_{\lambda} \in \Lambda_{q,t}^{n}, \qquad \tilde{K}_{\lambda\mu}(q,t) \in \mathbb{N}[q,t].$$

Several restrictions of these polynomials are known, see e.g. [Hai03, Section 3.5.3]. For us the restriction to t = 1 will be important.

Proposition 3.10 ([Mac15, p. 364, Example 7], see also [Hai03, Proposition 3.5.8] for this formulation). We have

$$\tilde{H}_{\lambda}(q,1) = c_{\lambda}(q) \cdot \prod_{i=1}^{l(\lambda)} h_{\lambda_i}\left[\frac{X}{1-q}\right]$$

where $c_{\lambda}(q) = \prod_{i=1}^{l(\lambda)} \prod_{j=1}^{\lambda_i} (1-q^j) \in \mathbb{Q}(q).$

Definition 3.11. Let λ be a partition of n. We use the notation

$$v_{\lambda} := h_{\lambda} \left[\frac{X}{1-q} \right] \in \Lambda_q^n.$$

Proposition 3.12. (1) Let λ and μ be partitions. We have

$$v_{\lambda \cdot \mu} = v_{\lambda} \cdot v_{\mu}$$
.

(2) The set $\{v_{\lambda}\}_{\lambda \vdash n}$ is a $\mathbb{Q}(q)$ -basis of Λ_{q}^{n} .

Proof. The proposition follows from analogous properties of the polynomials h_{λ} . We have

$$v_{\lambda \cdot \mu} = h_{\lambda \cdot \mu} \left[\frac{X}{1-q} \right] = (h_{\lambda} \cdot h_{\mu}) \left[\frac{X}{1-q} \right] = h_{\lambda} \left[\frac{X}{1-q} \right] \cdot h_{\mu} \left[\frac{X}{1-q} \right] = v_{\lambda} \cdot v_{\mu}$$

For the second point, note that the set $\{h_{\lambda}\}_{\lambda \vdash n}$ is a basis of Λ_q^n . The plethystic substitution $f \to f[\frac{X}{1-q}]$ is a $\mathbb{Q}(q)$ -linear automorphism of Λ_q^n , cf. Example 3.9 (2). Therefore the set $\{v_{\lambda}\}_{\lambda \vdash n}$ is also a basis.

3.5. Frobenius characteristic map. The conjugacy classes of the elements of the permutation group S_n are indexed by the partitions of n. Therefore, the representation ring $K_{S_n}(pt)$ is a Q-vector space with a basis corresponding to partitions. The space Λ^n is a Q-vector space of the same dimension. There is a standard choice of isomorphism between these spaces called the Frobenius characteristic map Fr: $K_{S_n}(pt) \to \Lambda^n$, see e.g. [Sta99, Section 7.18]. It is defined by

$$\operatorname{Fr}([V]) = \frac{1}{n!} \sum_{\sigma \in S_n} \chi_V(\sigma) p_{\lambda(\sigma)},$$

where χ_V is the character of the representation V and $\lambda(\sigma)$ is the partition induced by the permutation σ . Let \mathbb{T}_q be a one-dimensional torus and \mathbb{T} be a two-dimensional torus. We have isomorphisms

$$S_{\mathbb{T}_q}^{-1} \operatorname{K}_{\mathbb{T}_q \times \operatorname{S}_n}(\operatorname{pt}) \xrightarrow{\simeq} \operatorname{K}_{\operatorname{S}_n}(\operatorname{pt}) \otimes_{\mathbb{Q}} S_{\mathbb{T}_q}^{-1} \operatorname{K}_{\mathbb{T}_q}(\operatorname{pt}) \xrightarrow{\simeq} \operatorname{K}_{\operatorname{S}_n}(\operatorname{pt}) \otimes_{\mathbb{Q}} \mathbb{Q}(q) \xrightarrow{\operatorname{Fr} \otimes \operatorname{id}_{\mathbb{Q}(q)}} \Lambda_q^n ,$$

$$S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T} \times \operatorname{S}_n}(\operatorname{pt}) \xrightarrow{\simeq} \operatorname{K}_{\operatorname{S}_n}(\operatorname{pt}) \xrightarrow{\simeq} \operatorname{K}_{\operatorname{S}_n}(\operatorname{pt}) \xrightarrow{\simeq} \operatorname{K}_{\operatorname{S}_n}(\operatorname{pt}) \otimes_{\mathbb{Q}} \mathbb{Q}(q, t) \xrightarrow{\operatorname{Fr} \otimes \operatorname{id}_{\mathbb{Q}(q, t)}} \Lambda_{q, t}^n .$$

We call both these maps the Frobenius map and denote them by Fr.

4. HILBERT SCHEME

Let $\operatorname{Hilb}_n := \operatorname{Hilb}_n(\mathbb{A}^2_{\mathbb{C}})$ be the Hilbert scheme of n points in the complex plane $\mathbb{A}^2_{\mathbb{C}}$. We recall some standard facts about Hilb_n , for reference see [Nak99]. The Hilbert scheme Hilb_n parametrises zero-dimensional subschemes of $\mathbb{A}^2_{\mathbb{C}}$ of length n. It is a smooth irreducible quasi-projective variety of dimension 2n, [Fog68]. Set-theoretically, Hilb_n can be identified with the set of ideals $I \triangleleft \mathbb{C}[x, y]$ such that the \mathbb{C} -vector space $\mathbb{C}[x, y]/I$ has dimension n, i.e.,

$$\operatorname{Hilb}_{n} := \{ I \lhd \mathbb{C}[x, y] | \dim \mathbb{C}[x, y] / I = n \}.$$

Let $\mathbb{T} = (\mathbb{C}^*)^2$ be the algebraic torus. The standard \mathbb{T} -action on $\mathbb{A}^2_{\mathbb{C}}$ induces an action of \mathbb{T} on the Hilbert scheme Hilb_n. The fixed points of this action are monomial ideals contained in Hilb_n. They correspond bijectively to partitions of n. To a partition $\lambda = (\lambda_1, \ldots, \lambda_l)$ we associate an ideal

$$\langle x^{\lambda_1}, x^{\lambda_2}y, \dots, x^{\lambda_l}y^{l-1}, y^l \rangle \in \operatorname{Hilb}_{s(\lambda)}^{\mathbb{T}}$$

We do not distinguish in the notation between the partition λ and the associated fixed point. Denote by q and t the coordinate characters of the torus \mathbb{T} . We consider the one-dimensional coordinate subtorus $\mathbb{T}_q \subset \mathbb{T}$ acting on the first coordinate. Then

$$\mathbf{K}_{\mathbb{T}}(\mathbf{pt}) = \mathbb{Q}[q^{\pm 1}, t^{\pm 1}], \qquad \mathbf{K}_{\mathbb{T}_q}(\mathbf{pt}) = \mathbb{Q}[q^{\pm 1}].$$

The restriction map $K_{\mathbb{T}}(pt) \to K_{\mathbb{T}_q}(pt)$ is given by substitution t = 1.

The fixed point set $\operatorname{Hilb}_{n}^{\mathbb{T}_{q}}$ is a disjoint union of affine spaces, [Eva04, Section 1.2]. Corollary 2.2 gives the following description of the localised K-theory of Hilb_{n} .

Proposition 4.1. (1) The fundamental classes of the \mathbb{T} -fixed points form a basis of the localised K-theory $S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(\operatorname{Hilb}_n)$.

(2) The fundamental classes of the components of the fixed point set $\operatorname{Hilb}_{n}^{\mathbb{T}_{q}}$ form a basis of the localised K-theory $S_{\mathbb{T}_{q}}^{-1} \operatorname{K}_{\mathbb{T}_{q}}(\operatorname{Hilb}_{n})$.

Definition 4.2. Let F be a T-equivariant vector bundle on the Hilbert scheme Hilb_n and λ a partition of n. We write

$$F_{\lambda} \in \mathbf{K}^{\mathbb{T}}(\mathrm{pt}) \simeq \mathbb{Q}[q^{\pm 1}, t^{\pm 1}],$$

for the class of the restriction of F to the fixed point λ . We write

$$F^q_{\lambda} \in \mathcal{K}^{\mathbb{T}_q}(\mathrm{pt}) \simeq \mathbb{Q}[q^{\pm 1}]$$

for the restriction to \mathbb{T}_q -equivariant K-theory, i.e. $F_{\lambda}^q = (F_{\lambda})_{t=1}$.

The Hilbert scheme $Hilb_n$ comes with a universal flat family

$$\mathcal{F}_n \subset \operatorname{Hilb}_n \times \mathbb{A}^2_{\mathbb{C}},$$

whose fibre over a subscheme $Z \in \text{Hilb}_n$ is Z, treated as a subscheme in $\mathbb{A}^2_{\mathbb{C}}$. The pushforward of the sheaf of regular functions on \mathcal{F}_n along the projection $\pi : \mathcal{F}_n \to \text{Hilb}_n$ defines (a sheaf of sections of) a rank *n* vector bundle on Hilb_n ,

$$B_n = \pi_* \mathcal{O}_{\mathcal{F}_n} \in \operatorname{Vect}(\operatorname{Hilb}_n)$$

It is called the *tautological vector bundle* on $Hilb_n$. We are interested in the Adams powers of the tautological bundle, due to the following instance of Kirwan surjectivity.

Theorem 4.3 ([MN18, Corollary 1.4]). The classes of the bundles $\psi^k B_n$ for $k \in \mathbb{N}$ generate K(Hilb_n) as a Q-algebra.

The tautological bundle B_n admits a natural \mathbb{T} -linearisation. Its fibre over a fixed point $I \in \operatorname{Hilb}_n^{\mathbb{T}}$ is isomorphic to the *n*-dimensional vector space $\mathbb{C}[x, y]/I$ as a \mathbb{T} -representation. This allows for an explicit computation of the restrictions $(\psi^k B_n)_{\lambda}$.

Example 4.4. Let $\lambda = (\lambda_1, \ldots, \lambda_l)$ be a partition of *n*. We have

$$(B_n)_{\lambda} = \sum_{i=1}^{l} \sum_{j=1}^{\lambda_i} t^{i-1} q^{j-1}.$$

More generally, for $k \in \mathbb{N}$,

$$(\psi^k B_n)_{\lambda} = \sum_{i=1}^l \sum_{j=1}^{\lambda_i} (t^{i-1}q^{j-1})^k, \qquad (\psi^k B_n)_{\lambda}^q = \sum_{i=1}^l \sum_{j=1}^{\lambda_i} (q^{j-1})^k.$$

The last formula in the above example implies the following fact.

Corollary 4.5. Let λ be a partition of n, μ be a partition of m and $k \in \mathbb{N}$ be a natural number. We have

$$(\psi^k B_{n+m})^q_{\lambda \cdot \mu} = (\psi^k B_n)^q_{\lambda} + (\psi^k B_m)^q_{\mu}.$$

Remark 4.6. The above fact is false if we restrict to q = 1 instead of t = 1. In this case, we have

$$(\psi^k B_{n+m})^t_{(\lambda^\perp \cdot \mu^\perp)^\perp} = (\psi^k B_n)^t_{\lambda} + (\psi^k B_m)^t_{\mu},$$

where \perp denotes the transpose partition.

5. MCKAY CORRESPONDENCE

5.1. Additive correspondence. The permutation group S_n acts on the vector space $\mathbb{C}^{2n} = (\mathbb{C}^2)^n$ by permutation of the \mathbb{C}^2 factors. The Hilbert-Chow morphism $\operatorname{Hilb}_n \to \mathbb{C}^{2n} / S_n$ is a resolution of the quotient singularity. In [Hai01, Definition 3.2.4], the isospectral Hilbert scheme X_n is defined as the reduced fiber product

(3)
$$\begin{array}{c} X_n \xrightarrow{\rho} \mathbb{C}^{2n} \\ p \downarrow \qquad \qquad \downarrow \\ \text{Hilb}_n \longrightarrow \mathbb{C}^{2n} / S_n \end{array}$$

We use the names of the maps from the above diagram. The McKay correspondence relates the derived category of the Hilbert scheme with the S_n -equivariant derived category of the affine space \mathbb{C}^{2n} .

Proposition 5.1 ([BKR01, Corollary 1.3]). The composition

 $R\rho_* \circ p^* \colon D(\mathrm{Hilb}_n) \to D^{\mathrm{S}_n}(\mathbb{C}^{2n})$

is an equivalence of categories.

As a consequence, we have an isomorphism of K-groups.

Corollary 5.2. The map

$$\rho_* \circ p^* \colon \mathrm{K}(\mathrm{Hilb}_n) \to \mathrm{K}_{\mathrm{S}_n}(\mathbb{C}^{2n})$$

is an isomorphism of \mathbb{Q} -vector spaces.

Corollary 5.3. The maps

$$\rho_* \circ p^* \colon S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(\operatorname{Hilb}_n) \to S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T} \times \operatorname{S}_n}(\mathbb{C}^{2n}), \qquad \rho_* \circ p^* \colon S_{\mathbb{T}_q}^{-1} \operatorname{K}_{\mathbb{T}_q}(\operatorname{Hilb}_n) \to S_{\mathbb{T}_q}^{-1} \operatorname{K}_{\mathbb{T}_q \times \operatorname{S}_n}(\mathbb{C}^{2n})$$

are isomorphisms of vector spaces over $\mathbb{Q}(q,t)$ and $\mathbb{Q}(q)$, respectively.

The torus-equivariant version is an easy consequence of the non-equivariant one. It is widely known, see e.g. [Hai03, Section 5.4.3], yet we were unable to find a proof in the literature. For the sake of completeness we present a simple argument in Appendix A.

Definition 5.4 (cf. [Boi06b, Section 3.2]). Let $i: \{0\} \hookrightarrow \mathbb{C}^{2n}$ be the inclusion of the origin and Fr the Frobenius characteristic map (see Section 3.5). We consider the composition of isomorphisms of $\mathbb{Q}(q, t)$ -vector spaces:

$$\Theta_{q,t} \colon \Lambda_{q,t}^{n} \xrightarrow{\operatorname{Fr}^{-1}} S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T} \times \operatorname{S}_{n}}(\operatorname{pt}) \xrightarrow{(i^{*})^{-1}} S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T} \times \operatorname{S}_{n}}(\mathbb{C}^{2n}) \xrightarrow{(q_{*} \circ p^{*})^{-1}} S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(\operatorname{Hilb}_{n}) \xrightarrow{(i^{*})^{-1}} S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(\operatorname{Hilb}_{n})$$

Analogously, we consider isomorphisms

$$\Theta: \Lambda^n \xrightarrow{\simeq} \mathcal{K}(\mathrm{Hilb}_n), \qquad \Theta_q: \Lambda^n_q \xrightarrow{\simeq} S_{\mathbb{T}_q}^{-1} \mathcal{K}^{\mathbb{T}_q}(\mathrm{Hilb}_n)$$

5.2. Multiplication by the class of a vector bundle. In this section we recall the multiplicative properties of the McKay correspondence obtained in [Boi06b]. We describe the operation on the symmetric functions induced by the multiplication by the class of a vector bundle in the K-theory of Hilb_n.

Definition 5.5. Let F be a T-equivariant vector bundle on the Hilbert scheme Hilb_n. We consider a $\mathbb{Q}(q,t)$ -linear map $\mathcal{E}_{F}^{q,t} : \Lambda_{q,t}^{n} \to \Lambda_{q,t}^{n}$ defined by $\Theta_{q,t}(\mathcal{E}_{F}^{q,t}(x)) = \Theta_{q,t}(x) \otimes F$, i.e. the following diagram commutes:

$$\begin{array}{ccc} \Lambda_{q,t}^{n} & \stackrel{\Theta_{q,t}}{\longrightarrow} & S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(\operatorname{Hilb}_{n}) \\ \varepsilon_{F}^{q,t} & & & \downarrow^{-\otimes F} \\ \Lambda_{q,t}^{n} & \stackrel{\Theta_{q,t}}{\longrightarrow} & S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(\operatorname{Hilb}_{n}) \end{array}$$

Analogous definitions may be given when F is a vector bundle or a \mathbb{T}_q -equivariant vector bundle. We consider the \mathbb{Q} -linear map $\mathcal{E}_F : \Lambda^n \to \Lambda^n$, or the $\mathbb{Q}(q)$ -linear map $\mathcal{E}_F^q : \Lambda_q^n \to \Lambda_q^n$ defined by

$$\Theta_q(\mathcal{E}_F^q(x)) = \Theta_q(x) \otimes F, \qquad \Theta(\mathcal{E}_F(x)) = \Theta(x) \otimes F$$

As a straightforward consequence of the above definition we obtain that the maps \mathcal{E}_F^q and \mathcal{E}_F are restrictions of $\mathcal{E}_F^{q,t}$ to Λ_q^n and Λ^n , respectively.

Proposition 5.6. Let F be a \mathbb{T} -equivariant vector bundle on the Hilbert scheme Hilb_n and $f \in \Lambda_{q,t}^n$ be a symmetric function. Suppose that the restriction $f_{|t=1} \in \Lambda_q^n$ (or $f_{|t=1,q=1} \in \Lambda^n$) is defined. Then $\mathcal{E}_F^{q,t}(f)_{|t=1}$ (or $\mathcal{E}_F^{q,t}(f)_{|t=1,q=1}$) is also defined. Moreover,

$$\mathcal{E}_{F}^{q}(f_{|t=1}) = \mathcal{E}_{F}^{q,t}(f)_{|t=1}, \qquad \mathcal{E}_{F}(f_{|t=1,q=1}) = \mathcal{E}_{F}^{q,t}(y)_{|t=1,q=1}$$

Definition 5.7. Let F be a T-equivariant vector bundle on the Hilbert scheme Hilb_n. Let $\nabla_F^{q,t} : \Lambda_{q,t}^n \to \Lambda_{q,t}^n$ be the $\mathbb{Q}(q,t)$ -linear map defined by

$$\nabla_F^{q,t} \tilde{H}_{\lambda} = F_{\lambda} \tilde{H}_{\lambda} \text{ for all } \lambda \vdash n.$$

Remark 5.8. The operators $\nabla_F^{q,t}$ were studied in a purely combinatorial way in [BGHT99].

Theorem 5.9 ([Boi06b, Theorem 4.1 and its proof]). Let F be a \mathbb{T} -equivariant vector bundle over Hilb_n . We have

$$\mathcal{E}_F^{q,t} = \omega (\nabla_F^{q,t})^* \omega$$

where the dual is taken with respect to the standard scalar product.

Remark 5.10. There is a clear intuition behind the above theorem. The fundamental classes of the fixed points basis $[\lambda]$ form a basis of K-theory, cf. Proposition 4.1 (1). It is an eigenbasis of the map $-\otimes F$. Therefore $\Theta^{-1}([\lambda])$ is an eigenbasis of the operator $\mathcal{E}_F^{q,t}$. The essence of the theorem is a description of the element $\Theta^{-1}([\lambda])$ in terms of Macdonald polynomials.

Theorem 5.9 and Proposition 5.6 imply that the map $\nabla_F^{q,t}$ has well-defined restrictions to Λ_q^n and Λ^n .

Definition 5.11. Let F be a \mathbb{T} -equivariant vector bundle over Hilb_n . We define operators $\nabla_F^q : \Lambda_q^n \to \Lambda_q^n$ and $\nabla_F : \Lambda^n \to \Lambda^n$ such that

$$abla_F^q := (\omega \mathcal{E}_F^q \omega)^*, \qquad \nabla_F := (\omega \mathcal{E}_F \omega)^*.$$

Corollary 5.12. Let F be a \mathbb{T} -equivariant vector bundle over Hilb_n and $x \in \Lambda_{at}^n$.

(1) Suppose that $x_{|t=1} \in \Lambda_q^n$ is defined. Then $\nabla_F^{q,t}(x)_{|t=1}$ is also defined. Moreover,

$$\nabla_F^q(x_{|t=1}) = \nabla_F^{q,t}(x)_{|t=1}$$
.

(2) Suppose that $x_{|t=1,q=1} \in \Lambda^n$ is defined. Then $\nabla_F^{q,t}(x)_{|t=1,q=1}$ is also defined. Moreover,

$$\nabla_F(x_{|t=1,q=1}) = \nabla_F^{q,t}(x)_{|t=1,q=1}.$$

Proof. Theorem 5.9 and Definition 5.11 imply that

$$\nabla_F^{q,t}(x) = (\omega \mathcal{E}_F^{q,t} \omega)^*(x) = \sum_{\lambda \vdash n} \frac{\langle x, \omega \mathcal{E}_F^{q,t} \omega(p_\lambda) \rangle}{z_\lambda} p_\lambda, \qquad \nabla_F^q(x_{|t=1}) = \sum_{\lambda \vdash n} \frac{\langle x_{|t=1}, \omega \mathcal{E}_F^q \omega(p_\lambda) \rangle}{z_\lambda} p_\lambda.$$

The restriction $x_{t=1}$ is well defined, thus for any partition $\lambda \vdash n$ we have

$$\langle x, \omega \mathcal{E}_F^{q,t} \omega(p_\lambda) \rangle_{|t=1} = \langle x_{|t=1}, (\omega \mathcal{E}_F^{q,t} \omega(p_\lambda))_{|t=1} \rangle = \langle x_{|t=1}, \omega \mathcal{E}_F^q \omega(p_\lambda) \rangle.$$

We used Proposition 5.6 for $f = \omega(p_{\lambda})$. This proves the first point. The proof of the second point is analogous.

The K-theory of the Hilbert scheme is generated as a Q-algebra by the Adams powers of the tautological bundle, cf. Theorem 4.3. Therefore in order to describe the map \mathcal{E}_F for every F it is enough to describe it for $F = \psi^k B$, $k \in \mathbb{N}$.

Definition 5.13. We write $\nabla_k^{q,t}$ (or ∇_k^q , ∇_k) for $\nabla_{\psi^k B}^{q,t}$ (or $\nabla_{\psi^k B}^q$, $\nabla_{\psi^k B}$ respectively). In this definition we intentionally omit the number of points in the Hilbert scheme.

The operator $\nabla_1^{q,t}$ can be expressed using plethystic substitution. The Macdonald polynomial $\tilde{H}_{\lambda}(q,t)$ is an eigenvector of the operator $D: \Lambda_{q,t}^n \to \Lambda_{q,t}^n$ defined by

(4)
$$D(f) = \operatorname{Coeff}\left(z^{0}; f\left[X + \frac{z}{(1-q)(1-t)}\right] \cdot \exp\left(\sum_{k \ge 1} \frac{-z^{k}p_{k}}{k}\right)\right)$$

with eigenvalue $1 - (1 - q)(1 - t)B_{\lambda}$. Therefore, $\nabla_1^{q,t} = \frac{1 - D}{(1 - q)(1 - t)}$.

5.3. The main result. We are now ready to state the main theorem, which gives a closed formula for the multiplication induced by the tensoring with the k'th Adams power of the tautological bundle.

Theorem 5.14 (cf. [Boi06b, Conjecture 8.1]). Let $k \in \mathbb{N}$ be a natural number. We have

$$\mathcal{E}_{\psi^k B} = \sum_{\lambda} \frac{k^{l(\lambda)-1} s(\lambda)}{\prod_{i \in \mathbb{N}} a_i(\lambda)!} p_{s(\lambda)} \partial_{\lambda} \,.$$

For k = 1 the above theorem is [Boi06b, Theorem 5.1]. Using $\omega \mathcal{E}_F \omega = \nabla_F^*$ (Definition 5.11) and $\omega(p_{\lambda}) = (-1)^{l(\lambda)} p_{\lambda}$ (Definition 3.3), the above theorem can be reformulated in terms of the operator ∇_k^* , as follows.

Proposition 5.15. Let $k \in \mathbb{N}$ be a natural number. We have

(5)
$$\nabla_k^* = \sum_{\lambda} \frac{(-k)^{l(\lambda)-1} s(\lambda)}{\prod_{i \in \mathbb{N}} a_i(\lambda)!} p_{s(\lambda)} \partial_{\lambda} \, .$$

The next section is devoted to the proof of the above result.

Remark 5.16. Our proof is purely combinatorial, yet it has a geometric motivation. It is based on the substitution t = 1, which corresponds to the restriction to \mathbb{T}_q -equivariant K-theory. The fixed point set $\operatorname{Hilb}_n^{\mathbb{T}_q}$ is still relatively simple. Proposition 4.1 (2) provides an eigenbasis of the operator $-\otimes F$ acting on $S_{\mathbb{T}_q}^{-1} \operatorname{K}_{\mathbb{T}_q}(\operatorname{Hilb}_n)$. On the other hand, the fixed point set $\operatorname{Hilb}_n^{\mathbb{T}_q}$ is much bigger than $\operatorname{Hilb}_n^{\mathbb{T}}$. This implies that the combinatorics of the tangent weights simplifies after the restriction to \mathbb{T}_q . Therefore, the formulas for \mathcal{E}_F^q and ∇_F^q should also simplify.

Remark 5.17. The main step of our proof is showing that the maps ∇_k and ∇_k^q satisfy the Leibniz rule in Proposition 6.2. For the non-equivariant map ∇_k this can be predicted from Formula (5). Multiplication by $\frac{p_m}{m}$ and differentiation $\frac{\partial}{\partial p_m}$ are adjoint. Therefore, we expect that every summand of ∇_k contains a single differentiation.

Remark 5.18. The \mathbb{T} -equivariant map $\nabla_k^{q,t}$ does not satisfies the Leibniz rule even for p_1^2 . This confirms that restriction to the subtorus \mathbb{T}_q simplifies the problem, not only the proofs.

Remark 5.19. In [Boi06b], Theorem 5.14 for k = 1 is stated in the equivalent version

$$\mathcal{E}_B = \operatorname{Coeff}\left(t^0; \sum_{r \ge 1} r p_r t^r \cdot \exp\left(\sum_{r \ge 1} \frac{\partial}{\partial p_r} t^{-r}\right)\right) \,.$$

It is possible to write the result for an arbitrary k in this form

$$\mathcal{E}_{\psi^k B} = k^{-1} \cdot \operatorname{Coeff}\left(t^0; \sum_{r \ge 1} r p_r t^r \cdot \exp\left(k \cdot \sum_{r \ge 1} \frac{\partial}{\partial p_r} t^{-r}\right)\right) \,.$$

Remark 5.20. Contrary to the cohomological case, the formula for $\mathcal{E}_{\psi^k B}$ does not follow directly from the formula for \mathcal{E}_B , as the K-theory is not graded. It is tempting to use the Chern character to transfer the results from cohomology to K-theory, yet the relation between the Chern character and the McKay correspondence is surprisingly complicated, [Boi06a, Theorem 1.1].

6. Restriction to the one-dimensional torus

In this section we study the restriction of the operators $\nabla_F^{q,t}$ to t = 1. It turns out that the restricted operator is much simpler than the original one. For $F = \psi^k B$ it satisfies the Leibniz rule (Proposition 6.2), which allows us to obtain a closed formula for the operators ∇_k (Proposition 6.8). We use Proposition 3.6 to compute the dual ∇_k^* . Recall that we defined $v_{\lambda} = h_{\lambda} \left[\frac{X}{1-q} \right]$ (Definition 3.11).

Proposition 6.1. Let λ be a partition. The element v_{λ} is an eigenvector of ∇_F^q with eigenvalue F_{λ}^q .

Proof. Let $c_{\lambda}(q) \in \mathbb{Q}(q)$ be the constant from Proposition 3.10. Then

$$\nabla_F^q(c_\lambda(q)v_\lambda) = \nabla_F^q(\tilde{H}_\lambda(q,1)) = \nabla_F^{q,t}(\tilde{H}_\lambda(q,t))|_{t=1} = (F_\lambda \cdot \tilde{H}_\lambda(q,t))|_{t=1} = F_\lambda^q \cdot c_\lambda(q)v_\lambda.$$

Thus, $c_{\lambda}(q)v_{\lambda}$ is an eigenvector of ∇_F^q with eigenvalue F_{λ}^q . It follows that v_{λ} is also an eigenvector with the same eigenvalue.

The multiplicative behaviour of the basis v_{λ} implies that the operators ∇_k^q satisfy the Leibniz rule.

Proposition 6.2. Fix a positive integer k. The operator ∇_k^q satisfies the Leibniz rule with regard to the multiplication in Λ_q , i.e. for $f, g \in \Lambda_q$, we have

$$\nabla_k^q(fg) = \nabla_k^q(f) \cdot g + f \cdot \nabla_k^q(g) \,.$$

Proof. Let $F = \psi^k B$. It is enough to check the equation on the basis elements v_{λ} . We have

$$\begin{aligned} \nabla_k^q (v_{\lambda} \cdot v_{\mu}) &= \nabla_k^q (v_{\lambda \cdot \mu}) = F_{\lambda \cdot \mu}^q v_{\lambda \cdot \mu} = (F_{\lambda}^q + F_{\mu}^q) \cdot v_{\lambda,\mu} \\ &= F_{\lambda}^q v_{\lambda} \cdot v_{\mu} + v_{\lambda} \cdot F_{\mu}^q v_{\mu} \\ &= \nabla_k^q (v_{\lambda}) \cdot v_{\mu} + v_{\lambda} \cdot \nabla_k^q (v_{\mu}) \end{aligned}$$

The first and the fourth equalities follow from Proposition 3.12, the second and the fifth from Proposition 6.1 and the third from Corollary 4.5. $\hfill \Box$

Corollary 6.3. Let $\lambda = (\lambda_1, \dots, \lambda_l)$ be a partition and $k \in \mathbb{N}$ a natural number. Denote by $\hat{\lambda}_i$ the partition λ without the *i*-th part. We have

$$\nabla_k^q(p_\lambda) = \sum_{i=1}^l \nabla_k^q(p_{\lambda_i}) p_{\widehat{\lambda}_i} \,.$$

The above corollary implies that in order to provide a formula for the operator ∇_k^q it is enough to compute $\nabla_k^q(p_n)$ for every $n \in \mathbb{N}$. We start with the case k = 1.

Proposition 6.4. We have

$$\nabla_1^q(p_n) = \sum_{\lambda \vdash n} (-1)^{l(\lambda)-1} \cdot \frac{1-q^n}{1-q} \cdot \frac{n}{z_\lambda} \cdot p_\lambda, \qquad \nabla_1(p_n) = \sum_{\lambda \vdash n} (-1)^{l(\lambda)-1} \cdot \frac{n^2}{z_\lambda} \cdot p_\lambda.$$

Proof. Let D be the operator defined by Equation (4). By definition

$$D(p_n) = \operatorname{Coeff}\left(z^0; \left(p_n + \frac{(1-q^n)(1-t^n)}{z^n}\right) \cdot \exp\left(\sum_{k \ge 1} \frac{-z^k p_k}{k}\right)\right)$$
$$= p_n \operatorname{Coeff}\left(z^0; \exp\left(\sum_{k \ge 1} \frac{-z^k p_k}{k}\right)\right) + (1-q^n)(1-t^n) \operatorname{Coeff}\left(z^n; \exp\left(\sum_{k \ge 1} \frac{-z^k p_k}{k}\right)\right)$$
$$= p_n + (1-q^n)(1-t^n) \cdot \sum_{\lambda \vdash n} \frac{(-1)^{l(\lambda)}}{z_\lambda} p_\lambda$$

We have $\nabla_1^{q,t} = \frac{1-D}{(1-q)(1-t)}$, therefore

$$\nabla_1^{q,t}(p_n) = \frac{(1-q^n)(1-t^n)}{(1-q)(1-t)} \cdot \sum_{\lambda \vdash n} \frac{(-1)^{l(\lambda)-1}}{z_\lambda} p_\lambda$$

To obtain the result, we compute the limit at $t \to 1$ and then at $q \to 1$.

It turns out that the coefficients of $\nabla_1^q(p_n)$ determine the operators ∇_k^q for an arbitrary $k \in \mathbb{N}$, as the next two propositions show.

Proposition 6.5. There exist polynomials $C_{\lambda}(q) \in \mathbb{Q}[q]$ such that for every $k \in \mathbb{N}$ we have

$$\nabla_k^q(p_n) = \sum_{\lambda \vdash n} \frac{(1-q^n) \cdot C_\lambda(q^k)}{\prod_{i=1}^{l(\lambda)} (1-q^{\lambda_i})} p_\lambda.$$

Proof. There exist rational numbers $a_{\mu}, b_{\mu\lambda}$ such that

$$p_n = \sum_{\mu \vdash n} a_\mu h_\mu , \qquad h_\mu = \sum_{\mu \vdash n} b_{\mu\lambda} p_\lambda .$$

Applying the plethystic substitution $f \to f[\frac{X}{1-q}]$, we get

$$\frac{p_n}{1-q^n} = \sum_{\mu \vdash n} a_\mu v_\mu$$

It follows that

$$\nabla_k(p_n) = (1-q^n) \sum_{\mu \vdash n} a_\mu \nabla_k(v_\mu) = (1-q^n) \sum_{\mu \vdash n} a_\mu B^q_\mu(q^k) v_\mu = (1-q^n) \sum_{\mu \vdash n} \sum_{\lambda \vdash n} \frac{a_\mu b_{\mu\lambda} B^q_\mu(q^k)}{\prod_{i=1}^{l(\lambda)} (1-q^{\lambda_i})} p_\lambda$$

Here we used $(\psi^k B)^q_\mu(q) = B^q_\mu(q^k)$. To obtain the result we set

$$C_{\lambda}(q) = \sum_{\mu \vdash n} a_{\mu} b_{\mu\lambda} B^{q}_{\mu}(q) \,. \qquad \Box$$

Corollary 6.6. The polynomial $C_{\lambda}(q)$ is given by the formula

$$C_{\lambda}(q) = (-1)^{l(\lambda)-1} \cdot \frac{\prod_{i=1}^{l(\lambda)} (1-q^{\lambda_i})}{(1-q)} \cdot \frac{n}{z_{\lambda}}$$

Proof. We compute the coefficient of p_{λ} in $\nabla_1^q(p_n)$ using Propositions 6.4 and 6.5. Comparing the results we obtain

$$\frac{(1-q^n) \cdot C_{\lambda}(q)}{\prod_{i=1}^{l(\lambda)} (1-q^{\lambda_i})} = (-1)^{l(\lambda)-1} \cdot \frac{1-q^n}{(1-q)} \cdot \frac{n}{z_{\lambda}}.$$

This implies the statement.

Corollary 6.7. We have

$$\nabla_k^q(p_n) = \sum_{\lambda \vdash n} (-1)^{l(\lambda)-1} \cdot \frac{1-q^n}{1-q^k} \cdot \left(\prod_{i=1}^{l(\lambda)} \frac{1-q^{k\lambda_i}}{1-q^{\lambda_i}}\right) \cdot \frac{n}{z_\lambda} \cdot p_\lambda \,.$$

Moreover,

$$\nabla_k(p_n) = \sum_{\lambda \vdash n} (-1)^{l(\lambda)-1} \cdot \frac{n^2 \cdot k^{l(\lambda)-1}}{z_\lambda} \cdot p_\lambda.$$

Proof. The first part is a direct consequence of Corollary 6.6 and Proposition 6.5. The second is obtained by computing the limit at $q \rightarrow 1$.

Proposition 6.8. We have

(6)
$$\nabla_k = \sum_{\lambda} \frac{(-k)^{l(\lambda)-1} s(\lambda)^2}{z_{\lambda}} \cdot p_{\lambda} \frac{\partial}{\partial p_{s(\lambda)}}.$$

Proof. Both sides of Equation (6) are endomorphisms of Λ which satisfy the Leibniz rule. Therefore, it is enough to check this equality evaluated on the ring generators of Λ . Corollary 6.7 imply that both sides have the same value on the elements p_n .

As a consequence we obtain a proof of Proposition 5.15, which in turn is a reformulation of the main result, Theorem 5.14, i.e. for any $k \in \mathbb{N}$ we have

$$\nabla_k^* = \sum_{\lambda} \frac{(-k)^{l(\lambda)-1} s(\lambda)}{\prod_{i \in \mathbb{N}} a_i(\lambda)!} p_{s(\lambda)} \partial_{\lambda} \,.$$

Proof of Proposition 5.15. By Proposition 3.6 (2), for any partition λ we have

$$\left(\frac{s(\lambda)^2}{z_{\lambda}} \cdot p_{\lambda} \frac{\partial}{\partial p_{s(\lambda)}}\right)^* = \frac{s(\lambda)^2}{z_{\lambda}} \cdot \frac{\prod_{i=1}^{l(\lambda)} \lambda_i}{s(\lambda)} \cdot p_{s(\lambda)} \partial_{\lambda} = \frac{s(\lambda)}{\prod_{i \in \mathbb{N}} a_i(\lambda)!} \cdot p_{s(\lambda)} \partial_{\lambda},$$

here we treat p_{λ} as a multiplication map $\Lambda \to \Lambda$. Multiply the above expression by $(-k)^{l(\lambda)-1}$ and sum over all partitions λ . Using Proposition 6.8, the left hand side is equal to ∇_k^* . The right hand side gives the desired formula.

Remark 6.9. An analogous reasoning may be repeated for the operator ∇_k^q . We obtain formulas

$$\nabla_k^q = \sum_{\lambda} \frac{(-1)^{l(\lambda)-1} s(\lambda)}{z_{\lambda}} \cdot \frac{1-q^{s(\lambda)}}{1-q^k} \cdot \left(\prod_{i=1}^{l(\lambda)} \frac{1-q^{k\lambda_i}}{1-q^{\lambda_i}}\right) \cdot p_{\lambda} \frac{\partial}{\partial p_{s(\lambda)}},$$
$$(\nabla_k^q)^* = \sum_{\lambda} \frac{(-1)^{l(\lambda)-1}}{\prod_{i\in\mathbb{N}} a_i(\lambda)!} \cdot \frac{1-q^{s(\lambda)}}{1-q^k} \cdot \left(\prod_{i=1}^{l(\lambda)} \frac{1-q^{k\lambda_i}}{1-q^{\lambda_i}}\right) \cdot p_{s(\lambda)} \partial_{\lambda}.$$

7. Multiplicative structure

The tensor product in K-theory of the $Hilb_n$ induces a product on the space of symmetric functions. In this section we provide an explicit formula for this product in the power sum basis.

Definition 7.1. The multiplication $\odot : \Lambda^n \times \Lambda^n \to \Lambda^n$ is defined as the map induced by the tensor product in K(Hilb_n), i.e.

$$f \odot g = \Theta^{-1}(\Theta(f) \otimes \Theta(g))$$
 .

The multiplication $\odot_q : \Lambda_q^n \times \Lambda_q^n \to \Lambda_q^n$ is defined as the map induced by the tensor product in $S_{\mathbb{T}_q}^{-1} \mathrm{K}^{\mathbb{T}_q}(\mathrm{Hilb}_n)$, i.e.

$$f \odot_q g = \Theta_q^{-1}(\Theta_q(f) \otimes \Theta_q(g))$$
 .

Definition 7.2. Let λ_1, λ_2 and μ be partitions of n. The structure constants $c^{\mu}_{\lambda_1,\lambda_2} \in \mathbb{Q}$ and the equivariant structure constants $c^{\mu}_{\lambda_1,\lambda_2}(q) \in \mathbb{Q}(q)$ are defined by

$$p_{\lambda_1} \odot p_{\lambda_2} = \sum_{\mu \vdash n} c^{\mu}_{\lambda_1, \lambda_2} p_{\mu} , \qquad p_{\lambda_1} \odot_q p_{\lambda_2} = \sum_{\mu \vdash n} c^{\mu}_{\lambda_1, \lambda_2}(q) p_{\mu} .$$

The product \odot is fully determined by its structure constants. We give a formula for the constant $c^{\mu}_{\lambda_1,\lambda_2}$ as a coefficient of a certain power series. First we introduce some notation. For a power series $W(q) \in \mathbb{Q}[[q-1]]$ and a natural number m, we write $\operatorname{Coeff}_{q-1}(m; W(q))$ for the coefficient of W(q) at $(q-1)^m$. We use the function

$$W_{\lambda}(q) = (q-1)^n \left\langle s_n, h_{\lambda} \left[\frac{X}{q-1} \right] \right\rangle \,.$$

An easy computation (cf. Example 3.9(4)) shows that

$$W_{\lambda}(q) = \sum_{\nu \vdash n} (q-1)^{n-l(\nu)} \frac{\langle p_{\nu}, h_{\lambda} \rangle}{z_{\nu}} \cdot [\nu]_q^{-1}.$$

The inner product $\langle p_{(1)^n}, h_\lambda \rangle$ is non-zero for any λ , thus $W_\lambda(q)$ has a non-zero restriction at q = 1.

Theorem 7.3. Let λ_1, λ_2 and μ be partitions of n. Then

$$c_{\lambda_1,\lambda_2}^{\mu} = \operatorname{Coeff}_{q-1}\left(l(\lambda_1) + l(\lambda_2) - n - l(\mu); \frac{[\mu]_q}{z_{\mu}[\lambda_1]_q[\lambda_2]_q} \cdot \sum_{\nu \vdash n} \frac{\langle p_{\lambda_1}, h_{\nu} \rangle \langle p_{\lambda_2}, h_{\nu} \rangle \langle p_{\mu}, m_{\nu} \rangle}{W_{\nu}(q)}\right)$$

Corollary 7.4. Suppose that $l(\lambda_1) + l(\lambda_2) - n - l(\mu)$ is negative, then $c^{\mu}_{\lambda_1,\lambda_2} = 0$.

Proof. The power series in Theorem 7.3 has a well-defined restriction at q = 1.

Remark 7.5. The above corollary agrees with [Boi06a, Theorem 1.1]. The condition that $l(\lambda_1) + l(\lambda_2) - n - l(\mu)$ is negative is equivalent to

$$(n - l(\lambda_1)) + (n - l(\lambda_2)) > n - l(\mu)$$

By [Boi06a, Theorem 1.1], for every partition λ the class ch($\Theta(p_{\lambda})$) has non-zero graded components in $\mathrm{H}^{2(n-l(\lambda))}(\mathrm{Hilb}_n)$ and is equal to zero in smaller gradations. Corollary 7.4 may be deduced from this theorem and the fact that the cup product in cohomology preserves gradation.

The rest of this section is devoted to the proof of Theorem 7.3. First, we study the \mathbb{T}_q -equivariant case. The tensor product in $S_{\mathbb{T}_q}^{-1} K_{\mathbb{T}_q}(\operatorname{Hilb}_n)$ has an easy presentation in the fixed point basis, cf. Proposition 4.1 (2). For a partition λ , let $[X_{\lambda}]$ be the class of the corresponding component of the fixed point set $\operatorname{Hilb}_n^{\mathbb{T}_q}$. Then

(7)
$$[X_{\lambda}] \otimes [X_{\mu}] = \delta_{\lambda,\mu} a_{\lambda} [X_{\lambda}],$$

for some rational functions $a_{\lambda} \in \mathbb{Q}(q)$. This observation allows us to describe the product \odot_q in Propositions 7.8 and 7.9. Next, we study the connection between the products \odot_q and \odot in Proposition 7.10. To obtain a formula for \odot we need to change the basis.

Lemma 7.6. Let V be a finite-dimensional vector space equipped with a non-degenerate bilinear form $\langle -, - \rangle$. Let A be an endomorphism of V. Suppose that $\alpha_1, \ldots, \alpha_k$ is an eigenbasis for A with eigenvalues a_1, \ldots, a_k . Then $\alpha_1^*, \ldots, \alpha_k^*$ is an eigenbasis for the adjoint map A^* with the same eigenvalues.

Proof. It is enough to show that for any $i, j \in \{1, ..., k\}$ we have $\langle A^* \alpha_i^*, \alpha_j \rangle = \langle a_i \alpha_i^*, \alpha_j \rangle$. This is straightforward, as

$$\langle A^* \alpha_i^*, \alpha_j \rangle = \langle \alpha_i^*, A \alpha_j \rangle = a_j \cdot \langle \alpha_i^*, \alpha_j \rangle = a_j \delta_{ij} = a_i \delta_{ij} = \langle a_i \alpha_i^*, \alpha_j \rangle.$$

Lemma 7.7. The set $\{m_{\lambda}[X(q-1)]\}_{\lambda \vdash n}$ is an eigenbasis of the map $\mathcal{E}_{B_n}^q$.

Proof. The set $\{h_{\lambda}[\frac{X}{1-q}]\}_{\lambda \vdash n}$ is an eigenbasis of ∇_1^q by Proposition 6.1. Therefore,

$$\left\{\omega\left(h_{\lambda}\left[\frac{X}{1-q}\right]\right)\right\}_{\lambda\vdash n} = \left\{h_{\lambda}\left[\frac{X}{q-1}\right]\right\}_{\lambda\vdash n}$$

is an eigenbasis of $\omega \nabla_1^q \omega$. We have that $(\mathcal{E}_B^q)^* = \omega \nabla_1^q \omega$, so by Lemma 7.6 it is enough to prove that the bases $\{h_{\lambda}[\frac{X}{q-1}]\}_{\lambda \vdash n}$ and $\{m_{\lambda}[X(q-1)]\}_{\lambda \vdash n}$ are mutually dual. This is the case, as

$$\left\langle h_{\lambda}\left[\frac{X}{q-1}\right], m_{\mu}[X(q-1)] \right\rangle = \left\langle h_{\lambda}, (m_{\mu}[X(q-1)])\left[\frac{X}{q-1}\right] \right\rangle = \left\langle h_{\lambda}, m_{\mu} \right\rangle = \delta_{\lambda\mu}.$$

The first equality follows from Example 3.9 (3), the second from Example 3.9 (2). \Box

Proposition 7.8. Let λ and μ be partitions of n. We have

$$m_{\lambda}[X(q-1)] \odot_q m_{\mu}[X(q-1)] = \delta_{\lambda\mu} b_{\lambda} m_{\mu}[X(q-1)]$$

for some rational function $b_{\lambda} \in \mathbb{Q}(q)$.

Proof. The fundamental classes $[X_{\lambda}] \in K^{\mathbb{T}_q}(\operatorname{Hilb}_n)$ form an eigenbasis of the map $- \otimes B_n$. Therefore, the elements $\Theta^{-1}[X_{\lambda}]$ form an eigenbasis of $\mathcal{E}_{B_n}^q$. The eigenvalues of $\mathcal{E}_{B_n}^q$ are pairwise different, therefore Lemma 7.7 implies that

$$\Theta^{-1}[X_{\lambda}] = d_{\lambda} \cdot m_{\lambda}[X(q-1)]$$

for some nonzero $d_{\lambda} \in \mathbb{Q}(q)$. The proposition now follows from Equation (7).

Proposition 7.9. For arbitrary symmetric functions $f, g \in \Lambda_q^n$, we have

$$f \odot_q g = \sum_{\mu \vdash n} p_\mu \cdot \frac{(q-1)^n}{z_\mu} \cdot \sum_{\nu \vdash n} \frac{\langle f, h_\nu[\frac{X}{q-1}] \rangle \langle g, h_\nu[\frac{X}{q-1}] \rangle \langle p_\mu, m_\nu[X(q-1)] \rangle}{W_\nu(q)}$$

Proof. Proposition 7.8 implies that

(8)
$$f \odot_q g = \sum_{\nu \vdash n} \left\langle f, h_{\nu} \left[\frac{X}{q-1} \right] \right\rangle \left\langle g, h_{\nu} \left[\frac{X}{q-1} \right] \right\rangle b_{\nu} \cdot m_{\nu} [X(q-1)]$$

for some $b_{\nu} \in \mathbb{Q}(q)$. It is a standard fact that $\Theta_q(s_n)$ is the trivial bundle. Therefore s_n is the neutral element for \odot_q . For $g = s_n$ and $f = m_{\lambda}[X(q-1)]$, Equation (8) gives

$$m_{\lambda}[X(q-1)] = \sum_{\nu \vdash n} \delta_{\lambda\nu} \left\langle s_n, h_{\nu} \left[\frac{X}{q-1} \right] \right\rangle b_{\nu} \cdot m_{\nu}[X(q-1)].$$

It follows that

$$b_{\lambda} = \left\langle s_n; h_{\lambda} \left[\frac{X}{q-1} \right] \right\rangle^{-1} = \frac{(q-1)^n}{W_{\lambda}(q)}.$$

Moreover, $m_{\nu}[X(q-1)] = \sum_{\mu \vdash n} \langle p_{\mu}, m_{\nu}[X(q-1)] \rangle_{z_{\mu}}^{\underline{p}_{\mu}}$. After substituting these two equations into (8) we obtain the desired formula.

Proposition 7.10. Suppose that the symmetric functions $f, g \in \Lambda_q^n$ have well-defined restrictions at q = 1. Then $f \odot_q g$ also has a well-defined restriction at q = 1. Moreover,

$$(f \odot_q g)_{|q=1} = f_{|q=1} \odot g_{|q=1}$$

Proof. An analogous property is satisfied by the products in $S_{\mathbb{T}_q}^{-1} \mathrm{K}^{\mathbb{T}_q}(\mathrm{Hilb}_n)$ and $\mathrm{K}(\mathrm{Hilb}_n)$.

Proof of Theorem 7.3. Proposition 7.9 and Example 3.9 (4) imply that the formula for the equivariant structure constants is the following:

$$c_{\lambda_{1},\lambda_{2}}^{\mu}(q) = (q-1)^{n+l(\mu)-l(\lambda_{1})-l(\lambda_{2})} \frac{[\mu]_{q}}{z_{\mu}[\lambda_{1}]_{q}[\lambda_{2}]_{q}} \cdot \sum_{\nu \vdash n} \frac{\langle p_{\lambda_{1}}, h_{\nu} \rangle \langle p_{\lambda_{2}}, h_{\nu} \rangle \langle p_{\mu}, m_{\nu} \rangle}{W_{\nu}(q)}.$$

By Proposition 7.9 the above expression has a well-defined restriction at q = 1.

Appendix A. Equivariant McKay correspondence

Definition A.1. Let k be a field and V be a k-vector space of finite dimension. The restriction

$$\operatorname{res}_{q=1}: V \otimes_k k(q) \dashrightarrow V$$

is defined on a submodule $V \otimes k[q]_{\mathfrak{m}}$, where \mathfrak{m} is the ideal spanned by (q-1).

Definition A.2. Let k be a field, V and W be k-vector spaces of finite dimension and $\psi: V \to W$ be a k-linear map. We say that a k(q)-linear map $\psi_q: V \otimes_k k(q) \to W \otimes_k k(q)$ restricts to ψ if the following diagram commutes:

$$\begin{array}{cccc} V \otimes_k k(q) & \stackrel{\psi_q}{\longrightarrow} & W \otimes_k k(q) \\ \stackrel{\operatorname{res}_{q=1}}{\stackrel{\downarrow}{\longrightarrow}} & \stackrel{\downarrow}{\underset{\psi}{\longrightarrow}} & \stackrel{\operatorname{res}_{q=1}}{\overset{\psi}{\longrightarrow}} & , \\ V & \stackrel{\psi}{\longrightarrow} & W \end{array}$$

in the sense that if for $v \in V \otimes_k k(q)$ the restriction $\operatorname{res}_{q=1} v$ is defined, then the restriction $\operatorname{res}_{q=1} \psi_q(v)$ is also defined and equal to $\psi(\operatorname{res}_{q=1} v)$.

Lemma A.3. Let k be a field, let V and W be k-vector spaces of finite dimension, and let $\psi : V \to W$ be a k-linear map. Let $\psi_q : V \otimes_k k(q) \to W \otimes_k k(q)$ be a k(q)-linear map which restricts to ψ . Suppose that ψ is an isomorphism. Then ψ_q is also isomorphism. *Proof.* Let v_1, \ldots, v_k be a basis of V. Then $v_1 \otimes 1, \ldots, v_k \otimes 1$ is a k(q)-basis of $V \otimes k(q)$. It is enough to prove that the vectors $\psi_q(v_1 \otimes 1), \ldots, \psi_q(v_k \otimes 1)$ are k(q)-linearly independent. Suppose otherwise that there exist rational functions $a_k(q) \in k(q)$ such that

(9)
$$\sum_{i=1}^{k} a_k(q)\psi_q(v_i \otimes 1) = 0$$

Without loss of generality we can assume that all $a_k(q)$ are non-zero. Let val_{q-1} be the valuation associated with (q-1) and

$$t = \min\{\operatorname{val}_{q-1}(a_1(q)), \dots, \operatorname{val}_{q-1}(a_k(q))\}$$

Multiplying Equation (9) by $(q-1)^{-t}$ and applying $\operatorname{res}_{q=1}$ yields a non-trivial k-linear relation between elements $\operatorname{res}_{q=1}(\psi_q(v_i \otimes 1))$. The map ψ_q restricts to ψ , therefore

$$\operatorname{res}_{q=1}(\psi_q(v_i\otimes 1)) = \psi(v_i)$$

This contradicts that ψ is an isomorphism.

Proof of Corollary 5.3. Let p and ρ denote the maps from the diagram (3). Let ρ_* , $\rho_*^{\mathbb{T}_q}$, $\rho_*^{\mathbb{T}}$, p^* , $p_{\mathbb{T}_q}^*$,

Set $k = \mathbb{Q}, V = K(\operatorname{Hilb}_n), W = \operatorname{K}_{\operatorname{S}_n}(\operatorname{pt})$. We have

$$V \otimes_k k(q) = S_{\mathbb{T}_q}^{-1} \operatorname{K}_{\mathbb{T}_q}(\operatorname{Hilb}_n), \qquad W \otimes_k k(q) = S_{\mathbb{T}_q}^{-1} \operatorname{K}_{\mathbb{T}_q \times \operatorname{S}_n}(\operatorname{pt}).$$

The map $\rho_*^{\mathbb{T}_q} \circ p_{\mathbb{T}_q}^*$ restricts to $\rho_* \circ p^*$ which is an isomorphism. Therefore $\rho_*^{\mathbb{T}_q} \circ p_{\mathbb{T}_q}^*$ is also an isomorphism.

Set
$$k = \mathbb{Q}(q)$$
, $V = S_{\mathbb{T}_q}^{-1} \operatorname{K}_{\mathbb{T}_q}(\operatorname{Hilb}_n)$, $W = S_{\mathbb{T}_q}^{-1} \operatorname{K}_{\mathbb{T}_q \times \operatorname{S}_n}(\operatorname{pt})$. We have
 $V \otimes_k k(t) = S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T}}(\operatorname{Hilb}_n)$, $W \otimes_k k(t) = S_{\mathbb{T}}^{-1} \operatorname{K}_{\mathbb{T} \times \operatorname{S}_n}(\operatorname{pt})$.

The map $\rho_*^{\mathbb{T}} \circ p_{\mathbb{T}}^*$ restricts to $\rho_*^{\mathbb{T}_q} \circ p_{\mathbb{T}_q}^*$ which is an isomorphism. Therefore $\rho_*^{\mathbb{T}} \circ p_{\mathbb{T}}^*$ is also an isomorphism.

References

- [AB84] M. F. Atiyah and R. Bott. The moment map and equivariant cohomology. *Topology*, 23(1):1–28, 1984.
- [Ada62] J. F. Adams. Vector fields on spheres. Ann. of Math. (2), 75:603-632, 1962.
- [Bea00] A. Beauville. Symplectic singularities. *Invent. Math.*, 139(3):541–549, 2000.
- [BGHT99] F. Bergeron, A. M. Garsia, M. Haiman, and G. Tesler. Identities and positivity conjectures for some remarkable operators in the theory of symmetric functions. volume 6, pages 363–420. 1999. Dedicated to Richard A. Askey on the occasion of his 65th birthday, Part III.
- [BKR01] T. Bridgeland, A. King, and M. Reid. The McKay correspondence as an equivalence of derived categories. J. Amer. Math. Soc., 14(3):535–554, 2001.
- [BL05] L. Borisov and A. Libgober. McKay correspondence for elliptic genera. Ann. of Math. (2), 161(3):1521–1569, 2005.
- [BNW07] S. Boissière and M. A. Nieper-Wisskirchen. Generating series in the cohomology of Hilbert schemes of points on surfaces. LMS J. Comput. Math., 10:254–270, 2007.
- [Boi06a] S. Boissière. On the McKay correspondences for the Hilbert scheme of points on the affine plane. *Math. Ann.*, 334(2):419–438, 2006.
- [Boi06b] S. Boissière. Towards the multiplicative behavior of the K-theoretical McKay correspondence. Math. Z., 252(3):533–555, 2006.
- [CE15] P.-E. Chaput and L. Evain. On the equivariant cohomology of Hilbert schemes of points in the plane. Ann. Inst. Fourier (Grenoble), 65(3):1201–1250, 2015.
- [CG10] N. Chriss and V. Ginzburg. Representation theory and complex geometry. Modern Birkhäuser Classics. Birkhäuser Boston, Ltd., Boston, MA, 2010. Reprint of the 1997 edition.
- [DBW18] M. Donten-Bury and A. Weber. Equivariant Hirzebruch classes and Molien series of quotient singularities. *Transform. Groups*, 23(3):671–705, 2018.

- [ES87] G. Ellingsrud and S. A. Strømme. On the homology of the Hilbert scheme of points in the plane. *Invent. Math.*, 87(2):343–352, 1987.
- [ES93] G. Ellingsrud and S. A. Strømme. Towards the Chow ring of the Hilbert scheme of P². J. Reine Angew. Math., 441:33–44, 1993.
- [Eva04] L. Evain. Irreducible components of the equivariant punctual Hilbert schemes. Adv. Math., 185(2):328–346, 2004.
- [FL85] W. Fulton and S. Lang. Riemann-Roch algebra, volume 277 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, New York, 1985.
- [Fog68] J. Fogarty. Algebraic families on an algebraic surface. American Journal of Mathematics, 90(2):511–521, 1968.
- [GH96] A. M. Garsia and M. Haiman. A remarkable q, t-Catalan sequence and q-Lagrange inversion. J. Algebraic Combin., 5(3):191–244, 1996.
- [Hai01] M. Haiman. Hilbert schemes, polygraphs and the Macdonald positivity conjecture. J. Amer. Math. Soc., 14(4):941–1006, 2001.
- [Hai02] M. Haiman. Vanishing theorems and character formulas for the Hilbert scheme of points in the plane. *Invent. Math.*, 149(2):371–407, 2002.
- [Hai03] M. Haiman. Combinatorics, symmetric functions, and Hilbert schemes. In Current developments in mathematics, 2002, pages 39–111. Int. Press, Somerville, MA, 2003.
- [IN00] Y. Ito and H. Nakajima. McKay correspondence and Hilbert schemes in dimension three. Topology, 39(6):1155–1191, 2000.
- [Kal02] D. Kaledin. McKay correspondence for symplectic quotient singularities. Invent. Math., 148(1):151–175, 2002.
- [Kal09] D. Kaledin. Geometry and topology of symplectic resolutions. In Algebraic geometry—Seattle 2005. Part 2, volume 80 of Proc. Sympos. Pure Math., pages 595–628. Amer. Math. Soc., Providence, RI, 2009.
- [Kru14] A. Krug. Tensor products of tautological bundles under the Bridgeland-King-Reid-Haiman equivalence. *Geom. Dedicata*, 172:245–291, 2014.
- [Kru18] A. Krug. Remarks on the derived McKay correspondence for Hilbert schemes of points and tautological bundles. *Math. Ann.*, 371(1-2):461–486, 2018.
- [KZ23] J. Koncki and M. Zielenkiewicz. Nakajima's creation operators and the Kirwan map, 2023. arXiv:2304.07241.
- [Leh99] M. Lehn. Chern classes of tautological sheaves on Hilbert schemes of points on surfaces. Invent. Math., 136(1):157–207, 1999.
- [LS01] M. Lehn and C. Sorger. Symmetric groups and the cup product on the cohomology of Hilbert schemes. *Duke Math. J.*, 110(2):345–357, 2001.
- [Mac15] I. G. Macdonald. Symmetric functions and Hall polynomials. Oxford Classic Texts in the Physical Sciences. The Clarendon Press, Oxford University Press, New York, second edition, 2015. With contribution by A. V. Zelevinsky and a foreword by Richard Stanley.
- [MN18] K. McGerty and T. Nevins. Kirwan surjectivity for quiver varieties. *Invent. Math.*, 212(1):161–187, 2018.
- [MW21] M. Mikosz and A. Weber. Elliptic classes, McKay correspondence and theta identities. J. Algebraic Combin., 53(3):701–727, 2021.
- [Nak99] H. Nakajima. Lectures on Hilbert schemes of points on surfaces, volume 18 of University Lecture Series. American Mathematical Society, Providence, RI, 1999.
- [Ohm08] T. Ohmoto. Generating functions of orbifold Chern classes. I. Symmetric products. Math. Proc. Cambridge Philos. Soc., 144(2):423–438, 2008.
- [Sca09] L. Scala. Cohomology of the Hilbert scheme of points on a surface with values in representations of tautological bundles. *Duke Math. J.*, 150(2):211–267, 2009.
- [Smi20] A. Smirnov. Elliptic stable envelope for Hilbert scheme of points in the plane. Selecta Math. (N.S.), 26(1):Paper No. 3, 57, 2020.
- [Sta99] R. P. Stanley. Enumerative combinatorics. Vol. 2, volume 62 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1999. With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin.
- [SV13] O. Schiffmann and E. Vasserot. The elliptic Hall algebra and the K-theory of the Hilbert scheme of \mathbb{A}^2 . Duke Math. J., 162(2):279–366, 2013.
- [Tho92] R. W. Thomason. Une formule de Lefschetz en K-théorie équivariante algébrique. *Duke Math. Journal*, 68:447–462, 1992.

[Wae08] R. Waelder. Equivariant elliptic genera and local McKay correspondences. Asian J. Math., 12(2):251–284, 2008.

INSTITUTE OF MATHEMATICS OF THE POLISH ACADEMY OF SCIENCES, POLAND *Email address*: j.koncki@mimuw.edu.pl

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, POLAND *Email address*: magdaz@mimuw.edu.pl