

---

# SURVEY ON BIOMARKERS IN HUMAN VOCALIZATIONS

---

A PREPRINT

Aki Harma<sup>1\*</sup>, Bert den Brinker<sup>2</sup>, Ulf Grossekathefer<sup>2</sup>, Okke Ouweltjes<sup>2</sup>,  
Srikanth Nallanthighal<sup>3</sup>, Sidharth Abrol<sup>3</sup> and Vibhu Sharma<sup>2</sup>

July 7, 2024

ABSTRACT

Recent years has witnessed an increase in technologies that use speech for the sensing of the health of the talker. This survey paper proposes a general taxonomy of the technologies and a broad overview of current progress and challenges. Vocal biomarkers are often secondary measures that are approximating a signal of another sensor or identifying an underlying mental, cognitive, or physiological state. Their measurement involve disturbances and uncertainties that may be considered as noise sources and the biomarkers are coarsely qualified in terms of the various sources of noise involved in their determination. While in some proposed biomarkers the error levels seem high, there are vocal biomarkers where the errors are expected to be low and thus are more likely to qualify as candidates for adoption in healthcare applications.

Index Terms: Pathological speech sensing, speech breathing, respiratory health and fitness, automatic speech recognition

## 1 Introduction

Human language is said to go back in time at least 1.5 million years [1] but animal vocalization has naturally a much longer history [2]. Animal vocalization and human speech can be seen as adaptations of the respiratory system to the communication. Vocalization requires a high level of neural control of the muscles in the vocal track consisting of lungs, trachea, and the mouth. Speech, with a communicative message, results from cognitive processes in the brain which reflect all aspects of human life. Conversely, mental, cognitive, and physiological changes may have impact on vocalization. Vocal biomarkers, acoustic signatures extracted from speech and other vocalizations, are therefore expected to reflect these.

Surveys on the use of voice for health and fitness monitoring have been published, for example, by Fagrehazzi et al. [3] and Ferro et al. [4]. Kraus defines that a vocal biomarker is a signature, a feature, or a combination of features from the audio signal of the voice that is associated with a clinical outcome and can be used to monitor patients, diagnose a condition, or grade the severity or the stages of a disease or for drug development [5]. For the current report we use a broader definition of vocal biomarkers technology as systems that aim at using human vocalization, captured with a microphone, to detect a condition of the talker, or estimate the measurement values of another sensory modality.

Auscultation based on listening to sounds of heart, lungs, and bowel, using stethoscope, for example, has a long history in medicine ranging to 1800's [6,7]. In this report, we focus on vocalizations captured from air using a microphone. We leave out, for example, breathing sounds captured using microphones of various kinds and other bodily sounds not produced by air flow passing through the vocal tract. We also leave out ingressive vocalizations [8] and sounds captured by bone conduction, e.g., by in-ear headphones or eyeglasses.

Survey on biomarkers in human vocalizations

In vocalizations, the main interest is in conversational speech, for example, captured during a telephone call or in using a speech user interface in a smart speaker. However, there are several other types of vocalizations we want to cover. Some of those have been listed in Table 1, which also shows some of the detection targets such as asthma,

<sup>\*1</sup> DACS, Maastricht University, The Netherlands. <sup>2</sup> Philips Research, Eindhoven, The Netherlands. <sup>3</sup> Philips Research, Bangalore, India




Figure 1: Different aspects of speech communication research

depression, or alcohol intoxication. In estimating measurement values of another sensing modality our own case is Virtual Respiratory Belt (VRB) processing where we estimate from speech a waveform of a respiratory belt sensor signal of the talker. The opposite cases of sensing speech from other modalities, e.g., silent-speech interfaces based on speech movements or BMI [9], or methods that focus on speech reception testing [10] are out of the scope of this review.

Vocalization	Breathing sounds	Involuntary non-verbal vocalizations	Other nonverbal vocalization	Spontaneous speech	Protocolled speech
Examples	Normal breathing, wheezing, crackles	Snoring, groaning, grunting, tics	Coughing, sneezing, hiccups, backchanneling	Conversational speech, free speech	Read speech, therapeutic protocols
Condition	Asthma, COPD, covi d19	Apneas, Tourette’s syndrome	Infections, COPD, alcohol, depression	Respiratory health, AD, PD, depression	Dementia, arrythmi as

Table 1: Types of vocalizations

Speech communication is a central characteristic of humanity, and it is influenced by every aspect of our lives. It is also a multidimensional concept which we can study from many different perspectives, as illustrated in Fig. 1. The content of speech is the information we deliver to the listeners using a language. The vocal characteristics of speech results from our speech production system consisting of neural control and acoustics of the lungs, trachea, and larynx, and the vocal tract. The prosodic elements of intonation, tone, and rhythm are often modulated by various environmental and intrinsic factors. Prosodics is often thought to be a part of paralinguistic communication which also contain body-language and other behaviors. In conversational behavior, the speech is influenced by other

participants. Typical research topics in this area are conversational turn-taking and back-channeling behaviors which may reflect intrinsic mental and cognitive processes, but also habits, social conventions, and environmental constraints [11].

All aspects of speech communication illustrated in Fig. 1 and described below may be influenced by relevant health factors. In the following sections, the goal is to give an overview and focus on areas where the relationship has been established, and where it has been demonstrated that the condition can be detected using computational speech analytics and machine learning.

At the level of the dialogue, we may also consider the roles of the participants, their goals, values, and strategies. Often a dialogue can be seen as a collaborative problem solving or as a negotiation between the respondent and the caller. A dialogue may have measurable outcomes. For example, in a sales call center, the negotiation with a customer may lead to the conversion from free to a paying subscriber of the service. However, often the outcomes are not directly available but may influence the customer experience and give trust in the provided service on a longer term.




Figure 2: Sensing targets and interventions in different time-scales

Speech is personal learned behaviour and reflecting all aspects of the individual talker. Consequently, many of the sensing targets discussed in this report are only available at the level of an individual. For example, a mild cognitive decline or increase in the roughness of a voice can only be assessed in relation to a historical data from the same subject. This may also look problematic because of natural aging and changes in the environment of the talker. Nevertheless, we demonstrated in [12], that it is possible to train a multi-task classification system for independent prediction of the age and the level of cognitive decline in a talker.

We may look at the area of vocal biomarkers research on different time scales of sensing and intervention, see Fig. 2. Longitudinal speech recordings can be used to detect background conditions and gradual changes in various personality [13] and health factors. In a shorter time-scale, speech can be used as an unobtrusive measurement for various factors which are traditionally measured by sensors attached to the body, e.g., the blood glucose level [14]. Finally, we can use speech and vocalizations for alarms and precise control of, for example, a ventilatory support device used by a patient [15].

The outline of the paper is as follows. We recapture various aspects of vocalizations regarding speech and respiration in Section 2. The survey continues with an overview of the determination of physiological health conditions (Section 3) from vocal biomarkers followed by cognitive and mental health conditions (Section 4). Section 5 is devoted to applications and Section 6 is a first attempt to arrive at a qualitative performance overview of various vocal biomarkers. The paper closes with a conclusion and discussion of some of the future challenges in vocal biomarkers research.

## 2 Vocalization aspects: speech and respiration

In this section, aspects around speech respiration are recaptured. We start with considering aspects of speech content (Section 2.1) and the neural control (Section 2.1). We follow up with respiratory aspects (Section 2.3), paralinguistic (Section 2.4), and insights from conversational behavior (Section 2.5).

### 2.1 Content of speech

The analysis of transcripts of counseling sessions has a long history especially in psychiatry [16]. Traditionally, the text analysis was performed by a human expert, but a similar performance for a computational method was already demonstrated 20 years ago [17]. Recent work on text-based detection of depression in a popular data set shows an accuracy of 66% for single conversation turn and close to 100% when accumulated over the entire conversation [18]. The performance is robust for manual and automatic transcriptions of the original clean speech recordings [19]. In the last two years we have seen fast progress in the performance of large language models in understanding of human speech [20]. This is influencing all tasks where the speech content is relevant.

Aphasia is a common neurodegenerative disorder often following a stroke, head injury, or a brain tumor [21]. While the disorder primarily influences the speaker's vocabulary, it naturally has also an impact on paralinguistics, the fluency of speech production, and conversation. There are distinct types of aphasia. Fraser et al. demonstrated over 90% accuracy in detecting agrammatic aphasia, characterized by the omission of words, based on the analytics of transcriptions of 300-400 word long free narratives [22].

Aphasia patients often suffer from anomia, characterized by difficulties in naming objects. Peintner et al. [23] compared the detectability of aphasia patients from healthy controls in an automatic detection task using pure linguistic, prosodic, or a combination of both types of features. The accuracy when using only linguistic or prosodic features gave an accuracy of around 80% in their data set of 3-5 minutes speech recordings. The combination of the two feature types increased the accuracy performance. A recent paper by Le et al. [24] studied the separate detectability of different paraphasias in aphasia patients. Typical paraphasias are semantic and phonetic errors where target words are replaced semantically or phonetically by similar words, respectively. In neologistic errors, the replacement word is not related to the original target word. The detection of these categories requires a high level of understanding of the content of speech, which was in this case achieved using conventional speech features and deep recurrent neural networks (RNNs). The best results were obtained in the case of phonetic paraphasias, but the results are moderate. The F1-score, which is a commonly used measure to characterize the performance of a classifier gets a value in 0.5-0.7, on a scale from 0.0 to 1.0.

The difficulties in finding words in aphasia caused by stroke (Wernicke's aphasia) or by dementia is different [25]. Dementia patients, especially in the early phase of the disease, are less impaired in spontaneous speech. However, the deterioration of lexical-semantic processing in the brain starts and becomes detectable, already in an early phase of the disease [26].

### 2.2 Neural control of speech production

Dimauro et al. [27] compared the word error rates of an ASR system in Parkinson's Disease (PD) patients and healthy controls. The word error rates were significantly higher in the PD population. In PD, the lower intelligibility of speech is not necessarily related to cognitive functions but reflects typical symptoms of PD including tremors and rigidity and slow movement of the muscles, bradykinesia. The severity of PD is often characterized using the Unified Parkinson's Disease Rating Scale, UPDRS [28]. The detection target in PD speech is typically the UPDRS score of the patient.

Dysarthria is a common name for the speech impairments caused by difficulties in the muscular control of speech caused by a neurological problem in the motor control of speech muscles. Common conditions include Parkinson's, amyotrophic lateral sclerosis (ALS), and multiple sclerosis (MS) diseases. The severity of dysarthria can be described on a subjective four-level Dysarthria Severity Degree (DSD) scale. Laaridh et al. [29] have demonstrated that the DSD can be predicted from one minute of text reading, but the accuracy is not very high. It has been shown that the changes in the larynx caused by rheumatoid arthritis changes the quality of the voice of the patient [30].

Voice disorder, according to the definition of American Speech-Language-Hearing Association (ASHA), occurs when the voice quality, pitch, and loudness differ or are inappropriate for an individual's age, gender, cultural background, or geographic location [29]. Typical characterizations of voice disorders are listed in Table 1. Voice disorders are categorized into Organic or Functional disorders depending on whether there is a clear physiological reason for the disorder or not, respectively. One example of a functional disorder is temporary vocal fatigue of the talker. Organic disorders may be caused by a structural or neurogenetic reason. In this terminology, dysarthria is an organic neurogenetic condition.

The detection of a dysarthric condition is often performed in a protocolled speech task. For example, in [31], dysarthric speech due to Parkinson's disease was detected in a task that consisted of fast repetition of pa/ta/ka syllables and a text reading task. In the study, the accuracy in the detection task ranged from 60 to 99%.

Damage to muscles or other structures involved in speech production, for example caused by an injury or a cancer, may lead to changes in articulation. In some rare cases, such as the cardio-vocal syndrome, the characteristics of voice may

Name	Description
roughness	perception of aberrant vocal fold vibration
breathiness	perception of audible air escape in the sound ...
strained quality	perception of increased effort; tense or harsh...
strangled quality	as if talking with breath held
abnormal pitch	too high, too low, pitch breaks, decreased pit...
abnormal loudness/volume	too high, too low, decreased range, unsteady v...
abnormal resonance	hypernasal, hyponasal, cul de sac resonance
aphonia	loss of voice
phonation breaks	breaks in speech voice
asthenia	weak voice
gurgly/wet sounding voice	saliva effects
hoarse voice	raspy, audible aperiodicity in sound
pulsed voice	fry register, audible creaks or pulses in sound
shrill voice	high, piercing sound, as if stifling a scream
tremulous voice	shaky voice; rhythmic pitch and loudness undul...
high effort	increased vocal effort associated with speaking;
low fatigue	decreased vocal endurance or onset of fatigue ...
variability	variable vocal quality throughout the day or d...
coughing	frequent coughing or throat clearing (may wors...
tensioned	excessive throat or laryngeal tension/pain/ten...

Table 2: Voice disorders by American Speech-Language-Hearing Association (ASHA)

---

Survey on biomarkers in human vocalizations

---

Non-verbal tics	Verbal tics
Throat clearing	Repeating parts of words, or phrases
Sniffing	Coprolalia
Coughing	Prosodic changes
Yelling	Stuttering
Hiccupping	Talking to oneself (different identities)
Snorting	Derogatory phrases or statements
Animal sounds (e.g., barking)	Repetitive sounds
Belching	Silent palilalia

Table 3: Typical speech effects characterized by Tourette’s syndrome

change due to heart disease [32] and this can be detected even from smartphone recordings [33]. Tourette’s syndrome is characterized by various effects on speech motoric and language [34]. Some of the typical speech and language symptoms, see Table 3, are non-verbal motoric behaviors and some changes in the prosody or the linguistic content. The use of speech as a modality of triaging potential TS patients could be particularly useful for young children [35].

### 2.3 Respiratory functions

The speech production is also influenced by the infection of the lungs and the respiratory tract. In a recent challenge, the task was to detect a subject who has a cold [36]. The data set had 630 talkers, and the data was labeled according to the Wisconsin Upper Respiratory Symptom Survey (WURSS-24) score which ranks the illness severity into seven levels ranging from not sick to severely sick [37]. The data contains scripted and free speech. The best results in the detection of respiratory infection from speech had the Unweighted Average Recall (UAR) over 90% [38]. UAR measure is often used to compare classifiers in speech detection tasks, and it represents the mean of the recall values computed for the two classes.

The cough detection is a widely studied subject [39] as it is a symptom of many respiratory diseases, see also Section 3.4. Cough is used in different ways. One is the detection or monitoring of cough and is a longitudinal measurement. The other one is information extraction from cough and is a spot check. Obviously, these can be combined in a two-step approach. Cough sounds have been studied in terms of its character (wet/dry, productive/nonproductive, voluntary/spontaneous), its source (male/female), health status (healthy/unhealthy), lung condition (obstructive/restrictive/normal), and disease diagnosis (croup, COPD, asthma, pneumonia, pertussis, tuberculosis, bronchitis, COVID-19, cold, heart failure, lower/upper respiratory tract disease). Various devices can be used, like wearables (acceleration sensor where various areas of the body have been used), dedicated tube systems, or microphone recordings. It is generally accepted that cough is characterized by large inter-person and intra-person variability. This may be the reason that reported performance varies quite a lot. ResAppHealth is the major proponent of disease diagnosis, see Section 3.4. Cough detection and classification has received quite some attention, e.g., [40–42]. The latest deep learning algorithms achieve very good performance for the detection of single cough sounds [43,44]. Di Perna et al. [45] reported a system for unobtrusive, privacy-preserving, long-term monitoring of cough. However, a good classifier developed in the lab does not immediately imply success for clinical applications. The gap between cough detection systems and clinical applications is still not fully bridged: what is missing are trials showing the effect of cough monitoring combined with clinical intervention, see Section 3.4.

Detection of a risk of sleep apnea from speech signals [46,47] has been considered as well. However, the results seem mixed [48]. During the severe acute respiratory syndrome coronavirus 2, SARS-CoV-2, or the COVID-19 epidemic there has been a lot of attention to symptom tracking from speech recordings. Shortness of breath, or dyspnea, is one of the key indicators detectable from early disease phases. The current authors have developed technology for deep respiratory sensing from speech [49] and detection of mild dyspnea from pairs of speech recordings from the same individual [50] and describe this work in more detail in Section 4.6. Regarding COVID-19, it has also been shown that shortness of breath, and self-reported hoarse voice, see Table 1, are among the most important symptoms predicting

the long COVID-19 condition [51] where the symptoms continue often for several months after the initial infection. A recent study using convolutional networks for the COVID-19 classification from crowd-sourced speech and cough recordings demonstrated convincing results with AUC of 0.846 [52]. The paper also gives a good overview of various recent COVID-19 speech data collection studies.

## 2.4 Paralinguistics

Complex cognitive functions related to affect and emotion may influence speech communication at all levels of Fig. 1, see [53] for an overview. In particular, detection of depression [54] and bipolar disorder [55] has been a popular research topic in areas of speech, language, and Human-Computer Interaction (HCI) studies. A recent review by Morales et al. [56] aimed at comparing the literature results in the detection of depression from different detection modalities including speech, text analysis, and visual communication data. The results were not conclusive due to incomparability. Typical accuracy values ranged from 0.6 to 0.85, and the authors gave a general recommendation to use multi-modality features in the detection task. In an overview of the detectability of depression and suicide risk, Cummins et al. [57] concluded that the state-of-the-art methods for depression detection reaches an accuracy of 79%.

The detection of distress or arousal in a crisis hotline services [58] and other medical call center services [59] is an important application for speech analytics. Many commercial call center systems also provide detection of arousal based on voice and sentiment analysis of transcribed speech. While dementia has less impact on the fluency of speech, mild dementia may already lead to changes in the paralinguistics that can be detected from the speech signal. For example, a study showed 85% accuracy in the detection of mild dementia based on prosodic features [60]. In our recent work combining prosodic and linguistic for the detection of dementia condition, we obtained slightly higher performance [61].

It has been shown that the duration of pauses during a speech in ALS patients differs significantly from healthy controls [62]. The changes in speech rate are also characteristic of various forms of aphasia such as the primary progressive aphasia [63]. The articulation rate may reduce from an average of 5 syllables per second in healthy controls to 1-2 syllables per second. The duration of pauses also increases in the same proportion.

## 2.5 Conversational behavior

Typical phenomena in conversational behavior are turn-taking organization, backchanneling, accommodation, and response times. Turn-taking organization refers to the way the target talker takes a turn in the conversation. Backchanneling contains, for example, verbal and non-verbal acknowledgments during the speech of the other participant. Accommodation refers to various ways on how a talker adapts the vocabulary and speaking style to the other. These phenomena are naturally influenced by various health, social and environmental factors, see [64]. One example of accommodation is entrainment where a talker adopts the speech rhythm of another talker. There is evidence that entraining is less accurate in Parkinson's patients than in healthy controls [65]. Otherwise, it seems that the area of automatic conversation analytics for health conditions is not widely explored. Research so far has mostly focused on applications in psychiatry, certain brain diseases, and developmental problems in children. Clinical interviewing is a




Figure 3: Virtual Respiratory Belt sensing method where the goal is to predict the signal of a RIP belt from normal conversational speech.

popular way of diagnosing various health conditions [66]. The interviewing techniques such as motivational interviewing [67] or Cognitive Behavioral Therapy [68] in psychology, or Clinical Dementia Rating [69] or psycholinguistic assessment for aphasia [70] have been shown to be effective as diagnostic methods or interventions. The interview protocols mostly focus on the content, but practitioners naturally use also nonverbal cues to guide the interviewing. The use of automatic speech analytics has often been proposed as an additional tool to help in clinical interviews, yet the author of this report is not aware of established protocols for it. Scherer et al. [71] studied the changes in the speech in a conversation between an interviewer and patients with different severity of depression. Their results showed that the accommodation measured by the correlation between the speech of the patient and the practitioner was reduced in more depressed patients. A high level of accommodation is often suggested to relate to an increased positive attitude and liking, and therefore the result is understandable [72]. Stolar et al. [73] also concluded that there was a clear change in the speech parameters of the interviewer, i.e., more breathy voice, with more depressed patients. They reported a 94% classification accuracy in the detection of a depressed or non-depressed adolescent in conversation dynamics in parent-adolescent conversations, but the data set was small.

### 3 Physiological health conditions

In this overview report we make a separation between the use of speech for physiological, cognitive, and mental conditions. Obviously, there is a lot of overlap between the domains, and we could also talk about impact of systemic health on speech. In this section the focus is on sensing targets that relate to the health of organs such as lungs, heart, kidneys, or tissues forming the vocal tract.

This section is devoted to information in vocal biomarkers regarding physiological health. The first three sections are related to breathing covering respiratory health (Section 3.1), breathing behavior (Section 3.2), and spirometry (Section 3.3). The next sections cover conditions that are regularly encountered: cough (Section 3.4), heart health (Section 3.5), blood sugar level (Section 3.6), Dehydration (Section 3.7) and oncology (Section 3.8).

#### 3.1 Sensing of respiratory health

Speech and respiration are closely related in the sense that speech is produced by organs evolved for the respiratory function of the body. It implies that speech production process interferes with the standard way of regular breathing. Evolution mitigated this issue by an unconscious planning process for respiration being in place while speaking [74–76]. When the respiratory condition is impaired, it means that the normal respiratory planning is shortened or fails to operate resulting in frequent breaks during an utterance because of the need of air. Measuring the respiratory



phases during speech is possible with a respiration belt or an airflow sensor. However, such approach is in almost all cases rather awkward. The Virtual Respiratory Belt (VRB) [49] offers a solution here as these phases can be identified from the speech and an estimate of the belt signal can be obtained.

### 3.2 Breathing behavior

The details of the breathing pattern during speech may be indicative of various physiological health conditions of the talker. The lung diseases are often categorized as obstructive and restrictive diseases. In obstructive lung diseases, patient has difficulties in exhaling the air from the lungs. The most common causes of obstructive lung disease are:

- Chronic obstructive pulmonary disease (COPD), which includes emphysema and chronic bronchitis
- Asthma
- Bronchiectasis • Cystic fibrosis

In restrictive lung diseases where the patient has difficulties in filling lungs with air some of the typical conditions are:

- Interstitial lung disease, such as idiopathic pulmonary fibrosis
- Sarcoidosis, an autoimmune disease
- Obesity, including obesity hypoventilation syndrome
- Scoliosis
- Neuromuscular disease, such as muscular dystrophy or amyotrophic lateral sclerosis (ALS)

Analysis of the breathing pattern and looking for deviant patterns in the inhalation/exhalation phase of the breathing pattern could help to diagnose the nature of the lung disease. The information from the estimated breathing pattern during speech may also be used in different ways. For example, our recent paper proposed a smart control of a portable oxygen concentration device based on the forecasting of the next inspiration event from speech [15]. Respiratory behavior reflects the speech planning [74,77] and consequently, the observed respiratory pattern will demonstrate the planned structural organization of speech content. For example, an inspiration event is often placed at the end of the sentence and, therefore, the detection of an inspiration event indicates the intended end of the sentence of the talker. This helps to resolve ambiguities in sentence parsing in automatic speech recognition and detect parts of audio where the subject is not talking for background noise estimation.

### 3.3 Spirometry

Spirometry is considered the gold standard for evaluating pulmonary function for diagnosis of obstructive and restrictive respiratory disorders. A typical spirometer consists of predictive equations which are calibrated based on patient's age, gender, ethnicity, height, and weight. The test involves a deep maximal inhale by the patient and a forceful and rapid exhalation through a mouthpiece. The spirometer measures the volume flow of exhaled breath, and estimates parameters like Forced Expiratory Volume in 1 second (FEV1), Forced Vital Capacity (FVC), FEV1/FVC, Peak Expiratory Flow (PEF), Flow-Volume Curve, Pressure-Volume Curve and Work of Breathing. These parameters are used to assess pulmonary health of the patient and diagnose any respiratory disorders, e.g., based on the shape of the flow-volume curve.

In scenarios where spirometry outcomes are uncertain, broncho-provocative tests are used to confirm diagnosis. E.g., methacholine, which causes narrowing of airways in lungs, is used for diagnosis of asthma along with spirometry [78]. As an alternative to medical-grade spirometers which are expensive and have limited availability, portable spirometers are offered by several vendors (GoSpiro, SpiroLab, SpiroMax, etc.). These are shown to be as accurate as medical-grade spirometers, e.g., in diagnosis of COPD [79]. Audio- or speech-based spirometry has been attempted in the past. Samsung Research in US in collaboration with Brigham and Women's Hospital (BWH) have done clinical studies to collect data from healthy subjects and patients with respiratory disorder (COPD/Asthma) to build audio-based models to estimate spirometry parameters [80,81]. The team collected data on 211 subjects from 3 studies, where participants were asked to loudly read the 'rainbow' paragraph, and the audio was recorded using the Samsung Galaxy Note 8 smartphone mic. Besides the read speech data, data on exhalation sounds while performing the spirometry protocol, voluntary cough sounds and vowel sounds (Aa, ...) were recorded. One of their studies with BWH involved giving several doses of Methacholine to participants and recording their spirometry and audio data

after each dose. They have shown that the voice loudness, pitch and pauses are correlated to the flow-volume curves obtained from spirometry, and can be used to estimate parameters like FEV1, FVC, etc. They show that use of exhalation sounds, while following the spirometry protocol, results in better accuracy in estimating the spirometry parameters from audio. They also quantify the quality of the exhalation effort of the user, which is informed by the American Thoracic Society guidelines.

Similar work has been done by researchers at University of Washington showing forced exhalation sounds from multiple microphones with compressed or uncompressed audio data can be used to estimate spirometry parameters with




Figure 4: Flow-Volume curve from spirometry with inhalation (bottom) and exhalation (top) curves. Morphology of the curve indicate type of respiratory disorder.

reasonable accuracy [82,83]. They also propose use of a 3D printed vortex whistle (for which pitch is proportional to airflow), specifically for patients with low lung function. A solution based on whistle sound is shown to reduce false negatives in determining lung function and is more accurate in estimating flow-volume curves. It is also shown that the loss in mean performance from a call-in service (over a GSM network), compared to locally available uncompressed audio data is only around 1% while estimating some of the spirometry parameters from audio.

### 3.4 Cough screening

Cough sound acquisition comes in two major variants. One is detection or monitoring of cough and is a longitudinal measurement. The other one is information extraction from cough and is a spot check. Obviously, these can be combined in a two-step approach.

Cough sounds have been studied in terms of its character (wet/dry, productive/non-productive, voluntary/spontaneous), its source (male/female), health status (healthy/unhealthy), lung condition (obstructive/restrictive/normal), and disease diagnosis (croup, COPD, asthma, pneumonia, pertussis, tuberculosis, bronchitis, COVID-19, cold, heart failure, lower/upper respiratory tract disease). Various devices can be used, like wearables (acceleration sensor where various areas of the body have been used), dedicated tube systems, or microphone recordings. It is generally accepted that cough is characterized by large inter-person and intra-person

variability. This may be the reason that reported performance varies quite a lot. ResApp Health reported quite successful performance for disease differentiation [84]); results reported by other parties are less convincing [85].

Monitoring can be done by mobile or stationary systems, on-body accelerometers or microphones. There is a clear desire from clinicians for insight into (amount of) coughing for chronic coughers. Also pharmaceutical companies need cough monitors for testing/confirming effects of, e.g., antitussive medication. From the perspective of a chronic patient, unobtrusiveness is desired for obvious reasons; for the caregiver this is also a desired property as it greatly increases adherence. Promises here are early warning systems (leading to less/shorter hospital admissions), medication control (leading to better medication usage) and early discharge by deploying convalescence monitoring. Hye proposed, among others, TB [86] and COVID [87] as application areas based on results of cough frequency at a cohort level. Translation to a single-patient system in real-life situations (patient homes), covering aspects of proper metrics, personalization, cough prevalence, medical outcomes (e.g., exacerbation) and early warnings (lead time) are still under development. A first instance of an unobtrusive night-time cough monitor where such aspects were considered was reported in [45,88–90] using a set-up allowing personalization without privacy intrusion.

### 3.5 Heart health sensing from vocalizations

Beyond Verbal, currently a part of Vocalis health, has worked with the Mayo clinic on detection of certain heart conditions from speech signal. With a cohort of 101 patients, Maor et al. [91] were able to predict the mortality of CHF patients from a 20s speech recording. Certain heart diseases may also lead to a rare condition of cardio-vocal syndrome, an audible hoarseness [92] the voice due to a structural change in the arteries that impinges on the left recurrent laryngeal nerve controlling the muscles in the larynx [32]. Cardiokol has technology and an app for sensing arrhythmias from sustained vowels. In their 2021 paper [93] the accuracy in the detection of arterial fibrillation, AF, from a sustained “Ahh” vowel in a group of 143 patients was reported to be 95%. Gouda et al. [94] have proposed speech sensing and voice assistants as useful tools for diagnostic and health lifestyle management of cardiac patients.

### 3.6 Blood sugar level

The sensing of blood glucose level is very important especially for Type I diabetics and the possibility of using speech has been seen as a very attractive way to monitor glucose level unobtrusively. Some studies, see [14], report 79% accuracy in the detection of different levels of blood glucose concentration from short speech utterances collected using a smartphone and a continuous glucose monitoring device. In a meta study [95] it was concluded that the detection seems to be possible at an individual level but the interpretation of the results is often problematic, for example, because the subjects are aware of their blood glucose level. It is usually thought that there are two mechanisms in blood glucose level that may influence speech:

- Chance in the blood flowing in the larynx and the glottis causes changes in the elastic properties of the organs which would change the glottis in voiced speech.
- The mood changes related to hypoglycaemia, feeling of anxiety, and hyperglycaemia, which may cause arousal, respectively, are reflected in changes in the prosody and content of speech.

### 3.7 Dehydration and fluid accumulation

The body fluid balance may have various effects on speech. It is known that speech itself causes dehydration of vocal cords which may cause hoarseness in the voice after a long speech [96]. Dehydration increases the respiratory rate. Systemic dehydration also influences the speech in a similar way [97]. The dryness of the tongue, lips, and other tissues of the vocal tract may result from systemic dehydration of the body. The current authors are not aware of literature on using speech audio for for the detection or quantification of the level of dehydration. However, it has been shown, for example, that drug-induced dehydration increases the Phonation Threshold Pressure, PTP, that is, the required subglottal pressure needed to start vocal utterances [98]. Indirect estimation of PTP from speech would be potentially an interesting detection target, for example, in the assessment of adverse medication side-effects [99].

### 3.8 Oncology

Lung cancer is the type of cancer that is largest in terms of burden and deaths globally and is typically diagnosed with a CT scan. It is known that in certain populations, COPD (detected through spirometry) can be a precursor for lung

cancer, and its sub-typing can be used for effective lung cancer surveillance [100]. Also, respiration patterns show changes in patients undergoing radiotherapy [101] and chemotherapy [102]. Some of these lung functions can be ascertained with high accuracy using spirometry. Vocal biomarker technology can be integrated in any speech interface, for example, in conversational health counseling services for cancer patients [103].

As described in Section 3.3, virtual spirometry can be performed using audio or speech data with good accuracy. Audio/speech data from different protocols (paragraph read, exhalation, vowel and cough sounds), patient info (age, gender, height, weight) and other vitals (HR, SpO<sub>2</sub>, HRV) can be used as inputs to AI algorithms to predict RIP and spirometry data. The direct estimation of spirometry parameters using speech can potentially offer a safer (radiationfree), accessible and cheaper option to detect and stage lung cancer patients. Also, patients with similar audio signatures can be matched to clinical trials conducted for lung cancer treatment. The tracking of changes in audio can enable monitoring and clinical surveillance of patients undergoing radiotherapy or chemotherapy.

The cancers in the area of the vocal tract, including trachea, larynx, and mouth, have effect on voice of the talker. Halpern et al. [104] reported 88% accuracy in the detection of the speech of an oral cancer patient from a healthy control.

Brain cancers may lead to changes in cognitive and motor functions that may effect the voice and speech content.

## 4 Cognitive and mental health conditions

This section is devoted to information in vocal biomarkers regarding cognitive and mental health. The first sections are related to three diseases: Alzheimer (Section 4.1), Parkinson (Section 4.2), and autism (Section 4.3). Section 4.4 considers emotion and sentiments. The following sections address the states of depression (Section 4.5), intoxication (Section 4.6) and fatigue and sleepiness (Section 4.7).

### 4.1 Alzheimer's disease

The detection of AD from speech was already discussed in Chapter 2. According to WHO, about 55M people globally suffer from dementia and about 60-70% of them likely have AD. The expected global market size in 2029 is \$6.3B and a large part of it is in medication, although, there is no drug to cure the condition. Many patients are undiagnosed, and the progression of their disease could be potentially slowed down by nutrition, lifestyle changes, and medications. Therefore, early detection is seen as particularly important.

### 4.2 Parkinson's disease

The expected market size of Parkinson's disease, PD, is also increasing rapidly and expected to reach globally \$6.7B by 2030. About 70% of the market is medication costs and the rest, various therapies and care settings. Medications are known to be helpful in reducing symptoms and slowing down the progression of the disease, but the common medications have significant side effects. Also in PD, for the same reason as in AD, early detection, and various therapies in the early phase of the disease would significantly reduce the costs and improve the quality of life of the patients.

### 4.3 Autism spectrum disorder

Autism is frequently accompanied by communication difficulties, including delayed speech development, language comprehension difficulties, and social communication challenges. The market size of early detection, and treatment is expected to reach \$5B market size in 2030. Diagnostic evaluations commonly deploy speech assessments that measure speech production (e.g., articulation, fluency), language comprehension, and social communication skills (e.g., initiating and maintaining conversation, interpreting nonverbal cues). Assessments of pragmatic language, which capture the use of language in social contexts, can further assist in identifying autism. Given their efficacy in detecting autism, speech and language assessments constitute valuable tools in the diagnostic process. Several studies have reported that children with autism exhibit distinctive breathing patterns, which can include slower or shallower breathing or differences in respiratory rate variability [105]. These breathing pattern deviations may be linked to other autism symptoms, such as sensory processing difficulties or anxiety. Specifically, children with autism have been observed to have lower respiratory sinus arrhythmia (RSA) values, indicating atypical regulation of the

autonomic nervous system. Furthermore, compared to typically developing children, children with autism exhibit reduced respiratory variability, which can imply a decreased flexibility in respiratory control. Additionally, children with autism may demonstrate lower respiratory sinus arrhythmia and a flatter breathing pattern, indicating difficulties in emotional regulation and social engagement [106,107]. Taken together, these findings suggest that breathing pattern assessments may be useful in identifying and characterizing autism. McDaniel et al. [108] give a recent overview of measuring preverbal vocalization in infants with Autism Spectrum Disorder (ASD). Preverbal means here the phase in development where the child has no, or few, words. Some of the typical differences between typical development and children with ASD are related to the lower rate of intentional vocal communication, smaller consonant inventories, atypical vocalizations, and decreased responsivity and attention to the vocalizations of others. The detection of ASD from child vocalization has been studied using the De-Enigma data set. Typical performance for the detection of an ASD is around 70% [109,110].

#### 4.4 Emotion, state, and sentiment sensing

Emotion sensing from speech is a very popular area of research [53] where most of the applications are related to the sensing of caller emotion and arousal in call centres and public spaces, for example. The detection of various mental diseases has been studied a lot but the performance is usually low. For example, a recent paper using speech data for the detection of various psychiatric conditions [54] reported a low  $R^2$  value, i.e., in less than a half of the cases the condition was detectable by the speech data. The problems stem already from vague definitions of conditions such as schizophrenia, depression or general anxiety, and their volatility.

#### 4.5 Depression

Depression is globally a large problem that causes high healthcare costs for the societies. The global anxiety disorder and depression market was estimated to be around \$12B in 2019 and growing to reach \$16B by 2027. Approximately one half of the market is in medication and the rest for various therapies. Speech-based sensing can be used for triage and monitoring of the changes in the condition of the patient, for example, in response to medication. Speech-based interfaces can also be used to deliver therapeutic interventions to the patients. Dyspnea can be associated with anxiety and panic, which are often comorbid with depression. Chronic illnesses such as heart failure or chronic obstructive pulmonary disease (COPD) can lead to both dyspnea and depression due to their impact on physical health and quality of life. People experiencing depression may have changes in their physical health, including fatigue and altered respiratory patterns, which can contribute to a feeling of dyspnea.

#### 4.6 Intoxication

Intoxicated speech detection challenge was organized as a part of the Interspeech conference in 2011. The winning team received a 73% accuracy in detection of a sober or alcohol intoxicated ( $> 0.5\text{mg/L}$ ) speaker from a single speech utterance.

#### 4.7 Fatigue and sleepiness

It is generally assumed that fatigue and sleepiness affect the character of the voice. For example, it has been demonstrated that the sleepiness level of a subject can be detected from a single sustained vowel [111] with reasonable accuracy.

## 5 Applications

An overview of the various applications that were found in the survey are structured in 13 sections running well-known tests to applications in respiratory support and speech therapies.

### 5.1 Active monitoring or protocolled speech-based health tests

There is a wide array of speech-based health tests that are regularly used in clinical practise. Many of the tests that are currently performed in a clinic can be potentially automated and turned to a home test which can be expected

to improve the patient experience, health outcomes, and cost of care. Automation of routine testing also saves time of the health care professionals.

## 5.2 Diadochokinetic speech tests

Diadochokinetic (DDK) tests are regularly used to assess neurological speech motor impairments [112]. There are two common variants of the test: alternating motion rate (AMR) test and sequential motion rate (SMR) task. In AMR the subjects repeat as quickly as possible sequence of identical syllables, e.g., “pa-pa-pa-..” or “ta-ta-ta-...” and in SMR variant sequences of syllables, that is, “pa-ta-ka-pa-ta-...”. Conventionally the recordings are analyzed manually but automated techniques based on deep learning have been recently demonstrated [113].

## 5.3 Cognitive retrieval tests

Boston Naming Test (BNT) is a widely used test for children and adults with aphasia [114]. In the test, the patient is requested name entities shown in 60 drawings which are ordered by difficulty. The examiner scores each answer based on the responses of the patient. BNT test has been criticized for being culturally biased and lacking in accuracy [115] and proposed to be replaced with a more comprehensive test such as the Neuropsychological Assessment Battery, NAB [116]. NAB has 33 different tests for cognitive capability, learning, and memory. Montreal Cognitive Assessment, MoCA [117], is a tool for assessing mild cognitive impairment and contains a memory recall test and speech fluency tests. In typical speech fluency tests the patient is asked to list, as fast as possible, words starting with a particular letter, or names of entities from a category, e.g., names of animals. Fluency tests have been validated in many studies, see, e.g., [118]. Controlled Oral Word Association Test, COWAT [119] is a variant of the verbal fluency test where the subject lists in one-minute words starting with one letter and then repeats the task with two other letters.

## 5.4 Tests on vocal development in children

There are several tests that help to assess child development. Early Communication Indicator, ECI [120], is a test for toddlers where the parent is playing with a particular toy with the child and communicating with the child. The score is based on count of various kinds of vocalizations and gestures in a six-minute test session. Autism Diagnostic Interview (ADI-R) is a structured interview with a practitioner and a parent of a child with 93 questions for helping to distinguish ASD from other developmental disorders [121]. The interview and scoring of the results require extensive training and the interview takes approximately two hours.

## 5.5 Passive Monitoring

The main use case considered in this report can be called passive monitoring of the health condition of a patient. The sensing can be performed during any calls with the subscriber and use the natural speech content of the call. It is also possible to set up calls for speech monitoring and have protocolled conversations with the patient, which could be considered as an example of active monitoring. Active speech-based sensing is common practice in many brain diseases and forms also the basis for the diagnosing of, for example, aphasias.

Typically, the results of passive speech monitoring could be given to the caregivers in a monitoring dashboard. For example, this information may show some trends in the development of the cognitive state, communication capabilities, or respiratory functions of the patient. The caregiver can use the information to plan the care of the patient. The derived health information may also be used as features for risk prediction.

## 5.6 Assisting technologies for call centers

The information conveyed in speech may relate directly to the health of a person. For example, if a person says that she has fallen or has chest pain, this information may be extracted from the speech using Automatic Speech Recognition (ASR) and Natural Language Understanding (NLU) technologies. A typical task in NLU would be to detect that a patient reports a certain health condition, for example, “I have breathing difficulties.” In using vocal biomarkers, the respiratory health condition can be detected directly from the way the person is breathing or talking during the conversation. In the final application, of course, both NLU and speech-based sensing can be used. NLU technology and speech analytics, in general, can be for example used for automatic generation of case notes from calls.

Automatic note generation may reduce the time of a call significantly. The optimization of the average speed of answer and handling times in call centers, or contact centers [122], by using computational techniques is an active area of research, see, e.g., [123].

In the recent years, we have seen significant developments in ASR performance [124] which has led to the introduction of spoken dialogue systems for home use. Smart assistants such as Amazon Alexa, Google Home, and various others already offer various home health services that help to monitor and find solutions to health problems.

### 5.7 Automation of call center

The development of assisting technology for better logging of the call outcomes can be seen as the first step towards partial automation of the call center services [125] with the help of automated agents. In the first phase, the automation can be introduced as an assistive technology that would give the PRA reminders on what to ask and how to run the conversation the most effectively way the given customer. A large number of PERS calls are not emergency calls but relatively simple technical questions, for example, related to changing of batteries. Those calls could be gradually replaced by automated call agents. The use of automated conversational agents for outbound calls are also being introduced, for example, this is one of the proposed use cases for Google Duplex.

The development of automated health counseling systems [126] based on conversational interfaces is an area which grows very rapidly. The effectiveness of automated agents in depression counseling has been recently demonstrated in a random controlled trial [127]. The results of chatbot interventions in clinical psychiatry are still inconclusive, see, e.g., [128,129], one can anticipate their use in contact center applications in the future. A recent meta-analysis [130] concluded that chatbots are already effective for dietary, physical activity, and sleep counseling.

### 5.8 Home healthcare and tele-health

Speech-based sensing makes it possible to monitor the state and changes in the patient's condition. Listening to a patient for health monitoring purposes is nothing new in health care. In many home healthcare applications and services, the communication between the caregivers and the patient takes place over a telephone or video conference link. The caregiver gets information about the patient's health status from the patient's responses and from various non-verbal cues. In particular, the cognitive and mental state or, for example, respiratory conditions of a patient may lead to changes in spoken content, intonation, voice, and non-verbal conversational behavior [131]. When the caregiver knows the patient well, the monitoring of speech and non-verbal cues in conversation is a natural and instinctive part of the care process and does not necessarily require a technological solution based on speech analytics. In a call center where each respondent may have thousands of patients, this is more difficult, and in a case where the interaction takes place with a spoken dialog system, the sensing can only be performed using speech technology.

### 5.9 Elderly care and chronic disease management

The elderly care market size in the US alone is estimated to be around \$400B and expected to grow globally to \$2.4T by 2028. The market consists of medications, devices, and services. Vocal Biomarker technology has many potential use cases in elderly care. A large part of the market is in elderly care facilities where passive and active speech sensing, discussed in this report, can be integrated into the infrastructure and care services for the inhabitants. In addition to screening and longitudinal tracking health conditions of individual patients, passive sensing of vocalizations, for example coughing and sneezing, can be used for detecting and monitoring respiratory infections spreading in the facility.

In chronic disease management the patient education, screenings, check-ups, monitoring, and coordination of care are expected to have a market size of \$10B by 2028. This may consist of telehealth services, speech assistants in home, phones, and in-car applications where the availability of speech content opens possibilities for active and passive conversational health sensing.

Medication has various effects on speech and therefore it is possible to use vocal biomarkers to track the effectiveness of a medication to the health of the patient and detect possible medication side-effects such as fatigue or cognitive and mental effects.

In tele-health services and Personal Emergency Response Systems (PERS) for home dwelling frail elderly speech information is naturally available for speech-based sensing applications. In our analysis of the former Philips Lifeline

data it was found that a large number of PERS patients with a pendant and a speakerphone were very regularly, daily or even multiple times a day, in contact with the call center.

### 5.10 Cardiac health

The respiratory belt signal measures the expansion and contraction of the chest as a result of breathing. The ECG signal, on the other hand, measures the electrical activity of the heart. The ECG signal provides information about the rhythm and function of the heart, while the respiratory belt signal provides information about the rate and depth of breathing. The relationship between these two signals is that changes in the respiratory rate and depth can affect the heart rate and rhythm. For example, when we inhale, the heart rate typically slows down, and when we exhale, the heart rate typically speeds up. This phenomenon is known as respiratory sinus arrhythmia (RSA) and is a normal physiological response. By analysing how changes in respiratory activity affect the heart rate and rhythm, we can gain a better understanding of how the cardiovascular system is functioning. This information can be used in the diagnosis and treatment of various cardiovascular conditions.

### 5.11 Respiratory support

Many lung disease patients need additional oxygen support which is often based on an oxygen concentrator device which brings oxygen-rich air to the nose of the patient via a nasal cannula [132]. The POCs typically work either in a continuous flow mode or in pulse dose mode where the air flow pulse, bolus, is delivered during the inhale phase triggered by a pressure sensor. While the cannula allows the patient to talk freely, the flow of oxygen during speech is disturbed in both POC modes. In non-speech breathing during light activity the average inspiration and expiration duration in COPD patients are 1s and 1.7s, respectively [133]. During speech, the average respiratory rate is approximately half of the non-speech breathing and the duration of inhaling is significantly reduced [49,77,134]. In a continuous flow model only a fraction of oxygen gets into lungs during normal speech. In pulse dose mode, the latency of the air delivery in most devices is 100-200ms and the duration of the bolus is 200-800ms [135]. Moreover, the pulse dose mode is activated by nasal inhalation while it is common during speech to inhale through mouth, which would not trigger the pressure sensor. In [136] it was demonstrated that it is possible to forecast the inhalation event and therefore control the oxygen administration such that it is aligned with the brief inspiration events in continuous speech.

The forecasting of inhale moments from speech can be useful in various applications including open types of air masks for scuba divers, firemen, or astronauts, who need to talk naturally while wearing the mask for oxygen support, and for breathing support devices for lung patients. The professional safety mask market is expected to reach \$7.6B by 2026,




Figure 5: Vocal biomarker may represent a specific root factor, but often it is a secondary measure approximating a more intrusive primary measurement of the underlying root factor.



medical mask market \$35B in 2030, and scuba diving equipment market size was already \$1.95B in 2019. Moreover, it has very interesting opportunities in enabling more natural communication in various Metaverse applications, which has an expected value of tens of trillions of dollars [137].

### 5.12 Speech Therapies

Speech sensing can be used to support speech therapy techniques. Speech Entrainment for Aphasia Recovery (SpARc) is a therapeutic technique used in stroke rehabilitation of patients with speech disorders [138]. In SpARc the patient is asked to mimic the speech of talker presented by video or audio. The performance of the subject can be assessed by the percentage of correctly repeated words and by the change in various aphasia tests, such as the Western Aphasia Battery [139] or the Boston Naming Test [114].

### 5.13 Automotive applications

The size of driver health monitoring market size is around \$1B and it is expected to rise to \$2.4B by 2027, in some estimates reaching globally \$25B by 2030 [140]. The main driver is the fact that many accidents occur due to fatigue and sudden incidents such as the cardiac arrest of the driver, for overview, see, . In an automotive environment, speechbased sensing can be used to sense various driver and passenger conditions. Intoxication and fatigue are useful in all scenarios. Emotions such as anger or arousal, for example, can be a part of safety monitoring in public transportation systems. The sensing of driver’s health may use various health sensing techniques discussed in this report. Speechbased sensing has significant benefits over several other sensing modalities. Sensors attached to the body are difficult to use in a car environment, and inertial sensors mounted into the structure of the car suffer from a high level of vibrations. Cameras and IR systems suffer from continuously changing lighting conditions.

## 6 Performance of vocal biomarkers

In this review the goal was to give a broad overview of a range of vocal biomarkers proposed in the literature. In the Section 1 we defined a vocal biomarker as a method to detect a condition of the talker, or estimate the measurement values of another sensory modality. The detection of a specific characteristic of a vocalization, such as a cough or the roughness of voice, is subject to acoustic measurement error. The detection of cough is a *primary measurement* of the condition for the root factor influencing of the patient. As illustrated in Fig. 5 there is often another primary measure for the underlying root factor, for example, a respiratory rate of a patient, or the Hamilton Depression Rating Scale (HDRS) for the diagnosis of clinical depression [141]. The primary measure has some error which may be caused by sensor noise or ambiguities in the rating protocol. A vocal biomarker is typically developed and validated so that it represents a translation of the primary measure. The translation is subject to a translation error. Obviously, the acoustic measurement of the biomarker is influenced by noise and distortion.

Class	Root factor	Primary measure	Meas. error	Trans. Error	Acoust. Error
Cardiopulm.	Respiratory rate	RIP/Air flow	Low [142]	Low	Low
Cardiopulm.	Dyspnea	Spirometry	Medium	Medium	Low
Cardiopulm.	COPD/asthma	Clinician	Medium	Medium	Low
Cardiopulm.	Breathing sounds	Stethoscope	Medium	Medium	High
Cardiopulm.	Cough	Microphone	N/A	N/A	Medium
Cardiopulm.	Arrythmias	ECG	Low	High	High
Cardiopulm.	cardio-vocal syndrome	Imaging study	Medium	High	High
Infection	Flu/cold	WURSS	Low [37]	Low	Medium
Infection	Covid-19	PCR test	Low [143]	Medium	Medium
Mental	Depression	HDRS	Medium [141]	Medium	Low
Mental	Schizotype	DSM-5	High	High	Low
Cognitive	Aphasia	WAB	Medium [144]	Medium	Low

Survey on biomarkers in human vocalizations

Cognitive	Parkinson’s Disease	UPDRS	Medium [28]	Medium	Low
Cognitive	Dysarthia	DSD	Medium [112]	Medium	Low
Cognitive	Tourette’s syndrome	Clinician	Medium [145]	High	Low
Cognitive	Dementia	CDR	Medium	High	Low
Cognitive	ALS	EMG, US	Low	High	Low
Cognitive	Autism Spectrum Disorder	ASQ	Medium	High	Low
Systemic	Intoxication	Blood: alcohol/drug	Low	Medium	Low
Systemic	Blood Glucose Level	Blood: sugar level	Low	High	High
Systemic	Systemic dehydration	Blood: serum osmolality	Low	High	High
Systemic	Fatigue/sleepiness	Self-reporting	Medium	Medium	Low
Oncology	Oral cancer	Imaging, pathology	low	High	Medium

Table 4: Characterization of noise sources in vocal biomarkers

Table 4 gives a qualitative overview of different targets for vocal markers and the impact of the measurement noise, or lack of the reliability of the primary measure, the anticipated translation noise from the primary measure to the vocal biomarker, and the likelihood for the impact of acoustic error caused by the expected low level of the acoustic signature. In most cases the acoustic error can be expected low because the condition influences the content and prosody of speech. The translation error is often ranked medium to high because the acoustic measure is a very indirect indicator of the primary measure. Finally, the measurement error of the primary measure is in some cases a direct indication of the condition, for example, respiratory rate and the RIP belt measurement, where we can expect that the noise influencing the measurement is generally low. In cases where the primary measure is a holistic score, for example, Montreal Cognitive Assessment, MoCA [117] based on subjective assessment of the responses of a patient, the measure error may often be considered medium or high due to low reliability and repeatability.

One could conclude that the vocal biomarkers for the detection targets, or root factors, that have at least one High noise source are probably very challenging to develop into a useful and versatile measure. The easiest cases seem to be basic respiratory measurements, detection of infection, intoxication, and fatigue detection. However, it should be noted that the progress in the area of vocal biomarkers is fast and we can expect to see significant progress in the coming years in the development of new algorithms and primary measure targets.

## 7 Conclusions and discussion

This paper aims at giving a broad and updated survey on the recent progress in vocal biomarker technology and its applications. Over the past few years, the rapid advancement of machine learning techniques has expanded the horizons of potential applications and significantly enhanced the performance in conventional problem domains. However, it is important to note that the performance of vocal biomarkers still falls short of the standards required for their utilization in healthcare applications. A basic qualitative overview of noise sources determining the accuracy has been presented in order to identify promising directions.

Two challenges in this research area are particularly worth mentioning here. One challenge is that a vocal biomarker, due to training data, typically serve as secondary approximations of primary measures. In many instances, the machine learning models for vocal biomarkers are trained using the primary measure rather than the ground truth. This approach introduces measurement errors into the primary measure, complicates the interpretation of vocal biomarkers in relation to the primary measure, and is susceptible to acoustic noise and distortion, which can adversely affect the performance of vocal biomarkers.

Another challenge relates to the use of vocal biomarker technology in a practical application. Various biomarkers (e.g., protocolled speech, features from sustained vowels) require a triage set-up realizable as apps or websites that collect recordings and may require trending over longer periods of time. As such it requires (prolonged) adherence from the user or patient. **References**

- [1] D. L. Everett, *How Language Began: The Story of Humanity's Greatest Invention*. Liveright Publishing, Nov. 2017, google-Books-ID: RtpDDQAAQBAJ.
- [2] M. Garcia and L. Favaro, "Animal vocal communication: function, structures, and production mechanisms," *Current Zoology*, vol. 63, no. 4, pp. 417–419, Aug. 2017, publisher: Oxford Academic. [Online]. Available: <https://dx.doi.org/10.1093/cz/zox040>
- [3] G. Fagherazzi, A. Fischer, M. Ismael, and V. Despotovic, "Voice for Health: The Use of Vocal Biomarkers from Research to Clinical Practice," *Digital Biomarkers*, vol. 5, no. 1, pp. 78–88, 2021.
- [4] M. Ferro and R. Farmanfarmaian, *How AI Can Democratize Healthcare*. MV, Sep. 2022.
- [5] V. B. Kraus, "Biomarkers as drug development tools: discovery, validation, qualification and use," *Nature Reviews Rheumatology*, vol. 14, no. 6, pp. 354–362, Jun. 2018.
- [6] R. Laennec, "De l'auscultation mediate; de l'exploration de la poitrine : Laennec, R. T. H. (Rene' Theophile Hyacinthe), 1781-1826 : Free Download, Borrow, and Streaming," 1893. [Online]. Available: <https://archive.org/details/delauscultationm00laen>
- [7] P. Kligfield, "Laennec and the discovery of mediate auscultation," *The American Journal of Medicine*, vol. 70, no. 2, pp. 275–278, Feb. 1981. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S002934381907622>
- [8] R. Eklund, "Pulmonic ingressive phonation: Diachronic and synchronic characteristics, distribution and function in animal and human sound production and in human speech," *Journal of the International Phonetic Association*, vol. 38, no. 3, pp. 235–324, Dec. 2008, publisher: Cambridge University Press.
- [9] G. N. Jayabhavani and N. R. Raajan, "Silent speech generation using Brain Machine Interface," in *2013 IEEE International Conference ON Emerging Trends in Computing, Communication and Nanotechnology (ICECCN)*, Mar. 2013, pp. 630–633.
- [10] D. S. Brungart, B. Walden, M. Cord, S. Phatak, S. M. Theodoroff, S. Griest, and K. W. Grant, "Development and validation of the Speech Reception in Noise (SPRINT) Test," *Hearing Research*, vol. 349, pp. 90–97, Jun. 2017.
- [11] J. Sidnell and T. Stivers, Eds., *The Handbook of Conversation Analysis*, 1st ed. Chichester: Wiley-Blackwell, Oct. 2014.
- [12] Y. Pan, V. S. Nallanthighal, D. Blackburn, H. Christensen, and A. Harma, "Multi-Task Estimation of Age and Cognitive Decline from Speech," in *ICASSP 2021 - 2021 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*. Toronto, ON, Canada: IEEE, Jun. 2021, pp. 7258–7262. [Online]. Available: <https://ieeexplore.ieee.org/document/9414642/>
- [13] S. Lee, J. Park, and D. Um, "Speech Characteristics as Indicators of Personality Traits," *Applied Sciences*, vol. 11, no. 18, p. 8776, Jan. 2021, number: 18 Publisher: Multidisciplinary Digital Publishing Institute. [Online]. Available: <https://www.mdpi.com/2076-3417/11/18/8776>
- [14] J. Jeon, A. Palanica, S. Sarabadani, M. Lieberman, and Y. Fossat, "Biomarker potential of real-world voice signals to predict abnormal blood glucose levels," *bioRxiv*, p. 2020.09.25.314096, Sep. 2020, publisher: Cold Spring Harbor Laboratory Section: New Results. [Online]. Available: <https://www.biorxiv.org/content/10.1101/2020.09.25.314096v1>
- [15] A. Harma, U. Grossekofer, O. Ouweltjes, and S. Nallanthighal, "Forecasting of breathing events from speech for respiratory support," in *IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, Rhodes, Greece, Jun. 2023.
- [16] A. Abbe, C. Grouin, P. Zweigenbaum, and B. Falissard, "Text mining applications in psychiatry: a systematic literature review," *International Journal of Methods in Psychiatric Research*, vol. 25, no. 2, pp. 86–100, 2016.
- [17] L. A. Gottschalk, M. K. Stein, and D. H. Shapiro, "The application of computerized content analysis of speech to the diagnostic process in a psychiatric outpatient clinic," *Journal of Clinical Psychology*, vol. 53, no. 5, pp. 427–441, Aug. 1997.
- [18] J. Correia, I. Trancoso, and B. Raj, "Detecting Psychological Distress in Adults Through Transcriptions of Clinical Interviews," in *Advances in Speech and Language Technologies for Iberian Languages*, ser. Lecture Notes in

- Computer Science. Springer, Cham, Nov. 2016, pp. 162–171. [Online]. Available: [https://link.springer.com/chapter/10.1007/978-3-319-49169-1\\_16](https://link.springer.com/chapter/10.1007/978-3-319-49169-1_16)
- [19] P. Lopez-Otero, L. Docio-Fernandez, A. Abad, and C. Garcia-Mateo, “Depression Detection Using Automatic Transcriptions of De-Identified Speech,” in *Interspeech 2017*. ISCA, Aug. 2017, pp. 3157–3161. [Online]. Available: [https://www.isca-speech.org/archive/interspeech\\_2017/lopezotero17b\\_interspeech.html](https://www.isca-speech.org/archive/interspeech_2017/lopezotero17b_interspeech.html)
- [20] T. B. Brown, B. Mann, N. Ryder, M. Subbiah, J. Kaplan, P. Dhariwal, A. Neelakantan, P. Shyam, G. Sastry, A. Askell, S. Agarwal, A. Herbert-Voss, G. Krueger, T. Henighan, R. Child, A. Ramesh, D. M. Ziegler, J. Wu, C. Winter, C. Hesse, M. Chen, E. Sigler, M. Litwin, S. Gray, B. Chess, J. Clark, C. Berner, S. McCandlish, A. Radford, I. Sutskever, and D. Amodei, “Language Models are Few-Shot Learners,” *arXiv:2005.14165 [cs]*, Jul. 2020, arXiv: 2005.14165. [Online]. Available: <http://arxiv.org/abs/2005.14165>
- [21] R. H. P. Brookshire and M. R. McNeil, *Introduction to Neurogenic Communication Disorders, 8e*, 8th ed. St. Louis, Missouri: Mosby, Oct. 2014.
- [22] K. C. Fraser, G. Hirst, J. A. Meltzer, J. E. Mack, and C. K. Thompson, “Using statistical parsing to detect agrammatic aphasia.” Association for Computational Linguistics, 2014, pp. 134–142. [Online]. Available: <http://aclweb.org/anthology/W14-3420>
- [23] B. Peintner, W. Jarrold, D. Vergyri, C. Richey, M. L. G. Tempini, and J. Ogar, “Learning diagnostic models using speech and language measures,” in *2008 30th Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, Aug. 2008, pp. 4648–4651.
- [24] D. Le, K. Licata, and E. M. Provost, “Automatic Paraphasia Detection from Aphasic Speech: A Preliminary Study.” ISCA, Aug. 2017, pp. 294–298. [Online]. Available: [http://www.isca-speech.org/archive/Interspeech\\_2017/abstracts/0626.html](http://www.isca-speech.org/archive/Interspeech_2017/abstracts/0626.html)
- [25] J. M. Ogar, J. V. Baldo, S. M. Wilson, S. M. Brambati, B. L. Miller, N. F. Dronkers, and M. L. Gorno-Tempini, “Semantic dementia and persisting Wernicke’s aphasia: linguistic and anatomical profiles,” *Brain and Language*, vol. 117, no. 1, pp. 28–33, Apr. 2011.
- [26] K. E. Forbes, A. Venneri, and M. F. Shanks, “Distinct patterns of spontaneous speech deterioration: an early predictor of Alzheimer’s disease,” *Brain and Cognition*, vol. 48, no. 2-3, pp. 356–361, Apr. 2002.
- [27] G. Dimauro, V. D. Nicola, V. Bevilacqua, D. Caivano, and F. Girardi, “Assessment of Speech Intelligibility in Parkinson’s Disease Using a Speech-To-Text System,” *IEEE Access*, vol. 5, pp. 22199–22208, 2017.
- [28] C. G. Goetz, B. C. Tilley, S. R. Shaftman, G. T. Stebbins, S. Fahn, P. Martinez-Martin, W. Poewe, C. Sampaio, M. B. Stern, R. Dodel, B. Dubois, R. Holloway, J. Jankovic, J. Kulisevsky, A. E. Lang, A. Lees, S. Leurgans, P. A. LeWitt, D. Nyenhuis, C. W. Olanow, O. Rascol, A. Schrag, J. A. Teresi, J. J. van Hilten, N. LaPelle, and for the Movement Disorder Society UPDRS Revision Task Force, “Movement Disorder Society-sponsored revision of the Unified Parkinson’s Disease Rating Scale (MDS-UPDRS): Scale presentation and clinimetric testing results,” *Movement Disorders*, vol. 23, no. 15, pp. 2129–2170, Nov. 2008. [Online]. Available: <http://doi.wiley.com/10.1002/mds.22340>
- [29] I. Laaridh, C. Fredouille, and C. Meunier, “Automatic Anomaly Detection for Dysarthria Across Two Speech Styles: Read vs Spontaneous Speech,” May 2016. [Online]. Available: <https://hal.archives-ouvertes.fr/hal-01420956/document>
- [30] B. Kosztyła-Hojna, D. Moskal, and A. Kuryliszyn-Moskal, “Parameters of the assessment of voice quality and clinical manifestation of rheumatoid arthritis,” *Advances in Medical Sciences*, vol. 60, no. 2, pp. 321–328, Sep. 2015.
- [31] J. R. Orozco-Arroyave, F. Honig, J. D. Arias-Londoño, J. F. Vargas-Bonilla, K. Daqrouq, S. Skodda, J. Ruzs, and E. Noth, “Automatic detection of Parkinson’s disease in running speech spoken in three different languages,” *The Journal of the Acoustical Society of America*, vol. 139, no. 1, pp. 481–500, Jan. 2016. [Online]. Available: <https://asa.scitation.org/doi/full/10.1121/1.4939739>
- [32] I. I. Berekat and A. Azzu, “A rare cardiac cause of hoarseness of voice,” *The Libyan Journal of Medicine*, vol. 5, Jan. 2010. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3066782/>

- [33] E. Maor, J. D. Sara, D. M. Orbelo, L. O. Lerman, Y. Levanon, and A. Lerman, "Voice Signal Characteristics Are Independently Associated With Coronary Artery Disease," *Mayo Clinic Proceedings*, Apr. 2018. [Online]. Available: <http://linkinghub.elsevier.com/retrieve/pii/S0025619618300302>
- [34] L. Burd, "Language and Speech in Tourette Syndrome: Phenotype and Phenomenology," *Current Developmental Disorders Reports*, vol. 1, no. 4, pp. 229–235, Dec. 2014. [Online]. Available: <https://doi.org/10.1007/s40474-014-0027-1>
- [35] K. K. Set and J. N. Warner, "Tourette syndrome in children: An update," *Current Problems in Pediatric and Adolescent Health Care*, vol. 51, no. 7, p. 101032, Jul. 2021.
- [36] G. Gosztolya, R. Busa-Fekete, T. Grosz, and L. Tóth, "DNN-Based Feature Extraction and Classifier Combination for Child-Directed Speech, Cough and Snoring Identification." ISCA, Aug. 2017, pp. 3522–3526. [Online]. Available: [http://www.isca-speech.org/archive/Interspeech\\_2017/abstracts/0905.html](http://www.isca-speech.org/archive/Interspeech_2017/abstracts/0905.html)
- [37] Y. Wang, Z. He, S. Chen, Y. Liu, F. Li, B. Barrett, Z. Zhang, G. Su, and C. Stalsby Lundborg, "Validation of the Wisconsin upper respiratory symptom survey-24, Chinese version," *Annals of Medicine*, vol. 54, no. 1, pp. 655–665, Dec. 2022, publisher: Taylor & Francis eprint: <https://doi.org/10.1080/07853890.2022.2043559>. [Online]. Available: <https://doi.org/10.1080/07853890.2022.2043559>
- [38] M. Albes, Z. Ren, B. W. Schuller, and N. Cummins, "Squeeze for Sneeze: Compact Neural Networks for Cold and Flu Recognition," in *Interspeech 2020*. ISCA, Oct. 2020, pp. 4546–4550. [Online]. Available: [http://www.isca-speech.org/archive/Interspeech\\_2020/abstracts/2531.html](http://www.isca-speech.org/archive/Interspeech_2020/abstracts/2531.html)
- [39] S. Takahashi, T. Morimoto, S. Maeda, and N. Tsuruta, "Cough detection in spoken dialogue system for home health care." Jan. 2004.
- [40] R. Pramono, S. Bowyer, and E. Rodriguez-Villegas, "Automatic adventitious respiratory sound analysis: A systematic review," *PLoS ONE*, vol. 12(5), p. e0177926, 2017.
- [41] Y. Shi, H. Liu, Y. Wang, M. Cai, and W. Xu, "Theory and application of audio-based assessment of cough," *Journal of Sensors*, vol. 2018, 2018. [Online]. Available: <https://doi.org/10.1155/2018/9845321>
- [42] A. Serrurier, C. Neuschaefer-Rube, and R. Rohrig, "Past and trends in cough sound acquisition, automatic detection and automatic classification: A comparative review," *Sensors*, vol. 22, no. 8, 2022. [Online]. Available: <https://www.mdpi.com/1424-8220/22/8/2896>
- [43] J. Amoh and K. Odame, "Deep Neural Networks for Identifying Cough Sounds," *IEEE Transactions on Biomedical Circuits and Systems*, vol. 10, no. 5, pp. 1003–1011, Oct. 2016. [Online]. Available: <http://ieeexplore.ieee.org/document/7570164/>
- [44] Hui-Hui Wang, Jia-Ming Liu, Mingyu You, and Guo-Zheng Li, "Audio signals encoding for cough classification using convolutional neural networks: A comparative study." IEEE, Nov. 2015, pp. 442–445. [Online]. Available: <http://ieeexplore.ieee.org/document/7359724/>
- [45] L. D. Perna, G. Spina, S. Thackray-Nocera, M. G. Crooks, A. H. Morice, P. Soda, and A. C. d. Brinker, "An automated and unobtrusive system for cough detection," in *2017 IEEE Life Sciences Conference (LSC)*, Dec. 2017, pp. 190–193.
- [46] A. W. Fox, P. K. Monoson, and C. D. Morgan, "Speech dysfunction of obstructive sleep apnea. A discriminant analysis of its descriptors," *Chest*, vol. 96, no. 3, pp. 589–595, Sep. 1989.
- [47] E. Goldshtein, A. Tarasiuk, and Y. Zigel, "Automatic Detection of Obstructive Sleep Apnea Using Speech Signals," *IEEE Transactions on Biomedical Engineering*, vol. 58, no. 5, pp. 1373–1382, May 2011.
- [48] F. Espinoza-Cuadros, R. Fernandez-Pozo, D. T. Toledano, J. D. Alcázar-Ramírez, E. Lopez-Gonzalo, and L. A. Hernández-Gómez, "Reviewing the connection between speech and obstructive sleep apnea," *BioMedical Engineering Online*, vol. 15, Feb. 2016. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4761156/>
- [49] V. S. Nallanthighal, A. Harmata, and H. Strik, "Deep Sensing of Breathing Signal During Conversational Speech," in *Interspeech 2019*. ISCA, Sep. 2019, pp. 4110–4114. [Online]. Available: [http://www.isca-speech.org/archive/Interspeech\\_2019/abstracts/1796.html](http://www.isca-speech.org/archive/Interspeech_2019/abstracts/1796.html)

- [50] S. Boelders, V. S. Nallanthighal, V. Menkovski, and A. Harm“ a, “Detection of Mild Dyspnea from Pairs of” Speech Recordings,” in *ICASSP 2020 - 2020 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, May 2020, pp. 4102–4106, iSSN: 2379-190X.
- [51] C. H. Sudre, B. Murray, T. Varsavsky, M. S. Graham, R. S. Penfold, R. C. Bowyer, J. C. Pujol, K. Klaser, M. Antonelli, L. S. Canas, E. Molteni, M. Modat, M. Jorge Cardoso, A. May, S. Ganesh, R. Davies, L. H. Nguyen, D. A. Drew, C. M. Astley, A. D. Joshi, J. Merino, N. Tsereteli, T. Fall, M. F. Gomez, E. L. Duncan, C. Menni, F. M. K. Williams, P. W. Franks, A. T. Chan, J. Wolf, S. Ourselin, T. Spector, and C. J. Steves, “Attributes and predictors of long COVID,” *Nature Medicine*, pp. 1–6, Mar. 2021, publisher: Nature Publishing Group. [Online]. Available: <https://www.nature.com/articles/s41591-021-01292-y>
- [52] H. Coppock, A. Gaskell, P. Tzirakis, A. Baird, L. Jones, and B. Schuller, “End-to-end convolutional neural network enables COVID-19 detection from breath and cough audio: a pilot study,” *BMJ Innovations*, vol. 7, no. 2, Apr. 2021, publisher: BMJ Specialist Journals Section: Artificial intelligence. [Online]. Available: <https://innovations.bmj.com/content/7/2/356>
- [53] B. Schuller and A. Batliner, *Computational Paralinguistics: Emotion, Affect and Personality in Speech and Language Processing*, Sep. 2013.
- [54] J. Olah, K. Diederer, T. Gibbs-Dean, M. J. Kempton, R. Dobson, T. Spencer, and N. Cummins, “Online speech assessment of the psychotic spectrum: Exploring the relationship between overlapping acoustic markers of schizotypy, depression and anxiety,” *Schizophrenia Research*, p. S092099642300138X, Apr. 2023. [Online]. Available: <https://linkinghub.elsevier.com/retrieve/pii/S092099642300138X>
- [55] O. Schleusing, P. Renevey, M. Bertschi, S. Dasen, J. M. Koller, and R. Paradiso, “Monitoring physiological and behavioral signals to detect mood changes of bipolar patients,” in *2011 5th International Symposium on Medical Information and Communication Technology*, Mar. 2011, pp. 130–134.
- [56] M. Morales, S. Scherer, and R. Levitan, “A Cross-modal Review of Indicators for Depression Detection Systems,” *Proceedings of the Fourth Workshop on Computational Linguistics and Clinical Psychology — From Linguistic Signal to Clinical Reality*, pp. 1–12, 2017. [Online]. Available: <https://aclanthology.info/papers/W17-3101/w17-3101>
- [57] N. Cummins, S. Scherer, J. Krajewski, S. Schnieder, J. Epps, and T. F. Quatieri, “A review of depression and suicide risk assessment using speech analysis,” *Speech Communication*, vol. 71, pp. 10–49, Jul. 2015. [Online]. Available: <http://www.sciencedirect.com/science/article/pii/S0167639315000369>
- [58] M. Pacula, T. Meltzer, M. Crystal, A. Srivastava, and B. Marx, “Automatic detection of psychological distress indicators and severity assessment in crisis hotline conversations,” in *2014 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, May 2014, pp. 4863–4867.
- [59] D. Galanis, S. Karabetos, M. Koutsombogera, H. Papageorgiou, A. Esposito, and M. T. Riviello, “Classification of emotional speech units in call centre interactions,” in *2013 IEEE 4th International Conference on Cognitive Infocommunications (CogInfoCom)*, Dec. 2013, pp. 403–406.
- [60] E. Gonzalez-Moreira, D. Torres-Boza, H. A. Kairuz, C. Ferrer, M. Garcia-Zamora, F. Espinoza-Cuadros, and L. A. Hernandez-Gomez, “Automatic Prosodic Analysis to Identify Mild Dementia,” *BioMed Research International*, vol. 2015, p. 916356, 2015.
- [61] N. Cummins, Y. Pan, Z. Ren, J. Fritsch, V. Nallanthighal, H. Christensen, D. Blackburn, B. Schuller, M. Magimai-Doss, H. Strik, and A. Harm“ a, “A comparison of acoustic and linguistics methodologies for” Alzheimer’s dementia recognition,” Oct. 2020.
- [62] J. R. Green, D. R. Beukelman, and L. J. Ball, “Algorithmic Estimation of Pauses in Extended Speech Samples of Dysarthric and Typical Speech,” *Journal of medical speech-language pathology*, vol. 12, no. 4, pp. 149–154, Dec. 2004. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2902000/>
- [63] C. Cordella, B. C. Dickerson, M. Quimby, Y. Yunusova, and J. R. Green, “Slowed articulation rate is a sensitive diagnostic marker for identifying non-fluent primary progressive aphasia,” *Aphasiology*, vol. 31, no. 2, pp. 241–260, 2017. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5531197/>
- [64] J. Sidnell and T. Stivers, *The handbook of conversation analysis*. John Wiley & Sons, 2012.

- [65] M. Spath, I. Aichert, A. O. Ceballos-Baumann, E. Wagner-Sonntag, N. Miller, and W. Ziegler, “Entraining with” another person’s speech rhythm: Evidence from healthy speakers and individuals with Parkinson’s disease,” *Clinical Linguistics & Phonetics*, vol. 30, no. 1, pp. 68–85, 2016.
- [66] J. Sommers-Flanagan and R. Sommers-Flanagan, *Clinical Interviewing, with Video Resource Center*, 5th ed. Wiley, Jun. 2015.
- [67] S. Rubak, A. Sandbæk, T. Lauritzen, and B. Christensen, “Motivational interviewing: a systematic review and meta-analysis,” *The British Journal of General Practice*, vol. 55, no. 513, pp. 305–312, Apr. 2005. [Online]. Available: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1463134/>
- [68] S. G. Hofmann, A. Asnaani, I. J. Vonk, A. T. Sawyer, and A. Fang, “The Efficacy of Cognitive Behavioral Therapy: A Review of Meta-analyses,” *Cognitive therapy and research*, vol. 36, no. 5, pp. 427–440, Oct. 2012. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3584580/>
- [69] O. I. Okereke, N. Pantoja-Galicia, M. Copeland, B. T. Hyman, T. Wanggaard, M. S. Albert, R. A. Betensky, and D. Blacker, “The SIST-M: Predictive validity of a brief structured Clinical Dementia Rating interview,” *Alzheimer disease and associated disorders*, vol. 26, no. 3, pp. 225–231, Jul. 2012. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3257375/>
- [70] J. Kay, R. Lesser, and M. Coltheart, “Psycholinguistic assessments of language processing in aphasia (PALPA): An introduction,” *Aphasiology*, vol. 10, no. 2, pp. 159–180, Feb. 1996. [Online]. Available: <https://doi.org/10.1080/02687039608248403>
- [71] S. Scherer, Z. Hammal, Y. Yang, L.-P. Morency, and J. F. Cohn, “Dyadic Behavior Analysis in Depression Severity Assessment Interviews,” *Proceedings of the ... ACM International Conference on Multimodal Interaction . ICMI (Conference)*, vol. 2014, pp. 112–119, Nov. 2014. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5365085/>
- [72] H. Giles, A. Mulac, J. J. Bradac, and P. Johnson, “Speech Accommodation Theory: The First Decade and Beyond,” *Communication Yearbook*, vol. 10, Jan. 1987.
- [73] M. N. Stolar, M. Lech, and N. B. Allen, “Detection of depression in adolescents based on statistical modeling of emotional influences in parent-adolescent conversations,” in *2015 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, Apr. 2015, pp. 987–991.
- [74] A. L. Winkworth, P. J. Davis, E. Ellis, and R. D. Adams, “Variability and consistency in speech breathing during reading: lung volumes, speech intensity, and linguistic factors,” *Journal of Speech and Hearing Research*, vol. 37, no. 3, pp. 535–556, Jun. 1994.
- [75] M. Włodarczak and M. Heldner, “Breathing in Conversation,” *Frontiers in Psychology*, vol. 11, 2020. [Online]. Available: <https://www.frontiersin.org/articles/10.3389/fpsyg.2020.575566>
- [76] O. Egorow, T. Mrech, N. Weißkirchen, and A. Wendemuth, “Employing Bottleneck and Convolutional Features for Speech-Based Physical Load Detection on Limited Data Amounts,” in *Interspeech 2019*. ISCA, Sep. 2019, pp. 1666–1670. [Online]. Available: [http://www.isca-speech.org/archive/Interspeech\\_2019/abstracts/2502.html](http://www.isca-speech.org/archive/Interspeech_2019/abstracts/2502.html)
- [77] M. Włodarczak and M. Heldner, “Respiratory Constraints in Verbal and Non-verbal Communication,” *Frontiers in Psychology*, vol. 8, May 2017. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5434352/>
- [78] H.-J. Seo, P.-H. Lee, B.-G. Kim, S.-H. Lee, J.-S. Park, J. Lee, S.-W. Park, D.-J. Kim, C.-S. Park, and A.S. Jang, “Methacholine bronchial provocation test in patients with asthma: serial measurements and clinical significance,” *The Korean Journal of Internal Medicine*, vol. 33, no. 4, p. 807, 2018.
- [79] M. Ball, *The metaverse: and how it will revolutionize everything*. Liveright Publishing, 2022.
- [80] V. S. Nallanthighal, Z. Mostaani, A. Harm” a, H. Strik, and M. Magimai-Doss, “Deep learning architectures for” estimating breathing signal and respiratory parameters from speech recordings,” *Neural Networks*, vol. 141, pp. 211–224, 2021.
- [81] J. Z. Chen, I. M. Katz, M. Pichelin, K. Zhu, G. Caillibotte, M. L. Noga, W. H. Finlay, and A. R. Martin, “Comparison of pulsed versus continuous oxygen delivery using realistic adult nasal airway replicas,” *International journal of chronic obstructive pulmonary disease*, pp. 2559–2571, 2017.

- [82] E. C. Larson, M. Goel, G. Boriello, S. Heltshe, M. Rosenfeld, and S. N. Patel, "Spirosmart: using a microphone to measure lung function on a mobile phone," in *Proceedings of the 2012 ACM Conference on ubiquitous computing*, 2012, pp. 280–289.
- [83] M. Goel, E. Saba, M. Stiber, E. Whitmire, J. Fromm, E. C. Larson, G. Borriello, and S. N. Patel, "Spirocall: Measuring lung function over a phone call," in *Proceedings of the 2016 CHI conference on human factors in computing systems*, 2016, pp. 5675–5685.
- [84] ResAppHealth, "Positive results from australian prospective clinical study confirm ResApp's algorithms deliver high accuracy when diagnosing respiratory disease in adults," Press release, Brisbane, Australia, 2019. [Online].
- [85] G. P. Kafentzis, S. Tetsing, J. Brew, L. Jover, M. Galvosas, C. Chaccour, and P. M. Small, "Predicting tuberculosis from real-world cough audio recordings and metadata," arXiv:2307.04842, 2023.
- [86] S. Huddart, L. Asege, D. Jaganath, M. Golla, H. Dang, L. Lovelina, B. Derendinger, A. Andama, D. J. Christopher, N. V. Nhung, G. Theron, C. M. Denkinger, P. Nahid, A. Cattamanchi, and C. Yu, "Continuous cough monitoring: a novel digital biomarker for TB diagnosis and treatment response monitoring," *Int. J. Tuberc. Lung Dis.*, vol. 27(3), pp. 221–222, Mar 2023.
- [87] E. Altshuler, B. Tannir, G. Jolicoeur, M. Rudd, C. Saleem, K. Cherabuddi, D. H. Dore, P. Nagarsheth, J. Brew, P. M. Small, J. Glenn Morris, and S. Grandjean Lapierre, "Digital cough monitoring – a potential predictive acoustic biomarker of clinical outcomes in hospitalized COVID-19 patients," *Journal of Biomedical Informatics*, vol. 138, p. 104283, 2023. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S1532046423000047>
- [88] M. G. Crooks, A. den Brinker, Y. Hayman, J. D. Williamson, A. Innes, C. E. Wright, P. Hill, and A. H. Morice, "Continuous cough monitoring using ambient sound recording during convalescence from a COPD exacerbation," *Lung*, vol. 195, no. 3, pp. 289–294, Jun 2017. [Online]. Available: <https://doi.org/10.1007/s00408-017-9996-2>
- [89] M. G. Crooks, A. C. den Brinker, S. Thackray-Nocera, R. van Dinther, C. E. Wright, and A. H. Morice, "Domiciliary cough monitoring for the prediction of COPD exacerbations," *Lung*, vol. 199, no. 2, pp. 131–137, 2021. [Online]. Available: <https://doi.org/10.1007/s00408-021-00435-9>
- [90] A. C. den Brinker, R. van Dinther, M. G. Crooks, S. Thackray-Nocera, and A. H. Morice, "Alert system design based on experimental findings from long-term unobtrusive monitoring in COPD," *Biomedical Signal Processing and Control*, vol. 63, p. 102205, 2021. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S1746809420303396>
- [91] E. Maor, D. Perry, D. Mevorach, N. Taiblum, Y. Luz, I. Mazin, A. Lerman, G. Koren, and V. Shalev, "Vocal Biomarker Is Associated With Hospitalization and Mortality Among Heart Failure Patients," *Journal of the American Heart Association*, vol. 9, no. 7, p. e013359, Apr. 2020. [Online]. Available: <https://www.ahajournals.org/doi/10.1161/JAHA.119.013359>
- [92] D. Robertson, N. Patel, C. Chirculescu, J. Manfredonia, L. Peltzer, and S. Dogra, "HOARSE VOICE OF A FAILING HEART: CARDIO-VOCAL SYNDROME," *Chest*, vol. 160, no. 4, p. A111, Oct. 2021. [Online]. Available: <https://linkinghub.elsevier.com/retrieve/pii/S0012369221015889>
- [93] G. Golovchiner, M. Glikson, M. Swissa, Y. Sela, A. Abelov, O. Morelli, and A. Beker, "Automated detection of atrial fibrillation based on vocal features analysis," *Journal of Cardiovascular Electrophysiology*, vol. 33, no. 8, pp. 1647–1654, Aug. 2022.
- [94] P. Gouda, E. Ganni, P. Chung, V. K. Randhawa, G. Marquis-Gravel, R. Avram, J. A. Ezekowitz, and A. Sharma, "Feasibility of Incorporating Voice Technology and Virtual Assistants in Cardiovascular Care and Clinical Trials," *Current Cardiovascular Risk Reports*, vol. 15, no. 8, p. 13, 2021.
- [95] J. Sidorova, P. Carbonell, and M. Cukić, "Blood Glucose Estimation From Voice: First Review of Successes' and Challenges," *Journal of Voice*, vol. 36, no. 5, pp. 737.e1–737.e10, Sep. 2022, publisher: Elsevier. [Online]. Available: [https://www.jvoice.org/article/S0892-1997\(20\)30334-9/fulltext](https://www.jvoice.org/article/S0892-1997(20)30334-9/fulltext)
- [96] L. Wu and Z. Zhang, "A Computational Study of Vocal Fold Dehydration During Phonation," *IEEE Transactions on Biomedical Engineering*, vol. 64, no. 12, pp. 2938–2948, Dec. 2017, conference Name: IEEE Transactions on Biomedical Engineering.



- [97] K. Verdolini, Y. Min, I. R. Titze, J. Lemke, K. Brown, M. van Mersbergen, J. Jiang, and K. Fisher, "Biological mechanisms underlying voice changes due to dehydration," *Journal of speech, language, and hearing research: JSLHR*, vol. 45, no. 2, pp. 268–281, Apr. 2002.
- [98] S. S. Azar and D. K. Chhetri, "Phonation Threshold Pressure Revisited: Effects of Intrinsic Laryngeal Muscle Activation," *The Laryngoscope*, vol. 132, no. 7, pp. 1427–1432, 2022, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/lary.29944>. [Online]. Available: <https://onlinelibrary.wiley.com/doi/abs/10.1002/lary.29944>
- [99] T. B. Agbabiaka, J. Savovic, and E. Ernst, "Methods for Causality Assessment of Adverse Drug Reactions," *Drug Safety*, vol. 31, no. 1, pp. 21–37, Jan. 2008. [Online]. Available: <https://doi.org/10.2165/00002018-200831010-00003>
- [100] G. Golovchiner, M. Glikson, M. Swissa, Y. Sela, A. Abelow, O. Morelli, and A. Beker, "Automated detection of atrial fibrillation based on vocal features analysis," *Journal of Cardiovascular Electrophysiology*, vol. 33, no. 8, pp. 1647–1654, 2022.
- [101] P. Gouda, E. Ganni, P. Chung, V. K. Randhawa, G. Marquis-Gravel, R. Avram, J. A. Ezekowitz, and A. Sharma, "Feasibility of incorporating voice technology and virtual assistants in cardiovascular care and clinical trials," *Current Cardiovascular Risk Reports*, vol. 15, pp. 1–8, 2021.
- [102] J. Jeon, A. Palanica, S. Sarabadani, M. Lieberman, and Y. Fossat, "Biomarker potential of real-world voice signals to predict abnormal blood glucose levels," *bioRxiv*, pp. 2020–09, 2020.
- [103] A. Hassoon, Y. Baig, D. Q. Naiman, D. D. Celentano, D. Lansley, V. Stearns, J. Coresh, J. Schrack, S. S. Martin, H.-C. Yeh, H. Zeilberger, and L. J. Appel, "Randomized trial of two artificial intelligence coaching interventions to increase physical activity in cancer survivors," *npj Digital Medicine*, vol. 4, no. 1, pp. 1–6, Dec. 2021, number: 1 Publisher: Nature Publishing Group. [Online]. Available: <https://www.nature.com/articles/s41746-021-00539-9>
- [104] B. Halpern, R. van Son, M. Brekel, and O. Scharenborg, *Detecting and Analysing Spontaneous Oral Cancer Speech in the Wild*, Oct. 2020, pages: 4830.
- [105] X. Ming, R. Patel, V. Kang, S. Chokroverty, and P. O. Julu, "Respiratory and autonomic dysfunction in children with autism spectrum disorders," *Brain & Development*, vol. 38, no. 2, pp. 225–232, Feb. 2016.
- [106] D. H. Heck, B. L. Correia, M. B. Fox, Y. Liu, M. Allen, and S. Varga, "Recent insights into respiratory modulation of brain activity offer new perspectives on cognition and emotion," *Biological Psychology*, vol. 170, p. 108316, Apr. 2022.
- [107] M. Gevezova, D. Minchev, I. Pacheva, T. Todorova, R. Yordanova, E. Timova, I. Ivanov, and V. Sarafian, "Association of NGF and Mitochondrial Respiration with Autism Spectrum Disorder," *International Journal of Molecular Sciences*, vol. 23, no. 19, p. 11917, Jan. 2022, number: 19 Publisher: Multidisciplinary Digital Publishing Institute. [Online]. Available: <https://www.mdpi.com/1422-0067/23/19/11917>
- [108] J. McDaniel and C. M. Schuele, "When Will He Talk? An Evidence-Based Tutorial for Measuring Progress Toward Use of Spoken Words in Preverbal Children With Autism Spectrum Disorder," *American Journal of Speech-Language Pathology*, vol. 30, no. 1, pp. 1–18, Jan. 2021. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8740598/>
- [109] A. Baird, S. Amiriparian, N. Cummins, A. M. Alcorn, A. Batliner, S. Pugachevskiy, M. Freitag, M. Gerczuk, and B. Schuller, "Automatic Classification of Autistic Child Vocalisations: A Novel Database and Results," in *Interspeech 2017*. ISCA, Aug. 2017, pp. 849–853. [Online]. Available: [https://www.isca-speech.org/archive/interspeech\\_2017/baird17\\_interspeech.html](https://www.isca-speech.org/archive/interspeech_2017/baird17_interspeech.html)
- [110] J. H. Lee, G. W. Lee, G. Bong, H. J. Yoo, and H. K. Kim, "End-to-End Model-Based Detection of Infants with Autism Spectrum Disorder Using a Pretrained Model," *Sensors*, vol. 23, no. 1, p. 202, Dec. 2022. [Online]. Available: <https://www.mdpi.com/1424-8220/23/1/202>
- [111] J. Krajewski, R. Wieland, and A. Batliner, "An Acoustic Framework for Detecting Fatigue in Speech Based Human-Computer-Interaction," in *Computers Helping People with Special Needs*, ser. Lecture Notes in Computer Science. Springer, Berlin, Heidelberg, Jul. 2008, pp. 54–61. [Online]. Available: [https://link.springer.com/chapter/10.1007/978-3-540-70540-6\\_7](https://link.springer.com/chapter/10.1007/978-3-540-70540-6_7)

- [112] M. Gadesmann and N. Miller, "Reliability of speech diadochokinetic test measurement," *International Journal of Language & Communication Disorders*, vol. 43, no. 1, pp. 41–54, 2008.
- [113] Y. Segal, K. Hitczenko, M. Goldrick, A. Buchwald, A. Roberts, and J. Keshet, "DDKtor: Automatic Diadochokinetic Speech Analysis," Jun. 2022, arXiv:2206.14639 [cs, eess]. [Online]. Available: <http://arxiv.org/abs/2206.14639>
- [114] E. Kaplan, H. Goodglass, S. Weintraub, O. Segal, and A. v. Loon-Vervoorn, *Boston naming test*, 2nd ed. Austin, TX: Pro-ed, 2001, oCLC: 778264850.
- [115] A. Harry and S. F. Crowe, "Is the Boston Naming Test still fit for purpose?" *The Clinical Neuropsychologist*, vol. 28, no. 3, pp. 486–504, 2014.
- [116] B. E. Gavett, "Neuropsychological Assessment Battery," in *Encyclopedia of Clinical Neuropsychology*, J. S. Kreutzer, J. DeLuca, and B. Caplan, Eds. New York, NY: Springer, 2011, pp. 1761–1765. [Online]. Available: [https://doi.org/10.1007/978-0-387-79948-3\\_2250](https://doi.org/10.1007/978-0-387-79948-3_2250)
- [117] Z. S. Nasreddine, N. A. Phillips, V. Bedirian, S. Charbonneau, V. Whitehead, I. Collin, J. L. Cummings, and H. Chertkow, "The Montreal Cognitive Assessment, MoCA: A Brief Screening Tool For Mild Cognitive Impairment," *Journal of the American Geriatrics Society*, vol. 53, no. 4, pp. 695–699, 2005, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1532-5415.2005.53221.x>. [Online]. Available: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1532-5415.2005.53221.x>
- [118] M. G. Pendleton, R. K. Heaton, R. A. W. Lehman, and D. Hulihan, "Diagnostic utility of the thurstone word fluency test in neuropsychological evaluations," *Journal of Clinical Neuropsychology*, vol. 4, no. 4, pp. 307–317, Dec. 1982, publisher: Routledge eprint: <https://doi.org/10.1080/01688638208401139>. [Online]. Available: <https://doi.org/10.1080/01688638208401139>
- [119] J. Patterson, "Controlled Oral Word Association Test," in *Encyclopedia of Clinical Neuropsychology*, J. S. Kreutzer, J. DeLuca, and B. Caplan, Eds. Cham: Springer International Publishing, 2018, pp. 958–961. [Online]. Available: [https://doi.org/10.1007/978-3-319-57111-9\\_876](https://doi.org/10.1007/978-3-319-57111-9_876)
- [120] C. R. Greenwood, J. Buzhardt, D. Walker, F. Jia, and J. J. Carta, "Criterion Validity of the Early Communication Indicator for Infants and Toddlers," *Assessment for Effective Intervention*, vol. 45, no. 4, pp. 298–310, Sep. 2020, publisher: SAGE Publications Inc. [Online]. Available: <https://doi.org/10.1177/1534508418824154>
- [121] S. H. S. Kim, V. Hus, and C. Lord, "Autism Diagnostic Interview-Revised," in *Encyclopedia of Autism Spectrum Disorders*, F. R. Volkmar, Ed. New York, NY: Springer, 2013, pp. 345–349. [Online]. Available: [https://doi.org/10.1007/978-1-4419-1698-3\\_894](https://doi.org/10.1007/978-1-4419-1698-3_894)
- [122] P. Liston, J. Byrne, O. Keogh, and P. J. Byrne, "Beyond calls: Modeling the connection center," in *2017 Winter Simulation Conference (WSC)*, Dec. 2017, pp. 3804–3815.
- [123] G. Koole, *Call Center Optimization*. Amsterdam: MG books, Jan. 2013.
- [124] D. Amodei, S. Ananthanarayanan, R. Anubhai, J. Bai, E. Battenberg, C. Case, J. Casper, B. Catanzaro, Q. Cheng, G. Chen, J. Chen, J. Chen, Z. Chen, M. Chrzanowski, A. Coates, G. Diamos, K. Ding, N. Du, E. Elsen, J. Engel, W. Fang, L. Fan, C. Fougner, L. Gao, C. Gong, A. Hannun, T. Han, L. V. Johannes, B. Jiang, C. Ju, B. Jun, P. LeGresley, L. Lin, J. Liu, Y. Liu, W. Li, X. Li, D. Ma, S. Narang, A. Ng, S. Ozair, Y. Peng, R. Prenger, S. Qian, Z. Quan, J. Raiman, V. Rao, S. Satheesh, D. Seetapun, S. Sengupta, K. Srinet, A. Sriram, H. Tang, L. Tang, C. Wang, J. Wang, K. Wang, Y. Wang, Z. Wang, Z. Wang, S. Wu, L. Wei, B. Xiao, W. Xie, Y. Xie, D. Yogatama, B. Yuan, J. Zhan, and Z. Zhu, "Deep Speech 2: End-to-end Speech Recognition in English and Mandarin," in *Proceedings of the 33rd International Conference on International Conference on Machine Learning - Volume 48*, ser. ICML'16. New York, NY, USA: JMLR.org, 2016, pp. 173–182. [Online]. Available: <http://dl.acm.org/citation.cfm?id=3045390.3045410>
- [125] A. Tripathy, "The Contact Center Is Having Its Model T Moment." [Online]. Available: <https://www.forbes.com/sites/forbestechcouncil/2017/12/07/the-contact-center-is-having-its-model-t-moment/>
- [126] T. Bickmore and T. Giorgino, "Health dialog systems for patients and consumers," *Journal of Biomedical Informatics*, vol. 39, no. 5, pp. 556–571, Oct. 2006.
- [127] K. K. Fitzpatrick, A. Darcy, and M. Vierhile, "Delivering Cognitive Behavior Therapy to Young Adults With Symptoms of Depression and Anxiety Using a Fully Automated Conversational Agent (Woebot): A Randomized

- Controlled Trial," *JMIR Mental Health*, vol. 4, no. 2, p. e19, 2017. [Online]. Available: <https://mental.jmir.org/2017/2/e19/>
- [128] S. Provoost, H. M. Lau, J. Ruwaard, and H. Riper, "Embodied Conversational Agents in Clinical Psychology: A Scoping Review," *Journal of Medical Internet Research*, vol. 19, no. 5, p. e151, May 2017.
- [129] A. A. Abd-Alrazaq, A. Rababeh, M. Alajlani, B. M. Bewick, and M. Househ, "Effectiveness and Safety of Using Chatbots to Improve Mental Health: Systematic Review and Meta-Analysis," *Journal of Medical Internet Research*, vol. 22, no. 7, p. e16021, Jul. 2020.
- [130] B. Singh, T. Olds, J. Brinsley, D. Dumuid, R. Virgara, L. Matricciani, A. Watson, K. Szeto, E. Eglitis, A. Miatke, C. E. M. Simpson, C. Vandelanotte, and C. Maher, "Systematic review and meta-analysis of the effectiveness of chatbots on lifestyle behaviours," *npj Digital Medicine*, vol. 6, no. 1, pp. 1–10, Jun. 2023, number: 1 Publisher: Nature Publishing Group. [Online]. Available: <https://www.nature.com/articles/s41746-023-00856-1>
- [131] R. J. Baken and R. F. Orlikoff, *Clinical Measurement of Speech & Voice*, 2nd ed. San Diego: Cengage Learning, Dec. 1999.
- [132] G. Hardavella, I. Karampinis, A. Frille, K. Sreter, and I. Rousalova, "Oxygen devices and delivery systems," *Breathe*, vol. 15, no. 3, pp. e108–e116, 2019.
- [133] W. Chatila, T. Nugent, G. Vance, J. Gaughan, and G. J. Criner, "The effects of high-flow vs low-flow oxygen on exercise in advanced obstructive airways disease," *Chest*, vol. 126, no. 4, pp. 1108–1115, 2004.
- [134] V. S. Nallanthighal, Z. Mostaani, A. Harm` a, H. Strik, and M. Magimai-Doss, "Deep learning architectures for estimating breathing signal and respiratory parameters from speech recordings," *Neural Networks*, vol. 141, pp. 211–224, Sep. 2021. [Online]. Available: <https://linkinghub.elsevier.com/retrieve/pii/S0893608021001179>
- [135] J. Z. Chen, I. M. Katz, M. Pichelin, K. Zhu, G. Caillibotte, M. L. Noga, W. H. Finlay, and A. R. Martin, "Comparison of pulsed versus continuous oxygen delivery using realistic adult nasal airway replicas," *International Journal of Chronic Obstructive Pulmonary Disease*, vol. 12, pp. 2559–2571, Aug. 2017. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5574700/>
- [136] A. Harm` a, U. Großekath` ofer, O. Ouweltjes, and V. S. Nallanthighal, "Forecasting of Breathing Events from Speech for Respiratory Support," in *ICASSP 2023 - 2023 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, Jun. 2023, pp. 1–5.
- [137] Matthew Ball, *The Metaverse and how it will revolutionize everything*. Liveright Publishing, 2022.
- [138] C. Cassarly, A. Doyle, T. Ly, J. Horn, M. Aitchison, J. Elm, J. Fridriksson, and L. Bonilha, "Speech Entrainment for Aphasia Recovery (SpARc) phase II trial design," *Contemporary Clinical Trials Communications*, vol. 24, p. 100876, Nov. 2021. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8606333/>
- [139] "Western Aphasia Battery–Revised." [Online]. Available: <https://psycnet.apa.org/doiLanding?doi=10.1037%2F15168-000>
- [140] G. M. Insights, *Vehicle Health Monitoring Market Size By Component (Hardware [Sensors, GPS Systems, OBD Ports], Software, Services), By Health Management (Diagnostics, Prognostics), By Vehicle Type (Passenger, Commercial), By Sales Channel (OEM, Aftermarket), Industry Analysis Report, Regional Outlook, Growth Potential, Price Trends, Competitive Market Share Forecast, 2021*.
- [141] G. Trajkovic, V. Star` cevi` c, M. Latas, M. Le` starevi` c, T. Ille, Z. Bukumiri` c, and J. Marinkovi` c, "Reliability of the hamilton rating scale for depression: a meta-analysis over a period of 49 years," *Psychiatry research*, vol. 189, no. 1, pp. 1–9, 2011.
- [142] C. Heyde, H. Leutheuser, B. Eskofier, K. Roecker, and A. Gollhofer, "Respiratory Inductance Plethysmography—A Rationale for Validity during Exercise," *Medicine & Science in Sports & Exercise*, vol. 46, no. 3, pp. 488–495, Mar. 2014. [Online]. Available: <https://journals.lww.com/00005768-201403000-00008>
- [143] L. E. Brummer, S. Katzenschlager, M. Gaeddert, C. Erdmann, S. Schmitz, M. Bota, M. Grilli, J. Larmann,` M. A. Weigand, N. R. Pollock, A. Mace, S. Carmona, S. Ongarello, J. A. Sacks, and C. M. Denkinge, "Accuracy of novel antigen rapid diagnostics for SARS-CoV-2: A living systematic review and meta-analysis," *PLOS Medicine*, vol. 18, no. 8, p. e1003735, Aug. 2021, publisher: Public Library of Science. [Online]. Available: <https://journals.plos.org/plosmedicine/article?id=10.1371/journal.pmed.1003735>

- [144] A. Kertesz, "The Western Aphasia Battery: a systematic review of research and clinical applications," *Aphasiology*, vol. 36, no. 1, pp. 21–50, Jan. 2022, publisher: Routledge eprint: <https://doi.org/10.1080/02687038.2020.1852002>. [Online]. Available: <https://doi.org/10.1080/02687038.2020.1852002>
- [145] N. Szejko and K. R. Muller-Vahl, "Challenges in the Diagnosis and Assessment in Patients with Tourette" Syndrome and Comorbid Obsessive-Compulsive Disorder," *Neuropsychiatric Disease and Treatment*, vol. 17, pp. 1253–1266, Apr. 2021. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8096634/>