

On Asymptotic Analysis of Perturbed Sweeping Processes with Application to Optimization

Zhaoyue Xia, *Student Member, IEEE*, Jun Du, *Senior Member, IEEE*, Chunxiao Jiang, *Fellow, IEEE*,
H. Vincent Poor, *Life Fellow, IEEE*, and Yong Ren, *Senior Member, IEEE*

Abstract—Convergence analysis of constrained optimization methods from the dynamical systems viewpoint has attracted considerable attention because it provides a geometric demonstration towards the shadowing trajectory of a numerical scheme. In this work, we establish a tight connection between a continuous-time nonsmooth dynamical system called a perturbed sweeping process (PSP) and a proximal stochastic approximation scheme. Theoretical results are obtained by analyzing the asymptotic pseudo trajectory of a PSP. We show that under mild assumptions a proximal stochastic approximation scheme converges to an internally chain transitive invariant set of the corresponding PSP. Furthermore, given the existence of a Lyapunov function V with respect to a set Λ , convergence to Λ can be established if $V(\Lambda)$ has an empty interior. Based on these theoretical results, we are able to provide a useful framework for convergence analysis of proximal gradient methods. Illustrative examples are provided to determine the convergence of proximal variants of gradient methods (including accelerated gradient methods). Finally, numerical simulations are conducted to confirm the validity of theoretical analysis.

Index Terms—Constrained optimization, Lyapunov analysis, perturbed sweeping process, convergence analysis.

I. INTRODUCTION

Constrained optimization is a fundamental problem in mathematical programming [1]–[3], where the objective is to minimize a function subject to a set of constraints. These constraints are usually nonlinear and they often reflect real-world limitations such as resource availability, physical laws, or operational boundaries. The complexity of constrained optimization stems from the interplay between the objective function and these constraints, enforcing the trajectory of iterations produced by an optimization scheme to move along the boundaries of a constrained set or within the set.

As a classical and significant topic, convergence analysis of constrained optimization methods has been of interest due to its essential differences from that of unconstrained schemes. To be specific, a constrained method iteratively seeks a proximal point within a set, which yields a nonsmooth part in the iteration. From a geometric viewpoint, given a convex set $C \subseteq \mathbb{R}^m$ and a point $x \in C$, any vector $v \in \mathbb{R}^m$ can be decomposed into two orthogonal components in the tangent

cone $\mathcal{T}_C(x)$ and the normal cone $N_C(x)$, respectively. Moreover, the projection into the set C actually means canceling the component in the normal cone. Therefore, any movement in free space will be projected into the tangent cone.

To analyze the convergence properties of an optimization scheme, two principal methodologies have emerged: numerical analysis and dynamical systems theory. Numerical analysis [4]–[6] offers a straightforward depiction of the concrete convergence rates, providing a clear understanding of the speed at which an algorithm approaches its optimal solution. However, this approach often lacks the deeper geometric insights that can be gleaned from a dynamical systems perspective. This latter approach, grounded in the study of continuous-time systems, enriches the analysis by revealing the underlying geometric structures and dynamics that influence the convergence behavior of optimization schemes.

In recent years, there has been a significant surge in research on constrained optimization from a dynamical systems perspective. We present a synthesis of some of the most recent and representative findings in this domain. In [7], the authors analyzed differential inclusions associated with accelerated variants of the alternating direction method of multipliers (ADMM) and illustrated a tradeoff between the convergence rate and the damping factor. A primal-dual dynamical system approach was proposed to track an inequality constrained time-varying convex optimization problem in [8]. For online time-varying optimization of linear time-invariant systems, a linear dynamical system was applied to develop a convergent projected primal-dual gradient flow method [9]. Accelerated methods were developed under the framework of fixed-time stability of nonlinear dynamical systems for functions under Polyak-Ljapunov inequality conditions in [10]. In [11], dynamic optimization theory was established for nonlinear complementarity systems. The second-order dynamical system was extended to constrained distributed optimization in [12].

According to the classical result [13] in stochastic approximation, the continuous-time dynamics of an unconstrained iterative discrete method can be demonstrated by an ordinary differential equation (ODE). To be specific, a stochastic approximation scheme given by

$$x_{n+1} = x_n + \alpha_{n+1}(\psi(x_n) + \xi_{n+1}), \quad (1)$$

converges to an internally chain transitive set of the dynamical system expressed by the ODE $\dot{x} = \psi(x)$, where \dot{x} means the derivative with respect to time, $\{\alpha_n\}$ are vanishing step sizes, ψ is Lipschitz continuous, and $\{\xi_n\}$ is a sequence of martingale difference noise.

Z. Xia, J. Du, and Y. Ren are with the Department of Electronic Engineering, Tsinghua University, Beijing, 100084, China, (e-mail: xiazy19@mails.tsinghua.edu.cn; {jundu, reny}@tsinghua.edu.cn).

C. Jiang is with Tsinghua Space Center, Tsinghua University, Beijing, 100084, China. (e-mail: jchx@tsinghua.edu.cn).

H. V. Poor is with the Department of Electrical and Computer Engineering, Princeton University, Princeton, NJ 08544 USA, (e-mail: poor@princeton.edu).

Although it is straightforward to show that a gradient descent method converges simply by replacing ψ by $-\nabla f$ in (1), the underlying relationship between the internally chain transitive set and the critical point set of f is not immediately apparent. Bridging this gap is the concept of a Lyapunov function V (total energy of the system), which plays a pivotal role in the stability analysis of dynamical systems. By incorporating the objective function f into the Lyapunov function, the substantial dissipation of energy leads to a local minimum of V and hence f .

As a counterpart to the gradient descent method, the proximal gradient method is intrinsically linked to a PSP with a constraint set¹, as established in [14]. However, it is not evident whether the convergence conclusions drawn for unconstrained stochastic approximations remain valid in the context of constrained problems. In fact, this uncertainty arises from the nonsmooth characteristics inherent to proximal gradient methods for constrained optimization.

Furthermore, there is a natural inclination to employ a nonsmooth Lyapunov function that encapsulates the complexity of the problem. Ideally, such a function would be decomposed into two components: a smooth part that corresponds to the vector field ψ , and a lower-semicontinuous part that accounts for the constraints. Unfortunately, this approach often encounters difficulties, as the nonsmoothness can impede the straightforward application of traditional Lyapunov theory.

In fact, optimal control of a PSP has been a well-studied problem, which comes from the application to the crowd motion model. A number of theoretical results have been developed [15]–[17]. Numerical analysis [18], [19] on discretization of a continuous-time PSP is aimed at deriving the convergence order of the numerical scheme towards the continuous dynamics within a finite time. These results, however, do not provide the Lyapunov properties of ω -limit sets of a PSP and fail in the infinite-time asymptotic analysis.

In this article, we develop dynamical systems theory with respect to constrained optimization schemes, aimed at providing a general framework for convergence analysis. Specifically, the contributions of this work can be summarized as follows:

- We provide a Lyapunov analysis for a PSP with a fixed constraint set. We show that if a PSP is a gradient-like dynamical system with a compact convex set, the ω -limit set of any initial point x is contained in the fixed point set of the corresponding Lyapunov function.
- We establish the connection between a PSP and its Euler discretization and show that the discrete iterations converge to an internally chain transitive set of the PSP, which is similar to the behavior of unconstrained stochastic approximation. Furthermore, we develop the Lyapunov theory for such an iterative method.
- By utilizing the theory of Lyapunov pairs, we provide several examples of convergence analysis of constrained variants of popular gradient-based methods.
- Numerical simulations are conducted to verify the validity of the theoretical analysis. The results show that all the

constrained algorithms succeed in converging to local minima within the constraint set.

The rest of the article is organized as follows. Basic concepts and notation are introduced in Section II. Subsequently, the primary theoretical results are demonstrated and derived in Section III. We provide examples of applications to optimization in Section IV. Numerical simulation results are presented in Section V and Section VI concludes this article.

II. BASIC CONCEPTS AND NOTATION

In this section, we provide some notation and basic concepts (especially in the theory of dynamical systems) to be used throughout the article.

Let X be a topological space, \mathbb{R}^+ be the semigroup of nonnegative real numbers and $\mathbb{T} \subseteq \mathbb{R}^+$ be a subsemigroup of the additive group. A triplet (X, \mathbb{T}, π) , where $\pi : \mathbb{T} \times X \rightarrow X$ is a continuous mapping satisfying $\pi(0, x) = x$ and $\pi(s, \pi(t, x)) = \pi(s + t, x)$ for all $x \in X$ and $s, t \in \mathbb{T}$, is called a (continuous) dynamical system. Given $x \in X$, the set $\Upsilon_x := \pi(\mathbb{T}, x)$ is called a trajectory (associated with x). A point $x \in X$ is called a fixed point of (X, \mathbb{T}, π) if $\pi(t, x) = x$ for all $t \in \mathbb{T}$. A discrete dynamical system where $\mathbb{T} \subseteq \mathbb{Z}$ is called a cascade.

A nonempty set $M \subseteq X$ is called (positively) invariant with respect to a dynamical system (X, \mathbb{T}, π) if $\pi(t, M) \subset M$ for every $(t \geq 0) t \in \mathbb{T}$. Let $J \subseteq X$. The set

$$\omega(J) := \bigcap_{t \geq 0} \overline{\bigcup_{s \geq t} \pi(s, J)}, \quad (2)$$

where \bar{A} denotes the closure of a set A , is called the ω -limit set for J . An equivalent definition of the ω -limit set is

$$\omega(J) = \{u \in X : \exists x \in J, \exists t_n \rightarrow \infty, \pi(t_n, x) \rightarrow u\}. \quad (3)$$

Let $\Sigma \subseteq X$ be a compact positively invariant subset of a metric space (X, d) , $\varepsilon > 0$, and $t > 0$. The collection $\{x = x_0, x_1, x_2, \dots, x_k = y; t_0, t_1, \dots, t_k\}$ of points $x_i \in \Sigma$ and the numbers $t_i \in \mathbb{T}$ such that $t_i \geq t$ and the distance $d(\pi(t_i, x_i), x_{i+1}) < \varepsilon$, ($i = 0, 1, \dots, k - 1$) is called an (ε, t, π) -chain joining the points x and y . The set Σ is called *internally chain transitive* if for all $a, b \in \Sigma$, $\varepsilon > 0$ and $t > 0$, there exists an (ε, t, π) -chain in Σ connecting a and b .

A dynamical system (X, \mathbb{T}, π) is said to be a gradient-like dynamical system if it has a global Lyapunov function $V : X \rightarrow \mathbb{R}$, i.e., V is continuous and satisfies $V(\pi(t, x)) \leq V(x)$ for all $x \in X$ and $t \in \mathbb{T}$.

Let S be a nonempty subset of a Hilbert space \mathcal{H} , and $x \in \mathcal{H}$. The distance between x and S is expressed by

$$d(x; S) := \inf_{y \in S} \|x - y\|. \quad (4)$$

The set of nearest points of x in S is defined by

$$P_S(x) := \{u \in S : \|x - u\| = d(x; S)\}. \quad (5)$$

For a convex subset $S \subseteq \mathcal{H}$ and $x \in \mathcal{H}$, the normal cone to S at x is $N_S(x) = \{v \in S : \langle v, y - x \rangle \leq 0, \forall y \in S\}$. Correspondingly, we use $\mathcal{T}_S(x)$ to represent the tangent cone. Given a constrained optimization problem $\min_{x \in C} f(x)$ for a

¹Note that since the constraint set is time-independent for a standard constrained optimization problem, we will restrict our discussion to a PSP with a fixed set.

closed and convex set $C \subseteq \mathbb{R}^m$ and a differentiable function f , the set of Karush-Kuhn-Tucker (KKT) points is defined as $\mathcal{L} := \{x \in C : 0 \in \nabla f(x) + N_C(x)\}$.

A sequence $\{y(t)\}_{t \in \mathbb{R}}$ of elements in \mathcal{H} is said to converge to a set J if $d(y(t); J) \rightarrow 0$ as $t \rightarrow +\infty$, denoted by $y(t) \rightarrow J$.

Given a nonempty set C , we use \mathcal{I}_C to represent the indicator function of C , i.e., $\mathcal{I}_C(x) = 0$ if $x \in C$ and $\mathcal{I}_C(x) = +\infty$ otherwise. For a lower semi-continuous function $\varphi : \mathcal{H} \rightarrow \mathbb{R}$ on a Hilbert space \mathcal{H} , a vector $\xi \in \mathcal{H}$ is called a Fréchet subgradient, written $\xi \in \partial_F \varphi(x)$, at x if

$$\varphi(y) \geq \varphi(x) + \langle \xi, y - x \rangle + o(\|y - x\|), \quad \forall y \in \mathcal{H}. \quad (6)$$

We use $\mathbb{B}(x, r)$ to denote a closed ball in a metric space centered at x with radius r .

Let $\Upsilon(t, x)$ be a trajectory of a dynamical system and Λ be a subset of a metric space X . A continuous function $V : X \rightarrow \mathbb{R}$ is called a Lyapunov function for a set Λ , if $V(y) < V(x)$ for all $x \in X \setminus \Lambda$, $y \in \Upsilon(t, x)$, $t > 0$, and $V(y) \leq V(x)$ for all $x \in \Lambda$, $y \in \Upsilon(t, x)$, and $t \geq 0$.

Throughout the paper, two forms of the well-known Gronwall's inequalities [20] will be used.

- **The classical differential form.** Assume that $u : [0, T) \rightarrow \mathbb{R}$ is continuously differentiable, $T \in (0, \infty)$, and satisfies the differential inequality

$$\frac{du}{dt} \leq a(t)u(t) + b(t), \quad (7)$$

for some integrable functions a, b on $(0, T)$. Then, u satisfies the pointwise bound

$$u(t) \leq e^{A(t)}u(0) + \int_0^t b(s)e^{A(t)-A(s)}ds, \quad (8)$$

where $A(t) := \int_0^t a(s)ds$ for all $t \in [0, T)$.

- **The discrete form.** Consider a sequence of real numbers $\{u_n\}$ such that

$$u_{n+1} \leq a_{n+1}u_n + b_{n+1}, \quad \forall n \geq 0, \quad (9)$$

where $\{a_n\}$ and $\{b_n\}$ are two given sequences of real numbers and $\{a_n\}$ is furthermore positive. Then

$$u_n \leq A_n u_0 + \sum_{k=1}^n A_{k,n} b_k, \quad \forall n \geq 0, \quad (10)$$

where $A_n := \prod_{k=1}^n a_k$, $A_{k,n} := A_n / A_k$.

III. FROM CONTINUOUS DYNAMICS TO CASCADES

In this section, a constrained system is considered associated with a closed and convex set \mathcal{K} , which projects the dynamics into \mathcal{K} . The projected dynamics are expressed by

$$\frac{dz}{dt} \in -\psi(t, z) - N_{\mathcal{K}}(z). \quad (11)$$

This differential inclusion cannot be viewed as a variational inequality problem due to its non-autonomous nature. Indeed, the dynamics are covered by a topic termed the *PSP*, which will be discussed in detail below.

A. Perturbed Sweeping Processes

We first present sufficient conditions for the existence and uniqueness of a PSP.

Condition 1 (sweeping-regular). *Let \mathcal{H} and \mathcal{F} be Hilbert spaces. A function $\psi : \mathbb{R} \times \mathcal{H} \rightarrow \mathcal{F}$ is said to be sweeping-regular on a pair (I, C) for $I \subseteq \mathbb{R}$ and $C \subseteq \mathcal{H}$ if*

- $\forall \eta > 0$, there exists an integrable nonnegative function $L_\eta(t) : I \rightarrow \mathbb{R}$ such that, for all t and for all $\max\{\|x\|, \|y\|\} < \eta$,

$$\|\psi(t, x) - \psi(t, y)\|_{\mathcal{F}} \leq L_\eta(t)\|x - y\|_{\mathcal{H}}; \quad (12)$$

- there exists an integrable nonnegative function $\beta : I \rightarrow \mathbb{R}$ such that, for all t and for all $x \in C$, $\|\psi(t, x)\|_{\mathcal{F}} \leq \beta(t)(1 + \|x\|_{\mathcal{H}})$.

We begin with a useful lemma ensuring that a composite function is sweeping-regular.

Lemma 1. *Let \mathcal{H}_n be a sequence of Hilbert spaces, $\psi_n : \mathbb{R} \times \mathcal{H} \rightarrow \mathcal{H}_n$ for $n = 1, 2, \dots, N$, and $\mathcal{H} = \mathcal{H}_1 \times \mathcal{H}_2 \times \dots \times \mathcal{H}_N$. If each ψ_n is sweeping-regular on (I, C) , the composite function $\psi(t, x) = (\psi_1(t, x_1), \psi_2(t, x_2), \dots, \psi_N(t, x_N))$ is sweeping-regular on (I, C) , where $x = (x_1, x_2, \dots, x_N) \in \mathcal{H}$.*

Proof. The result is a straightforward consequence of the triangle inequality for Hilbert spaces. \square

Given this condition, we have the following lemma ensuring the existence and uniqueness of a solution to (11):

Lemma 2. [21, Theorem 2.1] *Let \mathcal{H} be a Hilbert space, C be a closed and convex subset of \mathcal{H} , I be a subset of \mathbb{R} , and $\psi : \mathbb{R} \times \mathcal{H} \rightarrow \mathcal{H}$ satisfying Condition 1 for (I, C) . Then the PSP with $x(0) \in C$*

$$-\frac{dx}{dt} \in \psi(t, x) + N_C(x), \quad \text{a.e. } t \in I, \quad (13)$$

has a unique absolutely continuous solution $x(t)$ defined on I . Moreover, for almost everywhere $t \in I$,

$$\|\dot{x}(t) + \psi(t, x(t))\| \leq D\beta(t), \quad \|\psi(t, x(t))\| \leq D\beta(t), \quad (14)$$

for some constant $D = D(x(0), \int_I \beta(s)ds) > 0$.

Note that an absolutely continuous function $x(t)$ is said to be a solution to the sweeping process (13) on an interval $I \subseteq \mathbb{R}$ if $x(t) \in C$ for a.e. $t \in I$ and $\dot{x}(t)$ satisfies (13). Since we will discuss properties of ω -limit sets of a PSP, it is necessary to extend the solution to the entire real line \mathbb{R} (or at least \mathbb{R}^+). By [22, Corollary 2], the differential inclusion (13) is equivalent to the ODE

$$\dot{x}(t) = P_{\mathcal{T}_C(x(t))}[-\psi(t, x)], \quad \text{a.e. } t \in I, \quad (15)$$

where $P_{\mathcal{T}_C(x)}$ denotes the projection into the tangent cone of C at x . By standard procedure to extend a solution of an ODE, we have the following lemma.

Lemma 3. *Suppose for every $\tau > 0$, ψ is sweeping-regular on $([-\tau, \tau], C)$ or $([0, \tau], C)$. Then the solution of (13) is defined for all $t \in \mathbb{R}$ or $t \in \mathbb{R}^+$, respectively.*

Proof. Using the bound from Condition 1, we have

$$\|x(t)\| \leq \|x(0)\| + \int_0^t \|P_{\mathcal{T}_C(x(t))}[-\psi(s, x(s))]\| ds. \quad (16)$$

Since projection into a closed and convex set is nonexpansive, it follows that

$$\begin{aligned} & \|P_{\mathcal{T}_C(x(t))}[-\psi(s, x(s))]\| \\ &= \|P_{\mathcal{T}_C(x(t))}[-\psi(s, x(s))] - P_{\mathcal{T}_C(x(t))}[0]\| \\ &\leq \|\psi(s, x(s))\| \leq \beta(s)(1 + \|x(s)\|). \end{aligned} \quad (17)$$

Hence we obtain

$$\|x(t)\| \leq \|x(0)\| + \int_0^t \beta(s)(1 + \|x(s)\|) ds. \quad (18)$$

Using the above variant of Gronwall's inequality implies

$$\|x(t)\| \leq \|x(0)\| e^{B(t)} + \int_0^t \beta(s) e^{B(t)-B(s)} ds, \quad (19)$$

where $B(t) := \int_0^t \beta(s) ds$. By the integrability of $\beta(t)$ as presented in Condition 1, $x(t)$ lies in a compact ball and the result follows by [23, p. 52, Corollary 2.15]. \square

In the subsequent analysis, we assume that the solution to the PSP is defined on the entire nonnegative real line \mathbb{R}^+ . We firstly consider the straightforward case where ψ is *strongly monotone*.

Lemma 4. *Let the conditions of Lemma 3 hold. Assume that ψ satisfies the condition for strong monotonicity, i.e.,*

$$\langle \psi(t, x) - \psi(t, y), x - y \rangle \geq \gamma(t) \|x - y\|^2, \quad (20)$$

for all $t \in \mathbb{R}$, $x, y \in \mathcal{H}$ and a nonnegative continuous function $\gamma : \mathbb{R} \rightarrow \mathbb{R}^+$ satisfying $\int_T^{+\infty} \gamma(\tau) d\tau = +\infty$ for any fixed T . Then the solution to the sweeping process (13) is globally stable, i.e., $\|x(t) - y(t)\| \rightarrow 0$ for two trajectories with arbitrary initial values $x_0, y_0 \in C$ as $t \rightarrow +\infty$.

Proof. Let $x(t)$ and $y(t)$ be two solutions of (13). Consider a domain $I = [s, t]$ such that both $x(t)$ and $y(t)$ are defined and the derivatives exist. By definition of the normal cone to a convex set, we have

$$\langle u - v, x - y \rangle \geq 0, \quad \forall u \in N_C(x), v \in N_C(y). \quad (21)$$

By definition of a sweeping process, it follows that

$$-\dot{x}(t) - \psi(t, x(t)) \in N_C(x(t)), \quad (22a)$$

$$-\dot{y}(t) - \psi(t, y(t)) \in N_C(y(t)). \quad (22b)$$

Hence we have

$$-\langle \psi(t, x) - \psi(t, y), x - y \rangle \geq \langle \dot{x}(t) - \dot{y}(t), x - y \rangle. \quad (23)$$

Using the strong monotonicity condition, we obtain

$$\langle \dot{x}(t) - \dot{y}(t), x(t) - y(t) \rangle \leq -\gamma(t) \|x(t) - y(t)\|^2. \quad (24)$$

This is equivalent to

$$\frac{d}{dt} \|x(t) - y(t)\|^2 \leq -\gamma(t) \|x(t) - y(t)\|^2. \quad (25)$$

Using Gronwall's inequality, we have

$$\|x(t) - y(t)\|^2 \leq \exp\left(-\int_s^t \gamma(\tau) d\tau\right) \|x(s) - y(s)\|^2 \quad (26)$$

for all $t > s$. Consider two different initial points $x(T_0) = x_0$ and $y(T_0) = y_0$ for some fixed T_0 and $x_0, y_0 \in C$. Letting $t \rightarrow +\infty$, we obtain

$$\lim_{t \rightarrow \infty} \|x(t) - y(t)\|^2 = 0, \quad (27)$$

which completes the proof. \square

Remark. *Lemma 4 reveals that the dynamical system associated with the sweeping process has a unique ω -limit set independent of the choice of initial points under certain conditions. Furthermore, if the closed subset $C \subseteq \mathcal{H}$ is bounded, the positive semitrajectory of the associated dynamical system will be precompact, and hence the ω -limit set will be internally chain transitive [24].*

Theorem 1. *Let $(\mathcal{H}, \mathbb{R}^+, \pi)$ be the dynamical system associated with a unique global solution (under the conditions of Lemma 3) to (13). Assume that ψ is strongly monotone. If $(\mathcal{H}, \mathbb{R}^+, \pi)$ is a gradient-like dynamical system with a Lyapunov function $V : \mathcal{H} \rightarrow \mathbb{R}$, the ω -limit set $\Omega(x)$ of any point $x \in C$ for any closed and convex subset $C \subseteq \mathcal{H}$ satisfies*

$$V(\pi(t, y)) = V(y), \quad \forall y \in \Omega(x), \quad \forall t \geq 0. \quad (28)$$

Proof. Denote the non-wandering set \mathcal{J}_x^+ of $x \in \mathcal{H}$ by

$$\mathcal{J}_x^+ := \{y \in \mathcal{H} \mid \exists t_n \rightarrow \infty, x_n \rightarrow x, \text{ s.t. } \pi(t_n, x_n) \rightarrow y\}.$$

We first show that if x is contained in its own non-wandering set, i.e., $x \in \mathcal{J}_x^+$, then $V(\pi(t, x)) = V(x)$ for all $t \geq 0$. In fact, since $\mathcal{J}_x^+ \subseteq \mathcal{J}_{\pi(t, x)}^+$ for all $t \geq 0$ by definition, there exists $\tilde{x}_n \rightarrow \pi(t, x)$, $t_n \rightarrow \infty$, such that $\pi(t_n, \tilde{x}_n) \rightarrow x$. Hence it follows that

$$V(x) = \lim_{n \rightarrow \infty} V(\pi(t_n, \tilde{x}_n)) \leq \lim_{n \rightarrow \infty} V(\tilde{x}_n) = V(\pi(t, x)),$$

for all $t \geq 0$. Since $(\mathcal{H}, \mathbb{R}^+, \pi)$ is a gradient-like dynamical system, for all $t \geq 0$ and $x \in \mathcal{H}$ we have $V(\pi(t, x)) \leq V(x)$. Therefore, it can be concluded that $V(\pi(t, x)) = V(x)$. It is also sufficient to observe that $\Omega(x) \subseteq \mathcal{J}_x^+$. We can conclude that if $x \in \Omega(x)$, then $V(\pi(t, x)) = V(x)$ for all $t \geq 0$.

By Lemma 4 and the remark following the lemma, $\Omega(x) = \Omega(y)$ for all $x, y \in C$. Therefore, the ω -limit set can be denoted by Ω_C . Since C is closed in a complete space, it follows that $\Omega(x) \subseteq C$ for all $x \in C$. Hence we have $\Omega(x) = \Omega(v)$ for all $x \in C$ and $v \in \Omega_C$. For any $x \in C$ and any $u \in \Omega(x)$, we have $u \in \Omega(x) = \Omega_C = \Omega(u)$. Therefore, $V(u) = V(\pi(t, u))$ for all $t \geq 0$. \square

To look closer at the fixed point set of the Lyapunov function V , it is sufficient to take derivatives with respect to time, i.e.,

$$\frac{dV(\pi(t, x))}{dt} = \left\langle \nabla V(\pi(t, x)), \frac{d\pi(t, x)}{dt} \right\rangle = 0, \quad (29)$$

for all $t \geq 0$. Letting $t = 0$, we obtain

$$\langle \nabla V(x), \dot{x}(0) \rangle = 0. \quad (30)$$

If we consider $\psi(0, x) = \nabla V(x)$ in (13), it follows [25, p. 266, Proposition 2] that

$$\|\mathcal{P}_{\mathcal{T}_C(x)}[-\nabla V(x)]\|^2 = 0. \quad (31)$$

Hence, $0 \in \nabla V(x) + N_C(x)$. This means that x is a stationary point of the constrained optimization problem $\min_{y \in C} V(y)$.

In fact, the strong monotonicity of the time-dependent vector field $\psi(t, x)$ implicitly indicates some kind of convexity in x of the time-varying vector field. To further investigate the general case where the vector field is non-convex, it is necessary to consider the case where strong monotonicity is not satisfied. In this case, the ω -limit set is not unique compared to the conditions of Lemma 4, while it is still possible to generalize this result.

Theorem 2. *Let $(\mathcal{H}, \mathbb{R}^+, \pi)$ be the dynamical system associated with a unique global solution (under the conditions of Lemma 3) to the sweeping process (13). If $(\mathcal{H}, \mathbb{R}^+, \pi)$ is a gradient-like dynamical system with a Lyapunov function $V : \mathcal{H} \rightarrow \mathbb{R}$, the ω -limit set $\Omega(x)$ of any point $x \in C$ for any closed, **bounded** and convex subset $C \subseteq \mathcal{H}$ satisfies*

$$V(\pi(t, y)) = V(y), \quad \forall y \in \Omega(x), \forall t \geq 0. \quad (32)$$

Proof. For an arbitrary point $x \in C$, we can define a continuous function $\phi_x : \mathbb{R} \rightarrow \mathbb{R}, t \mapsto V(\pi(t, x))$. Clearly, we have $\phi_x(s) \leq \phi_x(t)$ for all $s \geq t$. Since C is compact, the positive semitrajectory of $\pi(t, x)$ is precompact and hence $V(\pi(t, x))$ is bounded. Hence $\phi_x(t)$ is a continuous bounded monotonically decreasing function of t . Therefore, there exists $\sigma_x \in \mathbb{R}$ such that $\lim_{t \rightarrow \infty} \phi_x(t) = \sigma_x$. Now consider $y \in \Omega(x)$. Then by definition, there exists $\tilde{t}_n \rightarrow \infty$ such that $\pi(\tilde{t}_n, x) \rightarrow y$. Consequently, $V(y) = \lim_{n \rightarrow \infty} V(\pi(\tilde{t}_n, x)) = \sigma_x$. This indicates that $\forall y \in \Omega(x)$, we have $V(y) = \sigma_x$. Since the ω -limit set is invariant, we have $\pi(t, y) \in \Omega(x)$ for all $t \geq 0$. It then follows that $V(\pi(t, y)) = \sigma_x = V(y)$ for all $t \geq 0$. \square

In the subsequent analysis, we consider a hybrid sweeping process which can be decomposed into two components: a constrained part and the other unconstrained component. The implicit physical intuition is directly linked with a Hamiltonian system, of which the coordinates of the particle are restricted within a compact and convex space. Formally, we consider the following hybrid sweeping process defined as

$$\begin{cases} -\frac{dx}{dt} = \phi(t, x, y), \\ -\frac{dy}{dt} \in \varphi(t, x, y) + N_{\mathcal{Y}}(y), \end{cases} \quad (33)$$

where $x \in \mathcal{H}_1$ and $y \in \mathcal{H}_2$ for two Hilbert spaces \mathcal{H}_1 and \mathcal{H}_2 . $\mathcal{Y} \subseteq \mathcal{H}_2$ is a compact and convex subset, and both $\phi : \mathbb{R} \times \mathcal{H}_1 \times \mathcal{H}_2 \rightarrow \mathcal{H}_1$ and $\varphi : \mathbb{R} \times \mathcal{H}_1 \times \mathcal{H}_2 \rightarrow \mathcal{H}_2$ satisfy the conditions of Lemma 3 by treating (x, y) together as a whole. Recall that the Cartesian product of two convex sets remains convex. Therefore, (33) has a unique solution defined on a corresponding set I by taking $\mathcal{H}_1 \times \mathcal{Y}$ in Lemma 3.

Since $\mathcal{H}_1 \times \mathcal{Y}$ is not compact, the conditions of Theorem 2 are violated. Furthermore, we usually consider the Lyapunov function only for the coordinates in practice. To address the issue, it is sufficient to consider a dynamical systems $(\mathcal{H}_1 \times$

$\mathcal{H}_2, \mathbb{R}^+, \pi)$ for (33) and a function $\mu : \mathbb{R}^+ \times \mathcal{H}_1 \times \mathcal{H}_2 \rightarrow \mathcal{H}_2$ along with a map $\alpha : \mathcal{H}_1 \times \mathcal{H}_2 \rightarrow \mathcal{H}_2, (x, y) \mapsto y$, which is a projection onto the second argument. The function μ is defined as $\mu(t, x, y) := \alpha(\pi(t, (x, y)))$ for all $t \geq 0$ and $\forall (x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$. With these settings, we have the following corollary.

Corollary 1. *Consider the dynamical system given by (33). Assume that there exists a continuous function $V : \mathcal{H}_2 \rightarrow \mathbb{R}$ such that $V(\mu(t, x, y)) \leq V(\mu(s, x, y))$ for all $s < t$ and $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$. The ω -limit set $\Omega(x, y)$ of any $(x, y) \in \mathcal{H}_1 \times \mathcal{Y}$ for any compact and convex subset $\mathcal{Y} \subseteq \mathcal{H}_2$ satisfies*

$$V(\mu(t, u, w)) = V(w), \quad \forall (u, w) \in \Omega(x, y), \forall t \geq 0. \quad (34)$$

Proof. It is sufficient to consider a continuous function $\xi_{x,y} : \mathbb{R} \rightarrow \mathbb{R}, t \mapsto V(\mu(t, x, y))$. The proof follows a direct repetition of the proof of Theorem 2. \square

Remark. *Corollary 1 establishes a useful theoretical result on constrained continuous-time dynamical systems. Nevertheless, a discrete iteration is not guaranteed to remain stable under a general discretization scheme. Therefore, it is necessary to apply integrators which preserve certain structures of the continuous-time dynamics (especially the asymptotic behavior), as will be discussed in the next subsection.*

B. Explicit Euler Scheme with Decaying Step Sizes

Without loss of generality, we assume the conditions of Lemma 3 are satisfied by the PSP (13). Therefore, a unique solution is defined for the entire nonnegative real line \mathbb{R}^+ given any initial point.

To discretize the continuous-time process, we apply a time-decaying positive step size $h_k > 0$ ($\forall k \in \mathbb{N}^+$) which satisfies

$$h_0 = 0, \quad \lim_{k \rightarrow \infty} h_k = 0, \quad \sum_{k=1}^{\infty} h_k = \infty. \quad (35)$$

Correspondingly, the numerical scheme is given by

$$\bar{z}_k = P_{\mathcal{K}}[\bar{z}_{k-1} - h_k \psi(t_{k-1}, \bar{z}_{k-1})], \quad \forall k \in \mathbb{N}, \quad (36)$$

where $\bar{z}_0 = z(0) = z_0 \in \mathcal{K}$ and $t_k = \sum_{\ell=0}^k h_{\ell}$. Recall that $x - \bar{x} \in N_{\mathcal{K}}(\bar{x})$ for all $x \in \mathcal{H}$ and $\bar{x} = P_{\mathcal{K}}(x)$. Hence the numerical scheme (36) can be viewed as

$$-\frac{\bar{z}_k - \bar{z}_{k-1}}{h_k} \in \psi(t_{k-1}, \bar{z}_{k-1}) + N_{\mathcal{K}}(\bar{z}_k), \quad \forall k \in \mathbb{N}, \quad (37)$$

which is a discrete explicit Euler scheme of the continuous-time dynamics (13) with step size $h_k > 0$ for all k . Furthermore, it is sufficient to consider a linear interpolation process $u(t)$ for estimation, i.e., for all $k \in \mathbb{N}^+$

$$u(t) = \bar{z}_{k-1} + \frac{\bar{z}_k - \bar{z}_{k-1}}{h_k}(t - t_{k-1}), \quad \forall t_{k-1} \leq t < t_k. \quad (38)$$

Let $z^s(t)$ represent the unique solution to the PSP (13) starting at s , i.e., for $z^s(s) = u(s)$

$$-\dot{z}^s(t) \in \psi(t, z^s(t)) + N_{\mathcal{K}}(z^s(t)), \quad t \geq s. \quad (39)$$

Likewise, denote by $z_s(t)$ the unique solution to the PSP (13) ending at s , i.e., for $z_s(s) = u(s)$

$$-\dot{z}_s(t) \in \psi(t, z_s(t)) + N_{\mathcal{K}}(z_s(t)), \quad t \leq s. \quad (40)$$

To derive the convergence results, the following common assumption is introduced:

Assumption 1. *The following conditions hold:*

- The sequence $\{z_n\}$ is bounded;
- The function ψ is sweeping-regular (cf. Condition 1) on $([0, t], \mathcal{K})$, $\forall t \geq 0$;
- The step size sequence $\{h_k\}$ satisfies (35);
- ψ satisfies the weak monotonicity condition:

$$\langle \psi(t, x) - \psi(t, y), x - y \rangle \geq \gamma(t) \|x - y\|^2, \quad (41)$$

for all $t \in \mathbb{R}$, $x, y \in \mathcal{H}$ and a continuous function $\gamma : \mathbb{R} \rightarrow \mathbb{R}$ satisfying $\int_s^t \gamma(\tau) d\tau > -\infty$ for any $s, t \in \mathbb{R}$, $s \leq t$;

- Bounded variations: $(\forall M > 0, \forall k \in \mathbb{N}^+)$

$$\sup_{\|z\| \leq M} \|\psi(t_k, z) - \psi(t_{k-1}, z)\| \leq B_k < \infty, \quad (42)$$

for some $B_k = B_k(M) > 0$.

We then have the following lemma:

Lemma 5. *Let Assumption 1 hold. For all $\tau > 0$,*

$$\lim_{s \rightarrow \infty} \sup_{s \leq t \leq s+\tau} \|u(t) - z^s(t)\| = 0, \quad (43a)$$

$$\lim_{s \rightarrow \infty} \sup_{s-\tau \leq t \leq s} \|u(t) - z_s(t)\| = 0. \quad (43b)$$

Proof. It is sufficient to prove the claim for $z^s(t)$ as arguments for the other claim are completely analogous. Furthermore, if the following alternative claim:

$$\lim_{\ell \rightarrow \infty} \sup_{t_\ell \leq t \leq t_\ell + \tau} \|u(t) - z^{t_\ell}(t)\| = 0, \quad \forall \tau > 0, \quad (44)$$

holds, the other direction holds by analogy. Then, for all $s > 0$, there exists some sufficiently large $\ell > 0$ such that $t_\ell \leq s < s + \tau \leq t_\ell + T_s$ for some $T_s > 0$ and

$$\sup_{s \leq t \leq s+\tau} \|u(t) - z^s(t)\| \leq \sup_{t_\ell \leq t \leq t_\ell + T_s} \|u(t) - z^{t_\ell}(t)\|. \quad (45)$$

The desired result will be obtained by taking limit.

To begin with, we first show that $\dot{u}(t)$ is bounded in $[t_k, t_k + \tau]$ for all $\tau > 0$ and $k \in \mathbb{N}$. Without loss of generality, we assume that $N = N(\tau) = \sup\{m : t_m \leq \tau\} \geq k + 1$. Using the numerical scheme (37), we obtain

$$-v_k - \psi(t_k, \bar{z}_k) \in N_{\mathcal{K}}(\bar{z}_{k+1}), \quad \forall k \in \mathbb{N}, \quad (46)$$

where $v_k = (\bar{z}_{k+1} - \bar{z}_k)/h_{k+1}$. Applying the geometric characteristics of normal cones and making difference between v_ℓ and $v_{\ell-1}$, we find that

$$\langle v_\ell - v_{\ell-1}, v_\ell \rangle \leq -\langle \psi(t_\ell, \bar{z}_\ell) - \psi(t_{\ell-1}, \bar{z}_{\ell-1}), v_\ell \rangle. \quad (47)$$

By Assumption 1, it follows that

$$\|\psi(t_\ell, \bar{z}_\ell) - \psi(t_{\ell-1}, \bar{z}_{\ell-1})\| \leq S_\ell, \quad (48)$$

for some $S_\ell > 0$, and

$$\|\psi(t_{\ell-1}, \bar{z}_\ell) - \psi(t_{\ell-1}, \bar{z}_{\ell-1})\| \leq h_\ell L_\eta(t_{\ell-1}) \|v_{\ell-1}\|. \quad (49)$$

Using the arithmetic mean inequality, i.e.,

$$ab \leq \frac{1}{2}(b^2c + a^2/c), \quad \forall a, b \in \mathbb{R}, c > 0, \quad (50)$$

and taking some $0 < \varepsilon < 1$, we conclude that

$$(1 - \varepsilon) \|v_\ell\|^2 \leq \frac{1 + (L_\eta(t_{\ell-1}))^2 h_\ell^2}{2\varepsilon} \|v_{\ell-1}\|^2 + \frac{S_\ell^2}{\varepsilon}. \quad (51)$$

Letting $\varepsilon = 1/2$, we have for all $\ell \geq 0$

$$\|v_\ell\|^2 \leq 2[1 + (L_\eta(t_{\ell-1}))^2 \bar{h}^2] \|v_{\ell-1}\|^2 + 2S_\ell^2, \quad (52)$$

where $\bar{h} = \sup_{k \in \mathbb{N}} h_k$ is an upper bound of $\{h_k\}$. By the discrete Gronwall inequality, it follows that for all $\ell > k$

$$\|v_\ell\|^2 \leq (2 + 2L^2 \bar{h}^2)^{\ell-k} \|v_k\|^2 + 2S_\ell^2 \sum_{m=k}^{\ell} (2 + 2L^2 \bar{h}^2)^{\ell-m},$$

for some $L > 0$ due to the integrability of $L_\eta(t)$. Therefore, we conclude that for any fixed $\tau > 0$, v_ℓ is bounded for all $\ell \leq N$ and $\dot{u}(t)$ is bounded as a direct result.

Next we estimate $\|u(t) - z^{t_\ell}(t)\|$. Let $t_\ell \leq t < t_{\ell+1}$. It is clear that we have the following truncated dynamics:

$$-\dot{u}(t) - \psi(t_\ell, \bar{z}_\ell) \in N_{\mathcal{K}}(\bar{z}_{\ell+1}), \quad (53a)$$

$$-z^{t_\ell}(t) - \psi(t, z^{t_\ell}(t)) \in N_{\mathcal{K}}(z^{t_\ell}(t)). \quad (53b)$$

Applying the geometric characteristics of normal cones, it is straightforward to conclude that

$$\frac{1}{2} \frac{d}{dt} \|u(t) - z^{t_\ell}(t)\|^2 \leq -\langle \psi(t_\ell, \bar{z}_\ell) - \psi(t, z^{t_\ell}(t)), u(t) - z^{t_\ell}(t) \rangle. \quad (54)$$

It follows from the boundedness of $\dot{u}(t)$ that

$$\|u(t) - \bar{z}_\ell\| = \left\| \frac{\bar{z}_{\ell+1} - \bar{z}_\ell}{h_{\ell+1}} (t - t_\ell) \right\| \leq \|\dot{u}(t)\| h_{\ell+1} \leq M h_{\ell+1}.$$

Therefore, we have for some $K > 0$ by Assumption 1

$$\begin{aligned} & \|\psi(t, u(t)) - \psi(t_\ell, \bar{z}_\ell)\| \\ & \leq \|\psi(t, u(t)) - \psi(t_\ell, u(t))\| + \|\psi(t_\ell, u(t)) - \psi(t_\ell, \bar{z}_\ell)\| \\ & = S_\ell + K h_{\ell+1}. \end{aligned}$$

Recall that ψ satisfies the weak monotonicity condition (41). Estimating the right-hand side of (54), we conclude that

$$\begin{aligned} & -\langle \psi(t_\ell, \bar{z}_\ell) - \psi(t, z^{t_\ell}(t)), u(t) - z^{t_\ell}(t) \rangle \\ & \leq \frac{(S_\ell + K h_{\ell+1})^2}{2} - \frac{1}{2} (\gamma(t) - 1) \|u(t) - z^{t_\ell}(t)\|^2, \end{aligned} \quad (55)$$

Then (54) can be written as

$$\begin{aligned} & \frac{d}{dt} \|u(t) - z^{t_\ell}(t)\|^2 \\ & \leq -(\gamma(t) - 1) \|u(t) - z^{t_\ell}(t)\|^2 + (S_\ell + K h_{\ell+1})^2. \end{aligned} \quad (56)$$

Using Gronwall's inequality and $u(t_\ell) = z^{t_\ell}(t_\ell)$, we obtain

$$\|u(t) - z^{t_\ell}(t)\|^2 \leq \sum_{k=\ell+1}^{N(\tau)} (S_k + K h_{k+1})^2 h_{k+1}. \quad (57)$$

Since $h_k \rightarrow 0$ as $k \rightarrow \infty$, it follows that

$$\lim_{\ell \rightarrow \infty} \sup_{t_\ell \leq t \leq t_\ell + \tau} \|u(t) - z^{t_\ell}(t)\|^2 = 0, \quad (58)$$

which completes the proof. \square

Based on this lemma, via a straightforward application of [26, p. 17, Theorem 2.1], we obtain the desired convergence result as follows:

Theorem 3. *Under Assumption 1, the sequence $\{\bar{z}_n\}$ generated by (36) converges to a connected internally chain transitive invariant set of (13).*

In general, Theorem 3 is the best result one can obtain on convergence of the numerical scheme (37) corresponding to a PSP. Unfortunately, the result presented in Corollary 1 for a continuous-time hybrid sweeping process cannot be simply extended to the numerical case. The primary obstacle lies in the unboundedness of the trajectory as discussed in [13]. Besides, some alternatives for the assumption that $\{\bar{z}_k\}$ is bounded are provided in [26, Chap. 4]. Furthermore, the following corollary is immediate.

Corollary 2. *If the only internally chain transitive invariant sets for (13) are isolated critical points, then $\{z_n\}$ converges to a critical point under Assumption 1.*

In the previous subsection, we have characterized the ω -limit set of the continuous-time sweeping process on a compact and convex subset given the existence of a Lyapunov function. The question is whether this result can be extended to the numerical case. Such extensions are never straightforward since the ω -limit set of $u(t)$ only coincides with an internally chain transitive set of (13) as presented in Theorem 3. Although the ω -limit set of any precompact positive orbit with respect to a continuous semiflow is internally chain transitive [24, Lemma 2.1'], the opposite is not true in general. Fortunately, by introducing the concept of Lyapunov functions for a PSP, we can obtain a similar conclusion to that of the continuous-time case.

Corollary 3. *Let $\mathcal{L} \subset \mathbb{R}^m$ be a nonempty compact set, $U \subset \mathcal{K} \subset \mathbb{R}^m$ be a bounded open neighborhood of \mathcal{L} , and $V : \mathcal{K} \rightarrow \mathbb{R}^+$ be continuously differentiable. Let the following hold:*

- $u(t) \in U$ for all $t \geq 0$;
- $V^{-1}(0) = \mathcal{L}$;
- The Lie derivative $\frac{dV}{dt} \leq 0$ along (13) holds for all $t \geq 0$ and $x \in \mathcal{K}$ with equality if and only if $x \in \mathcal{L}$.

Then $\{z_n\}$ converges to an internally chain transitive set contained in \mathcal{L} under Assumption 1.

Proof. Note that the corollary is inspired by [26, p. 19, Corollary 2.1]; we reproduce the proof for the sake of completeness. Let $M = \sup_n \|z_n\| < \infty$ and $C = \sup_{\|z\| \leq M} V(z)$. For any constant $0 < b \leq C$, we define $Z^b := \{x \in U : V(x) < b\}$. For $0 < \epsilon < C/2$, we have

$$-\zeta := \sup_{t \geq 0, x \in \bar{Z}^C \setminus Z^\epsilon} \frac{dV}{dt}(t, x) < 0, \quad (59)$$

where \bar{Z}^C denotes the closure of Z^C . It then follows that

$$V(z(t)) = V(z(0)) + \int_0^t \frac{dV}{ds}(s, z(s)) ds \leq V(z(0)) - t\zeta.$$

Let τ be an upper bound on the time required for a solution to (13) starting from \bar{Z}^C to reach Z^ϵ . Hence, we can pick $C/\zeta < \tau < \infty$. Since \mathcal{K} is compact and V is continuously differentiable, V is Lipschitz continuous in \mathcal{K} . Then there exists some $\delta > 0$ such that for all $x \in \bar{Z}^C$ and $y \in \mathcal{K}$ with $\|x - y\| < \delta$, we have $|V(x) - V(y)| < \epsilon$. By

Lemma 5, there exists t_0 such that for all $t \geq t_0$, we have $\sup_{t \leq s \leq t+\tau} \|u(s) - z^t(s)\| < \delta$. Since $u(s) \in \bar{Z}^C$, it follows that $|\bar{V}(u(t+\tau)) - V(z^t(t+\tau))| < \epsilon$, and hence $u(t+\tau) \in Z^{2\epsilon}$ for $z^t(t+\tau) \in Z^\epsilon$. Therefore, $u(t) \in Z^{2\epsilon}$ for all $t \geq t_0 + \tau$. Letting $\epsilon \downarrow 0$, we have $u(t) \rightarrow \mathcal{L}$ as $t \rightarrow \infty$. \square

Although this corollary provides a useful tool to address optimization problems in smooth analysis, it fails to apply to composite optimization problem where V is nonsmooth. By contrast, the subsequent theorem offers a general framework.

Theorem 4. *Let $\Lambda \subset \mathbb{R}^m$ be any subset. Suppose that $V : \mathbb{R}^m \rightarrow \mathbb{R}$ is a Lyapunov function for Λ with respect to the trajectory of (13). Assume that $V(\Lambda)$ has empty interior. Then $\{z_n\}$ converges to an internally chain transitive set \mathcal{L} contained in Λ under Assumption 1 and V is constant in \mathcal{L} .*

Proof. The results follow from [27, Proposition 3.27] and Theorem 3. \square

In terms of the hybrid sweeping process (33), it is sufficient to let $\mathcal{K} = \mathcal{H}_1 \times \mathcal{C}$ for a compact and convex subset $\mathcal{C} \subset \mathcal{H}_2$. In this case, by Lemma 2 it follows that for all $t \in [T_s, T_e]$ where a solution is defined

$$\begin{aligned} \|p(t)\| &\leq \|p(T_s)\| + \int_{T_s}^{T_e} \|\phi(s, p(s), q(s))\| ds \\ &\leq \|p(T_s)\| + D \int_{T_s}^{T_e} \beta(\tau) d\tau < \infty, \end{aligned} \quad (60)$$

which indicates that the momentum $p(t)$ is uniformly bounded, and hence $z(t) = (p(t), q(t))$ lies in a fixed compact ball for all $t \in [T_s, T_e]$. Therefore, the first item in Assumption 1 can be replaced by compactness of \mathcal{C} .

Indeed, the conclusions regarding the numerical scheme, which are derived from an initial continuous-time PSP, can be viewed from the reversed direction. Specifically, given a numerical scheme

$$z_{k+1} = P_{\mathcal{K}}[z_k + h_{k+1}\phi_k(z_k)], \quad (61)$$

along with its corresponding continuous-time dynamics

$$-\dot{z}(t) \in \psi(t, z) + N_{\mathcal{K}}(z), \quad (62)$$

where $\psi(t_k, z) = \phi_k(z)$ for all $k \in \mathbb{N}$ and $z \in \mathcal{H}$, it follows that the aforementioned conclusions still hold.

Remark. *Consider a perturbed stochastic scheme, i.e.,*

$$\tilde{z}_{k+1} = P_{\mathcal{K}}[\tilde{z}_k - h_{k+1}\psi(t_k, \tilde{z}_k) - h_{k+1}(U_{k+1} + r_{k+1})], \quad (63)$$

where $\{U_k\}$ and $\{r_k\}$ are sequences of random perturbations. Assume that for each $T > 0$,

$$\lim_{n \rightarrow \infty} \sup_{\{k: 0 \leq t_k - t_n \leq T\}} \left\| \sum_{\ell=n}^{k-1} h_\ell U_\ell \right\| = 0, \text{ a.s.}, \quad (64)$$

and $\lim_{k \rightarrow \infty} r_k = 0$ a.s. Then the conclusions and corresponding analysis above for a deterministic scheme hold almost surely for the stochastic scheme, following standard analysis of the classical result [13].

Via straightforward application of this remark, we immediately obtain the following useful theorem:

Theorem 5. Consider an asymptotic numerical scheme

$$\bar{z}_{k+1} = P_{\mathcal{K}}[\phi_k(\bar{z}_k) - h_{k+1}(\psi(t_k, \bar{z}_k) + \xi_{k+1})], \quad (65)$$

where $\phi_k : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is continuous for all k , and $\{\xi_k\}$ is a sequence of random perturbations satisfying (64). Let Assumption 1 hold. Assume that

$$\lim_{n \rightarrow \infty} \sup_{x \in \mathbb{R}^m} \|\phi_n(x) - x\| = 0. \quad (66)$$

The conclusion of Theorem 4 holds for (65).

C. Lyapunov Pairs

The stability analysis of the nonsmooth dynamics of the PSP naturally requires nonsmooth Lyapunov functions. In this subsection, we introduce a useful theoretical result of nonsmooth Lyapunov analysis.

Definition 1 (A variant of Definition 1 in [28]). Let \mathcal{H} be a Hilbert space. Let functions $V, W : \mathbb{R} \times \mathcal{H} \rightarrow \mathbb{R}$ be lower semi-continuous, with $W \geq 0$. We say that (V, W) is a time-dependent Lyapunov pair for a dynamical system (X, \mathbb{R}^+, π) ($X \subseteq \mathcal{H}$) if for all $x_0 \in X$ and $\forall t \geq 0$,

$$V(t, x(t)) + \int_0^t W(\tau, x(\tau)) d\tau \leq V(0, x_0), \quad (67)$$

where $x(t) = \pi(t, x_0)$.

Identifying a suitable Lyapunov pair for nonsmooth dynamical systems is inherently complex, primarily due to the difficulty in determining the supremum of the Lie derivatives of potential Lyapunov functions. The challenge arises from the requirement to evaluate the supremum within the context of the Fréchet subdifferential, which encapsulates a broader set of candidates than the traditional derivative would allow.

Fortunately, the following theorem provides a powerful tool to settle the problem for a PSP as (13).

Theorem 6. Let \mathcal{H} be a Hilbert space. Let functions $V, W : \mathbb{R} \times \mathcal{H} \rightarrow \mathbb{R}$ be lower semi-continuous, with $W \geq 0$. (V, W) is a time-dependent Lyapunov pair if and only if for all $t \geq 0$, $x \in \mathcal{H}$ and $\xi \in \partial_F V(t, x)$, we have

$$\min_{v \in N_C(x) \cap \mathbb{B}(0, \|\psi(t, x)\|)} \langle \xi, -\psi(t, x) - v \rangle + W(t, x) \leq 0. \quad (68)$$

Proof. A combination of [29, Theorem 5.1] and [25, pp. 300-301, Proposition 5]. \square

IV. APPLICATION TO OPTIMIZATION

In this section, some examples are provided of convergence analysis of popular optimization methods by applying the above established theoretical results.

A. Centralized Optimization

We begin with the variants of two popular symplectic methods for unconstrained optimization. As demonstrated in Assumption 1, the time-varying vector field is not restricted to be continuous with respect to time (the first argument). Therefore, it is feasible to add countable bounded jump discontinuities to the vector field. This, in turn, supports numerical

schemes with a constant step size as we can add a cofactor to cancel the vanishing step size.

Example 1: Nesterov accelerated gradient (NAG). Consider the unconstrained dynamics given by

$$\begin{cases} \frac{dp}{dt} = \kappa(t)((\mu - 1)p - \gamma \nabla f(q + \mu p)), \\ \frac{dq}{dt} = \kappa(t)(\mu p - \gamma \nabla f(q + \mu p)), \end{cases} \quad (69)$$

where $\kappa(t)$ is defined by $\kappa(t) := \sup\{k + 1 : k \in \mathbb{N}, \tau_k \leq t\}$, $\tau_k = \sum_{\ell=0}^k \ell$ for all $k \in \mathbb{N}$ and $\gamma > 0, 0 < \mu < 1$ are positive constants, and $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is lower-bounded and has a Lipschitz-continuous gradient. Here we select \mathbb{R}^m as the constraint set in the perturbed sweeping process. Applying the explicit Euler scheme with step size $h_k = 1/k$, we have

$$\begin{cases} p_{n+1} = \mu p_n - \gamma \nabla f(q_n + \mu p_n), \\ q_{n+1} = q_n + p_{n+1}. \end{cases} \quad (70)$$

Assume that $\{p_n\}$ and $\{q_n\}$ are bounded. It is straightforward to verify that such dynamics and the numerical scheme satisfy the conditions of Lemma 3 and Assumption 1. In fact, it is the well-known Nesterov accelerated gradient method. An application of Theorem 4 yields the following result:

Corollary 4. Denote the critical point set of f by A . Assume that $f(A)$ has empty interior. The numerical scheme (70) converges to A (in terms of q_n).

Proof. To show convergence, it is sufficient to introduce a Lyapunov function $V : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}$, $V(p, q) = \frac{1}{2}\|p\|^2 + \frac{\gamma}{\mu^2}f(q + \mu p)$. Taking derivatives with respect to t , we have

$$\frac{dV}{dt} = \kappa(t) \left[(\mu - 1)\|p\|^2 - \frac{\gamma^2}{\mu^2}(1 + \mu)\|\nabla f(q + \mu p)\|^2 \right] \leq 0.$$

Denote $\Lambda = \{(p, q) : p = 0, \nabla f(q + \mu p) = 0\}$ by the critical point set of V . Clearly, $V(\Lambda)$ has empty interior by assumption. By Theorem 4, the proof is complete. \square

Remark. To be specific, (70) can be obtained from the following more familiar form:

$$\begin{cases} x_{n+1} = y_n - \gamma \nabla f(y_n), \\ y_{n+1} = x_{n+1} + \mu(x_{n+1} - x_n), \end{cases} \quad (71)$$

by letting $q_n = x_n$ and $p_n = x_n - x_{n-1}$ and writing the updates in terms of q_n and p_n .

In general, it is not natural to obtain convergence results for momentum based methods by Theorem 4 since the corresponding continuous-time dynamics is linked with a second-order ODE according to Polyak's pioneering work [30].

Example 2: Relativistic gradient descent (RGD). Consider the dynamics given by

$$\begin{cases} \frac{dx}{dt} = \frac{cp}{\sqrt{\|p\|^2 + m^2c^2}}, \\ \frac{dp}{dt} = -\nabla f(x) - \gamma p, \end{cases} \quad (72)$$

where c denotes the speed of light, m denotes the mass of the particle, $\gamma > 0$, and the other parameters follow the settings in Example 1. The numerical scheme is given by

$$\begin{cases} p_{k+1} = (1 - \gamma h_{k+1})p_k - h_{k+1}\nabla f(x_k), \\ x_{k+1} = x_k + \frac{h_{k+1}cp_k}{\sqrt{\|p_k\|^2 + m^2c^2}}. \end{cases} \quad (73)$$

This scheme is a variant of relativistic gradient descent [31] with decaying step sizes. Its convergence can be established by selecting the Lyapunov function V as

$$V(p, x) = c\sqrt{\|p\|^2 + m^2c^2} + f(x). \quad (74)$$

Then it follows that

$$\frac{dV}{dt} = -\frac{\gamma c}{\sqrt{\|p\|^2 + m^2c^2}}\|p\|^2 \leq 0. \quad (75)$$

Consider the set $\Lambda = \{(0, x) : \nabla f(x) = 0\}$. Clearly, Λ is the critical point set of V and V is a Lyapunov function for Λ . Assume that $V(\Lambda)$ has empty interior. By Theorem 5, $\{x_k\}$ in (73) converges to the critical point set of f .

Example 3: Stochastic projected gradient descent (PGD). Consider the dynamics given by

$$\frac{dz}{dt} \in -\nabla f(z) - N_{\mathcal{C}}(z(t)), \quad (76)$$

where $\mathcal{C} \subset \mathbb{R}^m$ is a compact convex set, $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is lower-bounded and has a Lipschitz-continuous gradient. Taking the step size $h_k = 1/k$, we have for a martingale difference noise sequence $\{\xi_k\}$ with bounded variance:

$$z_{n+1} = P_{\mathcal{C}}[z_n - h_{n+1}(\nabla f(z_n) + \xi_{n+1})], \quad (77)$$

which is the classical form of a stochastic projected gradient descent method. Its convergence can be established immediately by selecting f as the Lyapunov function and using the facts from [22, Proposition 2] according to Theorem 4.

Example 4-A: Projected Nesterov accelerated gradient (PNAG). Consider the constrained dynamics given by

$$\begin{cases} \frac{dp}{dt} = \kappa(t)[(\mu - 1)p - \gamma\nabla f(q + \mu p)], \\ \frac{dq}{dt} \in \kappa(t)[\mu p - \gamma\nabla f(q + \mu p) - N_C(q)], \end{cases} \quad (78)$$

where $C \subset \mathbb{R}^m$ is a closed and convex subset and the other parameters adopt the same settings as Example 1. The corresponding Euler discretization yields

$$\begin{cases} p_{n+1} = \mu p_n - \gamma\nabla f(q_n + \mu p_n), \\ q_{n+1} = P_C[q_n + p_{n+1}]. \end{cases} \quad (79)$$

Then we have the following corollary which coincides with the existing result in [32].

Corollary 5. *Let \mathcal{L} be the set of KKT points of f on C associated with the constrained optimization problem $\min_{x \in C} f(x)$. If $f(\mathcal{L})$ has empty interior, the numerical scheme (q_n) converges to \mathcal{L} .*

Proof. Denote the composite vector field $\psi(t, p, q)$ by

$$\psi(t, p, q) := \begin{pmatrix} \kappa(t)[(\mu - 1)p - \gamma\nabla f(q + \mu p)] \\ \kappa(t)[\mu p - \gamma\nabla f(q + \mu p)] \end{pmatrix}. \quad (80)$$

Take the differentiable function $V : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}$ as

$$V(p, q) = \frac{1}{2}\|p\|^2 + \frac{\gamma}{\mu^2}f(q + \mu p). \quad (81)$$

Take $W(t, p, q) = \kappa(t)((1 - \mu)\|p\|^2 + R(p, q))$ for

$$R(p, q) := \frac{\gamma^2}{\mu^2}(1 + \mu)\langle -\nabla f(q + \mu p) - v_0, -\nabla f(q + \mu p) \rangle, \quad (82)$$

where $v_0 = P_{N_C(q)}[-\nabla f(q + \mu p)]$. Therefore, we have

$$R(p, q) = \frac{\gamma^2}{\mu^2}(1 + \mu)\|P_{\mathcal{T}_C(q)}[-\nabla f(q + \mu p)]\|^2. \quad (83)$$

Compared to Example 1, the primary difference lies in the terms imposed by the nonsmooth part, i.e., the vectors in normal cones. In fact, the normal cone term in the nonsmooth dynamics can be cancelled by Theorem 6. To be specific, for $\xi = \nabla V(p, q)$ and $\forall v \in N_C(q)$,

$$\begin{aligned} & \langle \xi, -\psi(t, p, q) - (0, v)^T \rangle + (1 - \mu)\|p\|^2 \\ &= \kappa(t) \left[-R(p, q) - \left\langle v - \gamma(1 + \mu)v_0, \frac{\gamma}{\mu^2}\nabla f(q + \mu p) \right\rangle \right] \\ &= \kappa(t) \left[-R(p, q) - \frac{\gamma}{\mu^2}\langle v, \nabla f(q + \mu p) \rangle - \frac{\gamma^2}{\mu^2}(1 + \mu)\|v_0\|^2 \right], \end{aligned}$$

where we have used the fact that any vector x can be decomposed into $x_1 + x_2$ for $x_1 \in N_C(q)$ and $x_2 \in \mathcal{T}_C(q)$ and $\langle x_1, x_2 \rangle = 0$. Since $0 \in N_C(q)$, it follows that

$$\min_{\|v\| \leq \|\psi(t, p, q)\|} \langle \xi, -\psi(t, p, q) - (0, v)^T \rangle + W(t, p, q) \leq 0. \quad (84)$$

By Theorem 6, (V, W) is a Lyapunov pair. Observe that only if $W = 0$, $V(p(t), q(t)) \leq V(p(s), q(s))$ for all $t > s$. Therefore, it is sufficient to consider the set $\Lambda = \{(p, q) : p = 0, P_{\mathcal{T}_C(q)}[-\nabla f(q + \mu p)] = 0, q \in C\}$ such that $W(\Lambda) = 0$. It is clear that $\Lambda|_q$ coincides with the KKT point set of the constrained problem $\min_{x \in C} f(x)$. By assumption, Λ is nonempty and $V(\Lambda)$ has empty interior. Clearly, V is a Lyapunov function for Λ by definition and the desired result can be obtained via Theorem 4. \square

Example 4-B: PNAG (the other form). Consider the following perturbed sweeping process:

$$\begin{cases} \frac{dx}{dt} = \kappa(t)[y - x - \gamma\nabla f(y)], \\ \frac{dy}{dt} \in \kappa(t)[\mu y - \mu x - \nu\nabla f(y) - N_C(y)], \end{cases} \quad (85)$$

where $\nu = \gamma(1 + \mu)$. The parameters are set in accordance with Example 4-A. The Euler discretization produces

$$\begin{cases} x_{n+1} = y_n - \gamma\nabla f(y_n), \\ y_{n+1} = P_C[x_{n+1} + \mu(x_{n+1} - x_n)]. \end{cases} \quad (86)$$

Corollary 6. *Let \mathcal{L} be the set of KKT points of f on C associated with the constrained optimization problem $\min_{x \in C} f(x)$. If $f(\mathcal{L})$ has empty interior, (y_n) of the numerical scheme (86) converges to \mathcal{L} .*

Proof. Denote the composite vector field $\psi(t, p, q)$ by

$$\psi(t, x, y) := \begin{pmatrix} \kappa(t)[y - x - \gamma\nabla f(y)] \\ \kappa(t)[\mu y - \mu x - \nu\nabla f(y)] \end{pmatrix}. \quad (87)$$

Consider the differentiable function $V : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}$ as

$$V(x, y) = \frac{1}{2} \|x - y\|^2 + \gamma f(y). \quad (88)$$

Take $W(t, x, y) = \kappa(t)((1 - \mu)\|x - y\|^2 + R(y))$ for

$$\begin{aligned} R(y) &= \gamma \nu \langle -\nabla f(y) - u_0, -\nabla f(y) \rangle \\ &= \gamma \nu \|P_{\mathcal{T}_C(y)}[-\nabla f(y)]\|^2, \end{aligned} \quad (89)$$

where $u_0 = P_{N_C(y)}[-\nabla f(y)]$. For all $u \in N_C(y)$,

$$\begin{aligned} \langle \nabla V(x, y), -\psi(t, x, y) - (0, u)^T \rangle &= -W(t, x, y) \\ &\quad - \gamma \mu \|u_0\|^2 - \langle y - x + \gamma \nabla f(y), u \rangle. \end{aligned} \quad (90)$$

Since $0 \in N_C(y)$, we conclude that for $\zeta = \nabla V(x, y)$

$$\min_{\|u\| \leq \psi(t, x, y)} \langle \zeta, -\psi(t, x, y) - (0, u)^T \rangle + W(t, x, y) \leq 0. \quad (91)$$

The rest of the proof follows. \square

Remark. In practice, NAG is often utilized with time-varying step sizes, implying that the parameter μ_k evolves throughout the iterative process. As a consequence, the corresponding continuous-time model must be formulated to account for this temporal variability. Notably, the analysis presented herein remains valid in this context, given that the Lyapunov function V is not explicitly dependent on time, i.e., $\partial V / \partial t = 0$. However, this is not generally the case as will be discussed in the next example.

Example 5: Projected optimized gradient (POGM). Consider the following dynamical system:

$$\begin{cases} \frac{dx}{dt} = \kappa(t)[y - x - \gamma \nabla f(y)], \\ \frac{dy}{dt} \in \kappa(t)[\mu(t)y - \mu(t)x - \beta(t)\nabla f(y) - N_C(y)], \end{cases} \quad (92)$$

where $\beta(t) = \gamma(1 + \mu(t) + \lambda(t))$ and the time-varying step sizes are defined as for all $n \in \mathbb{N}$ and $t_n \leq t < t_{n+1}$

$$1 > \mu(t) = \mu(t_n) = \mu_n > 0, \quad \lambda(t) = \lambda(t_n) = \lambda_n > 0, \quad (93)$$

where $t_n = \sum_{\ell=0}^n h_\ell$. The other parameters are set according to Example 1. By Euler discretization, we obtain the following projected variant of the optimized gradient method [33]:

$$\begin{aligned} x_{n+1} &= y_n - \gamma \nabla f(y_n), \\ y_{n+1} &= P_C[x_{n+1} + \mu_{n+1}(x_{n+1} - x_n) - \gamma \lambda_{n+1} \nabla f(y_n)], \end{aligned} \quad (94)$$

Letting λ_k / μ_k decrease with respect to k , we have the following convergence result:

Corollary 7. Let \mathcal{L} be the set of KKT points of f on C associated with the constrained optimization problem $\min_{x \in C} f(x)$. If $f(\mathcal{L})$ has empty interior, (y_n) of the numerical scheme (94) converges to \mathcal{L} .

Proof. Denote the composite vector field $\psi(t, p, q)$ by

$$\psi(t, x, y) := \begin{pmatrix} \kappa(t)[y - x - \gamma \nabla f(y)] \\ \kappa(t)[\mu(t)y - \mu(t)x - \beta(t)\nabla f(y)] \end{pmatrix}. \quad (95)$$

Consider the function $V : \mathbb{R} \times \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}$ as

$$V(t, x, y) = \frac{1}{2} \|x - y\|^2 + \gamma \left(1 + \frac{\lambda(t)}{\mu(t)}\right) f(y). \quad (96)$$

Recall that V is piecewisely explicitly independent of time in the interval $[t_k, t_{k+1})$ for all $k \geq 0$. Since λ_k / μ_k decreases with respect to k , it is sufficient to discuss the variation of V piecewisely. For all $t \in [t_k, t_{k+1})$, we have

$$V(t, x, y) = V_k(x, y) = \frac{1}{2} \|x - y\|^2 + \sigma_k f(y), \quad (97)$$

where $\sigma_k := \gamma(1 + \lambda_k / \mu_k)$. Take $W(t, x, y) = \kappa(t)((1 - \mu(t))\|x - y\|^2 + R(y))$ for

$$\begin{aligned} R(y) &= \sigma_k \beta_k \langle -\nabla f(y) - w_0, -\nabla f(y) \rangle \\ &= \sigma_k \beta_k \|P_{\mathcal{T}_C(y)}[-\nabla f(y)]\|^2, \end{aligned} \quad (98)$$

where $w_0 = P_{N_C(y)}[-\nabla f(y)]$. For all $w \in N_C(y)$,

$$\begin{aligned} \langle \nabla V_k(x, y), -\psi(t, x, y) - (0, w)^T \rangle &= -W(t, x, y) \\ &\quad - \sigma_k \beta_k \|w_0\|^2 - \langle y - x + \sigma_k \nabla f(y), w \rangle. \end{aligned} \quad (99)$$

Since $0 \in N_C(y)$, we conclude that for $\zeta = \nabla V_k(x, y)$

$$\min_{\|w\| \leq \psi(t, x, y)} \langle \zeta, -\psi(t, x, y) - (0, w)^T \rangle + W(t, x, y) \leq 0. \quad (100)$$

The rest of the proof follows. \square

B. Decentralized Optimization

If asymptotic consensus can be achieved, an unconstrained decentralized optimization scheme can be viewed as a combination of a centralized vector (the consensus vector) and a vanishing random perturbation (the consensus error or the random shuffling). Therefore, such a method is intentionally a variant of a centralized stochastic approximation scheme.

The primary obstacle in analyzing constrained decentralized optimization methods results from the nonsmooth part of the iteration of the consensus vector. Consequently, it is significant to associate the iteration with a projected stochastic approximation scheme, of which the convergence can be established with the theoretical results in the previous section.

We take the classical multiagent projected method presented in [34] for example. Under certain assumptions on time-varying weighted graphs, the multiagent projected method can be transformed into the following projected stochastic approximation scheme in terms of the consensus vector θ_k :

$$\theta_{k+1} = P_{\mathcal{E}}[\theta_k - \gamma_{k+1} \nabla f(\theta_k) + \gamma_{k+1}(\xi_{k+1} + r_{k+1})], \quad (101)$$

where $\{\xi_k\}$ and $\{r_k\}$ are random perturbations satisfying

$$\limsup_{k \rightarrow \infty} \sup_{\ell \geq k} \left| \sum_{n=k}^{\ell} \gamma_n \xi_n \right| = 0, \quad \lim_{k \rightarrow \infty} r_k = 0, \quad \text{a.s.}, \quad (102)$$

and $\{\gamma_k\}$ is the positive time-varying step size such that $\sum_k \gamma_k = \infty$ and $\sum_k \gamma_k^2 < \infty$. Let \mathcal{E} be a nonempty convex and compact subset in \mathbb{R}^m and \mathcal{L} be the KKT point set of f on \mathcal{E} . Assume that $f(\mathcal{L})$ has empty interior. The convergence of this scheme can be immediately established by Theorem 5.




Fig. 1. Results of constrained convex optimization.

V. NUMERICAL SIMULATION

In this section, we provide the outcomes of the numerical simulations, which serve to validate the theoretical convergence analysis presented in Section IV. The convergence behavior of the algorithms is demonstrated under randomized initial conditions, highlighting the stability of the methods and the effectiveness of the developed theoretical results.

Specifically, we examine the standard PGD, PNAG with fixed parameters (FPNAG), PNAG with tuned time-dependent parameters (PNAG) and POGM. For PGD, we use vanishing step sizes as $h_k = 1/(k+1)$. For FPNAG, we let $\gamma = 0.1$ and $\mu = 0.5$. The parameters of PNAG and POGM follow the standard treatment as the unconstrained versions in the original articles (see [33], [35] for more detail).

We begin with a classical constrained convex optimization problem formulated as

$$\min_{x \in \mathcal{D}} f(x) = \frac{1}{2} \sum_{i=1}^M \|x - a_i\|^2, \quad (103)$$

where $\mathcal{D} = [-1, 1]^M$, $M = 10$ and each independent a_i is randomly selected from a uniform distribution $U(-1, 1)$. Since each $a_i \in [-1, 1]$, it is straightforward to conclude that the critical point must lie in the interior of \mathcal{D} . Hence, the KKT point x^* must satisfy $\nabla f(x^*) = 0$. This, indeed, matches the simulation result presented in Fig. 1. Moreover, since both NAG and OGM are momentum-based methods, they inherently possess a convergence rate $O(1/k^2)$ compared to PGD with convergence rate $O(1/k)$. Further, the result implies that projection will not slow down the convergence rate, which can be deduced from the nonexpansiveness of projection in a geometric perspective.

Next we consider a smooth nonconvex function called six-hump camel back function given by

$$g(x, y) = (4 - 2.1x^2 + x^{4/3})x^2 + xy + (-4 + 4y^2)y^2. \quad (104)$$

As presented in Fig. 2, the function has a global minimum $f^* = -1.0316$ for $(x^*, y^*) = \pm(0.0898, -0.7126)$ within the area $[-1, 1]^2$. While it is in general NP-hard to find the global minimum for a constrained nonconvex optimization problem, PGD (FPNAG) succeeds to find this point (or oscillates around





Fig. 2. Six-hump camel back function.

Fig. 3. Results of optimizing the six-hump camel back function within the area $[-1, 1]^2$.

the neighborhood of the minimum) as presented in Fig. 3. Both POGM and PNAG fall into the trap of the saddle point. Especially, PGD is able to find a “better” local minimum (actually the best) than the other methods. This phenomenon is quite interesting since the standard gradient-based method outperforms momentum-based methods in both stability and final precision. Such a result indicates that the projected method may possess different convergence behavior from the unconstrained counterpart.

In contrast to the smooth nonconvex function above, we now discuss the results of optimizing a nonconvex function with many sharp spikes as Fig. 4, which is given by

$$A(x) = -20e^{-0.2\|x\|/\sqrt{m}} - e^{\frac{\sum_{i=1}^m \cos(2\pi x_i)}{m}} + 20 + e, \quad (105)$$

where $m = 10$ is the dimension of x and x_i is the i th coordinate of x . It is clear that the function has a global minimum $A^*(0) = 0$ and plenty of local minima. As presented in Fig. 5, PNAG and POGM are successful to find the global minimum while PGD and FPNAG fall into some local sharp spike. Compared with Fig. 3, it can be deduced that all the





Fig. 4. 2D Ackley's path function.

Fig. 5. Results of optimizing Ackley's path function within the area $[-33, 33]^{10}$.

algorithms are able to converge to a moderate local minimum after a few iterations. In addition, the “quality” of the local minimum heavily depends on the optimization function and could be affected by the curvature properties of the graph.

Finally, we discuss the effects of adding random perturbations to gradients in the iterations. The perturbation ξ_k is a Gaussian stochastic vector with distribution $\mathcal{N}(0, \epsilon)$ for $\epsilon = 0.001$. The corresponding stochastic scheme is modified at gradients ($\nabla f(y_k) \rightarrow \nabla f(y_k) + \xi_k$) and the learning rate factor before the gradient term ($\gamma \rightarrow \gamma/k$). The results are demonstrated in Figs. 6, 7 and 8. It is clear that adding random perturbations will deteriorate the performance in convex optimization. However, both PNAG and POGM benefit from the perturbation as for optimizing the six-hump camel back function as presented in Fig. 7 compared with Fig. 3. Adding random perturbations to optimizing Ackley's path function barely makes effects. To summarize, it could be beneficial to add random perturbations when the current local minimum is not “good” enough, while it is infeasible if the local minimum




Fig. 6. Results of constrained convex optimization with methods incorporating random perturbations.




Fig. 7. Results of optimizing the six-hump camel back function with methods incorporating random perturbations.

is satisfactory but unstable.

VI. CONCLUSION

In this article, we have explored constrained optimization through the lens of dynamical systems theory. By establishing the connection between a PSP and its Euler discretization, we have obtained theoretical results similar to counterpart results of unconstrained dynamics. Notably, we have developed a novel framework for convergence analysis that transcends traditional numerical methods. Several examples of convergence analysis of proximal gradient methods have been provided to demonstrate the effectiveness of the framework. In the future, we plan to focus on time-varying constrained optimization problems and providing convergence guarantees for finite-time solutions, which require analysis of time-varying constraints.

REFERENCES

- [1] T. Vu, R. Raich, and X. Fu, “On local linear convergence of projected gradient descent for unit-modulus least squares,” *IEEE Trans. Signal Process.*, vol. 71, pp. 3883–3897, 2023.
- [2] Z. Yang, F.-q. Xia, K. Tu, and M.-C. Yue, “Variance reduced random relaxed projection method for constrained finite-sum minimization problems,” *IEEE Trans. Signal Process.*, vol. 72, pp. 2188–2203, 2024.




Fig. 8. Results of optimizing Ackley's path function with methods incorporating random perturbations.

- [3] S. T. Thomdapu, H. Vardhan, and K. Rajawat, "Stochastic compositional gradient descent under compositional constraints," *IEEE Trans. Signal Process.*, vol. 71, pp. 1115–1127, 2023.
- [4] R. Nassif, S. Vlaski, M. Carpentiero, V. Matta, M. Antonini, and A. H. Sayed, "Quantization for decentralized learning under subspace constraints," *IEEE Trans. Signal Process.*, vol. 71, pp. 2320–2335, 2023.
- [5] F. Han, X. Cao, and Y. Gong, "Decentralized stochastic optimization with pairwise constraints and variance reduction," *IEEE Trans. Signal Process.*, vol. 72, pp. 1960–1973, 2024.
- [6] Y. Cui, Y. Li, and C. Ye, "Sample-based and feature-based federated learning for unconstrained and constrained nonconvex optimization via mini-batch scca," *IEEE Trans. Signal Process.*, vol. 70, pp. 3832–3847, 2022.
- [7] G. França, D. P. Robinson, and R. Vidal, "A nonsmooth dynamical systems perspective on accelerated extensions of adm," *IEEE Trans. Autom. Control*, vol. 68, no. 5, pp. 2966–2978, May. 2023.
- [8] R. Raveendran, A. D. Mahindrakar, and U. Vaidya, "Dynamical system approach for time-varying constrained convex optimization problems," *IEEE Trans. Autom. Control*, vol. 69, no. 6, pp. 3822–3834, Jun. 2024.
- [9] G. Bianchin, J. Cortés, J. I. Poveda, and E. Dall'Anese, "Time-varying optimization of lti systems via projected primal-dual gradient flows," *IEEE Trans. Control Netw. Syst.*, vol. 9, no. 1, pp. 474–486, Mar. 2022.
- [10] X. Shi, X. Xu, G. Wen, and J. Cao, "Fixed-time gradient flows for solving constrained optimization: A unified approach," *IEEE/CAA J. Autom. Sinica*, vol. 11, no. 8, pp. 1849–1864, Aug. 2024.
- [11] P. Stechliniski, "Dynamic optimization of complementarity systems," *IEEE Trans. Autom. Control*, vol. 68, no. 2, pp. 1122–1129, Feb. 2023.
- [12] M. Tao, L. Guo, J. Cao, and L. Rutkowski, "A second-order primal-dual dynamics for set constrained distributed optimization problems," *IEEE Trans. Circuits Syst. II*, vol. 71, no. 3, pp. 1316–1320, Mar. 2024.
- [13] M. Benaïm, "A dynamical system approach to stochastic approximations," *SIAM J. Control Optim.*, vol. 34, no. 2, pp. 437–472, 1996.
- [14] A. M. Dalila Azzam-Laouir and L. Thibault, "On perturbed sweeping process," *Applicable Anal.*, vol. 95, no. 2, pp. 303–322, 2016.
- [15] T. H. Cao and B. Mordukhovich, "Optimal control of a nonconvex perturbed sweeping process," *J. Diff. Equations*, vol. 266, no. 2, pp. 1003–1050, Jan. 2019.
- [16] C. Hermosilla and M. Palladino, "Optimal control of the sweeping process with a nonsmooth moving set," *SIAM J. Control Optim.*, vol. 60, no. 5, pp. 2811–2834, 2022.
- [17] L. N. Wadippuli, I. Gudoshnikov, and O. Makarenkov, "Global asymptotic stability of nonconvex sweeping processes," *Discrete Contin. Dyn. Syst. Ser. B*, vol. 25, no. 3, pp. 1129–1139, Mar. 2020.
- [18] J. Venel, "A numerical scheme for a class of sweeping processes," *Numer. Math.*, vol. 118, no. 2, Jun. 2011.
- [19] F. Bernicot and J. Venel, "Convergence order of a numerical scheme for sweeping process," *SIAM J. Control Optim.*, vol. 51, no. 4, pp. 3075–3092, 2013.
- [20] H. Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, 1st ed. Springer New York, NY, 2011.
- [21] M. Kamenskii, O. Makarenkov, L. Niwanthi Wadippuli, and P. Raynaud de Fitte, "Global stability of almost periodic solutions to monotone sweeping processes and their response to non-monotone perturbations," *Nonlinear Anal. Hybrid Syst.*, vol. 30, pp. 213–224, 2018.
- [22] B. Brogliato, A. Daniilidis, C. Lemaréchal, and V. Acary, "On the equivalence between complementarity systems, projected systems and differential inclusions," *Syst. Control Lett.*, vol. 55, no. 1, pp. 45–51, Jan. 2006.
- [23] G. Teschl, *Ordinary Differential Equations and Dynamical Systems*. American Mathematical Society, 2012.
- [24] M. W. Hirsch, H. L. Smith, and X. Q. Zhao, "Chain transitivity, attractivity, and strong repellers for semidynamical systems," *J. Dyn. Diff. Equations*, vol. 13, no. 1, pp. 107–131, Jan. 2001.
- [25] J.-P. Aubin and A. Cellina, *Differential Inclusions: Set-Valued Maps and Viability Theory*, 1st ed. Berlin, Heidelberg: Springer, 1984.
- [26] V. S. Borkar, *Stochastic Approximation: A Dynamical Systems Viewpoint*, 2nd ed. Springer Singapore, 2023.
- [27] M. Benaïm, J. Hofbauer, and S. Sorin, "Stochastic approximations and differential inclusions," *SIAM J. Control Optim.*, vol. 44, no. 1, pp. 328–348, Jan. 2005.
- [28] S. Adly, A. Hantoute, and M. Théra, "Nonsmooth lyapunov pairs for infinite-dimensional first-order differential inclusions," *Nonlinear Anal. Theory Methods Appl.*, vol. 75, no. 3, pp. 985–1008, Feb. 2012.
- [29] S. Adly, A. Hantoute, and B. T. Nguyen, "Lyapunov stability of differential inclusions involving prox-regular sets via maximal monotone operators," *J. Optim. Theory Appl.*, vol. 182, no. 3, pp. 906–934, Sept. 2019.
- [30] B. Polyak, "Some methods of speeding up the convergence of iteration methods," *USSR Comput. Math. Math. Phys.*, vol. 4, no. 5, pp. 1–17, 1964.
- [31] G. Franca, J. Sulam, D. P. Robinson, and R. Vidal, "Conformal symplectic and relativistic optimization," in *Adv. Neural Inf. Process. Syst.*, Dec. 6–12, 2020, Vancouver, BC, Canada.
- [32] S. Ghadimi and G. Lan, "Accelerated gradient methods for nonconvex nonlinear and stochastic programming," *Math. Program.*, vol. 156, no. 1, pp. 59–99, Mar. 2016.
- [33] D. Kim and J. A. Fessler, "Optimized first-order methods for smooth convex minimization," *Math. Program.*, vol. 159, no. 1, Sept. 2016.
- [34] P. Bianchi and J. Jakubowicz, "Convergence of a multi-agent projected stochastic gradient algorithm for non-convex optimization," *IEEE Trans. Autom. Control*, vol. 58, no. 2, pp. 391–405, Feb. 2013.
- [35] Y. Nesterov, "A method of solving a convex programming problem with convergence rate $O(1/k^2)$," *Sov. Math. Dokl.*, vol. 27, no. 2, pp. 372–376, 1983.