

Measuring the speed of quantum particles without a round-trip under non-synchronized quantum clocks

Tomer Shushi

Center for Quantum Science and Technology
& Department of Business Administration,
Guilford Glazer Faculty of Business and Management,
Ben-Gurion University of the Negev, Beer-Sheva, Israel
Email Address: tomershu@bgu.ac.il

August 2, 2024

Abstract

One of the main issues in measuring the speed of light when it only travels from one spatial position into another position, known as the one-way speed of light, is that the clocks belonging to each separated spatial position are not and, in principle, cannot be synchronized with sufficient precision. This issue is the main reason why all of the measurements of the speed of light until now have measured the two-way speed of light, i.e., measuring the speed of light that travels from a source to another location and back to the source, and so there is a need for only one clock to measure the speed. Here, we show that it is possible, in principle, to measure the velocity of particles that travel at the speed of light without assuming a round-trip once we adopt a quantum mechanical description under two boundary conditions to the state of the quantum system followed by the two-state-vector formalism while assuming non-synchronized quantum clocks with unknown time dilation. We show that the weak value of velocity can be measured for a test particle that has a clock that is not synchronized with the clock of the quantum particle. Following the proposed setup, when the weak value of the velocity is known even without knowing the time states of the system, such a weak velocity is the two-way speed of light. Otherwise, one has to impose assumptions regarding the time states of the quantum clocks, which then give weak velocities that can be slower or even faster than the two-way speed of light. We further explore some fundamental implications of the setup. The proposed approach opens a new avenue toward measuring the velocities of quantum particles while overcoming relativistic issues regarding the synchronization of clocks.

Keywords: quantum clocks, quantum foundations, relativistic causality, time dilation, two-state-vector-formalism

1 Introduction

The speed of light is conventionally measured as a two-way speed, meaning that light is sent from a source to another location and back to the source, allowing for the round-trip time to be divided by the total distance traveled [1-5]. The one-way speed of light means that we measure the speed of light that only travels from a source to another location without a round-trip. This would require perfectly synchronized clocks at both the departure and arrival points. While there are no theoretical restrictions for measuring the one-way speed of light, no experiment currently succeeds in measuring the one-way speed of light. Various experiments have been claimed to measure the one-way speed of light [6-8]. However, all of these experiments have detected the two-way speed of light using different setups. For instance, in a 2009 paper [7], the authors proposed a new way for such a measurement. However, only a year after publishing their paper in [8], it has been shown that such experiments do not measure the one-way speed of light but, in fact, the two-way speed of light, and the problem remained open.

According to Einstein's theory of relativity, the synchronization of clocks depends on the relative motion of the observers and the frames of reference in which the measurements are taken. Any attempt to synchronize clocks across different locations without taking into account these relativistic effects would introduce errors, making the measurement unreliable. In the original paper of Einstein proposing special relativity in 1905 [9], Einstein postulated that the speed of light from spatial point A to point B is the same from point B to point A , following the quote from his paper: "We have not defined a common "time" for A and B , for the latter cannot be defined at all unless we establish by *definition* that the "time" required by light to travel from A to B equals the "time" it requires to travel from B to A ". While this postulate seems pleasurable, there is currently no evidence that this is indeed the case, i.e., that the speed of light is the same for every direction in space. Every experiment that aims to measure the speed of light takes the following general scheme:




Figure 1. The process for measuring the speed of light, starting from the spatial position A at time t_1 to separated spatial position B (a mirror) at time t_2 and back to the source at time t_3 .

Then, following Einstein's original idea, the speed of light is calculated by

$$c = \frac{2AB}{t_3 - t_1}, \quad (1)$$

i.e., it is the time that took the light to travel from point A to B and back to A . This is known as the *Einstein synchronization*. Following Einstein's original paper, the time of arrival of the light follows the equation $t_2 = t_1 + (t_3 - t_1)/2$. As shown by Reichenbach [1], we can consider an arbitrary rule for the time of arrival followed by $t_2 = t_1 + \epsilon(t_3 - t_1)$ for some $0 < \epsilon < 1$. This essentially corresponds to the speed of light in each direction

$$c_{\rightarrow} = \frac{1}{2} \frac{c}{\epsilon}, \quad (2)$$

and

$$c_{\leftarrow} = \frac{1}{2} \frac{c}{1 - \epsilon}, \quad (3)$$

so, in principle, it is possible that the speed of light in one direction does not equal the speed of light in the opposite direction, i.e., $c_{\rightarrow} \neq c_{\leftarrow}$. This may be interpreted such that the one-way speed of light is not isotropic in space, in the sense that when light travels in one direction, it possesses a different speed than when it travels in the opposite direction. While $c_{\rightarrow}, c_{\leftarrow}$ can have different values, they are mutually dependent. In particular, they obey the equation

$$\frac{c_{\rightarrow} \cdot (t_2 - t_1) + c_{\leftarrow} \cdot (t_3 - t_2)}{t_3 - t_1} = \frac{2l}{t_3 - t_1} = c. \quad (4)$$

If c_{\rightarrow} is faster/slower than the two-way speed of light c , c_{\leftarrow} compensates it with slower/faster speed, such that the average of c_{\rightarrow} and c_{\leftarrow} describes the usual (two-way) speed of light.

In the following, we demonstrate how the problem of measuring particle velocity without assuming a round-trip is resolved by adopting a quantum mechanical description based on the two-state-vector formalism (TSVF). In the proposed setup, we show that the weak value of velocity can be measured for a test particle with a clock that is not synchronized with the clock of the quantum particle, even when the time dilation between the clocks is entirely unknown. We examine some basic implications and show that the proposed setup does not break relativistic causality, and also examine the case of position-dependent speed of light.

2 Results

Before we introduce the proposed setup, we review the basic elements in consideration of time as a dynamical quantum variable and not a mere parameter. Let us consider a quantum system that contains a clock B and the rest of the system denoted by R , where each of the sub-systems is described by vectors

in the Hilbert spaces \mathcal{H}_A and \mathcal{H}_R , respectively. The entire system is then described by the quantum state $|\Psi\rangle \in \mathcal{H}_B \otimes \mathcal{H}_R$. Assuming that we have a closed quantum system, it follows the Wheeler-DeWitt equation

$$(H_B + H_R) |\Psi\rangle = 0, \quad (5)$$

which suggests we have a (stationary) state of the system that does not change with respect to an external time, with a zero eigenvalue of energy (see [10-18]). We can describe the time observable that is associated with clock B by proposing a time operator T_B in the Hilbert space \mathcal{H}_B , which is observable in units of time. We then define the time eigenvectors, $|t_B\rangle$, and time eigenvalues, t_B , of T_B as the solutions of the equation $T_B |t_B\rangle = t_B |t_B\rangle$. Similar to a standard quantum system, we follow the usual commutation relation $[T_B, H_B] = i\hbar I$, where the Hamiltonian of the clock is $H_B = -i\hbar \partial / \partial t_B$ (see, again, [12-13]). The unitary evolution of our time states is then given by

$$|t_0 + t_B\rangle = e^{-iH_B t_B / \hbar} |t_0\rangle, \quad (6)$$

for an initial time state $|t_0\rangle$. Similar to [13], we define an interaction $H_{int}(T_B)$, which is a time-dependent term of the evolution of system R set by clock B , and it quantifies the interaction between the clock B and the rest of the system R . Then, the total Hamiltonian of our quantum system is given by $H_{total} = H_B + H_R + H_{int}(T_B)$. Following the Wheeler-DeWitt equation, we have $H_{total} |\Psi\rangle = 0$, and by applying the time eigenstate $|t_B\rangle$ of T_B on the left, we can obtain the time-dependent Schrödinger equation for the quantum state $|\psi(t_B)\rangle := \langle t_B | \Psi \rangle$,

$$i\hbar \frac{\partial}{\partial t_B} |\psi(t_B)\rangle = [H_R + H_{int}(t_B)] |\psi(t_B)\rangle, \quad (7)$$

where the equation provides us the evolution of the wavefunction of system R with respect to clock B (see, [14-15,18]).

Let us now consider a system containing two synchronized clocks, A and B , where A is the internal clock of the rest of the system, i.e., $R = A + S$. The Hamiltonian of the system is then $H_{total} = H_A + H_S + H_B + H_{int}(T_B)$, and the Schrödinger equation is $i\hbar \frac{\partial}{\partial t_B} |\psi(t_B)\rangle = [H_A + H_S + H_{int}(T_B)] |\psi(t_B)\rangle$.

A key element in the one-way speed of light problem is the break of synchronicity between the clocks. We can model such a break by adding to the Hamiltonian the term

$$g(T_B) H_A, \quad (8)$$

where $g(u)$, $u \geq 0$, is some positive function, where $\int_0^\infty g(t) dt > 0$. Then, following [12-13], our Schrödinger equation is given by

$$i\hbar \frac{\partial}{\partial t_B} |\psi(t_B)\rangle = (I + g(t_B)) H_A |\psi(t_B)\rangle. \quad (9)$$

The Heisenberg picture allows us to get the exact break of synchronicity between clocks A and B , followed by the time observable of clock A , T_A , with respect to

the ticks of clock B ,

$$\frac{d}{dt_B} T_A = -\frac{i}{\hbar} [T_A, H_R + g(t_B) H_A] = I + g(t_B). \quad (10)$$

We note that when g vanishes, the flow of time in both clocks is the same, and in case $g \neq 0$, we see that clock A ticks faster than clock B .

We can now propose the setup demonstrating how to measure the velocity of quantum particles without a round-trip using the TSVF. For the setup of the weak measurement of velocity, we adopt the framework proposed in [19], which originally provided a way to obtain Cherenkov radiation in a vacuum based on the TSVF and the weak velocity of the quantum particles.

TSVF proposes a time-symmetric picture for the evolution of quantum systems based on two states: a state that is prepared and evolves unitarily forward from the past and a post-selected state. It has been shown that the TSVF allows for the measurement between the pre- and post-selected states, such that the quantum state of the particle is not disturbed, and so hidden information and odd quantum effects of the quantum particle are detected [20-26].

We consider a system that contains a quantum particle S with an internal clock A and an external clock B , followed by the total Hamiltonian

$$H_{Tot} = H_A + H_S + H_B + H_{int}(T_B) + g(T_B) H_A, \quad (11)$$

where H_A/H_B is the (time) Hamiltonian of the quantum clock that generates the translations in time T_A/T_B . H_S is the Hamiltonian of the quantum particle and is given by

$$H_S = p_z v_z, \quad (12)$$

where $p_z = -i\hbar\partial/\partial z$, and

$$v_z = c \cdot \sigma_z \quad (13)$$

acts on the internal Hilbert space of the particle, where c is the speed of light, and $N > 0$ is some integer. The Pauli matrix σ_z operates on the internal Hilbert space. We note that it does not represent spin, and so the particle has no electric or magnetic dipole moment. The eigenvalues of the velocity along the z direction v_z are $-c, +c$, and so if the only allowed values of v_z are its eigenvalues, the quantum particle can only travel at the speed of light c .

The particle moves with velocity v_z in the z direction, i.e., which can be described in the Heisenberg picture, $\frac{d}{dt_A} x = -\frac{i}{\hbar} [x, H_S] = 0$, $\frac{d}{dt_A} y = -\frac{i}{\hbar} [y, H_S] = 0$, and

$$\frac{d}{dt_A} z = -\frac{i}{\hbar} [z, H_S] = v_z. \quad (14)$$

We assume that the system is initially prepared in the form

$$|\Psi_{in}\rangle\rangle = |\Psi_{in}^S, \mathcal{T}_{in}\rangle \Phi(\mathbf{x}, 0), \quad (15)$$

where the quantum particle is prepared in the pre-selected state

$$|\Psi_{in}^S, \mathcal{T}_{in}\rangle = \frac{1}{\sqrt{2}} (|\uparrow\rangle |\mathcal{T}_{in}^+\rangle + |\downarrow\rangle |\mathcal{T}_{in}^-\rangle), \quad (16)$$

for initial time states $|\mathcal{T}_{in}^{-/+}\rangle$, and Φ describes an additional (test) particle that is approximately localized in the origin, $\mathbf{x} = (x, y, z) = 0$, and it takes the Gaussian form $\Phi(\mathbf{x}, 0) = (\varepsilon^2 \pi)^{-3/4} e^{-\mathbf{x}^T \mathbf{x} / 2\varepsilon^2}$ for $\varepsilon > 0$ which gives the dispersion of the Gaussian, and it evolves according to clock B .

We post-select the quantum particle and the clocks' states and find it in the state

$$|\Psi_{fin}^S, \mathcal{T}_{fin}\rangle = \alpha |\uparrow\rangle |\mathcal{T}_{fin}^+\rangle + \beta |\downarrow\rangle |\mathcal{T}_{fin}^-\rangle, \quad (17)$$

for (known) real-valued coefficients α, β , such that $\alpha^2 + \beta^2 = 1$, and final time states $|\mathcal{T}_{fin}^{-/+}\rangle$. Since we do not have any knowledge about the time dilation between the clocks, i.e., about g , we assume that we also do not have knowledge about the time states $|\mathcal{T}_{in/fin}^{-/+}\rangle$, we do, however, assume that they are normalized similar to any other standard quantum state and that $\langle \mathcal{T}_{fin}^{-/+} | \mathcal{T}_{in}^{-/+} \rangle \neq 0$.

The function g generates the break in synchronicity between the clocks, as shown in (10), implying a break in their synchronicity, as illustrated by (10)

$$\frac{d}{dt_B} T_A = -\frac{i}{\hbar} [T_A, H_A + H_S + H_B + H_{int}(T_B) + g(T_B) H_A] = I + g(t_B), \quad (18)$$

where the internal clock of the particle, clock A , ticks faster than clock B , the clock of the test particle.

Similar to the setup proposed in [19], $\Phi(\mathbf{x}, t_B)$ provides the quantum state of the test particle with respect to clock B , and is given by

$$\begin{aligned} \Phi(\mathbf{x}, t_B) &= \langle \Psi_{fin}^S, \mathcal{T}_{fin} | e^{-iH_{Tot} t_B / \hbar} | \Psi_{in}^S, \mathcal{T}_{in} \rangle \Phi(\mathbf{x}, 0) \\ &= \langle \Psi_{fin}^S, \mathcal{T}_{fin} | e^{-i(H_{clocks} + g(t_B) H_A + p_z v_z) t_B / \hbar} | \Psi_{in}^S, \mathcal{T}_{in} \rangle \Phi(\mathbf{x}, 0) \end{aligned} \quad (19)$$

where $H_{clocks} := H_A + H_B + H_{int}(T_B)$. We recall that g is unknown since we do not know the time dilation between clocks A and B .

Then, for a short enough time t_B and under the condition

$$g(t_B) \ll \frac{1}{t_B},$$

we have

$$\begin{aligned} \Phi(\mathbf{x}, t_B) &\approx \langle \Psi_{fin}^S, \mathcal{T}_{fin} | 1 - \frac{i}{\hbar} (H_{clocks} + g(t_B) H_A + p_z v_z) t_B | \Psi_{in}^S, \mathcal{T}_{in} \rangle \Phi(\mathbf{x}, 0) \\ &= \langle \Psi_{fin}^S, \mathcal{T}_{fin} | 1 - \frac{i}{\hbar} (\langle H_{clocks} \rangle_w + g(t_B) \langle H_A \rangle_w + p_z \langle v_z \rangle_w) t_B | \Psi_{in}^S, \mathcal{T}_{in} \rangle \Phi(\mathbf{x}, 0) \\ &\approx \kappa \langle \Psi_{fin}^S, \mathcal{T}_{fin} | e^{-ip_z \langle v_z \rangle_w t_B / \hbar} | \Psi_{in}^S, \mathcal{T}_{in} \rangle \Phi(\mathbf{x}, 0) \\ &= \kappa \langle \Psi_{fin}^S, \mathcal{T}_{fin} | \Psi_{in}^S, \mathcal{T}_{in} \rangle \cdot \Phi(x, y, z - \langle v_z \rangle_w t_B, 0), \end{aligned} \quad (20)$$

where $\kappa = e^{-i(\langle H_{clocks} \rangle_w + g(t_B) \langle H_A \rangle_w) t_B / \hbar}$, and $\langle A \rangle_w$ is the weak-value of an ob-

servable A ,

$$\langle A \rangle_w := \frac{\langle \Psi_{fin}^S, \mathcal{T}_{fin} | A | \Psi_{in}^S, \mathcal{T}_{in} \rangle}{\langle \Psi_{fin}^S, \mathcal{T}_{fin} | \Psi_{in}^S, \mathcal{T}_{in} \rangle}. \quad (21)$$

Eq. (20) implies that the wavefunction of the test particle is displaced by $\langle v_z \rangle_w t_B$.

The velocity of the quantum particle is directly related to the pre- and post-selected states, followed by the weak velocity $\langle v_z \rangle_w$,

$$\langle v_z \rangle_w = \frac{(\langle \mathcal{T}_{fin}^+ | \langle \uparrow | \alpha + \langle \mathcal{T}_{fin}^- | \langle \downarrow | \beta \rangle v_z (|\uparrow\rangle |\mathcal{T}_{in}^+ + |\downarrow\rangle |\mathcal{T}_{in}^-))}{(\langle \mathcal{T}_{fin}^+ | \langle \uparrow | \alpha + \langle \mathcal{T}_{fin}^- | \langle \downarrow | \beta \rangle (|\uparrow\rangle |\mathcal{T}_{in}^+ + |\downarrow\rangle |\mathcal{T}_{in}^-))} = \frac{\langle \mathcal{T}_{fin}^+ | \mathcal{T}_{in}^+ \rangle \alpha - \langle \mathcal{T}_{fin}^- | \mathcal{T}_{in}^- \rangle \beta}{\langle \mathcal{T}_{fin}^+ | \mathcal{T}_{in}^+ \rangle \alpha + \langle \mathcal{T}_{fin}^- | \mathcal{T}_{in}^- \rangle \beta} \cdot c, \quad (22)$$

which can, in principle, take different values than the eigenvalues of v_z , i.e., the two-way speed of light.

The following Figure illustrates the proposed setup of the gedanken experiment.




Figure 2. A scheme of the proposed setup. The experiment is conducted along the z -axis, where the quantum particle with internal clock A is pre- and post-selected followed by a test particle with internal clock B . The test particle is then shifted by $\langle v_z \rangle_w \cdot t_B$ in the z -axis.

It is important to emphasize that, unlike the expected value of velocity, $\langle v_z \rangle$, which is a statistical quantity of the system, the weak values are actual physical quantities, and so the measurement that gives $\langle v_z \rangle_w$ essentially gives the velocity of the quantum particle under the specified pre- and post-selected states of the quantum system. We note that we have an isotropy of the speed of light (of the particle) only statistically, following the expectation value of v_z , $\langle v_z \rangle = 0$, and the eigenvalues $-c, c$.

We can define a quantity, τ , for the ratio between the time amplitudes

$$\tau = \frac{\langle \mathcal{T}_{fin}^- | \mathcal{T}_{in}^- \rangle}{\langle \mathcal{T}_{fin}^+ | \mathcal{T}_{in}^+ \rangle}, \quad (23)$$

leading to the weak velocity of the form

$$\langle v_z \rangle_w = \frac{\alpha - \tau \beta}{\alpha + \tau \beta} \cdot c. \quad (24)$$

Here, τ is the unknown parameter, since we do not have the knowledge about the time states, and so we also do not have the knowledge about the transition amplitudes $\langle \mathcal{T}_{fin}^{-/+} | \mathcal{T}_{in}^{-/+} \rangle$. We can, however, find the value of $\langle v_z \rangle_w$ in specific cases. In the case where $\beta = 0$, the weak velocity becomes the speed of light,

$$\langle v_z \rangle_w = c. \quad (25)$$

The proposed setup reveals that when $\langle v_z \rangle_w$ is known and is independent on the time states, the speed is the two-way speed of light. When the initial and final time states are the same, i.e., $|\mathcal{T}_{in}^{-/+}\rangle = |\mathcal{T}_{fin}^{-/+}\rangle$, from the orthonormality property of quantum states we have $\langle \mathcal{T}_{fin}^{-/+} | \mathcal{T}_{in}^{-/+} \rangle = \langle \mathcal{T}_{in}^{-/+} | \mathcal{T}_{in}^{-/+} \rangle = 1$, and the weak velocity also becomes a known quantity with by having $\tau \equiv 1$, and a potential violation of the two-way speed of light, with $\langle v_z \rangle_w = \frac{\alpha - \beta}{\alpha + \beta} \cdot c$.

Unlike standard (strong) measurements of the velocity, where the measurements only give the eigenvalues of v_z , weak values can give values out of the spectrum of the eigenvalues. We considered a particle that can only travel at the speed of light, with eigenvalues $-c, c$ along the z -axis; however, the weak velocity $\langle v_z \rangle_w$ can be slower and even faster than c . We can now relate the ϵ in the model of the one-way speed of light with the weak velocity, following formula (2), with having

$$\epsilon = \frac{1}{2} \frac{\alpha + \tau\beta}{\alpha - \tau\beta} \quad (26)$$

and since $0 < \epsilon < 1$, we assume that τ is real-valued, and we are restricted to the coefficients of the post- selected state that satisfies

$$\tau < \frac{\alpha}{3\beta}. \quad (27)$$

The proposed setup can also be used for particles that have a range of possible speeds bounded by the speed of light. In particular, instead of only eigenvalues $-c, c$ of v_z we can consider the case where the eigenvalues of the velocity along the z direction v_z are $-c, -c + 2c/N, \dots, c - 2c/N, c$. This corresponds with the velocity observable

$$v_z = \frac{c}{N} \sum_{i=1}^N \sigma_z^{(i)}, \quad (28)$$

where, again, the Pauli matrices $\sigma_z^{(i)}$ operate on the internal Hilbert space. Then, we can obtain the same experiment as the previous one while considering the pre- and post- selected states of the quantum system

$$|\Psi_{in}^S, \mathcal{T}_{in}\rangle = 2^{-N/2} \otimes_{i=1}^N (|\uparrow_i\rangle |\mathcal{T}_{in}^+\rangle + |\downarrow_i\rangle |\mathcal{T}_{in}^-\rangle), \quad (29)$$

and

$$|\Psi_{fin}^S, \mathcal{T}_{fin}\rangle = \otimes_{i=1}^N \left(\alpha |\uparrow_i\rangle |\mathcal{T}_{fin}^+\rangle + \beta |\downarrow_i\rangle |\mathcal{T}_{fin}^-\rangle \right), \quad (30)$$

respectively, then similar to the original setup, the weak velocity is given by $\langle v_z \rangle_w = \frac{\alpha - \tau\beta}{\alpha + \tau\beta} \cdot c$.

2.1 Relativistic causality

The ϵ parameter in (2) and (3) implies a speed that is higher than c in at least one direction. However, it does not violate relativistic causality, since we can only measure two-way speed of light in the original classical setup. We provided a setup that can also exceed the speed of light whenever

$$\alpha \cdot \tau\beta < 0, \quad (31)$$

since then $\langle v_z \rangle_w > c$. However, this also does not violate relativistic causality, as mentioned in [19].

In the case

$$\left| \frac{\alpha - \tau\beta}{\alpha + \tau\beta} \right| < 1, \quad (32)$$

i.e.

$$|\langle v_z \rangle_w| < c, \quad (33)$$

we do not have any problem with relativistic causality since the speed is less than the speed of light. In the case where $|\langle v_z \rangle_w| > c$, we may have a problem with relativistic causality. However, no violation of relativistic causality is taking place. The reason is that since Φ is an analytic function with respect to z , we have a change in Φ , $\Phi(x, y, z - \langle v_z \rangle_w t_B, 0)$, which does not transmit any information, because the same message is passed to all \mathbf{x} and t (see, [19]). Thus, there is no way in which relativistic causality can be violated.

2.2 The case of position-dependent speed of light

Let us now examine the case of position-dependent speed of light. Since the beginning days of relativity theory, the position-dependent speed of light, $c(\mathbf{x})$, has been explored. Such consideration even appeared in Einstein's paper in 1911 [27]. Such models of varying speed of light (VSL), however, have not gained much attention in the literature. In recent years, there has been a relative rise in exploring such hypothesized possibilities under certain various conditions [28-31]. Such theories mainly aim at exploring new cosmological models for exploring, for example, the beginning of the universe. While the position-dependent speed of light is still merely a theoretical, speculative theory, it may provide a new pathway toward a possible theory that links concepts from quantum mechanics and general relativity into a single coherent scheme. There are also various attempts to establish experimental setups to test the possibility of position-dependent speed of light (see, e.g., [28]). When considering the position-dependent speed of light, one may also naturally consider other extensions of physical constants that become variables for getting a physically meaningful model of reality. Various models have been proposed, one of which is the gravitational constant and the Planck constant, which depend on the speed of light. Let us consider how $c(\mathbf{x})$ and a reduced Planck constant that depends on $c(\mathbf{x})$, $\hbar(\mathbf{x}) > 0 \forall \mathbf{x}$, are considered in the proposed model.

The standard momentum operator is given by $\hat{p}_j = -i\hbar_0\partial/\partial x_j, j = 1, 2, 3$, where \hbar_0 is the (reduced) Planck constant. When considering \hbar to be variable, and in particular, $\hbar(\mathbf{x})$, there is no unique form for the Hamiltonian of the system. While in the standard non-relativistic case, the Hamiltonian is $H = \hat{\mathbf{p}}^2/2m + V(\mathbf{x})$ where $\hat{\mathbf{p}} = (\hat{p}_1, \hat{p}_2, \hat{p}_3)$. Here, since

$$[\hbar(\mathbf{x}), \hat{p}_j] \neq 0, \quad j = 1, 2, 3, \quad (34)$$

there is no particular way of how to substitute $\hbar(\mathbf{x})$ in the momentum operator.

Following [32,33], we define a general form for the Hamiltonian

$$H_S = \Pi_{z,\hbar} v_z, \quad (35)$$

for the deformed momentum operator

$$\Pi_{z,\hbar} = \sqrt{\hbar} \left(-i \frac{\partial}{\partial z} \right) \sqrt{\hbar}. \quad (36)$$

We note that by adopting the deformed momentum operator, we also have a deformed commutation relation,

$$[z, \Pi_{z,\hbar}] = i\hbar(\mathbf{x}). \quad (37)$$

Going back to our experiment, we have

$$\begin{aligned} \Phi(\mathbf{x}, t_B) &\approx \langle \Psi_{fin}^S, \mathcal{T}_{fin} | 1 - \frac{i}{\hbar} (H_{\text{clocks}} + g(t_B) H_A + \Pi_{z,\hbar} v_z) t_B | \Psi_{in}^S, \mathcal{T}_{in} \rangle \Phi(\mathbf{x}, 0) \\ &= \langle \Psi_{fin}^S, \mathcal{T}_{fin} | 1 - \frac{i}{\hbar} (\langle H_{\text{clocks}} \rangle_w + g(t_B) \langle H_A \rangle_w + p_z \langle v_z \rangle_w) t_B | \Psi_{in}^S, \mathcal{T}_{in} \rangle \Phi(\mathbf{x}, 0) \\ &\approx \kappa \cdot \langle \Psi_{fin}^S, \mathcal{T}_{fin} | \exp \left\{ -i \frac{1}{\hbar(\mathbf{x})} \Pi_{z,\hbar} \langle v_z \rangle_w t_B \right\} | \Psi_{in}^S, \mathcal{T}_{in} \rangle \Phi(\mathbf{x}, 0), \end{aligned} \quad (38)$$

where

$$\langle v_z \rangle_w = \frac{\alpha - \tau\beta}{\alpha + \tau\beta} \cdot c(\mathbf{x}), \quad (39)$$

and

$$\frac{1}{\hbar(c(\mathbf{x}))} \Pi_{z,\hbar} \langle v_z \rangle_w = \frac{\alpha - \tau\beta}{\alpha + \tau\beta} \frac{1}{\sqrt{\hbar(c(\mathbf{x}))}} \left(-i \frac{\partial}{\partial z} \right) \sqrt{\hbar(\mathbf{x})} \cdot c(\mathbf{x}). \quad (40)$$

If we assume that the shift of Φ is similar to the case of constant speed of light and Planck constant, then we have to assume that

$$\sqrt{\hbar(\mathbf{x})} \cdot c(\mathbf{x}) = \sqrt{\Lambda}, \quad (41)$$

for some constant $\Lambda > 0$, and so we establish the relation

$$\hbar(\mathbf{x}) = \frac{\Lambda}{c(\mathbf{x})^2}, \quad (42)$$

implying that \hbar is a function of \mathbf{x} through $c(\mathbf{x})$. In this case, when we assume that $\Lambda = \sqrt{\hbar_0}c_0$ where \hbar_0 is the original (reduced) Planck constant and c_0 is the (two-way) speed of light in vacuum, we have

$$\Phi \left(x, y, z - \sqrt{\hbar_0/\hbar} (c(\mathbf{x})) \langle v_z^0 \rangle_w t_B, 0 \right) \quad (43)$$

where $\langle v_z^0 \rangle_w = \frac{\alpha - \tau\beta}{\alpha + \tau\beta} c_0$ is the weak-velocity in case the (two-way) speed of light is a constant c_0 .

3 Discussion

While all of the experiments for measuring the speed of light have measured the two-way speed of light, there is still a quest for obtaining adequate setups that allow the measurement without the consideration of a round-trip. We have shown that it is possible, in principle, to measure the velocity of particles that travel at the speed of light without assuming a round-trip once we adopt a quantum mechanical description when considering two boundary conditions to the quantum system followed by the two-state-vector formalism while assuming non-synchronized quantum clocks with unknown time dilation. Followed by the proposed setup, a test particle is shifted by the amount of $\langle v_z \rangle_w t_B$, which allows finding the weak velocity of the quantum particle, $\langle v_z \rangle_w$. The weak velocity takes, in general, different values than the usual eigenvalues of the velocity observable. There is a challenge in finding a theoretical prediction for $\langle v_z \rangle_w$ since the time states of the quantum clocks are unknown. However, when imposing $(\alpha = 0, \beta \neq 0)$ or $(\alpha \neq 0, \beta = 0)$, the weak velocity becomes independent from the time states, and it follows the speed of light with $\langle v_z \rangle_w = c$ and $\langle v_z \rangle_w = -c$ respectively. The weak velocity of the speed of light can also be found by knowing the interference between the time states, which boils down to knowing τ appearing in (24). Following the basic description of the one-way speed of light suggests a possible break in the isotropy of space in the sense that the speed of light is different for different spatial directions. The proposed setup provides a different interpretation, where the different speeds are derived from the given pre- and post- selected states of the quantum system. In particular, We have related the weak velocity with the ϵ parameter in the theory of the one-way speed of light, which allows us to re-interpret the well-known idea in which ϵ is merely a convention assumed by an observer by its own freedom of choice of ϵ , with a physical meaning where ϵ is determined by the weak value of the velocities governed by the pre- and post- selected states of the system. And so the freedom to choose ϵ is converted to the freedom of choosing the pre- and post-selected states of the system. This connection provides a new way to link quantum mechanical behavior into relativity theory, where instead of having a special treatment of space and time, assuming non-isotropic space, ϵ comes from the two boundary conditions of quantum mechanics that fully determines the velocity in between the initial and final states of the quantum system. Thus, the proposed model suggests that space can be, in fact, isotropic while, at the

same time, $\epsilon \neq 1/2$. For future research, we propose to explore whether one can establish a technique to synchronize the quantum clocks A and B by having the detected information on the shift by the test particle. When we do not know such synchronization, τ is, in general, unknown, and so in actual experiments, the shift of Φ will be determined by random outcomes of $\langle v_z \rangle_w$. Thus, one may provide a way to statistically estimate τ . We propose to explore it in future research. For future research, we also propose to extend the model to explore the case of measuring the speed of quantum particles that travel near strong gravitational fields under some curved spacetime. This may provide a new way to gain knowledge of the link between quantum mechanics and general relativity.

References

- [1] Reichenbach, H. (2012). The philosophy of space and time. Courier Corporation.
- [2] Anderson, R., Vetharaniam, I., & Stedman, G. E. (1998). Conventionality of synchronisation, gauge dependence and test theories of relativity. *Physics reports*, **295**, 93-180.
- [3] Blaney, T. G., Bradley, C. C., Edwards, G. J., Jolliffe, B. W., Knight, D. J. E., Rowley, W. R. C., ... & Woods, P. T. (1974). Measurement of the speed of light. *Nature*, **251**, 46-46.
- [4] Cao, S., Biesiada, M., Jackson, J., Zheng, X., Zhao, Y., & Zhu, Z. H. (2017). Measuring the speed of light with ultra-compact radio quasars. *Journal of Cosmology and Astroparticle Physics*, **2017**, 012.
- [5] Salzano, V., Dabrowski, M. P., & Lazkoz, R. (2015). Measuring the speed of light with Baryon Acoustic Oscillations. *Physical Review Letters*, **114**, 101304.
- [6] Zhang, Y. Z. (1997). Special relativity and its experimental foundation (Vol. 4). World Scientific.
- [7] Greaves, E. D., Rodríguez, A. M., & Ruiz-Camacho, J. (2009). A one-way speed of light experiment. *American Journal of Physics*, **77**, 894-896.
- [8] Finkelstein, J. (2010). Comment on “A one-way speed of light experiment” by ED Greaves, An Michel Rodríguez, and J. Ruiz-Camacho [Am. J. Phys. 77 (10), 894–896 (2009)]. *American Journal of Physics*, **78**, 877-877.
- [9] Einstein, A. (1905). Zur elektrodynamik bewegter körper. *Annalen der physik*, **17**, 891-921.
- [10] Page, D. N., & Wootters, W. K. (1983). Evolution without evolution: Dynamics described by stationary observables. *Physical Review D*, **27**, 2885.
- [11] Angelo, R. M., Brunner, N., Popescu, S., Short, A. J., & Skrzypczyk, P. (2011). Physics within a quantum reference frame. *Journal of Physics A: Mathematical and Theoretical*, **44**, 145304.
- [12] Paiva, I. L., Te'eni, A., Peled, B. Y., Cohen, E., & Aharonov, Y. (2022). Non-inertial quantum clock frames lead to non-Hermitian dynamics. *Communications Physics*, **5**, 298.
- [13] Cohen, E. (2023). Quantum clock frames: Uncertainty relations, non-

- Hermitian dynamics and nonlocality in time. *Journal of Physics: Conference Series*, **2533**. IOP Publishing.
- [14] Bužek, V., Derka, R., & Massar, S. (1999). Optimal quantum clocks. *Physical Review Letters*, **82**, 2207.
- [15] Komar, P., Kessler, E. M., Bishof, M., Jiang, L., Sørensen, A. S., Ye, J., & Lukin, M. D. (2014). A quantum network of clocks. *Nature Physics*, **10**, 582-587.
- [16] Castro-Ruiz, E., Giacomini, F., Belenchia, A., & Brukner, Č. (2020). Quantum clocks and the temporal localisability of events in the presence of gravitating quantum systems. *Nature Communications*, **11**, 2672.
- [17] Smith, A. R., & Ahmadi, M. (2020). Quantum clocks observe classical and quantum time dilation. *Nature communications*, **11**, 5360.
- [18] Castro-Ruiz, E., Giacomini, F., Belenchia, A., & Brukner, Č. (2020). Quantum clocks and the temporal localisability of events in the presence of gravitating quantum systems. *Nature Communications*, **11**, 2672.
- [19] Rohrlich, D., & Aharonov, Y. (2002). Cherenkov radiation of superluminal particles. *Physical Review A*, **66**, 042102.
- [20] Aharonov, Y., Albert, D. Z., & Vaidman, L. (1988). How the result of a measurement of a component of the spin of a spin-1/2 particle can turn out to be 100. *Physical Review Letters*, **60**, 1351.
- [21] Aharonov, Y., & Vaidman, L. (2008). The two-state vector formalism: an updated review. *Time in quantum mechanics*, 399-447.
- [22] Aharonov, Y., & Vaidman, L. (1991). Complete description of a quantum system at a given time. *Journal of Physics A: Mathematical and General*, **24**, 2315.
- [23] Aharonov, Y., & Shushi, T. (2022). Complex-valued classical behavior from the correspondence limit of quantum mechanics with two boundary conditions. *Foundations of Physics*, **52**, 56.
- [24] Aharonov, Y., Popescu, S., Rohrlich, D., & Skrzypczyk, P. (2013). Quantum cheshire cats. *New Journal of Physics*, **15**, 113015.
- [25] Aharonov, Y., Colombo, F., Popescu, S., Sabadini, I., Struppa, D. C., & Tollaksen, J. (2016). Quantum violation of the pigeonhole principle and the nature of quantum correlations. *Proceedings of the National Academy of Sciences*, **113**, 532-535.
- [26] Bliokh, K. Y., Bekshaev, A. Y., Kofman, A. G., & Nori, F. (2013). Photon trajectories, anomalous velocities and weak measurements: a classical interpretation. *New Journal of Physics*, **15**, 073022.
- [27] Einstein, A. (1911). Über den Einfluß der Schwerkraft auf die Ausbreitung des Lichtes. *Annalen der Physik*, **35**, 898-908.
- [28] Magueijo, J. (2003). New varying speed of light theories. *Reports on Progress in Physics*, **66**, 2025.
- [29] Kragh, H. S. (2006). Cosmologies with varying speed of light: A historical perspective. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics*, **37**, 726-737.
- [30] Unzicker, A. (2009). A look at the abandoned contributions to cosmology of Dirac, Sciama, and Dicke. *Annalen der Physik*, **521**, 57-70.

- [31] Albrecht, A., & Magueijo, J. (1999). Time varying speed of light as a solution to cosmological puzzles. *Physical Review D*, **59**, 043516.
- [32] Quesne, C., & Tkachuk, V. M. (2004). Deformed algebras, position-dependent effective masses and curved spaces: an exactly solvable Coulomb problem. *Journal of Physics A: Mathematical and General*, **37**, 4267.
- [33] Shushi, T. (2024). A Non-Standard Coupling Between Quantum Systems Originated From Their Kinetic Energy. *Annalen der Physik*, **536**, 2300363.