

# Main functions and the spectrum of super graphs

G. Arunkumar

Department of Mathematics, Indian Institute of Technology Madras, Chennai - 600036, India

Email: arun.maths123@gmail.com

Peter J. Cameron

School of Mathematics and Statistics, University of St Andrews, Fife, UK

Email: pjc20@st-andrews.ac.uk

R. Ganeshbabu

Department of Mathematics, Indian Institute of Technology Madras, Chennai - 600036, India

Email: ma22d011@smail.iitm.ac.in

Rajat Kanti Nath\*

Department of Mathematical Sciences, Tezpur University, Sonitpur, Assam 784028, India.

Email: rajatkantinath@yahoo.com

## Abstract

Let  $A$  be a graph type and  $B$  an equivalence relation on a group  $G$ . Let  $[g]$  be the equivalence class of  $g$  with respect to the equivalence relation  $B$ . The  $B$  super $A$  graph of  $G$  is an undirected graph whose vertex set is  $G$  and two distinct vertices  $g, h \in G$  are adjacent if  $[g] = [h]$  or there exist  $x \in [g]$  and  $y \in [h]$  such that  $x$  and  $y$  are adjacent in the  $A$  graph of  $G$ . In this paper, we compute spectrum of equality/conjugacy supercommuting graphs of dihedral/dicyclic groups and show that these graphs are not integral.

**Keywords** Supercommuting graph, spectrum, main function, dihedral group, dicyclic group

**Mathematics Subject Classification** 05C25, 20D99

---

\*Corresponding author

# 1 Introduction

This is a continuation of our work on Super graphs on groups done in [2] and [3]. In these earlier works, we introduced and studied three types of graphs, and three equivalence relations defined on a group, viz. the power graph, enhanced power graph, commuting graph, and the relations of equality, conjugacy, and same order; for each choice of a graph type A and an equivalence relation B, there is a graph, the B *superA graph* defined on  $G$ . Two distinct vertices  $g, h \in G$  are adjacent in B superA graph if  $[g] = [h]$  or there exist  $x \in [g]$  and  $y \in [h]$  such that  $x$  and  $y$  are adjacent in the A graph of  $G$ , where  $[g]$  is the equivalence class of  $g$  with respect to the equivalence relation B. For other terminologies and notations regarding graphs and supergraphs related to groups, we refer [2] and [6]. In the present work, we shall study the spectra of B *superA graphs* defined on a group  $G$ . The study of spectra of graphs associated with various algebraic structures is a widely explored area. For instance, see [5, 10, 18] for commuting graphs, commuting conjugacy class graphs and power graphs of finite groups; [4, 21] for Cayley graphs of finite groups; [19] for zero-divisor graphs of finite commutative rings; [11] for commuting graphs of finite rings; [13] for 1-point fixing graph etc. It is worth mentioning that Dalal et al. [9] also computed the spectrum and Laplacian spectrum of order/conjugacy supercommuting graphs of certain groups.

In [3], it was observed that super graphs are generalized compositions of complete graphs. Thus, to study the spectra of a B *superA graph*, results on the spectra of generalized compositions of complete graphs are useful. In [7], Cardoso et al. express the adjacency spectra of a generalized composition of regular graphs in terms of the adjacency spectra of the factor graphs and the determinant of a quotient matrix. Despite this result, explicit computations of the spectra of generalized compositions, especially in the quotient matrix, tend to get difficult.

One important concept in the study of spectra is that of a *main* eigenvalue of a graph, introduced by Cvetković in [8]. An eigenvalue  $\lambda$  of a graph  $\Gamma$  is a main eigenvalue if it has an eigenvector which is not orthogonal to the all-one vector  $\mathbf{1}_n$ , where  $n = |V(\Gamma)|$ , where  $V(\Gamma)$  is the set of vertices of  $\Gamma$ . For a survey on the main eigenvalues of a graph, see [22]. Closely related to main eigenvalues of a graph  $\Gamma$  is the function  $\mathbf{1}_n^t(\lambda I_n - A(\Gamma))^{-1}\mathbf{1}_n$  introduced as the *coronal* by Mcleman and McNicholas in [17]. The coronal plays a crucial role in determining the spectra of graphs arising from various graph operations [14, 15, 20, 26]. In [23], Saravanan et al. generalize the coronal of a graph to the concept of a *main* function: The main function associated with an  $n \times n$  matrix  $M$ , corresponding to two  $n \times 1$  vectors  $u, v$  is the function  $v^t(\lambda I_n - M)^{-1}u$ . They proved that the spectra of an arbitrary generalized composition are determined completely by the spectra of the factor graphs and the associated main functions [23, Theorem 4]. These main functions pave the way for the effective computation of the spectra of interest in this paper.

In Section 2, we recall results related to spectrum of generalized composition of regular graphs and main functions. In Section 3, using main function techniques, we explicitly compute the adjacency spectrum of the class of equality supercommuting and conjugacy supercommuting graphs of dihedral groups and dicyclic groups. It can be seen that these graphs are not integral.

## 2 Notation and auxiliary results

Let  $\mathcal{H}$  be a graph and  $V(\mathcal{H}) = \{1, \dots, k\}$ . Let  $\Gamma_1, \dots, \Gamma_k$  be a collection of graphs with  $V(\Gamma_i) = \{v_i^1, \dots, v_i^{n_i}\}$  for  $1 \leq i \leq k$ . Then  $\mathcal{H}$ -join (also known as generalized composition) of the graphs  $\Gamma_1, \dots, \Gamma_k$ , denoted by  $\mathcal{G} =: \mathcal{H}[\Gamma_1, \dots, \Gamma_k]$ , is a graph whose vertex set is  $V(\Gamma_1) \sqcup \dots \sqcup V(\Gamma_k)$  and two vertices  $v_i^p$  and  $v_j^q$  of  $\mathcal{G}$  are adjacent if one the following conditions is satisfied:

- (a)  $i = j$ , and  $v_i^p$  and  $v_j^q$  are adjacent vertices in  $\Gamma_i$ .
- (b)  $i \neq j$  and  $i$  and  $j$  are adjacent in  $\mathcal{H}$ .

This generalized composition of graphs introduced by Schwenk [24] which is also known as generalized lexicographic product and joined union (see [1, 12, 16, 25]). Let  $\mathcal{A}(\Gamma)$  be the adjacency matrix of  $\Gamma$  and  $\text{Spec}(\Gamma) := \text{Spec}(\mathcal{A}(\Gamma))$  be the spectrum of  $\Gamma$ , the multiset of eigenvalues of  $\mathcal{A}(\Gamma)$ . The following result of Cardoso [7] gives spectrum of  $\mathcal{H}[\Gamma_1, \dots, \Gamma_k]$ , to some extent.

**Theorem 2.1.** [7] *Let  $\mathcal{G} = \mathcal{H}[\Gamma_1, \dots, \Gamma_k]$  where  $V(\mathcal{H}) = \{1, \dots, k\}$ ,  $|V(\Gamma_i)| = n_i$  and  $\Gamma_i$  is  $r_i$ -regular for  $1 \leq i \leq k$ . For  $1 \leq i \leq j \leq k$ , define  $\rho_{i,j} = 1$  if  $\{i, j\}$  is an edge in  $\mathcal{H}$  and 0 otherwise. Then  $r_i$  is an eigenvalue of  $\mathcal{A}(\Gamma_i)$  and*

$$\text{Spec}(\mathcal{G}) = \left( \bigcup_{i=1}^k (\text{Spec}(\Gamma_i) \setminus \{r_i\}) \right) \cup \text{Spec}(\tilde{A}(\mathcal{G})),$$

$$\text{where } \tilde{A}(\mathcal{G}) = \begin{bmatrix} r_1 & \sqrt{n_1 n_2} \rho_{1,2} & \cdots & \sqrt{n_1 n_k} \rho_{1,k} \\ \sqrt{n_2 n_1} \rho_{2,1} & r_2 & \cdots & \sqrt{n_2 n_k} \rho_{2,k} \\ \vdots & \vdots & \ddots & \vdots \\ \sqrt{n_k n_1} \rho_{k,1} & \sqrt{n_k n_2} \rho_{k,2} & \cdots & r_k \end{bmatrix}.$$

We shall use Theorem 2.1 while computing spectrum of various super graphs of dihedral and dicyclic groups. We write  $\Gamma_A^B$  to denote the B superA graph on  $G$  associated to B, A and  $G$ . In [3, Proposition 4.1], it was shown that  $\Gamma_A^B$  isomorphic to  $\Delta[K_{n_1}, \dots, K_{n_r}]$  for some graph

$\Delta$  on  $r$  vertices, where  $n_1, \dots, n_r$  are the sizes of the equivalence classes of  $B$ . Note that the graph  $\Delta$  is the induced subgraph of  $\Gamma_A^B$  on the set of equivalence class representatives of  $B$ . Since  $K_n$  is  $(n-1)$ -regular and

$$\text{Spec}(K_n) \setminus \{n-1\} = \underbrace{\{-1, \dots, -1\}}_{(n-1) \text{ times}},$$

applying Theorem 2.1 to  $\Gamma_A^B$  we get the spectrum of  $\Gamma_A^B$  as given below.

**Theorem 2.2.** *The spectrum  $\Gamma_A^B = \Delta[K_{n_1}, \dots, K_{n_k}]$ , where  $V(\Delta) = \{1, \dots, k\}$  and  $n_i \geq 1$  for  $1 \leq i \leq k$ , is given by*

$$\text{Spec}(\Gamma_A^B) = \left( \bigcup_{\substack{1 \leq i \leq k \\ n_i \geq 1}} \underbrace{\{-1, \dots, -1\}}_{(n_i-1) \text{ times}} \right) \cup \text{Spec}(\tilde{\mathcal{A}}(\Gamma_A^B)),$$

$$\text{where } \tilde{\mathcal{A}}(\Gamma_A^B) = \begin{bmatrix} n_1 - 1 & \sqrt{n_1 n_2} \rho_{1,2} & \cdots & \sqrt{n_1 n_k} \rho_{1,k} \\ \sqrt{n_2 n_1} \rho_{2,1} & n_2 - 1 & \cdots & \sqrt{n_2 n_k} \rho_{2,k} \\ \vdots & \vdots & \ddots & \vdots \\ \sqrt{n_k n_1} \rho_{k,1} & \sqrt{n_k n_2} \rho_{k,2} & \cdots & n_k - 1 \end{bmatrix}.$$

In general it is difficult to compute  $\text{Spec}(\tilde{\mathcal{A}}(G))$ . However, in Section 3, we shall calculate  $\text{Spec}(\tilde{\mathcal{A}}(G))$  explicitly when  $\Gamma_A^B$  is equality *superA graph* and conjugacy *superA graph* for dihedral groups and dicyclic group where the graph  $A$  is the commuting graph. In this process, the concept of *main function* associated with a matrix and a few essential lemmas from [23] are useful and these are listed below.

**Definition 1.** Let  $M$  be an  $n \times n$  complex matrix, and let  $u$  and  $v$  be  $n \times 1$  complex vectors. The main function associated to the matrix  $M$  corresponding to the vectors  $u$  and  $v$ , denoted by  $\Gamma_M(u, v)$ , is defined to be  $\Gamma_M(u, v) = v^t(\lambda I - M)^{-1}u \in \mathbb{C}(\lambda)$ . When  $u = v$ , we denote  $\Gamma_M(u, v)$  by  $\Gamma_M(u)$ .

**Lemma 2.3.** *Let  $A, B, C$  and  $D$  be matrices such that  $M = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$ . If  $D$  is invertible, then  $\det(M) = \det(D) \det(A - BD^{-1}C)$ .*

**Lemma 2.4.** *Let  $A$  be an  $n \times n$  invertible matrix, and let  $u$  and  $v$  be any two  $n \times 1$  vectors such that  $1 + v^t A^{-1}u \neq 0$ . Then*

$$(a) \det(A + uv^t) = (1 + v^t A^{-1}u) \det(A).$$

$$(b) (A + uv^t)^{-1} = A^{-1} - \frac{A^{-1}uv^tA^{-1}}{1 + v^tA^{-1}u}.$$

**Lemma 2.5.** *Let  $M$  be a matrix of order  $n$  with an eigenvector  $u$  corresponding to the eigenvalue  $\mu$ . Then  $\Gamma_M(u; \lambda) = \frac{\|u\|^2}{\lambda - \mu}$ .*

We write  $\text{ESCom}(G)$  and  $\text{CSCom}(G)$  to denote the equality supercommuting and conjugacy supercommuting graphs of a group  $G$ . In [3],  $\text{ESCom}(G)$  and  $\text{CSCom}(G)$  were realized for dihedral groups and dicyclic groups, as described in the following theorem.

**Theorem 2.6.** *Let  $D_{2n} = \langle a, b : a^n = b^2 = e, bab^{-1} = a^{-1} \rangle$  be the dihedral group and  $Q_{4n} = \langle a, b : a^{2n} = e, a^n = b^2, bab^{-1} = a^{-1} \rangle$  be the dicyclic group. Then*

$$\text{ESCom}(D_{2n}) \cong \begin{cases} K_1 \vee (\underbrace{K_1 \sqcup \cdots \sqcup K_1}_{n \text{ times}} \sqcup K_{n-1}), & \text{if } n \text{ is odd} \\ K_2 \vee (\underbrace{K_2 \sqcup \cdots \sqcup K_2}_{\frac{n}{2} \text{ times}} \sqcup K_{n-2}), & \text{if } n \text{ is even,} \end{cases}$$

$$\text{CSCom}(D_{2n}) \cong \begin{cases} K_1 \vee (K_1 \sqcup K_{\frac{n-1}{2}}) [\underbrace{K_1, K_2, \dots, K_2}_{(\frac{n-1}{2} \text{ times})}, K_n], & \text{if } n \text{ is odd} \\ K_2 \vee (K_1 \sqcup K_1 \sqcup K_{\frac{n}{2}-1}) [\underbrace{K_1, K_1, K_2, \dots, K_2}_{(\frac{n}{2}-1 \text{ times})}, K_{\frac{n}{2}}, K_{\frac{n}{2}}], & \text{if } n \text{ and } \frac{n}{2} \text{ are even} \\ K_2 \vee (K_2 \sqcup K_{\frac{n}{2}-1}) [\underbrace{K_1, K_1, K_2, \dots, K_2}_{(\frac{n}{2}-1 \text{ times})}, K_{\frac{n}{2}}, K_{\frac{n}{2}}], & \text{if } n \text{ is even and } \frac{n}{2} \text{ is odd,} \end{cases}$$

$$\text{ESCom}(Q_{4n}) \cong K_2 \vee (\underbrace{K_2 \sqcup \cdots \sqcup K_2}_{n \text{ times}} \sqcup K_{2n-2})$$

and

$$\text{CSCom}(Q_{4n}) \cong \begin{cases} K_2 \vee (K_1 \sqcup K_1 \sqcup K_{n-1}) [\underbrace{K_1, K_1, K_2, \dots, K_2}_{(n-1 \text{ times})}, K_n, K_n], & \text{if } n \text{ is even} \\ K_2 \vee (K_2 \sqcup K_{n-1}) [\underbrace{K_1, K_1, K_2, \dots, K_2}_{(n-1 \text{ times})}, K_n, K_n], & \text{if } n \text{ is odd.} \end{cases}$$

### 3 Adjacency spectrum of supergraphs

In this section, we compute the spectrum of  $\text{ESCom}(G)$  and  $\text{CSCom}(G)$ , where  $G$  is a dihedral or dicyclic group. In our proofs we shall use Theorem 2.6, Theorem 2.1 along with Schur com-

plement and the main function technique ([23, Theorem 2]). We begin with the computation of  $\text{Spec}(\text{ESCom}(G))$ , where  $G$  is the dihedral group of order  $2n$ .

**Theorem 3.1.** *Let  $D_{2n} = \langle a, b : a^n = b^2 = e, bab^{-1} = a^{-1} \rangle$  be the dihedral group of order  $2n$ .*

(a) *If  $n$  is odd then  $\text{Spec}(\text{ESCom}(D_{2n})) = \{ \underbrace{0, \dots, 0}_{(n-1) \text{ times}}, \underbrace{-1, \dots, -1}_{(n-2) \text{ times}}, \alpha, \beta, \gamma \}$ , where  $\alpha, \beta, \gamma$  are roots of the equation  $x^3 - (n-2)x^2 - (2n-1)x + n(n-2) = 0$ .*

(b) *If  $n$  is even then  $\text{Spec}(\text{ESCom}(D_{2n})) = \{ \underbrace{1, \dots, 1}_{(\frac{n}{2}-1) \text{ times}}, \underbrace{-1, \dots, -1}_{(\frac{3n}{2}-2) \text{ times}}, \alpha, \beta, \gamma \}$ , where  $\alpha, \beta, \gamma$  are roots of the equation  $x^3 - (n-1)x^2 - (2n+1)x + 2n^2 - 5n - 1 = 0$ .*

*Proof.* (a) If  $n$  is odd then by Theorem 2.6, we have

$$\text{ESCom}(D_{2n}) \cong K_1 \vee \underbrace{(K_1 \sqcup \dots \sqcup K_1 \sqcup K_{n-1})}_{n \text{ times}}.$$

We identify the vertex set  $\{e, a, a^2, \dots, a^{n-1}, b, ab, \dots, a^{n-1}b\}$  of  $\text{ESCom}(D_{2n})$  with the set  $\{1, 2, \dots, 2n\}$  preserving the order and observe that

$$\mathcal{A}(\text{ESCom}(D_{2n})) = \begin{bmatrix} 0 & 1 & \cdots & 1 & 1 & 1 & \cdots & 1 \\ 1 & 0 & \cdots & 1 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ 1 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \end{bmatrix}. \quad (1)$$

We use the Schur complement and the main function technique from [23, Theorem 2] to completely describe the spectrum of the matrix  $\mathcal{A}(\text{ESCom}(D_{2n}))$ .

Let  $A$  be the adjacency matrix of the complete graph on  $n$  vertices. Let  $A * e_1$  be the  $(n+1) \times (n+1)$  matrix obtained from the matrix  $A$  as follows.

$$A * e_1 = \left[ \begin{array}{cccc|c} 0 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 1 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & \cdots & 0 & 0 \\ \hline 1 & 0 & \cdots & 0 & 0 \end{array} \right].$$

We observe that the matrix  $\mathcal{A}(\text{ESCom}(D_{2n}))$  is equal to the matrix

$$A_{n*n} := \underbrace{(((A * e_1) * e_1) \cdots) * e_1}_{n \text{ times}}.$$

For example, the matrix  $A_{2*4}$  is given below

$$A_{2*4} = \left[ \begin{array}{cc|cccc} 0 & 1 & | & 1 & 1 & 1 & 1 \\ 1 & 0 & | & 0 & 0 & 0 & 0 \\ \hline & & & \dots & & & \\ 1 & 0 & | & 0 & 0 & 0 & 0 \\ 1 & 0 & | & 0 & 0 & 0 & 0 \\ 1 & 0 & | & 0 & 0 & 0 & 0 \\ 1 & 0 & | & 0 & 0 & 0 & 0 \end{array} \right].$$

Note that  $A_{n*0}$  is the adjacency matrix of the graph  $K_n$ . We observe that

$$P_{A_{n*n}}(\lambda) := \det(\lambda I - A_{n*n}) = \det \begin{bmatrix} \lambda I - A_{n*(n-1)} & -e_1 \\ -e_1^t & \lambda \end{bmatrix}.$$

Now, by using Lemma 2.3 and Lemma 2.4, we have

$$\begin{aligned} \det(\lambda I - A_{n*n}) &= \det \begin{bmatrix} \lambda I - A_{n*(n-1)} & -e_1 \\ -e_1^t & \lambda \end{bmatrix} \\ &= \lambda \det(\lambda I - A_{n*(n-1)} - \frac{1}{\lambda} e_1 e_1^t) \\ &= \lambda(1 - \frac{1}{\lambda} T) P_{A_{n*(n-1)}}(\lambda), \end{aligned}$$

where  $T = e_1^t (\lambda I - A_{n*(n-1)})^{-1} e_1$ . Simplifying this expression for  $\Gamma$ , we get

$$T = \frac{(\text{Adj}(\lambda I - A_{n*(n-1)}))_{1,1}}{P_{A_{n*(n-1)}}(\lambda)} = \frac{\lambda^{n-1} P_{A_{(n-1)*0}}(\lambda)}{P_{A_{n*(n-1)}}(\lambda)}.$$

Substituting the value of  $\Gamma$  in the previous equation, we get the following recurrence relations.

$$P_{A_{n*n}}(\lambda) = \lambda \left( 1 - \frac{\lambda^{n-1} P_{A_{(n-1)*0}}(\lambda)}{\lambda P_{A_{n*(n-1)}}(\lambda)} \right) P_{A_{n*(n-1)}}(\lambda) = \lambda P_{A_{n*(n-1)}}(\lambda) - \lambda^{n-1} P_{A_{(n-1)*0}}$$

i.e.,

$$P_{A_{n*n}}(\lambda) = \lambda P_{A_{n*(n-1)}}(\lambda) - \lambda^{n-1} P_{A_{(n-1)*0}}(\lambda). \quad (2)$$

Expanding the middle term recursively, we get

$$P_{A_{n**n}}(\lambda) = \lambda^n P_{A_{n*0}}(\lambda) - n\lambda^{n-1} P_{A_{(n-1)*0}}(\lambda). \quad (3)$$

Equation (3) gives relation between the characteristic polynomial of the adjacency matrix of  $\text{ESCom}(D_{2n})$  and the characteristic polynomial of the adjacency matrix of the complete graphs of smaller size. Now, by substituting the values for  $P_{A_{n*0}}(\lambda)$  and  $P_{A_{(n-1)*0}}$  and simplifying, we get

$$\begin{aligned} P_{A_{n**n}}(\lambda) &= \lambda^n((\lambda - (n - 1))(\lambda + 1)^{n-1}) - n\lambda^{n-1}((\lambda - (n - 2))(\lambda + 1)^{n-2}) \\ &= \lambda^{n-1}(\lambda + 1)^{n-2}(\lambda(\lambda - (n - 1))(\lambda + 1) - n(\lambda - (n - 2))) \\ &= \lambda^{n-1}(\lambda + 1)^{n-2}(\lambda^3 - (n - 2)\lambda^2 - (2n - 1)\lambda + n(n - 2)). \end{aligned}$$

(b) If  $n$  is even then by Theorem 2.6, we have

$$\text{ESCom}(D_{2n}) \cong K_2 \vee \underbrace{(K_2 \sqcup \cdots \sqcup K_2)}_{\frac{n}{2} \text{ times}} \sqcup K_{n-2}.$$

We identify the vertex set  $\{e, a^{\frac{n}{2}}, a, a^2, \dots, a^{\frac{n}{2}-1}, a^{\frac{n}{2}+1}, \dots, a^{n-1}, b, a^{\frac{n}{2}}b, ab, a^{\frac{n}{2}+1}b, \dots, a^{\frac{n}{2}-1}b, a^{n-1}b\}$  of  $\text{ESCom}(D_{2n})$  with the set  $\{1, 2, \dots, 2n\}$  preserving the order and observe that

$$\mathcal{A}(\text{ESCom}(D_{2n})) = \begin{bmatrix} 0 & 1 & 1 & \cdots & 1 & 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & 1 & \cdots & 1 & 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & 0 & \cdots & 1 & 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & 1 & \cdots & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 1 & 0 & \cdots & 0 & 0 & 1 & \cdots & 0 & 0 \\ 1 & 1 & 0 & \cdots & 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 & 1 \\ 1 & 1 & 0 & \cdots & 0 & 0 & 0 & \cdots & 1 & 0 \end{bmatrix}. \quad (4)$$

Again, we use the Schur complement and the main function technique as in the previous case. Since the calculation is the same, we skip it. This will give us the following characteristic polynomial of  $\mathcal{A}(\text{ESCom}(D_{2n}))$

$$(x - 1)^{\frac{n}{2}-1}(x + 1)^{\frac{3n}{2}-2}(x^3 - (n - 1)x^2 - (2n + 1)x + 2n^2 - 5n - 1).$$

This completes the proof.  $\square$



**Remark 1.** In the proof of the above theorem, the characteristic polynomial of the matrix given in Equation 1 can be calculated directly by expanding the determinant of  $(\lambda I - A)$  along the last column. But the same method gets complicated in the case of the matrix given in Equation 4. So we use Schur complement and the main function technique, which works for odd and even cases uniformly, and the resulting calculations are much simpler.

**Theorem 3.2.** Let  $Q_{4n} = \langle a, b : a^{2n} = e, a^n = b^2, bab^{-1} = a^{-1} \rangle$  the dicyclic group. Then

$$\text{Spec}(\text{ESCom}(Q_{4n})) = \{ \underbrace{1, \dots, 1}_{(n-1) \text{ times}}, \underbrace{-1, \dots, -1}_{(3n-2) \text{ times}}, \alpha, \beta, \gamma \},$$

where  $\alpha, \beta, \gamma$  are roots of the equation  $x^3 - (2n-1)x^2 - (4n+1)x + 8n^2 - 10n - 1 = 0$ .

*Proof.* By Theorem 2.6 we have  $\text{ESCom}(Q_{4n}) \cong \text{ESCom}(D_{2 \times 2n})$ . Hence, the result follows from Theorem 2.6.  $\square$

**Theorem 3.3.** Let  $D_{2n} = \langle a, b : a^n = b^2 = e, bab^{-1} = a^{-1} \rangle$  be the dihedral group of order  $2n$ .

(a) If  $n$  is odd then  $\text{Spec}(\text{CSCCom}(D_{2n})) = \{ \underbrace{-1, \dots, -1}_{(2n-3) \text{ times}}, \alpha, \beta, \gamma \}$ , where  $\alpha, \beta, \gamma$  are the roots of the equation  $x^3 + (3-2n)x^2 + (n^2 - 5n + 3)x + 2n^2 - 4n + 1$ .

(b) If  $n$  and  $\frac{n}{2}$  are even then  $\text{Spec}(\text{CSCCom}(D_{2n})) = \{ \underbrace{-1, \dots, -1}_{(2n-4) \text{ times}}, \frac{n}{2} - 1, \alpha, \beta, \gamma \}$ , where  $\alpha, \beta, \gamma$  are the roots of the equation  $x^3 + (3 - \frac{3n}{2})x^2 + (\frac{n^2}{2} - 5n + 3)x + \frac{5n^2}{2} - \frac{15n}{2} + 1 = 0$ .

(c) If  $n$  is even and  $\frac{n}{2}$  is odd then  $\text{Spec}(\text{CSCCom}(D_{2n})) = \{ \underbrace{-1, \dots, -1}_{(2n-4) \text{ times}}, \alpha, \beta, \gamma, \delta \}$ , where  $\alpha, \beta, \gamma, \delta$  are the roots of the equation  $x^4 + (4-2n)x^3 + (n^2 - 8n + 6)x^2 + (4n^2 - 14n + 4)x + 3n^2 - 8n + 1 = 0$ .

*Proof.* (a) If  $n$  is odd then, by Theorem 2.6, we have

$$\text{CSCCom}(D_{2n}) \cong \Delta[K_1, \underbrace{K_2, \dots, K_2}_{(\frac{n-1}{2}) \text{ times}}, K_n],$$

where  $\Delta = K_1 \vee (K_1 \sqcup K_{\frac{n-1}{2}}) = K_1 \vee (K_{\frac{n-1}{2}} \sqcup K_1)$ . Therefore,

$$\mathcal{A}(\Delta) = \begin{bmatrix} 0 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 1 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \end{bmatrix} \text{ and so } \rho_{i,j} = \begin{cases} 0, & \text{if } i = j \text{ or} \\ & 2 \leq i \leq \frac{n+1}{2} \text{ and } j = \frac{n+3}{2} \text{ or} \\ & i = \frac{n+3}{2} \text{ and } 2 \leq j \leq \frac{n+1}{2} \\ 1, & \text{otherwise.} \end{cases}$$

Hence, by Theorem 2.2, it follows that

$$\text{Spec}(\text{CSCom}(D_{2n})) = \underbrace{\{-1, \dots, -1\}}_{\binom{3n-3}{2} \text{ times}} \cup \text{Spec}(\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))),$$

where  $\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))$  is a matrix of size  $\frac{n+3}{2}$  given by

$$\begin{aligned} \tilde{\mathcal{A}}(\text{CSCom}(D_{2n})) &= \begin{bmatrix} 0 & \sqrt{2} & \sqrt{2} & \cdots & \sqrt{2} & \sqrt{n} \\ \sqrt{2} & 1 & 2 & \cdots & 2 & 0 \\ \sqrt{2} & 2 & 1 & \cdots & 2 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \sqrt{2} & 2 & 2 & \cdots & 1 & 0 \\ \sqrt{n} & 0 & 0 & \cdots & 0 & n-1 \end{bmatrix} \\ &= \left[ \begin{array}{c|cccccc|c} 0 & \sqrt{2} & \sqrt{2} & \cdots & \sqrt{2} & \sqrt{2} & \sqrt{n} \\ \hline \sqrt{2} & 1 & 2 & \cdots & 2 & 2 & 0 \\ \sqrt{2} & 2 & 1 & \cdots & 2 & 2 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & 0 \\ \sqrt{2} & 2 & \vdots & \cdots & 1 & 2 & \vdots \\ \sqrt{2} & 2 & \cdots & \cdots & 2 & 1 & 0 \\ \hline \sqrt{n} & 0 & \cdots & 0 & 0 & 0 & n-1 \end{array} \right]. \end{aligned}$$

Note that the middle block of  $\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))$  is  $2J_{\frac{n-1}{2}} - I_{\frac{n-1}{2}}$  whose eigenvalues are  $-1$  with multiplicity  $\frac{n-3}{2} = \frac{n-1}{2} - 1$  and  $n-2$  with multiplicity 1. These eigenvalues are useful in obtaining the eigenvalues of  $\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))$ . We obtain the characteristic polynomial of  $\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))$  by expanding along the last column as given below:

$$\begin{aligned}
& P_{\tilde{A}(\text{CSCom}(D_{2n}))}(\lambda) \\
&= (-1)^{2+\frac{n+3}{2}} \sqrt{n} \det \begin{bmatrix} -\sqrt{2} & | & \lambda-1 & -2 & \cdots & \cdots & -2 \\ -\sqrt{2} & | & -2 & \lambda-1 & \cdots & \cdots & \vdots \\ -\sqrt{2} & | & -2 & -2 & \cdots & \cdots & \vdots \\ \vdots & | & \vdots & \vdots & \ddots & \vdots & \vdots \\ -\sqrt{2} & | & -2 & \cdots & \cdots & -2 & \lambda-1 \\ \hline -\sqrt{n} & | & 0 & \cdots & \cdots & 0 & 0 \end{bmatrix} \\
&\quad + (-1)^{n+3}(n-1) \det \begin{bmatrix} \lambda & | & -\sqrt{2} & -\sqrt{2} & \cdots & -\sqrt{2} \\ \hline -\sqrt{2} & | & \lambda-1 & -2 & \cdots & -2 \\ \vdots & | & \vdots & \vdots & \vdots & \vdots \\ -\sqrt{2} & | & -2 & \cdots & -2 & \lambda-1 \end{bmatrix} \\
&= (-1)^{(2+\frac{n+3}{2}+2+\frac{n+1}{2})} n(\lambda+1)^{\frac{n-3}{2}} \cdot (\lambda - (n-2)) + (-1)^{n+3}(\lambda - (n-1)) \det(\lambda I - \bar{A}) \\
&= (-1) \cdot n \cdot (\lambda+1)^{\frac{n-3}{2}} \cdot (\lambda - (n-2)) + (\lambda - (n-1)) \det(\lambda I - \bar{A}),
\end{aligned}$$

where

$$\bar{A} := \begin{bmatrix} 0 & | & \sqrt{2} & \sqrt{2} & \cdots & \sqrt{2} \\ \hline \sqrt{2} & | & 1 & 2 & \cdots & 2 \\ \vdots & | & \vdots & \vdots & \vdots & \vdots \\ \sqrt{2} & | & 2 & \cdots & 2 & 1 \end{bmatrix}.$$

To calculate the characteristic polynomial of  $\bar{A}$ , we consider  $\bar{A}$  as follows.

$$\bar{A} = \begin{bmatrix} 0 & | & \sqrt{2} & \sqrt{2} & \cdots & \sqrt{2} \\ \hline \sqrt{2} & | & 1 & 2 & \cdots & 2 \\ \vdots & | & \vdots & \vdots & \vdots & \vdots \\ \sqrt{2} & | & 2 & \cdots & 2 & 1 \end{bmatrix} := \begin{bmatrix} A & | & B \\ \hline C & | & D \end{bmatrix} \text{ (say)}.$$

By Lemma 2.3, we have

$$\det(\lambda I - \bar{A}) = \det(\lambda I - (2J_{\frac{n-1}{2}} - I_{\frac{n-1}{2}}))(\lambda - \Gamma_D(\sqrt{2} \cdot \mathbf{1}_{\frac{n-1}{2}})).$$

Since  $\sqrt{2} \cdot \mathbf{1}_{\frac{n-1}{2}}$  is an eigenvector of the matrix  $2J_{\frac{n-1}{2}} - I_{\frac{n-1}{2}}$  corresponding to the eigenvalue  $n-2$ , by Lemma 2.5, we have

$$\Gamma_D(\sqrt{2} \cdot \mathbf{1}_{\frac{n-1}{2}}) = \frac{\|\sqrt{2} \cdot \mathbf{1}_{\frac{n-1}{2}}\|^2}{\lambda - (n-2)} = \frac{n-1}{\lambda - (n-2)}.$$

Substituting this value of the main function in the above equation, we get

$$\begin{aligned} \det(\lambda I - \bar{A}) &= \det(\lambda I - (2J_{\frac{n-1}{2}} - I_{\frac{n-1}{2}})) \left( \lambda - \frac{n-1}{\lambda - (n-2)} \right) \\ &= \det(\lambda I - (2J_{\frac{n-1}{2}} - I_{\frac{n-1}{2}})) \left( \frac{\lambda^2 - \lambda(n-2) - (n-1)}{\lambda - (n-2)} \right) \\ &= (\lambda+1)^{\binom{n-3}{2}} (\lambda - (n-2)) \left( \frac{\lambda^2 - \lambda(n-2) - (n-1)}{\lambda - (n-2)} \right) \\ &= (\lambda+1)^{\binom{n-3}{2}} (\lambda^2 - \lambda(n-2) - (n-1)). \end{aligned}$$

which is the required characteristic polynomial of the matrix  $\bar{A}$ . Therefore,

$$\begin{aligned} P_{\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))}(\lambda) &= (-1) \cdot n \cdot (\lambda+1)^{\frac{n-3}{2}} \cdot (\lambda - (n-2)) + (\lambda - (n-1)) (\lambda+1)^{\binom{n-3}{2}} (\lambda^2 - \lambda(n-2) - (n-1)) \\ &= (\lambda+1)^{\frac{n-3}{2}} \cdot (\lambda^3 + (3-2n)\lambda^2 + (n^2 - 5n + 3)\lambda + (2n^2 - 4n + 1)). \end{aligned}$$

(b) If  $n$  and  $\frac{n}{2}$  are even then, by Theorem 2.6, we have

$$\text{CSCom}(D_{2n}) \cong \Delta[K_1, K_1, \underbrace{K_2, \dots, K_2}_{\binom{n}{2}-1 \text{ times}}, K_{\frac{n}{2}}, K_{\frac{n}{2}}],$$

where  $\Delta = K_2 \vee (K_1 \sqcup K_1 \sqcup K_{\frac{n}{2}-1})$ . Therefore,

$$\mathcal{A}(\Delta) = \begin{bmatrix} 0 & 1 & 1 & \cdots & 1 & 1 & 1 \\ 1 & 0 & 1 & \cdots & 1 & 1 & 1 \\ 1 & 1 & 0 & \cdots & 1 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 1 & 1 & 1 & \cdots & 0 & 0 & 0 \\ 1 & 1 & 0 & \cdots & 0 & 0 & 0 \\ 1 & 1 & 0 & \cdots & 0 & 0 & 0 \end{bmatrix} \text{ and so } \rho_{i,j} = \begin{cases} 0, & \text{if } i = j \text{ or} \\ & 3 \leq i \leq \frac{n}{2} + 2 \text{ and } j = \frac{n}{2} + 2, \frac{n}{2} + 3 \text{ or} \\ & i = \frac{n}{2} + 2, \frac{n}{2} + 3 \text{ and } 3 \leq j \leq \frac{n}{2} + 2 \\ 1, & \text{otherwise.} \end{cases}$$

Hence, by Theorem 2.2, it follows that

$$\text{Spec}(\text{CSCom}(D_{2n})) = \underbrace{\{-1, \dots, -1\}}_{\left(\frac{3n}{2} - 3\right) \text{ times}} \cup \text{Spec}(\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))),$$

where  $\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))$  is a matrix of size  $\frac{n}{2} + 3$  given by

$$\tilde{\mathcal{A}}(\text{CSCom}(D_{2n})) = \begin{bmatrix} 0 & 1 & \sqrt{2} & \sqrt{2} & \cdots & \sqrt{2} & \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ 1 & 0 & \sqrt{2} & \sqrt{2} & \cdots & \sqrt{2} & \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ \sqrt{2} & \sqrt{2} & 1 & 2 & \cdots & 2 & 0 & 0 \\ \sqrt{2} & \sqrt{2} & 2 & 1 & \cdots & 2 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \sqrt{2} & \sqrt{2} & 2 & \cdots & 2 & 1 & 0 & 0 \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & \cdots & 0 & \frac{n}{2} - 1 & 0 \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & \cdots & 0 & 0 & \frac{n}{2} - 1 \end{bmatrix}.$$

By taking the bottom right corner submatrix of size  $2 \times 2$  as  $D$ , and applying Lemma 2.3, we get

$$P_{\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))}(\lambda) = \det \begin{bmatrix} \lambda - \left(\frac{n}{2} - 1\right) & 0 \\ 0 & \lambda - \left(\frac{n}{2} - 1\right) \end{bmatrix} \times \\ \det \left( A - \begin{bmatrix} \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{\lambda - \left(\frac{n}{2} - 1\right)} & 0 \\ 0 & \frac{1}{\lambda - \left(\frac{n}{2} - 1\right)} \end{bmatrix} \begin{bmatrix} \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & 0 \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & 0 \end{bmatrix} \right).$$

Letting  $a = \frac{1}{\lambda - \left(\frac{n}{2} - 1\right)}$ , we get

$$P_{\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))}(\lambda) = \frac{1}{a^2} \det \left( A - \begin{bmatrix} \sqrt{\frac{n}{2}}a & \sqrt{\frac{n}{2}}a \\ \sqrt{\frac{n}{2}}a & \sqrt{\frac{n}{2}}a \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & 0 \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & 0 \end{bmatrix} \right)$$

$$\begin{aligned}
&= \frac{1}{a^2} \det \left( A - \begin{bmatrix} na & na & 0 & \cdots & 0 \\ na & na & 0 & \cdots & 0 \\ 0 & 0 & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & \cdots & 0 \end{bmatrix} \right) \\
&= \frac{1}{a^2} \det \left( \begin{bmatrix} \lambda - na & -1 - na & -\sqrt{2} & -\sqrt{2} & \cdots & -\sqrt{2} \\ -1 - na & \lambda - na & -\sqrt{2} & -\sqrt{2} & \cdots & -\sqrt{2} \\ -\sqrt{2} & -\sqrt{2} & \lambda - 1 & -2 & \cdots & -2 \\ -\sqrt{2} & -\sqrt{2} & -2 & \lambda - 1 & \cdots & -2 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ -\sqrt{2} & -\sqrt{2} & -2 & -2 & \cdots & \lambda - 1 \end{bmatrix} \right).
\end{aligned}$$

Again using Lemma 2.3, this time taking the bottom right corner submatrix of size  $\frac{n}{2}-1 \times \frac{n}{2}-1$  as  $D$ , we get

$$\begin{aligned}
P_{\tilde{\mathcal{A}}(\text{CSCCom}(D_{2n}))}(\lambda) &= \frac{1}{a^2} (\lambda + 1)^{\frac{n}{2}-2} (\lambda - (n-3)) \times \\
&\det \left( \begin{bmatrix} \lambda - na & -1 - na \\ -1 - na & \lambda - na \end{bmatrix} - \begin{bmatrix} \Gamma_{(\lambda I_{\frac{n}{2}-1} - D)}(\sqrt{2} \cdot \mathbf{1}_{\frac{n}{2}-1}) & \Gamma_{(\lambda I_{\frac{n}{2}-1} - D)}(\sqrt{2} \cdot \mathbf{1}_{\frac{n}{2}-1}) \\ \Gamma_{(\lambda I_{\frac{n}{2}-1} - D)}(\sqrt{2} \cdot \mathbf{1}_{\frac{n}{2}-1}) & \Gamma_{(\lambda I_{\frac{n}{2}-1} - D)}(\sqrt{2} \cdot \mathbf{1}_{\frac{n}{2}-1}) \end{bmatrix} \right).
\end{aligned}$$

Since  $\sqrt{2} \cdot \mathbf{1}_{\frac{n}{2}-1}$  is an eigenvector of  $(\lambda I_{\frac{n}{2}-1} - D)$  corresponding to the eigenvalue  $n-3$ , by Lemma 2.5, we have

$$\begin{aligned}
&P_{\tilde{\mathcal{A}}(\text{CSCCom}(D_{2n}))}(\lambda) \\
&= \frac{1}{a^2} (\lambda + 1)^{\frac{n}{2}-2} (\lambda - (n-3)) \det \left( \begin{bmatrix} \lambda - na & -1 - na \\ -1 - na & \lambda - na \end{bmatrix} - \begin{bmatrix} \frac{n-2}{\lambda-(n-3)} & \frac{n-2}{\lambda-(n-3)} \\ \frac{n-2}{\lambda-(n-3)} & \frac{n-2}{\lambda-(n-3)} \end{bmatrix} \right) \\
&= \frac{1}{a^2} (\lambda + 1)^{\frac{n}{2}-2} (\lambda - (n-3)) \left[ \left( (\lambda - na) - \frac{(n-2)}{\lambda - (n-3)} \right)^2 - \left( (-1 - na) - \frac{(n-2)}{\lambda - (n-3)} \right)^2 \right] \\
&= \frac{1}{a^2} (\lambda + 1)^{\frac{n}{2}-2} (\lambda - (n-3)) \left( \lambda - 1 - 2na - \frac{2(n-2)}{\lambda - (n-3)} \right) (\lambda + 1) \\
&= \frac{1}{a^2} (\lambda + 1)^{\frac{n}{2}-1} \left( (\lambda - 1 - 2na)(\lambda - (n-3)) - 2(n-2) \right) \\
&= (\lambda - (\frac{n}{2} - 1)) (\lambda + 1)^{\frac{n}{2}-1} \left( (\lambda^3 + (3 - \frac{3n}{2})\lambda^2 + (\frac{n^2}{2} - 5n + 3)\lambda + (\frac{5n^2}{2} - \frac{15n}{2} + 1)) \right).
\end{aligned}$$

Hence, the result follows.

(c) If  $n$  even and  $\frac{n}{2}$  is odd then, by Theorem 2.6, we have

$$\text{CSCom}(D_{2n}) \cong \Delta[K_1, K_1, \underbrace{K_2, \dots, K_2}_{(\frac{n}{2}-1) \text{ times}}, K_{\frac{n}{2}}, K_{\frac{n}{2}}],$$

where  $\Delta = K_2 \vee (K_2 \sqcup K_{\frac{n}{2}-1})$ . Therefore,

$$\mathcal{A}(\Delta) = \begin{bmatrix} 0 & 1 & 1 & \cdots & 1 & 1 & 1 \\ 1 & 0 & 1 & \cdots & 1 & 1 & 1 \\ 1 & 1 & 0 & \cdots & 1 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 1 & 1 & 1 & \cdots & 0 & 0 & 0 \\ 1 & 1 & 0 & \cdots & 0 & 0 & 1 \\ 1 & 1 & 0 & \cdots & 0 & 1 & 0 \end{bmatrix} \text{ and so } \rho_{i,j} = \begin{cases} 0, & \text{if } i = j \text{ or} \\ & 3 \leq i \leq \frac{n}{2} + 1 \text{ and } j = \frac{n}{2} + 2, \frac{n}{2} + 3 \text{ or} \\ & i = \frac{n}{2} + 2, \frac{n}{2} + 3 \text{ and } 3 \leq j \leq \frac{n}{2} + 1 \\ 1, & \text{otherwise.} \end{cases}$$

Hence, by Theorem 2.2, it follows that

$$\text{Spec}(\text{CSCom}(D_{2n})) = \underbrace{\{-1, \dots, -1\}}_{(\frac{3n}{2}-3) \text{ times}} \cup \text{Spec}(\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))),$$

where  $\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))$  is a matrix of size  $\frac{n}{2} + 3$  given by

$$\tilde{\mathcal{A}}(\text{CSCom}(D_{2n})) = \begin{bmatrix} 0 & 1 & \sqrt{2} & \sqrt{2} & \cdots & \sqrt{2} & \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ 1 & 0 & \sqrt{2} & \sqrt{2} & \cdots & \sqrt{2} & \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ \sqrt{2} & \sqrt{2} & 1 & 2 & \cdots & 2 & 0 & 0 \\ \sqrt{2} & \sqrt{2} & 2 & 1 & \cdots & 2 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \sqrt{2} & \sqrt{2} & 2 & \cdots & 2 & 1 & 0 & 0 \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & \cdots & 0 & \frac{n}{2} - 1 & \frac{n}{2} \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & \cdots & 0 & \frac{n}{2} & \frac{n}{2} - 1 \end{bmatrix}.$$

As before, by taking  $b = \frac{n}{2} - 1$  and the bottom right corner submatrix of size  $2 \times 2$  as  $D$ , and applying Lemma 2.3, we get

$$\begin{aligned} & P_{\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))}(\lambda) \\ &= \det \begin{bmatrix} \lambda - b & -\frac{n}{2} \\ -\frac{n}{2} & \lambda - b \end{bmatrix} \det \left( A - \begin{bmatrix} \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \lambda - b & -\frac{n}{2} \\ -\frac{n}{2} & \lambda - b \end{bmatrix}^{-1} \begin{bmatrix} \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & 0 \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & 0 \end{bmatrix} \right), \end{aligned}$$

$$\begin{aligned}
&= \left( (\lambda - b)^2 - \frac{n^2}{4} \right) \det \left( A - \begin{bmatrix} \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{bmatrix} \frac{1}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} \begin{bmatrix} \lambda - b & \frac{n}{2} \\ \frac{n}{2} & \lambda - b \end{bmatrix} \begin{bmatrix} \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & 0 \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & 0 \end{bmatrix} \right) \\
&= \left( (\lambda - b)^2 - \frac{n^2}{4} \right) \det \left( A - \frac{1}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} \begin{bmatrix} \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \lambda - b & \frac{n}{2} \\ \frac{n}{2} & \lambda - b \end{bmatrix} \begin{bmatrix} \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & 0 \\ \sqrt{\frac{n}{2}} & \sqrt{\frac{n}{2}} & 0 & \cdots & 0 \end{bmatrix} \right) \\
&= \left( (\lambda - b)^2 - \frac{n^2}{4} \right) \times \\
&\quad \det \left( A - \frac{1}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} \begin{bmatrix} 2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right) & 2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right) & 0 & \cdots & 0 \\ 2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right) & 2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right) & 0 & \cdots & 0 \\ 0 & 0 & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & \cdots & 0 \end{bmatrix} \right) \\
&= \left( (\lambda - b)^2 - \frac{n^2}{4} \right) \det \left( \begin{bmatrix} \lambda - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} & -1 - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} & -\sqrt{2} & -\sqrt{2} & \cdots & -\sqrt{2} \\ -1 - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} & \lambda - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} & -\sqrt{2} & -\sqrt{2} & \cdots & -\sqrt{2} \\ -\sqrt{2} & -\sqrt{2} & \lambda - 1 & -2 & \cdots & -2 \\ -\sqrt{2} & -\sqrt{2} & -2 & \lambda - 1 & \cdots & -2 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ -\sqrt{2} & -\sqrt{2} & -2 & -2 & \cdots & \lambda - 1 \end{bmatrix} \right).
\end{aligned}$$

Again using Lemma 2.3, this time taking the bottom right corner submatrix of size  $\frac{n}{2} - 1 \times \frac{n}{2} - 1$  as  $D$ , we get



$$\begin{aligned}
& P_{\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))}(\lambda) \\
&= \left( (\lambda - b)^2 - \frac{n^2}{4} \right) (\lambda + 1)^{\frac{n}{2}-2} (\lambda - (n - 3)) \det \left( \begin{bmatrix} \lambda - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} & -1 - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} \\ -1 - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} & \lambda - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} \end{bmatrix} \right. \\
&\quad \left. - \begin{bmatrix} \Gamma_{\lambda I_{\frac{n}{2}-1} - D}(\sqrt{2} \cdot \mathbf{1}_{\frac{n}{2}-1}) & \Gamma_{\lambda I_{\frac{n}{2}-1} - D}(\sqrt{2} \cdot \mathbf{1}_{\frac{n}{2}-1}) \\ \Gamma_{\lambda I_{\frac{n}{2}-1} - D}(\sqrt{2} \cdot \mathbf{1}_{\frac{n}{2}-1}) & \Gamma_{\lambda I_{\frac{n}{2}-1} - D}(\sqrt{2} \cdot \mathbf{1}_{\frac{n}{2}-1}) \end{bmatrix} \right).
\end{aligned}$$

Since  $\sqrt{2} \cdot \mathbf{1}_{\frac{n}{2}-1}$  is an eigenvector of  $\lambda I_{\frac{n}{2}-1} - D$  corresponding to the eigenvalue  $n - 3$ , By Lemma 2.5, we have

$$\begin{aligned}
P_{\tilde{\mathcal{A}}(\text{CSCom}(D_{2n}))}(\lambda) &= \left( (\lambda - b)^2 - \frac{n^2}{4} \right) (\lambda + 1)^{\frac{n}{2}-2} (\lambda - (n - 3)) \\
&\quad \times \det \left( \begin{bmatrix} \lambda - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} & -1 - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} \\ -1 - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} & \lambda - \frac{2 \left( (\lambda - b) \frac{n}{2} + \left( \frac{n^2}{4} \right) \right)}{\left( (\lambda - b)^2 - \frac{n^2}{4} \right)} \end{bmatrix} - \begin{bmatrix} \frac{n-2}{\lambda-(n-3)} & \frac{n-2}{\lambda-(n-3)} \\ \frac{n-2}{\lambda-(n-3)} & \frac{n-2}{\lambda-(n-3)} \end{bmatrix} \right) \\
&= (\lambda + 1)^{\frac{n}{2}-1} (\lambda^4 + (4 - 2n)\lambda^3 + (n^2 - 8n + 6)\lambda^2 + (4n^2 - 14n + 4)\lambda + (3n^2 - 8n + 1)).
\end{aligned}$$

Hence, the result follows.  $\square$

**Theorem 3.4.** *Let  $Q_{4n} = \langle a, b : a^{2n} = e, a^n = b^2, bab^{-1} = a^{-1} \rangle$  be the dicyclic group of order  $4n$ .*

- (a) *If  $n$  is even then  $\text{Spec}(\text{CSCom}(Q_{4n})) = \{ \underbrace{-1, \dots, -1}_{(4n-4) \text{ times}}, n-1, \alpha, \beta, \gamma \}$ , where  $\alpha, \beta, \gamma$  are the roots of the equation  $x^3 + (3 - 3n)x^2 + (2n^2 - 10n + 3)x + 10n^2 - 15n + 1 = 0$ .*
- (b) *If  $n$  is odd then  $\text{Spec}(\text{CSCom}(Q_{4n})) = \{ \underbrace{-1, \dots, -1}_{(4n-4) \text{ times}}, \alpha, \beta, \gamma, \delta \}$ , where  $\alpha, \beta, \gamma, \delta$  are the roots of the equation  $x^4 + (4 - 4n)x^3 + (4n^2 - 16n + 6)x^2 + (16n^2 - 28n + 4)x + 12n^2 - 16n + 1 = 0$ .*

*Proof.* By Theorem 2.6 we have  $\text{CSCom}(Q_{4n}) \cong \text{CSCom}(D_{2 \times 2n})$ . Hence, the result follows from Theorem 3.3.  $\square$

Notice that the cubic and quartic equations appearing in Theorems 3.1 – 3.4 have non-integral roots. For instances, the equations  $x^3 - 3x^2 - 11x + 15 = 0$  has roots  $\alpha \approx -2.81114, \beta \approx 1.14307, \gamma \approx 4.66807$ ;  $x^3 - 5x^2 - 13x + 41 = 0$  has roots  $\alpha \approx -3.17226, \beta \approx 2.14399, \gamma \approx 6.02827$ ;  $x^3 - 7x^2 + 3x + 31 = 0$  has roots  $\alpha \approx -1.72119, \beta \approx 3.35861, \gamma \approx 5.36258$ ;  $x^4 - 8x^3 - 6x^2 + 64x + 61 = 0$  has roots  $\alpha = -1, \beta \approx -2.2034, \gamma = 3.6798, \delta = 7.5236$  etc.

## References

- [1] N. Abreu, D. Cardoso, P. Carvalho and C. Vinagre, Spectra and Laplacian spectra of arbitrary powers of lexicographic products of graphs, *Discrete Math.* **340** (2017), 3235–3244.
- [2] G. Arunkumar, P.J. Cameron, R.K.Nath and L. Selvaganesh, Super graphs on groups, I, *Graphs Combin.* **38** (2022), 100 (14 pages).
- [3] G. Arunkumar, P.J. Cameron, R.K. Nath, Super graphs on groups, II, *Discrete Appl. Math.* Accepted, <https://doi.org/10.48550/arXiv.2401.09912>.
- [4] L. Babai, Spectra of Cayley graphs, *J. Combin. Theory Ser. B* **27** (1979), 180–189.
- [5] P. Bhowal and R.K. Nath, Spectral aspects of commuting conjugacy class graph of finite groups, *Algebr. Struct. their Appl.* **8** (2021), 67–118.
- [6] P.J. Cameron, Graphs defined on groups, *Int. J. Group Theory* **11** (2022), 43–124.
- [7] D.M. Cardoso, M.A. de Freitas, E.A. Martins, and M. Robbiano, Spectra of graphs obtained by a generalization of the join graph operation, *Discrete Math.* **313** (2013), 733–741.
- [8] D. Cvetković, The generating function for variations with restrictions and paths of the graph and self-complementary graphs, *Univ Beograd Publ Elektrotehn Fak Ser Mat Fiz*, 320–328 (1970), 27–34.
- [9] S. Dalal, S. Mukherjee and K. L. Patra, Spectrum of conjugacy and order super commuting graphs of some finite groups, <https://doi.org/10.48550/arXiv.2312.08930>
- [10] J. Dutta and R.K. Nath, Spectrum of commuting graphs of some classes of finite groups, *Matematika* **33** (2017), 87–95.

- [11] W.N.T. Fasfous, R.K. Nath and R. Sharafadini, Various spectra and energies of commuting graphs of finite rings, *Hacet. J. Math. Stat.* **49** (2020), 1915–1925.
- [12] A. Gerbaud, Spectra of generalized compositions of graphs and hierarchical networks, *Discrete Math.* **310** (2010), 2824–2830.
- [13] C.Y. Ku, T. Lau, K.B. Wong, The smallest eigenvalues of the 1-point fixing graph, *Linear Algebra Appl.* **493** (2016), 433–446.
- [14] J. Lan, B. Zhou, Spectra of graph operations based on  $R$ -graph, *Linear Multilinear Algebra* **63**, (2015), 1401–1422.
- [15] X. Liu, P. Lu, Spectra of subdivision-vertex and subdivision-edge neighbourhood coronae, *Linear Algebra Appl.* **438** (2013), 3547–3559.
- [16] Y. Mao, Z. Wang and I. Gutman, Steiner Wiener index of graph products, *Trans. Comb.* **5** (2016), 39–50.
- [17] C. McLeman, E. McNicholas, Spectra of coronae, *Linear Algebra Appl.* **435** (2011), 998–1007.
- [18] Z. Mehranian, A. Gholami, A. R. Ashrafi, The Spectra of power graphs of certain finite groups, *Linear Multilinear Algebra* **65** (2017), 1003–1010.
- [19] K. Mönius, Eigenvalues of zero-divisor graphs of finite commutative rings, *J. Algebraic Combin.* **54**, (2011), 787–802.
- [20] R. Rajkumar and R. Pavithra, Spectra of  $M$ -rooted product of graphs, *Linear Multilinear Algebra* **70** (2022), 1–26.
- [21] P. Renteln, The distance spectra of Cayley graphs of Coxeter groups, *Linear Algebra Appl.* **311** (2011), 738–755.
- [22] P. Rowlinson, The main eigenvalues of a graph: a survey, *Appl. Anal. Discrete Math.* **1** (2007), 455–471.
- [23] M. Saravanan, S. P. Murugan, G. Arunkumar, A generalization of Fiedler’s lemma and the spectra of  $H$ -join of graphs, *Linear Algebra Appl.* **625** (2021), 20–43.
- [24] A.J. Schwenk, Computing the characteristic polynomial of a graph, in: R. Bary, F. Harary (Eds.), *Graphs Combinatorics, Lecture Notes in Mathematics*, **406** (1974), 153–172, Springer-Verlag, Berlin.

- [25] Z. Wang and D. Wong, The characteristic polynomial of lexicographic product of graphs, *Linear Algebra Appl.* **541** (2018), 177–184.
- [26] S. Wang and B. Zhou, The signless Laplacian spectra of the corona and edge corona of two graphs, *Linear Multilinear Algebra* **61** (2013), 197–204.