Elementary equivalences for blocks with normal elementary abelian defect group of rank 2

Mary Schaps and Zehavit Zvi

August 5, 2024

Abstract

We consider the effect of performing an elementary equivalence as defined by Okuyama on a group block of form $F(C_p \times C_p) \rtimes C_r$, for a field F of characteristic p. If $I = \{0, 1, 2, \ldots r - 1\}$ is the set of residues corresponding to the simple modules of FC_r , the elementary equivalence is determined by a proper, non-empty subset $I_0 \subset I$, and the corresponding elementary tilting complex is completely determined by I_0 .

We give a catalog of homogeneous maps between irreducible components of the elementary tilting complex. When the subset I_0 is an interval, we prove that the maps in the catalog are sufficient to describe all homogeneous maps between two irreducible components.

1 INTRODUCTION

This work concerns blocks over a field F of odd finite characteristic p , with elementary abelian defect group $D = C_p \times C_p$. We restrict to odd characteristic because the case of $C_2 \times C_2$ has been intensively studied. The defect group $C_p \times C_p$ is the defect group with the next largest rank after the well-studied case of cyclic defect group.

We are interested in the tilting theory of these blocks, trying to apply some of the insights gathered from the cyclic defect case [\[R1,](#page-41-0) [SZ2,](#page-41-1) [SZ\]](#page-41-2). In that case, every block can be obtained from a block with normal defect group by a sequence of elementary tiltings known as mutations. Okuyama [\[O\]](#page-41-3) found some important group blocks as the result of a sequence of more general elementary equivalences in the case $p = 3$. The Broué conjecture for principal blocks when $p = 3$ was eventually proven by Koshitani and Kunugi in [\[KK\]](#page-40-0) but is still open for larger primes. Unfortunately, one of Okuyama's

central lemmas is less useful for primes $p \geq 5$, so his method had not found ready application to other primes.

A study of the blocks of subgroups of the groups in the Atlas of Simple Groups [\[C\]](#page-40-1) shows that most of the abelian defect groups are either cyclic or of the form $D = C_p \times C_p$, and all have a property we will define later as being balanced. Furthermore, the Brauer correspondents of these blocks are usually of the form $D \rtimes C_r$ or $D \rtimes D_n$. In this work, we are taking up the first case, $D \rtimes C_r$ and asking what sort of algebras can be reached by a single elementary equivalence, as a function of the prime p and the number r of simple modules in the original block of cyclic defect.

2 DEFINITIONS AND NOTATION

2.1 Blocks with normal defect groups

The blocks with normal defect groups were studied in depth by several authors [\[Lan\]](#page-41-4), [\[RR\]](#page-41-5), [\[Sw\]](#page-41-6) and also [\[K\]](#page-40-2). We will review that part of the theory which we need. By Brauer's First Main Theorem, we know that there is a one-to-one correspondence between the blocks B of FG with defect group D, and the blocks b of $FN_G(D)$ with defect group D [\[CR\]](#page-40-3). In order to investigate the relationship between B and b we must know more about the structure of the blocks of $FN_G(D)$. We restricted ourselves to Brauer correspondents which are Morita equivalent as blocks to $F[(C_p \times C_p) \rtimes H]$, with $H = C_r$ for r prime to p. We will show that we can choose a basis for the radical of $F[D]$ which are eigenvectors for the generator of C_r and we will assume that the eigenvalues of the generators of this radical are inverse to each other.

2.2 The quiver of a block

Let A be a K-algebra, where K is an algebraically closed field with $A/\operatorname{Rad}(A) \stackrel{\sim}{\to}$ $\bigoplus M_{n_i}(K)$. Let f_1, \ldots, f_t be a set of primitive idempotents, one from each matrix block, and set

$$
n'_{ij} = \dim(f_i(J/J^2)f_j)
$$

for $J = \text{Rad}(A)$.

Definition 2.1. The quiver of A is the directed graph with t vertices, where t is the number of simple modules, and n'_{ij} arrows from j to i. Then n'_{ij} is the number of irreducible morphisms from j to i .

In [\[S\]](#page-41-7), results were quoted showing that:

- 1. The block idempotents of FH embed in FG .
- 2. Rad(*FG*) is generated by $J = \text{Rad}(FD)$.
- 3. The action of H on D induces an action of H on J/J^2 with character χ , and there is an H-submodule N of J isomorphic to J/J^2 .
- 4. The quiver of G is the McKay graph $D(H, \chi)$ for the character χ of H.

Gerstenhaber and Schaps go on to show in [\[GS\]](#page-40-4) that the H-module J/J^2 can be embedded as a submodule of J of dimension d over F . If H is abelian, it decomposes as a direct sum of one-dimensional submodules, so that we can find a basis for J of idempotents for the action of H .

2.3 The McKay graph

In [\[M\]](#page-41-8), McKay introduced a graph that he called the representation graph and which later was called the McKay graph by Schaps, Shapira and Shlomo in [\[SSS\]](#page-41-9).

Let H be a finite group, and let F be a field of characteristic zero or of positive characteristic p, $p \nmid |H|$. Let K be an extension of F which is a splitting field for H . Let the character table of H be considered as having entries in K. Let χ be a character of H, with kernel N. If $\theta_1, \ldots, \theta_t$ is the complete set of irreducible characters of H, and $\chi \cdot \theta_j$ is defined by pointwise multiplication of the values for the various conjugacy classes, define the natural number n_{ij} as the multiplicity of the constituent θ_i in the product character $\chi \cdot \theta_j$. Since the θ_i form an orthonormal basis, with respect to an inner product (-, -), this is given by $n_{ij} = (\theta_i, \chi \cdot \theta_j)$.

Definition 2.2. If H is a finite group, K is a sufficiently large field of characteristic prime to $|H|$, and χ is a K character of H, the McKay graph $D(H,\chi)$ is the directed graph with

- (i) t vertices labelled by the irreducible characters $\theta_1, \ldots, \theta_t$ of H.
- (ii) For each pair (θ_i, θ_j) of characters, n_{ij} arrows from θ_j to θ_i , where

$$
n_{ij} = (\theta_i, \chi \cdot \theta_j).
$$

In ([\[SSS\]](#page-41-9), §3), it was shown that for a graph algebra of the form $(C_p^d) \rtimes H$, where $p \nmid |H|$, the quiver was given by the McKay graph, and a method was also given in [\[SSS\]](#page-41-9) for writing down the relations on the Morita equivalent basic algebra.

2.4 The group algebra $F[(C_p \times C_p) \rtimes C_r]$

We now restrict to the case most common for non-cyclic defect, the case of $d = 2$ and $H = C_r$, with r not divisible by p. By the results we quoted from $[GS]$, we can find two generators x and y for the radical J which are eigenvectors for the action of C_r . There are many ways to choose a pair of eigenvalues of the action of C_r , but what usually occurs in the examples is that there is an automorphism that interchanges x with y , and the generator of the C_r with its inverse, which means that the idempotent of y is the inverse of the idempotent of x, so that the action of C_r on xy is trivial.

Definition 2.3. Let $G = (C_p \times C_p) \rtimes C_r$, and let x and y be generators of J which are eigenvectors for the action of C_r . We will say that the action is balanced if the eigenvalue of the action of C_r on x is the inverse of the eigenvalue of y.

Example 1. If H is $C_4 = \langle a \rangle$, $G = (C_5 \times C_5) \rtimes C_4$, where the generator a of C_4 acts on the generators c and d of $C_5 \times C_5$ by sending c to c^2 and d to d^3 . Let $i \in F$ be a primitive 4th root of unity, with irreducible characters $\theta_j(a^k) = i^{(j-1)k}$, where $j = 1, ..., 4$ and $k = 0, ..., 3$. Let $e_u, u = 1, ..., 4$, be the idempotents of KC_4 . Set $\chi = \theta_2 + \theta_4$. Then,

$$
\chi \cdot \theta_j = \theta_{j+1} + \theta_{j-1}
$$

where the indices are taken mod 4. Since 2 and 3 are primitive fourth roots of unity modulo 5, we can choose the generators as follows

$$
x = c + 3c2 + 2c3 + 4c4
$$

$$
y = d + 2d3 + 3d2 + 4d4.
$$

Then $a^{-1}xa = c^2 + 3c^4 + 2c + 4c^3 = 2x$ and $a^{-1}ya = d^3 + 2d^4 + 3d + 4d^2 = 3y$ The quiver of FG is given by

with the relations:

$$
x_{u(u+1)} = e_u x e_{u+1}
$$

\n
$$
y_{u(u-1)} = e_u y e_{u-1}
$$

\n
$$
x = x_{01} + x_{12} + x_{23} + x_{30}
$$

\n
$$
y = y_{10} + x_{03} + x_{32} + x_{21}
$$

\n
$$
xy = yx
$$

\n
$$
x^5 = 0, y^5 = 0
$$

2.5 The structure of the projective indecomposable modules P_t for $F[(C_p \times C_p) \rtimes C_r]$

Since C_r is abelian, the simple modules of FG , which correspond to the simple modules of FC_r , are one-dimensional. They bijectively correspond to the idempotents of FC_r . Using the fact that powers of ξ and a can be taken modulo r, we recall that the central idempotents of FC_r are given by

$$
e_t = \frac{1}{r} \sum_{i=0}^{r-1} \xi^{-it} a^i, t = 0, \dots, r-1.
$$

We consider one particular idempotent e_t and its projective module P_t . We first consider the vector space $S_t = \langle e_t \rangle$, and show that it is a one-dimensional C_r -module under multiplication in FC_r .

$$
a \cdot e_t = \frac{1}{r} \sum_{i=0}^{r-1} \xi^{-it} a^{i+1}
$$

= $\xi^t \cdot \frac{1}{r} \sum_{i=0}^{r-1} \xi^{-(i+1)t} a^{i+1}$
= $\xi^t e_t$.

By §2.3, for $A = FG$, the projective modules are $P_t = FG_{t}$. If $J =$ Rad(FD) and $I = \text{Rad}(FG) = JFG$ then $\text{Rad}(P_t) = I \cap P_t$ and $S_t =$ $P_t/\text{Rad}(P_t)$ is the simple module corresponding to e_t .

We have assumed that the characteristic of F is p. Since $D \nightharpoonup C_p \times C_p$, $FD = F[x, y]/(x^p, y^p)$. Since C_r is an abelian group and $p \nmid r$ we set $FC_r =$ $\oplus Fe_t$. The structure of these blocks is treated in [\[Lan\]](#page-41-4), [\[Sw\]](#page-41-6), [\[K\]](#page-40-2), [\[RR\]](#page-41-5). Our notation follows that of [\[GS\]](#page-40-4) in the simpler case where there is no inertial subgroup. We know that the dimension of FD is p^2 and there is a basis $x^i y^j$. The module can be represented as a diamond of length p on each side.

2.6 Derived equivalence

We fix an abelian category $\mathcal C$ and we denote by $Ch(\mathcal C)$ the category of cochain complexes of objects of C . The differentials of the complex are morphisms ${d_n : C_n \to C_{n+1}}$ satisfying $d_{n+1} \circ d_n = 0$. For any given complex C^{\bullet} , the cocycles $Z^n(C^{\bullet})$, coboundaries $B^n(C^{\bullet})$, and cohomology modules $H^n(C^{\bullet}) =$ $Z^{n}(C^{\bullet})/B^{n}(C^{\bullet})$ are defined as in such standard texts as [\[W\]](#page-41-10).

Definition 2.4. A morphism $f_{\bullet}: C^{\bullet} \to D^{\bullet}$ of cochain complexes is called a quasi-isomorphism if the induced maps $H^n(C^{\bullet}) \to H^n(D^{\bullet})$ are all isomorphisms.

Definition 2.5. A cochain complex C is called *bounded* if almost all the $Cⁿ$ are zero. The complex C is *bounded below* if there is a bound a such that $C^n = 0$ for all $n < a$. The cochain complexes which are partially or fully bounded form full subcategories Ch^b , Ch^+ , Ch^- of $Ch(\mathcal{C})$.

Definition 2.6. The *derived category* of an abelian category \mathcal{C} is the category obtained from $Ch(\mathcal{C})$ by adding formal inverses to all the quasi-isomorphisms between chain complexes. It is called the *bounded derived category* and denoted $D^b(\mathcal{C})$ if we consider only bounded complexes. A *derived equivalence* between two abelian categories is an equivalence of categories between their derived categories.

Definition 2.7. We say that a chain map $f : C \rightarrow D$ is null homotopic if there are maps $s_n: C^n \to D^{n+1}$ such that $f = ds + sd$.

Definition 2.8. Two chain maps $f, g: C \to D$ are *chain homotopic* if their difference $f - g$ is null homotopic, in other words, if $f - g = sd + ds$ for some s. The maps $\{s_n\}$ are called a *homotopy* from f to g. We say that $f: C \to D$ is a homotopy equivalence if there is a map $g: D \to C$ such that $g \circ f$ is chain homotopic to the identity map id_C and $f \circ g$ is chain homotopic to the identity map id_D .

2.7 Tilting complexes

We now consider complexes T of projective left modules over an associative ring R. The notation $T[n]$ denotes the complex that is isomorphic to T as a module but in which the gradation has been shifted n places to the left and the differential is the shift of the differential multiplied by $(-1)^n$.

For any ring R, let $D^b(R)$ be the derived category of bounded complexes of R-modules.

Definition 2.9. Let R be a Noetherian ring. A bounded complex T of finitely generated projective R -modules is called a *tilting complex* if:

- (i) $\text{Hom}_{D^b(R)}(T,T[n])=0$ whenever $n \neq 0$.
- (ii) For any indecomposable projective P , define the stalk complex to be the complex $P: 0 \to P \to 0$. Then every such P is in the triangulated category generated by the direct summands of direct sums of copies of T.

A complex T satisfying only (i) is called a partial tilting complex.

Definition 2.10. Fix an abelian category \mathcal{C} and the category of cochain complexes $Ch(\mathcal{C})$. For two complexes X and Y denote by $Z(X, Y)$ the set of morphisms from X to Y which are homotopic to zero. The collection of all $Z(X,Y)$ forms a subgroup of $\text{Hom}_{C(\mathcal{C})}(X,Y)$. Denote by $K(\mathcal{C})$ the quotient category, i.e. $K(\mathcal{C})$ is the category having the same objects as $Ch(\mathcal{C})$ but with morphisms

$$
\operatorname{Hom}_{K(\mathcal{C})}(X,Y) = \operatorname{Hom}_{Ch(\mathcal{C})}(X,Y)/Z(X,Y),
$$

so that two homotopic maps are identified. The quotient category $K(\mathcal{C})$ is called the homotopy category, and a homotopy equivalence between complexes is an isomorphism in the homotopy category. For an abelian category $\mathcal{C}, K^-(\mathcal{C})$ is the homotopy category of right bounded complexes in \mathcal{C} , and similarly one can define $K^+(\mathcal{C})$.

The derived category is not an abelian category, but it is a triangulated category. The original theory of tilting concerned modules called tilting modules. Happel, in [\[H\]](#page-40-5) showed that if there was a tilting between two algebras Λ and Γ , it induced a functor which was an equivalence of their derived bounded categories. Rickard [\[R1\]](#page-41-0) then proved a converse when tilting

modules were replaced by tilting complexes, namely, that there is a tilting complex T over Λ with endomorphism ring $\text{End}_{D^b(\Lambda)}(T)^{op} \cong \Gamma$.

Let $\Lambda - Proj$ be the abelian category of projective Λ -modules. Let $Sum-T$ be the additive category of all the direct sums of copies of T (as a full subcategory of $K^{-}(\Lambda - Proj)$). From Lemma 3.3.1 of [\[KZ2\]](#page-40-6) we have that $K^{-}(\Gamma - Proj)$ is equivalent to $K^{-}(Sum - T)$ as triangulated categories. Thus, every tilting complex can be transformed into a projective complex, and to calculate the required endomorphism ring of the tilting complex, it will be sufficient to find chain maps between the components of the tilting complex which are not homotopic to zero.

2.8 Elementary equivalences

Of particular importance in this work are the elementary equivalences, which go back to the work of Rickard [\[R2\]](#page-41-11) and Okuyama [\[O\]](#page-41-3).

Definition 2.11. Let A be a finite-dimensional basic algebra, with projective modules $P_i, j \in I$.

Fix a subset $I_0 \subset I$. For any $i \in I$ we define a complex by

$$
T_i = \begin{cases} (0th) & (1st) \\ P_i & \longrightarrow & 0 \\ Q_i & \xrightarrow{\pi_i} & P_i \quad & i \in I - I_0 \end{cases}
$$

where $Q_i \stackrel{\pi_i}{\longrightarrow} P_i$ is a minimal projective presentation of the minimal kernel with quotient having only simples from $I - I_0$. Now we define $T := \bigoplus_{i \in I} T_i$. The algebra *elementary eqivalent* of A through T is is $A' \cong (\text{End}_{D^b(A)} T)^{op}$.

2.9 Homogeneous maps

To each $i \in I$, we can associate an idempotent \tilde{e}_i with $1_A = \sum^e$ $i=1$ \tilde{e}_i .

Definition 2.12. For $i, j \in \{0, ..., r-1\}$ let $P_i = Ae_iP_j = Ae_j$. The map $f: P_i \to P_j$ is homogeneous if $f(e_i) = \epsilon x^k y^\ell e_j$ for integers $0 \leq k, \ell \leq p-1$, $\epsilon \in F$.

Definition 2.13. Let

$$
f:\bigoplus_{i=1}^r P_i\to \bigoplus_{j=1}^r P_k.
$$

Then f will be called **homogeneous** if every component in $\bigoplus_{i,k=1}^r Hom(P_i, P_k)$ is a homogeneous map $f_{ik}: P_i \to P_k$ as in Definition [2.12.](#page-7-0)

Definition 2.14. Let l^{\bullet} be the chain map:

We say that l^{\bullet} is **homogeneous** if every vertical and horizontal map is homogeneous as in Definition [2.13](#page-7-1), and for every pair (i, u) and for all j and t we get

$$
\sum_{t=1}^{k} f_{tu} \circ d_{it} = \sum_{j=1}^{n} c_{ju} \circ b_{ij}
$$

Definition 2.15. A tilting complex $T = \bigoplus_{i=1}^{r} T_i$ over A is called homogeneous if each component of the differentials is ± 1 times a monomial in x, y times the appropriate idempotent.

3 MAPS BETWEEN COMPONENTS OF AN EL-EMENTARY TILTING COMPLEX

In order to describe the result of acting on the group algebra $A = F[C_p \times$ (C_p)) $\rtimes C_r$ by an elementary equivalence, we must calculate the endomorphism ring of the tilting complex $T = \bigoplus_{i=1}^r T_i$. To simplify all calculations, we will assume that the tilting complex T is homogeneous, a condition which can be achieved using Lemma [3.2](#page-9-0) below.

Let $I = \{0, 1, 2, \ldots, r - 1\}$ be arranged in a circle in the clockwise direction. Since I_0 is non-empty, removing the points of I_0 separates the circle into a finite number of intervals J_1, \ldots, J_m which we will call arcs. For a particular J_i , where $1 \leq i \leq m$, we let u_i be the first element of I_0 in the counter-clockwise direction and let v_i be the first element of I_0 in the clockwise direction. If I_0 contains a single point w, we will have $m = 1$ and $u_1 = v_1 = w$.

We denote the r-residue of $n \in \mathbb{Z}$ by \bar{n} . Any $t \in I - I_0$ belongs to an arc $J_i = \overline{[u_i+1,\ldots,\overline{t-1},t,\overline{t+1},\ldots,\overline{v_i-1}]}$ with the understanding that 0 follows $r-1$. The arc J_i is preceded by an interval I_0^i = $[\overline{v_{i-1}}, \overline{v_{i-1}+1}, \ldots, \overline{u_i-1}, \overline{u_i}] \subseteq I_0$ and followed by the interval I_0^{i+1} .

Definition 3.1. If $t, \tilde{t} \in I$ both lie in one of the intervals I_0^i , I_0^{i+1} or J_i , we will write $t \preceq \tilde{t}$ to indicate \tilde{t} either equals t or is subsequent to t in the interval.

Lemma 3.1. If $t, \tilde{t} \in I$ both lie in J_i and s lies in I_0^i , then

$$
\overline{\tilde{t}-t}+\overline{t-s}=\overline{\tilde{t}-s}.
$$

Similarly, if $s \in I_0^{i+1}$, then

$$
\overline{\tilde{t}-t}+\overline{s-\tilde{t}}=\overline{s-t}.
$$

Proof. Using the notation above, $I_0^i \cup J_i$ can be written as an ordered interval of length less than or equal to r. If $v = v_{i-1}$ is the first element of I_0^i , then we renumber I by replacing each $k \in I$ by $k' = \overline{k - v}$. If we identify $s', u', \tilde{t}',$ which are residues, with the corresponding integers between 0 and $r-1$, then we have $s' < t' \leq \tilde{t}'$, so that $t'-s' = \overline{t'-s'}, \tilde{t}'-t' = \tilde{t}'-t'$ and $\tilde{t}'-s' = \tilde{t}'-s'$. Since the renumbering is just a cyclic permutation and does not affect the residue of the differences between two residues, and since, as integers,

$$
(\tilde{t}' - t') + (t' - s') = \tilde{t}' - s',
$$

 \Box

we obtain the desired result.

In this article, we fix one arc J_i and we determine all the irreducible maps internal to the sets $I_0^i \cup J_i \cup I_0^{i+1}$. If $m = 1$, there is just a single arc J_1 , and a single complementary interval I_0 . Then the maps we will construct will generate the entire endomorphism ring. We will leave to a sequel the case where $m > 1$. For simplicity of notations, we will write $J = J_i$, $u=u_i, v=v_i.$

We first determine the maps between the projective modules in the block of normal defect group.

Lemma 3.2. We have a group block $F[(C_p \times C_p) \rtimes C_r]$ with indecomposible projective modules $P_0, P_1, \ldots P_{r-1}$. Suppose we are given a homogeneous map $P_i \rightarrow P_k$, for $0 \leq i, k \leq r-1$.

- 1. If $\overline{k-i} < p$ then there are homogeneous maps $f(e_i) = x^{k-i+hr}(xy)^q$, where h is an integer satisfying $0 \leq h \leq \frac{p-1-k-i}{r}$ $\frac{-\kappa - i}{r}$ and $0 \le q \le p - 1$ – $(\overline{k-i} + hr)$. If $\overline{i-k}$ < p then there are homogeneous maps $f(e_i) = y^{i-k+hr}(xy)^q$, where h is an integer satisfying $0 \leq h \leq \frac{p-1-i-k}{r}$ $\frac{-i-k}{r}$ and $0 \leq q \leq p-1$ – $(\overline{i-k}+hr)$.
- 2. The homogeneous maps give a basis for the homomorphisms from $P_i \rightarrow$ P_k thus we get

$$
f(e_i) = 1_{\overline{k-i} < p} \sum_{h=0}^{\lfloor \frac{p-1-\overline{k-i}}{r} \rfloor} \sum_{q=0}^{\lfloor p-1-(\overline{k-i}+hr) \rfloor} a_q^h \cdot x^{\overline{k-i}+hr} (xy)^q e_k + 1_{\overline{i-k} < p} \sum_{h=0}^{\lfloor \frac{p-1-\overline{i-k}}{r} \rfloor} \sum_{q=0}^{\lfloor p-1-(\overline{i-k}+hr) \rfloor} b_q^h \cdot y^{\overline{i-k}+hr} (xy)^q e_k.
$$

Note that if $r > p$ then $h = 0$.

Proof. 1. By standard arguments of representation theory, the image of e_i in P_k must be a linear combination of basis elements $e_i x^s y^t e_k$. Starting with e_k , if we multiply by x we get a basis element $e_{k-1}xe_k$. If we repeat this s times we get $e_{k-s}x^s e_k$. The first time that $e_{k-s} = e_i$ will be when $s = \overline{k - i}$. This can only occur if $\overline{k - i} < p$; otherwise we get 0. If we continue multiplying h times by x^r we keep on returning to e_i as long as $s = \overline{k - i} + hr < p$. Multiplication by y is dual, giving the elements $e_{k+t}y^t e_k$. For the first time that $e_{k+t} = e_i$ is when $t = \overline{i-k}$ and this can only occur if $\overline{i-k} < p$. If we continue multiplying h times by y^r we keep on returning to e_i as long as $t = \overline{i - k} + hr < p$.

If $e_i x^s e_k \neq 0$ then also $e_i x^s (xy)^q e_k \neq 0$ as long as $s + q \leq p - 1$. Dually if $e_i y^t e_k \neq 0$ then also $e_i y^t (xy)^q e_k \neq 0$ as long as $t + q \leq p - 1$. Note that if $i = k$ there are maps $f(e_i) = \sum_{q=0}^{p-1} c_q(xy)^q e_k$.

2. The general homomorphism is a linear combination of the homogeneous maps we found in 1. We added the two characteristic functions $1_{\overline{k-i}}$, $1_{\overline{i-k}}$ to indicate when part of the sum is zero.

 \Box

We recall from Definition [2.11](#page-7-2) that the irreducible components T_i are of two different types dependent on i. If $i \in I_0$, then T_i is a stalk complex,

whereas if $i \in I-I_0$, it is a complex with nonzero components in two degrees. We will be working in the homotopy category of complexes of projective modules and we will denote T_i by P'_i when regarded as an element of the homotopy category. What we will do is propose a catalog of irreducible maps between the various P'_i , and then demonstrate that a general homogenous map from P'_i to P'_j can be written as a composition of maps in the catalog.

3.1 Catalog of intra-arc maps

We now list the irreducible maps from P'_t to $P'_{t'}$ for t, t' in the interval $I_0^i \cup J_i \cup I_0^{i+1}$. The maps will generally be described by powers of x and y, but we will also need the following special maps in the catalog:

Definition 3.2. Let $t \in J$ in cyclic order, then the chain maps (ξ_1, x) , (ξ_2, y) are as follows

$$
\xi_1 = \begin{pmatrix} id & 0 \\ 0 & xy \end{pmatrix} \downarrow \begin{cases} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{cases} \longrightarrow P_t \longrightarrow P_t \longrightarrow P_t \xi_2 = \begin{pmatrix} xy & 0 \\ 0 & id \end{pmatrix} \downarrow \qquad P_u \oplus P_v \xrightarrow{p_{\overline{t-u}}} P_t \longrightarrow P_t \longrightarrow P_t \downarrow \qquad P_u \oplus P_v \xrightarrow{p_{\overline{t-u}}} P_u \oplus P_v \xrightarrow{p_{\overline{t-u}}} P_t \downarrow \qquad P_u \oplus P_v \xrightarrow{p_{\overline{t-u}}} P_{t+1} \longrightarrow P_{t+1}
$$

• A Adjacent maps.

For each arc J , as defined at the beginning of $\S3$ $\S3$, there is a sequence of maps, whose properties depend on the relative sizes of r and p and the location of $I_0 \subset I$. The maps we will give are all well-defined and not homotopic to zero.

Case 1: If $|J| \geq 2(p-1)$, maps as in Diagram 1. Case 2: If $p-1 \leq |J| < 2(p-1)$, maps as in Diagram 2. Case 3: If $1 \leq |J| < p-1$, maps as in Diagram 3.

If $j = p$, the horizontal maps $x, x^2 \dots x^{j-p}$ as well as $y, y^2 \dots y^{j-p}$ vanish and we get the left map of Diagram 3.

Diagram 3: $j = |J| + 1; 2 \le j \le p - 1$

We recall that we denoted T_i in Definition [2.11](#page-7-2) by P'_i . For $i, i + 1 \in J$ we denote by $\alpha_{i,i+1} : P'_i \to P'_{i+1}$ the appropriate adjacent map from the diagrams, and by $\alpha_{i+1,i} : P'_{i+1} \to P'_{i}$ the corresponding adjacent map in the opposite direction.

We denote by $\alpha_{t,\tilde{t}}$ the composition of maps $\alpha_{\tilde{t}-1,\tilde{t}} \circ \ldots \circ \alpha_{t+1,t+2} \circ$ $\alpha_{t,t+1}$ and by $\alpha_{\tilde{t},t}$ the composition $\alpha_{t+1,t} \circ \cdots \circ \alpha_{\tilde{t}-1,\tilde{t}-2} \circ \alpha_{\tilde{t},\tilde{t}-1}$. For $t, t+1 \in I_0$ we denote by $\gamma_{t,t+1} : P_t \to P_{t+1}$ and by $\gamma_{t+1,t} : P_{t+1} \to$ P_t multiplication by x or y respectively. When $|I_0^i|=1$ or $|I_0^{i+1}|=1$,

we have a map $\epsilon_t: P'_t \to P'_t$ inherited from the self-map xy from P_t to itself.

• B In addition, we have another set of self-maps which are maps from $Q_t \to P_t$ to itself which are not homotopic to zero. For $t \in J$,

$$
\begin{aligned}\n\textbf{(B.1)}\\
\begin{pmatrix}\n-d_0(xy)^{\overline{v-t}} & d_0x^{\overline{v-u}} \\
0 & 0\n\end{pmatrix} & \downarrow_{v} \frac{\left(\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\right)}{P_t} & P_t\\
P_u \oplus P_v \xrightarrow{f} P_t\\
P_t \oplus P_v \xrightarrow{f^{-u}} P_t\\
\begin{pmatrix}\n\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\n\end{pmatrix} & P_t\\
\begin{pmatrix}\n0 & 0 \\
c_0y^{\overline{v-u}} & -c_0(xy)^{\overline{t-u}}\n\end{pmatrix} & \downarrow_{v} \frac{\left(x^{\overline{t-u}}\right)}{P_t} & P_t\\
P_u \oplus P_v \xrightarrow{f^{-u}} P_t\n\end{aligned}
$$

We will show in the main theorem that when $d_0 + c_0 = 0$, the sum of B.1 and B.2 is homotopic to zero. We have left the dual pair in the catalog for convenience in decomposing other maps.

• C. Additional self-maps not necessarily irreducible

.

 $(C.1)$ Case of single projective in Q_t : By Lemma [3.2,](#page-9-0) a complex with Q_t a single projetive can exist anly if $j > p$. It is usually a composition of $\alpha_{t,t\pm1}$ and $\alpha_{t\pm1,t}$ but it is convenient to refer to, and is actually necessary to the catalog when $j = p + 1$.

$$
P_u \xrightarrow{\overline{x^{t-u}}} P_t \xrightarrow{\overline{y^{v-t}}} P_t
$$

\n
$$
\delta_{u,t}^q: (xy)^q
$$

\n
$$
P_u \xrightarrow{\overline{x^{t-u}}} P_t
$$

\n
$$
P_v \xrightarrow{\overline{y^{v-t}}} P_t
$$

\n
$$
\delta_{v,t}^w: (xy)^w
$$

\n
$$
P_v \xrightarrow{\overline{y^{v-t}}} P_t
$$

\n
$$
P_v \xrightarrow{\overline{y^{v-t}}} P_t
$$

The chain map $\delta_{u,t}^q$ is homotopic to zero when $q \geq \overline{u-t}$ and $\delta_{v,t}^w$ is homotopic to zero when $w \geq \frac{1}{v-t}$.

(C.2) Case of double projective in Q_t : $j \leq 2(p-1)$

$$
\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}
$$

$$
\begin{pmatrix} (xy)^q & 0 \\ (xy)^q & 0 \\ (xy)^q & 0 \end{pmatrix} \downarrow \begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix} \downarrow (xy)^q
$$

$$
P_u \oplus P_v \xrightarrow{p} P_t
$$

As in the case of C.1, this is usually a composition of $\alpha_{t,t\pm 1}$ and $\alpha_{t\pm 1,t}$ which we include as a convenience, but it is actually irreducible and needed in the catalog when $j = 2$. In order to consider the homotopy relations, it is useful to consider the following linear combination of B.1, B.2 and C.2:

$$
\begin{pmatrix}\n\begin{pmatrix}\nx^{\overline{t-u}} \\
y^{\overline{v-t}}\n\end{pmatrix} \\
\begin{pmatrix}\n(h_0 - d_0)(xy)^q & d_0(xy)^{q - (\overline{v-t})}x^{\overline{v-u}} \\
c_0(xy)^{q - (\overline{t-u})}y^{\overline{v-u}} & (h_0 - c_0)(xy)^q\n\end{pmatrix} \begin{pmatrix}\nx^{\overline{t-u}} \\
\begin{pmatrix}\nx^{\overline{t-u}} \\
y^{\overline{v-t}}\n\end{pmatrix} \begin{pmatrix}\nh_0(xy)^q \\
h_0(xy)^q\n\end{pmatrix} \\
P_u \oplus P_v \xrightarrow{P_v} P_t\n\end{pmatrix}
$$

If $q < \overline{v-t}$ then $d_0 = 0$ and if $q < \overline{t-u}$ then $c_0 = 0$. The map is homotopic to zero only if $b_0 + c_0 = h_0$ and then the homotopy is

$$
T = (c_0(xy)^{q - (\overline{t-u})} y^{\overline{t-u}}, d_0(xy)^{q - (\overline{v-t})} x^{\overline{v-t}}).
$$

(C.3) Case of double projective in Q_t : $j = \overline{v - u} \le r$ and $\overline{u - v} = r - j$.

$$
\begin{pmatrix}\n0 & e_0 y^{\overline{u-v}+hr}(xy)^{q_2} \\
0 & 0\n\end{pmatrix}\n\begin{pmatrix}\nx^{\overline{t-u}} \\
y^{\overline{v-t}}\n\end{pmatrix} P_t\n\begin{pmatrix}\n0 & e_0 y^{\overline{u-v}+hr}(xy)^{q_2} \\
0 & 0\n\end{pmatrix}\n\begin{pmatrix}\nx^{\overline{t-u}} \\
y^{\overline{v-\overline{t}}}\n\end{pmatrix}\n\begin{pmatrix}\n0 \\
0 \\
0\n\end{pmatrix}
$$
\n
$$
P_u \oplus P_v \xrightarrow{P_{\overline{t}}} P_{\overline{t}}
$$

For $e_0 \neq 0$, the map is non-zero if $0 \leq q_1 \leq p-1-\overline{u-v} - hr$ and well-defined if $q_1 \geq p - \overline{t} - v - hr$. For $f_0 \neq 0$, it is non-zero if $0 \le q_2 \le p - 1 - \overline{u - v} - hr$ and well-defined if $q_2 \ge p - \overline{u - t} - hr$. These occur for the maximal values of h

• **D**. When $r < p$, self-maps with double projectives.

$$
\begin{array}{ll}\n\textbf{(D.1)} & P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t \\
\begin{pmatrix} x^r & 0 \\ 0 & x^r \end{pmatrix} & \begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix} & \begin{pmatrix} x^r \\ x^r \end{pmatrix} \\
P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t \\
\textbf{(D.2)} & \begin{pmatrix} y^r & 0 \\ 0 & y^r \end{pmatrix} & \begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix} & \begin{pmatrix} y^r \\ y^r \end{pmatrix} \\
P_u \oplus P_v \xrightarrow{P_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t \\
P_u \oplus P_v \xrightarrow{P_t \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t\n\end{array}
$$

(D.3) This is a special case, occuring only when $r = 2j < p$ and h is maximal with $hr < p$. We combine two linearly independent maps. The map is well-defined for $e_0 \neq 0$ if $hr + \overline{v - t} \geq p$ and well-defined for $f_0 \neq 0$ if $hr + \overline{t - u} \geq p$.

$$
\begin{pmatrix}\n0 & e_0 y^{hr-j} (xy)^{\overline{t-u}} \\
\int_{0}^{0} f_0 x^{hr-j} (xy)^{\overline{v-t}} & 0\n\end{pmatrix}\n\downarrow\n\begin{pmatrix}\n\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\n\downarrow & P_t \\
\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\n\downarrow & \frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\n\end{pmatrix}\n\downarrow\n\begin{pmatrix}\ne_0 y^{hr} + f_0 x^{hr} \\
e_0 y^{hr} + f_0 x^{hr}\n\end{pmatrix}
$$

• E. Chain maps with one vertical term zero.

(E.1) Chain maps with degree 0 term zero. Case of single projective in Q_t : which can exist just if $j > p$.

(E.2) Chain maps with degree 1 term zero, single projective. We assume $j > p$. For every $w \in I_0^i, t \in J$, such that $\overline{t-u} \leq j-p$, $\overline{t-w} \leq p$, we have the following, which we denote by $\beta_w^u : P'_w \to P'_t$.

$$
P_w
$$

 $x^{\overline{u-w}}(xy)^{p-(\overline{t-w})}$
 $P_u \xrightarrow{\overline{x^{t-u}}} P_t$

For every $w \in I_0^{i+1}, t \in J$, such that $\overline{v-t} \leq j-p$, $\overline{w-t} \leq p$, we have the following map, which we denote by $\beta_w^v : P'_w \to P'_t$.

$$
P_w
$$
\n
$$
y^{\overline{w-v}}(xy)^{p-(w-t)} \Big|_{P_v} \longrightarrow P_t
$$
\n
$$
P_v \longrightarrow P_t
$$

(E.3) Chain maps with degree 1 term zero, double projective. For $hr < p$ and every $t \in J$, such that $hr + \overline{t - u} \geq p$, or in the dual, $hr + \overline{v - t} \geq p.$

$$
P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t
$$

\n
$$
\begin{pmatrix} x^{hr} & 0 \\ 0 & 0 \end{pmatrix} \downarrow \begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-\overline{t}}} \end{pmatrix} \rightarrow P_t
$$

\n
$$
P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t
$$

\n
$$
\begin{pmatrix} 0 & 0 \\ 0 & y^{hr} \end{pmatrix} \downarrow \begin{pmatrix} x^{\overline{\overline{t-u}}} \\ y^{\overline{v-\overline{t}}} \end{pmatrix} \rightarrow P_t
$$

\n
$$
P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{\overline{\overline{t-u}}} \\ y^{\overline{v-\overline{t}}} \end{pmatrix}} P_{\overline{t}}
$$

(E.4) Chain maps with degree 1 term zero, stalk complex to double projective.

$$
\begin{pmatrix} P_u \\ (x^{hr} & 0) \end{pmatrix}
$$

$$
P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t
$$

Well-defined if $hr + \overline{t - u} \geq p$.

$$
\begin{pmatrix} & & & P_v \\ (0 & y^{hr}) & & & \\ & & \downarrow & & \\ & P_u \oplus P_v & & \xrightarrow{p_v} & P_t \end{pmatrix}
$$

Well-defined if $hr + \overline{v - t} \geq p$.

(E.5) Maps of stalk complexes. Suppose the $I-I_0$ contains more than a single connected component, that $r < p$ and that $s = \overline{v_i - u_{i+1}} < p$. Maps from $P_{u_{i+1}} P_{v_{i-1}}$ given by x^s or y^s must be included in the catalog in order that we can obtain self maps of P_w given by x^r and y^r . These are similar to the maps x^j and y^j appearing in Diagram 3. If the interval in I_0 containing w consists of a single index, then the self-map of P_w given by xy is irreducible.

4 Maps between components of an elementary tilting complex

In this section, we will prove the proposition justifying the listed maps given in the catalog. In $\S 3.1$ $\S 3.1$ $\S 3.1$ the adjacent maps in A are irreducible, as are the self-maps in B. Of the maps in C, the maps with the smallest powers of xy are sometimes irreducible. When $u + 1 = v - 1 = t$, the map C.1 with $q = 1$ is irreducible.

Proposition 4.1. We are given a group algebra $F[C_p \times C_p) \times C_r$ and an elementary equivalence determined by a subset I_0 of the set of residues I. For each interval J_i in $I-I_0$, the set of maps in the catalog, whose nature depends on the relative sizes of r and p and the location of I_0 , are all well-defined and not homotopic to zero. All the maps internal to the set $I_0^i \cup J_i \cup I_0^{i+1}$ can be written a linear combinations of compositions of maps in the catalog.

Proof. 1. The maps in the catalog are well-defined and not homotopic to zero

We frequently use $x^p = y^p = 0$ as we mentioned before. All the maps between two stalk complexes are well-defined by the definition of the Brauer correspondent. Also, all the maps with zero at one end of the main diagonal are well-defined.

A. Diagram 1 We want to prove that all maps in Diagram 1 are welldefined and to check homotopy to zero. Maps between stalk complexes are already well-defined. Now we will check all maps between stalk complexes and 2-complexes as follows:

The long arrow.

$$
\begin{array}{ccc}\nP_u & & P_v \\
xy & & xy & xy \\
P_u & & x^{p-1} \n\end{array}\n\longrightarrow P_{u+(p-1)} \qquad\n\begin{array}{ccc}\nP_v & & & \\
xy & & y^{p-1} \n\end{array}\n\longrightarrow P_{v-(p-1)}
$$

both maps are well-defined because

$$
(xy) \cdot x^{p-1} = y \cdot x^p = 0
$$
 and $(xy) \cdot y^{p-1} = x \cdot y^p = 0$.

These maps are not homotopic to zero because they end with zero at the upper end of the off-diagonal.

The following maps occur only in Diagram 1:

$$
P_{u+p} \qquad P_{v-p}
$$
\n
$$
P_u \xrightarrow{p-1} P_{u+(p-1)} \qquad P_v \xrightarrow{y^{p-1}} P_{v-(p-1)}
$$

both maps are well-defined because one of the modules on the main diagonal is zero. These maps are not homotopic to zero because the diagonal maps are zero.

The following maps occur only in Diagram 1:

$$
P_u \xrightarrow{\quad x^{p-1}} P_{u+(p-1)} \qquad P_v \xrightarrow{\quad y^{p-1}} P_{v-(p-1)} \downarrow^x
$$

\n
$$
P_{u+p} \qquad P_{v-p}
$$

These maps are well-defined because $x^p = y^p = 0$ and they are not homotopic to zero because one of the off-diagonal modules is the zero module.

This pair of dual maps occurs also in Diagram 2.

$$
P_u \xrightarrow{~~x~~} P_{u+1} \qquad P_v \xrightarrow{~~y~~} P_{v-1} \n~\downarrow id \qquad \qquad P_u
$$
\n
$$
P_u \qquad \qquad P_v
$$

Both maps are automatically well-defined because one main diagonal module is zero and they are not homotopic to zero because of the identity map.

Now we will consider all maps between two 2-complexes in Diagram 1. These occur also in Diagram 2 when $j > p + 1$. We start with $\alpha_{t,\overline{t+1}}$ and $\alpha_{\overline{t+1},t}$ for $Q_t = P_u$ and dually, for $Q_t = P_v$, where $t, \overline{t+1} \in J$

$$
P_u \xrightarrow{\quad x^{\overline{t-u}}} P_t \qquad P_v \xrightarrow{\quad y^{\overline{v-t-1}}} P_{t+1}
$$

\n
$$
P_u \xrightarrow{\quad x^{\overline{t-u+1}}} P_{t+1} \qquad P_v \xrightarrow{\quad y^{\overline{v-t}}} P_t
$$

\n
$$
P_u \xrightarrow{\quad x^{\overline{t-u+1}}} P_{t+1} \qquad P_v \xrightarrow{\quad y^{\overline{v-t}}} P_t
$$

They are well-defined because $x^{t-u+1} \circ id = x \circ x^{\overline{t-u}}$ and dually $y^{\overline{v-t}} \circ$ $id = y \circ y^{v-t-1}$. Maps that have a square containing a vertical identity can never be homotopic to zero. In the second case:

$$
P_u \xrightarrow{\quad x^{t-u+1}} P_{t+1} \qquad P_v \xrightarrow{\quad y^{v-t}} P_t
$$

\n
$$
\downarrow xy \qquad \qquad \downarrow y \qquad \qquad \downarrow xy \qquad \qquad \downarrow xy \qquad \qquad \downarrow x
$$

\n
$$
P_u \xrightarrow{\quad x^{t-u}} P_t \qquad \qquad P_v \xrightarrow{\quad y^{v-t-1}} P_{t+1}
$$

the chain maps are well-defined because $x^{\overline{t-u}} \circ xy = y \circ x^{\overline{t-u+1}}$ and dually $(xy) \cdot y^{v-t-1} = y^{\overline{v-t}} \cdot x$. We want to prove that these maps are not homotopic to zero. If we can find $T: P_{t+1} \to P_u$, then since the horizontal map $x^{\overline{t-u}}$ has a non-zero power of x, the composition with T does not give y, and the case on the right is dual.

A. Diagram 2. Now we will look at Diagram 2 and consider maps that do not occur in Diagram 1. We start with the long arrows in

Diagram 2.

$$
\begin{pmatrix} p_u & & & & P_v \\ (xy & 0) & & & \\ & & (y^{j-p+1}) & & \\ & P_u \oplus P_v \xrightarrow{\hspace{0.5cm}} & & & \end{pmatrix} \xrightarrow{\hspace{0.5cm}} \begin{pmatrix} p_v & & & & \\ (0 & xy) & & & \\ & (y & xy) & & \\ & & P_u \oplus P_v \xrightarrow{\hspace{0.5cm}} & & & \end{pmatrix} \xrightarrow{\hspace{0.5cm}} \begin{pmatrix} x^{j-p+1} \\ y^{p-1} \end{pmatrix}
$$

They are well-defined because $x^{p-1} \circ xy = 0$ and dually $y^{p-1} \circ xy = 0$. Maps that have a zero module at one end of the off-diagonal can never be homotopic to zero.

$$
\begin{pmatrix} P_u \oplus P_v & \xrightarrow{\binom{x^{p-1}}{y^{j-p+1}}} P_{v-j+p-1} \\ \begin{pmatrix} 0 \\ xy \end{pmatrix}_P & \xrightarrow{T_1} \qquad \qquad P_v \oplus P_v \xrightarrow{\binom{x^{j-p+1}}{y^{p-1}}} P_{u+j-p+1} \\ \begin{pmatrix} x \\ xy \end{pmatrix}_P & \xrightarrow{T_2} \qquad \qquad P_{u+j-p+1} \\ \begin{pmatrix} xy \\ y \end{pmatrix}_P & \xrightarrow{T_2} \qquad \qquad P_{u+j-p} \\ \begin{pmatrix} x \\ y \end{pmatrix}_P & \xrightarrow{T_2} \qquad \qquad P_{u+j-p} \end{pmatrix}
$$

In this case $j > p$. The chain maps are well-defined because (xy) . $y^{j-p} = xy^{j-p+1}$ and dually $(xy) \cdot x^{j-p} = yx^{j-p+1}$. If $j > p$ the maps are not homotopic to zero because if $T_1 = x^{j-p+1}$ then $x^{j-p+1}y^{j-p} \neq x$ and dually, if $T_2 = y^{j-p+1}$, then $y^{j-p+1}x^{j-p} \neq y$.

$$
\begin{array}{ccc}\nP_u & \xrightarrow{\quad x^{j-p}} P_{u+j-p} & P_v \xrightarrow{\quad y^{j-p}} P_{v-j+p} \\
(id & 0) & \xrightarrow{\quad x^{j-p+1}} P_u \xrightarrow{\quad x^{j-p+1}} P_u & (0 & id) & \xrightarrow{\quad x^{p-1}} P_u \xrightarrow{\quad y^{p-1}} P_u \xrightarrow{\quad y^{p-1
$$

Again $j > p$. The chain maps are well-defined because $x^{j-p} \cdot x = x^{j-p+1}$ and dually $y^{j-p} \cdot y = y^{j-p+1}$. Maps that have a square containing a vertical identity map can never be homotopic to zero. Let $j = \overline{v - u}$ and $t \in J$.

P^u ⊕ P^v ξ1= id 0 0 xy x t−u y v−t /P^t x v♠♠ ♠♠♠♠♠♠♠♠♠♠♠♠♠♠♠♠ P^u ⊕ P^v x t−u+1 y v−t−1 /Pt+1 P^u ⊕ P^v x t−u y v−t /P^t P^u ⊕ P^v ♠6 ♠♠ ♠♠ ♠♠ ♠♠ ♠♠ ♠♠ ♠♠ ♠♠ ♠ ξ2= xy 0 0 id OO x t−u+1 y v−t−1 /Pt+1 y OO

They are well-defined because $\begin{pmatrix} id & 0 \\ 0 & xy \end{pmatrix}$. $\int x^{\overline{t-u+1}}$ $y^{\overline{v-t-1}}$ \setminus = $\int x^{\overline{t-u}}$ $y^{\overline{v-t}}$ \setminus $\cdot x$

as well as the dual case. They are not homotopic to zero because if in the map on the left, the first component of the homotopy is nonzero, the composition with the horizontal map contains x^2 and if the second component is non-zero, the composition has a power of y . The other map is dual.

A. Diagram 3. These maps occur in Diagram 3 when $j \leq p-1$ or in Diagram 2 when $j = p$.

$$
\begin{pmatrix} P_u & & & & P_v \\ (xy & -x^j) \Big| & & \left(x^{j-1} \right) & \left(-y^j & xy\right) \Big| & & \left(x \right) \\ P_u \oplus P_v \xrightarrow{\qquad \qquad} P_v \longrightarrow P_{v-1} & & P_u \oplus P_v \xrightarrow{\qquad \qquad} P_{u+1} \end{pmatrix}
$$

The chain maps are well-defined because $(xy) \cdot x^{j-1} - x^j \cdot y = 0$ and dually $-y^j \cdot x + xy \cdot y^{j-1} = 0$. Maps which have a zero module at one end of the off-diagonal can never be homotopic to zero.

Self-maps B. We now check well-defined for the maps in B.

For $t \in J$

$$
\begin{array}{ccc}\n\left(-b_0(xy)^{\overline{v-t}} & b_0x^{\overline{v-u}}\right) & \left(\frac{x^{\overline{t-u}}}{y^{v-t}}\right) & P_t \\
\left(-b_0(xy)^{\overline{v-t}} & b_0x^{\overline{v-u}}\right) & \left(\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\right) & \left(\frac{0}{y^{v-t}}\right) & P_t \\
\left(\frac{0}{c_0y^{\overline{v-u}}} & 0 & P_t\right) & \left(\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\right) & P_t \\
\left(c_0y^{\overline{v-u}} & -c_0(xy)^{\overline{t-u}}\right) & \left(c_0x^{\overline{v-u}}\right) & P_t\n\end{array}
$$

For the first map, let $a \in P_u$ and $b \in P_v$. We check the commutativity of the square by diagram chasing.

$$
(a,b)\cdot \begin{pmatrix} -b_0(xy)^{\overline{v-t}} & b_0x^{\overline{v-u}} \\ 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix} = -ab_0x^{\overline{t-u}}(xy)^{\overline{v-t}} + ab_0x^{\overline{v-u}}y^{\overline{v-t}} = 0
$$

For the second map

$$
(a,b) \cdot \begin{pmatrix} 0 & 0 \ c_0 y^{\overline{v-u}} & -c_0(xy)^{\overline{t-u}} \end{pmatrix} \cdot \begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix} = bc_0 x^{\overline{t-u}} y^{\overline{v-u}} - bc_0(xy)^{\overline{t-u}} y^{\overline{v-t}} = 0
$$

Now we will prove that B.1 is not homotopic to zero, and we will prove while dealing with the maps of C that B.2 is homotopic to B.1. For $t \in J$,

We need to find $T: P_t \to P_u \oplus P_v$ and to check if

$$
\begin{aligned}\n\left(x^{\overline{t-u}}\right) \circ T &= 0\\
T \circ \left(x^{\overline{t-u}}\right) &= \left(-d_0(xy)^{\overline{v-t}} & d_0x^{\overline{v-u}}\right),\\
T \circ \left(y^{\overline{v-t}}\right) &= \left(-d_0(xy)^{\overline{v-t}} & d_0x^{\overline{v-u}}\right),\n\end{aligned}
$$

Let us take as our candidate for homotopy the map $T: P_t \to P_u \oplus P_v$ given by $\left(e_0y^{\overline{t-u}}(xy)^{q_1}$ $f_0x^{\overline{v-t}}(xy)^{q_2}\right)$. By Lemma [3.2,](#page-9-0) if $r > p$, this is the only possibilty. From our calculations, it will be clear that even if $r < p$, multiplying the components of T by various power of x^r or y^r could not give the desired composition.

So, if we look at the bottom triangle $\left(x\frac{\overline{t} - u}{\overline{u} - t}\right)$ $y^{\overline{v-t}}$ \setminus $\circ T \neq 0$ unless $e_0+f_0 = 0$ and $q_1 + \overline{t - u} = q_2 + \overline{v - t}$. If $\overline{v - t} \geq \overline{t - u}$, we set $q_2 = q$, $q_1 =$

 $q + v + u - 2t$. If $\overline{t - u} \ge \overline{v - t}$, we set $q_1 = q$, $q_2 = q + 2t - u - v$. Now we check the second condition for the case $\overline{t - u} \geq \overline{v - t}$.

$$
-d_0(xy)^{\overline{v-t}} = x^{\overline{t-u}}e_0y^{\overline{v-t}}(xy)^q
$$

which can only hold if $e_0 = -d_0, \overline{t - u} = \overline{v - t}$ and $q = 0$. However, if all those conditions hold, then we must have $0 = y^{\overline{v-t}} e_0 y^{\overline{t-u}} (xy)^0$, which cannot be true because $\overline{t-u} + \overline{v-t} = j < p$. The case $\overline{v-t} \geq$ $\overline{t-u}$ is similar, using the second column of the matrix.

Self-maps C.

Most of the additional self-maps $C.1$ are compositions of adjacent maps and therefore well-defined. If $j = 2$, the maps in C.1 are irreducible but can be checked by inspection. In the case of a single projective, we have compositions of adjacent maps up and down or down and up, in the case of double projectives we have compositions of the maps in Definition [3.2.](#page-11-1)

C.1: Case of single projective: $j > p$. For $q < \overline{t - u}$, $w < \overline{v - t}$, $t \in J$.

When $q \geq \overline{t-u}$, we get a homotopy to the map on the left given by $T = (xy)^{q-\overline{t-u}}y^{\overline{t-u}}$. When $w \geq \overline{v-t}$, we get a homotopy for the second map given by $T = (xy)^{w-\overline{v-t}}y^{\overline{v-t}}$

C.2: Case of double projective: $j \leq 2(p-1)$ We only have to check our claim that the map below is homotopic to zero if $h_0 = d_0 + c_0$ and not all the parameters are zero. We assume $c_0 \neq 0$, in which case $q \geq \overline{t - u}$ or else $d_0 \neq 0$, in which case $q \geq \overline{v - t}$.

$$
\begin{pmatrix}\n\begin{pmatrix}\nx^{t-u} \\
y^{v-t}\n\end{pmatrix} \\
\begin{pmatrix}\n(h_0 - d_0)(xy)^q & d_0(xy)^{q - (v-t)}x^{v-u} \\
c_0(xy)^{q - (t-u)}y^{v-u} & (h_0 - c_0)(xy)^q\n\end{pmatrix}\n\begin{pmatrix}\nx^{t-u} \\
y^{v-t}\n\end{pmatrix} \\
P_u \oplus P_v \xrightarrow{p} P_t\n\end{pmatrix}
$$

We define the homotopy map $T: P_t \to P_u \otimes P_v$ by \overline{c}

$$
(c_0y^{(t-u)}(xy)^{q-\overline{t-u}}, d_0x^{\overline{v-t}}(xy)^{q-\overline{v-t}})
$$

Then the bottom triangle yields $(c_0 + d_0)(xy)^q$ and the upper triangle yields the desired matrix, with c_0 written for $h_0 - d_0$ and d_0 written for $h_0 - c_0$.

Self-maps D: When $r < p$, maps with double projective.

$$
\begin{array}{c}\n\textbf{(D.1)}\\
\begin{pmatrix} x^r & 0 \\ 0 & x^r \end{pmatrix} \downarrow R_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t\\
P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t\\
P_t \end{array}
$$

well-defined because

$$
(a,b) \cdot \left(\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\right) \cdot x^r = ax^{\overline{t-u}} \cdot x^r + by^{\overline{v-t}} \cdot x^r
$$

On the other hand

$$
(a,b)\cdot \begin{pmatrix} x^r & 0 \\ 0 & x^r \end{pmatrix} \cdot \begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix} = ax^r \cdot x^{\overline{t-u}} + bx^r \cdot y^{\overline{v-t}}.
$$

The case of $(D.2)$ is dual.

(D.3) This is a special case, occuring only when $r = 2j < p$ and h is maximal with $hr < p$. The map is well-defined for $f_0 \neq 0$ if $hr + \overline{t - u} \ge p$ and is well-defined for $e_0 \neq 0$ if $hr + \overline{v - t} \ge p$.

$$
\begin{pmatrix}\n0 & e_0 y^{hr-j} (xy)^{\overline{v-t}} \\
\int_{0}^{1} e_0 y^{hr-j} (xy)^{\overline{v-t}} & 0\n\end{pmatrix}\n\begin{pmatrix}\n\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\n\end{pmatrix}\n\begin{pmatrix}\n\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\n\end{pmatrix}\n\begin{pmatrix}\ne_0 y^{hr} \\
\frac{y^{hr}}{y^{\overline{v-t}}}\n\end{pmatrix}\n\begin{pmatrix}\ne_0 y^{hr} \\
\frac{y^{hr}}{y^{\overline{v-t}}}\n\end{pmatrix}\n\begin{pmatrix}\ne_0 y^{hr} \\
\frac{y^{hr}}{y^{\overline{v-t}}}\n\end{pmatrix}
$$

A homotopy for the bottom triangle would be $(f_0x^{hr-\overline{t-u}}, e_0y^{hr-\overline{v-t}})$, but this would not give the proper matrix when composing the upper triangle.

E. Chain maps with vertical term zero.

 $(E.1)$ Case of single projective in Q_t : By Lemma [3.2,](#page-9-0) it can exist just if $j > p$.

The maps are well-defined because all the powers of x , and, in the dual case, y, vanish and we get zero as desired.

(E.2) Chain maps with degree 1 term zero, stalk comples to single projective.

In the first of these, the horizontal map exists if $j > p$ and $0 \leq \overline{t - u} \leq$ $j - p$. The vertical map exists and is nonzero for every $w \in I_0^i, t \in J$, such that $\overline{t-w} \leq p$. The map is well-defined because the composition of horizontal and vertical maps contains x to the power p . Maps that have a zero module at one end of the off-diagonal can never be homotopic to zero. The second map is dual, where w lies in I_0^{i+1} .

(E.3) Chain maps with degree 1 term zero, double projective. Welldefined for $hr < p$ and every $t \in J$, such that $hr + \tilde{t} - u \geq p$, or in the dual, $hr + v - \tilde{t} \geq p$.

$$
P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t
$$

$$
\begin{pmatrix} x^{hr} & 0 \\ 0 & 0 \end{pmatrix} \downarrow \begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-\overline{t}}} \end{pmatrix} \searrow P_t
$$

$$
P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-\overline{t}}} \end{pmatrix}} P_{\overline{t}}
$$

$$
P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{pmatrix}} P_t
$$

\n
$$
\begin{pmatrix} 0 & 0 \\ 0 & y^{hr} \end{pmatrix} \downarrow \qquad \begin{pmatrix} x^{\overline{t-u}} \\ y^{\overline{v-\overline{t}}} \end{pmatrix} \downarrow \qquad \begin{pmatrix} 0 \\ \downarrow \\ \downarrow \\ P_t \end{pmatrix}
$$

(E.4) Chain maps with degree 1 term zero, stalk complex to double projective.

$$
\begin{pmatrix} P_u & & & P_v \\ (x^{hr} & 0) \Big| & & (x^{\overline{t-u}}) \\ P_u \oplus P_v \xrightarrow{(y^{\overline{v-t}})} P_t & & P_u \oplus P_v \xrightarrow{(y^{\overline{v-t}})} P_t \end{pmatrix}
$$

The first chain map is well-defined if $hr + \overline{t - u} \geq p$. Dually, the second chain map is well-defined if $hr + \overline{v - t} \geq p$. Maps that have a zero module at one end of the off-diagonal can never be homotopic to zero.

(E.5) Maps between stalk complexes allowed by Lemma [3.2](#page-9-0) must be well-defined and of course cannot be homotopic to zero, since there is no square.

2. All maps internal to $I_0^i \cup J_i \cup I_0^{i+1}$ are linear combinations of compositions of the maps in the catalog

In this section of the proof, we will consider pairs P'_t and P'_t of indecomposable components of the tilting complex of the elementary equivalence determined by the subinterval I_0 . The irreducible components are of two kinds, stalk complexes when $t \in I_0$ and complexes of length two when $t \in I - I_0$. We will first determine the general well-defined chain map, not necessarily homogeneous, between this pair of complexes, having constant recourse to Lemma [3.2,](#page-9-0) which gives the possible maps between the indecomposable projectives P_i . We then attempt to write the general map as a linear combination of non-zero, well-defined homogeneous chain maps. Finally, we will decompose each homogeneous chain map as a composition of homogeneous chain maps from the catalog. In Case 2.1, we will deal with all maps between stalk complexes, in Cases 2.2, 2.3, and 2.4 we will deal with maps between two-term complexes, and in Case 2.5, with maps between stalk complexes and two-term complexes. We establish a general notation for maps between two-term complexes P'_t and $P'_\tilde{t}$ where $t, \tilde{t} \in J, t \leq \tilde{t}$.

where Lemma [3.2](#page-9-0) gives the maps ℓ_1 and k_1 .

Case 2.1: First, we check the maps between stalk complexes, where $t,\tilde{t}\in I_0$. In general, we assume that if t,\tilde{t} in from the same interval then $t \leq \tilde{t}$, as in Def. [3.1.](#page-9-1) We get the maps ℓ_0 and k_0 rrom Lemma [3.2.](#page-9-0)

Suppose $t, \tilde{t} \in I_0^i$ or $t, \tilde{t} \in I_0^{i+1}$. Each individual map $x^{\tilde{t}-t}(xy)^q$ or $y^{\tilde{t}-t}(xy)^q$ is a composition of maps $\gamma_{\tilde{t},\tilde{t}+1}$ in the catalog, where $\bar{t}=t,t+1$ $1, t + 2, \ldots, \tilde{t} - 1$, followed by q pairs of maps xy, also compositions of the γ or the maps ϵ_t . In the special cases listed in E.5, one of the maps from E.5 might be needed. The general map is then a linear combination of these homogeneous maps with coefficients a_q or b_q .

Suppose that $t \in I_0^i$, $\tilde{t} \in I_0^{i+1}$, and $\tilde{t} - t < p$, which implies $j < p$, the general map given above is a linear combination with coefficients a_q or b_q of the homogeneous maps in the sums, each of which is a composition of the $\gamma_{k,k+1}$, sometimes including ϵ_t , with the jump map from Diagram 3, given by x^j or y^j , followed by q self-maps xy , each of which is a composition of two maps $\gamma_{t,t+1}$ and $\gamma_{t+1,t}$ or a copy of ϵ_t maps.

Case 2.2: Single projective, where $Q_t, Q_{\tilde{t}} = P_u$. We see from the diagrams in the catalog that such $Q_t, Q_{\tilde{t}}$ occur only when $j > p$, where we have $u+1 \leq t \leq \tilde{t} \leq u+(j-p)$. By Lemma [3.2,](#page-9-0) in a special case where $i = k$ and under the assumption $j > p$, all the non-zero maps l_0, k_0 between $P_u \to P_u$ have the form $\sum_{q=0}^{p-1} c_q(xy)^q$ and $\sum_{w=0}^{p-1} d_w(xy)^w$ respectively.

Now we try to simplify these sums by subtracting maps that are homotopic to zero. Considering the left-hand square, we are in the case given in Lemma [3.1](#page-9-2) with $s = u$, and conclude that $\overline{t - u} + \tilde{t} - t = \tilde{t} - u$. If we have a homogeneous map in degree zero of $(xy)^q$, then the chain map is well-defined if the map in degree 1 is $x^{\tilde{t}-t}(xy)^q$.

When $q \geq \overline{t-u}$ then we then have a homotopy to zero $T: P_t \to P_u$ given by $(y^{\overline{t-u}}(xy)^{q-\overline{t-u}})$. The composition of the upper horizontal map and T in the upper triangle gives $(xy)^q$ as needed, while the composition of T with the lower horizontal maps, using Lemma [3.1](#page-9-2) gives $(xy)^{q-\overline{t-u}}y^{\overline{t-u}}x^{\overline{t-u}+\tilde{t}-t}$. Pulling out the maximal power of xy and moving it to the end using commutativity, we have the desired vertical map $x^{\tilde{t}-t}(xy)^q$. Subtracting off these homotopic-to-zero chain maps, we can assume that $\ell_0 = \sum_{q=0}^{\min(p-1,\overline{t-u})} c_q(xy)^q$.

For each q in this range, we subtract off c_q times the composition of maps from the catalog $\delta^q \circ \alpha_{t,\tilde{t}}$, and we get the situation that $\ell_0 = 0$. For such a map to be well-defined, we must have $0 = x^{\overline{t-u}} \circ \ell_1$. These maps are compositions of the maps in E.1, composed with δ_t^q t^q . Similarly for the map k_1 or δ_t^w . The case of $Q_t, Q_{\tilde{t}} = P_v$ is dual.

Case 2.3: Double projectives, where $Q_t = Q_{\tilde{t}} = P_u \oplus P_v$. Of all the five cases, this is the most difficult, since we may have $r < p$, in which case we get the full sums in Lemma [3.2.](#page-9-0)

Looking at the powers of x that occur in the horizontal maps in Diagram 2, where $p \le j \le 2(p-1)$ and recalling that $j = \overline{v - u}$, then $\overline{(j-p)+1} \le$ $\overline{t-u} \leq \overline{\tilde{t}-u} \leq \overline{p-1}$. According to Diagram 3, where $2 \leq j \leq p-1$ then $1 \leq \overline{t-u} \leq \overline{\tilde{t}-u} \leq \overline{j-1}.$

We recall that $u \in I_0^i$, $v \in I_0^{i+1}$. Assume that $t \preceq \tilde{t} \in J$. Let $a \in P_u$ and $b \in P_v$. From our calculation of maps among projectives in Lemma [3.2,](#page-9-0) we have that all maps $P'_t \to P'_{\tilde{t}}$ are of the following form

$$
\left\{ \begin{aligned}\n &\quad P_u \oplus P_v \xrightarrow{\left(\begin{smallmatrix} x^{\overline{t-u}} \\ y^{\overline{v-t}} \end{smallmatrix} \right)} P_t \\
 &\quad \left(\begin{smallmatrix} a_{11}(x,y) & a_{12}(x,y) \\ a_{21}(x,y) & a_{22}(x,y) \end{smallmatrix} \right) \right\} \\
 &\quad P_u \oplus P_v \xrightarrow{\left(\begin{smallmatrix} x^{\overline{t-u}} \\ y^{\overline{v-\overline{t}}} \end{smallmatrix} \right)} P_{\overline{t}} \\
 &\quad P_{\overline{t}}\n\end{aligned}
$$

where, by Lemma [3.2,](#page-9-0)

$$
a_{ii}(x,y) = \sum_{h=0}^{\lfloor \frac{p-1}{r} \rfloor} \sum_{q_{ii}=0}^{(p-1-hr)} a_{q_{ii}}^{iih} \cdot x^{hr}(xy)^{q_{ii}}
$$

$$
+\sum_{h=0}^{\lfloor \frac{p-1}{r} \rfloor} \sum_{q'_{ii}=0}^{(p-1-hr)} b_{q'_{ii}}^{iih} \cdot y^{hr}(xy)^{q'_{ii}}
$$

$$
a_{12}(x,y) = 1_{\overline{v-u}} \sum_{h=0}^{\lfloor \frac{p-1-\overline{v-u}}{r} \rfloor} \sum_{q_{12}=0}^{p-1-(\overline{t-t}+hr)} a_{q_{12}}^{12h} \cdot x^{\overline{v-u}} + hr(xy)^{q_{12}}
$$

$$
+1_{\overline{u-v}} \sum_{h=0}^{\lfloor \frac{p-1-\overline{t-\overline{t}}}{r} \rfloor} (p-1-(\overline{v-u}+hr)) +1_{\overline{u-v}} \sum_{h=0}^{\lfloor \frac{p-1-\overline{t-\overline{t}}}{r} \rfloor} \sum_{q'_{12}=0}^{(p-1-(\overline{v-u}+hr))} b_q^{12h} \cdot y^{\overline{u-v}} + hr(xy)^{q_{21}}
$$

$$
a_{21}(x,y) = 1_{\overline{u-v}} \sum_{h=0}^{\lfloor \frac{p-1-\overline{u-v}}{r} \rfloor} \sum_{q_{21}=0}^{(p-1-(\overline{u-v}+hr))} a_q^{21h} \cdot x^{\overline{u-v}+hr}(xy)^{q'_{21}}
$$

$$
+ 1_{\overline{v-u}} \sum_{h=0}^{\lfloor \frac{p-1-\overline{u-v}}{r} \rfloor} \sum_{q_{21}=0}^{(p-1-(\overline{u-v}+hr))} b_q^{21h} \cdot y^{+hr}(xy)^{q'_{21}}
$$

and, finally,

$$
\ell_1(x, y) = 1 \frac{\left[\frac{p-1-\overline{t}-t}{r}\right](p-1-(\overline{t}-t+hr))}{\overline{t}-t} \sum_{q=0}^{q-1} c_q^h \cdot x^{\overline{t}-t+hr}(xy)^q
$$

$$
+ 1 \frac{\left[\frac{p-1-\overline{t}-\overline{t}}{r}\right](p-1-(\overline{t}-\overline{t}+hr))}{\sum_{h=0}^{q-1-\overline{t}-\overline{t}} d_{q'}^h \cdot y^{\overline{t}-\overline{t}+hr}(xy)^{q'}
$$

We now do diagram chasing on our diagram. When calculating the diagonal map using the upper triangle, we get

$$
(a,b) \cdot \left(\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\right) \cdot \ell_1(x,y) = (ax^{\overline{t-u}} + by^{\overline{v-t}}) \cdot \ell_1(x,y)
$$

When we use the lower triangle, we have

$$
(a,b)\cdot \begin{pmatrix} a_{11}(x,y) & a_{12}(x,y)\\ a_{21}(x,y) & a_{22}(x,y) \end{pmatrix} \cdot \begin{pmatrix} x^{\overline{\overline{t}-u}} \\ y^{\overline{v-\overline{t}}} \end{pmatrix}
$$

$$
= (aa_{11}(x,y) + ba_{21}(x,y), aa_{12}(x,y) + ba_{22}(x,y)) \cdot \left(\frac{x^{\overline{\overline{t}-u}}}{y^{\overline{v-\overline{t}}}}\right)
$$

The coefficients of a and the coefficients of b must match in these expressions, giving us two equations

$$
(a_{11}(x,y)x^{\overline{\overline{t}-u}} + a_{12}(x,y)y^{\overline{v-\overline{t}}}) = x^{\overline{t-u}} \cdot \ell_1(x,y)
$$
\n(4.1)

$$
(a_{21}(x,y)x^{\tilde{t}-u} + a_{22}(x,y)y^{v-\tilde{t}}) = y^{\overline{v-t}} \cdot \ell_1(x,y)
$$
\n(4.2)

We want to show, under the assumption that this general map is welldefined, that the map is a linear combination of compositions of maps from the catalog. We first try to diagonalize the matrix, using maps from the catalog whose right vertical map is zero. The first sum in a_{12} is eliminated by subtracting copies of B.1 composed with D.1 and C.2. When these have been removed, we are left with

$$
1_{\overline{u-v}
$$

The case of the double projective occurs only when $j = \overline{v - u} \leq 2(p -$ 1) and for this second sum to be non-zero we need $r - j = \overline{u - v} < p$. Furthermore, if $r > p$, then the upper limit of the first summation is zero, so we must have $h = 0$.

Whenever $q'_{12} \geq \overline{v - u}$ we can remove compositions with B.1, and we will assume that this has been done. We now use the map in C.2 with parameters b_0, c_0, h_0 , where we assume $c_0 = 0$ and $h_0 = b_0$. Since this means that $h_0 = c_0 + b_0$, the map is homotopic to zero, and thus its compositions with $\alpha_{t,\tilde{t}}$ and with powers of D.2 are also homotopic to zero.

$$
\begin{pmatrix}\n\begin{pmatrix}\n\frac{x^{t-u}}{y^{v-t}} \\
y^{v-t}\n\end{pmatrix} \\
\begin{pmatrix}\n0 & h_0(xy)^{q-(v-t)}x^{v-u}y^{hr} \\
0 & h_0(xy)^qy^{hr}\n\end{pmatrix} \downarrow \begin{pmatrix}\nx^{t-u} \\
y^{v-t} \\
P_u \oplus P_v \longrightarrow P_t\n\end{pmatrix} h_0(xy)^q y^{hr}
$$

Using Lemma [3.1,](#page-9-2) $\overline{u-v} + \overline{v-u} = r$ and $\overline{v-u} = \overline{v-t} + \overline{t-u}$, giving us that the upper right-hand entry in the matrix becomes $h_0(xy)^{q+t-u}y^{\overline{v-u}}+(h-1)r$. Composing with $\alpha_{t,\tilde{t}}$ will only multiply by a power of xy. Thus for proper choice of q'_{12} , we can use this map to remove the remaining terms of $a_{12}(x, y)$. A dual procedure for eliminating $a_{21}(x, y)$ will successfully diagonalize the matrix.

Once the matrix is diagonal, equations 4.1 and 4.2 assume a simple form that allows us to solve immediately for a_{11} and a_{22} whenever the map is welldefined. We separate ℓ_1 into the two possible sums and treat each separately. We start with the first sum, assuming $\tilde{t} - t < p$ and solve

$$
\begin{array}{ccc}\nP_u \oplus P_v & \xrightarrow{\left(\frac{x^{\overline{t-u}}}{y^{\overline{v-t}}}\right)} P_t \\
\begin{pmatrix} a_{11}(x,y) & 0 \\ 0 & a_{22}(x,y) \end{pmatrix} & \xrightarrow{\left(\frac{x^{\overline{t-u}}}{y^{\overline{v-\overline{t}}}}\right)} P_t \\
P_u \oplus P_v & \xrightarrow{\left(\frac{x^{\overline{t-u}}}{y^{\overline{v-\overline{t}}}}\right)} P_{\tilde{t}}\n\end{array}
$$

Using 4.1 and the fact that $\overline{\tilde{t} - u} = \overline{\tilde{t} - t} + \overline{t - u}$, we get

$$
a_{11}(x,y) = \sum_{h=0}^{\lfloor \frac{p-1-\overline{t-t}}{r} \rfloor} \sum_{q=0}^{(p-1-\overline{t-t}+hr)} c_q^h \cdot x^{hr}(xy)^q
$$

Using 4.2 and the fact that $\overline{v-t} = \overline{\tilde{t} - t} + \overline{v - \tilde{t}}$ we get

$$
a_{22}(x,y) = \sum_{h=0}^{\lfloor \frac{p-1-\bar{t}-t}{r} \rfloor} \sum_{q=0}^{(p-1-(\bar{t}-t+hr))} c_q^h \cdot x^{hr}(xy)^{\bar{t}-t+q}
$$

The final map obtained from the first sum is a linear combination for various values of h of C.2 composed with D.1 h times, followed by a composition with $\alpha_{t,\tilde{t}}$.

Now, assuming that $\overline{v-t} < p$, we must do something similar with the second sum. Instead of writing the chain map again as in the case of the first sum, we will go directly to equations 4.1 and 4.2, using the second sum in ℓ_1

$$
(a_{11}(x,y)x^{\overline{\overline{t}-u}}) = x^{\overline{t-u}} \cdot \sum_{h=0}^{\lfloor \frac{p-1-\overline{t-\overline{t}}}{r} \rfloor} \sum_{q'=\overline{\overline{t}-t}}^{(p-1-(\overline{t-\overline{t}}+hr))} d_{q'}^h \cdot y^{\overline{t-\overline{t}}+hr}(xy)^{q'} \qquad (4.3)
$$

$$
(a_{22}(x,y)y^{\overline{v-\overline{t}}}) = y^{\overline{v-t}} \sum_{h=0}^{\lfloor \frac{p-1-\overline{t-\overline{t}}}{r} \rfloor} \sum_{h=0}^{(p-1-(\overline{t-\overline{t}}+hr))} d_{q'}^h \cdot y^{\overline{t-\overline{t}}+hr}(xy)^{q'} \qquad (4.4)
$$

 $q' = \tilde{t} - t$

Since we cannot simply cancel the power of x in equation (4.3) , we must make up the missing power of x from the powers of xy , which produces an extra copy of y^r . The final values are

 $h=0$

$$
a_{11}(x,y) = \sum_{h=0}^{\lfloor \frac{p-1-\overline{t-\overline{t}}}{r} \rfloor} \sum_{q'=\overline{t-\overline{t}}}^{(p-1-(\overline{t-\overline{t}}+hr))} d_{q'}^h \cdot y^{(h+1)r}(xy)^{q'-\overline{t-\overline{t}}} \qquad (4.5)
$$

$$
a_{22}(x,y) = \sum_{h=0}^{\lfloor \frac{p-1-\overline{t-\overline{t}}}{r} \rfloor} \sum_{q'=\overline{t-\overline{t}}}^{(p-1-(\overline{t-\overline{t}}+hr))} d_{q'}^h \cdot y^{hr}(xy)^{q'}
$$
(4.6)

The final map obtained from the first sum is a linear combination for various values of h of C.2 composed with D.2 h times, followed by a composition with $\alpha_{t,\tilde{t}}$.

Case 2.4: In this case we want to combine single and double projectives. Assume first that $Q_t = P_u$, $Q_{\tilde{t}} = P_u \oplus P_v$. For the map $Q_t \to P_t$ to be nonzero, we need $j > p$, as we see from Diagram 2. Since $r \geq j$, for all the maps in Lemma [3.2](#page-9-0) we have $h = 0$. Since $j = \overline{v - u}$, there are no direct maps from P_u to P_v , so the most general map is

$$
(a_1(x, y) \quad 0) \begin{cases} P_u \xrightarrow{x \overline{t-u}} P_t \\ Q_1(x,y) \end{cases} P_u \oplus P_v \xrightarrow{p \overline{t-u}} P_t \longrightarrow P_t
$$

where $a_1 = (xy)^q$ is the only possibility compatible with $r > p$, and then, in order for the map to be well-defined, we must have $\ell_1 = x^{\tilde{t}-t}(xy)^q$. We must have $q \leq p-1$ for the map to be non-zero. This is simply a special case of the map $\alpha_{t,\tilde{t}}$ composed with q copies of C.1 at the beginning or q copies of C.2 at the end. The case where the map goes from P_v to $P_u \otimes P_v$ is entirely dual because again the diagonal map must be zero.

Assume now $Q_{\tilde{t}} = P_u \oplus P_v$, $Q_t = P_u$, $j > p$. Again all $h = 0$ as in Case 3, and there are no direct maps from P_u to P_v . However, in order to give the most general well-defined map we must have the coefficient of the generator of P_v vanishing in the upper triangle as it vanishes in the lower triangle, and for this, we need a high power of xy in $\ell_1(x, y)$, so also in $a_1(x, y)$.

where $a_1(x, y) = \sum_{q=0}^{p-1} a_q^1(xy)^q$ and $\ell_1 = \sum_{q=0}^{p-1} c_{q_1}^1 y^{\tilde{t}-t}(xy)^{q_1}$. Since $\overline{v-\tilde{t}}$ + $\overline{\tilde{t}-t}=\overline{v-t}$, in order to have the composition of the upper triangle equal to zero, we must have $q_1 \geq p-\overline{v-t}$. We have $\tilde{t} - u > \overline{t-u}$ because $t \preceq \tilde{t}$. Now to get the coefficient of the generator of P_u to be equal in both triangles, we require $a^1x^{\overline{t-u}}(xy)^q = c^1x^{\overline{t-u}}(xy)^{\tilde{t}-t+q_1}$. The total power of x will be less than p if $q_1 < p - \tilde{t} - u$. In that case, we take $q = q_1 + \tilde{t} - t$ and $a_q^1 = c_{q_1}^1$. For q and q_1 greater than these critical values, the coefficients can be taken arbitrarily. The case with P_v is dual.

To show that this map can be decomposed as a composition of maps in the catalog, let us start with the case $t = t_0 = u+(j-p)$ and $\tilde{t} = u+(j-p)+1$. In this case, we have the map in Diagram 2 designated by μ_u , where $q_1 = 0$ and $\tilde{t} - t = 1$. We can multiply by a constant to get the general $a_1(x, y)$ and we can compose with maps of type C.1 or C.2 to get higher values of q_1 . To get more general cases, we use compositions with $\alpha_{t_0,t}$ to get $q = \overline{t_0 + 1 - t}$ while still having $q_1 = 0$, and then compose with C.1 and C.2. Finally, to get a general value of \tilde{t} , we compose with $\alpha_{\tilde{t}, t_0+1}$, which increases q to $\tilde{t} - t$ and then compose with an additional q_1 copies of C.1 or C.2.

Case 2.5: It remains to consider the cases from $P'_w \to P'_t$ or $P'_t \to P'_w$ where $w \in I_0$, $t \in J$. If $Q_t = P_u$, then $j > p$, so we do not have to worry about the sums in Lemma [3.2.](#page-9-0)

$$
\begin{array}{ll}\nP_w & & P_u \xrightarrow{\quad x^{\overline{t-u}} \qquad} P_t \\
\text{(5.a)} & \downarrow \text{a}x^{\overline{u-w}}(xy)^q \\
P_u \xrightarrow{\quad x^{\overline{t-u}} \qquad} P_t & P_w \\
P_w & & P_w\n\end{array}
$$

The map (5.a) is non-zero if $\overline{u-w}+q < p$ and well-defined if $\overline{t-w}+q \geq$ p. There is no possibility of a non-zero power of x reaching P_v because $j =$ $\overline{v-u} > p$. The map (5.b) is non-zero if $\overline{u-w} + q < p$ and is automatically well-defined. It is not homotopic to zero as long as $q < \overline{t - u}$. It is irreducible when $q = 0$. There is a dual case where we replace x by y and u by v.

$$
\begin{array}{ll}\nP_w \\
\text{(5.c)} \\
P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{t-u} \\ y^{v-t} \end{pmatrix}} P_t\n\end{array}} P_t \longrightarrow P_t\n\begin{array}{ll}\n\text{(5.d)} \ P_u \oplus P_v \xrightarrow{\begin{pmatrix} x^{t-u} \\ y^{v-t} \end{pmatrix}} P_t \\
P_w\n\end{array}} P_w
$$

The map 5.c will be non-zero when one of the summands in a_1 or a_2 will be non-zero, where

$$
a_1(x,y) = \sum_{h=0}^{\lfloor \frac{p-1}{r} \rfloor} \sum_{q_1=0}^{(p-1-hr)} a_q^{1h} \cdot x^{\overline{u-w}+hr}(xy)^q
$$

+
$$
\sum_{h=0}^{\lfloor \frac{p-1}{r} \rfloor} \sum_{q'_{ii}=0}^{(p-1-hr)} b_{q'_{ii}}^{iih} \cdot x^{\overline{u-w}} y^{hr}(xy)^{q'_{ii}}
$$

$$
a_2(x,y) = 1_{\overline{v-u}} \sum_{h=0}^{\lfloor \frac{p-1-\overline{v-u}}{r} \rfloor} \sum_{q_{12}=0}^{p-1-(\overline{t-t}+hr)} a_{q_{12}}^{12h} \cdot x^{\overline{v-u}+\overline{u-q}+hr}(xy)^{q_{12}}
$$

$$
+ 1_{\overline{u-v}} \sum_{h=0}^{\lfloor \frac{p-1-\overline{t-\overline{t}}{r} \rfloor}{r}} (p-1-(\overline{v-u}+hr)) + 1_{\overline{u-v}} \sum_{h=0}^{\lfloor \frac{p-1-\overline{t-\overline{t}}{r} \rfloor}{r}} b_q^{12h} \cdot y^{\overline{u-v}+\overline{u-w}+hr}(xy)^{q_{21}}
$$

 $q'_{12}=0$

It is well-defined only when the composition of the vertical and horizontal arrows is zero.

In summary, we have found all possible maps between P'_t and P'_t where $u < t, \tilde{t} < v$, and shown how to factor them using intra-arc maps from the catalog. \Box

5 An example of intra-arc maps: $(C_5 \times C_5) \rtimes C_4$

We return to an example we mentioned before in Example [1](#page-3-0) with $I =$ ${0, 1, 2, 3}$. According to what we did in § [2.5](#page-4-0) and Lemma [3.2,](#page-9-0) the composition factors of the projective modules form diamonds. This case gives us an example of the extra maps that can appear when $r < p$. We choose the projective P_2 arbitrarily and make an elementary equivalence with $I_0 = \{0, 1, 3\}.$ We used a mutation for our first example because the resulting tilting complex is simpler. This gives a tilting complex $T' = \bigoplus T'_j$ as in [§2.8](#page-7-3)

$$
T'_0: 0 \rightarrow P_0 \rightarrow 0
$$

\n
$$
T'_1: 0 \rightarrow P_1 \rightarrow 0
$$

\n
$$
T'_2: 0 \rightarrow P_1 \oplus P_3 \rightarrow P_2 \rightarrow 0
$$

\n
$$
T'_3: 0 \rightarrow P_3 \rightarrow 0
$$

In order to compute the new projective module P'_2 in $\text{End}_{D^b(A)}(T',T')$, we want to calculate a basis for $\text{Hom}_{D^b(A)}(T', T'_i)$, $i = 0, \ldots, 3$, by finding all the maps between all the projectives, as already done in the previous Proposition [4.1.](#page-19-0) Since $r < p$ we have a reducible maps involving $x⁴$ from $P_i \to P_i$ and y^4 from $P_i \to P_i$ for $\{i = 0, 1, 3\}$. These maps commute with each other and their product is the socle of the corresponding projective. There are also reducible loops xy obtained by going to a neighbor by x and returning by y, or vice versa. Then $(xy)^4$ also gives the socle.

Since $J = \{2\}, j = 2$ and $p = 5$ we are in the case $j \leq p - 1$ and the adjacent maps come from Diagram 3 as below, where $u = 1, v = 3$ all the complexes in the middle section contract to a single line.

$$
P_0
$$
\n
$$
y \begin{pmatrix} P_0 \\ y \\ z \end{pmatrix} x
$$
\n
$$
e_u = (xy - x^2) \begin{pmatrix} P_1 \\ \uparrow & 1 \\ P_1 \oplus P_3 \end{pmatrix} \begin{pmatrix} q \\ y \\ z \end{pmatrix}
$$
\n
$$
P_1 \oplus P_3 \begin{pmatrix} q \\ z \end{pmatrix} \begin{pmatrix} q \\ z \end{pmatrix}
$$
\n
$$
y \begin{pmatrix} q \\ q \\ z \end{pmatrix} x
$$
\n
$$
y \begin{pmatrix} q \\ q \\ z \end{pmatrix} x
$$
\n
$$
P_0
$$
\n
$$
P_1
$$
\n
$$
x^2 \begin{pmatrix} q \\ q \\ z \end{pmatrix} y^2
$$

We compute the quiver, given in Fig. [5.1.](#page-39-0) We now go through the catalog. From A, Diagram 3, we have various maps shown between P'_0, P'_1, P'_3 . For chain maps involving P'_2 , we let ℓ_0 be the term in degree zero and ℓ_1 be the term in degree one. We start with maps between P'_2 and P'_1, P'_3 given by

$$
\epsilon_1 : P'_1 \to P'_2, \ell_1 = 0. \qquad \epsilon_3 : P'_3 \to P'_2, \ell_1 = 0. \n\ell_0 = (xy, -x^2) \qquad \qquad \ell_0 = (-y^2, xy) \n\pi_1 : P'_2 \to P'_1, \ell_1 = 0. \qquad \qquad \pi_3 : P'_2 \to P'_3, \ell_1 = 0. \n\ell_0 = \begin{pmatrix} id \\ 0 \end{pmatrix} \qquad \qquad \ell_0 = \begin{pmatrix} 0 \\ id \end{pmatrix}
$$

The self-maps in B are in this case reducible, The map in B.1 is the composition $\epsilon_1 \circ \pi_1$ and the map in B.2 is $\epsilon_3 \circ \pi_3$. Since the sum is homotopic to zero, we get a relation that $\epsilon_1 \circ \pi_1 + \epsilon_3 \circ \pi_3 = 0$. Denote by γ the map from

Figure 5.1

C.2 with $q = 1$. It satisfies a relation $\gamma^5 = 0$. It is not reducible because of the contraction of the middle section of Diagram 3 to a single complex P'_3 , which means that we cannot go back and forth using x and y .

From D.1 and D.2 we get self-maps η_1, η_3 and from D.3 we get two selfmaps ι_1, ι_3 on P'_2 , as below

$$
\eta_1 : P'_2 \to P'_2, \ell_1 = x^4.
$$
\n
$$
\eta_3 : P'_2 \to P'_2, \ell_1 = y^4.
$$
\n
$$
\ell_0 = \begin{pmatrix} x^4 & 0 \\ 0 & x^4 \end{pmatrix}
$$
\n
$$
\ell_1 : P'_2 \to P'_2, \ell_1 = x^4.
$$
\n
$$
\ell_0 = \begin{pmatrix} 0 & 0 \\ x^3y & 0 \end{pmatrix}
$$
\n
$$
\ell_1 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}
$$
\n
$$
\ell_2 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}
$$
\n
$$
\ell_3 = \begin{pmatrix} 0 & xy^3 \\ 0 & 0 \end{pmatrix}
$$

where η_1 is the difference between the map in D.1 and the first map in E.3, and η_3 is the difference between D.2 and the second map in E.3, while ι_1, ι_3 are the special cases in D.3. We also get a relation, because $\eta_1 \circ \eta_3 = \eta_3 \circ \eta_1 =$

 γ^4 . In addition, from E.4, we have maps from P'_i to P'_2 :

$$
\tau_3 : P'_3 \to P'_2, \ell_1 = 0. \qquad \tau_1 : P'_1 \to P'_2, \ell_1 = 0. \n\ell_0 = \begin{pmatrix} 0 & y^4 \end{pmatrix} \qquad \qquad \ell_0 = \begin{pmatrix} x^4 & 0 \end{pmatrix}
$$

Besides the relations already listed, we can derive relations from $x^5 = 0$, $y^5 = 0$ and $xy = yx$, and in addition, we also get $(\gamma)^5 = 0$. We believe that these are not yet all relations. There is a loop $\epsilon_3 \circ y^2 \circ \pi_1$ and its dual, which may combine with some of the self-maps to give a relation.

6 REFERENCES

- [C] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson, The ATLAS of Finite Groups. Maximal Subgroups and Characters of Simmple Groups, Clarendon Press, Oxford (1985).
- [CR] C. Curtis and I. Reiner, Representation Theory of Finite Groups and Associative Algebras, American Mathematical Society, Chelsea Publishing, (1962).
- [GS] M. Gerstenhaber, M. Schaps, The modular version of Maschke's theorem for normal abelian p-Sylows Journal of Pure and Applied Algebra 109 (1996), 257-264.
- [H] D. Happel, Triangulated Categories in the Representation Theory of Finite Dimensional Algebras, London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, (1988).
- [K] B. Külshammer, Crossed products and blocks with normal defect groups, Comm. Algebra 13 (1985), no. 1, 147-168.
- [KK] S. Koshitani and N. Kunugi, Broué conjecture holds for principal 3 blocks with elementary abelian defect group of order 9, Journal of Algebra, 248, (2002), 575-604.
- [KZ2] S. König nd A. Zimmermann, Derived equivalences for group rings. With contributions by Bernhard Keller, Markus Linckelmann, Jeremy Rickard and Raphaël Rouquier. Lecture Notes in Mathematics, 1685. Springer-Verlag, Berlin, (1998).
- [Lan] P. Landrock, Finite Group Algebras and their Modules, London Mathematical Society Lecture Note Series 84, Cambridge University Press, Cambridge, (1983), 78-80.
- [M] J. McKay, Graphs singularities and finite groups, Proc. Symp. Pure Math.37 (1980), 183-186.
- [O] T. Okuyama, Some examples of derived equivalent blocks of finite groups, (Japanese) Proceedings of Conference on Group Theory, (1996): English translation available as a preprint.
- [R1] J. Rickard, Morita theory for derived categories, Math. Soc., vol 39, no. 2, (1989), 436–456.
- [R2] J. Rickard, Derived equivalence and stable equivalence, J. Pure and Appl. Alg. 61, 303-317, (1989).
- [RR] I. Reiten and C. Riedtmanm, Skew group algebras in the representation theory of artin algebras, J. Algebra, 92 (1985), 224-282.
- [S] M. Schaps, A modular version of Maschke's theorem for groups with cyclic p-Sylow subgroup, J. of Algebra, 163 (1994), 623-635.
- [SSS] M. Schaps, D. Shapira and O. Shlomo, Quivers of blocks with normal defect group, American Mathematical Society Proc. Symp. Pure Math. 63 (1998), 497-510.
- [SZ2] M. Schaps, E. Zakay-Illouz,Pointed Brauer trees. J. Algebra 246, no. 2, (2001), 647–672.
- [SZ] M. Schaps and Z. Zvi, Mutations and pointing for Brauer trees algebras, arXiv 1606.04341, (2016), Osaka J. Math. 57,(2020), 689-709.
- [Sw] W. Schwarz, Die struktur modularer gruppenringe endlicher gruppen der p-Länge 1 , J. Algebra, 60 (1979), 51-75.
- [W] C.A. Weibel, An Introduction to Homological Algebra, Cambridge Studies in Advanced Mathematics, 38. Cambridge University Press, Cambridge, (1994).