

# On the max min of the algebraic degree and the nonlinearity of a Boolean function on an affine subspace

Jan Kristian Haugland

September 6, 2024

## Abstract

We investigate the max min of the algebraic degree and the nonlinearity of a Boolean function in  $n$  variables when restricted to a  $k$ -dimensional affine subspace of  $\mathbb{F}_2^n$ . Previous authors have focused on the cases when the max min of the algebraic degree is 0 or 1. Upper bounds, lower bounds and a conjecture on the exact value in special cases are presented.

## 1 Introduction

Let  $f$  be a Boolean function in  $n$  variables. With  $k \leq n$ , let  $\alpha(f, k)$  denote the minimal algebraic degree of  $f$  when restricted to a  $k$ -dimensional affine subspace of  $\mathbb{F}_2^n$  (subsequently referred to as a  $k$ -dimensional *flat*) and let  $g(n, k) = \max_f \alpha(f, k)$ .

Similarly, let  $\alpha'(f, k)$  denote the minimal nonlinearity of  $f$  when restricted to a  $k$ -dimensional flat, and let  $g'(n, k) = \max_f \alpha'(f, k)$ . The nonlinearity of a Boolean function is the minimal Hamming distance to an affine function.

$f$  is said to be  $k$ -normal if  $\alpha(f, k) = 0$  and weakly  $k$ -normal if  $\alpha(f, k) \leq 1$  (which is equivalent to  $\alpha'(f, k) = 0$ ), and these cases have been of particular interest. Dubuc [1] proved that for  $n \leq 7$ , any Boolean function in  $n$  variables is  $\lfloor \frac{n}{2} \rfloor$ -normal, i.e.,  $g(n, k) = 0$  if  $n \leq 7$  and  $k = \lfloor \frac{n}{2} \rfloor$ . In this note, we give an upper bound (Theorem 2.5) and a lower bound (Theorem 2.3) for  $g(n, k)$ , as well as considering exact values when  $n - k \in \{1, 2\}$  (Theorem 2.2 and the conjecture in Section 5) and in some single cases (Section 3).

We focus mostly on  $g(n, k)$ , but we also give a lower bound for  $g'(n, k)$  and values in some single cases.

## 2 General bounds

**Lemma 2.1.** *The algebraic degree of  $f$  when restricted to a  $k$ -dimensional flat ( $k \geq 1$ ) is equal to  $k$  if and only if the sum of  $f(x)$  over all vertices of the flat is 1.*

*Proof.* Confer [2], Theorem 2.5.  $\square$

**Theorem 2.2.** *If  $n \geq 2$ , then  $g(n, n-1) = n-2$ .*

*Proof.* First, we prove that  $g(n, n-1) < n-1$ . This follows almost directly from Lemma 2.1. We only need to identify an  $(n-1)$ -dimensional flat on which the sum of  $f(x)$  is 0.

Let  $S$  be any  $(n-2)$ -dimensional flat.  $S$  must have three additional cosets which are also flats. There must be two of them on which  $f$  has the same sum, and so we can simply combine them into an  $(n-1)$ -dimensional flat on which  $\sum f(x) = 0$  as required.

Second, we verify that there exists an example of a function  $f$  for which  $\alpha(f, n-1) \geq n-2$ . We can use the function that maps  $x$  to 1 if the weight of  $x$  is 0 or 1, and 0 otherwise. A flat of dimension  $n-1$  must contain at least one  $x$  such that  $f(x) = 1$ , otherwise the complement would contain all of them and generate the whole of  $\mathbb{F}_2^n$ . Then the intersection with either  $x_i = 1$  (if  $x$  is 1 in the  $i$ th coordinate) or  $\sum x_i = 0$  (if  $x$  is  $(0, 0, \dots, 0)$ ) contains exactly one vertex  $y$  for which  $f(y) = 1$ , and  $f$  must have maximal degree  $\geq n-2$  here. Therefore,  $f$  is also of degree  $\geq n-2$  on the original flat.  $\square$

**Theorem 2.3.** *If  $n, k, d$  are integers with  $n > k > d \geq 0$  satisfying*

$$(k+1)(n-k) + 2 \leq \binom{k}{0} + \dots + \binom{k}{k-d-1}$$

*then  $g(n, k) > d$ . Similarly, if  $m < 2^{k-2}$  and*

$$\binom{2^k}{0} + \dots + \binom{2^k}{m} \leq 2^{2^k - (k+1)(n-k+1) - 2}$$

*then  $g'(n, k) > m$ .*

*Proof.* We begin by observing that  $(1 - \frac{1}{2^i})^{2^{i-1}} \geq \frac{1}{2}$  for  $i \geq 1$ , and thus  $\prod_{i \geq 1} (1 - \frac{1}{2^i}) \geq (\frac{1}{2})^{1 + \frac{1}{2} + \frac{1}{4} + \dots} = \frac{1}{4}$ . (We could have used the exact value 0.288788... for the infinite product, but  $\geq \frac{1}{4}$  is good enough here.) We can now estimate the number of  $k$ -dimensional flats in  $\mathbb{F}_2^n$ , which is given by

$$2^{n-k} \frac{(2^n - 1)(2^{n-1} - 1) \dots (2^{n-k+1} - 1)}{(2^k - 1)(2^{k-1} - 1) \dots (2^1 - 1)}$$

We have

$$\begin{aligned} (2^n - 1)(2^{n-1} - 1) \dots (2^{n-k+1} - 1) &< 2^{k(n - \frac{k-1}{2})} \\ (2^k - 1)(2^{k-1} - 1) \dots (2^1 - 1) &> \left( \prod_{i \geq 1} \left(1 - \frac{1}{2^i}\right) \right) 2^{k \frac{k+1}{2}} \geq \frac{1}{4} \times 2^{k \frac{k+1}{2}} \\ \implies 2^{n-k} \frac{(2^n - 1)(2^{n-1} - 1) \dots (2^{n-k+1} - 1)}{(2^k - 1)(2^{k-1} - 1) \dots (2^1 - 1)} &< 2^{(n-k) + k(n-k) + 2} \end{aligned}$$

$$= 2^{(k+1)(n-k)+2}$$

For each of these flats, the proportion of functions of algebraic degree  $\leq d$  on it is  $2^{-\binom{k}{0} + \dots + \binom{k}{k-d-1}}$ . By the condition of the Theorem, the proportion of functions of algebraic degree  $\leq d$  on at least one flat must be smaller than 1, and it follows that there must exist a function  $f : \mathbb{F}_2^n \rightarrow \mathbb{F}_2$  such that  $\alpha(f, k) > d$ .

The argument only uses the scarceness of functions of low degree, and a similar argument applies to nonlinearity (or any other property, for that matter). Since  $m < 2^{k-2}$ , the proportion of functions of nonlinearity  $\leq m$  on any given  $k$ -dimensional flat is

$$\left( \binom{2^k}{0} + \dots + \binom{2^k}{m} \right) 2^{k+1-2^k}$$

and the bound for  $g'(n, k)$  follows.  $\square$

**Corollary 2.3.1.** *Suppose  $n, k$  are integers and that  $k \geq 5$ . If  $k+2 \leq n \leq \frac{3k-1}{2}$ , then  $g(n, k) \geq k-2$ . If  $\frac{3k}{2} \leq n \leq \frac{(k+1)(k+4)}{6}$ , then  $g(n, k) \geq k-3$ .*

*Proof.* Regarding the first claim, we can apply Theorem 2.3 with  $d = k-3$ . Since  $n \leq \frac{3k-1}{2}$ , we have

$$(k+1)(n-k) + 2 \leq \frac{k^2+3}{2} < \frac{k^2+k+2}{2} = \binom{k}{0} + \binom{k}{1} + \binom{k}{2}$$

For the second claim, we put  $d = k-4$  instead. Since  $n \leq \frac{(k+1)(k+4)}{6}$ , we have

$$(k+1)(n-k) + 2 \leq \frac{k^3+3k+16}{6} \leq \frac{k^3+5k+6}{6} = \binom{k}{0} + \binom{k}{1} + \binom{k}{2} + \binom{k}{3}$$

In both cases, the condition of Theorem 2.3 is satisfied.  $\square$

We can also give an upper bound by simple means.

**Lemma 2.4.** *If  $a, b$  and  $c$  are nonnegative integers satisfying*

$$\max(a+1, b) \leq c \leq a+b$$

*then*

$$\lceil 2^a \frac{2^b-1}{2^c-1} \rceil = 2^{a+b-c}$$

*Proof.* We have

$$2^a \frac{2^b-1}{2^c-1} = 2^a \left( \frac{1}{2^{c-b}} - \frac{1-2^{b-c}}{2^c-1} \right) > 2^a \left( \frac{1}{2^{c-b}} - \frac{1}{2^c-1} \right)$$

and since  $a < c$ , it follows that

$$2^a \frac{2^b-1}{2^c-1} > 2^{a+b-c} - 1$$

On the other hand,  $b \leq c$  implies that

$$2^a \frac{2^b - 1}{2^c - 1} \leq 2^{a+b-c}$$

Lastly, since  $c \leq a + b$ , it is clear that  $2^{a+b-c}$  is an integer, and the conclusion follows.  $\square$

**Theorem 2.5.** *If  $k > d \geq 0$  and  $n \geq 2^{k-1} + k - \lfloor 2^{d-1} \rfloor$ , then  $g(n, k) \leq d$ .*

*Proof.* Two  $\delta$ -dimensional flats  $\{x_i\}$  and  $\{y_j\}$  are said to be *parallel* if there exists a vector  $z \in \mathbb{F}_2^n$  such that  $\{x_i + z\} = \{y_j\}$ . Note that there are exactly  $2^{n-\delta}$  flats that are parallel to  $\{x_i\}$  (including itself), all mutually disjoint.

First, we prove the case  $d = 0$ . It suffices to assume that  $n = 2^{k-1} + k$  and verify that  $\alpha(f, k) = 0$  for any function  $f$  in  $n$  variables.

For  $0 \leq \delta \leq k$ , let  $h(\delta)$  denote the maximal number of parallel  $\delta$ -dimensional flats on which  $f$  is constant. Trivially,  $h(0) = 2^n$ .

For  $\delta = 1, 2, \dots$ , there are at least  $\lceil \frac{h(\delta-1)(h(\delta-1)-2)}{4} \rceil$  pairs of disjoint parallel  $(\delta-1)$ -dimensional flats on which  $f$  is equal to the same constant, while there are  $2^{n-\delta+1} - 1$  ways to pair up all  $(\delta-1)$ -dimensional flats parallel to a given one into parallel  $\delta$ -dimensional flats. Thus we have

$$h(\delta) \geq \lceil \frac{h(\delta-1)(h(\delta-1)-2)}{4(2^{n-\delta+1}-1)} \rceil$$

We can now show that  $h(\delta) \geq 2^{n-(2^\delta+\delta)+1}$  for  $0 \leq \delta \leq k-1$  by induction on  $\delta$ . By Lemma 2.4 with  $a = b = n - (2^{\delta-1} + \delta) + 1$ ,  $c = n - \delta + 1$  we have the induction step

$$\lceil \frac{h(\delta-1)}{2} \frac{\frac{h(\delta-1)}{2} - 1}{2^{n-\delta+1}-1} \rceil = 2^{n-(2^\delta+\delta)+1}$$

In particular,  $h(k-1) \geq 4$ , so we have at least two different parallel  $(k-1)$ -dimensional flats on which  $f$  is the same constant, and we can combine them into a  $k$ -dimensional flat. This completes the case  $d = 0$ . (It is straightforward to verify that the conditions on  $a$ ,  $b$  and  $c$  are satisfied.)

Second, when  $d > 0$ , the idea is to not distinguish between the admissible  $(\delta-1)$ -dimensional flats when forming  $\delta$ -dimensional flats for  $\delta > k-d$ . Then  $f$  will have algebraic degree at most 1 on the  $(k-d+1)$ -dimensional flats, at most 2 on the  $(k-d+2)$ -dimensional flats, and so on.

With  $n = 2^{k-1} + k - 2^{d-1}$ , the proof that  $h(k-d) \geq 2^{n-(2^{k-d}+k-d)+1}$  is similar as above, and we omit the details. Then, for  $k-d+1 \leq \delta \leq k$ , let  $h'(\delta)$  denote the maximal number of parallel  $\delta$ -dimensional flats formed by the admissible  $(\delta-1)$ -dimensional flats, and put  $h'(k-d) = h(k-d)$ . We have

$$h'(\delta) \geq \lceil \frac{h(\delta-1)(h(\delta-1)-1)}{2(2^{n-\delta+1}-1)} \rceil$$

and as above, we can show that  $h'(\delta) \geq 2^{n-(2^\delta+\delta)+2^{\delta-k+d}}$  for  $k-d+1 \leq \delta \leq k-1$ . By Lemma 2.4 with  $a = n - (2^{\delta-1} + \delta) + 2^{\delta-k+d-1}$ ,  $b = a + 1$ ,  $c = n - \delta + 1$ , we have the induction step

$$\left\lceil \frac{h'(\delta-1)}{2} \frac{h'(\delta-1)-1}{2^{n-\delta+1}-1} \right\rceil = 2^{n-(2^\delta+\delta)+2^{\delta-k+d}}$$

Thus we have  $h'(k-1) \geq 2$ , i.e., two different parallel  $(k-1)$ -dimensional flats that we can combine into a  $k$ -dimensional flat on which  $f$  has algebraic degree at most  $d$ . □

### 3 Single cases

We have verified that  $g(5, 3) = 1$ ,  $g(6, 4) = 2$  and  $g(7, 5) = 3$  with the help of Langevin's classification of Boolean functions in up to six variables [3] and in seven variables [4]. (A few more details are given in Section 5.) Polujan *et al.* [5] found a bent function in 8 variables that is not 4-normal, and Canteaut *et al.* [6] found a bent function in 10 variables that is not weakly 5-normal. In other words,  $g(8, 4) \geq 1$  and  $g(10, 5) \geq 2$ .

Given positive integers  $k$  and  $d$ , an algorithm for searching for a function  $f$  such that  $\alpha(f, k) \geq d$  is as follows. Start with a random function  $f_0$  in  $k$  variables. For each  $i$ ,  $1 \leq i \leq N$ , let  $f_i$  be given by the same truth table as that of  $f_{i-1}$  except at one or two randomly chosen points. If the number of flats on which the degree is  $< d$  is higher for  $f_i$  than for  $f_{i-1}$ , discard the new  $f_i$  and let  $f_i \leftarrow f_{i-1}$  instead. If the number of such flats has dropped to zero within  $N$  steps, we are done. If not, we may be stuck in a local dead end and can try again from the top. In this manner we were able to find a function  $f^{(7,4)} : \mathbb{F}_2^7 \rightarrow \mathbb{F}_2$  such that  $\alpha(f^{(7,4)}, 4) = 2$  and a function  $f^{(8,5)} : \mathbb{F}_2^8 \rightarrow \mathbb{F}_2$  such that  $\alpha(f^{(8,5)}, 5) = 3$ . Langevin found equivalent functions with fewer terms in the algebraic normal form (private communication), specifically

$$\begin{aligned} f^{(7,4)}(x_1, \dots, x_7) \sim & x_1x_2x_3x_4x_5x_7 \oplus x_1x_2x_3x_4x_5 \oplus x_1x_2x_3x_4x_6x_7 \oplus x_1x_2x_3x_4 \oplus \\ & x_1x_2x_4x_5x_6 \oplus x_1x_2x_4x_5x_7 \oplus x_1x_2x_6x_7 \oplus x_1x_2x_6 \oplus x_1x_2x_7 \oplus x_1x_3x_4x_5x_6 \oplus x_1x_3x_4x_5 \oplus \\ & x_1x_3x_4x_7 \oplus x_1x_3x_5x_7 \oplus x_1x_4x_5x_7 \oplus x_1x_4x_5 \oplus x_1x_4x_6x_7 \oplus x_1x_4x_6 \oplus x_1x_6x_7 \oplus \\ & x_1x_7 \oplus x_2x_3x_4x_5 \oplus x_2x_3x_5x_6x_7 \oplus x_2x_3x_5x_7 \oplus x_2x_4x_6x_7 \oplus x_2x_4x_6 \oplus x_2x_5x_6x_7 \oplus \\ & x_3x_4x_5x_6 \oplus x_3x_4x_5x_7 \oplus x_3x_4x_6 \oplus x_3x_4 \oplus x_3x_5 \oplus x_3x_7 \oplus x_4x_5x_7 \oplus x_5x_7 \oplus x_6x_7 \end{aligned}$$

and

$$\begin{aligned} f^{(8,5)}(x_1, \dots, x_8) \sim & x_1x_2x_3x_4x_5x_6x_8 \oplus x_1x_2x_3x_4x_5x_6 \oplus x_1x_2x_3x_4x_6x_7x_8 \oplus \\ & x_1x_2x_3x_4x_6x_7 \oplus x_1x_2x_3x_4x_6 \oplus x_1x_2x_3x_5x_6x_7 \oplus x_1x_2x_3x_5x_8 \oplus x_1x_2x_3x_6 \oplus x_1x_2x_4x_5x_6x_8 \oplus \\ & x_1x_2x_4x_5x_6 \oplus x_1x_2x_4x_6 \oplus x_1x_2x_4x_7 \oplus x_1x_2x_5x_6x_7 \oplus x_1x_2x_6x_7 \oplus x_1x_2x_6 \oplus x_1x_3x_4x_5x_8 \oplus \\ & x_1x_3x_4x_5 \oplus x_1x_3x_4x_6 \oplus x_1x_3x_4x_7x_8 \oplus x_1x_3x_4x_7 \oplus x_1x_3x_5x_7x_8 \oplus x_1x_3x_5x_7 \oplus \\ & x_1x_3x_5x_8 \oplus x_1x_3x_6x_7x_8 \oplus x_1x_3x_8 \oplus x_1x_4x_5x_6x_7x_8 \oplus x_1x_4x_5x_6x_7 \oplus x_1x_4x_6 \oplus \\ & x_1x_4x_7 \oplus x_1x_4x_8 \oplus x_1x_5x_7 \oplus x_1x_7x_8 \oplus x_2x_3x_4x_5x_6x_7x_8 \oplus x_2x_3x_4x_5x_7x_8 \oplus x_2x_3x_4x_5x_8 \oplus \end{aligned}$$

$$\begin{aligned}
& x_2x_3x_4x_6x_7x_8 \oplus x_2x_3x_4x_6x_8 \oplus x_2x_3x_4x_7 \oplus x_2x_3x_4x_8 \oplus x_2x_3x_5x_6x_7x_8 \oplus x_2x_3x_5x_6x_7 \oplus \\
& x_2x_3x_5 \oplus x_2x_3x_6x_7x_8 \oplus x_2x_3x_6x_8 \oplus x_2x_3x_7 \oplus x_2x_4x_5x_6x_7x_8 \oplus x_2x_4x_5x_6x_7 \oplus \\
& x_2x_4x_5x_7x_8 \oplus x_2x_4x_6x_7x_8 \oplus x_2x_4x_6x_8 \oplus x_2x_4x_6 \oplus x_2x_4x_7x_8 \oplus x_2x_5x_6x_7 \oplus x_2x_5x_7x_8 \oplus \\
& x_2x_5x_8 \oplus x_2x_6x_7x_8 \oplus x_3x_4x_5x_6x_7x_8 \oplus x_3x_4x_5x_6x_7 \oplus x_3x_4x_5x_8 \oplus x_3x_4x_6x_8 \oplus \\
& x_3x_4x_6 \oplus x_3x_4x_7 \oplus x_3x_5x_6x_7 \oplus x_3x_5x_6x_8 \oplus x_3x_5x_7x_8 \oplus x_3x_5x_8 \oplus x_3x_6x_7x_8 \oplus \\
& x_3x_6x_7 \oplus x_4x_5x_6x_7x_8 \oplus x_4x_5x_6x_8 \oplus x_5x_7x_8 \oplus x_6x_7x_8
\end{aligned}$$

Thus, since  $g(7, 4) \leq g(6, 4)$  and  $g(8, 5) \leq g(7, 5)$ , we have  $g(7, 4) = 2$  and  $g(8, 5) = 3$ .

As for nonlinearity, the optimal functions are of course bent or almost bent when  $k = n$ , which gives  $g'(2, 2) = 1$ ,  $g'(3, 3) = 2$ ,  $g'(4, 4) = 6$ ,  $g'(5, 5) = 12$  and  $g'(6, 6) = 28$ . Using Langevin's classifications, we found that this is common also in other cases with small  $n$  and  $k$  and nonzero value of  $g'(n, k)$ . We have  $g'(4, 3) = 2$ ,  $g'(5, 4) = 4$ ,  $g'(6, 4) = 4$  and  $g'(6, 5) = 12$ .

## 4 Tabulated values

Taking all the above results and single cases into account, we get the following values and lower bounds for  $g(n, k)$  when  $n \leq 12$  and  $k \leq 6$  (upper bounds, where not specified, can be inferred from lower values for  $n$ ).

$k \backslash n$	1	2	3	4	5	6	7	8	9	10	11	12
1	1	0	0	0	0	0	0	0	0	0	0	0
2		2	1	0	0	0	0	0	0	0	0	0
3			3	2	1	0	0	0	0	0	0	0
4				4	3	2	2	$\geq 1$	$\geq 0$	$\geq 0$	0 or 1	0
5					5	4	3	3	$\geq 2$	$\geq 2$	$\geq 0$	$\geq 0$
6						6	5	$\geq 4$	$\geq 3$	$\geq 3$	$\geq 3$	$\geq 2$

Here are the corresponding values and bounds for  $g'(n, k)$ . As previously indicated,  $g(n, k) \leq 1 \iff g'(n, k) = 0$  and  $g(n, k) \geq 2 \iff g'(n, k) \geq 1$ .

$k \backslash n$	1	2	3	4	5	6	7	8	9	10	11	12
1	0	0	0	0	0	0	0	0	0	0	0	0
2		1	0	0	0	0	0	0	0	0	0	0
3			2	2	0	0	0	0	0	0	0	0
4				6	4	4	$\geq 1$	$\geq 0$	$\geq 0$	$\geq 0$	0	0
5					12	12	$\geq 3$	$\geq 2$	$\geq 1$	$\geq 1$	$\geq 0$	$\geq 0$
6						28	$\geq 16$	$\geq 12$	$\geq 9$	$\geq 7$	$\geq 5$	$\geq 3$

## 5 A conjecture

We think the following statement is true.

**Conjecture:** Let  $k$  be an integer  $\geq 2$ . Then  $g(k+2, k) = k-2$ .

For  $k=2$ , the claim follows from the result of Dubuc, and for  $k \in \{3, 4, 5\}$ , this has also been verified, as mentioned in Section 3.

For  $k \geq 6$ , Corollary 2.3.1 says that  $k-2$  is indeed a valid lower bound for  $g(k+2, k)$ . On the other hand, we can give a heuristic argument for why  $g(k+2, k)$  is probably not as high as  $k-1$ . The algebraic degree of the restriction of a random function  $f: \mathbb{F}_2^{k+2} \rightarrow \mathbb{F}_2$  to a given  $k$ -dimensional flat is  $\geq k-1$  with probability  $1 - \frac{1}{2^{\binom{k}{6} + \binom{k}{1}}} = 1 - \frac{1}{2^{k+1}}$ , and so the probability that this holds for all  $k$ -dimensional flats should be approximately

$$\left(1 - \frac{1}{2^{k+1}}\right)^{2^2 \frac{(2^{k+2}-1)(2^{k+1}-1)}{3}}$$

(of course, the probabilities are not actually independent, which is why this is a heuristic argument and not a proof). For  $k$  sufficiently large, this is approximately  $e^{-\frac{4}{3}2^{k+2}}$ . There are "only"  $2^{2^{k+2}}$  Boolean functions in  $k+2$  dimensions, which suggests that the expected number of functions for which every restriction to a  $k$ -dimensional flat is of degree  $\geq k-1$  should be about  $(\frac{2}{e^{4/3}})^{2^{k+2}}$  which tends to zero quite rapidly. Actually, we only need to check one representative from each of a much smaller number of classes of functions, which diminishes the chances of the existence of such a function even further.

For  $k \in \{2, 3, 4, 5\}$ , a computer search for the Boolean functions in  $k+2$  variables given in [3], [4] of algebraic degree  $\leq k-2$  on the *smallest* number of  $k$ -dimensional flats gave the following results.

$k$	Function	No. of flats
2	$x_1x_2x_3 \oplus x_1x_4 \oplus x_2$	10 / 140
3	$x_1x_2x_3x_4 \oplus x_1x_2x_5 \oplus x_1x_4 \oplus x_2x_3$	15 / 620
4	$x_1x_2x_3x_4 \oplus x_1x_3x_5 \oplus x_1x_4x_6$ $\oplus x_2x_3x_5 \oplus x_2x_3x_6 \oplus x_2x_4x_5$	20 / 2604
5	$x_1x_2x_3x_4x_5x_6 \oplus x_1x_2x_3x_4x_7 \oplus x_1x_2x_4x_5 \oplus x_1x_2x_6x_7$ $\oplus x_1x_3x_4x_5 \oplus x_1x_3x_4x_6 \oplus x_1x_3x_5x_7 \oplus x_2x_3x_5x_6$	73 / 10668

## References

- [1] S. Dubuc. Etude des propriétés de dégénérescence et de normalité des fonctions booléennes et construction de fonctions q-aires parfaitement non-linéaires, Ph.D. Thesis, Université de Caen, 2001.
- [2] T. W. Cusick and Y. Cheon. Counting balanced Boolean functions in  $n$  variables with bounded degree. *Experimental Math.* 16:1 (2007), 101-105.

- [3] P. Langevin. Classification of Boolean functions under the affine group.  
<http://langevin.univ-tln.fr/project/agl/agl.html>
- [4] P. Langevin. Classification of Boolean functions in dimension 7.  
<https://langevin.univ-tln.fr/project/agl7/aglclass.html>
- [5] A. Polujan, L. Mariot and S. Picek. Normality of Boolean bent functions in eight variables, revisited.  
[https://boolean.w.uib.no/files/2023/08/BFA2023\\_paper\\_10.pdf](https://boolean.w.uib.no/files/2023/08/BFA2023_paper_10.pdf)
- [6] A. Canteaut, M. Daum, H. Dobbertin and G. Leander. Normal and Non Normal Bent Functions. Workshop on Coding and Cryptography '03, pages 91 - 100.

NORWEGIAN NATIONAL SECURITY AUTHORITY (NSM), NORWAY  
*E-mail address:* [admin@neutreeko.net](mailto:admin@neutreeko.net)