

# Projective (or spin) representations of finite groups. II

Tatsuya Tsurii<sup>1</sup>, Satoe Yamanaka<sup>2</sup>, Itsumi Mikami<sup>3</sup>, and Takeshi Hirai<sup>4 5</sup>

## Abstract

In the previous paper, we proposed a practical method of constructing explicitly representation groups  $R(G)$  for finite groups  $G$ , and apply it to certain typical finite groups  $G$  with Schur multiplier  $M(G)$  containing prime number 3. In this paper, we construct a complete list of irreducible projective (or spin) representations of  $G$  and compute their characters (called spin characters). It is a continuation of our study of spin representations in the cases where  $M(G)$  contains prime number 2 to the cases where other prime  $p$  appears, firstly  $p = 3$ . We classify irreducible spin representations and calculate spin characters according to their spin types.

**2020 Mathematics Subject Classification:** Primary 20C25; Secondary 20F05, 20E99.

**Key Words:** one-step efficient central extension, projective representation, representation group of finite group, spin representation, spin character.

**1. Introduction.** Let  $G$  be a finite group. Representation group  $R(G)$  is a certain central extension of  $G$  by  $M(G)$  as its central subgroup, and any irreducible spin representation  $\pi$  of  $G$  can be induced naturally from a linear representation  $\tilde{\pi}$  of  $R(G)$ . There exists a one-dimensional character  $\chi$  of  $M(G)$  such that  $\tilde{\pi}(z) = \chi(z)I$  ( $z \in M(G)$ ), called *spin type* of  $\pi$ , where  $I$  denotes the identity operator.

Our practical method of constructing representation group  $R(G)$  of  $G$ , named as “*Efficient stairway up to the Sky*”, is to start with a presentation of  $G$  by a pair of **Set of generators** and **Set of fundamental relations**. A central extension

$$(1) \quad 1 \longrightarrow A \longrightarrow H \xrightarrow{\alpha} G \longrightarrow 1 \quad (\text{exact})$$

of  $G$  by an abelian group  $A$  is called an *efficient covering* if  $A$  is contained in the commutator subgroup  $[H, H]$  of  $H$ , i.e.,

$$(2) \quad A \subset Z(H) \cap [H, H].$$

Take a pair  $x, y \in G$  commuting with each other and of the same order  $d > 1$ . Then try to take a *one-step efficient covering* given by the commuting relation  $[x, y] = 1$ , in such a way that,  $A = \langle z \rangle$  is a cyclic group generated by a central element  $z$  of order  $d$  which is given under the homomorphism  $\alpha$  as

$$(3) \quad [\xi, \eta] = z, \quad \xi \xrightarrow{\alpha} x, \eta \xrightarrow{\alpha} y, z \xrightarrow{\alpha} 1.$$

When we know Schur multiplier  $M(G)$  of  $G$ , we may arrive up to  $R(G)$  by repeating one-step efficient

central extensions step by step because  $R(G)$  is characterized [1, §2] as an efficient central extension  $H$  of  $G$  such that

$$M(G) \subset Z(H) \cap [H, H], \quad H/M(G) \cong G.$$

In cases we are treating here, for every one-step efficient central extensions, the resulting central extension  $H$  is expressed as semidirect product  $H = U \rtimes W$ , and so, construction of its irreducible representations is realized by Mackey’s induced representations when  $U$  is abelian (cf. [2, Theorem 5.1]), and by repeated application of the classical method in [3] for general semidirect products.

**2. Preliminaries.** Let  $G'$  be a central extension of  $G$  by an abelian group  $Z$  as

$$1 \longrightarrow Z \longrightarrow G' \xrightarrow{\alpha} G \longrightarrow 1.$$

Take a section  $s : G \rightarrow S_G \subset G'$  for the canonical homomorphism  $\alpha$ . Then, we have a  $Z$ -valued cocycle  $z_{g,h} \in Z$  ( $g, h \in G$ ) such that

$$s(g)s(h) = z_{g,h}s(gh), \quad \exists z_{g,h} \in Z.$$

For an irreducible linear representation  $\Pi$  of  $G'$ , put

$$\pi(g) := \Pi(s(g)) \quad (g \in G).$$

Let  $\chi \in \hat{Z}$  be the *spin type* of  $\Pi$  with respect to  $Z$ , that is,  $\Pi(z) = \chi(z)I$  ( $z \in Z$ ), then  $\pi$  is a spin representation of  $G$  associated with a factor set

$$r_{g,h} = \chi(z_{g,h}) \in \mathbf{T}^1 := \{\lambda \in \mathbf{C}; |\lambda| = 1\}.$$

<sup>1</sup>Tokyo University of Information Sciences, Onaridai 4-1, Wakaba-ku, Chiba-shi, Chiba, 265-8501, JAPAN

<sup>2</sup>Department of Liberal Studies, National Institute of Technology, Nara College, 22 Yata-cho, Yamatokoriyama, Nara 639-1080, JAPAN

<sup>3</sup>Hirai Mathematics Institute

<sup>4</sup>Kyoto University, Yoshida-honmachi, Sakyo-ku, Kyoto 606-8501, JAPAN

<sup>5</sup>Hirai Mathematics Institute, 22-5 Nakazaichi-cho, Iwakura, Sakyo, Kyoto 606-0027, JAPAN

In fact, for  $g, h \in G$ , we have

$$\begin{aligned}\pi(g)\pi(h) &= \Pi(s(g))\Pi(s(h)) = \Pi(s(g)s(h)) \\ &= \Pi(z_{g,h}s(gh)) = \chi(z_{g,h})\Pi(s(gh)) = r_{g,h}\pi(gh).\end{aligned}$$

We call  $\pi$  a *sectional restriction* (onto  $G$ ) of  $\Pi$ .

If we take another section  $s' : G \rightarrow G'$ , then it induces a spin representation  $\pi'(g) := \Pi(s'(g))$  of  $G$  associated to an equivalent factor set  $r'_{g,h}$  ( $g, h \in G$ ).

We know by Schur the following.

**Theorem 2.1.** (Schur [4]) *As a central extension  $G'$  above, take a representation group  $R(G)$ . Then, for any factor set  $r_{g,h}$  for  $G$ , there exists an irreducible linear representation  $\Pi$  of  $R(G)$  which induces as above spin representation  $\pi$  of  $G$  associated to a factor set which is equivalent to the given  $r_{g,h}$ .*

**3. Irreducible representations of semidirect product groups.** Let  $G$  be a compact group of semidirect product type  $G = U \rtimes W$ , where  $U$  is a compact group, normal in  $G$ , and  $W$  is a finite group. We gave in [3] the so-called *classical method* to construct a complete list of representatives of the dual  $\widehat{G}$ , or of all the equivalence classes of IRs of  $G$ . We will apply it in the case of finite groups.

So, starting from now on, we assume  $G$  to be finite, and explain briefly about the method.

**3.1. Construction of IRs of semidirect product groups.** Take an IR  $\rho$  of  $U$  and consider its equivalence class  $[\rho] \in \widehat{U}$ . Every  $w \in W$  acts on  $u \in U$  as  $u \rightarrow w(u)$ , and on  $\rho$  as

$$({}^w\rho)(u) := \rho(w^{-1}(u)) \quad (u \in U),$$

and therefore on equivalent classes as  $[\rho] \rightarrow [{}^w\rho]$ . Denote by  $\widehat{U}/W$  the set of  $W$ -orbits in the dual  $\widehat{U}$  of  $U$ .

**(Step 1)** Take a stationary subgroup  $W([\rho])$  in  $W$  of an equivalence class  $[\rho]$ , that is,

$$W([\rho]) := \{w \in W; {}^w\rho \cong \rho\},$$

and put  $H := U \rtimes W([\rho])$ .

**(Step 2)** For  $w \in W([\rho])$ , we determine explicitly an intertwining operator  $J_\rho(w)$  given as

$$(4) \quad \rho(w(u)) = J_\rho(w)\rho(u)J_\rho(w)^{-1} \quad (u \in U).$$

Then it is determined up to a non-zero scalar factor. Hence we have a spin representation  $W([\rho]) \ni w \mapsto J_\rho(w)$ . Let  $\alpha_{v,w}$  be its factor set given as

$$J_\rho(v)J_\rho(w) = \alpha_{v,w}J_\rho(vw) \quad (v, w \in W([\rho])).$$

**(Step 3)** Let  $W([\rho])'$  be a central extension of  $W([\rho])$  sufficiently large compared to the cocycle

$\alpha_{v,w}$ :

$$1 \rightarrow Z \rightarrow W([\rho])' \xrightarrow{\Phi} W([\rho]) \rightarrow 1 \quad (\text{exact}),$$

where  $\Phi$  denotes the canonical homomorphism. Then, by Theorem 2.1,  $J_\rho$  can be lifted up to a linear representation  $J'_\rho$  of  $W([\rho])'$ . Put  $H' := U \rtimes W([\rho])'$  with the action

$$w'(u) := w(u), \quad w' \in W([\rho])', \quad w = \Phi(w').$$

Put also

$$\pi^0((u, w')) := \rho(u) \cdot J'_\rho(w') \quad (u \in U, w' \in W([\rho])'),$$

then  $\pi^0 = \rho \cdot J'_\rho$  is an IR of  $H'$ .

**(Step 4)** Take an IR  $\pi^1$  of  $W([\rho])'$  and consider it as a representation of  $H'$  through the homomorphism  $H' \rightarrow W([\rho])' \cong H'/U$ , and consider inner tensor product

$$\pi := \pi^0 \square \pi^1$$

as a representation of  $H'$ . Let the factor set of  $\pi^1$ , viewed as a spin representation of the base group  $W([\rho])$ , be  $\beta_{v,w}$ , then that of  $\pi$  is  $\alpha_{v,w}\beta_{v,w}$ .

To get an IR of  $G$ , we should pick up  $\pi^1$  with the factor set  $\beta_{v,w} = \alpha_{v,w}^{-1}$ . This is possible by Theorem 2.1. Then  $\pi$  becomes a linear representation of the base group  $H = U \rtimes W([\rho])$ . Thus we obtain a representation of  $G$  by inducing it up as

$$(5) \quad \Pi(\pi^0, \pi^1) := \text{Ind}_H^G \pi = \text{Ind}_H^G(\pi^0 \square \pi^1).$$

**Theorem 3.1.** ([3, Theorem 3.1]) *Let  $G = U \rtimes W$  be as above a semidirect product group. Then the induced representation  $\Pi(\pi^0, \pi^1) = \text{Ind}_H^G(\pi^0 \square \pi^1)$  of  $G$  in (5) is irreducible.*

**3.2. Character of  $\Pi(\pi^0, \pi^1)$ .** Put  $\Pi = \Pi(\pi^0, \pi^1)$ , and let  $\chi_\Pi$  be the trace character of  $\Pi$ . Since  $\Pi = \text{Ind}_H^G \pi$ , it is expressed as

$$\chi_\Pi(g) = \int_{H \setminus G} \chi_\pi(kgk^{-1}) d\nu_{H \setminus G}(\dot{k})$$

where the character  $\chi_\pi$  of  $\pi$  is extended from  $H$  to  $G$  by putting 0 outside  $H$ , and  $\nu_{H \setminus G}$  is the invariant measure on  $H \setminus G$  giving mass 1 to each point, and  $\dot{k} = Hk$ . We can rewrite this as

**Theorem 3.2.** ([3, §3.1]) *The character of*

$$\Pi = \Pi(\pi^0, \pi^1) = \text{Ind}_H^G \pi, \quad \pi = \pi^0 \square \pi^1$$

*is expressed as*

$$\chi_\Pi(g) = |H \setminus G| \int_G \chi_\pi(kgk^{-1}) d\mu_G(k),$$

with the normalized Haar measure  $\mu_G$  on  $G$ , where for  $(u, w) \in H = U \rtimes W([\rho])$ ,

$$\chi_\pi((u, w)) = \chi_{\pi^0}((u, w')) \chi_{\pi^1}(w'),$$

with a preimage  $w' \in W([\rho])'$  of  $w$ , i.e.,  $w = \Phi(w')$ .

**3.3. A complete list of IRs of a semidirect product group.** For  $G = U \rtimes W$ , let

$$\{\rho_i; \text{IR of } U, i \in I_{U,W}\}$$

be a complete set of representatives for the orbit space  $\tilde{U}/W$ , and for each  $i \in I_{U,W}$ , let

$$\{\pi_{i,j}^1; j \in J_i\}$$

be a complete set of representatives of equivalence classes of IRs of  $W([\rho_i])'$  with factor set inverse to that of  $J_{\rho_i}$ . Put

$$\begin{aligned} H_i &:= U \rtimes W([\rho_i]), \quad H_i' := U \rtimes W([\rho_i])', \\ \pi_{i,j} &:= \pi_i^0 \square \pi_{i,j}^1, \\ \Pi_{i,j} &:= \Pi(\pi_i^0, \pi_{i,j}^1) = \text{Ind}_{H_i}^G \pi_{i,j}. \end{aligned}$$

Define a set of IRs of  $G$  as

$$\Omega(G) := \{\Pi_{i,j} = \Pi(\pi_i^0, \pi_{i,j}^1); i \in I_{U,W}, j \in J_i\}.$$

**Theorem 3.3.** ([3, Theorems 3.3 and 4.1]) *The above set  $\Omega(G)$  of IRs of  $G$  gives a complete set of representatives of the dual  $\widehat{G}$  of  $G$ , that is, it consists of mutually inequivalent IRs, and covers all equivalence classes of IRs exactly once.*

For the proof of this result, the character theory plays a decisive roll.

**4. Structures of  $G = G_{20}$  and representation group  $R(G)$ .** Let  $G = G_{20}$  be a non-abelian group of order 18, presented by the pair

**Set of generators:**  $\{a, b, c\}$ ;

**Set of fundamental relations:**

$$(6) \quad \begin{cases} a^2 = b^2 = c^2 = 1, \\ (abc)^2 = 1, \quad (ab)^3 = (ac)^3 = 1. \end{cases}$$

Its Schur multiplier is  $M(G_{20}) = \mathbb{Z}_3$ . The subgroup  $A = \langle a \rangle$  acts on  $G$  as  $\varphi(a)g := aga^{-1}$  ( $g \in G$ ).

We know by [6] that  $G$  is isomorphic to the group  $G_{18}^4$  with GAPIdentity [18,4].

**4.1. Another presentation of  $G = G_{18}^4$ .** We looked for another presentation of this group, convenient to our method of constructing *efficient one-step central extension*, and we found a good presentation as follows:

**Set of generators:**  $\{x_1 = ab, x_2 = ca, a\}$ ;

**Set of fundamental relations:**

$$(7) \quad \begin{cases} a^2 = 1, \quad x_1^3 = x_2^3 = 1, \\ x_1 x_2 = x_2 x_1, \quad \varphi(a)x_i = x_i^{-1} \quad (i = 1, 2). \end{cases}$$

**Lemma 4.1.** *Structure of the group  $G = G_{20} \cong G_{18}^4$  is given as follows: put  $X_i := \langle x_i \rangle$  ( $i = 1, 2$ ), and denote by  $C_k$  the cyclic group of order  $k$ , then,*

$$(8) \quad G_{20} = (X_1 \times X_2) \rtimes_{\varphi} A \cong (C_3 \times C_3) \rtimes_{\varphi} C_2.$$

**4.2. Representation group  $R(G_{18}^4)$ .** Taking commuting pair  $x_1$  and  $x_2$ , we construct a one-step efficient central extension  $H \xrightarrow{\delta} G$  coming from commuting relation  $[x_1, x_2] = 1$ . Let  $\xi_1, \xi_2, z_{12}$  and  $w$  be elements of  $H$  covering  $G$  as

$$\xi_i \xrightarrow{\delta} x_i \quad (i = 1, 2), \quad w \xrightarrow{\delta} a, \quad [\xi_1, \xi_2] = z_{12} \xrightarrow{\delta} [x_1, x_2] = 1.$$

The set of all elements of  $H$  is

$$h = z_{12}^{\beta} \xi_1^{\gamma_1} \xi_2^{\gamma_2} w^{\sigma},$$

$$0 \leq \beta, \gamma_1, \gamma_2 \leq 2 \pmod{3}, \quad 0 \leq \sigma \leq 1 \pmod{2}.$$

The set of fundamental relations is

$$(9) \quad \begin{cases} \xi_1^3 = \xi_2^3 = 1, \quad z_{12}^3 = 1, \quad w^2 = 1, \\ z_{12} \text{ central}, \quad \xi_2 \xi_1 = z_{12}^{-1} \xi_1 \xi_2, \\ \varphi(w)\xi_i = \xi_i^{-1} \quad (i = 1, 2). \end{cases}$$

**Theorem 4.2.** *The cyclic subgroup  $Z_{12} := \langle z_{12} \rangle$  is the center  $Z(H)$  of  $H$ , and isomorphic to Schur multiplier  $M(G)$  of  $G = G_{20} \cong G_{18}^4$ . The group  $H$  is a representation group, denoted by  $R(G)$ .*

*Structure of  $R(G)$  is given with  $\Xi_i := \langle \xi_i \rangle$  ( $i = 1, 2$ ) as follows and its GAPIdentity is [54,8]:*

$$R(G) = [(\Xi_1 \times Z_{12}) \rtimes \Xi_2] \rtimes W \cong [(C_3 \times C_3) \rtimes C_3] \rtimes C_2.$$

**Corollary 4.3.** *Spin type of an irreducible projective (or spin) representation of  $G \cong G_{18}^4$  is given as follows: with  $\omega := \exp(2\pi i/3)$ ,  $i = \sqrt{-1}$ ,*

$$\chi_{\varepsilon}(z_{12}) = \omega^{\varepsilon}, \quad \varepsilon = 0, \pm 1 \quad (z_{12} \in Z_{12} \cong M(G)).$$

**Remark 4.4.** In the list in [6], it is noted that the group  $G_{54}^5$ , with GAPIdentity [54, 5], has also the same structure  $((C_3 \times C_3) \rtimes_{\varphi} C_3) \rtimes_{\varphi} C_2$ . Later in the last section, we give more detailed data on this point, and show the dual (the set of equivalence classes of IRs) of  $G_{54}^5$ .

**5. Irreducible representations (=IRs) of  $G$  itself.** Put  $U := X_1 \times X_2$  and  $W := A$ , then  $G$  is semidirect product as  $G = U \rtimes W$ . Since  $U$  is abelian, we can apply Mackey's theory here. First, the dual  $\tilde{U}$  consists of linear characters  $\rho_{m_1, m_2}$  as follows:

for  $(m_1, m_2)$ ,  $m_1, m_2 \in \{-1, 0, 1\} \pmod{3}$ ,

$$(10) \quad \rho_{m_1, m_2}(x_k) = \omega^{m_k} \quad (k = 1, 2).$$

The subgroup  $W = \langle w \rangle$ ,  $w := a$ , acts on  $\widehat{U}$  as

$$(11) \quad {}^w \rho_{m_1, m_2} \cong \rho_{-m_1, -m_2},$$

and  $\widehat{U}/W$  has two kinds of orbits with parameters

(1\*) one-point orbit:  $m_1 = m_2 = 0$ ;

(2\*) two-points orbits:  $m_1 \neq 0$  or  $m_2 \neq 0$ .

Correspondingly the stationary subgroups are

$$W([\rho_{0,0}]) = W \text{ and } W([\rho_{m_1, m_2}]) = \{1\}.$$

**Case (1)**  $\Pi_{0,0;k} := \mathbf{1}_U \cdot (\text{sgn}_W)^k$  ( $k = 0, 1$ ),

where  $\mathbf{1}_U$  is the trivial representation of  $U$ , and  $\text{sgn}_W$  is the sign character of  $W \cong \{\pm 1\}$ .

**Case (2)** Take a representative  $\rho = \rho_{m_1, m_2}$  and its stationary subgroup  $W([\rho]) = \{1\}$ . Then,

$$\Pi_{m_1, m_2} := \text{Ind}_{U \rtimes W([\rho])}^G \rho_{m_1, m_2}$$

is an IR looked for. This is realised by using a section of  $S$  for  $U \backslash G \cong S \subset G$  as follows.

Space  $V(\Pi)$ ,  $\Pi = \Pi_{m_1, m_2}$ , consists of  $V(\rho)$ -valued function  $\mathbf{f}(s)$ ,  $s \in S$ , and operation by  $g_0 \in G$  is given by using decomposition as

$$\begin{aligned} s g_0 &= u(s, g_0) s \overline{g_0}, \quad u(s, g_0) \in U, \quad s \overline{g_0} \in S, \\ (\Pi(g_0) \mathbf{f})(s) &:= \rho(u(s, g_0))(\mathbf{f}(s \overline{g_0})), \end{aligned}$$

where  $s \rightarrow s \overline{g_0}$  expresses the  $G$ -action on the homogeneous space  $U \backslash G$ .

**5.1. Matrix realization of  $\Pi_{m_1, m_2}$ .** Now take  $S = W$ . Then, with transposed row vectors,

$$V(\Pi) \ni \mathbf{f} \rightarrow {}^t(\mathbf{f}(1), \mathbf{f}(w)) \in \mathbf{C}^2 =: V_2$$

is an isomorphism. On the space  $V_2$ , the operators  $\Pi(u)$ ,  $u \in U$ , and  $\Pi(w)$  are expressed by matrices as

$$\begin{aligned} \Pi_{m_1, m_2}(u) &= \begin{bmatrix} \rho_{m_1, m_2}(u) & 0 \\ 0 & \rho_{-m_1, -m_2}(u) \end{bmatrix}, \\ \Pi_{m_1, m_2}(w) &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}. \end{aligned}$$

**Theorem 5.1.** *A complete list of representatives of the dual of  $G \cong G_{18}^4$  is given as follows:*

(1\*) *two of one-dimensional IRs,  $\Pi_{0,0;k}$  ( $k = 0, 1$ );*

(2\*) *four of two-dimensional IRs,  $\Pi_{m_1, m_2}$  with  $(m_1, m_2) \in \{(1, 1), (1, 0), (1, -1), (0, 1)\}$ .*

**5.2. Characters of IRs  $\Pi = \Pi_{m_1, m_2}$ .** We list up trace characters of 2-dimensional IRs  $\Pi = \Pi_{m_1, m_2}$ . The elements of  $G$  is expressed as

$$(12) \quad g = x_1^{\beta_1} x_2^{\beta_2} w^\sigma \quad (0 \leq \beta_1, \beta_2 \leq 2, \sigma = 0, 1).$$

**Theorem 5.2.** *The character  $\chi_\Pi(g)$  is given as*

$$\chi_\Pi(x_1^{\beta_1} x_2^{\beta_2} w^\sigma) = \begin{cases} \omega^{\beta_1 m_1 + \beta_2 m_2} & \\ + \omega^{-\beta_1 m_1 - \beta_2 m_2} & (\sigma = 0), \\ 0 & (\sigma = 1). \end{cases}$$

**6. Construction of spin IRs of  $G = G_{18}^4$ .** Take the representation group  $H := R(G)$  of  $G$ , and utilize the canonical isomorphism  $M(G) \cong Z(H) \cong Z_{12}$ . The set of elements of  $H$  is

$$z_{12}^\alpha \xi_1^{\beta_1} \xi_2^{\beta_2} w^\sigma \quad (0 \leq \alpha, \beta_1, \beta_2 \leq 2, \sigma = 0, 1).$$

Consider spin IRs  $\Pi$  with non-trivial spin type

$$(13) \quad \chi_\varepsilon(z_{12}) = \omega^\varepsilon \text{ with } \varepsilon = \pm 1.$$

To construct  $\Pi$ , we apply the method explained in §3 in four steps and realize it as  $\Pi(\pi^0, \pi^1)$  in (5).

**6.1. Subgroup  $H_0$  of  $H$  and their IRs.** To begin with, first put  $U_0 = Z_{12} \times \Xi_1$  and take a semidirect product subgroup

$$H_0 = U_0 \rtimes \Xi_2 \subset H = R(G).$$

Then,  $H = H_0 \rtimes W$  and we consider two step semidirect product operation to arrive until  $H = R(G)$ .

First, to construct IRs of  $H_0$ , we apply Mackey's method. The action of  $\xi_2 \in \Xi_2$  on  $z_{12}, \xi_1 \in U_0$  is given as

$$\varphi(\xi_2) z_{12} = z_{12}, \quad \varphi(\xi_2) \xi_1 = z_{12}^{-1} \xi_1.$$

The dual of  $U_0$  consists of one-dimensional characters given as follows: for  $(\varepsilon, m_1)$ ,  $\varepsilon, m_1 \in \{-1, 0, 1\}$ ,

$$(14) \quad \rho_{\varepsilon, m_1}(z_{12}) = \omega^\varepsilon, \quad \rho_{\varepsilon, m_1}(\xi_1) = \omega^{m_1}.$$

Actions of  $\xi_2$  on them are

$$\begin{aligned} \varphi(\xi_2)^{-1} z_{12} &= z_{12}, \quad \varphi(\xi_2)^{-1} \xi_1 = z_{12} \xi_1, \\ (\xi_2 \rho_{\varepsilon, m_1})(\xi_1) &= \omega^{\varepsilon + m_1} = \rho_{\varepsilon, \varepsilon + m_1}(\xi_1), \\ (\xi_2 \rho_{\varepsilon, m_1})(z_{12}) &= \omega^\varepsilon = \rho_{\varepsilon, \varepsilon + m_1}(z_{12}). \end{aligned}$$

**Lemma 6.1.** *The orbit space  $\widehat{U}_0/\Xi_2$  consists of two kinds of orbits as*

(3\*) *three of one point orbits  $\{\rho_{0, m_1}\}$  with  $\varepsilon = 0$ ;*

(4\*) *two of three points orbits with  $\varepsilon = \pm 1$ :*

$$\mathcal{O}(\varepsilon) := \{\rho_{\varepsilon, m_1}; m_1 = -1, 0, 1\}.$$

Orbits in (3\*) relate with non-spin IRs of  $G$ , since they will produce representations with trivial spin type  $\chi_0(z_{12}) = 1$ . So we pick up orbits in (4\*).

**6.2. Induced representation from  $U_0 \rtimes \Xi_2([\rho_{\varepsilon, 0}])$  to  $H_0$ .** Take  $\rho_{\varepsilon, 0}$  as a representative of orbits  $\mathcal{O}(\varepsilon)$ , then as is seen from the above, the stationary subgroup  $\Xi_2([\rho_{\varepsilon, 0}])$  in  $\Xi_2$  is trivial, i.e.,  $\Xi_2([\rho_{\varepsilon, 0}]) = \{1\} \subset \Xi_2$ . Therefore, to get IRs of the

group  $H_0 = U_0 \rtimes \Xi_2$ , we should take induced representation from  $U_0 \rtimes \{1\}$  up to  $H_0 = U_0 \rtimes \Xi_2$  as

$$P_\varepsilon := \text{Ind}_{U_0 \rtimes \{1\}}^{U_0 \rtimes \Xi_2} (\rho_{\varepsilon,0} \rtimes \mathbf{1}_{\{1\}}) \text{ for } H_0 = U_0 \rtimes \Xi_2.$$

We realize this induced representation on the space of functions  $f(h)$  of  $h \in H_0$  satisfying

$$(15) \quad f(uh) = \rho_{\varepsilon,0}(u)f(h) \quad (u \in U_0, h \in H_0),$$

with right translation by  $h_0 \in H_0$  given as

$$f(h) \xrightarrow{h_0} f(hh_0).$$

Take a section  $S \subset H_0$  for the quotient map  $H_0 \rightarrow U_0 \backslash H_0$ , and restriction map  $R : f \rightarrow (f(s))_{s \in S}$  onto the space  $\mathcal{F}(S)$  of functions on  $S$ . Then  $P_\varepsilon$  is expressed as follows: for  $h_0 \in H_0$ ,

$$Q_\varepsilon(h_0) := R \cdot P_\varepsilon(h_0) \cdot R^{-1} : (f(s))_{s \in S} \rightarrow (f(sh_0))_{s \in S}.$$

Choose  $S = \Xi_2$  and express  ${}^t(f(\xi_2^{-1}), f(1), f(\xi_2))$  as a row vector (by transposition), then the matrix expression of  $Q_\varepsilon$  is given as follows.

**Lemma 6.2.** *IRs  $Q_\varepsilon$  is given by the following formula: with  $I_3$  the identity matrix of order 3,*

$$Q_\varepsilon(z_{12}) = \omega^\varepsilon I_3, \quad Q_\varepsilon(\xi_2) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix},$$

$$Q_\varepsilon(\xi_1) = \begin{bmatrix} \omega^\varepsilon & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \omega^{-\varepsilon} \end{bmatrix}.$$

*Proof.* For  $h_0 = \xi_2$ , we have transformation

$${}^t(f(\xi_2^{-1}), f(1), f(\xi_2)) \rightarrow {}^t(f(1), f(\xi_2), f(\xi_2^{-1})),$$

and similarly for  $h_0 = \xi_1$ .  $\square$

**6.3. From  $H_0$  to  $H = H_0 \rtimes W = R(G)$ .** Now we apply classical method explained in §3, and follow from Step 1 to Step 4 in §3.1.

For Step 1, we study action of  $w \in W$  to spin IR  $\rho = Q_\varepsilon$  of  $H_0$ . Since

$$\varphi(w)z_{12} = z_{12}, \quad \varphi(w)\xi_j = \xi_j^{-1} \quad (j = 1, 2),$$

we have  ${}^w\rho(z_{12}) = \rho(z_{12})$  and

$${}^w\rho(\xi_j) = \rho(\varphi(w)^{-1}\xi_j) = \rho(\xi_j^{-1})$$

and we see that  ${}^wQ_\varepsilon$  is equivalent to  $Q_\varepsilon$  and the stationary subgroup is  $W([\rho]) = W$ .

For Step 2, an intertwining operator

$$J_\rho(w) : \delta_{\xi_2^q} \rightarrow \delta_{\xi_2^{-q}} \quad \text{or} \quad J_\rho(w) = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

between them is given as

$$\rho(w(h_0)) = J_\rho(w) \rho(h_0) J_\rho(w)^{-1} \quad (h_0 \in H_0).$$

For Step 3, put for  $(h_0, w^\sigma) \in H_0 \rtimes W = H$ ,

$$\pi^0((h_0, w^\sigma)) := \rho(h_0) \cdot J_\rho(w)^\sigma \quad (\sigma = 0, 1),$$

then  $\pi^0 = \rho \cdot J_\rho$ , with  $\rho = Q_\varepsilon$ , is an IR of  $H$ .

For Step 4, take an IR

$$\pi^1 = (\text{sgn}_W)^k \quad (k = 0, 1)$$

of  $W([\rho]) = W$  and consider it as a representation of  $H$  through the homomorphism  $H \rightarrow H/H_0 \cong W([\rho]) = W$ , and consider inner tensor product

$$\pi := \pi^0 \boxtimes \pi^1$$

as a representation of  $H$ . Since the semidirect product  $H_0 \rtimes W([\rho])$  is just equal to  $H = H_0 \rtimes W$ , the inducing process  $\text{Ind}_{H_0 \rtimes W([\rho])}^{H_0 \rtimes W} \pi$  for  $\pi$  is trivial, and so we get in this special case

$$(16) \quad \Pi(\pi^0, \pi^1) = \pi^0 \boxtimes \pi^1 =: R_{\varepsilon;k}.$$

**Proposition 6.3.** *Representations  $R_{\varepsilon;k}$ ,  $k = 0, 1$ , of  $R(G)$ , with  $G = G_{18}^4$ , is irreducible and of spin type  $\chi_\varepsilon$ . Their representation matrices are given as follows:*

$$(17) \quad \begin{cases} R_{\varepsilon;k}(h_0) &= Q_\varepsilon(h_0) \quad (h_0 \in H_0) \\ R_{\varepsilon;k}(w) &= (-1)^k J_\rho(w). \end{cases}$$

By Theorems 3.1 and 3.2, we obtain

**Theorem 6.4.** *IRs of spin type  $\chi_\varepsilon$ , for  $\varepsilon = 1$  or  $\varepsilon = -1$ , of the group  $G = G_{18}^4$  is equivalent to one of  $R_{\varepsilon;k}$ ,  $k = 0, 1$ .*

**6.4. Characters of IRs of  $R(G_{18}^4)$ .** All spin IRs  $\Pi$  of spin type  $\chi_\varepsilon$ ,  $\varepsilon = \pm 1$ , are 3-dimensional. Therefore their trace characters can be calculated by explicit representation matrices  $\Pi(h)$ ,  $h \in H$ , and we obtain the following formula.

All elements of  $H = R(G_{18}^4)$  are expressed as

$$h = z_{12}^\alpha \xi_1^{\beta_1} \xi_2^{\beta_2} w^\sigma \quad (0 \leq \alpha, \beta_1, \beta_2 \leq 2, \quad \sigma = 0, 1),$$

and the matrix expression of  $R_{\varepsilon;k}(h)$  is

$$\omega^{\alpha\varepsilon} (-1)^{\sigma k} \begin{bmatrix} \omega^\varepsilon & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \omega^{-\varepsilon} \end{bmatrix}^{\beta_1} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}^{\beta_2} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}^\sigma.$$

**Theorem 6.5.** *Character  $\chi_\Pi$  for spin IR  $\Pi = R_{\varepsilon;k}$  of spin type  $\chi_\varepsilon$  is given for  $h = z_{12}^\alpha \xi_1^{\beta_1} \xi_2^{\beta_2} w^\sigma$  as follows: In case  $\sigma = 0$ ,*

$$\chi_{\Pi}(h) = \begin{cases} 3\omega^{\alpha\varepsilon} & \text{if } \beta_1 = \beta_2 = 0, \\ 0 & \text{otherwise;} \end{cases}$$

$$\text{in case } \sigma = 1, \\ \chi_{\Pi}(h) = \begin{cases} \omega^{\alpha\varepsilon}(-1)^k & \text{if } \beta_2 = 0, \\ \omega^{\alpha\varepsilon}(-1)^k\omega^{-\beta_1\varepsilon} & \text{if } \beta_2 = 1, \\ \omega^{\alpha\varepsilon}(-1)^k\omega^{\beta_1\varepsilon} & \text{if } \beta_2 = 2. \end{cases}$$

**7. Normalized form of  $\mathbf{A} = \Pi(a), \mathbf{B} = \Pi(b), \mathbf{C} = \Pi(c)$ .** The group  $G = G_{18}^4$  is defined firstly with set of generators  $\{a, b, c\}$  and set of fundamental relations (6) in §4. Then we introduced a new set of generators as  $\{x_1 = ab, x_2 = ca, a\}$ . Now, for a spin IR  $\Pi$  of  $G$ , put  $\mathbf{A} = \Pi(a), \mathbf{B} = \Pi(b), \mathbf{C} = \Pi(c)$ . In the previous paper [1, Theorem 4.1], it is proved that, under certain isomorphisms, the set of these matrices can be normalized as follows:

$$\mathbf{A}^2 = \mathbf{B}^2 = \mathbf{C}^2 = I, (\mathbf{ABC})^2 = \tau I, \\ (\mathbf{AB})^3 = (\mathbf{AC})^3 = I,$$

where  $\tau, \tau^3 = 1$ , is a multiplicative constant corresponding to the spin type of  $\Pi$  (hence  $M(G) \cong C_3$ ).

Here we list up these matrices  $\mathbf{A}, \mathbf{B}, \mathbf{C}$ , in this order, for every purely-spin IRs  $\Pi$  of  $G$ , calculating

$$\mathbf{A} = \Pi(w), \mathbf{B} = \Pi(w^{-1}\xi_1), \mathbf{C} = \Pi(\xi_2w^{-1}).$$

For 1-dim. IR  $\Pi_{0,0;k}$ :  $\tau = 1$ ,

$$\mathbf{A} = (-1)^k, \mathbf{B} = (-1)^k, \mathbf{C} = (-1)^k;$$

For 2-dim. IR  $\Pi_{m_1, m_2}$ :  $\tau = 1$ ,

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & \omega^{-m_1} \\ \omega^{m_1} & 0 \end{bmatrix}, \begin{bmatrix} 0 & \omega^{m_2} \\ \omega^{-m_2} & 0 \end{bmatrix};$$

For 3-dim. IRs  $R_{\varepsilon;k}$ :  $\tau = \omega^{\varepsilon}$ ,  $\varepsilon = 1$  or  $-1$ ,

$$(-1)^k \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, (-1)^k \begin{bmatrix} 0 & 0 & \omega^{-\varepsilon} \\ 0 & 1 & 0 \\ \omega^{\varepsilon} & 0 & 0 \end{bmatrix}, (-1)^k \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Out of explicit matrix calculations, we see that

$$\mathbf{ABC} = \Pi(w)\Pi(w^{-1}\xi_1)\Pi(\xi_2w^{-1}) = \Pi(\xi_1\xi_2w^{-1}), \\ \therefore (\mathbf{ABC})^2 = \Pi(\xi_1\xi_2w^{-1}\xi_1\xi_2w^{-1}) \\ = \Pi(\xi_1\xi_2)\Pi(w^{-1}\xi_1\xi_2w) = \Pi(\xi_1\xi_2\xi_1^{-1}\xi_2^{-1}) \\ = \Pi(z_{12}) = \omega^{\varepsilon}I_3.$$

**8. Study on the dual of the group  $G_{54}^5$ .** In the list in [6], it is noted that the structure of the group  $G_{54}^5$ , with GAPIdentity [54, 5], is expressed as  $((C_3 \times C_3) \rtimes_{\varphi} C_3) \rtimes_{\varphi} C_2$ , the same as for the group  $G_{54}^8$ . So, it would be better or necessary to study group structure of the former and its dual (the set of equivalence classes of IRs), to compare with the dual of the latter. Rewriting data from GAP system we have the following presentation (cf. [5, p.4]):

**Set of generators:**  $\{h_1, h_2, h_3, h_4\}$ ;

**Set of fundamental relations:**

$$\begin{cases} h_1^2 = 1, h_2^3 = h_3^3 = h_4^3 = 1, \\ \varphi(h_1)h_2 = h_2, \varphi(h_2)h_4 = h_4, \varphi(h_3)h_4 = h_4, \\ \varphi(h_1)h_3 = h_3^{-1}, \varphi(h_1)h_4 = h_4^{-1}, \varphi(h_2)h_3 = h_3h_4, \end{cases}$$

where  $\varphi(h)x = h_xh^{-1}$ . Denote by  $H_j := \langle h_j \rangle$  ( $1 \leq j \leq 4$ ) the cyclic group generated by  $h_j$ , and

$$W := H_1 \times H_2 \cong C_2 \times C_3, U := H_3 \times H_4 \cong C_3 \times C_3,$$

then  $U$  is an abelian normal subgroup and  $W$  acts on  $U$  as shown in the fundamental relations. So the structure of  $G_{54}^5$  is described more clearly as

$$G_{54}^5 \cong (C_3 \times C_3) \rtimes (C_2 \times C_3).$$

To construct a complete set of representatives (IRs) of the dual of this group, we can also apply Mackey's induced representations, since  $U$  is abelian.

## References

- [1] T. Hirai, I. Mikami, T. Tsurii, and S. Yamanaka, Projective (or spin) representations of finite groups. I, arXiv:2407.08116.
- [2] S. Yamanaka, T. Tsurii, I. Mikami, and T. Hirai, —. III, arXiv:2408.14727.
- [3] T. Hirai, Classical method of constructing a complete set of irreducible representations of semidirect of a compact group with a finite group, Prob. and Math. Statist., **33**(2013), pp.353-362.
- [4] J. Schur, Über die Darstellung der endlichen Gruppen durch gebrochene lineare Substitutionen, J. für die reine und angew. Math., **127**(1904), 20–50.
- [5] T. Tsurii, On the structures of groups of orders 54 and 81, Printed Note, 2023/6/18-19, in [https://researchmap.jp/7000028076/published\\_works](https://researchmap.jp/7000028076/published_works)
- [6] A. Wingerter and K. Parattu, Tribimaximal Mixing from small groups, additional material. <http://lpsc.in2p3.fr/theorie/akin/documents/listof100smallgroups.pdf>

## e-mail addresses:

Tatsuya Tsurii: t3tsuri23@rsch.tuis.ac.jp  
 Satoe Yamanaka: yamanaka@libe.nara-k.ac.jp  
 Itsumi Mikami: kojirou@kcn.ne.jp  
 Takeshi Hirai: hirai.takeshi.24e@st.kyoto-u.ac.jp