# A dichotomy for the Hofer growth of area preserving maps on the sphere via symmetrization

Lev Buhovsky<sup>1</sup>, Ben Feuerstein<sup>1</sup>, Leonid Polterovich<sup>1</sup>, and Egor Shelukhin<sup>2</sup>

Aug 16, 2024

#### Abstract

We prove that autonomous Hamiltonian flows on the two-sphere  $S^2$  exhibit the following dichotomy: the Hofer norm either grows linearly in time or is bounded in time by a universal constant C = 19Area $(S^2)$ . Our approach involves a new technique, Hamiltonian symmetrization. Essentially, we prove that every autonomous Hamiltonian diffeomorphism is conjugate to an element C-close in the Hofer metric to one generated by a function of the height.

# Contents

1	Intr	oduction and main results	2		
<b>2</b>	The	e symmetrization map	4		
	2.1	Symmetrization of flat quasi-Morse functions	5		
	2.2	Symmetrization of continuous functions	8		
	2.3	A combinatorial point of view on the symmetrization	10		
~	-		12		
3	Bounded growth control				
	3.1	Enhanced Dichotomy	12		
	3.2	The uniform control property	13		
	3.3	The Hölder inequality property	19		
	3.4	The Flattening control property	20		
	3.5	Conjugation control and Bounded growth control	22		
	3.6	The proof of the conjugation lemma	23		

<sup>1</sup>School of Mathematical Sciences, Tel Aviv University, Israel.

<sup>2</sup>Département de Mathématiques et de Statistique, Université de Montréal, Canada.

4	Disc	cussion	26
	4.1	Surfaces other than $S^2$	26
	4.2	Quasi-morphisms and autonomous Hamiltonian diffeomorphisms	27

### **1** Introduction and main results

Let  $(M, \omega)$  be a closed connected symplectic manifold, i.e., an even dimensional manifold M equipped with a closed non-degenerate 2-form  $\omega$ . Every smooth function  $F: M \times \mathbb{R} \to \mathbb{R}$  defines a time-dependent vector field  $X_F^t$  which solves a linear equation  $i_{X_F^t}\omega = -dF_t$  for  $F_t: M \to \mathbb{R}$  given by  $F_t(x) = F(x, t)$ . This field defines a flow of diffeomorphisms  $f_t: M \to M$ . We say that the Hamiltonian function F generates the Hamiltonian diffeomorphism  $\phi = f_1$ . Hamiltonian diffeomorphisms form a group denoted by  $\operatorname{Ham}(M, \omega)$ . The interest in Hamiltonian diffeomorphisms is due to the fact that they model motions of classical mechanics on the phase space M, and that they preserve the symplectic form  $\omega$ . In fact, if  $H^1(M, \mathbb{R}) = 0$  (e.g. for  $M = S^2$ ),  $\operatorname{Ham}(M, \omega)$  is simply the identity component of the symplectomorphism group of  $(M, \omega)$ .

A remarkable discovery of modern symplectic topology is that  $\operatorname{Ham}(M, \omega)$  carries a biinvariant Finsler metric with non-degenerate distance function  $d(\phi, \psi)$ . It is called *the Hofer metric* and is defined by  $d(\mathbf{1}, \phi) = \inf \int_0^1 ||F_t|| dt$ , where the infimum is taken over all Hamiltonian functions F generating  $\phi$ . Here  $||H|| = \max_{x \in M} |H(x)|$  is the uniform norm of a function on M. This metric was introduced by Hofer [8], who proved its non-degeneracy for the linear symplectic space (see also [21]). This was extended in [13] for symplectic manifolds whose symplectic form represents a rational cohomology class, and finally the non-degeneracy of dwas established in [10] for general symplectic manifolds. We refer to [14] for more details on Hofer's metric.

Given a one-parameter subgroup  $\{f_t\}$  of Ham generated by a *time-independent* Hamiltonian  $F : M \to \mathbb{R}$ , we note that the function  $t \mapsto d(\mathbf{1}, f_t)$  is subadditive and hence the following limit exists:

$$\rho(F) := \lim_{t \to +\infty} \frac{d(\mathbf{1}, f_t)}{t} \,. \tag{1}$$

If  $\rho(F) > 0$ , we say that  $\{f_t\}$  has *linear growth*. Linear growth takes place for  $C^{\infty}$ -generic functions F, at least on symplectically aspherical symplectic manifolds  $(M, \omega)$ , see [15, Chapter 6.3].

Interestingly enough, in all examples known to us, one-parameter subgroups enjoy the following dichotomy: either  $\rho(F) > 0$  so the growth is linear, or  $d(\mathbf{1}, f_t)$  is a bounded function of t and  $\rho(F) = 0$ . The latter option takes place, for instance, if  $\psi^*F = -F$  for some  $\psi \in \text{Ham}(M, \omega)$ ; such Hamiltonian flows are of interest in dynamical systems and have a special name - reversible systems.

**Question 1.1.** Do there exist one-parameter subgroups of intermediate growth?

Our main result proves the following enhanced dichotomy for  $M = S^2$ , and hence provides the negative answer to Question 1.1.

**Theorem 1.2** (Enhanced dichotomy). Consider the two dimensional sphere  $S^2$  equipped with a smooth area form. Then for every  $F \in C^{\infty}(M)$  with  $\rho(F) = 0$  and all t in  $\mathbb{R}$  we have

$$d(\mathbf{1}, f_t) \leqslant 19Area(S^2) . \tag{2}$$

A few remarks are in order. The fact that the upper bound in (2) does not depend on the choice of F is unexpected to us. The constant 19 is clearly not optimal - it would be interesting to explore this issue. We expect that our method will yield the (not necessarily enhanced) dichotomy for all surfaces, either closed or open. In the open case one deals with compactly supported Hamiltonians only. Let us mention that in the case of *open surfaces of infinite area* the dichotomy was established in [18] by different tools.

In order to describe our approach to the enhanced dichotomy, let's look at  $\operatorname{Ham}(S^2)$ as at an infinite-dimensional Lie group. Let us mention that this viewpoint highlights the following genuinely infinite-dimensional feature: this group is simple by a result due to Banyaga, it carries a bi-invariant Finsler metric, and it is by no means compact. For finitedimensional Lie groups, this is impossible due to (a minor modification of) a result by Milnor, see [15, Section 1.3.3] for a detailed discussion. Let  $z: S^2 \to \mathbb{R}$  be the height function with max z = 1/2, min z = -1/2. Denote by  $\mathcal{T} \subset \operatorname{Ham}(S^2)$  the group of diffeomorphisms generated by Hamiltonians of the form u(z), and by  $\mathcal{T}_{ev}$  the subgroup of  $\mathcal{T}$  generated by even functions, i.e., u(z) = u(-z). It is instructive (see [1, Proposition (2.2)] to think of  $\mathcal{T}$  as the maximal torus of  $\operatorname{Ham}(S^2)$ , and of  $\mathcal{T}_{ev}$  as its subgroup of invariant elements with respect to the action of the Weyl group (as defined in [3, p. 158]). (In our particular situation, the Weyl group is isomorphic to  $\mathbb{Z}/2\mathbb{Z}$ .)

Next, denote by  $(\widehat{\operatorname{Ham}}(S^2), \widehat{d})$  the completion of  $\operatorname{Ham}(S^2)$  with respect to Hofer's metric d, and write  $\widehat{\mathcal{T}_{ev}}$  for the closure of  $\mathcal{T}_{ev}$  in this completion. We shall prove the following result.

#### Theorem 1.3.

(i) For every non-trivial one-parameter subgroup  $\{f_t\}$  in Ham $(S^2)$  there exists unique oneparameter subgroup  $\{g_t\}$  in  $\widehat{\mathcal{T}_{ev}}$  and a family of elements  $\{\psi_t\}$  in Ham $(S^2)$  such that

$$\widehat{d}(\psi_t f_t \psi_t^{-1}, g_t) \leqslant 19 \cdot Area(S^2).$$
(3)

(ii) The subgroup  $\{g_t\}$  is either trivial (i.e.,  $g_t = 1$  for all t) or it has linear growth:

$$\lim_{t \to +\infty} \frac{\hat{d}(\mathbf{1}, g_t)}{t} > 0 \; .$$

Recall that in a compact simple finite-dimensional Lie group every element is conjugate to an element from a given maximal torus, see [3, p. 159]. Theorem 1.3(i) can be considered to be a substitute of this result for autonomous diffeomorphisms in the group  $\operatorname{Ham}(S^2)$ : the conjugation exists up to a bounded error and passing to the completion. Its proof is based on a quite substantial generalization of the Sikorav's trick see [4, 20]. The proof of Theorem 1.3(ii) involves the technique of Lagrangian estimators coming from Lagrangian Floer theory in symmetric products [5, 17], and in particular on quasi-morphisms on  $\widehat{\operatorname{Ham}}(S^2)$  constructed in [5].

The enhanced dichotomy (Theorem 1.2) readily follows from Theorem 1.3.

ORGANIZATION OF THE PAPER: In Section 2 we define the key notion of the paper, Hamiltonian symmetrization, prove a few of its basic properties, and discuss three points of view on it: symmetrization of suitably adapted, so-called "flattened quasi-Morse", Hamiltonian functions (Section 2.1), symmetrization by means of quasi-morphisms which applies to arbitrary continuous function (Section 2.2), and a combinatorial description in terms of the Reeb graph (Section 2.3). In Section 3 we prove Theorem 1.3 by means of establishing a number of more refined properties of the symmetrization which we summarize in Section 3.1: uniform control as  $z \to \pm 1/2$  for sufficiently smooth Hamiltonians (Section 3.2), a Hölder style inequality for comparing symmetrizations of nearby functions (Section 3.3), estimate the effect of flattening a Morse function on the symmetrization (Section 3.4), prove the theorem assuming it holds for flattened quasi-Morse functions (Section 3.5), and finally prove it for these functions (Section 3.6). In Section 4 we outline a generalization of our results to other surfaces and present a further direction of research.

### Acknowledgments

LB was partially supported by the Israel Science Foundation grant 1918/23. BF and LP were partially supported by the ISF-NSFC grant 3231/23. ES was partially supported by the NSERC, Fondation Courtois, and Alfred P. Sloan Foundation.

# 2 The symmetrization map

Our ultimate goal in this section is to define a map

$$\Sigma: C_0^0(S^2) \to C_{\text{even},0}^0(I)$$

for the interval  $I = \left(-\frac{1}{2}, \frac{1}{2}\right)$ , which we call the **symmetrization map**, which enjoys various useful properties. Here  $C_0^0(S^2)$  is the set of mean-zero  $C^0(S^2)$  functions, and  $C_{\text{even},0}^0(I)$  is the set of continuous mean-zero even functions on I.

For instance, we prove the following bounded growth control property, which shows how the symmetrization of an autonomous Hamiltonian  $H \in C_0^2(S^2)$  is related to the Hofer growth of the one-parametric group  $\{\phi_H^t\}_{t\in\mathbb{R}}$ :

$$\rho(H) = 0 \iff \Sigma(H) \equiv 0 \iff \forall t, \quad \widehat{d}(\mathbf{1}, \varphi_H^t) < 19 \operatorname{Area}(S^2).$$

Unfortunately,  $\Sigma$  has the disadvantage of having a somewhat involved definition, and the rest of this section is dedicated to it. In order to be as concise as possible, the definition of  $\Sigma$  is split into two parts, and followed by a quick discussion. In the first part, we define "(flat) quasi-Morse" functions to be functions of Morse functions of the sphere (constant near critical levels), and give a rather intuitive definition of  $\Sigma$  on these functions. In the second part we use a density argument to define the symmetrization for  $C_0^0(S^2)$  functions. This requires rather heavy machinery, notably quasi-morphisms and quasi-states from [5] (in the spirit of [17]). A running motif in these parts is that  $\Sigma$  can be defined for functions of trees, and extended to functions on  $S^2$  using the theory of Reeb Graphs (see [9]). This idea is expanded upon in the last subsection.

#### 2.1 Symmetrization of flat quasi-Morse functions

We will start with some auxiliary definitions and notations.

**Definition 2.1.** A function  $H \in C^{\infty}(S^2)$  is called **quasi-Morse** (QM) if it is the pullback of a function on the Reeb graph of a Morse function: that is, if there exists a Morse function  $f: M \to \mathbb{R}$  and a function

$$H: \Gamma_f \to \mathbb{R}$$

such that

$$H(q) = H(\pi(q))$$

for all  $q \in S^2$ , where  $\Gamma_f$  is the Reeb graph of f and  $\pi : M \to \Gamma_f$  is the natural projection. Furthermore, it is called **flat quasi-Morse** (FQM) if for every vertex v of  $\Gamma_f$  of degree at least 2 there is a neighborhood of  $\pi^{-1}(v)$  on which H is constant.

We will use the following terminology:

- We will say that the FQM function H is subordinate to the Morse function f. Note that such a function f is not unique.
- Denote by  $\operatorname{FQM}_0(M) \subset C^{\infty}(M)$  the set of mean-zero FQM functions on M subordinate to a Morse function with distinct critical values.
- For H Morse, and  $\varepsilon > 0$ , let  $r : \mathbb{R} \to \mathbb{R}$  be a smooth function which is constant near the critical values of H corresponding to saddles which satisfies  $||r - \mathbf{1}_{\mathbb{R}}||_{C^0} < \varepsilon$ . We call the FQM function  $\widetilde{H} = r \circ H$  the  $\varepsilon$ -flattening of H.

**Theorem 2.2.** There exists a map  $\widetilde{\Sigma}$  : FQM<sub>0</sub>( $S^2$ )  $\rightarrow C^{\infty}_{even,0}(I)$  for  $I = \left(-\frac{1}{2}, \frac{1}{2}\right)$  with the following properties:

1. (Homogeneity and symmetry) For all  $H \in FQM_0(S^2)$  and  $t \in \mathbb{R}$  we have

$$\widetilde{\Sigma}(tH) = t\widetilde{\Sigma}(H),$$

and if H(z) is a function of the height, then

$$\widetilde{\Sigma}(H) = \frac{H(z) + H(-z)}{2}$$

2. (FQM Hofer control) Every  $H \in FQM_0(S^2)$  has a constant  $C(H) \in \mathbb{R}_{\geq 0}$  such that

$$\forall t \in \mathbb{R}, \quad d\left(\varphi_{H}^{t}, \varphi_{\widetilde{\Sigma}(H)}^{t}\right) \leqslant C(H).$$

In the sequel we will identify functions H on I and functions  $H \circ z$  on  $S^2$  depending on the height only. Before proceeding with the proof of Theorem 2.2, we note that FQM Hofer control along with the fact that  $\widetilde{\Sigma}(H) \neq 0$  implies  $\rho\left(\widetilde{\Sigma}(H)\right) > 0$  (we will show this implication a bit later) proves the (not yet enhanced) dichotomy theorem for  $FQM_0(S^2)$  functions.

Proof. Let  $H \in \operatorname{FQM}_0(S^2)$  be subordinate to a Morse function f. We will construct  $\widetilde{\Sigma}$  directly as follows. Recall that  $\Gamma_f$  denotes the Reeb graph of f and  $\pi : S^2 \to \Gamma_f$  is the natural projection. Let  $\{A_i\}$  be the edges of  $\Gamma_f$  and  $S_i = \pi^{-1}(A_i) \subset S^2$ . Then the  $\{S_i\}$  are disks/annuli which are invariant under the flow of H, and whose closures form a cover of  $S^2$  with non-empty intersections consisting of figure-eights lying on critical levels of f. For every i, let  $H_i$  be the extension by constants of  $H|_{S_i}$  and  $\varphi_i^t := \varphi_{H_i}^t \in \operatorname{Ham}(S^2)$ . Note that as H is an FQM, all the  $H_i$  are smooth functions. Indeed, there exist disks or annuli  $S'_i$  whose closures are contained in  $S_i$ , such that H is constant on the connected components of  $S^2 \setminus \bigcup_i S'_i$  so the extension by constants of  $H|_{S'_i}$  is smooth and coincides with that of  $H|_{S_i}$ . Moreover,  $\varphi_H^t = \prod_i \varphi_i^t$ . We want to map  $S'_i$  to a standard cap/annulus with a Hamiltonian diffeomorphism  $\theta_i$  such that  $H_i \circ \theta_i^{-1}$  is a function of the height. We also want to split  $H_i \circ \theta_i^{-1}$  into two Hamiltonians, on opposite sides of the sphere so that  $\widetilde{\Sigma}(H)$  is an even function. Consider  $R \in \operatorname{Ham}(S^2)$  given via the embedding  $S^2 \subset \mathbb{R}^3$  as a sphere of radius 1/2, via  $(x, y, z) \mapsto (-x, y, -z)$ .

To construct  $\theta_i$  we use [19, Lemma 2.5], which we recall here along with relevant definitions for the reader's convenience.

**Definition 2.3.** A family  $\mathcal{L}$  of loops in a simply connected domain  $U \subset \mathbb{R}^2$  is called admissible if there is a diffeomorphism  $\beta : D(U) \setminus \{0\} \to U \setminus \{p\}$ , for some  $p \in U$ , where D(U) is an open disk of area Area(U) around the origin, such that:

- 1. Concentric circles are mapped to elements of  $\mathcal{L}$ .
- 2. In a neighborhood of the origin  $\beta$  is a translation.

**Lemma 2.4.** Let U and V be bounded and simply connected domains in  $\mathbb{R}^2$  of equal area and let  $\mathcal{L}_U$  and  $\mathcal{L}_U$  be admissible families of loops in U and V, respectively. Then there is a symplectomorphism between U and V mapping loops to loops.

Our admissible family of loops will be the regular level sets of  $H_i$  extended to a family of smooth loops on a neighborhood of the union of these regular level sets. We will map this family of loops to concentric circles  $\{h = c\}$  in  $W = \{(h, \vartheta) \in S^2 | h \in (c_i, d_i)\} \subset S^2$ . Here  $c_i < d_i$  are given as follows. Let  $\operatorname{Im}(H_i) = [a_i, b_i]$  and recall that  $H_i$  is constant outside  $S'_i$ . Then  $c_i = \operatorname{Area}\{H_i \leq a_i\}$ , and  $d_i = \operatorname{Area}\{H_i \leq b_i\}$ . This choice of W allows us to construct  $\theta_i$  as an area-preserving diffeomorphism on a disk or an annulus, which by [7, Lemmas 2 and 4] extends to a Hamiltonian diffeomorphism  $\theta_i : S^2 \to S^2$ . Now, we are ready to define  $\widetilde{\Sigma}(H)$ .

**Definition 2.5.** Let  $H \in FQM_0(S^2)$  and let  $\{\theta_i\} \subset Ham(S^2)$  be defined as above. The symmetrization of H is defined as

$$\widetilde{\Sigma}(H) = \frac{1}{2} \sum_{i} \left( H_i \circ \theta_i^{-1} + H_i \circ \theta_i^{-1} \circ R \right).$$

It is an even function of the height, which corresponds to a function  $\widetilde{\Sigma}(H): \left[-\frac{1}{2}, \frac{1}{2}\right] \to \mathbb{R}.$ 

This definition does not depend on the specific choices of  $\theta_i$  and satisfies the *Homogeneity and symmetry* property of Theorem 2.2. In order to show the *FQM Hofer control* property of the same theorem, consider the following.

Since all  $H_i \circ \theta_i^{-1}$ ,  $H_i \circ \theta_i^{-1} \circ R$  are functions of the height we can decompose  $\varphi_{\widetilde{\Sigma}(H)}^t$  to the commuting flows

$$\varphi_{\widetilde{\Sigma}(H)}^{t} = \prod_{i} \varphi_{H_{i} \circ \theta_{i}^{-1}}^{\frac{t}{2}} \varphi_{H_{i} \circ \theta_{i}^{-1} \circ R}^{\frac{t}{2}}$$

Now, by inductively using the triangle inequality, together with the bi-invariance of Hofer's metrics, one gets

$$d\left(\varphi_{H}^{t},\varphi_{\widetilde{\Sigma}(H)}^{t}\right) = d\left(\prod_{i}\varphi_{i}^{t},\prod_{i}\varphi_{H_{i}\theta_{i}^{-1}}^{t}\varphi_{H_{i}\theta_{i}^{-1}R}^{t}\right) \leqslant \sum_{i}d\left(\varphi_{H_{i}}^{t},\varphi_{H_{i}\theta_{i}^{-1}}^{t}\varphi_{H_{i}\theta_{i}^{-1}R}^{t}\right)$$

Set  $\|\psi\| = d(\mathbf{1}, \psi)$  and  $\phi_i = \varphi_{H_i}^{t/2}$ , and using the well-known fact that  $\|[A, B]\| = \|ABA^{-1}B^{-1}\| \leq C$ 

 $2\min\{||A||, ||B||\},$  we have

$$d\left(\varphi_{H_{i}}^{t},\varphi_{H_{i}\theta_{i}^{-1}}^{\frac{t}{2}}\varphi_{H_{i}\theta_{i}^{-1}R}^{\frac{t}{2}}\right) = d\left(\left(\varphi_{H_{i}}^{\frac{t}{2}}\right)^{2},\theta_{i}\varphi_{H_{i}}^{\frac{t}{2}}\theta_{i}^{-1}R^{-1}\theta_{i}\varphi_{H_{i}}^{\frac{t}{2}}\theta_{i}^{-1}R\right)$$
$$= d\left(\varphi_{i}^{2},\theta_{i}\phi_{i}\theta_{i}^{-1}R^{-1}\theta_{i}\phi_{i}\theta_{i}^{-1}R\right)$$
$$\leqslant d\left(\phi_{i},\theta_{i}\phi_{i}\theta_{i}^{-1}\right) + d\left(\phi_{i},R^{-1}\theta_{i}\phi_{i}\theta_{i}^{-1}R\right)$$
$$= \left\|\phi_{i}^{-1}\theta_{i}\phi_{i}\theta_{i}^{-1}\right\| + \underbrace{\left\|R^{-1}\theta_{i}\phi_{i}\theta_{i}^{-1}R\phi_{i}^{-1}\right\|}_{\left\|\left[R^{-1}\theta_{i},\phi_{i}\right]\right\|}$$
$$\leqslant 2\|\theta_{i}\| + 2\left\|R^{-1}\theta_{i}\right\| \leqslant 4\|\theta_{i}\| + 2\|R\|.$$

Thus, for all  $t \in \mathbb{R}$ ,

$$\left| d(\mathbf{1}, \varphi_H^t) - d\left(\mathbf{1}, \varphi_{\widetilde{\Sigma}(H)}^t\right) \right| \leq d\left(\varphi_H^t, \varphi_{\widetilde{\Sigma}(H)}^t\right) \leq \sum_i \left(4\|\theta_i\| + 2\|R\|\right) =: C(H).$$

#### 2.2 Symmetrization of continuous functions

We now want to define  $\Sigma$  as a map  $\Sigma : C_0^0(S^2) \to C_{\text{even},0}^0(I)$ . Clearly  $\text{FQM}_0(S^2)$  is a  $C^0$ -dense subset of  $C_0^0(S^2)$ . Thus a natural way to define  $\Sigma$  would be via continuity. Indeed, set  $I_k = \left[-\frac{1}{2} + \frac{1}{k+1}, \frac{1}{2} - \frac{1}{k+1}\right]$  for  $k \ge 2$ , and let  $r_k : C_{\text{even}}^0(I) \to C_{\text{even}}^0(I_k)$  be the restriction map. We will prove the following result.

**Theorem 2.6.** There exists a unique map  $\Sigma : C_0^0(S^2) \to C_{even,0}^0(I)$  that coincides with  $\widetilde{\Sigma}$  on  $\operatorname{FQM}_0(S^2)$ , such that the map  $r_k \circ \Sigma$  is (k-1) Lipschitz.

The proof of this statement requires the **link quasi-morphisms**  $\mu_{k,B}$ : Ham $(S^2) \to \mathbb{R}$ from [5] (cf. invariants introduced in [17]). We proceed as follows. Fix k > 1 and  $B \in (\frac{1}{k+1}, \frac{1}{2})$  and set  $C = \frac{1-2B}{k-1}$ . Let  $H \in \text{FQM}_0(S^2)$ . Recall the commutative-additivity of  $\mu_{k,B}$  (see [5]) and consider  $\{\theta_i\} \subset \text{Ham}(S^2)$  from the definition of  $\widetilde{\Sigma}$ . Since both  $\{\varphi_{H_i}\}$  and  $\{\theta_i\varphi_{H_i}\theta_i^{-1}\}$  are collections of commuting maps, and as  $\widetilde{\Sigma}(H)$  is a map of the height, we have

$$\mu_{k,B}(\varphi_H) = \mu_{k,B}\left(\prod_i \varphi_{H_i}\right) = \sum_i \mu_{k,B}(\varphi_{H_i}) = \sum_i \mu_{k,B}(\theta_i \varphi_{H_i} \theta_i^{-1})$$
$$= \mu_{k,B}\left(\varphi_{\widetilde{\Sigma}(H)}\right) = \frac{1}{k} \sum_{j=0}^{k-1} \widetilde{\Sigma}(H)\left(-\frac{1}{2} + B + jC\right).$$

Consider the set  $L_{\ell,B} := \left\{-\frac{1}{2} + B + jC\right\}_{j=0}^{\ell-1} \subset I$ , where  $1/(\ell+1) < B < 1/2$ . Then  $L_{k,B}$  and  $L_{k-2,B+C}$  coincide on all but two points  $\pm \left(-\frac{1}{2} + B\right)$ . (Note that indeed 1/(k-1) < 1)

B + C < 1/2.) This gives us the following explicit expression for  $\widetilde{\Sigma}$ :

$$\widetilde{\Sigma}(H)(z) = \frac{k}{2}\mu_{k,B}(H) - \frac{k-2}{2}\mu_{k-2,B+C}(H), \qquad \begin{aligned} |z| &< \frac{1}{2} - \frac{1}{k+1}, \\ B &= \frac{1}{2} - z, \\ C &= \frac{1-2B}{k-1} \end{aligned}$$
(4)

We can now use the Hofer-Lipschitz property of  $\mu_{k,B}$  to deduce our Lipschitz condition. For all  $H, H' \in FQM_0(S^2)$ 

$$\begin{split} |\Sigma(H')(z) - \Sigma(H)(z)| &= \left| \frac{k}{2} \mu_{k,B}(H') - \frac{k-2}{2} \mu_{k-2,B+C}(H') - \frac{k}{2} \mu_{k,B}(H) + \frac{k-2}{2} \mu_{k-2,B+C}(H) \right| \\ &\leqslant \frac{k}{2} |\mu_{k,B}(H') - \mu_{k,B}(H)| + \frac{k-2}{2} |\mu_{k-2,B+C}(H') - \mu_{k-2,B+C}(H)| \\ &\leqslant \left( \frac{k}{2} + \frac{k-2}{2} \right) \|H' - H\|_{C^0} = (k-1)\|H' - H\|_{C^0}. \end{split}$$

This proves the Lipschitz-continuity condition on  $\tilde{\Sigma}$  and finishes the proof of the theorem.

**Remark 2.7.** It should be noted that the explicit formula (4) directly extends to  $\Sigma$ , and inherits the independence of k from  $\tilde{\Sigma}$ . The latter can also be shown directly as follows. For k, B as above, let  $\delta_{k,B} = \frac{1}{k} \sum_{x \in L_{k,B}} \delta_x$ . If  $\delta_i = \delta_{k_i,B_i}$  satisfy  $\sum a_i \delta_i = 0$  for certain  $a_i \in \mathbb{R}$ , then the Lagrangian control property yields that the quasi-morphism  $\mu = \sum a_i \mu_{k_i,B_i}$  vanishes on  $\mathcal{T}$  and  $\hat{\mathcal{T}}$ . This immediately implies the independence of the above formula for  $\Sigma(H)(z)$ on k as long as |z| < 1/2 - 1/(k+1). (We also note that the values k = 1 and B = 1/2 are admissible for this argument.)

The map  $\Sigma : C_0^0(S^2) \to C_{\text{even},0}^0(I)$  inherits the algebraic properties of  $\widetilde{\Sigma}$ , satisfies Lipschitz continuity, and a new property: independence of Hamiltonian, as described below.

Recall that to every Hamiltonian H in  $C_0^0(S^2)$  there corresponds a one-parameter subgroup  $\phi_H = \{\phi_H^t\}$  in  $\widehat{\operatorname{Ham}}(S^2)$ . It is obtained by approximating H by smooth  $H_i$  and considering the limit  $\phi_H^t$  in  $\widehat{\operatorname{Ham}}(M)$  of the Hofer Cauchy sequence  $\phi_{H_i}^t$ . It can be checked that  $\phi_H^t = \phi_{tH}^1$ . We define  $\operatorname{Aut}^0(M)$  as the image of the map  $C_0^0(S^2) \to \widehat{\operatorname{Ham}}(S^2), H \mapsto \phi_H^1$ .

**Remark 2.8.** A related subspace  $\operatorname{Aut}(S^2)$  of  $\operatorname{Ham}(S^2)$  is defined as the closure inside  $\operatorname{Ham}(S^2)$  of  $\operatorname{Aut}(S^2)$ . In other words, it consists of the limits in  $\operatorname{Ham}(S^2)$  of all Hofer Cauchy sequences  $\phi_i$  in  $\operatorname{Aut}(S^2)$ . Clearly  $\operatorname{Aut}^0(S^2) \subset \operatorname{Aut}(S^2)$ . It would be very interesting to study the extent to which this inclusion is strict, as  $\operatorname{Aut}(S^2)$  is known to contain interesting elements: for instance the Anosov-Katok pseudo-rotations.

Recall that for k > 1,  $I_k = \left[-\frac{1}{2} + \frac{1}{k+1}, \frac{1}{2} - \frac{1}{k+1}\right]$ , while  $r_k : C^0_{\text{even}}(I) \to C^0_{\text{even}}(I_k)$  is the restriction map.

**Theorem 2.9.** The map  $\Sigma : C_0^0(S^2) \to C_{even,0}^0(I)$  satisfies the following properties.

- 1. (Independence of Hamiltonian) If  $H, H' \in C_0^0(S^2)$  generate the same elements  $\varphi_H^1 = \varphi_{H'}^1 \in \operatorname{Aut}^0(M)$ , then  $\Sigma(H) = \Sigma(H')$ , i.e.  $\Sigma$  descends to a map from  $\operatorname{Aut}^0(M)$  to  $C_{even,0}^0(I)$ . We denote this map by  $\sigma$ . Note that  $\sigma(\varphi_H^1) = \Sigma(H)$ .
- 2. (Lipschitz continuity) Both symmetrization maps  $\Sigma$  and  $\sigma$  are (k-1) Lipschitz after restriction to  $I_k$ . That is,  $r_k \circ \sigma$  :  $\left(\operatorname{Aut}^0(M), \widehat{d}\right) \to (C^0_{even}(I_k), d_{C_0})$ , and  $r_k \circ \Sigma$  :  $(C^0_0(S^2), d_{C_0}) \to (C^0_{even}(I_k), d_{C_0})$  are (k-1) Lipschitz.
- 3. (Algebraic properties) For all  $H \in C_0^0(S^2)$  and  $t \in \mathbb{R}$

$$\Sigma(tH) = t\Sigma(H).$$

If Hamiltonians F, G have commuting flows, then

$$\Sigma(F) + \Sigma(G) = \Sigma(F + G),$$

and if H(z) is a function of the height, then

$$\Sigma(H) = \frac{H(z) + H(-z)}{2}$$

*Proof.* The only new point to prove is independence of Hamiltonian, which follows on smooth functions by its expression (4) in terms of the link quasi-morphisms, as well as the fact that these quasi-morphisms vanish on  $\pi_1(\text{Ham}(S^2)) \cong \mathbb{Z}/2\mathbb{Z}$ . This extends to continuous functions by Lipschitz continuity of the link quasi-morphisms in Hofer's metric.

**Remark 2.10.** It is not hard to see that the Lipschitz continuity property of Theorem 2.9 implies that  $\sigma$  extends to  $\widehat{Aut}(S^2)$ . It would be very interesting to see if further results in this paper extend to this setting.

#### 2.3 A combinatorial point of view on the symmetrization

We conclude this section by formalizing the observation that while  $\Sigma$  is defined on  $C_0^0(S^2)$ , the underlying definition actually has to do with continuous function on trees.

Given a finite tree  $\Gamma$ , and  $\mu$  a probability measure on  $\Gamma$ , which is Lebesgue on every edge and such that all the vertices have measure 0, we want to define a map

$$\Sigma^{\Gamma}: C_0(\Gamma) \to C_{\text{even},0}(I), \quad h \mapsto \Sigma_h^{\Gamma},$$

where  $C_0(\Gamma)$  is the set of mean-zero (with respect to  $\mu$ ) functions, which is related to  $\Sigma$  as follows: if f is a Morse function on  $S^2$  and  $H = h \circ \pi_{\Gamma_f}$  is an QM function subordinate to fthen  $\Sigma(H) = \Sigma_h^{\Gamma_f}$ . (Recall that  $\pi_{\Gamma_f} : S^2 \to \Gamma_f$  is the natural projection.) Let us start with some notation. Consider  $\Gamma$  as a simplicial complex, denote it's edges by e = [v, w], where v, w are vertices. For an edge e we denote by  $e^{\circ}$  its interior. Given a vertex v of e, denote by  $T_{e,v}$  the connected component of  $\Gamma \setminus e^{\circ}$  containing v.

Define the intervals

$$I_{e,v}^{-} = (-1/2, -1/2 + \mu(T_{e,v})] = -I_{e,w}^{+} ,$$
  

$$I_{e,v} = (-1/2 + \mu(T_{e,v}), -1/2 + \mu(T_{e,v}) + \mu(e)) = -I_{e,w} ,$$
  

$$I_{e,v}^{+} = [-1/2 + \mu(T_{e,v}) + \mu(e), 1/2) = -I_{e,w}^{-} .$$

**Definition 2.11.** A function  $h \in C_0(\Gamma)$  is called elementary with respect to edge e = [v, w]if it is possibly non-constant on an edge e, and is equal to the constant h(v) on  $T_{e,v}$ , and to the constant h(w) on  $T_{e,w}$ .

**Theorem 2.12.** There is a unique  $\mathbb{R}$ -linear map  $\Sigma^{\Gamma} : C_0(\Gamma) \to C_{even,0}(I)$ , such that for elementary h, if we put

$$\Sigma_{h,v}^{\Gamma}(z) = h(v), \quad \text{if } z \in I_{e,v}^{-};$$
  

$$\Sigma_{h,v}^{\Gamma}(z) = h(x), \quad \text{if } z = -1/2 + \mu(T_{e,v}) + \mu([v,x]) \in I_{e,v} \text{ (here } x \in e);$$
  

$$\Sigma_{h,v}^{\Gamma}(z) = h(w), \quad \text{if } z \in I_{e,v}^{+},$$

then

$$\Sigma_{h}^{\Gamma}(z) = \frac{1}{2} \left( \Sigma_{h,v}^{\Gamma}(z) + \Sigma_{h,w}^{\Gamma}(z) \right) \; .$$

*Proof.* We have a couple of things to show. First, since

$$\Sigma_{h,w}^{\Gamma}(-z) = \Sigma_{h,v}^{\Gamma}(z) ,$$

we have that  $\Sigma_h^{\Gamma}(z) = \Sigma_h^{\Gamma}(-z)$ . Second, since the map

$$e \rightarrow I_{e,v}, \ x \mapsto -1/2 + \mu(T_{e,v}) + \mu([v, x])$$

is a homeomorphism sending measure dz to  $\mu$ , one readily checks that

$$\int_{-1/2}^{1/2} \Sigma_{h,v}^{\Gamma}(z) dz = \int_{\Gamma} h d\mu = 0.$$

Lastly, we must show that every  $h \in C_0(\Gamma)$  has a unique decomposition to a sum of elementary functions with respect to each edge. Indeed, denote by  $\tilde{h}_e$  the function which is equal to h on e = [v, w], equal to h(v) on  $T_{e,v}$  and equal to h(w) on  $T_{e,w}$ , denote

$$h_e = \tilde{h}_e - \int \tilde{h}_e d\mu.$$

Note  $a := h - \sum_e h_e$  is constant on each edge, and hence constant on  $\Gamma$ , and has zero mean, hence  $a \equiv 0$ , and the decomposition  $h = \sum_e h_e$  follows, uniqueness is clear.

Note that the fact that  $\Sigma^{\Gamma}$  coincides with  $\Sigma$  in the way discussed in the beginning of this section is evident by direct comparison between the description of  $\Sigma^{\Gamma}$  for elementary h from Theorem 2.12 and Equation (6) in Section 3.2.

# 3 Bounded growth control

#### 3.1 Enhanced Dichotomy

Our ultimate goal in this section is to prove the following "enhanced dichotomy" theorem.

**Theorem 3.1.** Let  $H \in C_0^2(S^2)$ , a mean-zero  $C^2$  function on  $S^2$ . Then:

- 1. If  $\Sigma(H) \neq 0$ , then H has linear growth type, i.e.  $\rho(H) > 0$ .
- 2. If  $\Sigma(H) \equiv 0$ , then  $\widehat{d}(\mathbf{1}, \varphi_H^t) \leq 19 \operatorname{Area}(S^2)$ .

Our proof proceeds by establishing the following useful properties of the symmetrization map  $\Sigma$ . Note that Theorem 3.1 is a direct consequence of the fifth property, *Bounded growth control*.

**Theorem 3.2.** The symmetrization map  $\Sigma$  has the following properties:

1. (Uniform control) There exists  $C \in \mathbb{R}_{>0}$  such that every  $H \in C_0^2(S^2)$  satisfies

$$\|\Sigma(H)\|_{C^0}\leqslant C\|H\|_{C^2}$$

Furthermore, for  $H \in C_0^2(S^2)$ ,  $\Sigma(H)$  extends to a continuous function on  $\left[-\frac{1}{2}, \frac{1}{2}\right]$ .

2. (Hölder's inequality) Every pair  $H, H' \in C_0^2(S^2)$  with  $||H - H'||_{C^0} < 1$  satisfies

$$\widehat{d}(\varphi_{\Sigma(H)}^{1},\varphi_{\Sigma(H')}^{1}) \leqslant C' \sqrt{\|H - H'\|_{C^{0}}} \left(\sqrt{1 + \|H\|_{C^{2}} + \|H'\|_{C^{2}}}\right)$$

for a universal constant C' > 0.

3. (Flattening control) Let H be a Morse function,  $\varepsilon > 0$ , and  $\widetilde{H}$  an  $\varepsilon$ -flattening of H, then

$$\left\|\Sigma(\widetilde{H}) - \Sigma(H)\right\|_{C^0} \leq (1 + 2e(H))\varepsilon$$

where e(H) is the number of edges of the Reeb graph of H (which is the same as the number of critical points of H minus one).

4. (Conjugation control) For every  $H \in C_0^2(S^2)$  there exists  $\psi \in \widehat{Ham}(M)$  such that

$$d(\varphi_H^1, \psi^{-1}\varphi_{\Sigma(H)}^1\psi) \leqslant 19\operatorname{Area}(S^2).$$

5. (Bounded growth control) For every  $H \in C_0^2(M)$ ,

$$\rho(H) = 0 \iff \Sigma(H) \equiv 0 \iff \forall t, \quad \widehat{d}(\mathbf{1}, \varphi_H^t) \leqslant 19 \operatorname{Area}(S^2).$$

#### **3.2** The uniform control property

In this section we will prove the *Uniform control* property of Theorem 3.2, and show that it implies a part of the *Bounded growth control* property of the same theorem.

An important tool in this and subsequent sections is the "improved Sikorav trick" [4, Lemma 2.1] (see also [20]), which we repeat here for the reader's convenience.

**Lemma 3.3.** Let  $(M, \omega)$  be a closed and connected symplectic surface. Let  $\varepsilon > 0$ , let m be a positive integer, let  $\mathcal{D}_0, ..., \mathcal{D}_m \subset M$  be disjoint topological open disks of area  $\varepsilon$  each, and let  $\phi_j : \mathcal{D}_0 \to \mathcal{D}_j$  for all  $1 \leq j \leq m$  be symplectic diffeomorphisms. Consider  $f_0, ..., f_m \in$  $\operatorname{Ham}(M, \omega)$  with supp  $(f_j) \subset \mathcal{D}_j$ . Define  $\Phi, \Phi' \in \operatorname{Ham}(M, \omega)$  by

$$\Phi = f_0 f_1 \cdots f_m$$

and

$$\Phi' = f_0 \prod_{j=1}^m \phi_j^* f_j$$

where  $\phi_j^* f_j$  is given by  $\phi_j^* f_j = (\phi_j)^{-1} f_j \phi_j$  on  $\mathcal{D}_0$  and  $\phi_j^* f_j = \mathbf{1}$  on  $M \setminus \mathcal{D}_0$ . Then

 $d(\Phi, \Phi') < 3\varepsilon.$ 

In this chapter, we will mainly use the following consequence of Lemma 3.3. Let  $F \in C^{\infty}(S^2)$  be supported in a disk  $\mathcal{D}$  of area  $B < \frac{1}{k}$ . Record the following identity  $\varphi_F^1 = (\varphi_F^{\frac{1}{k}})^k = \left(\varphi_{\frac{1}{k}F}^1\right)^k$ . Let  $\{\phi_i\}_{i=1}^k$  be symplectic diffeomorphisms, taking  $\mathcal{D}$  to k pairwise disjoint disks of area B, denote  $\Phi' = \prod_i \phi_i^* \varphi_{\frac{1}{k}F}^1$ , then  $\Phi'$  is generated by  $F' = \sum_{i=1}^k \frac{1}{k}F \circ \phi_i$ , hence

$$d(\mathbf{1}, \Phi') \leq \|F'\|_{C^0} \leq \frac{1}{k} \|F\|_{C^0}.$$

Lemma 3.3 implies that

$$d(\mathbf{1},\varphi_F^1) \leqslant d(\mathbf{1},\Phi') + 3\frac{1}{k+1} \leqslant \frac{1}{k} \|F\|_{C^0} + 3\frac{1}{k+1} \leqslant \frac{\|F\|_{C^0} + 3}{k}.$$
(5)

Let us state the following lemma (the *Uniform control* property of Theorem 3.2):

**Lemma 3.4.** Let  $H \in C_0^2(M, \mathbb{R})$ , then

$$\|\Sigma(H)\|_{C_0} \leq C \|H\|_{C^2}$$

for a constant C > 0 independent of H. In particular,  $\Sigma(H)$  is a bounded function in  $C^{0}(I)$ .

Before proving the lemma, let us show it implies an important part of the *Bounded growth control* property of Theorem 3.2.

Corollary 3.5. Let  $H \in C_0^2(S^2)$ , then

$$\Sigma(H) \equiv 0 \iff \rho(\varphi_H) = 0.$$

The first direction is quite immediate; in fact we can prove it independently of the lemma. Indeed  $\Sigma(H) \neq 0$  implies  $\mu_{k,B}(H) \neq 0$  for some k, B, and from the independence of Hamiltonian and Hofer-Lipschitz properties of  $\mu_{k,B}$  (see [5, 17]) we have

$$0 < |\mu_{k,B}(H)| \le \rho(H),$$

where by abuse of notation  $\rho(H)$  denotes the asymptotic Hofer growth of  $\{\phi_H^t\}$  inside  $\widehat{\operatorname{Ham}}(S^2)$ .

Now, for the other direction, assume by contradiction that both  $\rho(\varphi_H) = 2\varepsilon > 0$  and  $\Sigma(H) = 0$ . Let  $H_k$  be a sequence of Morse functions with  $||H - H_k||_{C_2} < \min(\frac{1}{k^2}, \varepsilon)$ . First, we have

$$\left|\frac{d(\mathbf{1},\varphi_{H}^{t})}{t} - \frac{d(\mathbf{1},\varphi_{H_{k}}^{t})}{t}\right| \leqslant \left|\frac{d(\varphi_{H}^{t},\varphi_{H'}^{t})}{t}\right| < \frac{t\varepsilon}{t} = \varepsilon \implies \rho(\varphi_{H_{k}}) > \varepsilon.$$

Second, for an interval  $J \subset I$  write  $J^c = I \setminus J$ , and define  $\Sigma(H_k)^-, \Sigma(H_k)^+$  as

 $\Sigma(H_k)^-|_{I_k} = \Sigma(H_k), \qquad \Sigma(H_k)^+|_{I_k^c} = \Sigma(H_k)$ 

extended by constants. Recall that  $r_k$  denotes the restriction of a function on I to  $I_k$ .

By Lipschitz continuity of Theorem 2.9 we have

$$\left\|\Sigma(H_k)^-\right\|_{C^0} = \|r_k \circ \Sigma(H_k)\|_{C^0} \leqslant k \frac{1}{k^2} \implies \rho(\varphi_{\Sigma(H_k)^-}) < \frac{1}{k}$$

Note,  $\Sigma(H_k)^+$  is supported on two disjoint disks of area  $\frac{1}{k+1}$  then we can use the improved Sikorav trick to get

$$\rho(\varphi_{\Sigma(H_k)^+}) \leqslant \frac{2\|H_k\|_{C^0} + 6}{k} \leqslant \frac{2\left(\frac{1}{k^2} + \|H\|_{C_0}\right) + 6}{k} \leqslant \frac{2\|H\|_{C_0} + 10}{k}.$$

In total, we get

$$\rho(\varphi_{H_k}) = \rho(\varphi_{\Sigma(H_k)}) = \rho(\varphi_{\Sigma(H_k)^+} \circ \varphi_{\Sigma(H_k)^+}) \leqslant \rho(\varphi_{\Sigma(H_k)^+}) + \rho(\varphi_{\Sigma(H_k)^-})$$
$$\leqslant \frac{1}{k} + \frac{2\|H\|_{C_0} + 10}{k},$$

and for k large enough this contradicts that  $\rho(\varphi_{H_k}) > \varepsilon$ .

Now we prove Lemma 3.4.

*Proof.* By Lipschitz continuity, it is easy to see that it is enough to prove the  $C^0$  bound for H Morse with distinct critical values. Like before, decompose H into  $H = \sum_i H|_{U_i}$ , where  $U_i$  are the disks/annuli of it's level sets, let  $H_i$  be the extension by constants of  $H|_{U_i}$ , denote the end points as  $a_i \leq b_i$ , i.e.  $\operatorname{Im}(H_i) = \overline{H(U_i)} = [a_i, b_i]$ , furthermore, denote

$$\begin{aligned} H'_i &= H_i - \langle H_i \rangle := H_i - \frac{1}{\operatorname{Area}(\mathbb{S}^2)} \int_{\mathbb{S}^2} H_i, \\ a'_i &:= a_i - \langle H_i \rangle, \\ b'_i &:= b_i - \langle H_i \rangle, \end{aligned}$$


This allows us to choose area coordinates on  $\Sigma(H'_i)$ , i.e. if  $B(x) = \text{Area}(\{H'_i < x\})$ , then

$$\Sigma(H'_i) = \frac{1}{2} (\widehat{\Sigma}(H'_i) + \widehat{\Sigma}(H'_i) \circ R),$$

where

$$\widehat{\Sigma}(H'_i)\left(-\frac{1}{2} + B(x)\right) = \begin{cases} a'_i & B(x) < A_i \\ x & A_i < B(x) < B_i \\ b'_i & B(x) > B_i \end{cases}$$
(6)

where  $A_i = \text{Area}\left(\{H'_i \leq a'_i\}\right) = \text{Area}\left(\{H'_i = a'_i\}\right), \ B_i = \text{Area}\left(\{H'_i \leq b'_i\}\right).$ Let  $B(x) \in (A_i, B_i)$ , then  $\frac{\partial}{\partial B} \Sigma(H'_i) = \frac{\partial x}{\partial B} = \frac{1}{\frac{\partial B}{\partial x}}.$ 



Via the Leibniz integral rule,

$$\frac{\partial B}{\partial x} = \frac{\partial}{\partial x} \int_{\{H'_i < x\}} \omega = \int_{\{H'_i = x\}} i_v \omega, \quad v = \frac{\nabla H'_i}{\left|\nabla H'_i\right|^2},$$

then

$$\frac{\partial B}{\partial x} = \int_{\left\{H'_i = x\right\}} i_v \omega = \int_0^{T(x)} \underbrace{\omega\left(\frac{\nabla H'_i}{\left|\nabla H'_i\right|^2}, X_{H'_i}\right)}_{=1} = T(x)$$

where T(x) is the period of  $\{H'_i = x\}$  as a trajectory of the flow of  $X_{H_i}$ . A result from [22] gives us the existence of some constant  $\tilde{C} > 0$  with the following bound,

$$T(x) \geqslant \frac{\tilde{C}}{\|H\|_{C^2}},$$

so,

$$\left|\frac{\partial}{\partial B}\Sigma(H'_i)\right| = \left|\frac{\partial x}{\partial B}\right| = \left|\frac{1}{\frac{\partial B}{\partial x}}\right| \leqslant C \|H\|_{C^2}, \qquad C = \frac{1}{\tilde{C}} > 0.$$

i.e.  $\Sigma(H'_i)$  is L-Lipschitz on  $(A_i, B_i)$  with  $L \leq C ||H||_{C^2}$ . Recall that the oscillation of a function  $G \in C^0(S^2)$  is defined as  $\operatorname{osc}(G) = \max(f) - \min(f)$ . Now from our normalization,  $0 \in (a'_i, b'_i)$  and hence

$$\begin{aligned} |\Sigma(H)| &\leq \sum_{i} \max |\Sigma(H'_{i})| \leq \sum_{i} \operatorname{osc}(H'_{i}) = \sum_{i} \operatorname{osc}(H_{i}) \\ &\leq \sum_{i} L \operatorname{Area}(U_{i}) = L \leq C ||H||_{C^{2}}. \end{aligned}$$

**Remark 3.6.** A more involved proof based on the theory of barcodes of functions (see [12]) allows one to replace the  $C_0^2(S^2)$  function space by the Sobolev  $W_0^{2,2}(S^2)$  function space, including the norms.

Let us now prove that  $\Sigma(H)$  for  $H \in C_0^2(S^2)$  is continuous at the endpoints  $\pm \frac{1}{2}$ . It will be convenient to denote the two-sphere by  $(M, \omega)$  in the following arguments. The proof of Lemma 3.4 yields the following:

**Lemma 3.7.** Let  $H \in C^2(M)$ . Denote by  $K \subset \mathbb{R}$  the set of critical values of H. The open set  $M \setminus H^{-1}(K)$  decomposes into at most countable union of disjoint open annuli  $A_j \subset M \setminus H^{-1}(K)$ , s.t. on each  $A_j$  the function H has a "standard" form, and such that

$$\sum_{j} \operatorname{osc}_{A_{j}} H \leqslant C \|H\|_{C^{2}}.$$

**Remark 3.8.** An equivalent way to formulate Lemma 3.7 is the following. Let  $H \in C^2(M)$ , let  $K \subset \mathbb{R}$  be the set of critical values of H, and denote  $I = [\min H, \max H]$ . Write  $I \setminus K = \bigcup_j I_j$  as at most countable disjoint union of open intervals. For every  $I_j$  there exists some  $m_j \in \mathbb{N}$  such that for each  $t \in I_j$ , the preimage  $H^{-1}(t)$  is a disjoint union of  $m_j$ embedded loops in M, and the preimage  $H^{-1}(I_j)$  is a union of  $m_j$  disjoint open annuli. Then  $\sum_{i=1}^{\infty} m_j \ell(I_j) \leq C ||H||_{C^2}$ , where  $\ell(I_j)$  is the length of  $I_j$ .

**Definition 3.9.** Let  $H \in C^2(M)$  and let  $\delta > 0$ . Denote by K the set of critical values of H, and denote  $I = [\min H, \max H]$ . A smooth function  $r : I \to \mathbb{R}$  is called  $(H, \delta)$ -admissible if the following holds:

- 1. r is locally constant on a neighborhood of K.
- 2.  $0 \leq r'(t) \leq 1$  for every  $t \in I$ .

- 3.  $|r(t) t| < \delta$  for every  $t \in I$ .
- 4.  $r(\min H) = \min H$ .

**Claim 1.** For every  $H \in C^2(M)$  and every  $\delta > 0$  there exists an  $(H, \delta)$ -admissible function.

*Proof.* Denote by K the set of critical values of H, then K has measure zero by the Sard theorem. Let  $I = [\min H, \max H]$ . Choose a neighborhood U of K in I such that the measure of U is less than  $\delta$ . Pick a smooth function  $h: I \to [0, 1]$  such that supp  $(h) \subset U$  and h = 1 on a neighborhood of K. Then the function  $r: I \to \mathbb{R}$  given by  $r(t) = \min H + \int_{\min H}^{t} (1-h(s)) ds$  satisfies all the properties.

**Lemma 3.10.** Let  $H \in C^2(S^2)$ . Then for every  $\epsilon > 0$  there exists  $\delta > 0$  such that if  $r: I \to \mathbb{R}$  is  $(H, \delta)$ -admissible then

$$\|\Sigma(r \circ H) - \Sigma(H)\|_{L_{\infty}(-1/2, 1/2)} < \epsilon.$$

**Corollary 3.11.** For every  $H \in C^2(S^2)$ , the function  $\Sigma(H)$  is continuous on the closed interval [-1/2, 1/2].

Proof. Take a sequence  $\delta_j > 0$  that converges to 0, and then choose a sequence  $r_j : I \to \mathbb{R}$  of smooth functions, where  $r_j$  is  $(H, \delta_j)$ -admissible for every j. Then it is not difficult to see that the functions  $\Sigma(r_j \circ H)$  are continuous on [-1/2, 1/2], and by Lemma 3.10,  $\Sigma(r_j \circ H)$  uniformly converges to  $\Sigma(H)$  on (-1/2, 1/2) when  $j \to \infty$ . Therefore  $\Sigma(H)$  extends to a continuous function on [-1/2, 1/2].

Proof of Lemma 3.10. Let  $H \in C^2(M)$ , let  $\epsilon > 0$ , and let  $\delta > 0$  be sufficiently small. We will show that if  $r, \tilde{r} : I \to \mathbb{R}$  are  $(H, \delta)$ -admissible then

$$\|\Sigma(r \circ H) - \Sigma(\tilde{r} \circ H)\|_{L_{\infty}(-1/2, 1/2)} < \epsilon \tag{7}$$

This would be enough for concluding the lemma. Indeed, assuming (7), for a given  $\epsilon > 0$ choose  $\delta > 0$  such that (7) holds. For any  $(H, \delta)$ -admissible function  $r : I \to \mathbb{R}$ , choose a sequence  $\tilde{r}_l : I \to \mathbb{R}$  such that  $\tilde{r}_l$  is  $(H, \delta/l)$ -admissible for every l. Choose any integer k > 2. Then by (7) we in particular get

$$\|\Sigma(r \circ H) - \Sigma(\tilde{r}_l \circ H)\|_{L_{\infty}(-1/2+1/k, 1/2-1/k)} < \epsilon.$$

The sequence of functions  $\tilde{r}_l \circ H$  uniformly converges to H when  $l \to \infty$ , therefore by the Lipschitz property, the sequence of functions  $\Sigma(\tilde{r}_l \circ H)$  uniformly converges to  $\Sigma(H)$  on (-1/2 + 1/k, 1/2 - 1/k) when  $l \to \infty$ . Hence we get

$$\|\Sigma(r \circ H) - \Sigma(H)\|_{L_{\infty}(-1/2+1/k, 1/2-1/k)} \leq \epsilon,$$

and since this holds for any k > 2, we finally conclude

$$\|\Sigma(r \circ H) - \Sigma(H)\|_{L_{\infty}(-1/2, 1/2)} \leq \epsilon,$$

and the lemma follows.

It remains to show (7). So let  $H \in C^2(M)$ , let  $\epsilon > 0$ , choose  $\delta > 0$  small enough, and let  $r, \tilde{r} : I \to \mathbb{R}$  be  $(H, \delta)$ -admissible, where  $I = [\min H, \max H]$ . Denote by K the set of critical values of H, and then write  $I \setminus K = \bigcup_j I_j$  as at most countable disjoint union of open intervals.

Now recall the Remark 3.8. For every  $I_j$  there exists  $m_j \in \mathbb{N}$  such that for each  $t \in I_j$ ,  $H^{-1}(t)$  is a union of  $m_j$  embedded loops on  $S^2$ , and such that  $H^{-1}(I_j)$  is a union of  $m_j$  disjoint open annuli on  $S^2$ . Write I = [a, b] and  $I_j = (a_j, b_j)$ . Lemma 3.7 and Remark 3.8 imply that

$$\sum_{j} m_j (b_j - a_j) \leqslant C \|H\|_{C^2} < +\infty.$$
(8)

**Claim 2.** If  $f: I \to \mathbb{R}$  is a smooth function such that f = 0 on a neighborhood of  $[a, a_j]$  and f = const on a neighborhood of  $[b_j, b]$ , then  $\|\Sigma(f \circ H)\|_{\infty} \leq m_j \operatorname{osc} f$ .

Proof. For each j, the pre-image  $H^{-1}(I_j)$  is a union of  $m_j$  annuli  $A_{j1}, \ldots, A_{jm_j}$ . For each  $1 \leq j \leq m_j$  let  $H_{jl}: M \to \mathbb{R}$  be the unique smooth function such that  $H_{jl} = f \circ H$  on  $A_{jl}$  and such that  $H_{jl}$  is constant on each of the connected components of  $M \setminus (A_{jl} \cap H^{-1}(I'_{jl}))$  where  $I'_{jl} \subset I$  is some closed sub-interval. Then the functions  $f \circ H$  and  $\sum_{l=1}^{m_j} H_{jl}$  differ by a constant (their difference is locally constant on M and hence globally constant), and comparing the functions at a point outside the union  $\bigcup_{l=1}^{m_j} A_{jl}$ , we conclude that the constant equals to some  $q_j f(b_j)$ , where  $0 \leq q_j \leq m_j$  is an integer. Hence we can write  $f \circ H = \sum_{l=1}^{m_j} \tilde{H}_{jl}$ , where  $\tilde{H}_{jl} = H_{jl} - f(b_j)$  for  $1 \leq l \leq q_j$  and  $\tilde{H}_{jl} = H_{jl}$  for  $q_j < l \leq m_j$ . Then for every l,  $\|\tilde{H}_{jl}\|_{\infty} \leq \operatorname{osc} f$ , and moreover  $\tilde{H}_{jl}$  is elementary and therefore  $\|\Sigma(\tilde{H}_{jl})\|_{\infty} \leq \|\tilde{H}_{jl}\|_{\infty} \leq \operatorname{osc} f$ . Since the functions  $\tilde{H}_{j1}, \ldots, \tilde{H}_{jm_j}$  Poisson commute, we get

$$\Sigma(f \circ H) = \Sigma(\sum_{l=1}^{m_j} \widetilde{H}_{jl}) = \sum_{l=1}^{m_j} \Sigma(\widetilde{H}_{jl})$$

and hence  $\|\Sigma(f \circ H)\|_{\infty} \leq m_j \underset{I}{\operatorname{osc}} f$ .

Now return to our functions  $r, \tilde{r} : I \to \mathbb{R}$ . Since they are  $(H, \delta)$ -admissible, each of them is locally constant on a neighborhood of K, hence there exists  $q \in \mathbb{N}$  such that r and  $\tilde{r}$ are locally constant on a neighborhood of  $I \setminus \bigcup_{j=1}^{q} I_j$ , in particular the difference  $r - \tilde{r}$  is locally constant on a neighborhood of  $I \setminus \bigcup_{j=1}^{q} I_j$ . Moreover,  $r(a) - \tilde{r}(a) = 0$ . Hence we can write  $r - \tilde{r} = \sum_{j=1}^{q} f_j$  where each  $f_j$  satisfies the assumptions of the claim (that is,  $f_j$  is smooth,  $f_j = 0$  on a neighborhood of  $[a, a_j]$ , and  $f_j = const$  on a neighborhood of  $[b_j, b]$ ), and  $\underset{I}{\operatorname{osc}} f_j = \underset{I_j}{\operatorname{osc}} (r - \tilde{r})$ . Since  $r \circ H$  and  $\tilde{r} \circ H$  Poisson commute, and since all  $f_j \circ H$  pairwise Poisson commute, we get

$$\Sigma(r \circ H) - \Sigma(\tilde{r} \circ H) = \Sigma((r - \tilde{r}) \circ H) = \Sigma(\sum_{j=1}^{q} f_j \circ H) = \sum_{j=1}^{q} \Sigma(f_j \circ H).$$

Hence by the claim,

$$\|\Sigma(r \circ H) - \Sigma(\tilde{r} \circ H)\|_{\infty} \leqslant \sum_{j=1}^{q} \|\Sigma(f_j \circ H)\|_{\infty} \leqslant \sum_{j=1}^{q} m_j \underset{I}{\operatorname{osc}} f_j = \sum_{j=1}^{q} m_j \underset{I_j}{\operatorname{osc}} (r - \tilde{r}).$$

The functions r and  $\tilde{r}$  are  $(H, \delta)$ -admissible, in particular  $|r(t) - t| < \delta$  and  $|\tilde{r}(t) - t| < \delta$ for every  $t \in I$ , and hence  $|r(t) - \tilde{r}(t)| < 2\delta$  for every  $t \in I$ . This implies that for every j we have  $\underset{I_j}{\operatorname{osc}}(r - \tilde{r}) < 4\delta$ . But also, since  $0 \leq r'(t), \tilde{r}'(t) \leq 1$  for every  $t \in I$ , we get  $\underset{I_j}{\operatorname{osc}}(r - \tilde{r}) \leq \ell(I_j) = b_j - a_j$ . Therefore  $\underset{I_j}{\operatorname{osc}}(r - \tilde{r}) \leq \min(4\delta, b_j - a_j)$  for every j, and hence we get

$$\|\Sigma(r \circ H) - \Sigma(\tilde{r} \circ H)\|_{\infty} \leq \sum_{j=1}^{q} m_j \operatorname{osc}_{I_j}(r - \tilde{r}) \leq \sum_j m_j \min(4\delta, b_j - a_j).$$

But because of (8), for  $\delta > 0$  small enough we have  $\sum_j m_j \min(4\delta, b_j - a_j) < \epsilon$ . We have proved (7).

#### 3.3 The Hölder inequality property

Now, let us prove the *Hölder's inequality* property of Theorem 3.2. Before we do that, we wish to extend the bound from the "improved Sikorav trick" to continuous functions. Note the following notation. For  $u \in C^0_{\text{even},0}(I)$  where  $I = \left(-\frac{1}{2}, \frac{1}{2}\right)$ , let us denote

$$||u||_k = \max_{I_k} |u| + \frac{2}{k} \max_{I \setminus I_k} |u|$$

If v is smooth and compactly supported, denote  $v_{-} = v|_{I_k}, v_{+} = v|_{I_k^c}$  continued by a constant, then like before, we can use the improved Sikorav trick as in estimate (5) to get

$$d\left(\mathbf{1},\varphi_{v}^{1}\right) \leq d\left(\mathbf{1},\varphi_{v_{-}}^{1}\right) + d\left(\mathbf{1},\varphi_{v_{+}}^{1}\right) \leq \left\|v\right\|_{k} + \frac{6}{k}.$$

For some general  $u \in C^0_{\text{even},0}(I)$ , we can take a smooth and compactly supported sequence  $\{v_i\}$  converging uniformly to u on any compact subsets of I, with supp  $(v_i) \subset I_i, \max_{I_k} |u - v_i| \leq \frac{1}{k}$  for all i > i(k), and  $\max_I |v_i| < C_1$ , denote by  $\psi_i$  the Hamiltonian diffeomorphism generated by  $v_i$ , and  $\psi_{i,j}$  the one generated by  $v_i - v_j$ , then

$$d(\psi_i, \psi_j) = d(\mathbf{1}, \psi_{ij}) \leqslant ||v_i - v_j||_k + \frac{6}{k} = O(1/k)$$

Therefor,  $\{\psi_i\}$  is a Cauchy sequence defining  $\varphi_u^1 \in \widehat{\operatorname{Ham}}(S^2)$ , and it follows that for all k,

$$\widehat{d}(\mathbf{1},\varphi_u^1) \leqslant \|u\|_k + \frac{6}{k}.$$

We will use this to prove the *Hölder's inequality* property of Theorem 3.2.

**Lemma 3.12.** Every pair  $H, H' \in C_0^2(S^2)$  with  $||H - H'||_{C^0} < 1$  has

$$\widehat{d}\left(\varphi_{\Sigma(H)}^{1},\varphi_{\Sigma(H')}^{1}\right) \leqslant C'\sqrt{\|H-H'\|_{C^{0}}}\left(\sqrt{1+\|H\|_{C^{2}}+\|H'\|_{C^{2}}}\right),$$

for a universal constant C' > 0.

Proof. Let  $H, H' \in C_0^2(S^2)$ . Set  $\varepsilon = ||H - H'||_{C^0}$ ,  $E = ||H||_{C^2} + ||H'||_{C^2}$ ,  $w = \Sigma(H) - \Sigma(H')$ . First, by the Uniform control property of Theorem 3.2 one has

$$\|w\|_{C^0} \leqslant \|\Sigma(H)\|_{C^0} + \|\Sigma(H')\|_{C^0} \leqslant CE,$$

and by the Lipschitz continuity property of Theorem 2.9, for every k,

$$\max_{I_k} |w| \leqslant k\varepsilon$$

Hence

$$\widehat{d}(\mathbf{1},\varphi_w^1) \leqslant \|w\|_k + \frac{6}{k} = \max_{I_k} |w| + \frac{2}{k} \max_{I \setminus I_k} |w| + \frac{6}{k} \leqslant k\varepsilon + \frac{2}{k}CE + \frac{6}{k} \leqslant k\varepsilon + \frac{C''(E+1)}{k},$$

for  $C'' \ge \max\{6, 2C\}$ . Note that by hypothesis, the set  $K = \left[\sqrt{\frac{C''(E+1)}{\varepsilon}}, 2\sqrt{\frac{C''(E+1)}{\varepsilon}}\right] \cap \mathbb{N}$  is non-empty. For an arbitrary  $k \in K$  we have

$$\begin{aligned} \widehat{d}(\mathbf{1},\varphi_w^1) \leqslant k\varepsilon + \frac{C''(E+1)}{k} \leqslant 2\sqrt{\frac{C''(E+1)}{\varepsilon}}\varepsilon + \frac{C''(E+1)}{\sqrt{\frac{C''(E+1)}{\varepsilon}}} \\ &= \underbrace{3\sqrt{C''}}_{:=C'}\sqrt{\varepsilon}\sqrt{E+1} = C'\sqrt{\|H-H'\|_{C^0}}\sqrt{1+\|H\|_{C^2} + \|H'\|_{C^2}} \end{aligned}$$

yielding our result.

### 3.4 The Flattening control property

Here we prove the *Flattening control* property of Theorem 3.2. For a Morse function  $H: S^2 \to \mathbb{R}$ , denote by e(H) the number of edges of the Reeb graph  $\Gamma_H$  of H.

**Claim 3.** If  $H: S^2 \to \mathbb{R}$  is a Morse function, and  $G: S^2 \to \mathbb{R}$  is an associated quasi-Morse function, then

$$\operatorname{osc} \Sigma(G) \leqslant e(H) \operatorname{osc} G.$$
(9)

Proof. We can find a sequence  $G_i: S^2 \to \mathbb{R}$  of smooth functions which uniformly converges to G, such that each  $G_i$  descends to  $\widehat{G_i}: \Gamma_H \to \mathbb{R}$  which is constant near the vertices of  $\Gamma_H$ of degree  $\geq 2$ . If (9) holds for every  $G_i$ , then from the Lipschitz continuity property for  $\Sigma$  it would follow that (9) holds also for G. Hence, without loss of generality we can assume that the induced function  $\widehat{G}: \Gamma_H \to \mathbb{R}$  is constant near all vertices of  $\Gamma_H$  of degree  $\geq 2$ .

We can decompose  $G = \sum_{e \in E(\Gamma_H)} G_e$ , where  $E(\Gamma_H)$  is the set of the edges of  $\Gamma_H$ , and for each  $e \in E(\Gamma_H)$ ,  $G_e$  is smooth and descends to  $\widehat{G_e} : \Gamma_H \to \mathbb{R}$  such that  $\widehat{G_e}$  is locally constant on  $\Gamma_H \setminus \widetilde{e}$  where  $\widetilde{e} \subset int(e)$  is a compact subset (this determines the functions  $G_e$  uniquely up to an additive constant). Notice that  $\operatorname{osc} G_e = \operatorname{osc}_e \widehat{G} \leq \operatorname{osc} G$ . Moreover, notice that for each  $G_e$  there exists  $\psi_e \in \operatorname{Ham}(S^2)$  such that  $F_e := \psi_e^* G_e$  depends only on the z-coordinate. We have  $\Sigma(G_e)(z) = \Sigma(F_e)(z) = (F_e(z) + F_e(-z))/2$ , hence

$$\operatorname{osc} \Sigma(G_e) \leq \operatorname{osc} F_e = \operatorname{osc} G_e \leq \operatorname{osc} G_e$$

In addition, we have  $\Sigma(G) = \Sigma(\sum_{e \in E(\Gamma_H)} G_e) = \sum_{e \in E(\Gamma_H)} \Sigma(G_e)$ . We finally conclude

osc 
$$\Sigma(G) \leq \sum_{e \in E(\Gamma_H)} \text{osc } \Sigma(G_e) \leq e(H) \text{ osc } G.$$

**Corollary 3.13.** Let  $H : S^2 \to \mathbb{R}$  be a Morse function, let  $\varepsilon > 0$ , and let  $r : \mathbb{R} \to \mathbb{R}$  be a smooth function such that  $|r(t) - t| \leq \varepsilon$  for every  $t \in \mathbb{R}$ . Denote  $\widetilde{H} = r \circ H$ . Then  $\|\Sigma(\widetilde{H}) - \Sigma(H)\|_{C^0} \leq (1 + 2e(H))\varepsilon$ .

*Proof.* As  $\widetilde{H}$ , H commute, in the sense that their Poisson bracket satisfies  $\{\widetilde{H}, H\} = 0$ , we have  $\Sigma(\widetilde{H}) - \Sigma(H) = \Sigma(\widetilde{H} - H)$ . Moreover

$$\left| \int_{S^2} \Sigma(\widetilde{H} - H) \omega \right| = \left| \int_{S^2} (\widetilde{H} - H) \omega \right| \leqslant \varepsilon \operatorname{Area}(S^2).$$

Hence, by the lemma

$$\begin{aligned} \|\Sigma(\widetilde{H}) - \Sigma(H)\|_{C^0} &= \|\Sigma(\widetilde{H} - H)\|_{C^0} \leqslant \varepsilon + \operatorname{osc} \Sigma(\widetilde{H} - H) \\ &\leqslant \varepsilon + e(H) \operatorname{osc} (\widetilde{H} - H) \leqslant \varepsilon + 2e(H)\varepsilon = (1 + 2e(H))\varepsilon. \end{aligned}$$

#### 3.5 Conjugation control and Bounded growth control

In this section we will state the "conjugation lemma", Lemma 3.14, and use it to prove *Conjugation control* of Theorem 3.2 and finish the proof of the *Bounded growth control* property of the same theorem. The proof of Lemma 3.14 is postponed to the next section.

**Lemma 3.14.** (The Conjugation lemma) Let H' be a Morse function on  $S^2$ . Let  $\tilde{H} = r \circ H'$ be a sufficiently small  $\varepsilon$ -flattening, such that r is constant near the median of H'. Then

$$d(\varphi^1_{\widetilde{H}}, \psi^{-1}\varphi^1_{\Sigma(\widetilde{H})}\psi) \leqslant 18\operatorname{Area}(S^2)$$

for some  $\psi \in \operatorname{Ham}(S^2)$ .

We will now prove that the lemma implies the following (*Conjugation control* property of Theorem 3.2).

**Corollary 3.15.** For every  $H \in C_0^2(S^2)$  there exists  $\psi \in \widehat{Ham}(M)$  such that

$$\widehat{d}(\varphi_H^1, \psi^{-1}\varphi_{\Sigma(H)}^1\psi) \leqslant 19\operatorname{Area}(S^2).$$

*Proof.* Let  $H \in C_0^2(S^2)$ . Let H' be Morse with distinct critical values such that

$$\|H-H'\|_{C^2} < \varepsilon_1 < 1,$$

where  $\varepsilon_1$  is an arbitrarily small constant which we will choose later. Let us record that

$$d(\varphi_H^1, \varphi_{H'}^1) < \varepsilon_1.$$

By flattening control (see Corollary 3.13 with H replaced by H'), we may choose a flattening  $\tilde{H}$  of H' such that  $\left\|\tilde{H} - H'\right\|_{C^0} \leq \varepsilon_2 = \frac{1}{k^2}$  where k > 1 + 2e(H') and

$$|\Sigma(H') - \Sigma(\widetilde{H})|_{C^0} < \frac{1}{k}$$

This implies that

$$\widehat{d}\left(\varphi_{\Sigma(H')}^{1},\varphi_{\Sigma(\widetilde{H})}^{1}\right)\leqslant\frac{1}{k}.$$

Let us also record that in this case

$$\widehat{d}\left(\varphi_{H'}^{1},\varphi_{\widetilde{H}}^{1}\right) \leqslant \frac{1}{k^{2}}$$

Finally, we estimate that

$$\begin{split} \widehat{d}\left(\varphi_{H}^{1},\psi^{-1}\varphi_{\Sigma(H)}^{1}\psi\right) &\leqslant \widehat{d}\left(\varphi_{H}^{1},\varphi_{\widetilde{H}}^{1}\right) + \widehat{d}\left(\varphi_{\widetilde{H}}^{1},\psi^{-1}\varphi_{\Sigma(\widetilde{H})}^{1}\psi\right) \\ &+ \underbrace{\widehat{d}\left(\psi^{-1}\varphi_{\Sigma(\widetilde{H})}^{1}\psi,\psi^{-1}\varphi_{\Sigma(H)}^{1}\psi\right)}_{\widehat{d}\left(\varphi_{\Sigma(\widetilde{H})}^{1},\varphi_{\Sigma(H)}^{1}\right)}. \end{split}$$

But of course,

$$\widehat{d}(\varphi_{H}^{1},\varphi_{\widetilde{H}}^{1}) \leqslant \widehat{d}(\varphi_{H}^{1},\varphi_{H'}^{1}) + \widehat{d}(\varphi_{H'}^{1},\varphi_{\widetilde{H}}^{1}) \leqslant \varepsilon_{1} + \frac{1}{k^{2}},$$

and,

$$\begin{split} \widehat{d}\left(\varphi_{\Sigma(\widetilde{H})}^{1},\varphi_{\Sigma(H)}^{1}\right) &\leqslant \widehat{d}\left(\varphi_{\Sigma(\widetilde{H})}^{1},\varphi_{\Sigma(H')}^{1}\right) + \widehat{d}\left(\varphi_{\Sigma(H')}^{1},\varphi_{\Sigma(H)}^{1}\right) \\ &\leqslant \widehat{d}\left(\varphi_{\Sigma(H')}^{1},\varphi_{\Sigma(H)}^{1}\right) + \frac{1}{k}. \end{split}$$

By the *Hölder's inequality* property of Theorem 3.2,

$$\widehat{d}\left(\varphi_{\Sigma(H')}^{1},\varphi_{\Sigma(H)}^{1}\right) \leqslant C'\varepsilon_{1}^{\frac{1}{2}}\left(\varepsilon_{1}+2\|H\|_{C_{2}}\right)^{\frac{1}{2}}.$$

In total, we obtain

$$\widehat{d}(\varphi_{H}^{1}, \psi^{-1}\varphi_{\Sigma(H)}^{1}\psi) \leq 18\operatorname{Area}(S^{2}) + \varepsilon_{1} + \frac{1}{k^{2}} + \frac{1}{k} + C'\varepsilon_{1}^{\frac{1}{2}}\left(\varepsilon_{1} + 2\|H\|_{C_{2}}\right)^{\frac{1}{2}}.$$

Note that  $\psi$  depends on  $\widetilde{H}$ , ultimately depending on  $\varepsilon_1$  and k. Taking  $\varepsilon_1$  arbitrarily small and k arbitrarily large we get that

$$\forall \varepsilon > 0, \ \exists \psi \in \widehat{\operatorname{Ham}}(S^2), \qquad \widehat{d}(\varphi_H^1, \psi^{-1} \varphi_{\Sigma(H)}^1 \psi) \leqslant 18 \operatorname{Area}(S^2) + \varepsilon.$$

Taking  $\varepsilon < \operatorname{Area}(S^2)$  we obtain the *Conjugation control* property of Theorem 3.2, but also, if  $\Sigma(H) \equiv 0$ , we have


$$\widehat{d}\left(\varphi_{H}^{1},\underbrace{\psi^{-1}\varphi_{\Sigma(H)}^{1}\psi}_{1}\right) \leqslant 19\operatorname{Area}(S^{2}),$$

and since  $\Sigma(tH) = t\Sigma(H)$ , this proves what remained to show of the *Bounded growth control* property.

### 3.6 The proof of the conjugation lemma

The following more flexible version of Lemma 3.3 will be needed in the proof of the conjugation lemma:

**Lemma 3.16.** Let  $(M, \omega)$  be a closed and connected symplectic surface. Let  $\varepsilon > 0$ , let m be a positive integer, and let  $\mathcal{D}_0, ..., \mathcal{D}_m \subset M$  be disjoint topological open disks, such that  $\mathcal{D}_0$  is of area  $\varepsilon$ , and for each  $1 \leq i \leq m$ , the area of  $\mathcal{D}_i$  lies in  $(\varepsilon/2, \varepsilon]$ . Finally, let  $\psi_j$ :



 $\mathcal{D}_j \to \mathcal{D}_0$  for all  $1 \leq j \leq m$  be symplectic embeddings. Consider  $f_0, ..., f_m \in \operatorname{Ham}(M, \omega)$  with supp  $(f_j) \subset \mathcal{D}_j$ . Define  $\Phi, \Phi' \in \operatorname{Ham}(M, \omega)$  by

$$\Phi = f_0 f_1 \cdots f_n$$

and

$$\Phi' = f_0 \prod_{j=1}^m (\psi_j)_* f_j$$

where  $(\psi_j)_* f_j$  is given by  $(\psi_j)_* f_j = \psi_j f_j(\psi_j)^{-1}$  on  $\psi_j(\mathcal{D}_j) \subset \mathcal{D}_0$  and  $(\psi_j)_* f_j = \mathbf{1}$  on  $M \setminus \psi_j(\mathcal{D}_j)$ . Then

$$d(\Phi, \Phi') < 7\varepsilon.$$

*Proof.* Without loss of generality we may assume that each  $\mathcal{D}_j$  has a smooth boundary and its area lies in  $(\varepsilon/2, \varepsilon)$ , and moreover each  $\psi_j$  extends to a smooth symplectic embedding  $\psi_j : \overline{\mathcal{D}_j} \to \mathcal{D}_0$ .

First consider the case when m = 2k is even. Look at the disks  $\mathcal{D}_0, \ldots, \mathcal{D}_k$ . The area of the complement to the union of these disks is greater than or equal to the sum of areas of  $\mathcal{D}_{k+1}, \ldots, \mathcal{D}_m$ , which in turn is greater than  $k\varepsilon/2$ . Moreover, recall that the area of each disk  $\mathcal{D}_j$  lies in  $(\varepsilon/2, \varepsilon]$ . Hence for each  $1 \leq j \leq k$  we can find an open disk  $\mathcal{D}'_j \supset \overline{\mathcal{D}}_j$  of area  $\varepsilon$ , such that each  $\psi_j$  can be extended to a symplectomorphism  $\psi_j : \mathcal{D}'_j \to \mathcal{D}_0$ , and such that moreover the disks  $\mathcal{D}_0, \mathcal{D}'_1, \ldots, \mathcal{D}'_k$  are pairwise disjoint. Applying Lemma 3.3, we conclude that for  $\Phi_0 := f_0 \cdots f_k$  and  $\Phi'_0 = f_0 \prod_{j=1}^k (\psi_j)_* f_j$  we have  $d(\Phi_0, \Phi'_0) < 3\varepsilon$ . Now, looking at the disks  $\mathcal{D}_0, \mathcal{D}_{k+1}, \ldots, \mathcal{D}_m$ , and applying a similar reasoning, for  $\Phi_1 := f_{k+1} \cdots f_m$  and  $\Phi'_1 = \prod_{j=k+1}^m (\psi_j)_* f_j$  we get  $d(\Phi_1, \Phi'_1) < 3\varepsilon$ . Finally, we conclude  $d(\Phi, \Phi') = d(\Phi_0 \Phi_1, \Phi'_0 \Phi'_1) \leq$  $d(\Phi_0, \Phi'_0) + d(\Phi_1, \Phi'_1) < 6\varepsilon < 7\varepsilon$ .

It remains to verify the case when m = 2k + 1 is odd. Denote  $\Psi := f_0 f_1 \cdots f_{m-1}$ and  $\Psi' = f_0 \prod_{j=1}^{m-1} (\psi_j)_* f_j$ . We have shown that  $d(\Psi, \Psi') < 6\varepsilon$ . Since the area of  $\mathcal{D}_m$ is less than  $\varepsilon$  and  $\psi_m(\overline{\mathcal{D}}_m) \cap \overline{\mathcal{D}}_m = \emptyset$ , we can find a Hamiltonian diffeomorphism  $\phi \in$  $\operatorname{Ham}(M, \omega)$  with  $d(\mathbf{1}, \phi) < \varepsilon/2$ , such that  $\phi = \psi_m$  on  $\mathcal{D}_m$ , and in particular  $(\psi_m)_* f_m =$  $\phi^{-1} f_m \phi$ , which implies  $d(f_m, (\psi_m)_* f_m) = d(f_m, \phi^{-1} f_m \phi) \leq 2d(\mathbf{1}, \phi) < \varepsilon$ . We conclude  $d(\Phi, \Phi') = d(\Psi f_m, \Psi'(\psi_m)_* f_m) < d(\Psi, \Psi') + d(f_m, (\psi_m)_* f_m) < 7\varepsilon$ .

Proof of the conjugation lemma. Decompose  $\widetilde{H} = \sum_A \widetilde{H}_A$ , like before, where  $\{A\}$  are annuli/disks corresponding to the edges of the Reeb graph of H', and  $\widetilde{H}_A$  is locally constant outside A. Crucially, since we assumed r is flat near the median level, each A lies in a disk of area  $< \frac{1}{2} \operatorname{Area}(S^2)$ . Let  $\partial A$  be the union of two disjoint circles,  $C_+, C_-$ . We denote by  $D_+(A)$ the disk of area  $< \frac{1}{2} \operatorname{Area}(S^2)$  containing A, whose boundary is  $C_+$ . Similarly, we denote by  $D_-(A)$  the disk of area  $< \frac{1}{2} \operatorname{Area}(S^2)$  bounded by  $C_-$ . If A is a disk,  $D_-(A) = \emptyset$ . Let mbe the median level of H', and let  $\varepsilon > 0$  such that  $r|_{[m-2\varepsilon,m+2\varepsilon]} = \text{const.}$  The connected components of  $S^2 \setminus (H')^{-1}(m-\varepsilon, m+\varepsilon)$  are displaceable, hence one can find a collection of disjoint disks  $D_1, ..., D_M$  with smooth boundary, each of Area  $< \frac{1}{2} \operatorname{Area}(S^2)$  such that

$$S^2 \setminus (H')^{-1}(m-\varepsilon,m+\varepsilon) \subset \bigcup_j D_j.$$


Order them such that  $\operatorname{Area}(D_1) \ge \operatorname{Area}(D_j)$  for all j, and by conjugating with some  $\psi \in \operatorname{Ham}(S^2)$ , we can assume  $D_1$  is the standard cap

$$D_1 = \{(x, y, z) \in \mathbb{S}^2 | z > z_1\}, \qquad z_1 \in \left(0, \frac{1}{2}\right).$$

Note that the time-1 map is  $\varphi_{\widetilde{H}}^1 = \prod_i \varphi_{\widetilde{H}_i}^1$ . We will now deal the annuli which lie inside and outside  $D_1$  separately. Before starting let us define the following.

**Definition 3.17.** A collection of annuli  $C = (A_1, ..., A_l)$  is called <u>ordered</u> if

 $D_{-}(A_i) \supset D_{+}(A_{i+1}), \quad \forall 1 \leq i < l.$ 



**Remark 3.18.** Note that an ordered collection of annuli is a chain in the partially ordered set of annuli contained in a disk of area  $< \frac{1}{2}$ Area $(S^2)$ , with partial order  $A_1 < A_2$  iff  $D_-(A_1) \supset D_+(A_2)$ .

Let us start with the annuli lying on the complement of  $D_1$ : Chose  $\varepsilon = \frac{1}{2^N} \operatorname{Area}(S^2)$  such that  $\varepsilon < \operatorname{Area}(D_+(A))$  for all annuli  $A \subset S^2 \setminus D_1$ . We will iterate the following step N - 1 times: For each j = 0, ..., N - 2, consider all annuli  $A \subset S^2 \setminus D_1$  such that  $\operatorname{Area}(D_+(A)) \in (2^j \varepsilon, 2^{j+1} \varepsilon]$ . Denote the collection of these annuli by  $C_j$ . From the constraint on the area of  $D_+(A)$ , we can partition  $C_j$  to be  $C_j = C_{j1} \cup ... \cup C_{jk_j}$ , such that each  $C_{jk}$  is an ordered collection of annuli, and the disks  $\{D_+(C_{jk})|k=1,...,k_j\}$  are pairwise disjoint, and disjoint from  $D_1$ . Now we may use Lemma 3.16 in the following way. Denote  $\widetilde{H}_{jk} = \sum_{A \in C_{jk}} \widetilde{H}_A$ , and for each  $1 \leq k \leq k_j$ , find  $\varphi_{k,j} \in \text{Ham}(\mathbb{S}^2)$  such that  $\varphi_{k,j}^* \widetilde{H}_{jk}$  is a function of z, constant outside  $D_1$ . Then if we denote

$$\widetilde{H}^j := \sum_{k=1}^{k_j} \widetilde{H}_{jk}, \qquad \widehat{H}^j := \sum_{k=1}^{k_j} \varphi_{k,j}^* \widetilde{H}_{jk},$$

then we have

$$d\left(\varphi_{\widetilde{H}^{j}}^{1},\varphi_{\widehat{H}^{j}}^{1}\right) < 7 \cdot 2^{j+1}\varepsilon < 2^{j+4}\varepsilon.$$

Hence in total, after N-1 steps, the functions

$$\widetilde{H}_{\text{out}} := \sum_{j=0}^{N-2} \widetilde{H}^j, \qquad \widehat{H}_{\text{out}} := \sum_{j=0}^{N-2} \widehat{H}^j,$$

satisfy

$$d(\varphi_{\widehat{H}_{\text{out}}}^1, \varphi_{\widehat{H}_{\text{out}}}^1) \leqslant \sum_{j=0}^{N-2} 2^{j+4} \varepsilon < 2^{N+3} \varepsilon = 8 \operatorname{Area}(\mathbb{S}^2).$$

For the annuli lying in  $D_1$ :

Here we do similarly as we did before, only first moving the annuli to  $S^2 \setminus D_1$ . Recall that  $D_1$  is the standard cap inside a hemisphere, so this additional movement will "cost" us at most  $\frac{1}{2}$  Area $(S^2)$  Hofer's energy.

In total, we will have  $\widetilde{H}_{in} = \sum_{A \subset D_1}^{\infty} \widetilde{H}_A$ , and some  $\widehat{H}_{in}(z)$  which is constant outside  $D_1$ , and has

$$d(\varphi_{\widetilde{H}_{\text{in}}}^1, \varphi_{\widehat{H}_{\text{in}}}^1) < 9 \operatorname{Area}(S^2).$$

Since

$$\widetilde{H} = \widetilde{H}_{out} + \widetilde{H}_{in}, \qquad \widehat{H} = \widehat{H}_{out} + \widehat{H}_{in}, \qquad \Sigma\left(\widetilde{H}\right) = \frac{1}{2}\widehat{H} + \frac{1}{2}\varphi^*\widehat{H}$$

where  $\varphi$  is a rotation of the sphere "upside-down", the calculation gives us,

$$d\left(\varphi_{\widehat{H}}^1, \varphi_{\Sigma(\widetilde{H})}^1\right) < \operatorname{Area}(S^2).$$

Then in total

$$d\left(\varphi_{\widetilde{H}}^{1},\varphi_{\Sigma(\widetilde{H})}^{1}\right) < 18\operatorname{Area}(S^{2}).$$

Recall that we conjugated to get  $D_1$  to be the standard cap, thus completing the proof.  $\Box$ 

# 4 Discussion

### 4.1 Surfaces other than $S^2$

The case of the two-disk and compactly supported Hamiltonian diffeomorphisms is very similar to that of the sphere. Furthermore, we expect our methods to prove a growth dichotomy statement for all compact surfaces (with or without boundary). We plan to attack this question by means of symmetrization based on partial quasi-morphisms from [11], which should apply to Hamiltonians with all regular components of level sets contractible. Hamiltonians H without this property are easily seen to have linear Hofer growth  $\rho(H) > 0$ for instance by the energy capacity inequality in the universal cover (cf. [14, Exercise 7.2.E]).

To make this more concrete, take a closed symplectic surface  $(M, \omega)$  of total area 1. We shall use the following "Cartan subalgebra" in  $C^0(M)$ . Fix a symplectic embedding of the open Euclidean disk of area 1,

$$i: (D^2, dp \wedge dq) \to (M, \omega),$$

where  $M \setminus i(D^2)$  is a collection of closed simple curves on M. Think of the interior of the standard fundamental domain of M on the universal cover. Write  $z = (p^2 + q^2)/2$  for the symplectic polar radius. Denote by  $\mathcal{F} \subset C_0^0(M)$  the set of continuous mean zero functions on M of the form  $i_*F - \int_{D^2} F\omega$ , where F runs over the set of smooth compactly supported functions on  $D^2$  depending only on z.

Denote by  $\mathcal{C}$  the set of mean zero autonomous Hamiltonians on M whose regular level sets are contractible. Define the symmetrization map  $\Sigma : \mathcal{C} \to \mathcal{F}$ . For suitably flattened functions (as in Lemma 3.10, see also Remark 4.1 below) it is defined analogously to the case of  $S^2$ , and in the general case one uses the local quasi-morphisms constructed in [11] similarly to what we did in Section 2.2 above.

The conclusion is that for  $H \in \mathcal{C}$  the growth is linear if  $\Sigma(H) \neq 0$ , and it is bounded if  $\Sigma(H) = 0$ . As explained above, this proves a growth dichotomy statement for arbitrary autonomous Hamiltonians on M. We expect that an enhanced dichotomy also holds in this setting. The details will appear in a forthcoming paper.

**Remark 4.1.** Crucially, the flattening of Lemma 3.10 would allow us to proceed with the proof of the soft part of the dichotomy on M, since given H with contractible regular level sets, we can directly pass to  $r \circ H$  without the need to approximate H by a Morse function with contractible regular level sets.

#### 4.2 Quasi-morphisms and autonomous Hamiltonian diffeomorphisms

We note that given measures  $\delta_i = \delta_{k_i,B_i}$  on (-1/2, 1/2) such that  $\sum a_i \delta_i = 0$  for certain  $a_i \in \mathbb{R}$ , the quasimorphism

$$\mu = \sum a_i \mu_{k_i, B_i} : \operatorname{Ham}(S^2) \to \mathbb{R}$$
(10)

vanishes on the subgroup  $\mathcal{T}$  of  $\operatorname{Ham}(S^2)$  given by autonomous Hamiltonians generated by functions of the height z only. As  $\mu$  is Lipschitz in the Hofer metric d, it extends to the Hofer completion  $\widehat{\operatorname{Ham}}(S^2)$  and vanishes on  $\widehat{\mathcal{T}}$ . Therefore it follows from Theorem 1.3 that  $\mu$  vanishes on the subset  $\operatorname{Aut}(S^2) \subset \operatorname{Ham}(S^2)$  of all autonomous Hamiltonian diffeomorphisms (those generated by time-independent Hamiltonians  $H \in C^{\infty}(S^2, \mathbb{R})$ ). It is also follows immediately from [7, Theorem 1.7] that  $\mu$  is continuous with respect to the  $C^0$  topology on  $\operatorname{Ham}(S^2)$  and extends to the  $C^0$ -closure  $\overline{\operatorname{Ham}}(S^2)$  of  $\operatorname{Ham}(S^2)$  inside  $\operatorname{Homeo}(S^2)$ . A particular example of such a quasimorphism  $\mu$  is given by

$$\mu = 3\mu_{3,B} - 2\mu_{2,B} - \mu_{1,1/2}$$

for 1/3 < B < 1/2, where  $\mu_{1,1/2}$  is the Entov-Polterovich quasimorphism [6].

In future work joint with P. Haim-Kislev we plan to attack the following question.

Question 4.2. Do all the quasimorphisms  $\mu : \operatorname{Ham}(S^2) \to \mathbb{R}$  from (10) vanish identically?

We note that if one such  $\mu$  does not vanish, then the Hofer distance

$$\operatorname{aut}(S^2) = \sup_{\phi \in \operatorname{Ham}(S^2)} d(\phi, \operatorname{Aut}(S^2))$$

is infinite, which would settle the last open case of a conjecture of Polterovich-Shelukhin [16] for surfaces. Moreover, for all natural k,

$$\operatorname{aut}_k(S^2) = \sup_{\phi \in \operatorname{Ham}(S^2)} d(\phi, \operatorname{Aut}(S^2)^k) = \infty,$$

similar to the case of  $L^p$ -metrics as in [2].

If on the other hand all such  $\mu$  vanish, then the symmetrization map

$$\operatorname{Aut}(S^2) \to \mathcal{T}_{ei}$$

extends to all  $\operatorname{Ham}(S^2)$ , and it would be very interesting to study its properties.

### References

- A. M. Bloch, H. Flaschka, and T. Ratiu. A Schur-Horn-Kostant convexity theorem for the diffeomorphism group of the annulus. *Invent. Math.*, 113(3):511–529, 1993.
- [2] M. Brandenbursky and E. Shelukhin. On the L<sup>p</sup>-geometry of autonomous Hamiltonian diffeomorphisms of surfaces. *Math. Res. Lett.*, 22(5):1275–1294, 2015.
- [3] T. Bröcker and T. tom Dieck. Representations of compact Lie groups, volume 98 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1995. Translated from the German manuscript, Corrected reprint of the 1985 translation.
- [4] L. Buhovsky. On two remarkable groups of area-preserving homeomorphisms. Zurnal matematiceskoj fiziki, analiza, geometrii, 19(2):339–373, 2023.

- [5] D. Cristofaro-Gardiner, V. Humilière, C. Y. Mak, S. Seyfaddini, and I. Smith. Quantitative Heegaard Floer cohomology and the Calabi invariant. *Forum Math. Pi*, 10:Paper No. e27, 59, 2022.
- [6] M. Entov and L. Polterovich. Calabi quasimorphism and quantum homology. Int. Math. Res. Not., (30):1635–1676, 2003.
- [7] M. Entov, L. Polterovich, and P. Py. On continuity of quasimorphisms for symplectic maps. In *Perspectives in analysis, geometry, and topology*, volume 296 of *Progr. Math.*, pages 169–197. Birkhäuser/Springer, New York, 2012. With an appendix by M. Khanevsky.
- [8] H. Hofer. On the topological properties of symplectic maps. Proc. Roy. Soc. Edinburgh Sect. A, 115(1-2):25–38, 1990.
- [9] A. Izosimov, B. Khesin, and M. Mousavi. Coadjoint orbits of symplectic diffeomorphisms of surfaces and ideal hydrodynamics. Ann. Inst. Fourier (Grenoble), 66(6):2385–2433, 2016.
- [10] F. Lalonde and D. McDuff. The geometry of symplectic energy. Ann. of Math. (2), 141(2):349–371, 1995.
- [11] C. Y. Mak and I. Trifa. Hameomorphism groups of positive genus surfaces. Preprint arXiv:2306.06377, 2023.
- [12] I. Polterovich, L. Polterovich, and V. Stojisavljević. Persistence barcodes and Laplace eigenfunctions on surfaces. *Geom. Dedicata*, 201:111–138, 2019.
- [13] L. Polterovich. Symplectic displacement energy for Lagrangian submanifolds. Ergodic Theory Dynam. Systems, 13(2):357–367, 1993.
- [14] L. Polterovich. The geometry of the group of symplectic diffeomorphisms. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2001.
- [15] L. Polterovich and D. Rosen. Function theory on symplectic manifolds, volume 34 of CRM Monograph Series. American Mathematical Society, Providence, RI, 2014.
- [16] L. Polterovich and E. Shelukhin. Autonomous Hamiltonian flows, Hofer's geometry and persistence modules. *Selecta Math.* (N.S.), 22(1):227–296, 2016.
- [17] L. Polterovich and E. Shelukhin. Lagrangian configurations and Hamiltonian maps. Compos. Math., 159(12):2483–2520, 2023.

- [18] L. Polterovich and K. F. Siburg. On the asymptotic geometry of area-preserving maps. Math. Res. Lett., 7(2-3):233-243, 2000.
- [19] F. Schlenk. Symplectic embeddings of ellipsoids. Israel Journal of Mathematics, 138(1):215-252, 2003.
- [20] J.-C. Sikorav. Systèmes hamiltoniens et topologie symplectique. 1990. Università di Pisa.
- [21] C. Viterbo. Symplectic topology as the geometry of generating functions. *Math. Ann.*, 292(4):685–710, 1992.
- [22] J. A. Yorke. Periods of periodic solutions and the Lipschitz constant. Proceedings of the American Mathematical Society, 22(2):509–512, 1969.