

Chow rings of quasi-split geometrically almost simple algebraic groups

Alexey Ananyevskiy and Nikita Geldhauser

Dedicated to Professor Nikolai Vavilov (1952 – 2023)
 who taught us the numerology of algebraic groups.
 Ἦθος ἀνθρώπων δαίμων.

Abstract

We compute the Chow ring $\mathrm{Ch}^*(G)$ of a quasi-split geometrically almost simple algebraic group G assuming the coefficients to be a field. This extends the classical computation for split groups done by Kac to the non-split quasi-split case. For the proof we introduce and study equivariant conormed Chow rings, which are well adapted to the study of quasi-split groups and their homogeneous varieties.

Contents

1	Introduction	2
2	Conormed Chow ring of a quasi-split simple group	7
2.1	Conormed Chow groups	8
2.2	Equivariant conormed Chow groups	10
2.3	Eilenberg–Moore isomorphism for quasi-trivial tori	14
2.4	Some equivariant conormed Chow rings of a point	18
2.5	Conormed characteristic sequence for quasi-split simple groups	20
2.6	Conormed Chow ring of a quasi-split group G via G/B	23
2.7	Conormed Chow ring of a quasi-split group G via $\mathcal{C}_K^*(G)$	30
2.8	Conormed Chow ring of a quasi-split group: the answer	31
2.9	Cokernel in the conormed characteristic sequence	35
3	Chow ring of a quasi-split simple group	36
3.1	Quasi-split submaximal orthogonal Grassmannian	36
3.2	Chow ring of an adjoint quasi-split simple group of type ${}^2\mathrm{D}_{2r}$	43
3.3	Chow ring of a quasi-split group at the splitting prime	45
3.4	Chow ring of a quasi-split group away from the splitting primes	47
3.5	Chow ring of the group of type ${}^6\mathrm{D}_4$ at $p = 2, 3$	49
A	Recollection on algebraic groups	53
A.1	Some groups of multiplicative type and their representations	53
A.2	Quasi-split simple groups	54
A.3	Vector bundles on homogeneous varieties	57
A.4	Fundamental group and Picard group of a quasi-split simple group	58
A.5	Chow ring of a split simple group	59

1 Introduction

Given a linear algebraic group G one can associate with it two rings of cohomological origin, $\mathrm{CH}^*(BG)$ and $\mathrm{CH}^*(G)$, both of which carrying important information about possible twisted forms of G , its torsors, homogeneous varieties and related motives. Classical topological analogues of these rings are the singular cohomology ring $H^*(BG)$ of the classifying space of a Lie group \mathcal{G} and the singular cohomology ring $H^*(\mathcal{G})$ of the Lie group itself, and these rings are intimately related by the Leray–Serre spectral sequence for the fibration $E\mathcal{G} \rightarrow B\mathcal{G}$.

The ring $\mathrm{CH}^*(BG)$ is the Chow ring of algebraic cycles of the classifying space BG introduced by Totaro [To99], see also [MV99, Section 4] for a motivic homotopy point of view, and it hosts the universal characteristic classes of principal homogeneous spaces (torsors) under G . This ring is in general rather hard to compute even for a split simple group except the special cases of SL_n and Sp_{2n} , see [To99, Pa98, MRV06, KM22] for the known answers.

The ring $\mathrm{CH}^*(G)$ is the Chow ring of algebraic cycles of G itself, considered as an algebraic variety forgetting the multiplicative structure. The problem of computing $\mathrm{CH}^*(G)$ for a split semisimple group G , and the closely related problem of computing the singular cohomology of a compact Lie group attracted a lot of attention in 1960s and was done case by case in the works of Miller, Borel, Baum, Browder, Araki, Shikata, Ishitoya, Kono, Toda and others, culminating with the celebrated article [Kac85] of Victor Kac, who summarized the previously known computations and gave a uniform conceptual answer for $\mathrm{CH}^*(G)$ assuming the coefficients to be a finite field (see [Kac85, Theorem 6] and especially [Kac85, Table 2]).

The computations by Victor Kac were used 20 years later by Geldhauser, Petrov, and Zainoulline to establish the structure of Chow motives of generically split twisted flag varieties and to introduce a discrete motivic invariant of algebraic groups of inner type, called the J -invariant (see [PSZ08, PS10, PS12]). In the case of quadratic forms an equivalent invariant was introduced previously by Vishik in [Vi05]. The J -invariant was an important tool to solve several long-standing problems, for example, it played an important role in the progress on the Kaplansky problem about possible values of the u -invariant of fields by Vishik [Vi07] and in the solution of a problem of Serre about groups of type E_8 and its finite subgroups [S16, GS10]. Besides, Garibaldi, Geldhauser, and Petrov used the J -invariant to relate the rationality of some parabolic subgroups of groups of type E_7 with the Rost invariant, proving a conjecture of Rost and solving a question of Springer in [GPS16]. The concept of the J -invariant is built up on the first three columns of [Kac85, Table 2] encoding the Chow rings of split semisimple algebraic groups. In fact, an important step in the construction of the J -invariant was an observation that the parameters in the Kac’s table coincide in certain cases with the parameters related to the generalized Rost motives, which were used by Rost and Voevodsky in the proofs of the Milnor and Bloch–Kato conjectures. The last column of Kac’s table also has a motivic interpretation in terms of G -equivariant motives given in [PS17].

In order to generalize the J -invariant to arbitrary linear algebraic groups (i.e. possibly of outer type) one has to study the Chow rings of non-split quasi-split semisimple groups. Progress in this direction was recently obtained in [GZ22]. However, there are several substantial difficulties when dealing with non-split quasi-split groups. First of all, quasi-split groups G are equipped with a finite Galois field extension K/k , which is the minimal splitting field of G , i.e., come together with some arithmetic information. A priori it is not clear whether their Chow rings depend on K , and we show in the current article that it is not the case. Besides, contrary to the split situation, our computations show that the Chow rings of non-split quasi-split groups do not have, in general, the structure of Hopf algebras. For example, the lacunary relation appearing additionally to the nilpotency of generators in the non-split quasi-split adjoint group of type ${}^2D_{2r}$ (case 2 with $l = 2$ of Theorem 1 below) is not possible in any Hopf algebra over a finite field. Moreover, contrary to the split situation, the characteristic sequence of Grothendieck (see [Gro58, Remark 2°]), which is a useful tool, is not right exact in general for quasi-split groups.

To overcome these difficulties we develop a theory of equivariant conormed Chow rings and apply it to quasi-split groups. The non-equivariant version of these, i.e. Chow rings modulo pushforwards (norms) coming from a fixed field extension, have already been used in the study of varieties homogeneous under an action of a quasi-split group in e.g. [Ka12, KaZ13, KM22] and were studied explicitly in [Fi19, GZ22]. We would also like to note that the category of conormed Chow motives shares some properties with the category of isotropic motives introduced and studied by Vishik [Vi24], in particular, in the latter category

motives of anisotropic varieties are zero while in the former category the motive of an irreducible k -variety is zero when the variety becomes reducible over the (Galois) field extension K/k .

The main result of this article is the following theorem:

Theorem 1. *Let G be a quasi-split geometrically almost simple group over a field k . For a field \mathbb{F} put $p := \text{char } \mathbb{F}$ and $\text{Ch}^*(-) := \text{CH}^*(-) \otimes \mathbb{F}$. Then, depending on the type $\Delta(G)$, the fundamental group $\pi_1(G)$ (see Sections A.1 and A.2 for the notation) and p the following holds.*

1. *If the triple $(\Delta(G), \pi_1(G), p)$ is in the Table 1, then*

$$\text{Ch}^*(G) \cong \mathbb{F}[e_1, e_2, \dots, e_s] / (e_1^{p^{k_1}}, e_2^{p^{k_2}}, \dots, e_s^{p^{k_s}}), \quad \deg e_i = d_i,$$

with the parameters s, d_i, k_i given in the table.

Table 1: $\text{Ch}^*(G)$ for a quasi-split simple algebraic group G

$\Delta(G)$	$\pi_1(G)$	p	s	$d_i, i = 1, 2, \dots, s$	$k_i, i = 1, 2, \dots, s$
A_n	$\mu_l, l \mid (n+1)$	$p \mid l$	1	1	$v_p(n+1)$
${}^2A_n, n \geq 2$	$2\mu_l, l \mid (n+1), l \text{ is odd}$	2	$\lfloor \frac{n}{2} \rfloor$	$2i+1$	1
${}^2A_{2r-1}, r \geq 2$	$2\mu_{2m}, m \mid r, m \text{ is odd}$	2	r	$2i-1$	$v_2(r)+1, i=1$ $1, i \geq 2$
${}^2A_{2r-1}, r \geq 2$	$2\mu_{2m}, m \mid r, m \text{ is even}$	2	$r+1$	$2, i=1$ $2i-3, i \geq 2$	$v_2(r), i=1$ $1, i \geq 2$
B_n	1	2	$\lfloor \frac{n-1}{2} \rfloor$	$2i+1$	$\lceil \log_2 \frac{2n}{2i+1} \rceil$
	μ_2	2	$\lfloor \frac{n+1}{2} \rfloor$	$2i-1$	$\lceil \log_2 \frac{2n}{2i-1} \rceil$
C_n	μ_2	2	1	1	$v_2(n)+1$
$D_n, n \geq 3$	1	2	$\lfloor \frac{n}{2} \rfloor - 1$	$2i+1$	$\lceil \log_2 \frac{2n-1}{2i+1} \rceil$
	μ_2^{so}	2	$\lfloor \frac{n}{2} \rfloor$	$2i-1$	$\lceil \log_2 \frac{2n-1}{2i-1} \rceil$
	μ_4 or $\mu_2 \times \mu_2$	2	$\lfloor \frac{n}{2} \rfloor + 1$	$1, i=1$ $2i-3, i \geq 2$	$v_2(n), i=1$ $\lceil \log_2 \frac{2n-1}{2i-3} \rceil, i \geq 2$
$D_{2r}, r \geq 2$	μ_2^{hs}	2	r	$1, i=1$ $2i-1, i \geq 2$	$v_2(r)+1, i=1$ $\lceil \log_2 \frac{4r-1}{2i-1} \rceil, i \geq 2$
${}^2D_n, n \geq 3$	1	2	$\lfloor \frac{n+1}{2} \rfloor - 1$	$2i+1$	$\lceil \log_2 \frac{2n}{2i+1} \rceil$
	μ_2	2	$\lfloor \frac{n+1}{2} \rfloor$	$1, i=1$ $2i-1, i \geq 2$	$\lceil \log_2 n \rceil + 1$ $\lceil \log_2 \frac{2n}{2i-1} \rceil$
${}^2D_{2r+1}, r \geq 1$	$2\mu_4$	2	$r+2$	$1, i=1$	$1, i=1$
				$2, i=2$	$\lceil \log_2(2r+1) \rceil, i=2$
				$2i-3, i \geq 3$	$\lceil \log_2 \frac{4r+2}{2i-3} \rceil, i \geq 3$
${}^3D_4, {}^6D_4$	1	2	1	3	1
	$1, {}^3\mu_{2,2}, {}^6\mu_{2,2}$	3	1	4	1
E_6	$1, \mu_3$	2	1	3	1

2E_6	1	3	1	4	1
	μ_3	3	2	1, 4	2, 1
	$1, {}^2\mu_3$	2	3	3, 5, 9	1, 1, 1
	1	3	1	4	1
E_7	1	2	3	3, 5, 9	1, 1, 1
	μ_2	2	4	1, 3, 5, 9	1, 1, 1, 1
	$1, \mu_2$	3	1	4	1
E_8	1	2	4	3, 5, 9, 15	3, 2, 1, 1
		3	2	4, 10	1, 1
		5	1	6	1
F_4	1	2	1	3	1
		3	1	4	1
G_2	1	2	1	3	1

Here and below v_p is the p -adic valuation.

2. If $(\Delta(G), \pi_1(G), p) = ({}^2D_{2r}, {}^2\mu_{2,2}, 2)$, then

$$\mathrm{Ch}^*(G) \cong \begin{cases} \mathbb{F}[e_1, \dots, e_s] / (e_1^{2^{k_1}}, \dots, e_s^{2^{k_s}}, e_1 \cdot (\sum_{j=0}^{k_1-1} e_1^{2^{k_1}-2^j+1} \cdot e_2^{2^j})), & l = 2, \\ \mathbb{F}[e_1, \dots, e_s] / (e_1^{2^{k_1}}, \dots, e_s^{2^{k_s}}, e_1 \cdot e_l^{2^{k_l-1}}), & l \geq 3, \end{cases}$$

where $l \in \mathbb{N}$ is such that $2r = 2^m(2l - 3)$ and

$$s = r + 1, \deg e_1 = 1, \deg e_2 = 2, k_1 = v_2(2r), k_2 = \lceil \log_2 2r \rceil,$$

$$\deg e_i = 2i - 3, k_i = \left\lceil \log_2 \frac{4r}{2i - 3} \right\rceil, 3 \leq i \leq s.$$

3. If $(\Delta(G), \pi_1(G), p)$ is in the table below, then $\mathrm{Ch}^*(G)$ is as in the table.

$\Delta(G)$	$\pi_1(G)$	p	$\mathrm{Ch}^*(G)$	$\deg e_i$
2A_n	${}^2\mu_l, l \mid n + 1$	$2 \neq p \mid l$	$\mathbb{F}[e_1] / (e_1^m), m = \frac{p^{v_p(n+1)} + 1}{2}$	2
3D_4	${}^3\mu_{2,2}$	2	$\mathbb{F}[e_1, e_2, e_3, e_4] / (e_1^4, e_2^2, e_3^2, e_4^2, e_1e_2, e_1e_3, e_1^3 + e_2e_3)$	2, 3, 3, 3
6D_4	${}^6\mu_{2,2}$	2	$\mathbb{F}[e_1, e_2, e_3] / (e_1^4, e_2^2, e_1e_2, e_3^2)$	2, 3, 3
2E_6	${}^2\mu_3$	3	$\mathbb{F}[e_1, e_2] / (e_1^5, e_2^3)$	1, 4

4. For all the remaining cases $(\Delta(G), \pi_1(G), p)$ one has $\mathrm{Ch}^*(G) \cong \mathbb{F}$.

It is clear that one can assume \mathbb{F} to be a prime field, that is \mathbb{Q} or a finite field \mathbb{F}_p . If $\mathbb{F} = \mathbb{Q}$, then it is well-known and goes back to [Gro58, Remark 2 on p. 21] that $\mathrm{Ch}^*(G) \cong \mathbb{Q}$ (see also Section A.5.1 for the details). The case of a split group G , i.e. $\Delta(G) = A_n, B_n, C_n, D_n, E_6, E_7, E_8, F_4$, or G_2 , is also classical [Kac85, Theorem 6] (see also Section A.5.2 for a recollection). All the remaining cases, that is when G is a non-split quasi-split simple group, are, to the best of our knowledge, completely new. Theorem 1 is

obtained as a combination of the above classical results and Theorems 3.2.2, 3.3.3, 3.4.3, and 3.5.1 of the present article.

Outline of the proof. We assume \mathbb{F} to be a prime field. Furthermore, as it was said above, using the classical results (see Section A.5 for references) we can assume that G is a non-split quasi-split geometrically almost simple group. Let K/k be the splitting field of G . The cases are naturally divided into three groups: (1) $p = [K : k]$, (2) p is coprime to $[K : k]$ and (3) $\Delta(G) = {}^6D_4$ with $p = 2$ or 3 . The first case is the most interesting one, while the latter two are rather straightforward.

Case (1), $p = [K : k]$. Let $\pi: G_K \rightarrow G$ be the projection and consider the exact sequence

$$\mathrm{Ch}^*(G_K) \xrightarrow{\pi_*} \mathrm{Ch}^*(G) \xrightarrow{q} \mathrm{CH}_K^*(G) \rightarrow 0 \quad (*)$$

with the last algebra being the cokernel of π_* . First we analyse the algebra $\mathrm{CH}_K^*(G)$, which fits into the setting of *conormed Chow groups* $\mathrm{CH}_K^*(-)$, i.e. Chow groups modulo the image of the norm map. Such groups proved to be a valuable invariant in the study of quasi-split homogeneous varieties and implicitly occurred in e.g. [Ka12, KaZ13, KM22] and were studied explicitly in [Fi19, GZ22]. As it was shown in [GZ22] one can define $\mathrm{CH}_K^*(-)$ for an arbitrary separable field extension K/k (see Definition 2.1.1), yielding an oriented cohomology theory on $\mathcal{S}m_k$ in the sense of [LM07] whose basic properties we study in Section 2.1. The most important property is that $\mathrm{CH}_K^*(X) = 0$ for a connected variety X that becomes disconnected over an intermediate Galois extension $K/L/k$ (see Corollary 2.1.7). This, in particular, means that Artin–Tate motives (motives of quasi-split projective homogeneous varieties are among the examples [CM06, Lemma 29]) behave like Tate motives from the conormed point of view.

Following the approach by Totaro [To99] and Edidin–Graham [EG98] for the ordinary Chow groups we extend the conormed Chow groups to the equivariant setting (Section 2.2) defining $\mathrm{CH}_{K,T}^*(X)$ for an algebraic group T over k and a T -variety X . If T is a torus which is (a) *quasi-trivial* (also known as *induced*), i.e. $T \cong R_{L_1/k}\mathbb{G}_m \times \dots \times R_{L_r/k}\mathbb{G}_m$ for Weil restrictions of split tori, and such that (b) T_K is split, then $\mathrm{CH}_{K,T}^*(-)$ behave very much like equivariant Chow groups for split tori, in particular, one can compute $\mathrm{CH}_{K,T}^*(X)$ for a smooth variety X with the trivial T -action (see Proposition 2.3.2) as

$$\mathrm{CH}_{K,T}^*(X) \cong \mathrm{CH}_K^*(X)[b_1, b_2, \dots, b_r], \quad \deg b_i = [L_i : k],$$

and for a smooth variety X with an arbitrary T -action one has an isomorphism (Corollary 2.3.3)

$$\mathrm{CH}_{K,T}^*(X) \otimes_{\mathrm{CH}_{K,T}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(X). \quad (**)$$

A similar result was also obtained in [Sal22, Theorem 3.1], but in a slightly different language.

Now let $T \leq B \leq G$ be a maximal torus and a Borel subgroup, and K be the splitting field of G as before. If T is quasi-trivial, which is the case if G is simply connected or adjoint (Lemma A.2.3), then the isomorphism (**) for $X := G$ yields that the characteristic sequence (Definition 2.5.5)

$$\mathrm{CH}_{K,T}^*(\mathrm{Spec} k) \xrightarrow{c} \mathrm{CH}_K^*(G/B) \xrightarrow{\phi^*} \mathrm{CH}_K^*(G) \rightarrow 0$$

is an exact sequence of graded algebras in the sense that ϕ^* is surjective and $c(\mathrm{CH}_{K,T}^{>0}(\mathrm{Spec} k))$ generates the kernel of ϕ^* as an ideal. This is the conormed version of the classical characteristic sequence from [Gro58] which since its introduction has been one of the main tools for the computations related to the Chow rings of split reductive groups. The cokernel of the conormed characteristic map

$$\mathcal{C}_K^*(G) := \mathrm{CH}_K^*(G/B) / c(\mathrm{CH}_{K,T}^{>0}(\mathrm{Spec} k)) \cdot \mathrm{CH}_K^*(G/B)$$

was computed in the adjoint case in [GZ22, Section 8]. This gives an answer for $\mathrm{CH}_K^*(G)$ in the adjoint case, since the exactness of the characteristic sequence yields $\mathrm{CH}_K^*(G) \cong \mathcal{C}_K^*(G)$.

If G is neither adjoint nor simply connected, then in general the characteristic sequence is not exact, since ϕ^* fails to be surjective already for $\mathrm{CH}_K^1(-)$, i.e. for the conormed Picard groups. In order to circumvent this issue we construct an *extended characteristic sequence* in the following way. Let $\tilde{G} \rightarrow G$ be the simply connected cover and $\tilde{T} \leq \tilde{B} \leq \tilde{G}$ be the maximal torus and the Borel subgroup over $T \leq B$. Then \tilde{T} acts on G with $\pi_1(G)$ being the kernel of the action and we have isomorphisms

$$\mathrm{CH}_{K,\tilde{T}}^*(G) \cong \mathrm{CH}_{K,\pi_1(G)}^*(\tilde{G}/\tilde{T}) \cong \mathrm{CH}_K^*(\tilde{G}/\tilde{B}) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k),$$

where the first one arises from the span of torsors $G \leftarrow \tilde{G} \rightarrow \tilde{G}/\tilde{T}$ and the second one is a combination of a Künneth property (Lemma 2.5.1) and homotopy invariance for $\tilde{G}/\tilde{T} \rightarrow \tilde{G}/\tilde{B}$. Using this and the isomorphism (**) for $X := G$ and the quasi-trivial torus \tilde{T} we obtain an extended characteristic sequence

$$\mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \xrightarrow{\hat{c}} \mathrm{CH}_K^*(\tilde{G}/\tilde{B}) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_K^*(G) \rightarrow 0$$

which is an exact sequence of graded algebras (Theorem 2.5.6). The algebras $\mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k)$ arising here can be computed (Proposition 2.4.1) using resolutions by quasi-trivial tori, yielding an explicit presentation of $\mathrm{CH}_K^*(G)$ as a quotient of $\mathrm{CH}_K^*(\tilde{G}/\tilde{B})$ or $\mathrm{CH}_K^*(\tilde{G}/\tilde{B})[x]$ (Theorem 2.6.3). In particular, this gives a precise relation between $\mathrm{CH}_K^*(G)$ and already known $\mathrm{CH}_K^*(G)$ for the adjoint quotient $G \rightarrow \tilde{G}$, allowing us to compute $\mathrm{CH}_K^*(G)$ (Theorem 2.8.2).

We also give a presentation of $\mathrm{CH}_K^*(G)$ as a quotient of $\mathcal{C}_K^*(G)[x]$ for the cokernel $\mathcal{C}_K^*(G)$ of the ordinary characteristic map (Theorem 2.7.2), and use it to compute $\mathcal{C}_K^*(G)$ explicitly (Theorem 2.9.1). Moreover, as a by-product of the isomorphism (**) we show that G enjoys the conormed Künneth property (Proposition 2.5.3), endowing $\mathrm{CH}_K^*(G)$ with a natural structure of a Hopf algebra, which in turn explains the structure of the answer for $\mathrm{CH}_K^*(G)$ obtained in Theorem 2.8.2. Note that in general the Künneth homomorphism

$$\mathrm{CH}^*(X) \otimes \mathrm{CH}^*(G) \rightarrow \mathrm{CH}^*(X \times G)$$

fails to be an isomorphism for a quasi-split group G , which could be seen from Table 1 taking $X := \mathrm{Spec} K$ with K being the splitting field.

Having computed $\mathrm{CH}_K^*(G)$ in the sequence (*) we turn to the study of π_* . It turns out (Proposition 3.3.2) that $\pi_* = 0$ for a simple group G except when G is an adjoint group of type $\Delta(G) = {}^2\mathrm{D}_{2r}$. This is proved on a case-by-case basis. The case of $\Delta(G) = {}^3\mathrm{D}_4$ is immediate, since the algebra $\mathrm{CH}^*(G_K)$ is trivial. The case of $\Delta(G) = {}^2\mathrm{A}_n$ is also straightforward, since $\mathrm{CH}^*(G_K)$ is rather simple. For $\Delta(G) = {}^2\mathrm{D}_n$ we compute the algebra

$$\mathrm{CH}^*(\mathrm{Gr}(n-1; q)) \cong \mathrm{CH}^*(\mathrm{Gr}(n-1; q)_K)^{\mathrm{Gal}(K/k)}$$

for a quasi-split submaximal isotropic Grassmannian (Proposition 3.1.1) using the answer for the ring $\mathrm{CH}^*(\mathrm{Gr}(n-1; q)_K)$ given by [Vi07] and [EKM08, Section 86] and calculating explicitly the action of the Galois group. Since the pullback

$$\mathrm{CH}^*(\mathrm{Gr}(n-1; q)_K) \rightarrow \mathrm{CH}^*(G_K)$$

is well-known to be surjective, this allows us to show that $\pi_* = 0$ when $\Delta(G) = {}^2\mathrm{D}_n$ and G is not an adjoint group of type $\Delta(G) = {}^2\mathrm{D}_{2r}$. In the last case $\Delta(G) = {}^2\mathrm{E}_6$ we have $\mathrm{Ch}^*(G_K) \cong \mathbb{F}_2[e_1]/(e_1^2)$ with $\deg e = 3$ by [Kac85, Theorem 6], so we need only to show $\pi_*(e_1) = 0$. For this we lift the problem to G/B , showing that for an explicit Schubert cycle $Z \in \mathrm{Ch}^3((G/B)_K)$ which goes to e_1 on G_K its pushforward $\rho_*(Z) \in \mathrm{Ch}^*(G/B)$ is in the image of the characteristic map. This is done via explicit computer-assisted calculations with Schubert cycles.

The vanishing of π_* in sequence (*) yields an answer for $\mathrm{CH}^*(G)$ (Theorem 3.3.3), since we already know $\mathrm{CH}_K^*(G)$. In the last remaining case of the adjoint group of type $\Delta(G) = {}^2\mathrm{D}_{2r}$ a more detailed analysis of $\mathrm{CH}^*(\mathrm{Gr}(2r-1; q))$ gives an answer for $\mathrm{CH}^*(G)$ (Theorem 3.2.2), with a crucial ingredient being $c_1(\mathcal{L}(\bar{\omega}_{2r-1} + \bar{\omega}_{2r}))^{2v_2(2r)} = 0$ for the first Chern class as in the split case (Proposition 3.1.1, see also Section A.3.3 for the notation).

Case (2), p is coprime to $[K : k]$. In this case we show that $\mathrm{Ch}^*(G) = \mathrm{Ch}^*(G_K)^{\mathrm{Gal}(K/k)}$ (Lemma 3.4.2). If $\Delta(G) \neq {}^2\mathrm{E}_6$, then a straightforward description of the $\mathrm{Gal}(K/k)$ -action on $\mathrm{Pic}(G_K)$ provides enough information to compute $\mathrm{Ch}^*(G_K)^{\mathrm{Gal}(K/k)}$, while in the case of $\Delta(G) = {}^2\mathrm{E}_6$ we again resort to explicit computer-assisted computations with Schubert cycles (Theorem 3.4.3).

Case (3), $\Delta(G) = {}^6\mathrm{D}_4$ with $p = 2$ or 3 (Theorem 3.5.1). This case is treated via a mixture of the methods used in the first two cases. In order to compute $\mathrm{Ch}^*(G)$ for $p = 2$ we partially split the group via a degree 3 subextension $K/L/k$, obtaining $\Delta(G_L) = {}^2\mathrm{D}_4$. We already know $\mathrm{Ch}^*(G_L)$ by the case (1), and we further analyse which cycles descent to $\mathrm{Ch}^*(G)$ computing the $\mathrm{Gal}(K/k)$ -action on $\mathrm{Ch}^*(G_K)$ by computer-assisted calculations with Schubert cycles. The case of $p = 3$ is treated similarly, using the degree

2 subextension $K/L/k$ such that $\Delta(G_L) = {}^3D_4$ and again computer-assisted calculations with Schubert cycles.

Acknowledgements. The work of the authors is supported by the DFG research grant AN 1545/4-1, the work of the first author is additionally supported by DFG Heisenberg grant AN 1545/1-1 and the work of the second author is additionally supported by DFG research grant SE 1721/4-1. The authors would like to thank Philippe Gille, Fabien Morel, Andrei Lavrenov, Burt Totaro, Alois Wohlschlager, Maksim Zhykhovich and Egor Zolotarev for valuable discussions.

Notation. Throughout the article we employ the following assumptions and notations.

- A *variety* over a field k is a reduced scheme separated and of finite type over k .
- An *algebraic group* over a field k is an algebraic group in the sense of [Mi17, Definition 1.1], i.e. a group object in the category of schemes of finite type over k , in particular, an algebraic group is not necessary reduced.
- A *group action* is a left group action. At the same time, somehow inconsistently, for a group G acting on X we denote by X/G the quotient.
- A *simple group* is a geometrically almost-simple group in the sense of [Mi17, Definition 19.7].
- By *the splitting field* of a group G we always mean the minimal splitting field. Note that the minimal splitting field is always a Galois extension.
- We use some standard terminology and notation related to quasi-split algebraic groups and algebraic groups in general, and also to vector bundles on (projective) homogeneous varieties, see Appendix A for a partial recollection.

v_p	p -adic valuation
k	a field
k^{sep}	separable closure of k
\mathbb{F}_p	finite field with p elements
$\text{Ch}^*(-)$	$\text{CH}^*(-) \otimes \mathbb{F}$ for a field \mathbb{F}
Sm_k	the category of smooth varieties over k
X_F	$X \times_{\text{Spec } k} \text{Spec } F$ for a field extension F/k and $X \in \text{Sm}_k$
$\mathcal{X}^*(S)$	the $\text{Gal}(k^{sep}/k)$ -module $\text{Hom}(S_{k^{sep}}, \mathbb{G}_m)$ of characters of an algebraic group S over k
$\pi_1(G)$	the fundamental group of G
$\Delta(G)$	Dynkin diagram of a quasi-split semisimple group G with the action of $\text{Gal}(k^{sep}/k)$
μ_l	the algebraic group of l -th roots of unity
${}^2\mu_l$	a non-trivial (if $l \geq 3$) form of μ_l split by a quadratic Galois extension
${}^2\mu_{2,2}, {}^3\mu_{2,2}$	a non-trivial form of $\mu_2 \times \mu_2$ split by a degree 2 (degree 3) Galois extension
${}^6\mu_{2,2}$	a form of $\mu_2 \times \mu_2$ with the splitting field K/k such that $\text{Gal}(K/k) \cong S_3$
Sm_k^T	the category of smooth T -varieties over k for an algebraic group T over k
$\text{CH}_{K,T}^*(X)$	equivariant conormed Chow ring of $X \in \text{Sm}_k^T$ (Definition 2.2.1)
$\Phi_\rho, \Phi_T, \Phi_T^c$	restriction homomorphisms on equivariant conormed Chow rings (Definition 2.2.3)
Res_ρ	restriction homomorphism $\text{Res}_\rho: \text{Rep}(T) \rightarrow \text{Rep}(S)$ on the representation rings for a homomorphism $\rho: S \rightarrow T$ of algebraic groups over a field k

2 Conormed Chow ring of a quasi-split simple group

In this part of the article we develop the theory of *conormed Chow rings* and its equivariant version, compute some equivariant conormed Chow rings of a point and compute the conormed Chow rings of quasi-split simple groups. Conormed Chow rings form an oriented cohomology theory, which is especially well adapted to the study of cohomological properties of quasi-split algebraic groups and their homogeneous varieties. It turns out that from the conormed point of view many non-split algebraic groups (e.g. quasi-trivial tori, twisted forms of μ_l and others) behave very much like their split counterparts.

2.1 Conormed Chow groups

In this section we recall the definition of conormed Chow rings and explore their basic properties. In particular, we show that for a Galois field extension K/k the conormed Chow group $\mathrm{CH}_K^*(-)$ is generated by irreducible cycles that remain irreducible over K .

Definition 2.1.1 (see also [GZ22, § 5] and [Fi19, § 2]). Let K/k be a separable extension of fields. For $X \in \mathcal{S}m_k$ the *conormed Chow ring of X* is

$$\mathrm{CH}_K^*(X) := \mathrm{CH}^*(X) \Big/ \mathrm{Im} \left(\bigoplus_{\substack{k \subseteq F \subseteq K \\ F/k \text{ finite}}} \mathrm{CH}^*(X_F) \xrightarrow{(\pi_{F/k})_*} \mathrm{CH}^*(X) \right),$$

where $\mathrm{CH}^*(X)$ is the Chow ring of X , $\pi_{F/k}: X_F \rightarrow X$ is the projection and $(\pi_{F/k})_*$ is the pushforward homomorphism. By the projection formula the above image is an ideal in $\mathrm{CH}^*(X)$. This ideal is referred to as the *norm ideal*. Note that in the case of $K = k$ one has $\mathrm{CH}_K^*(X) = \mathrm{CH}^*(X)$, since there are no nontrivial intermediate field extensions.

One can easily check that since the Chow rings form an oriented cohomology theory on $\mathcal{S}m_k$ in the sense of [LM07, Definition 1.1.2], the conormed Chow rings $\mathrm{CH}_K^*(-)$ for a given separable extension of fields K/k also form an oriented cohomology theory on $\mathcal{S}m_k$. In particular, following [Ma68] (see also [EKM08, Chapter 12]), for a separable extension of fields K/k one can consider the category of conormed Chow motives \mathcal{M}_K built from the category of conormed correspondences, i.e. from the category with the objects given by smooth proper varieties over k and the morphisms between irreducible proper varieties given by $\mathrm{CH}_K^{\dim Y}(X \times Y)$. The transformation of oriented cohomology theories $\mathrm{CH}^*(-) \rightarrow \mathrm{CH}_K^*(-)$ induces a functor $\mathcal{M} \rightarrow \mathcal{M}_K$ from the category of Chow motives to the category of conormed Chow motives.

Lemma 2.1.2. *Let K/k be a separable field extension. Suppose that there exists an intermediate field extension $K \supseteq L \supseteq k$ with L/k being finite Galois. Then the following holds.*

- If $[L : k]$ is not a power of a prime, then $\mathrm{CH}_K^*(\mathrm{Spec} k) = 0$.
- If $[L : k] = p^n$ for a prime p and for every intermediate extension $K/F/k$ the degree $[F : k]$ is either infinite or a power of p , then $\mathrm{CH}_K^*(\mathrm{Spec} k) \cong \mathbb{F}_p$.

Proof. It is straightforward to see from the definition that $\mathrm{CH}_K^*(\mathrm{Spec} k) \cong \mathbb{Z}/m\mathbb{Z}$ with

$$m := m(K, k) := \gcd_{\substack{k \subseteq F \subseteq K \\ F/k \text{ finite}}} ([F : k]).$$

Put $\mathcal{G} := \mathrm{Gal}(L/k)$. Suppose that $|\mathcal{G}| = [L : k] = l$ is not a power of a prime. For $p \mid l$ consider a Sylow p -subgroup $\mathcal{G}'_p \leq \mathcal{G}$, then for $F := L^{\mathcal{G}'_p}$ one has $[F : k] = l'_p$ for the prime to p part l'_p of l . Since the set of all l'_p is coprime, it follows that $m(K, k) = 1$ and $\mathrm{CH}_K^*(\mathrm{Spec} k) = 0$.

If $|\mathcal{G}| = p^n$ for a prime $p \in \mathbb{N}$, then there exists a subgroup $\mathcal{G}' \leq \mathcal{G}$ such that $[\mathcal{G} : \mathcal{G}'] = p$, thus there is an intermediate field extension $F := L^{\mathcal{G}'}$ such that $[F : k] = p$, yielding $m(K, k) \mid p$. On the other hand, it follows from the assumptions that $p \mid m(K, k)$, thus $m(K, k) = p$ and $\mathrm{CH}_K^*(\mathrm{Spec} k) \cong \mathbb{F}_p$. \square

Remark 2.1.3. If K/k has no intermediate extensions $k \subsetneq L \subseteq K$ with L/k being finite Galois, then $\mathrm{CH}_K^*(\mathrm{Spec} k)$ may fail to be a field, e.g. for a separable extension K with $[K : k] = n$ and without nontrivial intermediate extensions one has $\mathrm{CH}_K^*(\mathrm{Spec} k) \cong \mathbb{Z}/n\mathbb{Z}$.

Lemma 2.1.4. *Let $K/F/k$ be separable field extensions with F/k being finite. Then for $X \in \mathcal{S}m_F$ viewed as a variety over k one has $\mathrm{CH}_K^*(X) = 0$.*

Proof. The projection onto the first factor

$$\mathrm{Spec} F \times_{\mathrm{Spec} k} \mathrm{Spec} F \rightarrow \mathrm{Spec} F$$

has a section. Taking the base change along the morphism $X \rightarrow \text{Spec } F$ we see that the projection

$$X_F = X \times_{\text{Spec } k} \text{Spec } F \xrightarrow{\pi_{F/k}} X$$

also has a section. Let $s: X \rightarrow X_F$ be such section which means that s satisfies $\pi_{F/k} \circ s = \text{id}_X$. Then

$$1 = (\text{id}_X)_* 1 = (\pi_{F/k})_*(s_* 1) \in (\pi_{F/k})_*(\text{CH}^*(X_F)) \subseteq \text{CH}^*(X)$$

and the norm ideal coincides with $\text{CH}^*(X)$. The claim follows. \square

Lemma 2.1.5. *Let X be a normal irreducible variety over a field k and L/k be a finite Galois field extension. Suppose that X_L is reducible. Then there exists an intermediate field extension $L \supseteq F \supsetneq k$ such that the structure morphism $X \rightarrow \text{Spec } k$ factors as $X \rightarrow \text{Spec } F \rightarrow \text{Spec } k$.*

Proof. Let k' be the separable closure of k in the function field $k(X)$. Since X is normal, it follows that for every open affine subscheme $\text{Spec } A \subseteq X$ we have $k' \subseteq A \subseteq k(X)$, thus $k' \subseteq H^0(X, \mathcal{O}_X)$ yielding a factorization $X \rightarrow \text{Spec } k' \rightarrow \text{Spec } k$.

It suffices to show that there exists a subfield $k' \supseteq F \supsetneq k$ isomorphic to a subfield of L . Suppose that no such F exists. We claim that then $k' \otimes_k L$ is a field. Choose embeddings $k' \subseteq k^{\text{sep}}$ and $L \subseteq k^{\text{sep}}$, and put

$$\mathcal{G} := \text{Gal}(k^{\text{sep}}/k), \quad \mathcal{G}_L := \text{Gal}(k^{\text{sep}}/L), \quad \mathcal{G}_{k'} := \text{Gal}(k^{\text{sep}}/k'), \quad \mathcal{G}_{Lk'} := \text{Gal}(k^{\text{sep}}/Lk'),$$

where Lk' is the compositum of L and k' inside k^{sep} . These groups have the following properties:

- \mathcal{G}_L is a normal subgroup of \mathcal{G} , since L/k is Galois,
- the subgroup $\langle \mathcal{G}_L, \mathcal{G}_{k'} \rangle \leq \mathcal{G}$ generated by \mathcal{G}_L and $\mathcal{G}_{k'}$ is \mathcal{G} , since k' and L have no common subfields except k ,
- $\mathcal{G}_{Lk'} = \mathcal{G}_L \cap \mathcal{G}_{k'}$ since Lk' is the compositum of L and k' inside k^{sep} .

The first two properties yield that $\mathcal{G} = \mathcal{G}_{k'} \cdot \mathcal{G}_L$, thus the monomorphism

$$\mathcal{G}_{k'}/(\mathcal{G}_L \cap \mathcal{G}_{k'}) \rightarrow \mathcal{G}/\mathcal{G}_L$$

induced by the embedding is an isomorphism. Note that k'/k is an extension of finite degree, being an algebraic subextension of a finitely generated extension $k(X)/k$. Thus we have

$$[Lk' : k] = [\mathcal{G} : \mathcal{G}_{Lk'}] = [\mathcal{G} : \mathcal{G}_L \cap \mathcal{G}_{k'}] = [\mathcal{G} : \mathcal{G}_{k'}] \cdot [\mathcal{G}_{k'} : \mathcal{G}_L \cap \mathcal{G}_{k'}] = [\mathcal{G} : \mathcal{G}_{k'}] \cdot [\mathcal{G} : \mathcal{G}_L] = [k' : k] \cdot [L : k].$$

By the dimension count it follows that the homomorphism $k' \otimes_k L \rightarrow Lk'$ is an isomorphism, and $k' \otimes_k L$ is a field. Then $k(X) \otimes_k L \cong k(X) \otimes_{k'} (k' \otimes_k L)$ is also a field, since $k(X)$ is primary over k' (the algebraic closure of k' in $k(X)$ is purely inseparable over k') and $k' \otimes_k L$ is separable over k' . Thus X_L is irreducible which contradicts the assumptions. \square

Remark 2.1.6. Some normality assumptions on X and L/k in Lemma 2.1.5 are indeed necessary, as the following examples show:

- Let $X := \text{Spec } \mathbb{R}[x, y]/(x^2 + y^2)$ and $L/k = \mathbb{C}/\mathbb{R}$. Then X is an irreducible variety over \mathbb{R} and $X_{\mathbb{C}}$ is reducible being the union of two lines, but the structure morphism $X \rightarrow \text{Spec } \mathbb{R}$ does not factor through $\text{Spec } \mathbb{C}$, since X has a closed point with the residue field \mathbb{R} .
- Let K/k be a Galois field extension with the Galois group being $\text{Gal}(K/k) = S_4$, and $L_1, L_2 \subseteq K$ be the subfields with $\text{Gal}(K/L_1) = \langle (12) \rangle$ and $\text{Gal}(K/L_2) = \langle (1234) \rangle$. Put $X := \text{Spec } L_1$ and $L := L_2$. For the compositum $L_1 L_2$ we have

$$\dim_k L_1 \otimes_k L_2 = 12 \cdot 6 > 24 = [K : k] \geq [L_1 L_2 : k],$$

hence $L_1 \otimes_k L_2$ is not a field and X_L is not connected. Suppose that $X \rightarrow \text{Spec } k$ factors through $\text{Spec } F \rightarrow \text{Spec } k$ for a subfield $k \subseteq F \subseteq L = L_2$. Then F is isomorphic to a subfield of L_1 and there exists $\sigma \in \text{Gal}(K/k)$ such that $F \subseteq \sigma(L_1)$. We have $\text{Gal}(K/\sigma(L_1)) \cong C_2$ with the generator being the transposition $\sigma(12)\sigma^{-1}$. A transposition and a 4-cycle generate S_4 yielding $\langle \text{Gal}(K/\sigma(L_1)), \text{Gal}(K/L_2) \rangle = \text{Gal}(K/k)$. It follows that $\sigma(L_1) \cap L_2 = k$ and $F = k$.

Corollary 2.1.7. *Let $X \in \mathcal{S}m_k$ and $K/L/k$ be separable field extensions with L/k being a finite Galois extension. Suppose that X is connected and X_L is not connected. Then $\mathrm{CH}_K^*(X) = 0$.*

Proof. Follows from Lemma 2.1.4 and Lemma 2.1.5. □

Remark 2.1.8. Using normalization and an argument similar to the above one can show that for a (not necessary smooth) variety X and a Galois field extension K/k the group

$$\mathrm{CH}_n^K(X) := \mathrm{CH}_n(X) \Big/ \mathrm{Im} \left(\bigoplus_{\substack{k \subseteq F \subseteq K \\ F/k \text{ finite}}} \mathrm{CH}_n(X_F) \xrightarrow{(\pi_{F/k})^*} \mathrm{CH}_n(X) \right),$$

is generated by irreducible cycles that remain to be irreducible over K .

2.2 Equivariant conormed Chow groups

In this section we use the approach by Totaro [To99] and Edidin-Graham [EG98] to extend the conormed Chow groups to the equivariant setting, obtaining a T -equivariant ring cohomology theory $\mathrm{CH}_{K,T}^*(-)$.

Definition 2.2.1. Let T be an affine algebraic group over k and K/k be a separable field extension. Following [EG98, To99] we introduce the T -equivariant conormed Chow ring $\mathrm{CH}_{K,T}^*(X)$ of $X \in \mathcal{S}m_k^T$ as follows. Let $n \in \mathbb{N}_0$, let V be a representation of T and let $U \subseteq V$ be an open subset such that

1. $V \setminus U$ is of codimension greater than n in V ,
2. T acts freely on U ,
3. the quotient U/T exists as a scheme.

Such V and $U \subseteq V$ always exist, see e.g. [EG98, Lemma 9]. We refer to U as an n -th approximation to the universal torsor ET and usually denote such U as $E_n T$. We put

$$\mathrm{CH}_{K,T}^n(X) := \mathrm{CH}_K^n((X \times E_n T)/T),$$

where $(X \times E_n T)/T$ is considered as an algebraic space [EG98, Proposition 22] and the conormed Chow groups on the right are the Chow groups of an algebraic space as in [EG98, Section 6] modulo the respective norm ideal. If the quotient $(X \times E_n T)/T$ is represented by a smooth variety, e.g. if X and T satisfy the assumptions of [EG98, Proposition 23], then these are the conormed Chow groups of a variety introduced in Definition 2.1.1. It is straightforward to check (see [EG98, Definition-Proposition 1]) that, up to a canonical isomorphism, the group $\mathrm{CH}_{K,T}^n(X)$ does not depend on the choice of V and $U = E_n T$. Moreover, one clearly has

$$\mathrm{CH}_{K,T}^*(X) = \mathrm{CH}_T^*(X) \Big/ \mathrm{Im} \left(\bigoplus_{\substack{k \subseteq F \subseteq K \\ F/k \text{ finite}}} \mathrm{CH}_T^*(X_F) \xrightarrow{(\pi_{F/k})^*} \mathrm{CH}_T^*(X) \right),$$

where $\mathrm{CH}_T^*(-)$ are the equivariant Chow groups of [EG98]. This gives rise to an oriented T -equivariant ring cohomology theory on $\mathcal{S}m_k^T$, in particular, $\mathrm{CH}_{K,T}^*(X)$ is a ring for $X \in \mathcal{S}m_k^T$ and one has

- *pullback* ring homomorphisms

$$f^*: \mathrm{CH}_{K,T}^*(X) \rightarrow \mathrm{CH}_{K,T}^*(Y)$$

for a T -equivariant morphism $f: Y \rightarrow X$, $X, Y \in \mathcal{S}m_k^T$,

- *pushforward* homomorphism of $\mathrm{CH}_{K,T}^*(X)$ -modules

$$f_*: \mathrm{CH}_{K,T}^{*-n}(Y) \rightarrow \mathrm{CH}_{K,T}^*(X)$$

for a projective T -equivariant morphism $Y \rightarrow X$ of codimension n with $X, Y \in \mathcal{S}m_k^T$,

- an exact *localization sequence*

$$\mathrm{CH}_{K,T}^{*-n}(Z) \xrightarrow{i_*} \mathrm{CH}_{K,T}^*(X) \xrightarrow{j^*} \mathrm{CH}_{K,T}^*(X - Z) \rightarrow 0$$

for a closed codimension n embedding of smooth T -varieties $Z \rightarrow X$,

- *homotopy invariance* isomorphism

$$f^*: \mathrm{CH}_{K,T}^*(X) \xrightarrow{\cong} \mathrm{CH}_{K,T}^*(\mathcal{V})$$

for a T -equivariant vector bundle $f: \mathcal{V} \rightarrow X$,

- *Chern class* $c_n(\mathcal{V}) \in \mathrm{CH}_{K,T}^n(X)$ for a T -equivariant vector bundle \mathcal{V} over $X \in \mathrm{Sm}_k^T$ and $n \in \mathbb{N}$, in particular, $c_n(V) \in \mathrm{CH}_{K,T}^n(\mathrm{Spec} k)$ for a representation V of T and $n \in \mathbb{N}$,

- *normalization*

$$c_n(\mathcal{V}) = z^* z_*(1) \in \mathrm{CH}_{K,T}^n(X)$$

for the zero section $z: X \rightarrow \mathcal{V}$ of a rank n T -equivariant vector bundle $\mathcal{V} \rightarrow X$.

Combining the localization sequence for the zero section $z: X \rightarrow \mathcal{V}$ of a rank n T -equivariant vector bundle $\mathcal{V} \rightarrow X$, the homotopy invariance isomorphism and the normalization property one obtains the *Gysin exact sequence*

$$\mathrm{CH}_{K,T}^{*-n}(X) \xrightarrow{c_n(\mathcal{V})} \mathrm{CH}_{K,T}^*(X) \xrightarrow{j^*} \mathrm{CH}_{K,T}^*(\mathcal{V} - z(X)) \rightarrow 0.$$

Lemma 2.2.2. *Let T be an affine connected algebraic group over k , $X \in \mathrm{Sm}_k^T$ and let $K/L/k$ be separable extensions of fields with L/k being a finite Galois extension. Suppose that X_L is not connected. Then*

$$\mathrm{CH}_{K,T}^*(X) = 0.$$

Proof. Let $L \supseteq F \supseteq k$ and $X \rightarrow \mathrm{Spec} F \rightarrow \mathrm{Spec} k$ be a factorization of the structure morphism given by Lemma 2.1.5. The factorization arises from an embedding $F \subseteq H^0(X, \mathcal{O}_X)$, thus the projection $X \rightarrow \mathrm{Spec} F$ is T -equivariant with the T -action on $\mathrm{Spec} F$ induced by the one on $H^0(X, \mathcal{O}_X)$. Since the automorphism group of F over k is finite and T is connected, it follows that the T -action on $\mathrm{Spec} F$ is trivial. It follows that the splitting of the projection $X_F \rightarrow X$ constructed in the proof of Lemma 2.1.4 is T -equivariant, so the claim follows as in loc. cit. \square

Definition 2.2.3. Let $\rho: S \rightarrow T$ be a homomorphism of affine algebraic groups over a field k . For $n \in \mathbb{N}_0$ choose n -th approximations $E_n S$ and $E_n T$ to the universal torsors ES and ET and a morphism $f: E_n S \rightarrow E_n T$ equivariant with respect to ρ , i.e. such that the diagram

$$\begin{array}{ccc} S \times E_n S & \longrightarrow & E_n S \\ \rho \times f \downarrow & & \downarrow f \\ T \times E_n T & \longrightarrow & E_n T, \end{array}$$

commutes. Here the horizontal maps are the respective actions. Then for $X \in \mathrm{Sm}_k^T$ we have a morphism

$$\bar{f}: (X \times E_n S)/S \rightarrow (X \times E_n T)/T$$

induced by $\mathrm{id}_X \times f$ and for a separable field extension K/k we obtain a pullback homomorphism

$$\bar{f}^*: \mathrm{CH}_K^n((X \times E_n T)/T) \rightarrow \mathrm{CH}_K^n((X \times E_n S)/S).$$

As in [EG98, Definition-Proposition 1], one can show that this homomorphism does not depend on the choices of approximations and on f yielding a *restriction homomorphism*

$$\Phi_\rho: \mathrm{CH}_{K,T}^n(X) \rightarrow \mathrm{CH}_{K,S}^n(X)$$

with the action of S on X given by the composition of ρ and the action of T . If $S = 1$ is the trivial group, then $\mathrm{CH}_{K,S}^*(-) \cong \mathrm{CH}_K^*(-)$ and we put

$$\Phi_T := \Phi_\rho: \mathrm{CH}_{K,T}^n(X) \rightarrow \mathrm{CH}_{K,S}^n(X) \cong \mathrm{CH}_K^n(X).$$

If $T = 1$ is the trivial group, then $\mathrm{CH}_{K,T}^*(-) \cong \mathrm{CH}_K^*(-)$ and we put

$$\Phi_S^c := \Phi_\rho: \mathrm{CH}_K^n(X) \cong \mathrm{CH}_{K,T}^n(X) \rightarrow \mathrm{CH}_{K,S}^n(X),$$

where $X \in \mathcal{S}m_k$ is equipped with the trivial action of S .

Definition 2.2.4. Let K/k be a separable field extension.

Let $\rho: S \rightarrow T$ be a homomorphism of affine algebraic groups over a field k and $X \in \mathcal{S}m_k^T$. For $n \in \mathbb{N}_0$ choose n -th approximations $E_n S$ and $E_n T$ to the universal torsors ES and ET and consider the following cartesian square induced by the projections:

$$\begin{array}{ccc} (X \times E_n S \times E_n T)/S & \longrightarrow & (E_n S \times E_n T)/S \\ \downarrow & & \downarrow \\ (X \times E_n T)/T & \longrightarrow & E_n T/T \end{array}$$

Here the actions of S and T on the respective products are the diagonal ones, and the action of S on $E_n T$ is induced by the homomorphism ρ . Note that $E_n S \times E_n T$ with the diagonal action of S may also be considered as an n -th approximation to the universal torsor ES . Thus this gives rise to a homomorphism

$$\mathrm{CH}_{K,T}^*(X) \otimes_{\mathrm{CH}_{K,T}^*(\mathrm{Spec} k)} \mathrm{CH}_{K,S}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,S}^*(X), \quad (x, \alpha) \mapsto \alpha \cdot \Phi_\rho(x), \quad (2.2.4.1)$$

which we refer to as *Eilenberg–Moore homomorphism*. One can check that it does not depend on the choice of $E_n S$ and $E_n T$. In particular, for $S = 1$ being the trivial group the Eilenberg–Moore homomorphism looks as

$$\mathrm{CH}_{K,T}^*(X) \otimes_{\mathrm{CH}_{K,T}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_K^*(X), \quad (x, \alpha) \mapsto \alpha \cdot \Phi_T(x). \quad (2.2.4.2)$$

Let $S \xrightarrow{\rho} T \xrightarrow{\rho'} R$ be a short exact sequence of affine algebraic groups over a field k and $X \in \mathcal{S}m_k^T$. For $n \in \mathbb{N}_0$ choose n -th approximations $E_n T$ and $E_n R$ to the universal torsors ET and ER and similarly to the above consider the following cartesian squares:

$$\begin{array}{ccccc} (X \times E_n T \times E_n R)/S & \longrightarrow & (E_n T \times E_n R)/S & \longrightarrow & E_n R \\ \downarrow & & \downarrow & & \downarrow \\ (X \times E_n T \times E_n R)/T & \longrightarrow & (E_n T \times E_n R)/T & \longrightarrow & E_n R/R \end{array}$$

Here the action of S on $E_n T$ is restricted from T , the action of S on $E_n R$ is trivial and the action of T on $E_n R$ is induced by the homomorphism ρ' . With the specified actions $E_n T \times E_n R$ can be viewed as n -th approximations both to the universal torsors ES and ET . Thus we obtain the following version of the Eilenberg–Moore homomorphism:

$$\mathrm{CH}_{K,T}^*(X) \otimes_{\mathrm{CH}_{K,R}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,S}^*(X), \quad (x, \alpha) \mapsto \alpha \cdot \Phi_\rho(x), \quad (2.2.4.3)$$

with the structure of $\mathrm{CH}_{K,R}^*(\mathrm{Spec} k)$ -module on $\mathrm{CH}_{K,T}^*(X)$ induced by the restriction homomorphism $\Phi_{\rho'}: \mathrm{CH}_{K,R}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,T}^*(\mathrm{Spec} k)$.

Remark 2.2.5. The name for the homomorphisms in Definition 2.2.4 refers to the classical Eilenberg–Moore spectral sequence which computes the singular cohomology groups of a pullback over a fibration.

Lemma 2.2.6 (see [EG98, Proposition 8(a)]). *Let K/k be a separable field extension, let T be an affine algebraic group over k , $X \in \mathcal{S}m_k^T$ and suppose that the action of T on X is free and X/T is represented by a scheme. Then there is a canonical isomorphism*

$$\mathrm{CH}_{T,K}^*(X) \cong \mathrm{CH}_K^*(X/T).$$

Proof. The same reasoning as in the proof of [EG98, Proposition 8(a)] applies, but since we will make computations using this isomorphism later in the article, we provide the construction for a completeness.

Let $n \in \mathbb{N}_0$ and let $E_n T \subseteq V$ be an n -th approximation to the universal torsor ET . Since $X \rightarrow X/T$ a T -torsor, it follows that

$$(X \times V)/T \rightarrow X/T$$

is a vector bundle with

$$(X \times E_n T)/T \subseteq (X \times V)/T$$

being an open subset with the complement of codimension greater than n . Thus

$$\mathrm{CH}_{T,K}^n(X) = \mathrm{CH}_K^n((X \times E_n T)/T) \cong \mathrm{CH}_K^n((X \times V)/T) \cong \mathrm{CH}_K^n(X/T). \quad \square$$

Lemma 2.2.7. *Let K/k be a separable field extension, let N be a finite group scheme over k and let $X \in \mathcal{S}m_k$ be equipped with the trivial action of N . Suppose that the order of N is invertible in $\mathrm{CH}_K^*(\mathrm{Spec} k)$. Then the restriction homomorphism*

$$\Phi_N: \mathrm{CH}_{K,N}^*(X) \rightarrow \mathrm{CH}_K^*(X)$$

is an isomorphism.

Proof. For $n \in \mathbb{N}_0$ let $E_n N$ be an n -th approximation to the universal torsor EN . Then

$$f: X \times E_n N \rightarrow (X \times E_n N)/N \cong X \times (E_n N)/N$$

is a finite morphism of degree equal to the order of N and the composition

$$f_* \circ f^*: \mathrm{CH}_K^n((X \times E_n N)/N) \rightarrow \mathrm{CH}_K^n(X \times E_n N) \rightarrow \mathrm{CH}_K^n((X \times E_n N)/N),$$

is the multiplication by the order of N [Fu98, Example 1.7.4]. It follows that $f_* \circ f^*$ an isomorphism. Recall that

$$\mathrm{CH}_{K,N}^n(X) = \mathrm{CH}_K^n((X \times E_n N)/N), \quad \mathrm{CH}_K^n(X) = \mathrm{CH}_K^n(X \times E_n N),$$

and under this identifications we have $\Phi_N = f^*$. Thus the composition

$$\mathrm{CH}_{K,N}^n(X) \xrightarrow{\Phi_N} \mathrm{CH}_K^n(X) \xrightarrow{f_*} \mathrm{CH}_{K,N}^n(X)$$

is an isomorphism. On the other hand, since X has a trivial action of N , we have a homomorphism

$$\Phi_N^c: \mathrm{CH}_K^n(X) \rightarrow \mathrm{CH}_{K,N}^n(X)$$

induced by the projection $g: (X \times E_n N)/N \rightarrow X$, and the composition

$$\mathrm{CH}_K^n(X) \xrightarrow{\Phi_N^c} \mathrm{CH}_{K,N}^n(X) \xrightarrow{\Phi_N} \mathrm{CH}_K^n(X)$$

is also an isomorphism. The claim follows. □

2.3 Eilenberg–Moore isomorphism for quasi-trivial tori

In this section we investigate the relations between equivariant and non-equivariant conormed Chow groups for quasi-trivial tori, with the main results being the computation of the equivariant conormed Chow ring of the point and a receipt how to recover the non-equivariant conormed Chow groups from the equivariant ones. Here and below we use some standard notation and well-known facts about quasi-trivial tori, see Section A.1 for a recollection.

Proposition 2.3.1. *Let $K/L/k$ be separable extensions of fields with L/k being finite, let T be a connected algebraic group over k , let $R := R_{L/k}\mathbb{G}_m$ be the Weil restriction of the one-dimensional split torus and let $\rho: T \rightarrow R$ be a homomorphism. Suppose that R_K is split. Then for $X \in \mathcal{S}m_k^T$ the sequence*

$$\mathrm{CH}_{K,T}^{*-n}(X) \xrightarrow{c_n(V_R)} \mathrm{CH}_{K,T}^*(X) \xrightarrow{j^*} \mathrm{CH}_{K,T}^*(X \times R) \rightarrow 0$$

is exact. Here

- $n = [L : k]$,
- the first homomorphism is given by the multiplication with the Chern class $c_n(V_R)$ where the standard vector representation V_R of R is viewed as a representation of T via ρ ,
- T acts on $X \times R$ diagonally,
- $j: X \times R \rightarrow X$ is the projection.

Proof. We have a canonical open embedding $R \subseteq V_R \setminus \{0\} \subseteq V_R$ (see (A.1.2)) and the Gysin exact sequence for the vector bundle $X \times V_R \rightarrow X$ implies that it is sufficient to check that the pullback homomorphism

$$\mathrm{CH}_{K,T}^*(X \times (V_R \setminus \{0\})) \rightarrow \mathrm{CH}_{K,T}^*(X \times R)$$

is an isomorphism. Below we will omit X from the notation.

Since R_K is split by the assumption, it follows that K contains the Galois closure \bar{L}/k of L/k which is the splitting field of R . Choose a basis $\{e_1, e_2, \dots, e_n\}$ for L over k and fix the corresponding isomorphism $V_R \cong \mathbb{A}_k^n = \mathrm{Spec} k[x_1, x_2, \dots, x_n]$. Denote by

$$\mathcal{H} = Z(x_1e_1 + x_2e_2 + \dots + x_ne_n) \leq \mathbb{A}_{\bar{L}}^n \cong (V_R)_{\bar{L}}$$

the corresponding hyperplane. Put $\mathcal{G} := \mathrm{Gal}(\bar{L}/k)$, $\mathcal{G}' := \mathrm{Gal}(\bar{L}/L)$, $\mathcal{S} := \mathcal{G}/\mathcal{G}'$ and consider the following filtration of $\mathbb{A}_{\bar{L}}^n \cong (V_R)_{\bar{L}}$ by closed subsets.

$$\{0\} = \bar{V}_n \leq \bar{V}_{n-1} \leq \dots \leq \bar{V}_0 = \mathbb{A}_{\bar{L}}^n \cong (V_R)_{\bar{L}}, \quad \bar{V}_i := \bigcup_{\substack{I \subseteq \mathcal{S} \\ |I|=i}} \mathcal{H}_I, \quad \mathcal{H}_I := \bigcap_{\sigma \in I} \sigma(\mathcal{H}), \quad \mathcal{H}_\emptyset := \mathbb{A}_{\bar{L}}^n.$$

Note that \mathcal{H} is stable under the action of \mathcal{G}' , so we may define $\sigma(\mathcal{H})$ for $\sigma \in \mathcal{S}$ using an arbitrary representative of the corresponding coset. Since for every i the variety \bar{V}_i is clearly \mathcal{G} -stable, it follows that there exist closed subvarieties

$$\{0\} = V_n \leq V_{n-1} \leq \dots \leq V_0 = \mathbb{A}_k^n \cong V_R$$

such that $(V_i)_{\bar{L}} = \bar{V}_i$. For $0 \leq i \leq n-1$ we have

$$\bar{V}_i \setminus \bar{V}_{i+1} = \bigsqcup_{\substack{I \subseteq \mathcal{S} \\ |I|=i}} \mathcal{H}_I^\circ = \bigsqcup_{\mathcal{A} \in \mathrm{Orb}_{\mathcal{G}}(\mathcal{S}_i)} \left(\bigsqcup_{I \in \mathcal{A}} \mathcal{H}_I^\circ \right), \quad \mathcal{H}_I^\circ := \mathcal{H}_I \setminus \bigcup_{\tau \in \mathcal{S} \setminus I} (\mathcal{H}_I \cap \tau(\mathcal{H})).$$

Here $\mathrm{Orb}_{\mathcal{G}}(\mathcal{S}_i)$ is the set of \mathcal{G} -orbits under the natural action of \mathcal{G} on the set $\{I \subseteq \mathcal{S} \mid |I| = i\}$ induced by the action of \mathcal{G} on $\mathcal{G}/\mathcal{G}' = \mathcal{S}$. For $0 \leq i \leq n-1$ and $\mathcal{A} \in \mathrm{Orb}_{\mathcal{G}}(\mathcal{S}_i)$ put

$$\bar{W}_{\mathcal{A}} := \bigsqcup_{I \in \mathcal{A}} \mathcal{H}_I^\circ.$$

Note that every $\overline{W}_{\mathcal{A}}$ is \mathcal{G} -stable and smooth being a disjoint union of open subsets of different $\mathcal{H}_I \cong \mathbb{A}_{\overline{L}}^{n-i}$, hence it corresponds to some smooth locally closed subvariety $W_{\mathcal{A}} \subseteq \mathbb{A}_k^n \cong V_R$ satisfying $(W_{\mathcal{A}})_{\overline{L}} = \overline{W}_{\mathcal{A}}$. Since \mathcal{G} acts transitively on \mathcal{A} , it follows that $W_{\mathcal{A}}$ is connected, and since $|\mathcal{A}| > 1$ for $1 \leq i \leq n-1$, it follows that $(W_{\mathcal{A}})_{\overline{L}} = \overline{W}_{\mathcal{A}}$ is not connected for such i . Applying Lemma 2.2.2 to $W_{\mathcal{A}}$ and $K/\overline{L}/k$ we get

$$\mathrm{CH}_{K,T}^*(W_{\mathcal{A}}) = 0$$

for $1 \leq i \leq n-1$. Successively applying localization sequences

$$\mathrm{CH}_{K,T}^{*-i}\left(\bigsqcup_{\mathcal{A} \in \mathrm{Orb}_{\mathcal{G}}^i} W_{\mathcal{A}}\right) \rightarrow \mathrm{CH}_{K,T}^*(V_R \setminus V_{i+1}) \rightarrow \mathrm{CH}_{K,T}^*(V_R \setminus V_i) \rightarrow 0$$

for $i = n-1, \dots, 1$ we obtain isomorphisms

$$\mathrm{CH}_{K,T}^*(V_R \setminus V_n) \xrightarrow{\cong} \mathrm{CH}_{K,T}^*(V_R \setminus V_{n-1}) \xrightarrow{\cong} \dots \xrightarrow{\cong} \mathrm{CH}_{K,T}^*(V_R \setminus V_1).$$

Recall that $V_R \setminus V_n = V_R \setminus \{0\}$ while recollection (A.1.2) yields $V_R \setminus V_1 = R$. The claim follows. \square

Proposition 2.3.2 (see also [Ka12, § 3]). *Let K/k be a separable field extension and let $T \cong R_{L_1/k} \mathbb{G}_m \times \dots \times R_{L_r/k} \mathbb{G}_m$ be a quasi-trivial torus over k . Suppose that T_K is split. Then for $X \in \mathrm{Sm}_k$ equipped with the trivial action of T one has an isomorphism*

$$\mathrm{CH}_{K,T}^*(X) \cong \mathrm{CH}_K^*(X)[b_1, b_2, \dots, b_r], \quad b_i \mapsto c_{n_i}(V_i), \quad n_i := [L_i : k],$$

where $V_i := V_{R_i}$ is the standard vector representation of $R_i := R_{L_i/k} \mathbb{G}_m$ viewed as a representation of T via the projection $T \rightarrow R_i$.

Proof. The Weil restrictions $R_{L_i/k} \mathbb{P}_{L_i}^n$ are models for the n -th approximations $(E_n R_i)/R_i$ to the classifying spaces $(E R_i)/R_i$ by [Ka12, § 3] (see also [KM22, Lemma 3.1]), in particular, one has

$$\mathrm{CH}_{K,T}^*(X) \cong \varprojlim_n \mathrm{CH}_K^*(X \times R_{L_1/k} \mathbb{P}_{L_1}^n \times \dots \times R_{L_r/k} \mathbb{P}_{L_r}^n).$$

It follows from [Ka00, Proposition 5.6] that the (integral) Chow motive $\mathcal{M}(R_{L_i/k} \mathbb{P}_{L_i}^n)$ splits as a direct sum of motives $\mathcal{M}(\mathrm{Spec} F, l)$ for some $l \in \mathbb{N}_0$ and intermediate fields $\overline{L}_i/F/k$ with L_i/k being a Galois closure of L_i/k . Since T_K is split and \overline{L}_i is the splitting field of R_i , it follows that \overline{L}_i/k embeds into K/k . Lemma 2.1.4 yields that in the conormed motives one has $\mathcal{M}_K(\mathrm{Spec} F, l) = 0$ for $K \supseteq F \supsetneq k$, thus

$$\mathcal{M}_K(R_{L_i/k} \mathbb{P}_{L_i}^n) \cong \mathcal{M}_K(\mathrm{Spec} k, l_0) \oplus \dots \oplus \mathcal{M}_K(\mathrm{Spec} k, l_d)$$

for some integers l_0, \dots, l_d . Then the Künneth formula

$$\mathrm{CH}_K^*(X \times R_{L_1/k} \mathbb{P}_{L_1}^n \times \dots \times R_{L_r/k} \mathbb{P}_{L_r}^n) \cong \mathrm{CH}_K^*(X) \otimes \mathrm{CH}_K^*(R_{L_1/k} \mathbb{P}_{L_1}^n) \otimes \dots \otimes \mathrm{CH}_K^*(R_{L_r/k} \mathbb{P}_{L_r}^n)$$

holds and it suffices to show that for every i there is an isomorphism

$$\mathrm{CH}_{K,R_i}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[b_i], \quad b_i \mapsto c_{n_i}(V_i).$$

If $L_i = k$, then

$$\mathrm{CH}_{K,R_i}^*(\mathrm{Spec} k) = \mathrm{CH}_K^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[x], \quad x \mapsto c_1(V_{\mathbb{G}_m}),$$

by the usual projective bundle theorem. So we can assume that $L_i \supsetneq k$. Proposition 2.3.1 together with the isomorphism $\mathrm{CH}_{K,R_i}^*(R_i) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)$ of Lemma 2.2.6 yields that $\mathrm{CH}_{K,R_i}^{\geq 0}(\mathrm{Spec} k)$ is generated by $c_{n_i}(V_i)$. Since $\mathrm{CH}_{K,R_i}^0(\mathrm{Spec} k) = \mathrm{CH}_K^0(\mathrm{Spec} k)$ is either zero or a field by Lemma 2.1.2, it follows that $b_i \mapsto c_{n_i}(V_i)$ induces an isomorphism $\mathrm{CH}_{K,R_i}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[b_i]$ or $\mathrm{CH}_{K,R_i}^*(\mathrm{Spec} k) \cong$

$\mathrm{CH}_K^*(\mathrm{Spec} k)[b_i]/(b_i^m)$ for some $m \in \mathbb{N}$. In order to see that $c_{n_i}(V_i)$ is not nilpotent consider the homomorphism $\rho: \mathbb{G}_m \rightarrow R_i$ given by the unit of the adjunction between the extension of scalars and the Weil restriction. This gives rise to a homomorphism

$$\Phi_\rho: \mathrm{CH}_{K,R_i}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,\mathbb{G}_m}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[x].$$

The representation V_i of R_i restricted to \mathbb{G}_m is isomorphic to a direct sum of n_i copies of the standard linear representation of \mathbb{G}_m . Thus

$$\Phi_\rho(c_{n_i}(V_i)) = c_{n_i}(V_{\mathbb{G}_m}^{\oplus n_i}) = c_1(\mathcal{O}(-1))^{n_i},$$

so $\Phi_\rho(c_{n_i}(V_i))$ is not nilpotent and it follows that $c_{n_i}(V_i)$ is not nilpotent as well. \square

Corollary 2.3.3. *Let K/k be a separable field extension and let T be a quasi-trivial torus over a field k . Suppose that T_K is split. Then for $X \in \mathrm{Sm}_k^T$ the Eilenberg–Moore homomorphism (2.2.4.2)*

$$\mathrm{CH}_{K,T}^*(X) \otimes_{\mathrm{CH}_{K,T}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_K^*(X)$$

is an isomorphism.

Proof. Choose an isomorphism $T \cong R_{L_1/k}\mathbb{G}_m \times \dots \times R_{L_r/k}\mathbb{G}_m$ with L_i/k being finite separable field extensions. Put $n_i := [L_i : k]$ and $R_i := R_{L_i/k}\mathbb{G}_m$. Inductively applying Proposition 2.3.1 we obtain

$$\mathrm{CH}_{K,T}^*(X \times T) \cong \mathrm{CH}_{K,T}^*(X)/(c_{n_1}(V_{R_1}), \dots, c_{n_r}(V_{R_r}))$$

where V_{R_i} is viewed as a representation of T via the projection $T \rightarrow R_i$. Lemma 2.2.6 yields an isomorphism $\mathrm{CH}_{K,T}^*(X \times T) \cong \mathrm{CH}_K^*(X)$ while Proposition 2.3.2 yields an isomorphism $\mathrm{CH}_{K,T}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[b_1, \dots, b_r]$, $b_i \mapsto c_{n_i}(V_{R_i})$, so the claim follows. \square

Remark 2.3.4. A similar result was obtained in [Sal22, Theorem 3.1] claiming $\mathrm{CH}^*(X)$ to be the quotient of $\mathrm{CH}_T^*(X)$ modulo the pushforwards of characteristic classes defined over field extensions F/k .

Definition 2.3.5. Let N be an affine algebraic group over a field k and let K/k be a separable field extension. We say that BN satisfies the Künneth formula for K/k , if for every $X \in \mathrm{Sm}_k$ equipped with the trivial action of N the Künneth homomorphism

$$\mathrm{CH}_K^*(X) \otimes \mathrm{CH}_{K,N}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,N}^*(X), \quad (y, \alpha) \mapsto \alpha \cdot \Phi_N^c(y),$$

is an isomorphism.

Proposition 2.3.6. *Let K/k be a separable field extension, let T be a quasi-trivial torus over k , let $N \leq T$ be a subgroup and $X \in \mathrm{Sm}_k^T$. Suppose that T_K is split and that BN satisfies the Künneth formula for K/k . Then the homomorphism*

$$\begin{aligned} (\mathrm{CH}_{K,T}^*(X) \otimes \mathrm{CH}_{K,N}^*(\mathrm{Spec} k)) \otimes_{\mathrm{CH}_{K,T}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) &\rightarrow \mathrm{CH}_{K,N}^*(X), \\ (x, \alpha, \beta) &\mapsto \alpha \cdot \beta \cdot \Phi_\rho(x), \end{aligned}$$

is an isomorphism. Here $\rho: N \rightarrow T$ is the embedding and the structure of a $\mathrm{CH}_{K,T}^*(\mathrm{Spec} k)$ -module on $\mathrm{CH}_{K,T}^*(X) \otimes \mathrm{CH}_{K,N}^*(\mathrm{Spec} k)$ arises from the ring homomorphism

$$\mathrm{CH}_{K,T}^*(\mathrm{Spec} k) \xrightarrow{\Phi_\nu} \mathrm{CH}_{K,T \times N}^*(\mathrm{Spec} k) \cong \mathrm{CH}_{K,T}^*(\mathrm{Spec} k) \otimes \mathrm{CH}_{K,N}^*(\mathrm{Spec} k)$$

for the homomorphism $\nu: T \times N \rightarrow T$, $\nu(t, s) = t \cdot s^{-1}$, composed with the componentwise module structure.

Proof. Let $n \in \mathbb{N}_0$ and let $E_n T$ and $E_n N$ be n -th approximations to the universal torsors ET and EN respectively and equip $X \times E_n T \times E_n N$ with an action of $T \times N$ given by $(t, s) \cdot (x, u, v) := (tsx, tsu, sv)$.

Let $\tilde{X} := (X \times E_n T \times E_n N)/N$ and equip it with an action of T compatible with the projection $X \times E_n T \times E_n N \rightarrow \tilde{X}$. Corollary 2.3.3 applied to \tilde{X} yields an isomorphism

$$\mathrm{CH}_{K,T}^*(\tilde{X}) \otimes_{\mathrm{CH}_{K,T}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \xrightarrow{\cong} \mathrm{CH}_K^*(\tilde{X}).$$

Since $E_n T$ and $E_n N$ are n -th approximations to the universal torsors ET and EN , it follows that $E_n T \times E_n N$ with the diagonal action of N is also an n -th approximation to the universal torsor EN , and $\mathrm{CH}_K^n(\tilde{X}) \cong \mathrm{CH}_{K,N}^n(X)$. We have isomorphisms

$$\tilde{X}/T \cong (X \times E_n T \times E_n N)/(T \times N) \xrightarrow{\cong} (X \times E_n T)/T \times E_n N/N$$

with the second one induced by the respective projections. Applying further Lemma 2.2.6 we obtain isomorphisms

$$\mathrm{CH}_{K,T}^*(\tilde{X}) \cong \mathrm{CH}_K^*(\tilde{X}/T) \cong \mathrm{CH}_K^*((X \times E_n T)/T \times E_n N/N) \cong \mathrm{CH}_{K,T}^*(X) \otimes \mathrm{CH}_{K,N}^*(\mathrm{Spec} k) \quad (2.3.6.1)$$

in degrees up to n . Combining all the above we obtain the desired isomorphism

$$(\mathrm{CH}_{K,T}^*(X) \otimes \mathrm{CH}_{K,N}^*(\mathrm{Spec} k)) \otimes_{\mathrm{CH}_{K,T}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \xrightarrow{\cong} \mathrm{CH}_{K,N}^*(X), \quad (2.3.6.2)$$

It is straightforward to check that the isomorphism is given by $(x, \alpha, \beta) \mapsto \alpha \cdot \beta \cdot \Phi_\rho(x)$ as claimed. In order to identify the module structure consider the following diagram.

$$\begin{array}{ccc} E_n T/T & \xleftarrow{q_3} & (E_n T \times E_n N \times E_n T)/(T \times N) \\ p_4 \uparrow & \nearrow p_{234} & \downarrow q_{12} \\ (X \times E_n T \times E_n N \times E_n T)/(T \times N) & & (E_n T \times E_n N)/(T \times N) \\ \downarrow (p_{12}, p_3) & & \cong \downarrow (s_1, s_2) \\ (X \times E_n T)/T \times E_n N/N & \xrightarrow{(r_2, \mathrm{id})} & E_n T/T \times E_n N/N \end{array}$$

Here the action in the middle of the left column is given by $(t, s) \cdot (x, u, v, w) := (tsx, tsu, sv, tw)$, in the bottom-left corner T acts diagonally on $X \times E_n T$ and N acts on $E_n N$ in the given way, the action in the top-right corner is given by $(t, s) \cdot (u, v, w) := (tu, sv, ts^{-1}w)$, in the middle of the right column the action is the componentwise one and in the remaining corners it is the standard one. All the numbered morphisms are induced by the projections on the respective factors, e.g. p_4 is induced by the projection $X \times E_n T \times E_n N \times E_n T \rightarrow E_n T$ on the last $E_n T$, and p_{234} is induced by the projection $X \times E_n T \times E_n N \times E_n T \rightarrow E_n T \times E_n N \times E_n T$; it is straightforward to check that these projections descend to the quotients under the given group actions. The diagram clearly commutes. Morphism (s_1, s_2) is an isomorphism, and, moreover, morphisms (p_{12}, p_3) and q_{12} induce isomorphisms up to degree n on the respective conormed Chow groups by the codimension reasons and homotopy invariance. Unwinding the definition of the Eilenberg–Moore homomorphism (2.2.4.2) and the isomorphisms (2.3.6.1) we see that the $\mathrm{CH}_{K,T}^*(\mathrm{Spec} k)$ -module structure on $\mathrm{CH}_{K,T}^*(X) \otimes \mathrm{CH}_{K,N}^*(\mathrm{Spec} k)$ in the isomorphism (2.3.6.2) up to degree n is induced by the ring homomorphism

$$((p_{12}, p_3)^*)^{-1} \circ p_4^*: \mathrm{CH}_K^*(E_n T/T) \rightarrow \mathrm{CH}_K^*((X \times E_n T)/T \times E_n N/N) \cong \mathrm{CH}_{K,T}^*(X) \otimes \mathrm{CH}_{K,N}^*(\mathrm{Spec} k).$$

Commutativity of the diagram yields that up to degree n we have

$$((p_{12}, p_3)^*)^{-1} \circ p_4^* = (r_2, \mathrm{id})^* \circ ((s_1, s_2)^*)^{-1} \circ (q_{12}^*)^{-1} \circ q_3^*.$$

The pullback homomorphism q_3^* is a representative (up to degree n) for the restriction homomorphism

$$\Phi_\nu: \mathrm{CH}_{K,T}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,T \times N}^*(\mathrm{Spec} k),$$

and q_{12}^* is the canonical isomorphism between the models of the equivariant conormed Chow groups based on the different approximations to the universal torsor $E(T \times N)$, so the pullback homomorphism $(r_2, \mathrm{id})^* \circ ((s_1, s_2)^*)^{-1} \circ (q_{12}^*)^{-1} \circ q_3^*$ is the ring homomorphism giving precisely the claimed module structure. \square

2.4 Some equivariant conormed Chow rings of a point

In this section we study some equivariant conormed Chow rings of a point, in particular, we compute the conormed rings of a point $\mathrm{CH}_{K, \pi_1(G)}^*(\mathrm{Spec} k)$ for the fundamental groups of non-split quasi-split simple groups, assuming that K splits the group in question. From now on we use some further notation related to groups of multiplicative type introduced in Section A.1. In particular, for a degree 2 Galois field extension L/k we have the following groups:

- ${}^2\mathbb{G}_m$ is the non-trivial form of \mathbb{G}_m split by L ,
- ${}^2\mu_l$ is the non-trivial form of μ_l split by L (if $l \geq 3$), and ${}^2\mu_2 = \mu_2$,
- ${}^2\mu_{2,2}$ is the non-trivial form of $\mu_2 \times \mu_2$ split by L .

These groups come with canonical embeddings into the Weil restriction $R := R_{L/k}\mathbb{G}_m$ of the split one-dimensional torus. We denote the restrictions of the standard dimension 2 vector representation V_R of R to these groups by ${}^2V_{\mathbb{G}_m}$, ${}^2V_{\mu_l}$ and ${}^2V_{\mu_{2,2}}$ respectively. Furthermore, if $l = 2m$ is even, then there is also a nontrivial linear representation of ${}^2\mu_{2m}$ denoted by ${}^2\Lambda_{\mu_{2m}}^\pm$.

Proposition 2.4.1. *Let $K/L/k$ be separable field extensions with $[L : k] = 2$ and let $X \in \mathcal{S}m_k$. Then we have the following isomorphisms.*

1. $\mathrm{CH}_{K, {}^2\mathbb{G}_m}^*(X) \cong \mathrm{CH}_K^*(X)[b]$, $b \mapsto c_2({}^2V_{\mathbb{G}_m})$,
2. $\mathrm{CH}_{K, {}^2\mu_{2m+1}}^*(X) \cong \mathrm{CH}_K^*(X)$,
3. $\mathrm{CH}_{K, {}^2\mu_{2m}}^*(X) \cong \mathrm{CH}_K^*(X)[x, b]/(x^2 + m^2 \cdot b)$, $x \mapsto c_1({}^2\Lambda_{\mu_{2m}}^\pm)$, $b \mapsto c_2({}^2V_{\mu_{2m}})$,
4. $\mathrm{CH}_{K, {}^2\mu_{2,2}}^*(X) \cong \mathrm{CH}_K^*(X)[b]$, $b \mapsto c_2({}^2V_{\mu_{2,2}})$.

Here X is considered as a variety with the trivial action.

Proof. Put $R := R_{L/k}\mathbb{G}_m$ to be the Weil restriction of the split one-dimensional torus and let V_R be its standard vector representation of dimension 2.

- (1) Consider the Eilenberg–Moore homomorphism (2.2.4.3)

$$\mathrm{CH}_{K,R}^*(X) \otimes_{\mathrm{CH}_{K, \mathbb{G}_m}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K, {}^2\mathbb{G}_m}^*(X)$$

associated with the exact sequence of groups

$$1 \rightarrow {}^2\mathbb{G}_m \rightarrow R \rightarrow \mathbb{G}_m \rightarrow 1.$$

By the construction, this homomorphism arises from a \mathbb{G}_m -torsor

$$(X \times E_n R \times E_n \mathbb{G}_m) / {}^2\mathbb{G}_m \rightarrow (X \times E_n R \times E_n \mathbb{G}_m) / R$$

with $E_n R$ and $E_n \mathbb{G}_m$ being n -th approximations to the corresponding universal torsors. Corollary 2.3.3 applied to \mathbb{G}_m acting on $(X \times E_n R \times E_n \mathbb{G}_m) / {}^2\mathbb{G}_m$ yields that this Eilenberg–Moore homomorphism is an isomorphism. Furthermore, Proposition 2.3.2 yields isomorphisms

$$\mathrm{CH}_{K,R}^*(X) \cong \mathrm{CH}_K^*(X) \otimes \mathrm{CH}_{K,R}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(X) \otimes \mathrm{CH}_K^*(\mathrm{Spec} k)[b]$$

with $b \mapsto c_2(V_R)$. The action of $\mathrm{CH}_{K, \mathbb{G}_m}^*(\mathrm{Spec} k)$ on $\mathrm{CH}_{K,R}^*(X)$ in the Eilenberg–Moore homomorphism factors through the homomorphism

$$\mathrm{CH}_K^*(\mathrm{Spec} k)[x] \cong \mathrm{CH}_{K, \mathbb{G}_m}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,R}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[b]$$

which maps x to 0, since the right-hand side has no elements of degree 1. Combining all the above we obtain

$$\mathrm{CH}_K^*(X)[b] \cong \mathrm{CH}_{K,R}^*(X) \cong \mathrm{CH}_{K,R}^*(X) \otimes_{\mathrm{CH}_{K, \mathbb{G}_m}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \cong \mathrm{CH}_{K, {}^2\mathbb{G}_m}^*(X).$$

(2) We have $\mathrm{CH}_K^*(\mathrm{Spec} k) = \mathbb{F}_2$ or $\mathrm{CH}_K^*(\mathrm{Spec} k) = 0$ by Lemma 2.1.2. The claim follows from Lemma 2.2.7.

(3) Consider the short exact sequence of algebraic groups

$$1 \rightarrow {}^2\mu_{2m} \xrightarrow{f} {}^2\mathbb{G}_m \times \mathbb{G}_m \xrightarrow{g} R \rightarrow 1$$

The Cartier dual to the short exact sequence of $\mathrm{Gal}(L/k)$ -modules is

$$0 \rightarrow \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{\hat{g}} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{\hat{f}} \mathbb{Z}/2m\mathbb{Z} \rightarrow 0,$$

$$\hat{g}(x, y) = (mx - my, x + y), \quad \hat{f}(x, y) = x + my,$$

with the action of the nontrivial element $\tau \in \mathrm{Gal}(L/k) \cong C_2$ given by

$$\tau(x, y) = (y, x), \quad \tau(x, y) = (-x, y), \quad \tau(x) = -x$$

on the respective abelian groups. As in part (1) we obtain that the Eilenberg–Moore homomorphism (2.2.4.3)

$$\mathrm{CH}_{K, {}^2\mathbb{G}_m \times \mathbb{G}_m}^*(X) \otimes_{\mathrm{CH}_{K, R}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K, {}^2\mu_{2m}}^*(X) \quad (2.4.1.1)$$

is an isomorphism. Proposition 2.3.2 together with part (1) of the current lemma yields isomorphisms

$$\mathrm{CH}_{K, {}^2\mathbb{G}_m \times \mathbb{G}_m}^*(X) \cong \mathrm{CH}_K^*(X)[x, b], \quad \mathrm{CH}_{K, R}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[b'],$$

$$\mathrm{CH}_{K, {}^2\mathbb{G}_m \times \mathbb{G}_m}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[x, b]$$

with $b' \mapsto c_2(V_R)$, $x \mapsto c_1(V_{\mathbb{G}_m})$, $b \mapsto c_2({}^2V_{\mathbb{G}_m})$, and the $\mathrm{CH}_K^*(\mathrm{Spec} k)[b']$ -module structure on $\mathrm{CH}_K^*(X)[x, b]$ factors through the homomorphism

$$\mathrm{CH}_K^*(\mathrm{Spec} k)[b'] \cong \mathrm{CH}_{K, R}^*(\mathrm{Spec} k) \xrightarrow{\Phi_g} \mathrm{CH}_{K, {}^2\mathbb{G}_m \times \mathbb{G}_m}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[x, b].$$

In the representation ring $\mathrm{Rep}(R) \cong \mathbb{Z}[\mathcal{X}^*(R)]^{\mathrm{Gal}(L/k)} \cong \mathbb{Z}[\mathbb{Z} \oplus \mathbb{Z}]^{\mathrm{Gal}(K/k)}$ we have $[V_R] = x^{(1,0)} + x^{(0,1)}$. Then

$$\mathrm{Res}_g([V_R]) = \mathrm{Res}_g(x^{(1,0)} + x^{(0,1)}) = x^{\hat{g}(1,0)} + x^{\hat{g}(0,1)} = x^{(m,1)} + x^{(-m,1)} = (x^{(m,0)} + x^{(-m,0)}) \cdot x^{(0,1)}$$

in $\mathrm{Rep}({}^2\mathbb{G}_m \times \mathbb{G}_m) \cong (\mathbb{Z}[\mathcal{X}^*({}^2\mathbb{G}_m)] \otimes \mathbb{Z}[\mathcal{X}^*(\mathbb{G}_m)])^{\mathrm{Gal}(L/k)}$ with $(1, 0)$ and $(0, 1)$ being the generators for $\mathcal{X}^*({}^2\mathbb{G}_m)$ and $\mathcal{X}^*(\mathbb{G}_m)$ respectively. Then

$$\mathrm{Res}_g([V_R]) = \psi^m[{}^2V_{\mathbb{G}_m}] \otimes [V_{\mathbb{G}_m}],$$

where ψ^m is the m -th Adams operation. Using the standard properties of the Chern classes and that $c_1({}^2V_{\mathbb{G}_m}) = 0$, since $\mathrm{CH}_{K, {}^2\mathbb{G}_m}^1(\mathrm{Spec} k) = 0$, we obtain

$$\begin{aligned} \Phi_g(c_2(V_R)) &= c_2(\mathrm{Res}_g([V_R])) = c_2(\psi^m[{}^2V_{\mathbb{G}_m}] \otimes [V_{\mathbb{G}_m}]) = \\ &= c_1(V_{\mathbb{G}_m})^2 + m \cdot c_1(V_{\mathbb{G}_m}) \cdot c_1({}^2V_{\mathbb{G}_m}) + m^2 \cdot c_2({}^2V_{\mathbb{G}_m}) = c_1(V_{\mathbb{G}_m})^2 + m^2 \cdot c_2({}^2V_{\mathbb{G}_m}). \end{aligned}$$

Putting everything together in isomorphism (2.4.1.1) we obtain

$$\mathrm{CH}_{K, {}^2\mu_{2m}}^*(X) \cong \mathrm{CH}_K^*(X)[x, b]/(x^2 + m^2 \cdot b), \quad x \mapsto \Phi_f(c_1(V_{\mathbb{G}_m})), \quad b \mapsto \Phi_f(c_2({}^2V_{\mathbb{G}_m})).$$

We have

$$\Phi_f(c_1(V_{\mathbb{G}_m})) = c_1(\mathrm{Res}_f([V_{\mathbb{G}_m}])) = c_1({}^2\Lambda_{\mu_{2m}}^\pm), \quad \Phi_f(c_2({}^2V_{\mathbb{G}_m})) = c_2(\mathrm{Res}_f([{}^2V_{\mathbb{G}_m}])) = c_2({}^2V_{\mu_{2m}}),$$

thus x maps to $c_1({}^2\Lambda_{\mu_{2m}}^\pm)$ and b maps to $c_2({}^2V_{\mu_{2m}})$ as claimed.

(4) Consider the standard embedding $f: {}^2\mu_{2,2} \rightarrow R$. As in part (1), the Eilenberg–Moore homomorphism (2.2.4.3)

$$\mathrm{CH}_{K,R}^*(X) \otimes_{\mathrm{CH}_{K,R}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,{}^2\mu_{2,2}}^*(X) \quad (2.4.1.2)$$

associated with the short exact sequence of groups

$$1 \rightarrow {}^2\mu_{2,2} \xrightarrow{f} R \xrightarrow{g} R \rightarrow 1, \quad g(x) = x^2,$$

is an isomorphism. Proposition 2.3.2 yields

$$\mathrm{CH}_{K,R}^*(X) \cong \mathrm{CH}_K^*(X) \otimes \mathrm{CH}_{K,R}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(X)[b], \quad b \mapsto c_2(V_R),$$

and the $\mathrm{CH}_{K,R}^*(\mathrm{Spec} k)$ -module structure in isomorphism (2.4.1.2) is induced by the homomorphism

$$\Phi_g: \mathrm{CH}_{K,R}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,R}^*(\mathrm{Spec} k).$$

Computing the respective characters we obtain $\mathrm{Res}_g([V_R]) = \psi^2[V_R]$ for the second Adams operation ψ^2 , thus

$$\Phi_g(c_2(V_R)) = c_2(\mathrm{Res}_g([V_R])) = c_2(\psi^2[V_R]) = 4c_2(V_R) = 0$$

with the last equality arising from $2 = 0$ in $\mathrm{CH}_K^*(\mathrm{Spec} k)$ by Lemma 2.1.2. It follows that

$$\mathrm{CH}_K^*(X)[b] \cong \mathrm{CH}_{K,R}^*(X) \xrightarrow{\Phi_f} \mathrm{CH}_{K,{}^2\mu_{2,2}}^*(X)$$

is an isomorphism. We have $\Phi_f(c_2(V_R)) = c_2(\mathrm{Res}_f([V_R])) = c_2({}^2V_{\mu_{2,2}})$, so b maps to $c_2({}^2V_{\mu_{2,2}})$ as claimed. \square

Lemma 2.4.2. *Let $K/L/k$ be separable field extensions with $[L:k] = 2$, put $R := R_{L/k}\mathbb{G}_m$, $N := {}^2\mu_{2m}$ or $N := {}^2\mu_{2,2}$ and $V_N := {}^2V_{\mu_{2m}}$ or $V_N := {}^2V_{\mu_{2,2}}$ respectively. Let W be a rank 2 representation of $R \times N$ such that for its restrictions to the factors $\rho_1: R \rightarrow R \times N$ and $\rho_2: N \rightarrow R \times N$ we have*

$$\mathrm{Res}_{\rho_1}([W]) = [V_R], \quad \mathrm{Res}_{\rho_2}([W]) = \psi^i[V_N]$$

for the i -th Adams operation ψ^i . Then

$$c_2(W) = c_2(V_R) \otimes 1 + i^2 \cdot 1 \otimes c_2(V_N) \in \mathrm{CH}_{K,R}^*(\mathrm{Spec} k) \otimes \mathrm{CH}_{K,N}^*(\mathrm{Spec} k) \cong \mathrm{CH}_{K,R \times N}^*(\mathrm{Spec} k).$$

Proof. By Propositions 2.3.2 and 2.4.1 the degree 2 part of $\mathrm{CH}_{K,R}^*(\mathrm{Spec} k) \otimes \mathrm{CH}_{K,N}^*(\mathrm{Spec} k)$ is spanned by $c_2(V_R) \otimes 1$ and $1 \otimes c_2(V_N)$ and since $2 = 0$ in $\mathrm{CH}_K^*(\mathrm{Spec} k)$ by Lemma 2.1.2, it follows that the map

$$(\Phi_{\rho_1}, \Phi_{\rho_2}): \mathrm{CH}_{K,R \times N}^2(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,R}^2(\mathrm{Spec} k) \times \mathrm{CH}_{K,N}^2(\mathrm{Spec} k)$$

is bijective. We have

$$\begin{aligned} \Phi_{\rho_1}(c_2(W)) &= c_2(\mathrm{Res}_{\rho_1}([W])) = c_2(V_R) = \Phi_{\rho_1}(c_2(V_R) \otimes 1 + i^2 \cdot 1 \otimes c_2(V_N)), \\ \Phi_{\rho_2}(c_2(W)) &= c_2(\mathrm{Res}_{\rho_2}([W])) = c_2(\psi^i[V_N]) = i^2 \cdot c_2(V_N) = \Phi_{\rho_2}(c_2(V_R) \otimes 1 + i^2 \cdot 1 \otimes c_2(V_N)). \end{aligned}$$

The claim follows. \square

2.5 Conormed characteristic sequence for quasi-split simple groups

In this section we introduce the main computational tools for the conormed Chow rings of a quasi-split simple group, namely the conormed characteristic sequence and the conormed extended characteristic sequence, and show that the latter one is exact while the former one is exact provided that the groups is either simply connected or adjoint. We also show that G satisfies the conormed Künneth formula for the direct product with an arbitrary $X \in \mathcal{S}m_k$. Here and below we use some standard notation related to quasi-split algebraic groups, see Sections A.1 and A.2 for a recollection.

Lemma 2.5.1. *Let G be a product of a finite number of quasi-split simple groups over a field k and let K/k be a separable field extension such that G_K is split. Then $B\pi_1(G)$ satisfies the Künneth formula for K/k in the sense of Definition 2.3.5.*

Proof. We can assume that G is a quasi-split simple group. The proof goes case by case on all the possible fundamental groups, see e.g. recollection (A.4.1) for a list.

If G is split, then $\pi_1(G) \cong \mu_l$ or $\pi_1(G) \cong \mu_2 \times \mu_2$ and it is well known that the classifying spaces of such groups satisfy Künneth formula for the Chow groups (see, e.g., [To14b, proof of Theorem 2.10]), thus also for the conormed Chow groups. So we can assume that G is not split.

Let L be the splitting field of G . Since G_K is split, we can assume that $L \subseteq K$. The field extension L/k is Galois and $[L : k] = 2$, if $\Delta(G) = {}^2A_n, {}^2E_6$ or 2D_n , and $[L : k] = 3$ or 6 , if $\Delta(G) = {}^3D_4$ or $\Delta(G) = {}^6D_4$ respectively.

If $[L : k] = 6$, then $\mathrm{CH}_K^*(\mathrm{Spec} k) = 0$ by Lemma 2.1.2 and there is nothing to prove.

If $[L : k] = 3$, then $\Delta(G) = {}^3D_4$ and $\pi_1(G)$ is either trivial or $\pi_1(G) \cong {}^3\mu_{2,2}$, so its order is invertible in $\mathrm{CH}_K^*(\mathrm{Spec} k)$ by Lemma 2.1.2. The claim follows from Lemma 2.2.7.

If $[L : k] = 2$, then $\pi_1(G) \cong {}^2\mu_l$ for some $l \in \mathbb{N}$ or $\pi_1(G) \cong {}^2\mu_{2,2}$. The claim follows from Proposition 2.4.1. \square

Remark 2.5.2. Let G be a quasi-split semisimple group over a field k and K/k be a separable field extension K/k such that G_K is split. We do not know whether $B\pi_1(G)$ satisfies the Künneth formula for K/k , but it seems highly plausible.

Proposition 2.5.3. *Let G be a quasi-split simple group over a field k and K/k be a separable field extension such that G_K is split. Then for $X \in \mathrm{Sm}_k$ the Künneth homomorphism*

$$\mathrm{CH}_K^*(X) \otimes \mathrm{CH}_K^*(G) \rightarrow \mathrm{CH}_K^*(X \times G), \quad x \otimes y \mapsto p_X^*(x) \cdot p_G^*(y),$$

is an isomorphism. Here $p_X : X \times G \rightarrow X$, $p_G : X \times G \rightarrow G$ are the projections. In particular,

$$\Delta : \mathrm{CH}_K^*(G) \xrightarrow{m^*} \mathrm{CH}_K^*(G \times G) \cong \mathrm{CH}_K^*(G) \otimes \mathrm{CH}_K^*(G)$$

for the multiplication $m : G \times G \rightarrow G$ equips $\mathrm{CH}_K^*(G)$ with the structure of a Hopf algebra.

Proof. Let $\tilde{G} \rightarrow G$ be the simply connected cover and $\tilde{T} \leq \tilde{B} \leq \tilde{G}$ be a maximal torus and a Borel subgroup. Lemma A.2.3 yields that \tilde{T} is a quasi-trivial torus and Lemma 2.5.1 yields that $B\pi_1(G)$ satisfies the Künneth formula for K/k . Then it follows from Proposition 2.3.6 that there are isomorphisms

$$\begin{aligned} & (\mathrm{CH}_{K,\tilde{T}}^*(\tilde{G}) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k)) \otimes_{\mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,\pi_1(G)}^*(\tilde{G}), \\ & (\mathrm{CH}_{K,\tilde{T}}^*(X \times \tilde{G}) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k)) \otimes_{\mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k)} \mathrm{CH}_K^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,\pi_1(G)}^*(X \times \tilde{G}) \end{aligned}$$

with X equipped with the trivial action. Lemma 2.2.6 yields

$$\mathrm{CH}_{K,\pi_1(G)}^*(\tilde{G}) \cong \mathrm{CH}_K^*(G), \quad \mathrm{CH}_{K,\pi_1(G)}^*(X \times \tilde{G}) \cong \mathrm{CH}_K^*(X \times G), \quad \mathrm{CH}_{K,\tilde{T}}^*(X \times \tilde{G}) \cong \mathrm{CH}_K^*(X \times \tilde{G}/\tilde{T}).$$

Note that

$$\mathrm{CH}_K^*(\tilde{G}/\tilde{T}) \cong \mathrm{CH}_K^*(\tilde{G}/\tilde{B}), \quad \mathrm{CH}_K^*(X \times \tilde{G}/\tilde{T}) \cong \mathrm{CH}_K^*(X \times \tilde{G}/\tilde{B})$$

by the homotopy invariance property. It follows from [CM06, Lemma 29] that the (integral) Chow motive $\mathcal{M}(\tilde{G}/\tilde{B})$ splits as a direct sum of $\mathcal{M}(\mathrm{Spec} F, l)$ for some $l \in \mathbb{N}_0$ and field extensions $K/F/k$. Lemma 2.1.4 yields $\mathcal{M}_K(\mathrm{Spec} F, l) = 0$ for $K \supseteq F \supsetneq k$, so we obtain a decomposition

$$\mathcal{M}_K(\tilde{G}/\tilde{B}) \cong \bigoplus_{i \in I} \mathcal{M}_K(\mathrm{Spec} k, l_i)$$

for some $l_i \in \mathbb{N}_0$, yielding a Künneth isomorphism

$$\mathrm{CH}_K^*(X) \otimes \mathrm{CH}_K^*(\tilde{G}/\tilde{B}) \cong \mathrm{CH}_K^*(X \times \tilde{G}/\tilde{B}).$$

Combining all the isomorphisms above we get the claim. \square

Definition 2.5.4. Consider a sequence

$$\mathcal{A}^* \xrightarrow{c} \mathcal{B}^* \xrightarrow{\phi^*} \mathcal{C}^* \rightarrow 0$$

of $\mathbb{Z}_{\geq 0}$ -graded commutative rings with c and ϕ^* being graded degree 0 homomorphisms. We say that this sequence is *exact*, if ϕ^* is surjective and $c(\mathcal{A}^{>0})$ generates $\ker \phi^*$ as an ideal.

Definition 2.5.5. Let K/k be a separable extension of fields, let E be a G -torsor for a quasi-split group G over k , and let $T \leq B \leq G$ be a maximal torus and a Borel subgroup. Then the *characteristic map* is the composition

$$c: \mathrm{CH}_{K,T}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,T}^*(E) \cong \mathrm{CH}_K^*(E/T) \cong \mathrm{CH}_K^*(E/B),$$

where the first map is the pullback for the structure map, the middle isomorphism is given by Lemma 2.2.6 and the last isomorphism is the homotopy invariance property for the vector bundle torsor $E/T \rightarrow E/B$. The *characteristic sequence* is the sequence

$$\mathrm{CH}_{K,T}^*(\mathrm{Spec} k) \xrightarrow{c} \mathrm{CH}_K^*(E/B) \xrightarrow{\phi^*} \mathrm{CH}_K^*(E) \rightarrow 0$$

where $\phi: E \rightarrow E/B$ is the projection.

Suppose that G is a quasi-split semisimple group such that $B\pi_1(G)$ satisfies the Künneth formula for K/k , let $\tilde{G} \rightarrow G$ be its simply connected cover and $\tilde{T} \leq \tilde{B} \leq \tilde{G}$ be the maximal torus and the Borel subgroup lying over $T \leq B$. Then the *extended characteristic sequence* is the sequence

$$\mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \xrightarrow{\hat{c}} \mathrm{CH}_K^*(\tilde{G}/\tilde{B}) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \xrightarrow{\hat{\phi}} \mathrm{CH}_K^*(G) \rightarrow 0$$

where

- \hat{c} is the composition

$$\begin{aligned} \mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) &\xrightarrow{\hat{\nu}} \mathrm{CH}_{K,\tilde{T} \times \pi_1(G)}^*(\mathrm{Spec} k) \cong \mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \xrightarrow{\tilde{c} \otimes \mathrm{id}} \\ &\xrightarrow{\tilde{c} \otimes \mathrm{id}} \mathrm{CH}_K^*(\tilde{G}/\tilde{B}) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \end{aligned}$$

with $\nu: \tilde{T} \times \pi_1(G) \rightarrow \tilde{T}$ being the group homomorphisms given by $\nu(t, s) = t \cdot s^{-1}$, and \tilde{c} being the characteristic map for \tilde{G} ,

- $\hat{\phi}(x, \alpha) = \phi_G^*(x) \cdot p(\alpha)$ with $\phi_G: G \rightarrow G/B \cong \tilde{G}/\tilde{B}$ being the projection combined with the isomorphism $\tilde{G}/\tilde{B} \cong G/B$, and p being the composition

$$\mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \xrightarrow{p_G^*} \mathrm{CH}_{K,\pi_1(G)}^*(\tilde{G}) \cong \mathrm{CH}_K^*(G)$$

for the structure map $p_G: \tilde{G} \rightarrow \mathrm{Spec} k$ and the isomorphism given by Lemma 2.2.6.

Theorem 2.5.6. Let G be a quasi-split algebraic group over a field k , let $T \leq B \leq G$ be a maximal torus and a Borel subgroup, and let K/k be a separable extension of fields such that T_K is split.

1. If T is quasi-trivial, then for a G -torsor E the characteristic sequence

$$\mathrm{CH}_{K,T}^*(\mathrm{Spec} k) \xrightarrow{c} \mathrm{CH}_K^*(E/B) \xrightarrow{\phi^*} \mathrm{CH}_K^*(E) \rightarrow 0$$

is exact. In particular, this applies to a split G and, by Lemma A.2.3, to a simply connected or an adjoint quasi-split semisimple group G .

2. Let G be a quasi-split semisimple group, $\tilde{G} \rightarrow G$ be its simply connected cover and $\tilde{T} \leq \tilde{B} \leq \tilde{G}$ be the maximal torus and the Borel subgroup lying over $T \leq B$. Suppose that $B\pi_1(G)$ satisfies the Künneth formula for K/k . Then the extended characteristic sequence

$$\mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \xrightarrow{\hat{c}} \mathrm{CH}_K^*(\tilde{G}/\tilde{B}) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \xrightarrow{\hat{\phi}} \mathrm{CH}_K^*(G) \rightarrow 0$$

is exact. In particular, by Lemma 2.5.1 this applies to G being a product of quasi-split simple groups.

Proof. (1) Corollary 2.3.3 yields an exact sequence

$$\mathrm{CH}_{K,T}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{T,K}^*(E) \xrightarrow{\Phi_T} \mathrm{CH}_K^*(E) \rightarrow 0$$

where the first morphism is the pullback for the projection $E \rightarrow \mathrm{Spec} k$. The claim follows via the identification $\mathrm{CH}_{K,T}^*(E) \cong \mathrm{CH}_K^*(E/B)$ as in Definition 2.5.5.

(2) Proposition 2.3.6 applied to $X := \tilde{G}$ with the action of \tilde{T} and the subgroup $\pi_1(G) \leq \tilde{T}$ (we use the notation of Definition 2.5.5 for the simply connected cover \tilde{G} of G and for its subgroups) yields an exact sequence

$$\mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,\tilde{T}}^*(\tilde{G}) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,\pi_1(G)}^*(\tilde{G}) \rightarrow 0$$

where the first homomorphism is the composition

$$\begin{aligned} \mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) &\xrightarrow{\Phi_\nu} \mathrm{CH}_{K,\tilde{T} \times \pi_1(G)}^*(\mathrm{Spec} k) \cong \mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \rightarrow \\ &\xrightarrow{p_{\tilde{G}}^* \otimes \mathrm{id}} \mathrm{CH}_{K,\tilde{T}}^*(\tilde{G}) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \end{aligned}$$

and the second homomorphism is given by $(x, \alpha) \mapsto p_G^*(\alpha) \cdot \Phi_\rho(x)$. Here $\nu: \tilde{T} \times \pi_1(G) \rightarrow \tilde{T}$ is the homomorphism given by $\nu(t, s) = ts^{-1}$, $\rho: \pi_1(G) \rightarrow \tilde{T}$ is the embedding, and $p_{\tilde{G}}: \tilde{G} \rightarrow \mathrm{Spec} k$ is the structure morphism. Identifying $\mathrm{CH}_{K,\tilde{T}}^*(\tilde{G}) \cong \mathrm{CH}_K^*(\tilde{G}/\tilde{B})$ and $\mathrm{CH}_{K,\pi_1(G)}^*(\tilde{G}) \cong \mathrm{CH}_K^*(G)$ as in Definition 2.5.5 we obtain the claim. \square

Remark 2.5.7. If G is a split group, then Theorem 2.5.6 is well-known and goes back to [Gro58].

2.6 Conormed Chow ring of a quasi-split group G via G/B

In this section we use the conormed extended characteristic sequence to give a presentation for $\mathrm{CH}_K^*(G)$ as a quotient of $\mathrm{CH}_K^*(G/B)$ or $\mathrm{CH}_K^*(G/B)[x]$ modulo some explicit ideal. Here and below we use some standard notation for vector bundles over homogeneous spaces recalled in Section A.3, in particular, for the maximal torus \tilde{T} in the simply connected cover $\tilde{G} \rightarrow G$ lying over a maximal torus $T \leq B$ we have

- line bundle $\mathcal{L}(\varpi)$ over G/B associated with a Galois-invariant weight $\varpi \in \mathcal{X}^*(\tilde{T})$,
- vector bundle $\mathcal{V}(\mathcal{S})$ over G/B associated with a Galois-invariant finite set of weights $\mathcal{S} \subseteq \mathcal{X}^*(\tilde{T})$,
- line bundle $\mathcal{L}(\bar{\varpi})$ over G associated with a Galois-invariant element $\bar{\varpi} \in \mathcal{X}^*(\tilde{T})/\mathcal{X}^*(T)$.

Furthermore, we use the standard numbering for simple roots α_i and fundamental weights ϖ_i recalled in Section A.2, and shorten the notation as $\mathcal{L}_i := \mathcal{L}(\varpi_i)$, $\mathcal{V}_{i_1, i_2, \dots, i_j} := \mathcal{V}(\{\varpi_{i_1}, \varpi_{i_2}, \dots, \varpi_{i_j}\})$.

Lemma 2.6.1 (see also [KaZ13, Example 3.12]). *Let G be a quasi-split reductive group over a field k , let K/k be the splitting field of G and let $P \leq G$ be a parabolic subgroup. Then for every $n \in \mathbb{Z}$ and the canonical projection $\rho: (G/P)_K \rightarrow G/P$ the pullback homomorphism*

$$\rho^*: \mathrm{CH}^*(G/P) \otimes \mathbb{Z}/n\mathbb{Z} \rightarrow \mathrm{CH}^*((G/P)_K) \otimes \mathbb{Z}/n\mathbb{Z}$$

is injective and $\rho^(\mathrm{CH}^*(G/P) \otimes \mathbb{Z}/n\mathbb{Z}) = (\mathrm{CH}^*((G/P)_K) \otimes \mathbb{Z}/n\mathbb{Z})^{\mathrm{Gal}(K/k)}$.*

Proof. It follows from [CM06, Lemma 29] that there is a motivic decomposition

$$\mathcal{M}(G/P) \cong \bigoplus_{i \in I} \mathcal{M}(\mathrm{Spec} F_i, l_i)$$

for some $l_i \in \mathbb{N}_0$ and field extensions $K/F_i/k$. At the same time we have

$$\mathcal{M}((G/P)_K) \cong \bigoplus_{i \in I} \mathcal{M}(\mathrm{Spec}(F_i \otimes_k K), l_i) \cong \bigoplus_{i \in I} \mathcal{M}(\mathrm{Spec} K, l_i)^{\oplus [F_i:k]}.$$

The pullback ρ^* is given by the direct sum of diagonal embeddings

$$\rho^* : \mathrm{CH}^*(G/P) \otimes \mathbb{Z}/n\mathbb{Z} \cong \bigoplus_{i \in I} \mathbb{Z}/n\mathbb{Z} \xrightarrow{\oplus \mathrm{diag}} \bigoplus_{i \in I} (\mathbb{Z}/n\mathbb{Z})^{\oplus [F_i:k]} \cong \mathrm{CH}^*((G/P)_K) \otimes \mathbb{Z}/n\mathbb{Z}.$$

The group $\mathrm{Gal}(K/k)$ acts on $\bigoplus_{i \in I} (\mathbb{Z}/n\mathbb{Z})^{\oplus [F_i:k]}$ simultaneously permuting the factors of each summand $(\mathbb{Z}/n\mathbb{Z})^{\oplus [F_i:k]}$, so the claim follows. \square

Lemma 2.6.2. *Let G be a quasi-split simple group over a field k of type $\Delta(G) = {}^2\mathbf{A}_{2r-1}$, let $B \leq G$ be a Borel subgroup and let K/k be the splitting field of G . Then in $\mathrm{CH}_K^*(G/B)$ we have the following:*

1. $\mathrm{CH}_K^1(G/B) = \mathbb{F}_2 \cdot c_1(\mathcal{L}_r)$,
2. $c_1(\mathcal{L}_r)^2 = 0$,
3. $\mathrm{CH}_K^2(G/B) = \mathbb{F}_2 \cdot c_2(\mathcal{V}_{1,2r-1}) \oplus \mathbb{F}_2 \cdot c_2(\mathcal{V}_{2,2r-2}) \oplus \dots \oplus \mathbb{F}_2 \cdot c_2(\mathcal{V}_{r-1,r+1})$.

Proof. We have $[K : k] = 2$ and $\mathrm{CH}_K^*(\mathrm{Spec} k) \cong \mathbb{F}_2$ by Lemma 2.1.2. Since G_K is split, $(G/B)_K$ is isomorphic to the variety of full flags in a vector space of dimension $2r$, and

$$\mathrm{CH}^*((G/B)_K) \cong \mathbb{Z}[x_1, x_2, \dots, x_{2r}] / (\sigma_1, \sigma_2, \dots, \sigma_{2r}), \quad x_i \mapsto c_1(\mathcal{T}_i),$$

where σ_i is the i -th elementary symmetric polynomial in x_1, x_2, \dots, x_{2r} and \mathcal{T}_i is the i -th tautological line bundle over the full flag variety [Fu98, Example 3.3.5]. In the notation of (A.2.5) we have $\mathcal{L}(\epsilon_i) = \mathcal{T}_i$, thus the nontrivial element $\tau \in \mathrm{Gal}(K/k)$ acts on $\mathrm{CH}^*((G/B)_K)$ via

$$\tau(\bar{x}_i) = \tau(c_1(\mathcal{L}(\epsilon_i))) = c_1(\mathcal{L}(\tau(\epsilon_i))) = c_1(\mathcal{L}(-\epsilon_{2r+1-i})) = -\bar{x}_{2r+1-i}$$

with \bar{x}_i being the image of x_i in the quotient ring. Lemma 2.6.1 yields that the pullback homomorphism for the morphism $(G/B)_K \rightarrow G/B$ induces an isomorphism

$$\mathrm{CH}^*(G/B) \cong (\mathbb{Z}[x_1, x_2, \dots, x_{2r}] / (\sigma_1, \sigma_2, \dots, \sigma_{2r}))^{\mathrm{Gal}(K/k)}.$$

Hence

$$\mathrm{CH}_K^*(G/B) \cong (\mathbb{Z}[x_1, x_2, \dots, x_{2r}] / (\sigma_1, \sigma_2, \dots, \sigma_{2r}))^{\mathrm{Gal}(K/k)} / (\mathrm{id} + \tau)(\mathbb{Z}[x_1, x_2, \dots, x_{2r}] / (\sigma_1, \sigma_2, \dots, \sigma_{2r})).$$

Furthermore, it follows from the formulae in (A.2.5) for the fundamental weights that

$$(\mathcal{L}_r)_K \cong \bigotimes_{i=1}^r \mathcal{T}_i, \quad (\mathcal{V}_{i,2r-i})_K \cong \bigotimes_{j=1}^i \mathcal{T}_j \oplus \bigotimes_{j=1}^{2r-i} \mathcal{T}_j$$

and in $\mathrm{CH}^*(G/B) \cong \mathrm{CH}^*((G/B)_K)^{\mathrm{Gal}(K/k)}$ we have

$$c_1(\mathcal{L}_r) = \bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_r, \quad c_2(\mathcal{V}_{i,2r-i}) = (\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_i) \cdot (\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_{2r-i}).$$

The claims of the lemma follow by straightforward calculations with polynomials which we provide below.

The first claim follows from the $\mathrm{Gal}(K/k)$ -invariant decomposition

$$\mathrm{CH}^1((G/B)_K) \cong \left(\bigoplus_{i=1}^{2r} \mathbb{Z}x_i \right) / \mathbb{Z}\sigma_1 = \mathbb{Z} \cdot (\bar{x}_1 + \dots + \bar{x}_r) \oplus \left(\bigoplus_{i=1}^{r-1} [\mathbb{Z}\bar{x}_i \oplus \mathbb{Z} \cdot (-\bar{x}_{2r-i})] \right),$$

here the first summand is a rank 1 submodule with the trivial action of $\mathrm{Gal}(K/k)$, and then there are $r-1$ rank 2 submodules with the permutation action of $\mathrm{Gal}(K/k)$. Then $\mathrm{CH}_K^1(G/B)$ is of rank 1 with the generator given by the image of $c_1(\mathcal{L}_r) = \bar{x}_1 + \dots + \bar{x}_r$.

For the second claim note that

$$(x_1 + x_2 + \dots + x_r)^2 = (x_1 + x_2 + \dots + x_r) \cdot \sigma_1 - \sigma_2 + (\mathrm{id} + \tau)(\alpha)$$

for some $\alpha \in \mathbb{Z}[x_1, x_2, \dots, x_{2r}]$, thus $c_1(\mathcal{L}_r)^2 = 0$ in $\text{CH}_K^*(G/B)$.

For the last claim consider the following decomposition of the degree 2 part of $\mathbb{Z}[x_1, x_2, \dots, x_{2r}]$ into $\text{Gal}(K/k)$ -invariant submodules:

$$\begin{aligned} \mathbb{Z}[x_1, x_2, \dots, x_{2r}]^{(2)} = & \left(\bigoplus_{i=1}^r [\mathbb{Z}x_i\sigma_1 \oplus \mathbb{Z}x_{2r-i+1}\sigma_1] \right) \oplus [\mathbb{Z}\sigma_2] \oplus \\ & \oplus \left(\bigoplus_{i=1}^{r-1} [\mathbb{Z}x_ix_{2r-i+1}] \right) \oplus \left(\bigoplus_{1 \leq i < j < 2r-i+1} [\mathbb{Z}x_ix_j \oplus \mathbb{Z}x_{2r-i+1}x_{2r-j+1}] \right), \end{aligned}$$

here the square brackets group the $\text{Gal}(K/k)$ -invariant submodules, i.e. we have r submodules of rank 2 with the permutation action of $\text{Gal}(K/k)$, then one of rank 1 with the trivial action of $\text{Gal}(K/k)$, then $r-1$ submodules of rank 1 again with the trivial action, and then $r(r-1)$ submodules of rank 2 with the permutation action. It follows that

$$\text{CH}^2((G/B)_K) \cong \left(\bigoplus_{i=1}^{r-1} [\mathbb{Z}\bar{x}_i\bar{x}_{2r-i+1}] \right) \oplus \left(\bigoplus_{1 \leq i < j < 2r-i+1} [\mathbb{Z}\bar{x}_i\bar{x}_j \oplus \mathbb{Z}\bar{x}_{2r-i+1}\bar{x}_{2r-j+1}] \right),$$

and $\text{CH}_K^2(G/B)$ has an additive basis given by the images in the respective quotient ring of $\bar{x}_i\bar{x}_{2r-i+1}$ for $1 \leq i \leq r-1$. We have

$$\begin{aligned} c_2(\mathcal{V}_{i,2r-i}) = & (\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_i) \cdot (\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_{2r-i}) = (\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_i) \cdot (\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_{2r-i} - \bar{\sigma}_1) = \\ = & -(\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_i) \cdot (\bar{x}_{2r-i+1} + \bar{x}_{2r-i+2} + \dots + \bar{x}_{2r}) \end{aligned}$$

in $\text{CH}^2(G/B)$ yielding that in $\text{CH}_K^2(G/B)$ we have $c_2(\mathcal{V}_{i,2r-i}) = \sum_{j=1}^i \bar{x}_j\bar{x}_{2r-j+1}$. The claim follows. \square

Theorem 2.6.3. *Let G be a non-split quasi-split simple algebraic group over a field k , let $T \leq B \leq G$ be a maximal torus and a Borel subgroup, and let K/k be a separable field extension such that T_K is split. Put $\mathcal{R}^* := \text{CH}_K^*(G/B)$. If $\Delta(G) = {}^6\text{D}_4$, then $\text{CH}_K^*(G) = 0$, otherwise, depending on $(\Delta(G), \pi_1(G))$, we have $\text{CH}_K^*(G) \cong \mathcal{R}^*/I$ or $\text{CH}_K^*(G) \cong \mathcal{R}^*[x]/I$ with the generators of the ideal I given in the following table.*

$\Delta(G)$	$\pi_1(G)$	\mathcal{R}^* or $\mathcal{R}^*[x]$	generators of I
${}^2\text{A}_{2r}, r \geq 1$	$2\mu_l, l \mid 2r+1$	\mathcal{R}^*	$\{c_2(\mathcal{V}_{i,2r+1-i})\}_{i=1}^r$
${}^2\text{A}_{2r-1}, r \geq 2$	$2\mu_{2m+1}, 2m+1 \mid 2r$	\mathcal{R}^*	$\{c_1(\mathcal{L}_r)\} \cup \{c_2(\mathcal{V}_{i,2r-i})\}_{i=1}^{r-1}$
	$2\mu_{2m}, m \text{ even}, \frac{r}{m} \text{ odd}$	\mathcal{R}^*	\mathcal{S}_{2r-1}
	$2\mu_{2m}, m \text{ odd}, \frac{r}{m} \text{ odd}$	\mathcal{R}^*	$\{c_2(\mathcal{V}_{i,2r-i})\}_{i=1}^{r-1}$
	$2\mu_{2m}, m \text{ even}, \frac{r}{m} \text{ even}$	$\mathcal{R}^*[x]$	$\{c_1(\mathcal{L}_r), x^2\} \cup \mathcal{S}_{2r-1}$
	$2\mu_{2m}, m \text{ odd}, \frac{r}{m} \text{ even}$	$\mathcal{R}^*[x]$	$\{c_1(\mathcal{L}_r), x^2 + c_2(\mathcal{V}_{1,2r-1})\} \cup \mathcal{S}_{2r-1}$
${}^2\text{D}_n, n \geq 3$	1	\mathcal{R}^*	$\{c_1(\mathcal{L}_i)\}_{i=1}^{n-2} \cup \{c_2(\mathcal{V}_{n-1,n})\}$
	μ_2	$\mathcal{R}^*[x]$	$\{c_1(\mathcal{L}_i)\}_{i=1}^{n-2} \cup \{x^2 + c_2(\mathcal{V}_{n-1,n})\}$
${}^2\text{D}_{2r}, r \geq 2$	$2\mu_{2,2}$	\mathcal{R}^*	$\{c_1(\mathcal{L}_i)\}_{i=1}^{2r-2}$
${}^2\text{D}_{2r+1}, r \geq 1$	$2\mu_4$	\mathcal{R}^*	$\{c_1(\mathcal{L}_1)^2\} \cup \{c_1(\mathcal{L}_{2i})\}_{i=1}^{r-1} \cup \{c_1(\mathcal{L}_{2i+1}) + c_1(\mathcal{L}_1)\}_{i=1}^{r-1}$
${}^3\text{D}_4$	$1, {}^3\mu_{2,2}$	\mathcal{R}^*	$\{c_1(\mathcal{L}_2), c_3(\mathcal{V}_{1,3,4})\}$
${}^2\text{E}_6$	$1, 2\mu_3$	\mathcal{R}^*	$\{c_1(\mathcal{L}_2), c_1(\mathcal{L}_4), c_2(\mathcal{V}_{1,6}), c_2(\mathcal{V}_{3,5})\}$

Here $\mathcal{S}_{2r-1} = \{c_2(\mathcal{V}_{2i,2r-2i})\}_{i=1}^{\lfloor (r-1)/2 \rfloor} \cup \{c_2(\mathcal{V}_{2i+1,2r-2i-1}) + c_2(\mathcal{V}_{1,2r-1})\}_{i=1}^{\lfloor r/2 \rfloor - 1}$. The isomorphisms are induced by the pullback homomorphisms $\mathcal{R}^* = \mathrm{CH}_K^*(G/B) \rightarrow \mathrm{CH}_K^*(G)$ for the projections $G \rightarrow G/B$ and by $x \mapsto c_1(\mathcal{L}(\bar{\omega}_m))$ and $x \mapsto c_1(\mathcal{L}(\bar{\omega}_{n-1}))$ in the cases $({}^2\mathbf{A}_{2r-1}, {}^2\mu_{2m})$ and $({}^2\mathbf{D}_n, \mu_2)$ respectively.

Proof. Let $\tilde{G} \rightarrow G$ be the simply connected cover and $\tilde{T} \leq \tilde{B}$ be the maximal torus and the Borel subgroups lying over $T \leq B$. Theorem 2.5.6 yields an isomorphism

$$\mathrm{CH}_K^*(G) \cong (\mathcal{R}^* \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k))/I \quad (2.6.3.1)$$

where I denotes the ideal generated by $\hat{c}(\mathrm{CH}_{K,\tilde{T}}^{>0}(\mathrm{Spec} k))$ for the extended characteristic map

$$\begin{aligned} \hat{c}: \mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) &\xrightarrow{\Phi_\nu} \mathrm{CH}_{K,\tilde{T} \times \pi_1(G)}^*(\mathrm{Spec} k) \xrightarrow{\cong} \mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \rightarrow \\ &\xrightarrow{\tilde{c} \otimes \mathrm{id}} \mathrm{CH}_K^*(\tilde{G}/\tilde{B}) \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) = \mathcal{R}^* \otimes \mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k), \end{aligned}$$

where $\nu: \tilde{T} \times \pi_1(G) \rightarrow \tilde{T}$ is the homomorphism given by $(t, s) \mapsto t \cdot s^{-1}$. The torus \tilde{T} is quasi-trivial and decomposes into a product of the Weil restrictions of tori corresponding to the orbits of the Galois action on $\Delta(G)$. The rings $\mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k)$ and $\mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k)$ were computed in Propositions 2.3.2 and 2.4.1 respectively. The claim of the theorem then follows from an explicit computation of the homomorphism \hat{c} plugged into isomorphism (2.6.3.1), we provide below the details of these computations for each $(\Delta(G), \pi_1(G))$ case by case. Put

$$L_i := L(\varpi_i), \quad V_{i,j} := V(\{\varpi_i, \varpi_j\}), \quad V_{i,j,r} := V(\{\varpi_i, \varpi_j, \varpi_r\})$$

for the respective representations of \tilde{T} in the notation of (A.1.4) and note that $\tilde{c}(c_1(L_i)) = c_1(\mathcal{L}_i)$, $\tilde{c}(c_2(V_{i,j})) = c_2(\mathcal{V}_{i,j})$ and $\tilde{c}(c_3(V_{i,j,r})) = c_3(\mathcal{V}_{i,j,r})$, so it suffices to compute Φ_ν . In order to do this we use the formulae from (A.2.5) for the homomorphism of lattices $\mathcal{X}^*(\tilde{T}) \rightarrow \mathcal{X}^*(\pi_1(G))$ dual to the embedding $\pi_1(G) \leq \tilde{T}$. We denote by $L \subseteq K$ the splitting field of \tilde{T} which is a Galois extension L/k with $[L:k] = 2$ for $\Delta(G) = {}^2\mathbf{A}_n, {}^2\mathbf{D}_n$ or ${}^2\mathbf{E}_6$, $[L:k] = 3$ for $\Delta(G) = {}^3\mathbf{D}_4$ and $[L:k] = 6$ for $\Delta(G) = {}^6\mathbf{D}_4$.

⁶D₄. We have $[L:k] = 6$. Lemma 2.1.2 yields $\mathrm{CH}_K^*(\mathrm{Spec} k) = 0$ and $\mathrm{CH}_K^*(X) = 0$ for every $X \in \mathcal{S}m_k$.
²A_{2r}. For $1 \leq i \leq 2r$ we have $\tau(\varpi_i) = \varpi_{2r+1-i}$ for the nontrivial element $\tau \in \mathrm{Gal}(L/k)$, and

$$\mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[b_1, b_2, \dots, b_r], \quad b_i \mapsto c_2(V_{i,2r+1-i}). \quad (2.6.3.2)$$

The order of $\pi_1(G)$ is odd, thus $\mathrm{CH}_{K,\pi_1(G)}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)$ by Lemmas 2.1.2 and 2.2.7 and the claim follows.

²A_{2r-1}. For the nontrivial element $\tau \in \mathrm{Gal}(L/k)$ we have

$$\tau(\varpi_r) = \varpi_r, \quad \tau(\varpi_i) = \varpi_{2r-i}, \quad 1 \leq i \leq 2r-1, \quad i \neq r.$$

It follows that

$$\mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[y, b_1, \dots, b_{r-1}], \quad y \mapsto c_1(L_r), \quad b_i \mapsto c_2(V_{i,2r-i}). \quad (2.6.3.3)$$

If $\pi_1(G) = {}^2\mu_{2m+1}$, then $\mathrm{CH}_{K,{}^2\mu_{2m+1}}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)$ by Lemmas 2.1.2 and 2.2.7 and the claim follows.

Now suppose $\pi_1(G) = {}^2\mu_{2m}$. We need to compute the homomorphism

$$\Phi_\nu: \mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \rightarrow \mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \otimes \mathrm{CH}_{K,{}^2\mu_{2m}}^*(\mathrm{Spec} k)$$

with $\mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k)$ given by isomorphism (2.6.3.3) and

$$\mathrm{CH}_{K,{}^2\mu_{2m}}^*(\mathrm{Spec} k) \cong \mathrm{CH}_K^*(\mathrm{Spec} k)[x, b]/(x^2 + m^2 \cdot b), \quad x \mapsto c_1({}^2\Lambda_{\mu_{2m}}^\pm), \quad b \mapsto c_2({}^2V_{\mu_{2m}}),$$

by Proposition 2.4.1. The homomorphism

$$\mathrm{Res}_\nu: \mathrm{Rep}(\tilde{T}) \cong \mathbb{Z}[\mathcal{X}^*(\tilde{T})]^{\mathrm{Gal}(L/k)} \rightarrow \mathbb{Z}[\mathcal{X}^*(\tilde{T}) \oplus \mathcal{X}^*({}^2\mu_{2m})]^{\mathrm{Gal}(L/k)} \cong \mathrm{Rep}(\tilde{T} \times {}^2\mu_{2m})$$

is induced by the morphism of lattices

$$\begin{aligned} \mathcal{X}^*(\nu): \mathcal{X}^*(\tilde{T}) &= \bigoplus_{i=1}^{2r-1} \mathbb{Z} \cdot \varpi_i \rightarrow \bigoplus_{i=1}^{2r-1} \mathbb{Z} \cdot \varpi_i \oplus \mathbb{Z}/2m\mathbb{Z} \cdot \bar{\varpi}_1 = \mathcal{X}^*(\tilde{T}) \oplus \mathcal{X}^*({}^2\mu_{2m}), \\ \varpi_i &\mapsto \varpi_i - i\bar{\varpi}_1, \quad 1 \leq i \leq 2r-1. \end{aligned}$$

Then $\text{Res}_\nu([L_r]) = x^{\varpi_r - r\bar{\varpi}_1} = x^{\varpi_r} \cdot (x^{m\bar{\varpi}_1})^{\frac{r}{m}} = [L_r] \otimes [{}^2\Lambda_{\mu_{2m}}^\pm]^{\otimes \frac{r}{m}}$ and

$$\Phi_\nu(y) = y \otimes 1 + \frac{r}{m} \cdot 1 \otimes x.$$

For $1 \leq i \leq r-1$ we have

$$\text{Res}_\nu([V_{i,2r-i}]) = \text{Res}_\nu(x^{\varpi_i} + x^{\varpi_{2r-i}}) = x^{\varpi_i - i\bar{\varpi}_1} + x^{\varpi_{2r-i} - (2r-i)\bar{\varpi}_1}.$$

This element corresponds to a rank 2 representation W of $R \times {}^2\mu_{2m}$, where $R := R_{L/k} \mathbb{G}_m$. Restricting W to R and ${}^2\mu_{2m}$ respectively we obtain $W|_R \cong V_R$ and $W|_{{}^2\mu_{2m}} \cong \psi^i({}^2V_{\mu_{2m}})$ for the i -th Adams operation ψ^i . Then Lemma 2.4.2 yields

$$c_2(W) = c_2(V_R) \otimes 1 + i^2 \cdot 1 \otimes c_2({}^2V_{\mu_{2m}}) \in \text{CH}_{K,R}^*(\text{Spec } k) \otimes \text{CH}_{K,{}^2\mu_{2m}}^*(\text{Spec } k) \cong \text{CH}_{K,R \times {}^2\mu_{2m}}^*(\text{Spec } k).$$

Hence we have

$$\Phi_\nu(b_i) = c_2(\text{Res}_\nu([V_{i,2r-i}])) = c_2(V_{i,2r-i}) \otimes 1 + i^2 \cdot 1 \otimes c_2({}^2V_{\mu_{2m}}) = b_i \otimes 1 + i^2 \cdot 1 \otimes b.$$

Combining all the above we obtain $\text{CH}_K^*(G) \cong \mathcal{R}^*[x, b]/I$ with

$$I = I \left(\left\{ x^2 + m^2 \cdot b, c_1(\mathcal{L}_r) + \frac{r}{m} \cdot x \right\} \cup \left\{ c_2(\mathcal{V}_{2i,2r-2i}) \right\}_{i=1}^{\lfloor (r-1)/2 \rfloor} \cup \left\{ b + c_2(\mathcal{V}_{2i+1,2r-2i-1}) \right\}_{i=0}^{\lfloor r/2 \rfloor - 1} \right).$$

Then $I = I(\{x^2 + m^2 \cdot b, c_1(\mathcal{L}_r) + \frac{r}{m} \cdot x, b + c_2(\mathcal{V}_{1,2r-1})\} \cup \mathcal{S}_{2r-1})$. If $\frac{r}{m}$ is even, then the claim follows immediately, and if $\frac{r}{m}$ is odd, then the claim follows from Lemma 2.6.2.(2).

2D_n. For the nontrivial element $\tau \in \text{Gal}(L/k)$ we have

$$\tau(\varpi_i) = \varpi_i, \quad 1 \leq i \leq n-2, \quad \tau(\varpi_{n-1}) = \varpi_n, \quad \tau(\varpi_n) = \varpi_{n-1}.$$

It follows that

$$\text{CH}_{K,\tilde{T}}^*(\text{Spec } k) \cong \text{CH}_K^*(\text{Spec } k)[x_1, \dots, x_{n-2}, b], \quad x_i \mapsto c_1(L_i), \quad b \mapsto c_2(V_{n-1,n}). \quad (2.6.3.4)$$

If $\pi_1(G) = 1$, then the claim follows immediately.

Suppose $\pi_1(G) = \mu_2$. Then we need to compute the homomorphism

$$\Phi_\nu: \text{CH}_{K,\tilde{T}}^*(\text{Spec } k) \rightarrow \text{CH}_{K,\tilde{T}}^*(\text{Spec } k) \otimes \text{CH}_{K,\mu_2}^*(\text{Spec } k)$$

with $\text{CH}_{K,\tilde{T}}^*(\text{Spec } k)$ given by isomorphism (2.6.3.4) and

$$\text{CH}_{K,\mu_2}^*(\text{Spec } k) \cong \text{CH}_K^*(\text{Spec } k)[x], \quad x \mapsto c_1(\Lambda_{\mu_2}^\pm),$$

by Proposition 2.4.1 (we have ${}^2\mu_2 = \mu_2$), where $\Lambda_{\mu_2}^\pm$ is the nontrivial linear representation of μ_2 . The restriction homomorphism

$$\text{Res}_\nu: \text{Rep}(\tilde{T}) \cong \mathbb{Z}[\mathcal{X}^*(\tilde{T})]^{\text{Gal}(L/k)} \rightarrow \mathbb{Z}[\mathcal{X}^*(\tilde{T}) \oplus \mathcal{X}^*(\mu_2)]^{\text{Gal}(L/k)} \cong \text{Rep}(\tilde{T} \times \mu_2)$$

is induced by the morphism of lattices

$$\begin{aligned} \mathcal{X}^*(\nu): \mathcal{X}^*(\tilde{T}) &= \bigoplus_{i=1}^n \mathbb{Z} \cdot \varpi_i \rightarrow \bigoplus_{i=1}^n \mathbb{Z} \cdot \varpi_i \oplus \mathbb{Z}/2\mathbb{Z} \cdot \bar{\varpi}_{n-1} = \mathcal{X}^*(\tilde{T}) \oplus \mathcal{X}^*(\mu_2), \\ \varpi_i &\mapsto \varpi_i, \quad 1 \leq i \leq n-2, \quad \varpi_{n-1} \mapsto \varpi_{n-1} + \bar{\varpi}_{n-1}, \quad \varpi_n \mapsto \varpi_n + \bar{\varpi}_{n-1}. \end{aligned}$$

Then

$$\begin{aligned} \text{Res}_\nu([L_i]) &= [L_i], \quad 1 \leq i \leq n-2, \\ \text{Res}_\nu([V_{n-1,n}]) &= \text{Res}_\nu(x^{\varpi_{n-1}} + x^{\varpi_n}) = x^{\varpi_{n-1} + \bar{\varpi}_{n-1}} + x^{\varpi_n + \bar{\varpi}_{n-1}} = (x^{\varpi_{n-1}} + x^{\varpi_n}) \cdot x^{\bar{\varpi}_{n-1}} = \\ &= [V_{n-1,n}] \otimes [\Lambda_{\mu_2}^\pm]. \end{aligned}$$

Hence $\Phi_\nu(x_i) = x_i \otimes 1$, $1 \leq i \leq n-2$, and

$$\begin{aligned} \Phi_\nu(b) &= c_2(\text{Res}_\nu([V_{n-1,n}])) = c_2(V_{n-1,n} \otimes \Lambda_{\mu_2}^\pm) = \\ &= c_2(V_{n-1,n}) \otimes 1 + c_1(V_{n-1,n}) \otimes c_1(\Lambda_{\mu_2}^\pm) + 1 \otimes c_1(\Lambda_{\mu_2}^\pm)^2 = b \otimes 1 + 1 \otimes x^2. \end{aligned}$$

The claim follows.

Now suppose $(\Delta(G), \pi_1(G)) = ({}^2D_{2r}, {}^2\mu_{2,2})$. We need to compute the homomorphism

$$\Phi_\nu: \text{CH}_{K, \tilde{T}}^*(\text{Spec } k) \rightarrow \text{CH}_{K, \tilde{T}}^*(\text{Spec } k) \otimes \text{CH}_{K, {}^2\mu_{2,2}}^*(\text{Spec } k)$$

with $\text{CH}_{K, \tilde{T}}^*(\text{Spec } k)$ given by isomorphism (2.6.3.4) and

$$\text{CH}_{K, {}^2\mu_{2,2}}^*(\text{Spec } k) \cong \text{CH}_K^*(\text{Spec } k)[d], \quad d \mapsto c_2({}^2V_{\mu_{2,2}}),$$

by Proposition 2.4.1. The restriction homomorphism

$$\text{Res}_\nu: \text{Rep}(\tilde{T}) \cong \mathbb{Z}[\mathcal{X}^*(\tilde{T})]^{\text{Gal}(L/k)} \rightarrow \mathbb{Z}[\mathcal{X}^*(\tilde{T}) \oplus \mathcal{X}^*({}^2\mu_{2,2})]^{\text{Gal}(L/k)} \cong \text{Rep}(\tilde{T} \times {}^2\mu_{2,2})$$

is induced by the homomorphism of lattices

$$\begin{aligned} \mathcal{X}^*(\nu): \mathcal{X}^*(\tilde{T}) &= \bigoplus_{i=1}^{2r} \mathbb{Z} \cdot \varpi_i \rightarrow \bigoplus_{i=1}^{2r} \mathbb{Z} \cdot \varpi_i \oplus \mathbb{Z}/2\mathbb{Z} \cdot \bar{\varpi}_{2r-1} \oplus \mathbb{Z}/2\mathbb{Z} \cdot \bar{\varpi}_{2r} = \mathcal{X}^*(\tilde{T}) \oplus \mathcal{X}^*({}^2\mu_{2,2}), \\ \varpi_i &\mapsto \varpi_i + i \cdot (\bar{\varpi}_{2r-1} + \bar{\varpi}_{2r}), \quad 1 \leq i \leq 2r-2, \quad \varpi_{2r-1} \mapsto \varpi_{2r-1} + \bar{\varpi}_{2r-1}, \quad \varpi_{2r} \mapsto \varpi_{2r} + \bar{\varpi}_{2r}, \end{aligned}$$

Then

$$\text{Res}_\nu([L_{2i}]) = [L_{2i}], \quad \text{Res}_\nu([L_{2i+1}]) = [L_{2i+1}] \otimes [\det {}^2V_{\mu_{2,2}}], \quad 1 \leq 2i \leq 2r-3.$$

Hence

$$\Phi_\nu(x_i) = x_i \otimes 1, \quad 1 \leq i \leq 2r-2,$$

since $c_1(\det {}^2V_{\mu_{2,2}}) = 0$ because $\text{CH}_{K, {}^2\mu_{2,2}}^1(\text{Spec } k) = 0$. For $V_{2r-1, 2r}$ we have

$$\text{Res}_\nu([V_{2r-1, 2r}]) = \text{Res}_\nu(x^{\varpi_{2r-1}} + x^{\varpi_{2r}}) = x^{\varpi_{2r-1} + \bar{\varpi}_{2r-1}} + x^{\varpi_{2r} + \bar{\varpi}_{2r}}.$$

This element corresponds to a rank 2 representation W of $R \times {}^2\mu_{2,2}$, where $R := R_{L/k} \mathbb{G}_m$. Restricting the representation W to R and ${}^2\mu_{2,2}$ respectively we obtain $W|_R \cong V_R$ and $W|_{{}^2\mu_{2,2}} \cong {}^2V_{\mu_{2,2}}$. Then Lemma 2.4.2 yields

$$c_2(W) = c_2(V_R) \otimes 1 + 1 \otimes c_2({}^2V_{\mu_{2,2}}) \in \text{CH}_{K, R}^*(\text{Spec } k) \otimes \text{CH}_{K, {}^2\mu_{2,2}}^*(\text{Spec } k) \cong \text{CH}_{K, R \times {}^2\mu_{2,2}}^*(\text{Spec } k).$$

Hence we have

$$\Phi_\nu(b) = c_2(\text{Res}_\nu([V_{2r-1, 2r}])) = c_2(V_{2r-1, 2r}) \otimes 1 + 1 \otimes c_2({}^2V_{\mu_{2,2}}) = b \otimes 1 + 1 \otimes d.$$

The claim follows.

Now suppose $(\Delta(G), \pi_1(G)) = ({}^2D_{2r+1}, {}^2\mu_4)$. We need to compute the homomorphism

$$\Phi_\nu: \text{CH}_{K, \tilde{T}}^*(\text{Spec } k) \rightarrow \text{CH}_{K, \tilde{T}}^*(\text{Spec } k) \otimes \text{CH}_{K, {}^2\mu_4}^*(\text{Spec } k)$$

with $\text{CH}_{K, \tilde{T}}^*(\text{Spec } k)$ given by isomorphism (2.6.3.4) and

$$\text{CH}_{K, {}^2\mu_4}^*(\text{Spec } k) \cong \text{CH}_K^*(\text{Spec } k)[x, d]/(x^2), \quad x \mapsto c_1({}^2\Lambda_{\mu_4}^\pm), \quad d \mapsto c_2({}^2V_{\mu_4})$$

by Proposition 2.4.1. The restriction homomorphism

$$\text{Res}_\nu: \text{Rep}(\tilde{T}) \cong \mathbb{Z}[\mathcal{X}^*(\tilde{T})]^{\text{Gal}(L/k)} \rightarrow \mathbb{Z}[\mathcal{X}^*(\tilde{T}) \oplus \mathcal{X}^*(2\mu_4)]^{\text{Gal}(L/k)} \cong \text{Rep}(\tilde{T} \times 2\mu_4)$$

is induced by the morphism of lattices

$$\begin{aligned} \mathcal{X}^*(\nu): \mathcal{X}^*(\tilde{T}) &= \bigoplus_{i=1}^{2r+1} \mathbb{Z} \cdot \varpi_i \rightarrow \bigoplus_{i=1}^{2r+1} \mathbb{Z} \cdot \varpi_i \oplus \mathbb{Z}/4\mathbb{Z} \cdot \bar{\varpi}_{2r} = \mathcal{X}^*(\tilde{T}) \oplus \mathcal{X}^*(2\mu_4), \\ \varpi_i &\mapsto \varpi_i + 2i\bar{\varpi}_{2r}, \quad 1 \leq i \leq 2r-1, \quad \varpi_{2r} \mapsto \varpi_{2r} - \bar{\varpi}_{2r}, \quad \varpi_{2r+1} \mapsto \varpi_{2r+1} + \bar{\varpi}_{2r}. \end{aligned}$$

Then

$$\text{Res}_\nu([L_{2i}]) = [L_{2i}], \quad \text{Res}_\nu([L_{2i+1}]) = [L_{2i+1}] \otimes [\Lambda_{\mu_4}^\pm], \quad 1 \leq 2i \leq 2r-2,$$

and it follows that

$$\Phi_\nu(x_{2i}) = x_{2i} \otimes 1, \quad \Phi_\nu(x_{2i+1}) = x_{2i+1} \otimes 1 + 1 \otimes x, \quad 1 \leq 2i \leq 2r-2.$$

For $V_{2r,2r+1}$ we have

$$\text{Res}_\nu([V_{2r,2r+1}]) = \text{Res}_\nu(x^{\varpi_{2r}} + x^{\varpi_{2r+1}}) = x^{\varpi_{2r}+3\bar{\varpi}_{2r}} + x^{\varpi_{2r+1}+\bar{\varpi}_{2r}}.$$

This element corresponds to a rank 2 representation W of $R \times 2\mu_4$, where $R := R_{L/k}\mathbb{G}_m$. Restricting the representation W to R and $2\mu_4$ respectively we obtain $W|_R \cong V_R$ and $W|_{(2\mu_4)} \cong {}^2V_{\mu_4}$. Then Lemma 2.4.2 yields

$$c_2(W) = c_2(V_R) \otimes 1 + 1 \otimes c_2({}^2V_{\mu_4}) \in \text{CH}_{K,R}^*(\text{Spec } k) \otimes \text{CH}_{K,2\mu_4}^*(\text{Spec } k) \cong \text{CH}_{K,R \times 2\mu_4}^*(\text{Spec } k).$$

Hence we have

$$\Phi_\nu(b) = c_2(\text{Res}_\nu([V_{2r,2r+1}])) = c_2(V_{2r,2r+1}) \otimes 1 + 1 \otimes c_2({}^2V_{\mu_4}) = b \otimes 1 + 1 \otimes d$$

and the claim follows.

³D₄ and **²E₆**. The order of $\pi_1(G)$ is coprime to $[L:k]$ in both cases, and Lemmas 2.1.2 and 2.2.7 yield $\text{CH}_{K,\pi_1(G)}^*(\text{Spec } k) \cong \text{CH}_K^*(\text{Spec } k)$. Thus

$$\text{CH}_K^*(G) \cong \text{CH}_K^*(\tilde{G}) \cong \mathcal{R}^*/(\tilde{c}(\text{CH}_{K,\tilde{T}}^{>0}(\text{Spec } k))).$$

We have

$$\text{CH}_{K,\tilde{T}}^*(\text{Spec } k) \cong \text{CH}_K^*(\text{Spec } k)[x, w], \quad x \mapsto c_1(L_2), \quad w \mapsto c_3(V_{1,3,4}),$$

in the case of ³D₄ while in the case of ²E₆

$$\begin{aligned} \text{CH}_{K,\tilde{T}}^*(\text{Spec } k) &\cong \text{CH}_K^*(\text{Spec } k)[x_1, x_2, b_1, b_2], \\ x_1 &\mapsto c_1(L_2), \quad x_2 \mapsto c_1(L_4), \quad b_1 \mapsto c_2(V_{1,6}), \quad b_2 \mapsto c_2(V_{3,5}). \end{aligned}$$

The claim follows. \square

Remark 2.6.4. One can extend Theorem 2.6.3 also to the split case in a straightforward way, obtaining analogous formulae. We do not do this because in the split case the characteristic sequence is exact and supplies us with all the necessary information.

Remark 2.6.5. Let G be a simply connected quasi-split simple group over a field k of type $\Delta(G) = {}^2A_{2r-1}$ and let K/k be the splitting field of G . Let $m_1 \mid m_2 \mid r$ be integers with $\frac{r}{m_2}$ and $\frac{m_2}{m_1}$ being even. Consider the pullback homomorphism

$$f^*: \text{CH}_K^*(G/{}^2\mu_{2m_2}) \rightarrow \text{CH}_K^*(G/{}^2\mu_{2m_1})$$

for the quotient morphism $f: G/{}^2\mu_{2m_1} \rightarrow G/{}^2\mu_{2m_2}$. Then although these rings may be isomorphic (i.e. if m_1 is even), f^* fails to be an isomorphism, since $f^*(x) = 0$ in the notation of Theorem 2.6.3 by Lemma A.4.3.

2.7 Conormed Chow ring of a quasi-split group G via $\mathcal{C}_K^*(G)$

In this section we show how to recover $\mathrm{CH}^*(G)$ out of the cokernel $\mathcal{C}_K^*(G)$ of the conormed characteristic map. If G is either simply connected or adjoint, then the conormed characteristic sequence is exact and $\mathrm{CH}^*(G) \cong \mathcal{C}_K^*(G)$, otherwise one has to add one generator to cover the conormed Picard group $\mathrm{CH}_K^1(G)$ and also one relation on the square of this element, with the answer being $\mathrm{CH}^*(G) \cong \mathcal{C}_K^*(G)[x]/(x^2 + a)$ for a certain element $a \in \mathcal{C}_K^*(G)$. We continue to use the standard notation for quasi-split groups and vector bundles recalled in Sections A.2 and A.3.

Definition 2.7.1. In the notation of Definition 2.5.5 one has $\phi^* \circ c = 0$, e.g. by the construction of the Eilenberg–Moore homomorphism (2.2.4.2). We put

$$\mathcal{C}_K^*(G) := \mathrm{CH}_K^*(G/B) / (c(\mathrm{CH}_{K,T}^{\geq 0}(\mathrm{Spec} k)) \cdot \mathrm{CH}_K^*(G/B))$$

and denote by

$$\Theta_G: \mathcal{C}_K^*(G) \rightarrow \mathrm{CH}_K^*(G)$$

the induced homomorphism.

Theorem 2.7.2. *Let G be a quasi-split group over a field k , let $T \leq B \leq G$ be a maximal torus and a Borel subgroup, and let K/k be a separable extension of fields such that T_K is split. Then the following holds.*

1. *If T is quasi-trivial, then*

$$\Theta_G: \mathcal{C}_K^*(G) \rightarrow \mathrm{CH}_K^*(G)$$

is an isomorphism. In particular, this applies to a split G and, by Lemma A.2.3, to a simply connected or an adjoint quasi-split semisimple group G .

2. *If $(\Delta(G), \pi_1(G)) = ({}^2A_n, {}^2\mu_l)$ with l or $\frac{n+1}{l}$ being odd, then*

$$\Theta_G: \mathcal{C}_K^*(G) \rightarrow \mathrm{CH}_K^*(G)$$

is an isomorphism.

3. *If $(\Delta(G), \pi_1(G)) = ({}^2A_{2r-1}, {}^2\mu_{2m})$ with $\frac{r}{m}$ being even, then Θ_G induces an isomorphism*

$$\mathcal{C}_K^*(G)[x]/(x^2 + m^2 \cdot \bar{c}_2(\mathcal{V}_{1,2r-1})) \xrightarrow{\cong} \mathrm{CH}_K^*(G), \quad x \mapsto c_1(\mathcal{L}(\bar{\omega}_m)),$$

where $\bar{c}_2(\mathcal{V}_{1,2r-1}) \in \mathcal{C}_K^(G)$ is the image of $c_2(\mathcal{V}_{1,2r-1}) \in \mathrm{CH}_K^*(G/B)$ under the quotient homomorphism.*

4. *If $(\Delta(G), \pi_1(G)) = ({}^2D_n, \mu_2)$, then Θ_G induces an isomorphism*

$$\mathcal{C}_K^*(G)[x]/(x^2 + \bar{c}_2(\mathcal{V}_{n-1,n})) \xrightarrow{\cong} \mathrm{CH}_K^*(G), \quad x \mapsto c_1(\mathcal{L}(\bar{\omega}_{n-1})),$$

where $\bar{c}_2(\mathcal{V}_{n-1,n}) \in \mathcal{C}_K^(G)$ is the image of $c_2(\mathcal{V}_{n-1,n}) \in \mathrm{CH}_K^*(G/B)$ under the quotient homomorphism.*

Proof. (1) follows from Theorem 2.5.6.(1).

In cases (2)–(4) the considered homomorphisms are well-defined and surjective by the formulae obtained in Theorem 2.6.3, and it is sufficient to show that the generators of the ideals given in Theorem 2.6.3 that do not contain terms with x^2 , belong to the ideal $I(c)$ generated by the image of the characteristic map

$$c: \mathrm{CH}_{K,T}^{\geq 0}(\mathrm{Spec} k) \rightarrow \mathrm{CH}_K^*(G/B).$$

Let $f: \tilde{G} \rightarrow G$ be the simply connected cover and $g: G \rightarrow \bar{G}$ be the adjoint quotient, put

$$\tilde{T} := f^{-1}(T), \quad \tilde{B} := f^{-1}(B), \quad \bar{T} := g(T), \quad \bar{B} := g(B).$$

The following diagram commutes.

$$\begin{array}{ccccc}
\mathrm{CH}_{K,\bar{T}}^*(\mathrm{Spec} k) & \xrightarrow{\Phi_g} & \mathrm{CH}_{K,T}^*(\mathrm{Spec} k) & \xrightarrow{\Phi_f} & \mathrm{CH}_{K,\tilde{T}}^*(\mathrm{Spec} k) \\
\bar{c} \downarrow & & \downarrow c & & \downarrow \tilde{c} \\
\mathrm{CH}_K^*(\tilde{G}/\tilde{B}) & \xrightarrow{=} & \mathrm{CH}_K^*(G/B) & \xrightarrow{=} & \mathrm{CH}_K^*(\tilde{G}/\tilde{B})
\end{array} \tag{2.7.2.1}$$

Here \bar{c} , c and \tilde{c} are the characteristic maps, Φ_g and Φ_f are induced by the homomorphisms $g: T \rightarrow \bar{T}$ and $f: \tilde{T} \rightarrow T$ respectively, and the bottom maps are induced by the isomorphism $\tilde{G}/\tilde{B} \xrightarrow{\cong} G/B \xrightarrow{\cong} \tilde{G}/\tilde{B}$. Below we use the same notation as in Theorem 2.6.3.

(2), $n = 2r$. We need to show that $c_2(\mathcal{V}_{i,2r+1-i}) \in I(c)$ for $1 \leq i \leq r$. It follows from the formulae obtained in Theorem 2.6.3 in the adjoint case (i.e. $l = 2r$) and by the exactness of the characteristic sequence in the adjoint case proved in Theorem 2.5.6 that $c_2(\mathcal{V}_{i,2r+1-i}) \in I(\bar{c})$, $1 \leq i \leq r$. We have $I(\bar{c}) \subseteq I(c)$ by the commutativity of Diagram 2.7.2.1. The claim follows.

(2), $n = 2r - 1$, $l = 2m$, $\frac{r}{m}$ is odd. The claim follows from the adjoint case as above.

(2), $n = 2r - 1$, $l = 2m + 1$. We need to show that $c_1(\mathcal{L}_r) \in I(c)$ and $c_2(\mathcal{V}_{i,2r-i}) \in I(c)$ for $1 \leq i \leq r - 1$. Using the adjoint case as above we obtain $c_2(\mathcal{V}_{2i,2r-2i}) \in I(c)$ for $1 \leq i \leq \lfloor \frac{r-1}{2} \rfloor$ and $c_2(\mathcal{V}_{2i+1,2r-2i-1}) + c_2(\mathcal{V}_{1,2r-1}) \in I(c)$ for $1 \leq i \leq \lfloor \frac{r}{2} \rfloor - 1$, thus it suffices to show $c_1(\mathcal{L}_r), c_2(\mathcal{V}_{2m+1,2r-2m-1}) \in I(c)$. Since $2m+1 \mid 2r$, it follows that $2m+1 \mid r$ and $\varpi_{2m+1}, \varpi_r, \varpi_{2r-2m-1} \in \mathcal{X}^*(T)$, thus $c_1(\mathcal{L}_r), c_2(\mathcal{V}_{2m+1,2r-2m-1}) \in \mathrm{CH}_{K,\bar{T}}^*(\mathrm{Spec} k)$ belong to the image of Φ_f . Then the claim follows from the commutativity of diagram (2.7.2.1), since $c_1(\mathcal{L}_r) = \tilde{c}(c_1(\mathcal{L}_r))$ and $c_2(\mathcal{V}_{2m+1,2r-2m-1}) = \tilde{c}(c_2(\mathcal{V}_{2m+1,2r-2m-1}))$.

(3) We need to show that $c_1(\mathcal{L}_r) \in I(c)$ and $\mathcal{S}_{2r-1} \subseteq I(c)$. It follows from the adjoint case as above that $\mathcal{S}_{2r-1} \subseteq I(c)$. Since $\frac{r}{m}$ is even, it follows that $2m \mid r$ and $\varpi_r \in \mathcal{X}^*(T)$, thus $c_1(\mathcal{L}_r)$ belongs to the image of Φ_f . The claim follows as above.

(4) We need to show that $c_1(\mathcal{L}_i) \in I(c)$, $1 \leq i \leq 2r - 2$. We have $\varpi_i \in \mathcal{X}^*(T)$, $1 \leq i \leq n - 2$, thus $c_1(\mathcal{L}_i)$ belongs to the image of Φ_f . The claim follows as above. \square

2.8 Conormed Chow ring of a quasi-split group: the answer

In this section we combine the previous computations and results from [GZ22] to compute the conormed Chow ring $\mathrm{CH}_K^*(G)$ of a non-split quasi-split simple group assuming K to be the splitting field. If the type of G is $\Delta(G) = {}^6\mathrm{D}_4$, then $\mathrm{CH}_K^*(G) = 0$, while in the other types it follows from Proposition 2.5.3 and a Borel's result on the structure of Hopf algebras [MM65, Theorem 7.11 and Proposition 7.8] that

$$\mathrm{CH}_K^*(G) \cong \mathbb{F}_p[e_1, e_2, \dots, e_s] / (e_1^{p^{k_1}}, e_2^{p^{k_2}}, \dots, e_s^{p^{k_s}}), \quad \deg e_i = d_i, \quad p = [K : k],$$

for some parameters s, d_i, k_i , and we give a table for them depending on $(\Delta(G), \pi_1(G))$.

We continue to use the notation from Sections A.1, A.2 and A.3.

Lemma 2.8.1. *Let G be a quasi-split simple group over a field k of type $\Delta(G) = {}^2\mathrm{D}_n$, let $B \leq G$ be a Borel subgroup and let K/k be the splitting field of G . Then in $\mathrm{CH}_K^*(G/B)$ we have the following:*

- $\mathrm{CH}_K^1(G/B) = \bigoplus_{i=1}^{n-2} \mathbb{F}_2 \cdot c_1(\mathcal{L}_i)$,
- $\mathrm{CH}_K^2(G/B) = \mathrm{CH}_K^1(G/B) \cdot \mathrm{CH}_K^1(G/B) + \mathbb{F}_2 \cdot c_2(\mathcal{V}_{n-1,n})$.

Proof. We have $[K : k] = 2$, thus $\mathrm{CH}_K^*(\mathrm{Spec} k) \cong \mathbb{F}_2$ by Lemma 2.1.2. Lemma 2.6.1 yields that

$$\mathrm{CH}_K^*(G/B) \cong (\mathrm{CH}^*((G/B)_K))^{\mathrm{Gal}(K/k)} / (\mathrm{id} + \tau)(\mathrm{CH}^*((G/B)_K))$$

with $\tau \in \mathrm{Gal}(K/k)$ being the non-trivial element. Since G_K is split, $\mathrm{CH}^*((G/B)_K)$ has a basis given by the Schubert cycles $Z_w \in \mathrm{CH}^{l(w)}((G/B)_K)$, $w \in W(G_K)$, [Che94, Proposition 8], here $l(w)$ is the length of w . Put s_i to be the reflection for the simple root α_i . The involution τ acts on the Weyl group $W(G_K)$ via the action on the generators $\tau(s_i) = s_i$, $1 \leq i \leq n - 2$, $\tau(s_{n-1}) = s_n$, $\tau(s_n) = s_{n-1}$, which yields a

permutation action on the Schubert cycles compatible with the action of τ on $\text{CH}_K^*(G/B)$. In particular, we have the following $\text{Gal}(K/k)$ -invariant decompositions

$$\text{CH}^1((G/B)_K) = \left(\bigoplus_{i=1}^{n-2} \mathbb{Z} \cdot Z_{s_i} \right) \oplus [\mathbb{Z} \cdot Z_{s_{n-1}} \oplus \mathbb{Z} \cdot Z_{s_n}],$$

$$\begin{aligned} \text{CH}^2((G/B)_K) = & \left(\bigoplus_{1 \leq i < j \leq n-2} \mathbb{Z} \cdot Z_{s_i s_j} \right) \oplus \left(\bigoplus_{i=1}^{n-3} \mathbb{Z} \cdot Z_{s_{i+1} s_i} \right) \oplus \left(\bigoplus_{i=1}^{n-2} [\mathbb{Z} \cdot Z_{s_i s_{n-1}} \oplus \mathbb{Z} \cdot Z_{s_i s_n}] \right) \oplus \\ & \oplus [\mathbb{Z} \cdot Z_{s_{n-1} s_{n-2}} \oplus \mathbb{Z} \cdot Z_{s_n s_{n-2}}] \oplus \mathbb{Z} \cdot Z_{s_{n-1} s_n}. \end{aligned}$$

Here τ acts on the modules in the square brackets permuting the elements of the bases, and τ acts trivially on all the other summands. Then

$$\begin{aligned} \text{CH}_K^1(G/B) &= \bigoplus_{i=1}^{n-2} \mathbb{F}_2 \cdot \bar{Z}_{s_i}, \\ \text{CH}_K^2(G/B) &= \left(\bigoplus_{1 \leq i < j \leq n-2} \mathbb{F}_2 \cdot \bar{Z}_{s_i s_j} \right) \oplus \left(\bigoplus_{i=1}^{n-3} \mathbb{F}_2 \cdot \bar{Z}_{s_{i+1} s_i} \right) \oplus \mathbb{F}_2 \cdot \bar{Z}_{s_{n-1} s_n}. \end{aligned}$$

In $\text{CH}^*((G/B)_K)$ we have $Z_{s_i} = c_1(\mathcal{L}_i)$, so the first claim of the lemma follows. For the second claim note that the Chevalley's formula [Che94, Proposition 10] yields

$$Z_{s_i} \cdot Z_{s_j} = \begin{cases} Z_{s_i s_j}, & i \neq j \text{ and } \alpha_i \text{ is not connected to } \alpha_j \text{ in the Dynkin diagram } D_n \\ Z_{s_i s_j} + Z_{s_j s_i}, & i \neq j \text{ and } \alpha_i \text{ is connected to } \alpha_j \text{ in the Dynkin diagram } D_n \\ \sum_{(\alpha_l, \alpha_i) \neq 0} Z_{s_l s_i}, & i = j, \text{ the sum is taken over all } l \text{ such that } \alpha_l \text{ is connected to } \alpha_i. \end{cases}$$

In particular, we have

$$Z_{s_{n-1} s_n} = Z_{s_{n-1}} \cdot Z_{s_n} = c_1(\mathcal{L}_{n-1}) \cdot c_1(\mathcal{L}_n) = c_2(\mathcal{V}_{n-1, n})$$

in $\text{CH}^*((G/B)_K)$. Hence $c_2(\mathcal{V}_{n-1, n})$ coincides with $\bar{Z}_{s_{n-1} s_n}$ in $\text{CH}_K^*(G/B)$. The claim follows. \square

Theorem 2.8.2. *Let G be a non-split quasi-split simple group over a field k and let K/k be the splitting field of G . Suppose $\Delta(G) \neq {}^6D_4$ and put $p := [K : k]$. Then*

$$\text{CH}_K^*(G) \cong \mathbb{F}_p[e_1, e_2, \dots, e_s] / (e_1^{p^{k_1}}, e_2^{p^{k_2}}, \dots, e_s^{p^{k_s}}), \quad \deg e_i = d_i,$$

with the parameters s, d_i, k_i given by the following table.

Table 2: $\text{CH}_K^*(G)$ for a non-split quasi-split simple group G

$\Delta(G)$	$\pi_1(G)$	s	$d_i, i = 1, 2, \dots, s$	$k_i, i = 1, 2, \dots, s$
${}^2A_n, n \geq 2$	$2\mu_l, l \mid (n+1), l \text{ is odd}$	$\lfloor \frac{n}{2} \rfloor$	$2i+1$	1
${}^2A_{2r-1}, r \geq 2$	$2\mu_{2m}, m \mid r, m \text{ is odd}$	r	$2i-1$	$v_2(r)+1, i=1$ $1, i \geq 2$
${}^2A_{2r-1}, r \geq 2$	$2\mu_{2m}, m \mid r, m \text{ is even}$	$r+1$	$2, i=1$ $2i-3, i \geq 2$	$v_2(r), i=1$ $1, i \geq 2$

${}^2D_n, n \geq 3$	1	$\lfloor \frac{n+1}{2} \rfloor - 1$	$2i + 1$	$\lfloor \log_2 \frac{2n}{2i+1} \rfloor$
	μ_2	$\lfloor \frac{n+1}{2} \rfloor$	$1, i = 1$ $2i - 1, i \geq 2$	$\lfloor \log_2 n \rfloor + 1$ $\lfloor \log_2 \frac{2n}{2i-1} \rfloor$
${}^2D_{2r+1}, r \geq 1$	${}^2\mu_4$	$r + 2$	$1, i = 1$	$1, i = 1$
			$2, i = 2$	$\lfloor \log_2(2r + 1) \rfloor, i = 2$
			$2i - 3, i \geq 3$	$\lfloor \log_2 \frac{4r+2}{2i-3} \rfloor, i \geq 3$
${}^2D_{2r}, r \geq 2$	${}^2\mu_{2,2}$	r	$2, i = 1$	$\lfloor \log_2 2r \rfloor, i = 1$
			$2i - 1, i \geq 2$	$\lfloor \log_2 \frac{4r}{2i-1} \rfloor, i \geq 2$
3D_4	$1, {}^3\mu_{2,2}$	1	4	1
2E_6	$1, {}^2\mu_3$	3	3, 5, 9	1, 1, 1

Here v_2 is the 2-adic valuation.

Proof. Let $\tilde{G} \rightarrow G$ and $G \rightarrow \bar{G}$ be the simply connected cover and the adjoint quotient respectively. We have $\text{CH}_K^*(\bar{G}) \cong \mathcal{C}_K^*(\bar{G})$ by Theorem 2.7.2 and the last ring was determined in [GZ22, Section 7] (see especially Table 5 of loc. cit.). Furthermore, it follows from Theorem 2.6.3 that $\text{CH}_K^*(G) \cong \text{CH}_K^*(\bar{G})$, if $(\Delta(G), \pi_1(G)) = ({}^2A_n, {}^2\mu_l)$ with $\frac{n+1}{l}$ being odd, or $({}^3D_4, {}^3\mu_{2,2})$, or $({}^2E_6, {}^2\mu_3)$. Then the remaining cases are $({}^2A_{2r-1}, {}^2\mu_l)$ with $\frac{2r}{l}$ being even, $({}^2D_n, 1)$ and $({}^2D_n, \mu_2)$. In each case we use Theorem 2.6.3 to derive the answers from the ones for the respective adjoint groups. Below we adopt the notation of Theorem 2.6.3 and repeatedly use its claims without additional references.

(1) $(\Delta(G), \pi_1(G)) = ({}^2A_{2r-1}, {}^2\mu_l)$ with $\frac{2r}{l}$ being even. Suppose that r is odd. We have

$$\mathcal{R}^*/I(\{c_2(\mathcal{V}_{i,2r-i})\}_{i=1}^{r-1}) \cong \text{CH}_K^*(\bar{G}) \cong \mathbb{F}_2[e_2, \dots, e_{r+1}]/(e_2^2, \dots, e_{r+1}^2) \quad (2.8.2.1)$$

with $\deg e_i = 2i - 3, i \geq 2$. The element $c_1(\mathcal{L}_r) \in \mathcal{R}^*$ is the generator of \mathcal{R}^1 by Lemma 2.6.2, thus its image in the quotient ring on the left-hand side of the formula corresponds under isomorphisms (2.8.2.1) to e_2 on the right-hand side, which is a unique non-trivial element of degree 1. Since $\frac{2r}{l}$ is even, it follows that l is odd and

$$\begin{aligned} \text{CH}_K^*(G) \cong (\mathcal{R}^*/I(\{c_2(\mathcal{V}_{i,2r-i})\}_{i=1}^{r-1}))/(\bar{c}_1(\mathcal{L}_r)) &\cong (\mathbb{F}_2[e_2, \dots, e_{r+1}]/(e_2^2, \dots, e_{r+1}^2))/(e_2) \cong \\ &\cong \mathbb{F}_2[e_3, \dots, e_{r+1}]/(e_3^2, \dots, e_{r+1}^2). \end{aligned}$$

Suppose r is even, then

$$\mathcal{R}^*/I(\mathcal{S}_{2r-1}) \cong \text{CH}_K^*(\bar{G}) \cong \mathbb{F}_2[e_1, e_2, \dots, e_{r+1}]/(e_1^{2^{v_2(r)}}, e_2^2, \dots, e_{r+1}^2) \quad (2.8.2.2)$$

with $\deg e_1 = 2, \deg e_i = 2i - 3, i \geq 2$. As above, the image of the element $c_1(\mathcal{L}_r) \in \mathcal{R}^*$ in the quotient ring on the left-hand side of the formula corresponds under isomorphisms (2.8.2.2) to e_2 on the right-hand side. Furthermore, Lemma 2.6.2 yields that the image of $c_2(\mathcal{V}_{1,2r-1})$ in the quotient ring on the left-hand side of the formula is the generator of the degree 2 part, thus it is mapped to e_1 under isomorphisms (2.8.2.2). If l is odd, then

$$\begin{aligned} \text{CH}_K^*(G) \cong (\mathcal{R}^*/I(\mathcal{S}_{2r-1}))/(\bar{c}_1(\mathcal{L}_r), \bar{c}_2(\mathcal{V}_{1,2r-1})) &\cong \\ &\cong (\mathbb{F}_2[e_1, e_2, \dots, e_{r+1}]/(e_1^{2^{v_2(r)}}, e_2^2, \dots, e_{r+1}^2))/(e_1, e_2) \cong \mathbb{F}_2[e_3, \dots, e_{r+1}]/(e_3^2, \dots, e_{r+1}^2). \end{aligned}$$

If $l = 2m$ and m is even, then

$$\begin{aligned} \text{CH}_K^*(G) \cong (\mathcal{R}^*/I(\mathcal{S}_{2r-1}))[x]/(\bar{c}_1(\mathcal{L}_r), x^2) &\cong \\ &\cong (\mathbb{F}_2[e_1, \dots, e_{r+1}]/(e_1^{2^{v_2(r)}}, e_2^2, \dots, e_{r+1}^2))[x]/(e_2, x^2) \cong \mathbb{F}_2[e_1, \dots, e_{r+1}]/(e_1^{2^{v_2(r)}}, e_2^2, \dots, e_{r+1}^2). \end{aligned}$$

If $l = 2m$ and m is odd, then

$$\begin{aligned} \mathrm{CH}_K^*(G) &\cong (\mathcal{R}^*/I(\mathcal{S}_{2r-1}))[x]/(\bar{c}_1(\mathcal{L}_r), x^2 + \bar{c}_2(\mathcal{V}_{1,2r-1})) \cong \\ &\cong (\mathbb{F}_2[e_1, \dots, e_{r+1}]/(e_1^{2^{v_2(r)}}, e_2^2, \dots, e_{r+1}^2))[x]/(e_2, x^2 + e_1) \cong \\ &\cong \mathbb{F}_2[x, e_3, \dots, e_{r+1}]/(x^{2^{v_2(r)+1}}, e_3^2, \dots, e_{r+1}^2), \end{aligned}$$

where $\deg x = 1$, $\deg e_i = 2i - 3$.

(2) $\Delta(G) = {}^2D_n$. Suppose that $n = 2r + 1$ and put

$$I := I(\{c_1(\mathcal{L}_1)^2\} \cup \{c_1(\mathcal{L}_{2i})\}_{i=1}^{r-1} \cup \{c_1(\mathcal{L}_{2i+1}) + c_1(\mathcal{L}_1)\}_{i=1}^{r-1}) \subseteq \mathcal{R}^*.$$

We have

$$\mathcal{R}^*/I \cong \mathrm{CH}_K^*(\bar{G}) \cong \mathbb{F}_2[e_1, \dots, e_{r+2}]/(e_1^2, e_2^{2^{k_2}}, \dots, e_{r+2}^{2^{k_{r+2}}}) \quad (2.8.2.3)$$

with $\deg e_1 = 1$, $\deg e_2 = 2$, $\deg e_i = 2i - 3$, $i \geq 3$, and k_i as in the statement of the theorem for the adjoint case. The image of the element $c_1(\mathcal{L}_1) \in \mathcal{R}^*$ in the quotient ring on the left-hand side of the formula is the generator in degree 1 by Lemma 2.8.1, thus it corresponds under isomorphisms (2.8.2.3) to e_1 on the right-hand side, which is a unique non-trivial element of degree 1. Then, since $e_1^2 = 0$ on the right-hand side, it follows that $\bar{c}_1(\mathcal{L}_1)^2 = 0$ on the left-hand side. Thus it follows from Lemma 2.8.1 and the above that the image of the element $c_2(\mathcal{V}_{2r,2r+1}) \in \mathcal{R}^*$ in the quotient ring on the left-hand side of the formula is the generator in degree 2, hence it corresponds under isomorphisms (2.8.2.3) to e_2 on the right-hand side, which is a unique non-trivial element of degree 2. Then we have

$$\begin{aligned} \mathrm{CH}^*(\tilde{G}/\mu_2) &\cong (\mathcal{R}^*/I)[x]/(c_1(\mathcal{L}_1), x^2 + c_2(\mathcal{V}_{2r,2r+1})) \cong \\ &\cong (\mathbb{F}_2[e_1, \dots, e_{r+2}]/(e_1^2, e_2^{2^{k_2}}, \dots, e_{r+2}^{2^{k_{r+2}}})) [x]/(e_1, x^2 + e_2) \cong \\ &\cong \mathbb{F}_2[x, e_3, \dots, e_{r+2}]/(x^{2^{k_2+1}}, e_3^{2^{k_3}}, \dots, e_{r+2}^{2^{k_{r+2}}}) \end{aligned}$$

with $\deg x = 1$. Similarly,

$$\begin{aligned} \mathrm{CH}^*(\tilde{G}) &\cong (\mathcal{R}^*/I)/(c_1(\mathcal{L}_1), c_2(\mathcal{V}_{2r,2r+1})) \cong \\ &\cong (\mathbb{F}_2[e_1, \dots, e_{r+2}]/(e_1^2, e_2^{2^{k_2}}, \dots, e_{r+2}^{2^{k_{r+2}}})) / (e_1, e_2) \cong \mathbb{F}_2[e_3, \dots, e_{r+2}]/(e_3^{2^{k_3}}, \dots, e_{r+2}^{2^{k_{r+2}}}). \end{aligned}$$

Suppose $n = 2r$. We have

$$\mathcal{R}^*/I(\{c_1(\mathcal{L}_i)\}_{i=1}^{2r-2}) \cong \mathrm{CH}_K^*(\bar{G}) \cong \mathbb{F}_2[e_1, \dots, e_r]/(e_1^{2^{k_1}}, \dots, e_r^{2^{k_r}}) \quad (2.8.2.4)$$

with $\deg e_1 = 2$, $\deg e_i = 2i - 1$, $i \geq 2$, and k_i as in the statement of the theorem for the adjoint case. The degree 1 part on the right-hand side of the formula is trivial, thus the same holds on the left-hand side. It follows from Lemma 2.8.1 that the image of the element $c_2(\mathcal{V}_{2r-1,2r}) \in \mathcal{R}^*$ in the quotient ring on the left-hand side of the formula is the generator in degree 2, so it corresponds under isomorphisms (2.8.2.4) to e_1 on the right-hand side, which is a unique non-trivial element of degree 2. Then we have

$$\begin{aligned} \mathrm{CH}^*(\tilde{G}/\mu_2) &\cong (\mathcal{R}^*/I(\{c_1(\mathcal{L}_i)\}_{i=1}^{2r-2}))[x]/(x^2 + c_2(\mathcal{V}_{2r-1,2r})) \cong \\ &\cong (\mathbb{F}_2[e_1, \dots, e_r]/(e_1^{2^{k_1}}, \dots, e_r^{2^{k_r}}))[x]/(x^2 + e_1) \cong \mathbb{F}_2[x, e_2, \dots, e_r]/(x^{2^{k_1+1}}, e_2^{2^{k_2}}, \dots, e_r^{2^{k_r}}) \end{aligned}$$

with $\deg x = 1$. Similarly,

$$\begin{aligned} \mathrm{CH}^*(\tilde{G}) &\cong (\mathcal{R}^*/I(\{c_1(\mathcal{L}_i)\}_{i=1}^{2r-2}))/ (c_2(\mathcal{V}_{2r-1,2r})) \cong \\ &\cong (\mathbb{F}_2[e_1, \dots, e_r]/(e_1^{2^{k_1}}, \dots, e_r^{2^{k_r}}))/ (e_1) \cong \mathbb{F}_2[e_2, e_3, \dots, e_r]/(e_2^{2^{k_2}}, \dots, e_r^{2^{k_r}}). \end{aligned}$$

□

Remark 2.8.3. Let G be a simply connected quasi-split simple group over a field k of type $\Delta(G) = {}^2\mathbf{A}_{2r-1}$ and let K/k be the splitting field of G . Let $m_1 \mid m_2 \mid r$ be integers. Consider the pullback homomorphism

$$f^* : \mathrm{CH}_K^*(G/{}^2\mu_{2m_2}) \rightarrow \mathrm{CH}_K^*(G/{}^2\mu_{2m_1})$$

for the quotient morphism $f : G/{}^2\mu_{2m_1} \rightarrow G/{}^2\mu_{2m_2}$. It follows from the proof of Theorem 2.8.2 that f^* is an isomorphism, if $\frac{m_2}{m_1}$ is odd, while for $\frac{m_2}{m_1}$ even one has $f^*(e_2) = 0$ for the degree 1 generator e_2 by Lemma A.4.3.

2.9 Cokernel in the conormed characteristic sequence

In this section we complement the previous results with a computation of the cokernel $\mathcal{C}_K^*(G)$ in the conormed characteristic sequence. This cokernel coincides with $\mathrm{CH}_K^*(G)$ for the simply connected and adjoint groups, while for intermediate groups it is only a subring of $\mathrm{CH}_K^*(G)$ in general. We continue to use the notation from Sections A.1, A.2 and A.3.

Theorem 2.9.1. *Let G be a non-split quasi-split simple group over a field k and let K/k be the splitting field of G . Suppose $\Delta(G) \neq {}^6\mathbf{D}_4$ and put $p := [K : k]$. Then*

$$\mathcal{C}_K^*(G) \cong \mathbb{F}_p[e_1, e_2, \dots, e_s] / (e_1^{p^{k_1}}, e_2^{p^{k_2}}, \dots, e_s^{p^{k_s}}), \quad \deg e_i = d_i,$$

with the parameters s, d_i, k_i given by the following table.

Table 3: $\mathcal{C}_K^*(G)$ for a non-split quasi-split simple group G

$\Delta(G)$	$\pi_1(G)$	s	$d_i, i = 1, 2, \dots, s$	$k_i, i = 1, 2, \dots, s$
${}^2\mathbf{A}_n, n \geq 2$	${}^2\mu_l, l \mid (n+1), l \text{ is odd}$	$\lfloor \frac{n}{2} \rfloor$	$2i+1$	1
${}^2\mathbf{A}_{2r-1}, r \geq 2 \text{ odd}$	${}^2\mu_{2m}, m \mid r$	r	$2i-1$	1
${}^2\mathbf{A}_{2r-1}, r \geq 2 \text{ even}$	${}^2\mu_{2m}, m \mid r, \frac{r}{m} \text{ is even}$	r	$2, i=1$ $2i-1, i \geq 2$	$v_2(r), i=1$ $1, i \geq 2$
${}^2\mathbf{A}_{2r-1}, r \geq 2 \text{ even}$	${}^2\mu_{2m}, m \mid r, \frac{r}{m} \text{ is odd}$	$r+1$	$2, i=1$ $2i-3, i \geq 2$	$v_2(r), i=1$ $1, i \geq 2$
${}^2\mathbf{D}_n, n \geq 3$	1 μ_2	$\lfloor \frac{n+1}{2} \rfloor - 1$ $\lfloor \frac{n+1}{2} \rfloor$	$2i+1$ $2, i=1$ $2i-1, i \geq 2$	$\lfloor \log_2 \frac{2n}{2i+1} \rfloor$ $\lfloor \log_2 n \rfloor$ $\lfloor \log_2 \frac{2n}{2i-1} \rfloor$
${}^2\mathbf{D}_{2r+1}, r \geq 1$	${}^2\mu_4$	$r+2$	$1, i=1$ $2, i=2$ $2i-3, i \geq 3$	$1, i=1$ $\lfloor \log_2(2r+1) \rfloor, i=2$ $\lfloor \log_2 \frac{4r+2}{2i-3} \rfloor, i \geq 3$
${}^2\mathbf{D}_{2r}, r \geq 2$	${}^2\mu_{2,2}$	r	$2, i=1$ $2i-1, i \geq 2$	$\lfloor \log_2 2r \rfloor, i=1$ $\lfloor \log_2 \frac{4r}{2i-1} \rfloor, i \geq 2$
${}^3\mathbf{D}_4$	$1, {}^3\mu_{2,2}$	1	4	1
${}^2\mathbf{E}_6$	$1, {}^2\mu_3$	3	3, 5, 9	1, 1, 1

Proof. If G is adjoint, then this was computed in [GZ22, Section 8] (see especially Table 5 of loc. cit.). If G is simply connected, i.e. $\pi_1(G) = 1$, or if $(\Delta(G), \pi_1(G)) = ({}^2\mathbf{A}_n, {}^2\mu_l)$ with $l \mid n+1$ and either

l or $\frac{n+1}{l}$ being odd, then the claim follows from Theorems 2.7.2 and 2.8.2. Thus the remaining cases are $(\Delta(G), \pi_1(G)) = ({}^2A_{2r-1}, {}^2\mu_{2m})$ with $m \mid r$ and $\frac{r}{m}$ being even, and $(\Delta(G), \pi_1(G)) = ({}^2D_n, \mu_2)$. Let $g: G \rightarrow \bar{G}$ be the adjoint quotient and consider the following commutative diagram.

$$\begin{array}{ccc} \mathcal{C}_K^*(\bar{G}) & \xrightarrow[\Theta_{\bar{G}}]{\cong} & \text{CH}_K^*(\bar{G}) \\ g_{\bar{C}}^* \downarrow & & \downarrow g^* \\ \mathcal{C}_K^*(G) & \xrightarrow{\Theta_G} & \text{CH}_K^*(G) \end{array} \quad (2.9.1.1)$$

Here the horizontal morphisms are the canonical ones from Definition 2.7.1, g^* is the pullback, and $g_{\bar{C}}^*$ is induced by the isomorphism $\text{CH}_K^*(\bar{G}/\bar{B}) \xrightarrow{\cong} \text{CH}_K^*(G/B)$ given by the isomorphism $G/B \xrightarrow{\cong} \bar{G}/\bar{B}$ with $\bar{B} := g(B)$. The homomorphism $g_{\bar{C}}^*$ is clearly surjective while the homomorphism $\Theta_{\bar{G}}$ is an isomorphism by Theorem 2.7.2. Furthermore, Theorem 2.7.2 yields that Θ_G is injective. It follows that

$$\mathcal{C}_K^* \cong \Theta_G(\mathcal{C}_K^*) = g^*(\text{CH}_K^*(\bar{G})).$$

$({}^2A_{2r-1}, {}^2\mu_{2m})$, $m \mid r$, $\frac{r}{m}$ is odd. Theorem 2.6.3 yields that g^* is an isomorphism, thus $g_{\bar{C}}^*$ is an isomorphism as well.

$({}^2A_{2r-1}, {}^2\mu_{2m})$, $m \mid r$, $\frac{r}{m}$ is even. There is a unique nontrivial element $e_2 = c_1(\mathcal{L}(\bar{\omega}_r)) \in \text{CH}_K^1(\bar{G}) \cong \mathbb{F}_2$, and we have $g^*(e_2) = 0$ by Lemma A.4.3. In this case r is even and Theorem 2.6.3 yields that g^* induces a monomorphism $\text{CH}_K^*(\bar{G})/(e_2) \rightarrow \text{CH}_K^*(G)$. Thus $\mathcal{C}_K^*(G) \cong \text{CH}_K^*(\bar{G})/(e_2)$ and the claim follows.

$({}^2D_{2r}, \mu_2)$. The homomorphism g^* is injective by Theorem 2.6.3, thus $g_{\bar{C}}^*$ is an isomorphism.

$({}^2D_{2r+1}, \mu_2)$. There is a unique nontrivial element $e_1 = c_1(\mathcal{L}(\bar{\omega}_1)) \in \text{CH}_K^1(\bar{G}) \cong \mathbb{F}_2$, we have $g^*(e_1) = 0$ by Lemma A.4.3, and Theorem 2.6.3 yields that g^* induces a monomorphism $\text{CH}_K^*(\bar{G})/(e_1) \rightarrow \text{CH}_K^*(G)$. Thus $\mathcal{C}_K^*(G) \cong \text{CH}_K^*(\bar{G})/(e_1)$ and the claim follows. \square

3 Chow ring of a quasi-split simple group

In this part of the article we compute the Chow ring $\text{Ch}^*(G)$ for a non-split quasi-split simple group G . Here and below we put $\text{Ch}^*(-) := \text{CH}^*(-) \otimes \mathbb{F}_p$ for a given prime number p . The computations proceed differently depending on the relation of p and G .

If $p = [K : k]$ for the splitting field K/k of G , then we analyse the exact sequence

$$\text{Ch}^*(G_K) \xrightarrow{\pi^*} \text{Ch}^*(G) \rightarrow \text{CH}_K^*(G) \rightarrow 0,$$

where $\pi: G_K \rightarrow G$ is the projection. We show that $\pi_* = 0$ unless $(\Delta(G), \pi_1(G)) = ({}^2D_{2r}, {}^2\mu_{2,2})$, thus in all but one cases the computation follows from the computation of $\text{CH}_K^*(G)$ carried out before. In the case of $({}^2D_{2r}, {}^2\mu_{2,2})$ we perform a detailed analysis of the Chow ring $\text{Ch}^*(\text{Gr}(2r-1; q))$ of a quasi-split submaximal isotropic Grassmannian which yields the computation of $\text{Ch}^*(G)$.

If p is coprime to $[K : k]$, then we show that $\text{Ch}^*(G) \cong \text{Ch}^*(G_K)^{\text{Gal}(K/k)}$ and explicitly compute the action of the Galois group and the subalgebra of invariant elements.

In the last remaining case of $\Delta(G) = {}^6D_4$ we combine both the above approaches and also use some explicit computer-assisted computations with Schubert classes.

3.1 Quasi-split submaximal orthogonal Grassmannian

In this section we compute the ring $\text{Ch}^*(\text{Gr}(n-1; q)) := \text{CH}^*(\text{Gr}(n-1; q)) \otimes \mathbb{F}_2$ for the Grassmanian of isotropic subspaces of dimension $n-1$ in a dimension $2n$ vector space equipped with a quadratic form corresponding to a quasi-split simple group G of type $\Delta(G) = {}^2D_n$. For this we use the isomorphism $\text{Ch}^*(\text{Gr}(n-1; q)) \cong \text{Ch}^*(\text{Gr}(n-1; q)_K)^{\text{Gal}(K/k)}$, where K is the splitting field of G , the calculations done in the split case in [Vi07, § 2] and [EKM08, Chapter XVI] and compute explicitly the action of the Galois group. We further use the identification of $\text{Gr}(n-1; q)$ as a G -homogeneous variety to show that

$$c_1(\mathcal{L}(\bar{\omega}_{n-1} + \bar{\omega}_n))^{2^{v_2(n)}} = 0$$

in $\text{Ch}^*(G)$ as in the split case.

We continue to use the notation of Sections A.2 and A.3, and in this section we put

$$\text{Ch}^*(-) := \text{CH}^*(-) \otimes \mathbb{F}_2.$$

Proposition 3.1.1. *Let G be a simply connected quasi-split simple group of type $\Delta(G) = D_n$ or $\Delta(G) = {}^2D_n$ over a field k with the splitting field K/k , and let $T \leq B \leq G$ be a maximal torus and a Borel subgroup. Consider the parabolic subgroup $B \leq P = P_{n-1,n} \leq G$ in the notation of (A.2.6) and the line bundle $\mathcal{L} := \mathcal{L}_P(\varpi_{n-1} + \varpi_n)$ over G/P . Then the following holds.*

1. For the adjoint quotient $G \rightarrow \bar{G}$ and the projection $\varphi: \bar{G} \rightarrow G/P$ we have

$$c_1(\mathcal{L}(\bar{\varpi}_{n-1} + \bar{\varpi}_n))^{2v_2(n)} = c_1(\varphi^* \mathcal{L})^{2v_2(n)} = 0$$

in $\text{Ch}^*(\bar{G})$, where v_2 is the 2-adic valuation.

2. Suppose $\Delta(G) = {}^2D_n$ and put $s := \lfloor \frac{n}{2} \rfloor + 1$, $k_i := \lfloor \log_2 \frac{2n-1}{2i-3} \rfloor$, $2 \leq i \leq s$. Then there is an isomorphism

$$\Theta: \mathbb{F}_2[\varepsilon_1, \varepsilon_2, \dots, \varepsilon_s] / (\varepsilon_1^n, \varepsilon_2^{2^{k_2}}, \dots, \varepsilon_s^{2^{k_s}}) \xrightarrow{\cong} \text{Ch}^*((G/P)_K)$$

such that $\Theta(\varepsilon_1) = c_1(\mathcal{L}_K)$ and for the nontrivial element $\tau \in \text{Gal}(K/k)$ we have

$$\tau(\Theta(\varepsilon_1)) = \Theta(\varepsilon_1), \quad \tau(\Theta(\varepsilon_i)) = \Theta(\varepsilon_i) + \Theta(\varepsilon_1)^{2^{i-3}}, \quad 2 \leq i \leq s.$$

Proof. First we recall the necessary geometric setting (see e.g. [EKM08, Chapter XVI] and [Vi07, § 2]). By [KMRT98, Example 27.10] we have $G \cong \text{Spin}(V, q)$, where (V, q) is a vector space of dimension $2n$ equipped with a non-degenerate quadratic form. For integers $1 \leq i_1 < i_2 < \dots < i_l \leq n$ we denote by $F(i_1, i_2, \dots, i_l; q)$ the variety of totally isotropic flags in (V, q) of the respective dimensions, and put $\text{Gr}(i; q) := F(i; q)$. The standard action of G on V yields an isomorphism

$$\text{Gr}(n-1; q) \cong G/P.$$

The projection

$$\alpha: F(n-1, n; q) \rightarrow \text{Gr}(n-1; q)$$

is a finite morphism of degree 2, and there is a G_K -equivariant isomorphism

$$F(n-1, n; q)_K \cong \text{Gr}(n-1; q)_K \sqcup \text{Gr}(n-1; q)_K \cong (G/P)_K \sqcup (G/P)_K \quad (3.1.1.1)$$

such that α_K is the obvious projection. The projection

$$\beta: F(n-1, n; q) \rightarrow \text{Gr}(n; q)$$

is canonically identified with the projective bundle

$$\beta: \mathbb{P}(\mathcal{E}^\vee) \rightarrow \text{Gr}(n; q)$$

for the tautological rank n vector bundle \mathcal{E} over $\text{Gr}(n; q)$. For the tautological line bundle $\mathcal{O}(-1)$ over $\mathbb{P}(\mathcal{E}^\vee) \cong F(n-1, n; q)$ one has

$$\mathcal{O}(-1)_K \cong \mathcal{L}_{P_K}(\varpi_n - \varpi_{n-1}) \sqcup \mathcal{L}_{P_K}(\varpi_{n-1} - \varpi_n) \quad (3.1.1.2)$$

under isomorphism (3.1.1.1). This decomposition immediately follows from a straightforward computation of the action of the torus T_K on the fibers of $\mathcal{O}(-1)_K$ over the two P_K -stable points of $F(n-1, n; q)_K$.

- (1) We claim that in $\text{Ch}^*(G/P)$ for every $i \in \mathbb{N}_0$ one has

$$c_i(\mathcal{L}^{\oplus n}) = c_i(\mathcal{T}_{G/P}) \quad (3.1.1.3)$$

for the tangent bundle $\mathcal{T}_{G/P}$. Suppose that equality (3.1.1.3) holds and put $N := 2^{v_2(n)}$. Then we have

$$c_N(\mathcal{T}_{G/P}) = c_N(\mathcal{L}^{\oplus n}) = \binom{n}{N} c_1(\mathcal{L})^N = c_1(\mathcal{L})^N$$

with the last equality arising from the fact that $\binom{n}{N}$ is odd by the choice of N . Let $\bar{P} \leq \bar{G}$ be the image of P in the adjoint quotient, then $G/P \cong \bar{G}/\bar{P}$. The tangent bundle $T_{\bar{G}/\bar{P}}$ clearly admits the structure of a \bar{G} -equivariant vector bundle over \bar{G}/\bar{P} , thus for the projection $\varphi: \bar{G} \rightarrow \bar{G}/\bar{P} \cong G/P$ the vector bundle $\varphi^*(T_{G/P})$ is trivial. Then

$$c_1(\varphi^*\mathcal{L})^N = \varphi^*(c_1(\mathcal{L})^N) = \varphi^*(c_N(\mathcal{T}_{G/P})) = c_N(\varphi^*\mathcal{T}_{G/P}) = 0.$$

Now we turn to the proof of the equality (3.1.1.3). The injectivity part of Lemma 2.6.1 yields that it suffices to prove the equality after a base change to K , so till the end of the proof of claim (1) we assume that G is split, i.e. $K = k$ and the quadratic form q is hyperbolic. By [EKM08, Example 104.20] we have a short exact sequence of vector bundles

$$0 \rightarrow \mathcal{O}_{\mathbb{P}(\mathcal{E}^\vee)} \rightarrow \mathcal{O}(-1) \otimes \beta^*\mathcal{E}^\vee \rightarrow \mathcal{T}_\beta \rightarrow 0,$$

where $\beta: \mathbb{P}(\mathcal{E}^\vee) \rightarrow \text{Gr}(n; q)$ is the projection and \mathcal{T}_β is the relative tangent bundle that is defined by the short exact sequence

$$0 \rightarrow \mathcal{T}_\beta \rightarrow \mathcal{T}_{\mathbb{P}(\mathcal{E}^\vee)} \rightarrow \beta^*\mathcal{T}_{\text{Gr}(n; q)} \rightarrow 0.$$

For a vector bundle \mathcal{V} over $\mathbb{P}(\mathcal{E}^\vee)$ put

$$c_t(\mathcal{V}) := 1 + c_1(\mathcal{V})t + c_2(\mathcal{V})t^2 + \dots \in \text{Ch}^*(\mathbb{P}(\mathcal{E}^\vee))[[t]]$$

to be the total Chern class mod 2. Multiplicativity of the total Chern classes with respect to the short exact sequences applied to the sequences above yields

$$c_t(\mathcal{T}_{\mathbb{P}(\mathcal{E}^\vee)}) = c_t(\beta^*\mathcal{T}_{\text{Gr}(n; q)}) \cdot c_t(\mathcal{O}(-1) \otimes \beta^*\mathcal{E}^\vee).$$

Furthermore, it follows from [EKM08, Proposition 86.13 and Exercise 87.8] that $c_t(\beta^*\mathcal{E}^\vee) = 1$ and $c_t(\beta^*\mathcal{T}_{\text{Gr}(n; q)}) = 1$. Hence

$$c_t(\mathcal{T}_{\mathbb{P}(\mathcal{E}^\vee)}) = c_t(\mathcal{O}(-1) \otimes \beta^*\mathcal{E}^\vee) = c_t(\mathcal{O}(-1)^{\oplus n}).$$

Restricting to a connected component of $\mathbb{P}(\mathcal{E}^\vee)$ via decomposition (3.1.1.1) and using (3.1.1.2) we obtain

$$c_i(\mathcal{T}_{G/P}) = c_i(\mathcal{L}_P(\varpi_n - \varpi_{n-1})^{\oplus n}).$$

In the mod 2 Chow groups we have $c_1(\mathcal{L}_P(\varpi_n - \varpi_{n-1})) = c_1(\mathcal{L}_P(\varpi_n + \varpi_{n-1})) = c_1(\mathcal{L})$, thus

$$c_i(\mathcal{T}_{G/P}) = c_i(\mathcal{L}_P(\varpi_n - \varpi_{n-1})^{\oplus n}) = c_i(\mathcal{L}^{\oplus n}).$$

(2) Consider the following diagram with all the morphisms being projections.

$$\begin{array}{ccccc}
& & \text{F}(1, n-1; q)_K & \xrightarrow{f_2} & \text{Gr}(n-1; q)_K & & \\
& & \swarrow f_1 & & \nwarrow \alpha_K & & \\
& \text{Gr}(1; q)_K & & & & \text{F}(n-1, n; q)_K & (3.1.1.4) \\
& & \swarrow g_1 & & \nwarrow \beta_K & & \\
& & \text{F}(1, n; q)_K & \xrightarrow{g_2} & \text{Gr}(n; q)_K & &
\end{array}$$

The variety $\text{Gr}(1; q)_K$ is a smooth projective hyperbolic quadric of dimension $2n - 2$, thus [EKM08, Proposition 68.1] yields

$$\text{Ch}^i(\text{Gr}(1; q)_K) = \begin{cases} \mathbb{F}_2 \cdot h^i, & 0 \leq i \leq n - 2, \\ \mathbb{F}_2 \cdot h^{n-1} \oplus \mathbb{F}_2 \cdot l_{n-1}, & i = n - 1, \\ \mathbb{F}_2 \cdot l_{2n-2-i}, & n \leq i \leq 2n - 2, \end{cases}$$

where $h \in \text{Ch}^1(\text{Gr}(1; q)_K)$ is the class of a hyperplane section and $l_{2n-2-i} \in \text{Ch}^i(\text{Gr}(1; q)_K)$ is the class of a totally isotropic projective subspace in $\text{Gr}(1; q)_K$ of dimension $2n - 2 - i$. Note that there are two different classes $l_{n-1}, l'_{n-1} \in \text{Ch}^{n-1}(\text{Gr}(1; q)_K)$ of totally isotropic projective subspaces in $\text{Gr}(1; q)_K$ of dimension $n - 1$ and one has $l_{n-1} + l'_{n-1} = h^{n-1}$. Put

$$\begin{aligned} z_0 &:= (g_2)_* g_1^*(l_{n-1}) \in \text{Ch}^0(\text{Gr}(n; q)_K), & z'_0 &:= (g_2)_* g_1^*(l'_{n-1}) \in \text{Ch}^0(\text{Gr}(n; q)_K), \\ z_i &:= (g_2)_* g_1^*(l_{n-1-i}) \in \text{Ch}^i(\text{Gr}(n; q)_K), & & 1 \leq i \leq n - 1, \end{aligned}$$

then [EKM08, Proposition 86.16] yields that

$$\begin{aligned} \mathbb{F}_2[\varepsilon_0, \varepsilon'_0, \varepsilon_2, \varepsilon_3, \dots, \varepsilon_s] / (\varepsilon_0^2 - \varepsilon_0, (\varepsilon'_0)^2 - \varepsilon'_0, \varepsilon_0 \varepsilon'_0, \varepsilon_2^{2^{k_2}}, \dots, \varepsilon_s^{2^{k_s}}) &\xrightarrow{\cong} \text{Ch}^*(\text{Gr}(n; q)_K), \\ \varepsilon_0 \mapsto z_0, \quad \varepsilon'_0 \mapsto z'_0, \quad \varepsilon_i \mapsto z_{2i-3}, \quad 2 \leq i \leq s, \end{aligned}$$

is an isomorphism. Applying the projective bundle formula to

$$\beta: \text{F}(n - 1, n; q)_K \cong \mathbb{P}(\mathcal{E}^\vee)_K \rightarrow \text{Gr}(n; q)_K$$

and using [EKM08, Proposition 86.13] we obtain an isomorphism

$$\begin{aligned} \Psi: \mathbb{F}_2[\varepsilon_0, \varepsilon'_0, \varepsilon_1, \varepsilon_1, \dots, \varepsilon_s] / (\varepsilon_0^2 - \varepsilon_0, (\varepsilon'_0)^2 - \varepsilon'_0, \varepsilon_0 \varepsilon'_0, \varepsilon_1^n, \varepsilon_2^{2^{k_2}}, \dots, \varepsilon_s^{2^{k_s}}) &\xrightarrow{\cong} \text{Ch}^*(\text{F}(n - 1, n; q)_K), \\ \varepsilon_0 \mapsto \beta_K^*(z_0), \quad \varepsilon'_0 \mapsto \beta_K^*(z'_0), \quad \varepsilon_1 \mapsto c_1(\mathcal{O}(-1)_K), \quad \varepsilon_i \mapsto \beta_K^*(z_{2i-3}), \quad 2 \leq i \leq s, \end{aligned}$$

where $c_1(\mathcal{O}(-1)_K)$ is the first Chern class of the tautological line bundle over $\text{F}(n - 1, n; q)_K$.

In view of decomposition (3.1.1.1) the composition

$$\begin{aligned} \text{Ch}^*((G/P)_K) &\cong \text{Ch}^*(\text{Gr}(n - 1; q)_K) \xrightarrow{\alpha_K^*} \text{Ch}^*(\text{F}(n - 1, n; q)_K) \xrightarrow{\Psi^{-1}} \\ &\xrightarrow{\Psi^{-1}} \mathbb{F}_2[\varepsilon_0, \varepsilon'_0, \varepsilon_1, \dots, \varepsilon_s] / (\varepsilon_0^2 - \varepsilon_0, (\varepsilon'_0)^2 - \varepsilon'_0, \varepsilon_0 \varepsilon'_0, \varepsilon_1^n, \varepsilon_2^{2^{k_2}}, \dots, \varepsilon_s^{2^{k_s}}) \xrightarrow{\Pi} \\ &\xrightarrow{\Pi} \mathbb{F}_2[\varepsilon_1, \varepsilon_2, \dots, \varepsilon_s] / (\varepsilon_1^n, \varepsilon_2^{2^{k_2}}, \dots, \varepsilon_s^{2^{k_s}}) \end{aligned}$$

with the projection Π given by $\Pi(\varepsilon_0) = 1, \Pi(\varepsilon'_0) = 0$, is an isomorphism. Denote by

$$\Theta: \mathbb{F}_2[\varepsilon_1, \varepsilon_2, \dots, \varepsilon_s] / (\varepsilon_1^n, \varepsilon_2^{2^{k_2}}, \dots, \varepsilon_s^{2^{k_s}}) \rightarrow \text{Ch}^*((G/P)_K)$$

the inverse isomorphism. Since

$$c_1(\mathcal{L}_K) = c_1(\mathcal{L}_{P_K}(\varpi_n - \varpi_{n-1})) = c_1(\mathcal{L}_{P_K}(\varpi_{n-1} - \varpi_n))$$

in $\text{Ch}^*((G/P)_K)$, it follows from decomposition (3.1.1.2) that $\alpha_K^*(c_1(\mathcal{L}_K)) = c_1(\mathcal{O}(-1)_K)$ and

$$\Theta(\varepsilon_1) = c_1(\mathcal{L}_K).$$

Furthermore, for the nontrivial element $\tau \in \text{Gal}(K/k)$ we have

$$\tau(\Theta(\varepsilon_1)) = \tau(c_1(\mathcal{L}_K)) = c_1(\mathcal{L}_K) = \Theta(\varepsilon_1),$$

since $c_1(\mathcal{L}_K)$ is obtained by a base change from k to K .

For the computation of the action of $\text{Gal}(K/k)$ on $\Theta(\varepsilon_i)$, $2 \leq i \leq s$, note that all the morphisms in diagram (3.1.1.4) and the isomorphism $(G/P)_K \cong \text{Gr}(n - 1; q)_K$ are given by a base change of the

corresponding morphisms over k , thus the action of $\text{Gal}(K/k)$ commutes with the respective pullbacks and pushforwards. For the nontrivial element $\tau \in \text{Gal}(K/k)$ we have

$$\tau(l_{n-1}) = l'_{n-1}, \quad \tau(l'_{n-1}) = l_{n-1}, \quad \tau(l_{n-1-i}) = l_{n-1-i}, \quad 1 \leq i \leq n-1,$$

with the latter equality arising from the fact that $\text{Ch}^{n-1+i}(\text{Gr}(1; q)) \cong \mathbb{F}_2$, $1 \leq i \leq n-1$, and \mathbb{F}_2 admits only a trivial action of $\text{Gal}(K/k)$. Thus

$$\tau(\beta_K^*(z_0)) = \beta_K^*(z'_0), \quad \tau(\beta_K^*(z_i)) = \beta_K^*(z_i), \quad 1 \leq i \leq n-1.$$

Furthermore, put $x := c_1(\mathcal{O}(-1)_K) \in \text{Ch}^1(\text{F}(n-1, n; q)_K)$, then we have $\tau(x) = x$, since x is obtained by a base change from k . Consider the elements

$$\begin{aligned} y_1 &:= (f_2)_* f_1^*(l_{n-1}) \in \text{Ch}^1(\text{Gr}(n-1; q)_K), & y'_1 &:= (f_2)_* f_1^*(l'_{n-1}) \in \text{Ch}^1(\text{Gr}(n-1; q)_K), \\ y_i &:= (f_2)_* f_1^*(l_{n-i}) \in \text{Ch}^i(\text{Gr}(n-1; q)_K), & & 2 \leq i \leq n-1. \end{aligned}$$

It follows from [Vi07, Lemma 2.6] (loc. cit. has a blanket assumption that $\text{char } k \neq 2$, but the same proof of the lemma works in general) that

$$\alpha_K^*(y'_1) = \beta_K^*(z_1) + x \cdot \beta_K^*(z'_0), \quad \alpha_K^*(y_i) = \beta_K^*(z_i) + x \cdot \beta_K^*(z_{i-1}), \quad 1 \leq i \leq n-1.$$

In particular, we have $x = \alpha_K^*(y_1 + y'_1)$. Then

$$\beta_K^*(z_i) + x^i \cdot \beta_K^*(z'_0) = \sum_{j=0}^{i-2} x^j \cdot \alpha_K^*(y_{i-j}) + x^{i-1} \cdot \alpha_K^*(y'_1) = \alpha_K^*(w_i), \quad w_i := \sum_{j=0}^{i-2} (y_1 + y'_1)^j y_{i-j} + (y_1 + y'_1)^{i-1} y'_1,$$

where $w_i \in \text{Ch}^*(\text{Gr}(n-1; q)_K)$. Thus for $2 \leq i \leq s$ we have

$$\begin{aligned} \Theta^{-1}(w_{2i-3}) &= \Pi \circ \Psi^{-1} \circ \alpha_K^*(w_{2i-3}) = \Pi \circ \Psi^{-1}(\beta_K^*(z_{2i-3}) + x^{2i-3} \cdot \beta_K^*(z'_0)) = \varepsilon_i, \\ \Theta^{-1}(\tau(w_{2i-3})) &= \Pi \circ \Psi^{-1} \circ \alpha_K^*(\tau(w_{2i-3})) = \Pi \circ \Psi^{-1} \circ \tau(\beta_K^*(z_{2i-3}) + x^{2i-3} \cdot \beta_K^*(z'_0)) = \\ &= \Pi \circ \Psi^{-1}(\beta_K^*(z_{2i-3}) + x^{2i-3} \cdot \beta_K^*(z_0)) = \varepsilon_i + \varepsilon_1^{2i-3}. \end{aligned}$$

It follows that $\tau(\Theta(\varepsilon_i)) = \Theta(\varepsilon_i) + \Theta(\varepsilon_1)^{2i-3}$ for $2 \leq i \leq s$. \square

Lemma 3.1.2. *For $n \in \mathbb{N}$ put*

$$s := \left\lfloor \frac{n}{2} \right\rfloor + 1, \quad k_1 := v_2(n), \quad k_i := \left\lfloor \log_2 \frac{2n-1}{2i-3} \right\rfloor, \quad 2 \leq i \leq s,$$

where v_2 is the 2-adic valuation. Consider the algebra

$$\mathcal{P} := \mathcal{P}_n := \mathbb{F}_2[\varepsilon_1, \varepsilon_2, \dots, \varepsilon_s] / (\varepsilon_1^n, \varepsilon_2^{2^{k_2}}, \dots, \varepsilon_s^{2^{k_s}}),$$

denote by $\bar{\varepsilon}_i$ the image of ε_i in the quotient algebra \mathcal{P} , let $\tau: \mathcal{P} \rightarrow \mathcal{P}$ be the involution given by $\tau(\bar{\varepsilon}_1) = \bar{\varepsilon}_1$, $\tau(\bar{\varepsilon}_i) = \bar{\varepsilon}_i + \bar{\varepsilon}_1^{2i-3}$, $2 \leq i \leq s$, and put

$$\mathcal{P}^\tau := \{x \in \mathcal{P} \mid \tau(x) = x\} \subseteq \mathcal{P}$$

to be the subalgebra of τ -invariant elements. For $1 \leq m \leq n$ consider the sequence

$$\mathcal{P}/(\bar{\varepsilon}_1^m) \xrightarrow{\text{id}+\tau} \mathcal{P}^\tau/(\bar{\varepsilon}_1^m) \xrightarrow{J} \mathcal{P}/(\bar{\varepsilon}_1^m),$$

where J is induced by the inclusion $\mathcal{P}^\tau \subseteq \mathcal{P}$. Then the following holds.

1. $(\text{id}+\tau)(\mathcal{P}/(\bar{\varepsilon}_1^m)) = \bar{\varepsilon}_1 \cdot (\mathcal{P}^\tau/(\bar{\varepsilon}_1^m))$.
2. J restricted to $(\text{id}+\tau)(\mathcal{P}/(\bar{\varepsilon}_1^m))$ is injective.

3. There is an isomorphism

$$\begin{aligned} \mathcal{P}^\tau / (\bar{\varepsilon}_1^m) &\cong \mathbb{F}_2[e_1, e_2, \dots, e_s, x] / (e_1^m, e_1 \cdot x, x^2, h_2, h_3, \dots, h_s), \\ h_2 &:= e_1^{2^{k_2-n}} \cdot x + \sum_{j=0}^{k_2-1} e_1^{2^{k_2-2^{j+1}}} \cdot e_2^{2^j}, \\ h_i &:= e_i^{2^{k_i}} + e_1^{(2i-3) \cdot 2^{k_i-n}} \cdot x + e_1^{(2i-4) \cdot 2^{k_i}} \cdot \left(\sum_{j=0}^{k_i-1} e_1^{2^{k_i-2^{j+1}}} \cdot e_2^{2^j} \right), \quad 3 \leq i \leq s, \end{aligned}$$

given by $e_1 \mapsto \bar{\varepsilon}_1$, $x \mapsto \bar{\varepsilon}_1^{n-1} \bar{\varepsilon}_2$, $e_2 \mapsto \bar{\varepsilon}_2^2 + \bar{\varepsilon}_1 \bar{\varepsilon}_2$, $e_i \mapsto \bar{\varepsilon}_i + \bar{\varepsilon}_1^{2i-4} \bar{\varepsilon}_2$, $3 \leq i \leq s$. In particular, we have

$$\mathcal{P}^\tau / (\bar{\varepsilon}_1^{2^{k_1}}) \cong \begin{cases} \mathbb{F}_2[e_2, \dots, e_s, x] / (e_2^{2^{k'_2}}, \dots, e_s^{2^{k'_s}}, x^2), & n \text{ odd}, \\ \mathbb{F}_2[e_1, \dots, e_s] / (e_1^{2^{k'_1}}, e_2^{2^{k'_2}}, \dots, e_s^{2^{k'_s}}, e_1 \cdot e_1^{2^{k'_1-1}}), & n \text{ even}, n \neq 2^{k_1}, \\ \mathbb{F}_2[e_1, \dots, e_s] / \left(e_1^{2^{k'_1}}, e_2^{2^{k'_2}}, \dots, e_s^{2^{k'_s}}, e_1 \cdot \left(\sum_{j=0}^{k'_2-1} e_1^{2^{k'_2-2^{j+1}}} \cdot e_2^{2^j} \right) \right), & n = 2^{k_1}, \end{cases}$$

where $l \in \mathbb{N}$ is such that $n = 2^k(2l-3)$ and

$$k'_i = \begin{cases} k_i, & i \neq 2, l, \\ k_i - 1, & i = 2 \neq l, \\ k_i + 1, & l = i \neq 1, \\ k_i, & i = l = 2. \end{cases}$$

Note that $k'_1 = v_2(n)$, $k'_2 = \lceil \log_2 n \rceil$ and $k'_i = \lceil \log_2 \frac{2n}{2i-3} \rceil$, $3 \leq i \leq s$.

Proof. Put

$$\bar{e}_1 := \bar{\varepsilon}_1, \quad \bar{e}_2 := \bar{\varepsilon}_2^2 + \bar{\varepsilon}_1 \bar{\varepsilon}_2 \in \mathcal{P}^\tau, \quad \bar{e}_i := \bar{\varepsilon}_i + \bar{\varepsilon}_1^{2i-4} \bar{\varepsilon}_2 \in \mathcal{P}^\tau, \quad 3 \leq i \leq s.$$

The monomials

$$\bar{\varepsilon}_1^{i_1} \bar{\varepsilon}_2^{i_2} \dots \bar{\varepsilon}_s^{i_s}, \quad 0 \leq i_1 < n, 0 \leq i_j < 2^{k_j}, 2 \leq j \leq s,$$

form a basis of \mathcal{P} . Inducting on the multidegree $(i_s i_{s-1} \dots i_1)$ of the monomial $\bar{\varepsilon}_1^{i_1} \bar{\varepsilon}_2^{i_2} \dots \bar{\varepsilon}_s^{i_s}$ in the lexicographical order it is straightforward to see that $\bar{\varepsilon}_1^{i_1} \bar{\varepsilon}_2^{i_2} \dots \bar{\varepsilon}_s^{i_s} \in \mathcal{P}^\tau + \mathcal{P}^\tau \bar{\varepsilon}_2$. It follows that 1 and $\bar{\varepsilon}_2$ generate \mathcal{P} as a module over \mathcal{P}^τ . The morphism $\text{id} + \tau$ is clearly \mathcal{P}^τ -linear and we have

$$(\text{id} + \tau)(1) = 0, \quad (\text{id} + \tau)(\bar{\varepsilon}_2) = \bar{\varepsilon}_1,$$

so the first claim follows.

Put $\bar{x} := \bar{\varepsilon}_1^{n-1} \bar{\varepsilon}_2$. Again inducting on the multidegree one obtains that the set

$$\bar{e}_1^{i_1} \bar{e}_2^{i_2} \dots \bar{e}_s^{i_s}, \quad \bar{e}_2^{i_2} \dots \bar{e}_s^{i_s} \bar{x}, \quad 0 \leq i_1 < n, 0 \leq i_2 < 2^{k_2-1}, 0 \leq i_j < 2^{k_j}, 3 \leq j \leq s, \quad (3.1.2.1)$$

generates \mathcal{P}^τ as a vector space over \mathbb{F}_2 . Furthermore, these elements are linearly independent in \mathcal{P} , thus (3.1.2.1) is a basis of \mathcal{P}^τ . We have $\bar{\varepsilon}_1 \bar{x} = \bar{\varepsilon}_1^n \bar{\varepsilon}_2 = 0$, $\bar{\varepsilon}_1 \bar{e}_1^{i_1} = \bar{e}_1^{i_1+1}$, $\bar{e}_1^n = 0$, thus multiplication by $\bar{\varepsilon}_1$ takes the elements of this basis either to 0 or to distinct elements of the basis. It follows that

$$\bar{e}_1^{i_1} \bar{e}_2^{i_2} \dots \bar{e}_s^{i_s}, \quad \bar{e}_2^{i_2} \dots \bar{e}_s^{i_s} \bar{x}, \quad 0 \leq i_1 < m, 0 \leq i_2 < 2^{k_2-1}, 0 \leq i_j < 2^{k_j}, 3 \leq j \leq s, \quad (3.1.2.2)$$

form a basis of $\mathcal{P}^\tau / (\bar{\varepsilon}_1^m)$. Then $(\text{id} + \tau)(\mathcal{P} / (\bar{\varepsilon}_1^m)) = \bar{\varepsilon}_1 \cdot (\mathcal{P}^\tau / (\bar{\varepsilon}_1^m))$ is generated by the elements

$$\bar{e}_1^{i_1} \bar{e}_2^{i_2} \dots \bar{e}_s^{i_s}, \quad 1 \leq i_1 < m, 0 \leq i_2 < 2^{k_2-1}, 0 \leq i_j < 2^{k_j}, 3 \leq j \leq s,$$

whose images under J are clearly linearly independent. This proves the second claim.

For the third claim we first check the multiplicative relations between \bar{e}_i and \bar{x} . The relation $\bar{e}_1^m = 0$ is clear, the relations $\bar{e}_1 \cdot \bar{x} = 0$ and $\bar{x}^2 = 0$ hold in \mathcal{P} , thus also in \mathcal{P}^τ and $\mathcal{P}^\tau / (\bar{\varepsilon}_1^m)$. The relations $\bar{h}_i = 0$,

where \bar{h}_i is the image of h_i in \mathcal{P} , also hold already in \mathcal{P} , thus in \mathcal{P}^τ and $\mathcal{P}^\tau/(\bar{\varepsilon}_1^m)$, which could be seen expanding

$$\bar{e}_2^{2^{k_2-1}} = (\bar{\varepsilon}_2^2 + \bar{\varepsilon}_1 \bar{\varepsilon}_2)^{2^{k_2-1}} = \bar{\varepsilon}_1^{2^{k_2-1}} \bar{\varepsilon}_2^{2^{k_2-1}}, \quad \bar{e}_i^{2^{k_i}} = (\bar{\varepsilon}_i + \bar{\varepsilon}_1^{2i-4} \bar{\varepsilon}_2)^{2^{k_i}} = \bar{\varepsilon}_1^{(2i-4) \cdot 2^{k_i}} \bar{\varepsilon}_2^{2^{k_i}}, \quad 3 \leq i \leq s,$$

and substituting $\bar{e}_1 = \bar{\varepsilon}_1$, $\bar{e}_2 = \bar{\varepsilon}_2 + \bar{\varepsilon}_1 \bar{\varepsilon}_2$ and $\bar{x} = \bar{\varepsilon}_1^{-1} \bar{\varepsilon}_2$ into the formula defining \bar{h}_i . Thus we have a homomorphism

$$\mathbb{F}_2[e_1, e_2, \dots, e_s, x]/(e_1^m, e_1 \cdot x, x^2, h_2, h_3, \dots, h_s) \rightarrow \mathcal{P}^\tau/(\bar{\varepsilon}_1^m)$$

induced by $e_i \mapsto \bar{e}_i$ and $x \mapsto \bar{x}$. This homomorphism is clearly surjective, and is an isomorphism by the dimension count using the basis (3.1.2.2) on the right-hand side.

Finally, let $m = 2^{k_1}$ and let $l \in \mathbb{N}$ be such that $n = 2^{k_1}(2l - 3)$. Note that the integers k_i , $2 \leq i \leq s$, are defined in such a way that the following inequalities hold:

$$(2i - 3)2^{k_i-1} < n \leq (2i - 3)2^{k_i}.$$

First suppose $l \neq 2$. Then we have

$$\begin{aligned} 2^{k_2} - n &\geq 2^{k_1}, & 2^{k_2} - 2^{j+1} &\geq 2^{k_2-1} \geq 2^{k_1}, & 0 \leq j &\leq k_2 - 2, \\ (2i - 4) \cdot 2^{k_i} &\geq 2^{k_1}, & 3 \leq i &\leq s, & (2i - 3) \cdot 2^{k_i} - n &\geq 1, & i \neq l, \end{aligned}$$

and it follows that for some $h'_i \in \mathbb{F}_2[e_1, e_2, \dots, e_s, x]$ we have

$$h_2 = e_2^{2^{k_2-1}} + e_1^{2^{k_1}} \cdot h'_2, \quad h_i = e_i^{2^{k_i}} + (e_1 \cdot x) \cdot e_1^{(2i-3) \cdot 2^{k_i} - n - 1} + e_1^{2^{k_1}} \cdot h'_i, \quad 3 \leq i \leq s, i \neq l.$$

If n is odd, then $l > s$, so it follows from the above that

$$\mathbb{F}_2[e_1, \dots, e_s, x]/(e_1^{2^{k_1}}, e_1 \cdot x, x^2, h_2, h_3, \dots, h_s) = \mathbb{F}_2[e_1, \dots, e_s, x]/(e_1^{2^{k_1}}, e_1 \cdot x, x^2, e_2^{2^{k_2-1}}, e_3^{2^{k_3}}, \dots, e_s^{2^{k_s}}),$$

since the ideals are the same. Furthermore, we have $k_1 = 0$ and

$$\mathbb{F}_2[e_1, \dots, e_s, x]/(e_1^{2^{k_1}}, e_1 \cdot x, x^2, e_2^{2^{k_2-1}}, e_3^{2^{k_3}}, \dots, e_s^{2^{k_s}}) \cong \mathbb{F}_2[e_2, \dots, e_s, x]/(e_2^{2^{k_2-1}}, e_3^{2^{k_3}}, \dots, e_s^{2^{k_s}}, x^2).$$

If n is even, then $l \leq s$ and we have $h_l = e_l^{2^{k_l}} + x + e_1^{2^{k_1}} \cdot h'_l$, thus

$$\begin{aligned} \mathbb{F}_2[e_1, \dots, e_s, x]/(e_1^{2^{k_1}}, e_1 \cdot x, x^2, h_2, h_3, \dots, h_s) &= \\ &= \mathbb{F}_2[e_1, \dots, e_s, x]/(e_1^{2^{k_1}}, e_1 \cdot x, x^2, e_2^{2^{k_2-1}}, e_3^{2^{k_3}}, \dots, e_{l-1}^{2^{k_{l-1}}}, e_l^{2^{k_l}} + x, e_{l+1}^{2^{k_{l+1}}}, \dots, e_s^{2^{k_s}}) \cong \\ &\cong \mathbb{F}_2[e_1, \dots, e_s]/(e_1^{2^{k_1}}, e_2^{2^{k_2-1}}, e_3^{2^{k_3}}, \dots, e_s^{2^{k'_s}}, e_1 \cdot e_l^{2^{k'_l-1}}), \end{aligned}$$

where $k'_i = k_i$, if $i \neq l$ and $k'_l = k_l + 1$. The claim follows.

Now suppose $l = 2$, i.e. $n = 2^{k_1}$, then

$$(2i - 3) \cdot 2^{k_i} - n \geq 1, \quad (2i - 4) \cdot 2^{k_i} \geq 2^{k_1}, \quad 3 \leq i \leq s,$$

and it follows that for some $h'_i \in \mathbb{F}_2[e_1, e_2, \dots, e_s, x]$ we have

$$h_i = e_i^{2^{k_i}} + (e_1 \cdot x) \cdot e_1^{(2i-3) \cdot 2^{k_i} - n - 1} + e_1^{2^{k_1}} \cdot h'_i, \quad 3 \leq i \leq s.$$

Then we have

$$\mathbb{F}_2[e_1, \dots, e_s, x]/(e_1^{2^{k_1}}, e_1 \cdot x, x^2, h_2, h_3, \dots, h_s) = \mathbb{F}_2[e_1, \dots, e_s, x]/(e_1^{2^{k_1}}, e_1 \cdot x, x^2, h_2, e_3^{2^{k_3}}, \dots, e_s^{2^{k_s}}),$$

since the ideals are the same. For h_2 we have

$$h_2 = e_2^{2^{k_2-1}} + x + \sum_{j=0}^{k_2-2} e_1^{2^{k_2-2j+1}} \cdot e_2^j, \quad h_2 = e_2^{2^{k_2}} + x^2 + \sum_{j=0}^{k_2-2} e_1^{2^{k_2+1-2j+2}} \cdot e_2^{j+1} = e_2^{2^{k_2}} + x^2 + e_1^{2^{k_1}} \cdot h'_2$$

for some $h'_2 \in \mathbb{F}_2[e_1, e_2, \dots, e_s, x]$, since

$$2^{k_2+1} - 2^{j+2} \geq 2^{k_2} = 2^{k_1}, \quad 0 \leq j \leq k_2 - 2.$$

Then using the equality expressing x via h_2 , e_1 and e_2 we obtain

$$\begin{aligned} \mathbb{F}_2[e_1, \dots, e_s, x] / \left(e_1^{2^{k_1}}, e_1 \cdot x, x^2, h_2, e_3^{2^{k_3}}, \dots, e_s^{2^{k_s}} \right) &\cong \\ &\cong \mathbb{F}_2[e_1, \dots, e_s] / \left(e_1^{2^{k_1}}, e_2^{2^{k_2}}, \dots, e_s^{2^{k_s}}, e_1 \cdot \left(\sum_{j=0}^{k_2-1} e_1^{2^{k_2}-2^{j+1}} \cdot e_2^{2^j} \right) \right). \end{aligned}$$

□

3.2 Chow ring of an adjoint quasi-split simple group of type ${}^2D_{2r}$

In this section we compute the ring $\text{Ch}^*(G) := \text{CH}^*(G) \otimes \mathbb{F}_2$ for an adjoint quasi-split simple group of type $\Delta(G) = {}^2D_{2r}$. For this we use the previous computations with the submaximal isotropic Grassmannian and the computation of the conormed Chow ring $\text{CH}_K^*(G)$ for the splitting field K of G .

We continue to use the notation of Sections A.2 and A.3. In this section we put

$$\text{Ch}^*(-) := \text{CH}^*(-) \otimes \mathbb{F}_2.$$

Lemma 3.2.1. *Let G be a split simple group of type $\Delta(G) = D_n$ over a field k and let $T \leq B \leq G$ be a maximal torus and a Borel subgroup. Consider the parabolic subgroup $B \leq P = P_{n-1, n} \leq G$ in the notation of (A.2.6). Then the pullback homomorphism*

$$\varphi^*: \text{CH}^*(G/P) \rightarrow \text{CH}^*(G)$$

for the projection $\varphi: G \rightarrow G/P$ is surjective.

Proof. The derived subgroup $[L(P), L(P)] \leq L(P)$ of the Levi subgroup $L(P) \leq P$ is isomorphic to SL_{n-1} , thus P is special and the claim follows from [PS17, Lemma 7.1]. □

Theorem 3.2.2. *Let G be an adjoint quasi-split simple group over a field k of type $\Delta(G) = {}^2D_{2r}$, $r \geq 2$.*

1. *Let $T \leq B \leq G$ be a maximal torus and a Borel subgroup, consider the parabolic subgroup $B \leq P = P_{2r-1, 2r} \leq G$ in the notation of (A.2.6) and put $\mathcal{L} := \mathcal{L}_P(\varpi_{2r-1} + \varpi_{2r})$ for the respective line bundle over G/P . Then the pullback for the projection $\varphi: G \rightarrow G/P$ induces an isomorphism*

$$\tilde{\varphi}^*: \text{Ch}^*(G/P) / (c_1(\mathcal{L})^{v_2(2r)}) \xrightarrow{\cong} \text{Ch}^*(G).$$

In particular, if $2r$ is a power of 2, then the pullback homomorphism

$$\varphi^*: \text{Ch}^*(G/P) \rightarrow \text{Ch}^*(G)$$

is an isomorphism.

2. *There is an isomorphism*

$$\text{Ch}^*(G) \cong \begin{cases} \mathbb{F}_2[e_1, \dots, e_s] / \left(e_1^{2^{k_1}}, \dots, e_s^{2^{k_s}}, e_1 \cdot \left(\sum_{j=0}^{k_1-1} e_1^{2^{k_1}-2^{j+1}} \cdot e_2^{2^j} \right) \right), & l = 2, \\ \mathbb{F}_2[e_1, \dots, e_s] / (e_1^{2^{k_1}}, \dots, e_s^{2^{k_s}}, e_1 \cdot e_1^{2^{k_1-1}}), & l \geq 3, \end{cases}$$

where $l \in \mathbb{N}$ is such that $2r = 2^m(2l - 3)$ and

$$\begin{aligned} s &= r + 1, \quad \deg e_1 = 1, \quad \deg e_2 = 2, \quad k_1 = v_2(2r), \quad k_2 = \lceil \log_2 2r \rceil, \\ \deg e_i &= 2i - 3, \quad k_i = \left\lceil \log_2 \frac{4r}{2i - 3} \right\rceil, \quad 3 \leq i \leq s. \end{aligned}$$

Here v_2 is the 2-adic valuation.

Proof. Let K/k be the splitting field of G , let \mathcal{P} be the algebra with involution τ defined in Lemma 3.1.2 and let the non-trivial element of $\text{Gal}(K/k)$ act on \mathcal{P} via the involution τ . Then Proposition 3.1.1.(2) combined with Lemma 2.6.1 yields isomorphisms

$$\Theta: \mathcal{P} \xrightarrow{\cong} \text{Ch}^*((G/P)_K), \quad \Theta^\tau: \mathcal{P}^\tau \xrightarrow{\cong} \text{Ch}^*(G/P).$$

Furthermore, Lemma 3.1.2.(1) yields an isomorphism

$$\text{Ch}^*(G/P)/(c_1(\mathcal{L})) \xrightarrow{\cong} \text{Ch}_K^*(G/P).$$

We have $\varphi^*(c_1(\mathcal{L})^{v_2(2r)}) = 0$ by Proposition 3.1.1.(1), thus we have the following commutative diagram with exact rows.

$$\begin{array}{ccccccc} \text{Ch}^*(G_K) & \xrightarrow{\pi_*} & \text{Ch}^*(G) & \longrightarrow & \text{Ch}_K^*(G) & \longrightarrow & 0 \\ \tilde{\varphi}_K^* \uparrow & & \tilde{\varphi}^* \uparrow & & \tilde{\varphi}^* \uparrow & & \\ \text{Ch}^*((G/P)_K)/(c_1(\mathcal{L}_K)^{v_2(2r)}) & \xrightarrow{\tilde{\rho}_*} & \text{Ch}^*(G/P)/(c_1(\mathcal{L})^{v_2(2r)}) & \longrightarrow & \text{Ch}_K^*(G/P) & \longrightarrow & 0 \\ \tilde{\Theta} \uparrow \cong & & \tilde{\Theta}^\tau \uparrow \cong & & \cong \uparrow & & \\ \mathcal{P}/(\bar{\varepsilon}_1^{v_2(2r)}) & \xrightarrow{\text{id}+\tau} & \mathcal{P}^\tau/(\bar{\varepsilon}_1^{v_2(2r)}) & \longrightarrow & \mathcal{P}/(\bar{\varepsilon}_1) & \longrightarrow & 0 \end{array}$$

Here $\tilde{\rho}_*$ is induced by the projection $\rho: (G/P)_K \rightarrow G/P$, the isomorphisms $\tilde{\Theta}$ and $\tilde{\Theta}^\tau$ are induced by Θ , the homomorphisms $\tilde{\varphi}^*$, $\tilde{\varphi}_K^*$, $\tilde{\varphi}^*$ are induced by the pullback for the projection $\varphi: G \rightarrow G/P$, and the unlabelled morphisms are the quotient morphisms. The homomorphism $\tilde{\varphi}^*$ is an isomorphism by [GZ22, case of orthogonal groups in Section 8] and Theorem 2.7.2.(1). The homomorphism $\tilde{\varphi}_K^*$ is surjective by Lemma 3.2.1 and is an isomorphism by the dimension count using the explicit presentation for \mathcal{P} and the computation of $\text{Ch}^*(G_K)$ recalled in (A.5.2). Then $\tilde{\varphi}^*$ is surjective. Furthermore, a simple diagram chase yields that for injectivity of $\tilde{\varphi}^*$ it suffices to check that if $\pi_*\tilde{\varphi}_K^*\tilde{\Theta}(\alpha) = 0$ for $\alpha \in \mathcal{P}/(\bar{\varepsilon}_1^{v_2(2r)})$, then $(\text{id}+\tau)(\alpha) = 0$. Consider the following commutative diagram:

$$\begin{array}{ccccc} \text{Ch}^*(G_K) & \xrightarrow{\pi_*} & \text{Ch}^*(G) & \xrightarrow{\pi^*} & \text{Ch}^*(G_K) \\ \tilde{\varphi}_K^* \circ \tilde{\Theta} \uparrow & & \tilde{\varphi}^* \circ \tilde{\Theta}^\tau \uparrow & & \tilde{\varphi}_K^* \circ \tilde{\Theta} \uparrow \\ \mathcal{P}/(\bar{\varepsilon}_1^{v_2(2r)}) & \xrightarrow{\text{id}+\tau} & \mathcal{P}^\tau/(\bar{\varepsilon}_1^{v_2(2r)}) & \xrightarrow{J} & \mathcal{P}/(\bar{\varepsilon}_1^{v_2(2r)}) \end{array}$$

Here J is induced by the inclusion $\mathcal{P}^\tau \subseteq \mathcal{P}$. We have

$$\tilde{\varphi}_K^* \circ \tilde{\Theta} \circ J \circ (\text{id}+\tau)(\alpha) = \pi^* \circ \pi_* \circ \tilde{\varphi}_K^* \circ \tilde{\Theta}(\alpha) = 0,$$

thus $J \circ (\text{id}+\tau)(\alpha) = 0$, since $\tilde{\varphi}_K^* \circ \tilde{\Theta}$ is an isomorphism. Then Lemma 3.1.2.(2) yields $(\text{id}+\tau)(\alpha) = 0$. Thus $\tilde{\varphi}^*$ is an isomorphism. The remaining claims of the Theorem follow from this, the isomorphism $\Theta^\tau: \mathcal{P}^\tau \xrightarrow{\cong} \text{Ch}^*(G/P)$ and Lemma 3.1.2.(3). \square

Corollary 3.2.3. *Let G be an adjoint quasi-split simple group over a field k of type $\Delta(G) = {}^2\text{D}_{2r}$, $r \geq 2$. Then $\text{Ch}^*(G)$ does not admit a structure of a Hopf algebra, in particular, the Künneth homomorphism*

$$\text{Ch}^*(G) \otimes \text{Ch}^*(G) \rightarrow \text{Ch}^*(G \times G), \quad x \otimes y \mapsto p_1^*(x) \cdot p_2^*(y),$$

fails to be an isomorphism.

Proof. It follows from Theorem 3.2.2.(2) that $\text{Ch}^*(G)$ admits a basis given by

$$\begin{aligned} e_1^{i_1} e_2^{i_2} \dots e_s^{i_s}, \quad & 0 \leq i_j < 2^{k_j}, 1 \leq j \leq s, j \neq l, 0 \leq i_l < 2^{k_l-1}, \\ e_2^{i_2} e_3^{i_3} \dots e_s^{i_s}, \quad & 0 \leq i_j \leq 2^{k_j}, 2 \leq j \leq s, j \neq l, 2^{k_l-1} \leq i_l < 2^{k_l}, \end{aligned}$$

in the notation of loc. cit. Then

$$\dim_{\mathbb{F}_2} \text{Ch}^*(G) = 2^{N-1} + 2^{N-1-k_1}, \quad N := \sum_{i=1}^s k_i.$$

Since $k_1 = v_2(2r) \geq 1$, the integer $2^{N-1} + 2^{N-1-k_1}$ is not a power of 2, and it could not be the dimension of a Hopf algebra by a result of Borel on the structure of finite dimensional connected Hopf algebras [MM65, Theorem 7.11 and Proposition 7.8]. \square

3.3 Chow ring of a quasi-split group at the splitting prime

In this section we show that if G is a non-split quasi-split simple group over a field k such that $\Delta(G) \neq {}^6\text{D}_4$ and $(\Delta(G), \pi_1(G)) \neq ({}^2\text{D}_{2r}, {}^2\mu_{2,2})$, then $\text{CH}^*(G) \otimes \mathbb{F}_p \cong \text{CH}_K^*(G)$, where K/k is the splitting field of G and $p = [K : k]$. This is done analysing the pushforward homomorphism $\pi_* : \text{CH}^*(G_K) \otimes \mathbb{F}_p \rightarrow \text{CH}^*(G) \otimes \mathbb{F}_p$, which we show to be trivial in this case.

We continue to use the notation of Sections A.2 and A.3. In this section we denote

$$\text{Ch}^*(-) := \text{CH}^*(-) \otimes \mathbb{F}_p$$

for some given prime number p .

Lemma 3.3.1. *Let G be a non-split quasi-split simple algebraic group over a field k with the splitting field K/k and let $\pi : G_K \rightarrow G$ be the projection. Then the following holds.*

1. *If $(\Delta(G), \pi_1(G)) \neq ({}^2\text{D}_{2r}, {}^2\mu_{2,2})$, then $\pi_* : \text{Pic}(G_K) \rightarrow \text{Pic}(G)$ is the zero homomorphism.*
2. *If $(\Delta(G), \pi_1(G)) = ({}^2\text{D}_{2r}, {}^2\mu_{2,2})$, then*

$$\pi_*([\mathcal{L}(\bar{\omega}_{2r-1})]) = \pi_*([\mathcal{L}(\bar{\omega}_{2r})]) = [\mathcal{L}(\bar{\omega}_1)], \quad \pi^*([\mathcal{L}(\bar{\omega}_1)]) = [\mathcal{L}(\bar{\omega}_{2r-1})] + [\mathcal{L}(\bar{\omega}_{2r})]$$

in $\text{Pic}(G)$ and $\text{Pic}(G_K)$ respectively.

Proof. Straightforward from the recollection in Section A.4. \square

Proposition 3.3.2. *Let G be a non-split quasi-split simple algebraic group over a field k with the splitting field K/k and let $\pi : G_K \rightarrow G$ be the projection. Suppose $\Delta(G) \neq {}^6\text{D}_4$ and $(\Delta(G), \pi_1(G)) \neq ({}^2\text{D}_{2r}, {}^2\mu_{2,2})$ and put $p := [K : k]$. Then $\pi_* : \text{Ch}^*(G_K) \rightarrow \text{Ch}^*(G)$ is the zero homomorphism.*

Proof. Let $T \leq B \leq G$ be a maximal torus and a Borel subgroup. We argue case by case on $\Delta(G)$.

²**A_n**. We have $p = 2$ and it follows from recollection (A.5.2) that the ring $\text{Ch}^*(G_K)$ is either trivial or $\text{Ch}^*(G_K) \cong \mathbb{F}_2[e_1]/(e_1^{2^{v_2(n+1)}})$ with $\deg e_1 = 1$. Consider the following commutative diagram.

$$\begin{array}{ccc} \text{Ch}^*((G/B)_K) & \xrightarrow{\rho_*} & \text{Ch}^*(G/B) \\ \phi_K^* \downarrow & & \downarrow \phi^* \\ \text{Ch}^*(G_K) & \xrightarrow{\pi_*} & \text{Ch}^*(G) \end{array}$$

Here $\phi : G \rightarrow G/B$, $\phi_K : G_K \rightarrow (G/B)_K$ and $\rho : (G/B)_K \rightarrow G/B$ are the projections. For the line bundle $\mathcal{L}(\varpi_1)$ over $(G/B)_K$ we have $\phi_K^*(c_1(\mathcal{L}(\varpi_1))) = e_1$ by recollection (A.4.2). Then for all $m \in \mathbb{N}$ we have

$$\pi_*(e_1^m) = \pi_*\phi_K^*(c_1(\mathcal{L}(\varpi_1))^m) = \phi^*\rho_*(c_1(\mathcal{L}(\varpi_1))^m).$$

Furthermore, for the non-trivial element $\tau \in \text{Gal}(K/k)$ we have

$$\begin{aligned} \rho^*\rho_*(c_1(\mathcal{L}(\varpi_1))^m) &= c_1(\mathcal{L}(\varpi_1))^m + \tau(c_1(\mathcal{L}(\varpi_1))^m) = c_1(\mathcal{L}(\varpi_1))^m + c_1(\mathcal{L}(\varpi_n))^m = \\ &= (c_1(\mathcal{L}(\varpi_1)) + c_1(\mathcal{L}(\varpi_n))) \cdot \left(\sum_{i=0}^{n-1} c_1(\mathcal{L}(\varpi_1))^i c_1(\mathcal{L}(\varpi_n))^{n-1-i} \right) \end{aligned}$$

with the last sum being $\text{Gal}(K/k)$ -invariant. Then Lemma 2.6.1 yields that

$$\rho_*(c_1(\mathcal{L}(\varpi_1))^m) = (c_1(\mathcal{L}(\varpi_1)) + c_1(\mathcal{L}(\varpi_n))) \cdot \alpha = \rho_*(c_1(\mathcal{L}(\varpi_1))) \cdot \alpha$$

for some $\alpha \in \text{Ch}^*(G/B)$. Thus

$$\pi_*(e_1^m) = \phi^* \rho_*(c_1(\mathcal{L}(\varpi_1))^m) = \phi^* \rho_*(c_1(\mathcal{L}(\varpi_1))) \cdot \phi^* \alpha = \pi_*(e_1) \cdot \phi^* \alpha = 0$$

with the last equality following from $\pi_*(e_1) = 0$ by Lemma 3.3.1.

²D_n. We have $p = 2$. Consider the parabolic subgroup $B \leq P := P_{n-1,n} \leq G$ in the notation of (A.2.6) and the following commutative diagram.

$$\begin{array}{ccc} \text{Ch}^*((G/P)_K) & \xrightarrow{\rho_*} & \text{Ch}^*(G/P) \\ \varphi_K^* \downarrow & & \downarrow \varphi^* \\ \text{Ch}^*(G_K) & \xrightarrow{\pi_*} & \text{Ch}^*(G) \end{array}$$

Here $\varphi: G \rightarrow G/P$, $\varphi_K: G_K \rightarrow (G/P)_K$ and $\rho: (G/P)_K \rightarrow G/P$ are the projections. The homomorphism φ_K^* is surjective by Lemma 3.2.1, thus

$$\pi_*(\text{Ch}^*(G_K)) = \varphi^* \rho_*(\text{Ch}^*(G/P)_K).$$

Lemma 2.6.1 combined with Proposition 3.1.1.(2) and Lemma 3.1.2.(1) yields

$$\rho_*(\text{Ch}^*(G/P)_K) = c_1(\mathcal{L}_P(\varpi_{n-1} + \varpi_n)) \cdot (\text{Ch}^*(G/P)).$$

Since $(\Delta(G), \pi_1(G)) \neq ({}^2D_{2r}, {}^2\mu_{2,2})$, it follows that $\bar{\varpi}_{n-1} + \bar{\varpi}_n = 0$ in $\mathcal{X}^*(\pi_1(G))$ and

$$\varphi^* c_1(\mathcal{L}_P(\varpi_{n-1} + \varpi_n)) = c_1(\varphi^* \mathcal{L}_P(\varpi_{n-1} + \varpi_n)) = c_1(\mathcal{L}(\bar{\varpi}_{n-1} + \bar{\varpi}_n)) = c_1(\mathcal{O}_G) = 0.$$

²E₆. We have $p = 2$. Consider the following commutative diagram.

$$\begin{array}{ccc} \text{Ch}^*((G/B)_K) & \xrightarrow{\rho_*} & \text{Ch}^*(G/B) \\ \phi_K^* \downarrow & & \downarrow \phi^* \\ \text{Ch}^*(G_K) & \xrightarrow{\pi_*} & \text{Ch}^*(G) \end{array}$$

Here $\phi: G \rightarrow G/B$, $\phi_K: G_K \rightarrow (G/B)_K$ and $\rho: (G/B)_K \rightarrow G/B$ are the canonical morphisms. By recollection (A.5.2) we have $\text{Ch}^*(G_K) \cong \mathbb{F}_2[e_1]/(e_1^2)$. A straightforward computer-assisted computation shows that for the Schubert cycle $Z_{s_3 s_4 s_2} \in \text{Ch}^3((G/B)_K)$ one has $\phi_K^*(Z_{s_3 s_4 s_2}) = e_1$, where s_i is the reflection for the simple root α_i . Then

$$\pi_*(e_1) = \pi_* \phi_K^*(Z_{s_3 s_4 s_2}) = \phi^* \rho_*(Z_{s_3 s_4 s_2}).$$

The Galois group $\text{Gal}(K/k)$ acts on $\text{Ch}^*((G/B)_K)$ permuting Schubert cycles accordingly to the permutation of simple roots, thus

$$\rho^* \rho_*(Z_{s_3 s_4 s_2}) = Z_{s_3 s_4 s_2} + Z_{s_5 s_4 s_2}.$$

Another straightforward computer-assisted computation shows that

$$c_1(\mathcal{L}(\varpi_2)_K)^3 = Z_{s_3 s_4 s_2} + Z_{s_5 s_4 s_2}.$$

Then the injectivity part of Lemma 2.6.1 yields

$$c_1(\mathcal{L}(\varpi_2))^3 = \rho_*(Z_{s_3 s_4 s_2}).$$

Since the fundamental weight ϖ_2 belongs to the root lattice, it follows that $\bar{\varpi}_2 = 0$ in $\mathcal{X}^*(\pi_1(G))$ and

$$\pi_*(e_1) = \phi^* \rho_*(Z_{s_3 s_4 s_2}) = \phi^* c_1(\mathcal{L}(\varpi_2))^3 = c_1(\mathcal{L}(\bar{\varpi}_2))^3 = 0.$$

³D₄. We have $p = 3$ and the claim is trivial, since $\text{Ch}^*(G_K) = \mathbb{F}_3$ by recollection (A.5.2). \square

Theorem 3.3.3. *Let G be a non-split quasi-split simple group over a field k with the splitting field K/k and suppose that $\Delta(G) \neq {}^6D_4$ and $(\Delta(G), \pi_1(G)) \neq ({}^2D_{2r}, 2\mu_{2,2})$. Then for $p = [K : k]$ there is an isomorphism*

$$\mathrm{Ch}^*(G) \cong \mathbb{F}_p[e_1, e_2, \dots, e_s] / (e_1^{p^{k_1}}, e_2^{p^{k_2}}, \dots, e_s^{p^{k_s}}), \quad \deg e_i = d_i,$$

with the parameters s, d_i, k_i given in the Table 1.

Proof. Proposition 3.3.2 yields that the quotient morphism $\mathrm{Ch}^*(G) \rightarrow \mathrm{CH}_K^*(G)$ is an isomorphism. Thus the claim follows from Theorem 2.8.2. \square

3.4 Chow ring of a quasi-split group away from the splitting primes

In this section we compute the Chow ring $\mathrm{Ch}^*(G) := \mathrm{CH}^*(G) \otimes \mathbb{F}_p$ for a non-split quasi-split simple group G over a field k assuming that p is coprime to $[K : k]$ where K is the splitting field of G . For this we show that for such p one has $\mathrm{Ch}^*(G) \cong \mathrm{Ch}^*(G_K)^{\mathrm{Gal}(K/k)}$ and then we compute the Galois action and the subalgebra of invariant elements.

We continue to use the notation of Sections A.2 and A.3. In this section we put

$$\mathrm{Ch}^*(-) := \mathrm{CH}^*(-) \otimes \mathbb{F}_p$$

for some given prime number p .

Lemma 3.4.1. *Let $X \in \mathcal{S}m_k$, K/k be a field extension of finite degree and p be a prime such that $p \nmid [K : k]$. Then for the projection $\pi: X_K \rightarrow X$ the pullback homomorphism*

$$\pi^*: \mathrm{Ch}^*(X) \rightarrow \mathrm{Ch}^*(X_K)$$

is injective.

Proof. This is well-known and follows from [Fu98, Example 1.7.4]. \square

Lemma 3.4.2. *Let G be a quasi-split group over a field k with the splitting field K/k and let $p \in \mathbb{N}$ be a prime such that $p \nmid [K : k]$. Then for the projection $\pi: G_K \rightarrow G$ the pullback homomorphism*

$$\pi^*: \mathrm{Ch}^*(G) \rightarrow \mathrm{Ch}^*(G_K)$$

is injective and $\pi^*(\mathrm{Ch}^*(G)) = (\mathrm{Ch}^*(G_K))^{\mathrm{Gal}(K/k)}$.

Proof. The homomorphism π^* is injective by Lemma 3.4.1. Furthermore, it is clear that

$$\pi^*(\mathrm{Ch}^*(G)) \subseteq (\mathrm{Ch}^*(G_K))^{\mathrm{Gal}(K/k)}.$$

Let $B \leq G$ be a Borel subgroup and consider the following commutative diagram.

$$\begin{array}{ccc} \mathrm{Ch}^*(G/B) & \xrightarrow{\rho^*} & \mathrm{Ch}^*((G/B)_K) \\ \phi^* \downarrow & & \downarrow \phi_K^* \\ \mathrm{Ch}^*(G) & \xrightarrow{\pi^*} & \mathrm{Ch}^*(G_K) \end{array}$$

Here $\phi: G \rightarrow G/B$, $\phi_K: G_K \rightarrow (G/B)_K$ and $\rho: (G/B)_K \rightarrow G/B$ are the projections. Since G_K is split, ϕ_K^* is surjective by Theorem 2.5.6. Let $\alpha \in (\mathrm{Ch}^*(G_K))^{\mathrm{Gal}(K/k)}$ and pick some $\beta \in \mathrm{Ch}^*((G/B)_K)$ such that $\phi_K^*(\beta) = \alpha$. Then the element

$$\tilde{\beta} := \frac{1}{[K : k]} \sum_{\sigma \in \mathrm{Gal}(K/k)} \sigma(\beta) \in \mathrm{Ch}^*((G/B)_K)$$

is $\mathrm{Gal}(K/k)$ -invariant and Lemma 2.6.1 yields that there exists $\gamma \in \mathrm{Ch}^*(G/B)$ such that $\rho^*(\gamma) = \tilde{\beta}$. Then

$$\pi^*(\phi^*(\gamma)) = \phi_K^*(\rho^*(\gamma)) = \frac{1}{[K : k]} \sum_{\sigma \in \mathrm{Gal}(K/k)} \sigma(\alpha) = \alpha.$$

It follows that $\pi^*(\mathrm{Ch}^*(G)) = (\mathrm{Ch}^*(G_K))^{\mathrm{Gal}(K/k)}$ yielding the claim. \square

Theorem 3.4.3. *Let G be a non-split quasi-split simple group over a field k with the splitting field K/k and let $p \in \mathbb{N}$ be a prime such that $p \nmid [K : k]$. If $(\Delta(G_K), \pi_1(G_K), p)$ is not in Table 6, then $\text{Ch}^*(G) \cong \mathbb{F}_p$, otherwise $\text{Ch}^*(G)$ is given by the following table.*

Table 4: $\text{Ch}^*(G)$ away from the splitting primes

$\Delta(G)$	$\pi_1(G)$	p	$\text{Ch}^*(G)$	$\deg e_i$
${}^2\text{A}_n$	$2\mu_l, l \mid n+1$	$2 \neq p \mid l$	$\mathbb{F}_p[e_1]/(e_1^m), m = \frac{p^{v_p(n+1)}+1}{2}$	2
${}^3\text{D}_4$	1	2	$\mathbb{F}_2[e_1]/(e_1^2)$	3
	${}^3\mu_{2,2}$	2	$\mathbb{F}_2[e_1, e_2, e_3, e_4]/(e_1^4, e_2^2, e_3^2, e_4^2, e_1e_2, e_1e_3, e_1^3 + e_2e_3)$	2, 3, 3, 3
${}^2\text{E}_6$	1	3	$\mathbb{F}_3[e_1]/(e_1^3)$	4
	${}^2\mu_3$	3	$\mathbb{F}_3[e_1, e_2]/(e_1^5, e_2^3)$	1, 4

Here v_p is the p -adic valuation.

Proof. The claim follows from Lemma 3.4.2, the computation of $\text{Ch}^*(G_K)$ recalled in (A.5.2) and the description of the action of $\text{Gal}(K/k)$ on $\text{Ch}^*(G_K)$.

If $(\Delta(G_K), \pi_1(G_K), p)$ is not in Table 6, then $\text{Ch}^*(G_K) \cong \mathbb{F}_p$ and the claim follows. Otherwise, since $p \nmid [K : k]$, one can easily check that the triple $(\Delta(G_K), \pi_1(G_K), p)$ is in Table 4, and we study them case by case.

$\Delta(G) = {}^2\text{A}_n$. We have

$$\text{Ch}^*(G_K) \cong \mathbb{F}_p[e]/(e^{p^{v_p(n+1)}}), \quad \deg e = 1.$$

In view of recollection (A.4.2) we can assume $e = c_1(\mathcal{L}(\bar{\omega}_1))$. Thus for the nontrivial element $\tau \in \text{Gal}(K/k)$ we have

$$\tau(e) = \tau(c_1(\mathcal{L}(\bar{\omega}_1))) = c_1(\mathcal{L}(\bar{\omega}_n)) = -e.$$

Hence $\text{Ch}^*(G_K)^{\text{Gal}(K/k)} \subseteq \text{Ch}^*(G_K)$ is the subalgebra generated by e^2 and the claim follows.

$(\Delta(G), \pi_1(G)) = ({}^3\text{D}_4, 1)$. We have

$$\text{Ch}^*(G_K) \cong \mathbb{F}_2[e]/(e^2), \quad \deg e = 3.$$

The vector space $\mathbb{F}_2 \cong \text{Ch}^3(G_K)$ admits only the trivial action of $\text{Gal}(K/k)$, thus e is invariant and

$$\text{Ch}^*(G) \cong \text{Ch}^*(G_K)^{\text{Gal}(K/k)} \cong \mathbb{F}_2[e]/(e^2).$$

$(\Delta(G), \pi_1(G)) = ({}^3\text{D}_4, {}^3\mu_{2,2})$. We have

$$\text{Ch}^*(G_K) \cong \mathbb{F}_2[e_1, e_2, e_3]/(e_1^4, e_2^4, e_3^2), \quad \deg e_1 = \deg e_2 = 1, \deg e_3 = 3.$$

In view of recollection (A.4.2) we can assume $e_1 = c_1(\mathcal{L}(\bar{\omega}_3))$ and $e_2 = c_1(\mathcal{L}(\bar{\omega}_4))$. For a generator $\sigma \in \text{Gal}(K/k) \cong C_3$ we have

$$\sigma(e_1) = e_2, \quad \sigma(e_2) = e_1 + e_2.$$

Let $\mathcal{A}^* \subseteq \text{Ch}^*(G_K)$ be the subalgebra generated by e_1, e_2 . Then it is straightforward to check that $(\mathcal{A}^*)^{\text{Gal}(K/k)}$ has a basis given by

$$1, \quad e_1^2 + e_1e_2 + e_2^2, \quad e_1^2e_2 + e_1e_2^2, \quad e_1^3 + e_1^2e_2 + e_2^3, \quad e_1^2e_2^2, \quad e_1^3e_2^3.$$

We have $\mathcal{A}^3 \subseteq \text{Ch}^3(G_K)$ a codimension one $\text{Gal}(K/k)$ -invariant subspace. Maschke's theorem yields that

$$\text{Ch}^3(G_K) = \mathcal{A}^3 \oplus \mathbb{F}_2\bar{e}_3$$

for some $\tilde{e}_3 \in \text{Ch}^3(G_K)$ with $\sigma(\tilde{e}_3) = \tilde{e}_3$. We have $\tilde{e}_3 = e_3 + p(e_1, e_2)$ for some degree 3 homogeneous polynomial. Then we obtain a $\text{Gal}(K/k)$ -invariant decomposition

$$\text{Ch}^*(G_K) = \mathcal{A}^* \oplus \mathcal{A}^* \tilde{e}_3,$$

and it follows that $\text{Ch}^*(G_K)^{\text{Gal}(K/k)}$ has a basis given by

$$\begin{aligned} &1, \quad e_1^2 + e_1 e_2 + e_2^2, \quad e_1^2 e_2 + e_1 e_2^2, \quad e_1^3 + e_1^2 e_2 + e_2^3, \quad e_1^2 e_2^2, \quad e_1^3 e_2^3, \\ &\tilde{e}_3, \quad (e_1^2 + e_1 e_2 + e_2^2) \tilde{e}_3, \quad (e_1^2 e_2 + e_1 e_2^2) \tilde{e}_3, \quad (e_1^3 + e_1^2 e_2 + e_2^3) \tilde{e}_3, \quad e_1^2 e_2^2 \tilde{e}_3, \quad e_1^3 e_2^3 \tilde{e}_3. \end{aligned}$$

Thus $\text{Ch}^*(G_K)^{\text{Gal}(K/k)}$ is generated as an algebra by $e_1^2 + e_1 e_2 + e_2^2$, $e_1^2 e_2 + e_1 e_2^2$, $e_1^3 + e_1^2 e_2 + e_2^3$ and \tilde{e}_3 , and the multiplicative relations are straightforward.

$(\Delta(G), \pi_1(G)) = ({}^2\text{E}_6, 1)$. We have

$$\text{Ch}^*(G_K) \cong \mathbb{F}_3[e]/(e^3), \quad \deg e = 4.$$

Let $T \leq B \leq G$ be a maximal torus and a Borel subgroup and consider the characteristic sequence

$$\text{Ch}_{T_K}^*(\text{Spec } K) \xrightarrow{c} \text{Ch}^*(G_K/B_K) \xrightarrow{\phi_K^*} \text{Ch}^*(G_K) \rightarrow 0.$$

A straightforward computer-assisted computation shows that for the Schubert cycle

$$Z_{s_2 s_4 s_3 s_1} \in \text{Ch}^4(G_K/B_K)$$

its image $\phi_K^*(Z_{s_2 s_4 s_3 s_1})$ is a generator of $\text{Ch}^4(G_K)$. We have $\tau(Z_{s_2 s_4 s_3 s_1}) = Z_{s_2 s_4 s_5 s_6}$ for the nontrivial element $\tau \in \text{Gal}(K/k)$, and a further computer-assisted computation shows that $Z_{s_2 s_4 s_3 s_1} - Z_{s_2 s_4 s_5 s_6}$ lies in $c(\text{Ch}_{T_K}^*(\text{Spec } K))$. Then

$$\tau(\phi_K^*(Z_{s_2 s_4 s_3 s_1})) = \phi_K^*(\tau(Z_{s_2 s_4 s_3 s_1})) = \phi_K^*(Z_{s_2 s_4 s_5 s_6}) = \phi_K^*(Z_{s_2 s_4 s_3 s_1}),$$

so $\text{Gal}(K/k)$ acts trivially on $\text{Ch}^4(G_K)$ and

$$\text{Ch}^*(G) \cong \text{Ch}^*(G_K)^{\text{Gal}(K/k)} \cong \mathbb{F}_3[e]/(e^3), \quad \deg e = 4.$$

$(\Delta(G), \pi_1(G)) = ({}^2\text{E}_6, 2\mu_3)$. We have

$$\text{Ch}^*(G_K) \cong \mathbb{F}_3[e_1, e_2]/(e_1^9, e_2^3), \quad \deg e_1 = 1, \deg e_2 = 4.$$

In view of recollection (A.4.2) we can assume $e_1 = c_1(\mathcal{L}(\bar{\omega}_1))$, thus for the nontrivial element $\tau \in \text{Gal}(K/k)$ we have

$$\tau(e_1) = \tau(c_1(\mathcal{L}(\bar{\omega}_1))) = c_1(\mathcal{L}(\bar{\omega}_6)) = c_1(\mathcal{L}(-\bar{\omega}_1)) = -e_1.$$

We have $\text{Ch}^4(G_K) = \mathbb{F}_3 e_1^4 \oplus \mathbb{F}_3 e_2$, and it follows from the above that $\tau(e_1^4) = e_1^4$. Furthermore, the simply connected case yields $\tau(e_2) = e_2 + a e_1^4$ for some $a \in \mathbb{F}_3$. Since

$$e_2 = \tau^2(e_2) = \tau(e_2 + a e_1^4) = e_2 + 2a e_1^4,$$

we have $a = 0$ and $\tau(e_2) = e_2$. Thus $\text{Ch}^*(G_K)^{\text{Gal}(K/k)} \subseteq \text{Ch}^*(G_K)$ is the subalgebra generated by e_1^2 and e_2 . The claim follows. \square

3.5 Chow ring of the group of type ${}^6\text{D}_4$ at $p = 2, 3$

In this section we compute the ring $\text{Ch}^*(G) := \text{CH}^*(G) \otimes \mathbb{F}_p$ in the last remaining cases, namely $\Delta(G) = {}^6\text{D}_4$ and $p = 2, 3$. This is done partially splitting G to G_L with $\Delta(G_L) = {}^3\text{D}_4$ or $\Delta(G_L) = {}^2\text{D}_4$, and analyzing the Galois action on $\text{Ch}^*(G_K)$, where K is the splitting field of G .

We continue to use the notation of Sections A.2 and A.3. In this section we put

$$\text{Ch}^*(-) := \text{CH}^*(-) \otimes \mathbb{F}_p$$

for $p = 2$ or $p = 3$.

Theorem 3.5.1. *Let G be a quasi-split simple group over a field k of type $\Delta(G) = {}^6D_4$. Then for $p = 2, 3$ the algebra $\text{Ch}^*(G)$ is given by the following table.*

$\Delta(G)$	$\pi_1(G)$	p	$\text{Ch}^*(G)$	$\deg e_i$
6D_4	1	2	$\mathbb{F}_2[e_1]/(e_1^2)$	3
	${}^6\mu_{2,2}$	2	$\mathbb{F}_2[e_1, e_2, e_3]/(e_1^4, e_2^2, e_1e_2, e_3^2)$	2, 3, 3
	1, ${}^6\mu_{2,2}$	3	$\mathbb{F}_3[e_1]/(e_1^3)$	4

Proof. Let K/k be the splitting field of G and $T \leq B \leq G$ be a maximal torus and a Borel subgroup. We argue case by case on $(\pi_1(G), p)$.

$(\pi_1(G), p) = ({}^6\mu_{2,2}, 2)$. Let $K/L/k$ be an intermediate extension with $[L : k] = 3$ such that $\text{Gal}(K/L)$ stabilizes the root α_1 on the Dynkin diagram of G_K , then $\Delta(G_L) = {}^2D_4$ with the nontrivial element of $\text{Gal}(K/L)$ permuting the roots α_3 and α_4 . Consider the parabolic subgroup $B \leq P := P_{3,4} \leq G_L$ in the notation of (A.2.6) and consider the following commutative diagram.

$$\begin{array}{ccccc}
& & \text{Ch}^*(G_L/P) & \xrightarrow{g^*} & \text{Ch}^*(G_K/P_K) \\
& & \downarrow \psi_L^* & & \downarrow \psi_K^* \\
\text{Ch}^*(G/B) & \xrightarrow{\rho_{L/k}^*} & \text{Ch}^*(G_L/B_L) & \xrightarrow{\rho_{K/L}^*} & \text{Ch}^*(G_K/B_K) \\
\downarrow \phi^* & & \downarrow \phi_L^* & & \downarrow \phi_K^* \\
\text{Ch}^*(G) & \xrightarrow{\pi_{L/k}^*} & \text{Ch}^*(G_L) & \xrightarrow{\pi_{K/L}^*} & \text{Ch}^*(G_K)
\end{array}$$

Here all the morphisms are pullbacks for the respective projections. We have the following:

- $g^*, \rho_{L/k}^*, \rho_{K/L}^*$ are injective by Lemma 2.6.1,
- $\pi_{L/k}^*$ is injective by Lemma 3.4.1,
- $\phi_L^* \circ \psi_L^*$ is an isomorphism by Theorem 3.2.2.(1),
- $\phi_K^* \circ \psi_K^*$ is an isomorphism, this is well-known, see e.g. the beginning of the proof of Theorem 3.2.2.

It follows that $\pi_{K/L}^* = \phi_K^* \circ \psi_K^* \circ g^* \circ (\phi_L^* \circ \psi_L^*)^{-1}$ is injective, so the composition

$$\pi_{K/k}^* := \pi_{K/L}^* \circ \pi_{L/k}^* : \text{Ch}^*(G) \rightarrow \text{Ch}^*(G_K)$$

is injective. It is clear that $\pi_{K/k}^*(\text{Ch}^*(G)) \subseteq \text{Ch}^*(G_K)^{\text{Gal}(K/k)}$, and we claim that this is in fact an equality. Indeed, pick $x \in \text{Ch}^*(G_K)^{\text{Gal}(K/k)}$ and consider

$$y' := \psi_K^* \circ (\phi_K^* \circ \psi_K^*)^{-1}(x), \quad y := y' + \sigma(y') + \sigma^2(y'),$$

where $\sigma \in \text{Gal}(K/k) \cong S_3$ is an element of order 3. It is clear that $\sigma(y) = y$. For the nontrivial element $\tau \in \text{Gal}(K/L) \subseteq \text{Gal}(K/k)$ we have $\tau(y') = y'$, since $\tau(x) = x$ by the assumption and ψ_K and ϕ_K are obtained by a base change from $\text{Spec } L$. Then

$$\tau(y) = \tau(y') + \tau\sigma(y') + \tau\sigma^2(y') = \tau(y') + \sigma^2\tau(y') + \sigma\tau(y') = y' + \sigma^2(y') + \sigma(y') = y,$$

and it follows that $y \in \text{Ch}^*(G_K/B_K)^{\text{Gal}(K/k)}$. Lemma 2.6.1 yields that there exists $z \in \text{Ch}^*(G/B)$ such that

$$(\rho_{L/k} \circ \rho_{K/L})^*(z) = y.$$

Therefore, we have

$$\pi_{K/k}^* \phi^*(z) = \phi_K^*(y) = \phi_K^*(y' + \sigma(y') + \sigma^2(y')) = \phi_K^*(y') + \sigma \phi_K^*(y') + \sigma^2 \phi_K^*(y') = 3x = x,$$

with the third equality following from the fact that ϕ_K is obtained by a base change from $\text{Spec } k$. Then

$$\pi_{K/k}^*(\text{Ch}^*(G)) = \text{Ch}^*(G_K)^{\text{Gal}(K/k)}$$

as claimed, and it remains to compute the algebra of invariants.

We have an isomorphism

$$\Theta: \text{Ch}^*(G_K) \xrightarrow{\cong} \mathbb{F}_2[e_1, e_2, e_3]/(e_1^4, e_2^4, e_3^2), \quad \deg e_1 = \deg e_2 = 1, \quad \deg e_3 = 3,$$

by recollection (A.5.2). A straightforward computer-assisted computation with the characteristic sequence

$$\text{Ch}_{T_K}^*(\text{Spec } K) \rightarrow \text{Ch}^*(G_K/B_K) \rightarrow \text{Ch}^*(G_K) \rightarrow 0$$

yields that there is a choice of the isomorphism Θ such that

$$\begin{aligned} \tau(e_1) &= e_2, & \tau(e_2) &= e_1, & \sigma(e_1) &= e_2, & \sigma(e_2) &= e_1 + e_2, \\ \tau(e_3) &= e_3 + e_1^2 e_2 + e_1 e_2^2, & \sigma(e_3) &= e_3 + e_1^2 e_2 + e_1 e_2^2 + e_3^3 \end{aligned}$$

for order 2 and order 3 elements $\tau, \sigma \in \text{Gal}(K/k) \cong S_3$. As in the proof of Theorem 3.4.3.(3), a basis of the σ -invariant subalgebra of $\text{Ch}^*(G_K)$ is given by

$$\begin{aligned} &1, \quad e_1^2 + e_1 e_2 + e_2^2, \quad e_1^2 e_2 + e_1 e_2^2, \quad e_1^3 + e_1^2 e_2 + e_2^3, \quad e_1^2 e_2^2, \quad e_1^3 e_2^3, \\ \tilde{e}_3, & \quad (e_1^2 + e_1 e_2 + e_2^2) \tilde{e}_3, \quad (e_1^2 e_2 + e_1 e_2^2) \tilde{e}_3, \quad (e_1^3 + e_1^2 e_2 + e_2^3) \tilde{e}_3, \quad e_1^2 e_2^2 \tilde{e}_3, \quad e_1^3 e_2^3 \tilde{e}_3. \end{aligned} \quad (3.5.1.1)$$

for some $\tilde{e}_3 = e_3 + p(e_1, e_2)$ with p being a homogeneous polynomial of degree 3. Using the above description for the action we see that one can choose

$$\tilde{e}_3 := e_3 + e_1 e_2^2.$$

It turns out that all the elements of the basis (3.5.1.1) except $e_1^3 + e_1^2 e_2 + e_2^3$ and $(e_1^3 + e_1^2 e_2 + e_2^3) \tilde{e}_3$ are also τ -invariant, while

$$\begin{aligned} \tau(e_1^3 + e_1^2 e_2 + e_2^3) &= (e_1^3 + e_1^2 e_2 + e_2^3) + (e_1^2 e_2 + e_1 e_2^2), \\ \tau((e_1^3 + e_1^2 e_2 + e_2^3) \tilde{e}_3) &= (e_1^3 + e_1^2 e_2 + e_2^3) \tilde{e}_3 + (e_1^2 e_2 + e_1 e_2^2) \tilde{e}_3. \end{aligned}$$

It follows that the remaining 10 elements form a basis of $\text{Ch}^*(G_K)^{\text{Gal}(K/k)}$. The claim of the theorem follows with $e_1^2 + e_1 e_2 + e_2^2$, $e_1^2 e_2 + e_1 e_2^2$ and $e_3 + e_1 e_2^2$ being the algebra generators of $\text{Ch}^*(G_K)^{\text{Gal}(K/k)} \cong \text{Ch}^*(G)$.

$(\pi_1(G), p) = (1, 2)$. Let $K/L/k$ be an intermediate extension with $[L : k] = 3$ and $f: G \rightarrow \bar{G}$ be the adjoint quotient. Consider the following commutative diagram.

$$\begin{array}{ccccc} \text{Ch}^*(\bar{G}) & \xrightarrow{\bar{\pi}_{L/k}^*} & \text{Ch}^*(\bar{G}_L) & \xrightarrow{\bar{\pi}_{K/L}^*} & \text{Ch}^*(\bar{G}_K) \\ \downarrow f^* & & \downarrow f_L^* & & \downarrow f_K^* \\ \text{Ch}^*(G) & \xrightarrow{\pi_{L/k}^*} & \text{Ch}^*(G_L) & \xrightarrow{\pi_{K/L}^*} & \text{Ch}^*(G_K) \end{array}$$

Here all the morphisms are pullbacks for the respective projections. By recollection (A.5.2) we have

$$\text{Ch}^*(G_K) \cong \mathbb{F}_2[e]/(e^2), \quad \text{Ch}^*(\bar{G}_K) \cong \mathbb{F}_2[e_1, e_2, e_3]/(e_1^4, e_2^4, e_3^2),$$

where for the second isomorphism we choose the same one as above in the proof. We have $f_K^*(e_1) = f_K^*(e_2) = 0$ by Lemma A.4.3. Furthermore, f_K^* is surjective by Theorem 2.5.6, since $G_K/B_K \cong \bar{G}_K/\bar{B}_K$

for the respective Borel subgroups, and thus $f_K^*(e_3) = e$. Then by the adjoint case discussed above there exists an element $\tilde{e} \in \text{Ch}^*(\bar{G})$ such that

$$f_K^* \bar{\pi}_{K/L}^* \bar{\pi}_{L/k}^*(\tilde{e}) = f_K^*(e_3 + e_1 e_2^2) = e.$$

Hence the composition $f_K^* \bar{\pi}_{K/L}^* \bar{\pi}_{L/k}^* = \pi_{K/L}^* \pi_{L/k}^* f^*$ is surjective, and it follows that $\pi_{K/L}^* \pi_{L/k}^*$ is surjective as well. Moreover, the homomorphism $\pi_{L/k}^*$ is injective by Lemma 3.4.1, and $\text{Ch}^*(G_L) \cong \text{Ch}^*(G_K)$ by Theorem 3.3.3 and recollection (A.5.2), thus $\pi_{K/L}^*$ and $\pi_{L/k}^*$ are isomorphisms. The claim follows.

$(\pi_1(G), p) = ({}^6\mu_{2,2}, 3)$. Let $K/L/k$ be the intermediate Galois extension with $[L : k] = 2$. Then $\Delta(G_L) = {}^3D_4$. Consider the following commutative diagram.

$$\begin{array}{ccccc} & & \text{Ch}^*(G_L/B_L) & \xrightarrow{\phi_L^*} & \text{Ch}^*(G_L) \\ & & \downarrow q_2 & & \downarrow q_1 \\ \text{CH}_{K,T_L}^*(\text{Spec } L) & \longrightarrow & \text{CH}_K^*(G_L/B_L) & \xrightarrow{\tilde{\phi}_L^*} & \text{CH}_K^*(G_L) \end{array}$$

Here the bottom row is the conormed characteristic sequence from Definition 2.5.5, G_L and G_L/B_L are considered as varieties over L , the vertical homomorphisms are the quotient morphisms, and $\phi_L^*, \tilde{\phi}_L^*$ are the pullbacks for the projection $\phi_L: G_L \rightarrow G_L/B_L$. The bottom sequence is exact in the sense of Definition 2.5.4. The algebra $\text{CH}_{K,T_L}^*(\text{Spec } L)$ was computed in Proposition 2.3.2 and $\text{CH}_K^*(G_L/B_L)$ is an explicit quotient of $\text{Ch}^*(G_L/B_L)$ which in turn can be identified by Lemma 2.6.1 with $\text{Ch}^*(G_K/B_K)^{\text{Gal}(K/L)}$ and admits a description via Schubert cycles. Let $\rho_{K/L}: G_K/B_K \rightarrow G_L/B_L$ be the projection. We have $\text{CH}_K^*(G_L) \cong \mathbb{F}_3[e]/(e^3)$ by Theorem 2.8.2 and a straightforward computer-assisted computation shows that for the cycle $Z \in \text{Ch}^*(G_L/B_L)$ such that $\rho_{K/L}^*(Z) = Z_{s_2 s_1 s_3 s_4}$ the element $\tilde{\phi}_L^*(Z)$ is a generator of $\text{CH}_K^4(G_L) \cong \mathbb{F}_3$. The homomorphism q_1 is an isomorphism

$$q_1: \text{Ch}^*(G_L) \xrightarrow{\cong} \text{CH}_K^*(G_L) \cong \mathbb{F}_3[e]/(e^3)$$

by Proposition 3.3.2. It follows that $\phi_L^*(Z)$ is a generator of $\text{Ch}^4(G_L) \cong \mathbb{F}_3$.

Consider the following diagram.

$$\begin{array}{ccc} \text{Ch}^*(G/B) & \xrightarrow{\phi^*} & \text{Ch}^*(G) \\ \downarrow \rho_{L/k}^* & & \downarrow \pi^* \\ \text{Ch}^*(G_L/B_L) & \xrightarrow{\phi_L^*} & \text{Ch}^*(G_L) \\ \downarrow \rho_{K/L}^* & & \\ \text{Ch}^*(G_K/B_K) & & \end{array}$$

Here all the morphisms are pullbacks for the respective projections. Since $\rho_{K/L}^*(Z) = Z_{s_2 s_1 s_3 s_4}$ is clearly $\text{Gal}(K/k)$ -invariant (the group permutes $\alpha_1, \alpha_3, \alpha_4$ and s_1, s_3, s_4 commute with each other), Lemma 2.6.1 yields that there exists $Z' \in \text{Ch}^4(G/B)$ such that $\rho_{L/k}^*(Z') = Z$. Then $\pi^* \phi^*(Z')$ is a generator of $\text{Ch}_K^4(G_L)$ and it follows that π^* is surjective. At the same time π^* is injective by Lemma 3.4.1, thus it is an isomorphism. The claim follows.

$(\pi_1(G), p) = (1, 3)$. Let $K/L/k$ be the intermediate Galois extension with $[L : k] = 2$, then $\Delta(G_L) = {}^3D_4$. Let $f: G \rightarrow \bar{G}$ be the adjoint quotient and consider the following diagram.

$$\begin{array}{ccc} \text{Ch}^*(\bar{G}) & \xrightarrow{\bar{\pi}^*} & \text{Ch}^*(\bar{G}_L) \\ f^* \downarrow & & \downarrow f_L^* \\ \text{Ch}^*(G) & \xrightarrow{\pi^*} & \text{Ch}^*(G_L) \end{array}$$

Here all the morphisms are pullbacks for the respective projections. The morphism f_L^* is surjective by a combination of Proposition 3.3.2 and Theorem 2.5.6, thus an isomorphism by the dimension count and Theorem 3.3.3. The morphism $\bar{\pi}^*$ is an isomorphism by the proof in the adjoint case above, and π^* is injective by Lemma 3.4.1. Thus f^* is an isomorphism yielding the claim. \square

A Recollection on algebraic groups

A.1 Some groups of multiplicative type and their representations

A.1.1. An affine algebraic group S over a field k is of *multiplicative type*, if $S_{k^{sep}}$ is diagonalizable [Mi17, Chapter 12]. Algebraic tori, their subgroups and quotient groups are of multiplicative type. Recall [Mi17, Theorem 12.23] that the category of groups of multiplicative type over a field k is dual to the category of finitely generated \mathbb{Z} -modules equipped with a continuous action of $\text{Gal}(k^{sep}/k)$ with the contravariant equivalence given by

$$S \mapsto \mathcal{X}^*(S) := \text{Hom}(S_{k^{sep}}, \mathbb{G}_m).$$

A.1.2. A *quasi-trivial* (or *induced*) torus T over a field k is the torus corresponding to a permutation $\text{Gal}(k^{sep}/k)$ -module. One can show [Mi17, Lemma 12.61 and below] that a quasi-trivial torus is a torus T admitting an isomorphism

$$T \cong R_{L_1/k} \mathbb{G}_m \times \dots \times R_{L_r/k} \mathbb{G}_m$$

for some finite separable field extensions $L_1, \dots, L_r/k$, where $R_{L_i/k} \mathbb{G}_m$, $1 \leq i \leq r$, are the respective Weil restrictions of a one-dimensional split torus.

Let L/k be a finite separable extension of fields and let $R := R_{L/k} \mathbb{G}_m$ be the Weil restriction of a one-dimensional split torus. Put $V_R := R_{L/k} \mathbb{A}_k^1$. Since Weil restrictions commute with products, the standard action $\mathbb{G}_m \times \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^1$ gives rise to the action

$$R \times V_R \cong R_{L/k}(\mathbb{G}_m \times \mathbb{A}_k^1) \rightarrow R_{L/k} \mathbb{A}_k^1 = V_R.$$

Recall that $V_R = R_{L/k} \mathbb{A}_k^1 \cong \mathbb{A}_k^n$ with $n = [L : k]$, and it is straightforward to see that the above action is linear. We refer to it as the *standard vector representation of R* . It is well-known that R can be identified with the open subset of V_R given by the condition $N_{L/k} \neq 0$, where $N_{L/k}$ is the norm form of the field extension L/k . We briefly recall some details. Let \bar{L}/k be a Galois closure of L/k , which also coincides with the splitting field of R . A choice of a basis $\{e_1, e_2, \dots, e_n\}$ for L over k gives rise to an isomorphism $V_R \cong \mathbb{A}_k^n = \text{Spec } k[x_1, x_2, \dots, x_n]$ and under this isomorphism the norm form can be described as

$$N_{L/k}(x_1, x_2, \dots, x_n) = \prod_{\sigma \in \mathcal{G}/\mathcal{G}'} \sigma(x_1 e_1 + \dots + x_n e_n) \in k[x_1, \dots, x_n] \subseteq \bar{L}[x_1, \dots, x_n],$$

where $\mathcal{G} = \text{Gal}(\bar{L}/k)$, $\mathcal{G}' = \text{Gal}(\bar{L}/L)$, and the product is taken over some chosen set of representatives of the cosets \mathcal{G}/\mathcal{G}' . Note that the formula does not depend on the choice of the set of representatives of \mathcal{G}/\mathcal{G}' , since $e_i \in L$, so all e_i are stable under the action of \mathcal{G}' .

A.1.3. In the article we repeatedly use the following groups of multiplicative type over a field k associated with a degree 2 Galois field extension L/k . These groups are naturally closed subgroups of $R := R_{L/k} \mathbb{G}_m$. Recall that there is a canonical isomorphism $\mathcal{X}^*(R) \cong \mathbb{Z} \oplus \mathbb{Z}$ with the nontrivial element $\tau \in \text{Gal}(L/k)$ permuting the elements of the basis, $\tau(x_1, x_2) = (x_2, x_1)$.

- ${}^2\mathbb{G}_m$ with $\mathcal{X}^*({}^2\mathbb{G}_m) = \mathbb{Z}$, $\tau(x) = -x$. The closed embedding ${}^2\mathbb{G}_m \leq R$ corresponds to the projection $\mathcal{X}^*(R) = \mathbb{Z} \oplus \mathbb{Z} \rightarrow \mathbb{Z} = \mathcal{X}^*({}^2\mathbb{G}_m)$, $(x_1, x_2) \mapsto x_1 - x_2$. The group ${}^2\mathbb{G}_m$ is a unique non-trivial twisted form of \mathbb{G}_m split by L .
- ${}^2\mu_l$ with $\mathcal{X}^*({}^2\mu_l) = \mathbb{Z}/l\mathbb{Z}$, $\tau(\bar{x}) = -\bar{x}$. The closed embedding ${}^2\mu_l \leq R$ corresponds to the projection $\mathcal{X}^*(R) = \mathbb{Z} \oplus \mathbb{Z} \rightarrow \mathbb{Z}/l\mathbb{Z} = \mathcal{X}^*({}^2\mu_l)$, $(x_1, x_2) \mapsto \bar{x}_1 - \bar{x}_2$. If $l \geq 3$, then the group ${}^2\mu_l$ is a unique non-trivial twisted form of the group of l -th roots of unity μ_l split by L . If $l = 2$, then ${}^2\mu_2 = \mu_2$.

- ${}^2\mu_{2,2}$ with $\mathcal{X}^*({}^2\mu_{2,2}) = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$, $\tau(\bar{x}_1, \bar{x}_2) = (\bar{x}_2, \bar{x}_1)$. The closed embedding ${}^2\mu_{2,2} \leq R$ corresponds to the projection $\mathcal{X}^*(R) = \mathbb{Z} \oplus \mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} = \mathcal{X}^*({}^2\mu_{2,2})$, $(x_1, x_2) \mapsto (\bar{x}_1, \bar{x}_2)$. The group ${}^2\mu_{2,2}$ is a unique non-trivial twisted form of the group $\mu_2 \times \mu_2$ split by L . Moreover, one has ${}^2\mu_{2,2} \cong R_{L/k}\mu_2$ for the Weil restriction $R_{L/k}\mu_2$ of μ_2 .

Restricting the standard vector representation V_R to the respective subgroups we obtain the following *standard vector representations* of dimension 2:

$${}^2V_{\mathbb{G}_m} := V_R|_{({}^2\mathbb{G}_m)}, \quad {}^2V_{\mu_l} := V_R|_{({}^2\mu_l)}, \quad {}^2V_{\mu_{2,2}} := V_R|_{({}^2\mu_{2,2})}.$$

Furthermore, if $l = 2m$ is even, then ${}^2\mu_l$ admits a linear *sign representation* ${}^2\Lambda_{\mu_{2m}}^\pm$ given by the Galois-invariant character $m \in \mathbb{Z}/2m\mathbb{Z} = \mathcal{X}^*({}^2\mu_{2m})$.

We also have the following groups associated with a degree 3 Galois field extension L/k and to a degree 6 Galois field extension with $\text{Gal}(L/k) \cong S_3$.

- ${}^3\mu_{2,2}$ with $\mathcal{X}^*({}^3\mu_{2,2}) = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ and $\text{Gal}(L/k) \cong C_3$ permuting the non-zero elements. The group ${}^3\mu_{2,2}$ is a unique non-trivial twisted form of the group $\mu_2 \times \mu_2$ split by L .
- ${}^6\mu_{2,2}$ with $\mathcal{X}^*({}^6\mu_{2,2}) = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ and $\text{Gal}(L/k) \cong S_3 \cong \text{Aut}(\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z})$ acting as the whole group of automorphisms. The group ${}^6\mu_{2,2}$ is a unique non-trivial twisted form of the group $\mu_2 \times \mu_2$ with the splitting field L .

A.1.4. Let S be a group of multiplicative type over a field k . Then the category $\mathcal{R}ep(S)$ of representations of S is a semisimple abelian category and the isomorphism classes of simple objects are classified by the orbits of $\text{Gal}(k^{sep}/k)$ acting on $\mathcal{X}^*(S)$ [Mi17, Theorem 12.30]. In particular, for the representation ring $\text{Rep}(S)$ there is a canonical isomorphism

$$\text{Rep}(S) \xrightarrow{\cong} \mathbb{Z}[\mathcal{X}^*(S)]^{\text{Gal}(k^{sep}/k)}. \quad (\text{A.1.4.1})$$

For $\lambda \in \mathcal{X}^*(S)$ we denote by x^λ the corresponding element in $\mathbb{Z}[\mathcal{X}^*(S)]$, and for a $\text{Gal}(k^{sep}/k)$ -invariant subset $\mathcal{Q} \subseteq \mathcal{X}^*(S)$ the element $\sum_{\lambda \in \mathcal{Q}} x^\lambda$ corresponds to a representation of S which we denote by $V(\mathcal{Q})$. The class of a representation $V \in \mathcal{R}ep(S)$ in $\text{Rep}(S)$ is denoted by $[V]$ and we freely use the above isomorphism, in particular, we have $[V(\mathcal{Q})] = \sum_{\lambda \in \mathcal{Q}} x^\lambda$. For a homomorphism $\rho: S_1 \rightarrow S_2$ we denote by $\text{Res}_\rho: \text{Rep}(S_2) \rightarrow \text{Rep}(S_1)$ the induced homomorphism of the representation rings. Isomorphism (A.1.4.1) is functorial, in particular, the following diagram commutes.

$$\begin{array}{ccc} \text{Rep}(S_2) & \xrightarrow{\cong} & \mathbb{Z}[\mathcal{X}^*(S_2)]^{\text{Gal}(k^{sep}/k)} \\ \downarrow \text{Res}_\rho & & \downarrow \mathbb{Z}[\mathcal{X}^*(\rho)]^{\text{Gal}(k^{sep}/k)} \\ \text{Rep}(S_1) & \xrightarrow{\cong} & \mathbb{Z}[\mathcal{X}^*(S_1)]^{\text{Gal}(k^{sep}/k)} \end{array}$$

Let L/k be a degree 2 Galois field extension. Then we have the following formulae for the classes of the standard vector representations introduced above:

$$[V_R] = x^{(1,0)} + x^{(0,1)}, \quad [{}^2V_{\mu_l}] = x^{\bar{1}} + x^{-\bar{1}}, \quad [{}^2V_{\mu_{2,2}}] = x^{(\bar{1},\bar{0})} + x^{(\bar{0},\bar{1})}, \quad [{}^2\Lambda_{\mu_{2m}}^\pm] = x^{\bar{m}}.$$

A.2 Quasi-split simple groups

A.2.1. Let G be a reductive group over a field k .

- [Mi17, Definition 17.67] G is *quasi-split*, if there exists a Borel subgroup $B \leq G$.
- [Mi17, Definition 18.7] Suppose that G is semisimple. The *fundamental group* of G is the kernel $\pi_1(G) := \ker(\tilde{G} \rightarrow G)$ of the simply connected cover. The group G is *adjoint*, if its center is trivial, or, equivalently, if its fundamental group coincides with the center of the simply connected cover.

A.2.2. Let G be a quasi-split semisimple group over a field k and let $T \leq B \leq G$ be a maximal torus and a Borel subgroup. Then $T_{k^{sep}} \leq B_{k^{sep}} \leq G_{k^{sep}}$ gives rise to a pinned root datum and since B is defined over k , the action of $\text{Gal}(k^{sep}/k)$ on $\mathcal{X}^*(T)$ stabilizes the set of simple roots $\Pi \subseteq \mathcal{X}^*(T)$ yielding an action of $\text{Gal}(k^{sep}/k)$ on the Dynkin diagram of $G_{k^{sep}}$. A simply connected quasi-split semisimple group over k is determined up to an isomorphism by the action of $\text{Gal}(k^{sep}/k)$ on the Dynkin diagram of $G_{k^{sep}}$ [KMRT98, Proposition 27.8], see also [Mi17, Theorem 25.33] and [Ti66, Theorem 2]. It follows that a quasi-split semisimple group is determined by its fundamental group and the action of $\text{Gal}(k^{sep}/k)$ on the Dynkin diagram of $G_{k^{sep}}$. We refer to the Dynkin diagram of $G_{k^{sep}}$ together with the action of $\text{Gal}(k^{sep}/k)$ as the *type of G* and denote it by $\Delta(G)$, so quasi-split semisimple groups are classified by the pairs $(\Delta(G), \pi_1(G))$ consisting of the type of G and the fundamental group of G . Note that $\Delta(G_{k^{sep}})$ is just the Dynkin diagram of $G_{k^{sep}}$.

Lemma A.2.3 ([Mi17, Exercise 25-4]). *Let G be a quasi-split semisimple group over a field k and let $T \leq B \leq G$ be a maximal torus and a Borel subgroup. Suppose that G is simply connected or adjoint. Then T is quasi-trivial.*

Proof. The set of simple roots associated with $T_{k^{sep}} \leq B_{k^{sep}} \leq G_{k^{sep}}$ is stable under the action of the Galois group $\text{Gal}(k^{sep}/k)$ on the lattice of characters $\mathcal{X}^*(T)$. In the adjoint case the simple roots form a basis of $\mathcal{X}^*(T)$ giving a permutation basis under the action of the Galois group. In the simply connected case a permutation basis of $\mathcal{X}^*(T)$ is given by the set of fundamental weights, which are permuted by the Galois group, since they are dual to the simple roots. \square

A.2.4. By a *simple group* in the present article we mean a geometrically almost-simple group in the sense of [Mi17, Definition 19.7], i.e. a semisimple group with $\Delta(G_{k^{sep}})$ being a connected Dynkin diagram.

A.2.5. Let G be a non-split quasi-split simple group over a field k and let $T \leq B \leq G$ be a maximal torus and a Borel subgroup. Then $\Delta(G_{k^{sep}}) = A_n, n \geq 2, D_n, n \geq 4, \text{ or } E_6$, since only these connected Dynkin diagrams admit non-trivial automorphisms. We recall some structural data in these cases.

A_n, n ≥ 2. The action of $\text{Gal}(k^{sep}/k)$ on the Dynkin diagram factors through an action of $\text{Gal}(L/k)$ for a degree 2 Galois field extension L/k which is the splitting field of the maximal torus T . The type of G in this case is denoted by $\Delta(G) = {}^2A_n$. The corresponding quasi-split simply connected group is denoted by $G^{sc}({}^2A_n)$ and it is isomorphic to the special unitary group $\text{SU}(V, h)$, where (V, h) is a non-degenerate hermitian form over L/k of dimension $n+1$ and of maximal Witt index [KMRT98, Example 27.9]. We use the following notation for roots and weights (see [Mi17, Example 22.34]):

- $\mathcal{X}_{\mathbb{R}} := \left(\bigoplus_{i=1}^{n+1} \mathbb{R}e_i \right) / \mathbb{R}(e_1 + e_2 + \dots + e_{n+1})$ and $\epsilon_i \in \mathcal{X}_{\mathbb{R}}$ denotes the image of e_i ,
- simple roots: $\alpha_i := \epsilon_i - \epsilon_{i+1} \in \mathcal{X}_{\mathbb{R}}, 1 \leq i \leq n$,
- fundamental weights: $\varpi_i := \epsilon_1 + \epsilon_2 + \dots + \epsilon_i, 1 \leq i \leq n$.

In the simply connected case we have $\mathcal{X}^*(T) = \bigoplus_{i=1}^n \mathbb{Z}\varpi_i = \bigoplus_{i=1}^n \mathbb{Z}\epsilon_i \leq \mathcal{X}_{\mathbb{R}}$ with the action of the nontrivial element $\tau \in \text{Gal}(L/k)$ given by $\tau(\epsilon_i) = -\epsilon_{n+2-i}$. For the center $Z \leq G^{sc}({}^2A_n)$ we have

$$\mathcal{X}^*(Z) = \mathcal{X}^*(T) / \bigoplus_{i=1}^n \mathbb{Z}\alpha_i \cong \mathbb{Z}/(n+1)\mathbb{Z}$$

and the embedding $Z \leq T$ corresponds to the projection $\mathcal{X}^*(T) \rightarrow \mathcal{X}^*(Z)$ given by $\epsilon_i \mapsto \bar{1} \in \mathbb{Z}/(n+1)\mathbb{Z}, 1 \leq i \leq n$, so $\varpi_i \mapsto \bar{i} \in \mathbb{Z}/(n+1)\mathbb{Z}$. The action of $\text{Gal}(L/k)$ on $\mathcal{X}^*(Z)$ is compatible with the projection, thus $\tau(x) = -x$ and $Z \cong {}^2\mu_{n+1}$ in the notation of Section A.1. In particular, for a quasi-split simple group G of type 2A_n one has $G \cong G^{sc}({}^2A_n)/{}^2\mu_l$ for some $l \mid n+1$. We denote by $G^{ad}({}^2A_n) := G^{sc}({}^2A_n)/{}^2\mu_{n+1}$ the adjoint group.

D_n, n ≥ 5. The action of $\text{Gal}(k^{sep}/k)$ on the Dynkin diagram factors through an action of $\text{Gal}(L/k)$ for a degree 2 Galois field extension L/k which is the splitting field of the maximal torus T . The type of G in this case is denoted by $\Delta(G) = {}^2D_n$. The corresponding quasi-split simply connected group is denoted by $G^{sc}({}^2D_n)$ and it is isomorphic to $\text{Spin}(V, q)$, where (V, q) is a non-degenerate quadratic form of dimension $2n$, of Witt index $n-1$ and with the discriminant quadratic field extension being L/k [KMRT98, Example 27.10]. We use the following notation for roots and weights (see [Bou81, Planche IV]):

- $\mathcal{X}_{\mathbb{R}} := \bigoplus_{i=1}^n \mathbb{R}e_i$,
- simple roots: $\alpha_i := e_i - e_{i+1}$, $1 \leq i \leq n-1$, $\alpha_n := e_{n-1} + e_n$,
- fundamental weights: $\varpi_i := e_1 + e_2 + \dots + e_i$, $1 \leq i \leq n-2$, $\varpi_{n-1} := \frac{1}{2}(e_1 + e_2 + \dots + e_{n-1} - e_n)$,
 $\varpi_n := \frac{1}{2}(e_1 + e_2 + \dots + e_{n-1} + e_n)$.

In the simply connected case we have $\mathcal{X}^*(T) = \bigoplus_{i=1}^n \mathbb{Z}\varpi_i \leq \mathcal{X}_{\mathbb{R}}$ with the action of the nontrivial element $\tau \in \text{Gal}(L/k)$ given by $\tau(\varpi_{n-1}) = \varpi_n$, $\tau(\varpi_n) = \varpi_{n-1}$ and $\tau(\varpi_i) = \varpi_i$ for $1 \leq i \leq n-2$. This action extends to $\mathcal{X}_{\mathbb{R}}$ via $\tau(e_n) = -e_n$ and $\tau(e_i) = e_i$ for $1 \leq i \leq n-1$. For the center $Z \leq G^{sc}({}^2D_n)$ we have

$$\mathcal{X}^*(Z) = \mathcal{X}^*(T) / \bigoplus_{i=1}^n \mathbb{Z}\alpha_i \cong \begin{cases} \mathbb{Z}/4\mathbb{Z}, & n \text{ is odd} \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}, & n \text{ is even.} \end{cases}$$

If n is odd, then the embedding $Z \leq T$ corresponds to the projection $\mathcal{X}^*(T) \rightarrow \mathcal{X}^*(Z)$ given by

$$\varpi_i \mapsto \bar{2}i \in \mathbb{Z}/4\mathbb{Z}, \quad 1 \leq i \leq n-2, \quad \varpi_{n-1} \mapsto \bar{1} \in \mathbb{Z}/4\mathbb{Z}, \quad \varpi_n \mapsto \bar{3} \in \mathbb{Z}/4\mathbb{Z}.$$

If n is even, then the embedding $Z \leq T$ corresponds to the projection $\mathcal{X}^*(T) \rightarrow \mathcal{X}^*(Z)$ given by

$$\varpi_i \mapsto (\bar{i}, \bar{i}) \in \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}, \quad 1 \leq i \leq n-2, \quad \varpi_{n-1} \mapsto (\bar{1}, 0), \quad \varpi_n \mapsto (0, \bar{1}).$$

The action of $\text{Gal}(L/k)$ on $\mathcal{X}^*(Z)$ is compatible with the projection, thus it is given by $x \mapsto -x$ on $\mathbb{Z}/4\mathbb{Z}$, if n is odd and by $(x_1, x_2) \mapsto (x_2, x_1)$ on $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, if n is even, and in the notation of Section A.1 we have $Z \cong {}^2\mu_4$ for n odd and $Z \cong {}^2\mu_{2,2}$ for n even. In both cases $\mathcal{X}^*(Z)$ admits a unique non-trivial quotient Galois module and this module is isomorphic to $\mathbb{Z}/2\mathbb{Z}$ with the trivial action. In particular, for a quasi-split simple group G with $\Delta(G) = {}^2D_n$, $n \geq 5$, one has $G \cong G^{sc}({}^2D_n)/\pi_1(G)$ with $\pi_1(G) = 1$, μ_2 or ${}^2\mu_4$ for odd n and $\pi_1(G) = 1$, μ_2 or ${}^2\mu_{2,2}$ for even n . We denote by $G^{ad}({}^2D_{2r+1}) := G^{sc}({}^2D_{2r+1})/{}^2\mu_4$ and by $G^{ad}({}^2D_{2r}) := G^{sc}({}^2D_{2r})/{}^2\mu_{2,2}$ the respective adjoint groups. Note that in contrast to the split situation in the non-split case there is a unique intermediate group between $G^{sc}({}^2D_{2r})$ and $G^{ad}({}^2D_{2r})$.

D₄. The action of $\text{Gal}(k^{sep}/k)$ on the Dynkin diagram factors through a subgroup of the automorphism group of the diagram which is isomorphic to S_3 . If the action factors through a subgroup of order 2, then the same reasoning (possibly renumbering the simple roots) as in the case $n \geq 5$ applies, yielding the groups $G^{sc}({}^2D_4)$, $G^{sc}({}^2D_4)/\mu_2$ and $G^{ad}({}^2D_4) = G^{sc}({}^2D_4)/{}^2\mu_{2,2}$. In the remaining cases the splitting field L of the maximal torus T is either of degree 3 over k or of degree 6 over k with the $\text{Gal}(L/k) \cong S_3$ in the latter case. The type of G in these cases is denoted by $\Delta(G) = {}^3D_4$ and by $\Delta(G) = {}^6D_4$ respectively, with the corresponding simply connected groups denoted by $G^{sc}({}^3D_4)$ and by $G^{sc}({}^6D_4)$. In both cases the Galois group acts irreducibly on $\mathcal{X}^*(Z) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ for the center $Z \leq G^{sc}({}^3D_4)$ (resp. $Z \leq G^{sc}({}^6D_4)$), yielding $Z \cong {}^3\mu_{2,2}$ (resp. $Z \cong {}^6\mu_{2,2}$) and the non-split quasi-split groups are given by $G^{sc}({}^3D_4)$ and by $G^{ad}({}^3D_4) := G^{sc}({}^3D_4)/{}^3\mu_{2,2}$ (resp. $G^{sc}({}^6D_4)$ and $G^{ad}({}^6D_4) := G^{sc}({}^6D_4)/{}^6\mu_{2,2}$) in the notation of Section A.1.

E₆. The action of $\text{Gal}(k^{sep}/k)$ on the Dynkin diagram factors through an action of $\text{Gal}(L/k)$ for a degree 2 Galois field extension L/k which is the splitting field of the maximal torus T . The type of G in this case is denoted by $\Delta(G) = {}^2E_6$, the corresponding quasi-split simply connected group is denoted by $G^{sc}({}^2E_6)$. We use the following notation for roots and weights (see [Bou81, Planche V]):

- $\mathcal{X}_{\mathbb{R}} := \{x_1e_1 + x_2e_2 + \dots + x_8e_8 \in \mathbb{R}^8 \mid x_6 = x_7 = -x_8\} \leq \mathbb{R}^8$,
- simple roots: $\alpha_1 := \frac{1}{2}(e_1 - e_2 - e_3 - e_4 - e_5 - e_6 - e_7 + e_8)$, $\alpha_2 := e_1 + e_2$, $\alpha_3 := e_2 - e_1$, $\alpha_4 := e_3 - e_2$,
 $\alpha_5 := e_4 - e_3$, $\alpha_6 := e_5 - e_4$,
- fundamental weights: $\varpi_1 := \frac{2}{3}(e_8 - e_7 - e_6)$, $\varpi_2 := \frac{1}{2}(e_1 + e_2 + e_3 + e_4 + e_5 - e_6 - e_7 + e_8)$, $\varpi_3 := \frac{1}{2}(-e_1 + e_2 + e_3 + e_4 + e_5) + \frac{5}{6}(e_8 - e_7 - e_6)$, $\varpi_4 := e_3 + e_4 + e_5 - e_6 - e_7 + e_8$, $\varpi_5 := e_4 + e_5 + \frac{2}{3}(e_8 - e_7 - e_6)$,
 $\varpi_6 := e_5 + \frac{1}{3}(e_8 - e_7 - e_6)$.

In the simply connected case we have $\mathcal{X}^*(T) = \bigoplus_{i=1}^6 \mathbb{Z}\varpi_i$ with the action of the nontrivial element $\tau \in \text{Gal}(L/k)$ given by

$$\tau(\varpi_1) = \varpi_6, \quad \tau(\varpi_2) = \varpi_2, \quad \tau(\varpi_3) = \varpi_5, \quad \tau(\varpi_4) = \varpi_4, \quad \tau(\varpi_5) = \varpi_3, \quad \tau(\varpi_6) = \varpi_1.$$

For the center $Z \leq G^{sc}({}^2E_6)$ we have

$$\mathcal{X}^*(Z) = \mathcal{X}^*(T) / \bigoplus_{i=1}^6 \mathbb{Z}\alpha_i \cong \mathbb{Z}/3\mathbb{Z}$$

and the embedding $Z \leq T$ corresponds to the projection $\mathcal{X}^*(T) \rightarrow \mathcal{X}^*(Z)$ given by

$$\varpi_2, \varpi_4 \mapsto \bar{0} \in \mathbb{Z}/3\mathbb{Z}, \quad \varpi_1, \varpi_5 \mapsto \bar{1} \in \mathbb{Z}/3\mathbb{Z}, \quad \varpi_3, \varpi_6 \mapsto \bar{2} \in \mathbb{Z}/3\mathbb{Z}.$$

The action of $\text{Gal}(L/k)$ on $\mathcal{X}^*(Z)$ is compatible with the projection, thus it is given by $\tau(x) = -x$ and $Z \cong {}^2\mu_3$ in the notation of Section A.1. We denote by $G^{ad}({}^2E_6) := G^{sc}({}^2E_6)/{}^2\mu_3$ the corresponding adjoint group. In particular, for a quasi-split simple group G with $\Delta(G) = {}^2E_6$ one has $G \cong G^{sc}({}^2E_6)$ or $G \cong G^{ad}({}^2E_6)$.

A.2.6. Let G be a quasi-split semisimple group over a field k with a maximal torus and a Borel subgroup $T \leq B \leq G$ and let $I \subseteq \Delta(G_{k^{sep}})$ be a subset of its Dynkin diagram stable under the action of $\text{Gal}(k^{sep}/k)$. Then the corresponding parabolic subgroup of $G_{k^{sep}}$ [Mi17, Theorem 21.91] is defined over k , and we denote it by $P_I \leq G$. In particular, $P_\emptyset = B$ while maximal proper $\text{Gal}(k^{sep}/k)$ -stable subsets $I \subseteq \Delta(G_{k^{sep}})$ correspond to maximal proper parabolic subgroups $P_I \leq G$. For $1 \leq i_1 < i_2 < \dots < i_j \leq \text{rank}(G_{k^{sep}})$ such that $J := \{\alpha_{i_1}, \alpha_{i_2}, \dots, \alpha_{i_j}\} \subseteq \Delta(G_{k^{sep}})$ is $\text{Gal}(k^{sep}/k)$ -stable we put

$$P_{i_1, i_2, \dots, i_j} := P_{\Delta(G_{k^{sep}}) \setminus J}.$$

A.3 Vector bundles on homogeneous varieties

A.3.1. Let G be an algebraic group over a field k and let $H \leq G$ be a closed subgroup. Descent gives rise to an equivalence between the category of representations of H and the category of G -equivariant vector bundles over G/H ,

$$\text{Rep}(H) \xrightarrow{\sim} \text{Vect}_G(G/H), \quad V \mapsto (G \times V)/H,$$

where H acts on G via $(h, g) \mapsto gh^{-1}$. Forgetting about the equivariant structure we obtain a vector bundle \mathcal{V} over G/H .

A.3.2. Let G be a quasi-split reductive group over a field k with a maximal torus and a Borel subgroup $T \leq B \leq G$. According to (A.1.4), a finite $\text{Gal}(k^{sep}/k)$ -invariant subset $\mathcal{Q} \subseteq \mathcal{X}^*(T)$ gives rise to $V(\mathcal{Q}) \in \text{Rep}(T)$ such that

$$[V(\mathcal{S})_{k^{sep}}] = \sum_{\lambda \in \mathcal{Q}} x^\lambda.$$

Since the embedding $T \leq B$ has a canonical splitting, it follows that $V(\mathcal{Q})$ can be viewed as a representation of B in a canonical way. We denote by $\mathcal{V}(\mathcal{Q})$ the associated vector bundle over G/B ,

$$\mathcal{V}(\mathcal{Q}) := (G \times V(\mathcal{Q}))/B.$$

If $\mathcal{Q} = \{\varpi\}$ consists of a single element, we usually denote by $L(\varpi) := V(\{\varpi\})$ the corresponding linear representation and by $\mathcal{L}(\varpi) := \mathcal{V}(\{\varpi\})$ the corresponding line bundle.

Suppose G is semisimple of rank n , let $f: \tilde{G} \rightarrow G$ be the simply connected cover and put $\tilde{T} := f^{-1}(T)$, $\tilde{B} := f^{-1}(B)$. Then for a finite $\text{Gal}(k^{sep}/k)$ -invariant subset $\mathcal{Q} \subseteq \mathcal{X}^*(\tilde{T})$ the vector bundle $\mathcal{V}(\mathcal{Q})$ over \tilde{G}/\tilde{B} can be considered as a vector bundle over G/B via the isomorphism $\tilde{G}/\tilde{B} \xrightarrow{\sim} G/B$ induced by f . For a $\text{Gal}(k^{sep}/k)$ -invariant fundamental weight $\varpi_i \in \mathcal{X}^*(\tilde{T})$ and a $\text{Gal}(k^{sep}/k)$ -invariant set $\{\varpi_{i_1}, \varpi_{i_2}, \dots, \varpi_{i_j}\} \subseteq \{\varpi_1, \varpi_2, \dots, \varpi_n\} \subseteq \mathcal{X}^*(\tilde{T})$ of fundamental weights we put

$$\mathcal{L}_i := \mathcal{L}(\varpi_i), \quad \mathcal{V}_{i_1, i_2, \dots, i_j} := \mathcal{V}(\{\varpi_{i_1}, \varpi_{i_2}, \dots, \varpi_{i_j}\})$$

viewed as vector bundles over G/B .

Let $1 \leq i_1 < i_2 < \dots < i_j \leq n$ be such that $J := \{\alpha_{i_1}, \alpha_{i_2}, \dots, \alpha_{i_j}\} \subseteq \Delta(G_{k^{sep}})$ is $\text{Gal}(k^{sep}/k)$ -stable, put $P := P_{i_1, i_2, \dots, i_j} \leq G$ in the notation of (A.2.6) and $\tilde{P} := f^{-1}(P) \leq \tilde{G}$. For a $\text{Gal}(k^{sep}/k)$ -invariant

$\varpi \in \mathbb{Z}\varpi_{i_1} \oplus \mathbb{Z}\varpi_{i_2} \oplus \dots \oplus \mathbb{Z}\varpi_{i_j} \subseteq \mathcal{X}^*(\tilde{T})$ it follows from [Mi17, Theorem 21.91] that there exists a unique linear representation $V_{\tilde{P}}(\varpi)$ of \tilde{P} such that $[V_{\tilde{P}}(\varpi)_{k^{sep}}|_{\tilde{T}_{k^{sep}}}] = x^\varpi$. We put

$$\mathcal{L}_P(\varpi) := (\tilde{G} \times V_{\tilde{P}}(\varpi))/\tilde{P}$$

for the associated line bundle over \tilde{G}/\tilde{P} which may also be viewed as a line bundle over G/P via the isomorphism $\tilde{G}/\tilde{P} \xrightarrow{\cong} G/P$ induced by f . For the projection $g: G/B \rightarrow G/P$ one has $g^*\mathcal{L}_P(\varpi) \cong \mathcal{L}(\varpi)$.

A.3.3. Let G be a semisimple group over a field k . Then a $\text{Gal}(k^{sep}/k)$ -invariant character $\lambda \in \mathcal{X}^*(\pi_1(G))$ gives rise to a line bundle

$$\mathcal{L}(\lambda) := (\tilde{G} \times V(\{\lambda\}))/\pi_1(G)$$

over $G \cong \tilde{G}/\pi_1(G)$. Suppose G is quasi-split and let $T \leq B \leq G$ be a maximal torus and a Borel subgroup, let $f: \tilde{G} \rightarrow G$ be the simply connected cover and put $\tilde{T} := f^{-1}(T)$. Consider $\varpi \in \mathcal{X}^*(\tilde{T})$ such that its image $\bar{\varpi} \in \mathcal{X}^*(\pi_1(G)) \cong \mathcal{X}^*(\tilde{T})/\mathcal{X}^*(T)$ is $\text{Gal}(k^{sep}/k)$ -invariant. Then we have a line bundle $\mathcal{L}(\bar{\varpi})$ over G .

A.4 Fundamental group and Picard group of a quasi-split simple group

A.4.1. Let G be a quasi-split simple group over a field k . Then its fundamental group $\pi_1(G)$ depending on its type $\Delta(G)$ is one of the following.

$\Delta(G)$	A_n	B_n, C_n, E_7	$D_{2r+1}, r \geq 1$	$D_{2r}, r \geq 2$	F_4, G_2, E_8	E_6
$\pi_1(G)$	$ \mu_l, l (n+1)$	$1, \mu_2$	$1, \mu_2, \mu_4$	$1, \mu_2^{so}, \mu_2^{hs}, \mu_2 \times \mu_2$	1	$1, \mu_3$
$\Delta(G)$	${}^2A_n, n \geq 2$	${}^2D_{2r+1}, r \geq 1$	${}^2D_{2r}, r \geq 2$	3D_4	6D_4	2E_6
$\pi_1(G)$	${}^2\mu_l, l (n+1)$	$1, \mu_2, {}^2\mu_4$	$1, \mu_2, {}^2\mu_{2,2}$	$1, {}^3\mu_{2,2}$	$1, {}^6\mu_{2,2}$	$1, {}^2\mu_3$

Here we use the notation of Sections A.1 and A.2. The list arises from the fact that $\pi_1(G)$ is a subgroup of the center $Z(\tilde{G})$ of the simply connected cover $\tilde{G} \rightarrow G$, from the identification of $\mathcal{X}^*(Z(\tilde{G}))$ with the weight lattice of $\tilde{G}_{k^{sep}}$ modulo its root lattice [Mi17, Proposition 21.8, Proposition 23.59] and from the description of the corresponding lattices [Bou81, Planche I-IX]. In the non-split quasi-split case see also the discussion of the Galois action on $\mathcal{X}^*(Z(\tilde{G}))$ given in Section A.2.

Let G be a split simply connected simple group with $\Delta(G) = D_{2r}$ and $T \leq G$ be a split maximal torus. Then there are three embeddings $\mu_2 \rightarrow \mu_2 \times \mu_2 \cong Z(G) \leq T$ with the dual homomorphisms $\mathcal{X}^*(T) \rightarrow \mathcal{X}^*(\mu_2) \cong \mathbb{Z}/2\mathbb{Z}$ given by

$$\begin{aligned} \varpi_i &\mapsto \bar{0} \in \mathbb{Z}/2\mathbb{Z}, \quad 1 \leq i \leq 2r-2, & \varpi_{2r-1}, \varpi_{2r} &\mapsto \bar{1} \in \mathbb{Z}/2\mathbb{Z}, \\ \varpi_{2i} &\mapsto \bar{0} \in \mathbb{Z}/2\mathbb{Z}, \quad 1 \leq i \leq r, & \varpi_{2i-1} &\mapsto \bar{1} \in \mathbb{Z}/2\mathbb{Z}, \quad 1 \leq i \leq r, \\ \varpi_{2i}, \varpi_{2r-1} &\mapsto \bar{0} \in \mathbb{Z}/2\mathbb{Z}, \quad 1 \leq i \leq r-1, & \varpi_{2i-1}, \varpi_{2r} &\mapsto \bar{1} \in \mathbb{Z}/2\mathbb{Z}, \quad 1 \leq i \leq r-1. \end{aligned}$$

For the last two embeddings the quotient groups G/μ_2 are isomorphic. We denote in the table the first and the second embeddings as $\mu_2^{so} \leq T \leq G$ and $\mu_2^{hs} \leq T \leq G$ respectively.

A.4.2. Let G be a semisimple group over a field k and adopt the notation of (A.3.3). The homomorphism

$$\mathcal{X}^*(\pi_1(G))^{\text{Gal}(k^{sep}/k)} \rightarrow \text{Pic}(G), \quad \lambda \mapsto \mathcal{L}(\lambda),$$

is an isomorphism [Mi17, Corollary 18.26]. Thus the Picard group of a quasi-split simple group G is given by the following table, and if G is not in the table, then $\text{Pic}(G) = 0$.

Table 5: Picard group of a quasi-split simple algebraic group

$\Delta(G)$	$\pi_1(G)$	$\text{Pic}(G)$
A_n	$\mu_l, l \mid (n+1)$	$\mathbb{Z}/l\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_1)]$
${}^2A_{2r-1}$	${}^2\mu_{2m}, m \mid r$	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_m)]$
B_n	μ_2	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_n)]$
C_n	μ_2	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_1)]$
$D_{2r}, r \geq 2$	μ_2^{so}	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_{2r-1})]$
$D_{2r+1}, r \geq 1$	μ_2	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_{2r})]$
${}^2D_n, n \geq 3$	μ_2	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_{n-1})]$
$D_{2r}, r \geq 2$	μ_2^{hs}	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_1)]$
$D_{2r+1}, r \geq 1$	μ_4	$\mathbb{Z}/4\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_{2r})]$
${}^2D_{2r+1}, r \geq 1$	${}^2\mu_4$	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_1)]$
$D_{2r}, r \geq 2$	$\mu_2 \times \mu_2$	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_{2r-1})] \oplus \mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_{2r})]$
${}^2D_{2r}, r \geq 2$	${}^2\mu_{2,2}$	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_1)]$
E_6	μ_3	$\mathbb{Z}/3\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_1)]$
E_7	μ_2	$\mathbb{Z}/2\mathbb{Z} \cdot [\mathcal{L}(\bar{\omega}_2)]$

Here we use the numbering of the fundamental weights as in [Bou81, Planche I-IX]. Other possible generators for $\text{Pic}(G)$ can be easily read off from the description of the projection $\mathcal{X}^*(T) \rightarrow \mathcal{X}^*(\pi_1(G))$.

Lemma A.4.3. *Let G be a simply connected quasi-split simple group over a field k and let $N_1 \leq N_2 \leq Z(G)$ be central subgroups with $N_1 \neq N_2$. Then the pullback*

$$\pi^*: \text{Pic}(G/N_2) \rightarrow \text{Pic}(G/N_1)$$

for the projection $\pi: G/N_1 \rightarrow G/N_2$ is zero unless $(\Delta(G), N_1, N_2)$ is one of the following

- $(A_n, \mu_{l_1}, \mu_{l_2})$ with $1 \neq l_1 \mid l_2 \mid n+1$,
- $({}^2A_{2r-1}, {}^2\mu_{2m_1}, {}^2\mu_{2m_2})$ with $m_1 \mid m_2 \mid r$ and $\frac{m_2}{m_1}$ being odd,
- $(D_{2r}, \mu_2^{so}, \mu_2 \times \mu_2)$, $(D_{2r}, \mu_2^{hs}, \mu_2 \times \mu_2)$, (D_{2r+1}, μ_2, μ_4) .

Proof. Straightforward from the above. □

A.5 Chow ring of a split simple group

A.5.1. Let G be a reductive group over a field k . It is well-known that $\text{CH}^*(G) \otimes \mathbb{Q} \cong \mathbb{Q}$. This could be seen via the the isomorphisms

$$\mathbb{Q} \cong \text{CH}^*(G_K) \otimes \mathbb{Q} \cong \text{CH}^*(G) \otimes \mathbb{Q},$$

with K being the splitting field of G , where the first isomorphism is given by [Gro58, Remark 2 on p. 21] while the second one then follows from the standard pushforward-pullback argument (as in Lemma 3.4.1). Alternatively, one has isomorphisms

$$\text{CH}^*(G) \otimes \mathbb{Q} \cong K_0(G) \otimes \mathbb{Q} \cong \mathbb{Q}$$

with the first one given by the Chern character [Fu98, Example 15.2.16] and the second one given by [Me97, Corollary 7.8].

A.5.2. Let G be a split reductive group over a field k . It follows from the Bruhat decomposition and [To14a, Proposition 1] that the Künneth formula $\mathrm{CH}^*(G) \otimes \mathrm{CH}^*(G) \xrightarrow{\cong} \mathrm{CH}^*(G \times G)$ holds, endowing $\mathrm{CH}^*(G)$ with the structure of a Hopf algebra. Let $p \in \mathbb{N}$ be a prime and put $\mathrm{Ch}^*(G) := \mathrm{CH}^*(G) \otimes \mathbb{F}_p$. Then it follows from a Borel's result on the structure of Hopf algebras [MM65, Theorem 7.11 and Proposition 7.8] that

$$\mathrm{Ch}^*(G) \cong \mathbb{F}_p[e_1, e_2, \dots, e_s] / (e_1^{p^{k_1}}, e_2^{p^{k_2}}, \dots, e_s^{p^{k_s}}), \quad \deg e_i = d_i,$$

for some integers $s, d_i, k_i \in \mathbb{N}$.

Suppose G is a simple group. For $k = \mathbb{C}$ the integers s, d_i, k_i were determined in [Kac85, Theorem 6], and the answer does not depend on k e.g. because of the combinatorial presentation for $\mathrm{Ch}^*(G)$ given by the characteristic sequence. One has $\mathrm{Ch}^*(G) \not\cong \mathbb{F}_p$ only for a finite list of primes p . These primes and the respective parameters are summarized in the following table, where v_p is the p -adic valuation.

Table 6: $\mathrm{Ch}^*(G)$ for a split simple algebraic group G

$\Delta(G)$	$\pi_1(G)$	p	s	$d_i, i = 1, 2, \dots, s$	$k_i, i = 1, 2, \dots, s$
A_n	$\mu_l, l \mid (n+1)$	$p \mid l$	1	1	$v_p(n+1)$
B_n	1	2	$\lfloor \frac{n-1}{2} \rfloor$	$2i+1$	$\lfloor \log_2 \frac{2n}{2i+1} \rfloor$
	μ_2	2	$\lfloor \frac{n+1}{2} \rfloor$	$2i-1$	$\lfloor \log_2 \frac{2n}{2i-1} \rfloor$
C_n	μ_2	2	1	1	$v_2(n) + 1$
$D_n, n \geq 3$	1	2	$\lfloor \frac{n}{2} \rfloor - 1$	$2i+1$	$\lfloor \log_2 \frac{2n-1}{2i+1} \rfloor$
$D_{2r}, r \geq 2$	μ_2^{so}	2	r	$2i-1$	$\lfloor \log_2 \frac{4r-1}{2i-1} \rfloor$
	μ_2^{hs}	2	r	$1, i=1$	$v_2(r) + 1, i=1$
				$2i-1, i \geq 2$	$\lfloor \log_2 \frac{4r-1}{2i-1} \rfloor, i \geq 2$
	$\mu_2 \times \mu_2$	2	$r+1$	$1, i=1$	$v_2(r) + 1, i=1$
				$2i-3, i \geq 2$	$\lfloor \log_2 \frac{4r-1}{2i-3} \rfloor, i \geq 2$
$D_{2r+1}, r \geq 1$	μ_2, μ_4	2	r	$2i-1$	$\lfloor \log_2 \frac{4r+1}{2i-1} \rfloor$
E_6	$1, \mu_3$	2	1	3	1
	1	3	1	4	1
	μ_3	3	2	1, 4	2, 1
E_7	1	2	3	3, 5, 9	1, 1, 1
	μ_2	2	4	1, 3, 5, 9	1, 1, 1, 1
	$1, \mu_2$	3	1	4	1
E_8	1	2	4	3, 5, 9, 15	3, 2, 1, 1
		3	2	4, 10	1, 1
		5	1	6	1
F_4	1	2	1	3	1
		3	1	4	1
G_2	1	2	1	3	1

References

- [Bou81] N. Bourbaki, *Groupes et algèbres de Lie*, ch. IV-VI, Masson, Paris, 1981. [55](#), [56](#), [58](#), [59](#)
- [CM06] V. Chernousov, A. Merkurjev, *Motivic decomposition of projective homogeneous varieties and the Krull-Schmidt theorem*, Transformation Groups 11 (2006), 371–386. [5](#), [21](#), [23](#)
- [Che94] C. Chevalley, *Sur les décompositions cellulaires des espaces G/B* , Proc. Sympos. Pure Math. 56 1–23, Amer. Math. Soc., Providence, 1994. [31](#), [32](#)
- [EG98] D. Edidin, W. Graham, *Equivariant intersection theory (with an appendix by A. Vistoli: The Chow ring of \mathcal{M}_2)*, Invent. Math. 131 (1998), 595–634. [5](#), [10](#), [11](#), [13](#)
- [EKM08] R. Elman, N. Karpenko, A. Merkurjev, *The algebraic and geometric theory of quadratic forms*, vol. 56 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 2008. [6](#), [8](#), [36](#), [37](#), [38](#), [39](#)
- [Fi19] R. Fino, *J-invariant of hermitian forms over quadratic extensions*, Pacific J. of Math. 300:2 (2019), 375–404. [2](#), [5](#), [8](#)
- [Fu98] W. Fulton, *Intersection theory* (2nd ed), vol. 2 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics. Springer-Verlag, Berlin, 1998. [13](#), [24](#), [47](#), [59](#)
- [GPS16] S. Garibaldi, V. Petrov, N. Semenov, *Shells of twisted flag varieties and the Rost invariant*, Duke Math. J. 165:2 (2016), 285–339. [2](#)
- [GS10] S. Garibaldi, N. Semenov, *Degree 5 invariant of E_8* , Int. Math. Res. Not. 2010, no. 19, 3746–3762. [2](#)
- [GZ22] N. Geldhauser, M. Zhykhovich, *J-invariant of linear algebraic groups of outer type*, arXiv:2205.02884 [2](#), [5](#), [8](#), [31](#), [33](#), [35](#), [44](#)
- [Gro58] A. Grothendieck, *Torsion homologique et sections rationnelles*, in Séminaire Claude Chevalley, tome 3 (1958), exp. 5, 1–29. [2](#), [4](#), [5](#), [23](#), [59](#)
- [Kac85] V. Kac, *Torsion in cohomology of compact Lie groups and Chow rings of reductive algebraic groups*, Invent. Math. 80 (1985), 69–79. [2](#), [4](#), [6](#), [60](#)
- [Ka00] N. A. Karpenko, *Weil transfer of algebraic cycles*, Indag. Math. (N.S.) 11:1 (2000), 73–86. [15](#)
- [Ka12] N. A. Karpenko, *Unitary grassmannians*, J. Pure Appl. Algebra 216 (2012), 2586–2600. [2](#), [5](#), [15](#)
- [KM22] N. A. Karpenko, A. S. Merkurjev, *The Chow ring of the classifying space of the unitary group*, Algebra Number Theory 16:1 (2022), 139–154. [2](#), [5](#), [15](#)
- [KMRT98] M.-A. Knus, A. Merkurjev, M. Rost, J.-P. Tignol, *The book of involutions*, Colloquium Publications, vol. 44, AMS 1998. [37](#), [55](#)
- [KaZ13] N. A. Karpenko, M. Zhykhovich, *Isotropy of unitary involutions*, Acta Math. 211:2 (2013), 227–253. [2](#), [5](#), [23](#)
- [LM07] M. Levine, F. Morel, *Algebraic cobordism*, Springer Monographs in Mathematics, Springer-Verlag Berlin Heidelberg, 2007. [5](#), [8](#)
- [Ma68] Y. Manin, *Correspondences, motifs and monoidal transformations*, Math. USSR Sbornik 6 (1968), 439–470. [8](#)
- [Me97] A.S. Merkurjev, *Comparison of the equivariant and the standard K-theory of algebraic varieties*, Algebra i Analiz 9:4 (1997), 175–214. [59](#)

- [Mi17] J. S. Milne, *Algebraic groups. The theory of group schemes of finite type over a field*, Cambridge Studies in Advanced Mathematics, 170. Cambridge University Press, Cambridge, 2017. [7](#), [53](#), [54](#), [55](#), [57](#), [58](#)
- [MM65] J. Milnor, J. Moore, *On the structure of Hopf algebras*, Ann. Math. 81:2 (1965), 211–264. [31](#), [45](#), [60](#)
- [MRV06] L. A. Molina Rojas, A. Vistoli, *On the Chow rings of classifying spaces for classical groups*, Rend. Sem. Mat. Univ. Padova 116 (2006), 271–298. [2](#)
- [MV99] F. Morel, V. Voevodsky, \mathbb{A}^1 -homotopy theory of schemes, Publ. Math. IHES 90 (1999), 45–143. [2](#)
- [Pa98] R. Pandharipande, *Equivariant Chow rings of $O(k)$, $SO(2k+1)$, and $SO(4)$* , J. Reine Angew. Math. 496 (1998), 131–148. [2](#)
- [PS10] V. Petrov, N. Semenov, *Generically split projective homogeneous varieties*, Duke Math. J. 152 (2010), 155–173. [2](#)
- [PS12] V. Petrov, N. Semenov, *Generically split projective homogeneous varieties. II*, J. of K-theory 10:1 (2012), 1–8. [2](#)
- [PS17] V. Petrov, N. Semenov, *Rost motives, affine varieties, and classifying spaces*, J. London Math. Soc. 95:2 (2017), 895–918. [2](#), [43](#)
- [PSZ08] V. Petrov, N. Semenov, K. Zainoulline, *J-invariant of linear algebraic groups*, Ann. Sci. École Norm. Sup. 41 (2008), 1023–1053. [2](#)
- [Sal22] F. Sala, *On the Chow ring of the classifying stack of algebraic tori*, Doc. Math. 27 (2022), 917–932. [5](#), [16](#)
- [S16] N. Semenov, *Motivic construction of cohomological invariants*, Comment. Math. Helv. 91:1 (2016), 163–202. [2](#)
- [Ti66] J. Tits, *Classification of algebraic semisimple groups*, in Algebraic Groups and Discontinuous Subgroups, pp. 33–62, Proceedings of Symposia in Pure Mathematics, Boulder, CO, 1965. American Mathematical Society, Providence, RI. [55](#)
- [To99] B. Totaro, *The Chow ring of a classifying space*, Algebraic K-Theory, Proceedings of Symposia in Pure Mathematics 67 (eds W. Raskind and C. Weibel; American Mathematical Society, Providence, RI, 1999) 249–281. [2](#), [5](#), [10](#)
- [To14a] B. Totaro, *Chow groups, Chow cohomology and linear varieties*, Forum Math. Sigma 2 (2014), e17. [60](#)
- [To14b] B. Totaro, *Group cohomology and algebraic cycles*, Cambridge (2014). [21](#)
- [Vi05] A. Vishik, *On the Chow groups of quadratic Grassmannians*, Doc. Math. 10 (2005), 111–130. [2](#)
- [Vi07] A. Vishik, *Fields of u -invariant $2r+1$* , in: Algebra, Arithmetic and Geometry, Manin Festschrift, Birkhäuser. (2007). [2](#), [6](#), [36](#), [37](#), [40](#)
- [Vi24] A. Vishik, *Isotropic and numerical equivalence for Chow groups and Morava K-theories*, Inventiones Math. 237 (2024), 779–808. [2](#)

Alexey Ananyevskiy, Mathematisches Institut der Universität München, Theresienstr. 39, D-80333 München, Germany.
E-mail address: alseang@gmail.com

Nikita Geldhauser, Mathematisches Institut der Universität München, Theresienstr. 39, D-80333 München, Germany.
E-mail address: geldhauser@math.lmu.de