

C^2 ESTIMATES FOR k -HESSIAN EQUATIONS AND A RIGIDITY THEOREM

RUIJIA ZHANG

ABSTRACT. We derive a concavity inequality for k -Hessian operators under the semi-convexity condition. As an application, we establish interior estimates for semi-convex solutions of the k -Hessian equations with vanishing Dirichlet boundary and obtain a Liouville-type result. Additionally, we provide new and simple proofs of Guan-Ren-Wang's work [12] on global curvature estimates for k -curvature equations.

1. INTRODUCTION

In this paper, we consider the following k -Hessian equations with Dirichlet boundary

$$\begin{cases} \sigma_k(D^2u) = f(x, u, \nabla u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where Ω is a bounded domain in \mathbb{R}^n , f is a positive function in $\overline{\Omega} \times R \times R^n$, u is a function defined in $\overline{\Omega}$. Here we denote by ∇u and D^2u the gradient of u and the Hessian of u , respectively. σ_k is the k -th elementary symmetric polynomial of $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n$, which is defined as

$$\sigma_k(x) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} \lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_k}, \quad \forall i = 1, \dots, n.$$

Let λ be the eigenvalues of D^2u . Then the k -Hessian operator of u is defined by $\sigma_k(D^2u) = \sigma_k(\lambda(D^2))$.

Research on the k -Hessian equation

$$\sigma_k(D^2u) = f(x, u, \nabla u) \quad (1.2)$$

is a vital area within PDE and geometric analysis. We note that (1.2) simplifies to a semi-linear equation when $k = 1$. For $2 \leq k \leq n$, establishing second-order a priori estimates is essential for proving the existence of solutions to these fully nonlinear equations. Pogorelov was the first to investigate the interior C^2 estimates for the Monge-Ampère equations, as referred to in [10, 17]. For general k , Chou-Wang [7] established the Pogorelov estimates for k -convex solutions of the k -Hessian equation

$$\begin{cases} \sigma_k(D^2u) = f(x, u) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.3)$$

where f is independent of ∇u . They proved that

$$(-u)^\beta \Delta u \leq C$$

2020 *Mathematics Subject Classification.* 35J60, 53C42.

Key words and phrases. semi-convex, interior estimates, Liouville problem.

for some constants $\beta > 1$ and $C > 0$ depending on $n, k, \inf f, \|f\|_{C^{1,1}}, \|u\|_{C^1}$ but independent of Ω . Following [3], we define the k -convex function by : For a domain $\Omega \in \mathbb{R}^n$, a function $u \in C^2(\Omega)$ is called k -convex (or k -admissible) if the eigenvalues $\lambda(X)$ of the Hessian D^2u are in Γ_k (defined by (2.1)) for all $X \in \Omega$. We note that for all k -convex functions u , the k -Hessian operator is elliptic, i.e.,

$$\dot{\sigma}_k^{ij} = \frac{\partial \sigma_k}{\partial u_{ij}}$$

is positive-definite. In the following text, we consider the k -Hessian equation in this realm.

We extend Theorem 1.5 by Chou-Wang [7] to cases where f depends on ∇u , under an additional convexity assumption for the function u . We refer to the function u as semi-convex if there exists a constant $A > 0$ such that the eigenvalues λ of D^2u satisfy

$$\lambda_i(D^2u(x)) \geq -A, \quad \text{for all } x \in \Omega$$

for any $1 \leq i \leq n$. We have obtained the following interior estimates for semi-convex solutions of Equation (1.1).

Theorem 1.1. *Assume that $f \in C^2(\Omega \times \mathbb{R} \times \mathbb{R}^n)$ satisfies $f > f_0 > 0$ for a constant $f_0 > 0$ in $\overline{\Omega}$. Then, for any semi-convex k -admissible solution of the Dirichlet problem of the equation (1.1) provided $2 < k \leq n - 1$, we have*

$$(-u)^\alpha \Delta u \leq C.$$

where $\alpha > 0$ and $C > 0$ depend on $n, k, f_0, \|f\|_{C^{1,1}}, \|u\|_{C^1}$ but are independent of Ω .

Remark 1.1. The interior estimates in Theorem 1.1 were derived by Li-Ren-Wang [15] under the assumption that u is in the $(k + 1)$ -convex cone. Tu proved the interior estimates in Theorem 1.1 by using a test function involving $|X|^2$ in [23]. Both of the results mentioned above depend on the diameter of the domain Ω . In the proof of Theorem 1.1, we carry on with the method in [7] by Chou-Wang and have obtained estimates independent of Ω .

Using this class of interior estimates, we further derive a rigidity theorem for entire solutions in \mathbb{R}^n to

$$\sigma_k(D^2u) = 1. \tag{1.4}$$

It has long been a question of concern whether an entire function u in n -dimensional Euclidean space satisfying (1.4) is quadratic. For $k = 1$, the Liouville property of harmonic functions implies rigidity results for solutions with bounded Hessian. For $k = n$, the Jörgens–Calabi–Pogorelov theorem confirms the rigidity results for all convex entire solutions. Cheng and Yau [6] provided an alternative geometric proof. For $k = 2$, Chang and Yuan [4] proved that the entire solution to 2-Hessian equation (1.4) in \mathbb{R}^n is quadratic if

$$D^2u \geq \delta - \sqrt{2}n(n-1)I$$

for any $\delta > 0$. For $2 \leq k \leq n - 1$, the authors also guess that the rigidity theorem holds true under the semi-convexity assumption

$$D^2u \geq -AI$$

with arbitrarily large constant $A > 0$. Warren [24] constructed a class of non-trivial solutions to (1.4), which are neither semi-convex nor exhibit quadratic growth. Shankar and Yuan [20] obtained the rigidity theorem for $k = 2$ under the semi-convexity assumption. For a general k , the rigidity problem for the k -Hessian equation remains open. Assuming a quadratic

growth condition, Bao-Chen-Guan-Ji [2] showed that strictly convex entire solutions to (1.4) are quadratic, Li-Ren-Wang [15] extended the convexity assumption to $(k + 1)$ -convexity, and Chu-Dinew [9] further relaxed the condition to

$$\sigma_{k+1} > -A \tag{1.5}$$

for some $A > 0$ (see also [23]). A function u is said to have quadratic growth if there exists constants $b, c > 0$ and sufficient large $R > 0$, such that the following inequality holds:

$$u(x) \geq c|x|^2 - b, \quad \text{for } |x| \geq R. \tag{1.6}$$

In this paper, we confirm Chang-Yang's conjecture under the quadratic growth condition.

Theorem 1.2. *Given $2 < k \leq n - 1$, let u be any semi-convex entire function in \mathbb{R}^n with $\lambda(D^2u) \in \Gamma_k$ that satisfies*

$$\sigma_k(D^2u) = 1 \tag{1.7}$$

and exhibits quadratic growth. Then, u is a quadratic polynomial.

To establish the curvature estimates and derive the rigidity theorem, the main difficulty to overcome is how to deal with the third order derivatives. It is well known that $\sigma_k^{\frac{1}{k}}$ is concave in the Γ_k cone, which is very useful to deal with these terms. It's worth noting that Ren and Wang [18] established the concavity inequality of the Hessian operator for $k = n - 1$. They proved that

Theorem 1.3. [18] *Given two positive constants $\delta < 1$, ε , for any index i , if $\kappa_i \geq \delta\kappa_1$ and the positive constant K depending only on δ and ε is sufficient large, then we have*

$$\kappa_i [K(\sigma_{n-1})_i^2 - \sigma_{n-1}^{pp,qq} h_{ppi} h_{qqi}] - \sigma_{n-1}^{ii} h_{iii}^2 + (1 + \varepsilon) \sum_{j \neq i} \sigma_{n-1}^{jj} h_{jji}^2 \geq 0. \tag{1.8}$$

In [18], they also pointed out that the inequality (1.8) is not true even if $i = 1$, $\sigma_k = 1$ and κ_1 is sufficient large for $k < n - 1$. Besides, Guan and Qiu [11] established the interior C^2 estimates of σ_2 -Hessian equation under the assumption $\sigma_3 \geq -A$ for some constant $A > 0$. In their work, it's crucial to observe that under the condition $\sigma_3 \geq -A$ and κ_1 large enough

$$-\sum_{i \neq j} u_{iii} u_{jj1} + C \frac{(\dot{\sigma}_2^{ii} u_{ii1})^2}{\sigma_2} \geq (1 + \delta_0) \frac{\dot{\sigma}_2^{11} u_{111}^2}{u_{11}^2} \tag{1.9}$$

for some constant $C > 0$ and $\delta_0 > 0$. (1.9) is basically the similar form of (1.8) and is derived by using Chen's [5] work about the optimal concavity of the σ_2 operators with the eigenvector decomposition method. We note that the method makes sense only for $k = 2$ since the Hessian operator is the identity metric. It's difficult to generalize this method to other cases. Moreover, it reminds us that the concavity inequality of the general k -Hessian operator may also hold true under some additional convexity assumption. we develop a method that builds on [12] by Guan-Ren-Wang and [13] by Hong-Zhang which is grounded in the crucial observation of the concavity of $q_k = \frac{\sigma_k}{\sigma_{k-1}}$, introduced by Huisken-Sinestrari in [14]. Under the semi-convexity condition, we derive a concavity inequality which plays a crucial role in establishing curvature estimates of the Dirichlet problem of k -Hessian equations (1.1) and (5.1).

Lemma 1.1. *Assume $\{\lambda_i\} \in \Gamma_k$, $\lambda_1 \geq \lambda_2 \cdots \geq \lambda_n$, and $\lambda_n > -A$ for a constant $A > 0$. Then there exists some constant $C > 0$ depending on n, k, σ_k, A and small constant $\delta_0 > 0$ depending on k such that if $k_1 \geq C$, then the following inequality holds*

$$-\sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + K \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} + 2 \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{(\lambda_1 + A + 1) \sigma_k} \geq (1 + \delta_0) \frac{\dot{\sigma}_k^{11} \xi_1^2}{\lambda_1} \quad (1.10)$$

for some sufficient large $K > 0$ (depending on δ_0) where $\xi = (\xi_1, \dots, \xi_n)$ is an arbitrary vector in \mathbb{R}^n .

In the following text, we note that $\sigma_{k+1} > -A$ implies that $\lambda_i > -C(n, k, \sigma_k, A)$ by using Lemma 2.2. In other words, the semi-convexity of $\{\lambda_i\}$ is a weaker condition, which means that Lemma 1.1 is useful in deriving Guan-Qiu's [11] interior curvature estimates. Besides, Lemma 1.1 removes the additional assumption that $|\kappa_n| \geq \delta_0$ for some constant $\delta_0 > 0$ from Hong and Zhang [13] by employing an enhanced and concise approach.

Using the proof in Lemma 1.1, we also improve Lemma 3.1 from Lu [16] and present it as Lemma 3.1. In [16], the concavity inequality, Lemma 3.1, plays a crucial role in obtaining curvature estimates for admissible solutions to prescribed curvature measure type problem in hyperbolic space (referencing [25]). We present Lemma 3.1 in [16] here for ease of comparison.

Lemma 1.2 ([16]). *Let $\lambda = (\lambda_1, \dots, \lambda_n) \in \Gamma_k$ with $\lambda_1 \geq \dots \geq \lambda_n$ and let $1 \leq l < k$. For any $\epsilon, \delta, \delta_0 \in (0, 1)$, there exists a constant $\delta' > 0$ depending only on $\epsilon, \delta, \delta_0, n, k$ and l such that if $\lambda_l \geq \delta \lambda_1$ and $\lambda_{l+1} \leq \delta' \lambda_1$, then we have*

$$-\sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} \geq (1 - \epsilon) \frac{\xi_1^2}{\lambda_1^2} - \delta_0 \sum_{i>l} \frac{\sigma_k^{ii} \xi_i^2}{\lambda_1 \sigma_k},$$

where $\xi = (\xi_1, \dots, \xi_n)$ is an arbitrary vector in \mathbb{R}^n .

As a matter of fact, it's clear that if $\lambda_n > -A$ for a constant $A > 0$ and λ_1 is large enough, then $\dot{\sigma}_k^{11} \geq \frac{\sigma_k}{\lambda_1}$, which implies that Lemma 1.1 is a development of Lemma 1.2.

Moreover, Lemma 1.1 can be regarded as an extension of Lemma 3.3 by Chu [8] under a weaker convexity assumption. By applying Lemma 3.3 from [8], Chu gave a simple proof of Theorem 1.1 by Guan, Ren and Wang [12], which established the curvature estimate for convex hypersurfaces satisfying the Weingarten curvature equation

$$\sigma_k(\kappa(X)) = f(X, \nu(X)), \quad \forall X \in M^n, \quad (1.11)$$

where $f \in C^2(\Gamma)$ is a positive function and Γ is an open neighborhood of the unit normal bundle of M in $\mathbb{R}^{n+1} \times \mathbb{S}^n$, $\kappa(X)$ denotes the principal curvatures, $\nu(X)$ is the unit outer normal vector at point X , and σ_k represents the k -th mean curvature.

In Remark 4.7 of Guan-Ren-Wang [12], they also deduced the following curvature estimates under a weaker convexity condition. Let $M^n \in \mathbb{R}^{n+1}$ be a closed hypersurface. We say a hypersurface M is semi-convex if there exists a constant $A > 0$ such that its principal curvatures satisfy

$$\kappa_i(X) \geq -A, \quad 1 \leq i \leq n, \quad \forall X \in M.$$

Theorem 1.4 (Guan-Ren-Wang, [12]). *For any closed semi-convex hypersurface M^n satisfying the curvature equation (1.11), there exists a constant $C > 0$ depending on n , k , $\|M\|_{C^1}$, $\inf f$, and $\|f\|_{C^2}$ such that*

$$\max_{X \in M} \kappa_i(X) \leq C.$$

In the specific case where $k = 2$, they completely resolved the problem.

Theorem 1.5 (Guan-Ren-Wang, [12]). *For the solution of the curvature equation (1.1) with $k = 2$, it holds that*

$$\max_{X \in M} \kappa_i(X) \leq C$$

where $C > 0$ is a constant that depends on n , k , $\|M\|_{C^1}$, $\inf f$ and $\|f\|_{C^2}$.

As an application of Lemma 1.1, we also offer new simple proofs of Theorem 1.4 and Theorem 1.5. As mentioned in Remark 4.7 of [12], the curvature estimates in Theorem 1.4 can be obtained by considering the test function

$$\log P(\kappa(X)) + g(u) + a|X|^2$$

where P is a symmetric polynomial of the principal curvatures. It should be noted that Spruck-Xiao [21] provided a simplified proof of Theorem 1.5. Chu proved Theorem 1.4 more simply by assuming further that M^n is a closed convex hypersurface. In both of these works, the maximum principle was applied to a quantity involving the maximal principal curvature rather than the symmetric curvature function $P(\kappa)$. In what follows, we continue with this approach and use Lemma 1.1 to prove these results.

The paper is organized as follows. In Section 2, we collect and prove some algebraic properties of symmetric polynomials. In Section 3, we prove the crucial Lemma concerning the superior concavity of k -Hessian operator. In Section 4, we derive the Pogorelov estimates for the k -Hessian equation with vanishing Dirichlet boundary and complete the proof of Theorem 1.1. In Section 5, we prove the rigidity theorem and complete the proof of Theorem 1.2. In Section 6, we provide a new and simple proof of Guan-Ren-Wang's famous work [12] by using Lemma 1.1 derived in Section 3.

Acknowledgments. The author would like to thank Prof. Haizhong Li for bringing many related works to our attention regarding this problem. We are also deeply grateful to Prof. Yuguang Shi for his ongoing support and encouragement. The author is supported by the National Key R&D Program of China 2020YFA0712800, the China Postdoctoral Science Foundation No.2023M740108, and the Postdoctoral Fellowship Program of CPSF under Grant Number GZC20230094.

2. PRELIMINARIES

2.1. Symmetric function. Here we state some algebraic properties of the elementary symmetric function $\sigma_m(\lambda)$, $m = 1, \dots, n$, where $\lambda = (\lambda_1, \dots, \lambda_n)$. We recall $\sigma_m(\lambda)$ defined by

$$\sigma_m(\lambda) = \sum_{1 \leq i_1 < \dots < i_m \leq n} \lambda_{i_1} \cdots \lambda_{i_m}, \quad m = 1, 2, \dots, n.$$

The Garding cone Γ_m is an open symmetric convex cone in \mathbb{R}^n with vertex at the origin, given by

$$\Gamma_m = \{(\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n \mid \sigma_j(\lambda) > 0, \forall j = 1, \dots, m\}. \quad (2.1)$$

Clearly $\sigma_m(\lambda) = 0$ for $\lambda \in \partial\bar{\Gamma}_m$ and

$$\Gamma_n \subset \dots \subset \Gamma_m \subset \dots \subset \Gamma_1.$$

In particular, $\Gamma^+ = \Gamma_n$ is called the positive cone,

$$\Gamma^+ = \{(\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n \mid \lambda_1 > 0, \dots, \lambda_n > 0\}.$$

Always assume $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. We collect some properties of $\sigma_k(\lambda)$ where $\lambda \in \Gamma_k$:

- (1) $\sum_{p=1}^n \frac{\partial \sigma_k}{\partial \lambda_p} \lambda_p^2 = \sigma_1 \sigma_k - (k+1)\sigma_{k+1}$.
- (2) $\sum_{p=1}^n \frac{\partial \sigma_k}{\partial \lambda_p} = (n-k+1)\sigma_{k-1}$.
- (3) $\frac{\partial \sigma_k}{\partial \lambda_n} \geq \dots \geq \frac{\partial \sigma_k}{\partial \lambda_1} > 0$.
- (4) $\frac{\partial \sigma_k}{\partial \lambda_1} \lambda_1 \geq \frac{k}{n} \sigma_k$.
- (5) $q_k := \frac{\sigma_k}{\sigma_{k-1}}$ is concave, i.e., $\partial_\xi^2 q_k \leq 0$ for any $\xi \in \mathbb{R}^n$.
- (6) For any $s < k$, $\sigma_s > \lambda_1 \lambda_2 \dots \lambda_s$.

Lemma 2.1. *Assume that $\{\lambda_i\} \in \Gamma_k$ and $\lambda_1 \geq \lambda_2 \dots \geq \lambda_n$. Then $\lambda_k > 0$ and $|\lambda_n| < (n-k)\lambda_k$.*

Proof. From Property (3), we have

$$\frac{\partial \sigma_k}{\partial \lambda_1 \dots \partial \lambda_{k-1}} = \lambda_k + \dots + \lambda_n > 0.$$

Suppose only λ_n is negative, then $(n-k)\lambda_k > |\lambda_n|$. \square

Lemma 2.2. *Assume that $\{\lambda_i\} \in \Gamma_k$, $\lambda_1 \geq \lambda_2 \dots \geq \lambda_n$, and $\sigma_k(\lambda)$ bounded from upper and below. If $\sigma_{k+1} > -A$ for any positive constant A , then λ is semi-convex, i.e., $\lambda_{\min} > -C(n, k, A)$.*

Proof. There exists constant $C > 0$, such that

$$\frac{\partial \sigma_k}{\partial \lambda_n} \lambda_n^2 \leq \sum_i \frac{\partial \sigma_k}{\partial \lambda_i} \lambda_i^2 = \sigma_1 \sigma_k - (k+1)\sigma_{k+1} < C\sigma_1 + (k+1)A. \quad (2.2)$$

Then by Property (3) we have

$$\sum_i \frac{\partial \sigma_k}{\partial \lambda_i} \lambda_n^2 < nC\sigma_1 + n(k+1)A.$$

Combining $(n-k+1)\sigma_{k-1} = \sum_i \frac{\partial \sigma_k}{\partial \lambda_i}$ and Property (6), Lemma 2.1, there exists constant $C(n, k) > 0$

$$C'(n, k)|\lambda_n|^{k+1} \leq C(n, k)|\lambda_n|^k \lambda_1 < Cn\lambda_1 + (k+1)A.$$

Thus we get $|\lambda_n| < C(n, k, A)$ by Lemma 2.1, completing the proof. \square

Lemma 2.3. *Assume $\{\lambda_i\} \in \Gamma_k$ and $\lambda_1 \geq \lambda_2 \dots \geq \lambda_n$. If $\lambda_n > -A$, then $\lambda_k < C(\sigma_k, n, A)$.*

Proof. Assume that $\lambda_{k-1} > A$, otherwise $\lambda_k \leq A$, then we are done. Since

$$\sigma_k = \sum_{1 \leq i_1 \leq i_2 \leq \dots \leq i_k \leq n} \lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_k} > 0.$$

We have

$$\lambda_1 \cdots \lambda_{k-1} \lambda_k \leq \sigma_k - C(n, k) \lambda_1 \cdots \lambda_{k-1} \lambda_n. \quad (2.3)$$

Thus

$$\begin{aligned} \lambda_k &\leq 2 \left(\frac{\sigma_k}{\lambda_1} \right)^{\frac{1}{k-1}} + 2C(n, k) |\lambda_n| \\ &\leq 2 \left(\frac{n\sigma_k}{\sigma_1} \right)^{\frac{1}{k-1}} + 2C(n, k) A \\ &\leq c(n, k) \sigma_k^{\frac{1}{k}} + C(n, k, A). \end{aligned} \quad (2.4)$$

Here we use Property (6) in the second inequality, the last inequality follows from Maclaurin's inequality. Thus there exists some constant $C > 0$, such that $\lambda_k \leq C$, which depends on the bounds of σ_k , n and A . □

Lemma 2.4. *If $\{\lambda_i\} \in \Gamma_k$ and $\lambda_1 \geq \lambda_2 \cdots \geq \lambda_n$, then $\sigma_{k-1} \geq C \lambda_1^{\frac{1}{k-1}}$.*

Proof. Using Newton inequality and Property (6), we have

$$\sigma_{k-1} \geq C(n, k) \sigma_1^{\frac{1}{k-1}} \sigma_k^{\frac{k-2}{k-1}} \geq C \lambda_1^{\frac{1}{k-1}}. \quad (2.5)$$

□

We point out that we will denote C , c or C_i and c_i for $i = 1, 2, \dots, n$ as some positive constants throughout the subsequent sections, and these constants may change from line to line. Now we state a crucial calculation by Huisken-Sinestrari [14].

Lemma 2.5. *Given $k \geq 2$, if $\{\lambda_i\} \in \Gamma_k$, then for any $\xi \in \mathbb{R}^n$,*

$$-\partial_\xi^2 q_2 = \sum_{i=1}^n \frac{\left(\xi_i - \frac{\lambda_i}{\sigma_1} \sigma_1(\xi) \right)^2}{\sigma_1} \quad (2.6)$$

and

$$\partial_\xi^2 q_{k+1} \leq \sum_{i=1}^n \frac{\lambda_i^2 \partial_\xi^2 q_{k;i}}{(k+1)(q_{k;i} + \lambda_i)^2} \quad (2.7)$$

where $q_k = \frac{\sigma_k}{\sigma_{k-1}}$, $q_{k;i} = \frac{\sigma_k(\lambda|i)}{\sigma_{k-1}(\lambda|i)}$, and $\sigma_k(\lambda|i) = \frac{\partial \sigma_{k+1}}{\partial \lambda_i}$.

If $\xi \perp \gamma$, then $-\sigma_1 \partial_\xi^2 q_2 \geq |\xi|^2$ by (2.6). More generally, we have the following lemmas that are crucial to the proof of Lemma 1.1.

Lemma 2.6. *If $\{\lambda_i\} \in \Gamma_k$, then for any $\xi \in \mathbb{R}^n$,*

$$-\partial_\xi^2 q_{2;i_1, \dots, i_{k-2}} \geq \frac{|\xi_{i_1, \dots, i_{k-2}}^\perp|^2}{\sigma_1(\lambda|i_1, \dots, i_{k-2})}. \quad (2.8)$$

Here for any $1 \leq \ell \leq k-2$, $[\xi]_{i_1, \dots, i_{k-2}}$ denotes the vector that is obtained by throwing out the i_1 th, \dots , i_ℓ th components of ξ and $[\xi]_{i_1, \dots, i_\ell}^\perp$ denotes orthogonal part of $[\xi]_{i_1, \dots, i_\ell}$ with respect to $[\lambda]_{i_1, \dots, i_\ell}$. Moreover,

$$-\partial_\xi^2 q_{k-1; i_1} \leq \sum_{i=\{1, \dots, n\} \setminus i_1} \frac{\lambda_i^2 \partial_\xi^2 q_{k-2; i_1, i}}{(k-1)(q_{k-1; i_1, i} + \lambda_i)^2} \quad (2.9)$$

Proof. From (2.6) in Lemma 2.5 it follows

$$-\partial_\xi^2 q_{2; i_1, \dots, i_{k-2}} = \sum_{i=\{1, \dots, n\} \setminus \{i_1, \dots, i_{k-2}\}} \frac{\left(\xi_i - \frac{\lambda_i}{\sigma_1(\lambda|i_1, \dots, i_{k-2})} \sigma_1(\xi|i_1, \dots, i_{k-2}) \right)^2}{\sigma_1(\lambda|i_1, \dots, i_{k-2})} \quad (2.10)$$

Then (2.8) holds by decomposing $[\xi]_{i_1, \dots, i_{k-2}}$ with respect to $[\lambda]_{i_1, \dots, i_{k-2}}$, i.e., write $[\xi]_{i_1, \dots, i_{k-2}} = c[\lambda]_{i_1, \dots, i_{k-2}} + [\xi]_{i_1, \dots, i_{k-2}}^\perp$.

(2.9) follows from (2.7). \square

Lemma 2.7. *If $\{\lambda_i\} \in \Gamma_k$, then for any $\gamma \in \mathbb{R}^n$,*

$$-\sigma_{k-1} \partial_\gamma^2 q_k \geq C \sum_{j=1}^{k-1} \lambda_1 \cdots \hat{\lambda}_j \cdots \lambda_{k-1} \left| [\gamma]_{1, \dots, \hat{j}, \dots, k-1}^\perp \right|^2. \quad (2.11)$$

Proof. We note that

$$q_{k; i} + \lambda_i = \frac{\sigma_k}{\sigma_{k-1}(\lambda|i)}. \quad (2.12)$$

By (2.7), (2.9) and (2.8), we have

$$\begin{aligned} -\sigma_{k-1} \partial_\gamma^2 q_k &\geq - \sum_{i_1=1}^n \frac{\lambda_{i_1}^2 \partial_\gamma^2 q_{k-1; i_1} \sigma_{k-2}(\lambda|i_1)}{k \frac{\sigma_{k-1}}{\sigma_{k-2}(\lambda|i_1)}} \\ &\geq - \sum_{i_1=1}^n \sum_{i_2=\{1, \dots, n\} \setminus \{i_1\}} \frac{\lambda_{i_1}^2 \lambda_{i_2}^2 \partial_\gamma^2 q_{k-2; i_1, i_2} \sigma_{k-3}(\lambda|i_1, i_2)}{k(k-1) \frac{\sigma_{k-1}}{\sigma_{k-2}(\lambda|i_1)} \frac{\sigma_{k-2}(\lambda|i_1)}{\sigma_{k-3}(\lambda|i_1, i_2)}} \\ &\geq \dots \\ &\geq - \sum_{i_1=1}^n \dots \sum_{i_{k-2}=\{1, \dots, n\} \setminus \{i_1, \dots, i_{k-3}\}} \frac{2\lambda_{i_1}^2 \lambda_{i_2}^2 \cdots \lambda_{i_{k-2}}^2 \partial_\gamma^2 q_{2; i_1, i_2, \dots, i_{k-2}} \sigma_1(\lambda|i_1, i_2, \dots, i_{k-2})}{k! \frac{\sigma_{k-1}}{\sigma_1(\lambda|i_1, i_2, \dots, i_{k-2})}} \\ &\geq \sum_{i_1=1}^n \dots \sum_{i_{k-2}=\{1, \dots, n\} \setminus \{i_1, \dots, i_{k-3}\}} \frac{2\lambda_{i_1}^2 \lambda_{i_2}^2 \cdots \lambda_{i_{k-2}}^2 |[\gamma]_{i_1, \dots, i_{k-2}}^\perp|^2}{k! \frac{\sigma_{k-1}}{\sigma_1(\lambda|i_1, i_2, \dots, i_{k-2})}} \end{aligned} \quad (2.13)$$

Taking $(i_1, i_2, \dots, i_{k-2})$ from $(1, \dots, k-1)$ and denote the remaining index by j , by using the facts that $\sigma_{k-1} \leq C(n, k)\lambda_1 \cdots \lambda_{k-1}$ and $\sigma_1(\lambda|_{i_1, i_2, \dots, i_{k-2}}) \geq \lambda_j$, we obtain

$$-\sigma_{k-1} \partial_\gamma^2 q_k \geq C \sum_{1 \leq i_1 \leq \dots \leq i_{k-2} \leq n} \lambda_{i_1} \cdots \lambda_{i_{k-2}} \left| [\gamma]_{i_1, i_2, \dots, i_{k-2}}^\perp \right|^2. \quad (2.14)$$

Summing over j completes the proof. \square

Lemma 2.8. *Assume that $\gamma = (0, \gamma_2, \dots, \gamma_{k-1}, \gamma_k, \dots, \gamma_n)$ satisfying*

$$(\gamma_k, \dots, \gamma_n) \perp (\lambda_k, \dots, \lambda_n). \quad (2.15)$$

If $\{\lambda_i\} \in \Gamma_k$, then for any $2 \leq i \leq k-1$ and $k \leq p \leq n$,

$$-\sigma_{k-1} \partial_\gamma^2 q_k \geq C \frac{\sum_{j=1}^{k-1} \lambda_1 \lambda_2 \cdots \lambda_{k-1} \lambda_k^2 \gamma_j^2}{\lambda_j^3} + C \lambda_1 \lambda_2 \cdots \lambda_{k-2} \sum_{p=k}^n \gamma_p^2. \quad (2.16)$$

Proof. Denote by

$$V_i = \{(\gamma_i, \gamma_k, \dots, \gamma_n) \in \mathbb{R}^{n-k+2}\}$$

for $2 \leq i \leq k-1$ and

$$V = \{(0, \gamma_k, \dots, \gamma_n) | (\gamma_k, \dots, \gamma_n) \perp (\lambda_k, \dots, \lambda_n)\}.$$

For convenience, we name $(0, \lambda_k, \dots, \lambda_n)$ as β . Let $\alpha_i = (\lambda_i, \lambda_k, \dots, \lambda_n)$, $\beta_i = (\sin \theta_i, \cos \theta_i \frac{\beta}{|\beta|})$ and $\tan \theta_i = -\frac{|\beta|}{\lambda_i}$. We note that for any $2 \leq i \leq k-1$, $\beta_i \perp \alpha_i$, $\beta_i \perp \beta$ and

$$V_i = V \oplus \mathcal{L}(\beta_i) \oplus \mathcal{L}(\alpha_i).$$

Then we can write any $\theta_i = (\lambda_i, \lambda_k, \dots, \lambda_n) \in V_i$ as

$$\theta_i = (0, \gamma_p, \dots, \gamma_n) - \frac{|\beta|}{|\alpha_i|} \gamma_i \beta_i + \frac{\lambda_i}{|\alpha_i|^2} \gamma_i \alpha_i \quad (2.17)$$

where $\gamma_i \in V$. Plugging (2.17) into (2.11), we obtain

$$-\sigma_{k-1} \partial_\gamma^2 q_k \geq C \sum_{j=2}^{k-1} \lambda_1 \cdots \hat{\lambda}_j \cdots \lambda_{k-1} \left(\sum_{k \leq p \leq n} \gamma_p^2 + \frac{|\beta|^2}{|\alpha_j|^2} \gamma_j^2 \right) \quad (2.18)$$

$$\begin{aligned} &\geq C \sum_{j=2}^{k-1} \lambda_1 \cdots \hat{\lambda}_j \cdots \lambda_{k-1} \left(\sum_{k \leq p \leq n} \gamma_p^2 + c_0 \frac{\lambda_k^2}{\lambda_j^2} \gamma_j^2 \right) \\ &\geq C \frac{\sum_{2 \leq j \leq k-1} \lambda_1 \lambda_2 \cdots \lambda_{k-1} \lambda_k^2 \gamma_j^2}{\lambda_j^3} + C \lambda_1 \lambda_2 \cdots \lambda_{k-2} \sum_{k \leq p \leq n} \gamma_p^2 \end{aligned} \quad (2.19)$$

where we use Lemma 2.1 in the second inequality. \square

For convenience, we perform some straightforward calculations for q_k here.

Lemma 2.9.

$$\frac{\partial^2 q_k}{\partial \lambda_p \partial \lambda_q} = \frac{\partial^2 \sigma_k}{\partial \lambda_p \partial \lambda_q} - \frac{\partial \sigma_k}{\partial \lambda_p} \frac{\partial \sigma_{k-1}}{\partial \lambda_q} - \frac{\partial \sigma_k}{\partial \lambda_q} \frac{\partial \sigma_{k-1}}{\partial \lambda_p} - \frac{\partial^2 \sigma_{k-1}}{\partial \lambda_p \partial \lambda_q} + 2 \frac{\partial \sigma_{k-1}}{\partial \lambda_p} \frac{\partial \sigma_{k-1}}{\partial \lambda_q} \sigma_{k-1}^{-2}. \quad (2.20)$$

Proof.

$$\begin{aligned}
\frac{\frac{\partial^2 q_k}{\partial \lambda_p \partial \lambda_q}}{q_k} &= \frac{\partial \left(\frac{\frac{\partial \sigma_k}{\partial \lambda_p}}{\sigma_{k-1}} - \frac{\frac{\partial \sigma_{k-1}}{\partial \lambda_p} \sigma_k}{\sigma_{k-1}^2} \right)}{q_k \partial \lambda_q} \\
&= \frac{\frac{\frac{\partial^2 \sigma_k}{\partial \lambda_p \partial \lambda_q}}{\sigma_{k-1}} - \frac{\frac{\partial \sigma_k}{\partial \lambda_p} \frac{\partial \sigma_{k-1}}{\partial \lambda_q}}{\sigma_{k-1}^2} - \frac{\frac{\partial \sigma_k}{\partial \lambda_q} \frac{\partial \sigma_{k-1}}{\partial \lambda_p}}{\sigma_{k-1}^2} - \frac{\frac{\partial^2 \sigma_{k-1}}{\partial \lambda_p \partial \lambda_q} \sigma_k}{\sigma_{k-1}^2} + \frac{\frac{\partial \sigma_{k-1}}{\partial \lambda_p} \frac{\partial \sigma_{k-1}}{\partial \lambda_q} \sigma_k}{\sigma_{k-1}^3}}{q_k} \\
&= \frac{\frac{\partial^2 \sigma_k}{\partial \lambda_p \partial \lambda_q}}{\sigma_k} - \frac{\frac{\partial \sigma_k}{\partial \lambda_p} \frac{\partial \sigma_{k-1}}{\partial \lambda_q}}{\sigma_k \sigma_{k-1}} - \frac{\frac{\partial \sigma_k}{\partial \lambda_q} \frac{\partial \sigma_{k-1}}{\partial \lambda_p}}{\sigma_k \sigma_{k-1}} - \frac{\frac{\partial^2 \sigma_{k-1}}{\partial \lambda_p \partial \lambda_q}}{\sigma_{k-1}} + 2 \frac{\frac{\partial \sigma_{k-1}}{\partial \lambda_p} \frac{\partial \sigma_{k-1}}{\partial \lambda_q}}{\sigma_{k-1}^2}.
\end{aligned} \tag{2.21}$$

□

3. A CONCAVITY INEQUALITY

In this section, we define

$$\dot{\sigma}_k^{pp}(\lambda) = \frac{\partial \sigma_k}{\partial \lambda_p}, \quad \ddot{\sigma}_k^{pp,qq}(\lambda) = \frac{\partial^2 \sigma_k}{\partial \lambda_p \partial \lambda_q}.$$

Subsequently, we provide the proof of the crucial Lemma 1.1.

Proof. First we note that $\log \sigma_k$ is concave. From Property (5), we have for any vector $\xi = (\xi_1, \dots, \xi_n)$ in \mathbb{R}^n

$$\begin{aligned}
-\partial_\xi^2 \log \sigma_k &= -\partial_\xi^2 (\log q_k + \log q_{k-1} + \dots + \log q_2 + \log \sigma_1) \\
&= -\frac{\partial_\xi^2 q_k}{q_k} + (\partial_\xi \log q_k)^2 - \frac{\partial_\xi^2 q_{k-1}}{q_{k-1}} + (\partial_\xi \log q_{k-1})^2 + \dots + (\partial_\xi \log \sigma_1)^2 \\
&\geq \sum_{i=1}^k (\partial_\xi \log q_i)^2 \geq 0.
\end{aligned} \tag{3.1}$$

Besides, we denote by $\hat{\xi} = (0, \xi_2, \dots, \xi_n)$ and choose $K \geq \frac{1}{\epsilon}$, then

$$\begin{aligned}
&-\partial_\xi^2 \log \sigma_k + (K-1) \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} \\
&\geq -\frac{\partial_\xi^2 q_k}{q_k} + (\partial_\xi \log \sigma_k - \partial_\xi \log \sigma_{k-1})^2 - \partial_\xi^2 \log \sigma_{k-1} + \left(\frac{1}{\epsilon} - 1 \right) (\partial_\xi \log \sigma_k)^2 \\
&\geq -\frac{\partial_\xi^2 q_k}{q_k} + (1-\epsilon) (\partial_\xi \log \sigma_{k-1})^2 - \partial_\xi^2 \log \sigma_{k-1} \\
&= -\frac{\partial_\xi^2 q_k}{q_k} + (1-\epsilon) \left(\frac{\dot{\sigma}_{k-1}^{11} \xi_1}{\sigma_{k-1}} + \partial_\xi \log \sigma_{k-1} \right)^2 - \partial_\xi^2 \log \sigma_{k-1} \\
&\quad + \left(\frac{\dot{\sigma}_{k-1}^{11} \xi_1}{\sigma_{k-1}} \right)^2 + 2 \sum_{p>1} \frac{\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{pp} - \ddot{\sigma}_{k-1}^{11,pp} \sigma_{k-1}}{\sigma_{k-1}^2} \\
&\geq -\frac{\partial_\xi^2 q_k}{q_k} + (1-\epsilon) \left(\frac{\dot{\sigma}_{k-1}^{11} \xi_1}{\sigma_{k-1}} + \sum_{i=1}^{k-1} \partial_\xi \log q_i \right)^2 + \sum_{i=1}^{k-1} (\partial_\xi \log q_i)^2
\end{aligned}$$

$$\begin{aligned}
 & + \left(\frac{\dot{\sigma}_{k-1}^{11} \xi_1}{\sigma_{k-1}} \right)^2 + 2 \sum_{\alpha > 1} \frac{\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{\alpha\alpha} - \ddot{\sigma}_{k-1}^{11, \alpha\alpha} \sigma_{k-1}}{\sigma_{k-1}^2} \xi_1 \xi_\alpha \\
 & \geq -\frac{\partial_\xi^2 q_k}{q_k} + \left(1 + (1 - \epsilon) \frac{1}{k} \right) \left(\frac{\dot{\sigma}_{k-1}^{11} \xi_1}{\sigma_{k-1}} \right)^2 + 2 \sum_{\alpha > 1} \frac{\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{\alpha\alpha} - \ddot{\sigma}_{k-1}^{11, \alpha\alpha} \sigma_{k-1}}{\sigma_{k-1}^2} \xi_1 \xi_\alpha \quad (3.2)
 \end{aligned}$$

by choosing $K = \frac{1}{\epsilon}$ Here we estimate the second and the third inequality in (3.2) as

$$\left(\frac{\dot{\sigma}_{k-1}^{11} \xi_1}{\sigma_{k-1}} + \sum_{i=1}^{k-1} \partial_\xi \log q_i \right)^2 + \sum_{i=1}^{k-1} \left(\partial_\xi \log q_i \right)^2 \geq \frac{1}{k} \left(\frac{\dot{\sigma}_{k-1}^{11} \xi_1}{\sigma_{k-1}} \right)^2. \quad (3.3)$$

By

$$\sigma_l = \dot{\sigma}_l^{11} \lambda_1 + \sigma_l(\lambda|1), \quad (3.4)$$

we have

$$\frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} = \frac{\sigma_{k-1} - \sigma_{k-1}(\lambda|1)}{\lambda_1 \sigma_{k-1}} \quad (3.5)$$

and

$$|\sigma_{k-1}(\lambda|1)| = C(n, k) \lambda_2 \lambda_3 \cdots \lambda_k \leq C(A) \frac{\sigma_{k-1}}{\lambda_1} \quad (3.6)$$

where we use Property (6) in the last inequality. Thus we obtain

$$\frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} \geq \frac{1}{\lambda_1} - \frac{C(A)}{\lambda_1^2}. \quad (3.7)$$

Since

$$\frac{(\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{\alpha\alpha} - \ddot{\sigma}_{k-1}^{11, \alpha\alpha} \sigma_{k-1}) \xi_1 \xi_\alpha}{\sigma_{k-1}^2} = \frac{(\sigma_{k-2}(\lambda|1, \alpha))^2 - \sigma_{k-1}(\lambda|1, \alpha) \sigma_{k-3}(\lambda|1, \alpha)}{\sigma_{k-1}^2} \xi_1 \xi_\alpha, \quad (3.8)$$

for any $1 < i \leq k-1$, we have

$$\sigma_{k-2}(\lambda|1, i) \leq C(n, k) \frac{\lambda_1 \lambda_2 \cdots \lambda_k}{\lambda_1 \lambda_i} \leq C(n, k) \frac{\sigma_{k-1} \lambda_k}{\lambda_1 \lambda_i} \quad (3.9)$$

and

$$|\sigma_{k-1}(\lambda|1, i)| \sigma_{k-3}(\lambda|1, i) \leq C(n, k) \frac{\lambda_1 \cdots \lambda_{k-1} \lambda_k^2}{\lambda_1 \lambda_i} \frac{\lambda_1 \cdots \lambda_{k-1}}{\lambda_1 \lambda_i} \leq C(n, k) \left(\frac{\sigma_{k-1} \lambda_k}{\lambda_1 \lambda_i} \right)^2. \quad (3.10)$$

For any $k \leq p \leq n$, we have

$$\sigma_{k-2}(\lambda|1, p) \leq C(n, k) \frac{\lambda_1 \lambda_2 \cdots \lambda_{k-1}}{\lambda_1} \leq C(n, k) \frac{\sigma_{k-1}}{\lambda_1} \quad (3.11)$$

and

$$|\sigma_{k-1}(\lambda|1, p)| \sigma_{k-3}(\lambda|1, p) \leq C(n, k) \frac{\lambda_1 \cdots \lambda_{k-1} \lambda_k}{\lambda_1} \frac{\lambda_1 \cdots \lambda_{k-2}}{\lambda_1} \leq C(n, k) \frac{\sigma_{k-1}}{\lambda_1^2}. \quad (3.12)$$

Here we use Property (6) and Lemma 2.1 in (3.9), (3.10), (3.11) and (3.12). Then for any $1 < i \leq k-1$, (3.8) becomes

$$-2 \frac{(\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{ii} - \ddot{\sigma}_{k-1}^{11, ii} \sigma_{k-1}) \xi_1 \xi_i}{\sigma_{k-1}^2} \geq -2 \frac{C \lambda_k^2 |\xi_1 \xi_i|}{\lambda_1^2 \lambda_i^2} \geq -\frac{\epsilon \xi_1^2}{4(n-1) \lambda_1^2} - \frac{4(n-1) C^2 \lambda_k^4 \xi_i^2}{\epsilon \lambda_1^2 \lambda_i^4}. \quad (3.13)$$

Then for any $k \leq p \leq n$, (3.8) becomes

$$-2 \frac{(\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{pp} - \ddot{\sigma}_{k-1}^{11,pp} \sigma_{k-1}) \xi_1 \xi_p}{\sigma_{k-1}^2} \geq -2 \frac{C |\xi_1 \xi_p|}{\lambda_1^2} \geq -\frac{\varepsilon \xi_1^2}{4(n-1)\lambda_1^2} - \frac{4(n-1)C^2 \xi_p^2}{\varepsilon \lambda_1^2}. \quad (3.14)$$

Here we use Cauchy-Schwartz inequality in (3.13) and (3.14). Recall that

$$\dot{\sigma}_k^{11} = \lambda_\alpha \ddot{\sigma}_k^{11,\alpha\alpha} + \sigma_{k-1}(\lambda|1, \alpha), \quad (3.15)$$

$$\dot{\sigma}_k^{\alpha\alpha} = \lambda_1 \ddot{\sigma}_k^{11,\alpha\alpha} + \sigma_{k-1}(\lambda|1, \alpha). \quad (3.16)$$

If $\sigma_{k-1}(\lambda|1, \alpha) \leq 0$, then due to (3.15) $\lambda_\alpha > 0$ and

$$\frac{\lambda_\alpha}{\lambda_1} \dot{\sigma}_k^{\alpha\alpha} = \lambda_\alpha \ddot{\sigma}_k^{11,\alpha\alpha} + \frac{\lambda_\alpha}{\lambda_1} \sigma_{k-1}(\lambda|1, \alpha) \geq \dot{\sigma}_k^{11}. \quad (3.17)$$

If $\sigma_{k-1}(\lambda|1, \alpha) > 0$, using the Maclaurin's inequality, we have

$$\sigma_{k-1}(\lambda|1, \alpha)^{\frac{k-2}{k-1}} \leq \sigma_{k-2}(\lambda|1, \alpha). \quad (3.18)$$

Since

$$\sigma_{k-1}(\lambda|1, \alpha) \leq C \lambda_2 \lambda_3 \cdots \lambda_{k-1} \lambda_k \leq C \lambda_1^{k-2} C(A),$$

multiplying it with (3.18) we obtain

$$\sigma_{k-1}(\lambda|1, \alpha) \leq C \lambda_1^{-\frac{1}{k-1}} (\lambda_1 \ddot{\sigma}_k^{11,\alpha\alpha}).$$

Then using Property (4), we have

$$\left(\frac{\lambda_\alpha}{\lambda_1} + C \lambda_1^{-\frac{1}{k-1}}\right) \dot{\sigma}_k^{\alpha\alpha} \geq \left(\frac{\lambda_\alpha}{\lambda_1} + C \lambda_1^{-\frac{1}{k-1}}\right) \lambda_1 \ddot{\sigma}_k^{11,\alpha\alpha} \geq \dot{\sigma}_k^{11} \geq \frac{k\sigma_k}{n\lambda_1}, \quad \text{for } \lambda_\alpha > 0, \quad (3.19)$$

and

$$C \lambda_1^{-\frac{1}{k-1}} \dot{\sigma}_k^{\alpha\alpha} \geq \left(\frac{\lambda_\alpha}{\lambda_1} + C \lambda_1^{-\frac{1}{k-1}}\right) \lambda_1 \ddot{\sigma}_k^{11,\alpha\alpha} \geq \dot{\sigma}_k^{11} \geq \frac{k\sigma_k}{n\lambda_1}, \quad \text{for } \lambda_\alpha \leq 0. \quad (3.20)$$

We now divide it into two cases. We assume that there exists some positive constant C .

If $\lambda_\alpha \leq C \lambda_1^{\frac{k-2}{k-1}}$, then by (3.17), (3.19) and (3.20), we obtain that

$$C \lambda_1^{-\frac{1}{k-1}} \dot{\sigma}_k^{\alpha\alpha} \geq \frac{1}{\lambda_1}. \quad (3.21)$$

For any $k \leq p \leq n$, $|\lambda_p| \leq C(A)$ by Lemma 2.3. Then (3.14) becomes

$$-2 \frac{(\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{pp} - \ddot{\sigma}_{k-1}^{11,pp} \sigma_{k-1}) \xi_1 \xi_p}{\sigma_{k-1}^2} \geq -\frac{\varepsilon \xi_1^2}{4(n-1)\lambda_1^2} - \frac{C \frac{\partial \sigma_k}{\partial \lambda_p} \xi_p^2}{\varepsilon \lambda_1^{\frac{k}{k-1}}}. \quad (3.22)$$

For any $1 < i \leq k-1$ and $\lambda_i \leq C \lambda_1^{\frac{k-2}{k-1}}$, (3.13) becomes

$$-2 \frac{(\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{ii} - \ddot{\sigma}_{k-1}^{11,ii} \sigma_{k-1}) \xi_1 \xi_i}{\sigma_{k-1}^2} \geq -\frac{\varepsilon \xi_1^2}{4(n-1)\lambda_1^2} - \frac{4(n-1)C^2 \xi_i^2}{\varepsilon \lambda_1^2} \geq -\frac{\varepsilon \xi_1^2}{4(n-1)\lambda_1^2} - \frac{C \dot{\sigma}_k^{ii} \xi_i^2}{(\varepsilon \lambda_1^{\frac{1}{k-1}}) \lambda_1}. \quad (3.23)$$

If $\lambda_\alpha \geq C\lambda_1^{\frac{k-2}{k-1}}$, then by (3.17) and (3.19), we obtain that

$$C \frac{\lambda_i}{\lambda_1} \dot{\sigma}_k^{ii} \geq \frac{\sigma_k}{\lambda_1}. \quad (3.24)$$

For any $1 < i \leq k-1$ and $\lambda_i \geq C\lambda_1^{\frac{k-2}{k-1}}$, (3.13) becomes

$$-2 \frac{(\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{ii} - \ddot{\sigma}_{k-1}^{11,ii} \sigma_{k-1}) \xi_1 \xi_i}{\sigma_{k-1}^2} \geq -\frac{\varepsilon \xi_1^2}{4(n-1)\lambda_1^2} - \frac{C(A) \dot{\sigma}_k^{ii} \xi_i^2}{\varepsilon \lambda_1^2 \lambda_i^3} \geq -\frac{\varepsilon \xi_1^2}{4(n-1)\lambda_1^2} - \frac{C \dot{\sigma}_k^{ii} \xi_i^2}{(\varepsilon \lambda_1^{\frac{1}{k-1}}) \lambda_1}. \quad (3.25)$$

By direct calculation, (3.2) implies that

$$\begin{aligned} & - \sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + K \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} + 2 \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{(\lambda_1 + A + 1) \sigma_k} \\ & \geq -\frac{\partial_\xi^2 q_k}{q_k} + \left(1 + (1-\varepsilon) \frac{1}{k}\right) \left(\frac{\dot{\sigma}_{k-1}^{11} \xi_1}{\sigma_{k-1}}\right)^2 - \frac{\varepsilon \xi_1^2}{4 \lambda_1^2} + \left(2 - \frac{C}{\varepsilon \lambda_1^{\frac{1}{k-1}}} - \frac{C(A)}{\lambda_1^2}\right) \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{\lambda_1 \sigma_k} \\ & \geq -\frac{\partial_\xi^2 q_k}{q_k} + \left(1 + (1-\varepsilon) \frac{1}{k} - \varepsilon\right) \left(1 - \frac{C_1(A)}{\lambda_1}\right)^2 \frac{\xi_1^2}{\lambda_1^2} + \left(2 - \frac{C_2}{\varepsilon \lambda_1^{\frac{1}{k-1}}} - \frac{C_3(A)}{\lambda_1^2}\right) \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{\lambda_1 \sigma_k} \end{aligned} \quad (3.26)$$

where we use (3.7) in the last inequality. We choose $\varepsilon = \frac{1}{(k+1)^2}$ and assume that $\lambda_1 > \max\left\{(k+3)C_1(A), \frac{C_3(A)}{C_2(k+1)^2}\right\}$. Then (3.26) becomes

$$\begin{aligned} & - \sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + K \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} + 2 \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{\lambda_1 \sigma_k} \\ & \geq \frac{k+2}{k+1} \left(1 - \frac{C_1(A)}{\lambda_1}\right)^2 \frac{\xi_1^2}{\lambda_1^2} + \left(2 - \frac{C_2}{\varepsilon \lambda_1^{\frac{1}{k-1}}} - \frac{C_3(A)}{\lambda_1^2}\right) \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{\lambda_1 \sigma_k} \\ & \geq \frac{(k+2)^2}{(k+1)(k+3)} \frac{\xi_1^2}{\lambda_1^2} + \left(2 - 2(k+1)^2 \frac{C_2}{\lambda_1^{\frac{1}{k-1}}}\right) \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{\lambda_1 \sigma_k}. \end{aligned} \quad (3.27)$$

Recall that

$$\sigma_k = \lambda_1 \dot{\sigma}_k^{11} + \sigma_k(\lambda|1). \quad (3.28)$$

We note that if

$$\sigma_k(\lambda|1) \geq -\frac{\sigma_k}{2(k+2)^2 - 1}, \quad (3.29)$$

then (3.27) implies the conclusion (1.10) by assuming $\lambda_1 \geq (C_2(k+1)^2)^{k-1}$ and choosing $\delta_0 = \frac{1}{2(k+1)(k+3)}$.

In the subsequent content, we denote by $c_0 = \frac{1}{2(k+2)^2 - 1}$ and suppose that

$$\sigma_k(\lambda|1) < -c_0 \sigma_k. \quad (3.30)$$

Under this assumption we have following claims.

Claim 1:

$$\lambda_k \leq C(n, k)|\lambda_n|.$$

From (2.3),

$$\lambda_1 \cdots \lambda_k \leq 2\sigma_k \quad (3.31)$$

or

$$\lambda_1 \cdots \lambda_k \leq 2C(n, k)\lambda_1 \cdots \lambda_{k-1}|\lambda_n|. \quad (3.32)$$

If we assume that (3.31) holds, then

$$|\sigma_k(\lambda|1)| \leq C(n, k)\lambda_2 \cdots \lambda_k^2 \leq C'(n, k, A)\frac{\sigma_k}{\lambda_1} < c_0\sigma_k \quad (3.33)$$

by assuming $\lambda_1 > \frac{C'(n, k, A)}{c_0}$, which is contradict to our assumption above. Thus we complete the proof of the first claim.

Claim 2: there exists some $c_1, c_2 > 0$ such that $-\sigma_{k+1} \geq c_1\lambda_1 \cdots \lambda_{k-1}\lambda_k^2 \geq c_2\lambda_1$.

By Claim 1 and (3.33), we have

$$\lambda_1 \cdots \lambda_{k-1}\lambda_n^2 \geq c\lambda_1 \cdots \lambda_{k-1}\lambda_k^2 \geq \frac{cc_0\sigma_k}{C(n, k)}\lambda_1 \quad (3.34)$$

and denote $c_2 = \frac{cc_0}{C(n, k)}$. Since $\sigma_k(\lambda|1) < 0$, we have $\lambda_n < 0$ and from (3.4)

$$\sigma_k(\lambda|n) = \sigma_k - \lambda_n \dot{\sigma}_k^{nn} > 0. \quad (3.35)$$

Using Property (6), we get

$$\dot{\sigma}_k^{nn} > \lambda_1 \cdots \lambda_{k-1}. \quad (3.36)$$

We suppose that there exists some large $M > 0$ (to be chosen later) and $1 < l \leq k-1$, such that $\lambda_1 > M^k$, $\lambda_l > M$ and $\lambda_{l+1} \leq M$ in the following proof. By (3.4), (3.36) and Property (1), we have

$$\begin{aligned} \lambda_1 \cdots \lambda_{k-1}\lambda_n^2 &\leq \dot{\sigma}_k^{nn}\lambda_n^2 \leq \sum_{p=l+1}^n \dot{\sigma}_k^{pp}\lambda_p^2 \\ &= \sum_{i=1}^l \lambda_i(\lambda_i \dot{\sigma}_k^{ii} + \sigma_k(\lambda|i)) + \sum_{p=l+1}^n \lambda_p \sigma_k - \sum_{i=1}^l (\lambda_i \dot{\sigma}_{k+1}^{ii} + \sigma_{k+1}(\lambda|i)) \\ &\quad - (k+1-l)\sigma_{k+1} - \sum_{i=1}^l \dot{\sigma}_k^{ii}\lambda_i^2 \\ &= \sum_{p=l+1}^n \lambda_p \sigma_k - \sum_{i=1}^l \sigma_{k+1}(\lambda|i) - (k+1-l)\sigma_{k+1}. \end{aligned} \quad (3.37)$$

We now show that by using Lemma 2.1 for any $1 \leq i \leq k-1$

$$|\sigma_{k+1}(\lambda|i)| \leq \frac{C(n, k)\lambda_1 \cdots \lambda_{k-1}\lambda_k^3}{\lambda_i} \leq \frac{C_1(A)}{M}\lambda_1 \cdots \lambda_{k-1}\lambda_n^2. \quad (3.38)$$

Applying (3.34), we have

$$\sum_{p=l+1}^n \lambda_p \sigma_k \leq c_1 M^2 \leq \frac{c_1 \lambda_1}{M} \leq \frac{\lambda_1 \cdots \lambda_{k-1} \lambda_n^2}{M}. \quad (3.39)$$

Then by choosing $M \geq \max\{\frac{\sigma_k}{c_1}, 2(1 + (k-1)C_1)\}$, we obtain

$$-\sigma_{k+1} \geq \frac{\lambda_1 \cdots \lambda_{k-1} \lambda_n^2}{4}, \quad (3.40)$$

which complete the proof of *Claim 2*.

Claim 3: There exists some $c_3, c_4 > 0$ such that $c_3 \frac{\lambda_1 \cdots \lambda_k^2}{\lambda_1^2} \leq \dot{\sigma}_k^{11} \leq c_4 \frac{\lambda_1 \cdots \lambda_k^2}{\lambda_1^2}$.

Combining Claim 2, we have

$$-\sigma_k(\lambda|1) = \frac{-\sigma_{k+1} - \sigma_{k+1}(\lambda|1)}{\lambda_1} \leq \frac{-\sigma_{k+1} + C\lambda_2 \cdots \lambda_k^3}{\lambda_1} \leq \frac{-\sigma_{k+1}(1 + \frac{C}{\lambda_1})}{\lambda_1}. \quad (3.41)$$

For the same reason

$$-\sigma_k(\lambda|1) \geq \frac{-\sigma_{k+1}(1 - \frac{C}{\lambda_1})}{\lambda_1}. \quad (3.42)$$

Then by using the assumption $\sigma_k(\lambda|1) < -c_0$ and Claim 2, we have

$$\begin{aligned} \dot{\sigma}_k^{11} &= \frac{\sigma_k - \sigma_k(\lambda|1)}{\lambda_1} \geq C \frac{-\sigma_k(\lambda|1)}{\lambda_1} \\ &\geq c_4 \frac{\lambda_2 \cdots \lambda_k^2}{\lambda_1^2}. \end{aligned} \quad (3.43)$$

For the same reason, we have

$$\dot{\sigma}_k^{11} \leq c_3 \frac{\lambda_2 \cdots \lambda_k^2}{\lambda_1^2} \quad (3.44)$$

which complete the proof of *Claim 3*.

We denote by $\alpha = (\lambda_1, \dots, \lambda_n)$, $e_1 = (1, 0, \dots, 0)$ and $\gamma = (0, \gamma_2, \dots, \gamma_k, \dots, \gamma_n)$. Now we assume that

$$\xi = (1+a) \frac{\alpha \xi_1}{\lambda_1} - a \xi_1 e_1 + \gamma \quad (3.45)$$

and

$$(\gamma_k, \dots, \gamma_n) \perp (\lambda_k, \dots, \lambda_n). \quad (3.46)$$

Then by Lemma 2.9 the first term of the right side of (3.26) becomes

$$\begin{aligned} -\frac{\partial_\xi^2 q_k}{q_k} &= -(-a \xi_1 e_1 + \gamma)_p \frac{\ddot{q}_k^{pp,qq}}{q_k} (-a \xi_1 e_1 + \gamma)_q \\ &= -2a^2 \xi_1^2 \frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} \left(\frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} - \frac{\dot{\sigma}_k^{11}}{\sigma_k} \right) - \frac{\partial_\gamma^2 q_k}{q_k} \\ &\quad + 2a \xi_1 \gamma_\alpha \left(\frac{\ddot{\sigma}_k^{11,\alpha\alpha}}{\sigma_k} - \frac{\dot{\sigma}_k^{\alpha\alpha}}{\sigma_k} \frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} - \frac{\dot{\sigma}_k^{11}}{\sigma_k} \frac{\dot{\sigma}_{k-1}^{\alpha\alpha}}{\sigma_{k-1}} + 2 \frac{\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{\alpha\alpha}}{\sigma_{k-1}^2} - \frac{\ddot{\sigma}_{k-1}^{11,\alpha\alpha}}{\sigma_{k-1}} \right). \end{aligned} \quad (3.47)$$

By direct calculation, we obtain following equations:

$$\begin{aligned} \frac{\ddot{\sigma}_k^{11,\alpha\alpha}}{\sigma_k} - \frac{\dot{\sigma}_k^{\alpha\alpha} \sigma_{k-1}^{11}}{\sigma_k \sigma_{k-1}} &= \frac{\dot{\sigma}_k^{\alpha\alpha} - \sigma_{k-1}(\lambda|1\alpha)}{\lambda_1 \sigma_k} - \frac{\dot{\sigma}_k^{\alpha\alpha} \sigma_{k-1} - \sigma_{k-1}(\lambda|1)}{\sigma_k \lambda_1 \sigma_{k-1}} \\ &= -\frac{\sigma_{k-1}(\lambda|1\alpha)}{\lambda_1 \sigma_k} + \frac{\dot{\sigma}_k^{\alpha\alpha} \dot{\sigma}_k^{11}}{\sigma_k \lambda_1 \sigma_{k-1}}, \end{aligned} \quad (3.48)$$

and

$$\frac{\ddot{\sigma}_{k-1}^{11,\alpha\alpha}}{\sigma_{k-1}} - \frac{\dot{\sigma}_{k-1}^{\alpha\alpha} \dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}^2} = \frac{\sigma_{k-2}^2(\lambda|1\alpha) - \sigma_{k-1}(\lambda|1\alpha)\sigma_{k-3}(\lambda|1\alpha)}{\sigma_{k-1}^2}. \quad (3.49)$$

Hence, for any $2 \leq i \leq k-1$, we have

$$\left| \frac{\ddot{\sigma}_{k-1}^{11,ii}}{\sigma_{k-1}} - \frac{\dot{\sigma}_{k-1}^{ii} \dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}^2} \right| \leq \frac{C(n, k) \lambda_k^2}{\lambda_1^2 \lambda_i^2} \leq C \frac{\dot{\sigma}_k^{11} \lambda_k^2}{\lambda_1 \lambda_i^2}. \quad (3.50)$$

For any $k \leq p \leq n$, we have

$$\left| \frac{\ddot{\sigma}_{k-1}^{11,pp}}{\sigma_{k-1}} - \frac{\dot{\sigma}_{k-1}^{pp} \dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}^2} \right| \leq \frac{C(n, k)}{\lambda_1^2} \leq C \frac{\dot{\sigma}_k^{11}}{\lambda_1}. \quad (3.51)$$

Here we use Property (4), Claim 2 and Claim 3 in (3.50) and (3.51). Furthermore, for any $2 \leq i \leq k-1$, we have

$$\begin{aligned} &\frac{\ddot{\sigma}_k^{11,ii}}{\sigma_k} - \frac{\dot{\sigma}_k^{ii} \sigma_{k-1}^{11}}{\sigma_k \sigma_{k-1}} - \frac{\dot{\sigma}_k^{11} \dot{\sigma}_{k-1}^{ii}}{\sigma_k \sigma_{k-1}} + \frac{\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{ii}}{\sigma_{k-1}^2} \\ &= \frac{-\sigma_k + \dot{\sigma}_k^{11} \lambda_1 + \sigma_k(\lambda|1i)}{\lambda_1 \lambda_i \sigma_k} + \frac{\dot{\sigma}_k^{ii} \dot{\sigma}_k^{11}}{\sigma_k \lambda_1 \sigma_{k-1}} - \left(\frac{1}{\lambda_i} - \frac{\dot{\sigma}_k^{ii}}{\lambda_i \sigma_{k-1}} \right) \frac{\dot{\sigma}_k^{11}}{\sigma_k} \\ &\quad + \frac{(\sigma_{k-1} - \dot{\sigma}_k^{11})(\sigma_{k-1} - \dot{\sigma}_k^{ii})}{\lambda_1 \lambda_i \sigma_{k-1}^2} \\ &= \frac{\sigma_k(\lambda|1i)}{\lambda_1 \lambda_i \sigma_k} - \frac{\dot{\sigma}_k^{ii} \sigma_k(k|1)}{\lambda_1 \lambda_i \sigma_{k-1} \sigma_k} + \frac{\dot{\sigma}_k^{11}}{\lambda_1} \left(\frac{\dot{\sigma}_k^{ii}}{\sigma_{k-1} \sigma_k} - \frac{\sigma_{k-2}(\lambda|i)}{\sigma_{k-1}^2} \right). \end{aligned} \quad (3.52)$$

Here

$$\left| \frac{\sigma_k(\lambda|1i)}{\lambda_1 \lambda_i \sigma_k} \right| \leq \frac{C \lambda_1 \dots \lambda_k^3}{\lambda_1^2 \lambda_i^2 \sigma_k} \leq \frac{C \dot{\sigma}_k^{11} \lambda_k}{\lambda_i^2}. \quad (3.53)$$

By using (3.42), we have

$$\left| \frac{\dot{\sigma}_k^{ii} \sigma_k(k|1)}{\lambda_1 \lambda_i \sigma_{k-1} \sigma_k} \right| \leq C \frac{\lambda_1 \dots \lambda_k^2 \dot{\sigma}_k^{11}}{\lambda_i^2 \sigma_{k-1} \sigma_k} \leq C \frac{\lambda_k^2 \dot{\sigma}_k^{11}}{\lambda_i^2}. \quad (3.54)$$

Similarly, combining Claim 2 we have

$$\left| \frac{\sigma_{k-1}(\lambda|i)}{\sigma_{k-1} \sigma_k} - \frac{\sigma_{k-2}(\lambda|i)}{\sigma_{k-1}^2} \right| \leq C \frac{\lambda_1 \dots \lambda_k}{\lambda_i \sigma_{k-1} \sigma_k} + C \frac{\lambda_1 \dots \lambda_{k-1}}{\lambda_i \sigma_{k-1}^2} \leq \frac{C \lambda_k}{\lambda_i} \left(1 + \frac{\lambda_k}{\lambda_1} \right), \quad (3.55)$$

since

$$\sigma_{k-1} > \lambda_1 \dots \lambda_{k-1} = \frac{\lambda_1 \dots \lambda_k^2}{\lambda_k^2} \geq C \frac{\lambda_1}{\lambda_k^2}. \quad (3.56)$$

Inserting (3.53), (3.54) and (3.55) into (3.52) and combining (3.50) we have

$$\begin{aligned} & 2a\xi_1\gamma_i \left(\frac{\ddot{\sigma}_k^{11,ii}}{\sigma_k} - \frac{\dot{\sigma}_k^{ii}}{\sigma_k} \frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} - \frac{\dot{\sigma}_k^{11}}{\sigma_k} \frac{\dot{\sigma}_{k-1}^{ii}}{\sigma_{k-1}} + 2 \frac{\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{ii}}{\sigma_{k-1}^2} - \frac{\ddot{\sigma}_{k-1}^{11,ii}}{\sigma_{k-1}} \right) \\ & \geq -2C|a\xi_1\gamma_i| \frac{\dot{\sigma}_k^{11} \lambda_k}{\lambda_i^2} \geq -\epsilon \frac{\dot{\sigma}_k^{11} \lambda_1^2}{\lambda_i^3} \gamma_i^2 - \frac{Ca^2 \dot{\sigma}_k^{11} \xi_1^2}{\epsilon \lambda_1^2} \end{aligned} \quad (3.57)$$

where we use $\lambda_i > \lambda_k$, $\lambda_1 \gg 1$ and $\lambda_k < C(A)$ by Lemma 2.3 in the last inequality.

For any $k \leq p \leq n$, we have

$$\begin{aligned} & \frac{\ddot{\sigma}_k^{11,pp}}{\sigma_k} - \frac{\dot{\sigma}_k^{pp}}{\sigma_k} \frac{\sigma_{k-1}^{11}}{\sigma_{k-1}} - \frac{\dot{\sigma}_k^{11}}{\sigma_k} \frac{\dot{\sigma}_{k-1}^{pp}}{\sigma_{k-1}} + \frac{\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{pp}}{\sigma_{k-1}^2} \\ & = \frac{-\sigma_k + \dot{\sigma}_k^{pp} \lambda_p + \sigma_k(\lambda|1p)}{\lambda_1^2 \sigma_k} + \frac{\dot{\sigma}_k^{pp}}{\sigma_k} \frac{\dot{\sigma}_k^{11}}{\lambda_1 \sigma_{k-1}} + \frac{\dot{\sigma}_{k-1}^{pp}}{\sigma_{k-1}} \left(\frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} - \frac{\dot{\sigma}_k^{11}}{\sigma_k} \right) \\ & = \frac{\sigma_k(\lambda|1p) - \sigma_k}{\lambda_1^2 \sigma_k} + \frac{\sigma_{k-1} \lambda_p - \sigma_{k-2}(\lambda|p) \lambda_p^2}{\lambda_1^2 \sigma_k} + \frac{\dot{\sigma}_k^{pp}}{\sigma_k} \frac{\dot{\sigma}_k^{11}}{\lambda_1 \sigma_{k-1}} + \frac{\dot{\sigma}_{k-1}^{pp}}{\sigma_{k-1}} \left(\frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} - \frac{\dot{\sigma}_k^{11}}{\sigma_k} \right). \end{aligned} \quad (3.58)$$

By using Claim 2 and Claim 3, we have

$$\frac{|\sigma_k(\lambda|1p) - \sigma_k|}{\lambda_1^2 \sigma_k} \leq \frac{C\lambda_2 \cdots \lambda_k^2}{\lambda_1^2} + \frac{1}{\lambda_1^2} \leq C' \frac{\dot{\sigma}_k^{11}}{\lambda_1}, \quad (3.59)$$

$$\frac{\sigma_{k-2}(\lambda|p) \lambda_p^2}{\lambda_1^2 \sigma_k} \leq \frac{C\lambda_1 \cdots \lambda_{k-2} \lambda_k^2}{\lambda_1^2 \sigma_k} \leq C' \frac{\dot{\sigma}_k^{11}}{\lambda_{k-1}}, \quad (3.60)$$

and

$$\frac{\dot{\sigma}_k^{pp}}{\sigma_k} \frac{\dot{\sigma}_k^{11}}{\lambda_1 \sigma_{k-1}} \leq C \frac{\lambda_1 \cdots \lambda_{k-1}}{\sigma_{k-1}} \frac{\dot{\sigma}_k^{11}}{\lambda_1} \leq C \frac{\dot{\sigma}_k^{11}}{\lambda_1}. \quad (3.61)$$

Besides,

$$\left| \frac{\dot{\sigma}_{k-1}^{pp}}{\sigma_{k-1}} \left(\frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} - \frac{\dot{\sigma}_k^{11}}{\sigma_k} \right) \right| \leq C \frac{\lambda_1 \cdots \lambda_{k-2} \dot{\sigma}_k^{11}}{\sigma_{k-1}} \leq C \frac{\dot{\sigma}_k^{11}}{\lambda_{k-1}}. \quad (3.62)$$

Inserting (2.15), (3.60), (3.59), (3.61) and (3.62) into (3.58) and combining (3.51) we obtain

$$2a\xi_1\gamma_p \left(\frac{\ddot{\sigma}_k^{11,pp}}{\sigma_k} - \frac{\dot{\sigma}_k^{pp}}{\sigma_k} \frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} - \frac{\dot{\sigma}_k^{11}}{\sigma_k} \frac{\dot{\sigma}_{k-1}^{pp}}{\sigma_{k-1}} + 2 \frac{\dot{\sigma}_{k-1}^{11} \dot{\sigma}_{k-1}^{pp}}{\sigma_{k-1}^2} - \frac{\ddot{\sigma}_{k-1}^{11,pp}}{\sigma_{k-1}} \right) \quad (3.63)$$

$$\geq -2C \frac{\dot{\sigma}_k^{11}}{\lambda_{k-1}} |a\gamma_p \xi_1| \geq -\epsilon \frac{\dot{\sigma}_k^{11} \lambda_1^2 \gamma_p^2}{\lambda_{k-1} \lambda_k^2} - \frac{Ca^2 \dot{\sigma}_k^{11} \xi_1^2}{\epsilon \lambda_1^2} \quad (3.64)$$

where we use the fact that $\lambda_1 > \lambda_{k-1} > \lambda_k$ and $\lambda_k < C(A)$ by Lemma 2.3 in the last inequality. Using (3.4), we estimate the first term of (3.47):

$$\begin{aligned} -2a^2 \xi_1^2 \frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} \left(\frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} - \frac{\dot{\sigma}_k^{11}}{\sigma_k} \right) & = -2a^2 \xi_1^2 \frac{\dot{\sigma}_{k-1}^{11}}{\sigma_{k-1}} \left(\frac{\sigma_{k-1} - \dot{\sigma}_k^{11}}{\lambda_1 \sigma_{k-1}} - \frac{\sigma_k - \sigma_k(\lambda|1)}{\lambda_1 \sigma_k} \right) \\ & \geq \frac{2a^2 \xi_1^2}{\lambda_1} \left(1 - \frac{C}{\lambda_1^2} \right) \left(\frac{\lambda_1 \dot{\sigma}_k^{11} - \sigma_k}{\lambda_1 \sigma_k} \right) \end{aligned} \quad (3.65)$$

where we use Claim 2, Claim 3 and

$$\frac{\dot{\sigma}_k^{11}}{\lambda_1 \sigma_{k-1}} \leq C \frac{\lambda_1 \dots \lambda_k^2}{\lambda_1^3 \sigma_{k-1}} \leq \frac{C(A)}{\lambda_1^3} \leq -C'(A) \frac{\sigma_k(\lambda|1)}{\lambda_1^2} \quad (3.66)$$

in the last inequality.

Applying Lemma 2.8 and inserting (3.2), (3.57) and (3.63) into (3.47), we obtain

$$\begin{aligned} -\frac{\partial_\xi^2 q_k}{q_k} &\geq \frac{2a^2 \xi_1^2}{\lambda_1} \left(1 - \frac{C}{\lambda_1^2}\right) \left(\frac{\lambda_1 \dot{\sigma}_k^{11} - \sigma_k}{\lambda_1 \sigma_k}\right) + \sum_{i=2}^{k-1} (C_1 - \epsilon) \frac{\dot{\sigma}_k^{11} \lambda_1^2}{\lambda_i^3} \gamma_i^2 \\ &\quad + \sum_{p=k}^n (C_1 - \epsilon) \frac{\dot{\sigma}_k^{11} \lambda_1^2 \gamma_p^2}{\lambda_{k-1} \lambda_k^2} - (n-1) \frac{Ca^2 \dot{\sigma}_k^{11} \xi_1^2}{\epsilon \lambda_1^2} \\ &\geq \frac{2a^2 \xi_1^2}{\lambda_1} \left(1 - \frac{C}{\lambda_1^2}\right) \left(\frac{\lambda_1 \dot{\sigma}_k^{11} - \sigma_k}{\lambda_1 \sigma_k}\right) + \sum_{i=2}^{k-1} \frac{C_1 \dot{\sigma}_k^{11} \lambda_1^2}{2 \lambda_i^3} \gamma_i^2 + \frac{C_1 \dot{\sigma}_k^{11} \lambda_1^2 \gamma_p^2}{2 \lambda_{k-1} \lambda_k^2} \\ &\quad - 2(n-1) \frac{Ca^2 \dot{\sigma}_k^{11} \xi_1^2}{C_1 M \lambda_1} \end{aligned}$$

by choosing $\epsilon = \frac{C_1}{2}$. Then combining (3.26), (3.27) and (3.67), we have

$$\begin{aligned} &-\sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + K \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} + 2 \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{\lambda_1 \sigma_k} \\ &\geq \frac{2a^2 \xi_1^2}{\lambda_1} \left(1 - \frac{C}{\lambda_1^2}\right) \left(\frac{\lambda_1 \dot{\sigma}_k^{11} - \sigma_k}{\lambda_1 \sigma_k}\right) + \sum_{i=2}^{k-1} \frac{C_1 \dot{\sigma}_k^{11} \lambda_1^2}{2 \lambda_i^3} \gamma_i^2 + \sum_{p=k}^n \frac{C_1 \dot{\sigma}_k^{11} \lambda_1^2 \gamma_p^2}{2 \lambda_{k-1} \lambda_k^2} \\ &\quad - 2(n-1) \frac{Ca^2 \dot{\sigma}_k^{11} \xi_1^2}{C_1 M \lambda_1} + \frac{(k+2)^2}{(k+1)(k+3)} \left(1 - \frac{C(A)}{\lambda_1}\right)^2 \frac{\xi_1^2}{\lambda_1^2} \\ &\quad + \left(2 - \frac{C(k, A)}{M}\right) \sum_{l+1 \leq p \leq n} \frac{\sigma_k^{pp} \xi_p^2}{\lambda_1 \sigma_k}. \end{aligned} \quad (3.67)$$

Under the assumptions in (3.37), we consider

$$\begin{aligned} \sum_{l+1 \leq p \leq n} \frac{\sigma_k^{pp} \xi_p^2}{\lambda_1 \sigma_k} &= \sum_{l+1 \leq p \leq n} \frac{\sigma_k^{pp} \left((1+a) \frac{\lambda_p \xi_1}{\lambda_1} + \gamma_p \right)^2}{\lambda_1 \sigma_k} \\ &\geq (1+a)^2 \xi_1^2 \sum_{l+1 \leq p \leq n} \frac{\sigma_k^{pp} \lambda_p^2}{\lambda_1^3 \sigma_k} - 2 \left| (1+a) \xi_1 \sum_{l+1 \leq p \leq n} \frac{\sigma_k^{pp} \lambda_p \gamma_p}{\lambda_1^2 \sigma_k} \right| \\ &\geq (1+a)^2 \xi_1^2 \sum_{l+1 \leq p \leq n} \frac{\sigma_k^{pp} \lambda_p^2}{\lambda_1^3 \sigma_k} - 2 \left| (1+a) \xi_1 \sum_{k \leq p \leq n} \frac{\sigma_{k-2}(\lambda|p) \lambda_p^2 \gamma_p}{\lambda_1^2 \sigma_k} \right| \\ &\quad - 2 \sum_{l+1 \leq i \leq k-1} \left| (1+a) \xi_1 \frac{\sigma_k^{ii} \lambda_i \gamma_i}{\lambda_1^2 \sigma_k} \right| \end{aligned} \quad (3.68)$$

where we use (3.4) and (3.46), which implies that

$$\sum_{k \leq p \leq n} \dot{\sigma}_k^{pp} \lambda_p \gamma_p = (\sigma_{k-1} - \sigma_{k-2}(\lambda|p) \lambda_p) \lambda_p \gamma_p = -\sigma_{k-2}(\lambda|p) \lambda_p^2 \gamma_p \quad (3.69)$$

in the last inequality. Then by (3.37), (3.38), (3.39), and (3.41), we have

$$\begin{aligned} \sum_{l+1 \leq p \leq n} \frac{\sigma_k^{pp} \lambda_p^2}{\lambda_1^3 \sigma_k} &= \frac{\sum_{p=l+1}^n \lambda_p \sigma_k - \sum_{i=1}^l \sigma_{k+1}(\lambda|i) - (k+1-l) \sigma_{k+1}}{\lambda_1^3 \sigma_k} \\ &\geq -(k+1-l) \frac{\sigma_{k+1}}{\lambda_1^3 \sigma_k} - \frac{C(A) \lambda_1 \cdots \lambda_k^2}{M \lambda_1^3 \sigma_k} \\ &\geq (k+1-l) \left(1 - \frac{c}{\lambda_1}\right) \frac{\lambda_1 \dot{\sigma}_k^{11} - \sigma_k}{\lambda_1 \sigma_k} - \frac{C(A)}{M} \frac{\dot{\sigma}_k^{11}}{\lambda_1 \sigma_k}. \end{aligned} \quad (3.70)$$

Furthermore,

$$\begin{aligned} 2 \left| (1+a) \xi_1 \sum_{k \leq p \leq n} \frac{\sigma_{k-2}(\lambda|p) \lambda_p^2 \gamma_p}{\lambda_1^2 \sigma_k} \right| &\leq 2 |(1+a) \xi_1 \gamma_p| \frac{C \lambda_1 \cdots \lambda_{k-2} \lambda_k^2}{\lambda_1^2 \sigma_k} \\ &\leq 2 |(1+a) \xi_1 \gamma_p| \frac{C \dot{\sigma}_k^{11}}{\lambda_{k-1} \sigma_k} \\ &\leq -\frac{C_1 \dot{\sigma}_k^{11} \lambda_1^2 \gamma_p^2}{2 \lambda_{k-1} \lambda_k^2} - \frac{2}{C_1} \frac{C'(1+a)^2 \dot{\sigma}_k^{11} \xi_1^2}{M \lambda_1} \end{aligned} \quad (3.71)$$

and for any $l+1 \leq i \leq k-1$

$$\begin{aligned} 2 \left| (1+a) \xi_1 \frac{\sigma_{k-1}(\lambda|i) \lambda_i \gamma_i}{\lambda_1^2 \sigma_k} \right| &= 2 \left| (1+a) \xi_1 \frac{(\sigma_k - \sigma_k(\lambda|i)) \gamma_i}{\lambda_1^2 \sigma_k} \right| \\ &\leq 2 |(1+a) \xi_1 \gamma_i| \frac{C \lambda_1 \lambda_2 \cdots \lambda_{k-1} \lambda_k^2}{\lambda_1^2 \lambda_i \sigma_k} \\ &\leq 2 |(1+a) \xi_1 \gamma_i| \frac{C \dot{\sigma}_k^{11}}{\lambda_i} \\ &\leq -\frac{C_1 \dot{\sigma}_k^{11} \lambda_1^2 \gamma_i^2}{2 \lambda_i^3} - \frac{2}{C_1} \frac{C'(1+a)^2 \dot{\sigma}_k^{11} \xi_1^2}{M \lambda_1} \end{aligned} \quad (3.72)$$

where we use $\lambda_{k-1} \geq \lambda_k$ and $\lambda_k < C(A)$ by Lemma 2.3, the assumptions $\kappa_i \leq M$ and $\kappa_1 > M^2$ in Claim 2. Thus (3.67) implies that

$$\begin{aligned}
& - \sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + K \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} + 2 \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{\lambda_1 \sigma_k} \\
& \geq \frac{2a^2 \xi_1^2}{\lambda_1} \left(1 - \frac{C}{M}\right) \left(\frac{\lambda_1 \dot{\sigma}_k^{11} - \sigma_k}{\lambda_1 \sigma_k}\right) - 2(n-1) \frac{C' a^2 \dot{\sigma}_k^{11} \xi_1^2}{C_1 M \lambda_1} \\
& \quad + \frac{(k+2)^2}{(k+1)(k+3)} \frac{\xi_1^2}{\lambda_1^2} + \left(2 - \frac{C(k,A)}{M}\right) (k+1-l)(1+a)^2 \frac{\lambda_1 \dot{\sigma}_k^{11} - \sigma_k}{\lambda_1 \sigma_k} \\
& \quad - \frac{C(n,k,A)(1+a)^2 \dot{\sigma}_k^{11} \xi_1^2}{M \lambda_1} \\
& \geq \frac{2(a^2 + 2(1+a)^2) \xi_1^2}{\lambda_1} \left(1 - \frac{C_2(k,A)}{M}\right) \left(\frac{\lambda_1 \dot{\sigma}_k^{11} - \sigma_k}{\lambda_1 \sigma_k}\right) + \frac{(k+2)^2}{(k+1)(k+3)} \frac{\xi_1^2}{\lambda_1^2} \\
& \quad - C_3(n,k,A)(1+a^2) \frac{\dot{\sigma}_k^{11} \xi_1^2}{M \lambda_1} \\
& \geq \frac{8(a^2 + 2(1+a)^2) \xi_1^2}{5\lambda_1} \left(\frac{\lambda_1 \dot{\sigma}_k^{11} - \sigma_k}{\lambda_1 \sigma_k}\right) + \frac{(k+2)^2}{(k+1)(k+3)} \frac{\xi_1^2}{\lambda_1^2} \\
& \quad + 2 \frac{(1+a)^2 \xi_1^2 \sigma_k(\lambda|1)}{5\lambda_1^2 \sigma_k} - C_3(n,k,A)(1+a^2) \frac{\dot{\sigma}_k^{11} \xi_1^2}{M \lambda_1} \\
& \geq \frac{8}{5} (a^2 + 2(1+a)^2) \frac{\dot{\sigma}_k^{11} \xi_1^2}{\lambda_1 \sigma_k} + \left(\frac{(k+2)^2}{(k+1)(k+3)} - \frac{8}{5} (a^2 + 2(1+a)^2)\right) \frac{\xi_1^2}{\lambda_1^2}
\end{aligned} \tag{3.73}$$

by assuming $M = \max\{10C_2, \frac{5C_3(c_0+1)}{2c_0}\}$ depending on n, k and A where c_0 obtained from (3.30).

Since under the assumption $\sigma_k(\lambda|1) < -c_0 \sigma_k$, (3.4) implies that

$$\frac{\dot{\sigma}_k^{11}}{\sigma_k} = \frac{1}{\lambda_1} - \frac{\sigma_k(\lambda|1)}{\lambda_1 \sigma_k} > \frac{c_0 + 1}{\lambda_1}. \tag{3.74}$$

Thus (3.73) implies that

$$- \sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + K \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} + 2 \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{(\lambda_1 + A + 1) \sigma_k} > \left(1 + \frac{1}{(k+1)(k+3)}\right) \frac{\dot{\sigma}_k^{11} \xi_1^2}{\lambda_1 \sigma_k} \tag{3.75}$$

or

$$- \sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + K \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} + 2 \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{(\lambda_1 + A + 1) \sigma_k} \geq \frac{16}{15} \frac{\dot{\sigma}_k^{11} \xi_1^2}{\lambda_1 \sigma_k}. \tag{3.76}$$

Hence we obtain (1.10) and complete the proof. \square

Following the proof in Lemma 1.1, we also improve the results in [13, 16].

Lemma 3.1. *Assume $\{\lambda_i\} \in \Gamma_k$, $\lambda_1 \geq \lambda_2 \cdots \geq \lambda_n$, and $\lambda_n > -A$ for some constant $A > 0$. For any $\delta_0 \in (0, 1)$, there exists a constant $\delta_0 > 0$ and a large constant $C > 0$ depending on*

n, k, σ_k , and A , such that if $k_1 \geq C$, then the following inequality holds

$$-\sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + K \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} + 2 \sum_{i>1} \frac{\sigma_k^{ii} \xi_i^2}{(\lambda_1 + A + 1)\sigma_k} \geq (1 + \delta_0) \frac{\xi_1^2}{\lambda_1^2}, \quad (3.77)$$

for some sufficient large K (depending on δ_0) where $\xi = (\xi_1, \dots, \xi_n)$ is an arbitrary vector in \mathbb{R}^n .

4. INTERIOR ESTIMATES OF k -HESSIAN EQUATIONS WITH DIRICHLET BOUNDARY CONDITIONS

Somewhere in the sequel, we denote σ_k by F for brevity and define

$$\dot{F}^{pq}(D^2u) := \frac{\partial F}{\partial u_{pq}}, \quad \ddot{F}^{pq,rs}(D^2u) := \frac{\partial^2 F}{\partial u_{qp} \partial u_{rs}}.$$

If we further assume that $\{u_{ij}\}$ is diagonal at a certain point, then we define

$$\frac{\partial F}{\partial \lambda_p}(\lambda(D^2u)) = \dot{F}^{pp}(D^2u), \quad \frac{\partial^2 F}{\partial \lambda_p \partial \lambda_q}(\lambda(D^2u)) = \ddot{F}^{pp,qq}(D^2u).$$

We first proof Theorem 1.1.

Proof. we consider the function for any $\varepsilon > 0, x \in \Omega, \xi \in \mathbb{S}^{n-1}$

$$\tilde{P}(x, \xi) = \beta \log(-u) + \log \max\{u_{\xi\xi}(x), 1\} + \frac{L}{2}|Du|^2,$$

where $L \gg 1, \beta > 0$ are constants to be determined later. Clearly, the maximum of \tilde{P} is attained in some interior point x_0 of Ω and some $\xi(x_0) \in \mathbb{S}^{n-1}$. Choose canonical coordinate frames e_1, \dots, e_n at x_0 such that $\xi(x_0) = e_1$ and $\{u_{ij}(x_0)\}$ is diagonal. Let the eigenvalues λ of D^2u be ordered as

$$\lambda_1(x_0) \geq \lambda_2(x_0) \geq \dots \geq \lambda_n(x_0).$$

We construct a unit vector field $v(s)$ near p such that $v(0) = e_1(x_0)$ and $v'(0) = \sum_{p \neq 1} \frac{u_{1pi}}{\lambda_1 + 1 - \lambda_p} e_p$. We may also assume that $\lambda_1(x_0) > 1$ is sufficiently large. Then we consider the function

$$P(x) = \beta \log(-u) + \log(u(v(s), v(s)) + (g(v(s), e_1(s)))^2 - 1) + \frac{L}{2}|Du|^2.$$

Note that x_0 is also a maximum point of P . At x_0 , we have

$$0 = \nabla_i P(x_0) = \beta \frac{u_i}{u} + \frac{u_{11i}}{u_{11}} + Lu_i \lambda_i \quad (4.1)$$

and

$$\begin{aligned} 0 &\geq \nabla_{ij} P(x_0) \\ &= \beta \frac{u_{ij}}{u} - \beta \frac{u_i u_j}{u^2} + \frac{u_{11ij}}{u_{11}} + \sum_{l>1} \frac{u_{1li}^2}{u_{11}(u_{11} - u_l + 1)} \\ &\quad - \frac{u_{11i} u_{11j}}{u_{11}^2} + Lu_{ki} u_{kj} + Lu_{kij} u_k. \end{aligned} \quad (4.2)$$

By $\sigma_k(D^2u) = f(x, u, \nabla u)$, we have at x_0

$$\nabla_l \sigma_k = f_l + f_u u_l + f_{u_l} u_l \quad (4.3)$$

and

$$\nabla_{11}^2 \sigma_k = \dot{F}^{ij} u_{ij11} + \ddot{F}^{pq,rs} u_{pq1} u_{rs1} = f_{11}. \quad (4.4)$$

Besides, there exists some constant $C > 0$ depending on $\|f\|_{C^2}$ and $\|u\|_{C^1}$ such that

$$|f_1| \leq C + C u_{11}, \quad f_{11} \geq -C - C u_{11}^2 + f_{u_l} u_{11l}. \quad (4.5)$$

Multiplying (4.2) by using \dot{F}^{ij} , by (4.1), (4.3) and (4.4) we obtain

$$\begin{aligned} 0 &\geq \beta \frac{kF}{u} - \beta \frac{\dot{F}^{ii} u_i^2}{u^2} + \frac{\dot{F}^{ii} u_{11ii}}{u_{11}} + 2 \sum_{p>1} \frac{\dot{F}^{ii} u_{1pi}}{u_{11}(u_{11} - \lambda_p + 1)} - \frac{\dot{F}^{ii} u_{11i}^2}{u_{11}^2} + L \dot{F}^{ii} u_{ii}^2 + L \nabla_l \sigma_k u_l \\ &\geq \beta \frac{kF}{u} - \beta \frac{\dot{F}^{ii} u_i^2}{u^2} - \frac{\dot{F}^{pq,rs} u_{pq1} u_{rs1}}{u_{11}} + \frac{f_{u_l} u_{11l}}{u_{11}} + 2 \sum_{p>1} \frac{\dot{F}^{ii} u_{1pi}}{u_{11}(u_{11} - u_{pp} + 1)} \\ &\quad - \frac{\dot{F}^{ii} u_{11i}^2}{u_{11}^2} + L \dot{F}^{ii} u_{ii}^2 - C u_{11} + L f_{u_l} \lambda_l u_l - CL \\ &\geq \beta \frac{kF}{u} - \beta \frac{\dot{F}^{ii} u_i^2}{u^2} - \frac{\dot{F}^{pq,rs} u_{pq1} u_{rs1}}{u_{11}} + 2 \sum_{p>1} \frac{\dot{F}^{ii} u_{1pi}}{u_{11}(u_{11} - u_{pp} + 1)} - \frac{\dot{F}^{ii} u_{11i}^2}{u_{11}^2} + L \dot{F}^{ii} u_{ii}^2 \\ &\quad - \beta \frac{f_{u_l} u_l}{u} - C u_{11} - CL. \end{aligned} \quad (4.6)$$

Referring to Andrews [1], we have

$$\begin{aligned} -\ddot{F}^{pq,rs} h_{pq1} h_{rs1} &= -\ddot{F}^{pp,qq} h_{pp1} h_{qq1} + 2 \sum_{p>q} \ddot{F}^{pp,qq} h_{pq1}^2 \\ &\geq -\ddot{F}^{pp,qq} h_{pp1} h_{qq1} + 2 \sum_{\lambda_p < \lambda_1} \frac{\dot{F}^{pp} - \dot{F}^{11}}{\lambda_1 - \lambda_p} h_{11p}^2. \end{aligned} \quad (4.7)$$

Besides,

$$\sum_{1 < p, 1 \leq i \leq n} \frac{2 \dot{F}^{ii} u_{1pi}^2}{(u_{11} + 1 - u_{pp})} \geq 2 \sum_{p>1} \frac{\dot{F}^{11} u_{11p}^2}{(u_{11} - u_{pp} + 1)} + 2 \sum_{i>1} \frac{\dot{F}^{ii} u_{1ii}^2}{(u_{11} - u_{ii} + 1)}. \quad (4.8)$$

Using the critical equation (4.1), for any $i \geq 2$ we have

$$-\beta \frac{\dot{F}^{ii} u_i^2}{u^2} = -\frac{1}{\beta} \left(\frac{u_{11i}}{u_{11}} + L u_i u_{ii} \right)^2 \geq -\frac{2}{\beta} \frac{\dot{F}^{ii} u_{11i}^2}{u_{11}^2} - 2 \frac{L^2}{\beta} \dot{F}^{ii} u_i^2 u_{ii}^2. \quad (4.9)$$

Then we choose $\beta = L^2 \gg 4$ and obtain

$$\begin{aligned}
 & -\frac{\dot{F}^{pq,rs}u_{pq1}u_{rs1}}{u_{11}} + 2\sum_{p>1}\frac{\dot{F}^{ii}u_{1pi}}{u_{11}(u_{11}-u_{pp}+1)} - \left(1 + \frac{2}{\beta}\right)\sum_{i\geq 2}\frac{\dot{F}^{ii}u_{11i}^2}{u_{11}^2} - \frac{\dot{F}^{11}u_{111}^2}{u_{11}^2} \\
 & \geq -\ddot{F}^{pp,qq}h_{pp1}h_{qq1} + 2\sum_{p>1}\frac{\dot{F}^{pp}}{u_{11}(u_{11}-u_{pp}+1)}h_{11p}^2 + 2\sum_{i>1}\frac{\dot{F}^{ii}u_{1ii}^2}{u_{11}(u_{11}-u_{ii}+1)} \\
 & \quad - \frac{3}{2}\sum_{i\geq 2}\frac{\dot{F}^{ii}u_{11i}^2}{u_{11}^2} - \frac{\dot{F}^{11}u_{111}^2}{u_{11}^2} \tag{4.10} \\
 & = -\ddot{F}^{pp,qq}h_{pp1}h_{qq1} + 2\sum_{i>1}\frac{\dot{F}^{ii}u_{1ii}^2}{u_{11}(u_{11}+A+1)} - \frac{\dot{F}^{11}u_{111}^2}{u_{11}^2} - \frac{\dot{F}^{ii}u_{11i}^2(\frac{1}{2}u_{11}-\frac{3}{2}A-\frac{3}{2})}{u_{11}^2(u_{11}-u_{pp}+1)} \\
 & \geq -\ddot{F}^{pp,qq}h_{pp1}h_{qq1} + 2\sum_{i>1}\frac{\dot{F}^{ii}u_{1ii}^2}{u_{11}(u_{11}+A+1)} - \frac{\dot{F}^{11}u_{111}^2}{u_{11}^2}
 \end{aligned}$$

by assuming $\lambda_1 \geq 3A + 3$ in the last inequality.

Due to Lemma 1.1, combining (4.7) and (4.8), (4.6) becomes

$$\begin{aligned}
 0 & \geq -L^2\frac{\dot{F}^{11}u_1^2}{u^2} - 2\dot{F}^{ii}u_i^2\lambda_i^2 - \frac{\dot{F}^{pp,qq}u_{pp1}u_{qq1}}{u_{11}} + K\frac{(\nabla_1\sigma_k)^2}{u_{11}\sigma_k} + 2\sum_{i>1}\frac{\dot{F}^{ii}u_{1ii}^2}{u_{11}(u_{11}+A+1)} - \frac{\dot{F}^{11}u_{111}^2}{u_{11}^2} \\
 & \quad + L\dot{F}^{ii}u_{ii}^2 - \frac{C_3L^2}{u} - C_4L - C_5(K)u_{11} \\
 & \geq \left(\frac{Lu_{11}^2}{5} - \frac{C_1L^2}{u^2}\right)\dot{F}^{11} + \left(\frac{L}{5} - C_2\right)\sum_{i=1}^n\dot{F}^{ii}u_{ii}^2 + L\left(\frac{k\sigma_k}{5n}u_{11} - C_3\frac{L}{u}\right) \\
 & \quad + L\left(\frac{k\sigma_k}{5n}u_{11} - C_4\right) + \left(\frac{Lk\sigma_k}{5n} - C_5(K)\right)u_{11}. \tag{4.11}
 \end{aligned}$$

where we use Property (4) in the last inequality. In (4.11), C_i , $i = 1, 2, \dots, 5$ all depends on n , k , $\|f\|_{C^2}$ and $\|u\|_{C^1}$. We choose $L = \max\{4, 5C_2, \frac{5nC_5}{kf}\}$ such that the second term and the last term in the last inequality of (4.11) are positive. Then we obtain $\max \lambda_1 u^\beta \leq \max\{\sqrt{5C_1L} \sup u^{\beta-1}, \frac{5nLC_3 \sup u^{\beta-1}}{kf_0}, \frac{5nC_4 \sup u^\beta}{kf_0}\}$ and complete the proof. \square

5. A LIOUVILLE PROBLEM FOR GENERAL k -HESSIAN EQUATION

We first consider the interior C^2 estimates for the following Dirichlet problem:

$$\begin{cases} \sigma_k(D^2u) = 1 & x \in \Omega, \\ u = 0 & x \in \partial\Omega. \end{cases} \tag{5.1}$$

Lemma 5.1. *Assume that $D^2u \geq -AI$ for some positive constant $A > 0$. Then, for any k -convex solution $u \in C^2(\Omega) \cap C^{0,1}(\bar{\Omega})$, there exists some $\alpha > 1$ and $C > 0$ depending on n , k , $|u|_{C^1}$ and the diameter of Ω , such that*

$$\max_{x \in \Omega} (-u(x))^\alpha |D^2u(x)| \leq C. \tag{5.2}$$

Proof. By using the Comparison principle (see [3, 10] for detail), there exists large a and b depending on the diameter of Ω , such that

$$w = a|x|^2 - b$$

and

$$w \leq u \leq 0.$$

Thus, $-u$ can be controlled by the diameter of the domain Ω .

Now we consider the function for $x \in \Omega$, $\xi \in \mathbb{S}^{n-1}$

$$\tilde{P}(x, \xi) = \alpha \log(-u) + \log \max\{u_{\xi\xi}(x), 1\} + \frac{1}{2}|x|^2,$$

where $\alpha > k - 1$ is a constant to be determined later. Clearly, the maximum of \tilde{P} is attained in some interior point x_0 of Ω and some $\xi(x_0) \in \mathbb{S}^{n-1}$. Choose canonical coordinate frames e_1, \dots, e_n at x_0 such that $\xi(x_0) = e_1$ and $\{u_{ij}(x_0)\}$ is diagonal. Let the eigenvalues λ of D^2u be ordered as

$$\lambda_1(x_0) \geq \lambda_2(x_0) \geq \dots \geq \lambda_n(x_0).$$

We construct a unit vector field $v(s)$ near p such that $v(0) = e_1(x_0)$ and $v'(0) = \sum_{p \neq 1} \frac{u_{1pi}}{\lambda_1 + 1 - \lambda_p} e_p$. We may also assume that $\lambda_1(x_0) > 1$ is sufficiently large. Then we consider the function

$$P(x) = \alpha \log(-u) + \log(u(v(s), v(s)) + (g(v(s), e_1(s)))^2 - 1) + \frac{1}{2}|x|^2.$$

Note that x_0 is also a maximum point of P . At the maximum point x_0 ,

$$0 = P_i = \frac{\alpha u_i}{u} + \frac{u_{11i}}{u_{11}} + x_i, \quad (5.3)$$

and

$$P_{ij} = \frac{\alpha u_{ij}}{u} - \frac{\alpha u_i u_j}{u^2} + \frac{u_{11ij}}{u_{11}} + \sum_{p>1} \frac{2u_{1pi}^2}{u_{11}(u_{11} + 1 - u_{pp})} - \frac{u_{11i}u_{11j}}{u_{11}^2} + \delta_{ij}. \quad (5.4)$$

Multiplying (5.4) by \dot{F}^{ij} , at x_0 we have

$$0 \geq \alpha \frac{kF}{u} - \alpha \frac{\dot{F}^{ii}u_i^2}{u^2} + \frac{\dot{F}^{ii}u_{11ii}}{u_{11}} + 2 \sum_{p>1} \frac{\dot{F}^{ii}u_{1pi}^2}{u_{11}(u_{11} + 1 - u_{pp})} - \frac{\dot{F}^{ii}u_{11i}^2}{u_{11}^2} + (n - k + 1)\sigma_{k-1}. \quad (5.5)$$

Similar to the proof in Theorem 1.1, using the critical equation (5.3) we obtain at x_0

$$-\alpha \frac{\dot{F}^{ii}u_i^2}{u^2} = -\frac{1}{\alpha} \dot{F}^{ii} \left(\frac{u_{11i}}{u_{11}} + x_i \right)^2 \geq -\frac{2}{\alpha} \frac{\dot{F}^{ii}u_{11i}^2}{u_{11}^2} - \frac{2}{\alpha} \dot{F}^{ii}x_i^2. \quad (5.6)$$

Taking the derivatives of the equation (5.1), we obtain

$$\dot{F}^{ij}u_{ijk} = 0,$$

and

$$\ddot{F}^{ij,kl}u_{kl1}u_{ij1} + \dot{F}^{ij}u_{ij11} = 0.$$

By (4.7) and (4.8), similar to (4.10) we have by choosing $\alpha \gg 4$

$$\begin{aligned} & -\frac{\dot{F}^{pq,rs}u_{pq1}u_{rs1}}{u_{11}} + 2\sum_{p>1}\frac{\dot{F}^{ii}u_{1pi}}{u_{11}(u_{11}-u_{pp}+1)} - \left(1 + \frac{2}{\alpha}\right)\sum_{i\geq 2}\frac{\dot{F}^{ii}u_{11i}^2}{u_{11}^2} - \frac{\dot{F}^{11}u_{111}^2}{u_{11}^2} \\ & \geq -\ddot{F}^{pp,qq}h_{pp1}h_{qq1} + 2\sum_{i>1}\frac{\dot{F}^{ii}u_{1ii}^2}{u_{11}(u_{11}+A+1)} - \frac{\dot{F}^{11}u_{111}^2}{u_{11}^2} \end{aligned} \quad (5.7)$$

by assuming $\lambda_1 \geq 3A + 3$ in the last inequality. Then applying Lemma 1.1, we obtain

$$\begin{aligned} 0 & \geq \alpha \frac{kF}{u} - \frac{2}{\alpha} \dot{F}^{ii} X_i^2 - \ddot{F}^{pp,qq} h_{pp1} h_{qq1} + 2 \sum_{i>1} \frac{\dot{F}^{ii} u_{1ii}^2}{u_{11}(u_{11} + A + 1)} \\ & \quad - \left(1 + \frac{2}{\alpha}\right) \frac{\dot{F}^{11} u_{111}^2}{u_{11}^2} + (n - k + 1) \sigma_{k-1} \\ & \geq \alpha \frac{kF}{u} + \left(\delta_0 - \frac{2}{\alpha}\right) \frac{\dot{F}^{11} u_{111}^2}{u_{11}^2} + \left(n - k + 1 - \frac{2 \sup_{\Omega} |X|^2}{\alpha}\right) \sigma_{k-1} \\ & \geq \frac{C_1 \alpha}{u} + \frac{n - k + 1}{2} \sigma_{k-1} \end{aligned} \quad (5.8)$$

by choosing $\alpha = \max\{4, k - 1, \frac{2}{\delta_0}, \frac{4 \sup_{\Omega} |X|^2}{n - k + 1}\}$. Thus, we employ Lemma 2.4 to obtain that there exists some $C > 0$ only depending on n and the diameter of Ω , such that

$$C(-u)u_{11}^{\frac{1}{k-1}} \leq (-u)\sigma_{k-1} \leq \frac{2C_1\alpha}{n - k + 1}. \quad (5.9)$$

Then we complete the proof. \square

We now begin to prove Theorem 1.2.

Proof. The proof is standard (refer to [15, 22] for details). Let u be an entire solution of equation (5.1). We consider the domain

$$\Omega_R = \{y \in \mathbb{R}^n : u(Ry) < R^2\}$$

for any large $R > 0$. Let

$$u_R(y) = \frac{u(Ry) - R^2}{R^2}.$$

Then

$$D_y^2 u_R = D_x^2 u. \quad (5.10)$$

Hence u_R satisfying the following Dirichlet problem:

$$\begin{cases} \sigma_2(D^2 u_R(x)) = 1 & x \in \Omega_R, \\ u_R = 0 & x \in \partial\Omega_R. \end{cases} \quad (5.11)$$

The quadratic growth condition in Theorem 1.2 implies that

$$a|Ry|^2 - b \leq u(Ry) \leq R^2.$$

Then

$$|y|^2 \leq \frac{1+b}{a},$$

which means that Ω_R is bounded. Applying Lemma 1.1, we obtain

$$(-u_R)^\alpha |D^2 u_R| \leq C \quad (5.12)$$

where C is independent of R . Now we consider the domain

$$\Omega'_R = \{y \in \mathbb{R}^n : u(Ry) < \frac{R^2}{2}\} \subset \Omega_R.$$

In Ω'_R , we have

$$u_R(y) \leq -\frac{1}{2}.$$

Due to (5.10) and (5.12),

$$D_x^2 u = |D^2 u_R| \leq 2^\alpha C.$$

The estimate above holds true for all R . Applying Evans-Krylov theory we have

$$|D^2 u|_{C^\alpha(B_R)} \leq C(n, \alpha) \frac{|D^2 u|_{C^0(B_{2R})}}{R^\alpha} \leq \frac{C(n, \alpha)}{R^\alpha}.$$

Hence, we obtain our theorem by letting $R \rightarrow +\infty$. \square

6. GLOBAL CURVATURE ESTIMATE FOR CONVEX HYPERSURFACES

In this section, we focus on the semi-convex hypersurfaces with the prescribed k -th mean curvature and complete the proof of Theorem 1.4 and Theorem 1.5.

Proof of Theorem 1.4 We consider the test function

$$Q = \log \kappa_1 - A \log u$$

where A is a large constant to be considered. Further, we assume Q attains its maximum at point X_0 . Similar as the proof in [8], also in the proof in Theorem 1.1 and Theorem 1.2, we consider the perturbed quantity

$$\tilde{Q} = \log \tilde{\kappa}_1 - Au,$$

which also attains its maximum at point X_0 .

At X_0 , we have

$$0 = \frac{h_{11i}}{\kappa_1} - Ah_{ii} \langle X, \partial_i \rangle \quad (6.1)$$

and due to Lemma 3.1 in [8]

$$\begin{aligned} 0 \geq & 2 \sum_{p>1} \frac{\dot{F}^{11,pp} h_{11p}^2}{\kappa_1^2} + 2 \sum_{i>1} \frac{\dot{F}^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i + 1)} - \frac{\ddot{F}^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} + 2 \sum_{p>1} \frac{\dot{F}^{pp} h_{1pp}^2}{\kappa_1(\kappa_1 - \kappa_p + 1)} \\ & - \sum_{p=1}^n \frac{\dot{F}^{pp} h_{11p}^2}{\kappa_1^2} + \sum_{i=1}^n \left(\frac{A}{C} - C \right) \dot{F}^{ii} \kappa_i^2 - CA. \end{aligned} \quad (6.2)$$

Then

$$\begin{aligned}
 & 2 \sum_{p>1} \frac{\dot{F}^{11,pp} h_{11p}^2}{\kappa_1^2} + 2 \sum_{i>1} \frac{\dot{F}^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i + 1)} - \sum_{p=1}^n \frac{\dot{F}^{pp} h_{11p}^2}{\kappa_1^2} \\
 & \geq 2 \sum_{i>1} \frac{\dot{F}^{ii} h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i + 1)} - \sum_{p=1}^n \frac{\dot{F}^{pp} h_{11p}^2}{\kappa_1^2} \\
 & \geq \sum_{i>1} \frac{\kappa_1 + \kappa_i - 1}{\kappa_1 - \kappa_i + 1} \frac{\dot{F}^{ii} h_{11i}^2}{\kappa_1} - \frac{\dot{F}^{11} h_{111}^2}{\kappa_1^2}.
 \end{aligned} \tag{6.3}$$

According to Lemma 11 in [19], we have

$$-\kappa_i \leq \frac{n-k}{k} \kappa_1$$

which implies that

$$\kappa_1 + \kappa_i - 1 = \frac{2k-n}{k} \kappa_1 - 1 > 0 \tag{6.4}$$

by assuming that $2k > n$ and $\kappa_1 \gg 1$. Or under the assumption of Theorem 1.4 that $\kappa_i \geq -A$, we have

$$\kappa_1 + \kappa_i - 1 \geq \kappa_1 - A - 1 > 0 \tag{6.5}$$

by assuming $\kappa_1 \geq A + 1$ rather than the condition $2k > n$. Plugging (6.3) and (6.5) into (6.2), we have

$$\begin{aligned}
 0 & \geq -\frac{\ddot{F}^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} + 2 \sum_{p>1} \frac{\dot{F}^{pp} h_{1pp}^2}{\kappa_1(\kappa_1 - \kappa_p + 1)} - \frac{\dot{F}^{11} h_{111}^2}{\kappa_1^2} + \sum_{i=1}^n \left(\frac{A}{C} - C \right) \dot{F}^{ii} \kappa_i^2 - CA \\
 & \geq \sum_{i=1}^n \left(\frac{A}{C} - C \right) \dot{F}^{ii} \kappa_i^2 - K \frac{(\dot{F}^{11} h_{111})^2}{\sigma_k \kappa_1} - CA.
 \end{aligned} \tag{6.6}$$

Here we use the crucial Lemma 1.1. Differentiating (1.11), we have

$$\sum_i \dot{F}^{ii} h_{ii1} = h_{11} d_\nu f(e_1) + d_x f(e_1) \tag{6.7}$$

which implies that

$$\frac{\sum_i (\dot{F}^{ii} h_{ii1})^2}{\sigma_k \kappa_1} \leq C \kappa_1. \tag{6.8}$$

Then combining Property (4), we have

$$0 \geq \sum_{i=1}^n \left(\frac{kA}{nC} - \frac{k}{n} C - CK \right) \kappa_1 - CA, \tag{6.9}$$

which implies κ_1 is bounded above by some constant $C > 0$ only depending on $n, k, \|M\|_{C^1}, \inf f$ and $\|f\|_{C^2}$. \square

Remark 6.1. If we further assume $2k > n$, following the proof in Theorem 1.4 we can derive the global C^2 estimates of the k -convex hypersurface satisfying (1.11) without the assumption $\kappa_i > -A$ under the following conjecture.

Conjecture 1.5 Suppose that $2k > n$, and $\kappa = (\kappa_1, \dots, \kappa_n) \in \Gamma_k$. If we assume that $\kappa_1 \geq \dots \geq \kappa_n$ and κ_1 is large enough, then for an arbitrary n -vector ξ in \mathbb{R}^n and some sufficient large $K > 0$, the following inequality holds

$$\kappa_1 \left(K(\dot{F}^{jj}\xi_j)^2 - \ddot{\sigma}_k^{pp,qq}\xi_p\xi_q \right) - \dot{F}^{11}\xi_1^2 + (\dot{F}^{jj} + (\kappa_1 + \kappa_j)\ddot{\sigma}_k^{11,jj}) \geq 0. \quad (6.10)$$

One can refer to [23] for detailed proof. We note that this conjecture is a weakened version of Conjecture 2 by in [19] but exhibits the same capability in deriving the global C^2 estimates of the curvature equation (1.11). Our crucial Lemma 1.1 is a semi-convexity version of Conjecture 1.5.

Finally, we provide a concise proof of Theorem 1.5 using the crucial Lemma 1.1 when $k = 2$.

Sketch proof of Theorem 1.5 We consider the test function for some $N \gg 1$ and small $\varepsilon > 0$

$$Q = \log \kappa_1 - \varepsilon u + \frac{N}{2}|x|^2 \quad (6.11)$$

in this case. We assume that Q attains its maximum at point X_0 . Similar to the proof in Theorem 1.4 using the idea in [8], we disturb Q as below

$$\tilde{Q} = \log \tilde{\kappa}_1 - \varepsilon u + \frac{N}{2}|x|^2 \quad (6.12)$$

which also attains its maximum at X_0 . At X_0 , we have

$$0 = \frac{h_{11i}}{\kappa_1} - \varepsilon \frac{h_{ii}\langle X, \partial_i \rangle}{u} + Nx_i \quad (6.13)$$

and

$$\begin{aligned} 0 &\geq 2 \sum_{p>1} \frac{\dot{F}^{11,pp}h_{11p}^2}{\kappa_1^2} + 2 \sum_{i>1} \frac{\dot{F}^{11}h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i + 1)} - \frac{\ddot{F}^{pp,qq}h_{pp1}h_{qq1}}{\kappa_1} + 2 \sum_{p>1} \frac{\dot{F}^{pp}h_{1pp}^2}{\kappa_1(\kappa_1 - \kappa_p + 1)} \\ &\quad - \sum_{p=1}^n \frac{\dot{F}^{pp}h_{11p}^2}{\kappa_1^2} + \sum_{i=1}^n \varepsilon \dot{F}^{ii}\kappa_i^2 + N \sum_i \dot{F}^{ii} - 2NuF - C\kappa_1 - C \\ &\geq - \frac{\ddot{F}^{pp,qq}h_{pp1}h_{qq1}}{\kappa_1} + 2 \sum_{p>1} \frac{\dot{F}^{pp}h_{1pp}^2}{\kappa_1(\kappa_1 - \kappa_p + 1)} + 2 \sum_{i>1} \frac{\dot{F}^{ii}h_{11i}^2}{\kappa_1(\kappa_1 - \kappa_i + 1)} - \sum_{p=1}^n \frac{\dot{F}^{pp}h_{11p}^2}{\kappa_1^2} \\ &\quad + \sum_{i=1}^n \varepsilon \dot{F}^{ii}\kappa_i^2 + N(n-1)\sigma_1 - C\kappa_1 - CN. \end{aligned} \quad (6.14)$$

If we further assume that $-\kappa_i < A$ for any $1 < i \leq n$, then applying Lemma 1.1 similar to the proof in (6.6) we have

$$\begin{aligned} 0 &\geq - \frac{\ddot{F}^{pp,qq}h_{pp1}h_{qq1}}{\kappa_1} + 2 \sum_{p>1} \frac{\dot{F}^{pp}h_{1pp}^2}{\kappa_1(\kappa_1 - \kappa_p + 1)} + K \frac{(\dot{F}^{ii}h_{ii1})^2}{\sigma_k \kappa_1} - \frac{\dot{F}^{11}h_{111}^2}{\kappa_1^2} \\ &\quad + \sum_{i=1}^n \varepsilon \dot{F}^{ii}\kappa_i^2 + N(n-1)\sigma_1 - (C + CK)\kappa_1 - CN \\ &\geq (N(n-1) - C - CK)\kappa_1 - CN. \end{aligned} \quad (6.15)$$

By choosing N large enough (independent of A) we obtain κ_1 is bounded above by some constant C . Both N and C depends only on $n, k, \|f\|_{C^2}$ and $\|M\|_{C^1}$. If $|\kappa_n| \geq A$, then (6.14) becomes

$$\begin{aligned}
 0 &\geq -\frac{\ddot{F}^{pp,qq}h_{pp1}h_{qq1}}{\kappa_1} + \frac{(\dot{F}^{ii}h_{ii1})^2}{\sigma_k\kappa_1} - \sum_{p=1}^n \frac{\dot{F}^{pp}h_{11p}^2}{\kappa_1^2} + \sum_{i=1}^n \varepsilon \dot{F}^{ii}\kappa_i^2 + N(n-1)\sigma_1 - C\kappa_1 - CN \\
 &\geq -2C\varepsilon^2 \sum_{p=1}^n \dot{F}^{pp}\kappa_p^2 - 2N^2 \sum_{p=1}^n \dot{F}^{pp}X_p^2 + \sum_{i=1}^n \varepsilon \dot{F}^{ii}\kappa_i^2 + N(n-1)\sigma_1 - C\kappa_1 - CN \\
 &\geq \left(-2C\varepsilon^2 + \frac{\varepsilon}{2}\right) \sum_{p=1}^n \dot{F}^{pp}\kappa_p^2 + (-2nN^2|X|^2 + \frac{\varepsilon}{2}\kappa_n^2)\dot{F}^{nn} + N(n-1)\sigma_1 - C\kappa_1 - CN.
 \end{aligned} \tag{6.16}$$

where we employ (3.1) in the first inequality and the critical equation (6.13) in the second inequality. By setting $\varepsilon = \frac{1}{4c}$ and $A = 4N\sqrt{nC} \sup |X|$, the proof can be completed in the way similar to the one presented above.

REFERENCES

- [1] Ben Andrews, *Pinching estimates and motion of hypersurfaces by curvature functions*, J. Reine Angew. Math. **608** (2007), 17–33.
- [2] Jiguang Bao, Jingyi Chen, Bo Guan, and Min Ji, *Liouville property and regularity of a Hessian quotient equation*, Amer. J. Math. **125** (2003), no. 2, 301–316.
- [3] L. Caffarelli, L. Nirenberg, and J. Spruck, *The Dirichlet problem for nonlinear second-order elliptic equations. III. Functions of the eigenvalues of the Hessian*, Acta Math. **155** (1985), no. 3-4, 261–301.
- [4] Sun-Yung Alice Chang and Yu Yuan, *A Liouville problem for the sigma-2 equation*, Discrete Contin. Dyn. Syst. **28** (2010), no. 2, 659–664.
- [5] Chuanqiang Chen, *Optimal concavity of some Hessian operators and the prescribed σ_2 curvature measure problem*, Sci. China Math. **56** (2013), no. 3, 639–651.
- [6] Shiu Yuen Cheng and Shing-Tung Yau, *Complete affine hypersurfaces. I. The completeness of affine metrics*, Comm. Pure Appl. Math. **39** (1986), no. 6, 839–866.
- [7] Kai-Seng Chou and Xu-Jia Wang, *A variational theory of the Hessian equation*, Comm. Pure Appl. Math. **54** (2001), no. 9, 1029–1064.
- [8] Jianchun Chu, *A simple proof of curvature estimate for convex solution of k -Hessian equation*, Proc. Amer. Math. Soc. **149** (2021), no. 8, 3541–3552.
- [9] Jianchun Chu and Sławomir Dinew, *Liouville theorem for a class of Hessian equations*, available at [arXiv:2306.13825](https://arxiv.org/abs/2306.13825).
- [10] Gilbarg David and Neil S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, second editon, Springer, 1998.
- [11] Pengfei Guan and Guohuan Qiu, *Interior C^2 regularity of convex solutions to prescribing scalar curvature equations*, Duke Math. J. **168** (2019), no. 9, 1641–1663.
- [12] Pengfei Guan, Changyu Ren, and Zhizhang Wang, *Global C^2 -estimates for convex solutions of curvature equations*, Comm. Pure Appl. Math. **68** (2015), no. 8, 1287–1325.
- [13] Han Hong and Ruijia Zhang, *Curvature estimates for semi-convex solutions of asymptotic Plateau problem in \mathbb{H}^{n+1}* , available at [arXiv:2408.09428](https://arxiv.org/abs/2408.09428).
- [14] Gerhard Huisken and Carlo Sinestrari, *Convexity estimates for mean curvature flow and singularities of mean convex surfaces*, Acta Math. **183** (1999), no. 1, 45–70.
- [15] Ming Li, Changyu Ren, and Zhizhang Wang, *An interior estimate for convex solutions and a rigidity theorem*, J. Funct. Anal. **270** (2016), no. 7, 2691–2714.
- [16] Siyuan Lu, *Curvature estimates for semi-convex solutions of Hessian equations in hyperbolic space*, Calc. Var. Partial Differential Equations **62** (2023), no. 9, Paper No. 257, 23.
- [17] A.V. Pogorelov, *The Minkowski Multidimensional Problem*, John Wiley, 1978.

- [18] Changyu Ren and Zhizhang Wang, *On the curvature estimates for Hessian equations*, Amer. J. Math. **141** (2019), no. 5, 1281–1315.
- [19] ———, *The global curvature estimate for the $n - 2$ Hessian equation*, Calc. Var. Partial Differential Equations **62** (2023), no. 9, Paper No. 239, 50.
- [20] Ravi Shankar and Yu Yuan, *Rigidity for general semiconvex entire solutions to the sigma-2 equation*, Duke Math. J. **171** (2022), no. 15, 3201–3214.
- [21] Joel Spruck and Ling Xiao, *A note on star-shaped compact hypersurfaces with prescribed scalar curvature in space forms*, Rev. Mat. Iberoam. **33** (2017), no. 2, 547–554.
- [22] Neil S. Trudinger and Xu-Jia Wang, *The Monge-Ampere equation and its geometric applications*, Handbook of geometric analysis, 2008, pp. 467-524.
- [23] Qiang Tu, *Pogorelov type estimates for semi-convex solutions of Hessian equations and related rigidity theorems*, available at [arXiv:2405.02939](https://arxiv.org/abs/2405.02939).
- [24] Miccah Warren, *Nonpolynomial entire solutions to σ_k equations*, Comm. Partial Differential Equations **41** (2016), no. 5, 848–853.
- [25] Fengrui Yang, *Prescribed curvature measure problem in hyperbolic space*, Comm. Pure Appl. Math. **77** (2024), no. 1, 863–898.

KEY LABORATORY OF PURE AND APPLIED MATHEMATICS SCHOOL OF MATHEMATICAL SCIENCES,
PEKING UNIVERSITY, BEIJING 100871, P. R. CHINA

Email address: zhangrj@pku.edu.cn