

Improved thresholds for e-values

Christopher Blier-Wong* Ruodu Wang†

August 22, 2024

Abstract

The rejection threshold used for e-values and e-processes is by default set to $1/\alpha$ for a guaranteed type-I error control at α , based on Markov’s and Ville’s inequalities. This threshold can be wasteful in practical applications. We discuss how this threshold can be improved under additional distributional assumptions on the e-values; some of these assumptions are naturally plausible and empirically observable, without knowing explicitly the form or model of the e-values. For small values of α , the threshold can roughly be improved (divided) by a factor of 2 for decreasing or unimodal densities, and by a factor of e for decreasing or unimodal-symmetric densities of the log-transformed e-value. Moreover, we propose to use the supremum of comonotonic e-values, which is shown to preserve the type-I error guarantee. We also propose some preliminary methods to boost e-values in the e-BH procedure under some distributional assumptions while controlling the false discovery rate. Through a series of simulation studies, we demonstrate the effectiveness of our proposed methods in various testing scenarios, showing enhanced power.

Keywords: Markov’s inequality, e-variables, unimodality, calibrators, false discovery rate

1 Introduction

In statistical hypothesis testing, p-values and e-values summarize the evidence from a test against a null hypothesis. E-values have recently gained attention as a promising alternative to p-values, with inherent advantages over methods based on p-values, including validity in optional sampling and stopping, the ability to provide an intuitive notion of statistical significance, robustness to model misspecification, and flexibility within multiple testing procedures under dependence. We refer to Wasserman et al. [28], Shafer [20], Vovk and Wang [24], Wang and Ramdas [26] and Grünwald et al. [9] for advantages and recent developments on e-values, and to Ramdas et al. [18] for a summary.

Classic testing methods typically use a p-value with a significance level $\alpha \in (0, 1)$, representing the allowed type-I error. A statistician often starts with a test statistic and designs a rejection region such that the probability of that test statistic belonging to the rejection region, under the null hypothesis, is at most α . A simple analogous rejection region for e-values is $[1/\alpha, \infty)$ to guarantee a type-I error control of α ; the lower bound of the rejection region is based on Markov’s inequality. We will refer to an e-test as a test that rejects when an e-value exceeds a certain threshold. Using a threshold of $1/\alpha$ for e-tests is often wasteful in practice since this threshold is typically too conservative, meaning that the type-I error will be much smaller than the desired level α , resulting in a loss of power. By selectively using known information about the distribution or dependence structure of e-values, we can improve the power of e-tests while preserving type-I error control, striking a balance between safety and efficiency.

The paper’s main objective is to provide valid thresholds for e-values, under some distributional assumptions, that control the type-I error of e-tests. Including some distributional assumptions enables one to construct larger rejection regions (lower rejection thresholds), improving the power of e-tests while retaining some desirable properties and putting e-values on a more comparable scale to p-values.

If one knows the true distribution of the e-variable under the null, one may design an e-test to obtain exact type-I error control by inverting the cumulative distribution function of the e-variable; this would be

*Department of Statistical Sciences, University of Toronto, Canada. christopher.blierwong@utoronto.ca.

†Department of Statistics and Actuarial Science, University of Waterloo, Canada. wang@uwaterloo.ca.




Figure 1: Histogram of null knockoffs, which are e-values up to a constant. The left is based on knockoffs where all the nulls in the dataset are true; the right is based on knockoffs where 10% of variables are non-nulls.

equivalent to an approach based on p-values. However, some e-variables are constructed in a complicated way where the precise distribution of the e-variable is difficult to study, but we may be confident about some features. We provide two examples below. In both cases, the detailed construction of the e-values is described in the appendices.

Suppose that we have a high-dimensional dataset and wish to identify informative variables for a target response through variable selection. A flexible approach to accomplish this task is the model-X knockoffs (see Barber and Candès [3] and Candès et al. [6]), which control the false discovery rate (FDR). There are many ways to construct knockoffs, but the resulting distribution of the knockoffs is not easy to identify. Ren and Barber [19] showed that knockoffs are e-values in a slightly more general sense, called compound e-values by Ignatiadis et al. [13]. We may improve the threshold of e-tests on knockoffs with distributional information on the shape of their distribution. In Figure 1, we present a histogram of 10,000 knockoffs from linear regression when the number of variables is 80 and the number of non-null variables is 0 (left) and 8 (right). In both cases, the knockoffs appear to have a single mode, implying that the e-values also have a single mode.

Universal inference, proposed by Wasserman et al. [28], is a powerful method for developing hypothesis tests that provide finite-sample guarantees on the type-I error. This technique is especially useful for situations where classical p-tests are not feasible, such as when dealing with composite null hypotheses in irregular statistical models, provided that a maximum likelihood estimator can be computed. Assume we have iid observations $X_1, \dots, X_n \sim Q$ and want to test the composite null $Q = N(0, 1)$ against the mixture $Q = 0.5N(\mu_1, 1) + 0.5N(\mu_2, 1)$ for unknown (μ_1, μ_2) . A split likelihood-ratio test statistic may be used to construct a valid e-test; see Wasserman et al. [28]. In Figure 2, we present a histogram of 10,000 e-values and log-transformed e-values. Even if the shape of the histogram of e-values seems complicated, we can observe that its density is decreasing for e-values larger than one, and the density of the log-transformed e-values is decreasing for log e-values larger than zero. With this distributional knowledge, we may improve the e-test's threshold without knowing the null e-variable's true distribution. Tse and Davison [22] also propose a method to improve the threshold of universal inference tests based on the asymptotic distribution of universal inference e-values, but the finite-sample type-I error control is not guaranteed theoretically.

The rest of the paper is dedicated to obtaining improved thresholds that can be useful for situations like the above two examples and many others. Section 2 provides preliminaries on e-values and their thresholds, as well as some basic results. Section 3 contains our main results on the improved thresholds for e-values under various conditions, including decreasing and unimodal densities, as well as conditions on densities or random variables after a log-transform. As most useful results, for small values of α , the threshold can roughly be improved (divided) by a factor of 2 for decreasing (only required on $[1, \infty)$) or unimodal densities,




Figure 2: Histogram of universal inference e-values and log e-values. The red line is at 1 on the left plot and 0 on the right plot, and the density decreases after the red lines.

and by a factor of $e = \exp\{1\}$ for decreasing (only required on $[0, \infty)$) or unimodal-symmetric densities of the log-transformed e-variable.

In Section 4, we propose tests with valid type-I errors based on taking the supremum over a set of e-variables. Section 5 provides some improved thresholds for combined e-variables. In Section 6, we discuss methods to boost e-values in the e-BH procedure for multiple hypothesis testing with distributional assumptions. We admit that results in this section are not very practical as they require quite strong assumptions; they mainly serve as an exploration of possible improvements in multiple testing, a topic of great challenge. Section 7 includes simulation studies that illustrate the improved power we obtain using the improved thresholds. Section 8 concludes. We keep a few intuitive proofs in the main text, whereas additional proofs and other results are presented in the appendices.

2 Thresholds for e-values: preliminaries

2.1 E-variables and thresholds for e-tests

We first describe the basic setting and terminology of our paper. Our terminology follows Vovk and Wang [23]. Throughout, fix a sample space. A hypothesis is a collection H of probability measures on the sample space. An *e-variable* E for a hypothesis H is a $[0, \infty]$ -valued random variable satisfying $\mathbb{E}^Q[E] \leq 1$ for all $Q \in H$. For the general background on e-values in hypothesis testing, see Shafer [20], Vovk and Wang [23] and Grünwald et al. [9]. E-variables are often obtained from stopping an *e-process* $(E_t)_{t \geq 0}$, which is a nonnegative stochastic process adapted to a pre-specified filtration such that $\mathbb{E}^Q[E_\tau] \leq 1$ for any stopping time τ and any $Q \in H$. Realizations of e-variables are referred to as e-values.

We will also encounter p-values and p-variables. A *p-variable* P for a hypothesis H is a random variable that satisfies $Q(P \leq \alpha) \leq \alpha$ for all $\alpha \in (0, 1)$ and all $Q \in H$.

We fix a probability measure \mathbb{P} . All e-variables and p-variables are for the hypothesis $\{\mathbb{P}\}$, and the expectation \mathbb{E} is with respect to \mathbb{P} . To study properties and thresholds of e-variables and p-variables, such an assumption is without loss of generality; see the explanation in Vovk and Wang [23]. Denote by \mathcal{E}_0 the set of all e-variables. All terms like “increasing” and “decreasing” are in the non-strict sense. We write $a \wedge b = \min\{a, b\}$ and $a \vee b = \max\{a, b\}$ for real numbers a, b .

Below, we describe the main quantity of interest, the worst-case type-I error for a given set \mathcal{E} of e-variables. The set \mathcal{E} will be assumed to satisfy some distributional conditions to be specified later. Our goal

is to find the quantity $R_\gamma(\mathcal{E})$ for $\gamma > 0$, defined by

$$R_\gamma(\mathcal{E}) = \sup_{E \in \mathcal{E}} \mathbb{P}(E \geq 1/\gamma),$$

that is, the largest probability that $\mathbb{P}(E \geq 1/\gamma)$ can attain for $E \in \mathcal{E}$. The Markov inequality gives

$$\mathbb{P}(E \geq 1/\alpha) \leq \alpha \quad \text{for all } \alpha \in (0, 1] \text{ and all e-variables } E,$$

and moreover, the inequality is attainable. Hence, for any set \mathcal{E} of e-variables, it holds that $R_\gamma(\mathcal{E}) \leq R_\gamma(\mathcal{E}_0) = \gamma$ for $\gamma \in (0, 1]$.

We are interested only in the case $\gamma \in (0, 1]$. For $\gamma > 1$, it is usually the case that $R_\gamma(\mathcal{E}) = 1$ because for most classes \mathcal{E} that we consider, either $1 \in \mathcal{E}$ or 1 is the limit of elements of \mathcal{E} . Further, the usual interpretation of e-values does not allow for a rejection of the null hypothesis with a threshold less than 1.

The Markov inequality is often seen as being quite conservative in statistical practice, because for an e-variable E , $\mathbb{P}(E \geq 1/\alpha) = \alpha$ if and only if E has a two-point distribution such that $\mathbb{P}(E = 1/\alpha) = \alpha = 1 - \mathbb{P}(E = 0)$. Such an e-variable is unlikely to be observed in practice unless it is obtained by an algorithm which stops an e-process when it reaches $1/\alpha$. This suggests that there is room to improve $R_\gamma(\mathcal{E})$ under additional conditions.

To run an e-test at level $\alpha \in (0, 1)$ for the e-variable E , one needs to find a threshold $t > 0$, the smaller the better, such that $\mathbb{P}(E \geq t) \leq \alpha$. For this, we intuitively should use the smallest t such that $R_{1/t}(\mathcal{E}) \leq \alpha$, which satisfies $R_{1/t}(\mathcal{E}) = \alpha$ in case $\gamma \mapsto R_\gamma(\mathcal{E})$ is continuous. We only consider $t \geq 1$, corresponding to $\gamma \leq 1$.

2.2 Basic properties

The following result gives an equivalent formulation of the smallest threshold t for a set of e-variables. Denote by $q_\beta(X)$ the left $(1 - \beta)$ -quantile function of X , that is,

$$q_\beta(X) = \inf\{x \in \mathbb{R} : \mathbb{P}(X \leq x) \geq 1 - \beta\} \quad \text{for } \beta \in (0, 1).$$

The intuition is that the threshold for a set of e-variables is the supremum of the corresponding quantiles, just like the threshold for one test statistic, which should be chosen as its quantile at a given level.

Lemma 1. *For $\alpha \in (0, 1)$, the quantity $T_\alpha(\mathcal{E}) := \inf\{t \geq 1 : R_{1/t}(\mathcal{E}) \leq \alpha\}$ satisfies*

$$T_\alpha(\mathcal{E}) = \left(\sup_{E \in \mathcal{E}} q_\alpha(E) \right) \vee 1.$$

If $\gamma \mapsto R_\gamma(\mathcal{E})$ is continuous, then $T_\alpha(\mathcal{E})$ is the smallest real number $t \geq 1$ such that $\mathbb{P}(E \geq t) \leq \alpha$ for all $E \in \mathcal{E}$.

Proof. Since $R_\gamma(\mathcal{E}) \leq \gamma$ for $\gamma \in (0, 1]$, we have that $\{t \geq 1 : R_{1/t}(\mathcal{E}) \leq \alpha\}$ is not empty. Note that

$$T_\alpha(\mathcal{E}) \geq \inf\{t \geq 1 : \mathbb{P}(E > t) \leq \alpha \text{ for all } E \in \mathcal{E}\} = \left(\sup_{E \in \mathcal{E}} q_\alpha(E) \right) \vee 1.$$

Take any $\epsilon \in (0, 1)$. We have

$$\begin{aligned} T_\alpha(\mathcal{E}) - \epsilon &= \inf\{t \geq 1 - \epsilon : R_{1/(t+\epsilon)}(\mathcal{E}) \leq \alpha\} \\ &= \inf\left\{t \geq 1 - \epsilon : \sup_{E \in \mathcal{E}} \mathbb{P}(E \geq t + \epsilon) \leq \alpha\right\} \\ &= \inf\{t \geq 1 - \epsilon : \mathbb{P}(E \geq t + \epsilon) \leq \alpha \text{ for all } E \in \mathcal{E}\} \\ &\leq \inf\{t \geq 1 - \epsilon : \mathbb{P}(E > t) \leq \alpha \text{ for all } E \in \mathcal{E}\} \\ &= \left(\sup_{E \in \mathcal{E}} q_\alpha(E) \right) \vee (1 - \epsilon) \leq \left(\sup_{E \in \mathcal{E}} q_\alpha(E) \right) \vee 1 \leq T_\alpha(\mathcal{E}). \end{aligned}$$

Since $\epsilon \in (0, 1)$ is arbitrary, we have $T_\alpha(\mathcal{E}) = (\sup_{E \in \mathcal{E}} q_\alpha(E)) \vee 1$, showing the first statement. If $\gamma \mapsto R_\gamma(\mathcal{E})$ is continuous, then

$$T_\alpha(\mathcal{E}) = \min\{t \geq 1 : R_{1/t}(\mathcal{E}) \leq \alpha\} = \min\{t \geq 1 : \mathbb{P}(E \geq t) \leq \alpha \text{ for all } E \in \mathcal{E}\},$$

showing the second statement. \square

Remark 1. In the last statement of Lemma 1, the condition that $\gamma \mapsto R_\gamma(\mathcal{E})$ is continuous is essential. In general, there may not exist a smallest $t \geq 1$ such that $\mathbb{P}(E \geq t) \leq \alpha$ for all $E \in \mathcal{E}$; this is the case even when \mathcal{E} is a singleton, such as $\mathcal{E} = \{1\}$, for which we can see that $T_\alpha(\mathcal{E}) = 1$ but $\mathbb{P}(E \geq T_\alpha(\mathcal{E})) = 1$ for each $\alpha \in (0, 1)$.

Vovk and Wang [23] studied e-to-p calibrators, which are mappings that convert any e-value into a p-value. Our function $\gamma \mapsto R_{1/\gamma}(\mathcal{E})$ refines this concept. For a subset \mathcal{E} of e-variables, we say that a function $f : [0, \infty] \rightarrow [0, \infty]$ is an e-to-p calibrator on \mathcal{E} if f is decreasing and $f(E)$ is a p-variable for all $E \in \mathcal{E}$. Vovk and Wang [23] showed that $x \mapsto (1/x) \wedge 1$ is the only useful e-to-p calibrator on \mathcal{E}_0 in the sense of being the smallest. For various subsets \mathcal{E} of \mathcal{E}_0 , we can find better e-to-p calibrators based on $R_{1/\gamma}(\mathcal{E})$ than $x \mapsto (1/x) \wedge 1$. Moreover, the following result implies that any class \mathcal{E} admits a smallest e-to-p calibrator. This is in sharp contrast to the set of calibrators from p-values to e-values, which does not admit a smallest element, well discussed by Shafer [20] and Vovk and Wang [23].

Theorem 1. *The function $x \mapsto R_{1/x}(\mathcal{E})$ on $[0, \infty]$ is an e-to-p calibrator on \mathcal{E} , and it is the smallest such calibrator.*

Proof. We first show that the function $g : x \mapsto R_{1/x}(\mathcal{E})$ is an e-to-p calibrator on \mathcal{E} . This means $\mathbb{P}(g(E) \leq \alpha) \leq \alpha$ for all $\alpha \in (0, 1)$ and $E \in \mathcal{E}$. By definition, for $x \in [0, \infty]$ and $E \in \mathcal{E}$,

$$g(x) \geq \mathbb{P}(E \geq x) = \mathbb{P}(-E \leq -x) = F_{-E}(-x),$$

where F_X is the cdf of a random variable X . Since $F_X(X)$ is a p-variable for any random variable X , we have $\mathbb{P}(F_{-E}(-E) \leq \alpha) \leq \alpha$ for all $E \in \mathcal{E}$. Therefore, $\mathbb{P}(g(E) \leq \alpha) \leq \mathbb{P}(F_{-E}(-E) \leq \alpha) \leq \alpha$ for all $\alpha \in (0, 1)$.

We now prove that g is the smallest calibrator on \mathcal{E} . Suppose that f is another calibrator on \mathcal{E} such that $f(x) < g(x)$ for at least one $x \in [1, \infty]$. By definition, there exists $E \in \mathcal{E}$ such that $f(x) < \mathbb{P}(E \geq x)$. This implies $\mathbb{P}(f(E) \leq f(x)) \geq \mathbb{P}(E \geq x) > f(x)$, which violates the definition of $f(E)$ as a p-variable. \square

Theorem 1 implies that by computing $R_\gamma(\mathcal{E})$ or an upper bound on it, we can find calibrators to convert e-values realized by elements of \mathcal{E} into p-values, which work better than the e-to-p calibrator $x \mapsto (1/x) \wedge 1$ on \mathcal{E}_0 . This can be useful in procedures that take p-values as input, in e.g., that of Benjamini and Hochberg [4].

3 Improved thresholds under distributional information

In this section, we study the improved e-thresholds under some conditions on the density function or distribution function of the e-variable by computing $R_\gamma(\mathcal{E})$ or an upper bound on it.

3.1 Decreasing and unimodal densities

We first consider e-variables that have a decreasing density on its support, a decreasing density for values above 1, or a unimodal density. Denote by

$$\begin{aligned} \mathcal{E}_D &= \{E \in \mathcal{E}_0 : E \text{ has a decreasing density on its support}\}, \\ \mathcal{E}_{D>1} &= \{E \in \mathcal{E}_0 : E \text{ has a decreasing density over } [1, \infty)\}, \\ \mathcal{E}_U &= \{E \in \mathcal{E}_0 : E \text{ has a unimodal density on } [0, \infty)\}. \end{aligned}$$

The set \mathcal{E}_D contains, for instance, exponential and Pareto distributions. Whenever we consider a density, we include its limit case; that is, we allow $E \in \mathcal{E}_D$ to have a point-mass at the left end-point of its support. This convention does not change any results but simplifies many arguments in the proofs.

The set $\mathcal{E}_{D>1}$ is the set of e-variables with a decreasing density over $[1, \infty)$, but arbitrary density over $[0, 1)$. Such e-variables are relevant to e-testing since e-values between 0 and 1 provide no evidence against the null; the e-values above 1 are the ones that cause type-I errors since they may appear in the rejection region of an e-test.

A random variable on $[0, \infty)$ is said to have a unimodal density, or a unimodal distribution, if it has a density function that is increasing on $[0, x]$ and decreasing on $[x, \infty)$ for some constant $x \in [0, \infty)$, and it can possibly have a point mass at x . Such x is called a mode of the random variable. The set \mathcal{E}_U contains, for instance, log-normal and gamma distributions. The set of all modes of $E \in \mathcal{E}_U$ may be a singleton or an interval. Note that $\mathcal{E}_D \subseteq \mathcal{E}_{D>1}$ and $\mathcal{E}_D \subseteq \mathcal{E}_U$ and hence $R_\gamma(\mathcal{E}_U) \geq R_\gamma(\mathcal{E}_D)$ and $R_\gamma(\mathcal{E}_{D>1}) \geq R_\gamma(\mathcal{E}_D)$ for all $\gamma \in (0, 1]$.

The proof techniques of the following result are similar to those in Li et al. [16], Bernard et al. [5] and Wang [25] for computing bounds on probabilities with given distributional information. Huber [11] has different results on halving the bound in Markov's inequality via smoothing.

Theorem 2. For $\gamma \in (0, 1]$,

(i) $R_\gamma(\mathcal{E}_D) = \gamma/2$ if $\gamma \neq 1$ and $R_1(\mathcal{E}_D) = 1$;

(ii) $R_\gamma(\mathcal{E}_{D>1}) = \gamma/(1 + \sqrt{1 - \gamma^2})$;

(iii) $R_\gamma(\mathcal{E}_U) = (\gamma/2) \vee (2\gamma - 1)$.

Proof. We only present the proof of part (i) here, which is the simplest, but it illustrates the basic techniques of proving more complicated statements. The proof of parts (ii) and (iii) are relegated to the appendices. We first show $\mathbb{P}(E \geq 1/\gamma) \leq \gamma/2$ for each $E \in \mathcal{E}_D$ and $\gamma \in (0, 1)$. Let f be the density function of E and let a be the left end-point of the support of E . For a fixed $z > 1$, set $b := z + \mathbb{P}(E \geq z)/f(z)$. We construct an e-variable E_0 with density g_0 such that (a) $g_0(x) = f(x)$ for $a < x < b$; (b) E_0 has a point-mass at a with probability $1 - \int_a^\infty g_0(x)dx$.




Figure 3: Proof sketch in part (i) of Theorem 2. Here, f and g_0 have the same area (probability) exceeding z , with g_0 having a smaller mean; g_1 has mean 1 and larger probability of exceeding z than g_0 .

We illustrate in Figure 3 how to construct g_0 from f . Since f has a decreasing density over $[a, \infty)$, we will have $f(x) \geq g_0(x)$ for $x \in [a, z)$ and $f(x) \leq g_0(x)$ for $x > z$. We can construct g_0 by shifting the area between $f(x)$ and $f(z)$, for $x \in (a, z)$ to a point mass at a and shifting the area between $f(x)$ and 0, for $x > b$ to the area between $f(x)$ and $f(z)$, for $x \in (z, b)$. Note that b is chosen so that $\mathbb{P}(E_0 \in [z, b)) = \mathbb{P}(E \geq z)$, so $\mathbb{P}(E_0 \geq z) = \mathbb{P}(E \geq z)$. Further, since we construct g_0 by shifting the density of f to the left, we have $\mathbb{E}[E_0] \leq \mathbb{E}[E]$.

Therefore, for any $E \in \mathcal{E}_D$ with left end-point of support a , there exists an e-variable $E' \in \mathcal{E}_D$ such that E' has a point-mass at a and a uniform density on $[a, b]$ for some constant b , and $\mathbb{P}(E' \geq z) = \mathbb{P}(E \geq z)$. To show $\mathbb{P}(E \geq z) \leq 1/(2z)$, it suffices to look at the collection of e-variables that have a point-mass at a and a uniform density on $[a, b]$ for any $b > 1$. Moreover, it suffices to consider the case $\mathbb{E}[E'] = 1$.

Note that $\mathbb{E}[E'] = (1-w)a + w(a+b)/2$ for some $w \in [0, 1]$ and $b \geq 1$. The requirement $\mathbb{E}[E'] = 1$ implies that $w = 2(1-a)/(b-a)$. Now, we seek b that maximizes $\mathbb{P}(E' \geq z)$. We have $\mathbb{P}(E' \geq z) = w(b-z)/(b-a) = 2(1-a)(b-z)/(b-a)^2$. Taking the derivative with respect to b and setting the result equal to zero, we find that $b^* := 2z - a$ for $z > 1$. Substituting this value of b^* back into $\mathbb{P}(E' \geq z)$, we find that $\mathbb{P}(E' \geq z) = (1-a)/(2(z-a))$. We illustrate the density of E' that maximizes $\mathbb{P}(E' \geq z)$, denoted by g_1 , in Figure 3. Since $z > a$, the latter probability is a decreasing function of a , hence its supremum occurs for $a = 0$ and we have $\mathbb{P}(E \geq z) \leq 1/(2z)$.

It remains to show $\mathbb{P}(E \geq 1/\gamma) = \gamma/2$ for some $E \in \mathcal{E}_D$. Take E following a mixture distribution of a uniform distribution on $[0, 2/\gamma]$ with weight γ and a point-mass at 0. It is easy to see that $\mathbb{P}(E \geq 1/\gamma) = \gamma/2$ and $\mathbb{E}[E] = 1$, showing the desired result. The last statement of part (i) is trivial by checking with $E = 1$. \square

We obtain the following result by combining Theorems 1 and 2.

Corollary 1. *If $E \in \mathcal{E}_D$ then $P := (2E)^{-1}\mathbb{1}_{\{E>1\}} + \mathbb{1}_{\{E\leq 1\}}$ is a p-variable.*

The p-variable P in Corollary 1 is not a continuous function of E at $E = 1$. This is because the quantity $R_\gamma(\mathcal{E}_D)$ is not continuous at $\gamma = 1$. Indeed, $R_1(\mathcal{E}_D) = 1$ and $R_\gamma(\mathcal{E}_D) < 1/2$ for $\gamma < 1$.

Remark 2. Proposition 6.4 of Wang [25] gives that $2E^{-1}$ is a p-variable for any e-variable E with a decreasing density on $[0, \infty)$, using the concept of p*-values. Contrasted to Corollary 1, the separate treatment of $E > 1$ and $E \leq 1$ is absent in that result. This is because if the left end-point of the support of E is 0, then $\mathbb{P}(E > 1/x) \leq x/2$ for all $x > 0$; in part (i) of Theorem 2 we only have this probability bound for $x > 1$ due to our less restrictive assumption on the support of E .

The main message from Theorem 2 is that when an e-variable has a decreasing density, its threshold can be boosted by a factor of 2. That is, $T_\alpha(\mathcal{E}_D) = 1/(2\alpha)$. Further, $T_\alpha(\mathcal{E}_{D>1}) = 1/(2\alpha) + \alpha/2 = T_\alpha(\mathcal{E}_D) + \alpha/2$. Therefore, for small values of α , we can boost the threshold by a factor slightly less than 2 when $E \in \mathcal{E}_{D>1}$. This validates our message that, for small α , the distribution of uninformative e-values (between 0 and 1) has little impact on the rejection region; the shape of the distribution of large e-values has the most influence on the rejection threshold.

In part (iii) of Theorem 2, $R_\gamma(\mathcal{E}_U) = \gamma/2$, for $\gamma \in (0, 2/3]$ and $R_\gamma(\mathcal{E}_U) = 2\gamma - 1$ for $\gamma \in (2/3, 1]$. Since $\gamma \leq 2/3$ is the most practical situation (meaning a type-I error control of $1/3$), the main message from part (iii) is similar to that from part (i): when an e-variable has a unimodal density, its threshold can be boosted by a factor of 2 in practice. We provide in Figure 4 a comparison of different worst-case type-I errors and improved thresholds for $\mathcal{E}_0, \mathcal{E}_D, \mathcal{E}_{D>1}$ and \mathcal{E}_U .

3.2 Distributions of log-transformed e-variables

In many applications of e-values, the final e-variable E is the product of many e-variables. In such a setting, assuming $\log E$ has some simple distributional properties may be reasonable. We say that a random variable X has a symmetric distribution if there exists $c \in \mathbb{R}$ such that X and $c - X$ are identically distributed. We consider six different sets in this section:

$$\begin{aligned} \mathcal{E}_{LS} &= \{E \in \mathcal{E}_0 : \log E \text{ has a symmetric distribution}\}, \\ \mathcal{E}_{LU} &= \{E \in \mathcal{E}_0 : \log E \text{ has a unimodal density}\}, \\ \mathcal{E}_{LD>0} &= \{E \in \mathcal{E}_0 : \log E \text{ has a decreasing density over } [0, \infty)\}, \\ \mathcal{E}_{LD} &= \{E \in \mathcal{E}_0 : \log E \text{ has a decreasing density}\}, \\ \mathcal{E}_{LUS} &= \{E \in \mathcal{E}_0 : \log E \text{ has a unimodal and symmetric distribution}\}, \\ \mathcal{E}_{LN} &= \{E \in \mathcal{E}_0 : E \text{ has a log-normal distribution}\}. \end{aligned}$$

In all sets above, we require $\mathbb{P}(E = 0) = 0$, so that $\log E$ is a real-valued random variable. Note that $\mathcal{E}_{LN} \subseteq \mathcal{E}_{LUS} \subseteq \mathcal{E}_{LS}$ and $\mathcal{E}_{LD} \subseteq \mathcal{E}_{LD>0}$. The point-mass distributions $x \in (0, 1]$ are included in all sets above, which are degenerate cases of log-normal distributions.




Figure 4: Comparison of worst-case type-I errors and improved thresholds for decreasing and unimodal densities.

The following theorem provides the worst-case type-I errors for the six sets of e-variables. The proof is given in Appendix C.

Theorem 3. For $\gamma \in (0, 1]$,

(i) $R_\gamma(\mathcal{E}_{LS}) = \gamma \wedge (1/2)$ if $\gamma \neq 1$ and $R_1(\mathcal{E}_{LS}) = 1$;

(ii) $R_\gamma(\mathcal{E}_{LU}) = \gamma$;

(iii) $\mathcal{E}_{LD>0} \subseteq \mathcal{E}_{D>1}$ and

$$\frac{\gamma}{e - \gamma} \leq R_\gamma(\mathcal{E}_{LD>0}) = \frac{\gamma}{t_\gamma},$$

where t_γ is the unique solution of $t(1 - \log t) = \gamma$ for $t \in (1, e)$, and we have

$$t \uparrow e \quad \text{and} \quad \frac{R_\gamma(\mathcal{E}_{LD>0})}{\gamma} \downarrow \frac{1}{e} \quad \text{as } \gamma \downarrow 0;$$

(iv) $R_\gamma(\mathcal{E}_{LD}) = R_\gamma(\mathcal{E}_{LUS})$ and

$$\frac{\gamma}{e} \leq R_\gamma(\mathcal{E}_{LD}) = R_\gamma(\mathcal{E}_{LUS}) \leq \frac{\gamma}{e(1 - \gamma^2)} \wedge \left(\frac{\gamma}{1 + \sqrt{1 - \gamma^2}} \right);$$

(v) if $\gamma \neq 1$ then

$$R_\gamma(\mathcal{E}_{LN}) = \Phi\left(-\sqrt{-2 \log \gamma}\right) \leq \frac{\gamma}{2\sqrt{-\pi \log \gamma}},$$

where Φ is the standard normal cdf, and $R_1(\mathcal{E}_{LN}) = 1$.

The results in parts (i) and (ii) of Theorem 3 show that the assumption of log-transformed symmetric or log-transformed unimodal distributions do not improve the bound from Markov's inequality for the most useful cases where $\gamma \leq 1/2$.

The class $\mathcal{E}_{LD>0}$ offers a non-trivial threshold improvement by roughly a factor of e . As with $\mathcal{E}_{D>1}$, most evidence against the null comes from observing e-values larger than 1, which is equivalent to observing log e-values larger than zero. For γ small, such as 0.01 or 0.05, the lower and upper bounds in part (iv) of

Theorem 3 are very close; hence, using the upper bound is not wasteful. Part (v) shows that log-normality improves over $R_\gamma(\mathcal{E}_0)$ by at least $3.54\sqrt{-\log \gamma}$.

We summarize the bounds on worst-case type-I errors and improved thresholds from Theorem 3 in Figure 5.




Figure 5: Comparison of worst-case type-I errors and improved thresholds for log-transformed e-variables. Results for \mathcal{E}_{LUS} are conservative bounds since we only find an upper bound for $R_\gamma(\mathcal{E}_{LUS})$ and $T_\alpha(\mathcal{E}_{LUS})$.

3.3 Log-concave density, survival or distribution functions

In this section, we study distributional assumptions of log-concavity, a popular property for density functions, survival functions, and distributional functions. We will consider the sets

$$\begin{aligned}\mathcal{E}_{LCD} &= \{E \in \mathcal{E}_0 : E \text{ has a log-concave density on } \mathbb{R}\}, \\ \mathcal{E}_{LCS} &= \{E \in \mathcal{E}_0 : E \text{ has a log-concave survival function on } \mathbb{R}\}, \\ \mathcal{E}_{LCF} &= \{E \in \mathcal{E}_0 : E \text{ has a log-concave distribution function on } \mathbb{R}\}.\end{aligned}$$

The density of $E \in \mathcal{E}_{LCD}$ may take value 0 on some intervals like $(-\infty, a)$ or (a, ∞) for $a \geq 0$, or both, that is, the log-density function is $-\infty$ on such intervals. The assumption of log-concave density is satisfied by, for instance, normal, uniform, Laplace and gamma (with shape parameter ≥ 1) distributions, as well as their location-scale transforms. When considering e-variables, the support of these distributions should be confined to $[0, \infty)$.

All distributions with log-concave densities have log-concave survival functions and log-concave distribution functions, thus $\mathcal{E}_{LCD} \subseteq \mathcal{E}_{LCS}$ and $\mathcal{E}_{LCD} \subseteq \mathcal{E}_{LCF}$. Moreover, $\mathcal{E}_{LCD} \subseteq \mathcal{E}_U$ and $\mathcal{E}_D \subseteq \mathcal{E}_{LCF}$. The property of having a log-concave survival function is also known as having an increasing hazard rate or increasing failure rate. Log-normal and Pareto distributions belong to the class of distributions with log-concave cumulative distribution functions but do not have log-concave densities or survival functions. For many examples of distributions belonging to the above classes, see, e.g., Bagnoli and Bergstrom [2].

Theorem 4. For $\gamma \in (0, 1]$, we have

(i) $e^{-1/\gamma} \leq R_\gamma(\mathcal{E}_{LCS}) = e^{s_\gamma/\gamma} \leq e^{1-1/\gamma}$, where $s_1 = 0$ and for $\gamma < 1$, s_γ is the unique solution $s \in (-1, 0)$ to the equation $e^{s/\gamma} = s + 1$;

(ii) $\mathcal{E}_{LCD} \subseteq \mathcal{E}_{LCS}$ and $\mathcal{E}_{LCD} \subseteq \mathcal{E}_U$ and thus

$$e^{-1/\gamma} \leq R_\gamma(\mathcal{E}_{LCD}) \leq R_\gamma(\mathcal{E}_U) \wedge R_\gamma(\mathcal{E}_{LCS}) \leq e^{1-1/\gamma};$$

(iii) $R_\gamma(\mathcal{E}_{\text{LCF}}) = \gamma$.

The proof of Theorem 4 is given in Appendix D.

Part (i), dealing with log-concave survival functions, gives the most useful result since it is easy to verify visually and provides a useful bound. The upper bound in part (i) of Theorem 4 is convenient for applications due to its analytical form. Numerically, we find that $R_\gamma(\mathcal{E}_{\text{LCS}})$ is closer to the lower bound of $e^{-1/\gamma}$, but using the upper bound is not so wasteful since it is at most the constant e too large, while the bound grows exponentially. In part (ii), we could not obtain the distribution that attains the lower bound, implying that there may be room for improvement. For the class of log-concave distributions, we obtain a negative result: There is generally no better bound on $R_\gamma(\mathcal{E}_{\text{LCF}})$ other than the Markov bound.

We summarize the worst-case type-I error bounds and the improved thresholds in Figure 6. Recall that the bounds for \mathcal{E}_{LCD} do not hold with equality, so there may be better bounds available, although we have not identified them.




Figure 6: Comparison of worst-case type-I errors and improved thresholds for log-concave density, survival and distribution functions. Results for \mathcal{E}_{LCD} are conservative bounds since we only find an upper bound for $R_\gamma(\mathcal{E}_{\text{LCD}})$ and $T_\alpha(\mathcal{E}_{\text{LCD}})$.

3.4 Summary

For a class of e-variables \mathcal{E} , Lemma 1 allows one to design a more aggressive e-test based on the rejection threshold $T_\alpha(\mathcal{E})$. The rejection region is enlarged, leading to rejections with less evidence while keeping the type-I error controlled at α for $\alpha \in (0, 1)$. We have already provided plots of the worst-case type-I errors and improved rejection thresholds in Figures 4, 5, and 6. We summarize the results on the improved thresholds in Table 1.

There are some ad-hoc thresholds of e-values used in the literature, as it is clear that the standard calibrator $\min(1/p, 1)$ is too restrictive. For instance, Shafer [20] recommended using the conversion formula $1/\sqrt{p} - 1$. Vovk and Wang [23] suggested using the rule of thumb proposed in Appendix B of Jeffreys [14] to convert between p-values and Bayes factors, as Bayes factors align with e-values in the context of simple hypotheses.

According to Jeffreys' rule of thumb, a p-value threshold of 0.05 is equivalent to an e-value threshold of $\sqrt{10}$, while a p-value threshold of 0.01 is equivalent to an e-value threshold of 10. The correspondence between p-values and e-values generated from this rule of thumb can be arbitrary since it depends on the distribution of the e-variable, which may be unknown. The improved thresholds derived in this section provide a mathematical framework to compare the scales of e-values and p-values. It appears that Jeffreys'

Table 1: Improved thresholds $T_\alpha(\mathcal{E})$ for different distributional hypothesis.

	α					
	0.001	0.01	0.02	0.05	0.1	0.2
$\mathcal{E}_0, \mathcal{E}_{LS}$	1000	100	50	20	10	5
$\mathcal{E}_D, \mathcal{E}_U$	500	50	25	10	5	2.5
$\mathcal{E}_{D>1}$	500	50.01	25.01	10.03	5.05	2.60
$\mathcal{E}_{LCD}, \mathcal{E}_{LCS}$	6.91	4.65	4	3.15	2.56	2
$\mathcal{E}_{LUS}, \mathcal{E}_{LD}$	367.88	36.82	18.45	7.49	3.93	2.25
$\mathcal{E}_{LD>0}$	368.25	37.16	18.77	7.73	4.07	2.25
\mathcal{E}_{LN}	118	14.97	8.24	3.87	2.27	1.42

rule of thumb based on Bayes factors is slightly less conservative than the class \mathcal{E}_{LN} but more conservative than \mathcal{E}_{LCS} . More importantly, in contrast to the ad-hoc choices, the thresholds obtained in this paper have a provable type-I error guarantee under the respective assumptions.

4 Supremum of comonotonic e-values

In this section, we will introduce a testing strategy when we have access to a family of comonotonic e-variables. This strategy can be used in conjunction with the improved thresholds.

4.1 Threshold for comonotonic e-values

In what follows, each Q_θ is a probability measure. To test $\{Q_{\theta_0}\}$ against $\{Q_\theta\}$, we usually use the e-variable as a likelihood ratio, that is,

$$E_\theta = \frac{dQ_\theta}{dQ_{\theta_0}}.$$

If we test $\{Q_{\theta_0}\}$ against $\{Q_\theta : \theta \in \Theta_1\}$ where $\Theta_1 \subseteq \Theta$, we can use a mixture

$$E_\nu = \int_{\Theta_1} \frac{dQ_\theta}{dQ_{\theta_0}} \nu(d\theta),$$

where ν is a prior probability on Θ_1 .

For instance, as in Vovk and Wang [23], the simplest example is to test $N(0, 1)$ against $N(\mu, 1)$ with $\mu \neq 0$ for iid observations X_1, \dots, X_n . The e-variable E_μ for $N(0, 1)$ is the likelihood ratio

$$E_\mu = \exp(\mu S_n - n\mu^2/2),$$

where $S_n = \sum_{i=1}^n X_i$. Suppose that the alternative hypothesis is $\mu > 0$. In this case, an interesting observation is that $E_\mu, \mu > 0$ are *comonotonic* random variables. Recall that random variables $E_\theta, \theta \in \Theta$ are comonotonic if there exists a common random variable Z such that E_θ is an increasing function of Z for each $\theta \in \Theta$.

Lemma 2. *Let $x \in \mathbb{R}$ and $\gamma \in [0, 1]$. Suppose that \mathcal{X} is a collection of comonotonic random variables satisfying $\mathbb{P}(X \geq x) \leq \gamma$ for each $X \in \mathcal{X}$. Then $\mathbb{P}(\sup_{X \in \mathcal{X}} X \geq x) \leq \gamma$.*

Proof. Let q_X be the right quantile function of X for each X , that is,

$$q_X(\alpha) = \inf\{x \in \mathbb{R} : \mathbb{P}(X \leq x) > \alpha\}, \quad \alpha \in (0, 1).$$

Note that $\mathbb{P}(X \geq x) \leq \gamma$ is equivalent to $q_X(1 - \gamma) \leq x$ (e.g., Lemma 1 of Guan et al. [10]). This gives $\sup_{X \in \mathcal{X}} q_X(1 - \gamma) \leq x$. Note that comonotonicity implies that $\sup_{X \in \mathcal{X}} q_X(1 - \gamma)$ is the right quantile function of $\sup_{X \in \mathcal{X}} X$. Using the above relation again we get $\mathbb{P}(\sup_{X \in \mathcal{X}} X > x) \leq \gamma$. \square

The next result follows directly.

Proposition 1. *Suppose that $\{E_\theta : \theta \in \Theta\} \subseteq \mathcal{E}_0$ is a collection of comonotonic e-variables for a hypothesis H . Then $\sup_{Q \in H} Q(\sup_{\theta \in \Theta} E_\theta \geq T_\alpha(\mathcal{E})) \leq \alpha$.*

Consequently, we can obtain a valid test using the supremum of e-variables instead of their mixture under comonotonicity. This will always have a higher power than using a mixture.

If we push this idea further, we can take the set \mathcal{E} of all e-variables for \mathbb{P} that are decreasing functions of a statistic. This will lead to a precise p-variable.

Proposition 2. *Let \mathcal{E} be the set of all e-variables for \mathbb{P} that are decreasing functions of a random variable X with distribution function F . Then, $(\sup_{E \in \mathcal{E}} E)^{-1} = F(X)$ almost surely, i.e., the natural p-variable built from X .*

Proof. For $s \in \mathbb{R}$, let

$$E_s = \frac{1}{F(s)} \mathbf{1}_{\{X \leq s\}},$$

which is an e-variable. It is clear that $\sup_{s \in \mathbb{R}} E_s = 1/F(X)$, and $\sup_{E \in \mathcal{E}} E \geq 1/F(X)$. To show the opposite, suppose that there exists $E \in \mathcal{E}$ such that

$$\mathbb{P}\left(E > \frac{1}{F(X)}\right) > 0.$$

Since E is decreasing in X , there exists $s \in \mathbb{R}$ such that $\mathbb{P}(E > E_s) > 0$ and $\mathbb{P}(E \geq E_s) = 1$. This contradicts $\mathbb{E}[E] \leq 1$. \square

Proposition 2 suggests that the supremum of comonotonic e-values can be seen as a conceptual middle ground between an e-value (\mathcal{E} is a singleton) and a p-value (\mathcal{E} in Proposition 2, which is a maximal comonotonic set).

The idea of taking a supremum does not generalize to optional stopping for the e-process because comonotonicity is not preserved in general. That is, for e-processes $n \mapsto E_\theta(n)$ indexed by θ , even if the e-variables $E_\theta(n)$ are comonotonic across θ for each fixed n (as in the normal example), the stopped e-variables $E_\theta(\tau)$ need not be comonotonic across θ for a stopping time τ .

4.2 Implications in the Gaussian setting

When testing $N(0, 1)$ against $\{N(\mu, 1) : \mu > 0\}$, the likelihood ratio e-variables are comonotonic. Instead of using

$$E_\mu(n) = \exp(\mu S_n - n\mu^2/2)$$

for a fixed $\mu > 0$ or its mixture, we can use

$$Y(n) = \sup_{\mu > 0} \exp(\mu S_n - n\mu^2/2),$$

which, although not being an e-variable, gives $\mathbb{P}(Y(n) \geq 1/\alpha) \leq \alpha$ under the null. We can explicitly compute

$$Y(n) = \exp\left(\frac{(S_n)_+^2}{2n}\right).$$

We can compare this with (23) of Vovk and Wang [23], which gives the e-variable

$$E_\Phi(n) = \frac{1}{\sqrt{n+1}} \exp\left(\frac{S_n^2}{2n+2}\right),$$

obtained via a mixture of $E_\mu(n)$ using a standard normal prior Φ (a two-sided alternative). Under the null, we can compute

$$\mathbb{P}(Y(n) \geq 1/\alpha) = \mathbb{P}\left(\frac{(S_n)_+^2}{2n} \geq -\log \alpha\right) = 1 - \Phi\left(\sqrt{-2 \log \alpha}\right),$$

where Φ is the standard normal cdf. Denote by ϕ the standard normal pdf. Since for $x > 0$,

$$\frac{x}{x^2 + 1} \leq \frac{1 - \Phi(x)}{\phi(x)} \leq \frac{1}{x},$$

we get

$$\mathbb{P}(Y(n) \geq 1/\alpha) \approx \frac{1}{\sqrt{-2 \log \alpha}} \frac{1}{\sqrt{2\pi}} \alpha.$$

Although this is still less than α , the magnitude has improved substantially from $E_\Phi(n)$, which roughly corresponds to replacing α by $\alpha^{(n+1)^{-1/2}}$.

Remark 3. We also note that $E_\mu, \mu > 0$ are log-normally distributed, so we can use the threshold $T_\alpha(\mathcal{E}_{\text{LN}}) = \exp(-(\Phi^{-1}(\alpha))^2/2)$, where Φ^{-1} is the inverse distribution function (quantile function) of a standard normal distribution. Then, it is easy to show $\mathbb{P}(Y(n) > T_\alpha(\mathcal{E}_{\text{LN}})) = \alpha$ for $\alpha \in (0, 1)$; that is, we get an exact type-I error.

4.3 Maximum likelihood e-values

The test constructed in the previous section is based on taking the supremum over a collection of Gaussian likelihood ratio e-variables. We now investigate how to generalize this to other likelihood ratio e-variables.

Suppose that we test Q_{θ_0} against $\{Q_\theta : \theta \in \Theta\}$ with a likelihood ratio e-variable,

$$E_\theta(n) := \prod_{i=1}^n \frac{dQ_\theta}{dQ_{\theta_0}}(x_i).$$

If $\{E_\theta(n) : \theta \in \Theta\}$ is a collection of comonotonic random variables, we can take the supremum according to Proposition 1. We have

$$\arg \max_{\theta \in \Theta} \prod_{i=1}^n \frac{dQ_\theta}{dQ_{\theta_0}}(x_i) = \arg \max_{\theta \in \Theta} \prod_{i=1}^n \left. \frac{dQ_\theta(x)}{dx} \right|_{x=x_i},$$

which corresponds to the maximum likelihood estimator (MLE) over Θ , denoted $\hat{\theta}$. Therefore,

$$\sup_{\theta \in \Theta} E_\theta(n) = E_{\hat{\theta}}(n),$$

and we can use an e-value constructed with a plug-in estimator using the entire dataset instead of splitting the data as in the split likelihood-ratio e-test of Wasserman et al. [28].

For $\{E_\theta(n) : \theta \in \Theta\}$ to be a collection of comonotonic random variables, we require that, for every $\theta \in \Theta$, $(dQ_\theta/dQ_{\theta_0})(x)$ increases in x over the union of the supports of Q_θ and Q_{θ_0} . The latter condition coincides with the definition of the likelihood ratio order (see, for instance, Section 1.C of Shaked and Shanthikumar [21] for properties).

One simple way to verify if a class of likelihood ratio e-variables is comonotonic is within the exponential family. For a parameter θ (possibly a vector), a family of distributions is said to belong to the exponential family if its density function or probability mass function can be written as

$$\frac{dQ_\theta(x)}{dx} = h(x) \exp \{ \eta(\theta)T(x) - A(\theta) \},$$

where $\eta(\theta)$ is the natural parameter, $T(x)$ is the summary statistic, and $A(\theta)$ is called the log-partition. When testing two distributions in the exponential family, the likelihood ratio e-value (for one observation) is

$$\frac{dQ_\theta}{dQ_{\theta_0}}(x) = \exp \{ (\eta(\theta) - \eta(\theta_0))T(x) - A(\theta) + A(\theta_0) \}.$$

The latter forms a comonotonic family if $\eta(\theta) - \eta(\theta_0)$ has the same sign for each θ . We will give examples of maximum likelihood e-values within the gamma family in Section 7.3.

5 Thresholds for combined e-values

Researchers using e-values frequently rely on independent or sequential data when collecting evidence for scientific discovery. To perform an e-test from these data, a common strategy is to compute an e-value for each observation and combine them to form an e-value or an e-process. Denote by $[T] = \{1, \dots, T\}$. We will assume that e-variables E_1, \dots, E_T are available. We will assume that the e-variables are either independent or sequential; e-variables are said to be sequential if $\mathbb{E}[E_t | E_1, \dots, E_{t-1}] \leq 1$ for $t \in [T]$. Vovk and Wang [23] studied merging functions of independent e-values and showed that the product merging function weakly dominates any merging function for independent e-values.

If one has access to sequential e-values, one can construct an e-process $\{M_t : t \in [T]\}$, accumulating evidence from individual sequential e-variables. E-processes have the property that, for any stopping time τ , the stopped e-process is an e-value. The property $\mathbb{E}[M_\tau] \leq 1$ for any stopping time is called the anytime validity property of e-processes, implying that a statistician can stop an experiment early or continue an experiment after peeking at the results from an e-test while retaining type-I error control. To test with an e-processes while controlling the type-I error at level α , it is common to use Ville's inequality, stating for $\alpha \in (0, 1]$ that

$$\mathbb{P}\left(\sup_{t \in [T]} M_t \geq \frac{1}{\alpha}\right) \leq \alpha.$$

If a researcher is not interested in anytime validity (for instance, if they have already performed experiments and do not have the option to gather more evidence), then the anytime validity property is too conservative and may contribute to a loss of power. In that case, we can construct an e-value by merging e-values and using an improved threshold that controls the type-I error for a fixed sample size.

We will now investigate how to improve the threshold of products of independent or sequential e-variables. To answer this question, we will first define the multiplicative strong unimodality (MSU) property proposed by Cuculescu and Theodorescu [7]. A random variable X is said to have the MSU property if, for any random variable Y with a unimodal density function, the random variable XY has a unimodal density function. Denote by

$$\mathcal{E}_{\text{MSU}} = \{E \in \mathcal{E}_0 : E \text{ is multiplicative strongly unimodal over } [0, \infty)\}.$$

Proposition 3.6 of Cuculescu and Theodorescu [7] states that if X is a unimodal random variable with a mode at zero, then the product of X with any other random variable defined on \mathbb{R} will also have a mode at zero; it follows that any e-variable with a decreasing density on $[0, \infty)$ will have the MSU property. Note that this does not coincide with the class \mathcal{E}_{D} since we require the mode to be zero, while the left end-point of supports for e-variables in \mathcal{E}_{D} could be larger than 0. Theorem 3.7 of Cuculescu and Theodorescu [7] provides the condition for which e-variables will have the MSU property. Letting f denote the density function of a random variable in \mathcal{E}_{MSU} , then

- (i) f has a decreasing density on $[0, \infty)$, implying it is unimodal with a mode at zero or
- (ii) $x \mapsto f(e^x)$ is log-concave on $[0, \infty)$.

Distributions that are log-concave with respect to $x \mapsto f(e^x)$ include beta, Cauchy, Fisher, gamma, inverse Gaussian, logistic, log-normal, Gaussian, Student, Pareto distributions, along with their location-scale transforms; see Alimohammadi et al. [1] for more examples.

The following is immediately apparent.

Proposition 3. *Let $S_T = \prod_{t=1}^T E_t$.*

- (i) *Let E_t be sequential e-values for $t \in [T-1]$ and let E_T be independent of S_{T-1} and have a decreasing density on $[0, \infty)$. Then, $S_T \in \mathcal{E}_{\text{D}}$.*
- (ii) *Let $E_t \in \mathcal{E}_{\text{MSU}}$ for $t \in [T]$ be independent e-variables. Then, $S_T \in \mathcal{E}_{\text{U}}$.*

Proposition 3 (i) tells us that, as long as we have one e-variable with a decreasing density on $[0, \infty)$ that is independent of other e-variables, then we can take the product and boost the threshold by 2.

In Appendix E, we provided improved thresholds for the weighted average of independent e-variables when each e-variable has a log-concave density.

When constructing e-processes based on betting strategies, one often uses, instead of E_t , $1 - \lambda_t + \lambda_t E_t$ for some λ_t , possibly depending on previous e-values, that takes values in $[0, 1]$; see Shafer [20] and Waudby-Smith and Ramdas [27]. For \mathcal{E} being any of \mathcal{E}_D , \mathcal{E}_U , \mathcal{E}_{LCD} and \mathcal{E}_{LCF} , we have the convenient property that if $E \in \mathcal{E}$, then $1 - \lambda + \lambda E \in \mathcal{E}$ for all constant $\lambda \in [0, 1]$. This holds true because properties such as monotonicity, unimodality, and log-concavity are invariant to changes in location and scale. However, part (i) of Proposition 3 requires that E_T is decreasing on $[0, \infty)$, so it is not valid to apply to $1 - \lambda + \lambda E$ in this case.

6 Boosting e-values in the e-BH procedure

The e-BH procedure of Wang and Ramdas [26] is introduced to test K hypotheses with a false discovery rate (Benjamini and Hochberg [4]) control. As with e-tests for hypothesis testing, the base e-BH procedure typically controls the FDR at a more conservative level than the desired (or target) FDR. To improve the power of the e-BH procedure, we can boost the e-values in the e-BH procedure such that it approaches the desired FDR level, while still controlling it. There are two known ways to accomplish this task. The first is to use distributional information on the marginal e-variables to boost the e-values individually. This was proposed already by Wang and Ramdas [26]. Another method to boost the e-values is to use a relaxed definition of e-values, as in Ren and Barber [19], possibly supplementing this with dependence information through conditional calibration as in Lee and Ren [15].

Since this paper aims to incorporate distributional assumptions through the marginal information on e-variables, we will focus on marginal boosting. We admit that the results in this section are mainly exploratory: The only positive results we obtained require strong assumptions (log-concave survival functions) and may not be easy to apply in practical situations. The proofs for some results in this section are given in Appendix F.

To boost e-values in the e-BH procedure through marginal boosting, we need to find constants b_k in Wang and Ramdas [26, Equations (8) and (9)]. That is, for an e-variable E , to find a constant $b > 0$ (the larger, the better) such that

$$\mathbb{E}[T(\alpha b E)] \leq \alpha, \quad (1)$$

where $\mathcal{K} := [K]$, $K/\mathcal{K} := \{K/k : k \in \mathcal{K}\}$, and $T : [0, \infty) \rightarrow [0, K]$ is defined by

$$T(x) = \frac{K}{\lceil K/x \rceil} \mathbf{1}_{\{x \geq 1\}} \quad \text{with } T(\infty) = K.$$

Noting that $\mathbb{E}[T(\alpha E)] \leq \mathbb{E}[\alpha E] \leq \alpha$, we know that $b = 1$ satisfies (1). If $b \mapsto \mathbb{E}[T(\alpha b E)]$ is continuous (it suffices if E has density on $(0, \infty)$), then we will take

$$b = \max\{c \geq 1 : \mathbb{E}[T(\alpha c E)] \leq \alpha\}$$

under arbitrary dependence (AD). If the e-values in K hypotheses satisfy the property of positive regression dependence on a subset (PRDS), then b can be improved to a constant satisfying

$$\max_{x \in K/\mathcal{K}} x \mathbb{P}(\alpha b E \geq x) \leq \alpha. \quad (2)$$

Since T depends on K , it may not be convenient to analyze. We can relax (1), by using $T(x) \leq x \mathbf{1}_{\{x \geq 1\}}$ for $x \in \mathbb{R}$, to

$$\mathbb{E}[\alpha b E \mathbf{1}_{\{\alpha b E \geq 1\}}] \leq \alpha, \quad (3)$$

and relax (2) to

$$\max_{x \geq 1} x \mathbb{P}(\alpha b E \geq x) \leq \alpha. \quad (4)$$

We define the quantity $B_\alpha^{\text{AD}}(\mathcal{E})$ as

$$B_\alpha^{\text{AD}}(\mathcal{E}) = \inf_{E \in \mathcal{E}} \sup \left\{ c \geq 1 : \mathbb{E} [\alpha c E \mathbf{1}_{\{\alpha c E \geq 1\}}] \leq \alpha \right\},$$

which is the boosting factor under AD for $E \in \mathcal{E}$ assuming continuity. Similarly, we define

$$B_\alpha^{\text{PR}}(\mathcal{E}) = \inf_{E \in \mathcal{E}} \sup \left\{ c \geq 1 : \max_{x \geq 1} x \mathbb{P}(\alpha c E \geq x) \leq \alpha \right\},$$

which is the boosting factor for $E \in \mathcal{E}$ assuming continuity for a PRDS multiple testing problem.

We first present a negative result on decreasing densities. This result suggests that getting a useful boosting factor for other similar distributional information may be difficult. In particular, there is no hope for any set containing \mathcal{E}_D , such as \mathcal{E}_U and $\mathcal{E}_{D>1}$.

Proposition 4. *For any $\alpha \in (0, 1)$, we have $B_\alpha^{\text{AD}}(\mathcal{E}_D) = 1$.*

Proof. It suffices to find a sequence of random variables within \mathcal{E}_D to satisfy (3) with equality, hence no boosting is possible (that is, $B_\alpha^{\text{AD}}(\mathcal{E}_D) = 1$). For any $w \in (0, 2\alpha)$, let E be a mixture random variable of a uniform distribution on $[0, 2/w]$ with weight w and a point-mass at 0. Then, we have

$$\mathbb{E} [\alpha E \mathbf{1}_{\{\alpha E \geq 1\}}] = \int_{1/\alpha}^{2/w} \alpha x \frac{w^2}{2} dx = \alpha - \frac{w^2}{4\alpha}.$$

Since we can make w arbitrarily small, we have that $B_\alpha^{\text{AD}}(\mathcal{E}_D) = 1$. \square

Remark 4. Consider again the e-variable defined in the proof of Proposition 4. If $w < \alpha/K$, then we have for any $b \geq 1$ that

$$\mathbb{E}[T(\alpha b E)] \leq K w \leq \alpha,$$

so, in this case, it is possible to boost the e-values by an arbitrary amount while still satisfying the criterion, but we could not boost the e-values at all using the simplified problem that does not rely on K .

Next, two positive results concern e-variables with log-concave survival functions.

Proposition 5. *For some $\alpha \in (0, 1)$, let $c_1^{\text{AD}}(\alpha)$ be the constant $b \geq 1$ such that $\exp\{-1/(ab)\}(1 + ab) = \alpha \exp\{-1\}$ and $c_2^{\text{AD}}(\alpha)$ be the constant $b' \geq 1$ such that $\exp\{-1/(ab')\}(1 + ab') = \alpha$. Then, we have*

$$c_1^{\text{AD}}(\alpha) \leq B_\alpha^{\text{AD}}(\mathcal{E}_{\text{LCS}}) \leq c_2^{\text{AD}}(\alpha).$$

Remark 5. For any $\alpha \in (0, 1)$, we may obtain an analytical lower bound for $c_1^{\text{AD}}(\alpha)$ and upper bound for $c_2^{\text{AD}}(\alpha)$ that are slightly looser than the bounds in Proposition 5. Note that since

$$\exp\{-1/(ab)\}(\alpha + 1) \leq \exp\{-1/(ab)\}(ab + 1),$$

we may obtain an upper bound for $c_2^{\text{AD}}(\alpha)$ by solving for b in $\exp\{-1/(ab)\}(\alpha + 1) = \alpha$, such that

$$c_2^{\text{AD}}(\alpha) \leq \frac{1}{\alpha \log\left(\frac{1+\alpha}{\alpha}\right)} \leq -\frac{1}{\alpha \log \alpha}. \quad (5)$$

Then, we have that $c_1^{\text{AD}}(\alpha)$ is the solution for $b \geq 1$ in

$$\exp\{-1/(ab)\}(1 + ab) = \alpha \exp\{-1\}. \quad (6)$$

We may obtain a lower bound for $c_1^{\text{AD}}(\alpha)$ by replacing the second b in (6) by its upper bound (5), hence a lower bound for $c_1^{\text{AD}}(\alpha)$ is the solution for $b \geq 1$ in $\exp\{-1/(ab)\}(1 - 1/\log(\alpha)) = \alpha \exp\{-1\}$, which leads to

$$(\alpha [\log \alpha - 1 + \log(-\log(\alpha)) - \log(1 - \log(\alpha))])^{-1} \leq c_1^{\text{AD}}(\alpha).$$

Finally, let us remark that the bounds in Proposition 5 are asymptotically tight as $\alpha \rightarrow 0$ since

$$\lim_{\alpha \rightarrow 0^+} \frac{c_2^{\text{AD}}(\alpha)}{c_1^{\text{AD}}(\alpha)} = \lim_{\alpha \rightarrow 0^+} \frac{\log \alpha - 1 + \log(-\log(\alpha)) - \log(1 - \log(\alpha))}{\log \alpha} = 1.$$

Proposition 6. Let $\alpha \in (0, 1)$. Define

$$c_1^{\text{PR}}(\alpha) = \frac{1}{\alpha - \alpha \log \alpha} \quad \text{and} \quad c_2^{\text{PR}}(\alpha) = \begin{cases} \exp\{1\}, & \alpha \geq \exp\{-1\}; \\ -\frac{1}{\alpha \log \alpha}, & \alpha \leq \exp\{-1\}. \end{cases}$$

We have

$$c_1^{\text{PR}}(\alpha) \leq B_\alpha^{\text{PR}}(\mathcal{E}_{\text{LCS}}) \leq c_2^{\text{PR}}(\alpha).$$

We can obtain numerically the bounds in Propositions 5 and 6. We present the lower and upper bounds for $B_\alpha^{\text{AD}}(\mathcal{E}_{\text{LCS}})$ and $B_\alpha^{\text{PR}}(\mathcal{E}_{\text{LCS}})$ in Table 2. We note that $c_1^{\text{AD}}(\alpha)$ is only slightly smaller than $c_1^{\text{PR}}(\alpha)$, so one doesn't gain much boosting power by assuming PRDS (and the p-BH procedure is valid under PRDS and typically makes more discoveries, so the e-BH procedure may not be the best procedure to use in this case).

Table 2: Lower and upper bounds for $B_\alpha^{\text{AD}}(\mathcal{E}_{\text{LCS}})$ and $B_\alpha^{\text{PR}}(\mathcal{E}_{\text{LCS}})$.

α	$c_1^{\text{AD}}(\alpha)$	$c_2^{\text{AD}}(\alpha)$	$c_1^{\text{PR}}(\alpha)$	$c_2^{\text{PR}}(\alpha)$
0.01	17.35	20.86	17.84	21.71
0.02	9.82	12.11	10.18	12.78
0.05	4.75	6.13	5.01	6.68
0.10	2.82	3.81	3.03	4.34

7 Simulation studies

In this section, we use the results derived in this paper to illustrate how the improved thresholds enhance the power of e-tests via simulation.

7.1 Tests within the Gaussian family

A simple example illustrates the difference between e-processes and the supremum process. Suppose we have a sequence of N iid observations X_1, \dots, X_N from $N(\delta, 1)$ with $\delta = 0.3$ and let $S_n = X_1 + \dots + X_n$ for $1 \leq n \leq N$. We want to test the null hypothesis that data are $N(0, 1)$ distributed using different test statistics: the likelihood ratio e-process $\{E_\mu(n)\}_{1 \leq n \leq N}$ with the true alternative $\mu = \delta$ and two likelihood ratio e-processes with misspecified alternatives $\mu = 0.2$ and $\mu = 0.4$. We further compare with the universal e-process $\{E_\Phi(n)\}_{1 \leq n \leq N}$ and the supremum process $\{Y(n)\}_{1 \leq n \leq N}$. Note that the supremum process controls the type-I error for fixed n but is not anytime valid.

In Figure 7, we present a simulated path for the process $\{S_n\}_{1 \leq n \leq N}$ (with its expected value $0.3 \times n$) along with the e-processes and the supremum process for this example with $N = 1000$. One observes for this sample path that the process $\{S_n\}_{1 \leq n \leq N}$ starts off above its mean but reverts to its mean for the second half of the process. Hence, for this simulated path, the supremum process for n under 300 is close to the e-process with $\mu = 0.4$ (since the empirical average is closer to 0.4 than 0.3). As the $\{S_n\}_{1 \leq n \leq N}$ approaches its mean for n above 500, the supremum process matches the e-process with $\mu = 0.3$.

In Figure 8, we present the average e-process and supremum process for 100 replications (where the average is taken over the log of the e-process and the supremum process). For the e-processes with the likelihood ratio, the average of the log is close to the expected value of the log e-process, which is $\delta n(\mu - \delta/2)$. The mixture e-process becomes larger than the likelihood ratio e-processes with a misspecified alternative for n larger than 600. The growth optimal e-process is the likelihood ratio e-process with the correct alternative, that is, $\{E_\delta(n)\}_{1 \leq n \leq N}$ (see, for instance, Grünwald et al. [9], Shafer [20]). The supremum process is larger than the growth-optimal e-process but does not lead to an anytime valid test.

Note that $E_\mu(n)$ is log-normally distributed; hence we have a valid test by using the thresholds $T_\alpha(\mathcal{E}_0)$, $T_\alpha(\mathcal{E}_U)$, $T_\alpha(\mathcal{E}_{\text{LUS}})$ or $T_\alpha(\mathcal{E}_{\text{LN}})$. Since $\mathcal{E}_0 \supseteq \mathcal{E}_U \supseteq \mathcal{E}_{\text{LUS}} \supseteq \mathcal{E}_{\text{LN}}$, we have $T_\alpha(\mathcal{E}_0) \geq T_\alpha(\mathcal{E}_U) \geq T_\alpha(\mathcal{E}_{\text{LUS}}) \geq T_\alpha(\mathcal{E}_{\text{LN}})$, so using $T_\alpha(\mathcal{E}_{\text{LN}})$ will yield the higher power out of the four sets of distributional hypotheses. Further, since $\{E_\mu : \mu > 0\}$ is a collection of comonotonic and log-normal e-variables, Lemma 2 states that we can also use the threshold of $T_\alpha(\mathcal{E}_{\text{LN}})$ on the supremum of the collection of e-variables. Finally, one may




Figure 7: Left: Simulated path for one run of S_n ; Right: E-processes and supremum process for the same run of S_n .

show that E_Φ has a decreasing density, so we may use the thresholds $T_\alpha(\mathcal{E}_0)$ or $T_\alpha(\mathcal{E}_D)$ to control the type-I error at $\alpha \in (0, 1)$. For this simulation study, we will use $\alpha = 0.05$; we recall that numerical values for the improved thresholds are provided in Table 1.

We next study the rejection rates for fixed n . Table 3 presents simulated rejection rates for 10,000 replications with type-I error control of $\alpha = 0.05$ when the simulated data comes from $N(0.3, 1)$. It also presents the number of observations required to obtain a rejection rate of β for $\beta \in \{0.5, 0.9, 0.95, 0.99\}$ (based on simulated data).

As expected, decreasing the threshold always improves the rejection rate. The improvement in rejection rate is especially large for low values of n when there is little evidence from data. It is usually the case that using improved thresholds gives more power than using the optional stopping property of e-processes (although the optional stopping property also enables one to continue gathering data, which the improved thresholds do not). Taking the supremum of comonotonic e-variables with the threshold $T_\alpha(\mathcal{E}_{LN})$ leads to the highest rejection rates among every test considered.

In Appendix G, we provide a table showing simulated rejection rates when data comes from the null. The type-I error is controlled at 0.05 for all tests. The test based on the supremum process using the improved threshold of $\mathcal{T}_\alpha(\mathcal{E}_{LN})$ achieves a simulated type-I error equal to the target level, implying that it is not wasteful at this level.

7.2 Improved threshold for universal inference

We next investigate the power increase we obtain by improving the threshold for universal inference e-tests. As in Wasserman et al. [28], we test the null hypothesis $N(0, 1)$, while the alternative hypothesis is that data comes from a mixture of two Gaussians, that is, $wN(\mu_1, \sigma_1) + (1 - w)N(\mu_2, \sigma_2)$ for unknown $(w, \mu_1, \mu_2, \sigma_1, \sigma_2)$. We construct the e-test using the universal inference e-variable, using 200 observations to estimate the parameters under the alternative and 200 observations to evaluate the likelihood ratio. In Figure 9, we present the histogram for 10,000 e-values under the null along with their log transforms. We observe that the e-values have a decreasing density; it is reasonable to assume that the corresponding e-variable belongs to \mathcal{E}_D . Also, since the density of positive log-transformed e-values is decreasing, we assume that e-variables belong to $\mathcal{E}_{LD>0}$.

In the following, we set the true model at $0.5N(-\mu, 1) + 0.5N(\mu, 1)$, where μ is a signal parameter. We control the type-I error at level $\alpha = 0.1$ using the thresholds $T_{0.1}(\mathcal{E}_0) = 10$, $T_{0.1}(\mathcal{E}_D) = 5$ and $T_{0.1}(\mathcal{E}_{LD>0}) = 4.07$. We simulate rejection rates from 10,000 replications and present rejection curves in Figure 10.




Figure 8: Average of the log e-process and supremum process.

The rejection rate for $\mu = 0$ is 0.0126 using a threshold of $T_{0.1}(\mathcal{E}_0)$, 0.0299 using a threshold of $T_{0.1}(\mathcal{E}_D)$, and 0.0365 for a threshold of $T_{0.1}(\mathcal{E}_{LD>0})$. For $\mu = 0.5$, we reject the null hypothesis 30% more often using the threshold $T_{0.1}(\mathcal{E}_D)$ compared with $T_{0.1}(\mathcal{E}_0)$, while the rejection rate is 40% higher when using the threshold $T_{0.1}(\mathcal{E}_{LD>0})$ compared with $T_{0.1}(\mathcal{E}_0)$.

In Appendix G.2 we conduct simulations for the alternative hypothesis $0.5N(\mu_1, 1) + 0.5N(\mu_2, 1)$ with unknown (μ_1, μ_2) , corresponding to the e-values in Figure 2 in the Introduction. The power improvement in that simulation study is similar to those discussed in this section.

7.3 Tests within the gamma family

We now illustrate using the technique derived in Section 4.3 to obtain maximum likelihood e-variables. Suppose we want to test between two gamma distributions (using the scale and rate parametrization). The parameter vector is $\theta = (\alpha, \beta)$ and the natural parameter (in the context of exponential families) is $\eta(\theta) = (\alpha - 1, -\beta)$.

For a null hypothesis $\alpha = \alpha_0$ and $\beta = \beta_0$, let us define the sets

- (a) $\Theta_1 = \{(\alpha, \beta) : \alpha > 0; \beta > 0\}$;
- (b) $\Theta_2 = \{(\alpha, \beta) : \alpha_0 < \alpha; 0 < \beta < \beta_0\}$;
- (c) $\Theta_3 = \{(\alpha, \beta) : 0 < \alpha < \alpha_0; \beta_0 < \beta\}$.

As mentioned in Section 4.3, since $\eta(\theta) - \eta(\theta_0)$ have the same sign for the choices $\Theta' = \Theta_2$ or $\Theta' = \Theta_3$, the collection of likelihood ratio e-variables $\{E_{\theta}(n) : \theta \in \Theta'\}$ is comonotonic. Letting $(\hat{\alpha}, \hat{\beta})$ denote the maximum likelihood estimate of (α, β) over Θ_2 or Θ_3 based on data (x_1, \dots, x_n) , we can use the statistic

$$Y(n) := \prod_{i=1}^n \frac{\Gamma(\alpha_0)}{\Gamma(\hat{\alpha})} \frac{\hat{\beta}^{\hat{\alpha}}}{\beta_0^{\alpha_0}} x_i^{\hat{\alpha} - \alpha_0} \exp\left\{-\left(\hat{\beta} - \beta_0\right)x_i\right\} \quad (7)$$

with rejection threshold $1/\alpha$ to construct a valid test, as discussed in Section 4.3. On the other hand, if $\Theta' = \Theta_1$, the collection of e-variables is no longer comonotonic and taking the supremum may not guarantee type-I error control.

Let us validate through simulations that the test based on the supremum of comonotonic e-variables controls the type-I error over different alternative hypotheses. We simulate observations from a gamma distribution with $\alpha_0 = 1$ and $\beta_0 = 1$ (this corresponds to an exponentially distributed random variable with mean 1). We test using the test statistic in (7) for n varying between 2 and 50. In Figure 11, we present

Table 3: Left-hand side: simulated rejection rates for tests up to n observations with type-I error control of $\alpha = 0.05$. Right-hand side: number of observations required to attain a simulated rejection rate of β with a type-I error control of 0.05. By default, tests are based on fixed n , while optional stopping (OS) tests consider the rolling maximum of the e-process.

Test	Threshold	n				β			
		10	50	100	500	0.5	0.9	0.95	0.99
Likelihood ratio $\mu = 0.2$	$\mathcal{T}_\alpha(\mathcal{E}_0)$	0	0.24	0.69	1	75	153	183	258
	$\mathcal{T}_\alpha(\mathcal{E}_U)$	0	0.41	0.80	1	58	129	160	234
	$\mathcal{T}_\alpha(\mathcal{E}_{LUS})$	0.01	0.49	0.84	1	51	120	150	226
	$\mathcal{T}_\alpha(\mathcal{E}_{LN})$	0.07	0.68	0.91	1	34	96	123	194
	OS+ $\mathcal{T}_\alpha(\mathcal{E}_0)$	0	0.29	0.77	1	67	135	162	229
Likelihood ratio $\mu = 0.3$	$\mathcal{T}_\alpha(\mathcal{E}_0)$	0	0.36	0.69	1	67	179	227	361
	$\mathcal{T}_\alpha(\mathcal{E}_U)$	0.03	0.49	0.77	1	52	158	206	321
	$\mathcal{T}_\alpha(\mathcal{E}_{LUS})$	0.05	0.54	0.80	1	45	145	193	312
	$\mathcal{T}_\alpha(\mathcal{E}_{LN})$	0.17	0.67	0.86	1	31	124	171	286
	OS+ $\mathcal{T}_\alpha(\mathcal{E}_0)$	0	0.46	0.80	1	54	138	177	272
Likelihood ratio $\mu = 0.4$	$\mathcal{T}_\alpha(\mathcal{E}_0)$	0.02	0.36	0.59	0.97	75	283	391	681
	$\mathcal{T}_\alpha(\mathcal{E}_U)$	0.07	0.46	0.66	0.98	59	257	362	647
	$\mathcal{T}_\alpha(\mathcal{E}_{LUS})$	0.10	0.50	0.69	0.98	51	247	347	633
	$\mathcal{T}_\alpha(\mathcal{E}_{LN})$	0.23	0.59	0.75	0.98	34	219	322	604
	OS+ $\mathcal{T}_\alpha(\mathcal{E}_0)$	0.03	0.49	0.75	0.99	51	179	246	451
Supremum	$\mathcal{T}_\alpha(\mathcal{E}_0)$	0.07	0.37	0.71	1	67	153	183	253
	$\mathcal{T}_\alpha(\mathcal{E}_U)$	0.12	0.49	0.80	1	52	128	159	221
	$\mathcal{T}_\alpha(\mathcal{E}_{LUS})$	0.15	0.54	0.84	1	45	120	147	207
	$\mathcal{T}_\alpha(\mathcal{E}_{LN})$	0.24	0.68	0.91	1	31	95	119	174
Mixture	$\mathcal{T}_\alpha(\mathcal{E}_0)$	0.02	0.14	0.39	1	122	240	280	371
	$\mathcal{T}_\alpha(\mathcal{E}_U)$	0.03	0.20	0.48	1	104	217	254	338
	OS+ $\mathcal{T}_\alpha(\mathcal{E}_0)$	0.04	0.25	0.53	1	95	211	249	333

simulated type-I errors at control levels 0.01 and 0.05 for 100,000 replications. When using the supremum test over Θ_2 or Θ_3 , we obtain valid type-I error control no matter the number of observations, validating Proposition 1. In that case, the simulated type-I error is about four times as small as the control level. On the other hand, the supremum test over Θ_1 is not valid since the collection $\{E_\theta(n) : \theta \in \Theta_1\}$ is not comonotonic. Indeed, we observe that the rejection thresholds for the supremum test over Θ_1 are above the control level for each value of n .

In Appendix G.3, we show that the supremum test based on the statistic in (7) achieves more power than the likelihood-ratio test for testing within the gamma family of distributions.

7.4 Multiple testing with boosted e-values

Next, we illustrate the boosting factors of the e-BH procedure in a simple setting. Suppose that each e-variable under the null is $\text{Exp}(1)$ distributed such that we can use the boosting factors derived in Section 6. Further, assume that each e-variable follows a gamma distribution if the null hypothesis is not true. We set the non-null distribution's shape parameter to $1 + \Theta$, and its rate parameter to $1/(1 + \Theta)$. To incorporate uncertainty into the alternative e-variable, we model Θ using an exponential distribution with a mean of b for $b \geq 0$. In other words, the e-variable under the alternative is given by the model

$$(E|\Theta = \theta) \stackrel{d}{\sim} \text{Gamma}(1 + \theta, 1/(1 + \theta)); \quad \Theta \stackrel{d}{\sim} \text{Exp}(1/b).$$

If $b = 0$ (i.e., $\Theta = 0$), then there is no signal. We have $\mathbb{E}[E] = \mathbb{E}[(1 + \Theta)^2] = 1 + 2b + 2b^2$.

We construct a set of K e-values (e_1, \dots, e_K) by simulating K_0 e-values from the null distribution and $K - K_0$ e-values from the alternative, where we set $K = 1000$ and $K_0 = 500$. We also construct (e_1, \dots, e_K) such that its dependence structure is given by a K -variate Gaussian copula where off-diagonal entries of the




Figure 9: Histogram of null universal inference e-values and log e-values in the Gaussian mixture example of Section 7.2.

correlation matrix are set to $-1/(K-1)$. For this dependence model, the assumption of PRDS does not hold.

We compare the base e-BH (Table 4a) and the boosted e-BH with the knowledge that null e-variables are log-concave (Table 4b) to understand how many more discoveries we can make by using the results from Section 6. Each number is an average from 1000 replications of the experiment. The base e-BH only makes discoveries for $\alpha = 0.1$ and $b = 5$; even in this case, the number of discoveries is very low. On the other hand, boosting the e-values by $c_1^{\text{AD}}(\alpha)$ makes discoveries when the signal or the acceptable FDR bound is high enough. Note that the realized FDP in Table 4b is still very low: the e-BH has a theoretical FDR guarantee $\alpha K_0/K$ (which is $\alpha/2$ in this example), but we observe a realized FDP much smaller. This observation on the conservative nature of the e-BH procedure is well known, because e-BH only has sharp FDR when the e-variables take certain two-point distributions (far away from our setting); see Wang and Ramdas [26] and Ren and Barber [19] for discussions. For further comparison, simulation results on the case of using full distribution of null e-variables and the corresponding p-BH procedure are given in Appendix G.4.

8 Conclusion

We have investigated methods to improve the power of e-tests in situations where we have some distributional knowledge of the e-variables. Distributional assumptions such as decreasing or unimodal densities can often be numerically verified or theoretically justified. In most relevant cases, the e-value threshold can be roughly improved by a factor of 2 or e . We also show that when a collection of e-variables is comonotonic, we can take the supremum over that collection and obtain a valid test. This enables us to improve upon split likelihood ratio tests and use the entire dataset to estimate and test. In simulation experiments, various improvements obtained in this paper lead to visibly more powerful tests than the standard methods.

References

- [1] Alimohammadi, M., Alamatsaz, M. H., and Cramer, E. (2016). Convolutions and generalization of logconcavity: Implications and applications. *Naval Research Logistics NRL*, **63**(2):109–123.
- [2] Bagnoli, M. and Bergstrom, T. (2005). Log-concave probability and its applications. *Economic Theory*, **26**(2):445–469.




Figure 10: Power for universal inference e-tests with and without improved thresholds.

- [3] Barber, R. F. and Candès, E. J. (2015). Controlling the false discovery rate via knockoffs. *Annals of Statistics*, **43**(5).
- [4] Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B*, **57**(1), 289–300.
- [5] Bernard, C., Kazzi, R. and Vanduffel, S. (2020). Range value-at-risk bounds for unimodal distributions under partial information. *Insurance: Mathematics and Economics*, **94**, 9–24.
- [6] Candès, E., Fan, Y., Janson, L. and Lv, J. (2018). Panning for gold: ‘Model-X’ knockoffs for high dimensional controlled variable selection. *Journal of the Royal Statistical Society Series B*, **80**(3):551–577.
- [7] Cuculescu, I. and Theodorescu, R. (1998). Multiplicative strong unimodality. *Australian & New Zealand Journal of Statistics*, **40**(2):205–214.
- [8] Gordon, R. D. (1941). Values of Mills’ ratio of area to bounding ordinate and of the normal probability integral for large values of the argument. *The Annals of Mathematical Statistics*, **12**(3):364–366.
- [9] Grünwald, P., de Heide, R. and Koolen, W. M. (2024). Safe testing. *Journal of the Royal Statistical Society Series B*, forthcoming, arXiv:1906.07801.
- [10] Guan, Y., Jiao, Z. and Wang, R. (2024). A reverse ES (CVaR) optimization formula. *North American Actuarial Journal*, forthcoming, arXiv:2203.02599.
- [11] Huber, M. (2019). Halving the bounds for the Markov, Chebyshev, and Chernoff Inequalities using smoothing. *American Mathematical Monthly*, **126**(10) 915–927.
- [12] Ibragimov, I. A. (1956). On the composition of unimodal distributions. *Theory of Probability & Its Applications*, **1**(2):255–260.
- [13] Ignatiadis, N., Wang, R. and Ramdas, A. (2024). E-values as unnormalized weights in multiple testing. *Biometrika*, **111**(2), 417–439.
- [14] Jeffreys, H. (1961). *Theory of Probability*. Oxford Classic Texts in the Physical Sciences. Oxford University Press, 3rd edition.
- [15] Lee, J. and Ren, Z. (2024). Boosting e-BH via conditional calibration. arXiv:2404.17562.




Figure 11: Simulated type-I errors for the gamma supremum test at levels 0.01 and 0.05. Full horizontal lines correspond to the type-I error control level.

- [16] Li, L., Shao, H., Wang, R. and Yang, J. (2018). Worst-case Range Value-at-Risk with partial information. *SIAM Journal on Financial Mathematics*, **9**(1), 190–218.
- [17] Marsiglietti, A. and Melbourne, J. (2022). Moments, concentration, and entropy of log-concave distributions. arXiv:2205.08293.
- [18] Ramdas, A., Grünwald, P., Vovk, V. and Shafer, G. (2023). Game-theoretic statistics and safe anytime-valid inference. *Statistical Science*, **38**(4), 576–601.
- [19] Ren, Z. and Barber, R. F. (2024). Derandomised knockoffs: Leveraging e-values for false discovery rate control. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, **86**(1):122–154.
- [20] Shafer, G. (2021). The language of betting as a strategy for statistical and scientific communication. *Journal of the Royal Statistical Society, Series A*, **184**(2), 407–431.
- [21] Shaked, M. and Shanthikumar, J. G. (2007). *Stochastic Orders*. Springer Series in Statistics.
- [22] Tse, T. and Davison, A. C. (2022). A note on universal inference. *Stat*, **11**(1), e501.
- [23] Vovk, V. and Wang, R. (2021). E-values: Calibration, combination, and applications. *Annals of Statistics*, **49**(3), 1736–1754.
- [24] Vovk, V. and Wang, R. (2024). Merging sequential e-values via martingales. *Electronic Journal of Statistics*, **18**(1), 1185–1205.
- [25] Wang, R. (2024). Testing with p*-values: Between p-values, mid p-values, and e-values. *Bernoulli*, **30**(2), 1313–1346.
- [26] Wang, R. and Ramdas, A. (2022). False discovery rate control with e-values. *Journal of the Royal Statistical Society Series B*, **84**(3), 822–852.
- [27] Waudby-Smith, I. and Ramdas, A. (2024). Estimating means of bounded random variables by betting. *Journal of the Royal Statistical Society Series B*, **86**(1), 1–27.
- [28] Wasserman, L., Ramdas, A. and Balakrishnan, S. (2020). Universal inference. *Proceedings of the National Academy of Sciences*, **117**(29), 16880–16890.

Table 4: Number of discoveries and realized FDP in percentage based on 1000 repetitions.

(a) The base e-BH procedure

α (%)	$b = 3$		$b = 4$		$b = 5$	
	Discov.	FDP (%)	Discov.	FDP (%)	Discov.	FDP (%)
1	0	0	0	0	0	0
2	0	0	0	0	0	0
5	0	0	0	0	0	0
10	0	0	0	0	6.91	0

(b) The boosted e-BH procedure with the knowledge of a log-concave survival function

α (%)	$b = 3$		$b = 4$		$b = 5$	
	Discov.	FDP (%)	Discov.	FDP (%)	Discov.	FDP (%)
1	0	0	65.91	0	208.70	0
2	0	0	133.38	0	233.66	0
5	5.55	0	196.16	0	269.32	0
10	65.11	0	234.48	0	294.65	0.001

Appendices

A Construction of e-values in the introduction

The null knockoffs in Figure 1 are constructed similarly to the simulation study in Section 4 of Ren and Barber [19]. To generate one e-value, we use a Gaussian linear model with 100 observations and 80 variables. The number of non-null variables is either 0 or 8. The joint distribution of the feature vector \mathbf{X} is Gaussian with mean 0 and covariance matrix Σ with entries $\Sigma_{ij} = 0.5^{|i-j|}$. The distribution of $Y|\mathbf{X}$ is given by a linear model with mean $\mathbf{X}^T\boldsymbol{\beta}$, where $\boldsymbol{\beta}$ is the vector of regression coefficients. In the case of zero non-nulls, we fix the vector $\boldsymbol{\beta} = \mathbf{0}$, a vector with 80 zeroes. When there are eight non-nulls, $\boldsymbol{\beta}$ contains eight values of one, that we distribute uniformly on the vector, with the remaining 72 entries set to zero. We construct knockoffs using the lasso coefficient difference statistic, with a target FDR level of 0.05. We draw 50 copies of knockoffs and report the average.

We construct the null universal inference e-values from Figure 2 similarly to the mixture model example in Section 3 of Wasserman et al. [28]. The null hypothesis is that data comes from $N(0, 1)$, while the alternate hypothesis is that data comes from $0.5N(\mu_1, 1) + 0.5N(\mu_2, 1)$, for unknown (μ_1, μ_2) . We use 400 observations: 200 to estimate the parameters under the alternative and 200 to compute the e-value from the likelihood ratio test statistic. We estimate the parameters from the alternate hypothesis using maximum likelihood estimation with the expectation-maximization algorithm.

B Proof of Theorem 2, parts (ii) and (iii)

We now prove parts (ii) and (iii) of Theorem 2. The proof arguments are similar to part (i) but more complicated.

- (ii) Take $E \in \mathcal{E}_{D>1}$ and $z \geq 1$. Let f be the density function of E . We construct an e-variable E_0 with density g_0 such that (a) $g_0(x) = f(z)$ for $1 \leq x < z$; (b) $g_0(x) = f(x)$ for $x \geq z$; (c) E_0 has a point-mass at 0 with probability $1 - \int_1^\infty g_0(x)dx$. Clearly, $\mathbb{E}[E_0] \leq \mathbb{E}[E]$ and $\mathbb{P}(E_0 \geq z) = \mathbb{P}(E \geq z)$. Construct another e-variable E_1 with density g_1 such that $g_1(x) = g_0(x)$ for $0 \leq x < z$ and $g_1(x) = f(z)$ for $z \leq x \leq z + \mathbb{P}(E \geq z)/f(z)$. Since f is decreasing for $x > z$, we have $\mathbb{E}[E_1] \leq \mathbb{E}[E_0]$ and $\mathbb{P}(E_1 \geq z) = \mathbb{P}(E_0 \geq z)$.

Therefore, for any $E \in \mathcal{E}_{D>1}$, there exists an e-variable $E' \in \mathcal{E}_{D>1}$ such that E' has a point-mass at 0 and a uniform density on $[1, b]$ for some constant b , and $\mathbb{P}(E' \geq z) = \mathbb{P}(E \geq z)$. To show $\mathbb{P}(E \geq z) \leq z - \sqrt{z^2 - 1}$, we can consider the collection of e-variables that have a point-mass at 0 and a uniform density on $[1, b]$ for any $b > 1$. Moreover, it suffices to consider the case $\mathbb{E}[E] = 1$.

Denote by c the constant density of E' over $[1, b]$ and note that $\mathbb{E}[E'] = \int_1^b cxdx = c(b^2 - 1)/2$. Therefore, $\mathbb{E}[E'] = 1$ implies $c = 2/(b^2 - 1)$. We have $\mathbb{P}(E' \geq z) = 0$ if $z > b$, and otherwise

$$\mathbb{P}(E' \geq z) = \int_z^b cdx = \int_z^b \frac{2}{b^2 - 1}dx = 2\frac{b - z}{b^2 - 1}. \quad (8)$$

Let us now find the constant b maximizing the right-hand side of (8). Taking the derivative with respect to b and setting the result equal to zero, we find that b is the solution to $b^2 - 2zb + 1 = 0$, or $b = z + \sqrt{z^2 - 1}$ (we take the positive solution to ensure $b > z$). Inserting the latter into the right hand side of (8), we have

$$\mathbb{P}(E' \geq z) \leq \frac{\sqrt{z^2 - 1}}{z^2 + z\sqrt{z^2 - 1} - 1} = z - \sqrt{z^2 - 1},$$

which, together with the attainability of this distribution, shows $R_\gamma(\mathcal{E}_{D>1}) = 1/\gamma - \sqrt{1/\gamma^2 - 1}$. The last equality is standard algebra.

- (iii) We first show $\mathbb{P}(E \geq 1/\gamma) \leq (\gamma/2) \vee (2\gamma - 1)$ for each $E \in \mathcal{E}_U$. Let m be the largest mode of E and f be the density function of E . Denote by $z = 1/\gamma$. We consider two cases separately.

First, consider the case that $z \leq m$. In this case, we construct an e-variable E_0 with density g_0 such that $g_0 = \int_0^z f(x)dx/z$ and E_0 has a point-mass at z with probability $1 - \int_0^z f(x)dx$. Since f is increasing on $[0, z]$, replacing f with g_0 on $[0, z]$ will decrease the mean of the distribution. Moreover, moving all probability mass of E on $[z, \infty)$ to the point z will decrease the mean. Hence, $\mathbb{E}[E_0] \leq \mathbb{E}[E]$. Note that $\mathbb{P}(E_0 \geq z) = \mathbb{P}(E \geq z)$, and $E_0 \in \mathcal{E}_U$.

Second, consider the case that $z > m$. In this case, consider another e-variable E_1 with density function g_1 such that (a) $g_1(x) = f(x)$ for $x \geq z$; (b) $g_1(x) = f(z) =: a$ for $x \in (m, z]$; (c) $g_1(x) = \int_0^m f(x)/m$ for $x \in [0, m)$; (d) E_1 has a point-mass at m with probability $p := 1 - \int_0^\infty g_1(x)dx$. Clearly, $\mathbb{P}(E \geq z) = \mathbb{P}(E_1 \geq z)$. We will verify that $\mathbb{E}[E_1] \leq \mathbb{E}[E] \leq 1$. This is similar to what has been argued above. Note that on $x \in [0, m)$, replacing the density f with the constant density g_1 will decrease the mean, since f is an increasing function on $[0, m)$. Moreover, on $[m, z)$, replacing f with the point-mass and g_1 will also decrease the mean. Putting both observations together, we have $\mathbb{E}[E_1] \leq \mathbb{E}[E]$. Further, E_1 also has a unimodal density.

Next, construct e-variable E_2 with density function g_2 such that (a) $g_2(x) = g_1(x)$ for $x > m$; (b) $g_2(x) = g_1(x) + p/m =: b$ for $x \in [0, m)$. In other words, the point-mass at m of E_1 is replaced by a uniform density on $[0, m)$. Certainly, $\mathbb{E}[E_2] \leq \mathbb{E}[E_1] \leq 1$. By construction, $\mathbb{P}(E_1 \geq z) = \mathbb{P}(E_2 \geq z)$.

If either $m = 0$ or $b \geq a$, then E_2 has a decreasing density. Hence, using part (i) of Theorem 2, we have $\mathbb{P}(E_2 \geq z) \leq 1/(2z)$. This gives $\mathbb{P}(E \geq z) = \mathbb{P}(E_2 \geq z) \leq 1/(2z)$.

Next, assume $b < a$. In this case, construct e-variable E_3 with density function g_3 such that (a) $g_3(x) = g_2(x)$ for $x \geq z$; (b) $g_3(x) = \int_0^z g_2(x)/z =: d$ for $x \in [0, z)$. Using the same argument above, we have $\mathbb{E}[E_3] \leq \mathbb{E}[E_2]$, and E_3 has a unimodal density with mode at z . By construction, $\mathbb{P}(E_2 \geq z) = \mathbb{P}(E_3 \geq z)$.

Construct e-variable E_4 with density function g_4 such that $g_4(x) = g_3(x)$ for $x \in [0, z)$ and E_4 has a point-mass at z with probability $\int_z^\infty g_3(x)dx$. We have $\mathbb{E}[E_4] \leq \mathbb{E}[E_3]$ and $E_4 \in \mathcal{E}_U$ and by construction, $\mathbb{P}(E_3 \geq z) = \mathbb{P}(E_4 \geq z)$.

The above arguments in two cases together show that for any $E \in \mathcal{E}_U$, either $\mathbb{P}(E \geq z) \leq 1/(2z)$ or there exists $E' \in \mathcal{E}_U$ such that E' has a uniform density on $[0, z)$ and a point-mass at z , and $\mathbb{P}(E' \geq z) = \mathbb{P}(E \geq z)$. This E' is E_0 in the case $z \leq m$ and E_4 in the case $z > m$. It suffices to analyze $q := \mathbb{P}(E' \geq z)$. Note that $\mathbb{E}[E'] = zq + (1 - q)z/2 \leq 1$, giving rise to $q \leq 2/z - 1$. Hence, in this case, $\mathbb{P}(E \geq 1/\gamma) = \mathbb{P}(E' \geq 1/\gamma) \leq 2\gamma - 1$. Putting the above two cases together we have $\mathbb{P}(E \geq 1/\gamma) \leq (\gamma/2) \vee (2\gamma - 1)$.

To show the converse statement $R_\gamma(\mathcal{E}_U) \geq (\gamma/2) \vee (2\gamma - 1)$, it suffices to verify the two inequalities separately, namely, $R_\gamma(\mathcal{E}_U) \geq \gamma/2$ for $\gamma \in (0, 2/3]$ and $R_\gamma(\mathcal{E}_U) \geq 2\gamma - 1$ for $\gamma \in (2/3, 1]$. The first inequality follows from part (i) of Theorem 2, which gives $R_\gamma(\mathcal{E}_U) \geq R_\gamma(\mathcal{E}_D) = \gamma/2$ for $\gamma \in (0, 1)$. To show the second inequality, for $z \in [1, 2]$, construct E with a uniform density on $[0, z)$ and a point-mass at z and with probability $q := 2/z - 1$, we get $\mathbb{E}[E] = zq + (1 - q)z/2 = 1$. Hence, $E \in \mathcal{E}_U$, and by choosing $\gamma = 1/z \in [1/2, 1]$, we get $\mathbb{P}(E \geq 1/\gamma) = 2\gamma - 1$, showing the second inequality.

C Proof of Theorem 3

In this section, we provide proofs for the five results in Theorem 3.

- (i) Let $x \geq \log 2$, $a > 0$ and $p < e^{-x} \leq 1/2$. Consider a three point distribution at x with probability p , at $x - a$ with probability $1 - 2p$ and at $x - 2a$ with probability p . Clearly, this distribution is symmetric, and we let X follow this distribution. We can compute $\mathbb{E}[e^X] = pe^x + pe^{x-2a} + (1-2p)e^{x-a}$. Sending $a \rightarrow \infty$ gives $\mathbb{E}[e^X] \rightarrow pe^x < 1$. Therefore, $e^X \in \mathcal{E}_{LS}$ for sufficiently large a . Note that $\mathbb{P}(e^X \geq e^x) = p$. Since p is an arbitrary number smaller than e^{-x} , we get $R_\gamma(\mathcal{E}_{LS}) \geq \gamma$ for all $\gamma \leq 1/2$, and it holds as an equality due to Markov's inequality. Next, suppose that X has any symmetric distribution and $\mathbb{E}[e^X] \leq 1$. Jensen's inequality gives $\mathbb{E}[e^X] \geq e^{\mathbb{E}[X]}$. Hence, $\mathbb{E}[X] \leq 0$. This implies $\mathbb{P}(e^X > 1) = \mathbb{P}(X > 0) \leq 1/2$. This gives $R_\gamma(\mathcal{E}_{LS}) = \gamma \wedge (1/2)$. The last statement $R_1(\mathcal{E}_{LS}) = 1$ is easily verified by taking $X = 0$.

- (ii) Let $m > 0$ be large, $z = -\log \gamma$ and $p \in (0, \gamma)$, and let X follow the distribution given by combining a uniform distribution on $[z - m, z]$ with density p/m and a point mass at z of probability $1 - p$. This distribution is unimodal, and $\mathbb{E}[e^X]$ can be easily computed as

$$\mathbb{E}[e^X] = (1 - p)e^z + \frac{p}{m} \int_{z-m}^z e^x dx = (1 - p)\frac{1}{\gamma} + \frac{p}{m}(e^z - e^{z-m}).$$

It is clear that for $\mathbb{E}[e^X] \leq 1$ by sending $m \rightarrow \infty$ $\mathbb{E}[e^X] < 1$. Note that $\mathbb{P}(e^X \geq 1/\gamma) = p$ and p can be any number smaller than γ . This shows $R_\gamma(\mathcal{E}_{\text{LU}}) \geq \gamma$. The other direction $R_\gamma(\mathcal{E}_{\text{LU}}) \leq \gamma$ is Markov's inequality.

- (iii) Let $E \in \mathcal{E}_{\text{LD}>0}$ and $X = \log E$. Let g denote the density function of X and f denote the density function of E . We have that $f(x) = g(\log x)/x$ for $x \geq 0$. Note that if $E \in \mathcal{E}_{\text{LD}>0}$, then $g'(x) < 0$ for $x > 0$. Further, for $x > 0$, we have that

$$f'(x) = g'(\log x)/x^2 - g(\log x)/x^2 < 0,$$

implying $\mathcal{E}_{\text{LD}>0} \subseteq \mathcal{E}_{\text{D}>1}$, which proves the upper bound.

Suppose we have a random variable Y with distribution function H . If we construct a new random variable Y' with distribution function H' by shifting some mass to the left, then $\mathbb{P}(Y' \leq x) \geq \mathbb{P}(Y \leq x)$, so Y dominates Y' under the usual stochastic order and $\mathbb{E}[\phi(Y')] \leq \mathbb{E}[\phi(Y)]$ for any increasing function ϕ .

Let us now show that $R_\gamma(\mathcal{E}_{\text{LD}>0}) \leq e^{-a\gamma}$. Start with a random variable $X \in \mathcal{E}_{\text{LD}>0}$ with density function g and let $z = 1/\gamma$. Let $m \leq \inf(\text{supp}(X))$, that is, m is smaller than the left-hand support of X (if $\inf(\text{supp}(X)) = -\infty$, then we mean that m approaches $-\infty$). Construct a new random variable X_0 with density function g_0 such that (a) $g_0(x) = g(x)$ for $x > \log z$; (b) $g_0(x) = g(\log z)$ for $0 \leq x \leq \log z$; (c) X_0 has a point-mass at m . Then $\mathbb{P}(X_0 \geq \log z) = \mathbb{P}(X \geq \log z)$ and X_0 is smaller than X under the usual stochastic order since we shifted mass to the left.

Construct another random variable X_1 with density function g_1 such that (a) $g_1(x) = g_0(x)$ for $x \leq \log z$; (b) $g_1(x) = g(\log z)$ for $\log z \leq x \leq \log z + \mathbb{P}(X \geq \log z)/g(\log z)$. Then $\mathbb{P}(X_1 \geq \log z) = \mathbb{P}(X_0 \geq \log z)$ and X_1 is smaller than X_0 under the usual stochastic order since we shifted mass to the left.

This shows that if $E \in \mathcal{E}_{\text{LD}>0}$, then we can construct a random variable $E' \in \mathcal{E}_{\text{LD}>0}$ such that $\log E'$ has a point mass m approaching $-\infty$ and a constant density over $[0, a]$ for some $a \geq \log z$, and $\log E'$ is smaller than $\log E$ under the usual stochastic order, implying that $\mathbb{E}[E'] \leq \mathbb{E}[E]$.

Denote by c the constant density of $\log E'$ over $[0, a]$ and $p = \mathbb{P}(\log E' = m)$. We have that $\mathbb{E}[E'] = \mathbb{E}[e^{\log E'}] = pe^m + c(e^a - 1)$. Since E' is an e-variable and since m approaches $-\infty$, we have $c \leq 1/(e^a - 1)$. This implies that, for any $E \in \mathcal{E}_{\text{LD}>0}$,

$$\mathbb{P}(E \geq z) = \int_{\log z}^a c dx \leq \int_{\log z}^a 1/(e^a - 1) dx = \frac{a - \log z}{e^a - 1}. \quad (9)$$

We obtain the supremum of the right-hand side of (9) over $a > \log z$, which is the solution to the equation $e^a(1 - a + \log z) = 1$. Alternatively, using the substitution $t = e^a/z$, we are seek the solution for t in $t(1 - \log t) = 1/z$ for $t > 1$, that we denote by $t_{1/z}$. Note that

$$\frac{d}{dt}(t(1 - \log t)) = -\log t,$$

implying that $t \mapsto t(1 - \log t)$ is monotonically decreasing and $t_{1/z}$ has a unique solution for $t > 1$. It is easy to verify that e is the solution to $t(1 - \log t) = 1/z$ when $1/z \downarrow 0$, hence $t_{1/z} \in (1, e)$. Replacing $t_{1/z}$ into (9) and simplifying yields $\mathbb{P}(E \geq z) \leq 1/(zt_{1/z})$.

To show the upper bound holds with equality, fix some $z > 1$. We construct an e-variable E such that $\log E$ has a constant density of $1/(t_{1/z}z - 1)$ over $[0, \log(t_{1/z}z)]$, where $t_{1/z}$ satisfies $t_{1/z}(1 - \log t) = 1/z$, and the remaining mass approaching $-\infty$. Then, $\mathbb{P}(E \geq z) = e^{-a_{1/z}}$.

To show the lower bound, we note that $R_\gamma(\mathcal{E}_{\text{LD}>0})$ maximizes $(a + \log \gamma)/(e^a - 1)$, so any $a \in (-\log \gamma, \infty)$ will give a lower bound for $R_\gamma(\mathcal{E}_{\text{LD}>0})$. We start by the observation that $\gamma \mapsto a_\gamma$ is decreasing, so e^{a_γ} is large for small γ , implying that $1 - a_\gamma - \log \gamma$ is a small positive number. We approximate a_γ by the solution for a in $1 - a - \log \gamma = 0$, which leads to $a = 1 - \log \gamma$. Replacing this in the right-hand side of (9) leads to the lower bound.

- (iv) Let $m \geq 1$ and $c > 0$. Consider the uniform distribution $F_{m,c}$ on $[m - 1/c, m]$, and let X follow this distribution. Clearly, $F_{m,c}$ is a unimodal and symmetric distribution, and it has a constant, thus decreasing, density. We can compute

$$\mathbb{E}[e^X] = \int_{m-1/c}^m ce^x dx = ce^m(1 - e^{-1/c}) \leq ce^m.$$

By taking $c = e^{-m}$, we know that $e^X \in \mathcal{E}_{\text{LUS}} \cap \mathcal{E}_{\text{LD}}$. For $x \geq 0$, we have $\mathbb{P}(e^X \geq e^x) = c(m - x) = e^{-m}(m - x)$. By taking $m = x + 1$ we get

$$\mathbb{P}(e^X \geq e^x) = e^{-x-1}.$$

Since $e^X \in \mathcal{E}_{\text{LUS}}$, we get $R_\gamma(\mathcal{E}_{\text{LUS}}) \geq \gamma/e$ and $R_\gamma(\mathcal{E}_{\text{LD}}) \geq \gamma/e$.

Next, we show the upper bound. We will first argue that both \mathcal{E}_{LUS} and \mathcal{E}_{LD} admit the same class of worst-case distributions. Fix $x > 0$. For any $E \in \mathcal{E}_{\text{LUS}}$ or $E \in \mathcal{E}_{\text{LD}}$, let $X = \log E$. Denote the distribution of X by F and its density function by g .

First, we argue the case of \mathcal{E}_{LUS} . Denote by x_0 the median of F , which is unique. We consider another distribution G , such that G has a point mass at x_0 and a uniform density on $[x_0 - a, x_0 + a]$ for some $a > 0$. Let g be the density function of G outside $\{x_0\}$. Let Y follow the distribution G . We choose a and the point mass at x_0 such that $f(x) = g(x)$ and $F(x) = G(x)$. It is clear that such G exists, and it is straightforward to check that G is dominated by F in increasing convex order (see e.g., Theorem 3.A.44 of Shaked and Shanthikumar [21]), that is,

$$\int_{\mathbb{R}} u dF \geq \int_{\mathbb{R}} u dG \quad \text{for all increasing convex functions } u : \mathbb{R} \rightarrow \mathbb{R}.$$

It follows that $\mathbb{E}[e^Y] \leq \mathbb{E}[e^X] \leq 1$, and thus $e^Y \in \mathcal{E}_{\text{LUS}}$. Moreover, $\mathbb{P}(Y \geq x) = \mathbb{P}(X \geq x)$. Therefore, it suffices to consider distributions of the form G . Keeping the density g at x fixed, we can increase a and remove the point-mass at x_0 . This will reduce the mean of Y and does not change $\mathbb{P}(Y \geq x)$. Therefore, it suffices to consider uniform distribution on some interval $[m - 1/c, m]$, thus the distribution $F_{m,c}$ above.

Next, consider the case of \mathcal{E}_{LD} , which is simpler. Let Y follow the distribution $G = F_{m,c}$ with density g , such that $G(x) = F(x)$ and $g(x) = f(x)$. It is straightforward to check that F dominates G in increasing convex order, and $e^Y \in \mathcal{E}_{\text{LD}}$. Hence, we, again, only need to consider $F_{m,c}$ above. Moreover, $\mathcal{E}_{\text{LUS}} = \mathcal{E}_{\text{LD}}$.

For $Y \stackrel{d}{\sim} F_{m,c}$, note that $\mathbb{P}(Y > x) = c(m - x)_+$. To find the largest value of this probability, it is clear that $m > x$. Since the median of Y is less than or equal to 0, as we argued above, we know $1/c \geq 2m \geq 2x$. Therefore, by using $ce^m(1 - e^{-1/c}) \leq 1$, we get $ce^m(1 - e^{-2x}) \leq 1$, and thus $c \leq e^{-m}(1 - e^{-2x})^{-1}$. This leads to

$$\mathbb{P}(Y > x) = c(m - x)_+ \leq \sup_{m \geq x} e^{-m}(1 - e^{-2x})^{-1}(m - x) = \frac{e^{-x-1}}{1 - e^{-2x}}.$$

This shows

$$R_\gamma(\mathcal{E}_{\text{LUS}}) \leq \frac{\gamma/e}{1 - \gamma^2}.$$

The remaining inequality $R_\gamma(\mathcal{E}_{\text{LD}}) \leq R_\gamma(\mathcal{E}_{\text{LD}})$ follows from $\mathcal{E}_{\text{LD}} \subseteq \mathcal{E}_{\text{LD}>0} \subseteq \mathcal{E}_{\text{D}>1}$.

- (v) Clearly, if $\gamma = 1$, then $R_1(\mathcal{E}_{\text{LN}}) = 1$ by choosing the trivial degenerate e-variable $E = 1$. Next, let $\sigma > 0$ and X_σ follow $\text{LN}(-\sigma^2/2, \sigma^2)$, which has mean 1. For $x > 1$, we have

$$\begin{aligned} \sup_{\sigma>0} \mathbb{P}(X_\sigma \geq x) &= \sup_{\sigma>0} \Phi\left(\frac{-\log x - \sigma^2/2}{\sigma}\right) \\ &= \Phi\left(\sup_{\sigma>0} \left(-\frac{\log x}{\sigma} - \frac{\sigma}{2}\right)\right) = \Phi\left(-\sqrt{2\log x}\right). \end{aligned}$$

This shows $R_\gamma(\mathcal{E}_{\text{LN}}) = \Phi(-\sqrt{-2\log \gamma})$.

We can provide an explicit upper bound by using the inequality,

$$\left(\frac{1}{x} - \frac{1}{x^3}\right) \phi(x) < \Phi(-x) < \frac{1}{x} \phi(x),$$

for $x > 0$, where ϕ is the standard normal pdf (see, e.g., Gordon [8]). It follows from part (v) of Theorem 3 that for $\gamma < 1$,

$$R_\gamma(\mathcal{E}_{\text{LN}}) = \Phi(-\sqrt{-2\log \gamma}) \leq \frac{\gamma}{2\sqrt{-\pi \log \gamma}}.$$

D Proof of Theorem 4

In this section, we provide proofs for the three results in Theorem 4.

- (i) Let $E \in \mathcal{E}_{\text{LCS}}$ and denote by G its survival function. Note that log-concavity implies for $x, y \geq 0$ and $\theta \in [0, 1]$,

$$G(\theta x + (1 - \theta)y) \geq G(x)^\theta G(y)^{1-\theta}.$$

Moreover, log-concave survival functions cannot jump at 0, hence $G(0) = 0$. Taking $y = 0$, $x > 1$ and $\theta = z/x$ with $z \in (0, x]$, we get

$$G(z) \geq G(x)^{z/x}.$$

For fixed $x > 1$, denote by $t_x = (\log G(x))/x$, which is negative by Markov's inequality. We have

$$\mathbb{E}[E] = \int_0^\infty G(z) dz \geq \int_0^x G(x)^{z/x} dz = \int_0^x e^{t_x z} dz = \frac{e^{x t_x} - 1}{t_x} \leq 1.$$

Therefore,

$$e^{x t_x} \geq t_x + 1.$$

Note that the above inequality always holds true if $t_x \leq -1$. Below, we consider the case $t_x > -1$. We note that

$$e^{x t} - t - 1 = 0.$$

has a unique solution for $t \in (-1, 0)$. Note that $e^{x t} - t - 1 > 0$ for $t = -1$ and $e^{x t} - t - 1 < 0$ for t close to 0. Strict convexity guarantees a unique solution on $(-1, 0)$, denoted by t_x^* . The above argument also shows $t_x \leq t_x^*$, and hence $G(x) = e^{x t_x} \leq e^{x t_x^*}$ for all $x > 1$. Taking $\gamma = 1/x$ and $s_\gamma = t_x^*$ proves $R_\gamma(\mathcal{E}_{\text{LCS}}) \leq e^{s_\gamma/\gamma}$.

To show that the above inequality holds as an equality, let $G(z) = e^{t_x^* z}$ for $z < x$, and $G(z) = 0$ for $z \in [0, x]$. It is clear that G is a log-concave survival function, and the corresponding random variable E has mean 1, and thus $E \in \mathcal{E}_{\text{LCS}}$. For any $z \in [0, x)$, we have $\mathbb{P}(E > z) = e^{t_x^* z}$. By continuity, with $\gamma = 1/x$, we get $R_\gamma(\mathcal{E}_{\text{LCS}}) \geq e^{t_x^* x} = e^{s_\gamma/\gamma}$. Thus, the first equality in the proposition holds.

To show the remaining inequality, by using

$$e^{-x t_x} (t_x + 1) \leq 1,$$

and $t_x \in (-1, 0)$, we get $(1 - x t_x)(t_x + 1) \leq 1$, which is $-x t_x^2 + t_x - x t_x \leq 0$, leading to $-x t_x \geq x - 1$. Thus $t_x \leq (1 - x)/x$, yielding $G(x) = e^{x t_x} \leq e^{1-x}$ for all $x > 1$. Taking $\gamma = 1/x$ proves the desired inequality.

- (ii) The first inequality follows by checking E following an exponential distribution with mean 1. The relations $\mathcal{E}_{\text{LCD}} \subseteq \mathcal{E}_{\text{U}}$ and $\mathcal{E}_{\text{LCD}} \subseteq \mathcal{E}_{\text{LCS}}$ give the second inequality. The last inequality is shown in part (i) of Theorem 4.
- (iii) Let $m > 1$, $b \in (0, 1)$ and $\lambda > 0$. Consider the cumulative distribution function F given by

$$F(x) = (1 - b) + be^{-\lambda(m-x)_+} \text{ for } x \geq 0.$$

It is straightforward to verify that the distribution function F is log-concave, and its survival function is given by $x \mapsto b(1 - e^{-\lambda(m-x)})$ on $[0, m]$. We further choose the parameters so that F has mean 1, that is,

$$1 = \int_0^m b(1 - e^{-\lambda(m-x)})dx = b \left(m - \frac{1 - e^{-\lambda m}}{\lambda} \right),$$

and this is equivalent to $b = \lambda/(m\lambda - 1 + e^{-\lambda m})$. For any fixed $m > 1$, if we send $\lambda \rightarrow \infty$, then $b \rightarrow 1/m \in (0, 1)$. Therefore, for any fixed m we can take λ large enough so that F is the distribution function of some $E \in \mathcal{E}_{\text{LCF}}$. Hence, for $x \in [0, m)$,

$$\mathbb{P}(E > x) = \frac{\lambda(1 - e^{-\lambda(m-x)})}{m\lambda - 1 + e^{-\lambda m}}.$$

Sending $\lambda \rightarrow \infty$ we get $\mathbb{P}(E > x) \rightarrow 1/m$. This implies that, for any $x \in [1, m)$, we have $R_{1/x}(\mathcal{E}_{\text{LCF}}) \geq 1/m$. Since m is arbitrary, this leads to $R_\gamma(\mathcal{E}_{\text{LCF}}) \geq \gamma$ for $\gamma \in (0, 1]$. Together with Markov's inequality, implying $R_\gamma(\mathcal{E}_{\text{LCF}}) \leq \gamma$, we get the desired statement.

E Weighted average of independent e-values

We now consider the weighted average of independent e-variables. Note that a weighted average of e-variables is an e-variable regardless of the dependence structure, and using the weighted average is the only admissible way of merging arbitrarily dependent e-values as shown by Vovk and Wang [23]. However, the following result relies on the independence assumption for an improved threshold.

Proposition 7. *Let $E_t \in \mathcal{E}_{\text{LCD}}$ for $t \in [T]$ be independent e-variables, (w_1, \dots, w_T) is in the standard simplex, and $M_T = \sum_{t=1}^T w_t E_t$. Denote by $w_{(1)} = \min_t w_t$. For $\gamma \in (0, 1]$,*

$$\mathbb{P}(M_T \geq 1/\gamma) \leq \left(\gamma^{-w_{(1)}} e^{-w_{(1)}(1/\gamma-1)} \right) \wedge R_\gamma(\mathcal{E}_{\text{U}}).$$

Proof. From Theorem 1.6 of Marsiglietti and Melbourne [17], we have $\mathbb{P}(M_T \geq t) \leq e^{-w_{(1)}(t-1-\log t)}$ for all $t \geq 1$. Replacing $t = 1/\gamma$ for $\gamma \in (0, 1]$ completes the first bound. If $E \in \mathcal{E}_{\text{LCD}}$ and w is a non-zero real constant, then $wE \in \mathcal{E}_{\text{LCD}}$. Further, since the convolution of log-concave density functions is unimodal (see, for instance, Ibragimov [12]), we have $M_t \in \mathcal{E}_{\text{U}}$, showing the second bound. \square

We provide in Table 5 some improved thresholds for the average of independent e-variables with log-concave densities (using equal weights for each e-variable). We provide thresholds based on Proposition 7, but note that for $T = 1$, the bound in Part (ii) of Theorem 3 holds and gives a smaller threshold in Table 1. The exponential bound provides the lower threshold for small values of T and α , while the unimodal bound takes over in other situations.

F Proofs for results in Section 6

We will require the following lemma for the proof of Proposition 5.

Lemma 3. *For $E \in \mathcal{E}_{\text{LCS}}$ and $t > 0$, we have $\mathbb{E}[E|E > t] \leq \mathbb{E}[E] + t$.*

Table 5: Improved thresholds for the average of T e-variables with log-concave densities.

T	α					
	0.001	0.01	0.02	0.05	0.1	0.2
1	10.23	7.64	6.83	5.74	4.9	2.5
2	17.69	12.76	11.24	9.21	5.0	2.5
5	39.21	27.33	23.73	10.00	5.0	2.5
10	74.39	50.00	25.00	10.00	5.0	2.5
20	144.13	50.00	25.00	10.00	5.0	2.5

Proof. Since $E \in \mathcal{E}_{\text{LCS}}$, we have from Theorem 3.B.19 (b) of Shaked and Shanthikumar [21] that $(E-t|E > t)$ is decreasing in t in the sense of the dispersive order. It follows from Theorem 3.B.13 (a) of Shaked and Shanthikumar [21] that $(E-t|E > t)$ is smaller than E under the usual stochastic order. Therefore, $\mathbb{E}[E-t|E > t] \leq \mathbb{E}[E] \leq 1$ and $\mathbb{E}[E|E > t] \leq 1+t$. \square

Proof of Proposition 5. Notice first that

$$\mathbb{E}[\alpha b E \mathbf{1}_{\{\alpha b E \geq 1\}}] = \alpha b \mathbb{P}\left(E \geq \frac{1}{\alpha b}\right) \mathbb{E}\left[E \mid E \geq \frac{1}{\alpha b}\right]. \quad (10)$$

We start with the upper bound. If E is exponentially distributed with mean 1, (10) becomes

$$\mathbb{E}[\alpha b E \mathbf{1}_{\{\alpha b E \geq 1\}}] = \alpha b \exp\{-1/(\alpha b)\} \left(1 + \frac{1}{\alpha b}\right).$$

It follows that the problem in (3) becomes

$$\exp\{-1/(\alpha b)\} (\alpha b + 1) = \alpha,$$

and solving for b leads to $c_2^{\text{AD}}(\alpha)$, which gives an attainable boosting factor for a random variable in \mathcal{E}_{LS} and proves the upper bound for $B_\alpha^{\text{AD}}(\mathcal{E}_{\text{LCS}})$.

For the lower bound, let us first find an upper bound for the quantity in (10). We may bound the probability term in (10) by the bound in part (i) of Theorem 4 and bound the conditional expectation term by its upper bound of $1 + 1/(\alpha b)$; the latter bound follows from Lemma 3. This gives us

$$\mathbb{E}[\alpha b E \mathbf{1}_{\{\alpha b E \geq 1\}}] \leq \alpha b \exp\left\{1 - \frac{1}{\alpha b}\right\} \left(1 + \frac{1}{\alpha b}\right).$$

Solving for b in

$$\alpha b \exp\left\{1 - \frac{1}{\alpha b}\right\} \left(1 + \frac{1}{\alpha b}\right) = \alpha$$

proves the lower bound for $B_\alpha^{\text{AD}}(\mathcal{E}_{\text{LCS}})$. \square

Proof of Proposition 6. We have for any $x \geq 1/(\alpha b)$ that

$$x \mathbb{P}(\alpha b E \geq x) = x \mathbb{P}\left(E \geq \frac{x}{\alpha b}\right) \leq x \exp\left\{1 - \frac{x}{\alpha b}\right\},$$

where the inequality is due to part (i) of Theorem 4. Note that $x \mapsto x \exp\left\{1 - \frac{x}{\alpha b}\right\}$ is unimodal with a mode at αb , so it is a decreasing function of x whenever $x \geq \alpha b$. Suppose that $1 \geq \alpha b$, then we have

$$\max_{x \geq 1} x \mathbb{P}(\alpha b E \geq x) \leq \max_{x \geq 1} x \exp\left\{1 - \frac{x}{\alpha b}\right\} \leq \exp\left\{1 - \frac{1}{\alpha b}\right\} \leq \alpha.$$

Solving the equation on the right-hand side with equality leads to $c_1^{\text{PR}}(\alpha)$. One may verify that $1 \geq \alpha c_1^{\text{AD}}(\alpha)$ whenever $\alpha \in (0, 1]$. Suppose that $1 \leq \alpha b$, then

$$\max_{x \geq 1} x \mathbb{P}(\alpha b E \geq x) \leq \max_{x \geq 1} x \exp\left\{1 - \frac{x}{\alpha b}\right\} \leq \alpha b \exp\left\{1 - \frac{\alpha b}{\alpha b}\right\} = \alpha b.$$

Table 6: Simulated rejection rates when testing within the Gaussian family when data comes from the null $N(0, 1)$, with a type-I error control of $\alpha = 0.05$.

Test	Threshold	n				
		10	50	100	500	
$\mu = 0.2$	$\mathcal{T}_\alpha(\mathcal{E}_0)$	0.00	0.00	0.01	0.00	
	Likelihood	$\mathcal{T}_\alpha(\mathcal{E}_U)$	0.00	0.01	0.02	0.00
	ratio	$\mathcal{T}_\alpha(\mathcal{E}_{LUS})$	0.00	0.02	0.02	0.00
		$\mathcal{T}_\alpha(\mathcal{E}_{LN})$	0.01	0.05	0.05	0.01
		OS+ $\mathcal{T}_\alpha(\mathcal{E}_0)$	0.00	0.01	0.02	0.04
$\mu = 0.3$	$\mathcal{T}_\alpha(\mathcal{E}_0)$	0.00	0.01	0.01	0.00	
	Likelihood	$\mathcal{T}_\alpha(\mathcal{E}_U)$	0.00	0.02	0.01	0.00
	ratio	$\mathcal{T}_\alpha(\mathcal{E}_{LUS})$	0.01	0.02	0.01	0.00
		$\mathcal{T}_\alpha(\mathcal{E}_{LN})$	0.03	0.04	0.02	0.00
		OS+ $\mathcal{T}_\alpha(\mathcal{E}_0)$	0.00	0.02	0.03	0.04
$\mu = 0.4$	$\mathcal{T}_\alpha(\mathcal{E}_0)$	0.00	0.01	0.00	0.00	
	Likelihood	$\mathcal{T}_\alpha(\mathcal{E}_U)$	0.01	0.01	0.01	0.00
	ratio	$\mathcal{T}_\alpha(\mathcal{E}_{LUS})$	0.01	0.02	0.01	0.00
		$\mathcal{T}_\alpha(\mathcal{E}_{LN})$	0.04	0.03	0.01	0.00
		OS+ $\mathcal{T}_\alpha(\mathcal{E}_0)$	0.00	0.03	0.04	0.04
Supremum comonotonic	$\mathcal{T}_\alpha(\mathcal{E}_0)$	0.01	0.01	0.01	0.01	
		$\mathcal{T}_\alpha(\mathcal{E}_U)$	0.02	0.02	0.02	0.02
		$\mathcal{T}_\alpha(\mathcal{E}_{LUS})$	0.02	0.02	0.02	0.02
		$\mathcal{T}_\alpha(\mathcal{E}_{LN})$	0.05	0.05	0.05	0.05
Mixture	$\mathcal{T}_\alpha(\mathcal{E}_0)$	0.00	0.00	0.00	0.00	
		$\mathcal{T}_\alpha(\mathcal{E}_U)$	0.01	0.00	0.00	0.00
		OS+ $\mathcal{T}_\alpha(\mathcal{E}_0)$	0.01	0.02	0.03	0.03

We, therefore, require $b = 1$ to satisfy (4). Note that we don't have $1 \geq \alpha b$ for $\alpha \in (0, 1]$, so a lower bound of $B_\alpha^{\text{AD}}(\mathcal{E}_{\text{LCS}})$ is $c_1^{\text{PR}}(\alpha)$ for $\alpha \in (0, 1]$. As for the upper bound, finding an attainable boosting factor for a random variable $E \in \mathcal{E}_{\text{LCS}}$ suffices. Letting E be exponentially distributed with mean 1, we must find b such that

$$\max_{x \geq 1} x \mathbb{P}(\alpha b E \geq x) = \max_{x \geq 1} x \exp\{-x/(\alpha b)\} = \alpha. \quad (11)$$

The function $x \mapsto x \exp\{-x/(\alpha b)\}$ is unimodal, where its mode is located at αb . When $1 \leq \alpha b$, the solution to (11) is $-1/(\alpha \log(\alpha))$, which is valid when $1 \leq -\alpha/(\alpha \log(\alpha))$, or equivalently, when $\alpha \leq \exp\{-1\}$. When $1 \leq \alpha b$, the solution to (11) is $b = \exp\{1\}$; this occurs when $\alpha > \exp\{-1\}$. \square

G Additional simulation results

G.1 Tests within the Gaussian family, continued

We showed in Section 7.1 that improved thresholds and taking the supremum of e-values lead to power improvements. Let us now validate that the tests control the type-I errors through simulations. In Table 6, we present simulated rejection rates when data comes from the null $N(0, 1)$ using the tests from Section 7.1. The type-I error is controlled at 0.05 for all tests. The type-I error is close to the control level for optional stopping strategies as n increases, while the type-I error for the likelihood ratio e-processes decreases to 0 as n increases. The test based on the supremum process using the improved threshold of $\mathcal{T}_\alpha(\mathcal{E}_{LN})$ achieves a simulated type-I error equal to the target level, implying that it is not wasteful at this level.

G.2 Improved threshold for universal inference, continued

In Section 7.2, we conducted a simulation study for universal inference e-tests when testing the null hypothesis $N(0, 1)$ against the alternative that data comes from a mixture of two Gaussians. The motivating example




Figure 12: Power for universal inference e-tests with and without improved thresholds for the simplified Gaussian mixture model.

in the Introduction uses a different setup, resulting in the e-values presented in Figure 2. The difference between the setup of Section 7.2 and in the Introduction is that the alternate hypothesis in the Introduction was $0.5N(\mu_1, 1) + 0.5N(\mu_2, 1)$ for unknown (μ_1, μ_2) , that is, the weights and variances in the Gaussian mixture was assumed known. We repeat the simulation study for this setup for completeness.

In Figure 2, the e-values have a decreasing density over $[1, \infty)$, while the log-transformed e-values have a decreasing density over its positive domain. It is, therefore, reasonable to assume that the e-variables belong to $\mathcal{E}_{D>1}$ and $\mathcal{E}_{LD>0}$.

Figure 12 shows power curves when the data comes from $0.5N(-\mu, 1) + 0.5N(\mu, 1)$, where μ is a signal parameter. For $\mu = 0.5$, using the threshold of $T_{0.1}(\mathcal{E}_{D>1})$ leads to 26% more power compared with $T_{0.1}(\mathcal{E}_0)$. Similarly, using the threshold $T_{0.1}(\mathcal{E}_{LD>0})$ gives 35% more power compared with the Markov bound.

G.3 Tests within the gamma family, continued

Let us return to the simulation study outlined in Section 7.3, where the null hypothesis is a gamma distribution with parameters $\alpha_0 = 1$ and $\beta_0 = 1$ such that the random variable's mean under the null is 1. Assume now that data comes from a gamma distribution with parameters $\alpha = 1.1$ and $\beta = 0.9$, such that the random variable's mean is $11/9$. We use two test statistics to test the null hypothesis $\alpha_0 = \beta_0 = 1$. The first is the likelihood ratio e-variable using the true alternative that $\alpha = 1.1$ and $\beta = 0.9$. The second is against the composite alternative hypothesis that $(\alpha, \beta) \in \Theta_2 = \{(\alpha, \beta) : \alpha_0 < \alpha; 0 < \beta < \beta_0\}$ using the supremum test statistic. We present power curves in Figure 13. Using the supremum test over the likelihood ratio test enhances power by a factor of 1.81 for $n = 50$, 1.17 for $n = 100$, 1.09 for $n = 200$ and 1.03 for $n = 500$. To achieve a power of 0.9, we require around 268 observations using the supremum test and 341 observations using the likelihood ratio e-test.

G.4 The boosted e-BH, continued

To understand how well the boosted e-BH procedure performs, we compare the method with another e-BH procedure, but with the knowledge that the null e-variables are exponentially distributed with mean 1 (Table 7a). We also compare with the p-BH procedure, where we construct the p-values p_k for the p-BH procedure as $\exp\{-e_k\}$ for $k \in \mathcal{K}$ (Table 7b). In this example, the PRDS property is not satisfied by the p-values (p_1, \dots, p_K) , so the p-BH procedure does not have a theoretically proven FDR control. Indeed, some numbers are slightly higher than $\alpha K_0/K$.




Figure 13: Simulated power when testing within the gamma family. The likelihood ratio e-test is against the true alternative, while the supremum test is against the alternative $\alpha > \alpha_0$ and $\beta_0 > \beta > 0$.

Table 7: Number of discoveries and realized FDP in percentage based on 1000 repetitions.

(a) The boosted e-BH procedure with the knowledge of the null e-variable distribution

α (%)	$b = 3$		$b = 4$		$b = 5$	
	Discov.	FDP (%)	Discov.	FDP (%)	Discov.	FDP (%)
1	0	0	157.99	0	246.13	0
2	7.25	0	200.71	0	271.78	0
5	118.05	0	252.11	0.001	309.00	0.003
10	204.29	0.002	290.73	0.024	336.71	0.062

(b) The p-BH procedure with the knowledge of the null e-variable distribution

α (%)	$b = 3$		$b = 4$		$b = 5$	
	Discov.	FDP (%)	Discov.	FDP (%)	Discov.	FDP (%)
1	298.58	0.503	340.52	0.488	368.48	0.490
2	316.79	0.997	355.90	0.999	381.70	1.018
5	346.01	2.562	381.84	2.479	405.95	2.533
10	379.14	5.045	410.86	4.973	432.12	5.001