

Data-Driven Parametrization of Molecular Mechanics Force Fields for Expansive Chemical Space Coverage

Tianze Zheng,^{*,†} Ailun Wang,[†] Xu Han,[†] Yu Xia,[†] Xingyuan Xu,[†] Jiawei Zhan,^{†,‡}
Yu Liu,[†] Yang Chen,[†] Zhi Wang,[†] Xiaojie Wu,[†] Sheng Gong,[†] and Wen Yan^{*,†}

[†]*ByteDance Research*

[‡]*work done as intern at ByteDance Research*

E-mail: zhengtianze@bytedance.com; wen.yan@bytedance.com

Abstract

A force field is a critical component in molecular dynamics simulations for computational drug discovery. It must achieve high accuracy within the constraints of molecular mechanics' (MM) limited functional forms, which offers high computational efficiency. With the rapid expansion of synthetically accessible chemical space, traditional look-up table approaches face significant challenges. In this study, we address this issue using a modern data-driven approach, developing ByteFF, an Amber-compatible force field for drug-like molecules. To create ByteFF, we generated an expansive and highly diverse molecular dataset at the B3LYP-D3(BJ)/DZVP level of theory. This dataset includes 2.4 million optimized molecular fragment geometries with analytical Hessian matrices, along with 3.2 million torsion profiles. We then trained an edge-augmented, symmetry-preserving molecular graph neural network (GNN) on this dataset, employing a carefully optimized training strategy. Our model predicts all bonded and non-bonded MM force field parameters for drug-like molecules simultaneously across a broad chemical space.

ByteFF demonstrates state-of-the-art performance on various benchmark datasets, excelling in predicting relaxed geometries, torsional energy profiles, and conformational energies and forces. Its exceptional accuracy and expansive chemical space coverage make ByteFF a valuable tool for multiple stages of computational drug discovery.

Introduction

Drug discovery involves identifying potential therapeutic candidates within the vast and intricate landscape known as chemical space, which encompasses all possible molecular structures.¹⁻³ As a pivotal tool in this process, molecular dynamics (MD) simulations offer insights into dynamical behaviors and physical properties of molecules, as well as interactions in the molecular systems at an atomic level.⁴⁻⁸ Central to the accuracy and reliability of these simulations is the force field, a mathematical model that describes the potential energy surface (PES) of the molecular system as a function of the positions of the atoms involved.⁹⁻¹² Recent advances in synthetic chemistry and high-throughput screening technologies have significantly expanded the chemical space for drug candidates,¹³⁻¹⁷ which necessitates the development of force fields that can provide accurate predictions of PES for diverse molecules in expansive chemical space.

Generally, force fields can be classified into two main categories: conventional molecular mechanics force fields (MMFFs),^{9-12,18,19} which parameterize a fixed analytical form to approximate the energy landscape, and machine learning force fields (MLFFs),²⁰⁻²³ that aim to map the atomistic and molecular features and coordinates to the PES using neural networks without being limited by the fixed functional forms. Most conventional MMFFs, such as Amber,^{24,25} GAFF⁹ and OPLS,^{11,12,26} describe the molecular PES by decomposing it into various degrees of freedom, including bonded (i.e., bonds, angles, and torsions) and non-bonded interactions (i.e., electrostatics and dispersion). These conventional MMFFs benefit from the computational efficiency of these terms, while suffering inaccuracies due to the inherent approximation, especially when non-pairwise additivity of non-bonded interac-

tions are of significant importance. Emerging recently, MLFFs have shown great promise for modeling PES due to their ability to capture subtle interactions and complex behaviors that may be overlooked or oversimplified by classical models.²⁰⁻²³ Despite their outstanding accuracies, several drawbacks limit their applications in drug discovery. Owing to the complexity of the machine learning models involved, the computational efficiency of MLFFs is relatively low. Meanwhile, the amount of data required to train an effective MLFF is extremely large, which imposes constraints on their ability to cover the chemical space comprehensively. Consequently, conventional MMFFs remain the most reliable and commonly used tool for MD simulations involving biological systems to this day.^{7,8}

In the past few years, several efforts have been made to improve the quality of MMFFs for predicting the PES of small molecules. Following the traditional look-up table approach, OPLS3e increased the number of torsion types to 146,669 to enhance accuracy and expand chemical space coverage.¹¹ Furthermore, OPLS3e and its successor OPLS4 were empowered with FFBuilder¹¹ to refine the torsion terms for molecules beyond the coverage of the pre-determined torsion list. OpenFF^{18,19,27} took a different approach by utilizing SMIRKS patterns to describe the chemical environment of both bonded and Lennard-Jones terms. However, these discrete descriptions of the chemical environment have inherent limitations that hamper the transferability and scalability of these force fields. To tackle this problem, Espaloma^{28,29} introduced a novel end-to-end workflow where the MMFF parameters are predicted by graph neural networks (GNN), opening new avenues for the advancement of MMFFs.³⁰ Despite promising results, these early attempts are constrained by the ML techniques and the training data, where significant improvements can be achieved.

In this work, we propose ByteFF, a data-driven MMFF trained on a large-scale, high-diversity, and high-quality quantum mechanics (QM) dataset with sophisticated ML techniques. ByteFF is designed to leverage both atom and bond features using a state-of-the-art GNN model, while preserving molecular symmetry. To construct the dataset for training ByteFF, we employ novel fragmentation methods and a rigorous QM calculation workflow,

generating 2.4 million optimized molecular fragment geometries with Hessian matrices, along with 3.2 million torsion profiles. Additionally, we introduce a differentiable partial Hessian loss and an iterative optimization-and-training procedure to effectively train ByteFF on the dataset. Finally, we demonstrate the performance of ByteFF on various benchmarks, showing its expansive chemical space coverage and exceptional accuracy on intra-molecular conformational PES.

Methods

Molecular Mechanics Force Field

In this work, we follow the analytical forms in GAFF⁹ and OpenFF:²⁷

$$\begin{aligned}
 E^{\text{MM}} &= E_{\text{bonded}}^{\text{MM}} + E_{\text{non-bonded}}^{\text{MM}} \\
 E_{\text{bonded}}^{\text{MM}} &= E_{\text{bond}}^{\text{MM}} + E_{\text{angle}}^{\text{MM}} + E_{\text{proper}}^{\text{MM}} + E_{\text{improper}}^{\text{MM}} \\
 &= \sum_{\text{bonds}} \frac{1}{2} k_{r,ij} (r_{ij} - r_{ij}^0)^2 + \sum_{\text{angles}} \frac{1}{2} k_{\theta,ijk} (\theta_{ijk} - \theta_{ijk}^0)^2 \\
 &\quad + \sum_{\text{probers}} \sum_{n_\phi} k_{\phi,ijkl}^{n_\phi} \left[1 + \cos(n_\phi \phi_{ijkl} - \phi_{ijkl}^{n_\phi,0}) \right] \\
 &\quad + \sum_{\text{improbers}} \sum_{n_\psi} k_{\psi,ijkl}^{n_\psi} \left[1 + \cos(n_\psi \psi_{ijkl} - \psi_{ijkl}^{n_\psi,0}) \right] \\
 E_{\text{non-bonded}}^{\text{MM}} &= E_{\text{vdW}}^{\text{MM}} + E_{\text{Coulomb}}^{\text{MM}} \\
 &= \sum_{i < j} \left[4\epsilon_{ij} \left(\frac{\sigma_{ij}^{12}}{r_{ij}^{12}} - \frac{\sigma_{ij}^6}{r_{ij}^6} \right) + \frac{q_i q_j}{4\pi\epsilon_0 r_{ij}} \right]
 \end{aligned} \tag{1}$$

where the bond lengths r , angles θ , proper torsion angles ϕ and improper torsion angles ψ are rotationally invariant internal coordinates. The force field parameters include bonded parameters (equilibrium values r^0 , θ^0 , ϕ^0 and ψ^0 , and force constants k_r , k_θ , k_ϕ and k_ψ), and non-bonded parameters (van der Waals (vdW) parameters σ and ϵ , and partial charges q). A well-trained transferrable MMFF shall predict these parameters accurately for any given

molecules. In this article, "torsion" refers to proper torsion unless otherwise specified. We also fix the phase angles $\phi_{ijkl}^{n_\phi,0}$ and $\psi_{ijkl}^{n_\psi,0}$ at 0° for odd n and 180° for even n , and therefore ensure that the torsional energy is independent of the order being $ijkl$ or $lkji$.

Force field parameters should adhere to several physical constraints: (1) Force field parameters should be permutationally invariant, e.g., the force constant of bond (i, j) should be equal to that of bond (j, i) . (2) Force field parameters should be in accordance with chemical symmetries of molecules, e.g., force constants of the two C-O bonds in a carboxyl group ($-\text{C}([\text{O}-])=\text{O}$) must be equal to each other, due to their chemical equivalency, even though they may have different bond orders assigned when written as a SMILES or SMARTS string.³¹ (3) Charge conservation should be guaranteed, i.e., the summation of assigned partial charges of atoms in one molecule should be consistent with the molecule’s net charge, which avoids net charge gain/loss for the molecule in the parameter determination. These constraints are naturally satisfied in the traditional look-up table approaches, which should also serve as essential guidelines when force field parameters are inferred by a machine-learning model.

Additionally, there are two key philosophies worth considering when building general small molecule MMFFs: (1) Force field parameters should be dominated by local structures, so that the parameters trained from small molecules can be consistently transferred to similar structures in relatively large molecules.^{9,11} (2) Torsional energy profiles should be accurately captured, because the quality of torsion parameters strongly affects the protein-ligand binding affinity predictions.^{11,32,33}

Dataset Construction

Molecular fragments generation

The curated dataset is mostly built from the ChEMBL database³⁴ with some additions from the ZINC20³⁵ datasets to further enhance the diversity. From these datasets, a subset of molecules was initially selected by several criteria, including number of aromatic rings, polar surface area (PSA), quantitative estimate of drug-likeness (QED), element types, and hy-

bridization types. These selected molecules were then cleaved into fragments with less than 70 atoms using our in-house graph-expansion algorithm (details in supporting information), such that the local chemical environments were well-preserved.^{11,36} In brief, this fragmentation algorithm traverses over each bond, angle, and non-ring torsion in a molecule, retains the relevant atoms and their conjugated partners, then trims the rest and caps the cleaved bonds. Next, these fragments were expanded to various protonation states within a pKa range of 0.0 to 14.0, calculated by Epik 6.5,³⁷ to cover most possible protonation states that might appear in aqueous solutions. Finally, 2.4 million unique fragments were selected for QM calculations after deduplication.

Quantum chemistry methods and workflow

From the 2.4 million fragments, we created two QM datasets, namely the *optimization dataset* and the *torsion dataset*, at the B3LYP-D3(BJ)/DZVP level of theory. The same QM method is also used by OpenFF to generate their training data.^{18,19} This method achieves a good balance between accuracy (relative to CCSD(T)/CBS) and computational cost in recent benchmarks.^{38,39} For molecular conformational PES, more advanced methods like ω B97M-v are significantly more expensive but only marginally more accurate.

The *optimization dataset* was generated from the entire 2.4 million fragments. The 3D conformations of each fragment was initially generated by RDKit from its SMILES string, and then optimized using the geomeTRIC⁴⁰ optimizer at the chosen QM level. The relaxed structure is verified where no accidental bond breaking or formation happened during the structural relaxation. Finally, the Hessian matrix is calculated for each fragment using Q-Chem 6.⁴¹ The hessian matrix is further verified by checking the six smallest eigenvalues to ensure that a true local minimum is captured. More details about the workflow and the filtering criteria are provided in the supporting information.

The *torsion dataset* consists of two subsets, non-ring torsions and in-ring torsions, curated separately. For each non-ring torsion fragment, we first rotate the torsion angle in 15°

increments from the optimized conformation, creating 24 initial frames that were then separately relaxed using the geomeTRIC optimizer with the rotated torsion angle constrained. While the in-ring torsions were scanned using a sequential frame-by-frame approach, with an early stopping strategy if the conformational energy is more than 20 kcal/mol higher than the unconstrained relaxed conformation, since those high-energy regimes are irrelevant in room-temperature molecular dynamics simulations. For both non-ring and in-ring subsets, the resultant conformers were also carefully filtered by the bond breaking or formation criteria. Due to the high computational cost of torsion scan, these calculations were performed with GPU4PySCF⁴² for significant acceleration and cost reduction. The results were also verified to be consistent with Q-Chem.

Diversity of Chemical Space

Evaluating the diversity of chemical space composed of numerous molecules often involves representing molecules with descriptors or fingerprints followed by measuring of their distances. One widely used fingerprint is the Morgan fingerprint, which captures the local environment of each atom within a specified radius defined by the number of covalent bonds. To emphasize the local structure of torsions, Morgan fingerprints were calculated with a radius of 2, focusing exclusively on the neighborhoods of the two central atoms in the selected torsion. This torsional fingerprint was used to evaluate the diversity of different datasets, including SPICE, GEOM, and our *torsion dataset*, as well as in the creation of a diverse torsion dataset (BDTorsion).

ByteFF Model

ByteFF predicts both the bonded and non-bonded force field terms in one-pass. It leverages the state-of-the-art Edge-augmented Graph Transformer (EGT)⁴³ as its backbone to achieve accuracy and transferability. The output of EGT is further processed by a series of MLPs to derive force field parameters while preserving both the total charge and the physical

symmetry of each force field term. The ML model structure is shown in Fig. 1 and consists of three major parts: featurization, GNN, and output layers. They are briefly described here, and more details are given in the supporting information.




Figure 1: **Model structure of ByteFF.** ByteFF predicts MMFF parameters in three steps. First, atom and bond features are extracted from the molecular graph and then projected into embeddings. Then, an edge-augmented graph transformer (EGT)⁴³ is used to synergize the edge embeddings with the node-based attention mechanism. Lastly, the output MLPs derive force field parameters while preserving the molecular symmetry and total charge.

Featurization layers. For each molecule, we first convert the molecule structure into a graph, and detect chemically equivalent atoms and bonds with bond orders ignored. Atomic features are then detected from this graph, including the element type, formal charge, ring connectivity and minimum ring size. Additionally, to better describe the chemical environ-

ment of each atom, bond features including bond order and whether the bond is in ring, are also extracted. Replacing the discrete atom-typing rules used in conventional MMFFs, atom and bond features are projected into continuous vectorial embedding spaces, and concatenated into node and edge feature embeddings, respectively. Importantly, embeddings of chemically equivalent atoms and bonds are averaged before passed to the next layers, which helps to preserve the chemical symmetry of the molecule.

GNN layers. In the GNN layers, a modified EGT architecture is employed to learn the chemical environment of molecules from both atom and bond features. This architecture synergizes the node and edge features in the attention mechanism to efficiently capture the structural information of the molecule. To speed up training and inference, the original global self-attention in EGT is localized such that attention is applied only to a local neighborhood of each node.

Output module. After a few layers of localized EGT, atom and bond hidden states generated by GNN layers are passed to the output layers to predict each force field term individually. For each bonded term, the corresponding output block is designed to preserve the necessary symmetry, respectively. For the partial charge term, the output block is designed to preserve the total charge of the molecule following a bond-charge-correction (BCC) style postprocessing.

Additionally, an ensemble of three models randomly initialized is trained for improved predictive performance and uncertainty quantification.

Training Procedures

To efficiently use the curated *optimization* and *torsion datasets*, while enhancing the robustness and performance of the ByteFF model, we took an ingenious training strategy with three stages: pre-training, training, and fine-tuning. The loss functions (details in supporting information) are carefully designed in each stage such that the improvement of the force field is properly reflected by the minimization of loss functions.

In the pre-training stage, the non-bonded parameters and force constants of proper torsions were fitted to GAFF2 with mean squared error (MSE) losses, while the equilibrium values of bonded parameters were fitted with energy-based loss functions, using the *optimization dataset*. Meanwhile, the force constants of bonded terms in Eq. 1 were trained using the partial Hessian loss, evaluated by the mean absolute percentage error (MAPE). The design of the partial Hessian loss combines the training accuracy with the computational efficiency, which takes the form of Hessian matrices but only fit the blocks corresponding to the target bonds, angles and improper torsions, namely partial Hessian blocks.

Given the significance of torsion profiles in the quality of force fields, in the training stage, we incorporated the curated *torsion dataset* to fit the force constants of proper torsions, while the other parameters were trained using *optimization dataset* following the same manner in the pre-training stage. Here, we employed the Boltzmann MSE loss functions⁴⁴ into an iterative optimization-and-training process to train the force constants of proper torsions using the *torsion dataset*. In each iteration, the QM-optimized conformations were refined by the force field, with the torsion angle constrained and atoms restrained. The parameters were then trained on both the *optimization* and *torsion datasets*. As the force field’s accuracy improved, the positional restraint force constant was gradually reduced. Additionally, L1-norm regularization was applied to the force constants of proper torsions to restrain redundant degrees of freedom. The combination of pre-training and training stages yields the ByteFF-gopt model.

In the final fine-tuning stage, we incorporated part of the training set in Espaloma-0.3.0,²⁹ named *off-equilibrium dataset* in this work, to refine the force field parameters with the QM energy and forces. Combining all the three stages (pre-training, training, and fine-tuning), the ByteFF-joint model was obtained.

The detailed settings of the three-stage training procedure are summarized in Table 1.

Table 1: Summary of training settings for different stages

	Pre-training	Training	Fine-tuning
Model parameters initialized	randomly initialized parameters	parameters from pre-training stage	parameters from training stage
Datasets and fitting targets	<i>optimization dataset:</i> - GAFF $\sigma/\epsilon/q/k_\phi$ - $E_{\text{bond}}^{\text{MM}} / E_{\text{angle}}^{\text{MM}} / E_{\text{improper}}^{\text{MM}}$ - QM Hessian matrix	<i>optimization dataset:</i> - GAFF $\sigma/\epsilon/q$ - $E_{\text{bond}}^{\text{MM}} / E_{\text{angle}}^{\text{MM}} / E_{\text{improper}}^{\text{MM}}$ - Hessian matrix <i>torsion dataset:</i> - QM energy - L1-norm of k_ϕ	<i>optimization dataset:</i> same as training <i>torsion dataset:</i> same as training <i>off-equilibrium dataset:</i> - QM energy/force
Optimization-and-training strategy	no force field optimization	three optimization-and-train iterations on <i>torsion dataset</i>	using coordinates from the last iteration of training stage
Optimizer	RAdam	RAdam	RAdam
Learning rate	10^{-4}	10^{-4}	2×10^{-5}
Scheduler	ReduceLROnPlateau factor: 0.2 patience: 10 threshold: 10^{-6}	ReduceLROnPlateau factor: 0.2 patience: 4 threshold: 10^{-6}	ReduceLROnPlateau factor: 0.2 patience: 4 threshold: 10^{-6}
Early stop patience	50	10	10
Yield	-	ByteFF-gopt	ByteFF-joint

Benchmark Datasets and Metrics

To be compatible with the Amber family of protein force fields, the non-bonded terms of ByteFF are trained to reproduce the am1bcc charges and GAFF vdW parameters. Therefore, in this work we focus on the intramolecular PES performance relative to QM references of ByteFF, and leave the further improvement of non-bonded terms as a future work.

Three different types of benchmarks are used to provide a thorough assessment of ByteFF. First, the large-scale industrial collaborative OpenFFBenchmark public dataset⁴⁵ is used to quantify how well the force field-relaxed geometries and energies reproduce the QM-relaxed references, without constraints. Three different metrics are used, namely root-mean-square deviation of atomic positions (RMSD), torsion fingerprint deviation (TFD),⁴⁶ and relative energy difference ($\Delta\Delta E$) defined by OpenFF team.⁴⁷ Second, the crucial torsional energy profile performance is benchmarked on two different datasets, namely the publicly available TorsionNet500 dataset⁴⁸ and the BDTorsion dataset curated in-house. For consistency,

energies of conformations in TorsionNet500⁴⁸ dataset were re-calculated with the B3LYP-D3(BJ)/DZVP level of theory. On these datasets, the root mean squared error (RMSE) and Boltzmann-weighted RMSE of each torsional energy profile are calculated and compared.

Beyond the constrained (torsional energy) and unconstrained (OpenFFBenchmark) optimization benchmarks, the performance of ByteFF is further benchmarked on the QM references energies and forces calculated on the *off-equilibrium dataset*, which includes subsets of the SPICE dataset⁴⁹ and the RNA Structure Atlas.⁵⁰ On this dataset, RMSE of energies and forces were calculated to quantify if the PES is properly captured beyond local minima.

More details of these benchmarks can be found in supporting information.

Results and Discussion

Diversity of Various Datasets

Using the Morgan-based torsional fingerprint defined above, we quantify the diversity of SPICE, GEOM and BDTorsion datasets in terms of torsional local environment. Specifically, in all three datasets, only torsions with circular standard deviation⁵¹ greater than 0.3 are included and used in fingerprint analysis. The diversity of each dataset was then visualized using the standard t-SNE (t-Distributed Stochastic Neighbor Embedding) algorithm. In Fig. 2, the scatter dots are colored by the element types of the two center atoms (the rotatable bond) in each torsion. The most frequent rotatable bond is C-C, followed by C-N rotatable bonds and other bonds including C-O, C-S, etc. It is evident that our *torsion dataset* excels in the diversity and comprehensiveness of its chemical space coverage, while the SPICE and GEOM datasets are more sparse. In comparison, both SPICE and GEOM datasets show obvious vacancies, corresponding to the absence of certain chemical torsion patterns. For example, the uncovered areas around (-45, -20) in Fig. 2 (a) and (b) correspond to in-ring torsions of protonated pyridine and imidazole, as well as their derivatives, which are indispensable for the training of force fields for bio-organic molecules. Such an expansive

coverage of the chemical space in the training dataset provides a solid foundation for the coverage and transferability of ByteFF.




Figure 2: **t-SNE analysis of different datasets.** The Morgan-based torsional fingerprint analysis results are illustrated using the t-SNE algorithm. Every scatter dot corresponds to a torsion profile being analyzed in the corresponding dataset, which is colored by the element types of the two center atoms in the torsion.

Torsional Potential Energy Surfaces

To evaluate the accuracy of ByteFF in predicting torsional PES, and directly compare with the performance of other force fields, we performed comprehensive benchmarks using the TorsionNet500 and BDTorsion datasets. TorsionNet500⁴⁸ is a benchmark dataset consisting of 500 chemically diverse fragments relevant to biological and pharmaceutical applications. The BDTorsion dataset is curated in this work. It is divided into Non-Ring and In-Ring torsion subsets and each contains 1000 chemically diverse fragments. For both TorsionNet500 and BDTorsion datasets, dozens of conformations are sampled by scanning torsion angle, then QM-relaxed, and tagged with QM energy labels for each fragment. Here, two metrics were used to evaluate the accuracy of the predictions from various force fields with respect to the ground truth from QM: the RMSE and Boltzmann RMSE, with RMSE assessing the overall discrepancy while Boltzmann RMSE emphasizing the deviations near the energy minimum. When calculating the Boltzmann RMSE, we used 2.0 kcal/mol for both cutoff and scaling factors.




Figure 3: **Histograms of the discrepancy of torsional PES between predictions of QM and force fields.** As a comprehensive benchmark, two metrics including Boltzmann RMSE (a-c) and RMSE (d-f) are used to assess the accuracy of force field-predicted torsional energy profiles with respect to the QM results. Three datasets were included in this benchmark: TorsionNet500 (a & d), BDTorsion-NonRing (b & e), and BDTorsion-InRing (c & f).

As illustrated in Fig. 3, both ByteFF-gopt and ByteFF-joint significantly outperform competitors on all three benchmark datasets. On all the three datasets, ByteFF-gopt exhibits the best performance on both metrics, with the highest population of predictions with lower RMSE and Boltzmann RMSE. When tested on TorsionNet500, the Boltzmann RMSE of most predictions by ByteFF-gopt are within 0.5 kcal/mol, while none of the Boltzmann RMSE exceeds 1.17 kcal/mol, demonstrating the exceptional accuracy of ByteFF-gopt in terms of predicting torsional energy profiles. On the more challenging BDTorsion datasets, both RMSE and Boltzmann RMSE are generally higher for all the force fields being tested. Nevertheless, ByteFF-gopt and ByteFF-joint still outperformed competitors significantly, with most Boltzmann RMSE values below 1.0 kcal/mol (chemical accuracy). In these torsion profile benchmarks, the performance of ByteFF-joint is compromised by the addition of off-

equilibrium energies and forces training data. This is inevitable due to the very limited fixed functional form of MMFFs.




Figure 4: **Example of the in-ring torsion prediction accuracy of various force fields.** (a) 2D representation of the example molecule. (b) The comparison of torsional energy profiles predicted by various force fields and calculated from QM. (c) The zoomed-in comparison of torsional energy profiles predicted with torsion angles between -175° and -75° .

One key feature of organic molecules is the presence of a great variety of rings,⁵² and in-ring torsion parameters have crucial impact on the conformational PES of rings. However, this is much less discussed in the development of MMFFs. In this work, we carefully constructed the in-ring torsion scan training dataset to improve the performance of ByteFF on in-ring torsions. We also curated the BDTorsion-InRing benchmark dataset, which consists of 1000 chemically diverse fragments with QM calculated in-ring torsion profiles. As shown in Fig. 3 (c) and (f), both ByteFF-gopt and ByteFF-joint significantly outperform competitors, with both RMSE and Boltzmann RMSE mostly better than chemical accuracy, respectively. As an example to illustrate the accuracy of ByteFF-gopt and ByteFF-joint in predicting in-ring torsional energy profile, we present a molecule from the BDTorsion-InRing dataset that contains a four-membered aliphatic ring (Fig. 4 (a)). On this molecule, we calculated the torsional energy profile for the in-ring torsion atoms highlighted in Fig. 4 (a), and compared PES predictions from various force fields with the QM references (Fig. 4 (b) and (c)). It is evident that both Espaloma-0.3.0 (green) and OpenFF-2.0.0 (blue) failed to predict the energy landscape in the vicinity of the local minimum, which could lead to incorrect predictions of ring conformations in geometric optimization or MD simulations. Though GAFF-2.2 captured the approximate shape of this torsional energy landscape, it overesti-

mated the barrier near -130° while significantly underestimated the energy near -160° , which led to significantly incorrect shifts of the predicted location of global energy minimum. In contrast, both ByteFF-gopt and ByteFF-joint predictions align well with QM references, which not only captured the shape of the energy landscape, but also accurately predicted the positions and energy differences of local minima.

The effectiveness and accuracy of ByteFF-gopt and ByteFF-joint stems from the synergy of two components: comprehensive in-house training datasets and carefully tweaked training strategy. The iterative optimization-and-training strategy allows the model to accurately capture the energy landscape along torsional degrees of freedom, minimizing interference from complex interactions such as bond-torsion coupling that is beyond the limited description of the chosen functional form. The predictions are further enhanced by the simple but effective model-ensemble averages.

Equilibrium and Off-Equilibrium Conformations

In addition to predict the torsional PES of molecules, the OpenFFBenchmark⁴⁵ dataset is used to benchmark the performance of various force fields in terms of predicting unconstrained equilibrium conformations and associated energies.




Figure 5: **Histograms of different metrics on OpenFFBenchmark.** The accuracy of force field-relaxed geometry relative to QM-relaxed references is quantified with (a) RMSD and (b) Torsion Fingerprint Deviation (TFD) scores. The energetic accuracy is quantified by (c) $\Delta\Delta E$ distributions. All benchmark results for “OPLS4 cst”, and the detailed protocols to calculate the benchmark results are obtained from ref 45.

As shown in Fig. 5 (a) and (b), the distribution of ByteFF-gopt and ByteFF-joint in terms of RMSD and TFD are both concentrated near zero, with higher peak values near zero and narrower distributions than other force fields, demonstrating superior consistencies with QM-optimized molecular conformations. Particularly, both ByteFF-gopt and ByteFF-joint significantly outperform the “OPLS4 cst” model (OPLS4 with refined parameters using the FFBuilder tool for each molecule individually), which is considered the state-of-the-art MM force field in current reports. Additionally, we calculated the relative energy differences ($\Delta\Delta E$) to assess energetic agreement between force field-optimized conformations and the QM-optimized ones (Fig. 5 (c)). Both ByteFF-gopt and ByteFF-joint exhibit sharp peaks around zero in their $\Delta\Delta E$ distribution, significantly outperform all other force fields, indicating that both of them reproduce correctly the QM relative energies between conformers. The accurate predictions of equilibrium conformations ensures the correct conformations being sampled in MD simulations, while the precise prediction of $\Delta\Delta E$ guarantees the Boltzmann distribution in the conformational space is properly reproduced.




Figure 6: **Example of the equilibrium conformation predictions of molecules.** (a) 2D representation of the example molecule. (b-d) The equilibrium conformations predicted by ByteFF-gopt (green), Espaloma-0.3.0 (yellow), and GAFF-2.2 (cyan), superimposed to the equilibrium conformations predicted by QM (grey).

Fig. 6 illustrates a more detailed example of the performance of reproducing equilib-

rium geometries for various force fields. The QM-relaxed reference geometry shows that the four-membered ring takes a “butterfly” conformation due to significant torsional strain. The predictions relaxed using various force fields, including ByteFF-gopt (green), Espaloma-0.3.0 (yellow), and GAFF-2.2 (cyan), are superimposed on the QM reference. It is evident that Espaloma-0.3.0 (yellow) predicts the four-member ring as a planer conformation, while GAFF-2.2 (cyan) overestimates the bending of the ring. Being at the center of the molecule, a minor misprediction of the four-member ring conformation results in more significant geometric deviations on the two ends of the molecule, leading to overall RMSD values of 0.9 Å for Espaloma-0.3.0 and 0.4 Å for GAFF-2.2, respectively. Notably, the geometry relaxed by ByteFF-gopt (green) achieved superior alignment with the QM prediction with an RMSD of 0.2 Å. With increasing ring sizes and increasing number of neighboring rings, the complexity of the conformational space expands, making precise predictions of in-ring torsion profiles increasingly critical.

In addition to the equilibrium conformations, it is equally important for the MD simulations to accurately predict the off-equilibrium conformations. We used several datasets to comprehensively assess the prediction accuracy of both energy and force predictions from various force fields, quantified by RMSE relative to QM references (Table 2). Among all five subsets, ByteFF-gopt achieved higher accuracy than both GAFF-2.2 and OpenFF-2.0, but less accurate than Espaloma-0.3.0. Importantly, ByteFF-gopt achieved this accuracy with relaxed geometry, energy, and partial hessian, without using any force labels in the training process. With additional off-equilibrium energy and forces training data, we further trained ByteFF-joint to improve the force prediction capabilities of ByteFF-gopt. Remarkably, after fine-tuning with just around 10k data in the *off-equilibrium dataset*—a significantly smaller dataset than that used by Espaloma-0.3.0—the performance of ByteFF-joint surpassed that of its competitors. Rigorous procedures were followed in the benchmarking of ByteFF models, which ensured no data leakage between the training and testing data.

As a result, ByteFF-joint achieves state-of-the-art performance in almost all subsets,

except for the SPICE-Pubchem dataset, where ByteFF-joint shows marginally higher force RMSE compared to Espaloma-0.3.0. As discussed earlier, it is inherently challenging of an MMFF to perfectly replicate the QM PES. There is a trade-off between predicting the PES of relaxed-geometry and off-equilibrium conformations, and we focused more on the relaxed-geometry performances during the training process, since this is more relevant to the sampling of local minima in molecular dynamics simulations and is more stressed in the latest large-scale collaborative industrial benchmark.⁴⁵ Despite this, ByteFF-joint achieves overall state-of-the-art performance in all benchmarks considered in this study.

Table 2: Comparison of energy and force RMSE for different dataset and force fields. SPICE,⁴⁹ RNA, peptide.

Dataset		GAFF-2.2	OpenFF-2.0	Espaloma-0.3	ByteFF-gopt	ByteFF-joint
SPICE-Pubchem	<i>Energy</i>	4.8	4.4	2.3	3.4	2.2
	<i>Force</i>	14.1	14.0	6.5	9.9	6.9
SPICE-DES-Monomers	<i>Energy</i>	2.7	2.7	1.4	1.9	1.3
	<i>Force</i>	10.7	12.5	6.0	8.0	5.5
SPICE-Dipeptide	<i>Energy</i>	5.2	4.5	3.3	3.8	2.3
	<i>Force</i>	12.5	12.4	8.1	9.6	5.5
RNA-Diverse	<i>Energy</i>	6.7	5.4	3.9	6.3	3.5
	<i>Force</i>	15.8	18.4	4.5	11.1	3.8
RNA-Trinucleotide	<i>Energy</i>	6.2	6.0	3.8	6.0	3.4
	<i>Force</i>	16.5	18.7	4.4	11.3	3.7

Limitations

Despite the excellent performance of ByteFF compared to other force fields, several limitations still exist. As an MMFF with a minimal analytical functional, ByteFF inherently lacks the capability of modeling more complicated effects in intramolecular PES, such as bond-torsion and angle-torsion couplings. Moreover, while intramolecular parameters are carefully trained against QM references, non-bonded parameters are directly inherited from GAFF to ensure Amber compatibility, leaving room for further refinement. Addressing these limitations presents a path forward for the further improvements of ByteFF.

Conclusion

In this work, we developed ByteFF, an Amber-compatible MMFF with broad chemical space coverage and state-of-the-art accuracy. Through Morgan-based torsional fingerprints and t-SNE analysis, we demonstrated that our dataset covers superior chemical space. Based on this strong data foundation, ByteFF leverages an edge-augmented, symmetry-preserving GNN model to predict MMFF parameters. We also carefully designed our training strategy, enhanced by techniques such as partial Hessian fitting, an iterative optimization-and-training scheme, and effective model ensembling for prediction improvements and uncertainty quantification. Benefitting from the comprehensive datasets and exquisitely designed training strategy, ByteFF outperforms existing MM force fields across multiple benchmarks, delivering accurate predictions of torsional energy profiles, equilibrium conformations, and off-equilibrium energies and forces. We believe ByteFF will significantly benefit biomolecular simulations for various purposes.

Acknowledgement

References

- (1) Reymond, J.-L.; Awale, M. Exploring chemical space for drug discovery using the chemical universe database. *ACS chemical neuroscience* **2012**, *3*, 649–657.
- (2) Warr, W. A.; Nicklaus, M. C.; Nicolaou, C. A.; Rarey, M. Exploration of ultralarge compound collections for drug discovery. *Journal of Chemical Information and Modeling* **2022**, *62*, 2021–2034.
- (3) Caron, G.; Kihlberg, J.; Goetz, G.; Ratkova, E.; Poongavanam, V.; Ermondi, G. Steering new drug discovery campaigns: permeability, solubility, and physicochemical properties in the bRo5 chemical space. *ACS Medicinal Chemistry Letters* **2021**, *12*, 13–23.

- (4) Liu, X.; Shi, D.; Zhou, S.; Liu, H.; Liu, H.; Yao, X. Molecular dynamics simulations and novel drug discovery. *Expert opinion on drug discovery* **2018**, *13*, 23–37.
- (5) De Vivo, M.; Masetti, M.; Bottegoni, G.; Cavalli, A. Role of molecular dynamics and related methods in drug discovery. *Journal of medicinal chemistry* **2016**, *59*, 4035–4061.
- (6) Das, P.; Sercu, T.; Wadhawan, K.; Padhi, I.; Gehrman, S.; Cipcigan, F.; Chenthamakshan, V.; Strobel, H.; Dos Santos, C.; Chen, P.-Y., et al. Accelerated antimicrobial discovery via deep generative models and molecular dynamics simulations. *Nature Biomedical Engineering* **2021**, *5*, 613–623.
- (7) Muegge, I.; Hu, Y. Recent advances in alchemical binding free energy calculations for drug discovery. *ACS Medicinal Chemistry Letters* **2023**, *14*, 244–250.
- (8) Song, L. F.; Merz Jr, K. M. Evolution of alchemical free energy methods in drug discovery. *Journal of Chemical Information and Modeling* **2020**, *60*, 5308–5318.
- (9) Wang, J.; Wolf, R. M.; Caldwell, J. W.; Kollman, P. A.; Case, D. A. Development and Testing of a General Amber Force Field. *Journal of Computational Chemistry* **2004**, *25*, 1157–1174.
- (10) Harder, E. et al. OPLS3: A Force Field Providing Broad Coverage of Drug-like Small Molecules and Proteins. *Journal of Chemical Theory and Computation* **2016**, *12*, 281–296.
- (11) Roos, K.; Wu, C.; Damm, W.; Reboul, M.; Stevenson, J. M.; Lu, C.; Dahlgren, M. K.; Mondal, S.; Chen, W.; Wang, L.; Abel, R.; Friesner, R. A.; Harder, E. D. OPLS3e: Extending Force Field Coverage for Drug-Like Small Molecules. *Journal of Chemical Theory and Computation* **2019**, *15*, 1863–1874.
- (12) Lu, C.; Wu, C.; Ghoreishi, D.; Chen, W.; Wang, L.; Damm, W.; Ross, G. A.; Dahlgren, M. K.; Russell, E.; Von Bargen, C. D.; Abel, R.; Friesner, R. A.; Harder, E. D.

- OPLS4: Improving Force Field Accuracy on Challenging Regimes of Chemical Space. *Journal of Chemical Theory and Computation* **2021**, *17*, 4291–4300.
- (13) Buskes, M. J.; Blanco, M.-J. Impact of cross-coupling reactions in drug discovery and development. *Molecules* **2020**, *25*, 3493.
- (14) Guillemard, L.; Kaplaneris, N.; Ackermann, L.; Johansson, M. J. Late-stage C–H functionalization offers new opportunities in drug discovery. *Nature Reviews Chemistry* **2021**, *5*, 522–545.
- (15) Brandenburg, N.; Hoehnel, S.; Kuttler, F.; Homicsko, K.; Ceroni, C.; Ringel, T.; Gjorevski, N.; Schwank, G.; Coukos, G.; Turcatti, G., et al. High-throughput automated organoid culture via stem-cell aggregation in microcavity arrays. *Nature biomedical engineering* **2020**, *4*, 863–874.
- (16) Mohammad, T.; Mathur, Y.; Hassan, M. I. InstaDock: A single-click graphical user interface for molecular docking-based virtual high-throughput screening. *Briefings in Bioinformatics* **2021**, *22*, bbaa279.
- (17) Touret, F.; Gilles, M.; Barral, K.; Nougairède, A.; Van Helden, J.; Decroly, E.; De Lamballerie, X.; Coutard, B. In vitro screening of a FDA approved chemical library reveals potential inhibitors of SARS-CoV-2 replication. *Scientific reports* **2020**, *10*, 13093.
- (18) Qiu, Y. et al. Development and Benchmarking of Open Force Field v1.0.0—the Parsley Small-Molecule Force Field. *Journal of Chemical Theory and Computation* **2021**, *17*, 6262–6280.
- (19) Boothroyd, S. et al. Development and Benchmarking of Open Force Field 2.0.0: The Sage Small Molecule Force Field. *Journal of Chemical Theory and Computation* **2023**, *19*, 3251–3275.

- (20) Batatia, I.; Kovacs, D. P.; Simm, G.; Ortner, C.; Csanyi, G. MACE: Higher Order Equivariant Message Passing Neural Networks for Fast and Accurate Force Fields. *Advances in Neural Information Processing Systems* **2022**, *35*, 11423–11436.
- (21) Batzner, S.; Musaelian, A.; Sun, L.; Geiger, M.; Mailoa, J. P.; Kornbluth, M.; Molinari, N.; Smidt, T. E.; Kozinsky, B. E(3)-Equivariant Graph Neural Networks for Data-Efficient and Accurate Interatomic Potentials. *13*, 2453.
- (22) Rezaee, M.; Ekrami, S.; Hashemianzadeh, S. M. Comparing ANI-2x, ANI-1ccx Neural Networks, Force Field, and DFT Methods for Predicting Conformational Potential Energy of Organic Molecules. *Scientific Reports* **2024**, *14*, 11791.
- (23) Pelaez, R. P.; Simeon, G.; Galvelis, R.; Mirarchi, A.; Eastman, P.; Doerr, S.; Thölke, P.; Markland, T. E.; De Fabritiis, G. TorchMD-Net 2.0: Fast Neural Network Potentials for Molecular Simulations. *Journal of Chemical Theory and Computation* **2024**, *20*, 4076–4087.
- (24) Hornak, V.; Abel, R.; Okur, A.; Strockbine, B.; Roitberg, A.; Simmerling, C. Comparison of multiple Amber force fields and development of improved protein backbone parameters. *Proteins: Structure, Function, and Bioinformatics* **2006**, *65*, 712–725.
- (25) Maier, J. A.; Martinez, C.; Kasavajhala, K.; Wickstrom, L.; Hauser, K. E.; Simmerling, C. ff14SB: improving the accuracy of protein side chain and backbone parameters from ff99SB. *Journal of chemical theory and computation* **2015**, *11*, 3696–3713.
- (26) Rizzo, R. C.; Jorgensen, W. L. OPLS all-atom model for amines: resolution of the amine hydration problem. *Journal of the American chemical society* **1999**, *121*, 4827–4836.
- (27) Wang, L.; Behara, P. K.; Thompson, M. W.; Gokey, T.; Wang, Y.; Wagner, J. R.; Cole, D. J.; Gilson, M. K.; Shirts, M. R.; Mobley, D. L. The Open Force Field Initiative: Open Software and Open Science for Molecular Modeling. *The Journal of Physical Chemistry B* **2024**,

- (28) Wang, Y.; Fass, J.; Kaminow, B.; Herr, J. E.; Rufa, D.; Zhang, I.; Pulido, I.; Henry, M.; Macdonald, H. E. B.; Takaba, K.; Chodera, J. D. End-to-End Differentiable Construction of Molecular Mechanics Force Fields. *Chemical Science* **2022**, *13*, 12016–12033.
- (29) Takaba, K.; Friedman, A.; Cavender, C.; Behara, P.; Pulido, I.; Henry, M.; MacDermott-Opeskin, H.; Iacovella, C.; Nagle, A.; Payne, A.; Shirts, M.; Mobley, D. L.; Chodera, J. D.; Wang, Y. Machine-Learned Molecular Mechanics Force Fields from Large-Scale Quantum Chemical Data. *Chemical Science* **2024**,
- (30) Chen, G.; Jaffrelot Inizan, T.; Plé, T.; Lagardère, L.; Piquemal, J.-P.; Maday, Y. Advancing Force Fields Parameterization: A Directed Graph Attention Networks Approach. *Journal of Chemical Theory and Computation* **2024**,
- (31) March, J., et al. *Advanced organic chemistry: reactions, mechanisms, and structure*; McGraw-Hill New York, 1977.
- (32) Lahey, S.-L. J.; Thien Phuc, T. N.; Rowley, C. N. Benchmarking Force Field and the ANI Neural Network Potentials for the Torsional Potential Energy Surface of Biaryl Drug Fragments. *Journal of Chemical Information and Modeling* **2020**, *60*, 6258–6268.
- (33) Horton, J. T.; Boothroyd, S.; Wagner, J.; Mitchell, J. A.; Gokey, T.; Dotson, D. L.; Behara, P. K.; Ramaswamy, V. K.; Mackey, M.; Chodera, J. D.; Anwar, J.; Mobley, D. L.; Cole, D. J. Open Force Field BespokeFit: Automating Bespoke Torsion Parametrization at Scale. *Journal of Chemical Information and Modeling* **2022**, *62*, 5622–5633.
- (34) Zdrazil, B. et al. The ChEMBL Database in 2023: A Drug Discovery Platform Spanning Multiple Bioactivity Data Types and Time Periods. *Nucleic Acids Research* **2024**, *52*, D1180–D1192.
- (35) Irwin, J. J.; Tang, K. G.; Young, J.; Dandarchuluun, C.; Wong, B. R.; Khurelbaatar, M.; Moroz, Y. S.; Mayfield, J.; Sayle, R. A. ZINC20—a free ultralarge-scale chemical

- database for ligand discovery. *Journal of chemical information and modeling* **2020**, *60*, 6065–6073.
- (36) Jinsong, S.; Qifeng, J.; Xing, C.; Hao, Y.; Wang, L. Molecular fragmentation as a crucial step in the AI-based drug development pathway. *Communications Chemistry* **2024**, *7*, 20.
- (37) Johnston, R. C.; Yao, K.; Kaplan, Z.; Chelliah, M.; Leswing, K.; Seekins, S.; Watts, S.; Calkins, D.; Chief Elk, J.; Jerome, S. V., et al. Epik: p K a and Protonation State Prediction through Machine Learning. *Journal of chemical theory and computation* **2023**, *19*, 2380–2388.
- (38) Kesharwani, M. K.; Karton, A.; Martin, J. M. L. Benchmark ab Initio Conformational Energies for the Proteinogenic Amino Acids through Explicitly Correlated Methods. Assessment of Density Functional Methods. *Journal of Chemical Theory and Computation* **2015**, *12*, 444–454.
- (39) Behara, P. K.; Jang, H.; Horton, J. T.; Gokey, T.; Dotson, D. L.; Boothroyd, S.; Bayly, C. I.; Cole, D. J.; Wang, L.-P.; Mobley, D. L. Benchmarking Quantum Mechanical Levels of Theory for Valence Parametrization in Force Fields. *The Journal of Physical Chemistry B* **2024**,
- (40) Wang, L.-P.; Song, C. Geometry optimization made simple with translation and rotation coordinates. *The Journal of Chemical Physics* **2016**, *144*.
- (41) Epifanovsky, E. et al. Software for the frontiers of quantum chemistry: An overview of developments in the Q-Chem 5 package. *The Journal of Chemical Physics* **2021**, *155*.
- (42) Wu, X.; Sun, Q.; Pu, Z.; Zheng, T.; Ma, W.; Yan, W.; Yu, X.; Wu, Z.; Huo, M.; Li, X.; Ren, W.; Gong, S.; Zhang, Y.; Gao, W. Enhancing GPU-acceleration in the Python-based Simulations of Chemistry Framework. 2024; <https://arxiv.org/abs/2404.09452>.

- (43) Hussain, M. S.; Zaki, M. J.; Subramanian, D. Global Self-Attention as a Replacement for Graph Convolution. Proceedings of the 28th ACM SIGKDD Conference on Knowledge Discovery and Data Mining. New York, NY, USA, 2022; pp 655–665.
- (44) Dahlgren, M. K.; Schyman, P.; Tirado-Rives, J.; Jorgensen, W. L. Characterization of Biaryl Torsional Energetics and Its Treatment in OPLS All-Atom Force Fields. *Journal of Chemical Information and Modeling* **2013**, *53*, 1191–1199.
- (45) D’Amore, L. et al. Collaborative Assessment of Molecular Geometries and Energies from the Open Force Field. *Journal of Chemical Information and Modeling* **2022**, *62*, 6094–6104.
- (46) Schulz-Gasch, T.; Schärfer, C.; Guba, W.; Rarey, M. TFD: Torsion Fingerprints As a New Measure To Compare Small Molecule Conformations. *Journal of Chemical Information and Modeling* **2012**, *52*, 1499–1512.
- (47) Lim, VT.; Hahn, DF.; Tresadern, G.; Bayly, CI.; Mobley, DL. Benchmark Assessment of Molecular Geometries and Energies from Small Molecule Force Fields [Version 1; Peer Review: 2 Approved]. *F1000Research* **2020**, *9*.
- (48) Rai, B. K.; Sresht, V.; Yang, Q.; Unwalla, R.; Tu, M.; Mathiowetz, A. M.; Bakken, G. A. TorsionNet: A Deep Neural Network to Rapidly Predict Small-Molecule Torsional Energy Profiles with the Accuracy of Quantum Mechanics. *Journal of Chemical Information and Modeling* **2022**, *62*, 785–800.
- (49) Eastman, P.; Behara, P. K.; Dotson, D. L.; Galvelis, R.; Herr, J. E.; Horton, J. T.; Mao, Y.; Chodera, J. D.; Pritchard, B. P.; Wang, Y.; De Fabritiis, G.; Markland, T. E. SPICE, A Dataset of Drug-like Molecules and Peptides for Training Machine Learning Potentials. *Scientific Data* **2023**, *10*, 11.
- (50) Parlea, L. G.; Sweeney, B. A.; Hosseini-Asanjan, M.; Zirbel, C. L.; Leontis, N. B.

The RNA 3D Motif Atlas: Computational methods for extraction, organization and evaluation of RNA motifs. *Methods* **2016**, *103*, 99–119.

- (51) Marida, K. V. *In Statistics of Directional Data*; Academic Press, 1972; pp 18–24.
- (52) Shearer, J.; Castro, J. L.; Lawson, A. D. G.; MacCoss, M.; Taylor, R. D. Rings in Clinical Trials and Drugs: Present and Future. *Journal of Medicinal Chemistry* **2022**, *65*, 8699–8712.