Structured Backward Error Analysis with Preserving Sparsity for Three-by-Three Block Saddle Point Problems

Sk. Safique Ahmad^{†*}

Pinki Khatun[‡]

Abstract

Backward error (BE) analysis emerges as a powerful tool for assessing the backward stability and strong backward stability of iterative algorithms. In this paper, we explore structured BEs for a class of three-by-three block saddle point problems (TBSPPs), aiming to assess the strong backward stability of iterative algorithms devised to find their solution. Our investigations preserve the inherent matrix structure and sparsity pattern in the corresponding perturbation matrices and derive explicit formulae for the structure BEs. Moreover, we provide formulae for the structure-preserving minimal perturbation matrices for which the structured BE is attained. Furthermore, using the obtained structured BEs, we propose three new termination criteria for the iterative algorithms to solve the TBSPP. Numerical experiments are performed to test the strong backward stability of various iterative algorithms, as well as the reliability and effectiveness of the proposed termination criteria.

Keywords. Backward error analysis, Three-by-three block saddle point problems, Termination criterion, Iterative algorithms, Sparse matrices. AMS subject classification. 15A06, 65F10, 65F99

1 Introduction

The three-by-three block saddle point problem (TBSPP) is a class of $(n+m+p) \times (n+m+p)$ block linear system, which has attracted the attention of many researchers in recent decades for its versatile application in the community of scientific computations. For example, the *TBSPP* arises in solving PDE-constrained optimization problem [24], liquid crystal director model [23], Stokes-Darcy equations [27], finite element discretization of Maxwell equations

^{*}Corresponding author.

[†]Department of Mathematics, Indian Institute of Technology Indore, Khandwa Road, Indore, 452020, Madhya Pradesh, India, safique@iiti.ac.in, safique@gmail.com.

[‡]Research Scholar, Department of Mathematics, Indian Institute of Technology Indore, Khandwa Road, Indore, 452020, Madhya Pradesh, India, phd2001141004@iiti.ac.in, pinki996.pk@gmail.com.

[10] and incompressible magnetohydrodynamics models [14], and so on. We consider the general form of TBSPP in the following form:

$$\mathcal{A}\boldsymbol{w} := \begin{bmatrix} A & B_1^T & \mathbf{0} \\ B_2 & -C & D_1^T \\ \mathbf{0} & D_2 & E \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} f \\ g \\ h \end{bmatrix} =: \boldsymbol{d}, \qquad (1.1)$$

where $A \in \mathbb{R}^{n \times n}$, $C \in \mathbb{R}^{m \times m}$, $E \in \mathbb{R}^{p \times p}$ are symmetric matrices, $B_1, B_2 \in \mathbb{R}^{m \times n}$, $D_1, D_2 \in \mathbb{R}^{p \times m}$, $x, f \in \mathbb{R}^n$, $y, g \in \mathbb{R}^m$ and $z, h \in \mathbb{R}^p$.

Many researchers have developed efficient iterative algorithms in past years for solving the *TBSPP* of the form (1.1); for example [4, 29, 12, 7]. To assess the accuracy and reliability of an approximate solution obtained using an iterative algorithm, backward error (BE)analysis is widely employed in numerical linear algebra [13]. It aims to identify a closely perturbed problem (having minimum norm perturbation), ensuring that the approximate solution of the original problem coincides with the actual solution of the perturbed problem. The minimal distance between the original and perturbed problem is referred to as the BE. Furthermore, BE is used to assess the quality of an approximate solution as well as serves as a reliable and effective termination criterion when solving a problem using some iterative algorithm [8]. For an approximate solution of a given problem, if the computed BE is of unit round-off error, then the corresponding algorithm is classified as backward stable [13]. Moreover, the algorithm is classified as strongly backward stable [5, 6] if the perturbation matrix (with small magnitude) retains the structure of the coefficient matrix. This leads to a natural inquiry: Whether a backward stable algorithm for solving (1.1)exhibit strong backward stability or not? In this paper, the notion of structured *BE* facilitates us in addressing the aforementioned question, where we study the BE by preserving the inherent structure of the coefficient matrix \mathcal{A} .

Note that a *TBSPP* of the form (1.1) can be converted into a two-by-two block saddle point problem [3]. For instance, if we partition the coefficient matrix \mathcal{A} in (1.1) in the following form:

$$\begin{bmatrix} H & F_1^T \\ F_2 & G \end{bmatrix} \begin{bmatrix} p \\ q \end{bmatrix} = \begin{bmatrix} u \\ v \end{bmatrix},$$

where $H = \begin{bmatrix} A & B_1^T \\ B_2 & -C \end{bmatrix}$, G = E, $F_i = \begin{bmatrix} \mathbf{0} & D_i \end{bmatrix}$, $i = 1, 2, p = \begin{bmatrix} x \\ y \end{bmatrix}$, $q = z, u = \begin{bmatrix} f \\ g \end{bmatrix}$, and v = h. Considerable research effort has been devoted to structured *BEs* for the two-by-two block saddle point problem in the past years; see [28, 9, 20, 30, 22]. Nevertheless, due to the special block structure of \mathcal{A} , these investigations do not provide exact structured *BEs* for the *TBSPP* (1.1). Recently, Lv and Zheng [19] have investigated the structured *BEs* of (1.1), when $A = A^T$, $C = \mathbf{0}$ and $E = \mathbf{0}$. Further, Lv [18] studied the structured *BEs* of the equivalent form of the *TBSPP* (1.1) given by

$$\widehat{\mathcal{A}}\widehat{\boldsymbol{w}} := \begin{bmatrix} A & \mathbf{0} & B^T \\ \mathbf{0} & E & D \\ -B & -D^T & C \end{bmatrix} \begin{bmatrix} x \\ z \\ y \end{bmatrix} = \begin{bmatrix} f \\ h \\ -g \end{bmatrix}, \qquad (1.2)$$

with $B_1 = B_2 =: B$, $D_1 = D_2 =: D$, A and E are non-symmetric, and C is symmetric. When $C = \mathbf{0}$ and $E = \mathbf{0}$, computable expressions for the structured *BEs* are obtained in [21] in three the cases: first, $A^T = A$, $B_1 \neq B_2$ and $D_1 = D_2$; second, $A^T = A$, $B_1 = B_2$ and $D_1 \neq D_2$; and third, $A^T = A$, $B_1 \neq B_2$ and $D_1 \neq D_2$. However, these studies lack the following investigations: (a) the coefficient matrix \mathcal{A} in (1.1) is generally sparse, and the existing studies do not preserve the sparsity pattern to the perturbation matrices, (b) existing research does not provide explicit formulae for the minimal perturbation matrices that preserve the inherent structures of original matrices for which an approximated solution becomes the actual solution of a nearly perturbed *TBSPP*.

To address the aforementioned drawbacks, in this paper, we investigate the structured *BEs* in three cases: (i) $A^T = A$, $B_1 = B_2 =: B$, $C^T = C$, $D_1 = D_2 =: D$, and $E^T = E$; (ii) $A^T = A$, $B_1 \neq B_2$, $C^T = C$, $D_1 = D_2 =: D$, and $E^T = E$; (iii) $A^T \neq A$, $B_1 = B_2 =: B$, $C^T = C$, $D_1 \neq D_2$, and $E^T = E$.

The main contributions of this study are as follows:

- We investigate the structured *BEs* when the perturbation matrices preserve the structures mentioned in the cases (i), (ii) and (iii) as well as preserve the sparsity patterns of the block matrices of the coefficient matrix \mathcal{A} .
- We derive explicit formulae for the minimal perturbation matrices for which the structured BE is attained. These perturbation matrices preserve the inherent structures of the original matrices as well as their sparsity pattern.
- Utilizing the derived structured BE formulae, we propose three novel termination criteria for iterative algorithms to find the effective solution of TBSPPs
- Numerical experiments are performed to validate our theoretical findings, to test the backward stability and strong backward stability of iterative algorithms for solving the *TBSPPs* and to investigate the effectiveness of proposed termination criteria.

The organization of this paper is as follows. In Section 2, we present some notation, definitions, and preliminary results. In Sections 3, 4 and 5, we derive explicit formulae for the structured *BEs* corresponding to cases (i), (ii) and (iii), respectively. In Section 6, we propose three effective and reliable termination criteria for iterative algorithms to solve *TBSPP* for the cases (i)-(iii). Section 7 includes extensive numerical experiments. Lastly, Section 8 presents some concluding remarks.

2 Notation, preliminaries and definitions

2.1 Notation

In this paper, we consistently utilize the following notation. We denote $\mathbb{R}^{m \times n}$ as the collection of all $m \times n$ real matrices and \mathcal{S}_n as the set of all $n \times n$ symmetric matrices. The symbols $\|\cdot\|_2$ and $\|\cdot\|_F$ stand for the Euclidean and Frobenius norms, respectively.

For any matrix $X = [x_{ij}] \in \mathbb{R}^{m \times n}$, the symbols X^T and X^{\dagger} refer to the transpose and the Moore-Penrose inverse of X, respectively. The notation $\mathbf{1}_{m \times n} \in \mathbb{R}^{m \times n}$ represents the matrix with all entries are set to 1. We set $\mu = \frac{n(n+1)}{2}$, $\sigma = \frac{m(m+1)}{2}$ and $\tau = \frac{p(p+1)}{2}$. For $X = [\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n] \in \mathbb{R}^{m \times n}$, set $\operatorname{vec}(X) := [\mathbf{x}_1^T, \mathbf{x}_2^T, \dots, \mathbf{x}_n^T]^T \in \mathbb{R}^{mn}$, where \mathbf{x}_i is the *i*-th column of X. For any symmetric matrix $X \in S_n$, we define its generator vector by $\operatorname{vec}_{\mathcal{S}}(X) :=$ $[\mathbf{x}_1^T, \mathbf{x}_2^T, \dots, \mathbf{x}_n^T]^T \in \mathbb{R}^{\mu}$, where $\mathbf{x}_1 = [x_{11}, x_{21}, \dots, x_{n1}]^T \in \mathbb{R}^n$, $\mathbf{x}_2 = [x_{22}, x_{32}, \dots, x_{n2}]^T \in$ $\mathbb{R}^{n-1}, \dots, \mathbf{x}_{n-1} = [x_{(n-1)(n-1)}, x_{n(n-1)}]^T \in \mathbb{R}^2$, $\mathbf{x}_n = [x_{nn}] \in \mathbb{R}$. The Hadamard product of $X, Y \in \mathbb{R}^{m \times n}$ is defined as $X \odot Y = [x_{ij}y_{ij}] \in \mathbb{R}^{m \times n}$. For $X \in \mathbb{R}^{m \times n}$, we define $\Theta_X := \operatorname{sgn}(X) = [\operatorname{sgn}(x_{ij})] \in \mathbb{R}^{m \times n}$, where

$$\operatorname{sgn}(x_{ij}) = \begin{cases} 1, & \text{for } x_{ij} \neq 0, \\ 0, & \text{for } x_{ij} = 0. \end{cases}$$

For $x = [x_i] \in \mathbb{R}^n$, \mathfrak{D}_x denotes the diagonal matrix with the *i*-th diagonal entry x_i . We use the notation $\mathbf{0}_{m \times n}$ to represent the zero matrix of size $m \times n$. For simplicity, we use **0** when the matrix size is clear. Let θ_i , $i = 1, 2, \ldots, 10$ be the nonnegative real numbers with the convention that $\frac{1}{\theta_i} = 0$, whenever $\theta_i = 0$.

2.2 Preliminaries and definitions

In this subsection, we recall the definition of unstructured BE and introduce the definitions of structured BEs for the three cases (i)-(iii). Furthermore, we establish two pivotal lemmas essential for deriving structured BEs. Throughout the paper, we assume that \mathcal{A} is nonsingular. Next, we recall the definition of unstructured BE for an approximate solution of the linear system $\mathcal{A}w = d$.

Definition 2.1. [17] Let $\tilde{\boldsymbol{w}} = [\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ be an approximate solution of the TBSPP (1.1). Then, the unstructured BE, denoted by $\boldsymbol{\eta}(\tilde{\boldsymbol{w}})$ is defined as:

$$\eta(\tilde{\boldsymbol{w}}) := \min_{\Delta \mathcal{A}, \Delta \boldsymbol{d}} \left\{ \left\| \begin{bmatrix} \frac{\|\Delta \mathcal{A}\|_F}{\|\mathcal{A}\|_F}, & \frac{\|\Delta \boldsymbol{d}\|_F}{\|\boldsymbol{d}\|_F} \end{bmatrix} \right\|_2 \left| (\mathcal{A} + \Delta \mathcal{A}) \tilde{\boldsymbol{w}} = \boldsymbol{d} + \Delta \boldsymbol{d} \right\} \\ = \frac{\|\boldsymbol{d} - \mathcal{A} \tilde{\boldsymbol{w}}\|_2}{\sqrt{\|\mathcal{A}\|_F^2 \|\tilde{\boldsymbol{w}}\|_2^2 + \|\boldsymbol{d}\|_2^2}}.$$
(2.1)

A small value of $\eta(\tilde{w})$ indicates that the approximated solution \tilde{w} is the actual solution of a slightly perturbed system $(\mathcal{A} + \Delta \mathcal{A})\tilde{w} = d + \Delta d$, where $\|\Delta \mathcal{A}\|_F$ and $\|\Delta d\|_2$ are relatively small. That is, the corresponding iterative algorithm is backward stable.

Next, we defined structured BE for an approximate solution of the TBSPP (1.1).

Definition 2.2. Let $\widetilde{\boldsymbol{w}} = [\widetilde{x}^T, \widetilde{y}^T, \widetilde{z}^T]^T$ be an approximate solution of the TBSPP (1.1). Then, we define the structured BEs, denoted by $\boldsymbol{\eta}^{\mathbb{S}_i}(\widetilde{x}, \widetilde{y}, \widetilde{z}), i = 1, 2, 3$, for the cases (i)-(iii) as follows:

$$\boldsymbol{\eta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) = \min_{\begin{pmatrix}\Delta A, \Delta B, \Delta C, \\ \Delta D, \Delta E, \Delta f, \\ \Delta g, \Delta h \end{pmatrix}} \left\| \begin{bmatrix} \theta_{1} \|\Delta A\|_{F} & \theta_{2} \|\Delta B\|_{F} & \theta_{4} \|\Delta C\|_{F} \\ \theta_{5} \|\Delta D\|_{F} & \theta_{7} \|\Delta E\|_{F} & 0 \\ \theta_{8} \|\Delta f\|_{2} & \theta_{9} \|\Delta g\|_{2} & \theta_{10} \|\Delta h\|_{2} \end{bmatrix} \right\|_{F},$$

$$\boldsymbol{\eta}^{\mathbb{S}_{2}}(\tilde{x}, \tilde{y}, \tilde{z}) = \min_{\begin{pmatrix}\Delta A, \Delta B_{1}, \Delta B_{2}, \\ \Delta C, \Delta D, \Delta E, \\ \Delta f, \Delta g, \Delta h \end{pmatrix}} \in \mathbb{S}_{2} \left\| \begin{bmatrix} \theta_{1} \|\Delta A\|_{F} & \theta_{2} \|\Delta B_{1}\|_{F} & \theta_{3} \|\Delta B_{2}\|_{F} \\ \theta_{4} \|\Delta C\|_{F} & \theta_{5} \|\Delta D\|_{F} & \theta_{7} \|\Delta E\|_{F} \\ \theta_{4} \|\Delta C\|_{F} & \theta_{5} \|\Delta D\|_{F} & \theta_{7} \|\Delta E\|_{F} \\ \theta_{8} \|\Delta f\|_{2} & \theta_{9} \|\Delta g\|_{2} & \theta_{10} \|\Delta h\|_{2} \end{bmatrix} \right\|_{F},$$

$$\boldsymbol{\eta}^{\mathbb{S}_{3}}(\tilde{x}, \tilde{y}, \tilde{z}) = \min_{\begin{pmatrix}\Delta A, \Delta B, \Delta C, \\ \Delta D_{1}, \Delta D_{2}, \Delta E, \\ \Delta f, \Delta g, \Delta h \end{pmatrix}} \in \mathbb{S}_{3} \left\| \begin{bmatrix} \theta_{1} \|\Delta A\|_{F} & \theta_{2} \|\Delta B\|_{F} & \theta_{4} \|\Delta C\|_{F} \\ \theta_{5} \|\Delta D_{1}\|_{F} & \theta_{6} \|\Delta D_{2}\|_{F} & \theta_{7} \|\Delta E\|_{F} \\ \theta_{8} \|\Delta f\|_{2} & \theta_{9} \|\Delta g\|_{2} & \theta_{10} \|\Delta h\|_{2} \end{bmatrix} \right\|_{F},$$

respectively, where

$$\mathbb{S}_{1} = \left\{ \left(\begin{array}{c} \Delta A, \Delta B, \Delta C\\ \Delta D, \Delta E, \Delta f\\ \Delta g, \Delta h \end{array} \right) \middle| \left[\begin{array}{c} A + \Delta A & (B + \Delta B)^{T} & \mathbf{0}\\ B + \Delta B & -(C + \Delta C) & (D + \Delta D)^{T}\\ \mathbf{0} & D + \Delta D & (E + \Delta E)^{T} \end{array} \right] \left[\begin{array}{c} \tilde{x}\\ \tilde{y}\\ \tilde{z} \end{array} \right] = \left[\begin{array}{c} f + \Delta f\\ g + \Delta g\\ h + \Delta h \end{array} \right],$$
$$\Delta A \in \mathcal{S}_{n}, \Delta C \in \mathcal{S}_{m}, \Delta E \in \mathcal{S}_{p}, \Delta B \in \mathbb{R}^{m \times n}, \Delta D \in \mathbb{R}^{p \times m},$$
$$\Delta f \in \mathbb{R}^{n}, \Delta g \in \mathbb{R}^{m}, \Delta h \in \mathbb{R}^{p} \right\}.$$
(2.2)

$$\mathbb{S}_{2} = \left\{ \left(\begin{array}{c} \Delta A, \Delta B_{1}, \Delta B_{2} \\ \Delta C, \Delta D, \Delta E, \\ \Delta f, \Delta g, \Delta h \end{array} \right) \middle| \left[\begin{array}{c} A + \Delta A & (B_{1} + \Delta B_{1})^{T} & \mathbf{0} \\ B_{2} + \Delta B_{2} & -(C + \Delta C) & (D + \Delta D)^{T} \\ \mathbf{0} & D + \Delta D & (E + \Delta E)^{T} \end{array} \right] \left[\begin{array}{c} \tilde{x} \\ \tilde{y} \\ \tilde{z} \end{array} \right] = \left[\begin{array}{c} f + \Delta f \\ g + \Delta g \\ h + \Delta h \end{array} \right], \\ \Delta A \in \mathcal{S}_{n}, \Delta C \in \mathcal{S}_{m}, \Delta E \in \mathcal{S}_{p}, \Delta B_{1}, \Delta B_{2} \in \mathbb{R}^{m \times n}, \Delta D \in \mathbb{R}^{p \times m}, \end{cases}$$

$$\Delta f \in \mathbb{R}^n, \Delta g \in \mathbb{R}^m, \Delta h \in \mathbb{R}^p \bigg\}.$$
(2.3)

$$\mathbb{S}_{3} = \left\{ \left(\begin{array}{c} \Delta A, \Delta B, \Delta C\\ \Delta D_{1}, \Delta D_{2}, \Delta E, \\ \Delta f, \Delta g, \Delta h \end{array} \right) \middle| \left[\begin{array}{c} A + \Delta A & (B + \Delta B)^{T} & \mathbf{0}\\ B + \Delta B & -(C + \Delta C) & (D_{1} + \Delta D_{1})^{T}\\ \mathbf{0} & D_{2} + \Delta D_{2} & (E + \Delta E)^{T} \end{array} \right] \left[\begin{array}{c} \tilde{x}\\ \tilde{y}\\ \tilde{z} \end{array} \right] = \left[\begin{array}{c} f + \Delta f\\ g + \Delta g\\ h + \Delta h \end{array} \right],$$
$$\Delta A \in \mathbb{R}^{n \times n}, \Delta C \in \mathcal{S}_{m}, \Delta E \in \mathcal{S}_{p}, \Delta B \in \mathbb{R}^{m \times n}, \Delta D_{1}, \Delta D_{2} \in \mathbb{R}^{p \times m},$$
$$\Delta f \in \mathbb{R}^{n}, \Delta g \in \mathbb{R}^{m}, \Delta h \in \mathbb{R}^{p} \right\}.$$
(2.4)

Next, we state the problem of finding structure-preserving minimal perturbation matrices for which the structured BE is attained.

Problem 2.1. Find out the minimal perturbation matrices

$$\left(\begin{array}{c} \widehat{B}, \widehat{\Delta}\widehat{C}, \\ \widehat{E}, \widehat{\Delta}\widehat{f}, \\ \widehat{\Delta}\widehat{h} \end{array} \right) \in \mathbb{S}_1,$$

$$\begin{pmatrix} \widehat{\Delta A}, \widehat{\Delta B}_{1}, \widehat{\Delta B}_{2}, \\ \widehat{\Delta C}, \widehat{\Delta D}, \widehat{\Delta E}, \\ \widehat{\Delta f}, \widehat{\Delta g}, \widehat{\Delta h} \end{pmatrix} \in \mathbb{S}_{2} \text{ and } \begin{pmatrix} \widehat{\Delta A}, \widehat{\Delta B}, \widehat{\Delta C}, \\ \widehat{\Delta D}_{1}, \widehat{\Delta D}_{2}, \widehat{\Delta E}, \\ \widehat{\Delta f}, \widehat{\Delta g}, \widehat{\Delta h} \end{pmatrix} \in \mathbb{S}_{3} \text{ such that} \\ \boldsymbol{\eta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \begin{bmatrix} \theta_{1} \| \widehat{\Delta A} \|_{F} & \theta_{2} \| \widehat{\Delta B} \|_{F} & \theta_{4} \| \widehat{\Delta C} \|_{F} \\ \theta_{5} \| \widehat{\Delta D} \|_{F} & \theta_{7} \| \widehat{\Delta E} \|_{F} & 0 \\ \theta_{8} \| \widehat{\Delta f} \|_{2} & \theta_{9} \| \widehat{\Delta g} \|_{2} & \theta_{10} \| \widehat{\Delta h} \|_{F} \end{bmatrix} \right\|_{F}, \\ \boldsymbol{\eta}^{\mathbb{S}_{2}}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \begin{bmatrix} \theta_{1} \| \widehat{\Delta A} \|_{F} & \theta_{2} \| \widehat{\Delta B} \|_{F} & \theta_{3} \| \widehat{\Delta B}_{2} \|_{F} \\ \theta_{4} \| \widehat{\Delta C} \|_{F} & \theta_{5} \| \widehat{\Delta D} \|_{F} & \theta_{7} \| \widehat{\Delta E} \|_{F} \\ \theta_{8} \| \widehat{\Delta f} \|_{2} & \theta_{9} \| \widehat{\Delta g} \|_{2} & \theta_{10} \| \widehat{\Delta h} \|_{F} \end{bmatrix} \right\|_{F}, \\ \boldsymbol{\eta}^{\mathbb{S}_{3}}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \begin{bmatrix} \theta_{1} \| \widehat{\Delta A} \|_{F} & \theta_{2} \| \widehat{\Delta B} \|_{F} & \theta_{4} \| \widehat{\Delta C} \|_{F} \\ \theta_{8} \| \widehat{\Delta f} \|_{2} & \theta_{9} \| \widehat{\Delta g} \|_{2} & \theta_{10} \| \widehat{\Delta h} \|_{F} \end{bmatrix} \right\|_{F}, \\ \mathbf{\eta}^{\mathbb{S}_{3}}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \begin{bmatrix} \theta_{1} \| \widehat{\Delta A} \|_{F} & \theta_{2} \| \widehat{\Delta B} \|_{F} & \theta_{4} \| \widehat{\Delta C} \|_{F} \\ \theta_{5} \| \widehat{\Delta D}_{1} \|_{F} & \theta_{6} \| \widehat{\Delta D}_{2} \|_{F} & \theta_{7} \| \widehat{\Delta E} \|_{F} \\ \theta_{8} \| \widehat{\Delta f} \|_{2} & \theta_{9} \| \widehat{\Delta g} \|_{2} & \theta_{10} \| \widehat{\Delta h} \|_{F} \end{bmatrix} \right\|_{F}, \\ \end{pmatrix}$$

respectively.

Remark 2.1. When $\theta_i = 0$ for any given i (i = 1, 2, ..., 10), it indicates that the corresponding block matrix has no perturbation.

Remark 2.2. Many researchers have studied the structured BE by preserving the sparsity pattern in the literature; see [1, 2]. Our specific interest lies in investigating structured BEs while the perturbation matrices preserve the sparsity pattern of the original matrices. To perform this investigation, we substitute the perturbation matrices ΔA , ΔB_1 , ΔB_2 ΔC , $\Delta D_1 \ \Delta D_2$ and ΔE by $\Delta A \odot \Theta_A$, $\Delta B_1 \odot \Theta_{B_1}$, $\Delta B_2 \odot \Theta_{B_2}$, $\Delta C \odot \Theta_C$, $\Delta D_1 \odot \Theta_{D_1}$, $\Delta D_2 \odot \Theta_{D_2}$ and $\Delta E \odot \Theta_E$, respectively. Within this framework, we denote the structured BEs as $\eta_{sps}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z})$, i = 1, 2, 3. Moreover, the minimal perturbation matrices are denoted by $\widehat{\Delta A}_{sps}$, $\widehat{\Delta B}_{1sps}$, $\widehat{\Delta B}_{2sps}$, $\widehat{\Delta C}_{sps}$, $\widehat{\Delta D}_{1sps}$, $\widehat{\Delta D}_{2sps}$, $\widehat{\Delta E}_{sps}$, $\widehat{\Delta f}_{sps}$, $\widehat{\Delta g}_{sps}$ and $\widehat{\Delta h}_{sps}$.

When the structured *BEs* $\boldsymbol{\eta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z})$ and $\boldsymbol{\eta}^{\mathbb{S}_i}_{sps}(\tilde{x}, \tilde{y}, \tilde{z})$ are around an order of unit roundoff error, then the approximate solution $\tilde{\boldsymbol{w}} = [\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ becomes an actual solution of nearly perturbed structure-preserving *TBSPP* of the form (1.1). Thus, the corresponding algorithm is referred to as strongly backward stable. To obtain the structured *BEs* formulae, the following lemmas play a pivotal role.

Lemma 2.1. Let $A, H \in S_n$ with generator vectors $\operatorname{vec}_{S}(A) = [\boldsymbol{a}_1^T, \boldsymbol{a}_2^T, \dots, \boldsymbol{a}_n^T]^T$ and $\operatorname{vec}_{S}(H) = [\boldsymbol{h}_1^T, \boldsymbol{h}_2^T, \dots, \boldsymbol{h}_n^T]^T$, respectively. Suppose $x = [x_1, \dots, x_n]^T \in \mathbb{R}^n$, $y = [y_1, \dots, y_n]^T \in \mathbb{R}^n$ and $\boldsymbol{d} = [d_1, \dots, d_n]^T \in \mathbb{R}^n$. Then $(A \odot \Theta_H)x = \boldsymbol{d}$ can be expressed as:

$$\mathcal{K}_x \Phi_H \operatorname{vec}_{\mathcal{S}}(A \odot \Theta_H) = \boldsymbol{d},$$

where $\Phi_H = \text{diag}(\text{vec}_{\mathcal{S}}(\Theta_H)), \ \mathcal{K}_x = \begin{bmatrix} K_x^1 & K_x^2 & \cdots & K_x^n \end{bmatrix} \in \mathbb{R}^{n \times \mu} \ and \ K_x^i \in \mathbb{R}^{n \times (n-i+1)} \ are$ given by

$$K_x^1 = \begin{bmatrix} x_1 & x_2 & \cdots & \cdots & x_n \\ 0 & x_1 & 0 & \cdots & 0 \\ 0 & 0 & x_1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & x_1 \end{bmatrix}, \quad K_x^2 = \begin{bmatrix} 0 & 0 & \cdots & \cdots & 0 \\ x_2 & x_3 & \cdots & \cdots & x_n \\ 0 & x_2 & 0 & \cdots & 0 \\ 0 & 0 & x_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & \cdots & 0 & x_2 \end{bmatrix}, \quad \dots, \quad K_x^n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ x_n \end{bmatrix}.$$

Proof. Since $H \in S_n$, we have $\Theta_H \in S_n$. Let $\operatorname{vec}_{S}(A) = [\boldsymbol{a}_1, \boldsymbol{a}_2, \dots, \boldsymbol{a}_n]^T$ and $\operatorname{vec}_{S}(\Theta_H) = [\Theta_{\boldsymbol{h}_1}^T, \Theta_{\boldsymbol{h}_2}^T, \dots, \Theta_{\boldsymbol{h}_n}^T]^T$, where $\Theta_{\boldsymbol{h}_i} = [\operatorname{sgn}(h_{ii}), \operatorname{sgn}(h_{i(i+1)}), \dots, \operatorname{sgn}(h_{in})]^T \in \mathbb{R}^{n-i+1}$. Then, $(A \odot \Theta_H)$ can be equivalently expressed as

$$(A \odot \Theta_{H})x = \begin{bmatrix} a_{11} \operatorname{sgn}(h_{11})x_{1} + a_{21} \operatorname{sgn}(h_{21})x_{2} + \dots + a_{n1} \operatorname{sgn}(h_{n1})x_{n} \\ a_{21} \operatorname{sgn}(h_{21})x_{1} + a_{22} \operatorname{sgn}(h_{22})x_{2} + \dots + a_{n2} \operatorname{sgn}(h_{n2})x_{n} \\ a_{31} \operatorname{sgn}(h_{31})x_{1} + a_{32} \operatorname{sgn}(h_{32})x_{2} + \dots + a_{n3} \operatorname{sgn}(h_{n3})x_{n} \\ \vdots \\ a_{n1} \operatorname{sgn}(h_{n1})x_{1} + a_{21} \operatorname{sgn}(h_{21})x_{2} + \dots + a_{1n} \operatorname{sgn}(h_{n1})x_{n} \\ a_{21} \operatorname{sgn}(h_{21})x_{1} \\ \vdots \\ a_{n1} \operatorname{sgn}(h_{11})x_{1} + a_{21} \operatorname{sgn}(h_{21})x_{2} + \dots + a_{1n} \operatorname{sgn}(h_{n1})x_{n} \\ \vdots \\ a_{n1} \operatorname{sgn}(h_{n1})x_{1} \\ \vdots \\ a_{n1} \operatorname{sgn}(h_{n1})x_{1} \\ \vdots \\ a_{n1} \operatorname{sgn}(h_{n1})x_{1} \\ \vdots \\ a_{n2} \operatorname{sgn}(h_{n2})x_{2} \\ \vdots \\ a_{n2} \operatorname{sgn}(h_{n2})x_{2} \\ \vdots \\ a_{n2} \operatorname{sgn}(h_{n2})x_{2} \\ \vdots \\ a_{n3} \operatorname{sgn}(h_{n3})x_{3} + \dots + a_{n3} \operatorname{sgn}(h_{n3})x_{n} \\ \vdots \\ a_{n3} \operatorname{sgn}(h_{n3})x_{3} + \dots + a_{n3} \operatorname{sgn}(h_{n3})x_{n} \\ \vdots \\ a_{n3} \operatorname{sgn}(h_{n3})x_{3} \\ = K_{x}^{1} \mathfrak{D}_{\Theta_{h_{1}}} (a_{1} \odot \Theta_{h_{1}}) + K_{x}^{2} \mathfrak{D}_{\Theta_{h_{2}}} (a_{2} \odot \Theta_{h_{2}}) + \dots + K_{x}^{n} \mathfrak{D}_{\Theta_{h_{n}}} (a_{n} \odot \Theta_{h_{n}}) \\ \vdots \\ a_{n} \odot \Theta_{h_{n}} \end{bmatrix}$$

Therefore, the proof is concluded. \blacksquare

Remark 2.3. When the matrix $\Theta_H = \mathbf{1}_{n \times n}$, from Lemma 2.1, the linear system $Ax = \mathbf{b}$, where $A \in S_n$ can be written as $\mathcal{K}_x \operatorname{vec}_{\mathcal{S}}(A) = \mathbf{b}$.

Lemma 2.2. Let $A, B, H \in \mathbb{R}^{m \times n}$ be three given matrices. Suppose that $x = [x_1, \ldots, x_n]^T \in \mathbb{R}^n$, $y = [y_1, \ldots, y_m]^T \in \mathbb{R}^m$, $d^1 \in \mathbb{R}^m$ and $d^2 \in \mathbb{R}^n$. Then $(A \odot \Theta_H)x = d^1$ and $(B \odot \Theta_H)^T y = d^2$ are equivalent to

$$\mathcal{M}_x^m \mathfrak{D}_{\operatorname{vec}(\Theta_H)}\operatorname{vec}(A \odot \Theta_H) = d^1 \quad and \quad \mathcal{N}_y^n \mathfrak{D}_{\operatorname{vec}(\Theta_H)}\operatorname{vec}(B \odot \Theta_H) = d^2,$$

respectively, where $\mathcal{M}_x^m = \begin{bmatrix} x_1 I_m & x_2 I_m & \cdots & x_n I_m \end{bmatrix} \in \mathbb{R}^{m \times mn}$ and

$$\mathcal{N}_{y}^{n} = \begin{bmatrix} y^{T} & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & y^{T} & \mathbf{0} & \cdots & \mathbf{0} \\ \vdots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ \mathbf{0} & \cdots & \cdots & \mathbf{0} & y^{T} \end{bmatrix} \in \mathbb{R}^{n \times mn}.$$
(2.6)

Proof. The proof proceeds using a method analogous to the proof of Lemma 2.1.

Lemma 2.3. [11] Let $A \in \mathbb{C}^{m \times n}$ and $b \in \mathbb{C}^m$. The system of linear equations Ax = b is consistent if and only if $AA^{\dagger}b = b$. Additionally, the minimum norm least-squares solution of this system is represented by $A^{\dagger}b$.

3 Derivation of structured *BEs* for case (i)

In this section, we discuss the structured *BEs* for the *TBSPP* (1.1) for the case (*i*), i.e., $A \in S_n, C \in S_m, E \in S_p, B_1 = B_2 =: B$ and $D_1 = D_2 =: D$, and perturbation matrices belongs to set S_1 . Prior to that, we construct the diagonal matrix $\mathfrak{D}_{S_n} \in \mathbb{R}^{n \times n}$, where

$$\begin{cases} \mathfrak{D}_{S_n}(k,k) = 1, & \text{for } k = \frac{(2n - (i-2))(i-1)}{2} + 1, i = 1, 2, \dots, n, \\ \mathfrak{D}_{S_n}(k,k) = \sqrt{2}, & \text{otherwise.} \end{cases}$$

The matrix $\mathfrak{D}_{\mathcal{S}_n}$ has the property, $||A||_F = ||\mathfrak{D}_{\mathcal{S}_n} \operatorname{vec}_{\mathcal{S}}(A)||_2$. Further, we introduce the following notation:

$$\Phi_A = \operatorname{diag}(\operatorname{vec}_{\mathcal{S}}(\Theta_A)), \ \Phi_B = \operatorname{diag}(\operatorname{vec}(\Theta_B)), \ \Phi_C = \operatorname{diag}(\operatorname{vec}_{\mathcal{S}}(\Theta_C)), \tag{3.1}$$

$$\Phi_D = \operatorname{diag}(\operatorname{vec}(\Theta_D)) \text{ and } \Phi_E = \operatorname{diag}(\operatorname{vec}_{\mathcal{S}}(\Theta_E)).$$
(3.2)

Theorem 3.1. Let $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ be an approximate solution of the TBSPP (1.1) with $A \in S_n, C \in S_m, E \in S_p$, and $\theta_8, \theta_9, \theta_{10} \neq 0$. Then, we have

$$\boldsymbol{\eta}_{\mathbf{sps}}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \mathcal{J}_{\mathbb{S}_1}^T (\mathcal{J}_{\mathbb{S}_1} \mathcal{J}_{\mathbb{S}_1}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \right\|_2,$$
(3.3)

where $\mathcal{J}_{\mathbb{S}_1} \in \mathbb{R}^{(n+m+p) \times l}$ is given by

$$\mathcal{J}_{\mathbb{S}_{1}} = \begin{bmatrix} \frac{1}{\theta_{1}} \mathcal{K}_{\tilde{x}} \Phi_{A} \mathfrak{D}_{\mathcal{S}_{n}}^{-1} & \frac{1}{\theta_{2}} \mathcal{N}_{\tilde{y}}^{n} \Phi_{B} & \mathbf{0} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{8}} I_{n} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \frac{1}{\theta_{2}} \mathcal{M}_{\tilde{x}}^{m} \Phi_{B} & -\frac{1}{\theta_{4}} \mathcal{K}_{\tilde{y}} \Phi_{C} \mathfrak{D}_{\mathcal{S}_{m}}^{-1} & \frac{1}{\theta_{5}} \mathcal{N}_{\tilde{x}}^{m} \Phi_{D} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{9}} I_{m} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \frac{1}{\theta_{5}} \mathcal{M}_{\tilde{y}}^{p} \Phi_{D} & \frac{1}{\theta_{7}} \mathcal{K}_{\tilde{x}} \Phi_{E} \mathfrak{D}_{\mathcal{S}_{p}}^{-1} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{10}} I_{p} \end{bmatrix}$$

 $R_f = f - A\tilde{x} - B^T\tilde{y}, R_g = g - B\tilde{x} + C\tilde{y} - D^T\tilde{z}, R_h = h - D\tilde{y} - E\tilde{z}, and \mathbf{l} = \mu + \sigma + \tau + \sigma$ mn + mp + m + n + p.

The minimal perturbation matrices for the Problem 2.1 are given by the following generating vectors:

$$\begin{aligned} \operatorname{vec}_{\mathcal{S}}(\widehat{\Delta A}_{\mathtt{sps}}) &= \theta_{1}^{-1}\mathfrak{D}_{\mathcal{S}_{n}}^{-1} \begin{bmatrix} I_{\mu} & \mathbf{0} \end{bmatrix} \mathcal{J}_{\mathbb{S}_{1}}^{T} (\mathcal{J}_{\mathbb{S}_{1}}\mathcal{J}_{\mathbb{S}_{1}}^{T})^{-1} \begin{bmatrix} R_{f}^{T}, \ R_{g}^{T}, \ R_{h}^{T} \end{bmatrix}^{T}, \\ \operatorname{vec}(\widehat{\Delta B}_{\mathtt{sps}}) &= \theta_{2}^{-1} \begin{bmatrix} \mathbf{0} & I_{mn} & \mathbf{0} \end{bmatrix} \mathcal{J}_{\mathbb{S}_{1}}^{T} (\mathcal{J}_{\mathbb{S}_{1}}\mathcal{J}_{\mathbb{S}_{1}}^{T})^{-1} \begin{bmatrix} R_{f}^{T}, \ R_{g}^{T}, \ R_{h}^{T} \end{bmatrix}^{T}, \\ \operatorname{vec}_{\mathcal{S}}(\widehat{\Delta C}_{\mathtt{sps}}) &= \theta_{4}^{-1}\mathfrak{D}_{\mathcal{S}_{m}}^{-1} \begin{bmatrix} \mathbf{0} & I_{\sigma} & \mathbf{0} \end{bmatrix} \mathcal{J}_{\mathbb{S}_{1}}^{T} (\mathcal{J}_{\mathbb{S}_{1}}\mathcal{J}_{\mathbb{S}_{1}}^{T})^{-1} \begin{bmatrix} R_{f}^{T}, \ R_{g}^{T}, \ R_{h}^{T} \end{bmatrix}^{T}, \\ \operatorname{vec}(\widehat{\Delta D}_{\mathtt{sps}}) &= \theta_{5}^{-1} \begin{bmatrix} \mathbf{0} & I_{mp} & \mathbf{0} \end{bmatrix} \mathcal{J}_{\mathbb{S}_{1}}^{T} (\mathcal{J}_{\mathbb{S}_{1}}\mathcal{J}_{\mathbb{S}_{1}}^{T})^{-1} \begin{bmatrix} R_{f}^{T}, \ R_{g}^{T}, \ R_{h}^{T} \end{bmatrix}^{T}, \\ \operatorname{vec}_{\mathcal{S}}(\widehat{\Delta E}_{\mathtt{sps}}) &= \theta_{7}^{-1} \mathfrak{D}_{\mathcal{S}_{p}}^{-1} \begin{bmatrix} \mathbf{0} & I_{\tau} & \mathbf{0} \end{bmatrix} \mathcal{J}_{\mathbb{S}_{1}}^{T} (\mathcal{J}_{\mathbb{S}_{1}}\mathcal{J}_{\mathbb{S}_{1}}^{T})^{-1} \begin{bmatrix} R_{f}^{T}, \ R_{g}^{T}, \ R_{h}^{T} \end{bmatrix}^{T}, \\ \operatorname{vec}_{\mathcal{S}}(\widehat{\Delta E}_{\mathtt{sps}}) &= \theta_{7}^{-1} \mathfrak{D}_{\mathcal{S}_{p}}^{-1} \begin{bmatrix} \mathbf{0} & I_{\tau} & \mathbf{0} \end{bmatrix} \mathcal{J}_{\mathbb{S}_{1}}^{T} (\mathcal{J}_{\mathbb{S}_{1}}\mathcal{J}_{\mathbb{S}_{1}}^{T})^{-1} \begin{bmatrix} R_{f}^{T}, \ R_{g}^{T}, \ R_{h}^{T} \end{bmatrix}^{T}, \\ \operatorname{vec}_{\mathcal{S}}(\widehat{\Delta E}_{\mathtt{sps}}) &= \theta_{7}^{-1} \mathfrak{D}_{\mathcal{S}_{p}}^{-1} \begin{bmatrix} \mathbf{0} & I_{\pi} & \mathbf{0} \end{bmatrix} \mathcal{J}_{\mathbb{S}_{1}}^{T} (\mathcal{J}_{\mathbb{S}_{1}}\mathcal{J}_{\mathbb{S}_{1}}^{T})^{-1} \begin{bmatrix} R_{f}^{T}, \ R_{g}^{T}, \ R_{h}^{T} \end{bmatrix}^{T}, \\ \operatorname{a}\widehat{\Delta f}_{\mathtt{sps}} &= \theta_{8}^{-1} \begin{bmatrix} \mathbf{0} & I_{m} & \mathbf{0} \end{bmatrix} \mathcal{J}_{\mathbb{S}_{1}}^{T} (\mathcal{J}_{\mathbb{S}_{1}}\mathcal{J}_{\mathbb{S}_{1}}^{T})^{-1} \begin{bmatrix} R_{f}^{T}, \ R_{g}^{T}, \ R_{h}^{T} \end{bmatrix}^{T}, \\ \operatorname{a}\widehat{\Delta h}_{\mathtt{sps}} &= \theta_{10}^{-1} \begin{bmatrix} \mathbf{0} & I_{p} \end{bmatrix} \mathcal{J}_{\mathbb{S}_{1}}^{T} (\mathcal{J}_{\mathbb{S}_{1}}\mathcal{J}_{\mathbb{S}_{1}}^{T})^{-1} \begin{bmatrix} R_{f}^{T}, \ R_{g}^{T}, \ R_{h}^{T} \end{bmatrix}^{T}. \end{aligned} \right]^{T}. \end{aligned}$$

Proof. For the approximate solution $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$, we need to construct perturbations $\Delta A \in \mathcal{S}_n, \ \Delta B \in \mathbb{R}^{m \times n}, \ \Delta C \in \mathcal{S}_m, \ \Delta D \in \mathbb{R}^{p \times m}, \ \Delta E \in \mathcal{S}_p$, which maintain the sparsity pattern of A, B, C, D, E, respectively, and the perturbations $\Delta f \in \mathbb{R}^n, \Delta g \in \mathbb{R}^m$, and sity pattern of A, D, C, D, L, D, L, D, L $\Delta h \in \mathbb{R}^p$. By (2.2), $\begin{pmatrix} \Delta A, \Delta B, \Delta C, \\ \Delta D, \Delta E, \Delta f, \\ \Delta g, \Delta h \end{pmatrix} \in \mathbb{S}_1$ if and only if $\Delta A, \Delta B, \Delta C, \Delta D, \Delta E, \Delta f, \Delta g$

$$\left. \begin{array}{l} \Delta A\tilde{x} + \Delta B^{T}\tilde{y} - \Delta f = R_{f}, \\ \Delta B\tilde{x} - \Delta C\tilde{y} + \Delta D^{T}\tilde{z} - \Delta g = R_{g}, \\ \Delta D\tilde{y} + \Delta E\tilde{z} - \Delta h = R_{h}, \end{array} \right\}$$

$$(3.4)$$

and $\Delta A \in \mathcal{S}_n, \Delta C \in \mathcal{S}_m, \Delta E \in \mathcal{S}_p$. To maintain the sparsity pattern of A, B, C, D and E to the perturbation matrices, we replace $\Delta A, \Delta B, \Delta C, \Delta D$ and ΔE by $\Delta A \odot \Theta_A, \Delta B \odot \Theta_B$, $\Delta C \odot \Theta_C, \Delta D \odot \Theta_D$ and $\Delta E \odot \Theta_E$, respectively. Thus (3.4) can be equivalently reformulated as:

$$\theta_1^{-1}\theta_1(\Delta A \odot \Theta_A)\tilde{x} + \theta_2^{-1}\theta_2(\Delta B \odot \Theta_B)^T \tilde{y} - \theta_8^{-1}\theta_8 \Delta f = R_f, \tag{3.5}$$

$$\theta_2^{-1}\theta_2(\Delta B \odot \Theta_B)\tilde{x} - \theta_4^{-1}\theta_4(\Delta C \odot \Theta_C)\tilde{y} + \theta_5^{-1}\theta_5(\Delta D \odot \Theta_D)^T - \theta_9^{-1}\theta_9\Delta g = R_g, \quad (3.6)$$

$$\theta_5^{-1}\theta_5(\Delta D \odot \Theta_D)\tilde{x} + \theta_7^{-1}\theta_7(\Delta E \odot \Theta_E)^T \tilde{y} - \theta_{10}^{-1}\theta_{10}\Delta h = R_h.$$
(3.7)

Using Lemma 2.1 in (3.5), we get

$$\theta_1^{-1} \mathcal{K}_{\tilde{x}} \Phi_A \theta_1 \operatorname{vec}_{\mathcal{S}}(A \odot \Theta_A) + \theta_2^{-1} \mathcal{N}_{\tilde{y}}^n \Phi_B \theta_2 \operatorname{vec}(B \odot \Theta_B) - \theta_8^{-1} \theta_8 \Delta f = R_f.$$
(3.8)

Further, (3.8) can be express as

$$\theta_1^{-1} \mathcal{K}_{\tilde{x}} \Phi_A \mathfrak{D}_{\mathcal{S}_n}^{-1} \mathfrak{D}_{\mathcal{S}_n} \theta_1 \operatorname{vec}_{\mathcal{S}} (A \odot \Theta_A) + \theta_2^{-1} \mathcal{N}_{\tilde{y}}^n \Phi_B \theta_2 \operatorname{vec}(B \odot \Theta_B) - \theta_8^{-1} \theta_8 \Delta f = R_f.$$
(3.9)

Equivalently, (3.9) can be written as follows:

$$\mathcal{J}^{1}_{\mathbb{S}_{1}}\Delta X = \begin{bmatrix} R_{f} \\ R_{g} \\ R_{h} \end{bmatrix}, \qquad (3.10)$$

where

$$\mathcal{J}_{\mathbb{S}_1}^1 = \begin{bmatrix} \theta_1^{-1} \mathcal{K}_{\tilde{x}} \Phi_A \mathfrak{D}_{\mathcal{S}_n}^{-1} & \theta_2^{-1} \mathcal{N}_{\tilde{y}}^n \Phi_B & \mathbf{0} & \mathbf{0} & -\theta_8^{-1} I_n & \mathbf{0} & \mathbf{0} \end{bmatrix} \in \mathbb{R}^{n \times l}$$

and

$$\Delta X = \begin{bmatrix} \theta_1 \mathfrak{D}_{S_n} \operatorname{vec}_{\mathcal{S}} (\Delta A \odot \Theta_A) \\ \theta_2 \operatorname{vec} (\Delta B \odot \Theta_B) \\ \theta_4 \mathfrak{D}_{S_m} \operatorname{vec}_{\mathcal{S}} (\Delta C \odot \Theta_C) \\ \theta_5 \operatorname{vec} (\Delta D \odot \Theta_D) \\ \theta_7 \mathfrak{D}_{S_p} \operatorname{vec}_{\mathcal{S}} (\Delta E \odot \Theta_E) \\ \theta_8 \Delta f \\ \theta_9 \Delta g \\ \theta_{10} \Delta h \end{bmatrix} \in \mathbb{R}^l.$$
(3.11)

Similarly, using Lemma 2.1 to (3.6) and (3.7), we obtain

$$\mathcal{J}_{\mathbb{S}_1}^2 \Delta X = \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \quad \text{and} \quad \mathcal{J}_{\mathbb{S}_1}^3 \Delta X = \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix}, \quad (3.12)$$

where $\mathcal{J}^2_{\mathbb{S}_1} \in \mathbb{R}^{m \times l}$ and $\mathcal{J}^3_{\mathbb{S}_1} \in \mathbb{R}^{p \times l}$ are given by

$$\mathcal{J}_{\mathbb{S}_1}^2 = \begin{bmatrix} \mathbf{0} \quad \theta_2^{-1} \mathcal{M}_{\tilde{x}}^m \Phi_B & -\theta_4^{-1} \mathcal{K}_{\tilde{y}} \Phi_C \mathfrak{D}_{\mathcal{S}_m}^{-1} & \theta_5^{-1} \mathcal{N}_{\tilde{z}}^m \Phi_D & \mathbf{0} & \mathbf{0} & -\theta_9^{-1} I_m & \mathbf{0} \end{bmatrix}$$
(3.13)

and

$$\mathcal{J}_{\mathbb{S}_1}^3 = \begin{bmatrix} \mathbf{0} & \mathbf{0} & \mathbf{0} & \theta_5^{-1} \mathcal{M}_{\tilde{y}}^p \Phi_D & \theta_7^{-1} \mathcal{K}_{\tilde{z}} \mathfrak{D}_{\mathcal{S}_p}^{-1} \Phi_E & \mathbf{0} & \mathbf{0} & -\theta_{10}^{-1} I_p \end{bmatrix},$$
(3.14)

respectively. Combining (3.10) and (3.12), we obtain the following equivalent system

$$\mathcal{J}_{\mathbb{S}_1} \Delta X = \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix}.$$
(3.15)

Clearly, for $\theta_8, \theta_9, \theta_{10} \neq 0$, $\mathcal{J}_{\mathbb{S}_1}$ has full row rank. Therefore, by Lemma 2.3, the minimum norm solution of (3.15) is given by

$$\Delta X_{\min} = \mathcal{J}_{\mathbb{S}_1}^{\dagger} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} = \mathcal{J}_{\mathbb{S}_1}^T (\mathcal{J}_{\mathbb{S}_1} \mathcal{J}_{\mathbb{S}_1}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix}.$$
(3.16)

On the other hand, the minimization problem in Definition 2.2 can be reformulated as:

$$\begin{aligned} \left[\boldsymbol{\eta}_{\mathsf{sps}}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) \right]^{2} &= \min \left\{ \theta_{1}^{2} \| \Delta A \odot \Theta_{A} \|_{F}^{2} + \theta_{2}^{2} \| \Delta B \odot \Theta_{B} \|_{F}^{2} + \theta_{4}^{2} \| \Delta C \odot \Theta_{C} \|_{F}^{2} + \theta_{5}^{2} \| \Delta D \odot \Theta_{D} \|_{F}^{2} \\ &+ \theta_{7}^{2} \| \Delta E \odot \Theta_{E} \|_{F}^{2} + \theta_{8}^{2} \| \Delta f \|_{2}^{2} + \theta_{9}^{2} \| \Delta g \|_{2}^{2} + \theta_{10}^{2} \| \Delta h \|_{2}^{2} \mid \\ &\left(\begin{array}{c} \Delta A \odot \Theta_{A}, \Delta B \odot \Theta_{B}, \Delta C \odot \Theta_{C}, \\ \Delta D \odot \Theta_{D}, \Delta E \odot \Theta_{E}, \Delta f, \\ \Delta g, \Delta h \end{array} \right) \in \mathbb{S}_{1} \right\} \\ &= \min \left\{ \theta_{1}^{2} \| \mathfrak{D}_{\mathcal{S}_{n}} \operatorname{vec}_{\mathcal{S}}(\Delta A \odot \Theta_{A}) \|_{2}^{2} + \theta_{2}^{2} \| \operatorname{vec}(\Delta B \odot \Theta_{B}) \|_{2}^{2} \\ &+ \theta_{4}^{2} \| \mathfrak{D}_{\mathcal{S}_{n}} \operatorname{vec}_{\mathcal{S}}(\Delta C \odot \Theta_{C}) \|_{2}^{2} + \theta_{5}^{2} \| \operatorname{vec}(\Delta D \odot \Theta_{D}) \|_{2}^{2} \\ &+ \theta_{7}^{2} \| \mathfrak{D}_{\mathcal{S}_{p}} \operatorname{vec}_{\mathcal{S}}(\Delta E \odot \Theta_{E}) \|_{2}^{2} + \theta_{5}^{2} \| \Delta f \|_{2}^{2} + \theta_{9}^{2} \| \Delta g \|_{2}^{2} + \theta_{10}^{2} \| \Delta h \|_{2}^{2} \right| \\ &\mathcal{J}_{\mathbb{S}_{1}} \Delta X = \begin{bmatrix} R_{f} \\ R_{g} \\ R_{h} \end{bmatrix} \right\} \tag{3.17} \\ &= \min \left\{ \left\| \Delta X \right\|_{2}^{2} \left| \mathcal{J}_{\mathbb{S}_{1}} \Delta X = \begin{bmatrix} R_{f} \\ R_{g} \\ R_{h} \end{bmatrix} \right\} = \| \Delta X_{\min} \|_{2}^{2}. \end{aligned}$$

Consequently, substituting (3.16) into (3.17), we obtain

$$\boldsymbol{\eta}_{\mathtt{sps}}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \mathcal{J}_{\mathbb{S}_1}^T (\mathcal{J}_{\mathbb{S}_1} \mathcal{J}_{\mathbb{S}_1}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \right\|_2.$$

From (3.11), we have $\theta_1 \mathfrak{D}_{S_n} \operatorname{vec}_{S}(\Delta A \odot \Theta_A) = \begin{bmatrix} I_{\mu} & \mathbf{0} \end{bmatrix} \Delta X$. Therefore, the generating vector for the minimal perturbation matrix $\widehat{\Delta A}_{sps}$ which also preserves the sparsity pattern is given by

$$\operatorname{vec}_{\mathcal{S}}(\widehat{\Delta A}_{\operatorname{sps}}) = \theta_1^{-1} \mathfrak{D}_{\mathcal{S}_n}^{-1} \begin{bmatrix} I_{\mu} & \mathbf{0} \end{bmatrix} \Delta X_{\min}.$$

Similarly, the generating vectors for other minimal perturbation matrices can be obtained. Hence, the proof is completed. \blacksquare

Corollary 3.1. Suppose the approximate solution of the TBSPP (1.1) with $A \in S_n$, $C \in S_m$, $E \in S_p$, and $\theta_8, \theta_9, \theta_{10} \neq 0$ is $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$. Then, we have

$$\boldsymbol{\eta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \widehat{\mathcal{J}}_{\mathbb{S}_1}^T (\widehat{\mathcal{J}}_{\mathbb{S}_1} \widehat{\mathcal{J}}_{\mathbb{S}_1}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \right\|_2,$$
(3.19)

where $\widehat{\mathcal{J}}_{\mathbb{S}_1} \in \mathbb{R}^{(n+m+p) \times l}$ is given by

$$\widehat{\mathcal{J}}_{\mathbb{S}_{1}} = \begin{bmatrix} \frac{1}{\theta_{1}} \mathcal{K}_{\tilde{x}} \mathfrak{D}_{\mathcal{S}_{n}}^{-1} & \frac{1}{\theta_{2}} \mathcal{N}_{\tilde{y}}^{n} & \mathbf{0} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{8}} I_{n} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \frac{1}{\theta_{2}} \mathcal{M}_{\tilde{x}}^{m} & -\frac{1}{\theta_{4}} \mathcal{K}_{\tilde{y}} \mathfrak{D}_{\mathcal{S}_{m}}^{-1} & \frac{1}{\theta_{5}} \mathcal{N}_{\tilde{z}}^{m} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{9}} I_{m} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \frac{1}{\theta_{5}} \mathcal{M}_{\tilde{y}}^{p} & \frac{1}{\theta_{7}} \mathcal{K}_{\tilde{z}} \mathfrak{D}_{\mathcal{S}_{p}}^{-1} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{10}} I_{p} \end{bmatrix}.$$

Proof. Since we are not preserving the sparsity pattern, the proof follows by considering $\Theta_A = \mathbf{1}_{n \times n}, \ \Theta_B = \mathbf{1}_{m \times n}, \ \Theta_C = \mathbf{1}_{m \times m}, \ \Theta_D = \mathbf{1}_{p \times m}, \ \text{and} \ \Theta_E = \mathbf{1}_{p \times p}$ in Theorem 3.1.

Remark 3.1. The structure-preserving minimal perturbation matrices ΔA , ΔB , ΔC , ΔD , ΔE , $\widehat{\Delta E}$, $\widehat{\Delta f}$, $\widehat{\Delta g}$, and $\widehat{\Delta h}$, for which $\eta^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ is attained are given by formulae presented in Theorem 3.1 with $\mathcal{J}_{\mathbb{S}_1} = \widehat{\mathcal{J}}_{\mathbb{S}_1}$.

In the next result, we present the formula of structured *BE* when $C = \mathbf{0}$ and $E = \mathbf{0}$. **Corollary 3.2.** Suppose $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ is an approximate solution of the TBSPP (1.1) with $A \in S_n, C = \mathbf{0}, E = \mathbf{0}, and \theta_8, \theta_9, \theta_{10} \neq 0$. Then, we have

$$\boldsymbol{\eta}_{\mathbf{sps}}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \widetilde{\mathcal{J}}_{\mathbb{S}_1}^T (\widetilde{\mathcal{J}}_{\mathbb{S}_1} \widetilde{\mathcal{J}}_{\mathbb{S}_1}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \right\|_2,$$
(3.20)

where $\widetilde{\mathcal{J}}_{\mathbb{S}_1} \in \mathbb{R}^{(n+m+p) \times l}$ is given by

$$\widetilde{\mathcal{J}}_{\mathbb{S}_1} = \begin{bmatrix} \frac{1}{\theta_1} \mathcal{K}_{\tilde{x}} \Phi_A \mathfrak{D}_{\mathcal{S}_n}^{-1} & \frac{1}{\theta_2} \mathcal{N}_{\tilde{y}}^n \Phi_B & \mathbf{0} & -\frac{1}{\theta_8} I_n & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \frac{1}{\theta_2} \mathcal{M}_{\tilde{x}}^m \Phi_B & \frac{1}{\theta_5} \mathcal{N}_{\tilde{z}}^m \Phi_D & \mathbf{0} & -\frac{1}{\theta_9} I_m & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \frac{1}{\theta_5} \mathcal{M}_{\tilde{y}}^p \Phi_D & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{10}} I_p \end{bmatrix},$$

 $R_f = f - A\tilde{x} - B^T\tilde{y}, R_g = g - B\tilde{x} - D^T\tilde{z}, R_h = h - D\tilde{y}, and \boldsymbol{l} = \mu + mn + mp + m + n + p.$

Proof. Since $C = \mathbf{0}$ and $E = \mathbf{0}$, the proof follows by considering $\theta_3 = \theta_5 = 0$.

Remark 3.2. When $C = \mathbf{0}$ and $E = \mathbf{0}$, Lv and Zheng [19] derive the structured BE for the TBSPP (1.1). However, their investigations do not take into account the sparsity pattern of the coefficient matrices.

4 Derivation of structured *BEs* for case (ii)

In this section, we derive explicit formulae for the structured *BEs* for the *TBSPP* for the case (*ii*), i.e., $A \in S_n$, $B_1 \neq B_2$, $C \in S_m$, $D_1 = D_2 =: D$ and $E \in S_p$. We use the Lemmas 2.1, 2.2 and 2.3, and apply a similar methodology used in Section 3 to derive the formulae for the structured *BEs*. In the next result, we present computable formulae for the structured *BE* $\eta_{sps}^{S_2}(\tilde{x}, \tilde{y}, \tilde{z})$ by preserving sparsity pattern of the original matrices to the perturbation matrices. Before proceeding, we set the following notations:

$$\Phi_{B_1} = \operatorname{diag}(\operatorname{vec}(\Theta_{B_1})), \ \Phi_{B_2} = \operatorname{diag}(\operatorname{vec}_{\mathcal{S}}(\Theta_{B_2})),$$

and Φ_A, Φ_C, Φ_D and Φ_E are same as defined in Section 3.

Theorem 4.1. Let $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ be an approximate solution of the TBSPP (1.1) with $A \in S_n, B_1 \neq B_2, C \in S_m, D_1 = D_2 = D, E \in S_p$ and $\theta_8, \theta_9, \theta_{10} \neq 0$. Then, we have

$$\boldsymbol{\eta}_{\mathbf{sps}}^{\mathbb{S}_2}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \mathcal{J}_{\mathbb{S}_2}^T (\mathcal{J}_{\mathbb{S}_2} \mathcal{J}_{\mathbb{S}_2}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \right\|_2,$$
(4.1)

where $\mathcal{J}_{\mathbb{S}_2} \in \mathbb{R}^{(n+m+p) \times l}$ is given by

$$\mathcal{J}_{\mathbb{S}_{2}} = \begin{bmatrix} \frac{1}{\theta_{1}} \mathcal{K}_{\bar{x}} \Phi_{A} \mathfrak{D}_{S_{n}}^{-1} & \frac{1}{\theta_{2}} \mathcal{N}_{\bar{y}}^{n} \Phi_{B_{1}} & \mathbf{0} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{3}} I_{n} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \frac{1}{\theta_{3}} \mathcal{M}_{\bar{x}}^{m} \Phi_{B_{2}} & -\frac{1}{\theta_{4}} \mathcal{K}_{\bar{y}} \Phi_{C} \mathfrak{D}_{S_{m}}^{-1} & \frac{1}{\theta_{5}} \mathcal{N}_{\bar{z}}^{m} \Phi_{D} & \mathbf{0} & -\frac{1}{\theta_{9}} I_{m} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \frac{1}{\theta_{5}} \mathcal{M}_{\bar{y}}^{p} \Phi_{D} & \frac{1}{\theta_{7}} \mathcal{K}_{\bar{z}} \Phi_{E} \mathfrak{D}_{S_{p}}^{-1} & \mathbf{0} & -\frac{1}{\theta_{10}} I_{p} \end{bmatrix},$$

 $R_f = f - A\tilde{x} - B_1^T \tilde{y}, R_g = g - B_2 \tilde{x} + C\tilde{y} - D^T \tilde{z}, R_h = h - D\tilde{y} - E\tilde{z} \text{ and } \boldsymbol{l} = n^2 + \sigma + \tau + 2mn + mp + m + n + p.$

Proof. Given that $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ is an approximate solution of the *TBSPP* (1.1) for the case (*ii*). Now, we are required to construct perturbations matrices $\Delta A \in \mathcal{S}_n$, $\Delta B_1, \Delta B_2 \in \mathbb{R}^{m \times n}$, $\Delta C \in \mathcal{S}_m$, $\Delta D \in \mathbb{R}^{p \times m}$, $\Delta E \in \mathcal{S}_p$, which maintain the sparsity pattern of A, B_1, B_2, C, D, E , respectively, and the perturbations $\Delta f \in \mathbb{R}^n$, $\Delta g \in \mathbb{R}^m$ and $\Delta h \in \mathbb{R}^p$. Using (2.3), $\begin{pmatrix} \Delta A, \Delta B_1, \Delta B_2, \\ \Delta C, \Delta D, \Delta E, \end{pmatrix} \in \mathbb{S}_2$ if and only if $\Delta A, \Delta B_1, \Delta B_2, \Delta C, \Delta D, \Delta E, \Delta f, \Delta g$ and

$$\sum_{\Delta h \text{ satisfy the following equations:}} \Delta f, \Delta g, \Delta h \quad \mathcal{J}$$

 Δh satisfy the following equations:

$$\left. \begin{array}{l} \Delta A\tilde{x} + \Delta B_{1}^{T}\tilde{y} - \Delta f = R_{f}, \\ \Delta B_{2}\tilde{x} - \Delta C\tilde{y} + \Delta D^{T}\tilde{z} - \Delta g = R_{g}, \\ \Delta D\tilde{y} + \Delta E\tilde{z} - \Delta h = R_{h}, \end{array} \right\}$$

$$(4.2)$$

and $\Delta A \in \mathcal{S}_n, \Delta C \in \mathcal{S}_m, \Delta E \in \mathcal{S}_p$.

By following a similar the proof methodology of Theorem 3.1 and applying Lemma 2.1, we get:

$$\mathcal{J}_{\mathbb{S}_2}^1 \Delta X = R_f, \ \mathcal{J}_{\mathbb{S}_2}^2 \Delta X = R_g \text{ and } \mathcal{J}_{\mathbb{S}_2}^3 \Delta X = R_h,$$
(4.3)

where

$$\mathcal{J}_{\mathbb{S}_{2}}^{1} = \begin{bmatrix} \theta_{1}^{-1} \mathcal{K}_{\tilde{x}} \Phi_{A} \mathfrak{D}_{\mathcal{S}_{n}}^{-1} & \theta_{2}^{-1} \mathcal{N}_{\tilde{y}}^{n} \Phi_{B_{1}} & \mathbf{0} & \mathbf{0} & \mathbf{0} & -\theta_{8}^{-1} I_{n} & \mathbf{0} & \mathbf{0} \end{bmatrix} \in \mathbb{R}^{n \times l}, \\ \mathcal{J}_{\mathbb{S}_{2}}^{2} = \begin{bmatrix} \mathbf{0} & \mathbf{0} & \theta_{3}^{-1} \mathcal{M}_{\tilde{x}}^{m} \Phi_{B_{2}} & -\theta_{4}^{-1} \mathcal{K}_{\tilde{y}} \Phi_{C} \mathfrak{D}_{\mathcal{S}_{m}}^{-1} & \theta_{5}^{-1} \mathcal{N}_{\tilde{y}}^{m} \Phi_{D} & \mathbf{0} & \mathbf{0} & -\theta_{9}^{-1} I_{m} & \mathbf{0} \end{bmatrix} \in \mathbb{R}^{m \times l}, \\ \mathcal{J}_{\mathbb{S}_{2}}^{3} = \begin{bmatrix} \mathbf{0} & \mathbf{0} & \mathbf{0} & \theta_{5}^{-1} \mathcal{M}_{\tilde{y}}^{p} \Phi_{D} & \theta_{7}^{-1} \mathcal{K}_{\tilde{z}} \Phi_{E} \mathfrak{D}_{\mathcal{S}_{p}}^{-1} & \mathbf{0} & \mathbf{0} & -\theta_{10}^{-1} I_{p} \end{bmatrix} \in \mathbb{R}^{p \times l},$$

and

$$\Delta X = \begin{bmatrix} \theta_1 \mathfrak{D}_{S_n} \operatorname{vec}_{\mathcal{S}} (\Delta A \odot \Theta_A) \\ \theta_2 \operatorname{vec} (\Delta B \odot \Theta_{B_1}) \\ \theta_3 \operatorname{vec} (\Delta B \odot \Theta_{B_2}) \\ \theta_4 \mathfrak{D}_{S_m} \operatorname{vec}_{\mathcal{S}} (\Delta C \odot \Theta_C) \\ \theta_5 \operatorname{vec} (\Delta D \odot \Theta_D) \\ \theta_7 \mathfrak{D}_{S_p} \operatorname{vec}_{\mathcal{S}} (\Delta E \odot \Theta_E) \\ \theta_8 \Delta f \\ \theta_9 \Delta g \\ \theta_{10} \Delta h \end{bmatrix} \in \mathbb{R}^l.$$

$$(4.4)$$

Combining the three equations in (4.3), we obtain

$$\mathcal{J}_{\mathbb{S}_2} \Delta X = \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix}.$$
(4.5)

Since, $\mathcal{J}_{\mathbb{S}_2}$ has full row rank for $\theta_8, \theta_9, \theta_{10} \neq 0$. Therefore, (4.5) is consistent and by Lemma 2.3, its minimum norm solution is given by

$$\Delta X_{\min} = \mathcal{J}_{\mathbb{S}_2}^T (\mathcal{J}_{\mathbb{S}_2} \mathcal{J}_{\mathbb{S}_2}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix}.$$
(4.6)

Now, applying a similar argument to the proof method of Theorem 3.1, the required structured BE is

$$\boldsymbol{\eta}_{\mathbf{sps}}^{\mathbb{S}_2}(\tilde{x}, \tilde{y}, \tilde{z}) = \|\Delta X_{\min}\|_2 = \left\| \mathcal{J}_{\mathbb{S}_2}^T (\mathcal{J}_{\mathbb{S}_2} \mathcal{J}_{\mathbb{S}_2}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \right\|_2.$$

Hence, the proof is completed. \blacksquare

Remark 4.1. The minimal perturbation matrices $\widehat{\Delta A}_{sps}$, $\widehat{\Delta C}_{sps}$, $\widehat{\Delta E}_{sps}$, $\widehat{\Delta f}_{sps}$, $\widehat{\Delta g}_{sps}$ and $\widehat{\Delta h}_{sps}$ for the Problem 2.1 can be computed using the formulae provided in Theorem 3.1 by replacing $\mathcal{J}_{\mathbb{S}_1}$ with $\mathcal{J}_{\mathbb{S}_2}$. The generating vectors for the minimal perturbation matrices $\widehat{\Delta B}_{1sps}$ and $\widehat{\Delta B}_{2sps}$ are given by

$$\operatorname{vec}(\widehat{\Delta B}_{1\,\operatorname{sps}}) = \frac{1}{\theta_2} \begin{bmatrix} \mathbf{0}_{\mu} & I_{mn} & \mathbf{0}_{l-\mu-mn} \end{bmatrix} \mathcal{J}_{\mathbb{S}_2}^T (\mathcal{J}_{\mathbb{S}_2} \mathcal{J}_{\mathbb{S}_2}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix},$$

$$\operatorname{vec}(\widehat{\Delta B}_{2_{\operatorname{sps}}}) = \frac{1}{\theta_3} \begin{bmatrix} \mathbf{0}_{\mu+mn} & I_{mn} & \mathbf{0}_{l-\mu-2mn} \end{bmatrix} \mathcal{J}_{\mathbb{S}_2}^T (\mathcal{J}_{\mathbb{S}_2} \mathcal{J}_{\mathbb{S}_2}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix}$$

In the subsequent result, we provide the structured BE while the sparsity pattern is not preserved.

Corollary 4.1. Let $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ be an approximate solution of the TBSPP (1.1) with $A \in S_n, C \in S_m, E \in S_p$, and $\theta_8, \theta_9, \theta_{10} \neq 0$. Then, we have

$$\boldsymbol{\eta}^{\mathbb{S}_2}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \widehat{\mathcal{J}}_{\mathbb{S}_2}^T (\widehat{\mathcal{J}}_{\mathbb{S}_2} \widehat{\mathcal{J}}_{\mathbb{S}_2}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \right\|_2,$$
(4.7)

where $\widehat{\mathcal{J}}_{\mathbb{S}_2} \in \mathbb{R}^{(n+m+p) \times l}$ is given by

$$\widehat{\mathcal{J}}_{\mathbb{S}_{2}} = \begin{bmatrix} \frac{1}{\theta_{1}}\mathcal{M}_{\tilde{x}}^{n} & \frac{1}{\theta_{2}}\mathcal{N}_{\tilde{y}}^{n} & \mathbf{0} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{8}}I_{n} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \frac{1}{\theta_{3}}\mathcal{M}_{\tilde{x}}^{m} & -\frac{1}{\theta_{4}}\mathcal{K}_{\tilde{y}}\mathfrak{D}_{\mathcal{S}_{m}}^{-1} & \frac{1}{\theta_{5}}\mathcal{N}_{\tilde{z}}^{m} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{9}}I_{m} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \frac{1}{\theta_{5}}\mathcal{M}_{\tilde{y}}^{p} & \frac{1}{\theta_{7}}\mathcal{K}_{\tilde{z}}\mathfrak{D}_{\mathcal{S}_{p}}^{-1} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{10}}I_{p} \end{bmatrix}$$

Proof. The proof proceeds by choosing $\Theta_A = \mathbf{1}_{n \times n}$, $\Theta_{B_1} = \Theta_{B_2} = \mathbf{1}_{m \times n}$, $\Theta_C = \mathbf{1}_{m \times m}$, $\Theta_D = \mathbf{1}_{p \times m}$, and $\Theta_E = \mathbf{1}_{p \times p}$ in the expression of the structured *BE* presented in Theorem 4.1.

Remark 4.2. Similar to Corollary 3.2, we can compute the structured BE for the case (ii) with $C = \mathbf{0}$ and $E = \mathbf{0}$. This specific instance of structured BE has also been addressed in [21]. However, our investigation additionally ensures the preservation of the sparsity pattern.

5 Derivation of structured BEs for case (*iii*)

This section deals with the structured BE of the TBSPP for the case (*iii*), i.e., $A \neq A^T$, $C \in S_m$, $E \in S_p$, $B_1 = B_2 =: B$, and $D_1 \neq D_2$. Using a similar technique as in Section 3, in the following theorem, we present the computable formula of the structured BE when sparsity pattern of the original matrices are preserved in the perturbation matrices. Before continuing, we introduce the following notations:

$$\Phi_{D_1} = \operatorname{diag}(\operatorname{vec}(\Theta_{D_1})), \ \Phi_{D_2} = \operatorname{diag}(\operatorname{vec}_{\mathcal{S}}(\Theta_{D_2})),$$

along with Φ_A , Φ_B , Φ_C and Φ_E as defined in Section 3.

Theorem 5.1. Let $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ be an approximate solution of the TBSPP (1.1) with $C \in S_m, E \in S_p$, and $\theta_8, \theta_9, \theta_{10} \neq 0, B_1 = B_2 =: B$ and $D_1 \neq D_2$. Then, we have

$$\boldsymbol{\eta}_{\mathbf{sps}}^{\mathbb{S}_3}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \mathcal{J}_{\mathbb{S}_3}^T (\mathcal{J}_{\mathbb{S}_3} \mathcal{J}_{\mathbb{S}_3}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \right\|_2,$$
(5.1)

where $\mathcal{J}_{\mathbb{S}_3} \in \mathbb{R}^{(n+m+p) \times l}$ is given by

$$\mathcal{J}_{\mathbb{S}_3} = \begin{bmatrix} \frac{1}{\theta_1} \mathcal{M}_{\bar{x}}^n \Phi_A & \frac{1}{\theta_2} \mathcal{N}_{\bar{y}}^n \Phi_B & \mathbf{0} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_8} I_n & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \frac{1}{\theta_2} \mathcal{M}_{\bar{x}}^m \Phi_B & -\frac{1}{\theta_4} \mathcal{K}_{\bar{y}} \Phi_C \mathfrak{D}_{S_m}^{-1} & \frac{1}{\theta_5} \mathcal{N}_{\bar{z}}^m \Phi_{D_1} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_7} \mathcal{K}_{\bar{z}} \Phi_E \mathfrak{D}_{S_p}^{-1} & \mathbf{0} & -\frac{1}{\theta_9} I_m & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \frac{1}{\theta_6} \mathcal{M}_{\bar{y}}^p \Phi_{D_2} & \frac{1}{\theta_7} \mathcal{K}_{\bar{z}} \Phi_E \mathfrak{D}_{S_p}^{-1} & \mathbf{0} & \mathbf{0} & -\frac{1}{\theta_{10}} I_p \end{bmatrix},$$

 $R_f = f - A\tilde{x} - B^T \tilde{y}, R_g = g - B\tilde{x} + C\tilde{y} - D_1^T \tilde{z}, R_h = h - D_2 \tilde{y} - E\tilde{z} \text{ and } \boldsymbol{l} = n^2 + \sigma + \tau + mn + 2mp + m + n + p.$

Proof. For the given approximate solution $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$, we are required to construct perturbations matrices $\Delta A \in \mathbb{R}^n$, $\Delta B \in \mathbb{R}^{m \times n}$, $\Delta C \in \mathcal{S}_m$, $\Delta D_1, \Delta D_2 \in \mathbb{R}^{p \times m}$, $\Delta E \in \mathcal{S}_p$, which maintain the sparsity pattern of A, B, C, D_1, D_2, E , respectively, and the perturba-

tions $\Delta f \in \mathbb{R}^n$, $\Delta g \in \mathbb{R}^m$ and $\Delta h \in \mathbb{R}^p$. By (2.4), $\begin{pmatrix} \Delta A, \Delta B, \Delta C, \\ \Delta D_1, \Delta D_2, \Delta E, \\ \Delta f, \Delta g, \Delta h \end{pmatrix} \in \mathbb{S}_3$ if and only

if $\Delta A, \Delta B, \Delta C, \Delta D_1, \Delta D_2 \Delta E, \Delta f, \Delta g$ and Δh satisfy the following equations:

$$\left. \begin{array}{l} \Delta A\tilde{x} + \Delta B^{T}\tilde{y} - \Delta f = R_{f}, \\ \Delta B\tilde{x} - \Delta C\tilde{y} + \Delta D_{1}^{T}\tilde{z} - \Delta g = R_{g}, \\ \Delta D_{2}\tilde{y} + \Delta E\tilde{z} - \Delta h = R_{h}, \end{array} \right\}$$
(5.2)

and $\Delta C \in \mathcal{S}_m, \Delta E \in \mathcal{S}_p$.

By following a similar the proof methodology of Theorem 3.1 and applying Lemmas 2.1 and 2.2, we get:

$$\mathcal{J}^{1}_{\mathbb{S}_{3}}\Delta X = R_{f}, \ \mathcal{J}^{2}_{\mathbb{S}_{3}}\Delta X = R_{g} \text{ and } \mathcal{J}^{3}_{\mathbb{S}_{3}}\Delta X = R_{h},$$
(5.3)

where

$$\begin{aligned} \mathcal{J}_{\mathbb{S}_{3}}^{1} &= \begin{bmatrix} \theta_{1}^{-1} \mathcal{M}_{\tilde{x}}^{n} \Phi_{A} & \theta_{2}^{-1} \mathcal{N}_{\tilde{y}}^{n} \Phi_{B} & \mathbf{0} & \mathbf{0} & \mathbf{0} & -\theta_{8}^{-1} I_{n} & \mathbf{0} & \mathbf{0} \end{bmatrix} \in \mathbb{R}^{n \times l}, \\ \mathcal{J}_{\mathbb{S}_{3}}^{2} &= \begin{bmatrix} \mathbf{0} & \theta_{2}^{-1} \mathcal{M}_{\tilde{x}}^{m} \Phi_{B} & -\theta_{4}^{-1} \mathcal{K}_{\tilde{y}} \Phi_{C} \mathfrak{D}_{\mathcal{S}_{m}}^{-1} & \theta_{5}^{-1} \mathcal{N}_{\tilde{y}}^{m} \Phi_{D_{1}} & \mathbf{0} & \mathbf{0} & -\theta_{9}^{-1} I_{m} & \mathbf{0} \end{bmatrix} \in \mathbb{R}^{m \times l}, \\ \mathcal{J}_{\mathbb{S}_{3}}^{3} &= \begin{bmatrix} \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \theta_{6}^{-1} \mathcal{M}_{\tilde{y}}^{p} \Phi_{D} & \theta_{7}^{-1} \mathcal{K}_{\tilde{z}} \Phi_{E} \mathfrak{D}_{\mathcal{S}_{p}}^{-1} & \mathbf{0} & \mathbf{0} & -\theta_{10}^{-1} I_{p} \end{bmatrix} \in \mathbb{R}^{p \times l}, \end{aligned}$$

and

$$\Delta X = \begin{bmatrix} \theta_1 \operatorname{vec}(\Delta A \odot \Theta_A) \\ \theta_2 \operatorname{vec}(\Delta B \odot \Theta_B) \\ \theta_4 \mathfrak{D}_{\mathcal{S}_m} \operatorname{vec}_{\mathcal{S}}(\Delta C \odot \Theta_C) \\ \theta_5 \operatorname{vec}(\Delta D \odot \Theta_{D_1}) \\ \theta_6 \operatorname{vec}(\Delta D \odot \Theta_{D_2}) \\ \theta_7 \mathfrak{D}_{\mathcal{S}_p} \operatorname{vec}_{\mathcal{S}}(\Delta E \odot \Theta_E) \\ \theta_8 \Delta f \\ \theta_9 \Delta g \\ \theta_{10} \Delta h \end{bmatrix} \in \mathbb{R}^l.$$
(5.4)

Combining the three equations in (5.3), we obtain

$$\mathcal{J}_{\mathbb{S}_3} \Delta X = \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix}.$$
(5.5)

Observed that, for $\theta_8, \theta_9, \theta_{10} \neq 0$, $\mathcal{J}_{\mathbb{S}_3}$ has full row rank. Therefore, (5.5) is consistent, and by Lemma 2.3, its minimum norm solution is given by

$$\Delta X_{\min} = \mathcal{J}_{\mathbb{S}_3}^T (\mathcal{J}_{\mathbb{S}_3} \mathcal{J}_{\mathbb{S}_3}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix}.$$
 (5.6)

Applying a similar proof technique of Theorem 3.1, the required structured BE is

$$\boldsymbol{\eta}_{\mathbf{sps}}^{\mathbb{S}_3}(\tilde{x}, \tilde{y}, \tilde{z}) = \|\Delta X_{\min}\|_2 = \left\| \mathcal{J}_{\mathbb{S}_3}^T (\mathcal{J}_{\mathbb{S}_3} \mathcal{J}_{\mathbb{S}_3}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \right\|_2.$$

Hence, the proof is completed. \blacksquare

Remark 5.1. The minimal perturbation matrices $\widehat{\Delta B}_{sps}$, $\widehat{\Delta C}_{sps}$, $\widehat{\Delta E}_{sps}$, $\widehat{\Delta f}_{sps}$, $\widehat{\Delta g}_{sps}$ and $\widehat{\Delta h}_{sps}$ can be computed using the formulae provided in Theorem 3.1 with $\mathcal{J}_{\mathbb{S}_1} = \mathcal{J}_{\mathbb{S}_3}$. The generating vector for the minimal perturbation matrices $\widehat{\Delta A}_{sps}$, $\widehat{\Delta D}_{1sps}$ and $\widehat{\Delta D}_{2sps}$ are given by

$$\operatorname{vec}(\widehat{\Delta A}_{\mathtt{sps}}) = \frac{1}{\theta_1} \begin{bmatrix} I_{n^2} & \mathbf{0}_{l-n^2} \end{bmatrix} \mathcal{J}_{\mathbb{S}_3}^T (\mathcal{J}_{\mathbb{S}_3} \mathcal{J}_{\mathbb{S}_3}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix}.$$
$$\operatorname{vec}(\widehat{\Delta D}_{1\,\mathtt{sps}}) = \frac{1}{\theta_5} \begin{bmatrix} \mathbf{0}_{n^2+\sigma+mn} & I_{mp} & \mathbf{0}_{\tau+mp+n+m+p} \end{bmatrix} \mathcal{J}_{\mathbb{S}_3}^T (\mathcal{J}_{\mathbb{S}_3} \mathcal{J}_{\mathbb{S}_3}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix},$$
$$\operatorname{vec}(\widehat{\Delta D}_{2\,\mathtt{sps}}) = \frac{1}{\theta_6} \begin{bmatrix} \mathbf{0}_{n^2+\sigma+mn+mp} & I_{mp} & \mathbf{0}_{\tau+n+m+p} \end{bmatrix} \mathcal{J}_{\mathbb{S}_3}^T (\mathcal{J}_{\mathbb{S}_3} \mathcal{J}_{\mathbb{S}_3}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix}.$$

In the next result, we present the structured BE when the sparsity pattern is not considered.

Corollary 5.1. For the approximate solution $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ of the TBSPP (1.1) with $C \in S_m$, $E \in S_p$, and $\theta_8, \theta_9, \theta_{10} \neq 0$, we have

$$\boldsymbol{\eta}^{\mathbb{S}_3}(\tilde{x}, \tilde{y}, \tilde{z}) = \left\| \widehat{\mathcal{J}}_{\mathbb{S}_3}^T (\widehat{\mathcal{J}}_{\mathbb{S}_3} \widehat{\mathcal{J}}_{\mathbb{S}_3}^T)^{-1} \begin{bmatrix} R_f \\ R_g \\ R_h \end{bmatrix} \right\|_2,$$
(5.7)

where, $\widehat{\mathcal{J}}_{\mathbb{S}_3} \in \mathbb{R}^{(n+m+p) \times l}$ is given by

$$\widehat{\mathcal{J}}_{\mathbb{S}_3} = egin{bmatrix} rac{1}{ heta_1}\mathcal{M}_{ ilde{x}}^n & rac{1}{ heta_2}\mathcal{N}_{ ilde{y}}^n & \mathbf{0} & \mathbf{0} & \mathbf{0} & -rac{1}{ heta_8}I_n & \mathbf{0} & \mathbf{0} \ \mathbf{0} & rac{1}{ heta_2}\mathcal{M}_{ ilde{x}}^n & -rac{1}{ heta_4}\mathcal{K}_{ ilde{y}}\mathfrak{D}_{\mathcal{S}_m}^{-1} & rac{1}{ heta_5}\mathcal{N}_{ ilde{x}}^m & \mathbf{0} & \mathbf{0} & -rac{1}{ heta_8}I_m & \mathbf{0} \ \mathbf{0} & \mathbf{0} & \mathbf{0} & -rac{1}{ heta_9}I_m & \mathbf{0} \ \mathbf{0} & \mathbf{0} & \mathbf{0} & rac{1}{ heta_6}\mathcal{M}_{ ilde{y}}^p & rac{1}{ heta_7}\mathcal{K}_{ ilde{z}}\mathfrak{D}_{\mathcal{S}_p}^{-1} & \mathbf{0} & \mathbf{0} & -rac{1}{ heta_{10}}I_p \end{bmatrix}.$$

Proof. The proof follows by taking $\Theta_A = \mathbf{1}_{n \times n}$, $\Theta_B = \mathbf{1}_{m \times n}$, $\Theta_C = \mathbf{1}_{m \times m}$, $\Theta_{D_1} = \Theta_{D_2} = \mathbf{1}_{p \times m}$, and $\Theta_E = \mathbf{1}_{p \times p}$ in the *BE* expression provided in Theorem 5.1.

Remark 5.2. Structured BEs for the TBSPP (1.1) when $A \in S_n$, $C \in S_m$, $E \in S_p$, $B_1 \neq B_2$, $D_1 \neq D_2$, or, $A \neq A^T$, $C \in S_m$, $E \in S_p$, $B_1 \neq B_2$, $D_1 \neq D_2$ can be derived in a similar technique used in this section and in Sections 3 and 4. As the derivation process is similar, we have not studied them here in detail.

6 Three reliable and effective termination criteria

The choice of an effective and reliable termination criterion plays a vital role when solving a linear system by using some iterative algorithms. Proper selection of effective termination criteria can improve the quality of the computed solution and can reduce the storage requirement for the iterative algorithm. For solving an unstructured linear system $\mathcal{A}w = d$, the commonly used termination criteria are given by [25]:

$$\boldsymbol{\eta}^{\texttt{term1}}(\mathcal{A}, \widetilde{\boldsymbol{w}}) := \min\left\{\delta : (\mathcal{A} + \Delta \mathcal{A})\widetilde{\boldsymbol{w}} = \boldsymbol{d} + \Delta \boldsymbol{d}, \|\Delta \mathcal{A}\|_{F} \leq \|\delta \mathcal{A}\|_{F}, \|\Delta \boldsymbol{d}\|_{2} \leq \delta \|\boldsymbol{d}\|_{2}\right\}$$
$$= \frac{\|\boldsymbol{d} - \mathcal{A}\widetilde{\boldsymbol{w}}\|_{2}}{\|\mathcal{A}\|_{F} \|\widetilde{\boldsymbol{w}}\|_{2} + \|\boldsymbol{d}\|_{2}} < \boldsymbol{tol},$$
(6.1)

$$\boldsymbol{\eta}^{\texttt{term2}}(\widetilde{\boldsymbol{w}}) := \min\left\{\delta : \mathcal{A}\widetilde{\boldsymbol{w}} = \boldsymbol{d} + \Delta \boldsymbol{d}, \|\Delta \boldsymbol{d}\|_{2} \le \delta \|\boldsymbol{d}\|_{2}\right\} = \frac{\|\boldsymbol{d} - \mathcal{A}\widetilde{\boldsymbol{w}}\|_{2}}{\|\boldsymbol{d}\|_{2}} < \boldsymbol{tol}, \qquad (6.2)$$

where "**tol**" is a given threshold tolerance. The above two termination criteria are the unstructured *BE* for the linear system $\mathcal{A}w = d$. This leads to the obtained solution satisfying a nearly perturbed linear system. However, when we use (6.1) and (6.2) to solve *TBSPP* in the form (1.1), the computed solution may not be an actual solution or a nearly perturbed *TBSPPs*, or the corresponding iterative algorithm may not be strongly backward stable. To overcome these shortcomings, we propose the following three termination criteria to solve the *TBSPP* with structure (*i*)-(*iii*), respectively:

$$\begin{split} \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) &= \min \left\{ \vartheta : \|\Delta A\|_{F} \leq \vartheta \|A\|_{F}, \|\Delta B\|_{F} \leq \vartheta \|B\|_{F}, \|\Delta C\|_{F} \leq \vartheta \|C\|_{F}, \\ \|\Delta D\|_{F} \leq \vartheta \|D\|_{F}, \|\Delta E\|_{F} \leq \vartheta \|E\|_{F}, \|\Delta f\|_{2} \leq \vartheta \|f\|_{2}, \\ \|\Delta g\|_{2} \leq \vartheta \|g\|_{2}, \|\Delta h\|_{2} \leq \vartheta \|h\|_{2}, \left(\begin{array}{c} \Delta A, \Delta B, \Delta C, \\ \Delta D, \Delta E, \Delta f, \\ \Delta g, \Delta h \end{array} \right) \in \mathbb{S}_{1} \right\} < \boldsymbol{tol}, \\ \boldsymbol{\zeta}^{\mathbb{S}_{2}}(\tilde{x}, \tilde{y}, \tilde{z}) &= \min \left\{ \vartheta : \|\Delta A\|_{F} \leq \vartheta \|A\|_{F}, \|\Delta B_{1}\|_{F} \leq \vartheta \|B_{1}\|_{F}, \|\Delta B_{2}\|_{F} \leq \vartheta \|B_{2}\|_{F}, \\ \|\Delta C\|_{F} \leq \vartheta \|C\|_{F}, \|\Delta D\|_{F} \leq \vartheta \|D\|_{F}, \|\Delta E\|_{F} \leq \vartheta \|E\|_{F}, \|\Delta f\|_{2} \leq \vartheta \|f\|_{2} \\ \|\Delta g\|_{2} \leq \vartheta \|g\|_{2}, \|\Delta h\|_{2} \leq \vartheta \|h\|_{2}, \left(\begin{array}{c} \Delta A, \Delta B_{1}, \Delta B_{2}, \\ \Delta C, \Delta D, \Delta E, \\ \Delta f, \Delta g, \Delta h \end{array} \right) \in \mathbb{S}_{2} \right\} < \boldsymbol{tol}, \end{split}$$

and

$$\begin{split} \boldsymbol{\zeta}^{\mathbb{S}_{3}}(\tilde{x}, \tilde{y}, \tilde{z}) &= \min \left\{ \vartheta : \|\Delta A\|_{F} \leq \vartheta \|A\|_{F}, \|\Delta B\|_{F} \leq \vartheta \|B\|_{F}, \|\Delta C\|_{F} \leq \vartheta \|C\|_{F}, \\ \|\Delta D_{1}\|_{F} \leq \vartheta \|D_{1}\|_{F}, \|\Delta D_{2}\|_{F} \leq \vartheta \|D_{2}\|_{F}, \|\Delta E\|_{F} \leq \vartheta \|E\|_{F}, \|\Delta f\|_{2} \leq \vartheta \|f\|_{2}, \\ \|\Delta g\|_{2} \leq \vartheta \|g\|_{2}, \|\Delta h\|_{2} \leq \vartheta \|h\|_{2}, \left(\begin{array}{c} \Delta A, \Delta B_{1}, \Delta C, \\ \Delta D_{1}, \Delta D_{2}, \Delta E, \\ \Delta f, \Delta g, \Delta h \end{array} \right) \in \mathbb{S}_{3} \right\} < \boldsymbol{tol}. \end{split}$$

Finding analytical expressions for $\boldsymbol{\zeta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z}), i = 1, 2, 3$, is quite a difficult task. For this reason, we use $\boldsymbol{\eta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z}) < \boldsymbol{tol}$ as the termination criteria with $\theta_1 = \frac{1}{\|A\|_F}, \theta_2 = \frac{1}{\|B_1\|_F}, \theta_3 = \frac{1}{\|B_2\|_F}, \theta_4 = \frac{1}{\|C\|_F}, \theta_5 = \frac{1}{\|D_1\|_F}, \theta_7 = \frac{1}{\|D_2\|_2}, \theta_8 = \frac{1}{\|f\|_2}, \theta_9 = \frac{1}{\|g\|_2} \text{ and } \theta_{10} = \frac{1}{\|h\|_2}.$ The values of $\boldsymbol{\eta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z})$ can be bounded above and below by $\boldsymbol{\zeta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z})$, which is presented in the next result.

Theorem 6.1. Let $[\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ be an approximate solution of the TBSPP (1.1). Then, we have

$$\boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z}) \le \boldsymbol{\eta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z}) \le \sqrt{8} \, \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z}), \tag{6.3}$$

$$\boldsymbol{\zeta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z}) \le \boldsymbol{\eta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z}) \le \sqrt{9} \, \boldsymbol{\zeta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z}), \text{ for } i = 2, 3.$$
(6.4)

Proof. Let $\widehat{\Delta A}, \widehat{\Delta B}, \widehat{\Delta C}, \widehat{\Delta D}, \widehat{\Delta E}, \widehat{\Delta f}, \widehat{\Delta g}$ and $\widehat{\Delta h}$ be the minimal perturbation matrices for $\eta^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$. However, these selections of matrices do not lead to an improvement in the value of $\boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$. Consequently, either $\|\widehat{\Delta A}\|_F \ge \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})\|A\|_F$ or $\|\widehat{\Delta B}\|_F \ge \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})\|B\|_F$ or $\|\widehat{\Delta C}\|_F \ge \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})\|C\|_F$ or $\|\widehat{\Delta D}\|_F \ge \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})\|D\|_F$ or $\|\widehat{\Delta E}\|_F \ge \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})\|B\|_F$ or $\|\widehat{\Delta f}\|_2 \ge \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})\|f\|_2$ or $\|\widehat{\Delta g}\|_2 \ge \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})\|g\|_2$ or $\|\widehat{\Delta h}\|_2 \ge \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})\|h\|_2$ or, all gives

$$\max\left\{\frac{\|\widehat{\Delta A}\|_F}{\|A\|_F}, \frac{\|\widehat{\Delta B}\|_F}{\|B\|_F}, \frac{\|\widehat{\Delta C}\|_F}{\|C\|_F}, \frac{\|\widehat{\Delta D}\|_F}{\|D\|_F}, \frac{\|\widehat{\Delta E}\|_F}{\|E\|_F}, \frac{\|\widehat{\Delta f}\|_2}{\|f\|_2}, \frac{\|\widehat{\Delta g}\|_2}{\|g\|_2}, \frac{\|\widehat{\Delta h}\|_2}{\|h\|_2}\right\} \ge \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z}).$$

Thus, we obtain

$$\boldsymbol{\eta}^{\mathbb{S}_{1}}(\tilde{x},\tilde{y},\tilde{z}) = \left\| \begin{bmatrix} \frac{\|\widehat{\Delta A}\|_{F}}{\|A\|_{F}}, & \frac{\|\widehat{\Delta B}\|_{F}}{\|B\|_{F}}, & \frac{\|\widehat{\Delta C}\|_{F}}{\|C\|_{F}}, & \frac{\|\widehat{\Delta D}\|_{F}}{\|D\|_{F}}, & \frac{\|\widehat{\Delta F}\|_{F}}{\|E\|_{F}}, & \frac{\|\widehat{\Delta f}\|_{2}}{\|f\|_{2}}, & \frac{\|\widehat{\Delta g}\|_{2}}{\|g\|_{2}}, & \frac{\|\widehat{\Delta h}\|_{2}}{\|h\|_{2}} \end{bmatrix} \right\|_{2}$$

$$\geq \max \left\{ \frac{\|\widehat{\Delta A}\|_{F}}{\|A\|_{F}}, \frac{\|\widehat{\Delta B}\|_{F}}{\|B\|_{F}}, \frac{\|\widehat{\Delta C}\|_{F}}{\|C\|_{F}}, \frac{\|\widehat{\Delta D}\|_{F}}{\|D\|_{F}}, \frac{\|\widehat{\Delta E}\|_{F}}{\|E\|_{F}}, \frac{\|\widehat{\Delta f}\|_{2}}{\|f\|_{2}}, \frac{\|\widehat{\Delta g}\|_{2}}{\|g\|_{2}}, \frac{\|\widehat{\Delta h}\|_{2}}{\|g\|_{2}}, \frac{\|\widehat{\Delta h}\|_{2}}{\|h\|_{2}} \right\}$$

$$\geq \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}). \tag{6.5}$$

On the contrary, for the minimal perturbation matrices $\widetilde{\Delta A}, \widetilde{\Delta B}, \widetilde{\Delta C}, \widetilde{\Delta D}, \widetilde{\Delta E}, \widetilde{\Delta f}, \widetilde{\Delta g}$ and $\widetilde{\Delta h}$ for which $\boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ is attained, we get

$$\begin{split} \|\widetilde{\Delta A}\|_{F} &\leq \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) \|A\|_{F}, \|\widetilde{\Delta B}\|_{F} \leq \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) \|B\|_{F}, \|\widetilde{\Delta C}\|_{F} \leq \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) \|C\|_{F}, \\ \|\widetilde{\Delta D}\|_{F} &\leq \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) \|D\|_{F}, \|\widetilde{\Delta E}\|_{F} \leq \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) \|E\|_{F}, \|\widetilde{\Delta f}\|_{2} \leq \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) \|f\|_{2}, \\ \|\widetilde{\Delta g}\|_{2} \leq \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) \|g\|_{2}, \text{ and } \|\widetilde{\Delta h}\|_{2} \leq \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) \|h\|_{2}. \end{split}$$

However, these minimal perturbations do not improve the value of $\eta^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ given in Definition 2.2. Consequently,

$$\boldsymbol{\eta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}) \leq \left\| \begin{bmatrix} \| \underline{\widetilde{\Delta A}} \|_{F} \\ \| A \|_{F} \end{bmatrix}, \frac{\| \underline{\Delta \widetilde{\Delta B}} \|_{F}}{\| B \|_{F}}, \frac{\| \underline{\Delta \widetilde{\Delta C}} \|_{F}}{\| C \|_{F}}, \frac{\| \underline{\widetilde{\Delta D}} \|_{F}}{\| D \|_{F}}, \frac{\| \underline{\widetilde{\Delta F}} \|_{F}}{\| E \|_{F}}, \frac{\| \underline{\widetilde{\Delta f}} \|_{2}}{\| f \|_{2}}, \frac{\| \underline{\widetilde{\Delta g}} \|_{2}}{\| g \|_{2}}, \frac{\| \underline{\widetilde{\Delta h}} \|_{2}}{\| h \|_{2}} \end{bmatrix} \right\|_{2} \leq \sqrt{8} \boldsymbol{\zeta}^{\mathbb{S}_{1}}(\tilde{x}, \tilde{y}, \tilde{z}).$$

$$(6.6)$$

Combining (6.5) and (6.6) yields the bounds of (6.3). Similarly, we can obtain the bounds of (6.4). Hence, the proof is completed. \blacksquare

Theorem 6.1 gives $\boldsymbol{\eta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z}) < \boldsymbol{tol} \Rightarrow \boldsymbol{\zeta}^{\mathcal{S}_i}(\tilde{x}, \tilde{y}, \tilde{z}) < \boldsymbol{tol}$ for $i = 1, 2, 3, \text{ and } \boldsymbol{\zeta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z}) < \boldsymbol{tol} \Rightarrow \boldsymbol{\eta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z}) < \sqrt{8} \boldsymbol{tol}$ and $\boldsymbol{\zeta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z}) < \boldsymbol{tol} \Rightarrow \boldsymbol{\eta}^{\mathbb{S}_i}(\tilde{x}, \tilde{y}, \tilde{z}) < \sqrt{9} \boldsymbol{tol}$, for i = 2, 3.

7 Numerical Experiments

In this section, we carry out several numerical experiments to validate our theoretical findings and the strong backward stability of iterative algorithms for solving the *TBSPP*. Moreover, an illustrative example is included to demonstrate the effectiveness and reliability of the proposed termination criteria when integrated into the GMRES method [26] to solve the *TBSPP*. In Examples 2-6, we consider $\theta_1 = \frac{1}{\|A\|_F}$, $\theta_2 = \frac{1}{\|B_1\|_F}$, $\theta_3 = \frac{1}{\|B_2\|_F}$, $\theta_4 = \frac{1}{\|C\|_F}$, $\theta_5 = \frac{1}{\|D_1\|_F}$, $\theta_7 = \frac{1}{\|D_2\|_2}$, $\theta_8 = \frac{1}{\|f\|_2}$, $\theta_9 = \frac{1}{\|g\|_2}$ and $\theta_{10} = \frac{1}{\|h\|_2}$. All numerical experiments are performed using MATLAB (version R2023b) on an Intel(R) Core(TM) i7-10700 CPU running at 2.90GHz with 16GB of memory. The machine precision is set to 2.2×10^{-16} .

Example 1. We consider the *TBSPP* (1.1) with the block matrices $A \in S_5$, $B \in \mathbb{R}^{3 \times 5}$, $C \in S_3$, $D \in \mathbb{R}^{2 \times 3}$, $E \in S_2$, $f \in \mathbb{R}^5$, $g \in \mathbb{R}^3$ and $h \in \mathbb{R}^2$; that are given by

$$A = \begin{bmatrix} -0.4083 & 0.3472 & 0 & 0.0636 & 0 \\ 0.3472 & -0.8593 & 0.0647 & 0.1433 & 0 \\ 0 & 0.0647 & 0 & 0 & 0.3129 \\ 0.0636 & 0.1433 & 0 & -0.4236 & -1.2123 \\ 0 & 0 & 0.3129 & -1.2123 & 0 \end{bmatrix},$$

$$B = \begin{bmatrix} 0 & 0 & 0 & 0 & 0.0961 \\ -2.2777 & 0 & -0.1180 & 0 & 0 \\ 1.0582 & 0.4363 & 0 & 1.4115 & -0.0146 \end{bmatrix}, C = \begin{bmatrix} 0 & 0 & -0.2299 \\ 0 & -0.7390 & 1.0800 \\ -0.2299 & 1.0800 & 0 \end{bmatrix},$$

$$D = \begin{bmatrix} 0 & 0 & -0.38734 \\ 0 & 0 & -0.31964 \end{bmatrix}, E = \begin{bmatrix} 0 & 0.5387 \\ 0.5387 & 0 \end{bmatrix}, f = \begin{bmatrix} -2.5245 \\ -1.0063 \\ -0.4242 \\ -0.6612 \\ 0.7276 \end{bmatrix}, g = \begin{bmatrix} 0.4566 \\ -0.5062 \\ -1.1846 \end{bmatrix},$$

and $h = \begin{bmatrix} 0.7818 \\ -0.0804 \end{bmatrix}$. We take the approximate solution $\widetilde{\boldsymbol{w}} = [\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$, where

$$\tilde{x} = \begin{bmatrix} -4.4871\\11.3517\\100.3742\\18.4213\\-1.6524 \end{bmatrix}, \quad \tilde{y} = \begin{bmatrix} -86.5918\\5.4127\\2.6903 \end{bmatrix} \text{ and } \tilde{z} = \begin{bmatrix} 1.4512\\3.3886 \end{bmatrix}$$

with residue $\|\mathcal{A}\widetilde{\boldsymbol{w}}-\boldsymbol{d}\|_2 = 3.4180 \times 10^{-02}$. Further, we take $\theta_i = 1$, for all i = 1, 2, ..., 10. We compute the unstructured $BE \ \boldsymbol{\eta}(\widetilde{\boldsymbol{w}})$ using formulae provided in (2.1), structured BE with preserving sparsity $\boldsymbol{\eta}_{sps}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ using Theorem 3.1, and structured BE without preserving sparsity $\boldsymbol{\eta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ using Corollary 3.1. The computed values are given as follows:

 $\boldsymbol{\eta}(\widetilde{\boldsymbol{w}}) = 5.0177 \times 10^{-05}, \ \boldsymbol{\eta}_{\mathtt{sps}}^{\mathbb{S}_1}(\widetilde{x}, \widetilde{y}, \widetilde{z}) = 2.8084 \times 10^{-03} \text{ and } \boldsymbol{\eta}^{\mathbb{S}_1}(\widetilde{x}, \widetilde{y}, \widetilde{z}) = 2.9142 \times 10^{-04}.$

The structure-preserving minimal perturbation matrices that preserve the sparsity pattern as well for the Problem 2.1 are given as follows:

$$\widehat{\Delta A}_{\rm sps} = 10^{-04} \times \begin{bmatrix} -2.1152 & 1.4013 & 0 & 4.4380 & 0 \\ 1.4013 & 6.4477 & -1.6479 & 4.98870 & \\ 0 & -1.6479 & 0 & 0 & 10.0133 \\ 4.4380 & 4.9887 & 0 & -0.7883 & 1.0675 \\ 0 & 0 & 10.0133 & 1.0675 & 0 \end{bmatrix},$$

$$\widehat{\Delta B}_{\rm sps} = 10^{-04} \times \begin{bmatrix} 0 & 0 & 0 & 0 & 10.9193 \\ 1.1455 & 0 & 2.6987 & 0 & 0 \\ 2.9280 & -2.6709 & 0 & -6.9292 & 0.9127 \end{bmatrix},$$

$$\widehat{\Delta C}_{\rm sps} = 10^{-05} \times \begin{bmatrix} 0 & 0 & 7.7234 \\ 0 & -16.9623 & 5.7956 \\ 7.7234 & 5.7956 & 0 \end{bmatrix}, \ \widehat{\Delta D}_{\rm sps} = 10^{-04} \times \begin{bmatrix} 0 & 0 & -2.3102 \\ 0 & 0 & -6.1506 \end{bmatrix},$$

$$\begin{split} \widehat{\Delta E}_{\rm sps} &= 10^{-04} \times \begin{bmatrix} 0 & -2.4376 \\ -2.4376 & 0 \end{bmatrix}, \ \widehat{\Delta f}_{\rm sps} = 10^{-05} \times \begin{bmatrix} -4.7141 \\ -5.6800 \\ 53.1269 \\ 0.4279 \\ -1.1206 \end{bmatrix} \\ \widehat{\Delta g}_{\rm sps} &= 10^{-05} \times \begin{bmatrix} 124.8013 \\ -3.1338 \\ 3.6990 \end{bmatrix} \text{ and } \widehat{\Delta h}_{\rm sps} = 10^{-04} \times \begin{bmatrix} 0.6591 \\ 1.8203 \end{bmatrix}. \end{split}$$

The above perturbation matrices are computed using the formulae presented in Theorem 3.1. Furthermore, the structure-preserving minimal perturbation matrices for which the

structured *BE* $\boldsymbol{\eta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ is attained are given by

$$\begin{split} \widehat{\Delta A} &= 10^{-06} \times \begin{bmatrix} -4.2896 & 5.6947 & 48.7625 & 8.8353 & -2.2268 \\ 5.6947 & -1.3596 & -7.9961 & -1.1793 & 3.7343 \\ 48.7625 & -7.9961 & -35.1043 & -3.8939 & 32.4331 \\ 8.8353 & -1.1793 & -3.8939 & -0.2468 & 5.9104 \\ -2.2268 & 3.7343 & 32.433 & 5.9104 & -1.0584 \end{bmatrix}, \\ \widehat{\Delta B} &= 10^{-06} \times \begin{bmatrix} -81.1245 & 6.1830 & -6.7514 & -5.6364 & -54.8512 \\ -3.4239 & 21.1045 & 190.4498 & 35.2275 & 0.3003 \\ 6.7250 & -10.8292 & -93.8457 & -17.0866 & 3.2526 \end{bmatrix}, \\ \widehat{\Delta C} &= 10^{-06} \times \begin{bmatrix} -31.9502 & 83.9647 & -39.5777 \\ 83.9647 & -10.3722 & -0.0727 \\ -39.5777 & -0.0727 & 2.4901 \end{bmatrix}, \ \widehat{\Delta D} &= 10^{-06} \times \begin{bmatrix} 17.4919 & 1.6540 & -1.9033 \\ 25.4210 & 4.8262 & -3.9650 \end{bmatrix}, \end{split}$$

$$\begin{split} \widehat{\Delta E} &= 10^{-06} \times \begin{bmatrix} -0.3021 & -0.5762 \\ -0.5762 & -1.0437 \end{bmatrix}, \ \widehat{\Delta f} = 10^{-07} \times \begin{bmatrix} -9.5598 \\ 1.1977 \\ 3.4973 \\ 0.1340 \\ -6.4049 \end{bmatrix}, \\ \widehat{\Delta g} &= 10^{-07} \times \begin{bmatrix} 3.6897 \\ -19.1626 \\ 9.2558 \end{bmatrix}, \text{ and } \widehat{\Delta h} = 10^{-07} \times \begin{bmatrix} 2.0819 \\ 3.0801 \end{bmatrix}. \end{split}$$

Example 2. To test the strong backward stability of numerical algorithms, we consider the TBSPP (1.1) with

$$A = GPG(1:3,1:3) \in \mathcal{S}_3, \ B = D = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 10^4 & 0 & 0 \end{bmatrix} \in \mathbb{R}^{3\times3}, \ C = \begin{bmatrix} 1 & -2 & 1 \\ -2 & 6 & 0 \\ 1 & 0 & 0 \end{bmatrix} \in \mathbb{R}^{3\times3},$$
$$E = GPG(4:6,4:6) \in \mathcal{S}_3, \ f = \begin{bmatrix} 10^8 \\ 10 \\ 0 \end{bmatrix} \in \mathbb{R}^3, \text{ and } g = h = \begin{bmatrix} 10^{-8} \\ 0 \\ 0 \end{bmatrix} \in \mathbb{R}^3,$$

where $G = 10^6 \times \text{diag}(1, 5, 10, 50, 100, 500)$ and $P = [p_{ij}] \in \mathbb{R}^{6 \times 6}$, $p_{ij} = \frac{(i+j-1)!}{(i-1)!(j-1)!}$. The approximate solution $\tilde{\boldsymbol{w}} = [\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$ of this *TBSPP* is obtained using Gaussian elimination with partial pivoting (GEP), where

$$\tilde{x} = 10^{-05} \times \begin{bmatrix} 60.0120 \\ -8.0016 \\ 1.0002 \end{bmatrix}, \quad \tilde{y} = \begin{bmatrix} 6.0012 \\ 2.0004 \\ -2.0004 \end{bmatrix}, \text{ and } \tilde{z} = 10^{-13} \times \begin{bmatrix} -1.7109 \\ 0.8556 \\ -0.0475 \end{bmatrix}.$$

We compute the unstructured $BE \ \boldsymbol{\eta}(\tilde{\boldsymbol{w}})$ and structured $BEs \ \boldsymbol{\eta}_{sps}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ and $\boldsymbol{\eta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ using the formulae given in (2.1), Theorem 3.1 and Corollary 3.1, respectively. The obtained BEs are given by

$$\boldsymbol{\eta}(\widetilde{\boldsymbol{w}}) = 6.9314 \times 10^{-27}, \ \boldsymbol{\eta}_{\text{sps}}^{\mathbb{S}_1}(\widetilde{x}, \widetilde{y}, \widetilde{z}) = 5.4649 \times 10^{-06}, \ \boldsymbol{\eta}^{\mathbb{S}_1}(\widetilde{x}, \widetilde{y}, \widetilde{z}) = 4.7907 \times 10^{-06}. \ (7.1)$$

From (7.1), we can observe that $\eta(\tilde{w})$ of $\mathcal{O}(10^{-27})$ indicated that GEP is backward stable for solving this *TBSPP*. On the other side $\eta_{\text{sps}}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ and $\eta^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ is much smaller than $\eta(\tilde{w})$ implies that GEP for solving this *TBSPP* is not strongly backward stable. This shows that a backward stable iterative algorithm for solving the *TBSPP* may not be strongly backward stable.

Example 3. In this example, we perform a comparison among our obtained structured BEs and the structured BE considered in [21]. For this, we consider the TBSPP (1.1) with

$$A = \begin{bmatrix} 0.0968 & 0 & -0.2438 & -0.2823 \\ 0 & 0 & 1.1180 & -1.1611 \\ -0.2438 & 1.1180 & 1.6014 & -0.8693 \\ -0.2823 & -1.1611 & -0.8693 & -0.4914 \end{bmatrix}, B1 = \begin{bmatrix} 0 & 0 & 0.7090 & 0 \\ 1.9046 & 0.0928 & -0.0430 & 0.0508 \end{bmatrix}, B2 = \begin{bmatrix} -0.2592 & 0 & 0.2543 & 0.1248 \\ 0.0876 & 1.1375 & 0 & 0.0766 \end{bmatrix}, C = \mathbf{0}_{2\times 2}, D = \begin{bmatrix} 0 & 1.8070 \\ 1.0365 & -1.5516 \end{bmatrix} \text{ and } E = \mathbf{0}_{2\times 2}.$$

Here, n = 4, m = 2 and p = 2. Further, we consider the right-hand side vector $\boldsymbol{d} = [f^T, g^T, h^T]^T \in \mathbb{R}^8$, where

$$f = \begin{bmatrix} -1.1251 \\ -1.9000 \\ -0.4320 \\ -1.1422 \end{bmatrix}, g = \begin{bmatrix} -0.5516 \\ 1.8738 \end{bmatrix} \text{ and } h = \begin{bmatrix} 0.4982 \\ 0.8347 \end{bmatrix}.$$

The computed solution using the the MATLAB 'blackshash' command is $\tilde{\boldsymbol{w}} = [\tilde{x}^T, \tilde{y}^T, \tilde{z}^T]^T$, where

$$\tilde{x} = \begin{bmatrix} -1.6927\\ -1.5778\\ 1.9746\\ 3.5598 \end{bmatrix}, g = \begin{bmatrix} 1.2180\\ 0.2757 \end{bmatrix} \text{ and } h = \begin{bmatrix} 0.3571\\ -1.8683 \end{bmatrix}.$$

The computed solution $\tilde{\boldsymbol{w}}$ has residue $\|\mathcal{A}\tilde{\boldsymbol{w}} - \boldsymbol{d}\| = 1.4864 \times 10^{-15}$. The unstructured *BE* computed using the formula (2.1) is 5.2700×10^{-17} , structured *BE* using the Theorem 3.2 of [21] is 4.1137×10^{-16} , structured *BE* with sparsity using Theorem 5.1 is 2.7992×10^{-16} and the structured *BE* without sparsity using Corollary 5.1 is 2.5525×10^{-16} . We observe that all the computed *BEs* are in unit round-off error and the structured *BEs* are one order larger than the unstructured ones. Furthermore, the structured *BEs* derived in our work and those obtained in the reference [21] exhibit uniform order. This shows the reliability of our derived structured *BEs* formulae. One notable advantage of our derived formulae lies in our ability to preserve the sparsity pattern within the perturbation matrices.

Example 4. To test the strong backward stability of the GMRES method, in this example, we consider the TBSPP (1.1) [15] with the block matrices

$$A = \begin{bmatrix} I \otimes Z + Z \otimes I & \mathbf{0} \\ \mathbf{0} & I \otimes Z + Z \otimes I \end{bmatrix} \in \mathbb{R}^{2r^2 \times 2r^2}, \ B = \begin{bmatrix} I \otimes H & H \otimes I \end{bmatrix} \in \mathbb{R}^{r^2 \times 2r^2}, \ (7.2)$$

 $D = G \otimes H \in \mathbb{R}^{r^2 \times r^2}$ and $C = E = \mathbf{0}_{r^2 \times r^2}$, where $Z = \frac{1}{(r+1)^2} \operatorname{tridiag}(-1, 2, -1) \in \mathbb{R}^{r \times r}$, $H = \frac{1}{r+1} \operatorname{tridiag}(0, 1, -1) \in \mathbb{R}^{r \times r}$ and $G = \operatorname{diag}(1, r+1, \ldots, r^2 - r + 1) \in \mathbb{R}^{r \times r}$. Here, $G \otimes H$ represents the Kronecker product of matrices G and H, and $\operatorname{tridiag}(d_1, d_2, d_3) \in \mathbb{R}^{r \times r}$ represents the $r \times r$ tridiagonal matrix with the subdiagonal entry d_1 , diagonal entry d_2 , and superdiagonal entry d_3 . For this problem, the dimension of \mathcal{A} is $4r^2$. We use GMRES method to solve this TBSPP with termination criteria $\frac{\|\mathcal{A}\boldsymbol{w}_k - \boldsymbol{d}\|_2}{\|\boldsymbol{d}\|_2} < \boldsymbol{tol}$, where \boldsymbol{w}_k is solution at each iterate and $\boldsymbol{tol} = 10^{-13}$ and the initial guess vector zero. We compute the structured and unstructured BEs for the solution at the final iteration. The computed BEs for different values of r are listed in Table 1.

Table 1: Values of structured and unstructured BEs of the approximate solution obtained using GMRES for Example 4.

r	$rac{\ \mathcal{A}oldsymbol{w}_k - oldsymbol{d}\ _2}{\ oldsymbol{d}\ _2}$	$oldsymbol{\eta}(ilde{x}, ilde{y}, ilde{z})$	$\boldsymbol{\eta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$	$oldsymbol{\eta}_{ extsf{sps}}^{\mathbb{S}_1}(ilde{x}, ilde{y}, ilde{z})$
4	1.0593e-15	4.1823e-17	1.3757e-16	4.9831e-16
6	2.4960e-14	5.3825e-16	1.8436e-15	7.3871e-15
8	2.0868e-14	3.0086e-16	9.6476e-16	5.4875e-15
10	3.2981e-14	3.4862e-16	1.3781e-15	9.1775e-15

From Table 1, we observe that unstructured $BE \ \boldsymbol{\eta}(\tilde{x}, \tilde{y}, \tilde{z})$, structured BE with preserving sparsity $\boldsymbol{\eta}_{sps}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ and structured $BE \ \boldsymbol{\eta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ are all around order of unit round-off error. Using our obtained structured BEs, we successfully demonstrate that the GMRES method for solving this TBSPP exhibits strong backward stability.

Example 5. To test the strong backward stability of GMRES method to solve the TBSPP for case (*ii*), in this example, we consider the data matrices for (1.1) as follows:

$$A = randn(n, n), \ B = sprandn(m, n, 0.5), \ C = 0.5(C_1 + C_1^T),$$

$$D1 = sprandn(p, m, 0.5), \ D2 = sprandn(p, m, 0.5), \ E = 0.5(E_1 + E_1^T),$$

where $C_1 = sprandn(m, m, 0.2)$, $E_1 = sprandn(p, p, 0.3)$, n = 2k, m = k, and p = k. Furthermore, we take $\boldsymbol{d} = [f^T, g^T, h^T]^T$ such that the actual solution of the *TBSPP* is $[1, 1, \ldots, 1]^T \in \mathbb{R}^{n+m+p}$. Here, the symbols 'randn(n, m)' and 'sprandn(n, m, ω)' stand for normally distributed random matrix and the sparse random matrix with density ω , respectively, of size $n \times m$.



Figure 1: Unstructured and structured BEs versus k

We employ the GMRES method [26] with an initial guess vector set to zero, termination criteria $\frac{\|\mathcal{A}\boldsymbol{w}_k-\boldsymbol{d}\|_2}{\|\boldsymbol{d}\|_2} < \boldsymbol{tol}$, where \boldsymbol{w}_k is solution at each iterate and the tolerance is $\boldsymbol{tol} = 10^{-13}$. We take values of k from 10 to 70 with step size 10. For these values of k, we plot the unstructured $BE \boldsymbol{\eta}(\tilde{\boldsymbol{w}})$ using the formula (2.1) (abbreviated as 'unstructured \boldsymbol{BE} '), structured BE $\boldsymbol{\eta}_{sps}^{\mathbb{S}_3}(\tilde{x}, \tilde{y}, \tilde{z})$ (abbreviated as 'structured \boldsymbol{BE} with sparsity') using Theorem 5.1, and $\boldsymbol{\eta}^{\mathbb{S}_3}(\tilde{x}, \tilde{y}, \tilde{z})$ (abbreviated as 'structured \boldsymbol{BE} ') using Corollary 5.1. From Figure 1, it is seen that, for all values of k, the unstructured $BE \boldsymbol{\eta}(\tilde{\boldsymbol{w}})$ are of order $\mathcal{O}(10^{-16})$ and the structured $BEs \boldsymbol{\eta}_{sps}^{\mathbb{S}_3}(\tilde{x}, \tilde{y}, z)$, and $\boldsymbol{\eta}_{sps}^{\mathbb{S}_3}(\tilde{x}, \tilde{y}, z)$ are of $\mathcal{O}(10^{-15})$, which are very small. Notably, both the structured BEs are only one order less than the unstructured BE. Thus, the utilization of our structured BE affirms that the GMRES method for solving the TBSPP is strongly backward stable. Furthermore, the resultant approximate solution corresponds to an actual solution of a nearly perturbed TBSPP of the form (1.1), which preserves the inherent matrix structure and sparsity pattern.

Example 6. In this example, we incorporate our proposed termination criteria into the direct Uzawa method [16] to solve the TBSPP (1.1) with

$$A = \text{diag}(2XX^{T} + D_{1}, D_{2}, D_{3}) \in \mathbb{R}^{n \times n}, \ B = \begin{bmatrix} G & -I_{2\tilde{r}} & I_{2\tilde{r}} \end{bmatrix} \in \mathbb{R}^{m \times n}, \ C = \mathbf{0}_{m \times m}$$
$$D = H, \ E = \mathbf{0}_{p \times p}, \text{ where } X = [x_{ij}] \in \mathbb{R}^{\hat{r} \times \hat{r}} \text{ with } x_{ij} = e^{-2((i/3)^{2} + (j/3)^{2})}, \ D_{1} = I_{\hat{r}},$$

and $D_k = \text{diag}(d_j^{(k)}) \in \mathbb{R}^{2r \times 2r}, \ k = 2, 3$, are diagonal matrices with

$$d_j^{(2)} = \begin{cases} 1, & \text{for } 1 \le j \le \tilde{r}, \\ 10^{-5}(j-\tilde{r})^2, & \text{for } \tilde{r}+1 \le j \le 2\tilde{r}, \end{cases}$$

 $d_j^{(3)} = 10^{-5}(j+\tilde{r})^2$, for $1 \le j \le 2\tilde{r}$, where $\hat{r} = r^2$ and $\tilde{r} = r(r+1)$. Further, $G = \begin{bmatrix} \widehat{G} \otimes I_r \\ I_r \otimes \widehat{G} \end{bmatrix} \in I_r \otimes \widehat{G}$



Figure 2: Characteristics curves of termination criteria $\eta^{\texttt{term1}}(\mathcal{A}, \widetilde{w}), \eta^{\texttt{term2}}(\widetilde{w})$ and $\eta^{\mathbb{S}_2}(\tilde{x}, \tilde{y}, \tilde{z})$ versus iteration numbers.

$$\begin{split} \mathbb{R}^{\hat{r} \times \tilde{r}}, \, \hat{G} &= tridiag(0, 2, -1) \in \mathbb{R}^{r \times (r+1)}, \, H = \left[\hat{H} \otimes I_r \quad I_r \otimes \hat{H} \right] \in \mathbb{R}^{\hat{r} \times \hat{r}}, \\ \hat{H} &= \begin{bmatrix} r+1 & -\frac{r-1}{r} & \frac{r-2}{r} & \dots & \frac{(-1)^{r-1}}{r} \\ -\frac{r-1}{r} & 2r+1 & \ddots & \ddots & \vdots \\ \frac{r-2}{r} & \ddots & \ddots & \ddots & \frac{r-2}{r} \\ \vdots & \ddots & \ddots & \ddots & \frac{r-2}{r} \\ \frac{(-1)^{r-1}}{r} & \vdots & \frac{r-2}{r} & -\frac{r-1}{r} & r^2 + 1 \\ 0 & 0 & \dots & 0 & 1 \end{bmatrix} \in \mathbb{R}^{(r+1) \times r}. \end{split}$$

Right hand side vector \boldsymbol{d} is chosen as so that the actual solution of the *TBSPP* is $\boldsymbol{w} = [1, 1, ..., 1]^T \in \mathbb{R}^{n+m+l}$. We choose r = 6 with the tolerance $\boldsymbol{tol} = 10^{-14}$ and use the termination criteria $\boldsymbol{\eta}^{\texttt{term1}}(\mathcal{A}, \tilde{\boldsymbol{w}})$ (abbreviated as 'term1'), $\boldsymbol{\eta}^{\texttt{term2}}(\tilde{\boldsymbol{w}})$ (abbreviated as 'term2'), and $\boldsymbol{\eta}^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ (abbreviated as 'term3'). The characteristic curves of these three termination criteria at each iteration step are displayed in Figure 2.

From Figure 2, we observe that, our proposed termination criterion $\eta^{\mathbb{S}_1}(\tilde{x}, \tilde{y}, \tilde{z})$ converges in fewer iteration numbers compare to the existing $\eta^{\texttt{term2}}(\tilde{w})$, whereas the termination criterion $\eta^{\texttt{term1}}(\mathcal{A}, \tilde{w})$ has competitive performance. Therefore, these observations show the superiority and reliability of our proposed termination criterion compared to the existing ones.

8 Conclusions

This paper investigates the structured BEs for a class of TBSPP by preserving the inherent matrix structure and sparsity pattern in the perturbation matrices. We derive explicit formulae for the structured BEs (in three cases) and present concrete formulae for

structure-preserving minimal perturbation matrices. These perturbation matrices yield a nearly perturbed TBSPP. Thereby, the approximate solution becomes its actual solution. Moreover, using the obtained structured BEs, we propose three effective and reliable termination criteria for iterative algorithms to find a more accurate solution of the TBSPP. Several numerical experiments are performed to validate our obtained theory and to test the strong backward stability of iterative algorithms. Our observations reveal that iterative algorithms (such as GEP) demonstrating backward stability do not always exhibit strong backward stability. Finally, we integrate our proposed termination criteria into iterative algorithms, and the corresponding numerical experiments show that the obtained solution satisfies a nearly perturbed structured-preserving TBSPP. Thus, the theoretical investigations and empirical findings affirm the superiority of the proposed termination criteria compared to the baseline ones.

Acknowledgments

Pinki Khatun expresses her gratitude to the Council of Scientific & Industrial Research (CSIR) in New Delhi, India, for providing financial support awarded as a fellowship (File no. 09/1022(0098)/2020-EMR-I).

References

- S. S. Ahmad and P. Kanhya. Structured perturbation analysis of sparse matrix pencils with s-specified eigenpairs. *Linear Algebra Appl.*, 602:93–119, 2020.
- [2] S. S. Ahmad and P. Kanhya. Backward error analysis and inverse eigenvalue problems for Hankel and Symmetric-Toeplitz structures. *Appl. Math. Comput.*, 406:126288, 15, 2021.
- [3] Z.-Z. Bai and J.-Y. Pan. Matrix Analysis and Computations. SIAM, Philadelphia, PA, 2021.
- [4] S. Bradley and C. Greif. Eigenvalue bounds for double saddle-point systems. IMA J. Numer. Anal., 43(6):3564–3592, 2023.
- [5] J. R. Bunch. The weak and strong stability of algorithms in numerical linear algebra. *Linear Algebra Appl.*, 88/89:49–66, 1987.
- [6] J. R. Bunch, J. W. Demmel, and C. F. Van Loan. The strong stability of algorithms for solving symmetric linear systems. *SIAM J. Matrix Anal. Appl.*, 10(4):494–499, 1989.
- [7] Y. Cao, L.-Q. Yao, and M.-Q. Jiang. A modified dimensional split preconditioner for generalized saddle point problems. J. Comput. Appl. Math., 250:70–82, 2013.

- [8] X.-W. Chang, C. C. Paige, and D. Titley-Peloquin. Stopping criteria for the iterative solution of linear least squares problems. SIAM J. Matrix Anal. Appl., 31(2):831–852, 2009.
- [9] X. S. Chen, W. Li, X. Chen, and J. Liu. Structured backward errors for generalized saddle point systems. *Linear Algebra Appl.*, 436(9):3109–3119, 2012.
- [10] Z. Chen, Q. Du, and J. Zou. Finite element methods with matching and nonmatching meshes for Maxwell equations with discontinuous coefficients. SIAM J. Numer. Anal., 37(5):1542–1570, 2000.
- [11] G. H. Golub and C. F. Van Loan. *Matrix Computations*. Johns Hopkins University Press, Baltimore, fourth edition, 2013.
- [12] C. Greif and Y. He. Block preconditioners for the marker-and-cell discretization of the Stokes-Darcy equations. SIAM J. Matrix Anal. Appl., 44(4):1540–1565, 2023.
- [13] N. J. Higham. Accuracy and Stability of Numerical Algorithms. SIAM, Philadelphia, PA, second edition, 2002.
- [14] K. Hu and J. Xu. Structure-preserving finite element methods for stationary MHD models. Math. Comp., 88(316):553–581, 2019.
- [15] N. Huang and C.-F. Ma. Spectral analysis of the preconditioned system for the 3×3 block saddle point problem. *Numer. Algor.*, 81(2):421-444, 2019.
- [16] N. Huang, Y.-H. Dai, and Q. Hu. Uzawa methods for a class of block three-by-three saddle-point problems. Numer. Linear Algebra Appl., 26(6):e2265, 26, 2019.
- [17] X. Li and X. Liu. Structured backward errors for structured KKT systems. J. Comput. Math., 22(4):605–610, 2004.
- [18] P. Lv. Structured backward errors analysis for generalized saddle point problems arising from the incompressible Navier-Stokes equations. AIMS Math., 8(12):30501– 30510, 2023.
- [19] P. Lv and B. Zheng. Structured backward error analysis for a class of block three-bythree saddle point problems. *Numer. Algorithms*, 90(1):59–78, 2022.
- [20] W. Ma. On normwise structured backward errors for the generalized saddle point systems. *Calcolo*, 54(2):503–514, 2017.
- [21] W. Ma, Y. Fan, and X. Xu. Structured backward errors for block three-by-three saddle point systems. *Linear and Multilinear Algebra*, pages 1–21, 2024.
- [22] L. Meng, Y. He, and S.-X. Miao. Structured backward errors for two kinds of generalized saddle point systems. *Linear Multilinear Algebra*, 70(7):1345–1355, 2022.

- [23] A. Ramage and E. C. Gartland, Jr. A preconditioned nullspace method for liquid crystal director modeling. SIAM J. Sci. Comput., 35(1):B226–B247, 2013.
- [24] T. Rees, H. S. Dollar, and A. J. Wathen. Optimal solvers for PDE-constrained optimization. SIAM J. Sci. Comput., 32(1):271–298, 2010.
- [25] J. L. Rigal and J. Gaches. On the compatibility of a given solution with the data of a linear system. J. Assoc. Comput. Mach., 14:543–548, 1967.
- [26] Y. Saad and M. H. Schultz. GMRES: a generalized minimal residual algorithm for solving nonsymmetric linear systems. SIAM J. Sci. Statist. Comput., 7(3):856–869, 1986.
- [27] U. Wilbrandt. Stokes-Darcy Equations: Analytic and Numerical Analysis. Springer, 2019.
- [28] H. Xiang and Y. Wei. On normwise structured backward errors for saddle point systems. SIAM J. Matrix Anal. Appl., 29(3):838–849, 2007.
- [29] A.-L. Yang, J.-L. Zhu, and W. Yu-Jiang. Multi-parameter dimensional split preconditioner for three-by-three block system of linear equations. *Numer. Algorithms*, 95 (2):721–745, 2024.
- [30] B. Zheng and P. Lv. Structured backward error analysis for generalized saddle point problems. Adv. Comput. Math., 46(2):34–27, 2020.