GENERALIZED CARLOS SCALES

ANDREW V. SILLS

Department of Mathematical Sciences Georgia Southern University Statesboro, Georgia, 30460, USA

Abstract. In 1986, composer Wendy Carlos introduced three unusual musical scales she called alpha, beta and gamma, equal temperament-inspired scales that de-emphasize the octave as the primary interval in favor of the major and minor thirds and the perfect fifth. A derivation of the alpha, beta, and gamma scales due to David Benson is generalized to produce many Carlos-type scales.

1. Introduction: equal temperament

For any musical scale where the spaces between available notes are equal, there is a positive real number x_0 , so that multiplying the frequency of a given note by 2^{x_0} results in the frequency of the next higher available note. In such a scale, we will refer to the interval between adjacent pitches as a $unit¹$ $unit¹$ $unit¹$. For example, in twelve tone equal temperament (12TET), $x_0 = 1/12$ and the unit is the equal-tempered semitone. Once one has decided that the octave must be a perfect $2:1$ ratio and that there will be 12 units per octave, then $x_0 = 1/12$ is the unique solution. With this solution, the 12TET perfect fifth has a ratio of $2^{7/12}$, which approximates the just perfect fifth of 3 : 2 quite well, falling just 1.96 cents flat. The minor third, at $2^{3/12}$, is a less satisfactory approximation of the just ratio 6 : 5, falling 15.64 cents flat. The major third, at $2^{4/12}$, is similarly not such a great approximation to the just ratio of 5 : 4, as it is 13.69 cents sharp.

A well trained human ear is perhaps sensitive to deviations in pitch at the 5 cent level. Given the centrality of the major and minor triad in Western music, the 12TET solution may seem unacceptable to those of us with a fine-tuned sense of pitch.

2. Deriving the original Carlos scales and some new Carlos-type scales

In the 1980s, Wendy Carlos effectively asked the following question: what if we design an equal unit scale giving priority to the major and minor third and the perfect fifth, so the major and minor triad built on the tonic approximates just intonation as closely as possible, and let the octave fall where it may? In [\[4\]](#page-4-0), Carlos reveals that she "wrote a computer program to perform a precise deep-search investigation" that produced three new interesting scales that she named alpha, beta, and gamma. These new musical scales were introduced to the public in Carlos' 1986 album Beauty in the Beast [\[2\]](#page-4-1). The methodology of the computer

E-mail address: asills@georgiasouthern.edu.

Date: August 29, 2024.

Key words and phrases. nonstandard scale; Wendy Carlos; Carlos scales; alpha scale; beta scale; gamma scale.

¹The author prefers the term "unit" to "step" since the latter could be confused with the whole step/half step terminology common in Western music theory. Our "unit" is a generalization of the semi-tone or "half-step".

search that Carlos used to derive the scales is not described in [\[4\]](#page-4-0), but guided by a method given by Benson [\[1,](#page-4-2) pp. 231–233], we can not only re-derive the Carlos alpha, beta, and gamma scales, but also many other scales guided by the same principle.

The goal of a "Carlos-type" scale is to choose positive integers a and b, with $a < b$, so that a minor third is a units above the tonic, the major third is b units above the tonic, and the perfect fifth is $a + b$ units above the tonic. Thus we want to find a unit size x that simultaneously approximates

(1)
$$
(2^x)^a \approx 6/5
$$
, $(2^x)^b \approx 5/4$, and $(2^x)^{a+b} \approx 3/2$.

The approximations given in line [\(1\)](#page-1-0) are equivalent to

(2)
$$
\frac{2^{ax}}{6/5} \approx 1, \frac{2^{bx}}{4/3} \approx 1, \frac{2^{(a+b)x}}{3/2} \approx 1,
$$

which, by taking the base-2 logarithm of both sides of each approximation, are equivalent to

(3)
$$
ax - \log_2(6/5) \approx 0
$$
, $bx - \log_2(5/4) \approx 0$, $(a+b)x - \log_2(3/2) \approx 0$.

To find the "best" x that satisfies all three of the approximations in [\(3\)](#page-1-1) simultaneously, we seek the absolute minimum of the function

(4)
$$
g(x) = (ax - \log_2(6/5))^2 + (bx - \log_2(5/4))^2 + ((a+b)x - \log_2(3/2))^2
$$

over the positive real numbers. Notice that $q(x)$ is the sum of the squared differences of the three approximations we seek to minimize; i.e., we will employ the principle of least squares to find our optimal x .

Since $q(x)$ is a quadratic function with a positive leading coëfficient, it attains its absolute minimum at $x = x_0$, where its derivative $g'(x_0) = 0$. Using elementary calculus, the x-value that minimizes $g(x)$ can be shown to be

(5)
$$
x_0^{(a,b)} = \frac{1}{a^2 + b^2 + (a+b)^2} \log_2 \left[\left(\frac{6}{5} \right)^a \left(\frac{5}{4} \right)^b \left(\frac{3}{2} \right)^{a+b} \right].
$$

Let us refer to the scale with unit size given by $x_0^{(a,b)}$ $\int_0^{(a, b)}$ in [\(11\)](#page-3-0) as the (a, b) -Carlos scale. To express the unit size in cents, just multiply $x_0^{(a,b)}$ by 1200. The ratio of the frequency of one note to the next available lower note is $2^{x_0^{(a,b)}}$.

Next, one needs to experiment with various values of a and b , and check to see if all the approximations are within an acceptable tolerance, say, 5 cents. The author programmed Mathematica to perform such a search. Recall that the just minor third, major third, and perfect fifth are 315.64, 381.61, and 701.96 cents, respectively. Thus, if

(6)
$$
\max\left\{|1200ax_0^{(a,b)} - 315.64|, |1200bx_0^{(a,b)} - 381.61|, |1200(a+b)x_0^{(a,b)} - 701.96|\right\} < 5,
$$

then we have identified an (a, b) -Carlos scale of interest.

The minimal pair (a, b) that satisfies [\(6\)](#page-1-2) is (4, 5). The (4, 5)-Carlos scale is the Carlos alpha scale, with

$$
x_0^{(4,5)} = \frac{1}{122} \log_2 \left((6/5)^4 (5/4)^5 (3/2)^9 \right) = \log_2 \left(\sqrt[122]{\frac{7971615}{32768}} \right),
$$

and which is equivalent to a unit size of about 77.965 cents. As is well known, an acceptably approximated octave does not exist in the alpha scale: it takes 15.3915 units to get a perfect octave, and 15 units falls about 30.5252 cents flat.

The (5, 6)-Carlos scale is the Carlos beta scale, and the (9, 11)-Carlos scale is the gamma scale.

Another "nearby" scale that was not described by Carlos herself is the (8, 10)-Carlos scale. Likely this is the case because it is simply the alpha scale with each unit cut in half (or equivalently, the number of units is doubled). Here the unit is about 38.9825 cents, half that of the alpha scale. Of course, the thirds and the fifth are identical to that of the alpha scale; but this "doubled alpha" scale has one advantage over the alpha scale: an octave requires about 30.783 units. Thus the octave approximated by the 31st note is about 8.45734 cents sharp, which is much better than the 15th note of the alpha scale. Nonetheless, if $gcd(a, b) > 1$, that (a, b) -Carlos scale could reasonably be considered a trivial variation of the $(a/gcd(a, b), b/gcd(a, b))$ -Carlos scale, and therefore not of genuine interest.^{[2](#page-2-0)}

Carlos [\[4\]](#page-4-0) only considered potential scales with between 10 and 40 units in her computer search. Of course, 40 is an arbitrary cutoff; Harry Partch, for example, created a well known 43-unit scale [\[6\]](#page-4-3). The 53TET scale is also of particular importance; see, e.g. [\[1,](#page-4-2) p. 224 ff.]. So let us consider possibilities beyond the 40-unit per octave limit.

Of particular interest is the (14, 17)-Carlos scale. Not only are the thirds and the fifth very close to just intonation, but serendipitously 53 units gives a note just 0.267329 cents sharp of a perfect octave! Thus the (14, 17)-Carlos scale is a very close approximation to 53TET scale, but with slightly sweeter thirds and fifths, and an ever-so-slightly flat octave.

Similarly of particular note for an excellent octave in addition to the thirds and the fifth is the (17, 21)-Carlos scale, as 65 units is 0.520 cents sharp of an octave. The (19, 23)-Carlos scale in addition to excellent thirds and fifths also has a very good octave: 72 units is 3.56 cents sharp of a perfect octave. Please see the table below for additional examples of (a, b) -Carlos scales of interest, all with less than 81 units per octave.

For further discussion of the alpha, beta, and gamma scales, along with further thoughts on tuning and temperament by Carlos, see [\[3\]](#page-4-4).

3. Further Generalizations with four or more parameters or with alternative intervals optimized

3.1. Three or more independent parameters. Rather than requiring the perfect fifth to be exactly $a + b$ units, we could instead allow it to be c units where $a < b < c$. Then the function to minimize is

(7)
$$
g(x) = (ax - \log_2(6/5))^2 + (bx - \log_2(5/4))^2 + (cx - \log_2(3/2))^2,
$$

²As usual, we denote the greatest common divisor of a and b by $gcd(a, b)$.

and its minimum value is

(8)
$$
x_0^{(a,b,c)} = \frac{1}{a^2 + b^2 + c^2} \log_2 \left[\left(\frac{6}{5} \right)^a \left(\frac{5}{4} \right)^b \left(\frac{3}{2} \right)^c \right].
$$

We will name a scale built from this unit $x_0^{(a,b,c)}$ $a_0^{(a,b,c)}$ an (a, b, c) -Carlos scale. If it so happens that $c = a + b$, then this is just the (a, b) -Carlos scale above. But notice that here no attempt is made to control the size of the minor seventh as $a + c$ units nor the major seventh as $b + c$ units.

An example of such a scale can be found in the Wikipedia article on the "Delta scale" [\[5\]](#page-4-5): what is described there is the (23, 28, 50)-Carlos scale in our nomenclature. Notice that $23 + 28 \neq 50$, and therefore is not the same as the (23, 28)-Carlos scale defined above.

3.2. Alternative intervals optimized. Carlos decided to optimize the intonation of the major and minor triads to the possible detriment of the octave, thus giving equal unit scales with very near just minor and minor thirds and perfect fifths. But one could alternatively choose to optimize other intervals. For example, if we allocate a units for the perfect fourth and b units for the perfect fifth, and therefore $a + b$ units for the perfect octave, the analog of Eq. [\(4\)](#page-1-3) becomes

(9)
$$
g(x) = (ax - \log_2(4/3))^2 + (bx - \log_2(3/2))^2 + ((a+b)x - 1)^2.
$$

The unique minimizer of [\(9\)](#page-3-1) is

(10)
$$
x_{\{P4, P5\}}^{(a,b)} = \frac{1}{a^2 + b^2 + (a+b)^2} \log_2 \left[\left(\frac{4}{3} \right)^a \left(\frac{3}{2} \right)^b 2^{a+b} \right]
$$

$$
= \frac{3a + (b-a) \log_2 3}{a^2 + b^2 + (a+b)^2},
$$

where we introduce the notation $x_{\{I_{\alpha}\}}^{(a,b)}$ ${I_{I_a,I_b}}$ to denote the unit size of the Carlos-type scale where the just interval I_k is approximated by k units, for $k = a$ or b, and the interval $I_a \oplus I_b$ is approximated by $a + b$ units. Addition of intervals is defined in the intuitive way, e.g. $m3 \oplus M3 = P5$, $P4 \oplus P5 = P8$, $M2 \oplus m3 = P4$, etc. In this way, the previously introduced $x_0^{(a,b)}$ $u_0^{(a,b)}$ is the same as $x_{\text{m3}}^{(a,b)}$ ${^{(a,b)}_{m3,M3}}$ in this more general notation.

Note that unit size $x_{\{P4, P5\}}^{(5,7)} = (15 + 2 \log_2 3)/218 \approx 0.0833483 \approx 0.083333333 \cdots = 1/12$ gives a very close approximation to 12TET.

Remark 1. One could argue, however, that the preceding example is an abuse of Carlos' original idea, as it could be considered wasteful to optimize for both the perfect fourth and fifth, as each is the inversion of the other. Indeed, the most simple solution $(a, b) = (5, 7)$ does not really get us anything new, but rather a slightly out of tune version of 12TET(!)

Letting J_I denote the just ratio for interval I, so that, e.g. $J_{\rm P5} = 3/2$, $J_{\rm M3} = 5/4$, etc., we can produce scales with optimized approximations for intervals I_a , I_b , and $I_a \oplus I_b$ using unit

(11)
$$
x_{\{I_a,I_b\}}^{(a,b)} = \frac{1}{a^2 + b^2 + (a+b)^2} \log_2 \left[(J_{I_a})^a (J_{I_b})^b (J_{I_a \oplus I_b})^{a+b} \right].
$$

To be careful, we may choose to exclude possibilities where intervals I_a and I_b are inversions of each other.

3.3. Example: a Carlos-type pentatonic optimized scale. We will provide one more example to demonstrate the flexibility of the proposed optimization scheme even with only two independent parameters $a < b$. Consider a pentatonic scale given by the consecutive intervals M2, M2, m3, M2. (Relative to the tonic, the intervals are M2, M3, P5, M6.) We wish to set up a Carlos-type scale with unit size determined so that the intervals in the pentatonic scale (rather than the major and minor triads) are as close to just as possible. Letting the major second be a units apart and the minor third b units apart, we would then want the major third to be 2a units, the perfect fifth to be $2a + b$ units. (In light of the earlier remark, we exclude the condition that the major sixth ought to be approximated by $3a + b$ units, as the major sixth is the inversion of the already included minor third.) Thus, following from the discussion earlier, we want find the unique minimizer of the function (12)

$$
g(x) = (ax - \log_2(9/8))^2 + (2ax - \log_2(5/4))^2 + (bx - \log_2(6/5))^2 + ((2a + b)x - \log_2(3/2))^2
$$

over the positive real numbers which is unit size

over the positive real numbers, which is unit size

(13)
$$
\frac{1}{a^2 + (2a)^2 + b^2 + (2a + b)^2} \log_2 \left[\left(\frac{9}{8} \right)^a \left(\frac{5}{4} \right)^{2a} \left(\frac{6}{5} \right)^b \left(\frac{3}{2} \right)^{2a+b} \right] = \frac{1}{9a^2 + 4ab + 2b^2} \log_2 \left(\frac{5^{2a - b} 9^{2a + b}}{512^a} \right).
$$

For $(a, b) = (7, 11)$, we obtain a scale where each of the major second, minor third, major third, and perfect fifth differs from its just counterpart by less than 7.25 cents. The 43-unit octave is 8.71617 cents sharp.

Another notable Carlos-type pentatonic-optimized scale occurs at $(a, b) = (17, 27)$, where each of the major second, minor third, major third, and perfect fifth differs from its just counterpart by less than 8 cents, and the 104-unit octave is only 1.3501 cents flat.

4. Conclusion

From the examples above, interested readers should be able to construct many additional variants that will fall under the general umbrella of Carlos-type scales.

ACKNOWLEDGMENTS

The author wishes to thank Robert Schneider for his assistance and encouragement, and for valuable suggestions. The author also thanks the Mathematics and Music Lab at Michigan Technological University for invaluable consultation work.

DISCLOSURE STATEMENT

The author reports that there are no competing interests to declare.

REFERENCES

- [1] D. Benson, Music: a Mathematical Offering, Cambridge University Press, 2006.
- [2] W. Carlos, Beauty in the Beast, studio album, Audion Records (1986). Reissued, East Side Digital ESD 81552 (2000).
- [3] W. Carlos, Tuning: at the crossroads, Computer Music J. 11 (1987) 29–43.
- [4] W. Carlos, Three asymmetric divisions of the octave, Pitch, Spring 1990, 2–3. https://www.wendycarlos.com/resources/pitch.html
- [5] Delta scale, Wikipedia, the Free Encyclopedia, Wikimedia Foundation, last edited 9 February 2022, https://en.wikipedia.org/wiki/Delta_scale .
- [6] H. Partch, Genesis of a Music, 2nd ed., Da Capo Press, 1974.