

BOUNDARY REGULARITY FOR THE DISTANCE FUNCTIONS, AND THE EIKONAL EQUATION

NIKOLAI NIKOLOV AND PASCAL J. THOMAS

ABSTRACT. We study the gain in regularity of the distance to the boundary of a domain in \mathbb{R}^m . In particular, we show that if the signed distance function happens to be merely differentiable in a neighborhood of a boundary point, it and the boundary have to be $\mathcal{C}^{1,1}$ regular. Conversely, we study the regularity of the distance function under regularity hypotheses of the boundary. Along the way, we point out that any solution to the eikonal equation, differentiable everywhere in a domain of the Euclidean space, admits a gradient which is locally Lipschitz.

1. INTRODUCTION

Let $D \subsetneq \mathbb{R}^m$ be a domain ($m \geq 2$). The distance to its boundary is denoted $\delta_D(x) := \min_{y \in \partial D} |x - y|$. The *signed distance function* to ∂D is defined by $d_D := \delta_D$ on D and $d_D := -\delta_D$ on $\mathbb{R}^m \setminus D$.

Throughout this note, we write $\langle \cdot, \cdot \rangle$ for the usual Euclidean inner product in \mathbb{R}^m , and $|x| := \langle x, x \rangle^{1/2}$ for the Euclidean norm.

Our main goal is to prove that the signed distance function has a bootstrap property: if d_D is differentiable in an open set U , it must be $\mathcal{C}^{1,1}$ regular (Corollary 6). This is a generalization of results obtained in a succession of previous works, notably [3, 7, 6, 1].

We achieve this in Section 2 by proving that d_D satisfies the *eikonal equation* $|\nabla d_D(x)| = 1$ where it is differentiable. This is easy and well-known away from ∂D , and we show how to extend it to the boundary in Proposition 2. We then bring to bear previous works about the eikonal equation, summed up in Proposition 4.

While the eikonal equation may seem a more general hypothesis, Caffarelli and Crandall [1] proved that, up to additive constant, solutions of the eikonal equation are actually distance functions [1, (1.9) and Proposition 4.4], a result recalled here in Lemma 5. Where it is defined, $\nabla d_D(x)$ is a divergence-free unit vector field. Gains in regularity also occur in this more general case of vector fields; for the case $m = 2$, [4, Theorem 1] shows that if D is a domain in \mathbb{R}^2 , any such vector field which is in the Sobolev space $W^{1/p,p}(D)$ for some $p \in [1, 2]$ must be locally Lipschitz continuous inside D (which means locally the gradient of a $\mathcal{C}^{1,1}$ potential) except at a locally finite number of singular points. One can also find in [4] a wealth of examples on those topics, including cases where the $\mathcal{C}^{1,1}$ potential cannot be of class \mathcal{C}^2 , and a simply connected domain which allows for an infinite number of said singular points.

Then in Section 3, we turn to results about the distance function under hypotheses on the regularity of ∂D . To conclude, we prove Proposition 8, a more precise estimate about the variation of d_D .

Key words and phrases. eikonal equation, distance function, defining function, gain in regularity.

The first named author was partially supported by the Bulgarian National Science Fund, Ministry of Education and Science of Bulgaria under contract KP-06-N52/3.

2. BOOTSTRAP PROPERTIES OF d_D

In general, d_D is a 1-Lipschitz function. It has been known for a long time that it inherits much of the smoothness of ∂D , see [6] and the references therein. Precisely, when ∂D is \mathcal{C}^k -smooth, $k \geq 2$, so is d_D in a neighborhood of ∂D ; the same is true for $k = 1$ under the additional hypothesis that ∂D is of *positive reach*, that is, that x admits a unique projection to ∂D whenever the distance from x to ∂D is less than a certain uniform positive number [3].

Recall the following basic fact (see e.g. [3, Theorem 4.8]).

Proposition 1. *Let $D \subsetneq \mathbb{R}^m$ be an open set. Then d_D is differentiable at $x \notin \partial D$ exactly when x admits a unique projection $\pi(x)$ to ∂D . In that case, $\nabla d_D(x) = \frac{x - \pi(x)}{d(x)}$.*

In particular, whenever d_D is differentiable at $x \notin \partial D$, d_D satisfies the eikonal equation. This equation has been much studied, see e.g. [1], [2]. It turns out to extend to the boundary wherever ∇d_D still makes sense.

Proposition 2. *Let $D \subsetneq \mathbb{R}^m$ be an open set. If d_D is differentiable at $p \in \partial D$ and in $U \cap D$, for a neighborhood U of p , then $|\nabla d_D(p)| = 1$.*

Proof. Suppose d_D is differentiable at a point $p \in \partial D$.

We first prove that if $\nabla d_D(p) \neq 0$, then $|\nabla d_D(p)| = 1$. Since d_D is 1-Lipschitz, $|\nabla d_D(p)| \leq 1$. Take coordinates in an orthonormal basis such that $p = 0$ and $\nabla d_D(p) = \alpha e_1$, $\alpha \in (0, 1]$.

Suppose $\alpha < 1$. Then for any small enough positive x_1 , $|\pi(x_1, 0) - (x_1, 0)| < (1 + \alpha)x_1/2$, where $\pi(x_1, 0)$ is any of the closest points to $(x_1, 0)$ on ∂D . Let $p \in D$ be such that $|\pi(x_1, 0) - p| < \alpha(1 - \alpha)x_1/4$; we must have

$$p_1 > x_1 - \frac{1 + \alpha}{2}x_1 - \alpha \frac{1 - \alpha}{4}x_1 = \left(1 - \frac{\alpha}{2}\right) \frac{1 - \alpha}{2}x_1,$$

and

$$|p| \leq x_1 + \frac{1 + \alpha}{2}x_1 + \alpha \frac{1 - \alpha}{4}x_1 = Cx_1.$$

Finally

$$\frac{\alpha}{4}(1 - \alpha)x_1 > d_D(p) = \alpha p_1 + o(|p|) \geq \alpha \left(1 - \frac{\alpha}{2}\right) \frac{1 - \alpha}{2}x_1 + o(x_1),$$

a contradiction, so $|\nabla d_D(p)| = 1$.

Finally we are reduced to excluding the case $|\nabla d_D(p)| = 0$. If that holds, for any $\varepsilon > 0$, there exists $\delta_\varepsilon > 0$ such that if $|x - p| < \delta_\varepsilon$, then $|d_D(x)| < \varepsilon|x - p|$.

For some $x \in D$ such that d_D is differentiable at x , consider the integral curve of the vector field ∇d_D with starting point at x , i.e. the map $\chi : [0, T) \rightarrow D$ such that $\chi'(t) = \nabla d_D(\chi(t))$, $\chi(0) = x$ (this is known as a *characteristic* of the function d_D and will turn out to be an affine map [1, Lemma 2.2], but we shall not need this fact at this point).

Let

$$t_x := \sup \{t : d_D \text{ is differentiable in a neighborhood of } \chi(s), 0 \leq s \leq t\}.$$

By construction, for any $\eta > 0$, there is a point $x' \in D$ where d_D is not differentiable and $|x' - \chi(t_x)| < \eta$. By the differential equation, for $0 \leq s \leq t_x$,

$$d_D(\chi(s)) = d_D(x) + \int_0^s \langle \nabla d_D(\chi(t)), \chi'(t) \rangle dt = d_D(x) + \int_0^s |\nabla d_D(\chi(t))|^2 dt = d_D(x) + s.$$

We claim that for any $\varepsilon > 0$, there exists $y \in B(p, \varepsilon) \cap D$ such that d_D is not differentiable at y . For any $\varepsilon \in (0, \frac{1}{4})$, choose δ small enough so that $\delta < \varepsilon$, $2\delta \leq \delta_\varepsilon$. Let $x \in B(p, \delta) \cap D$. If d_D is not differentiable at x , we can take $y = x$.

It is, choose x' as above with $\eta := \varepsilon/10$. We claim that $t_x \leq \frac{1}{2}|x - p|$. Indeed, if $t_x > \frac{1}{2}|x - p|$, then for $s = \frac{1}{2}|x - p|$, $d_D(\chi(s)) \geq d_D(x) + s$. On the other hand, $|\chi(s) - p| \leq \frac{3}{2}|x - p| < \delta_\varepsilon$, so that $d_D(\chi(s)) < \varepsilon|\chi(s) - p| < 3\varepsilon s < \frac{3}{4}s$, a contradiction.

Finally $|x' - p| \leq |x - p| + t_x + \eta < \varepsilon$, and we can take $y = x'$. \square

Notice that the case where $\nabla d_D(p) = 0$ can occur. We give an example in \mathbb{R}^2 , which we assimilate to \mathbb{C} in order to use polar coordinates.

Let $f : [0, \infty) \rightarrow (0, \infty)$ be a continuous strictly decreasing function satisfying $\lim_{x \rightarrow \infty} f(x) = 0$. Let

$$\Omega := \{re^{i\theta} : 0 < \theta, f(\theta + \pi) < r < f(\theta)\}.$$

The domain Ω looks like a thickened spiral and for any α ,

$$\Omega \cap \mathbb{R}_+ e^{i\alpha} = \{re^{i\alpha} : f(\alpha + (2k + 1)\pi) < r < f(\alpha + 2k\pi), k \in \mathbb{Z}\}.$$

Proposition 3. *If $\lim_{x \rightarrow \infty} \frac{f(\theta)}{f(\theta + \pi)} = 1$, then Ω is a connected, simply connected domain such that at the point $0 \in \partial\Omega$, the signed distance function d_Ω is differentiable and has gradient equal to 0.*

Proof. The domain $\Omega_0 := \{(r, \theta) \in (0, \infty) \times (0, \infty) : 0 < \theta, f(\theta + \pi) < r < f(\theta)\}$ is clearly connected and simply connected.

The map $(r, \theta) \mapsto re^{i\theta}$ is continuous, so Ω is connected. For a point $re^{i\theta} \in \Omega$, the condition $f(\theta + \pi) < r < f(\theta)$ means that there is a single possible choice of θ , therefore the inverse map is well defined and continuous, so Ω is homeomorphic to Ω_0 .

Let $z = re^{i\theta} \in \Omega$. Considering the line segment $[e^{i\theta}f(\theta + \pi); e^{i\theta}f(\theta)]$ we see that $d_\Omega(z) \leq f(\theta) - f(\theta + \pi)$ and $|z| \geq f(\theta + \pi)$. Therefore

$$0 \leq \frac{d_\Omega(z)}{|z|} \leq \frac{f(\theta)}{f(\theta + \pi)} - 1 \rightarrow 0,$$

as $|z| \rightarrow 0$ (which implies $\theta \rightarrow \infty$).

For points outside of Ω , the proof is similar. \square

In the above example, between the two spirals that constitute $\partial\Omega$, there is a spiral of points of Ω which are equidistant from the two components and where d_D is not differentiable.

The following is more or less implicit in [1].

Proposition 4. *Let $\Omega \subset \mathbb{R}^m$ be an open set and $\varphi : \Omega \rightarrow \mathbb{R}$ be an everywhere differentiable solution of the eikonal equation $|\nabla\varphi| = 1$ in Ω . Then $\varphi \in \mathcal{C}^{1,1}(\Omega)$, i.e. $\nabla\varphi$ is locally Lipschitz in Ω .*

As a consequence, if d_D is differentiable in a neighborhood U of ∂D , and $|\nabla d_D(x)| = 1$ for $x \in U \cap \partial D$, then $d_D \in \mathcal{C}^{1,1}(U)$.

Proof. We outline how to use the results of Caffarelli and Crandall [1] to prove this proposition in our more special case (they consider a more general family of norm and functions with a possible small singularity set).

By [1, Lemma 2.2], if $a < 0 < b$ and $x + t\nabla\varphi(x) \in U$ for $a < t < b$, then for those values of t , $\nabla\varphi(x + t\nabla\varphi(x)) = \nabla\varphi(x)$ and $\varphi(x + t\nabla\varphi(x)) = t$. Moreover, $\varphi \in \mathcal{C}^1(U)$.

The straight lines $\{x + t\nabla\varphi(x), t \in \mathbb{R}\}$ are called *characteristics* of φ . For any $a \in \varphi(U)$, we denote the level set by $S_a := \{x \in U : \varphi(x) = a\}$.

The following is a direct consequence of [1, Proposition 4.4] and its proof.

Lemma 5. *Let V be an open set of the form*

$$\{x + t\nabla\varphi(x), t \in (a(x), b(x)), x \in S_a \cap V\},$$

where $a \in \varphi(U)$ and $a(x) < 0 < b(x)$. Then for any $y \in V$, $d_{S_a}(y) = \varphi(y) - a$.

In order to apply it to our situation, we need to see that any $x \in U$ admits a neighborhood V of the form above. Let $x_0 \in U$. Choose affine local coordinates $(x', x^m) \in \mathbb{R}^{m-1} \times \mathbb{R}$ so that $x_0 = 0$ and $\nabla\varphi(x_0) = (0, \dots, 0, 1)$. Then $\Phi(x', x^m) = (x', \varphi(x))$ is a local \mathcal{C}^1 -diffeomorphism. In particular, for $r > 0$ small enough and $|x'| \leq r$, there exists a unique x^m such that $\varphi(x', x^m) = \varphi(0)$.

We can choose δ small enough so that

$$U_r := \{x + t\nabla\varphi(x) : x \in S_{\varphi(x_0)}, |x'| < r, |t| < r\} = \Phi^{-1}(B^{m-1}(0, r) \times (-r, r))$$

is a open neighborhood of x_0 contained in U for $r \leq 3\delta$, made up of a disjoint union of segments of characteristics. Let $a := \varphi(x_0) - 2\delta$, then if $x \in U_\delta$, we have $\varphi(x) = \text{dist}(x, S_a) - 2\delta + \varphi(x_0)$, so $\nabla\varphi(x) = \nabla d_{S_a}(x)$.

Using Proposition 1, for $x, y \in U_\delta$,

$$\begin{aligned} \nabla\varphi(x) - \nabla\varphi(y) &= \frac{x - \pi_{S_a}(x)}{d_{S_a}(x)} - \frac{y - \pi_{S_a}(y)}{d_{S_a}(y)} \\ &= \frac{1}{d_{S_a}(x)} (x - y - (\pi_{S_a}(x) - \pi_{S_a}(y))) - (x - \pi_{S_a}(x)) \frac{d_{S_a}(x) - d_{S_a}(y)}{d_{S_a}(x)d_{S_a}(y)}, \end{aligned}$$

so that, since $d_{S_a}(x), d_{S_a}(y) \geq \delta$, then $|\nabla\varphi(x) - \nabla\varphi(y)| \leq 2\delta^{-1}|x - y| + \delta^{-1}|x - y|$. \square

Corollary 6. *If d_D is differentiable in a neighborhood U of $p \in \partial D$, then $d_D \in \mathcal{C}^{1,1}(U)$.*

Proof. By Proposition 1, $|\nabla d_D| = 1$ on $U \setminus \partial D$. By Proposition 2, $|\nabla d_D| = 1$ on $U \cap \partial D$. Then Proposition 4 applies. \square

Remark. If we already know that $d_D \in \mathcal{C}^1(U)$, for U a neighborhood of ∂D , then the hypotheses of the second part of Proposition 4 are satisfied. With that hypothesis, ∂D is \mathcal{C}^1 -smooth (since $|\nabla d_D| = 1$ on U by continuity). In this case, as pointed out in [6, p. 120], K. Lucas's work [7, Section 2] implies that ∂D is $\mathcal{C}^{1,1}$ -smooth (see also [3, Theorem 4.18]) and hence, by Proposition 8 below, d_D is $\mathcal{C}^{1,1}$ -smooth near ∂D . But that proof is rather more roundabout than the arguments from [1].

3. BOUNDARY REGULARITY ASSUMPTIONS

We now turn to weaker hypotheses, involving only the regularity of ∂D itself.

Proposition 7. *Let p be a \mathcal{C}^1 -smooth boundary point of a domain D in \mathbb{R}^m . Then d_D is differentiable at p and $\nabla d_D(p)$ is the inner unit normal vector to ∂D at p .*

This does not necessarily extend to any neighborhood. For $0 < \alpha < 1$ the domain $D_\alpha = \{x \in \mathbb{R}^2 : x_1 > |x_2|^{1+\alpha}\}$ is $\mathcal{C}^{1,\alpha}$ - but not $\mathcal{C}^{1,1}$ -smooth. By Propositions 1 and 7, the function d_D is non-differentiable at x exactly when $x = (x_1, 0)$ with $x_1 > 0$.

Proof. We may assume that that $D = \{x_1 > f(x')\}$ near $p = 0$, where $x = (x_1, x') \in \mathbb{R} \times \mathbb{R}^{m-1}$ and $f(x') = o(|x'|)$.

Let $x \in D$. Since $\tilde{x} = (f(x'), x') \in \partial D$, then

$$d_D(x) - x_1 \leq |x - \tilde{x}| - x_1 = -f(x').$$

On the other hand, let $\hat{x} \in \partial D$ be such that $d_D(x) = |x - \hat{x}|$. Then

$$d_D(x) - x_1 \geq |x_1 - \hat{x}_1| - x_1 \geq -|\hat{x}_1| = -|f(\hat{x}')|.$$

Since $|\hat{x}| \leq |x| + d_D(x) \leq 2|x|$, it follows that $d_D(x) - x_1 = o(|x'|)$.

Similar arguments imply the same for $x \notin D$ which completes the proof. \square

We now give a sufficient condition for the $\mathcal{C}^{1,1}$ smoothness of d_D in terms of conditions on ∂D only, along with a more precise estimate of its second order variation.

Proposition 8. *Let p be a $\mathcal{C}^{1,1}$ -smooth boundary point of a domain D in \mathbb{R}^m . Then d_D is $\mathcal{C}^{1,1}$ -smooth near p . Moreover, for x, y near p one has that*

$$|d_D(x) - d_D(y) - \langle \nabla d_D(x), x - y \rangle| \leq (\chi_{D,p}/2 + o(1))(|x - y|^2 - (d_D(x) - d_D(y))^2).$$

Proof. Recall that when $q \in \partial D$, $\nabla d_D(q) = n_q$, the unit inner normal vector to ∂D at $q \in \partial D$. Set \tilde{x} to be the projection of x near p on ∂D (it is unique by $\mathcal{C}^{1,1}$ -smoothness). Then

$$\begin{aligned} \langle n_{\tilde{x}}, x - y \rangle &= \langle n_{\tilde{x}}, \tilde{x} + d_D(x)n_{\tilde{x}} - \tilde{y} - d_D(y)n_{\tilde{y}} \rangle = \\ &= d_D(x) - d_D(y) + \langle n_{\tilde{x}}, \tilde{x} - \tilde{y} \rangle + (1 - \langle n_{\tilde{x}}, n_{\tilde{y}} \rangle)d_D(y). \end{aligned}$$

Let $\chi_0 = \chi_{D,p}$. Since $|n_{\tilde{x}} - n_{\tilde{y}}| \leq (\chi_0 + o(1))|\tilde{x} - \tilde{y}|$, it follows that

$$1 - \langle n_{\tilde{x}}, n_{\tilde{y}} \rangle \leq (\chi_0^2/2 + o(1))|\tilde{x} - \tilde{y}|^2.$$

To estimate $\langle n_{\tilde{x}}, \tilde{x} - \tilde{y} \rangle$, we may assume that $p = 0$ and ∂D near 0 is given by $u_1 = f(u')$, where $u' := (u_2, \dots, u_m)$, for $|u'| < \varepsilon_0$, with $f(0) = 0$, and $\nabla f(0) = 0$.

For $\tilde{x} = (f(\tilde{x}'), \tilde{x}')$, $(1, -\nabla f(\tilde{x}')) = \sqrt{1 + |\nabla f(\tilde{x}')|^2} n_{\tilde{x}} = (1 + o(1))n_{\tilde{x}}$, so

$$(3.1) \quad |\nabla f(u') - \nabla f(v')| \leq (\chi_0 + o(1))|u' - v'|.$$

Then

$$\sqrt{1 + |\nabla f(\tilde{x}')|^2} \langle n_{\tilde{x}}, \tilde{x} - \tilde{y} \rangle = f(\tilde{x}') - f(\tilde{y}') - \langle \nabla f(\tilde{x}'), \tilde{x}' - \tilde{y}' \rangle.$$

Writing $g(t) := f((1-t)\tilde{x}' + t\tilde{y}')$, $g \in \mathcal{C}^{1,1}$, and we need to estimate $|g(1) - g(0) - g'(0)|$; by Taylor's formula with integral remainder, it is bounded by

$$\frac{1}{2}|\tilde{x}' - \tilde{y}'|^2 \sup_{[0,1]} |g''| = \frac{1}{2}(\chi_0 + o(1))|\tilde{x}' - \tilde{y}'|^2,$$

by (3.1). Finally

$$|\langle n_{\tilde{x}}, \tilde{x} - \tilde{y} \rangle| \leq (\chi_0/2 + o(1))|\tilde{x}' - \tilde{y}'|^2 \leq (\chi_0/2 + o(1))|\tilde{x} - \tilde{y}|^2,$$

thus

$$|d_D(x) - d_D(y) - \langle n_{\tilde{x}}, x - y \rangle| = |\langle n_{\tilde{x}}, \tilde{y} - \tilde{x} \rangle - (1 - \langle n_{\tilde{x}}, n_{\tilde{y}} \rangle)d_D(y)| \leq (\chi_0/2 + o(1))|\tilde{x} - \tilde{y}|^2.$$

It remains to compare $|\tilde{x} - \tilde{y}|^2$ with $|x - y|^2$. But

$$\begin{aligned} |x - y|^2 &= |(\tilde{x} - \tilde{y}) + (d_D(x) - d_D(y))n_{\tilde{x}} + d_D(y)(n_{\tilde{x}} - n_{\tilde{y}})|^2 \\ &= |\tilde{x} - \tilde{y}|^2 + (d_D(x) - d_D(y))^2 + 2[d_D(x)\langle n_{\tilde{x}}, \tilde{x} - \tilde{y} \rangle + d_D(y)\langle n_{\tilde{y}}, \tilde{y} - \tilde{x} \rangle + d_D(x)d_D(y)(1 - \langle n_{\tilde{x}}, n_{\tilde{y}} \rangle)] \\ &= (1 + o(1))|\tilde{x} - \tilde{y}|^2 + (d_D(x) - d_D(y))^2. \end{aligned}$$

So

$$|d_D(x) - d_D(y) - \langle n_{\tilde{x}}, x - y \rangle| \leq (\chi_0/2 + o(1))(|x - y|^2 - (d_D(x) - d_D(y))^2).$$

Hence d_D is $\mathcal{C}^{1,1}$ -smooth near p and $\nabla d_D(x) = n_{\tilde{x}}$. \square

Acknowledgements. We wish to thank our colleague Radu Ignat for useful discussions about the eikonal equation. He also has a direct proof for the $\mathcal{C}^{1,1}$ interior regularity of solutions of the eikonal equations in any dimension [5].

REFERENCES

- [1] L.A. Caffarelli, M. G. Crandall, *Distance functions and almost global solutions of eikonal equations*, Commun. Partail Differ. Equ. **35** (2010), 391–414.
- [2] C. De Lellis, R. Ignat, *A regularizing property of the 2D-eikonal equation*, Commun. Partail Differ. Equ. **40** (2015), 1543–1557.
- [3] H. Federer, *Curvature measures*, Trans. Amer. Math. Soc. **93** (1959), 418–491.
- [4] R. Ignat, *Two-dimensional unit-length vector fields of vanishing divergence*, J. Funct. Anal. **262** (2012), 3465–3494.
- [5] R. Ignat, *A short proof of the $\mathcal{C}^{1,1}$ regularity for the eikonal equation*, in preparation.
- [6] S.G. Krantz, H.R. Parks, *Distance to \mathcal{C}^k hypersurfaces*, J. Differ. Equ. **40** (1981), 116–120.
- [7] K.R. Lucas, *Submanifolds of dimension $n - 1$ in \mathcal{E}^n with normals satisfying a Lipschitz condition*, Studies in Eigenvalue Problems, Technical Report no. 18, University of Kansas, 1957.

N. NIKOLOV, INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, ACAD. G. BONCHEV 8, 1113 SOFIA, BULGARIA

FACULTY OF INFORMATION SCIENCES, STATE UNIVERSITY OF LIBRARY STUDIES AND INFORMATION TECHNOLOGIES, SHIPCHENSKI PROHOD 69A, 1574 SOFIA, BULGARIA

Email address: `nik@math.bas.bg`

P.J. THOMAS, INSTITUT DE MATHÉMATIQUES DE TOULOUSE; UMR5219, UNIVERSITÉ DE TOULOUSE; CNRS, UPS, F-31062 TOULOUSE CEDEX 9, FRANCE

Email address: `pascal.thomas@math.univ-toulouse.fr`