

Refined existence theorems for doubly degenerate chemotaxis-consumption systems with large initial data

DUAN WU*

Institut für Mathematik, Universität Paderborn, 33098 Paderborn, Germany

and

School of Mathematics and Statistics, Northwestern Polytechnical University, Xi'an 710129, PR China

Abstract

This work considers the doubly degenerate nutrient model

$$\begin{cases} u_t = \nabla \cdot (u^{m-1}v\nabla u) - \nabla \cdot (f(u)v\nabla v) + \ell uv, & x \in \Omega, t > 0, \\ v_t = \Delta v - uv, & x \in \Omega, t > 0, \end{cases}$$

under no-flux boundary conditions in a smoothly bounded convex domain $\Omega \subset \mathbb{R}^n$ ($n \leq 2$), where the nonnegative function $f \in C^1([0, \infty))$ is assumed to satisfy $f(s) \leq C_f s^\alpha$ with $\alpha > 0$ and $C_f > 0$ for all $s \geq 1$.

When $m = 2$, it was shown that a global weak solution exists, either in one-dimensional setting with $\alpha = 2$, or in two-dimensional version with $\alpha \in (1, \frac{3}{2})$. The main results in this paper assert the global existence of weak solutions for $1 \leq m < 3$ and classical solutions for $3 \leq m < 4$ to the above system under the assumption

$$\alpha \in \begin{cases} \left[m - 1, \min \left\{ m, \frac{m}{2} + 1 \right\} \right] & \text{if } n = 1, \quad \text{and} \\ \left(m - 1, \min \left\{ m, \frac{m}{2} + 1 \right\} \right) & \text{if } n = 2, \end{cases}$$

which extend the range $\alpha \in (1, \frac{3}{2})$ to $\alpha \in (1, 2)$ in two dimensions for the case $m = 2$. Our proof will be based on a new observation on the coupled energy-type functional and on an inequality with general form.

Keywords: degenerate diffusion; chemotaxis; global existence

AMS (2020) Subject Classification: 35K65; 92C17; 35A01

1 Introduction

Being motivated by some experimental indications when investigating the dynamical motility of *Bacillus subtilis* grown on thin agar plates ([4],[14],[15]), Kawasaki et.al in [6] introduced the fol-

*E-mail: duan_wu@126.com

lowing reaction-diffusion continuous system

$$\begin{cases} u_t = \nabla \cdot (D(u, v)\nabla u) - \nabla \cdot (S(u, v)\nabla v) + \ell uv, \\ v_t = \Delta v - uv \end{cases} \quad (1.1)$$

with the non-linear diffusion coefficient $D(u, v) = uv$ in line with the peculiarity that the bacteria would be immobile when exposed to regions with the low level of nutrients, and with the absence of taxis processes given by $S(u, v) = 0$. Subsequently, to comply with experimental phenomena discovered in [5] and [3], Leyva et.al in [9] proposed that the bacterial response function in the chemotactic flux term should be considered by $S(u, v) = \chi u^2 v$ with $\chi > 0$ measuring the intensity of the chemotaxis, which was supported by numerical simulations: the conformity between the experimentally gained observations and the behavior of numerical solutions is noticeably greater than the case taking $S(u, v) = 0$ (see [9, Sections 4 and 5]).

Within the realm of results from mathematical analysis, even in the case of the prototypical setting upon the choices $D(u, v) = 1$ and $S(u, v) = u$, to the best of our knowledge there has been sparse literature with respect to solvability and spatially homogeneous equilibria for (1.1) ([18], [20]), which are confined to the relatively lower-dimensional space.

For the doubly degenerate diffusion problem such as when $D(u, v) = uv$, virtually no mathematical findings regarding basic solution theory touched on it until Winkler in [21] revealed the existence of global weak solutions for arbitrary large initial data in one-dimensional space with the choice of $S(u, v) = u^2 v$ as suggested in [9], where the most stunning feature is that the first component of the solution asymptotically stabilizes towards a nontrivial function coinciding with the spatial profile of a solution to a scalar porous medium-type parabolic equation in stark contrast to the majority of previous results concerning the large time behavior. In the corresponding taxis-free framework, namely $S(u, v) = 0$, the global existence of weak solutions and the stabilization were asserted in convex domains of any dimension ([24]). For the same choices of $S(u, v)$ and $D(u, v)$ as studied in [21], Li and Winkler removed the integrability assumption $\int_{\Omega} \ln u_0 > -\infty$ to establish the similar results to system (1.1) in the spatially one-dimensional analogue by relying on a striking energy functional ([11]), while for its two-dimensional version the global solvability were substantiated in bounded convex domains with a smallness condition exclusively involving v_0 ([23]). When the cross-diffusion coefficient has the form $S(u, v) = \chi u^{\alpha} v$ with $\alpha > 0$, without any smallness restrictions on the size of initial data in bounded convex domains, Li identified that (1.1) possesses a global weak solution under the range $1 < \alpha < \frac{3}{2}$ in two-dimensional case, and $\frac{7}{6} < \alpha < \frac{13}{9}$ in three-dimensional case ([10]).

Motivated by the above precedent results, we have a natural question that when $D(u, v) = uv$ and $S(u, v) = \chi u^{\alpha} v$ in (1.1), if $\alpha = \frac{3}{2}$ is a critical existence exponent in two dimensions. Specifically, we attempt to figure out whether the smallness on initial data is necessary to guarantee the solvability in the flavor of [10] for the case $\frac{3}{2} \leq \alpha < 2$. Meanwhile, we are devoted to studying the corresponding solvability for system (1.1) with a more general nonlinear diffusion term, which plays an important role in the biological context (see [7], [2]).

More precisely, in the present work we will be concerned with the initial-boundary value problem

$$\begin{cases} u_t = \nabla \cdot (u^{m-1}v\nabla u) - \nabla \cdot (f(u)v\nabla v) + \ell uv, & x \in \Omega, t > 0, \\ v_t = \Delta v - uv, & x \in \Omega, t > 0, \\ (u^{m-1}v\nabla u - f(u)v\nabla v) \cdot \nu = \nabla v \cdot \nu = 0, & x \in \partial\Omega, t > 0, \\ u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), & x \in \Omega, \end{cases} \quad (1.2)$$

with $m \geq 1$, $\ell \geq 0$, under the overall hypotheses that f besides

$$f \in C^1([0, \infty)) \quad \text{and} \quad f(u) \geq 0 \quad \text{for all } u \geq 0 \quad (1.3)$$

is such that

$$f(u) \leq C_f u(u+1)^{\alpha-1} \quad \text{for all } u \geq 0 \quad (1.4)$$

or

$$f(u) \leq C_f u^\alpha \quad \text{for all } u \geq 0 \quad (1.5)$$

with $C_f > 0$ and $\alpha \geq 0$. In addition, the initial data (u_0, v_0) is throughout supposed to satisfy that

$$\begin{cases} u_0 \in W^{1,\infty}(\Omega) \text{ is nonnegative with } u_0 \not\equiv 0 & \text{and} \\ v_0 \in W^{1,\infty}(\Omega) \text{ is positive in } \bar{\Omega}. \end{cases} \quad (1.6)$$

Before stating our main results, we first introduce the definition of weak solutions pursued in this context.

Definition 1.1. *Let $n \in \{1, 2\}$ and $\Omega \subset \mathbb{R}^n$ be a bounded domain with smooth boundary, $m \geq 1$ and $\ell \geq 0$. Suppose that f satisfies (1.4) or (1.5) with $C_f > 0$ and $\alpha \geq 0$ in addition to (1.3), and that $u_0 \in L^1(\Omega)$ and $v_0 \in L^1(\Omega)$ are nonnegative. Then we call that a pair (u, v) of nonnegative functions*

$$\begin{cases} u \in L^1_{loc}(\bar{\Omega} \times [0, \infty)) & \text{and} \\ v \in L^\infty_{loc}(\bar{\Omega} \times [0, \infty)) \cap L^1_{loc}([0, \infty); W^{1,1}(\Omega)). \end{cases}$$

satisfying

$$u^m \nabla v \in L^1_{loc}(\bar{\Omega} \times [0, \infty); \mathbb{R}^n) \quad \text{and} \quad u^m v \in L^1_{loc}(\bar{\Omega} \times [0, \infty))$$

is a global weak solution of (1.2) if

$$\begin{aligned} - \int_0^\infty \int_\Omega u \varphi_t - \int_\Omega u_0 \varphi(\cdot, 0) &= \frac{1}{m} \int_0^\infty \int_\Omega u^m \nabla v \cdot \nabla \varphi + \frac{1}{m} \int_0^\infty \int_\Omega u^m v \Delta \varphi \\ &+ \int_0^\infty \int_\Omega f(u)v \nabla v \cdot \nabla \varphi + \ell \int_0^\infty \int_\Omega uv \varphi \end{aligned}$$

for all $\varphi \in C_0^\infty(\bar{\Omega} \times [0, \infty))$ fulfilling $\frac{\partial \varphi}{\partial \nu} = 0$ on $\partial\Omega \times (0, \infty)$, as well as

$$\int_0^\infty \int_\Omega v \varphi_t + \int_\Omega v_0 \varphi(\cdot, 0) = \int_0^\infty \int_\Omega \nabla v \cdot \nabla \varphi + \int_0^\infty \int_\Omega uv \varphi$$

for each $\varphi \in C_0^\infty(\bar{\Omega} \times [0, \infty))$.

The first result concerns one-dimensional space.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}$ be an open bounded interval. Suppose that $\ell \geq 0$ and that the initial data u_0 and v_0 satisfy (1.6). Then for all $K > 0$ with the property that*

$$\|u_0\|_{L^\infty(\Omega)} + \|v_0\|_{L^\infty(\Omega)} + \|\partial_x \ln v_0\|_{L^\infty(\Omega)} \leq K, \quad (1.7)$$

if one of the following cases holds:

- (i) $1 \leq m < 2$, f fulfills (1.3) and (1.4) with $m - 1 \leq \alpha \leq m$;
 - (ii) $2 \leq m < 3$, f fulfills (1.3) and (1.5) with $m - 1 \leq \alpha \leq \frac{m}{2} + 1$,
- the problem (1.2) admits a global weak solution (u, v) in the sense of Definition 1.1. Moreover, if
- (iii) $3 \leq m < 4$, f fulfills (1.3) and (1.5) with $m - 1 \leq \alpha \leq \frac{m}{2} + 1$ and $u_0 > 0$ in $\overline{\Omega}$,
- the problem (1.2) admits a global classical solution (u, v) .

Furthermore, $v > 0$ a.e. in $\Omega \times (0, \infty)$, and that there exists $C(K) > 0$ such that

$$\|u(\cdot, t)\|_{L^\infty(\Omega)} + \|v(\cdot, t)\|_{W^{1,\infty}(\Omega)} \leq C(K) \quad \text{for a.e. } t > 0. \quad (1.8)$$

Remark 1.1. In addition, with the aid of the Harnack type inequality valid in one-dimensional space ([25]), in much the same way as used in [11], we could further obtain the following asymptotic behavior result.

Corollary 1. *Let the assumptions in Theorem 1.1 hold and (u, v) be as accordingly given by Theorem 1.1, then there exists $u_\infty = w(\cdot, 1) \in C^0(\overline{\Omega})$ such that $u(\cdot, t) \rightarrow u_\infty$ in $L^\infty(\Omega)$ as $t \rightarrow \infty$. Here $w \in C^0(\overline{\Omega} \times [0, 1]) \cap L^2_{loc}([0, 1]; W^{1,2}(\Omega))$ is a weak solution of*

$$\begin{cases} w_\tau = (a(x, \tau)w^{m-1}w_x)_x - (b(x, \tau)f(w))_x + \ell a(x, \tau)w, & x \in \Omega, \tau \in (0, 1), \\ w_x = 0, & x \in \partial\Omega, \tau \in (0, 1), \\ w(x, 0) = u_0(x), & x \in \Omega, \end{cases}$$

where $a(x, \tau)$ and $b(x, \tau)$ are defined as that in [21, Theorem 1.3].

In two-dimensional setting, for convex domains, we have the following result.

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^2$ be a bounded convex domain with smooth boundary, and let $\ell \geq 0$. Suppose that the initial data u_0 and v_0 satisfy (1.6). Then for all $K > 0$ with the property that*

$$\|u_0\|_{L^\infty(\Omega)} + \|v_0\|_{L^\infty(\Omega)} + \|\nabla \ln v_0\|_{L^\infty(\Omega)} \leq K, \quad (1.9)$$

if one of the following cases holds:

- (i) $1 \leq m < 2$, f fulfills (1.3) and (1.4) with $m - 1 < \alpha < m$;
 - (ii) $2 \leq m < 3$, f fulfills (1.3) and (1.5) with $m - 1 < \alpha < \frac{m}{2} + 1$,
- the problem (1.2) admits a global weak solution (u, v) in the sense of Definition 1.1. Moreover, if
- (iii) $3 \leq m < 4$, f fulfills (1.3) and (1.5) with $m - 1 < \alpha < \frac{m}{2} + 1$ and $u_0 > 0$ in $\overline{\Omega}$,
- the problem (1.2) admits a global classical solution (u, v) .

Furthermore, $v > 0$ a.e. in $\Omega \times (0, \infty)$, and that for all $p > 2$, there exists $C(p, K) > 0$ such that

$$\|u(\cdot, t)\|_{L^p(\Omega)} + \|v(\cdot, t)\|_{W^{1,\infty}(\Omega)} \leq C(p, K) \quad \text{for a.e. } t > 0. \quad (1.10)$$

Remark 1.2. The large time behavior has not yet been solved because of the absence of the corresponding Harnack type inequality in two-dimensional space, which also results in that we can only obtain the L^p bounds for u in (1.10) instead of the L^∞ bounds.

Remark 1.3. When $m = 2$, the upper bound on α obtained in this paper seems natural and reasonable from the perspective that if $\alpha = 2$, the solvability requires imposing small conditions on the initial data as indicated in [23].

Remark 1.4. In a manner rather similar to proving Theorem 1.2, the corresponding results in higher-dimensional space is able to be derived: at least in the context of three dimensions with the case $m = 2$, we can check that the global solvability holds within the range $1 < \alpha < \frac{3}{2}$, which together with Theorem 1.2 extends the previous results in [10].

The foundation of our approach relies closely on the useful functional first discovered in [22] of the form

$$\int_{\Omega} \frac{|\nabla v|^q}{v^{q-1}}, \quad q \geq 2$$

with the favorable features

$$\frac{d}{dt} \int_{\Omega} \frac{|\nabla v|^q}{v^{q-1}} + \frac{q}{2} \int_{\Omega} \frac{|\nabla v|^{q-2}}{v^{q-3}} |D^2 \ln v|^2 + (q-1)^2 \int_{\Omega} u \frac{|\nabla v|^q}{v^{q-1}} \leq C_1(q) \int_{\Omega} u^{\frac{q+2}{2}} v,$$

in particular when $q = 4$, by a slightly different estimation used to get the above estimation and making use of the good term $\int_{\Omega} \frac{|\nabla v|^6}{v^5}$ (identified by Lemma 2.2),

$$\frac{d}{dt} \int_{\Omega} \frac{|\nabla v|^4}{v^3} + \int_{\Omega} \frac{|\nabla v|^2}{v} |D^2 \ln v|^2 + \int_{\Omega} u \frac{|\nabla v|^4}{v^3} \leq C_2 \int_{\Omega} v |\nabla u|^2, \quad (1.11)$$

where $C_1(q)$ and C_2 are positive constants (Lemma 2.3).

In the spatially one-dimensional case, we commence with a coupled functional expressed by

$$\mathcal{F}(t) := \int_{\Omega} u^p + \int_{\Omega} \frac{v_x^2}{v}, \quad (1.12)$$

where p is suitably taken by $p = 3 - m$, roughly speaking. For the peculiar cases of $m = 2$ and $m = 3$, we substitute 1 with the function $\ln u$ which exhibits the same scaling as 1, and correspondingly u is replaced by $u \ln u$ (Lemma 3.3). Then on the basis of L^4 bound on v_x (Lemma 3.2), by means of a functional inequality (Lemma 3.1), the ill-signed contributions of \mathcal{F} could be controlled by the dissipated integral $\int_{\Omega} u \frac{v_x^2}{v}$ and the favorable term $\int_{\Omega} v u_x^2$ produced by the evolution of first summand in (1.12). This furnishes some boundedness properties presented in Lemma 3.4, which will be utilized to derive bounds for u in L^p with any $p > 1$ (Lemma 3.5).

When the spatial setting is two-dimensional, it is noticeable from Lemma 4.9 that if we have a bound for u in L^{p_0} with some $p_0 > 1$, then using an energy-like functional of the form

$$\int_{\Omega} u^p + \int_{\Omega} \frac{|\nabla v|^q}{v^{q-1}} \quad (1.13)$$

with suitably chosen $q \in \left(\frac{2(p+m-1)}{p_0}, 2(p_0 + p + m - 2)\right) =: I$ is an effectively direct method to obtain the L^p bounds for u with arbitrary $p > 2$. This way is not applicable when we try to merely rely on the basic boundedness of $\|u\|_{L^1}$ because I is empty when $p_0 = 1$. Thus, the current difficulty consists in how to obtain an L^{p_0} bound for u with some $p_0 > 1$. To achieve this, the core of our analysis is based on the following quasi-energy functional, given by

$$\mathcal{G}(t) := c \int_{\Omega} u^p + \int_{\Omega} \frac{|\nabla v|^4}{v^3} + \int_{\Omega} u^{p^*}, \quad (1.14)$$

where $c > 0$, p is as in (1.12) and $p^* \in (2 - m, 3 - m)$. Some basic differential inequalities for the first two summands in (1.14) are shown in Lemma 4.4, while the key role in addressing the unfavorable expressions therein is played by the crucial observation that

$$\int_{\Omega} u^{\beta} v \leq \eta \int_{\Omega} u^{\kappa} v |\nabla u|^2 + \eta \int_{\Omega} \frac{|\nabla v|^2}{v} |D^2 \ln v|^2 + C_3(\beta, \eta) \int_{\Omega} uv$$

holds for $\beta \in [1, \kappa + 3)$ with $\kappa \in (-1, 0)$ (Lemmas 4.2). The integral $\int_{\Omega} \frac{|\nabla v|^2}{v} |D^2 \ln v|^2$ can be dissipated during the evolution of $\int_{\Omega} \frac{|\nabla v|^4}{v^3}$ as shown in (1.11), and the other integral $\int_{\Omega} u^{\kappa} v |\nabla u|^2$ will be absorbed by further testing u^{p^*} in a standard procedure with diffusion-induced quantity $\int_{\Omega} u^{p^* + m - 3} v |\nabla u|^2$ arising. Then we could get some essential boundedness properties, which serve as a crucial ingredient in our derivation of an L^{p_0} bound for u (Lemmas 4.5-4.7).

Based on these results, we are able to collect some higher regularities (cf. Lemma 5.1, Lemma 5.3 and Lemma 5.5) and further announce the conclusion on the global existence.

2 Some preliminaries

In this section, we introduce some basic lemmas. Similar with the approximating procedure used in [11], we consider the regularized variant of (1.2) given by

$$\begin{cases} u_{\varepsilon t} = \nabla \cdot (u_{\varepsilon}^{m-1} v_{\varepsilon} \nabla u_{\varepsilon}) - \nabla \cdot (f(u_{\varepsilon}) v_{\varepsilon} \nabla v_{\varepsilon}) + \ell u_{\varepsilon} v_{\varepsilon}, & x \in \Omega, t > 0, \\ v_{\varepsilon t} = \Delta v_{\varepsilon} - u_{\varepsilon} v_{\varepsilon}, & x \in \Omega, t > 0, \\ \frac{\partial u_{\varepsilon}}{\partial \nu} = \frac{\partial v_{\varepsilon}}{\partial \nu} = 0, & x \in \partial\Omega, t > 0, \\ u_{\varepsilon}(x, 0) = u_{0\varepsilon}(x), \quad v_{\varepsilon}(x, 0) = v_{0\varepsilon}(x) := v_0(x), & x \in \Omega \end{cases} \quad (2.1)$$

with $\varepsilon \in (0, 1)$, where $u_{0\varepsilon}(x)$ depending on m is defined by

$$u_{0\varepsilon}(x) := \begin{cases} u_0(x) + \varepsilon, & 1 \leq m < 3, \\ u_0(x), & 3 \leq m < 4. \end{cases} \quad (2.2)$$

In light of the positivity of $u_{0\varepsilon}$, this problem enjoys the following properties.

Lemma 2.1. *Suppose that the assumptions in Theorem 1.1 or Theorem 1.2 are satisfied. Then for each $\varepsilon \in (0, 1)$, there exist $T_{\max, \varepsilon} \in (0, \infty]$ and at least one pair $(u_{\varepsilon}, v_{\varepsilon})$ of functions*

$$\begin{cases} u_{\varepsilon} \in \cap_{q \geq 1} C^0([0, T_{\max, \varepsilon}]; W^{1, q}(\Omega)) \cap C^{2, 1}(\overline{\Omega} \times (0, T_{\max, \varepsilon})) \\ v_{\varepsilon} \in \cap_{q \geq 1} C^0([0, T_{\max, \varepsilon}]; W^{1, q}(\Omega)) \cap C^{2, 1}(\overline{\Omega} \times (0, T_{\max, \varepsilon})) \end{cases} \quad (2.3)$$

such that $u_\varepsilon > 0$ and $v_\varepsilon > 0$ in $\overline{\Omega} \times (0, T_{\max, \varepsilon})$, that $(u_\varepsilon, v_\varepsilon)$ solves (2.1) in the classical sense in $\Omega \times (0, T_{\max, \varepsilon})$, and that

$$\text{if } T_{\max, \varepsilon} < \infty, \text{ then } \limsup_{t \rightarrow T_{\max, \varepsilon}} \|u_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} = \infty. \quad (2.4)$$

In addition, this solution satisfies

$$\|v_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq \|v_\varepsilon(\cdot, t_0)\|_{L^\infty(\Omega)} \quad \text{for all } t_0 \in [0, T_{\max, \varepsilon}) \text{ and any } t \in (t_0, T_{\max, \varepsilon}), \quad (2.5)$$

$$\int_{\Omega} u_{0\varepsilon} \leq \int_{\Omega} u_\varepsilon(\cdot, t) \leq \int_{\Omega} u_{0\varepsilon} + \ell \int_{\Omega} v_{0\varepsilon} \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \quad (2.6)$$

and

$$\int_{t_0}^{T_{\max, \varepsilon}} \int_{\Omega} u_\varepsilon v_\varepsilon \leq \int_{\Omega} v_\varepsilon(\cdot, t_0) \quad \text{for all } t_0 \in [0, T_{\max, \varepsilon}). \quad (2.7)$$

Proof. By means of Theorem 14.4, Theorem 14.6 and Theorem 15.5 in [1], we can obtain the local existence and extensibility with the form that

if $T_{\max, \varepsilon} < \infty$, then

$$\limsup_{t \rightarrow T_{\max, \varepsilon}} \left\{ \|u_\varepsilon(\cdot, t)\|_{W^{1, \infty}(\Omega)} + \|v_\varepsilon(\cdot, t)\|_{W^{1, \infty}(\Omega)} + \left\| \frac{1}{u_\varepsilon(\cdot, t)} \right\|_{L^\infty(\Omega)} + \left\| \frac{1}{v_\varepsilon(\cdot, t)} \right\|_{L^\infty(\Omega)} \right\} = \infty. \quad (2.8)$$

We assert that (2.8) and (2.4) are equivalent within this framework. It is obvious that (2.4) implies (2.8). On the other hand, after assuming that for some $\varepsilon \in (0, 1)$ there exists $c_1(\varepsilon) > 0$ such that $\|u_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq c_1(\varepsilon)$ for all $t \in (0, T_{\max, \varepsilon})$ with $T_{\max, \varepsilon} < \infty$, we could find positive constants $c_2(\varepsilon)$ and $c_3(\varepsilon)$ such that

$$\|v_\varepsilon(\cdot, t)\|_{W^{1, \infty}(\Omega)} \leq c_2(\varepsilon) \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \quad (2.9)$$

and

$$v_\varepsilon \geq c_3(\varepsilon) \quad \text{in } \Omega \times (0, T_{\max, \varepsilon}) \quad (2.10)$$

by using the same way as the first part of [23, Lemma 2.1]. It is easy to verify that there exists $c_4(\varepsilon) > 0$ such that

$$f(u_\varepsilon) \leq c_4(\varepsilon) u_\varepsilon^{\frac{m-1}{2}} \quad (2.11)$$

either under the case $1 \leq m < 2$, f fulfills (1.3) and (1.4) with $m - 1 \leq \alpha \leq m$ or the case $2 \leq m < 4$, f fulfills (1.3) and (1.5) with $m - 1 \leq \alpha \leq \frac{m}{2} + 1$. Thus, rewriting the first equation in (2.1) in the following form

$$u_{\varepsilon t} = \nabla \cdot a_\varepsilon(x, t, u_\varepsilon, \nabla u_\varepsilon) + b_\varepsilon(x, t, u_\varepsilon, \nabla u_\varepsilon), \quad x \in \Omega, t \in (0, T_{\max, \varepsilon})$$

with

$$\begin{aligned} a_\varepsilon(x, t, u_\varepsilon, \nabla u_\varepsilon) &:= v_\varepsilon(x, t) u_\varepsilon^{m-1}(x, t) \nabla u_\varepsilon(x, t) - f(u_\varepsilon(x, t)) v_\varepsilon(x, t) \nabla v_\varepsilon(x, t) \quad \text{and} \\ b_\varepsilon(x, t) &= \ell u_\varepsilon(x, t) v_\varepsilon(x, t), \quad (x, t) \in \Omega \times (0, T_{\max, \varepsilon}), \end{aligned}$$

the Young inequality combined with (2.9)-(2.11) yields that there exists $c_5(\varepsilon) > 0$ such that

$$\begin{aligned}
a_\varepsilon(x, t, u_\varepsilon, \nabla u_\varepsilon) \cdot \nabla u_\varepsilon &= v_\varepsilon u_\varepsilon^{m-1} |\nabla u_\varepsilon|^2 - f(u_\varepsilon) v_\varepsilon \nabla v_\varepsilon \cdot \nabla u_\varepsilon \\
&\geq c_3(\varepsilon) u_\varepsilon^{m-1} |\nabla u_\varepsilon|^2 - c_4(\varepsilon) u_\varepsilon^{\frac{m-1}{2}} v_\varepsilon |\nabla v_\varepsilon| |\nabla u_\varepsilon| \\
&\geq c_3(\varepsilon) u_\varepsilon^{m-1} |\nabla u_\varepsilon|^2 - c_2^2(\varepsilon) c_4(\varepsilon) u_\varepsilon^{\frac{m-1}{2}} |\nabla u_\varepsilon| \\
&\geq \frac{c_3}{2}(\varepsilon) u_\varepsilon^{m-1} |\nabla u_\varepsilon|^2 - c_5(\varepsilon)
\end{aligned}$$

as well as

$$\begin{aligned}
|a_\varepsilon(x, t, u_\varepsilon, \nabla u_\varepsilon)| &\leq c_2(\varepsilon) u_\varepsilon^{m-1} |\nabla u_\varepsilon| + c_2^2(\varepsilon) c_4(\varepsilon) u_\varepsilon^{\frac{m-1}{2}}, \quad \text{and} \\
|b_\varepsilon(x, t)| &\leq \ell c_1(\varepsilon) c_2(\varepsilon) \quad \text{for all } (x, t) \in \Omega \times (0, T_{max, \varepsilon}),
\end{aligned}$$

which guarantee the existence of $\theta_1 = \theta_1(\varepsilon), \theta_2 = \theta_2(\varepsilon) \in (0, 1)$ such that $u_\varepsilon \in C^{\theta_1, \frac{\theta_1}{2}}(\overline{\Omega} \times [0, T_{max, \varepsilon}])$ and $v_\varepsilon \in C^{2+\theta_2, 1+\frac{\theta_2}{2}}(\overline{\Omega} \times [\frac{1}{4}T_{max, \varepsilon}, T_{max, \varepsilon}])$ provided by the Hölder estimates in [16] and parabolic Schauder theory in [8].

Now we arrange the first equation in (2.1) to the following form

$$u_{\varepsilon t} = A_\varepsilon(x, t) \Delta u_\varepsilon + B_\varepsilon(x, t) \cdot \nabla u_\varepsilon + D_\varepsilon(x, t) u_\varepsilon \quad x \in \Omega, t \in (0, T_{max, \varepsilon}),$$

where

$$\begin{aligned}
A_\varepsilon(x, t) &:= u_\varepsilon^{m-1}(x, t) v_\varepsilon(x, t), \\
B_\varepsilon(x, t) &:= (m-1) u_\varepsilon^{m-2}(x, t) v_\varepsilon(x, t) \nabla u_\varepsilon(x, t) + u_\varepsilon^{m-1}(x, t) \nabla v_\varepsilon(x, t) - f'(u_\varepsilon) v_\varepsilon(x, t) \nabla v_\varepsilon(x, t) \quad \text{and} \\
D_\varepsilon(x, t) &:= -\frac{f(u_\varepsilon(x, t))}{u_\varepsilon(x, t)} v_\varepsilon(x, t) \Delta v_\varepsilon(x, t) - \frac{f(u_\varepsilon(x, t))}{u_\varepsilon(x, t)} |\nabla v_\varepsilon(x, t)|^2 + \ell v_\varepsilon(x, t), \quad (x, t) \in \Omega \times (0, T_{max, \varepsilon}).
\end{aligned}$$

By the assumptions on m, f and α , one can find some positive constant $c_6(\varepsilon)$ satisfying

$$\frac{f(u_\varepsilon(x, t))}{u_\varepsilon(x, t)} \leq c_6(\varepsilon) \quad \text{for all } (x, t) \in \Omega \times (0, T_{max, \varepsilon}),$$

which by means of the boundedness of $v_\varepsilon, \nabla v_\varepsilon$ and Δv_ε entails that there exists $c_7(\varepsilon) > 0$ such that

$$D_\varepsilon \geq -c_7(\varepsilon) \quad \text{in } \Omega \times \left(\frac{1}{4}T_{max, \varepsilon}, T_{max, \varepsilon}\right).$$

Then by the comparison principle we get that

$$u_\varepsilon \geq \inf_{x \in \Omega} u_\varepsilon\left(x, \frac{1}{4}T_{max, \varepsilon}\right) e^{-c_7(\varepsilon) \cdot \frac{3}{4}T_{max, \varepsilon}} \quad \text{in } \Omega \times \left(\frac{1}{4}T_{max, \varepsilon}, T_{max, \varepsilon}\right). \quad (2.12)$$

Furthermore, applying the first order parabolic Hölder regularity theory ([12]), there exists $\theta_3 = \theta_3(\varepsilon) \in (0, 1)$ such that $u_\varepsilon \in C^{1+\theta_3, \frac{1+\theta_3}{2}}(\overline{\Omega} \times [\frac{1}{2}T_{max, \varepsilon}, T_{max, \varepsilon}])$. In particular, we have

$$\|u_\varepsilon(\cdot, t)\|_{W^{1, \infty}(\Omega)} \leq c_8 \quad \text{for all } t \in \left(\frac{1}{2}T_{max, \varepsilon}, T_{max, \varepsilon}\right)$$

with $c_8 = c_8(\varepsilon) > 0$. This together with (2.9), (2.10) and (2.12) shows that (2.8) fails, so that our assertion is identified. We thus obtain (2.4). Finally, (2.5)-(2.7) can be derived easily by some basic integration computations and maximum principle. \blacksquare

From now on, $\ell \geq 0$ is fixed. Without further explicit mentioning, we assume u_0 and v_0 always fulfill (1.6), and accordingly let $(u_\varepsilon, v_\varepsilon)$ and $T_{max,\varepsilon}$ be as yielded by Lemma 2.1.

Now we present two crucial inequalities from [22, Lemma 3.4], which will be utilized frequently in the following context.

Lemma 2.2. *Let $q \geq 2$. Then every $\varphi \in C^2(\bar{\Omega})$ fulfilling $\varphi > 0$ in Ω and $\frac{\partial \varphi}{\partial \nu} = 0$ on $\partial\Omega$ satisfies*

$$\int_{\Omega} \frac{|\nabla \varphi|^{q+2}}{\varphi^{q+1}} \leq (q + \sqrt{n})^2 \int_{\Omega} \frac{|\nabla \varphi|^{q-2}}{\varphi^{q-3}} |D^2 \ln \varphi|^2 \quad (2.13)$$

and

$$\int_{\Omega} \frac{|\nabla \varphi|^{q-2}}{\varphi^{q-1}} |D^2 \varphi|^2 \leq (q + \sqrt{n} + 1)^2 \int_{\Omega} \frac{|\nabla \varphi|^{q-2}}{\varphi^{q-3}} |D^2 \ln \varphi|^2. \quad (2.14)$$

The next lemma is related to the type of functional $\int_{\Omega} v_\varepsilon^{-q+1} |\nabla v_\varepsilon|^q$ for $q \geq 2$, which was first proved in [22, Lemma 3.3] for the general domain with boundary integral $\int_{\partial\Omega} v_\varepsilon^{-q+1} |\nabla v_\varepsilon|^{q-2} \cdot \frac{\partial |\nabla v_\varepsilon|^2}{\partial \nu}$ arising. It was not feasible to treat this boundary integral term as in [22, Lemma 3.5] because the term $\int_0^\infty \int_{\Omega} v_\varepsilon(\cdot, t)$ is unexpected to appear later. So, as previously mentioned in [10], the convexity on Ω in two-dimensional setting is imperative to allow us to disregard the boundary integral.

Lemma 2.3. *Let $\Omega \subset \mathbb{R}^n$ ($n \leq 2$) be a bounded convex domain and $q \geq 2$. Then for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$, there exist $C_1(q) > 0$ and $C_2 > 0$ such that*

$$\frac{d}{dt} \int_{\Omega} \frac{|\nabla v_\varepsilon|^q}{v_\varepsilon^{q-1}} + \frac{q}{2} \int_{\Omega} \frac{|\nabla v_\varepsilon|^{q-2}}{v_\varepsilon^{q-3}} |D^2 \ln v_\varepsilon|^2 + (q-1)^2 \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^q}{v_\varepsilon^{q-1}} \leq C_1(q) \int_{\Omega} u_\varepsilon^{\frac{q+2}{2}} v_\varepsilon \quad (2.15)$$

and

$$\frac{d}{dt} \int_{\Omega} \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_{\Omega} \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} \leq C_2 \int_{\Omega} v_\varepsilon |\nabla u_\varepsilon|^2. \quad (2.16)$$

Proof. From Lemma 3.3 in [22], we have

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \frac{|\nabla v_\varepsilon|^q}{v_\varepsilon^{q-1}} + q \int_{\Omega} \frac{|\nabla v_\varepsilon|^{q-2}}{v_\varepsilon^{q-3}} |D^2 \ln v_\varepsilon|^2 + (q-1)^2 \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^q}{v_\varepsilon^{q-1}} \\ & \leq q(q-2 + \sqrt{n}) \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^{q-2}}{v_\varepsilon^{q-2}} |D^2 v_\varepsilon| + \frac{q}{2} \int_{\partial\Omega} \frac{|\nabla v_\varepsilon|^{q-2}}{v_\varepsilon^{q-1}} \cdot \frac{\partial |\nabla v_\varepsilon|^2}{\partial \nu} \end{aligned} \quad (2.17)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Due to $\frac{\partial |\nabla v_\varepsilon|^2}{\partial \nu} \leq 0$ on $\partial\Omega$ by convexity of Ω ([13]), we can disregard the last term in (2.17). Using (2.13) and (2.14) and the Young inequality, the penultimate term in (2.17) could be estimated as follows

$$\begin{aligned} & q(q-2 + \sqrt{n}) \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^{q-2}}{v_\varepsilon^{q-2}} |D^2 v_\varepsilon| \\ & \leq \frac{q}{4(q + \sqrt{n} + 1)^2} \int_{\Omega} \frac{|\nabla v_\varepsilon|^{q-2}}{v_\varepsilon^{q-1}} |D^2 v_\varepsilon|^2 + c_1 \int_{\Omega} u_\varepsilon^2 \frac{|\nabla v_\varepsilon|^{q-2}}{v_\varepsilon^{q-3}} \end{aligned}$$

$$\begin{aligned}
&\leq \frac{q}{4} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 + c_1 \int_{\Omega} u_{\varepsilon}^2 \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} \\
&\leq \frac{q}{4} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 + \frac{q}{4(q + \sqrt{n})^2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q+2}}{v_{\varepsilon}^{q+1}} + c_2 \int_{\Omega} u_{\varepsilon}^{\frac{q+2}{2}} v_{\varepsilon} \\
&\leq \frac{q}{2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 + c_2 \int_{\Omega} u_{\varepsilon}^{\frac{q+2}{2}} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1)
\end{aligned}$$

with positive constants $c_1 = c_1(q)$ and $c_2 = c_2(q)$, which combined (2.17) gives (2.15).

According to (2.12) in [10], one has

$$\frac{d}{dt} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + 4 \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} \leq -4 \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}^2} (\nabla u_{\varepsilon} \cdot \nabla v_{\varepsilon}) \quad (2.18)$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. Again using the Young inequality and (2.13) with $q = 4$, implies the existence of $c_3 > 0$ such that

$$\begin{aligned}
-4 \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}^2} (\nabla u_{\varepsilon} \cdot \nabla v_{\varepsilon}) &\leq \frac{3}{(4 + \sqrt{n})^2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^6}{v_{\varepsilon}^5} + c_3 \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\
&\leq 3 \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + c_3 \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2
\end{aligned}$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. This inserting (2.18) ensures (2.16). \blacksquare

In the following, we show a boundedness result of a space-time integral derived by a basic calculation.

Lemma 2.4. *Let $\Omega \subset \mathbb{R}^n$ ($n \leq 2$) be a bounded convex domain. For all $K > 0$ there exists $C(K) > 0$ with the property that whenever (1.7) or (1.9) holds, we have*

$$\int_0^{T_{max, \varepsilon}} \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 \leq C(K) \quad \text{for all } \varepsilon \in (0, 1). \quad (2.19)$$

Proof. Testing the second equation in (2.1) by v_{ε}^2 and integrating it on Ω could yield this conclusion, where the boundedness of $\int_{\Omega} v_{0\varepsilon}^3$ ensured by (1.7) or (1.9) is necessary. \blacksquare

At the last of this section, we present the outcomes of some standard L^p testing procedures applied to the first equation in (2.1).

Lemma 2.5. *For all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$, the following estimations hold:*

(i) *If $p > 1$, then we have*

$$\frac{1}{p} \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^p + \frac{p-1}{2} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq \frac{p-1}{2} \int_{\Omega} u_{\varepsilon}^{p-m-1} f^2(u_{\varepsilon}) v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \ell \int_{\Omega} u_{\varepsilon}^p v_{\varepsilon}; \quad (2.20)$$

(ii) *if $0 < p < 1$, then we have*

$$-\frac{1}{p} \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^p + \frac{1-p}{2} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq \frac{1-p}{2} \int_{\Omega} u_{\varepsilon}^{p-m-1} f^2(u_{\varepsilon}) v_{\varepsilon} |\nabla v_{\varepsilon}|^2; \quad (2.21)$$

(iii) if $p < 0$, then we have

$$\frac{d}{dt} \int_{\Omega} u_{\varepsilon}^p + \frac{p(p-1)}{2} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq \frac{p(p-1)}{2} \int_{\Omega} u_{\varepsilon}^{p-m-1} f^2(u_{\varepsilon}) v_{\varepsilon} |\nabla v_{\varepsilon}|^2; \quad (2.22)$$

(iv) if $m = 2$, then we have

$$\frac{d}{dt} \int_{\Omega} u_{\varepsilon} \ln u_{\varepsilon} + \frac{1}{2} \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq \frac{1}{2} \int_{\Omega} \frac{f^2(u_{\varepsilon})}{u_{\varepsilon}^2} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \ln u_{\varepsilon}; \quad (2.23)$$

(v) if $m = 3$, then we have

$$-\frac{d}{dt} \int_{\Omega} \ln u_{\varepsilon} + \frac{1}{2} \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq \frac{1}{2} \int_{\Omega} \frac{f^2(u_{\varepsilon})}{u_{\varepsilon}^4} v_{\varepsilon} |\nabla v_{\varepsilon}|^2. \quad (2.24)$$

Proof. If $p > 1$ or $p < 0$, we simply use the first equation in (2.1) along with the boundary conditions, an integration by parts and the Young inequality to entail that

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^p &= p \int_{\Omega} u_{\varepsilon}^{p-1} \nabla \cdot (u_{\varepsilon}^{m-1} v_{\varepsilon} \nabla u_{\varepsilon}) - p \int_{\Omega} u_{\varepsilon}^{p-1} \nabla \cdot (f(u_{\varepsilon}) v_{\varepsilon} \nabla v_{\varepsilon}) + p\ell \int_{\Omega} u_{\varepsilon}^p v_{\varepsilon} \\ &= -p(p-1) \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + p(p-1) \int_{\Omega} u_{\varepsilon}^{p-2} f(u_{\varepsilon}) v_{\varepsilon} \nabla u_{\varepsilon} \cdot \nabla v_{\varepsilon} + p\ell \int_{\Omega} u_{\varepsilon}^p v_{\varepsilon} \\ &\leq -\frac{p(p-1)}{2} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{p(p-1)}{2} \int_{\Omega} u_{\varepsilon}^{p-m-1} f^2(u_{\varepsilon}) v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + p\ell \int_{\Omega} u_{\varepsilon}^p v_{\varepsilon} \end{aligned} \quad (2.25)$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. When $p > 1$, dividing (2.25) by p establishes (2.20). When $p < 0$, the negativity of the last term in (2.25) guarantees (2.22).

Noting that $p(1-p) > 0$ when $0 < p < 1$, similar to the procedures of getting (2.25), we have

$$\begin{aligned} -\frac{d}{dt} \int_{\Omega} u_{\varepsilon}^p &= -p(1-p) \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + p(1-p) \int_{\Omega} u_{\varepsilon}^{p-2} f(u_{\varepsilon}) v_{\varepsilon} \nabla u_{\varepsilon} \cdot \nabla v_{\varepsilon} - p\ell \int_{\Omega} u_{\varepsilon}^p v_{\varepsilon} \\ &\leq -\frac{p(1-p)}{2} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{p(1-p)}{2} \int_{\Omega} u_{\varepsilon}^{p-m-1} f^2(u_{\varepsilon}) v_{\varepsilon} |\nabla v_{\varepsilon}|^2 \end{aligned}$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$, and then dividing this inequality by p , we get (2.21).

The proof of (2.23) follows a similar approach to the proof of (2.20), and the proof of (2.24) is analogous to that of (2.21). We omit the detailed proofs here. \blacksquare

3 Uniform boundedness of u_{ε} when $n = 1$

In this section, we first introduce a crucial functional inequality coming from [11], which is very useful to address terms of the form $\int_{\Omega} u_{\varepsilon}^{\beta} v_{\varepsilon}$ appearing in Lemma 3.4 and Lemma 3.5 in one-dimensional framework, and is inaccessible in two-dimensional case.

Lemma 3.1. *Let $\Omega \subset \mathbb{R}$, $p \geq 1$ and $r > 1$. Then for all $\eta > 0$ there exists $C(\eta, p, r) > 0$ such that*

$$\|\varphi^{\frac{p+1}{2}} \sqrt{\psi}\|_{L^r(\Omega)}^2 \leq \eta \int_{\Omega} \varphi^{p-1} \psi \varphi_x^2 + \eta \cdot \left\{ \int_{\Omega} \varphi^p \right\} \cdot \int_{\Omega} \frac{\varphi}{\psi} \psi_x^2 + C(\eta, p, r) \cdot \left\{ \int_{\Omega} \varphi \right\}^p \cdot \int_{\Omega} \varphi \psi \quad (3.1)$$

is valid for arbitrary nonnegative function $\varphi \in C^1(\overline{\Omega})$ and positive function $\psi \in C^1(\overline{\Omega})$.

The next is about an a priori bound on the gradient of the second solution component in (2.1). As a direct application of this, in light of Lemma 3.1, the terms with the form $\int_{\Omega} u_{\varepsilon}^{\gamma} v_{\varepsilon} v_{\varepsilon x}^2$ appearing in Lemma 3.3 and Lemma 3.5 could be dealt.

Lemma 3.2. *Let $\Omega \subset \mathbb{R}$ be an open interval and $K > 0$. Then there exists $C(K) > 0$ with the property that whenever (1.7) holds, then*

$$\|v_{\varepsilon x}(\cdot, t)\|_{L^4(\Omega)} \leq C(K) \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1). \quad (3.2)$$

Proof. From (2.6), we have

$$\|u_{\varepsilon}(\cdot, t)\|_{L^1(\Omega)} \leq \int_{\Omega} u_{0\varepsilon} + \ell \int_{\Omega} v_{0\varepsilon} \leq \|u_0\|_{L^1(\Omega)} + |\Omega|(1 + \ell \cdot \|v_0\|_{L^{\infty}(\Omega)})$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. The variation-of-constants representation and well-known smoothing properties of Neumann heat semigroup $(e^{t\Delta})_{t \geq 0}$ ([19, Lemma 1.3]) indicate (3.2). \blacksquare

In the course of utilizing Lemma 3.1 to handle the integral $\int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon}$ arising from testing the second summand in (1.12), another unfavorable term $\int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2$ will appear, which suggests us how to choose appropriate p in (1.12) at a first stage. The following is a more concrete outcome based on Lemma 2.5.

Lemma 3.3. *Let $\Omega \subset \mathbb{R}$ be an open interval. Then if one of the following cases holds:*

(i) $1 \leq m < 2$, f fulfills (1.3) and (1.4) with $m - 1 \leq \alpha \leq m$;

(ii) $2 \leq m < 4$, f fulfills (1.3) and (1.5) with $m - 1 \leq \alpha \leq m$,

there exist positive constants c and C such that for each $\varepsilon \in (0, 1)$, the function $\mathcal{F}_{\varepsilon}$ defined on $t \in (0, T_{max, \varepsilon})$ by letting

$$\mathcal{F}_{\varepsilon}(t) := \begin{cases} \int_{\Omega} u_{\varepsilon}^{3-m} & \text{when } 1 \leq m < 2 \quad \text{or} \quad 3 < m < 4, \\ \int_{\Omega} u_{\varepsilon} \ln u_{\varepsilon} & \text{when } m = 2, \\ - \int_{\Omega} \ln u_{\varepsilon} & \text{when } m = 3, \\ - \int_{\Omega} u_{\varepsilon}^{3-m} & \text{when } 2 < m < 3 \end{cases}$$

satisfies

$$\mathcal{F}_{\varepsilon}'(t) + c \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 \leq C \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} v_{\varepsilon x}^2 + C \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} + C \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + C \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad (3.3)$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$.

Proof. We prove this assertion in five cases.

Case 1: $1 \leq m < 2$. Noting that $(1+s)^{2\alpha-2} \leq c_1 + c_1 s^{2\alpha-2}$ is valid for all $s \geq 0$ with some positive constant c_1 , it yields by taking $p = 3 - m > 1$ in (2.20) that

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^{3-m} + \frac{(2-m)(3-m)}{2} \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 \\ & \leq \frac{(2-m)(3-m)}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{4-2m} (u_{\varepsilon} + 1)^{2\alpha-2} v_{\varepsilon} v_{\varepsilon x}^2 + (3-m)\ell \int_{\Omega} u_{\varepsilon}^{3-m} v_{\varepsilon} \\ & \leq c_1 C_f^2 \int_{\Omega} u_{\varepsilon}^{4-2m} v_{\varepsilon} v_{\varepsilon x}^2 + c_1 C_f^2 \int_{\Omega} u_{\varepsilon}^{2-2m+2\alpha} v_{\varepsilon} v_{\varepsilon x}^2 + 2\ell \int_{\Omega} u_{\varepsilon}^{3-m} v_{\varepsilon} \end{aligned} \quad (3.4)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Because $0 < 4 - 2m \leq 2$ due to $1 \leq m < 2$ and $0 \leq 2 - 2m + 2\alpha \leq 2$ due to $m - 1 \leq \alpha \leq m$, using the Young inequality, the first two terms on the right side of (3.4) could be estimated by

$$c_1 C_f^2 \int_{\Omega} u_{\varepsilon}^{4-2m} v_{\varepsilon} v_{\varepsilon x}^2 + c_1 C_f^2 \int_{\Omega} u_{\varepsilon}^{2-2m+2\alpha} v_{\varepsilon} v_{\varepsilon x}^2 \leq 2c_1 C_f^2 \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} v_{\varepsilon x}^2 + 2c_1 C_f^2 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Another application of the Young inequality together with the fact $1 < 3 - m \leq 2$ shows that the last term in (3.4) has the following estimation

$$2\ell \int_{\Omega} u_{\varepsilon}^{3-m} v_{\varepsilon} \leq 2\ell \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} + 2\ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1).$$

Inserting the above two inequalities into (3.4) shows that if we take $c := \frac{(2-m)(3-m)}{2}$ and $C := \max\{2c_1 C_f^2, 2\ell\}$, then (3.3) holds.

Case 2: $m = 2$. Recalling (2.23), it follows from the fact $\ln u_{\varepsilon} \leq u_{\varepsilon}$ and the Young inequality in light of $0 \leq 2\alpha - 2 \leq 2$ that for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$,

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} u_{\varepsilon} \ln u_{\varepsilon} + \frac{1}{2} \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 & \leq \frac{1}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{2\alpha-2} v_{\varepsilon} v_{\varepsilon x}^2 + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \ln u_{\varepsilon} \\ & \leq \frac{1}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} v_{\varepsilon x}^2 + \frac{1}{2} C_f^2 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} + \ell \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon}, \end{aligned}$$

which indicates (3.3) by taking $c := \frac{1}{2}$ and $C := \max\{\frac{1}{2} C_f^2, \ell\}$.

Case 3: $2 < m < 3$. Taking $p := 3 - m \in (0, 1)$ in (2.21) and making use of the Young inequality once more with $0 \leq 2 - 2m + 2\alpha \leq 2$ due to $m - 1 \leq \alpha \leq m$, we get

$$\begin{aligned} -\frac{d}{dt} \int_{\Omega} u_{\varepsilon}^{3-m} + \frac{(m-2)(3-m)}{2} \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 & \leq \frac{(m-2)(3-m)}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{2-2m+2\alpha} v_{\varepsilon} v_{\varepsilon x}^2 \\ & \leq C_f^2 \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} v_{\varepsilon x}^2 + C_f^2 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 \end{aligned}$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Then (3.7) is obtained by taking $c := \frac{(2-m)(3-m)}{2}$ and $C := C_f^2$.

Case 4: $m = 3$. It is easy to verify from (2.24) that if $m = 3$, one has

$$-\frac{d}{dt} \int_{\Omega} \ln u_{\varepsilon} + \frac{1}{2} \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 \leq \frac{1}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{2\alpha-4} v_{\varepsilon} v_{\varepsilon x}^2 \leq \frac{1}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} v_{\varepsilon x}^2 + \frac{1}{2} C_f^2 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$, where the last inequality is provided by the Young inequality with the help of $0 \leq 2\alpha - 4 \leq 2$ due to $2 \leq \alpha \leq 3$. This implies (3.7) by letting $c := \frac{1}{2}$ and $C := \frac{1}{2}C_f^2$.

Case 5: $3 < m < 4$. The proof of this case can be derived in a very similar, even simpler, way as in case 1. We point out that the only difference is “taking $p = 3 - m < 0$ in (2.22)” instead of “taking $p = 3 - m > 1$ in (2.20)” when we use Lemma 2.5. \blacksquare

Based on Lemmas 3.1-3.3, we are able to derive the following conclusion by making use of the energy functional (1.12).

Lemma 3.4. *Let $\Omega \subset \mathbb{R}$ be an open interval, and $K > 0$ with the property that (1.7) is valid. Then if one of the following cases holds:*

- (i) $1 \leq m < 2$, f fulfills (1.3) and (1.4) with $m - 1 \leq \alpha \leq m$;
 - (ii) $2 \leq m < 3$, f fulfills (1.3) and (1.5) with $m - 1 \leq \alpha \leq m$;
 - (iii) $3 \leq m < 4$, f fulfills (1.3) and (1.5) with $m - 1 \leq \alpha \leq m$ and $u_0 > 0$ in $\bar{\Omega}$,
- one can find $C(K) > 0$ such that

$$\int_0^{T_{max,\varepsilon}} \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} + \int_0^{T_{max,\varepsilon}} \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 \leq C(K) \quad \text{for all } \varepsilon \in (0, 1). \quad (3.5)$$

Proof. Taking $q = 2$, (2.15) in Lemma 2.3 has the following one-dimensional version

$$\frac{d}{dt} \int_{\Omega} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} + \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} \leq c_1 \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1) \quad (3.6)$$

with $c_1 > 0$. Letting $\mathcal{F}_{\varepsilon}$ be as in Lemma 3.3, it follows from (3.3) and (3.6) that there exist $c_2 > 0$ and $c_3 > 0$ fulfilling

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} + \mathcal{F}'_{\varepsilon}(t) + c_2 \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 + \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} \\ & \leq c_3 \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} v_{\varepsilon x}^2 + c_3 \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} + c_3 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + c_3 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \end{aligned} \quad (3.7)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$, where the Cauchy-Schwarz inequality and (3.2) imply the existence of $c_4 = c_4(K) > 0$ such that

$$\begin{aligned} c_3 \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} v_{\varepsilon x}^2 + c_3 \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} & \leq c_3 \|v_{\varepsilon x}\|_{L^4(\Omega)}^2 \cdot \|u_{\varepsilon} v_{\varepsilon}^{\frac{1}{2}}\|_{L^4(\Omega)}^2 + c_3 |\Omega|^{\frac{1}{2}} \|u_{\varepsilon} v_{\varepsilon}^{\frac{1}{2}}\|_{L^4(\Omega)}^2 \\ & \leq c_4 \|u_{\varepsilon} v_{\varepsilon}^{\frac{1}{2}}\|_{L^4(\Omega)}^2 \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1), \end{aligned} \quad (3.8)$$

and we can further invoke Lemma 3.1 with $p := 1$ and $r := 4$ and use (2.6) to find $c_5 = c_5(K) := (K + 1 + \ell K)|\Omega|$, $c_6 = c_6(K) > 0$ satisfying

$$\begin{aligned} c_4 \|u_{\varepsilon} v_{\varepsilon}^{\frac{1}{2}}\|_{L^4(\Omega)}^2 & \leq \frac{c_2}{2} \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 + \frac{1}{2c_5} \cdot \left\{ \int_{\Omega} u_{\varepsilon} \right\} \cdot \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} + c_6 \cdot \left\{ \int_{\Omega} u_{\varepsilon} \right\} \cdot \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \\ & \leq \frac{c_2}{2} \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 + \frac{1}{2} \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} + c_5 c_6 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \end{aligned} \quad (3.9)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Substituting (3.8) and (3.9) into (3.7), we get

$$\frac{d}{dt} \int_{\Omega} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} + \mathcal{F}'_{\varepsilon}(t) + \frac{c_2}{2} \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 + \frac{1}{2} \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} \leq c_3 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + (c_3 + c_5 c_6) \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad (3.10)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$.

A combination of (1.7) with Lemma 2.4 and (2.7) entails that there exists $c_7 = c_7(K) > 0$ satisfying

$$c_3 \int_0^{T_{max,\varepsilon}} \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + (c_3 + c_5 c_6) \int_0^{T_{max,\varepsilon}} \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \leq c_7 \quad \text{for all } \varepsilon \in (0, 1),$$

as a consequence of which, upon an integration of (3.10), we claim from (1.7) that

$$\begin{aligned} & \frac{c_2}{2} \int_0^t \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 + \frac{1}{2} \int_0^t \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} \\ & \leq c_3 \int_0^t \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + (c_3 + c_5 c_6) \int_0^t \int_{\Omega} u_{\varepsilon} v_{\varepsilon} + \int_{\Omega} \frac{v_{0x}^2}{v_0} + \mathcal{F}_{\varepsilon}(0) - \mathcal{F}_{\varepsilon}(t) \\ & \leq c_7 + K^3 \cdot |\Omega| + \mathcal{F}_{\varepsilon}(0) - \mathcal{F}_{\varepsilon}(t) \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1). \end{aligned}$$

Thus, to complete the proof, it suffices to show that $\mathcal{F}_{\varepsilon}(0) - \mathcal{F}_{\varepsilon}(t)$ always has an upper bound. Indeed, when $1 \leq m < 2$, because of (1.7) and the positivity of $3 - m$, we see that

$$\mathcal{F}_{\varepsilon}(0) - \mathcal{F}_{\varepsilon}(t) = \int_{\Omega} u_{0\varepsilon}^{3-m} - \int_{\Omega} u_{\varepsilon}^{3-m} \leq \int_{\Omega} u_{0\varepsilon}^{3-m} \leq (K+1)^{3-m} \cdot |\Omega|.$$

When $m = 2$, the pointwise estimate $-u_{\varepsilon} \ln u_{\varepsilon} \leq \frac{1}{e}$ and (1.7) infer that

$$\mathcal{F}_{\varepsilon}(0) - \mathcal{F}_{\varepsilon}(t) = \int_{\Omega} u_{0\varepsilon} \ln u_{0\varepsilon} - \int_{\Omega} u_{\varepsilon} \ln u_{\varepsilon} \leq (K+1) \ln(K+1) \cdot |\Omega| + \frac{|\Omega|}{e}.$$

When $2 < m < 3$, it follows from (2.6) and the Young inequality with the fact $0 < 3 - m < 1$ that

$$\mathcal{F}_{\varepsilon}(0) - \mathcal{F}_{\varepsilon}(t) = - \int_{\Omega} u_{0\varepsilon}^{3-m} + \int_{\Omega} u_{\varepsilon}^{3-m} \leq \int_{\Omega} u_{\varepsilon}^{3-m} \leq \int_{\Omega} u_{\varepsilon} + |\Omega| \leq c_5 + |\Omega|.$$

When $m = 3$, the definition of $u_{0\varepsilon}$ and the strict positivity of u_0 entail the existence of $c_8 > 0$ independent on ε fulfilling $u_{0\varepsilon} > c_8$. Thus, when $m = 3$, due to the fact $\ln u_{\varepsilon} \leq u_{\varepsilon}$, we have

$$\mathcal{F}_{\varepsilon}(0) - \mathcal{F}_{\varepsilon}(t) = - \int_{\Omega} \ln u_{0\varepsilon} + \int_{\Omega} \ln u_{\varepsilon} \leq - \int_{\Omega} \ln u_{0\varepsilon} + \int_{\Omega} u_{\varepsilon} \leq - \ln c_8 \cdot |\Omega| + c_5$$

and when $3 < m < 4$, since $3 - m$ is negative, we have

$$\mathcal{F}_{\varepsilon}(0) - \mathcal{F}_{\varepsilon}(t) = \int_{\Omega} u_{0\varepsilon}^{3-m} - \int_{\Omega} u_{\varepsilon}^{3-m} \leq \int_{\Omega} u_{0\varepsilon}^{3-m} \leq c_8^{3-m} \cdot |\Omega|.$$

The proof is finished. ■

We are now prepared to derive the main result of this section.

Lemma 3.5. Let $\Omega \subset \mathbb{R}$ be an open interval, $p > 2$ and $K > 0$ with the property that (1.7) is valid.

Then there exists $C(p, K) > 0$ such that if one of the following cases holds:

(i) $1 \leq m < 2$, f fulfills (1.3) and (1.4) with $m - 1 \leq \alpha \leq m$;

(ii) $2 \leq m < 3$, f fulfills (1.3) and (1.5) with $m - 1 \leq \alpha \leq \frac{m}{2} + 1$;

(iii) $3 \leq m < 4$, f fulfills (1.3) and (1.5) with $m - 1 \leq \alpha \leq \frac{m}{2} + 1$ and $u_0 > 0$ in $\overline{\Omega}$,

for any choice of $\varepsilon \in (0, 1)$, one has

$$\int_{\Omega} u_{\varepsilon}^p(\cdot, t) \leq C(p, K) \quad \text{for all } t \in (0, T_{max, \varepsilon}). \quad (3.11)$$

Proof. According to Lemma 3.2, (2.6) and (1.7), we can fix $c_1 = c_1(K) > 0$ such that

$$\int_{\Omega} v_{\varepsilon x}^4 + \int_{\Omega} u_{\varepsilon} \leq c_1 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1), \quad (3.12)$$

and we note that it is sufficient to prove (3.11) for $p \geq 4$ because of the Young inequality.

Now we firstly prove it in the case $1 \leq m < 2$. From (2.20), there exists $c_2 = c_2(p) > 0$ such that

$$\begin{aligned} & \frac{1}{p} \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^p + \frac{p-1}{2} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} u_{\varepsilon x}^2 \\ & \leq \frac{p-1}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{p-m+1} (u_{\varepsilon} + 1)^{2\alpha-2} v_{\varepsilon} v_{\varepsilon x}^2 + \ell \int_{\Omega} u_{\varepsilon}^p v_{\varepsilon} \\ & \leq c_2 \int_{\Omega} u_{\varepsilon}^{p-m-1+2\alpha} v_{\varepsilon} v_{\varepsilon x}^2 + c_2 \int_{\Omega} u_{\varepsilon}^{p-m+1} v_{\varepsilon} v_{\varepsilon x}^2 + \ell \int_{\Omega} u_{\varepsilon}^p v_{\varepsilon} \end{aligned} \quad (3.13)$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. Making use of the Cauchy-Schwarz inequality, Lemma 3.1 and (3.12), we can find $c_3 = c_3(p, K) > 0$ satisfying

$$\begin{aligned} c_2 \int_{\Omega} u_{\varepsilon}^{p-m-1+2\alpha} v_{\varepsilon} v_{\varepsilon x}^2 & \leq c_2 \cdot \left\{ \int_{\Omega} v_{\varepsilon x}^4 \right\}^{\frac{1}{2}} \cdot \left\{ \int_{\Omega} u_{\varepsilon}^{2(p-m-1+2\alpha)} v_{\varepsilon}^2 \right\}^{\frac{1}{2}} \\ & \leq c_2 c_1^{\frac{1}{2}} \|u_{\varepsilon}^{\frac{p-m-1+2\alpha}{2}} v_{\varepsilon}^{\frac{1}{2}}\|_{L^4(\Omega)}^2 \\ & \leq \frac{p-1}{4} \int_{\Omega} u_{\varepsilon}^{p-m-3+2\alpha} v_{\varepsilon} u_{\varepsilon x}^2 + \left\{ \int_{\Omega} u_{\varepsilon}^{p-m-2+2\alpha} \right\} \cdot \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} \\ & \quad + c_3 \cdot \left\{ \int_{\Omega} u_{\varepsilon} \right\}^{p-m-2+2\alpha} \cdot \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \end{aligned}$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$, where $0 \leq p - m - 3 + 2\alpha \leq p + m - 3$ due to $m - 1 \leq \alpha \leq m$ and $p \geq 4$ enables us to use the Young inequality to estimate

$$\int_{\Omega} u_{\varepsilon}^{p-m-3+2\alpha} v_{\varepsilon} u_{\varepsilon x}^2 \leq \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} u_{\varepsilon x}^2 + \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1)$$

and another application of the Young inequality together with the fact $1 \leq p - m - 2 + 2\alpha < p$ because of $m - 1 \leq \alpha \leq m$ implies that

$$\int_{\Omega} u_{\varepsilon}^{p-m-2+2\alpha} \leq \int_{\Omega} u_{\varepsilon}^p + |\Omega| \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1),$$

which infer that with $c_4 = |\Omega|$ and $c_5 = c_3 c_1^{p-m-2+2\alpha}$, for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$ we have

$$\begin{aligned} c_2 \int_{\Omega} u_{\varepsilon}^{p-m-1+2\alpha} v_{\varepsilon} v_{\varepsilon x}^2 &\leq \frac{p-1}{4} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} u_{\varepsilon x}^2 + \frac{p-1}{4} \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 \\ &\quad + \left\{ \int_{\Omega} u_{\varepsilon}^p \right\} \cdot \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} + c_4 \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} + c_5 \int_{\Omega} u_{\varepsilon} v_{\varepsilon}. \end{aligned} \quad (3.14)$$

Similarly, thanks to $0 < p - m + 1, p + m - 2 < p \leq p + m - 1$ in the case $1 \leq m < 2$, we could invoke the Young inequality several times and Lemma 3.1 once more to obtain that

$$\begin{aligned} &c_2 \int_{\Omega} u_{\varepsilon}^{p-m+1} v_{\varepsilon} v_{\varepsilon x}^2 + \ell \int_{\Omega} u_{\varepsilon}^p v_{\varepsilon} \\ &\leq c_2 \int_{\Omega} u_{\varepsilon}^{p+m-1} v_{\varepsilon} v_{\varepsilon x}^2 + c_2 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + \ell \int_{\Omega} u_{\varepsilon}^{p+m-1} v_{\varepsilon} + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \\ &\leq \left(c_2 c_1^{\frac{1}{2}} + \ell |\Omega|^{\frac{1}{2}} \right) \cdot \|u_{\varepsilon}^{\frac{p+m-1}{2}} v_{\varepsilon}^{\frac{1}{2}}\|_{L^4(\Omega)}^2 + c_2 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \\ &\leq \frac{p-1}{4} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} u_{\varepsilon x}^2 + \left\{ \int_{\Omega} u_{\varepsilon}^{p+m-2} \right\} \cdot \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} \\ &\quad + c_6 \cdot \left\{ \int_{\Omega} u_{\varepsilon} \right\}^{p+m-2} \cdot \int_{\Omega} u_{\varepsilon} v_{\varepsilon} + c_2 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \\ &\leq \frac{p-1}{4} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} u_{\varepsilon x}^2 + \left\{ \int_{\Omega} u_{\varepsilon}^p + c_4 \right\} \cdot \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} + c_2 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + c_7 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \end{aligned} \quad (3.15)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$ with $c_6 = c_6(p) > 0$ and $c_7 = c_6 c_1^{p+m-2} + \ell$. This together with (3.13) and (3.14) shows that

$$\begin{aligned} \frac{1}{p} \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^p &\leq \frac{p-1}{4} \int_{\Omega} v_{\varepsilon} u_{\varepsilon x}^2 + 2 \left\{ \int_{\Omega} u_{\varepsilon}^p \right\} \cdot \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} + 2c_4 \int_{\Omega} u_{\varepsilon} \frac{v_{\varepsilon x}^2}{v_{\varepsilon}} \\ &\quad + c_2 \int_{\Omega} v_{\varepsilon} v_{\varepsilon x}^2 + (c_5 + c_7) \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1). \end{aligned}$$

If we write

$$y_{\varepsilon}(t) := \int_{\Omega} u_{\varepsilon}^p(\cdot, t), \quad t \in [0, T_{max,\varepsilon})$$

and

$$g_{\varepsilon}(t) := 2p \int_{\Omega} u_{\varepsilon}(\cdot, t) \frac{v_{\varepsilon x}^2(\cdot, t)}{v_{\varepsilon}(\cdot, t)}, \quad t \in (0, T_{max,\varepsilon})$$

as well as

$$\begin{aligned} h_{\varepsilon}(t) &:= \frac{p(p-1)}{4} \int_{\Omega} v_{\varepsilon}(\cdot, t) u_{\varepsilon x}^2(\cdot, t) + 2pc_4 \int_{\Omega} u_{\varepsilon}(\cdot, t) \frac{v_{\varepsilon x}^2(\cdot, t)}{v_{\varepsilon}(\cdot, t)} \\ &\quad + pc_2 \int_{\Omega} v_{\varepsilon}(\cdot, t) v_{\varepsilon x}^2(\cdot, t) + p(c_5 + c_7) \int_{\Omega} u_{\varepsilon}(\cdot, t) v_{\varepsilon}(\cdot, t), \quad t \in (0, T_{max,\varepsilon}), \end{aligned}$$

then we get that

$$y'_{\varepsilon}(t) \leq g_{\varepsilon}(t) y'_{\varepsilon}(t) + h_{\varepsilon}(t) \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1)$$

with the property that there is $c_8 = c_8(p, K) > 0$ such that

$$\int_0^t g_\varepsilon(s) ds \leq c_8 \quad \text{and} \quad \int_0^t h_\varepsilon(s) ds \leq c_8 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1)$$

provided by Lemma 2.4, Lemma 3.4, (2.7) and (1.7). An ODE comparison argument in conjunction with the fact that $y_\varepsilon(0) = \int_\Omega (u_0 + \varepsilon)^p \leq (K + 1)^p |\Omega|$ will indicate (3.11).

In the case $2 \leq m < 4$, invoking $f(u_\varepsilon) \leq C_f u_\varepsilon^\alpha$ into (2.20), then (3.13) becomes

$$\frac{1}{p} \frac{d}{dt} \int_\Omega u_\varepsilon^p + \frac{p-1}{2} \int_\Omega u_\varepsilon^{p+m-3} v_\varepsilon u_{\varepsilon x}^2 \leq c_2 \int_\Omega u_\varepsilon^{p-m-1+2\alpha} v_\varepsilon v_{\varepsilon x}^2 + \ell \int_\Omega u_\varepsilon^p v_\varepsilon$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$, where on the basis of $\frac{m}{2} + 1 \leq m$, the term $c_2 \int_\Omega u_\varepsilon^{p-m-1+2\alpha} v_\varepsilon v_{\varepsilon x}^2$ can be estimated in the same way used in obtaining (3.14), and we point out that the rightmost term $\ell \int_\Omega u_\varepsilon^p v_\varepsilon$ will be handled in a slightly different manner from (3.15). Indeed, if $2 \leq m < 4$, we have $0 < p-1 \leq p+m-3$, which allows us to use Lemma 3.1 and the Young inequality to find $c_9 = c_9(p) > 0$ such that for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$,

$$\begin{aligned} \ell \int_\Omega u_\varepsilon^p v_\varepsilon &\leq \ell \int_\Omega u_\varepsilon^{p+1} v_\varepsilon + \ell \int_\Omega u_\varepsilon v_\varepsilon \\ &\leq \ell |\Omega|^{\frac{1}{2}} \cdot \|u_\varepsilon^{\frac{p+1}{2}} v_\varepsilon^{\frac{1}{2}}\|_{L^4(\Omega)}^2 + \ell \int_\Omega u_\varepsilon v_\varepsilon \\ &\leq \frac{p-1}{4} \int_\Omega u_\varepsilon^{p-1} v_\varepsilon u_{\varepsilon x}^2 + \left\{ \int_\Omega u_\varepsilon^p \right\} \cdot \int_\Omega u_\varepsilon \frac{v_{\varepsilon x}^2}{v_\varepsilon} + c_9 \cdot \left\{ \int_\Omega u_\varepsilon \right\}^p \cdot \int_\Omega u_\varepsilon v_\varepsilon + \ell \int_\Omega u_\varepsilon v_\varepsilon \\ &\leq \frac{p-1}{4} \int_\Omega u_\varepsilon^{p+m-3} v_\varepsilon u_{\varepsilon x}^2 + \frac{p-1}{4} \int_\Omega v_\varepsilon u_{\varepsilon x}^2 + \left\{ \int_\Omega u_\varepsilon^p \right\} \cdot \int_\Omega u_\varepsilon \frac{v_{\varepsilon x}^2}{v_\varepsilon} + (c_9 c_1^p + \ell) \int_\Omega u_\varepsilon v_\varepsilon. \end{aligned}$$

Trivially applying the procedures after (3.15) thereby proves (3.11) in the case $2 \leq m < 4$. ■

4 Uniform boundedness of u_ε when $n = 2$

4.1 Some crucial estimations

As shown in Section 3, the differential inequality (3.1) plays a critical role in helping us utilize the degenerate diffusive action of the first equation in (2.1). However, this approach does not work in two-dimensional space because the embedding $W^{1,1}(\Omega) \hookrightarrow L^2(\Omega)$ is only continuous, not compact. To make sure that the term of the form $\int_\Omega \varphi^{p-1} \psi |\nabla \varphi|^2$ involves an arbitrarily small coefficient, we adopt the following functional inequality ([10, Lemma 3.1]).

Lemma 4.1. *Let $\Omega \subset \mathbb{R}^2$, $p > 0$ and $r \geq 2$. Then for all $\eta > 0$ there exists $C(\eta, p, r) > 0$ such that*

$$\left\| \varphi^{\frac{p+1}{2}} \sqrt{\psi} \right\|_{L^r(\Omega)}^2 \leq \eta \int_\Omega \varphi^{p-1} \psi |\nabla \varphi|^2 + \eta \int_\Omega \varphi^{p+1} \frac{|\nabla \psi|^2}{\psi} + C(\eta, p, r) \cdot \left\{ \int_\Omega \varphi \right\}^p \cdot \int_\Omega \varphi \psi \quad (4.1)$$

is valid for arbitrary nonnegative function $\varphi \in C^1(\overline{\Omega})$ and positive function $\psi \in C^1(\overline{\Omega})$.

Proof. When $p \geq 1$, for any given $\eta > 0$, we take

$$\eta_1 \equiv \eta_1(\eta, p) \leq \min \left\{ \frac{1}{1 + |\Omega|^{\frac{r-2}{r}}}, \frac{2\eta}{(p+1)^2} \right\}.$$

Because $W^{1,2}(\Omega)$ is compactly embedded into $L^r(\Omega)$ and the latter space is embedded into $L^1(\Omega)$ with the continuous sense, the Ehrling lemma implies that for η_1 , there exists a positive constant $C_1 = C_1(\eta_1, r) > 0$ such that

$$\begin{aligned} \left\| \varphi^{\frac{p+1}{2}} \sqrt{\psi} \right\|_{L^r(\Omega)}^2 &\leq \frac{\eta_1}{8} \left\| \nabla \left(\varphi^{\frac{p+1}{2}} \sqrt{\psi} \right) \right\|_{W^{1,2}(\Omega)}^2 + C_1 \left\| \varphi^{\frac{p+1}{2}} \sqrt{\psi} \right\|_{L^1(\Omega)}^2 \\ &\leq \frac{\eta_1}{4} \left\| \nabla \left(\varphi^{\frac{p+1}{2}} \sqrt{\psi} \right) \right\|_{L^2(\Omega)}^2 + \frac{\eta_1}{4} \left\| \left(\varphi^{\frac{p+1}{2}} \sqrt{\psi} \right) \right\|_{L^2(\Omega)}^2 + C_1 \left\| \varphi^{\frac{p+1}{2}} \sqrt{\psi} \right\|_{L^1(\Omega)}^2 \\ &\leq \frac{\eta_1}{4} \left\| \nabla \left(\varphi^{\frac{p+1}{2}} \sqrt{\psi} \right) \right\|_{L^2(\Omega)}^2 + \frac{1}{4} \left\| \left(\varphi^{\frac{p+1}{2}} \sqrt{\psi} \right) \right\|_{L^r(\Omega)}^2 + C_1 \left\| \varphi^{\frac{p+1}{2}} \sqrt{\psi} \right\|_{L^1(\Omega)}^2, \end{aligned} \quad (4.2)$$

where the last inequality is due to the Hölder inequality. Then we can use the Young inequality to interpolate the norm in $L^1(\Omega)$ between $L^r(\Omega)$ and $L^{\frac{2}{p+1}}(\Omega)$, that is, with $C_2 = C_2(\eta_1, p, r) > 0$ we have

$$C_1 \left\| \varphi^{\frac{p+1}{2}} \sqrt{\psi} \right\|_{L^1(\Omega)}^2 \leq \frac{1}{2} \left\| \varphi^{\frac{p+1}{2}} \sqrt{\psi} \right\|_{L^r(\Omega)}^2 + C_2 \left\| \varphi^{\frac{p+1}{2}} \sqrt{\psi} \right\|_{L^{\frac{2}{p+1}}(\Omega)}^2,$$

which combined with (4.2) indicates that for any $p \geq 1$, we have

$$\left\| \varphi^{\frac{p+1}{2}} \sqrt{\psi} \right\|_{L^r(\Omega)}^2 \leq \eta_1 \left\| \nabla \left(\varphi^{\frac{p+1}{2}} \sqrt{\psi} \right) \right\|_{L^2(\Omega)}^2 + 4C_2 \left\| \varphi^{\frac{p+1}{2}} \sqrt{\psi} \right\|_{L^{\frac{2}{p+1}}(\Omega)}^2$$

and thereby proves (4.1) by some basic calculus.

In the case when $0 < p < 1$, using the Ehrling lemma on the basis of the fact that $W^{1,2}(\Omega)$ is compactly embedded into $L^r(\Omega)$, while the latter space trivially lies in $L^{\frac{2}{p+1}}(\Omega)$, could directly derive the conclusion. \blacksquare

Parallel to that in Section 3, the key of our analysis in this section is also to control the unfavorable terms $\int_{\Omega} u_{\varepsilon}^{\beta} v_{\varepsilon}$ and $\int_{\Omega} u_{\varepsilon}^{\gamma} v_{\varepsilon} |\nabla v_{\varepsilon}|^2$. Fortunately, an observation is that in view of Lemma 4.1, this objective can be achieved when β and γ fall within the appropriate intermediate range, respectively. Additionally, we could catch some rationales why we take p^* in (1.14) arbitrarily close to $3 - m$, but not equal to the critical value here.

Lemma 4.2. *Let $\Omega \subset \mathbb{R}^2$ and $K > 0$ with the property that (1.9) is valid. If $\kappa \in (-1, 0)$, then for any $\beta \in [1, \kappa + 3)$ and $\eta > 0$, there exists $C(K, \beta, \eta) > 0$ such that*

$$\int_{\Omega} u_{\varepsilon}^{\beta} v_{\varepsilon} \leq \eta \int_{\Omega} u_{\varepsilon}^{\kappa} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \eta \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + C(K, \beta, \eta) \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad (4.3)$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$.

Proof. We begin with establishing the validity of (4.3) for $\beta \in [\kappa + 2, \kappa + 3)$. According to (2.6) and (1.9), we can find $c_1 = c_1(K) > 0$ such that

$$\|u_\varepsilon(\cdot, t)\|_{L^1(\Omega)} \leq c_1 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1). \quad (4.4)$$

Taking

$$\vartheta := \frac{1}{3 + \kappa - \beta} \quad \text{and} \quad \vartheta^* := \frac{\vartheta}{\vartheta - 1} = \frac{1}{\beta - \kappa - 2},$$

it follows from $\beta \in [\kappa + 2, \kappa + 3)$ that

$$\vartheta \geq 1 \quad \text{and} \quad \vartheta^* > 1.$$

Thus, for given $\eta > 0$, we may use the Hölder inequality, Lemma 4.1 and (4.4) to conclude that there exists $c_2 = c_2(K, \beta, \eta) > 0$ satisfying

$$\begin{aligned} \int_{\Omega} u_\varepsilon^\beta v_\varepsilon &= \int_{\Omega} \left(u_\varepsilon^{\frac{\kappa+2}{2}} v_\varepsilon^{\frac{1}{2}} \right)^2 u_\varepsilon^{\beta-\kappa-2} \\ &\leq \left\| u_\varepsilon^{\frac{\kappa+2}{2}} v_\varepsilon^{\frac{1}{2}} \right\|_{L^{2\vartheta}(\Omega)}^2 \cdot \left\| u_\varepsilon^{\beta-\kappa-2} \right\|_{L^{\vartheta^*}(\Omega)} \\ &\leq (c_1^{\frac{1}{\vartheta^*}} + 1) \left\| u_\varepsilon^{\frac{\kappa+2}{2}} v_\varepsilon^{\frac{1}{2}} \right\|_{L^{2\vartheta}(\Omega)}^2 \\ &\leq \frac{\eta}{2} \int_{\Omega} u_\varepsilon^\kappa v_\varepsilon |\nabla u_\varepsilon|^2 + \int_{\Omega} u_\varepsilon^{\kappa+2} \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} + c_2 \int_{\Omega} u_\varepsilon v_\varepsilon \end{aligned} \quad (4.5)$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. In treating the second term on the right side of (4.5), we first apply Lemma 2.3 with $q := 4$ to see the existence of $c_3 > 0$ such that

$$\int_{\Omega} \frac{|\nabla v_\varepsilon|^6}{v_\varepsilon^5} \leq c_3 \int_{\Omega} \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1).$$

Then let

$$\chi := -\frac{2}{\kappa} \quad \text{and} \quad \chi^* := \frac{\chi}{\chi - 1} = \frac{2}{\kappa + 2},$$

where our assumption $\kappa \in (-1, 0)$ ensures that

$$\chi > 1 \quad \text{and} \quad \chi^* > 1,$$

whence by means of the Young inequality and the Hölder inequality, Lemma 4.1 and (4.4) are applied again so as to deduce that with $c_i = c_i(K, \beta, \eta) > 0$, $i = 4, 5$, we have

$$\begin{aligned} \int_{\Omega} u_\varepsilon^{\kappa+2} \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} &\leq \frac{\eta}{2c_3} \int_{\Omega} \frac{|\nabla v_\varepsilon|^6}{v_\varepsilon^5} + c_4 \int_{\Omega} u_\varepsilon^{\frac{3}{2}\kappa+3} v_\varepsilon \\ &\leq \frac{\eta}{2} \int_{\Omega} \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + c_4 \int_{\Omega} \left(u_\varepsilon^{\frac{\kappa+2}{2}} v_\varepsilon^{\frac{1}{2}} \right)^2 u_\varepsilon^{\frac{1}{2}\kappa+1} \end{aligned}$$

$$\begin{aligned}
&\leq \frac{\eta}{2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + c_4 \left\| u_{\varepsilon}^{\frac{\kappa+2}{2}} v_{\varepsilon}^{\frac{1}{2}} \right\|_{L^{2\chi}(\Omega)}^2 \cdot \left\| u_{\varepsilon}^{\frac{1}{2}\kappa+1} \right\|_{L^{\chi^*}(\Omega)} \\
&\leq \frac{\eta}{2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + c_1^{\frac{1}{\chi^*}} c_4 \left\| u_{\varepsilon}^{\frac{\kappa+2}{2}} v_{\varepsilon}^{\frac{1}{2}} \right\|_{L^{2\chi}(\Omega)}^2 \\
&\leq \frac{\eta}{2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + \frac{\eta}{4} \int_{\Omega} u_{\varepsilon}^{\kappa} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{1}{2} \int_{\Omega} u_{\varepsilon}^{\kappa+2} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} + c_5 \int_{\Omega} u_{\varepsilon} v_{\varepsilon}
\end{aligned}$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$, which indicates that

$$\int_{\Omega} u_{\varepsilon}^{\kappa+2} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} \leq \eta \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + \frac{\eta}{2} \int_{\Omega} u_{\varepsilon}^{\kappa} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + 2c_5 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad (4.6)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Inserting (4.6) into (4.5) will yield (4.3).

Now we illustrate that (4.3) also holds for $\beta \in [1, \kappa + 2)$. Using the Young inequality, we see that

$$\int_{\Omega} u_{\varepsilon}^{\beta} v_{\varepsilon} \leq \int_{\Omega} u_{\varepsilon}^{\kappa+2} v_{\varepsilon} + \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1),$$

which in conjunction with the fact that (4.3) holds for $\kappa + 2$ completes the proof. \blacksquare

The following conclusion is a byproduct of the above lemma.

Lemma 4.3. *Let $\Omega \subset \mathbb{R}^2$ and $K > 0$ with the property that (1.9) is valid. If $\kappa \in (-1, 0)$, then for any $\gamma \in [0, \frac{\kappa}{2} + 2)$ and $\eta > 0$, there exists $C = C(K, \gamma, \eta) > 0$ such that*

$$\begin{aligned}
\int_{\Omega} u_{\varepsilon}^{\gamma} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 &\leq \eta \int_{\Omega} u_{\varepsilon}^{\kappa} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \eta \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \eta \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 \\
&\quad + \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + C(K, \gamma, \eta) \int_{\Omega} u_{\varepsilon} v_{\varepsilon}
\end{aligned} \quad (4.7)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$.

Proof. If $\gamma \in [0, 1]$, using the Young inequality twice and relying on (1.9), we see that

$$\begin{aligned}
\int_{\Omega} u_{\varepsilon}^{\gamma} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 &\leq \int_{\Omega} u_{\varepsilon} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\
&\leq \eta \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \frac{1}{4\eta} \int_{\Omega} u_{\varepsilon} v_{\varepsilon}^5 + \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\
&\leq \eta \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \frac{K^4}{4\eta} \int_{\Omega} u_{\varepsilon} v_{\varepsilon} + \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2
\end{aligned}$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. This shows that (4.7) holds for $\gamma \in [0, 1]$. On the other hand, when $\gamma \in (1, \frac{\kappa}{2} + 2)$, it is evident that $2\gamma - 1 \in (1, \kappa + 3)$. Therefore, it is available to make use of the Young inequality, (1.9) and Lemma 4.2 to see that with $c_1 = c_1(K, \gamma, \eta) > 0$, we have

$$\begin{aligned}
\int_{\Omega} u_{\varepsilon}^{\gamma} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 &\leq \eta \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \frac{K^4}{4\eta} \int_{\Omega} u_{\varepsilon}^{2\gamma-1} v_{\varepsilon} \\
&\leq \eta \int_{\Omega} u_{\varepsilon}^{\kappa} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \eta \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + \eta \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + c_1 \int_{\Omega} u_{\varepsilon} v_{\varepsilon}
\end{aligned}$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Thereby, we finish the proof. \blacksquare

4.2 Uniform L^{p_0} bounds on u_ε for some $p_0 > 1$

Let us first derive some basic information on the time evolution of the first two summands in (1.14).

Lemma 4.4. *Let $\Omega \subset \mathbb{R}^2$ be a bounded convex domain, and $K > 0$ with the property that (1.9) is valid. Then if one of the following cases holds:*

- (i) $1 \leq m < 2$, f fulfills (1.3) and (1.4) with $m - 1 < \alpha < m$;
 - (ii) $2 \leq m < 4$, f fulfills (1.3) and (1.5) with $m - 1 < \alpha < m$,
- one can find positive constants c and C such that for each $\varepsilon \in (0, 1)$, the function \mathcal{G}_ε defined on $t \in (0, T_{\max, \varepsilon})$ by letting

$$\mathcal{G}_\varepsilon(t) := \begin{cases} c \int_{\Omega} u_\varepsilon^{3-m} + \int_{\Omega} \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} & \text{when } 1 \leq m < 2 \text{ or } 3 < m < 4, \\ c \int_{\Omega} u_\varepsilon \ln u_\varepsilon + \int_{\Omega} \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} & \text{when } m = 2, \\ -c \int_{\Omega} \ln u_\varepsilon + \int_{\Omega} \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} & \text{when } m = 3, \\ -c \int_{\Omega} u_\varepsilon^{3-m} + \int_{\Omega} \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} & \text{when } 2 < m < 3 \end{cases}$$

satisfies for all $t \in (0, T_{\max, \varepsilon})$ and $\varepsilon \in (0, 1)$,

$$\begin{aligned} \mathcal{G}'_\varepsilon(t) + \int_{\Omega} \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_{\Omega} v_\varepsilon |\nabla u_\varepsilon|^2 \\ \leq C \int_{\Omega} u_\varepsilon^{2-2m+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 + C \int_{\Omega} u_\varepsilon^2 v_\varepsilon + C \int_{\Omega} u_\varepsilon v_\varepsilon \quad \text{when } m = 1 \text{ or } m = 2, \end{aligned} \quad (4.8)$$

$$\begin{aligned} \mathcal{G}'_\varepsilon(t) + \int_{\Omega} \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_{\Omega} v_\varepsilon |\nabla u_\varepsilon|^2 \\ \leq C \int_{\Omega} u_\varepsilon^{2-2m+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 + C \int_{\Omega} u_\varepsilon^{4-2m} v_\varepsilon |\nabla v_\varepsilon|^2 + C \int_{\Omega} u_\varepsilon^{3-m} v_\varepsilon \quad \text{when } 1 < m < 2 \end{aligned} \quad (4.9)$$

and

$$\begin{aligned} \mathcal{G}'_\varepsilon(t) + \int_{\Omega} \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_{\Omega} v_\varepsilon |\nabla u_\varepsilon|^2 \\ \leq C \int_{\Omega} u_\varepsilon^{2-2m+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 \quad \text{when } 2 < m < 4. \end{aligned} \quad (4.10)$$

Proof. We first apply (2.16) in Lemma 2.3 to find $c_1 > 0$ such that

$$\frac{d}{dt} \int_{\Omega} \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_{\Omega} \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} \leq c_1 \int_{\Omega} v_\varepsilon |\nabla u_\varepsilon|^2 \quad (4.11)$$

for all $t \in (0, T_{\max, \varepsilon})$ and $\varepsilon \in (0, 1)$. When $1 \leq m < 2$, taking $p := 3 - m > 1$ in (2.20) and dividing both sides by $\frac{2-m}{2(c_1+1)}$, it follows that

$$\frac{2(c_1+1)}{(2-m)(3-m)} \frac{d}{dt} \int_{\Omega} u_\varepsilon^{3-m} + (c_1+1) \int_{\Omega} v_\varepsilon |\nabla u_\varepsilon|^2$$

$$\leq (c_1 + 1)C_f^2 \int_{\Omega} u_{\varepsilon}^{4-2m}(u_{\varepsilon} + 1)^{2\alpha-2} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \frac{2\ell(c_1 + 1)}{2 - m} \int_{\Omega} u_{\varepsilon}^{3-m} v_{\varepsilon}$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Then if we write

$$\mathcal{G}_{\varepsilon}(t) := \frac{2(c_1 + 1)}{(2 - m)(3 - m)} \int_{\Omega} u_{\varepsilon}^{3-m} + \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3},$$

the above inequality together with (4.11) deduces that

$$\begin{aligned} \mathcal{G}'_{\varepsilon}(t) + \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\ \leq (c_1 + 1)C_f^2 \int_{\Omega} u_{\varepsilon}^{4-2m}(u_{\varepsilon} + 1)^{2\alpha-2} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \frac{2\ell(c_1 + 1)}{2 - m} \int_{\Omega} u_{\varepsilon}^{3-m} v_{\varepsilon} \end{aligned} \quad (4.12)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Obviously, (4.9) can be derived directly by using the fact that $(s + 1)^{2\alpha-2} \leq c_2 s^{2\alpha-2} + c_2$ holds for any $s \geq 0$ with some $c_2 > 0$.

When $m = 1$, the inequality $(s + 1)^{2\alpha-2} \leq s^{2\alpha-2}$ holds for any $s \geq 0$ due to our assumption $\alpha < 1$, which in conjunction with (4.12) proves (4.8) for the case $m = 1$. Moreover, for this $\mathcal{G}_{\varepsilon}$, it is easy to verify from (4.11) and (2.22) that (4.10) holds for the case $3 < m < 4$.

When $m = 2$, for the fixed c_1 , we could get from (2.23) that

$$\begin{aligned} 2(c_1 + 1) \frac{d}{dt} \int_{\Omega} u_{\varepsilon} \ln u_{\varepsilon} + (c_1 + 1) \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\ \leq (c_1 + 1)C_f^2 \int_{\Omega} u_{\varepsilon}^{2\alpha-2} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + 2(c_1 + 1)\ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} + 2(c_1 + 1)\ell \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} \end{aligned}$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$, which together with (4.11) implies (4.8) for the case $m = 2$ by taking

$$\mathcal{G}_{\varepsilon}(t) := 2(c_1 + 1) \int_{\Omega} u_{\varepsilon} \ln u_{\varepsilon} + \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3}.$$

Analogously, based on (4.11), we can prove (4.10) for the case $2 < m < 3$ by letting

$$\mathcal{G}_{\varepsilon}(t) := -\frac{2(c_1 + 1)}{(m - 2)(3 - m)} \int_{\Omega} u_{\varepsilon}^{3-m} + \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3},$$

taking $p := 3 - m \in (0, 1)$ in (2.21) and dividing both sides by $\frac{m-2}{2(c_1+1)}$. Relying on (2.24) and (4.11), we can present (4.10) for the case $m = 3$ with

$$\mathcal{G}_{\varepsilon}(t) := -2(c_1 + 1) \int_{\Omega} \ln u_{\varepsilon} + \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3}.$$

The proof is thereby finished. ■

From the above lemma, we are readily able to derive the L^{p_0} bounds for u_{ε} with some $p_0 > 1$ in the case $1 \leq m < 2$.

Lemma 4.5. *Let $\Omega \subset \mathbb{R}^2$ be a bounded convex domain and $1 \leq m < 2$. Suppose that f fulfills (1.3) and (1.4) with $m - 1 < \alpha < m$ and $K > 0$ with the property that (1.9) is valid. Then there exist $p_0 > 1$ and $C(K, p_0) > 0$ such that*

$$\int_{\Omega} u_{\varepsilon}^{p_0}(\cdot, t) \leq C(K, p_0) \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \text{ and } \varepsilon \in (0, 1)$$

and

$$\int_0^{T_{\max, \varepsilon}} \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_0^{T_{\max, \varepsilon}} \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq C(K, p_0) \quad \text{for all } \varepsilon \in (0, 1).$$

Proof. Let $\mathcal{G}_{\varepsilon}$ be defined as in Lemma 4.4. We see that there is $c_1 > 0$ such that for all $t \in (0, T_{\max, \varepsilon})$ and $\varepsilon \in (0, 1)$, when $m = 1$,

$$\begin{aligned} \mathcal{G}'_{\varepsilon}(t) &+ \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\ &\leq c_1 \int_{\Omega} u_{\varepsilon}^{2-2m+2\alpha} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + c_1 \int_{\Omega} u_{\varepsilon}^2 v_{\varepsilon} + c_1 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \end{aligned} \quad (4.13)$$

and when $1 < m < 2$,

$$\begin{aligned} \mathcal{G}'_{\varepsilon}(t) &+ \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\ &\leq c_1 \int_{\Omega} u_{\varepsilon}^{4-2m} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + c_1 \int_{\Omega} u_{\varepsilon}^{2-2m+2\alpha} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + c_1 \int_{\Omega} u_{\varepsilon}^{3-m} v_{\varepsilon}. \end{aligned} \quad (4.14)$$

Noting that $m - 1 < 1 < 3 - m$ due to $m < 2$, and that $\max\{m + 1 - 2\alpha, 3 - 5m + 4\alpha\} < 3 - m$ due to our assumption $m - 1 < \alpha < m$, thus it is possible to pick $p_0 > 1$ satisfying

$$\max\{m + 1 - 2\alpha, 3 - 5m + 4\alpha\} < p_0 < 3 - m, \quad (4.15)$$

from which we can check that

$$0 < p_0 - m + 2\alpha - 1 < 2 - 2m + 2\alpha < \frac{p_0}{2} + \frac{m}{2} + \frac{1}{2} \quad (4.16)$$

and moreover, together with the fact that $3 - m \leq 3m - 1$ because of $m \geq 1$, we have

$$0 < p_0 - m + 1 < \frac{p_0}{2} + \frac{m}{2} + \frac{1}{2}. \quad (4.17)$$

For the fixed $p_0 > 1$, we employ (2.20) to find $c_2 > 0$ fulfilling

$$\begin{aligned} &\frac{1}{p_0} \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^{p_0} + \frac{p_0 - 1}{2} \int_{\Omega} u_{\varepsilon}^{p_0+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\ &\leq \frac{p_0 - 1}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{p_0-m+1} (1 + u_{\varepsilon})^{2\alpha-2} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \ell \int_{\Omega} u_{\varepsilon}^{p_0} v_{\varepsilon} \\ &\leq c_2 \int_{\Omega} u_{\varepsilon}^{p_0-m+2\alpha-1} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + c_2 \int_{\Omega} u_{\varepsilon}^{p_0-m+1} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \ell \int_{\Omega} u_{\varepsilon}^{p_0} v_{\varepsilon} \end{aligned} \quad (4.18)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Observing that our restrictions $1 < p_0 < 3 - m$ and $m \geq 1$ warrant $\kappa := p_0 + m - 3 \in (-1, 0)$, thus we may invoke Lemma 4.3 along with (4.16) and (4.17) to infer the existence of $c_3 = c_3(K, p_0) > 0$ such that

$$\begin{aligned} & c_1 \int_{\Omega} u_{\varepsilon}^{2-2m+2\alpha} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + c_2 \int_{\Omega} u_{\varepsilon}^{p_0-m+2\alpha-1} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + c_2 \int_{\Omega} u_{\varepsilon}^{p_0-m+1} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 \\ & \leq \frac{p_0-1}{8} \int_{\Omega} u_{\varepsilon}^{p_0+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{1}{2} \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \frac{1}{4} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 \\ & \quad + \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + c_3 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1). \end{aligned} \quad (4.19)$$

For the taken κ , it is clear that $1 < p_0 < \kappa + 3$ and $1 < 3 - m < \kappa + 3$, which ensure that Lemma 4.2 is available to estimate

$$\begin{aligned} & \ell \int_{\Omega} u_{\varepsilon}^{p_0} v_{\varepsilon} + c_1 \int_{\Omega} u_{\varepsilon}^{3-m} v_{\varepsilon} \\ & \leq \frac{p_0-1}{8} \int_{\Omega} u_{\varepsilon}^{p_0+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{1}{4} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + c_4 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \end{aligned} \quad (4.20)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$ with $c_4 = c_4(K, p_0) > 0$.

Combining (4.13), (4.18), (4.19) and (4.20), it gives that when $m = 1$,

$$\begin{aligned} & \mathcal{G}'_{\varepsilon}(t) + \frac{1}{p_0} \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^{p_0} + \frac{1}{2} \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\ & \leq \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + (c_1 + c_3 + c_4) \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1). \end{aligned} \quad (4.21)$$

And we note that if $1 < m < 2$, the fact $7 - 5m < 3 - m$ warrants the possibility to pick p_0 not only satisfying (4.15), but also fulfilling

$$p_0 > 7 - 5m.$$

Therefore, we readily find that

$$0 < 4 - 2m < \frac{p_0}{2} + \frac{m}{2} + \frac{1}{2}.$$

Another application of Lemma 4.3 may furnish positive constant $c_5 = c_5(K, p_0) > 0$ such that

$$\begin{aligned} c_1 \int_{\Omega} u_{\varepsilon}^{4-2m} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 & \leq \frac{p_0-1}{8} \int_{\Omega} u_{\varepsilon}^{p_0+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{1}{4} \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \frac{1}{4} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 \\ & \quad + \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + c_5 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1), \end{aligned}$$

which combined with (4.14), (4.18), (4.19) and (4.20) gives that when $1 < m < 2$,

$$\begin{aligned} & \mathcal{G}'_{\varepsilon}(t) + \frac{1}{p_0} \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^{p_0} + \frac{1}{4} \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\ & \leq 2 \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + (c_3 + c_4 + c_5) \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1). \end{aligned} \quad (4.22)$$

Integrating (4.21) and (4.22) in time respectively, Lemma 2.4, (2.7) and (1.9) show at once that whenever $1 \leq m < 2$, with some $c_6 = c_6(K, p_0) > 0$ and $c_7 = c_7(K, p_0) > 0$, we have

$$\begin{aligned}
& \frac{1}{p_0} \int_{\Omega} u_{\varepsilon}^{p_0} + \frac{1}{4} \int_0^t \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_0^t \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\
& \leq 2 \int_0^t \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + (c_1 + c_3 + c_4 + c_5) \int_0^t \int_{\Omega} u_{\varepsilon} v_{\varepsilon} + \mathcal{G}_{\varepsilon}(0) - \mathcal{G}_{\varepsilon}(t) \\
& \leq c_6 + c_7 \int_{\Omega} (u_0 + 1)^{3-m} + \int_{\Omega} \frac{|\nabla v_0|^4}{v_0^3} \\
& \leq c_6 + c_7 (K + 1)^{3-m} |\Omega| + K^5 |\Omega| \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1).
\end{aligned}$$

Thereby the proof is finished. ■

Differing from the previous lemma, when $2 \leq m \leq 4$, p^* belongs to $(-2, 1)$ rather than $p^* > 1$. Therefore, merely relying on functional (1.14) is insufficient to obtain our desired estimate directly. For this, we additionally consider another component $\int_{\Omega} u^{p_0}$ with some $p_0 > 1$.

Lemma 4.6. *Let $\Omega \subset \mathbb{R}^2$ be a bounded convex domain and $2 \leq m < 3$. Suppose that f fulfills (1.3) and (1.5) with $m - 1 < \alpha < \frac{m}{2} + 1$ and $K > 0$ with the property that (1.9) is valid. Then there exist $0 < p_* < 1$, $p_0 > 1$ and $C(K, p_*, p_0) > 0$ such that*

$$\int_{\Omega} u_{\varepsilon}^{p_0}(\cdot, t) \leq C(K, p_*, p_0) \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1)$$

and

$$\int_0^{T_{max, \varepsilon}} \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_0^{T_{max, \varepsilon}} \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq C(K, p_*, p_0) \quad \text{for all } \varepsilon \in (0, 1).$$

Proof. Firstly, we note that our assumption $m - 1 < \alpha < \frac{m}{2} + 1$ makes it possible to find $p_* > 0$ satisfying

$$\max\{m + 1 - 2\alpha, 4\alpha - 3m - 1\} < p_* < 3 - m \leq 1. \tag{4.23}$$

Evidently, we have $\frac{p_*}{2} + \frac{3}{2}m + \frac{3}{2} - 2\alpha > 1$ and $p_* + m > 1$, so that it is also possible to fix p_0 such that

$$1 < p_0 < \min\left\{\frac{p_*}{2} + \frac{3}{2}m + \frac{3}{2} - 2\alpha, p_* + m\right\}. \tag{4.24}$$

Taking $\kappa := p_* + m - 3$, it is easy to verify from (4.23) that

$$\kappa \in (-1, 0). \tag{4.25}$$

Moreover, in view of $\alpha > m - 1$, it follows from (4.24) that

$$0 < p_0 - m - 1 + 2\alpha < \frac{p_*}{2} + \frac{m}{2} + \frac{1}{2} =: \frac{\kappa}{2} + 2, \tag{4.26}$$

and due to our choice of p_* in (4.23), we have

$$0 < p_* - m - 1 + 2\alpha < 2 - 2m + 2\alpha < \frac{p_*}{2} + \frac{m}{2} + \frac{1}{2} =: \frac{\kappa}{2} + 2, \tag{4.27}$$

where the last inequality uses that the restriction $m \geq 2$ warrants that $p_* > 4\alpha - 3m - 1 \geq 3 - 5m + 4\alpha$.

Next, let \mathcal{G}_ε be defined as in Lemma 4.4. From (4.8) and (4.10), we can conclude that when $2 \leq m < 3$, there is $c_1 > 0$ such that

$$\begin{aligned} \mathcal{G}'_\varepsilon(t) &+ \int_\Omega \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + \int_\Omega u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_\Omega v_\varepsilon |\nabla u_\varepsilon|^2 \\ &\leq c_1 \int_\Omega u_\varepsilon^{2-2m+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 + c_1 \int_\Omega u_\varepsilon^2 v_\varepsilon + c_1 \int_\Omega u_\varepsilon v_\varepsilon \end{aligned} \quad (4.28)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. For the above fixed p_* and p_0 , we draw on (2.21) and (2.20) respectively to see that for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$, we have

$$-\frac{1}{p_*} \frac{d}{dt} \int_\Omega u_\varepsilon^{p_*} + \frac{1-p_*}{2} \int_\Omega u_\varepsilon^{p_*+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 \leq \frac{1-p_*}{2} C_f^2 \int_\Omega u_\varepsilon^{p_*-m-1+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2$$

and

$$\frac{1}{p_0} \frac{d}{dt} \int_\Omega u_\varepsilon^{p_0} \leq \frac{p_0-1}{2} C_f^2 \int_\Omega u_\varepsilon^{p_0-m-1+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 + \ell \int_\Omega u_\varepsilon^{p_0} v_\varepsilon,$$

which in conjunction with (4.28) yield that

$$\begin{aligned} \mathcal{G}'_\varepsilon(t) &+ \frac{1}{p_0} \frac{d}{dt} \int_\Omega u_\varepsilon^{p_0} - \frac{1}{p_*} \frac{d}{dt} \int_\Omega u_\varepsilon^{p_*} + \frac{1-p_*}{2} \int_\Omega u_\varepsilon^{p_*+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 \\ &+ \int_\Omega \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + \int_\Omega u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_\Omega v_\varepsilon |\nabla u_\varepsilon|^2 \\ &\leq \frac{1-p_*}{2} C_f^2 \int_\Omega u_\varepsilon^{p_*-m-1+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 + \frac{p_0-1}{2} C_f^2 \int_\Omega u_\varepsilon^{p_0-m-1+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 \\ &+ c_1 \int_\Omega u_\varepsilon^{2-2m+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 + \ell \int_\Omega u_\varepsilon^{p_0} v_\varepsilon + c_1 \int_\Omega u_\varepsilon^2 v_\varepsilon + c_1 \int_\Omega u_\varepsilon v_\varepsilon \end{aligned} \quad (4.29)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Now in view of (4.25), (4.26) and (4.27), Lemma 4.3 becomes applicable to detect that there is $c_2 = c_2(K, p_*, p_0) > 0$ fulfilling

$$\begin{aligned} &\frac{1-p_*}{2} C_f^2 \int_\Omega u_\varepsilon^{p_*-m-1+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 + \frac{p_0-1}{2} C_f^2 \int_\Omega u_\varepsilon^{p_0-m-1+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 + c_1 \int_\Omega u_\varepsilon^{2-2m+2\alpha} v_\varepsilon |\nabla v_\varepsilon|^2 \\ &\leq \frac{1-p_*}{4} \int_\Omega u_\varepsilon^{p_*+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + \frac{1}{2} \int_\Omega u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \frac{1}{2} \int_\Omega \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 \\ &+ \int_\Omega v_\varepsilon |\nabla v_\varepsilon|^2 + c_2 \int_\Omega u_\varepsilon v_\varepsilon \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1). \end{aligned} \quad (4.30)$$

On the other hand, it is obvious from (4.24) that $1 < p_0 < p_* + m := \kappa + 3$ and $2 < p_* + m := \kappa + 3$, which together with (4.25) enable us to employ Lemma 4.2 to find $c_3 = c_3(K, p_*, p_0) > 0$ such that

$$\ell \int_\Omega u_\varepsilon^{p_0} v_\varepsilon + c_1 \int_\Omega u_\varepsilon^2 v_\varepsilon \leq \frac{1-p_*}{4} \int_\Omega u_\varepsilon^{p_*+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + \frac{1}{2} \int_\Omega \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + c_3 \int_\Omega u_\varepsilon v_\varepsilon \quad (4.31)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. Inserting (4.30) and (4.31) into (4.29) shows that

$$\mathcal{G}'_\varepsilon(t) + \frac{1}{p_0} \frac{d}{dt} \int_\Omega u_\varepsilon^{p_0} - \frac{1}{p_*} \frac{d}{dt} \int_\Omega u_\varepsilon^{p_*} + \frac{1}{2} \int_\Omega u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_\Omega v_\varepsilon |\nabla u_\varepsilon|^2$$

$$\leq \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + (c_1 + c_2 + c_3) \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \text{ and } \varepsilon \in (0, 1). \quad (4.32)$$

Before proceeding, similar to the assertions in the proof of Lemma 3.4, we claim that there is $c_4 = c_4(K) > 0$ such that $\mathcal{G}_{\varepsilon}(0) - \mathcal{G}_{\varepsilon}(t) \leq c_4$ for all $t \in (0, T_{\max, \varepsilon})$ and $\varepsilon \in (0, 1)$. Indeed when $m = 2$, with some positive constant $c_5 > 0$ we have

$$\mathcal{G}_{\varepsilon}(0) - \mathcal{G}_{\varepsilon}(t) = c_5 \int_{\Omega} u_{0\varepsilon} \ln u_{0\varepsilon} - c_5 \int_{\Omega} u_{\varepsilon} \ln u_{\varepsilon} + \int_{\Omega} \frac{|\nabla v_0|^4}{v_0^3} \leq c_5(K+1)^2 |\Omega| + \frac{c_5}{e} |\Omega| + K^5 |\Omega|,$$

and when $2 < m < 3$, using (2.6), (1.9) and the Young inequality along with $0 < 3 - m < 1$, there exists $c_6 > 0$ such that

$$\mathcal{G}_{\varepsilon}(0) - \mathcal{G}_{\varepsilon}(t) \leq c_6 \int_{\Omega} u_{\varepsilon}^{3-m} + \int_{\Omega} \frac{|\nabla v_0|^4}{v_0^3} \leq c_6 \left(\int_{\Omega} u_{\varepsilon} + |\Omega| \right) + K^5 |\Omega| \leq (c_6(\ell K + K + 2) + K^5) |\Omega|.$$

So, integrating (4.32) from 0 to t , by means of (2.7) and Lemma 2.4, once more using (2.6), (1.9) and the Young inequality along with the fact that $0 < p_* < 1$, we have

$$\begin{aligned} & \frac{1}{p_0} \int_{\Omega} u_{\varepsilon}^{p_0} + \frac{1}{2} \int_0^t \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_0^t \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\ & \leq \int_0^t \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + c_7 \int_0^t \int_{\Omega} u_{\varepsilon} v_{\varepsilon} + \mathcal{G}_{\varepsilon}(0) - \mathcal{G}_{\varepsilon}(t) + \frac{1}{p_*} \int_{\Omega} u_{\varepsilon}^{p_*} + \frac{1}{p_0} \int_{\Omega} u_{0\varepsilon}^{p_0} \\ & \leq c_8 + c_4 + \frac{\ell K + K + 2}{p_*} |\Omega| + \frac{K^{p_0}}{p_0} |\Omega| \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \text{ and } \varepsilon \in (0, 1), \end{aligned}$$

where $c_7 = c_1 + c_2 + c_3$ and $c_8 = c_8(K) > 0$. The proof is complete. \blacksquare

Lemma 4.7. *Let $\Omega \subset \mathbb{R}^2$ be a bounded convex domain, $3 \leq m < 4$ and $K > 0$. Suppose that f fulfills (1.3) and (1.5) with $m - 1 < \alpha < \frac{m}{2} + 1$. Then there exist $p_* < 0$, $p_0 > 1$, $C_1(K, p_*, p_0) > 0$ and $C_2(K, p_*) > 0$ such that if not only (1.9) holds but also $u_0 > 0$ in $\overline{\Omega}$, one has*

$$\int_{\Omega} u_{\varepsilon}^{p_0}(\cdot, t) \leq C_1(K, p_*, p_0) \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \text{ and } \varepsilon \in (0, 1)$$

and

$$\int_0^{T_{\max, \varepsilon}} \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_0^{T_{\max, \varepsilon}} \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq C_2(K, p_*) \quad \text{for all } \varepsilon \in (0, 1).$$

Proof. Let $\mathcal{G}_{\varepsilon}$ be defined as in Lemma 4.4 with $c_1 > 0$ when $m = 3$, and with $c_2 > 0$ when $3 < m < 4$. We firstly show that there exists some positive constant $c_3 = c_3(K)$ such that

$$\mathcal{G}_{\varepsilon}(0) - \mathcal{G}_{\varepsilon}(t) \leq c_3 \quad \text{for all } t \in (0, T_{\max, \varepsilon}) \text{ and } \varepsilon \in (0, 1). \quad (4.33)$$

Indeed, the strict positivity of u_0 in $\overline{\Omega}$ warrants the existence of $c_4 > 0$ independent on ε fulfilling $u_{0\varepsilon} \geq c_4$. Then in light of (2.6) and (1.9), when $m = 3$,

$$\mathcal{G}_{\varepsilon}(0) - \mathcal{G}_{\varepsilon}(t) = -c_1 \int_{\Omega} \ln u_{0\varepsilon} + c_1 \int_{\Omega} \ln u_{\varepsilon} + \int_{\Omega} \frac{|\nabla v_0|^4}{v_0^3}$$

$$\begin{aligned}
&\leq -c_1 \int_{\Omega} \ln u_{0\varepsilon} + c_1 \int_{\Omega} u_{\varepsilon} + \int_{\Omega} \frac{|\nabla v_0|^4}{v_0^3} \\
&\leq -c_1 \ln c_4 \cdot |\Omega| + c_1(\ell K + K + 1)|\Omega| + K^5|\Omega|,
\end{aligned}$$

and when $3 < m < 4$, since $3 - m$ is negative and u_{ε} is nonnegative, we have

$$\mathcal{G}_{\varepsilon}(0) - \mathcal{G}_{\varepsilon}(t) = c_2 \int_{\Omega} u_{0\varepsilon}^{3-m} - c_2 \int_{\Omega} u_{\varepsilon}^{3-m} + \int_{\Omega} \frac{|\nabla v_0|^4}{v_0^3} \leq c_2 c_4^{3-m} |\Omega| + K^5 |\Omega|.$$

Moreover, relying on (4.10), we can find $c_5 > 0$ satisfying

$$\mathcal{G}'_{\varepsilon}(t) + \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq c_5 \int_{\Omega} u_{\varepsilon}^{2+2\alpha-2m} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 \quad (4.34)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$.

Then similar to the discussions in the proof of Lemma 4.6, we can pick $p_* \in (2 - m, 3 - m)$ and $p_0 > 1$ such that the following three conditions

$$0 < p_* + 2\alpha - m - 1 < 2 + 2\alpha - 2m < \frac{p_*}{2} + \frac{m}{2} + \frac{1}{2}, \quad (4.35)$$

and

$$0 < p_0 + 2\alpha - m - 1 < \frac{p_*}{2} + \frac{m}{2} + \frac{1}{2} = \frac{p_* + m - 3}{2} + 2 \quad (4.36)$$

as well as

$$1 < p_0 < p_* + m = (p_* + m - 3) + 3 \quad (4.37)$$

are fulfilled simultaneously. For the fixed p_* , recalling (2.22), we have

$$\frac{d}{dt} \int_{\Omega} u_{\varepsilon}^{p_*} + \frac{p_*(p_* - 1)}{2} \int_{\Omega} u_{\varepsilon}^{p_*+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq \frac{p_*(p_* - 1)}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{p_*+2\alpha-m-1} v_{\varepsilon} |\nabla v_{\varepsilon}|^2$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. This in conjunction with (4.34) yields that

$$\begin{aligned}
&\mathcal{G}'_{\varepsilon}(t) + \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^{p_*} + \frac{p_*(p_* - 1)}{2} \int_{\Omega} u_{\varepsilon}^{p_*+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\
&\quad + \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 + \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\
&\leq c_5 \int_{\Omega} u_{\varepsilon}^{2+2\alpha-2m} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \frac{p_*(p_* - 1)}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{p_*+2\alpha-m-1} v_{\varepsilon} |\nabla v_{\varepsilon}|^2
\end{aligned}$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$, whereas thanks to (4.35), applying Lemma 4.3 will result in that there is $c_6 = c_6(K, p_*) > 0$ fulfilling

$$\begin{aligned}
&c_5 \int_{\Omega} u_{\varepsilon}^{2+2\alpha-2m} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \frac{p_*(p_* - 1)}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{p_*+2\alpha-m-1} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 \\
&\leq \frac{p_*(p_* - 1)}{4} \int_{\Omega} u_{\varepsilon}^{p_*+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{1}{2} \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3} + \frac{1}{2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} |D^2 \ln v_{\varepsilon}|^2 \\
&\quad + \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + c_6 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1),
\end{aligned}$$

so that

$$\begin{aligned}
& \mathcal{G}'_\varepsilon(t) + \frac{d}{dt} \int_\Omega u_\varepsilon^{p_*} + \frac{p_*(p_* - 1)}{4} \int_\Omega u_\varepsilon^{p_*+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 \\
& \quad + \frac{1}{2} \int_\Omega \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + \frac{1}{2} \int_\Omega u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_\Omega v_\varepsilon |\nabla u_\varepsilon|^2 \\
& \leq \int_\Omega v_\varepsilon |\nabla v_\varepsilon|^2 + c_6 \int_\Omega u_\varepsilon v_\varepsilon \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1).
\end{aligned}$$

Upon integration in time, in view of Lemma 2.4, (2.7), (4.33) and the fact that p_* is negative, this implies that there is $c_7 = c_7(K) > 0$ satisfying

$$\begin{aligned}
& \frac{p_*(p_* - 1)}{4} \int_0^t \int_\Omega u_\varepsilon^{p_*+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + \frac{1}{2} \int_0^t \int_\Omega \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 \\
& \quad + \frac{1}{2} \int_0^t \int_\Omega u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_0^t \int_\Omega v_\varepsilon |\nabla u_\varepsilon|^2 + \int_0^t \int_\Omega v_\varepsilon |\nabla v_\varepsilon|^2 + \int_0^t \int_\Omega u_\varepsilon v_\varepsilon \\
& \leq 2 \int_0^t \int_\Omega v_\varepsilon |\nabla v_\varepsilon|^2 + (c_6 + 1) \int_0^t \int_\Omega u_\varepsilon v_\varepsilon + \mathcal{G}_\varepsilon(0) - \mathcal{G}_\varepsilon(t) + \int_\Omega u_{0\varepsilon}^{p_*} \\
& \leq c_7 + c_3 + c_4^{p_*} |\Omega| \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1). \tag{4.38}
\end{aligned}$$

For the above p_0 , it follows from (2.20) that for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$,

$$\frac{1}{p_0} \frac{d}{dt} \int_\Omega u_\varepsilon^{p_0} \leq \frac{p_0 - 1}{2} C_f^2 \int_\Omega u_\varepsilon^{p_0+2\alpha-m-1} v_\varepsilon |\nabla v_\varepsilon|^2 + \ell \int_\Omega u_\varepsilon^{p_0} v_\varepsilon,$$

where in light of (4.36) and Lemma 4.3, we get

$$\begin{aligned}
& \frac{p_0 - 1}{2} C_f^2 \int_\Omega u_\varepsilon^{p_0+2\alpha-m-1} v_\varepsilon |\nabla v_\varepsilon|^2 \\
& \leq \int_\Omega u_\varepsilon^{p_*+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + \int_\Omega u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_\Omega \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 \\
& \quad + \int_\Omega v_\varepsilon |\nabla v_\varepsilon|^2 + c_8 \int_\Omega u_\varepsilon v_\varepsilon \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1)
\end{aligned}$$

with $c_8 = c_8(K, p_0) > 0$, and in view of (4.37), Lemma 4.2 says that with $c_9 = c_9(K, p_0) > 0$ we have

$$\ell \int_\Omega u_\varepsilon^{p_0} v_\varepsilon \leq \int_\Omega u_\varepsilon^{p_*+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + \int_\Omega \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 + c_9 \int_\Omega u_\varepsilon v_\varepsilon$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$. We therefore obtain that

$$\begin{aligned}
& \frac{1}{p_0} \frac{d}{dt} \int_\Omega u_\varepsilon^{p_0} \leq 2 \int_\Omega u_\varepsilon^{p_*+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + \int_\Omega u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + 2 \int_\Omega \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} |D^2 \ln v_\varepsilon|^2 \\
& \quad + \int_\Omega v_\varepsilon |\nabla v_\varepsilon|^2 + (c_8 + c_9) \int_\Omega u_\varepsilon v_\varepsilon \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1),
\end{aligned}$$

while once integrated in time, this together with (4.38) completes the lemma. \blacksquare

4.3 Uniform L^p bounds on u_ε for any $p > 1$

Now we are readily to get the bounds for u_ε with respect to the norm in $L^p(\Omega)$ for any $p > 1$. Before doing this, we first state the following conclusion, which will allow us to control some ill-contributions arising in Lemma 4.9.

Lemma 4.8. *Let $\Omega \subset \mathbb{R}^2$, $m \geq 1$ and $p > 1$. Suppose that $K > 0$ with the property that (1.9) holds, and that for some $p_0 > 1$ there exists $c(K, p_0) > 0$ satisfying*

$$\int_{\Omega} u_\varepsilon^{p_0}(\cdot, t) \leq c(K, p_0) \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1). \quad (4.39)$$

Then for any $\beta \in [p + m - 1, p_0 + p + m - 1)$ and $\eta > 0$, there exists $C(K, p_0, \beta, \eta) > 0$ such that

$$\int_{\Omega} u_\varepsilon^\beta v_\varepsilon \leq \eta \int_{\Omega} u_\varepsilon^{p+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + \eta \int_{\Omega} \frac{|\nabla v_\varepsilon|^{q-2}}{v_\varepsilon^{q-3}} |D^2 \ln v_\varepsilon|^2 + C(K, p_0, \beta, \eta) \int_{\Omega} u_\varepsilon v_\varepsilon \quad (4.40)$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$, where $q > \frac{2(p+m-1)}{p_0}$.

Proof. We first observe that

$$1 < \frac{(p+m-1)(q+2)}{q} < p_0 + p + m - 1 \quad (4.41)$$

due to our assumptions $q > \frac{2(p+m-1)}{p_0}$, $p > 1$ and $m \geq 1$, and note that for any $\beta > 1$ satisfying $\beta \in \left[p + m - 1, \frac{(p+m-1)(q+2)}{q} \right)$, the Young inequality implies that $\int_{\Omega} u_\varepsilon^\beta v_\varepsilon \leq \int_{\Omega} u_\varepsilon v_\varepsilon + \int_{\Omega} u_\varepsilon^{\frac{(p+m-1)(q+2)}{q}} v_\varepsilon$ for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. Therefore we claim that to prove this lemma, it is sufficient to concentrate our focus on

$$\beta \in \left[\frac{(p+m-1)(q+2)}{q}, p_0 + p + m - 1 \right), \quad (4.42)$$

which entails that

$$\vartheta^* := \frac{p_0}{\beta - p - m + 1} > 1 \quad \text{and} \quad \vartheta := \frac{\vartheta^*}{\vartheta^* - 1} > 1.$$

Then for given $\eta > 0$, using the Hölder inequality, Lemma 4.1 and (4.39), there is $c_1 = c_1(K, p_0, \beta, \eta) > 0$ such that

$$\begin{aligned} \int_{\Omega} u_\varepsilon^\beta v_\varepsilon &= \int_{\Omega} \left(u_\varepsilon^{\frac{p+m-1}{2}} v_\varepsilon^{\frac{1}{2}} \right)^2 u_\varepsilon^{\beta-(p+m-1)} \\ &\leq \left\| u_\varepsilon^{\frac{p+m-1}{2}} v_\varepsilon^{\frac{1}{2}} \right\|_{L^{2\vartheta}(\Omega)}^2 \cdot \left\| u_\varepsilon^{\beta-(p+m-1)} \right\|_{L^{\vartheta^*}(\Omega)} \\ &\leq \frac{\eta}{2} \int_{\Omega} u_\varepsilon^{p+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + \int_{\Omega} u_\varepsilon^{p+m-1} \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} + c_1 \int_{\Omega} u_\varepsilon v_\varepsilon \end{aligned} \quad (4.43)$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. Here Lemma 2.2 and two applications of the Young inequality along with the fact that $1 < \frac{(p+m-1)(q+2)}{q} < \beta$ from (4.41) and (4.42) show that there exist $c_2 = c_2(\eta) > 0$ and $c_3 = c_3(\eta, \beta) > 0$ such that

$$\begin{aligned} \int_{\Omega} u_{\varepsilon}^{p+m-1} \frac{|\nabla v_{\varepsilon}|^2}{v_{\varepsilon}} &\leq \frac{\eta}{2(q + \sqrt{2})^2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q+2}}{v_{\varepsilon}^{q+1}} + c_2 \int_{\Omega} u_{\varepsilon}^{\frac{(p+m-1)(q+2)}{q}} v_{\varepsilon} \\ &\leq \frac{\eta}{2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 + \frac{1}{2} \int_{\Omega} u_{\varepsilon}^{\beta} v_{\varepsilon} + c_3 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \end{aligned}$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$, which inserted into (4.43) yields that

$$\int_{\Omega} u_{\varepsilon}^{\beta} v_{\varepsilon} \leq \eta \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \eta \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 + 2(c_1 + c_3) \int_{\Omega} u_{\varepsilon} v_{\varepsilon}$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. ■

With the improved integrability of u_{ε} (Lemmas 4.5-4.7) at hand, the strategy of further deriving the announced boundedness of u_{ε} relies on analysing the energy functional with the form of (1.13) for conveniently large $p > 1$ and appropriately chosen $q > 2$. By suitably interpolating, some unfavorable summands with the expression $\int_{\Omega} u_{\varepsilon}^{\beta} v_{\varepsilon}$ for some β will rise on the right sides of (4.51) and (4.52). To cope with these terms, we hope that the corresponding parameters meet the assumption in Lemma 4.8, so q therein should be in an appropriate intermediate range. In these processes the condition $p_0 > 1$ is of critical importance.

Lemma 4.9. *Let $\Omega \in \mathbb{R}^2$ be a bounded convex domain and $K > 0$ with the property that (1.9) is valid. Then for all $p > 2$ there exists $C(K, p) > 0$ such that if one of the following cases holds:*

- (i) $1 \leq m < 2$, f fulfills (1.3) and (1.4) with $m - 1 < \alpha < m$;
 - (ii) $2 \leq m < 3$, f fulfills (1.3) and (1.5) with $m - 1 < \alpha < \frac{m}{2} + 1$;
 - (iii) $3 \leq m < 4$, f fulfills (1.3) and (1.5) with $m - 1 < \alpha < \frac{m}{2} + 1$ and $u_0 > 0$ in $\bar{\Omega}$,
- one has,

$$\int_{\Omega} u_{\varepsilon}^p(\cdot, t) \leq C(K, p) \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1), \quad (4.44)$$

$$\int_0^{T_{max, \varepsilon}} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq C(K, p) \quad \text{for all } \varepsilon \in (0, 1) \quad (4.45)$$

and

$$\int_0^{T_{max, \varepsilon}} \int_{\Omega} u_{\varepsilon}^{p+m-1} v_{\varepsilon} \leq C(K, p) \quad \text{for all } \varepsilon \in (0, 1). \quad (4.46)$$

Proof. We first recall Lemmas 4.5-4.7 to see that there exist some $p_0 > 1$ and $c_1 = c_1(K, p_0) > 0$ such that

$$\int_{\Omega} u_{\varepsilon}^{p_0}(\cdot, t) \leq c_1 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1), \quad (4.47)$$

and note that Lemma 2.4, (2.7) and (1.9) entail the existence of $c_2 = c_2(K) > 0$ fulfilling

$$\int_0^{T_{max, \varepsilon}} \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \int_0^{T_{max, \varepsilon}} \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \leq c_2 \quad \text{for all } \varepsilon \in (0, 1), \quad (4.48)$$

which in view of the Young inequality implies that it is sufficient to prove (4.44)-(4.46) for $p \geq 3$. As $\frac{2(p+m-1)}{p_0} < 2(p_0 + p + m - 2)$, we are able to pick $q := q(p) > 2(p + m - 2)$ in a way that

$$q \in \left(\frac{2(p+m-1)}{p_0}, 2(p_0 + p + m - 2) \right),$$

which clearly warrants

$$p + m - 1 < \frac{(p+m-1)(q+2)}{q} < p_0 + p + m - 1 \quad (4.49)$$

and

$$p + m - 1 < \frac{q+2}{2} < p_0 + p + m - 1. \quad (4.50)$$

Our assumptions on α and m yield that $0 < p-m+1 \leq p+m-1$ and $0 < p+2\alpha-m-1 < p+m-1$. Thus, we may once more invoke the Young inequality several times to deduce from (2.20) and (2.13) that there is $c_3 = c_3(p) > 0$ such that

$$\begin{aligned} & \frac{1}{p} \frac{d}{dt} \int_{\Omega} u_{\varepsilon}^p + \frac{p-1}{2} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \\ & \leq \frac{p-1}{2} \int_{\Omega} u_{\varepsilon}^{p-m-1} f^2(u_{\varepsilon}) v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \ell \int_{\Omega} u_{\varepsilon}^p v_{\varepsilon} \\ & \leq \frac{p-1}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{p+2\alpha-m-1} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \frac{p-1}{2} C_f^2 \int_{\Omega} u_{\varepsilon}^{p-m+1} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \ell \int_{\Omega} u_{\varepsilon}^p v_{\varepsilon} \\ & \leq (p-1) C_f^2 \int_{\Omega} u_{\varepsilon}^{p+m-1} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + (p-1) C_f^2 \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + \ell \int_{\Omega} u_{\varepsilon}^{p+m-1} v_{\varepsilon} + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \\ & \leq \frac{q}{2(q+\sqrt{2})^2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q+2}}{v_{\varepsilon}^{q+1}} + c_3 \int_{\Omega} u_{\varepsilon}^{\frac{(p+m-1)(q+2)}{q}} v_{\varepsilon}^{\frac{3q+4}{q}} + (p-1) C_f^2 \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 \\ & \quad + \ell \int_{\Omega} u_{\varepsilon}^{p+m-1} v_{\varepsilon} + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \\ & \leq \frac{q}{2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 + c_3 K^{\frac{2q+4}{q}} \int_{\Omega} u_{\varepsilon}^{\frac{(p+m-1)(q+2)}{q}} v_{\varepsilon} + (p-1) C_f^2 \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 \\ & \quad + \ell \int_{\Omega} u_{\varepsilon}^{p+m-1} v_{\varepsilon} + \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max,\varepsilon}) \text{ and } \varepsilon \in (0, 1). \end{aligned} \quad (4.51)$$

This in conjunction with the following inequality identified from Lemma 2.3

$$\frac{d}{dt} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^q}{v_{\varepsilon}^{q-1}} + q \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 \leq c_4 \int_{\Omega} u_{\varepsilon}^{\frac{q+2}{2}} v_{\varepsilon} \quad (4.52)$$

for all $t \in (0, T_{max,\varepsilon})$ and $\varepsilon \in (0, 1)$ with $c_4 = c_4(p) > 0$ implies that

$$\begin{aligned} & \frac{d}{dt} \left\{ \frac{1}{p} \int_{\Omega} u_{\varepsilon}^p + \frac{|\nabla v_{\varepsilon}|^q}{v_{\varepsilon}^{q-1}} \right\} + \frac{p-1}{2} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{q}{2} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 \\ & \leq c_3 K^{\frac{2q+4}{q}} \int_{\Omega} u_{\varepsilon}^{\frac{(p+m-1)(q+2)}{q}} v_{\varepsilon} + c_4 \int_{\Omega} u_{\varepsilon}^{\frac{q+2}{2}} v_{\varepsilon} + \ell \int_{\Omega} u_{\varepsilon}^{p+m-1} v_{\varepsilon} + (p-1) C_f^2 \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 \end{aligned}$$

$$+ \ell \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1). \quad (4.53)$$

Owing to (4.47), (4.49) and (4.50), Lemma 4.8 becomes applicable so as to ensure that there exists $c_5 = c_5(K, p) > 0$ such that

$$\begin{aligned} & c_3 K^{\frac{2q+4}{q}} \int_{\Omega} u_{\varepsilon}^{\frac{(p+m-1)(q+2)}{q}} v_{\varepsilon} + c_4 \int_{\Omega} u_{\varepsilon}^{\frac{q+2}{2}} v_{\varepsilon} + \ell \int_{\Omega} u_{\varepsilon}^{p+m-1} v_{\varepsilon} \\ & \leq \frac{p-1}{4} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{q}{4} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 + c_5 \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \end{aligned} \quad (4.54)$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. Inserting this into (4.53), we get that

$$\begin{aligned} & \frac{d}{dt} \left\{ \frac{1}{p} \int_{\Omega} u_{\varepsilon}^p + \frac{|\nabla v_{\varepsilon}|^q}{v_{\varepsilon}^{q-1}} \right\} + \frac{p-1}{4} \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{q}{4} \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 \\ & \leq (p-1) C_f^2 \int_{\Omega} v_{\varepsilon} |\nabla v_{\varepsilon}|^2 + (\ell + c_5) \int_{\Omega} u_{\varepsilon} v_{\varepsilon} \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1), \end{aligned}$$

which together with (4.48) and (1.9), upon an integration in time, entails that with $c_6 = c_6(K, p) > 0$, we have

$$\frac{1}{p} \int_{\Omega} u_{\varepsilon}^p + \frac{p-1}{4} \int_0^t \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + \frac{q}{4} \int_0^t \int_{\Omega} \frac{|\nabla v_{\varepsilon}|^{q-2}}{v_{\varepsilon}^{q-3}} |D^2 \ln v_{\varepsilon}|^2 \leq c_6 \quad (4.55)$$

for all $t \in (0, T_{max, \varepsilon})$ and $\varepsilon \in (0, 1)$. Thus (4.44) and (4.45) are included. Going back to integrating (4.54) in time and making use of (4.48) and (4.55) conclude (4.46). \blacksquare

5 Proof of Theorem 1.1 and Theorem 1.2

Now by making use of Lemma 3.5 and Lemma 4.9, we could derive the $W^{1, \infty}(\Omega)$ boundedness of v_{ε} from the standard well-known smoothing properties of Neumann heat semigroup once more.

Lemma 5.1. *Suppose that the assumptions in Theorem 1.1 or Theorem 1.2 are satisfied with $K > 0$. Then there exists $C(K) > 0$ fulfilling*

$$\|v_{\varepsilon}(\cdot, t)\|_{W^{1, \infty}(\Omega)} \leq C(K) \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1). \quad (5.1)$$

Proof. We rely on Lemma 3.5 and Lemma 4.9 to see that whenever $n = 1$ or $n = 2$, for any $p > n$ there is $c_1 = c_1(K, p) > 0$ such that

$$\|u_{\varepsilon}(\cdot, t)\|_{L^p(\Omega)} \leq c_1 \quad \text{for all } t \in (0, T_{max, \varepsilon}) \text{ and } \varepsilon \in (0, 1),$$

which guarantees that (5.1) can be deduced in a manner that closely resembles the reasoning in Lemma 3.2. \blacksquare

Having Lemmas 3.5, 4.9 and 5.1 at hand, we can conclude by using a method very similar to [23, Lemma 4.1] that the approximate solutions obtained in Lemma 2.1 are global. That is, we have the following lemma.

Lemma 5.2. *Suppose that the assumptions in Theorem 1.1 or Theorem 1.2 are satisfied. Then $T_{max,\varepsilon} = \infty$ for all $\varepsilon \in (0, 1)$.*

In the following, we will show that in one-dimensional setting, by means of the elliptic Harnack type inequality documented in [25], the L^∞ boundedness of u_ε can be derived by adapting a similar strategy in [11].

Lemma 5.3. *Suppose that the assumptions in Theorem 1.1 are satisfied with $K > 0$ such that (1.7) holds. Then there exists $C(K) > 0$ with the property that for all $\varepsilon \in (0, 1)$ we have*

$$\|u_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq C(K) \quad \text{for all } t > 0. \quad (5.2)$$

Proof. This result can be proved in much the same way as Theorems 5.3 and 5.4 in [11]. So we just show the sketch. Firstly, thanks to the Harnack type inequality

$$v_\varepsilon(x, t) \geq c_1 \|v_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \quad \text{for all } x \in \Omega \quad t > 0 \quad \text{and } \varepsilon \in (0, 1) \quad (5.3)$$

with $c_1 = c_1(K) > 0$, it is easy to verify from (2.6) and (2.7) that

$$L_\varepsilon := \int_0^\infty \|v_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} dt, \quad \varepsilon \in (0, 1)$$

is well-defined. Setting

$$\tau := \phi_\varepsilon(t) := \frac{1}{L_\varepsilon} \int_0^t \|v_\varepsilon(\cdot, s)\|_{L^\infty(\Omega)} ds, \quad t \geq 0$$

and

$$w_\varepsilon(x, \tau) := u_\varepsilon(x, \phi_\varepsilon^{-1}(\tau)), \quad x \in \bar{\Omega}, \quad \tau \in [0, 1),$$

it follows from (2.1) that for each $\varepsilon \in (0, 1)$

$$\begin{cases} w_{\varepsilon\tau} = (a_\varepsilon(x, \tau)w_\varepsilon^{m-1}w_{\varepsilon x})_x - (b_\varepsilon(x, \tau)f(w_\varepsilon))_x + \ell a_\varepsilon(x, \tau)w_\varepsilon, & x \in \Omega, \tau \in (0, 1), \\ w_{\varepsilon x} = 0, & x \in \partial\Omega, \tau \in (0, 1), \\ w_\varepsilon(x, 0) = u_{0\varepsilon}(x), & x \in \Omega \end{cases}$$

is valid with

$$a_\varepsilon(x, \tau) := L_\varepsilon \cdot \frac{v_\varepsilon(x, t)}{\|v_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)}} \quad \text{and} \quad b_\varepsilon(x, \tau) := L_\varepsilon \cdot \frac{v_\varepsilon(x, t)v_{\varepsilon x}(x, t)}{\|v_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)}}.$$

Additionally, there exists $c_2 = c_2(K) > 0$ such that

$$\frac{1}{c_2} \leq a_\varepsilon(x, \tau) \leq c_2 \quad \text{for all } x \in \Omega, \tau \in (0, 1) \quad \text{and } \varepsilon \in (0, 1),$$

and for any $p > 1$ there is $c_3 = c_3(p, K) > 0$ satisfying

$$\|w_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)} + \|b_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)} \leq c_3 \quad \text{for all } \tau \in (0, 1) \quad \text{and } \varepsilon \in (0, 1),$$

which combined with the Moser-type result ([17, Lemma A.1]) show the existence of $c_4 = c_4(K) > 0$ such that

$$\|w_\varepsilon(\cdot, \tau)\|_{L^\infty(\Omega)} \leq c_4 \quad \text{for all } \tau \in (0, 1) \text{ and } \varepsilon \in (0, 1).$$

By rescaling back to u_ε , we finish the proof. \blacksquare

Actually, up to this point if we recall how the regularized form behaves in (2.1) and (2.2), we can assert that the proofs of case (iii) in Theorem 1.1 and case (iii) in Theorem 1.2 have already been finished. Therefore, in the rest of this context we just concentrate on the case $1 \leq m < 3$.

When $n = 1$, similar to the arguments in [11, Theorems 5.4-5.7], we can proceed to identify a suitable sequence whose limit solution (u, v) is accurately a weak solution of (1.2), which simultaneously enjoys the additional properties announced in (1.8).

Lemma 5.4. *Let $n = 1$ and suppose that the assumptions in Theorem 1.1 are satisfied. Then there exist $(\varepsilon_j)_{j \in \mathbb{N}} \in (0, 1)$ and functions $u \geq 0$ and $v > 0$ a.e. in $\Omega \times (0, \infty)$ satisfying*

$$\begin{cases} u \in C^0(\overline{\Omega} \times [0, \infty)) \cap L^\infty(\Omega \times (0, \infty)) & \text{and} \\ v \in C^{2,1}(\overline{\Omega} \times (0, \infty)) \end{cases}$$

such that $\varepsilon = \varepsilon_j \searrow 0$ we have

$$\begin{aligned} u_\varepsilon &\rightarrow u && \text{in } C_{loc}^0(\overline{\Omega} \times [0, \infty)) \quad \text{and} \\ v_\varepsilon &\rightarrow v && \text{in } C_{loc}^{2,1}(\overline{\Omega} \times (0, \infty)), \end{aligned}$$

and that (u, v) forms a global weak solution of (1.2) in the sense of Definition 1.1.

When $n = 2$, we first state some additional spatiotemporal regularity properties.

Lemma 5.5. *Suppose that the assumptions in Theorem 1.2 are satisfied with $K > 0$ such that (1.9) holds. Then for any $p > 4$, there exists $C_1(p, K) > 0$ such that*

$$\int_0^t \left\| \nabla \left(u_\varepsilon^{\frac{p+m-1}{2}}(\cdot, s) v_\varepsilon(\cdot, s) \right) \right\|_{L^2(\Omega)}^2 ds \leq C_1(K, p) \quad \text{for all } t > 0 \text{ and } \varepsilon \in (0, 1) \quad (5.4)$$

and for any $T > 0$, there exist $C_2(K, T, p) > 0$ and $C_3(K, T) > 0$ fulfilling

$$\int_0^T \left\| \partial_t \left(u_\varepsilon^{\frac{p+m-1}{2}}(\cdot, t) v_\varepsilon(\cdot, t) \right) \right\|_{(W^{3,2}(\Omega))^*} dt \leq C_2(K, T, p) \quad \text{for all } \varepsilon \in (0, 1) \quad (5.5)$$

as well as

$$\int_\Omega \ln \frac{\|v_0\|_{L^\infty(\Omega)}}{v_\varepsilon(\cdot, t)} \leq C_3(K, T) \quad \text{for all } t \in (0, T) \text{ and } \varepsilon \in (0, 1). \quad (5.6)$$

Proof. Relying on (1.9) and (2.5), the Young inequality shows that

$$\int_0^t \left\| \nabla \left(u_\varepsilon^{\frac{p+m-1}{2}}(\cdot, s) v_\varepsilon(\cdot, s) \right) \right\|_{L^2(\Omega)}^2 ds$$

$$\begin{aligned}
&\leq \frac{(p+m-1)^2}{2} \int_0^t \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon}^2 |\nabla u_{\varepsilon}|^2 + 2 \int_0^t \int_{\Omega} u_{\varepsilon}^{p+m-1} |\nabla v_{\varepsilon}|^2 \\
&\leq \frac{(p+m-1)^2 K}{2} \int_0^t \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + K^2 \int_0^t \int_{\Omega} u_{\varepsilon}^{2p+2m-3} v_{\varepsilon} + \int_0^t \int_{\Omega} u_{\varepsilon} \frac{|\nabla v_{\varepsilon}|^4}{v_{\varepsilon}^3}
\end{aligned}$$

for all $t > 0$ and $\varepsilon \in (0, 1)$, which by means of (4.45), (4.46) and Lemmas 4.5-4.7 indicates (5.4).

To proceed to prove (5.5), we first fix $\psi \in W^{3,2}(\Omega)$ fulfilling $\|\psi\|_{W^{3,2}(\Omega)} \leq 1$. Drawing on the Sobolev inequality, one can find $c_1 > 0$ such that

$$\|\psi\|_{L^{\infty}(\Omega)} + \|\nabla \psi\|_{L^{\infty}(\Omega)} \leq c_1 \quad \text{for all } t > 0 \text{ and } \varepsilon \in (0, 1).$$

By computing directly and integrating by parts, we see that for all $t > 0$ and $\varepsilon \in (0, 1)$,

$$\begin{aligned}
&\int_{\Omega} \partial_t (u_{\varepsilon}^{\frac{p+m-1}{2}} v_{\varepsilon}) \cdot \psi \\
&= -\frac{p+m-1}{2} \int_{\Omega} \nabla (u_{\varepsilon}^{\frac{p+m-3}{2}} v_{\varepsilon} \psi) \cdot \{u_{\varepsilon}^{m-1} v_{\varepsilon} \nabla u_{\varepsilon} - f(u_{\varepsilon}) v_{\varepsilon} \nabla v_{\varepsilon}\} \\
&\quad + \frac{p+m-1}{2} \ell \int_{\Omega} u_{\varepsilon}^{\frac{p+m-1}{2}} v_{\varepsilon}^2 \psi - \int_{\Omega} \nabla (u_{\varepsilon}^{\frac{p+m-1}{2}} \psi) \cdot \nabla v_{\varepsilon} - \int_{\Omega} u_{\varepsilon}^{\frac{p+m+1}{2}} v_{\varepsilon} \psi \\
&= -\frac{(p+m-1)(p+m-3)}{4} \left\{ \int_{\Omega} u_{\varepsilon}^{\frac{p+3m-7}{2}} v_{\varepsilon}^2 |\nabla u_{\varepsilon}|^2 \psi - \int_{\Omega} u_{\varepsilon}^{\frac{p+m-5}{2}} f(u_{\varepsilon}) v_{\varepsilon}^2 (\nabla u_{\varepsilon} \cdot \nabla v_{\varepsilon}) \psi \right\} \\
&\quad - \frac{p+m-1}{2} \left\{ \int_{\Omega} u_{\varepsilon}^{\frac{p+3m-5}{2}} v_{\varepsilon} (\nabla u_{\varepsilon} \cdot \nabla v_{\varepsilon}) \psi + \int_{\Omega} u_{\varepsilon}^{\frac{p+3m-5}{2}} v_{\varepsilon}^2 (\nabla u_{\varepsilon} \cdot \nabla \psi) - \ell \int_{\Omega} u_{\varepsilon}^{\frac{p+m-1}{2}} v_{\varepsilon}^2 \psi \right. \\
&\quad \left. - \int_{\Omega} u_{\varepsilon}^{\frac{p+m-3}{2}} f(u_{\varepsilon}) v_{\varepsilon} |\nabla v_{\varepsilon}|^2 \psi - \int_{\Omega} u_{\varepsilon}^{\frac{p+m-3}{2}} f(u_{\varepsilon}) v_{\varepsilon}^2 (\nabla v_{\varepsilon} \cdot \nabla \psi) + \int_{\Omega} u_{\varepsilon}^{\frac{p+m-3}{2}} (\nabla u_{\varepsilon} \cdot \nabla v_{\varepsilon}) \psi \right\} \\
&\quad - \int_{\Omega} u_{\varepsilon}^{\frac{p+m-1}{2}} (\nabla v_{\varepsilon} \cdot \nabla \psi) - \int_{\Omega} u_{\varepsilon}^{\frac{p+m+1}{2}} v_{\varepsilon} \psi \\
&=: -\frac{(p+m-1)(p+m-3)}{4} \sum_{i=1}^2 I_i(\varepsilon) - \frac{p+m-1}{2} \sum_{i=3}^8 I_i(\varepsilon) + \sum_{i=9}^{10} I_i(\varepsilon). \tag{5.7}
\end{aligned}$$

Here, we see from $p > 4$ and $1 \leq m < 3$ that $0 < \frac{p+3m-7}{2} < p+m-3$ and $0 \leq 2m-2 < p+m-1$ to assert that by means of the Young inequality if we pick $c_2 = c_2(K) > 0$ satisfying $\|v_{\varepsilon}\|_{W^{1,\infty}(\Omega)} \leq c_2$, then for all $t > 0$ and $\varepsilon \in (0, 1)$,

$$|I_1| \leq c_1 c_2 \int_{\Omega} u_{\varepsilon}^{\frac{p+3m-7}{2}} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 \leq c_1 c_2 \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + c_1 c_2 \int_{\Omega} v_{\varepsilon} |\nabla u_{\varepsilon}|^2$$

and

$$\begin{aligned}
|I_3| + |I_4| &\leq c_1 c_2 \int_{\Omega} u_{\varepsilon}^{\frac{p+3m-5}{2}} v_{\varepsilon} |\nabla u_{\varepsilon}| \\
&\leq c_1 c_2 \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + c_1 c_2 \int_{\Omega} u_{\varepsilon}^{2m-2} v_{\varepsilon} \\
&\leq c_1 c_2 \int_{\Omega} u_{\varepsilon}^{p+m-3} v_{\varepsilon} |\nabla u_{\varepsilon}|^2 + c_1 c_2 \int_{\Omega} u_{\varepsilon}^{p+m-1} v_{\varepsilon} + c_1 c_2^2 |\Omega|.
\end{aligned}$$

When $0 \leq \alpha < 1$, we have $f(u_\varepsilon) \leq u_\varepsilon$, which infers that for all $t > 0$ and $\varepsilon \in (0, 1)$,

$$|I_2| \leq c_1 c_2^2 \int_{\Omega} u_\varepsilon^{\frac{p+m-3}{2}} v_\varepsilon |\nabla u_\varepsilon| \leq c_1 c_2^2 \int_{\Omega} u_\varepsilon^{p+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + c_1 c_2^3 |\Omega|.$$

When $\alpha \geq 1$, there exists $c_3 > 0$ such that $f(u_\varepsilon) \leq c_3 u_\varepsilon + c_3 u_\varepsilon^\alpha$. Then two applications of the Young inequality along with the facts that $0 < \frac{p+m-5}{2} + \alpha < p+m-3$ and $0 \leq 2\alpha - 2 < p+m-1$ reveal that for all $t > 0$ and $\varepsilon \in (0, 1)$,

$$\begin{aligned} |I_2| &\leq c_1 c_2^2 \int_{\Omega} u_\varepsilon^{\frac{p+m-5}{2}} f(u_\varepsilon) v_\varepsilon |\nabla u_\varepsilon| \\ &\leq c_1 c_2^2 c_3 \int_{\Omega} u_\varepsilon^{\frac{p+m-3}{2}} v_\varepsilon |\nabla u_\varepsilon| + c_1 c_2^2 c_3 \int_{\Omega} u_\varepsilon^{\frac{p+m-5}{2} + \alpha} v_\varepsilon |\nabla u_\varepsilon| \\ &\leq 2c_1 c_2^2 c_3 \int_{\Omega} u_\varepsilon^{p+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + c_1 c_2^3 c_3 |\Omega| + c_1 c_2^2 c_3 \int_{\Omega} u_\varepsilon^{2\alpha-2} v_\varepsilon \\ &\leq 2c_1 c_2^2 c_3 \int_{\Omega} u_\varepsilon^{p+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + c_1 c_2^2 c_3 \int_{\Omega} u_\varepsilon^{p+m-1} v_\varepsilon + 2c_1 c_2^3 c_3 |\Omega|. \end{aligned}$$

Similarly, since $0 < \frac{p+m-3}{2} + \alpha < p+m-1$ and $0 < \frac{p+m-1}{2} < \frac{p+m+1}{2} < p+m-1$, for all $t > 0$ and $\varepsilon \in (0, 1)$ we have

$$|I_6| + |I_7| \leq 2c_1 c_2^2 c_3 \int_{\Omega} u_\varepsilon^{\frac{p+m-1}{2}} v_\varepsilon + 2c_1 c_2^2 c_3 \int_{\Omega} u_\varepsilon^{\frac{p+m-3}{2} + \alpha} v_\varepsilon \leq 4c_1 c_2^2 c_3 \int_{\Omega} u_\varepsilon^{p+m-1} v_\varepsilon + 4c_1 c_2^3 c_3 |\Omega|$$

and

$$|I_5| + |I_{10}| \leq c_1 c_2 \int_{\Omega} u_\varepsilon^{\frac{p+m-1}{2}} v_\varepsilon + c_1 \int_{\Omega} u_\varepsilon^{\frac{p+m+1}{2}} v_\varepsilon \leq (c_1 c_2 + c_1) \int_{\Omega} u_\varepsilon^{p+m-1} v_\varepsilon + (c_1 c_2^2 + c_1 c_2) |\Omega|.$$

Using the Young inequality several times, we can treat I_8 and I_9 as follows

$$\begin{aligned} |I_8| + |I_9| &\leq c_1 \int_{\Omega} u_\varepsilon^{\frac{p+m-3}{2}} |\nabla u_\varepsilon| |\nabla u_\varepsilon| + c_1 \int_{\Omega} u_\varepsilon^{\frac{p+m-1}{2}} |\nabla u_\varepsilon| \\ &\leq c_1 \int_{\Omega} u_\varepsilon^{p+m-4} v_\varepsilon |\nabla u_\varepsilon|^2 + c_1 \int_{\Omega} u_\varepsilon^{p+m-2} v_\varepsilon + 2c_1 \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^2}{v_\varepsilon} \\ &\leq c_1 \int_{\Omega} u_\varepsilon^{p+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + c_1 \int_{\Omega} u_\varepsilon^{p+m-1} v_\varepsilon + 2c_1 \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + 3c_1 \int_{\Omega} u_\varepsilon v_\varepsilon + c_1 \int_{\Omega} v_\varepsilon |\nabla u_\varepsilon|^2. \end{aligned}$$

Substituting these estimations on $I_i, i = 1, 2, \dots, 10$ into (5.7), we obtain that for all $t > 0$ and $\varepsilon \in (0, 1)$, there is $c_4 = c_4(K) > 0$ such that

$$\begin{aligned} \left\| \partial_t \left(u_\varepsilon^{\frac{p+m-1}{2}} v_\varepsilon \right) \right\|_{(W^{3,2}(\Omega))^*} &\leq c_4 \left\{ \int_{\Omega} u_\varepsilon^{p+m-3} v_\varepsilon |\nabla u_\varepsilon|^2 + \int_{\Omega} u_\varepsilon^{p+m-1} v_\varepsilon \right. \\ &\quad \left. + \int_{\Omega} u_\varepsilon \frac{|\nabla v_\varepsilon|^4}{v_\varepsilon^3} + \int_{\Omega} v_\varepsilon |\nabla u_\varepsilon|^2 + \int_{\Omega} u_\varepsilon v_\varepsilon + 1 \right\}. \end{aligned} \quad (5.8)$$

Integrating (5.8) from 0 to T , we obtain (5.5) as a consequence of (2.7), (1.9), Lemmas 4.5-4.7 and Lemma 4.9. Finally, (5.6) can be proved by using the same way in [23, Lemma 4.4]. \blacksquare

With all the preparations, we are able to construct a weak solution to (1.2) when $n = 2$. The presentation takes the following form. One can refer to [23, Lemma 5.1] for the details.

Lemma 5.6. *Let $n = 2$ and $p > 2$. Suppose that the assumptions in Theorem 1.2 are satisfied. Then there exist $(\varepsilon_j)_{j \in \mathbb{N}} \in (0, 1)$ and functions $u \geq 0$ and $v > 0$ a.e. in $\Omega \times (0, \infty)$ satisfying*

$$\begin{cases} u \in L^\infty((0, \infty); L^p(\Omega)) \\ v \in L^\infty(\Omega \times (0, \infty)) \quad \text{and} \\ \nabla v \in L^\infty(\Omega \times (0, \infty)) \end{cases}$$

such that $\varepsilon = \varepsilon_j \searrow 0$ we have

$$\begin{aligned} u_\varepsilon &\rightarrow u && \text{a.e. in } \Omega \times (0, \infty) \text{ and in } L^p_{loc}(\overline{\Omega} \times [0, \infty)), \\ v_\varepsilon &\rightarrow v && \text{a.e. in } \Omega \times (0, \infty) \text{ and in } L^p_{loc}(\overline{\Omega} \times [0, \infty)) \quad \text{and} \\ \nabla v_\varepsilon &\overset{*}{\rightharpoonup} \nabla v && \text{in } L^\infty(\Omega \times (0, \infty)), \end{aligned}$$

and that (u, v) forms a global weak solution of (1.2) in the sense of Definition 1.1.

Our main results thereby in fact reduce to a mere summary:

Proofs of Theorems 1.1 and 1.2. The proofs of cases (i) and (ii) in Theorem 1.1 are consequences of Lemma 5.4, while Lemmas 5.1-5.3 together with Lemma 2.1 prove case (iii) in Theorem 1.1. Apart from that Lemma 5.6 indicates cases (i) and (ii) in Theorem 1.2, and Lemmas 4.9, 5.1, 5.2 in conjunction with Lemma 2.1 complete the proof of case (iii) in Theorem 1.2. \square

Acknowledgements. The author is very grateful to Prof. Michael Winkler and Dr. Genglin Li for their useful discussions and advice. This work was supported by the China Scholarship Council (No. 202206290145) and the Deutsche Forschungsgemeinschaft (No. 462888149).

References

- [1] Amann, H.: *Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems*. In: Functional space, differential operators and nonlinear analysis, Teubner-Texte Math. 133 (1993): 9-126.
- [2] Bellomo, N., Outada, N., Soler, J., Tao, Y., Winkler, M.: *Chemotaxis and cross-diffusion models in complex environments: models and analytic problems toward a multiscale vision*. Math. Models Methods Appl. Sci. 32 (2022): 713-792.
- [3] Ben-Jacob, E., Cohen, I., Levine, H.: *Cooperative self-organization of microorganisms*. Adv. Phys. 49 (2000): 395-554.
- [4] Fujikawa, H.: *Periodic growth of Bacillus subtilis colonies on agar plates*. Physica A 189 (1992): 15-21.
- [5] Golding, I., Kozlovsky, Y., Cohen, I., Ben-Jacob, E.: *Studies of bacterial branching growth using reaction-diffusion models for colonial development*. Physica A 260 (1998): 510-554.
- [6] Kawasaki, K., Mochizuki, A., Matsushita, M., Umeda, T., Shigesada, N.: *Modeling spatio-temporal patterns generated by Bacillus subtilis*. J. Theor. Biol. 188 (1997): 177-185.
- [7] Kitsunezaki, S.: *Interface dynamics for bacterial colony formation*. J. Phys. Soc. Jpn. 66 (1997): 1544-1550.

- [8] Ladyženskaja, O.A., Solonnikov, V.A., Ural'ceva, N.N.: *Linear and quasilinear equations of parabolic type*. Translations of Mathematical Monographs, vol. 23. American Mathematical Society, Providence, RI, 1968.
- [9] Leyva, J.F., Málaga, C., Plaza, R.G.: *The effects of nutrient chemotaxis on bacterial aggregation patterns with non-linear degenerate cross diffusion*. Physica A 392 (2013): 5644-5662.
- [10] Li G.: *Large-data global existence in a higher-dimensional doubly degenerate nutrient system*. J. Differential Equations 329 (2022): 318-347.
- [11] Li, G., Winkler, M.: *Nonnegative solutions to a doubly degenerate nutrient taxis system*. Commun. Pure Appl. Anal. 21 (2022): 687-704.
- [12] Lieberman, G.M.: *Hölder continuity of the gradient of solutions of uniformly parabolic equations with conormal boundary conditions*. Ann. Mat. Pura Appl. 148 (1987): 77-99.
- [13] Lions, P.L.: *Résolution de problèmes elliptiques quasilinéaires*. Arch. Ration. Mech. Anal. 74 (1980): 335-353.
- [14] Matsushita, M., Fujikawa, H.: *Diffusion-limited growth in bacterial colony formation*. Physica A 168 (1990): 498-506.
- [15] Ohgiwari, M., Matsushita, M., Matsuyama, T.: *Morphological changes in growth phenomena of bacterial colony patterns*. J. Phys. Soc. Jpn. 61 (1992): 816-822.
- [16] Porzio, M.M., Vespi, V.: *Hölder estimates for local solutions of some doubly nonlinear degenerate parabolic equations*. J. Differential Equations 103 (1993): 146-178.
- [17] Tao, Y., Winkler, M.: *Boundedness in a quasilinear parabolic-parabolic Keller-Segel system with subcritical sensitivity*. J. Differential Equations 252 (2012): 692-715.
- [18] Tao, Y., Winkler, M.: *Eventual smoothness and stabilization of large-data solutions in a three-dimensional chemotaxis system with consumption of chemoattractant*. J. Differential Equations 252 (2012): 2520-2543.
- [19] Winkler, M.: *Aggregation vs. global diffusive behavior in the higher-dimensional Keller-Segel model*. J. Differential Equations 248 (2010): 2889-2905.
- [20] Winkler, M.: *Asymptotic homogenization in a three-dimensional nutrient taxis system involving food-supported proliferation*. J. Differential Equations 263 (2017): 4826-4869.
- [21] Winkler, M.: *Does spatial homogeneity ultimately prevail in nutrient taxis systems? A paradigm for structure support by rapid diffusion decay in an autonomous parabolic flow*. Trans. Amer. Math. Soc. 374 (2021): 219-268.
- [22] Winkler, M.: *Approaching logarithmic singularities in quasilinear chemotaxis-consumption systems with signal-dependent sensitivities*. Discrete Contin. Dyn. Syst. Ser. B 27 (2022): 6565-6587.
- [23] Winkler, M.: *Small-signal solutions of a two-dimensional doubly degenerate taxis system modeling bacterial motion in nutrient-poor environments*. Nonlinear Anal. Real World Appl. 63 (2022): 103407.
- [24] Winkler, M.: *Stabilization of arbitrary structures in a doubly degenerate reaction-diffusion system modeling bacterial motion on a nutrient-poor agar*. Calc. Var. Partial Differential Equations 61 (2022): 108.
- [25] Winkler, M.: *Elliptic Harnack inequalities in linear parabolic equations and application to the asymptotics in a doubly degenerate nutrient taxis system*. Preprint.