

# SEMI-LOCAL BEHAVIOUR OF NON-LOCAL HYPOELLIPTIC EQUATIONS: BOLTZMANN

AMÉLIE LOHER

ABSTRACT. The purpose of this note is to demonstrate the announced result in [12] by filling the gap in the proof sketch. We prove the semi-local Strong Harnack inequality for the Boltzmann equation for moderately soft potentials without cutoff assumption. The non-local operator in the Boltzmann equation is in non-divergence form, and thus the method developed in [14] does not apply. However, we exploit that the Boltzmann equation is on average in divergence form, and we show that the non-divergent part of the collision operator is of lower order in a suitable sense, which proves to be sufficient to deduce the Strong Harnack inequality. Consequentially, we derive upper and lower bounds on the fundamental solution of the linearised Boltzmann equation.

## CONTENTS

1.	Introduction	1
2.	Setup	4
3.	Tail bound on upper level sets	7
4.	Linear $L^1$ to $L^\infty$ bound	15
5.	Brief note on bounds of the fundamental solution	21
	References	27

## 1. INTRODUCTION

The aim of this article is to establish the Strong Harnack inequality for the Boltzmann equation for moderately soft potentials without cutoff assumption. The Boltzmann equation models the dynamics of a dilute gas. It is given by

$$(1.1) \quad \partial_t f + v \cdot \nabla_x f = Q(f, f),$$

where  $f : \mathbb{R} \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow [0, \infty)$  encodes the density of the gas particles, which at any given time  $t \in \mathbb{R}$  have location  $x \in \mathbb{R}^d$  and velocity  $v \in \mathbb{R}^d$ . The right hand side  $Q$  denotes the Boltzmann collision operator, whose explicit form is given by

$$(1.2) \quad Q(f, f) = \int_{\mathbb{R}^d} \int_{\mathbb{S}^{d-1}} [f(w_*)f(w) - f(v_*)f(v)] B(|v - v_*|, \cos \theta) \, d\sigma \, dv_*,$$

where  $v, v_*$  are the post-collisional velocities, and  $w, w_*$  the pre-collisional velocities, so that

$$w = \frac{v + v_*}{2} + \frac{|v - v_*|}{2} \sigma, \quad w_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2} \sigma.$$

---

*Date:* Apr 25, 2024.

The rate of change in velocities is determined through the cross-section  $B$ , which reads

$$(1.3) \quad B(r, \cos \theta) = r^\gamma b(\cos \theta), \quad b(\cos \theta) \approx \left| \sin \left( \frac{\theta}{2} \right) \right|^{-(d-1)-2s},$$

with parameters  $\gamma \in (-d, 1]$  and  $s \in (0, 1)$ , and where  $\cos \theta$  is defined as

$$\cos \theta := \frac{v - v_*}{|v - v_*|} \cdot \sigma, \quad \sin \left( \frac{\theta}{2} \right) := \frac{w - v}{|w - v|} \cdot \sigma.$$

We restrict our analysis to moderately soft potentials, which means we consider  $0 \leq \gamma + 2s \leq 2$  throughout this article, and moreover, we are interested in long-range interactions, which occur when we do not cut off the singularity in the kernel at sufficiently small deviation angles  $\theta$ .

**1.1. Boltzmann collision operator.** The Boltzmann equation is a kinetic integro-differential equation with a bounded, non-negative source term. To see this, we split the collision operator  $Q$  into two parts

$$Q(f, f) = Q_1(f, f) + Q_2(f, f),$$

where

$$Q_1(f, f) = \int_{\mathbb{R}^d} \int_{\mathbb{S}^{d-1}} f(w_*) [f(w) - f(v)] B(|v - v_*|, \cos \theta) \, d\sigma \, dv_*,$$

and

$$Q_2(f, f) = f(v) \left( \int_{\mathbb{R}^d} \int_{\mathbb{S}^{d-1}} [f(w_*) - f(v_*)] B(|v - v_*|, \cos \theta) \, d\sigma \, dv_* \right).$$

Then, as shown in [15, Lemma 4.1],  $Q_1$  can be rewritten as

$$Q_1(f, f) = \int_{\mathbb{R}^d} [f(t, x, w) - f(t, x, v)] K_f(t, x, v, w) \, dw,$$

where the kernel  $K_f$  depends implicitly on the solution  $f$  and is given by

$$(1.4) \quad K_f(v, w) = 2^{d-1} |w - v|^{-1} \int_{w' \perp w - v} f(v + w') B(r, \cos \theta) r^{-d+2} \, dw',$$

with  $r^2 = |w - v|^2 + |w'|^2 = |v - v_*|^2$  and  $\cos \theta = \frac{w' - (v - w)}{|w' - (v - w)|} \cdot \frac{w' + (w - v)}{|w' + (w - v)|}$ ,  $w_* = v + w'$ , and  $v_* = w + w'$ . The proof is based on a change of variables. In particular,  $Q_1$  is a non-local operator with a non-negative kernel  $K_f$ .

Moreover, the term  $Q_2$  can be viewed as a non-negative source term due to cancellation effects. It is well known in the kinetic literature, as stated for example in [15, Lemma 5.1], that  $Q_2$  is of the form

$$Q_2(f, f) = f(v) \left( \int_{\mathbb{R}^d} f(v - w') \tilde{B}(|w'|) \, dw' \right),$$

where

$$(1.5) \quad \tilde{B}(r) := \int_{\mathbb{S}^{d-1}} \left( \frac{2^{\frac{d}{2}}}{(1 - \sigma \cdot e)^{\frac{d}{2}}} B \left( \frac{\sqrt{r}}{\sqrt{1 - \sigma \cdot e}}, \cos \theta \right) - B(r, \cos \theta) \right) \, d\sigma > 0.$$

Here  $e$  is any unit vector, and  $\cos \theta = \sigma \cdot e$ .

**1.1.1. Integro-differential equation with non-negative source.** We have therefore derived the integro-differential form of the Boltzmann equation with a non-negative source term: if  $f$  solves (1.1), then

$$(1.6) \quad \partial_t f(t, x, v) + v \cdot \nabla_x f(t, x, v) = \int_{\mathbb{R}^d} (f(t, x, w) - f(t, x, v)) K_f(t, x, v, w) \, dw + \Lambda_f f(t, x, v),$$

where  $K_f$  is given by (1.4), and  $\Lambda_f$  is determined by

$$(1.7) \quad \Lambda_f := \int_{\mathbb{R}^d} f(v - w') \tilde{B}(|w'|) \, dw' \geq 0,$$

where  $\tilde{B}$  is defined in (1.5).

1.1.2. *Integro-differential equation in divergence form on average.* There is, however, also another way we can think of the Boltzmann equation, which proves to be useful for the purpose of this article. If we reconsider the term  $Q_2$ , then we realise that the change of variables according to [6, Lemma A.9] together with (1.4) yields

$$\begin{aligned}
 Q_2(f, f) &= f(v) \left( \int_{\mathbb{R}^d} \int_{\mathbb{S}^{d-1}} (f(w_*) - f(v_*)) B(|v - v_*|, \cos \theta) \, d\sigma \, dv_* \right) \\
 (1.8) \quad &= f(v) \left( 2^{d-1} \int_{\mathbb{R}^d} \int_{w' \perp w - v} |w - v|^{-1} B(r, \cos \theta) r^{-d+2} (f(v + w') - f(w + w')) \, dw' \, dw \right) \\
 &= f(v) \left( \int_{\mathbb{R}^d} (K_f(v, w) - K_f(w, v)) \, dw \right)
 \end{aligned}$$

Thus any solution of (1.1) satisfies

$$\begin{aligned}
 \partial_t f(t, x, v) + v \cdot \nabla_x f(t, x, v) &= \int_{\mathbb{R}^d} (f(t, x, w) - f(t, x, v)) K_f(t, x, v, w) \, dw \\
 &\quad + f(t, x, v) \left( \int_{\mathbb{R}^d} (K_f(t, x, v, w) - K_f(t, x, w, v)) \, dw \right) \\
 (1.9) \quad &= \frac{1}{2} \int_{\mathbb{R}^d} (f(t, x, w) - f(t, x, v)) (K_f(t, x, v, w) + K_f(t, x, w, v)) \, dw \\
 &\quad + \frac{1}{2} \int_{\mathbb{R}^d} (f(t, x, w) + f(t, x, v)) (K_f(t, x, v, w) - K_f(t, x, w, v)) \, dw.
 \end{aligned}$$

The strength of this reformulation (1.9) is the emphasis on the fact that *on average, the Boltzmann equation is an equation in divergence form.*

**1.2. Conditional regime.** In either formulation, (1.6) and (1.9), we see that the coefficients  $K_f$  (1.4) and  $\Lambda_f$  (1.7) depend implicitly on the solution, stressing the non-linear character of the equation. Without any further assumptions on the solution itself, it seems out of reach to treat the non-linearity in the equation. In order to make quantitative statements on the coefficients, we need to work in a conditional regime [7]. We thus assume throughout this article that the solutions we consider satisfy the following hydrodynamic bounds: for any  $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$ , there exists  $m_0, M_0, E_0, H_0$ , such that

$$\begin{aligned}
 0 < m_0 &\leq \int_{\mathbb{R}^d} f(t, x, v) \, dv \leq M_0, \\
 (1.10) \quad &\int_{\mathbb{R}^d} f(t, x, v) |v|^2 \, dv \leq E_0, \\
 &\int_{\mathbb{R}^d} f(t, x, v) \log f(t, x, v) \, dv \leq H_0.
 \end{aligned}$$

In other words, the mass, the energy and the entropy are bounded. Then, Silvestre showed [15, Theorem 1.2], that there is a constant  $C_0 > 0$  depending only on  $m_0, M_0, E_0, H_0$  such that

$$\|f\|_{L^\infty(\mathbb{R}_+ \times \mathbb{R}^d \times \mathbb{R}^d)} \leq C_0.$$

This conditional a priori bound on the solution enables us to quantify an ellipticity class on the coefficients  $K_f$  and  $\Lambda_f$ , and thus the non-linearity is treated by absorbing it in the macroscopic quantities.

**1.3. Main result.** In this conditional regime, a series of works by Imbert-Silvestre(-Mouhot) [5–8] showed first that any solution of (1.1) satisfies the Weak Harnack inequality, and is thus Hölder continuous [6], then that the regularity of the kernel  $K$  can be transferred onto the solution in the sense of Schauder estimates [8], and finally that these local results can be globalised via a change of variables [7, Section 5] due to the decay properties of the solution [5], so that eventually the Schauder estimates can be bootstrapped to obtain smooth solutions [7]. Later a constructive proof of the Weak Harnack inequality and the Schauder estimates appeared in [11, 13]. One of the difficulties posed by (1.1) is its non-local operator, which means

that the behaviour of the solution inside a given domain is affected by the values attained in the whole velocity space. In contrast to equations with a local diffusion operator, we cannot deduce from these Hölder estimates a bound on the local supremum of the solution in terms of the local infimum. Such a bound is known as the Strong Harnack inequality. The problem is that the behaviour of the tail is encoded in the constant appearing in the bound. Intuitively, it might not even be clear that we should expect a fully local bound on a solution to a non-local equation. Even though it is known that such a statement does hold for parabolic non-local equations [16], it turns out to fail in the kinetic case [10]. We showed in [14], however, that a local Harnack inequality without tail terms holds for *global* solutions to non-local kinetic equations in divergence form. The key is to capture the behaviour of the tail. Here we show how to obtain the Strong Harnack inequality for global solutions to the Boltzmann equation.

**Theorem 1.1** (Strong Harnack for Boltzmann). *Let  $T > 0$ . Let  $f$  solve (1.1) in  $(0, T) \times \mathbb{R}^d \times \mathbb{R}^d$  in the sense of Definition 2.1 with a cross section  $B$  given by (1.3) with  $0 \leq \gamma + 2s \leq 2$ . If  $f$  satisfies (1.10), then for any  $0 < r_0 < \frac{1}{8}$ , there is  $C > 0$  such that*

$$\sup_{(\tau_0, \tau_1) \times Q_{\frac{r_0}{4}}^t} f \leq C \inf_{(\tau_2, \tau_3) \times Q_{\frac{r_0}{4}}^t} f,$$

where  $0 < \tau_0 < \tau_1 < \tau_2 < \tau_3$  such that  $\tau_1 - \tau_0 = \tau_2 - \tau_3 = \left(\frac{r_0}{4}\right)^{2s}$  and  $\tau_2 - \tau_1 \geq r_0^{2s}$ , and where  $Q_{\frac{r_0}{4}}^t := \{(x, v) \in \mathbb{R}^{2d} : |x| < \left(\frac{r_0}{4}\right)^{1+2s}, |v| < \frac{r_0}{4}\}$  is the kinetic cylinder sliced in time. The constant  $C$  depends only on  $s, d, \gamma, m_0, M_0, E_0, H_0$ , and in case that  $\gamma \leq 0$ , it depends on the decay of the initial data.

This is the first time a Strong Harnack inequality for the Boltzmann equation has been derived. Moreover, we can deduce consequences on bounds on the fundamental solution, similar to [14, Theorem 1.2 and Theorem 1.3]. We refer the reader to Section 5.

**Acknowledgements.** We are extremely grateful for the incredible support we get from Clément Mouhot, and for his confidence in our work that endures long after ours vanished.

## 2. SETUP

We first describe the notion of solutions that we work with.

**2.1. Notion of solutions.** Since we require further regularity properties of the solution to deduce that the skew-symmetric part of  $K_f$  is of lower order in the sense of (2.6), we need to work with the following notion of solutions.

**Definition 2.1** (Solutions). We say that  $f : (0, T) \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$  is a solution of (1.1), if

- (i)  $f$  is non-negative everywhere,
- (ii)  $f$  solves (1.1) classically for every  $(t, x, v)$ ,
- (iii)  $f \in C^\infty$  in time, space, and velocity,
- (iv) For each value of  $(t, x)$  the function  $f$  decays rapidly as  $|v| \rightarrow \infty$ , that is for any  $q > 0$

$$\lim_{|v| \rightarrow \infty} \frac{f(t, x, v)}{(1 + |v|)^q} = 0$$

locally uniformly in  $(t, x)$ .

- (v)  $f$  is periodic in  $x$ ,

(vi)  $f$  conserves mass:

$$\forall 0 \leq t_1 < t_2 \leq T : \iint_{\mathbb{R}^{2d}} f(t_2) \, dx \, dv \leq \iint_{\mathbb{R}^{2d}} f(t_1) \, dx \, dv.$$

Apart from the mass conservation, this notion of solutions was also used by Imbert-Silvestre in [7].

For any solution in the sense of Definition 2.1 satisfying the assumption on the macroscopic quantities (1.10), we can quantify a notion of ellipticity on the coefficients.

**2.2. Ellipticity class.** For any  $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$ , we can show that there exists  $\lambda_0, \Lambda_0, \mu_0 > 0$  depending on  $\gamma, s, d, m_0, M_0, E_0, H_0$  such that  $K_f$  given in (1.4) and  $\Lambda_f$  given in (1.7) satisfy the following statements.

We start by noting that  $\Lambda_f$  given in (1.7) is non-negative, and can be bounded by a constant depending only on  $\gamma, s, d$  and the macroscopic bounds (1.10), that is

$$(2.1) \quad 0 \leq \Lambda_f \leq \Lambda_0.$$

Concerning  $K_f$ , first, we assume a weak form of coercivity: for any  $\varphi \in C_c^\infty(\mathbb{R}^d)$  there holds

$$(2.2) \quad \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (\varphi(v) - \varphi(w))^2 K_f(v, w) \, dw \, dv \geq \lambda_0 \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{|\varphi(v) - \varphi(w)|^2}{|v - w|^{d+2s}} \, dv \, dw.$$

Second, we assume an upper bound on average: for any  $r > 0$

$$(2.3) \quad \forall v \in \mathbb{R}^d \quad \int_{B_r(v)} K_f(v, w) |v - w|^2 \, dw \leq \Lambda_0 r^{2-2s}.$$

Third, we discuss the symmetry of the kernel. Note that the kernel  $K_f$  satisfies a pointwise symmetry of the form  $K_f(v, v+w) = K_f(v, v-w)$ . This symmetry assumption gives rise to a non-local operator in non-divergence form. However, the kernel *does not satisfy a pointwise divergence form symmetry*:  $K_f(v, w) = K_f(w, v)$ . This implies that the non-local operator with kernel  $K_f$  is not self-adjoint. It does, however, satisfy a weak divergence form symmetry:

$$(2.4) \quad \forall v \in \mathbb{R}^d \quad \left| \text{PV} \int_{\mathbb{R}^d} (K_f(v, w) - K_f(w, v)) \, dw \right| \leq \Lambda_0,$$

and if  $s \geq \frac{1}{2}$  we assume that for all  $r > 0$

$$(2.5) \quad \forall v \in \mathbb{R}^d \quad \left| \text{PV} \int_{B_r(v)} (v - w)(K_f(v, w) - K_f(w, v)) \, dw \right| \leq \Lambda_0 r^{1-2s}.$$

These conditions mean that the anti-symmetric part is bounded. It seems that conditions (2.4) and (2.5) are not sufficient to deduce our main theorem. We also need to assume that the anti-symmetric part is of lower order with respect to the symmetric part: for any  $0 < r \leq 1$  there is  $\alpha > s$  such that

$$(2.6) \quad \sup_{v \in B_r} \int_{B_r} \frac{|K_f(v, w) - K_f(w, v)|^2}{(K_f(v, w) + K_f(w, v))} \, dw \leq \mu_0 r^{2(\alpha-s)}.$$

This condition has been used to derive Hölder continuity of solutions to parabolic non-local equations by Kassmann-Weidner [9].

The assumptions (2.1)-(2.6) are satisfied for any solution  $f$  of (1.1) in  $(0, T) \times \mathbb{R}^d \times \mathbb{R}^d$ , such that (1.10) holds, and in case that  $\gamma \leq 0$ , provided that for any  $q > 0$  there is  $N_q > 0$  such that  $f_0(x, v) = f(0, x, v) \leq N_q(1 + |v|)^{-q}$  for  $(x, v) \in \mathbb{R}^d \times \mathbb{R}^d$ . For the justifications, we refer the reader to [15, Lemma 5.2] for (2.1), to [3, Theorem 1.2] for (2.2), to [6, Lemma 3.5, Lemma 3.6, Lemma 3.7] for (2.3), (2.4) and (2.5), respectively. For a justification of the last assumption, we use further regularity properties of solutions to (1.1), derived by Imbert-Mouhot-Silvestre in [4, 6, 7].

**2.3. On the non-divergent part of the kernel.** We consider a solution  $f$  of (1.1) in  $(0, T) \times \mathbb{R}^d \times \mathbb{R}^d$ , periodic in  $x$ , such that (1.10) holds. If  $\gamma \leq 0$ , then we also assume that for any  $q > 0$  there is  $N_q > 0$  such that  $f_0(x, v) = f(0, x, v) \leq N_q(1 + |v|)^{-q}$  for  $(x, v) \in \mathbb{R}^d \times \mathbb{R}^d$ . Then for every  $q > 0$  there is a constant  $C_{s,q} > 0$  and  $\alpha > s$  depending only on  $s, d, \gamma, m_0, M_0, E_0, H_0$  and  $N_q$  in case that  $\gamma \leq 0$ , such that for any  $R > 0$ , any  $(t, x) \in (\tau_0, T) \times B_{R^{1+2s}}$  and any  $v, w \in \mathbb{R}^d$  with  $|v - w| < \min\{\frac{R}{2}, 1\}$  there holds

$$(2.7) \quad |f(v) - f(w)| \leq C_{s,q} |v - w|^\alpha (1 + |v|)^{-q} \approx C_{s,q} |v - w|^\alpha (1 + |w|)^{-q}.$$

An even stronger statement is derived in [7, Corollary 7.8]. We now consider  $(t, x, v) \in Q_R$  for  $0 < R \leq 1$ . We recall [15, Corollary 4.2]: for any  $v, w \in \mathbb{R}^d$

$$K_f(v, w) \approx |v - w|^{-(d+2s)} \int_{w' \perp v-w} f(v+w') |w'|^{\gamma+2s+1} dw'.$$

Thus for  $v \in B_R$  we take  $q > d + 2s + \gamma$  and use (2.7), so that

$$\begin{aligned} PV \int_{B_R} |K_f(v, w) - K_f(w, v)| dw &= 2^{d-1} \int_{B_R} \int_{w' \perp v-w} |f(v+w') - f(w+w')| \frac{r^\gamma b(\cos \theta)}{|v-w|^{d-2}} dw' dw \\ &\approx \int_{B_R} |v-w|^{-(d+2s)} \int_{w' \perp v-w} |f(v+w') - f(w+w')| |w'|^{\gamma+2s+1} dw' dw \\ &\leq C_{s,q} \int_{B_R} |v-w|^{-(d+2s)+\alpha} \int_{w' \perp v-w} (1+|v+w'|)^{-q} |w'|^{\gamma+2s+1} dw' dw \\ &\approx C_{s,q} \int_{B_R} |v-w|^{-(d+2s)+\alpha} \int_{w' \perp v-w} (1+|w'|)^{-q} |w'|^{\gamma+2s+1} dw' dw. \end{aligned}$$

In particular, we see for any  $0 < R \leq 1$  and any  $v \in B_R$

$$\begin{aligned} &\int_{B_R} \frac{|K_f(v, w) - K_f(w, v)|^2}{(K_f(v, w) + K_f(w, v))} dw \\ &\lesssim \int_{B_R} |v-w|^{-(d+2s)} \frac{\left( \int_{w' \perp v-w} |f(v+w') - f(w+w')| |w'|^{\gamma+2s+1} dw' \right)^2}{\int_{w' \perp v-w} (f(v+w') + f(w+w')) |w'|^{\gamma+2s+1} dw'} dw \\ &\leq C_{s,q}^2 \int_{B_R} |v-w|^{-(d+2s)+2\alpha} \frac{\left( \int_{w' \perp v-w} (1+|w'|)^{-q} |w'|^{\gamma+2s+1} dw' \right)^2}{\int_{w' \in \mathbb{R}^d} (f(v+w') + f(w+w')) |w'|^{\gamma+2s} dw'} dw \\ &\leq CC_{s,q}^2 \int_{B_R} |v-w|^{-d+2(\alpha-s)} \frac{1}{\int_{|w'| \geq 1} (f(v+w') + f(w+w')) dw'} dw \\ &\leq Cm_0^{-1} C_{s,q}^2 R^{2(\alpha-s)}. \end{aligned}$$

This proves (2.6) for any solution  $f$  of (1.1) in  $Q_R$  satisfying (1.10) and (2.7), or in general for a solution  $f$  of (1.1) in  $(0, T) \times \mathbb{R}^d \times \mathbb{R}^d$  satisfying (1.10), and  $f_0(x, v) \leq N_q(1 + |v|)^{-q}$  if  $\gamma \leq 0$ .

**2.4. Notation.** We define the kinetic domains for time, space and velocity respecting the scaling of the equations. On the one hand, equation (1.1) is scaling-invariant. Specifically, for any  $r \in [0, 1]$  the scaled function

$$f_r(t, x, v) = f(r^{2s}t, r^{1+2s}x, rv)$$

satisfies the Boltzmann equation with scaled coefficients. On the other hand, (1.1) is invariant under Galilean transformations

$$z \rightarrow z_0 \circ z = (t_0 + t, x_0 + x + tv_0, v_0 + v)$$

with  $z_0 = (t_0, x_0, v_0) \in \mathbb{R}^{1+2d}$ . If  $f$  is a solution of (1.1), then its Galilean transformation  $f_{z_0}(z) = f(z_0 \circ z)$  solves (1.1) with correspondingly translated coefficients. In view of these invariances we define kinetic cylinders

$$Q_r(z_0) := \{(t, x, v) : -r^{2s} \leq t - t_0 \leq 0, |v - v_0| < r, |x - x_0 - (t - t_0)v_0| < r^{1+2s}\},$$

for  $r > 0$  and  $z_0 = (t_0, x_0, v_0) \in \mathbb{R}^{1+2d}$ .

### 3. TAIL BOUND ON UPPER LEVEL SETS

In this section we establish the non-local-to-local bound from Proposition 3.1 in [14] for the Boltzmann equation. It is in this section that the weak divergence form structure of (1.9) proves useful. Just as in the divergence form case, we derive this non-local-to-local bound on upper level sets. To this end, we consider  $l \in \mathbb{R}_+$ . We check using (1.9) that if  $f$  solves (1.1), then  $f - l$  satisfies

$$\begin{aligned} \partial_t f + v \cdot \nabla_x f &= \frac{1}{2} \int_{\mathbb{R}^d} ((f - l)(w) - (f - l)(v)) (K_f(v, w) + K_f(w, v)) dw \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^d} ((f - l)(v) + (f - l)(w)) (K_f(v, w) - K_f(w, v)) dw \\ &\quad + l \int_{\mathbb{R}^d} (K_f(v, w) - K_f(w, v)) dw \\ (3.1) \quad &\geq \frac{1}{2} \int_{\mathbb{R}^d} ((f - l)(w) - (f - l)(v)) (K_f(v, w) + K_f(w, v)) dw \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^d} ((f - l)(v) + (f - l)(w)) (K_f(v, w) - K_f(w, v)) dw. \end{aligned}$$

The last inequality follows due to the classical cancellation from (1.8) and (1.5) provided that  $l \geq 0$ .

To derive a tail bound on upper level sets, we use the formulation in (3.1). The idea is to construct a concave test function that localises only on the upper level sets. Then, the weak divergence form structure proves to be sufficient to make the terms on the lower level sets vanish. Moreover, the cross terms end up with a sign.

**Proposition 3.1** (Non-local-to-local bound). *For given  $R > 0$ ,  $z_0 \in \mathbb{R}^{1+2d}$ ,  $l \in \mathbb{R}_+$ , and any non-negative super-solution  $f$  of (1.9) in  $[0, T] \times \mathbb{R}^d \times \mathbb{R}^d$  with a kernel  $K_f$  that satisfies (2.3), (2.4), (2.5), (2.6), and such that  $f$  conserves mass, we can find for any  $\zeta \in (0, 1)$ , a constant  $C > 0$  depending on  $\zeta, s, d, \Lambda_0, \mu_0$  such that*

$$\begin{aligned} (3.2) \quad &\int_{Q_{\frac{3R}{4}}(z_0)} \int_{\mathbb{R}^d \setminus B_R(v_0)} (f - l)_+(w) \chi_{f > l}(v) K_f(v, w) dw dv \\ &\leq CR^{-2s} \sup_{Q_R(z_0)} (f - l)_+^{1-\zeta} \int_{Q_R(z_0)} (f - l)_+^\zeta dz. \end{aligned}$$

The proof of this proposition relies on the following preliminary estimates.

**Lemma 3.2.** *Let  $R > 0$  and  $v_0 \in \mathbb{R}^d$ . Let  $\eta \in C_c^\infty(\mathbb{R}^d)$  be some smooth function with support in  $B_R(v_0)$ . Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be non-negative,  $l \geq 0$ ,  $f_{l, \varepsilon} = (f - l)_+ + \varepsilon$  for some  $\varepsilon > 0$  and  $\zeta \in (0, 1)$ . Then there holds:*

*i. First, we have for any  $v, w$  in  $f(v) > l \cap f(w) > l$*

$$(3.3) \quad [f(v) - f(w)] \left[ f_{l, \varepsilon}^{-(1-\zeta)}(v) - f_{l, \varepsilon}^{-(1-\zeta)}(w) \right] \leq -\frac{4(1-\zeta)}{\zeta^2} \left( f_{l, \varepsilon}^{\frac{\zeta}{2}}(v) - f_{l, \varepsilon}^{\frac{\zeta}{2}}(w) \right)^2.$$

ii. Second, for any  $v, w$  in  $f(v) > l \cap f(w) > l$

$$(3.4) \quad \min \left\{ f_{l,\varepsilon}^{-(1-\zeta)}(v), f_{l,\varepsilon}^{-(1-\zeta)}(w) \right\} |f(v) - f(w)| \leq \frac{2}{\zeta} \max \left\{ f_{l,\varepsilon}^{\frac{\zeta}{2}}(v), f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right\} \left| f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) - f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right|,$$

iii. Third, for any  $v, w$  in  $f(v) > l \cap f(w) > l$

$$(3.5) \quad \min\{(f-l)_+(v), (f-l)_+(w)\} \left| f_{l,\varepsilon}^{-(1-\zeta)}(v) - f_{l,\varepsilon}^{-(1-\zeta)}(w) \right| \\ \leq \frac{\zeta}{2(1-\zeta)} \left| f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) - f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right| \min \left\{ f_{l,\varepsilon}^{\frac{\zeta}{2}}(v), f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right\}.$$

iv. Fourth,

$$(3.6) \quad \frac{1}{2} \left| \left( \eta f_{l,\varepsilon}^{\frac{\zeta}{2}} \right) (v) - \left( \eta f_{l,\varepsilon}^{\frac{\zeta}{2}} \right) (w) \right|^2 - (\eta(v) - \eta(w))^2 f_{l,\varepsilon}^{\zeta}(w) \leq \min \{ \eta^2(v), \eta^2(w) \} \left| f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) - f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right|^2.$$

and

$$(3.7) \quad \frac{1}{2} \max \{ \eta^2(v), \eta^2(w) \} \left| f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) - f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right|^2 \leq \left| \left( \eta f_{l,\varepsilon}^{\frac{\zeta}{2}} \right) (v) - \left( \eta f_{l,\varepsilon}^{\frac{\zeta}{2}} \right) (w) \right|^2 + (\eta(v) - \eta(w))^2 f_{l,\varepsilon}^{\zeta}(v).$$

*Proof of Lemma 3.2.* For a proof of (3.4), (3.6), (3.7) we refer the reader to [14, Lemma 3.3]. It remains to establish (3.5). To prove (3.5), if we use

$$\tilde{\mathcal{F}}(\xi) := -\xi^{-\frac{\zeta}{2(1-\zeta)}}, \quad \text{with} \quad \tilde{\mathcal{F}}'(\xi) := \frac{\zeta}{2} \frac{1}{1-\zeta} \xi^{-\frac{(2-\zeta)}{2(1-\zeta)}},$$

so that for all  $v, w$  in  $f(v) > l \cap f(w) > l$ , we note by Cauchy's mean value theorem

$$\frac{\left| f_{l,\varepsilon}^{-(1-\zeta)}(v) - f_{l,\varepsilon}^{-(1-\zeta)}(w) \right|}{\left| f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) - f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right|} = \frac{\left| f_{l,\varepsilon}^{-(1-\zeta)}(v) - f_{l,\varepsilon}^{-(1-\zeta)}(w) \right|}{\left| \tilde{\mathcal{F}}(f_{l,\varepsilon}^{-(1-\zeta)}(w)) - \tilde{\mathcal{F}}(f_{l,\varepsilon}^{-(1-\zeta)}(v)) \right|} \\ \leq \max \left\{ \frac{1}{\tilde{\mathcal{F}}'(f_{l,\varepsilon}^{-(1-\zeta)}(v))}, \frac{1}{\tilde{\mathcal{F}}'(f_{l,\varepsilon}^{-(1-\zeta)}(w))} \right\} \\ \leq \frac{2(1-\zeta)}{\zeta} \max \left\{ f_{l,\varepsilon}^{-\frac{(2-\zeta)}{2}}(v), f_{l,\varepsilon}^{-\frac{(2-\zeta)}{2}}(w) \right\}.$$

Multiplied by  $\min\{(f-l)_+(v), (f-l)_+(w)\}$  yields (3.5).  $\square$

*Proof of Proposition 4.1.* Our aim is to get a local bound on the tail of a level set function. To this end we do a concavity estimate on level sets by testing the equation with the level set of the solution to some inverse power. This creates *cross* terms, which have the good sign due to the concavity of the test function. Moreover, we only localise on upper level sets, which makes the terms on lower level sets vanish due to the weak divergence structure.

Due to the translation invariance of (1.1), we may, with no loss in generality, consider  $z_0 = (0, 0, 0)$ . Then we consider  $0 \leq \psi$  given by

$$\psi_l(t, x, v) := \eta^2(t, x, v) \chi_{f>l}(t, x, v) + \chi_{f<l}(t, x, v),$$

for  $l \in \mathbb{R}_+$ , and where  $\eta \in C_c^\infty(\mathbb{R}^{2d+1})$  is such that  $\eta = 1$  in  $Q_{\frac{3R}{4}}$  and  $\eta = 0$  outside  $Q_{\frac{7R}{8}}$ . In particular, for any  $v, w \in \mathbb{R}^d$  there holds:

- i. If  $f(v), f(w) > l$  then  $\psi_l(v) = \eta^2(v)$  and  $\psi_l(w) = \eta^2(w)$ .
- ii. If  $f(v), f(w) < l$  then  $\psi_l(v) = \psi_l(w) = 1$ .



iii. If  $f(v) > l > f(w)$  then  $\psi_l(v) = \eta^2(v) \leq 1 = \psi_l(w)$ .

iv. If  $f(w) > l > f(v)$  then  $\psi_l(w) = \eta^2(w) \leq 1 = \psi_l(v)$ .

We test (3.1) with  $f_{l,\varepsilon}^{-(1-\zeta)} \psi_l$  for any  $l \in \mathbb{R}_+$  and  $\varepsilon > 0$ .

*Step 1: Non-local operator.*

We first consider for fixed  $(t, x)$  the right hand side of (3.1) after testing with  $f_{l,\varepsilon}^{-(1-\zeta)} \psi_l$ :

$$\begin{aligned} \mathcal{E} &:= \mathcal{E} \left( f - l, \psi f_{l,\varepsilon}^{-(1-\zeta)} \right) \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} [(f-l)(v)K_f(w, v) - (f-l)(w)K_f(v, w)] \psi_l(v) f_{l,\varepsilon}^{-(1-\zeta)}(v) \, dw \, dv \\ &= \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} [(f-l)(v)K_f(w, v) - (f-l)(w)K_f(v, w)] \left[ \psi_l(v) f_{l,\varepsilon}^{-(1-\zeta)}(v) - \psi_l(w) f_{l,\varepsilon}^{-(1-\zeta)}(w) \right] \, dw \, dv. \end{aligned}$$

We then split  $\mathcal{E}$  into three parts: we introduce

$$\begin{aligned} \chi_{up}(v, w) &:= \chi_{f(v) > l} \chi_{f(w) > l}, \\ \chi_{low}(v, w) &:= \chi_{f(v) < l} \chi_{f(w) < l}, \\ \chi_{cross}(v, w) &:= \chi_{f(v) > l} \chi_{f(w) < l} + \chi_{f(v) < l} \chi_{f(w) > l}, \end{aligned}$$

so that

$$\begin{aligned} \mathcal{E} &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} [(f-l)(v)K_f(w, v) - (f-l)(w)K_f(v, w)] \psi_l(v) f_{l,\varepsilon}^{-(1-\zeta)}(v) \chi_{up}(v, w) \, dw \, dv \\ &\quad + \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} [(f-l)(v)K_f(w, v) - (f-l)(w)K_f(v, w)] \psi_l(v) f_{l,\varepsilon}^{-(1-\zeta)}(v) \chi_{low}(v, w) \, dw \, dv \\ &\quad + \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} [(f-l)(v)K_f(w, v) - (f-l)(w)K_f(v, w)] \psi_l(v) f_{l,\varepsilon}^{-(1-\zeta)}(v) \chi_{cross}(v, w) \, dw \, dv \\ &=: \mathcal{E}_{up} + \mathcal{E}_{low} + \mathcal{E}_{cross}. \end{aligned} \tag{3.8}$$

*Step 1-i.: Non-locality on lower level sets.*

As a consequence of the weak divergence form structure of (1.9), we see that the choice of  $\psi_l$  implies through observation ii, that

$$\begin{aligned} \mathcal{E}_{low} &= \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} [(f-l)(v)K_f(w, v) - (f-l)(w)K_f(v, w)] \\ &\quad \times \left[ \psi_l(v) f_{l,\varepsilon}^{-(1-\zeta)}(v) - \psi_l(w) f_{l,\varepsilon}^{-(1-\zeta)}(w) \right] \chi_{low}(v, w) \, dw \, dv \\ &= 0. \end{aligned} \tag{3.9}$$

*Step 1-ii.: Cross non-locality.*

Moreover, we claim that the cross term has a sign:

$$\begin{aligned}
\mathcal{E}_{cross} &= \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \\
&\quad \times [\psi_l(v)f_{l,\varepsilon}^{-(1-\zeta)}(v) - \psi_l(w)f_{l,\varepsilon}^{-(1-\zeta)}(w)] \chi_{cross}(v,w) \, dw \, dv \\
(3.10) \quad &= \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \\
&\quad \times [\eta^2(v)(f-l+\varepsilon)^{-(1-\zeta)}(v) - \varepsilon^{-(1-\zeta)}] \chi_{f(v)>l>f(w)} \, dw \, dv \\
&\quad + \frac{1}{2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \\
&\quad \times [\varepsilon^{-(1-\zeta)} - \eta^2(w)(f-l+\varepsilon)^{-(1-\zeta)}(w)] \chi_{f(w)>l>f(v)} \, dw \, dv \\
&\leq 0,
\end{aligned}$$

where the last inequality uses if  $f(v) - l > 0$  then  $-(f(w) - l) > 0$ , and thus the concavity of the test function together with [iii](#) gives a signed integrand. Similarly, if  $-(f(w) - l) < 0$  then  $f(v) - l < 0$  and [iv](#) applies.

*Step 1-iii.: Non-locality on upper level sets.*

To bound  $\mathcal{E}_{up}$ , we first note that  $\mathcal{E}_{up}$  is localised in time and space due to the test function. In fact, if we integrate over time and space on the whole space, the support of  $\eta$  restricts it to a local domain. Thus, we find for any  $(t, x) \in [-R^{2s}, 0] \times B_{R^{1+2s}}$  due to [i](#)

$$\begin{aligned}
\mathcal{E}_{up} &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \psi_l(v)f_{l,\varepsilon}^{-(1-\zeta)}(v) \chi_{up}(v,w) \, dw \, dv \\
(3.11) \quad &= \int_{B_R} \int_{B_R} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \psi_l(v)f_{l,\varepsilon}^{-(1-\zeta)}(v) \chi_{up}(v,w) \, dw \, dv \\
&\quad + \int_{B_R} \int_{\mathbb{R}^d \setminus B_R} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \psi_l(v)f_{l,\varepsilon}^{-(1-\zeta)}(v) \chi_{up}(v,w) \, dw \, dv \\
&=: \mathcal{E}_{up}^{loc} + \mathcal{E}_{up}^{tail}.
\end{aligned}$$

For  $\mathcal{E}_{up}^{tail}$ , we note that there is no singularity, due to the choice of  $\eta$ , which vanishes outside  $B_{\frac{7R}{8}}$ . Thus we find due to [\(2.3\)](#)

$$\begin{aligned}
\mathcal{E}_{up}^{tail} &= \int_{B_R} \int_{\mathbb{R}^d \setminus B_R} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \eta^2(v)f_{l,\varepsilon}^{-(1-\zeta)}(v) \chi_{f(v)>l} \chi_{f(w)>l} \, dw \, dv \\
(3.12) \quad &= \int_{B_R} \int_{\mathbb{R}^d \setminus B_R} (f-l)_+(v) \eta^2(v) f_{l,\varepsilon}^{-(1-\zeta)}(v) K_f(w,v) \chi_{f(v)>l} \chi_{f(w)>l} \, dw \, dv \\
&\quad - \int_{B_R} \int_{\mathbb{R}^d \setminus B_R} (f-l)_+(w) \eta^2(w) f_{l,\varepsilon}^{-(1-\zeta)}(w) K_f(v,w) \chi_{f(v)>l} \chi_{f(w)>l} \, dw \, dv \\
&\leq \Lambda_0 \left(\frac{R}{8}\right)^{-2s} \int_{B_R} ((f-l)_+ + \varepsilon)^\zeta(v) \chi_{f>l}(v) \, dv \\
&\quad - \int_{B_R} \int_{\mathbb{R}^d \setminus B_R} (f-l)_+(w) \eta^2(w) f_{l,\varepsilon}^{-(1-\zeta)}(w) K_f(v,w) \chi_{f(v)>l} \, dw \, dv.
\end{aligned}$$

In particular, the tail of  $(f-l)_+$  has a good sign.

To bound the not-too-non-local non-locality of  $\mathcal{E}_{up}$ , we exploit the concavity of the test function again, due to which we get a signed coercive term, and we show that the terms that have no sign have a singularity of

lower order, and thus can be bounded. Concretely,

$$\begin{aligned}
\mathcal{E}_{up}^{loc} &= \frac{1}{2} \int_{B_R} \int_{B_R} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \\
&\quad \times [\eta^2(v)f_{l,\varepsilon}^{-(1-\zeta)}(v) - \eta^2(w)f_{l,\varepsilon}^{-(1-\zeta)}(w)] \chi_{up}(v,w) \, dw \, dv \\
&= \int_{B_R} \int_{B_R} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \\
&\quad \times [\eta^2(v)f_{l,\varepsilon}^{-(1-\zeta)}(v) - \eta^2(w)f_{l,\varepsilon}^{-(1-\zeta)}(w)] \chi_{up}(v,w) \chi_{f(w)>f(v)} \, dw \, dv \\
&= \int_{B_R} \int_{B_R} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \\
&\quad \times [f_{l,\varepsilon}^{-(1-\zeta)}(v) - f_{l,\varepsilon}^{-(1-\zeta)}(w)] \eta^2(v) \chi_{up}(v,w) \chi_{f(w)>f(v)} \, dw \, dv \\
(3.13) \quad &+ \int_{B_R} \int_{B_R} [(f-l)(v)K_f(w,v) - (f-l)(w)K_f(v,w)] \\
&\quad \times [\eta^2(v) - \eta^2(w)] f_{l,\varepsilon}^{-(1-\zeta)}(w) \chi_{up}(v,w) \chi_{f(w)>f(v)} \, dw \, dv \\
&= \int_{B_R} \int_{B_R} [f_l(v) - f_l(w)] [f_{l,\varepsilon}^{-(1-\zeta)}(v) - f_{l,\varepsilon}^{-(1-\zeta)}(w)] \eta^2(v) K_f(v,w) \chi_{up} \chi_{f(w)>f(v)} \, dw \, dv \\
&\quad + \int_{B_R} \int_{B_R} [K_f(w,v) - K_f(v,w)] f_l(v) [f_{l,\varepsilon}^{-(1-\zeta)}(v) - f_{l,\varepsilon}^{-(1-\zeta)}(w)] \eta^2(v) \chi_{up} \chi_{f(w)>f(v)} \, dw \, dv \\
&\quad + \int_{B_R} \int_{B_R} [f_l(v) - f_l(w)] [\eta^2(v) - \eta^2(w)] f_{l,\varepsilon}^{-(1-\zeta)}(w) K_f(w,v) \chi_{up} \chi_{f(w)>f(v)} \, dw \, dv \\
&\quad + \int_{B_R} \int_{B_R} [K_f(w,v) - K_f(v,w)] [\eta^2(v) - \eta^2(w)] f_l(w) f_{l,\varepsilon}^{-(1-\zeta)}(w) \chi_{up} \chi_{f(w)>f(v)} \, dw \, dv \\
&=: I_{concave}^f + I_{concave}^K + I_{cutoff}^f + I_{cutoff}^K,
\end{aligned}$$

where we denoted  $f_l = f - l$ .

For  $I_{concave}^f$  we use (3.3), (3.6), and (2.3)

$$\begin{aligned}
I_{concave}^f &\leq -\frac{4(1-\zeta)}{\zeta^2} \int_{B_R} \int_{B_R} \left( f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) - f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right)^2 \eta^2(v) K_f(v,w) \chi_{up}(v,w) \chi_{f(w)>f(v)} \, dw \, dv \\
&\leq \frac{4(1-\zeta)}{\zeta^2} \int_{B_R} \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) (\eta(v) - \eta(w))^2 K_f(v,w) \, dw \, dv \\
&\quad - \frac{2(1-\zeta)}{\zeta^2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 K_f(v,w) \chi_{up}(v,w) \chi_{f(w)>f(v)} \, dw \, dv.
\end{aligned}$$

In order to absorb the remaining terms by the signed term, we distinguish the symmetric and the anti-symmetric part, so that due to Young's inequality and (2.6) we find for  $\delta$  sufficiently small

$$\begin{aligned}
I_{concave}^f &\leq CR^{2-2s} \|D_v \eta\|_{L^\infty}^2 \int_{B_R} f_{l,\varepsilon}^\zeta(w) \, dw \\
&\quad - \frac{(1-\zeta)}{\zeta^2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(v,w) + K_f(w,v)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + \frac{(1-\zeta)}{\zeta^2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(w,v) - K_f(v,w)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\leq CR^{-2s} \int_{B_R} f_{l,\varepsilon}^\zeta(w) \, dw \\
&\quad - \frac{(1-\zeta)}{\zeta^2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(v,w) + K_f(w,v)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + \delta \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(w,v) + K_f(v,w)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
(3.14) \quad &\quad + C(\delta, \zeta) \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 \frac{|K_f(w,v) - K_f(v,w)|^2}{(K_f(v,w) + K_f(w,v))} \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\leq CR^{-2s} \int_{B_R} f_{l,\varepsilon}^\zeta(w) \, dw \\
&\quad - \frac{(1-\zeta)}{2\zeta^2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(v,w) + K_f(w,v)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + C(\delta, \zeta) \int_{B_R} \eta^2(v) f_{l,\varepsilon}^\zeta(v) \sup_{v \in B_R} \int_{B_R} \frac{|K_f(w,v) - K_f(v,w)|^2}{(K_f(v,w) + K_f(w,v))} \, dw \, dv \\
&\quad + C(\delta, \zeta) \int_{B_R} \eta^2(w) f_{l,\varepsilon}^\zeta(w) \sup_{w \in B_R} \int_{B_R} \frac{|K_f(w,v) - K_f(v,w)|^2}{(K_f(v,w) + K_f(w,v))} \, dv \, dw \\
&\leq CR^{-2s} \int_{B_R} f_{l,\varepsilon}^\zeta(w) \, dw \\
&\quad - \frac{(1-\zeta)}{2\zeta^2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(v,w) + K_f(w,v)) \chi_{f(w) > f(v) > l} \, dw \, dv.
\end{aligned}$$

To bound  $I_{cutoff}^K$  we use  $|\eta^2(v) - \eta^2(w)| \leq \max\{\eta(v), \eta(w)\} |\eta(v) - \eta(w)|$ ,  $(f-l)_+ \leq f_{l,\varepsilon}$ , Young's inequality, (2.6), and (2.3)

$$\begin{aligned}
I_{cutoff}^K &= \int_{B_R} \int_{B_R} [K_f(w,v) - K_f(v,w)] [\eta^2(v) - \eta^2(w)] (f-l)(w) f_{l,\varepsilon}^{-(1-\zeta)}(w) \chi_{f(w) > f(v) > l} \, dw \, dv \\
(3.15) \quad &\leq \int_{B_R} \int_{B_R} \frac{|K_f(w,v) - K_f(v,w)|^2}{(K_f(v,w) + K_f(w,v))} \max\{\eta^2(v), \eta^2(w)\} f_{l,\varepsilon}^\zeta(w) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + \int_{B_R} \int_{B_R} (K_f(w,v) + K_f(v,w)) |\eta(v) - \eta(w)|^2 f_{l,\varepsilon}^\zeta(w) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\leq C(\mu_0) \int_{B_R} f_{l,\varepsilon}^\zeta(w) \, dw + CR^{2-2s} \|D\eta\|_{L^\infty}^2 \int_{B_R} f_{l,\varepsilon}^\zeta(w) \, dw.
\end{aligned}$$

For  $I_{cutoff}^f$ , we use (3.4), Young's inequality twice, (3.7), (2.3), and (2.6)

$$\begin{aligned}
I_{cutoff}^f &= \int_{B_R} \int_{B_R} [(f-l)(v) - (f-l)(w)] [\eta^2(v) - \eta^2(w)] f_{l,\varepsilon}^{-(1-\zeta)}(w) K_f(w, v) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\leq \frac{2}{\zeta} \int_{B_R} \int_{B_R} \left| f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) - f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right| f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) |\eta^2(v) - \eta^2(w)| K_f(w, v) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\leq \frac{\delta}{2} \int_{B_R} \int_{B_R} \left| f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) - f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right|^2 \max\{\eta^2(v), \eta^2(w)\} K_f(w, v) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + C(\delta) \int_{B_R} \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) |\eta(v) - \eta(w)|^2 K_f(w, v) \chi_{up}(v, w) \chi_{f(w) > f(v)} \, dw \, dv \\
&\leq \delta \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 K_f(w, v) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + \delta \int_{B_R} \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) (\eta(v) - \eta(w))^2 K_f(w, v) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + C(\delta) \|D_v \eta\|_{L^\infty}^2 R^{2-2s} \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) \, dw \\
&\leq \frac{\delta}{2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(v, w) + K_f(w, v)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + \frac{\delta}{2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(w, v) - K_f(v, w)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + C(\delta) R^{-2s} \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) \, dw \\
&\leq \frac{\delta}{2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(v, w) + K_f(w, v)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + C(\delta) \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 \frac{|K_f(v, w) - K_f(w, v)|^2}{(K_f(v, w) + K_f(w, v))} \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + \frac{\delta}{2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(v, w) + K_f(w, v)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + C(\delta) R^{-2s} \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) \, dw \\
&\leq \delta \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(v, w) + K_f(w, v)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + C(\delta) R^{-2s} \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) \, dw.
\end{aligned}
\tag{3.16}$$

Finally, to bound  $I_{concave}^K$ , we use (3.5), Young's inequality, (2.6), (3.7), (2.3)

$$\begin{aligned}
I_{concave}^K &= \int_{B_R} \int_{B_R} [K_f(w, v) - K_f(v, w)] \left[ f_{l,\varepsilon}^{-(1-\zeta)}(v) - f_{l,\varepsilon}^{-(1-\zeta)}(w) \right] f_l(v) \eta^2(v) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\leq \frac{\zeta}{(1-\zeta)} \int_{B_R} \int_{B_R} |K_f(w, v) - K_f(v, w)| \left| f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) - f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right| f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) \eta^2(v) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\leq C(\delta) \int_{B_R} \int_{B_R} \frac{|K_f(w, v) - K_f(v, w)|^2}{(K_f(v, w) + K_f(w, v))} f_{l,\varepsilon}^{\zeta}(v) \eta^2(v) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + \frac{\delta}{2} \int_{B_R} \int_{B_R} (K_f(v, w) + K_f(w, v)) \left| f_{l,\varepsilon}^{\frac{\zeta}{2}}(v) - f_{l,\varepsilon}^{\frac{\zeta}{2}}(w) \right| \eta^2(v) \chi_{f(w) > f(v) > l} \, dw \, dv \\
(3.17) \quad &\leq C \int_{B_R} f_{l,\varepsilon}^{\zeta}(v) \, dv \\
&\quad + \delta \int_{B_R} \int_{B_R} (K_f(v, w) + K_f(w, v)) \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\quad + \delta \int_{B_R} \int_{B_R} (K_f(v, w) + K_f(w, v)) (\eta(v) - \eta(w))^2 f_{l,\varepsilon}^{\zeta}(v) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\leq C \int_{B_R} f_{l,\varepsilon}^{\zeta}(v) \, dv + CR^{2-2s} \|D_v \eta\|_{L^\infty}^2 \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) \, dw \\
&\quad + \delta \int_{B_R} \int_{B_R} (K_f(v, w) + K_f(w, v)) \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 \chi_{f(w) > f(v) > l} \, dw \, dv.
\end{aligned}$$

If we combine (3.13) with (3.14), (3.15), (3.16), (3.17), we find for  $\delta$  sufficiently small

$$\begin{aligned}
\mathcal{E}_{up}^{loc} &\leq CR^{-2s} \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) \, dw \\
&\quad - \frac{(1-\zeta)}{2\zeta^2} \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(v, w) + K_f(w, v)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
(3.18) \quad &\quad + 2\delta \int_{B_R} \int_{B_R} \left( (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(v) - (\eta f_{l,\varepsilon}^{\frac{\zeta}{2}})(w) \right)^2 (K_f(v, w) + K_f(w, v)) \chi_{f(w) > f(v) > l} \, dw \, dv \\
&\leq CR^{-2s} \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) \, dw.
\end{aligned}$$

In particular, this implies together with (3.11) and (3.12)

$$(3.19) \quad \mathcal{E}_{up} \leq CR^{-2s} \int_{B_R} f_{l,\varepsilon}^{\zeta}(w) \, dw - \int_{B_R} \int_{\mathbb{R}^d \setminus B_R} (f-l)_+(w) \eta^2(v) f_{l,\varepsilon}^{-(1-\zeta)}(v) K_f(v, w) \chi_{f(v) > l} \, dw \, dv.$$

*Step 1-iv.: Non-locality.*

We conclude from (3.8), (3.9), (3.10), (3.19),

$$\begin{aligned}
\mathcal{E} = \mathcal{E}_{up} + \mathcal{E}_{low} + \mathcal{E}_{cross} &\leq C(\zeta, \Lambda_0, \mu_0) R^{-2s} \int_{B_R} (f-l+\varepsilon)^\zeta(v) \chi_{f > l} \, dv \\
(3.20) \quad &\quad - \int_{B_R} \int_{\mathbb{R}^d \setminus B_R} (f-l)_+(w) \eta^2(v) f_{l,\varepsilon}^{-(1-\zeta)}(v) K_f(v, w) \chi_{f(v) > l} \, dw \, dv.
\end{aligned}$$

Note that this estimate holds true for any  $(t, x)$ . Moreover, as remarked above (3.11), since the only positive contribution stems from  $\mathcal{E}_{up}$ , the estimate in (3.20) is localised in time and space.

*Step 2.: Transport.*

Finally we use the equation. For the transport operator we use the divergence theorem, the fact that  $f$  conserves mass, integration by parts and use that  $|\mathcal{T}\eta| \sim R^{-2s}$ . We get

$$\begin{aligned}
\int_{[0,T] \times \mathbb{R}^{2d}} \mathcal{T}f(z)\psi_l(z)f_{l,\varepsilon}^{-(1-\zeta)}(z) dz &= \int_{[0,T] \times \mathbb{R}^{2d}} \mathcal{T}f(z)\eta^2(z)\chi_{f>l}(v)f_{l,\varepsilon}^{-(1-\zeta)}(z) dz \\
&\quad + \int_{[0,T] \times \mathbb{R}^{2d}} \mathcal{T}f(z)\chi_{f<l}(z)\varepsilon^{-(1-\zeta)} dz \\
&= \frac{1}{\zeta} \int_{\mathbb{R}^{1+2d}} \mathcal{T}((f-l)_+ + \varepsilon)^\zeta(z)\eta^2(z) dz \\
&\quad + \int_{[0,T] \times \mathbb{R}^{2d}} \mathcal{T}(f-l)_-(z)\varepsilon^{-(1-\zeta)} dz \\
(3.21) \quad &= -\frac{1}{\zeta} \int_{\mathbb{R}^{1+2d}} ((f-l)_+ + \varepsilon)^\zeta(z)\mathcal{T}\eta^2(z) dz \\
&\quad + \int_{\mathbb{R}^{2d}} \left( (f-l)_-(T, x, v) - (f-l)_-(0, x, v) \right) \varepsilon^{-(1-\zeta)} dx dv \\
&\leq C(\zeta)R^{-2s} \int_{Q_R} ((f-l)_+ + \varepsilon)^\zeta(z) dz.
\end{aligned}$$

Thus by (3.1), (3.21), and (3.20)

$$\begin{aligned}
0 &\leq \int_{[0,T] \times \mathbb{R}^{2d}} \mathcal{T}f(z)\psi_l(z)f_{l,\varepsilon}^{-(1-\zeta)}(z) dz + \int_{[0,T] \times \mathbb{R}^{2d}} \mathcal{E}(f-l, \psi_l f_{l,\varepsilon}^{-(1-\zeta)}) dx dt \\
&\leq C(\zeta)R^{-2s} \int_{Q_R} (f-l+\varepsilon)^\zeta(z)\eta^2(z)\chi_{f>l}(z) dz \\
&\quad - \int_{\mathbb{R}^{1+d}} \int_{B_R} \int_{\mathbb{R}^d \setminus B_R} (f-l)_+(w)\eta^2(z)f_{l,\varepsilon}^{-(1-\zeta)}(v)K_f(v,w)\chi_{f(v)>l} dw dz,
\end{aligned}$$

or rearranged

$$\int_{Q_{\frac{3R}{4}}} \int_{\mathbb{R}^d \setminus B_R} (f-l)_+(w)f_{l,\varepsilon}^{-(1-\zeta)}(v)K_f(v,w)\chi_{f(v)>l} dw dz \leq CR^{-2s} \int_{Q_R} (f-l+\varepsilon)^\zeta(z)\eta^2(v)\chi_{f>l}(v) dz.$$

*Step 3.: Conclusion.*

This implies (3.2), if we let  $\varepsilon \rightarrow 0$ , take out the infimum of  $(f-l)^{-(1-\zeta)}$  over  $z \in Q_{\frac{3R}{4}} \cap \chi_{f>l}$  on the left hand side, divide the estimate by the infimum (note  $f-l \in L^\infty(Q_R)$ ), and use  $(\inf g)^{-1} = \sup g^{-1}$ , so that

$$\int_{Q_{\frac{3R}{4}}} \int_{\mathbb{R}^d \setminus B_R} (f-l)_+(w)\chi_{f>l}(v)K_f(v,w) dw dv \leq CR^{-2s} \sup_{Q_{\frac{3R}{4}} \cap f>l} (f-l)^{1-\zeta} \int_{Q_R} (f-l)_+^\zeta dz.$$

□

#### 4. LINEAR $L^1$ TO $L^\infty$ BOUND

We use a De Giorgi argument in  $L^1$  to derive a linear  $L^1$ - $L^\infty$  bound for solutions to (1.1).

**Proposition 4.1** ( $L^1$ - $L^\infty$  bound). *Let  $0 < R \leq 1$ ,  $z_0 \in \mathbb{R}^{1+2d}$ , and let  $f \in L^2([-3, 0] \times B_1; L^\infty(\mathbb{R}^d) + H^s(B_1))$  be a non-negative solution to (1.6) in  $Q_R(z_0)$  with a kernel  $K_f$  and a coefficient  $\Lambda_f$  that satisfy (2.1), (2.2), (2.3), (2.4), (2.5), (2.6). Then there exists a large constant  $C > 1$  depending on  $s, d, \lambda_0, \Lambda_0, \mu_0$  such that*

$$\text{for a.e. } z_1 \in Q_{\frac{R}{8}}(z_0): \quad f(z_1) \leq CR^{-(2d(1+s)+2s)} \int_{Q_R(z_0)} f(z) dz.$$

The proof of this proposition makes use of the gain of integrability stemming from the fractional Kolmogorov equation.

**4.1. Fractional Kolmogorov.** We consider the fractional Kolmogorov equation given by

$$(4.1) \quad \partial_t f + v \cdot \nabla_x f + (-\Delta_v)^s f = h_1 + h_2$$

for some  $h_2 \in L^1([0, \tau] \times \mathbb{R}^{2d})$  and some  $h_1$  such that  $\left\| (-\Delta_v)^{-\frac{s+\epsilon}{2}} h_1 \right\|_{L^1([0, \tau] \times \mathbb{R}^{2d})} < +\infty$  for  $0 \leq \epsilon < s$ .

**Proposition 4.2.** *Let  $0 \leq f$  solve (or be a sub-solution of) (4.1) in  $[0, \tau] \times \mathbb{R}^{2d}$ , with  $f(0, x, v) = f_0(x, v) \in L^1 \cap L^2(\mathbb{R}^{2d})$ , and with  $h = h_1 + h_2$  where  $h_1, h_2 \in L^1 \cap L^2([-\tau, 0] \times \mathbb{R}^{2d})$  such that*

$$\left\| (-\Delta_v)^{-\frac{s+\epsilon}{2}} h_1 \right\|_{L^1([-\tau, 0] \times \mathbb{R}^{2d})} < +\infty.$$

*Then for any  $0 \leq \epsilon < s$  and any  $1 \leq p < 1 + \frac{s-\epsilon}{s+2d(s+1)+\epsilon}$  there holds*

$$\|f\|_{L^p([0, \tau] \times \mathbb{R}^{2d})} \lesssim \tau^{\frac{1}{p}-\alpha_\epsilon} \left\| (-\Delta_v)^{-\frac{s+\epsilon}{2}} h_1 \right\|_{L^1([-\tau, 0] \times \mathbb{R}^{2d})} + \tau^{\frac{1}{p}+\frac{1}{2}-\alpha_0} \|h_2\|_{L^1([0, \tau] \times \mathbb{R}^{2d})} + \tau^{\frac{1}{p}+\frac{1}{2}-\alpha_0} \|f_0\|_{L^1(\mathbb{R}^{2d})},$$

*where  $\alpha_\epsilon = d \left(1 + \frac{1}{s}\right) \left(1 - \frac{1}{p}\right) + \frac{s+\epsilon}{2s}$ .*

We refer the reader to [14, Proposition 4.1].

**4.2. Proof of Proposition 4.1.** We use the tail bound from Proposition 3.1 to get a local energy estimate. Then we use the gain of integrability in  $L^1$  from Proposition 4.2, by comparing the solution of (1.6) to the fractional Kolmogorov equation (4.1). We bound the right hand side by the local energy estimate, and the tail term again with Proposition 3.1. Finally, we iterate the so gained local a priori estimate on level set functions.

*Step 1: Energy estimate.* Let  $\phi \in C_c^\infty(\mathbb{R}^{1+2d})$  be such that  $\phi(t, x, v) = 0$  for  $(t, x, v)$  outside  $Q_{\frac{R}{4}}$  and  $\phi(t, x, v) = 1$  for  $(t, x, v) \in Q_{\frac{R}{8}}$ . We then test (1.6) with  $(f-l)_+ \phi^2$ , so that if we denote by

$$\mathcal{L}f(v) := \int_{\mathbb{R}^d} (f(w) - f(v)) K_f(v, w) dw,$$

we get

$$(4.2) \quad \begin{aligned} & \Lambda_f \int_{\mathbb{R}^{1+2d}} f(z) \phi^2(z) (f-l)_+(z) dz \\ & \geq \frac{1}{2} \int_{\mathbb{R}^{1+2d}} \mathcal{T}(f-l)_+^2(z) \phi^2(z) dz - \int_{\mathbb{R}^{1+2d}} \mathcal{L}f(z) (f-l)_+(z) \phi^2(z) dz. \end{aligned}$$



We observe that for any  $\frac{R}{4} < r$ , if we abbreviate  $(f-l)_+ =: f_{l+}$  and if we denote by  $\Omega_\rho := [-\rho^{2s}, 0] \times B_{\rho^{1+2s}}$ , then

$$\begin{aligned}
& - \int_{\mathbb{R}^{1+2d}} \mathcal{L}f(z)(f-l)_+(z)\phi^2(z) \, dz \\
&= \int_{\mathbb{R}^{1+2d}} \int_{\mathbb{R}^d} [f(t,x,v) - f(t,x,w)]((f-l)_+\phi^2)(t,x,v)K_f(t,x,v,w) \, dw \, dz \\
(4.3) \quad &= \frac{1}{2} \int_{\Omega_{\frac{R}{4}}} \int_{B_r} \int_{B_r} [f(v) - f(w)] [(f_{l+}\phi^2)(v) - (f_{l+}\phi^2)(w)] K_f(v,w) \, dw \, dz \\
&+ \frac{1}{2} \int_{\Omega_{\frac{R}{4}}} \int_{B_r} \int_{B_r} [f(v) - f(w)] [(f_{l+}\phi^2)(v) + (f_{l+}\phi^2)(w)] K_f(v,w) \, dw \, dx \, dt \\
&+ \int_{\Omega_{\frac{R}{4}}} \int_{B_r} \int_{\mathbb{R}^d \setminus B_r} [f(v) - f(w)] (f_{l+}\phi^2)(z) K_f(v,w) \, dw \, dz.
\end{aligned}$$

*Step 1-i.: Transport.* We integrate by parts the transport term, and use  $\mathcal{T}\phi \sim -R^{-2s}$ .

$$\begin{aligned}
(4.4) \quad & -\frac{1}{2} \int_{\mathbb{R}^{1+2d}} \mathcal{T}(f-l)_+^2(z)\phi^2(z) \, dz = \frac{1}{2} \int_{\mathbb{R}^{1+2d}} (f-l)_+^2(t,x,v)\mathcal{T}\phi^2(t,x,v) \, dz \\
&\leq CR^{-2s} \int_{Q_{\frac{R}{4}}} (f-l)_+^2(z) \, dz.
\end{aligned}$$

*Step 1-ii.: Tail bound.* We use Proposition 3.1 for  $f$ , which we assume to be a solution, in particular a super-solution, so that for  $r = \frac{R}{2}$

$$\begin{aligned}
(4.5) \quad & \int_{\Omega_{\frac{R}{4}}} \int_{B_r} \int_{\mathbb{R}^d \setminus B_r} [f(w) - f(v)] (f_{l+}\phi^2)(z) K_f(v,w) \, dw \, dz \\
&\leq \sup_{Q_{\frac{R}{4}}} (f-l)_+ \int_{Q_{\frac{R}{4}}} \int_{\mathbb{R}^d \setminus B_{\frac{R}{2}}} (f-l)_+(w) K_f(v,w) \chi_{f>i}(v) \, dw \, dz \\
&\leq CR^{-2s} \sup_{Q_{\frac{R}{4}}} (f-l)_+ \sup_{Q_{\frac{R}{2}}} (f-l)_+^{1-\zeta} \int_{Q_{\frac{R}{2}}} (f-l)_+^\zeta(z) \, dz \\
&\leq CR^{-2s} \left( \sup_{Q_{\frac{R}{2}}} (f-l)_+ \right)^{2-\zeta} \int_{Q_{\frac{R}{2}}} (f-l)_+^\zeta(z) \, dz.
\end{aligned}$$

We used  $-f(v) < -l$  in the first inequality, and Young's inequality in the last inequality.

*Step 1-iii.: Not-too-non-local operator.* What remains to be estimated is the local contribution of the non-local operator,

$$\begin{aligned}
\mathcal{E}_{loc} &:= \frac{1}{2} \int_{B_r} \int_{B_r} [f(v) - f(w)] [(f_{l+}\phi^2)(v) - (f_{l+}\phi^2)(w)] K_f(v,w) \, dw \, dv \\
&+ \frac{1}{2} \int_{B_r} \int_{B_r} [f(v) - f(w)] [(f_{l+}\phi^2)(v) + (f_{l+}\phi^2)(w)] K_f(v,w) \, dw \, dv =: \mathcal{E}_{loc}^{sym} + \mathcal{E}_{loc}^{skew},
\end{aligned}$$

where we recall  $f_{l+} = (f-l)_+$ .

*Claim 1.* For  $z = (t, x, v)$  and  $r = \frac{R}{2}$

$$(4.6) \quad \begin{aligned} \mathcal{E}_{loc}^{sym} &\geq \frac{1}{2} \int_{B_r} \int_{B_r} [(f_{i_+} \phi)(v) - (f_{i_+} \phi)(w)]^2 K_f(v, w) \, dv - Cr^{-2s} \int_{B_r} f_{i_+}^2(v) \, dv \\ &\quad + \frac{1}{2} \int_{B_r} \int_{B_r} [f_{i_-}(v) - f_{i_-}(w)] [(f_{i_+} \phi^2)(v) - (f_{i_+} \phi^2)(w)] K_f(v, w) \, dw \, dv. \end{aligned}$$

See [14, Claim (4.5)].

*Claim 2.* For any  $\delta_0 \in (0, 1)$ , there exists a constant  $C > 0$  depending on  $\delta_0$  such that

$$(4.7) \quad \begin{aligned} \mathcal{E}_{loc}^{skew} &\geq \frac{1}{2} \int_{B_r} \int_{B_r} [f_{i_-}(v) - f_{i_-}(w)] [(f_{i_+} \phi^2)(v) + (f_{i_+} \phi^2)(w)] K_f(v, w) \, dw \, dv \\ &\quad - CR^{-2s} \int_{B_{\frac{R}{4}}} f_{i_+}^2(v) \, dv - \left(\frac{1}{4} + \delta_0\right) \int_{B_r} \int_{B_r} [(f_{i_+} \phi)(v) - (f_{i_+} \phi)(w)]^2 K_f(v, w) \, dw \, dv. \end{aligned}$$

To prove (4.7), we note

$$\begin{aligned} [f_{i_+}(v) - f_{i_+}(w)] [(f_{i_+} \phi^2)(v) + (f_{i_+} \phi^2)(w)] \\ = [(f_{i_+}^2 \phi^2)(v) - (f_{i_+}^2 \phi^2)(w)] + f_{i_+}(v) f_{i_+}(w) [\phi^2(w) - \phi^2(v)]. \end{aligned}$$

Together with  $f_{i_-}(v) - f_{i_-}(w) = f(v) - f(w) - (f_{i_+}(v) - f_{i_+}(w))$ , this implies

$$(4.8) \quad \begin{aligned} &\int_{B_r} \int_{B_r} [f(v) - f(w)] [(f_{i_+} \phi^2)(v) + (f_{i_+} \phi^2)(w)] K_f(v, w) \, dw \, dv \\ &= \int_{B_r} \int_{B_r} [f_{i_-}(v) - f_{i_-}(w)] [(f_{i_+} \phi^2)(v) + (f_{i_+} \phi^2)(w)] K_f(v, w) \, dw \, dv \\ &\quad + \int_{B_r} \int_{B_r} \left\{ [(f_{i_+}^2 \phi^2)(v) - (f_{i_+}^2 \phi^2)(w)] + f_{i_+}(v) f_{i_+}(w) [\phi^2(w) - \phi^2(v)] \right\} K_f(v, w) \, dw \, dv. \end{aligned}$$

Then we use the cancellation (2.4)

$$(4.9) \quad \begin{aligned} &\int_{B_r} \int_{B_r} [(f_{i_+} \phi)^2(w) - (f_{i_+} \phi)^2(v)] K_f(v, w) \, dw \, dv \\ &= \int_{B_r} \int_{B_r} (f_{i_+} \phi)^2(w) [K_f(v, w) - K_f(w, v)] \, dw \, dv \leq \Lambda_0 \int_{B_r} (f_{i_+}^2 \phi^2)(w) \, dw. \end{aligned}$$

Moreover, we bound with Young's inequality, on the one hand, and using  $-2ab = (a-b)^2 - a^2 - b^2$ , on the other hand,

$$\begin{aligned} &f_{i_+}(v) f_{i_+}(w) [\phi^2(v) - \phi^2(w)] \\ &\leq \frac{1}{2} [f_{i_+}^2(v) + f_{i_+}^2(w)] \phi^2(v) + \frac{1}{2} (f_{i_+}(v) - f_{i_+}(w))^2 \phi^2(w) - \frac{1}{2} [f_{i_+}^2(v) + f_{i_+}^2(w)] \phi^2(w) \\ &= \frac{1}{2} [f_{i_+}^2(v) + f_{i_+}^2(w)] [\phi^2(v) - \phi^2(w)] + \frac{1}{2} (f_{i_+}(v) - f_{i_+}(w))^2 \phi^2(w). \end{aligned}$$

Then, using the proof of (3.7) and Young's inequality, we see for any  $\delta_0 \in (0, 1)$

$$\frac{1}{2} (f_{i_+}(v) - f_{i_+}(w))^2 \phi^2(w) \leq \left(\frac{1}{2} + \delta_0\right) [(f_{i_+} \phi)(v) - (f_{i_+} \phi)(w)]^2 + C(\delta_0) (\phi(v) - \phi(w))^2 f_{i_+}^2(v).$$

Thus, by doing a Taylor expansion of  $\phi$ , using the upper bound (2.3) for  $s \in (0, 1/2)$ , or the cancellation (2.5) and the upper bound (2.3) for  $s \in [1/2, 1)$ , and finally using the definition of  $\eta$ , we find

$$\begin{aligned}
& \int_{B_r} \int_{B_r} f_{l_+}(v) f_{l_+}(w) [\phi^2(v) - \phi^2(w)] K_f(v, w) \, dw \, dv \\
& \leq \frac{1}{2} \int_{B_r} \int_{B_r} [f_{l_+}^2(v) + f_{l_+}^2(w)] [\phi^2(v) - \phi^2(w)] K_f(v, w) \, dw \, dv \\
& \quad + \frac{1}{2} \int_{B_r} \int_{B_r} (f_{l_+}(v) - f_{l_+}(w))^2 \phi^2(w) K_f(v, w) \, dw \, dv \\
& \leq \frac{1}{2} \int_{B_r} \int_{B_r} [f_{l_+}^2(v) D_v \phi^2(v) + f_{l_+}^2(w) D_v \phi^2(w)] (v - w) K_f(v, w) \, dw \, dv \\
& \quad + \|D_v^2 \phi^2\|_{L^\infty} \int_{B_r} \int_{B_r} [f_{l_+}^2(v) + f_{l_+}^2(w)] |v - w|^2 K_f(v, w) \, dw \, dv \\
& \quad + \left(\frac{1}{2} + \delta_0\right) \int_{B_r} \int_{B_r} [(f_{l_+} \phi)(v) - (f_{l_+} \phi)(w)]^2 K_f(v, w) \, dw \, dv \\
& \quad + C(\delta_0) \int_{B_r} \int_{B_r} (\phi(v) - \phi(w))^2 f_{l_+}^2(v) K_f(v, w) \, dw \, dv \\
& \leq \frac{1}{2} \int_{B_r} \int_{B_r} f_{l_+}^2(v) D_v \phi^2(v) (v - w) [K_f(v, w) - K_f(w, v)] \, dw \, dv \\
& \quad + CR^{2-2s} \|D_v^2 \phi^2\|_{L^\infty} \int_{B_r} f_{l_+}^2(v) \, dv \\
& \quad + \left(\frac{1}{2} + \delta_0\right) \int_{B_r} \int_{B_r} [(f_{l_+} \phi)(v) - (f_{l_+} \phi)(w)]^2 K_f(v, w) \, dw \, dv \\
& \leq CR^{1-2s} \|D_v \phi^2\|_{L^\infty} \int_{B_r} f_{l_+}^2(v) \, dv + CR^{-2s} \int_{B_r} f_{l_+}^2(v) \, dv \\
& \quad + \left(\frac{1}{2} + \delta_0\right) \int_{B_r} \int_{B_r} [(f_{l_+} \phi)(v) - (f_{l_+} \phi)(w)]^2 K_f(v, w) \, dw \, dv
\end{aligned} \tag{4.10}$$

Combining (4.8) with (4.9) and (4.10) yields (4.7).

Thus, from the claims in (4.6) and (4.7), we infer for  $\delta_0$  sufficiently small ( $\delta_0 < \frac{1}{8}$ ) and for  $r = \frac{R}{2}$

$$\begin{aligned}
\mathcal{E}_{loc}^{sym} + \mathcal{E}_{loc}^{skew} & \geq \frac{1}{2} \int_{B_r} \int_{B_r} [(f_{l_+} \phi)(v) - (f_{l_+} \phi)(w)]^2 K_f(v, w) \, dv - Cr^{-2s} \int_{B_r} f_{l_+}^2(v) \, dv \\
& \quad + \frac{1}{2} \int_{B_r} \int_{B_r} [f_{l_-}(v) - f_{l_-}(w)] [(f_{l_+} \phi^2)(v) - (f_{l_+} \phi^2)(w)] K_f(v, w) \, dw \, dv \\
& \quad + \frac{1}{2} \int_{B_r} \int_{B_r} [f_{l_-}(v) - f_{l_-}(w)] [(f_{l_+} \phi^2)(v) + (f_{l_+} \phi^2)(w)] K_f(v, w) \, dw \, dv \\
& \quad - CR^{-2s} \int_{B_{\frac{R}{4}}} f_{l_+}^2(v) \, dv \\
& \quad - \left(\frac{1}{4} + \delta_0\right) \int_{B_r} \int_{B_r} [(f_{l_+} \phi)(v) - (f_{l_+} \phi)(w)]^2 K_f(v, w) \, dw \, dv \\
& \geq \frac{1}{8} \int_{B_r} \int_{B_r} [(f_{l_+} \phi)(v) - (f_{l_+} \phi)(w)]^2 K_f(v, w) \, dv - Cr^{-2s} \int_{B_r} f_{l_+}^2(v) \, dv.
\end{aligned} \tag{4.11}$$

*Step 1-iv.: Conclusion.* Combining (4.2) with (4.4), (4.5), and (4.11), we conclude

$$(4.12) \quad \begin{aligned} & \frac{1}{8} \int_{\Omega_{\frac{R}{4}}} \iint_{B_{\frac{R}{2}} \times B_{\frac{R}{2}}} \left[ ((f-l)_+\phi)(v) - ((f-l)_+\phi)(w) \right]^2 K_f(v, w) \, dw \, dv \, dx \, dt \\ & \leq CR^{-2s} \int_{Q_{\frac{R}{4}}} (f-l)_+^2(z) \, dz + CR^{-2s} (\sup_{Q_{\frac{R}{2}}} f)^{2-\zeta} \int_{Q_{\frac{R}{2}}} (f-l)_+^\zeta(z) \, dz + \mu_0 \sup_{Q_{\frac{R}{4}}} f \int_{Q_{\frac{R}{4}}} (f-l)_+ \, dz. \end{aligned}$$

Note that we used (2.1) to bound  $\Lambda_f$ .

*Step 2 & 3: Gain of Integrability & De Giorgi iteration.* These steps carry over verbally from [14, Proof of Prop. 3.1], if we work with the Boltzmann equation in the formulation (1.6). Then we have the following gain of integrability

$$(4.13) \quad \begin{aligned} \|(f-l)_+\phi\|_{L^p(Q_{\frac{R}{8}})} & \leq CR^{s+\epsilon} \|(f-l)_+(t_0)\|_{L^1(Q_{\frac{R}{4}})} + CR^{-s+\epsilon} \|(f-l)_+\|_{L^1(Q_{\frac{R}{2}})} \\ & \quad + CR^{-s+\epsilon} \left| \{f > l\} \cap Q_{\frac{R}{2}} \right|^{\frac{1}{2}} \|(f-l)_+\|_{L^2(Q_{\frac{R}{2}})} \\ & \quad + CR^{-s+\epsilon} \left( \sup_{Q_R} (f-l)_+ \right)^{\frac{2-\zeta}{2}} \left( \int_{Q_R} (f-l)_+^\zeta(z) \, dz \right)^{\frac{1}{2}} |\{f > l\} \cap Q_R|^{\frac{1}{2}} \\ & \quad + CR^{-s+\epsilon} \sup_{Q_R} (f-l)_+^{1-\zeta} \int_{Q_R} (f-l)_+^\zeta \, dz \\ & \quad + CR^{s+\epsilon} \sup_{Q_R} f |\{f > l\} \cap Q_R| + CR^\epsilon (\sup_{Q_R} f)^{\frac{1}{2}} |\{f > l\} \cap Q_R|^{\frac{1}{2}} \|(f-l)_+\|_{L^1}^{\frac{1}{2}}, \end{aligned}$$

with which we conclude for almost every  $(t, x, v) \in Q_{\frac{R}{8}}$

$$f(z) \leq L = \delta(1 + R^{2s}) \sup_{Q_R} f + R^{-\frac{(s-\epsilon)p}{p-1}} 2^{\frac{4p^2}{(p-1)^2}} \delta^{-\frac{2-\zeta}{\zeta} \frac{p}{p-1}} \int_{Q_R} f(z) \, dz.$$

For  $R \leq 1$ , we absorb the first term on the right hand side with a standard iteration argument, concluding the proof of Proposition 4.1.

### 4.3. Strong Harnack for Boltzmann.

*Proof of Theorem 1.1.* Let  $f$  solve (1.1) in  $(0, T) \times \mathbb{R}^d \times \mathbb{R}^d$ , and assume (1.10) holds. Then, in particular,  $K_f$  and  $\Lambda_f$  given in (1.4) and (1.7) satisfy (2.1)-(2.5). Moreover, due to our notion of solutions in Definition 2.1, also (2.6) is satisfied.

Let  $\mathcal{I}^- := (\tau_0, \tau_1)$  and  $\mathcal{I}^+ := (\tau_2, \tau_3)$  be two disjoint compactly contained subsets of  $(0, T)$  such that  $\tau_2 - \tau_1 \geq r_0^{2s}$  for sufficiently small  $r_0 < \frac{1}{6}$ . As a consequence of Proposition 4.1, Young's inequality and the Weak Harnack inequality [6, Theorem 1.6], we obtain for any  $\delta \in (0, 1)$  and for  $\zeta \in (0, 1)$  from the Weak Harnack inequality,

$$\begin{aligned} \sup_{\mathcal{I}^- \times Q_{\frac{r_0}{4}}} f & \leq Cr_0^{-(2d(1+s)+2s)} \|f\|_{L^1(\mathcal{I}^- \times Q_{2r_0}^t)} \leq \delta \sup_{\mathcal{I}^- \times Q_{\frac{r_0}{4}}} f + C(\delta) r_0^{-\frac{(2d(1+s)+2s)}{\zeta}} \|f\|_{L^\zeta(\mathcal{I}^- \times Q_{2r_0}^t)} \\ & \leq \delta \sup_{\mathcal{I}^- \times Q_{\frac{r_0}{4}}} f + C(\delta) r_0^{-\frac{(2d(1+s)+2s)}{\zeta}} \inf_{\mathcal{I}^+ \times Q_{2r_0}^t} f. \end{aligned}$$

We recall that there is no source term appearing after applying the Weak Harnack inequality, due to the positivity of  $Q_2$  (1.8), which implies that  $f$  is a non-negative super-solution to (1.6) with zero source term. Absorbing the first term on the left hand side with a standard iteration argument concludes the proof of Theorem 1.1.  $\square$

## 5. BRIEF NOTE ON BOUNDS OF THE FUNDAMENTAL SOLUTION

We end this article with a short remark on how to adapt Aronson's method for non-local hypoelliptic equations in divergence form to the case of more general kernels that satisfy the ellipticity assumptions (2.2)-(2.6) inspired from the Boltzmann collision kernel.

**5.1. Results.** As a consequence of Theorem 1.1, we derive polynomial upper and exponential lower bounds on the fundamental solution of (1.6) with coefficients  $K_h$  and  $\Lambda_h$  given by (1.4) and (1.7), respectively, for a fixed function  $h \geq 0$  that satisfies (1.10). To give sense to the next three theorems, we assume existence of a non-negative measurable function  $J$ , which is the fundamental solution of (1.6) linearised around a fixed function  $g$ , connecting a given point  $(t, x, v) \in \mathbb{R}^{1+2d}$  with  $(\tau, y, w) \in \mathbb{R}^{1+2d}$ , in the sequel denoted by

$$J(t, x, v; \tau, y, w) = J(t - \tau, x - y - (t - \tau)w, v - w) =: J((\tau, y, w)^{-1} \circ (t, x, v)),$$

where  $\circ$  denotes the Galilean translation, that is  $(t_0, x_0, v_0) \circ (t, x, v) = (t_0 + t, x_0 + x + tv_0, v_0 + v)$ , which respects the translation invariance of (1.1). Moreover, we assume that the fundamental solution  $J$  has the following properties:

i. For every  $t \in \mathbb{R}_+$  there holds the normalisation

$$(5.1) \quad \int_{\mathbb{R}^{2d}} J(t, x, v) dx dv = 1.$$

ii. There holds  $J \geq 0$ , and for all  $(t, x, v), (\tau, y, w) \in \mathbb{R}_+ \times \mathbb{R}^{2d}$  a form of symmetry

$$(5.2) \quad J((\tau, y, w)^{-1} \circ (t, x, v)) = J((\tau, x, v)^{-1} \circ (t, y, w)).$$

iii. For any  $0 \leq \tau < \sigma < t < T$  and any  $(x, v), (y, w) \in \mathbb{R}^{2d}$  the Chapman-Kolmogorov identity holds

$$(5.3) \quad J(t, x, v; \tau, y, w) = \int_{\mathbb{R}^{2d}} J(t, x, v; \sigma, \varphi, \xi) J(\sigma, \varphi, \xi; \tau, y, w) d\varphi d\xi.$$

Then, we deduce, on the one hand, polynomial upper bounds.

**Theorem 5.1** (Polynomial upper bounds on the fundamental solution). *Let  $x, v, y_0, w_0 \in \mathbb{R}^d$ , and  $0 \leq \tau_0 < \sigma < T$ . Let  $J$  be the fundamental solution of (1.6) in  $[0, T] \times \mathbb{R}^{2d}$  with coefficients  $K_h$  and  $\Lambda_h$  given by (1.4) and (1.7), respectively, for a fixed function  $h \geq 0$  that satisfies (1.10). Assume  $J$  satisfies (5.1), (5.2) and (5.3). Then there exists  $C > 0$  depending on  $s, \gamma, d, m_0, M_0, E_0, H_0$  (note that these constants refer to the mass, energy and entropy of  $h$ ), such that*

$$(5.4) \quad \begin{aligned} & J(\sigma, x, v; \tau_0, y_0, w_0) \\ & \leq C(\sigma - \tau_0)^{-\frac{2d(1+s)}{2s}} \left[ 1 + \frac{\max \left\{ |v - w_0|^{2s}, |x - y_0 - (\sigma - \tau_0)(v - w_0)|^{\frac{2s}{1+2s}} \right\}}{\sigma - \tau_0} \right]^{-\frac{s}{4s}}. \end{aligned}$$

On the other hand, we derive an exponential lower bound.

**Theorem 5.2** (Exponential lower bounds on the fundamental solution). *Let  $x, v, y_0, w_0 \in \mathbb{R}^d$ , and  $0 \leq \tau_0 < \sigma < T$ . Let  $J$  be the fundamental solution of (1.6) in  $[0, T] \times \mathbb{R}^{2d}$  with coefficients  $K_h$  and  $\Lambda_h$  given by (1.4) and (1.7), respectively, for a fixed function  $h \geq 0$  that satisfies (1.10). Assume  $J$  satisfies (5.1), (5.2) and (5.3). Then there exists  $C > 0$  depending on  $s, \gamma, d, m_0, M_0, E_0, H_0$  (note that these constants refer to the mass, energy and entropy of  $h$ ), such that*

$$J(\sigma, x, v; \tau_0, y_0, w_0) \geq C(\sigma - \tau_0)^{-\frac{2d(1+s)}{2s}} \exp \left\{ -C \left( \frac{|x - y_0 - (\sigma - \tau_0)w_0|^{2s}}{(\sigma - \tau_0)^{1+2s}} + \frac{|v - w_0|^{2s}}{\sigma - \tau_0} \right) \right\}.$$

*Remark 5.3.* (i) The existence of a fundamental solution operator for hypoelliptic non-local equations has recently been established by Auscher-Imbert-Niebel [1, 2].

(ii) We can also draw a connection to the Gaussian lower bounds for solutions to the Boltzmann equation by Imbert-Mouhot-Silvestre in [5]. The authors show that for any  $t \in [0, T]$  there exists  $a(t), b(t) > 0$  such that any non-negative solution of the Boltzmann equation satisfies

$$f(t, x, v) \geq a(t)e^{-b(t)|v|^2}.$$

The same authors establish decay estimates for solutions to (1.1) in [4].

**5.2. On the polynomial upper bounds.** The proof method remains the same as [14, Section 6]. The on-diagonal bound [14, Theorem 6.1] follows by Proposition 4.1. The off-diagonal bound [14, Theorem 6.2] follows if we are able to derive Aronson's bound in [14, Proposition 6.4] without using the divergence form symmetry of the kernel of the non-local operator. This is contained in Proposition 5.4 below, which is the weak divergence form analogue of [14, Proposition 6.4]. Once we have established Proposition 5.4 and constructed a decay function  $H$  satisfying Aronson's condition (5.5), the proof of Theorem 5.1 carries over almost verbally from the proof of [14, Theorem 1.2].

5.2.1. *Decay relation.* We aim to define a function that satisfies for some  $\rho > 0$

$$(5.5) \quad \mathcal{T}H(v) + \frac{1}{2} \int_{B_\rho(v)} [H(w) - H(v)] K_h(w, v) dw + \frac{H(v)}{2} \int_{\mathbb{R}^d} [K_h(v, w) - K_h(w, v)] dw \leq 0.$$

Then we can derive the following statement.

**Proposition 5.4** (Aronson's auxiliary proposition). *Let  $0 < \tau_0 < \sigma < T$  and  $0 < \rho$ . Let  $f \in L^\infty((\tau_0, T) \times \mathbb{R}^{2d})$  solve (1.9) in  $(\tau_0, T) \times \mathbb{R}^{2d}$  with a non-local operator whose kernel is non-negative and satisfies (2.3), (2.4), (2.5). Then for every bounded function  $H : [\tau_0, \sigma] \times \mathbb{R}^{2d} \rightarrow [0, \infty)$  such that  $H^{\frac{1}{2}} \in L^2((\tau_0, \sigma) \times \mathbb{R}^d; H_v^s(\mathbb{R}^d))$  and  $D_v H, D_v^2 H \in L^2((\tau_0, \sigma) \times \mathbb{R}^d; L^\infty(\mathbb{R}^d))$ , and, moreover, satisfying (5.5) in  $(\tau_0, \sigma) \times \mathbb{R}^{2d}$ , there exists a constant  $C = C(\lambda, \Lambda, s, d)$  such that*

$$(5.6) \quad \sup_{t \in (\tau_0, \sigma)} \int_{\mathbb{R}^{2d}} f^2(t, x, v) H(t, x, v) dx dv \leq \int_{\mathbb{R}^{2d}} f^2(\tau_0, x, v) H(\tau_0, x, v) dx dv + C\rho^{-2s} \|H\|_{L^\infty([\tau_0, \sigma] \times \mathbb{R}^{2d})} \|f\|_{L^2([\tau_0, \sigma] \times \mathbb{R}^{2d})}^2.$$

*Proof of Proposition 5.4.* For  $R \geq \max\{2, 2\rho + 1\}$  we consider  $\varphi_R \in C_c^\infty(\mathbb{R}^{2d})$  with  $0 \leq \varphi_R \leq 1$  such that  $\varphi_R \equiv 1$  for  $(x, v) \in B_{(R-1)^{1+2s}} \times B_{R-1}$  and  $\varphi_R \equiv 0$  for  $(x, v)$  outside  $B_{R^{1+2s}} \times B_R$ , with bounded derivatives and such that  $|v \cdot \varphi_R| \sim R^{-2s}$ . We test (1.9) with  $fH\varphi_R^2$  where  $H$  satisfies (5.5) over  $[\tau_0, \tau_1] \times \mathbb{R}^{2d}$  for  $0 \leq \tau_0 \leq \tau_1 \leq \sigma$  and get

$$\begin{aligned} \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} \mathcal{T}f(fH\varphi_R^2) dz &= \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} \int_{\mathbb{R}^d} (f(w) - f(v)) K_h(v, w) (fH\varphi_R^2)(v) dw dz \\ &\quad + \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} f(v) (fH\varphi_R^2)(v) \left( \int_{\mathbb{R}^d} (K_h(v, w) - K_h(w, v)) dw \right) dz. \end{aligned}$$

*Step 1: Transport operator.*

First we integrate by parts the transport operator

$$\begin{aligned}
(5.7) \quad & \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} \mathcal{T}f(fH\varphi_R^2) \, dz \\
&= \frac{1}{2} \int_{\mathbb{R}^{2d}} f^2 H\varphi_R^2 \Big|_{t=\tau_0}^{\tau_1} \, dx \, dv - \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} f^2 H\varphi_R v \cdot \nabla_x \varphi_R \, dz - \frac{1}{2} \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} f^2 \varphi_R^2 \mathcal{T}H \, dz \\
&\geq \frac{1}{2} \int_{\mathbb{R}^{2d}} f^2 H\varphi_R^2 \Big|_{t=\tau_0}^{\tau_1} \, dx \, dv - CR^{-2s} \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} f^2 H\varphi_R \, dz - \frac{1}{2} \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} f^2 \varphi_R^2 \mathcal{T}H \, dz.
\end{aligned}$$

*Step 2: Non-local operator.*

Now we deal with the non-local term. We write

$$\begin{aligned}
\int_{\mathbb{R}^d} (\tilde{\mathcal{L}}f) f H\varphi_R^2 \, dv &:= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (f(w) - f(v)) K_h(v, w) (fH\varphi_R^2)(v) \, dw \, dv \\
&\quad + \int_{\mathbb{R}^d} (f^2 H\varphi_R^2)(v) \left( \int_{\mathbb{R}^d} (K_h(v, w) - K_h(w, v)) \, dw \right) \, dv \\
&= \frac{1}{2} \int_{B_{2R}} \int_{B_{2R}} [f(w) - f(v)] [(fH\varphi_R^2)(v) - (fH\varphi_R^2)(w)] K_h(v, w) \, dw \, dv \\
&\quad + \frac{1}{2} \int_{B_{2R}} \int_{B_{2R}} [f(w) - f(v)] [(fH\varphi_R^2)(v) + (fH\varphi_R^2)(w)] K_h(v, w) \, dw \, dv \\
&\quad + \int_{\mathbb{R}^d} (f^2 H\varphi_R^2)(v) \left( \int_{\mathbb{R}^d} (K_h(v, w) - K_h(w, v)) \, dw \right) \, dv \\
&\quad + \int_{B_{2R}} \int_{\mathbb{R}^d \setminus B_{2R}} [f(w) - f(v)] (fH\varphi_R^2)(v) K_h(v, w) \, dw \, dv.
\end{aligned}$$

Then we note that

$$\begin{aligned}
& [f(w) - f(v)] [(fH\varphi_R^2)(v) + (fH\varphi_R^2)(w)] \\
&= [f(w) + f(v)] [(fH\varphi_R^2)(v) - (fH\varphi_R^2)(w)] - 2f(v)(fH\varphi_R^2)(v) + 2f(w)(fH\varphi_R^2)(w),
\end{aligned}$$

and moreover, by Young's inequality

$$\begin{aligned}
& f(w) [(fH\varphi_R^2)(v) - (fH\varphi_R^2)(w)] \\
&\leq \frac{1}{2} f^2(w) [(H\varphi_R^2)(v) - (H\varphi_R^2)(w)] + \frac{1}{2} f^2(v) (H\varphi_R^2)(v) - \frac{1}{2} f^2(w) (H\varphi_R^2)(w).
\end{aligned}$$

Thus

$$\begin{aligned}
\int_{\mathbb{R}^d} (\tilde{\mathcal{L}}f) f H \varphi_R^2 \, dv &\leq \frac{1}{2} \int_{B_{2R}} \int_{B_{2R}} f^2(w) \left[ (H \varphi_R^2)(v) - (H \varphi_R^2)(w) \right] K_h(v, w) \, dw \, dv \\
&\quad + \frac{1}{2} \int_{B_{2R}} \int_{B_{2R}} \left[ (f^2 H \varphi_R^2)(w) - (f^2 H \varphi_R^2)(v) \right] K_h(v, w) \, dw \, dv \\
&\quad + \int_{\mathbb{R}^d} (f^2 H \varphi_R^2)(v) \left( \int_{\mathbb{R}^d} (K_h(v, w) - K_h(w, v)) \, dw \right) \, dv \\
&\quad + \int_{B_{2R}} \int_{\mathbb{R}^d \setminus B_{2R}} [f(w) - f(v)] (f H \varphi_R^2)(v) K_h(v, w) \, dw \, dv \\
(5.8) \quad &\leq \frac{1}{2} \int_{B_{2R}} \int_{B_{2R}} f^2(v) \left[ (H \varphi_R^2)(w) - (H \varphi_R^2)(v) \right] K_h(w, v) \, dw \, dv \\
&\quad + \frac{1}{2} \int_{B_{2R}} \int_{B_{2R}} (f^2 H \varphi_R^2)(v) [K_h(w, v) - K_h(v, w)] \, dw \, dv \\
&\quad + \int_{\mathbb{R}^d} (f^2 H \varphi_R^2)(v) \left( \int_{\mathbb{R}^d} (K_h(v, w) - K_h(w, v)) \, dw \right) \, dv \\
&\quad + \int_{B_{2R}} \int_{\mathbb{R}^d \setminus B_{2R}} [f(w) - f(v)] (f H \varphi_R^2)(v) K_h(v, w) \, dw \, dv \\
&=: \mathcal{I}_{loc}^H + \mathcal{I}_{loc}^{skew} + \mathcal{I}_{nonloc}^{skew} + \mathcal{I}_{tail}.
\end{aligned}$$

First we note that the last integral in (5.8) tends to zero as  $R \rightarrow \infty$ :

$$\begin{aligned}
(5.9) \quad \mathcal{I}_{tail} &= \int_{B_{2R}} \int_{\mathbb{R}^d \setminus B_{2R}} [f(w) - f(v)] (f H \varphi_R^2)(v) K_h(v, w) \, dw \, dv \\
&\leq C \Lambda R^{-2s} \|H \varphi_R^2\|_{L^\infty} \|f\|_{L^1(B_R)} \|f\|_{L^\infty(\mathbb{R}^d \setminus B_{2R})} \xrightarrow{R \rightarrow \infty} 0.
\end{aligned}$$

Second, we distinguish the singular from the non-singular part to bound  $\mathcal{I}_{loc}^H$ . We have

$$\begin{aligned}
(5.10) \quad \mathcal{I}_{loc}^H &= \frac{1}{2} \int_{B_{2R}} \int_{B_{2R}} f^2(v) \left[ (H \varphi_R^2)(w) - (H \varphi_R^2)(v) \right] K_h(w, v) \, dw \, dv \\
&= \frac{1}{2} \int_{B_{2R}} \int_{B_{2R} \cap B_\rho(v)} \dots \, dw \, dv + \frac{1}{2} \int_{B_{2R}} \int_{B_{2R} \setminus B_\rho(v)} \dots \, dw \, dv =: \mathcal{I}_{loc}^{H,s} + \mathcal{I}_{loc}^{H,ns}.
\end{aligned}$$

Then we bound the non-singular part using (2.3)

$$\begin{aligned}
(5.11) \quad \mathcal{I}_{loc}^{H,ns} &= \frac{1}{2} \int_{B_{2R}} \int_{B_{2R} \setminus B_\rho(v)} f^2(v) \left[ (H \varphi_R^2)(w) - (H \varphi_R^2)(v) \right] K_h(w, v) \, dw \, dv \\
&\leq C \rho^{-2s} \|H \varphi_R\|_{L^\infty(B_{2R})} \|f\|_{L^2(B_{2R})}^2.
\end{aligned}$$

For the singular part, we observe that  $\varphi_R(v) = \varphi_R(w)$  for any  $w \in B_\rho(v)$  and  $v \in B_{\frac{R-1}{2}}$  if  $R \geq 1 + 2\rho$ , so that we further split

$$\begin{aligned}
(5.12) \quad \mathcal{I}_{loc}^{H,s} &= \frac{1}{2} \int_{B_{2R}} \int_{B_{2R} \cap B_\rho(v)} f^2(v) \left[ (H \varphi_R^2)(w) - (H \varphi_R^2)(v) \right] K_h(w, v) \, dw \, dv \\
&= \frac{1}{2} \int_{B_{\frac{R-1}{2}}} \int_{B_{2R} \cap B_\rho(v)} f^2(v) \varphi_R^2(v) \left[ H(w) - H(v) \right] K_h(w, v) \, dw \, dv \\
&\quad + \frac{1}{2} \int_{B_{2R} \setminus B_{\frac{R-1}{2}}} \int_{B_{2R} \cap B_\rho(v)} f^2(v) \left[ (H \varphi_R^2)(w) - (H \varphi_R^2)(v) \right] K_h(w, v) \, dw \, dv.
\end{aligned}$$



Then note that as  $R \rightarrow \infty$ , the last integral tends to zero as the integrand is bounded: Taylor expanding  $H\varphi_R$ , using (2.3) and (2.5)

$$\begin{aligned}
& \frac{1}{2} \int_{B_{2R} \setminus B_{\frac{R-1}{2}}} \int_{B_{2R} \cap B_\rho(v)} f^2(v) \left[ (H\varphi_R^2)(w) - (H\varphi_R^2)(v) \right] K_h(w, v) \, dw \, dv \\
&= \frac{1}{2} \int_{B_{2R} \setminus B_{\frac{R-1}{2}}} \int_{B_{2R} \cap B_\rho(v)} f^2(v) D_v(H\varphi_R^2)(v) \cdot (w - v) K_h(w, v) \, dw \, dv \\
&\quad + C \|D_v^2(H\varphi_R^2)\|_{L^\infty(B_{2R} \setminus B_{\frac{R-1}{2}})} \int_{B_{2R} \setminus B_{\frac{R-1}{2}}} \int_{B_{2R} \cap B_\rho(v)} f^2(v) |w - v|^2 K_h(w, v) \, dw \, dv \\
&\leq C \rho^{1-2s} \int_{B_{2R} \setminus B_{\frac{R-1}{2}}} f^2(v) D_v(H\varphi_R^2)(v) \, dv + C \rho^{2-2s} \|D_v^2(H\varphi_R^2)\|_{L^\infty(B_{2R} \setminus B_{\frac{R-1}{2}})} \int_{B_{2R} \setminus B_{\frac{R-1}{2}}} f^2(v) \, dv \\
&\xrightarrow{R \rightarrow \infty} 0.
\end{aligned}$$

We combine (5.10), (5.11), (5.12) and let  $R \rightarrow \infty$ , so that

$$(5.13) \quad \mathcal{I}_{loc}^H \leq C \rho^{-2s} \|H\|_{L^\infty(\mathbb{R}^d)} \|f\|_{L^2(\mathbb{R}^d)} + \frac{1}{2} \int_{\mathbb{R}^d} \int_{B_\rho(v)} f^2(v) \left[ H(w) - H(v) \right] K_h(w, v) \, dw \, dv.$$

Thus, by letting  $R \rightarrow \infty$ , we conclude from (5.8), (5.9), (5.13),

$$(5.14) \quad \begin{aligned} \int_{\mathbb{R}^d} (\tilde{\mathcal{L}}f) f H \, dv &\leq \frac{1}{2} \int_{\mathbb{R}^d} f^2(v) \int_{B_\rho(v)} \left[ H(w) - H(v) \right] K_h(w, v) \, dw \, dv \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^d} (f^2 H)(v) \int_{\mathbb{R}^d} \left[ K_h(v, w) - K_h(w, v) \right] \, dw \, dv + C \rho^{-2s} \|H\|_{L^\infty(\mathbb{R}^d)} \|f\|_{L^2(\mathbb{R}^d)}^2. \end{aligned}$$

*Step 3: Conclusion.*

We assemble the pieces. Equation (5.7) implies

$$\begin{aligned}
& \frac{1}{2} \int_{\mathbb{R}^{2d}} f^2(\tau_1, x, v) H(\tau_1, x, v) \varphi_R^2 \, dx \, dv \\
&\leq \frac{1}{2} \int_{\mathbb{R}^{2d}} f^2(\tau_0, x, v) H(\tau_0, x, v) \varphi_R^2 \, dx \, dv + C R^{-2s} \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} f^2(z) H(z) \varphi_R(x, v) \, dz \\
&\quad + \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} f^2(z) \varphi_R^2(x, v) \mathcal{T}H(z) \, dz + \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} (\tilde{\mathcal{L}}f)(z) f(z) H(z) \varphi_R^2(x, v) \, dz,
\end{aligned}$$

which as  $R \rightarrow \infty$  yields together with (5.14)

$$\begin{aligned}
& \frac{1}{2} \int_{\mathbb{R}^{2d}} f^2(\tau_1, x, v) H(\tau_1, x, v) \, dx \, dv \\
&\leq \frac{1}{2} \int_{\mathbb{R}^{2d}} f^2(\tau_0, x, v) H(\tau_0, x, v) \, dx \, dv + \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} f^2(z) \mathcal{T}H(z) \, dz \\
&\quad + \frac{1}{2} \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} f^2(z) \int_{B_\rho(v)} \left[ H(w) - H(v) \right] K_h(w, v) \, dw \, dz \\
&\quad + \frac{1}{2} \int_{[\tau_0, \tau_1] \times \mathbb{R}^{2d}} f^2(z) H(z) \int_{\mathbb{R}^d} \left[ K_h(v, w) - K_h(w, v) \right] \, dw \, dv + C \rho^{-2s} \|H\|_{L^\infty} \|f\|_{L^2([\tau_0, \tau_1] \times \mathbb{R}^{2d})}^2 \\
&\leq \frac{1}{2} \int_{\mathbb{R}^{2d}} f^2(\tau_0, x, v) H(\tau_0, x, v) \, dx \, dv + C \rho^{-2s} \|H\|_{L^\infty} \|f\|_{L^2([\tau_0, \tau_1] \times \mathbb{R}^{2d})}^2,
\end{aligned}$$

since by construction  $H$  satisfies (5.5). This concludes the proof of Proposition 5.4.  $\square$

It now only remains to show that we can construct a function  $H$  satisfying (5.5).

**Lemma 5.5.** *Let  $y_0, w_0 \in \mathbb{R}^d$ . Let  $0 < \rho$  and  $0 \leq \tau_0 < \sigma$ . Let  $k \geq 1$  and  $\alpha \geq 0$  be such that  $\sigma - \tau_0 \leq \frac{\rho^{2s}}{4k}$ . For  $(t, x, v) \in [\tau_0, \sigma] \times \mathbb{R}^{2d}$ , we define  $\delta(t) := 2(\sigma - \tau_0) - (t - \tau_0)$  and*

$$(5.15) \quad H(t, x, v) := e^{-\max\left\{1, \frac{1}{3\rho} \max\left(|v-w_0|, |x-y_0-(\sigma+t-2\tau_0)w_0|^{\frac{1}{1+2s}}\right)\right\} \log\left(\frac{\rho^{2s}}{k\delta(t)}\right)} e^{\alpha \frac{(\sigma-t)}{\rho^{2s}}}.$$

*Then there exist constants  $C_1, C_2 > 0$  depending only on  $s, d, \Lambda_0$  such that, if  $k > C_1$  and  $\alpha > C_2$ , then  $H$  satisfies (5.5), where  $K_h$  is a non-negative kernel satisfying (2.3), (2.4), (2.5).*

To check that  $H$  satisfies (5.5), we note that due to (2.4), we choose  $\alpha$  such that

$$(5.16) \quad -\frac{\alpha}{\rho^{2s}} H(v) + \frac{1}{2} H(v) \int_{\mathbb{R}^d} [K_h(v, w) - K_h(w, v)] dw \leq -\alpha H + \Lambda_0 H \leq 0.$$

Then, for ease of notation, we write

$$H(t, x, v) = H_0(t, x, v) e^{\alpha \frac{(\sigma-t)}{\rho^{2s}}},$$

so that if  $H_0$  satisfies

$$(5.17) \quad \mathcal{T}H_0 + \frac{1}{2} \int_{B_\rho(v)} [H_0(w) - H_0(v)] K_h(w, v) dw \leq 0,$$

then in particular  $H$  satisfies (5.5) due to (5.16). One can verify (5.17) by a case distinction, similar to [14, Lemma 6.5].

Note that the remainder of the proof of Theorem 6.2 and Theorem 6.3 in [14] carries over almost verbally, upon replacing Proposition 6.4 and Lemma 6.5 of [14] by Proposition 5.4, and Lemma 5.5. In particular, the Theorem 5.1 follows similarly to [14, Theorem 1.2].

**5.3. On the exponential lower bound.** The only part where we use the divergence form symmetry of the non-local operator in the derivation of the exponential lower bound on the fundamental solution is in [14, Section 7.3]. Instead, we argue as follows.

Fix  $(\tau_0, y_0, w_0) \in [0, T] \times \mathbb{R}^{2d}$  and let  $0 < \tau_0 < \sigma < T$ . Consider for  $\tau_1 < \sigma$  and for some  $\alpha$  large enough the function

$$f(\tau_1, y_1, w_1) = M \int_{\max\left\{|x-y_0-(\sigma-\tau_0)w_0|^{\frac{2s}{1+2s}}, |v-w_0|^{2s}\right\} < \alpha(\sigma-\tau_0)} J(\sigma, x, v; \tau_1, y_1, w_1) dx dv,$$

where  $M \geq 1$  is some constant such that  $f(\tau_1, y_1, w_1) \leq M$  for  $(\tau_1, y_1, w_1) \in [0, \sigma] \times \mathbb{R}^{2d}$ . Define  $g(\tau_1, y_1, w_1) := f(\tau_1, y_1, w_1) e^{-c_0(\tau_1-\sigma)}$  for  $\tau_1 < \sigma$ . Then  $g$  is a super-solution to the adjoint of (1.9) in  $(0, \sigma) \times \mathbb{R}^{2d}$ , with initial values

$$\begin{cases} g(\sigma, y_1, w_1) = M, & \text{if } \max\left\{|y_1 - y_0 - (\sigma - \tau_0)w_0|^{\frac{2s}{1+2s}}, |w_1 - w_0|^{2s}\right\} < \alpha(\sigma - \tau_0), \\ g(\sigma, y_1, w_1) = 0, & \text{if } \max\left\{|y_1 - y_0 - (\sigma - \tau_0)w_0|^{\frac{2s}{1+2s}}, |w_1 - w_0|^{2s}\right\} > \alpha(\sigma - \tau_0). \end{cases}$$

since

$$\mathcal{L}^* g(v) = \int_{\mathbb{R}^d} (g(w) - g(v)) K_h(v, w) dw + \left( \int_{\mathbb{R}^d} K_h(w, v) - K_h(v, w) dw \right) g(v),$$

so that using (2.4) and choosing  $c_0 > 0$  large enough, we get

$$\mathcal{T}^* g(v) - \mathcal{L}^* g(v) = -e^{-c_0(\tau_1-\sigma)} \mathcal{T}f - e^{-c_0(\tau_1-\sigma)} \mathcal{L}f - \Lambda_0 g + c_0 g \geq 0.$$

If we set

$$\tilde{g}(\tau_1, y_1, w_1) = \begin{cases} g(\tau_1, y_1, w_1), & \text{if } \tau_1 < \sigma, \\ M e^{-c_0(\tau_1-\sigma)}, & \text{if } \tau_1 > \sigma, \end{cases}$$

then  $\tilde{g}$  is a non-negative super-solution of the adjoint of (1.9) in  $(0, \infty) \times \mathbb{R}^{2d}$  since  $\tilde{g} \leq Me^{-c_0(\tau_1 - \sigma)}$  for  $(\tau_1, y_1, w_1) \in [0, \infty) \times \mathbb{R}^{2d}$ , so that for  $\tau_1 > \sigma$

$$\mathcal{T}^* \tilde{g} - \mathcal{L}^* \tilde{g} = c_0 Me^{-c_0(\tau_1 - \sigma)} - \mathcal{L} \tilde{g} - \Lambda_0 \tilde{g} = Me^{-c_0(\tau_1 - \sigma)} (c_0 - \Lambda_0) + \int \left[ Me^{-c_0(\tau_1 - \sigma)} - \tilde{g}(w) \right] K_h(v, w) dw \geq 0.$$

Thus by the Weak Harnack inequality [6, Theorem 1.6], we get using  $-c_0(\tau_1 - \sigma) \geq 0$  for  $c_0 \geq 0$  and the fact that  $\tilde{f}(\sigma, y, w) = M$  for some  $\delta > 0$  sufficiently small

$$\begin{aligned} \tilde{g}(\tau_0, y_0, w_0) &\geq \left( \int_{\sigma - \delta}^{\sigma + \delta} \int_{\max\{|y' - y_0 - (\tau' - \tau_0)w_0|^{\frac{2s}{1+2s}}, |w' - w_0|^{2s}\} < \alpha(\sigma - \tau_0)} \tilde{g}^\zeta(\tau', y', w') dy' dw' d\tau' \right)^{\frac{1}{\zeta}} \\ &\geq \left( \int_{\sigma - \delta}^{\sigma + \delta} \int_{\max\{|y' - y_0 - (\tau' - \tau_0)w_0|^{\frac{2s}{1+2s}}, |w' - w_0|^{2s}\} < \alpha(\sigma - \tau_0)} \tilde{f}^\zeta(\tau', y', w') dy' dw' d\tau' \right)^{\frac{1}{\zeta}} \\ &\geq c > 0, \end{aligned}$$

so that

$$f(\tau_0, y_0, w_0) \geq ce^{c_0(\tau_0 - \tau_1)}.$$

We conclude the proof of Theorem 5.2 with [14, Theorem 7.2].

#### REFERENCES

- [1] AUSCHER, P., IMBERT, C., AND NIEBEL, L. Fundamental solutions to Kolmogorov-Fokker-Planck equations with rough coefficients: Existence, uniqueness, upper estimates. arXiv:2403.17468, 2024.
- [2] AUSCHER, P., IMBERT, C., AND NIEBEL, L. Weak solutions to Kolmogorov-Fokker-Planck equations: Regularity, existence and uniqueness. arXiv:2403.17464, 2024.
- [3] CHAKER, J., AND SILVESTRE, L. Coercivity estimates for integro-differential operators. *Calc. Var.* 59, 106 (2020).
- [4] IMBERT, C., MOUHOT, C., AND SILVESTRE, L. Décroissance aux grandes vitesses pour les solutions de l'équation de Boltzmann sans troncature angulaire. *J. Éc. Polytech., Math.* 7 (2020), 143–184.
- [5] IMBERT, C., MOUHOT, C., AND SILVESTRE, L. Gaussian Lower Bounds for the Boltzmann Equation without Cutoff. *SIAM J. Math. Anal.* 52, 3 (2020), 2930–2944.
- [6] IMBERT, C., AND SILVESTRE, L. The weak Harnack inequality for the Boltzmann equation without cut-off. *J. Eur. Math. Soc.* 22, 2 (2020), 507–592.
- [7] IMBERT, C., AND SILVESTRE, L. Global regularity estimates for the Boltzmann equation without cut-off. *J. Amer. Math. Soc.* (2021).
- [8] IMBERT, C., AND SILVESTRE, L. The Schauder estimate for kinetic integral equations. *Anal. PDE* 14, 1 (2021), 171–204.
- [9] KASSMANN, M., AND WEIDNER, M. Nonlocal operators related to nonsymmetric forms I: Hölder estimates. arXiv:2203.07418, 2022.
- [10] KASSMANN, M., AND WEIDNER, M. The Harnack inequality fails for nonlocal kinetic equations. arXiv:2405.05223, 2024.
- [11] LOHER, A. Quantitative Schauder estimates for hypoelliptic equations. arXiv:2305.00463, 2023.
- [12] LOHER, A. The Strong Harnack inequality for the Boltzmann equation. *Séminaire Laurent Schwartz - EDP et applications*, 1 (2023-2024), 1–15.
- [13] LOHER, A. Quantitative De Giorgi methods in kinetic theory for non-local operators. *J. Funct. Anal.* 286, 6 (2024).
- [14] LOHER, A. Semi-local behaviour of non-local hypoelliptic equations: divergence form. arXiv:2404.05612, 2024.
- [15] SILVESTRE, L. A new regularization mechanism for the Boltzmann equation without cut-off. *Comm. Math. Phys.* 348, 1 (2016), 69–100.
- [16] ZHEN-QING CHEN, T. K. Heat kernel estimates for stable-like processes on  $d$ -sets. *Stochastic Process. Appl.* 108, 1 (2003), 27–62.

(Amélie Loher) DPMMS, UNIVERSITY OF CAMBRIDGE, WILBERFORCE ROAD, CAMBRIDGE CB3 0WA, UK

Email address: aj1221@cam.ac.uk