

Partitioning 2-edge-coloured bipartite graphs into monochromatic cycles*

Fabrício Siqueira Benevides[†] Arthur Lima Quintino[†] Alexandre Talon[‡]

Abstract

Given an r -colouring of the edges of a graph G , we say that it can be partitioned into p monochromatic cycles when there exists a set of p vertex-disjoint monochromatic cycles covering all the vertices of G . In the literature of this problem, an edge and a single vertex both count as a cycle.

We show that for every 2-colouring of the edges of a complete balanced bipartite graph, $K_{n,n}$, it can be partitioned into at most 4 monochromatic cycles. This type of question was first studied in 1970 for complete graphs and in 1983, by Gyárfás and Lehel, for $K_{n,n}$. In 2014, Pokrovskiy, has showed that any 2-colouring of the edges of $K_{n,n}$ can be partitioned into at most 3 paths. It turns out that finding monochromatic cycles instead of paths is a natural question that has also been asked for other graphs. In 2015, Schaudt and Stein have showed that at most 14 cycles are necessary.

1 Introduction

An r -colouring of a graph G is a colouring of its edges with up to r different colours. For such an r -colouring we say that *the coloured graph is partitioned into p monochromatic cycles* when there exist a set of p vertex-disjoint monochromatic cycles covering all the vertices of G . We are interested in the *cycle partitioning number of G* , that is the smallest number, $\text{cp}_r(G)$, such that, for every r -colouring of G , it can be partitioned into at most $\text{cp}_r(G)$ monochromatic cycles. In the literature of this problem, an edge and a single vertex both count as a cycle.

In the 1970's, Lehel made an influential conjecture stating that when the edges of a complete graph K_n are coloured with two colours, it can be partitioned into at most two monochromatic cycles. Łuczak, Rödl and Szemerédi [7] confirmed this conjecture for sufficiently large complete graphs, after preliminary work of Gyárfás [3]. In 2010, Bessy and Thomassé [1] proved it for every complete graph, with a surprisingly short proof. For fixed r , the best upper bound, to our knowledge, is $\text{cp}_r(K_n) \leq 100r \log r$, for n large enough (see [6]).

In the 2-colour case, if we replace cycles by paths, the problem becomes much easier. As noted by Gerencsér and Gyárfás [2] in 1967: *the vertex set of any 2-edge-coloured complete graph can be partitioned into two paths of different colours*. To prove it, take any red-blue colouring of K_n and suppose that R and B are vertex disjoint red and blue paths, with respective endpoints r and b . Let $v \notin V(R) \cup V(B)$. If vr is red or vb is blue then we can extend R or B accordingly. Otherwise, without loss of generality rb is red and $\{R \cup rb \cup bv, B - \{b\}\}$ is a pair of red-blue paths covering one extra vertex.

The general problem has also been studied for the case where the base graph G is a fair complete k -partite graph and for graphs with large minimum degree in general. We point to [4] for a (2016) survey and call attention to a couple of (older and newer) results.

Let us now define the basic concepts that we will use throughout this paper.

Definition 1. We say that a path is **simple** when it is the union of a blue path, $v_1v_2 \dots v_i$ and a red path, $v_iv_{i+1} \dots v_k$. We call v_i the **turning point** of the path.

Definition 2. We say that a bipartite graph is **balanced** when its partition classes have the same number of vertices.

*This work was funded by the Fog Research Institute under contract no. FRI-454, by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) – Brasil (Finance Code 001).

[†]Departamento de Matemática, Universidade Federal do Ceará, Brazil (fabricio@mat.ufc.br)

[‡](alexandre.talon@ens-lyon.fr)

Definition 3. Let G be a bipartite graph with partition classes X and Y , whose edges are coloured red and blue. The colouring on G is **split** when X and Y can each be partitioned into two non-empty sets, $X = X_1 \cup X_2$ and $Y = Y_1 \cup Y_2$, such that all edges between X_i and Y_j are red for $i = j$ and blue for $i \neq j$. (Figure 1.1).




Figure 1.1: A split colouring.

In 1983, Gyárfás and Lehel [3, 5] showed that every balanced complete bipartite graph with a two-colouring that is not split contains two vertex-disjoint monochromatic paths that cover all but at most one of its vertices. This result was later extended by Pokrovskiy.

Theorem 1 (Pokrovskiy [8], 2014). *Let G be a balanced complete bipartite graph whose edges are coloured red and blue. There is a vertex-partition of G into two monochromatic **paths with different colours** if, and only if, the colouring on G is not split.*

Corollary 2. *Let G be a complete bipartite graph G with a split colouring of its edges into red and blue. The vertices of G cannot be partitioned into two monochromatic cycles of different colours.*

Proof. By Theorem 1 (or by a very simple argument), the graph G cannot be partitioned into two paths of different colours. Therefore, it cannot be partitioned into two cycles of different colours either. \square

However, for split colourings the following is (trivially) true (but we include a proof for completeness).

Proposition 3. *Let G be a balanced complete bipartite graph whose edges are coloured red and blue. If the colouring on G is split, then G can be vertex-partitioned into at most 3 monochromatic cycles. Furthermore, two cycles suffice if and only if $|X_1| = |Y_1|$ or $|X_1| = |Y_2|$.*

Proof. Let X and Y be the partition classes of G . And $X = X_1 \cup X_2$, $Y = Y_1 \cup Y_2$ as in the definition of split colouring (Definition 3). We may assume without loss of generality that $|X_1| \geq |Y_1|$. As $|X| = |Y|$, we have $|X_1| - |Y_1| = |Y_2| - |X_2|$ and, therefore, $|X_2| \leq |Y_2|$. Hence, G can be vertex-partitioned into a red cycle that covers all vertices in Y_1 and $|Y_1|$ vertices in X_1 , a red cycle that covers all vertices in X_2 and $|X_2|$ vertices in Y_2 and a blue cycle that covers all the remaining vertices in X_1 and Y_2 . And clearly, if $|X_1| = |Y_1|$, then $|X_2| = |Y_2|$ and two cycles (of same colour) suffice.

On the contrary, suppose that two monochromatic cycles suffice. By Corollary 2 they must be of the same colour. Suppose, without loss of generality that both are red. Each cycle must be entirely contained in $X_1 \cup Y_1$ or $X_2 \cup Y_2$. And because all the sets X_1 , X_2 , Y_1 and Y_2 are non-empty, one cycle must cover vertices in $X_1 \cup Y_1$ and the other in $X_2 \cup Y_2$. And each cycle covers as many vertices in X as in Y . Therefore $|X_1| = |Y_1|$ and $|X_2| = |Y_2|$. Note: similarly blue cycles suffice if and only if $|X_1| = |Y_2|$ and $|X_2| = |Y_1|$. \square

A natural conjecture would be that every 2-colouring of a balanced bipartite graph that is not split can be partitioned into at most 2 monochromatic cycles. In Proposition 7 we give an example of a colouring that is not split and needs at least 3 cycles.

For upper bounds, using the regularity method, Schaudt and Stein [9] showed that for every k , a 2-edge-coloured sufficiently large k -partite graph G , such that no partition class of G contains more than half of its vertices, can be partitioned into at most 14 monochromatic cycles. For $k = 2$, they proved that this can be done with at most 12 monochromatic cycles.

Our main contribution is to reduce this number to only 4 monochromatic cycles, for any balanced bipartite graph. Also, note that our result does not require G to be large.

Theorem 4. *If G is a balanced complete bipartite graph whose edges are coloured red and blue, then G can be partitioned into at most 4 monochromatic cycles.*

2 Proving Theorem 4

Thanks to Proposition 3, we only have to prove Theorem 4 when the given colouring is not split. Our proof is self-contained apart from the following result.

Corollary 5 (Stein [10], 2022). *Let G be a balanced complete bipartite graph whose edges are coloured red and blue. If the colouring on G is not split, then G can be vertex-partitioned into a monochromatic path and a monochromatic cycle with different colours.*

We observe that the above is a corollary to Theorem 1, and follows from a very short case analysis on the parity of the paths given by Theorem 1. This is done in details in Section 4.1 of [10]. We also note that Section 4.2 of [10] presents an alternative proof of Theorem 1 that is also short and self-contained.

In view of Corollary 5, it is enough to prove the following lemma.

Lemma 6. *Let G be a balanced complete bipartite graph whose edges are coloured red and blue. If G has a monochromatic Hamiltonian path, then G can be vertex-partitioned into at most 3 monochromatic cycles.*

In fact, let H be any balanced complete bipartite graph with a two-colouring on its edges. When we apply Corollary 5 to H , the monochromatic cycle we obtain must be even because H is bipartite. And since H is balanced, the monochromatic path given by Corollary 5 is also even. So its vertices induce a balanced complete bipartite subgraph of H that contains a monochromatic Hamiltonian path.

Lemma 6 is best possible, in the sense that even with the extra condition that the colouring of the bipartite graph contains a Hamiltonian monochromatic path, there are 2-colourings that cannot be partitioned into 2 monochromatic cycles, as illustrated below.

Proposition 7. *Let G be a balanced complete bipartite graph on $2n$ vertices, with bipartition $S = \{x_1, \dots, x_n\} \cup \{y_1, \dots, y_n\}$. Take the colouring where:*

- x_1y_1, x_2y_2 and all edges of the form x_1y_i and y_1x_i for $i \geq 3$ are blue;
- the edges x_2y_i for $i \geq 4$ are blue;
- all the other edges are red.

Then G contains a Hamiltonian red path and cannot be partitioned into two monochromatic cycles.

Proof. Since the graph induced by $S_{\geq 3} = \{x_3, \dots, x_n\} \cup \{y_3, \dots, y_n\}$ has all edges coloured red, there is a path P starting in x_3 and ending in y_3 and passing through all vertices of $S_{\geq 3}$. The red path $x_1y_2x_3Py_3x_2y_1$ is a red Hamiltonian path of G .

Now we show that G cannot be partitioned into 2 monochromatic cycles. First, the vertex y_1 cannot belong to a cycle with more than two vertices: such a cycle would be blue because only one red edge is incident to y_1 , and it would then contain one x_i with $i \geq 2$ but such an x_i only has one incident blue edge, towards y_1 , hence it could not continue further. Now, if we make a cycle with y_1 only, given the fact that the graph is bipartite, we cannot make a single cycle with all the $2n - 1$ remaining vertices. Let us now assume we have a partition into two monochromatic cycles. One cycle must contain y_1 and exactly one of the x_i 's. The other cycle must contain y_2 , therefore it cannot be blue since only one blue edge is incident to that vertex. So it needs to be a red cycle, and then cannot contain x_1 (only one incident red edge), hence the first cycle is x_1y_1 . Then this second cycle, which is red, must contain both x_2 and y_2 but x_2 has only one incident red edge in $G \setminus \{x_1, y_1\}$ so this is a contradiction. \square

Next we present the structure of the proof of Theorem 4. In all that follows, we consider a balanced complete bipartite graph G whose edges are coloured red and blue **and contains a red Hamiltonian path**. We also consider that this colouring is not split, and we want to show that there is partition of G into at most 3 monochromatic cycles.

Our idea is to search for more structure in the above graph G . Let's say that each partition class of G has n vertices. We will label its vertices in a particular way. Then, in order to prove Lemma 6, we use induction with on a parameter k , $1 \leq k \leq n$, with a well chosen hypothesis. We will try to show the existence of a specific subgraph which we will define later as a *blue even plait*: the first $k/2$ vertices with even indices from both classes (of the bipartition) inducing a complete blue bipartite graph, and the same with the first $k/2$ vertices with odd indices. This exists for $k = 1$, and we will show that if G cannot be partitioned into three monochromatic cycles and the induction hypothesis holds with k , then

it also holds with $k + 1$. Iterating this will give us our result, because a blue even plait shall be trivially partitioned into 2 blue cycles. The proof will mostly consists in looking at specific edges and proving that they are blue (though we also need to argue that a few edges are red), with the goal to increase the order of the blue even plait.

We split this section into two subsections. Subsection 2.1 contains some preliminary results that we will use many times and help us to organise the other subsection in a concise way. Subsection 2.2 is devoted to the main proof of Lemma 6, which is longer. The preliminary results include a lemma with a short inductive proof which suffices to deduce a weakening of Lemma 6 with 4 cycles, which in turn (by the above arguments) implies a weakening of Theorem 4 with 5 cycles.

2.1 Preliminary structural results

Throughout the rest of this paper we deal with balanced complete bipartite graphs whose edges are coloured red and blue that have a monochromatic Hamiltonian path. Therefore, in order to avoid exhaustive repetitions, it will be convenient to establish the following convention for the order of the vertices in such a path.

Definition 4. A balanced complete bipartite graph with $2n$ vertices whose edges are coloured red and blue is a **red zigzag graph** when its partition classes are $X = \{x_1, \dots, x_n\}$ and $Y = \{y_1, \dots, y_n\}$ and the Hamiltonian path

$$P = (x_1, y_2, x_3, y_1, \dots, x_4, y_3, x_2, y_1)$$

is red. (See Figure 2.1.)

For every $k \leq n$, we also define the sets:

$$S_{\leq k} = \{x_i : i \leq k\} \cup \{y_i : i \leq k\},$$

$$S_{< k} = \{x_i : i < k\} \cup \{y_i : i < k\},$$

$$S_{\geq k} = \{x_i : i \geq k\} \cup \{y_i : i \geq k\},$$

$$S_{> k} = \{x_i : i > k\} \cup \{y_i : i > k\}.$$

We denote by $x_i P y_j$ the red Hamiltonian subpath of P starting from vertex x_i and ending at y_j , both end-vertices being included.




Figure 2.1: Labelling of the red Hamiltonian path $P = (x_1, y_2, x_3, y_1, \dots, x_4, y_3, x_2, y_1)$ of a red zigzag graph with $2n$ vertices. And the set $S_{\leq 2}$.

In addition to these and to other definitions that will appear later, we shall also establish some simple facts that will help us in the proofs of the main results of this section. Such facts will be just stated as remarks, since they are straightforward or can be easily checked.

Remark 8. Let G be a red zigzag graph with $2n$ vertices. Then, for $i \leq n$:

- (a) if the edge $x_i y_i$ is red, then the subgraph $G[S_{\geq i}]$ has a red Hamiltonian cycle, namely $(x_i P y_i)$. (See Figure 2.2a.)
- (b) if the edge $x_i y_{i+2}$ (resp. $x_{i+2} y_i$) is red, then the subgraph $G[S_{\geq i}]$ can be vertex-partitioned into two red cycles, namely into the red cycle $(x_i, y_{i+2} P x_{i+2}, y_{i+1})$ (resp. $(x_{i+1}, y_{i+2} P x_{i+2}, y_i)$) and the edge-cycle (x_{i+1}, y_i) (resp. (x_i, y_{i+1})). (See Figure 2.2b.)




Figure 2.2: The subgraph $G[S_{\geq i}]$ according to cases (a) and (b) of Remark 8.

In the above language, proving Lemma 6 is the same as proving that *every red zigzag graph can be vertex-partitioned into at most 3 monochromatic cycles*. Therefore, we see that the lemma below is a weaker version of Lemma 6, which allows one extra monochromatic cycle to be used. As compensation, this version gives us some additional information about the monochromatic cycles of the obtained partition.

Lemma 9. *If G is a red zigzag graph with $2n$ vertices, then G can be vertex-partitioned into at most t monochromatic cycles satisfying one of the following conditions.*

(i) $t = 2$.

(ii) $t = 3$ and the edge x_1y_1 is used in some blue cycle of the partition.

(iii) $t = 4$ and the edges x_1y_1 and x_2y_2 are used in different blue cycles of the partition.

Proof. The proof is by induction on n . For $n \leq 2$, the graph G satisfies condition (i) trivially. For $n \geq 3$, we may assume that G does not satisfy condition (i), since otherwise we would be done. Hence, by Remark 8, we know that the edges x_1y_1 , x_1y_3 and x_3y_1 are blue. Now, we apply the inductive hypothesis to the subgraph $G[S_{>1}]$. If $G[S_{>1}]$ satisfies conditions (i) or (ii), then we may simply take the edge x_1y_1 as a blue cycle and so we see that G satisfies conditions (ii) or (iii), respectively. On the other hand, if $G[S_{>1}]$ satisfies condition (iii), then we may use the edges x_1y_1 , x_1y_3 and x_3y_1 to extend the blue cycle of the partition of $G[S_{>1}]$ that uses the edge x_3y_3 and so we see that G satisfies condition (iii) as well. In any case, the result follows. \square

In view of the proof of Theorem 4, observe that the lemma above has a certain relevance by itself, since it implies that 5 monochromatic cycles are sufficient to vertex-partition any balanced complete bipartite graph whose edges are coloured red and blue, which already improves results from [9] (where it is proved that large balanced bipartite graphs can be partitioned into 12 cycles), with a significantly shorter proof.

However, Lemma 9 is just the first step towards Theorem 4. The next step is to introduce some tools to prove Lemma 6. We will also prove a weaker version of Lemma 6, which will make the proof of Lemma 6 clearer.

Definition 5. *Let G be a red zigzag graph with $2n$ vertices. For $2 \leq k \leq n$, the set $S_{\leq k}$ is a blue special set when the following two conditions hold:*

(i) *the subgraph $G[S_{\leq k-2}]$ is either empty or has a blue Hamiltonian cycle,*

(ii) *the subgraph $G[S_{\leq k-1}]$ has blue Hamiltonian cycle, and*

(iii) *the subgraph $G[S_{\leq k}]$ has a blue Hamiltonian cycle that uses the edges $x_{k-1}y_{k-1}$ and x_ky_k .*

Furthermore, we need the following remark.

Remark 10. *Let G be a complete bipartite graph and $C_1 = (u_1, \dots, u_s)$ and $C_2 = (v_1, \dots, v_t)$ be two disjoint blue cycles in G , where $u_1, v_1 \in X$. Then u_1v_t and v_1u_s are edges of G . If those edges are both blue, then there is a blue cycle in G that passes through all edges in C_1 and C_2 except u_1u_s and v_1v_t and covers all vertices in C_1 and C_2 , namely $(u_1, \dots, u_s, v_1, \dots, v_t)$. (See Figure 2.3.)*




Figure 2.3: The blue cycle $(u_1, \dots, u_s, v_1, \dots, v_t)$ represented in two ways. The one on the left shows the vertices in their appropriate class of the bi-partition.

The following lemma shows us that the existence of a blue special set in a red zigzag graph is a sufficient extra condition to make Lemma 6 valid.

Lemma 11. *Let G be a red zigzag graph with $2n$ vertices. If G has a **blue special set**, then G can be vertex-partitioned into at most 3 monochromatic cycles.*

Proof. Let $S_{\leq k}$ be a blue special set in G for some $2 \leq k \leq n$. By Definition 5 (iii), $G[S_{\leq k}]$ has a blue Hamiltonian cycle. Hence, we may assume that $n > k + 2$, since otherwise we would be done. By Remark 8(b) and Definition 5 (iii), we may also assume that the edges $x_{k-1}y_{k+1}$, $x_{k+1}y_{k-1}$, $x_k y_{k+2}$ and $x_{k+2}y_k$ are blue, since otherwise we would be done.

Now, we apply Lemma 9 to the subgraph $G[S_{>k}]$. If $G[S_{>k}]$ satisfies condition (i), then we may simply take a blue Hamiltonian cycle in $G[S_{\leq k}]$ to obtain a good partition of G . By Remark 10, if $G[S_{>k}]$ satisfies condition (ii), then we use the edges $x_{k-1}y_{k+1}$ and $x_{k+1}y_{k-1}$ to combine the blue cycle that uses the edge $x_{k+1}y_{k+1}$ from the 3-cycle partition of $G[S_{>k}]$ with the blue Hamiltonian cycle of $G[S_{\leq k}]$ (since it uses the edge $x_{k-1}y_{k-1}$), thereby obtaining a good partition of G . Observe that the previously built blue cycle also passes through the edge $x_k y_k$. Hence, if $G[S_{>k}]$ satisfies condition (iii), then first we do the same operation as in the previous case (using edges $x_{k-1}y_{k+1}$ and $x_{k+1}y_{k-1}$) and after that we use the edges $x_k y_{k+2}$ and $x_{k+2}y_k$ to build a blue cycle in G that covers all vertices in $G[S_{\leq k}]$ and all vertices of two different blue cycles of the partition of $G[S_{>k}]$ (remember that the edges $x_{k+1}y_{k+1}$ and $x_{k+2}y_{k+2}$ were in different cycles of the partition). Thereby obtaining a desired partition of G . In every case, the result follows. \square

To prove the next lemma, it will be necessary to make a more involved analysis of the edges compared to what we did in the proofs of Lemmas 9 and 11. Therefore, as a way to facilitate our work and the reader's understanding, we shall make some considerations first.

Definition 6. *An edge $x_i y_j$ in a red zigzag graph is **even** when i and j have the same parity, i.e., when $i + j$ is even.*

Usually, if an even edge with “low” values for its indices is red, we can then partition the edges into 3 monochromatic cycles. Otherwise if that edge is blue we can look at other edges. The fact that such an edge is blue helps us little by little to establish a nice blue structure.

Definition 7. *Let G be a red zigzag graph with $2n$ vertices. For $1 \leq k \leq n$, the subgraph $G[S_{\leq k}]$ is a blue even plait when all its even edges are blue. We denote by G_k^O the subgraph of G formed by vertices of odd indices in $S_{\leq k}$, and G_k^E the one formed by vertices of even indices.*

Remark 12. *Let G be a red zigzag graph with $2n$ vertices. If $G[S_{\leq k}]$ is a blue even plait, for $1 \leq k \leq n$, then G_k^O and G_k^E are complete balanced bipartite graphs whose edges are all blue. In particular, each of them is Hamiltonian, so $G[S_{\leq k}]$ can be partitioned into at most two monochromatic blue cycles.*

Notation 8. *A Hamiltonian path in G_k^E starting in x_i and ending in y_j is denoted $x_i \overset{E_k}{\rightsquigarrow} y_j$ (note that such a path needs to start and finish on different classes of the bi-partition).*

A path in G_k^E starting in x_i and ending in x_j that passes through all vertices of G_k^E except for a given y_ℓ is denoted $x_i \overset{E_k \setminus y_\ell}{\rightsquigarrow} x_j$ (note that such a path needs to start and finish on the same class of the bi-partition).

We use the same notations $(x_i \overset{O_k}{\rightsquigarrow} y_j$ and $x_i \overset{O_k \setminus y_\ell}{\rightsquigarrow} x_j)$ for G_k^O .

Such Hamiltonian or quasi-Hamiltonian paths with specific extremities will be extensively used in the proof of Proposition 3. Indeed, we will regularly be able to show that a few edges between G_k^O and G_k^E are blue too. Then by choosing carefully the end-vertices of two (quasi-)Hamiltonian paths we will be able to combine the two paths into a cycle passing through all the vertices in $G_k^O \cup G_k^E$ except at most two of them, helping us to get a decomposition of the red zigzag graph into at most 3 monochromatic cycles.

2.2 Main proof

After these considerations, we may advance to our next result. As we shall see hereafter, Lemma 6 follows almost directly from the lemma below.

Lemma 13. *Let G be a red zigzag graph with $2n$ vertices which cannot be vertex-partitioned into at most 3 monochromatic cycles. If the subgraph $G[S_{\leq k}]$ is a blue even plait for some $1 \leq k \leq n - 2$, then $G[S_{\leq k+1}]$ is a blue even plait as well (see Definition 7).*

Proof. First since G cannot be vertex-partitioned into 3 monochromatic cycles, we can easily see that $n \geq 5$. It is also easy to check that $G[S_{\leq 4}]$ is a blue even plait: if the edge x_1y_1 (resp. x_2y_2) were red, then G (resp. $G \setminus \{x_1y_1\}$) would be Hamiltonian; if the edge x_2y_2 , x_1y_3 or x_3y_1 were red we could decompose G into a red cycle and one edge; x_3y_3 being red would imply a decomposition into a cycle and two paths. Now if x_2y_4 or x_4y_2 were red we could decompose G into a red cycle and two edges; if x_4y_4 were red then we would decompose into the big red cycle using x_4y_4 , the blue cycle we just showed the existence ($x_1y_1x_3y_3$) and the edge x_2y_2 . We have just proved that the even edges within $G[S_{\leq 4}]$ are blue, hence $G[S_{\leq 4}]$ is a blue even plait as we announced. Therefore we can assume that $k \geq 4$, so that each of G_k^O and G_k^E have at least 4 vertices.

Also, Remarks 8(a) and 12, imply that **the edge $x_{k+1}y_{k+1}$ is blue.**

Now, suppose by way of contradiction that $G[S_{\leq k+1}]$ is not a blue even plait. Hence, there is some natural j less than $k + 1$ such that j and $k + 1$ have the same parity and at least one of the edges $x_{k+1}y_j$ or x_jy_{k+1} is red. We shall show that this implies that G has a blue special set, which is a contradiction by Lemma 11.

We divide the remainder of the proof into two cases depending on the parity of k . These cases are treated quite similarly, but at some points they are not absolutely analogous. Therefore, we will give the full details for both cases whenever necessary but we will just guide how to make the analogy whenever possible.

Case A: k is even and $k \geq 4$, therefore j is odd with $1 \leq j < k \leq n - 2$.

Note that some claims below will require $k < n - 4$, but we will see that these claims are only needed if $k < n - 4$: otherwise the proof would end before.

Claim 1A.

- (i) *If there exists $1 \leq l \leq k$ odd such that $x_{k+1}y_l$ is red, then x_1y_k and $x_{k-1}y_2$ are not both blue.*
- (ii) *If there exists $1 \leq l \leq k$ odd such that x_ly_{k+1} is red, then x_ky_1 and x_2y_{k-1} are not both blue.*

Proof. We will prove only (i) since (ii) is analogous. So $x_{k+1}y_l$ is red. Suppose by way of contradiction that the edges x_1y_k and $x_{k-1}y_2$ are blue (see Figure 2.4). In this case, we shall prove that $S_{\leq k+2}$ is a blue special set of G .

First, let us prove that the edge x_ky_l is blue. Indeed, otherwise we could decompose G into the red cycle $(x_k, y_{k+1}Px_{k+1}, y_l)$ that covers $S_{\geq k+1} \cup \{x_k, y_l\}$ and a blue cycle that covers $S_{\leq k} \setminus \{x_k, y_l\}$. More precisely:

$$y_2 \overset{E_k \setminus x_k}{\rightsquigarrow} y_k \rightarrow x_1 \overset{O_k \setminus y_l}{\rightsquigarrow} x_{k-1} \rightarrow y_2.$$

Now, we can now construct a blue Hamiltonian cycle for $G[S_{\leq k}]$: say $y_k \overset{E_k}{\rightsquigarrow} x_k \rightarrow y_l \overset{O_k}{\rightsquigarrow} x_1 \rightarrow y_k$. This is our first step towards proving that $S_{\leq k+2}$ is a blue special set of G .

The edge $x_{k+1}y_{k-1}$ is blue, otherwise G could be vertex-partitioned into the red cycle $(x_k, y_{k+1}Px_{k+1}, y_{k-1})$ (recall that x_ky_{k-1} is also an edge of P) and (similarly to above) a blue cycle that covers all vertices in $S_{\leq k}$ except x_k and y_{k-1} . The edge $x_{l+1}y_{k+1}$ is blue too, since otherwise G could similarly be vertex-partitioned into the red cycle $(x_{l+1}, y_{k+1}Px_{k+1}, y_l)$ and a blue cycle that covers all vertices in $S_{\leq k}$ except x_{l+1} and y_l . Therefore, we see that the subgraph $G[S_{\leq k+1}]$ has a blue Hamiltonian cycle:

$$C = (x_{k+1} \rightarrow y_{k+1} \rightarrow x_{l+1} \overset{E_k}{\rightsquigarrow} y_2 \rightarrow x_{k-1} \overset{O_k}{\rightsquigarrow} y_{k-1} \rightarrow x_{k+1}).$$




Figure 2.4: The case where the edges x_1y_k and $x_{k-1}y_2$ are blue.

By Remark 8(a), it follows that the edge $x_{k+2}y_{k+2}$ is blue.

Next, the edge $x_{k+2}y_k$ is blue, otherwise G could be vertex-partitioned into the edge-cycle (x_{l+1}, y_{k+1}) , the red cycle $(x_{k+1}, y_{k+2}Px_{k+2}, y_k)$, and a blue cycle that covers $S_{\leq k} \setminus \{x_{l+1}, y_k\}$. Note that such blue cycle does not use y_k so it does not benefit from the blue edge x_1y_k but rather x_ky_l . Use edges of G_k^E to go from y_2 to x_k passing through all its vertices except x_{l+1} and y_k , add edge x_ky_l go through all vertices in G_k^O finishing in x_{k-1} and add the edge $x_{k-1}y_2$. The edge x_ky_{k+2} is blue too, since otherwise G could be vertex-partitioned into the red cycle $(x_k, y_{k+2}Px_{k+2}, y_{k+1})$, the edge-cycle (x_{k+1}, y_{k-1}) and a blue cycle that covers all vertices of $S_{\leq k} \setminus \{x_k, y_{k-1}\}$ (using the edges x_1y_k and $x_{k-1}y_2$).

Hence, we see that the subgraph $G[S_{\leq k+2}]$ has a blue Hamiltonian cycle that uses the edges $x_{k+1}y_{k+1}$ and $x_{k+2}y_{k+2}$, which can be built by the following manner: take the blue Hamiltonian cycle C of $G[S_{\leq k+1}]$ that we have just built. We can require it to contain the edge x_ky_k . Then consider the cycle $C' = (C - x_ky_k) + x_ky_{k+2}x_{k+2}y_k$. Therefore $S_{\leq k+2}$ is a blue special set of G as needed. \square

Claim 2A. *The edges $x_{k+1}y_1$ and x_1y_{k+1} are blue.*

Proof. Suppose by way of contradiction that the edge $x_{k+1}y_1$ is red (see Figure 2.5). In this case, G can be vertex-partitioned into the red cycle $(x_{k+1}Py_1)$ and the red path (x_1, y_2Px_{k-1}, y_k) . By Claim 1A(i), with $l = 1$, we know that the edges x_1y_k and $x_{k-1}y_2$ are not both blue. Hence, G can be vertex-partitioned into at most 3 monochromatic cycles, a contradiction. Thus, the edge $x_{k+1}y_1$ is blue. Analogously, the edge x_1y_{k+1} is blue too, by Claim 1A(ii). Thus, the result follows.




Figure 2.5: The case where the edge $x_{k+1}y_1$ is red.

\square

Claim 3A. *The edge $x_{k+2}y_{k+2}$ is blue.*

Proof. By Claim 2A, we see that the subgraph $G[S_{\leq k+1}]$ can be vertex-partitioned into a blue Hamiltonian cycle of G_k^E and a blue cycle that covers x_{k+1}, y_{k+1} and all vertices in G_k^O , which can be built by the following manner: take a blue Hamiltonian cycle of G_k^O that uses the edge x_1y_1 ; and use the path $(x_1, y_{k+1}, x_{k+1}, y_1)$ to extend it. Hence, by Remark 8(a), the result follows. \square

Remember that there is some odd j such that $x_{k+1}y_j$ or x_jy_{k+1} is red. In what remains, without loss of generality, **we assume that the edge $x_{k+1}y_j$ is red.**

Note: with that we loose the symmetry between the sets X and Y . We have not made this assumption earlier, because we need Claim 2A to hold regardless of which $x_{k+1}y_j$ or x_jy_{k+1} is red.

By Claim 2A, we know that $j \neq 1$, hence all the following claims will be on the assumption that $2 \leq j$. Besides, if $k = n - 2$ then the edge $x_{k+2}y_{k+2} = x_ny_n$ belongs to the red path P , hence is not blue, contradicting Claim 3A. Thus if $k = n - 2$ the proof ends here. In the remainder, therefore, we have that:

$$2 \leq j < k \leq n - 3.$$

This allows us to use indices $j - 1$ and $k + 3$ (that we have not used it so far).

Claim 4A. *The edges $x_{j+1}y_1$ and x_2y_{j+2} are blue.*

In particular, when $j = k - 1$, this means that x_ky_1 and x_2y_{k+1} are blue.

Proof. Suppose by way of contradiction that the edge $x_{j+1}y_1$ is red (see Figure 2.6). In this case, G can be vertex-partitioned into the red cycle $(x_{k+1}Px_{j+1}, y_1Py_j)$ and the red path (x_1, y_2Px_{k-1}, y_k) . By Claim 1A(i), we know that the edges x_1y_k and $x_{k-1}y_2$ are not both blue. Hence, we see that G can be vertex-partitioned into at most 3 monochromatic cycles, a contradiction. Thus, the edge $x_{j+1}y_1$ is blue.




Figure 2.6: The case where the edge $x_{j+1}y_1$ is red.

Now, suppose by way of contradiction that the edge x_2y_{j+2} is red (see Figure 2.7). In this case, the edge x_1y_k is blue, since otherwise G could be vertex-partitioned into the red cycles $(x_{k+1}Py_{j+2}, x_2Py_j)$ and (x_1, y_2Px_{k-1}, y_k) and the edge-cycle (x_{j+1}, y_1) . By Claim 1A(i), it follows that the edge $x_{k-1}y_2$ is red. Observe that the edges x_1y_1 and $x_{j+1}y_k$ are even and, therefore, blue. Hence, we see that G can be vertex-partitioned into the red cycles $(x_{k+1}Py_{j+2}, x_2Py_j)$ and (y_2Px_{k-1}) and the blue cycle (x_1, y_k, x_{j+1}, y_1) , a contradiction. Thus, the edge x_2y_{j+2} is blue.




Figure 2.7: The case where the edge x_2y_{j+2} is red.

□

Claim 5A. *The edges x_jy_{k+2} and $x_{k+2}y_k$ are blue.*

Proof. Suppose by way of contradiction that the edge x_jy_{k+2} is red (see Figure 2.8). In this case, G can be vertex-partitioned into the red cycle $(x_jPx_{k+1}, y_jPy_{k+2})$ and the subgraph $G[S_{\leq j-1}]$. But, observe that the subgraph $G[S_{\leq j-1}]$ is a blue even plait. Hence, by Remark 12, we see that G can be vertex-partitioned into 3 monochromatic cycles, a contradiction. Thus, the edge x_jy_{k+2} is blue.

Now, suppose by way of contradiction that the edge $x_{k+2}y_k$ is red. Take the red cycle $(x_{k+1}, y_{k+2}Px_{k+2}, y_k)$, the cover $S_{\geq k+2} \cup \{x_{k+1}, y_k\}$. We claim that the remaining vertices of G can be vertex-partitioned into at most 3 blue cycles. Indeed, if $j = k - 1$, by Claim 4A the edges x_ky_1 and x_2y_{k+1} are blue and by Claim 2A x_1y_{k+1} is blue (see Figure 2.9a). Take only the blue cycle $x_2 \xrightarrow{E_k \setminus y_k} x_k \rightarrow y_1 \xrightarrow{O_k^k} x_1 \rightarrow y_{k+1} \rightarrow x_2$.

On the other hand, if $j < k - 1$, then $y_{j+2} \in G_k^O$ (see Figure 2.9b). Again by Claim 4A and Claim 2A, we can form the blue cycle $x_2 \xrightarrow{E_k \setminus y_k} x_{j+1} \rightarrow y_1 \xrightarrow{O_k \setminus x_1} y_{j+2} \rightarrow x_2$ and we also take the edge-cycle (x_1, y_{k+1}) . □

Claim 6A. *The subgraph $G[S_{\leq k+2}]$ has a blue Hamiltonian cycle that uses the edges $x_{k+1}y_{k+1}$, $x_{k+2}y_{k+2}$ and $x_{k+2}y_k$. Furthermore, the edge $x_{k+3}y_{k+3}$ is blue.*




Figure 2.8: The case where the edge $x_j y_{k+2}$ is red.




Figure 2.9: The case where the edge $x_{k+2} y_k$ is red.

Proof. Let us build such a cycle (see Figure 2.10). We will use the blue edges $x_{j+1} y_1$ (Claim 4A), $x_{k+1} y_1$ and $x_1 y_{k+1}$ (Claim 2A), $x_{k+1} y_{k+1}$ (Remark 8(a)), $x_{k+2} y_k$ (Claim 5A), $x_{k+2} y_{k+2}$ (Claim 3A), and $x_j y_{k+2}$ (Claim 5A).

First take the cycle $C = x_{j+1} \xrightarrow{E_k^E} y_k \rightarrow x_{k+2} \rightarrow y_{k+2} \rightarrow x_j \xrightarrow{O_k^O} y_1 \rightarrow x_1 \rightarrow y_{k+1}$. It is a blue cycle covering all vertices of $S_{\leq k} \cup \{x_{k+2}, y_{k+2}\}$ and it uses the edge $x_1 y_1$. So $C' = C - x_1 y_1 + x_1 y_{k+1} x_{k+1} y_1$ is a blue cycle which satisfies the first part of the claim. By Remark 8(a), it follows that the edge $x_{k+3} y_{k+3}$ is blue. \square

As before, if $k = n - 3$ the proof ends here since we have just proved that the edge $x_n y_n$ is blue, whereas it belongs to the red path P . This is a contradiction. Hence, in the remaining claims

$$k \leq n - 4.$$

This allows us to use indices up to $k + 4$.

Claim 7A. *The edge $x_j y_{k+1}$ is blue.*

Proof. Suppose by way of contradiction that the edge $x_j y_{k+1}$ is red. In this case, we shall prove that $S_{\leq k+2}$ is a blue special set of G . Now Claim 2A tells us that the edges $x_{k+1} y_1$ and $x_1 y_{k+1}$ are blue. Claim 4A tells us that the edges $x_{j+1} y_1$ and $x_2 y_{j+2}$ are blue too. Also, since here we assume that the edge $x_j y_{k+1}$ is red we can use the ‘alternate’ version of Claim 4A consisting in exchanging X and Y , therefore the edges $x_1 y_{j+1}$ and $x_{j+2} y_2$ are also blue. Figure 2.11 shows all those edges.

On the one hand, if $j = k - 1$ then this means that the edges $x_2 y_{k+1}$ and $x_{k+1} y_2$ are blue. Then $x_2 \xrightarrow{E_k^E} y_2 \rightarrow x_{k+1} \rightarrow y_1 \xrightarrow{O_k^O} x_1 \rightarrow y_{k+1} \rightarrow x_2$ is a blue Hamiltonian cycle of $G[S_{\leq k+1}]$. On the other hand if




Figure 2.10: A blue Hamiltonian cycle of $G[S_{\leq k+2}]$.

$j < k - 1$ then $x_{j+2}, y_{j+2} \in G_k^O$. Start with $C = x_2 \xrightarrow{E_k^k} y_2 \rightarrow x_{j+2} \xrightarrow{O_k^k} y_{j+2} \rightarrow x_2$. C is a blue Hamiltonian cycle of $G[S_{\leq k}]$ and we can require it to use the edge x_1y_1 (G_k^O is a blue bipartite complete graph, hence $x_{j+2} \xrightarrow{O_k^k} y_{j+2}$ can be required to use x_1y_1). Therefore $C' = C - x_1y_1 + x_1y_{x+1}x_{k+1}y_1$ is a blue Hamiltonian cycle of $G[S_{\leq k+1}]$.

In both cases, by Claim 6A, we see that $S_{\leq k+2}$ is a blue special set of G , a contradiction. Thus, the result follows. \square




Figure 2.11: The case where the edge $x_j y_{k+1}$ is red.

Claim 8A. *There is a blue cycle in G that covers all vertices in $S_{\leq k+1}$ except x_{k+1} and y_k .*

Proof. By Claim 4A and Claim 2A the edges $x_{j+1}y_1$, x_2y_{j+2} and x_1y_{k+1} are blue.

On the one hand if $j = k - 1$, this means that the edges $x_k y_1$ and $x_2 y_{k+1}$ are blue (see Figure 2.12a). Then $x_2 \xrightarrow{E_k^k \setminus y_k} x_k \rightarrow y_1 \xrightarrow{O_k^k} x_1 \rightarrow y_{k+1} \rightarrow x_2$ is a blue cycle covering y_{k+1} and all vertices of $S_{\leq k}$ except y_k .

On the other hand ($j < k - 1$), $y_{j+2} \in G_k^O$. We will also need the edge $x_j y_{k+1}$, which is blue by Claim 7A. Let $C = x_2 \xrightarrow{E_k^k \setminus y_k} x_{j+1} \rightarrow y_1 \rightarrow x_1 \rightarrow y_{k+1} \rightarrow x_j \rightarrow y_{j+2} \rightarrow x_2$. This cycle covers $G_k^E \setminus y_k$ and passes through the vertices x_1, y_1, x_j, y_{j+2} in G_k^O . Now, we can replace the edge x_1y_1 of C with a path that covers all remaining vertices of G_k^O . \square

Claim 9A. *The edge $x_{k+1}y_{k+3}$ is blue.*




Figure 2.12: A blue cycle that covers all vertices in $S_{\leq k+1}$ except x_{k+1} and y_k .

Proof. Suppose by way of contradiction that the edge $x_{k+1}y_{k+3}$ is red (see Figure 2.13). Hence, by Claim 8A, we see that G can be vertex-partitioned into the red cycle $(x_{k+1}, y_{k+3}Px_{k+3}, y_{k+2})$, the edge-cycle (x_{k+2}, y_k) and a blue cycle that covers all vertices in $S_{\leq k+1}$ except x_{k+1} and y_k , a contradiction. Thus, the result follows. \square




Figure 2.13: The case where the edge $x_{k+1}y_{k+3}$ is red.

Claim 10A. *The edges x_1y_k and $x_{k+1}y_2$ are not both blue.*

Proof. Suppose by way of contradiction that the edges x_1y_k and $x_{k+1}y_2$ are blue (see Figure 2.14). In this case, we shall prove that $S_{\leq k+2}$ is a blue special set of G , a contradiction. The edge x_1y_{k+1} is blue by Claim 2A and $x_{j+1}y_1$ is blue by Claim 4A. So, $x_{j+1} \xrightarrow{E_k^k} y_k \rightarrow x_1 \xrightarrow{Q_k^k} y_1 \rightarrow x_{j+1}$ is a blue Hamiltonian cycle of $G[S_{\leq k}]$. And $x_{j+1} \xrightarrow{E_k^k} y_2 \rightarrow x_{k+1} \rightarrow y_{k+1} \rightarrow x_1 \xrightarrow{Q_k^k} y_1 \rightarrow x_{j+1}$ is a blue Hamiltonian cycle $G[S_{\leq k+1}]$. Together with Claim 6A, we see that $S_{\leq k+2}$ is a blue special set of G . Thus, the result follows. \square

Claim 11A. *The edge x_2y_{k+2} is blue.*

Proof. Suppose by way of contradiction that the edge x_2y_{k+2} is red (see Figure 2.15). In this case, G can be vertex-partitioned into the red cycle (x_2Py_{k+2}) , the red path (x_1, y_2Py_k, x_{k+1}) and the vertex y_1 . By Claim 10A, we know that the edges x_1y_k and $x_{k+1}y_2$ are not both blue. Hence, we see that G can be vertex-partitioned into 3 monochromatic cycles (one of which is an edge-cycle), a contradiction. Thus, the result follows. \square

Claim 12A. *The edge $x_{k+3}y_{k+1}$ is blue.*




Figure 2.14: The case where the edges x_1y_k and $x_{k+1}y_2$ are blue.




Figure 2.15: The case where the edge x_2y_{k+2} is red.

Proof. Suppose by way of contradiction that the edge $x_{k+3}y_{k+1}$ is red (see Figure 2.16). By Claim 4A and Claim 11A the edges $x_{j+1}y_1$ and x_2y_{k+2} are blue. Let $C = x_{j+1} \xrightarrow{E_k \setminus y_k} x_2 \rightarrow y_{k+2} \rightarrow x_j \xrightarrow{O_k} y_1 \rightarrow x_{j+1}$, which is a blue cycle that covers $(S_{\leq k} \setminus y_k) \cup \{y_{k+2}\}$. Then G can be vertex-partitioned into the red cycle $(x_{k+2}, y_{k+3} P x_{k+3}, y_{k+1})$, the edge-cycle (x_{k+1}, y_k) and C , a contradiction. Thus, the result follows. \square




Figure 2.16: The case where the edge $x_{k+3}y_{k+1}$ is red.

Claim 13A. *The subgraph $G[S_{\leq k+3}]$ has a blue Hamiltonian cycle that uses the edges $x_{k+2}y_k$ and $x_{k+3}y_{k+3}$. Consequently, the edge $x_{k+4}y_{k+4}$ is blue.*

Proof. By Claim 6A, start with a blue Hamiltonian cycle C of $G[S_{\leq k+2}]$ that uses the edges $x_{k+1}y_{k+1}$ and $x_{k+2}y_k$. Claim 6A also tell us that $x_{k+3}y_{k+3}$ is blue. Now, extend C by replacing the edge $x_{k+1}y_{k+1}$ by the path $x_{k+1}y_{k+3}x_{k+3}y_{k+1}$ (using Claims 9A and 12A). This gives the desired blue Hamiltonian cycle of $G[S_{\leq k+3}]$. By Remark 8(a), it follows that the edge $x_{k+4}y_{k+4}$ is blue. \square

Claim 14A. *The edges $x_{k+4}y_k$ and $x_{k+2}y_{k+4}$ are blue.*

Proof. Suppose by way of contradiction that the edge $x_{k+4}y_k$ is red (see Figure 2.17a). Hence, by Claim 8A, we see that G can be vertex-partitioned into the red cycle $(x_{k+1}, y_{k+2}, x_{k+3}, y_{k+4}Px_{k+4}, y_k)$, the edge-cycle (x_{k+2}, y_{k+3}) and a blue cycle that covers all vertices in $S_{\leq k+1}$ except x_{k+1} and y_k , a contradiction. Thus, the edge $x_{k+4}y_k$ is blue.




Figure 2.17: The cases of Claim 14A.

Now, suppose by way of contradiction that the edge $x_{k+2}y_{k+4}$ is red (see Figure 2.17b). By Claim 4A and Claim 11A the edges $x_{j+1}y_1$ and x_2y_{k+2} are blue. Together with Claim 2A we build the blue cycle $C = x_{j+1} \xrightarrow{E_k \setminus y_k} x_2 \rightarrow y_{k+2} \rightarrow x_j \xrightarrow{O_k \setminus y_1} x_1 \rightarrow y_{k+1} \rightarrow x_{k+1} \rightarrow y_1 \rightarrow x_{j+1}$ that covers $(S_{\leq k+1} \setminus y_k) \cup \{y_{k+2}\}$. Thus G can be vertex-partitioned into the red cycle $(x_{k+2}, y_{k+4}Px_{k+4}, y_{k+3})$, the edge-cycle (x_{k+3}, y_k) and C . Hence, we get a contradiction. Thus, the edge $x_{k+2}y_{k+4}$ is blue. \square

Claim 15A. *The subgraph $G[S_{\leq k+4}]$ has a blue Hamiltonian cycle that uses the edges $x_{k+3}y_{k+3}$ and $x_{k+4}y_{k+4}$.*

Proof. Using Claim 13A, take a blue Hamiltonian cycle of $G[S_{\leq k+3}]$ that uses the edges $x_{k+2}y_k$ and $x_{k+3}y_{k+3}$ and extend it to a blue Hamiltonian cycle of $G[S_{\leq k+4}]$ that uses the edges $x_{k+3}y_{k+3}$ and $x_{k+4}y_{k+4}$ by replacing the edge $x_{k+2}y_k$ by the path $(x_{k+2}, y_{k+4}, x_{k+4}, y_k)$, which is blue by Claims 13A and 14A. \square

Thus, by Claims 6A, 13A and 15A, we see that $S_{\leq k+4}$ is a blue special set of G , which is our final contradiction for Case A.

Case B: k is odd and $k \geq 4$, therefore j is even with $2 \leq j < k \leq n - 2$.

Most proofs are similar to the ones in Case A. We will give the details on how to derive the A proofs to B proofs. Of course, we should also replace any reference to a Claim A to its Claim B counterpart.

Claim 1B.

- (i) *If there exists $1 \leq l \leq k$ odd such that $x_{k+1}y_l$ is red, then x_2y_k and $x_{k-1}y_1$ are not both blue.*
- (ii) *If there exists $1 \leq l \leq k$ odd such that $x_{k+1}y_l$ is red, then x_ky_2 and x_1y_{k-1} are not both blue.*

Proof. Interchange the roles of x_1 and x_2 , of y_1 and y_2 and of G_k^O and G_k^E in the proof of Claim 1A. \square

Claim 2B. *The edges $x_{k+1}y_2$ and x_2y_{k+1} are blue.*

Proof. Suppose by way of contradiction that the edge $x_{k+1}y_2$ is red (see Figure 2.18). In this case, G can be vertex-partitioned into the red cycle $(x_{k+1}Py_2)$, the red path (y_1, x_2Px_{k-1}, y_k) and the vertex x_1 (a difference here is that in Claim 2A we did not have this hanging vertex x_1). By Claim 1B(i), we know that the edges x_2y_k and $x_{k-1}y_1$ are not both blue. Hence, we see that G can be vertex-partitioned into 3 monochromatic cycles, a contradiction. Thus, the edge $x_{k+1}y_2$ is blue. Analogously, swapping the roles of X and Y and using Claim 1B(ii), the edge x_2y_{k+1} is blue too. Thus, the result follows. \square




Figure 2.18: The case where the edge $x_{k+1}y_2$ is red. \square

Claim 3B. *The edge $x_{k+2}y_{k+2}$ is blue.*

Proof. Just interchange the roles of x_1 and x_2 , of y_1 and y_2 and of G_k^O and G_k^E . \square

As in Case A, from this point, without loss of generality **we assume that the edge $x_{k+1}y_j$ is red.** By Claim 2B we know that $j \neq 2$. And as in Case A, if $k = n - 2$, then the proof ends here: we proved that x_ny_n is red, a contradiction. Hence from now:

$$2 < j \leq k \leq n - 3$$

Claim 4B. *The edges $x_{j+1}y_2$ and x_1y_{j+2} are blue. In particular, when $j = k - 1$, this means that x_ky_2 and x_1y_{k+1} are blue.*

Proof. This proof is similar to Claim 4A, with minor changes. First, to prove that the edge $x_{j+1}y_2$ is blue, we do as in Claim 2B, interchanging the roles of x_1 and x_2 , and of y_1 and y_2 .

Now, suppose by way of contradiction that the edge x_1y_{j+2} is red (see Figure 2.19). Then, G can be vertex-partitioned into the red cycle $(x_{k+1}Py_{j+2}, x_1Py_j)$, the red path (y_1, x_2Px_{k-1}, y_k) and the vertex x_{j+1} . By Claim 1B, we know that the edges x_2y_k and $x_{k-1}y_1$ are not both blue. Hence, G can be vertex-partitioned into 3 monochromatic cycles, a contradiction. Thus, the edge x_1y_{j+2} is blue. \square



Figure 2.19: The case where the edge x_1y_{j+2} is red. \square

Claim 5B. *The edges x_jy_{k+2} and $x_{k+2}y_k$ are blue.*

Proof. To show that x_jy_{k+2} is blue, use the proof of Claim 5A as is (see Figure 2.8). Then to show that $x_{k+2}y_k$ is blue, do as in Claim 5A, interchanging the roles of x_1 and x_2 , of y_1 and y_2 , of O_k and E_k , and of G_k^O and G_k^E . \square

Claim 6B. *The subgraph $G[S_{\leq k+2}]$ has a blue Hamiltonian cycle that uses the edges $x_{k+1}y_{k+1}$, $x_{k+2}y_{k+2}$ and $x_{k+2}y_k$. Furthermore, the edge $x_{k+3}y_{k+3}$ is blue.*

Proof. Interchange the roles of x_1 and x_2 , of y_1 and y_2 and of G_k^O and G_k^E in the proof of Claim 6A. \square

As in Case A, if $k = n - 3$ then the proof ends here: we proved that x_ny_n is red, a contradiction. Hence, in the remaining claims we have

$$k \leq n - 4.$$

Claim 7B. *The edge x_jy_{k+1} is blue.*

Proof. Interchange the roles of x_1 and x_2 , of y_1 and y_2 and of G_k^O and G_k^E in the proof of Claim 7A. \square

Claim 8B. *There is a blue cycle in G that covers all vertices in $S_{\leq k+1}$ except x_{k+1} and y_k .*

Proof. Interchange the roles of x_1 and x_2 , of y_1 and y_2 and of G_k^O and G_k^E in the proof of Claim 8A. \square

Claim 9B. *The edge $x_{k+1}y_{k+3}$ is blue.*

Proof. The proof of Claim 9A can be used as is. \square

Claim 10B. *The edges x_2y_k and $x_{k+1}y_1$ are not both blue.*

Proof. Interchange the roles of x_1 and x_2 , of y_1 and y_2 and of G_k^O and G_k^E in the proof of Claim 10A. \square

Claim 11B. *The edge x_1y_{k+2} is blue.*

Proof. Suppose by way of contradiction that the edge x_1y_{k+2} is red (see Figure 2.20). In this case, G can be vertex-partitioned into the red cycle (x_1Py_{k+2}) and the red path (x_{k+1}, y_kPx_2, y_1) . By Claim 10B, we know that the edges x_2y_k and $x_{k+1}y_1$ are not both blue. Hence, we see that G can be vertex-partitioned into at most 3 monochromatic cycles, a contradiction.




Figure 2.20: The case where the edge x_1y_{k+2} is red.

\square

Claim 12B. *The edge $x_{k+3}y_{k+1}$ is blue.*

Proof. Interchange the roles of x_1 and x_2 , of y_1 and y_2 and of G_k^O and G_k^E in the proof of Claim 12A. \square

Claim 13B. *The subgraph $G[S_{\leq k+3}]$ has a blue Hamiltonian cycle that uses the edges $x_{k+2}y_k$ and $x_{k+3}y_{k+3}$. Consequently, the edge $x_{k+4}y_{k+4}$ is blue.*

Proof. The proof of Claim 13A can almost be used as is: just replace Claim 6A by 6B. \square

Claim 14B. *The edges $x_{k+4}y_k$ and $x_{k+2}y_{k+4}$ are blue.*

Proof. Interchange the roles of x_1 and x_2 , of y_1 and y_2 and of G_k^O and G_k^E in the proof of Claim 14A. \square

Claim 15B. *The subgraph $G[S_{\leq k+4}]$ has a blue Hamiltonian cycle that uses the edges $x_{k+3}y_{k+3}$ and $x_{k+4}y_{k+4}$.*

Proof. We can use the proof of Claim 15A almost as is: just use Claim 13B instead of Claim 13A. \square

Thus, by Claims 6B, 13B and 15B, we see that $S_{\leq k+4}$ is a blue special set of G , which is our final contradiction for Case B.

This ends the proof of Lemma 13. \square

Now, we may finally prove Lemma 6.

Proof of Lemma 6. Let us take G as in the statement: a balanced complete bipartite graph whose edges are coloured red or blue, and which has a monochromatic Hamiltonian path. Without loss of generality, we can assume this path is red and that its vertices are labelled so that it is a red zigzag graph. Assume for contradiction that G cannot be partitioned into three monochromatic cycles. Let the number of vertices of G be $2n$.

By Remark 8(a), we may assume that the edges x_1y_1 and x_2y_2 are blue, since otherwise we would be done. Thus, the subgraph $G[S_{\leq 2}]$ is a blue even plait and so we may apply Lemma 13 iteratively until $k = n - 2$ inclusive. Therefore $S_{\leq n-1}$ is a blue plait, hence it can be decomposed into two blue cycles C_E and C_O by Remark 12. Finally, G can be decomposed into the red edge-cycle x_ny_n and the two blue cycles C_E and C_O , which is a contradiction and concludes the proof. \square

3 Conclusions and perspectives

In this paper, we proved a new bound on the number of monochromatic cycles needed to partition the vertices of any edge-bicoloured complete balanced bipartite graph. We made quite an improvement as the previous known bound was 12 and we reduced it to 4. Also we can point out that our work concerns all graphs, while many paper tackling similar problems focus on large graphs. For this reason our proofs do not resort to strong results, and is self contained apart from Stein's result (Corollary 5) for which we also had an alternate but very similar proof, hence not worthy of being shown here.

We wrote the proof of our main result in a very detailed way, with the goal to make it easy to be verified. So it ended up being somewhat long. Of course, one can make it shorter (but harder to check) by omitting some trivial details and remarks or not explicitly listing all cycles (the reader is free for not reading this details and search for the cycles in the pictures). We could also have omitted the “B claims” (when k is odd), asking the reader to believe that the proofs are similar to the ones of the case when k is even. However, we did not find a way to make the analysis intrinsically shorter. A good reason for this is that many of those cases have to be checked even to deal with small values of k (say $k \leq 10$).

We also showed that if the colouring is split, then we can partition the vertices into at most three monochromatic cycles and described exactly when 2 cycles are enough (and when are not). And we gave examples of a 2-colouring of a complete balanced bipartite graph that is not split and also cannot be partitioned into 2 cycles. Therefore, it remains to decide if the 2-colour cycle partition number of balanced bipartite graphs is 3 or 4.

References

- [1] BESSY, S., AND THOMASSÉ, S. Partitioning a graph into a cycle and an anticycle, a proof of lehel's conjecture. *Journal of Combinatorial Theory, Series B* 100, 2 (2010), 176–180.
- [2] GERENCSÉR, L., AND GYÁRFÁS, A. On ramsey-type problems. *Ann. Univ. Sci. Budapest. Eötvös Sect. Math* 10 (1967), 167–170.
- [3] GYÁRFÁS, A. Vertex coverings by monochromatic paths and cycles. *Journal of Graph Theory* 7, 1 (1983), 131–135.
- [4] GYÁRFÁS, A. Vertex covers by monochromatic pieces — a survey of results and problems. *Discrete Mathematics* 339, 7 (2016), 1970–1977.
- [5] GYÁRFÁS, A., AND LEHEL, J. A ramsey-type problem in directed and bipartite graphs. *Periodica Mathematica Hungarica* 3, 3-4 (1973), 299–304.
- [6] GYÁRFÁS, A., RUSZINKÓ, M., SÁRKÖZY, G. N., AND SZEMERÉDI, E. An improved bound for the monochromatic cycle partition number. *Journal of Combinatorial Theory, Series B* 96, 6 (2006), 855–873.
- [7] LUCZAK, T., RÖDL, V., AND SZEMERÉDI, E. Partitioning two-coloured complete graphs into two monochromatic cycles. *Combinatorics, Probability and Computing* 7, 04 (1998), 423–436.
- [8] POKROVSKIY, A. Partitioning edge-coloured complete graphs into monochromatic cycles and paths. *Journal of Combinatorial Theory, Series B* 106 (2014), 70–97.
- [9] SCHAUDT, O., AND STEIN, M. Partitioning two-coloured complete multipartite graphs into monochromatic paths and cycles. *Electronic Notes in Discrete Mathematics* 50 (2015), 313–318.
- [10] STEIN, M. Monochromatic paths in 2-edge-coloured graphs and hypergraphs. *Electron. J. Comb.* 30, 1 (2023).