

# A Refinement of the Multinomial Distribution with Application

Andrew V. Sills  
Department of Mathematical Sciences,  
Georgia Southern University, Statesboro and Savannah, Georgia, USA  
email: asills@georgiasouthern.edu

September 10, 2024

Key words: multinomial distribution;  $q$ -multinomial theorem, permutation inversion statistic  
MSC: 60E05, 05A05

## Abstract

A refinement of the multinomial distribution is presented where the number of inversions in the sequence of outcomes is tallied. This refinement of the multinomial distribution is its joint distribution with the number of inversions in the accompanying experiment. An application of this additional information is described in which the number of inversions acts as a proxy measure of homogeneity (or lack thereof) in the sequence of outcomes.

## Competing Interests Statement

The author has no competing interests to declare that are relevant to the content of this article. No funds, grants, or other support was received.

## 1 Preliminaries

The author sets out to establish a refinement of the classical multinomial probability distribution that generalizes his earlier work (Sills, 2023) and here draws inspiration from the non-commutative  $q$ -multinomial theorem of Farsi (2007). This multinomial distribution generalization is different that of Charalambides (2021), which makes use of the commutative  $q$ -multinomial theorem.

Recall the classical multinomial experiment:

- The experiment consists of  $n$  independent, identical trials;
- there are  $k$  possible outcomes on each trial, labeled outcome  $1, 2, \dots, k$ ;
- on a given trial, the probability that outcome  $i$  occurs is the constant  $p_i$ .

Define the random variable  $Y_i$  as the number of trials that result in outcome  $i$ . Let  $\mathbf{Y} = (Y_1, Y_2, \dots, Y_k)^T$  and  $\mathbf{y} = (y_1, y_2, \dots, y_k)^T$ .

The joint pmf for  $\mathbf{Y}$  is

$$P(\mathbf{Y} = \mathbf{y}) = \binom{y_1 + y_2 + \dots + y_k}{y_1, y_2, \dots, y_k} p_1^{y_1} p_2^{y_2} \dots p_k^{y_k},$$

for all nonnegative integers  $y_1, y_2, \dots, y_k$  that sum to  $n$ , and 0 otherwise; where

$$\binom{y_1 + y_2 + \dots + y_k}{y_1, y_2, \dots, y_k} = \frac{(y_1 + y_2 + \dots + y_k)!}{y_1! y_2! \dots y_k!} \quad (1.1)$$

is the multinomial coefficient and  $\sum_{i=1}^k p_i = 1$ . In this case, we may write  $\mathbf{Y} \sim \text{MULT}(n; p_1, p_2, \dots, p_{k-1})$  noting that once  $p_1, p_2, \dots, p_{k-1}$  are specified, it must be the case that  $p_k = 1 - \sum_{i=1}^{k-1} p_i$ .

A *weak  $k$ -composition of  $n$*  is a  $k$ -tuple of nonnegative integers whose components sum to  $n$ . For example, there are 10 weak 3-compositions of 3: namely

$$\begin{aligned} & (0, 0, 3) \quad (0, 1, 2) \quad (0, 2, 1) \quad (0, 3, 0) \quad (1, 0, 2) \\ & (1, 1, 1) \quad (1, 2, 0) \quad (2, 0, 1) \quad (2, 1, 0) \quad (3, 0, 0). \end{aligned}$$

In general, the number of weak  $k$ -compositions of  $n$  is  $\binom{n+k-1}{k-1}$ . Denote the set of weak  $k$ -compositions of  $n$  by  $\mathcal{C}_k(n)$ .

The weak  $k$ -compositions of  $n$  comprise the support of the random vector  $\mathbf{Y}$ , where  $\mathbf{Y} \sim \text{MULT}(n; p_1, p_2, \dots, p_{k-1})$ .

A standard reference on integer compositions is Heubach and Mansour (2009).

The  $q$ -multinomial coefficient (or *Gaussian multinomial coefficient*) is given by

$$\left[ \begin{matrix} y_1 + y_2 + \dots + y_k \\ y_1, y_2, \dots, y_k \end{matrix} \right]_q = \frac{(q)_{y_1 + y_2 + \dots + y_k}}{(q)_{y_1} (q)_{y_2} \dots (q)_{y_k}}, \quad (1.2)$$

where  $(q)_j := (1 - q)(1 - q^2) \dots (1 - q^j)$ . Note that the  $q$ -multinomial coefficient in (1.2) is a polynomial in  $q$  of degree

$$\sum_{1 \leq i < j \leq k} y_i y_j$$

and if  $q$  is set to 1 in (1.2), then the ordinary multinomial coefficient (1.1) is recovered.

Let us denote the multiset  $M = (a_1, a_2, \dots, a_n)$ , where the first  $m_1$  elements equal 1, the next  $m_2$  elements equal 2, the next  $m_3$  elements equal 3,  $\dots$ , the last  $m_k$  elements equal  $k$ , and  $n = m_1 + m_2 + \dots + m_k$ , by the notation  $\{1^{m_1} 2^{m_2} \dots k^{m_k}\}$ . A permutation  $(\xi_1, \xi_2, \dots, \xi_n)$  of  $M$  is said to have  $i$  *inversions* if there are exactly  $i$  pairs  $(\xi_i, \xi_j)$  such that  $i < j$  and  $\xi_i > \xi_j$ . The number of permutations of  $M$  having exactly  $i$  inversions is denoted  $\text{inv}(m_1, m_2, \dots, m_k; i)$ .

The  $q$ -multinomial coefficient figures prominently in the theory of permutations of the multiset  $\{1^{m_1} 2^{m_2} \dots k^{m_k}\}$ . In fact, MacMahon (1917) proved that

$$\sum_{i \geq 0} \text{inv}(y_1, y_2, \dots, y_k; i) q^i = \begin{bmatrix} y_1 + y_2 + \dots + y_k \\ y_1, y_2, \dots, y_k \end{bmatrix}_q, \quad (1.3)$$

i.e. that the  $q$ -multinomial coefficient is the ordinary power series generating function for the function that counts the number of inversions in the multiset  $M$  defined above.

By setting  $q = 1$  in (1.3), it is immediate that

$$\sum_{i \geq 0} \text{inv}(y_1, y_2, \dots, y_k; i) = \binom{y_1 + y_2 + \dots + y_k}{y_1, y_2, \dots, y_k}. \quad (1.4)$$

For more on the  $q$ -multinomial coefficients, see section 3.3 of Andrews (1976).

## 2 A refined multinomial distribution with mathematical motivation

### 2.1 A special case

To motivate the refinement about to be proposed, let us consider in detail the specific example where  $(Y_1, Y_2, Y_3) \sim \text{MULT}(3; p_1, p_2)$ ; of course,  $Y_3 = 3 - Y_1 - Y_2$  and  $p_3 = 1 - p_1 - p_2$ . That is, we are considering  $n = 3$  trials with  $k = 3$  possible outcomes on each trial. In the associated multinomial experiment, there are  $k^n = 3^3 = 27$  possible outcomes of the  $n = 3$  trials. Each outcome is enumerated explicitly in the following table:

$\mathbf{y}$	$P(\mathbf{Y} = \mathbf{y})$	outcome(s)	inv
(0, 0, 3)	$p_3^3$	333	0
(0, 1, 2)	$3p_2p_3^2$	233	0
		323	1
		332	2
(0, 2, 1)	$3p_2^2p_3$	223	0
		232	1
		322	2
(0, 3, 0)	$p_2^3$	222	0
(1, 0, 2)	$p_1p_3^2$	133	0
		313	1
		331	2
(1, 1, 1)	$6p_1p_2p_3$	123	0
		132	1
		213	1
		231	2
		312	2
		321	3
(1, 2, 0)	$3p_1p_2^2$	122	0
		212	1
		221	2
(2, 0, 1)	$3p_1^2p_3$	113	0
		131	1
		311	2
(2, 1, 0)	$3p_1^2p_2$	112	0
		121	1
		211	2
(3, 0, 0)	$p_1^3$	111	0

Table 1. The distribution  $\text{MULT}(3; p_1, p_2)$  refined with inversion count

If we ignore the last column, we have the classical multinomial distribution, where the number of each of the  $k$  possible outcomes is recorded over the  $n$  trials. However, the last column suggests that we could additionally keep track of the number of inversions, where we view the sequence of outcomes across the  $n = 3$  trials as some permutation of  $\{1^{y_1}2^{y_2}3^{y_3}\}$ .

## 2.2 The random variable that counts the number of inversions in the outcomes of the sequence of trials, and the resulting joint pmf

Let  $I$  denote the random variable that records the number of inversions in outcome of the multinomial experiment and let  $\mathbf{X} = (Y_1, Y_2, \dots, Y_k, I)$ , a random vector with  $k + 1$  components. Then it seems reasonable to define the joint pmf

$$P(\mathbf{X} = \mathbf{x}) = P(\mathbf{Y} = \mathbf{y}, I = i) = \text{inv}(y_1, y_2, \dots, y_k; i) p_1^{y_1} p_2^{y_2} \cdots p_k^{y_k}, \quad (2.1)$$

for  $\mathbf{y} \in \mathcal{C}_k(n)$  and nonnegative integers  $i$ ; and 0 otherwise.

*Remark 2.1.* Actually, the support of  $I$  is *not* all nonnegative integers  $i$ , although if a value of  $i$  that is out of bounds is selected, the value of  $\text{inv}(y_1, \dots, y_k; i)$  will equal 0, so no real harm done by extending  $i$  over all nonnegative integers. For a fixed  $n$  and  $k$ , it appears that the maximum  $i$  for which  $\text{inv}(y_1, \dots, y_k; i) > 0$  is  $\lfloor (k-1)n^2/(2k) \rfloor$ .

Notice that (2.1) is a mathematically legitimate joint pmf, as  $P(\mathbf{Y} = \mathbf{y}, I = i) \geq 0$  for all  $\mathbf{y}$  and  $i$ , and

$$\begin{aligned} \sum_{\mathbf{y} \in \mathcal{C}_k(n)} \sum_{i \geq 0} P(\mathbf{Y} = \mathbf{y}, I = i) &= \sum_{\mathbf{y} \in \mathcal{C}_k(n)} \sum_{i \geq 0} \text{inv}(y_1, \dots, y_k; i) p_1^{y_1} \cdots p_k^{y_k} \\ &= \sum_{\mathbf{y} \in \mathcal{C}_k(n)} p_1^{y_1} \cdots p_k^{y_k} \sum_{i \geq 0} \text{inv}(y_1, \dots, y_k; i) \\ &= \sum_{\mathbf{y} \in \mathcal{C}_k(n)} p_1^{y_1} \cdots p_k^{y_k} \binom{y_1 + \cdots + y_k}{y_1, \dots, y_k} \\ &= 1. \end{aligned}$$

From this perspective, the previously considered  $n = k = 3$  case can be tabulated as follows, where each cell contains the relevant sequence of outcomes, and the probability of that particular value of  $(Y_1, Y_2, Y_3)$  and  $I$ :

$\mathbf{y}$	$I$			
	0	1	2	3
(0,0,3)	333 $p_3^3$	— 0	— 0	— 0
(0,1,2)	233 $p_2p_3^2$	323 $p_2p_3^2$	332 $p_2p_3^2$	— 0
(0,2,1)	223 $p_2^2p_3$	232 $p_2^2p_3$	322 $p_2^2p_3$	— 0
(0,3,0)	222 $p_2^3$	— 0	— 0	— 0
(1,0,2)	133 $p_1p_3^2$	313 $p_1p_3^2$	331 $p_1p_3^2$	— 0
(1,1,1)	123 $p_1p_2p_3$	132, 213 $2p_1p_2p_3$	231, 312 $2p_1p_2p_3$	321 $p_1p_2p_3$
(1,2,0)	122 $p_1p_2^2$	212 $p_1p_2^2$	221 $p_1p_2^2$	— 0
(2,0,1)	113 $p_1^2p_3$	131 $p_1^2p_3$	311 $p_1^2p_3$	— 0
(2,1,0)	112 $p_1^2p_2$	121 $p_1^2p_2$	211 $p_1^2p_2$	— 0
(3,0,0)	111 $p_1^3$	— 0	— 0	— 0

Table 2. The joint distribution of  $(\mathbf{Y}, I)$ .

Notice how the coefficient of probability expression of the cell in row  $\mathbf{Y} = (y_1, y_2, y_3)$  and column  $I = i$  equals the coefficient of  $q^i$  in the expansion of the  $q$ -multinomial coefficient  $\begin{bmatrix} y_1+y_2+y_3 \\ y_1, y_2, y_3 \end{bmatrix}_q$ . For instance,  $\begin{bmatrix} 3 \\ 1, 1, 1 \end{bmatrix}_q = 1 + 2q + 2q^2 + q^3$ , as a demonstration of Eq. (1.3).

### 2.3 Farsi's noncommutative $q$ -multinomial theorem

For additional mathematical motivation, recall the non-commutative  $q$ -multinomial theorem of Farsi (2007). In particular, consider (Farsi, 2007, p. 1539, Example 2.8), with each  $\rho_j$  set equal to  $q$ , and notation adjusted for our present purposes: Let  $n$  denote a positive integer and let  $x_1, x_2, \dots, x_k$  denote multiplicatively noncommutative indeterminates that satisfy the  $q$ -commutation relation

$$x_j x_i = q x_i x_j \text{ whenever } i < j; \quad (2.2)$$

then

$$(x_1 + x_2 + \cdots + x_k)^n = \sum_{\mathbf{y} \in \mathcal{C}_k(n)} \begin{bmatrix} y_1 + y_2 + \cdots + y_k \\ y_1, y_2, \dots, y_k \end{bmatrix}_q x_1^{y_1} x_2^{y_2} \cdots x_k^{y_k}. \quad (2.3)$$

The left member of (2.3), upon expansion, encodes all possible permutations of  $\{1^{y_1} 2^{y_2} \cdots k^{y_k}\}$  for all ordered  $k$ -tuples of nonnegative integers  $(y_1, \dots, y_k)$  that sum to  $n$ , i.e. all weak  $k$ -compositions of  $n$ , as a sum of  $k^n$  terms, each consisting of a noncommutative product of  $n$   $x_i$ 's, in every possible order. The right member expresses an algebraically equivalent expression, taking into account the  $q$ -commutation relation (2.2), where the  $x_i$ 's appear in their "natural" order, i.e. with increasing subscripts. In the  $n = k = 3$  case, the left member of (2.3) expands as these 27 terms:

$$\begin{aligned} & x_3 x_3 x_3 + x_2 x_3 x_3 + x_3 x_2 x_3 + x_3 x_3 x_2 + x_2 x_2 x_3 + x_2 x_3 x_2 + x_3 x_2 x_2 + x_2 x_2 x_2 \\ & + x_1 x_3 x_3 + x_3 x_1 x_3 + x_3 x_3 x_1 + x_1 x_2 x_3 + x_1 x_3 x_2 + x_2 x_1 x_3 + x_2 x_3 x_1 + x_3 x_1 x_2 \\ & + x_3 x_2 x_1 + x_1 x_2 x_2 + x_2 x_1 x_2 + x_2 x_2 x_1 + x_1 x_1 x_3 + x_1 x_3 x_1 + x_3 x_1 x_1 \\ & + x_1 x_1 x_2 + x_1 x_2 x_1 + x_2 x_1 x_1 + x_1 x_1 x_1, \quad (2.4) \end{aligned}$$

where the set of possible outcomes is enumerated by following the subscripts. In contrast, the right member of (2.3) in the  $n = k = 3$  case expands as

$$\begin{aligned} & (1)x_3^3 + (1 + q + q^2)x_2 x_3^2 + (1 + q + q^2)x_2^2 x_3 + (1)x_2^2 + (1 + q + q^2)x_1 x_3^2 \\ & + (1 + 2q + 2q^2 + q^3)x_1 x_2 x_3 + (1 + q + q^2)x_1 x_2^2 + (1 + q + q^2)x_1^2 x_3 \\ & + (1 + q + q^2)x_1^2 x_2 + (1)x_1^3. \quad (2.5) \end{aligned}$$

Here, the number of inversions required to transform  $x_1^{y_1} x_2^{y_2} \cdots x_k^{y_k}$  into the permuted form as required by (2.2) is recorded in each exponent of  $q$ .

If (1.3) is applied to the right member of (2.3), the result is:

$$\begin{aligned} & \sum_{\mathbf{y} \in \mathcal{C}_k(n)} \begin{bmatrix} y_1 + y_2 + \cdots + y_k \\ y_1, y_2, \dots, y_k \end{bmatrix}_q x_1^{y_1} x_2^{y_2} \cdots x_k^{y_k} \\ & = \sum_{\mathbf{y} \in \mathcal{C}_k(n)} \sum_{i \geq 0} \text{inv}(y_1, \dots, y_k; i) q^i x_1^{y_1} x_2^{y_2} \cdots x_k^{y_k} \quad (2.6) \end{aligned}$$

Consider the generic summand of the right member of (2.6). Substitute  $x_i = p_i$  for  $i = 1, 2, \dots, k$ , and take  $q = 1$ . The resulting expression is the joint pmf of  $\mathbf{Y}$  and  $I$  summed over its support (2.1).

## 2.4 Some Marginal distributions

Clearly, we have  $\mathbf{Y} \sim \text{MULT}(n; p_1, p_2, \dots, p_{k-1})$  as this was our starting point before introducing the random variable  $I$ . Additionally,  $Y_j \sim \text{BIN}(n, p_j)$  for each  $j$ ,  $1 \leq j \leq k$ , and accordingly for any such  $i$ , the joint pmf of  $Y_j$  and  $I$  is

$$P(Y_j = y_j, I = i) = \text{inv}(y_j, n - y_j; i) p_j^{y_j} (1 - p_j)^{n - y_j}, \quad (2.7)$$

for  $0 \leq y_j \leq n$  and  $0 \leq i \leq \lfloor n^2/4 \rfloor$ , and 0 otherwise. This case was studied in Sills (2023).

## 3 A summation lemma to aid calculations of moments

**Lemma 3.1.** *Let  $y_1, y_2, \dots, y_k$  be fixed nonnegative integers.*

$$\sum_{i \geq 0} i \cdot \text{inv}(y_1, \dots, y_k; i) = \frac{1}{2} \binom{y_1 + \dots + y_k}{y_1, \dots, y_k} \sum_{1 \leq i < j \leq k} y_i y_j \quad (3.1)$$

$$\begin{aligned} & \sum_{i \geq 0} i^2 \cdot \text{inv}(y_1, \dots, y_k; i) \\ &= \binom{y_1 + \dots + y_k}{y_1, \dots, y_k} \left( \frac{1}{12} \sum_{1 \leq i < j \leq k} y_i y_j + \frac{1}{12} \sum_{1 \leq i < j \leq k} y_i^2 y_j + \frac{1}{12} \sum_{1 \leq i < j \leq k} y_i y_j^2 \right. \\ & \quad + \frac{1}{4} \sum_{1 \leq i < j \leq k} y_i^2 y_j^2 + \frac{1}{6} \sum_{1 \leq h < i < j \leq k} y_h y_i y_j + \frac{1}{2} \sum_{1 \leq h < i < j \leq k} y_h^2 y_i y_j \\ & \quad \left. + \frac{1}{2} \sum_{1 \leq h < i < j \leq k} y_h y_i^2 y_j + \frac{1}{2} \sum_{1 \leq h < i < j \leq k} y_h y_i y_j^2 + \frac{3}{2} \sum_{1 \leq g < h < i < j \leq k} y_g y_h y_i y_j \right) \\ &= \frac{1}{12} \binom{y_1 + \dots + y_k}{y_1, \dots, y_k} \left\{ \sum_{1 \leq i < j \leq k} y_i y_j + \left( \sum_{1 \leq h \leq i < j \leq k} + \sum_{1 \leq h < i \leq j \leq k} \right) y_h y_i y_j \right. \\ & \quad + 6 \left( \sum_{1 \leq g \leq h < i < j \leq k} + \sum_{1 \leq g < h \leq i < j \leq k} + \sum_{1 \leq g < h < i \leq j \leq k} \right) y_g y_h y_i y_j \\ & \quad \left. + 3 \sum_{1 \leq g = h < i = j \leq k} y_g y_h y_i y_j \right\}. \quad (3.2) \end{aligned}$$



For (3.1), the key is to note that the sum  $\sum_{i \geq 0} i \cdot \text{inv}(y_1, \dots, y_k; i)$  can be obtained by differentiating the  $q$ -multinomial coefficient with respect to  $q$ , and then setting  $q = 1$ :

$$\sum_{i \geq 0} i \cdot \text{inv}(y_1, \dots, y_k; i) = \frac{d}{dq} \left[ \begin{matrix} y_1 + \dots + y_k \\ y_1, \dots, y_k \end{matrix} \right]_q \Big|_{q=1}.$$

(MacMahon, 1915, p. 142) computed the  $k = 2, 3$  cases, and then he recorded the result for general  $k$ .

For (3.2), we analogously have

$$\sum_{i \geq 0} i^2 \cdot \text{inv}(y_1, \dots, y_k; i) = \frac{d}{dq} \left( q \frac{d}{dq} \left[ \begin{matrix} y_1 + \dots + y_k \\ y_1, \dots, y_k \end{matrix} \right]_q \right) \Big|_{q=1}.$$

A full proof is presented as an appendix to this paper, so as not to disrupt the flow of the current line of thought.

Notice that for  $k < 4$ , Eq. (3.2) simplifies considerably as some of the multisums are empty. Specifically,

$$\sum_{i \geq 0} i^2 \cdot \text{inv}(y_1, y_2; i) = \begin{pmatrix} y_1 + y_2 \\ y_1, y_2 \end{pmatrix} \left( \frac{1}{12} (y_1 y_2 + y_1^2 y_2 + y_1 y_2^2) + \frac{1}{4} y_1^2 y_2^2 \right) \quad (3.3)$$

$$\begin{aligned} & \sum_{i \geq 0} i^2 \cdot \text{inv}(y_1, y_2, y_3; i) \\ &= \begin{pmatrix} y_1 + y_2 + y_3 \\ y_1, y_2, y_3 \end{pmatrix} \left( \frac{1}{12} \left( \sum_{1 \leq i < j \leq 3} y_i y_j + \sum_{1 \leq i < j \leq 3} y_i^2 y_j + \sum_{1 \leq i < j \leq 3} y_i y_j^2 \right) \right. \\ & \quad \left. + \frac{1}{4} \sum_{1 \leq i < j \leq 3} y_i^2 y_j^2 + \frac{1}{6} y_1 y_2 y_3 + \frac{1}{2} (y_1^2 y_2 y_3 + y_1 y_2^2 y_3 + y_1 y_2 y_3^2) \right), \quad (3.4) \end{aligned}$$

## 4 The marginal distribution of $I$

### 4.1 The probability mass function and a possible limiting distribution

The marginal distribution of  $I$  is given by

$$P(I = i) = \sum_{(y_1, \dots, y_k) \in \mathcal{C}_k} \text{inv}(y_1, \dots, y_k; i) p_1^{y_1} \cdots p_k^{y_k},$$

for  $i = 0, 1, 2, \dots, \lfloor (k-1)n^2/(2k) \rfloor$ , (4.1)

and 0 otherwise.

One referee commented that it was not easy to compute directly with Eq. (4.1), and that it would be nice to have an asymptotic distribution for  $I$ . The author worked at this for a while and encountered difficulties, so he decided to consider an easier but related problem: that of finding an asymptotic distribution for  $I$  in the case where all  $p_j$  are equal, i.e.  $p_1 = p_2 = \dots = p_k = 1/k$ . In this case (4.1) simplifies (slightly) to

$$P(I = i) = \frac{1}{k^n} \sum_{(y_1, \dots, y_k) \in \mathcal{C}_k} \text{inv}(y_1, \dots, y_k; i) \quad \text{for } i = 0, 1, 2, \dots, \lfloor (k-1)n^2/(2k) \rfloor, \quad (4.2)$$

and 0 otherwise; but the problem of counting the inversions, which appear as the coefficients of  $q^i$  in the  $q$ -multinomial coefficient  $\begin{bmatrix} n \\ y_1, \dots, y_k \end{bmatrix}_q$  remains, and is far from trivial. Since the constant term in  $\begin{bmatrix} n \\ y_1, \dots, y_k \end{bmatrix}_q$  is 1 for all  $n$  and  $k$ , we may deduce that

$$P(I = 0) = \frac{1}{k^n} \binom{n+k-1}{n},$$

but for  $i > 0$ ,  $P(I = i)$  seems inaccessible for symbolic  $n$  and  $k$ . Of course, for specific numeric  $n$  and  $k$ , the computation of the full pmf for  $I$  presents no difficulty.

Extensive computations performed using Mathematica, and the resulting graphs for the pmf of  $I$  in the case of equal probabilities for all  $k$  outcomes of each trial strongly suggest that as  $n$  increases, while  $k$  is left fixed, the pmf of  $I$  tends toward a normal distribution. Specifically, it appears that

$$\frac{\sqrt{n} (I - \mu)}{\binom{n}{2} \sigma} \xrightarrow{d} N(0, 1), \quad (4.3)$$

where

$$\mu = \binom{n}{2} \binom{k}{2} \frac{1}{k^2} = \frac{n(n-1)(k-1)}{4k}, \quad (4.4)$$

$$\sigma^2 = \frac{(k-1)(k+1)(n-1)n(2n+5)}{72k^2}. \quad (4.5)$$

Of course, both  $\mu$  and  $\sigma$  tend to  $\infty$  as  $n$  increases without bound, but the preceding can be used to visualize an approximation to the distribution of  $I$  for a large, fixed  $n$ .

The same referee suggested that it may be possible to find a result along the lines of Theorem 5.2 in Charalambides (2021), when  $n$  and  $k$  are proportional.

## 4.2 Moments of $I$

**Theorem 4.1.** *Let  $y_1, y_2, \dots, y_k$  denote fixed nonnegative integers, and let  $n = y_1 + \dots + y_k$ . The first two raw moments of  $I$  are*

$$E(I) = \binom{n}{2} \sum_{1 \leq i < j \leq k} p_i p_j. \quad (4.6)$$

and

$$\begin{aligned} E(I^2) &= \binom{n}{2} \sum_{1 \leq i < j \leq k} p_i p_j + 2 \binom{n}{3} \sum_{1 \leq i < j \leq k} (p_i^2 p_j + p_i p_j^2) + 6 \binom{n}{4} \sum_{1 \leq i < j \leq k} p_i^2 p_j^2 \\ &\quad + 10 \binom{n}{3} \sum_{1 \leq h < i < j \leq k} p_h p_i p_j + 36 \binom{n}{4} \sum_{1 \leq g < h < i < j \leq k} p_g p_h p_i p_j \\ &\quad + 12 \binom{n}{4} \sum_{1 \leq h < i < j \leq k} (p_h^2 p_i p_j + p_h p_i^2 p_j + p_h p_i p_j^2). \end{aligned} \quad (4.7)$$

Thus the variance of  $I$  is given by

$$\begin{aligned} V(I) &= \binom{n}{2} \sum_{1 \leq i < j \leq k} p_i p_j + 2 \binom{n}{3} \sum_{1 \leq i < j \leq k} (p_i^2 p_j + p_i p_j^2) \\ &\quad - \binom{n}{2} (2n-3) \sum_{1 \leq i < j \leq k} p_i^2 p_j^2 + 10 \binom{n}{3} \sum_{1 \leq h < i < j \leq k} p_h p_i p_j \\ &\quad - 6 \binom{n}{2} (2n-3) \sum_{1 \leq g < h < i < j \leq k} p_g p_h p_i p_j \\ &\quad - 2 \binom{n}{2} (2n-3) \sum_{1 \leq h < i < j \leq k} (p_h^2 p_i p_j + p_h p_i^2 p_j + p_h p_i p_j^2). \end{aligned} \quad (4.8)$$

*Proof.*

$$\begin{aligned} E(I) &= \sum_{i \geq 0} i \cdot P(I = i) \\ &= \sum_{i \geq 0} \sum_{\mathbf{y} \in \mathcal{C}_k(n)} i \operatorname{inv}(y_1, \dots, y_k; i) p_1^{y_1} \cdots p_k^{y_k} \\ &= \sum_{\mathbf{y} \in \mathcal{C}_k(n)} p_1^{y_1} \cdots p_k^{y_k} \sum_{i \geq 0} i \operatorname{inv}(y_1, \dots, y_k; i) \\ &= \sum_{\mathbf{y} \in \mathcal{C}_k(n)} \frac{1}{2} p_1^{y_1} \cdots p_k^{y_k} \binom{n}{y_1, \dots, y_k} \sum_{1 \leq i < j \leq k} y_i y_j \\ &= \binom{n}{2} \sum_{1 \leq i < j \leq k} p_i p_j. \end{aligned}$$

□

$$\begin{aligned}
E(I^2) &= \sum_{i \geq 0} i^2 \cdot P(I = i) \\
&= \sum_{i \geq 0} \sum_{\mathbf{y} \in \mathcal{C}_k(n)} i^2 \operatorname{inv}(y_1, \dots, y_k; i) p_1^{y_1} \cdots p_k^{y_k} \\
&= \sum_{\mathbf{y} \in \mathcal{C}_k(n)} p_1^{y_1} \cdots p_k^{y_k} \sum_{i \geq 0} i^2 \operatorname{inv}(y_1, \dots, y_k; i) \\
&= \sum_{\mathbf{y} \in \mathcal{C}_k(n)} p_1^{y_1} \cdots p_k^{y_k} \binom{y_1 + \cdots + y_k}{y_1, \dots, y_k} \left( \frac{1}{12} \sum_{1 \leq i < j \leq k} y_i y_j + \frac{1}{12} \sum_{1 \leq i < j \leq k} y_i^2 y_j + \frac{1}{12} \sum_{1 \leq i < j \leq k} y_i y_j^2 \right. \\
&\quad \left. + \frac{1}{4} \sum_{1 \leq i < j \leq k} y_i^2 y_j^2 + \frac{1}{6} \sum_{1 \leq h < i < j \leq k} y_h y_i y_j + \frac{1}{2} \sum_{1 \leq h < i < j \leq k} y_h^2 y_i y_j \right. \\
&\quad \left. + \frac{1}{2} \sum_{1 \leq h < i < j \leq k} y_h y_i^2 y_j + \frac{1}{2} \sum_{1 \leq h < i < j \leq k} y_h y_i y_j^2 + \frac{3}{2} \sum_{1 \leq g < h < i < j \leq k} y_g y_h y_i y_j \right) \\
&= \binom{n}{2} \sum_{1 \leq i < j \leq k} p_i p_j + 2 \binom{n}{3} \sum_{1 \leq i < j \leq k} (p_i^2 p_j + p_i p_j^2) + 6 \binom{n}{4} \sum_{1 \leq i < j \leq k} p_i^2 p_j^2 \\
&\quad + 10 \binom{n}{3} \sum_{1 \leq h < i < j \leq k} p_h p_i p_j + 36 \binom{n}{4} \sum_{1 \leq g < h < i < j \leq k} p_g p_h p_i p_j \\
&\quad + 12 \binom{n}{4} \sum_{1 \leq h < i < j \leq k} (p_h^2 p_i p_j + p_h p_i^2 p_j + p_h p_i p_j^2).
\end{aligned}$$

Noting that

$$\begin{aligned}
&\left( \sum_{1 \leq i < j \leq k} p_i p_j \right)^2 \\
&= \sum_{1 \leq i < j \leq k} p_i^2 p_j^2 + 2 \sum_{1 \leq h < i < j \leq k} (p_h^2 p_i p_j + p_h p_i^2 p_j + p_h p_i p_j^2) + 6 \sum_{1 \leq g < h < i < j \leq k} p_g p_h p_i p_j,
\end{aligned} \tag{4.9}$$

$$(E(I))^2 = \binom{n}{2}^2 \left( \sum_{1 \leq i < j \leq k} p_i^2 p_j^2 + 2 \sum_{1 \leq h < i < j \leq k} (p_h^2 p_i p_j + p_h p_i^2 p_j + p_h p_i p_j^2) + 6 \sum_{1 \leq g < h < i < j \leq k} p_g p_h p_i p_j \right). \quad (4.10)$$

Thus,

$$\begin{aligned} V(I) &= E(I^2) - (E(I))^2 \\ &= \binom{n}{2} \sum_{1 \leq i < j \leq k} p_i p_j + 2 \binom{n}{3} \sum_{1 \leq i < j \leq k} (p_i^2 p_j + p_i p_j^2) \\ &\quad + \left( 6 \binom{n}{4} - \binom{n}{2}^2 \right) \sum_{1 \leq i < j \leq k} p_i^2 p_j^2 + 10 \binom{n}{3} \sum_{1 \leq h < i < j \leq k} p_h p_i p_j \\ &\quad + \left( 36 \binom{n}{4} - 6 \binom{n}{2}^2 \right) \sum_{1 \leq g < h < i < j \leq k} p_g p_h p_i p_j \\ &\quad + \left( 12 \binom{n}{4} - 2 \binom{n}{2}^2 \right) \sum_{1 \leq h < i < j \leq k} (p_h^2 p_i p_j + p_h p_i^2 p_j + p_h p_i p_j^2) \\ &= \binom{n}{2} \sum_{1 \leq i < j \leq k} p_i p_j + 2 \binom{n}{3} \sum_{1 \leq i < j \leq k} (p_i^2 p_j + p_i p_j^2) \\ &\quad - \binom{n}{2} (2n-3) \sum_{1 \leq i < j \leq k} p_i^2 p_j^2 + 10 \binom{n}{3} \sum_{1 \leq h < i < j \leq k} p_h p_i p_j \\ &\quad - 6 \binom{n}{2} (2n-3) \sum_{1 \leq g < h < i < j \leq k} p_g p_h p_i p_j \\ &\quad - 2 \binom{n}{2} (2n-3) \sum_{1 \leq h < i < j \leq k} (p_h^2 p_i p_j + p_h p_i^2 p_j + p_h p_i p_j^2). \end{aligned}$$

## 5 Conditional distribution of $I$ given $\mathbf{Y}$ .

Given a vector  $\mathbf{y} = (y_1, y_2, \dots, y_k)$ , with  $n = y_1 + y_2 + \dots + y_k$ , the conditional pmf of  $I$  given  $\mathbf{Y} = \mathbf{y}$  is

$$P(I = i \mid \mathbf{Y} = \mathbf{y}) = \frac{\text{inv}(y_1, y_2, \dots, y_k; i)}{\binom{n}{y_1, y_2, \dots, y_k}} \quad (5.1)$$

for  $i = 0, 1, 2, \dots, \sum_{1 \leq i < j \leq k} y_i y_j$ ; and 0 otherwise.

The conditional expectation is

$$\begin{aligned} E(I = i \mid \mathbf{Y} = \mathbf{y}) &= \binom{n}{y_1, \dots, y_k}^{-1} \sum_{i \geq 0} i \cdot \text{inv}(y_1, \dots, y_k; i) \\ &= \frac{1}{2} \sum_{1 \leq i < j \leq k} y_i y_j, \end{aligned} \quad (5.2)$$

by Eq. (3.1).

## 6 Application

Recording the counts of the  $k$  outcomes of a multinomial experiment provides valuable information, but does not provide *all* information we might need to know. For example, suppose that a standard six-sided die is rolled 60 times, and it turns out that exactly 10 of the sixty rolls showed  $j$  dots, for  $j = 1, 2, 3, 4, 5, 6$ . Using that information alone we would strongly suspect that the die was “fair.” However, if we looked further and noted that the sequence of rolls was in fact

$$\{1, 1, \dots, 1, 2, 2, \dots, 2, 3, 3, \dots, 3, 4, 4, \dots, 4, 5, 5, \dots, 5, 6, 6, \dots, 6\} \quad (6.1)$$

then we would certainly think that something unexpected was happening with the die after all. The preceding sequence of die rolls has an inversion number of  $I = 0$ . Of course, the opposite extreme

$$\{6, 6, \dots, 6, 5, 5, \dots, 5, 4, 4, \dots, 4, 3, 3, \dots, 3, 2, 2, \dots, 2, 1, 1, \dots, 1\} \quad (6.2)$$

would be just as suspicious. This latter sequence has an inversion number of  $I = 1500$ , the maximum possible.

In contrast, a sequence of die rolls (still assuming 10 of each of the six possible outcomes), of around 750, the expected value of  $I$  given  $\mathbf{Y} = (10, 10, 10, 10, 10, 10)$ , might provide some assurance that the occurrences of the various outcomes were “well-mixed” as we might expect in a truly fair die.

For a given value  $\mathbf{y}$  of the multinomial random variable  $\mathbf{Y}$ , the possible values of  $I$  are from 0 through  $\sum_{1 \leq i < j \leq k} y_i y_j$ ; it would be handy to adjust the value of  $I$  into a scale ranging from  $-1$  to  $1$ , with  $1$  indicating that all occurrences of outcome 1 occur first, followed by all occurrences of outcome 2, etc.; and with  $-1$  indicating the exact reverse. The simplest way to do this is via the transformation

$$H := 1 - \frac{2}{\sum_{1 \leq i < j \leq k} y_i y_j} I.$$

In this way, the sequence (6.1) corresponds to  $H = 1$  and the sequence (6.2) to  $H = -1$ . Clearly,  $E(H) = 0$ .

## Acknowledgments

The author thanks Doron Zeilberger for suggesting that he consider the major index of a permutation during a discussion related to Sills (2023). This in turn led the author to further consider the inversion statistic of a permutation, which in turn led to the refinement of the multinomial distribution presented in the present manuscript. The author further thanks the anonymous referees who patiently provided helpful suggestions and guidance which improved the paper, including catching a calculation error in an earlier version.

## References

- Andrews GE (1976) *The Theory of Partitions*. Addison–Wesley, Reading, Massachusetts (Reissued (1998) Cambridge University Press, Cambridge UK)
- Charalambides CA (2021)  $q$ -Multinomial and negative  $q$ -multinomial distributions. *Comm. Statist. Theory Methods* 50, 5873–5898
- Farsi C (2007) A non-commutative  $n$ -nomial formula. *Rocky Mountain J. Math.* 37: 1527–1540
- Gasper G, Rahman M (2004) *Basic Hypergeometric Series*, 2nd ed. Cambridge University Press, Cambridge
- Heubach S, Mansour T (2009) *Combinatorics of Compositions and Words*. CRC Press, Boca Raton, Florida
- MacMahon PA (1915) *Combinatory Analysis*, vol. I, Cambridge University Press, Cambridge UK
- MacMahon PA (1917) Two applications of general theorems in combinatory analysis: (1) to the theory of inversions of permutations; (2) to the



ascertainment of the numbers of terms in the development of a determinant which has amongst its elements an arbitrary number of zeros, Proc. London Math. Soc. Ser. 2 15: 314–321

Sills AV (2023) A refinement of the binomial distribution using the quantum binomial theorem. Comm. Statist. Theory Methods 52: 294–308.

## A Proof of Lemma 3.1

### A.1 Some notation and elementary results

In the theory of  $q$ -series, the  $q$ -analog  $[m]_q$  of a nonnegative integer  $m$  is given by

$$[m]_q = 1 + q + q^2 + \cdots + q^{m-1}; \quad (\text{A.1})$$

see Gasper and Rahman (2004, p. 7ff). Notice  $[m]_q \rightarrow m$  when  $q$  is set equal to 1. Note further each of the following consequences:

$$\frac{d}{dq}[m]_q = 1 + 2q + 3q^2 + \cdots + (m-1)q^{m-2}; \quad (\text{A.2})$$

$$\frac{d}{dq} \log[m]_q = \frac{1 + 2q + 3q^2 + \cdots + (m-1)q^{m-2}}{1 + q + q^2 + \cdots + q^{m-1}} \quad (\text{A.3})$$

$$\xrightarrow{q=1} \frac{\binom{m}{2}}{m} = \frac{1}{2}(m-1);$$

$$\frac{d}{dq} \left( q \frac{d}{dq} \log[m]_q \right) = \frac{d}{dq} \frac{q + 2q^2 + 3q^3 + \cdots + (m-1)q^{m-1}}{1 + q + q^2 + \cdots + q^{m-1}} \quad (\text{A.4})$$

$$= \frac{[m]_q \sum_{i=0}^{m-2} (i+1)^2 q^i - \left( \sum_{j=1}^{m-1} j q^j \right) \left( \sum_{r=0}^{m-2} (r+1) q^r \right)}{([m]_q)^2}$$

$$\xrightarrow{q=1} \left( m \frac{(2m-1)m(m-1)}{6} - \frac{m^2(m-1)^2}{4} \right) \frac{1}{m^2}$$

$$= \frac{1}{12}(m^2 - 1).$$

### A.2 Proof of Lemma 3.1

Let  $y_1, y_2, \dots, y_k$  denote fixed nonnegative integers, let  $n = y_1 + y_2 + \cdots + y_k$ , and let

$$f = f(q) = \left[ \begin{matrix} y_1 + \cdots + y_k \\ y_1, \dots, y_k \end{matrix} \right]_q = \prod_{i=1}^n [i]_q \div \prod_{r=1}^k \prod_{i_r=1}^{y_r} [i_r]_q.$$

This latter expression for the  $q$ -multinomial coefficient is equivalent to that which was given in (1.2), but is more convenient for our present purposes.

It follows that

$$\log f = \sum_{i=1}^n \log[i]_q - \sum_{r=1}^k \sum_{i_r=1}^{y_r} \log[i_r]_q \quad (\text{A.5})$$

and thus

$$\frac{d}{dq} \log f = \sum_{i=1}^n \frac{\frac{d}{dq}[i]_q}{[i]_q} - \sum_{r=1}^k \sum_{i_r=1}^{y_r} \frac{\frac{d}{dq}[i_r]_q}{[i_r]_q}. \quad (\text{A.6})$$

Using logarithmic differentiation, we find that

$$\frac{df}{dq} = f \frac{d}{dq} \log f, \quad (\text{A.7})$$

and then applying Leibniz' rule,

$$\frac{d}{dq} \left( q \frac{df}{dq} \right) = f \left( \frac{d}{dq} \left( q \frac{d}{dq} \log f \right) + q \left( \frac{d}{dq} \log f \right)^2 \right). \quad (\text{A.8})$$

We proceed to evaluate the expressions contained in the right members of (A.7) and (A.8) at  $q = 1$ .

$$\begin{aligned} \frac{d}{dq} \log f &= \sum_{i=1}^n \frac{d}{dq} \log[i]_q - \sum_{r=1}^k \sum_{i_r=1}^{y_r} \frac{d}{dq} \log[i_r]_q \\ &\xrightarrow{q=1} \frac{1}{2} \sum_{i=1}^n (i-1) - \frac{1}{2} \sum_{r=1}^k \sum_{i_r=1}^{y_r} (i_r-1) \\ &= \frac{1}{2} \left( \binom{n+1}{2} - n \right) - \frac{1}{2} \left( \sum_{r=1}^k \binom{y_r+1}{2} - y_r \right) \\ &= \frac{1}{4} (n(n+1) - y_1(y_1+1) - y_2(y_2+1) - \cdots - y_k(y_k+1)) \\ &= \frac{1}{2} \sum_{1 \leq i < j \leq k} y_i y_j. \end{aligned}$$

Thus, by (A.7),

$$\sum_{i \geq 0} i \cdot \text{inv}(y_1, \dots, y_k; i) = q \frac{df}{dq} = \frac{1}{2} \binom{n}{y_1, \dots, y_k} \sum_{1 \leq i < j \leq k} y_i y_j, \quad (\text{A.9})$$

which is (3.1) of Lemma 3.1.

Next, we need to compute  $\frac{d}{dq} \left( q \frac{d}{dq} \log f \right)$  at  $q = 1$ .

$$\begin{aligned}
& \frac{d}{dq} \left( q \frac{d}{dq} \log f \right) \\
&= \sum_{i=1}^n \frac{d}{dq} \left( q \frac{d}{dq} \log [i]_q \right) - \sum_{r=1}^k \sum_{i_r=1}^{y_r} \frac{d}{dq} \left( q \frac{d}{dq} \log [i_r]_q \right) \\
&\stackrel{q=1}{\rightarrow} \frac{1}{12} \left\{ \left( \frac{n(n+1)(2n+1)}{6} - n \right) - \sum_{r=1}^k \left( \frac{y_r(y_r+1)(2y_r+1)}{6} - y_r \right) \right\} \\
&= \frac{1}{12} \left\{ \sum_{1 \leq i < j \leq k} y_i y_j + \sum_{1 \leq i < j \leq k} y_i^2 y_j + \sum_{1 \leq i < j \leq k} y_i y_j^2 + 2 \sum_{1 \leq h < i < j \leq k} y_h y_i y_j \right\}.
\end{aligned}$$

So, we have

$$\begin{aligned}
\sum_{i \geq 0} i^2 \cdot \text{inv}(y_1, \dots, y_k; i) &= \frac{d}{dq} \left( q \frac{df}{dq} \right) \\
&= f \left( \frac{d}{dq} \left( q \frac{d}{dq} \log f \right) + q \left( \frac{d}{dq} \log f \right)^2 \right) \\
&= \begin{pmatrix} y_1 + \dots + y_k \\ y_1, \dots, y_k \end{pmatrix} \left( \frac{1}{12} \left\{ \sum_{1 \leq i < j \leq k} y_i y_j + \sum_{1 \leq i < j \leq k} y_i^2 y_j + \sum_{1 \leq i < j \leq k} y_i y_j^2 + \right. \right. \\
&\quad \left. \left. 2 \cdot \sum_{1 \leq h < i < j \leq k} y_h y_i y_j \right\} + \frac{1}{4} \left( \sum_{1 \leq i < j \leq k} y_i y_j \right)^2 \right) \\
&= \begin{pmatrix} y_1 + \dots + y_k \\ y_1, \dots, y_k \end{pmatrix} \left( \frac{1}{12} \sum_{1 \leq i < j \leq k} y_i y_j + \frac{1}{12} \sum_{1 \leq i < j \leq k} y_i^2 y_j + \frac{1}{12} \sum_{1 \leq i < j \leq k} y_i y_j^2 \right. \\
&\quad \left. + \frac{1}{4} \sum_{1 \leq i < j \leq k} y_i^2 y_j^2 + \frac{1}{6} \sum_{1 \leq h < i < j \leq k} y_h y_i y_j + \frac{1}{2} \sum_{1 \leq h < i < j \leq k} y_h^2 y_i y_j \right. \\
&\quad \left. + \frac{1}{2} \sum_{1 \leq h < i < j \leq k} y_h y_i^2 y_j + \frac{1}{2} \sum_{1 \leq h < i < j \leq k} y_h y_i y_j^2 + \frac{3}{2} \sum_{1 \leq g < h < i < j \leq k} y_g y_h y_i y_j \right), \tag{A.10}
\end{aligned}$$

which is (3.2) of Lemma 3.1.  $\square$