

Global well-posedness of the MHD boundary layer equations in the Sobolev Space

Wei-Xi Li, Zhan Xu and Anita Yang

Abstract. We study the two-dimensional MHD boundary layer equations. For small perturbation around a tangential background magnetic field, we obtain the global-in-time existence and uniqueness of solutions in Sobolev spaces. The proof relies on the novel combination of the well-explored cancellation mechanism and the idea of linearly-good unknowns, and we use the former idea to deal with the top tangential derivatives and the latter one admitting fast decay rate to control lower-order derivatives.

CONTENTS

1. Introduction and main result	2
2. <i>A priori</i> estimate	7
2.1. Cancellation mechanism and good unknowns	7
2.2. Methodology and statement of <i>a priori</i> estimate	8
2.3. Inequalities with time-dependent weights	10
3. Preliminary lemmas	12
4. Estimate on the auxiliary functions u_m and f_m	17
5. Estimate on the auxiliary functions \mathcal{U} and \mathcal{F}	22
Appendix A. Equations for \mathcal{U} and \mathcal{F}	27
Appendix B. Equations for u_m and f_m	27
References	31

2020 *Mathematics Subject Classification.* 35Q35, 76W05, 76D10.

Key words and phrases. Global well-posedness, Sobolev spaces, MHD boundary layer equations.

1. Introduction and main result

Magnetohydrodynamic (MHD) is the study of the interaction between electromagnetic fields and conducting fluids, describing the motion of conducting fluid under the influence of the magnetic fields. In this work we consider the two-dimensional (2D) case and assume the fluid moves in the half-space Ω . Then in the incompressible framework, the governing equations read

$$\begin{cases} \partial_t \mathbf{U} + (\mathbf{U} \cdot \nabla) \mathbf{U} - (\mathbf{B} \cdot \nabla) \mathbf{B} + \nabla P - \frac{1}{\text{Re}} \Delta \mathbf{U} = 0, \\ \partial_t \mathbf{B} - \nabla \times (\mathbf{U} \times \mathbf{B}) - \frac{1}{\text{Rm}} \Delta \mathbf{B} = 0, \\ \nabla \cdot \mathbf{U} = \nabla \cdot \mathbf{B} = 0, \\ \mathbf{U}|_{t=0} = \mathbf{U}_{in}, \quad \mathbf{B}|_{t=0} = \mathbf{B}_{in}, \end{cases} \quad (1.1)$$

where \mathbf{U} and \mathbf{B} are the velocity and magnetic fields, respectively, and Re and Rm stand for the hydrodynamic and magnetic Reynolds numbers, respectively. Moreover, suppose system (1.1) is complemented with an uniform background magnetic field and the no-slip boundary condition on the velocity field, saying

$$\mathbf{U}|_{\partial\Omega} = 0, \quad \mathbf{B}|_{\partial\Omega} = \mathbf{e},$$

for some constant vector \mathbf{e} . Formally, as the Reynolds numbers Re and Rm go to $+\infty$ we expect system (1.1) will trend to the ideal MHD system. However, in the case with physical boundaries, the mathematically rigorous justification of this process of high Reynolds limit is more complicated and interesting, due to the mismatched boundary conditions between (1.1) and the limiting ideal system which lead to the appearance of a boundary layer. In the specific case when $\text{Re} \approx \text{Rm}$, following the argument of Prandtl's ansatz, Gérard-Varet and Prestipino [9] derived the following governing equations for the boundary layer:

$$\begin{cases} \partial_t u + (u\partial_x + v\partial_y)u + \partial_x p - \partial_y^2 u = (b_1\partial_x + b_2\partial_y)b_1, \\ \partial_t b_1 + (u\partial_x + v\partial_y)b_1 - \partial_y^2 b_1 = (b_1\partial_x + b_2\partial_y)u, \\ \partial_x u + \partial_y v = 0, \quad \partial_x b_1 + \partial_y b_2 = 0, \end{cases} \quad (1.2)$$

which are posed on the half-space $\mathbb{R}_+^2 = \{(x, y) \in \mathbb{R}^2, y > 0\}$, and complemented with the boundary condition

$$(u, v)|_{y=0} = (0, 0) \quad \text{and} \quad (b_1, b_2)|_{y=0} = \mathbf{e}, \quad \lim_{y \rightarrow +\infty} (u, b_1) = (\alpha(t, x), \beta(t, x)), \quad (1.3)$$

and the initial condition

$$(u, b_1)|_{t=0} = (u_{in}, b_{in}). \quad (1.4)$$

Here (u, v) and (b_1, b_2) are the velocity and the magnetic fields, respectively, and $(\alpha(t, x), \beta(t, x), p(t, x))$ are given functions of (t, x) , standing for the outflow of velocity, magnetic and pressure, which satisfy the Bernoulli's law:

$$\begin{cases} (\partial_t + \alpha \cdot \partial_x)\alpha + \partial_x p - \beta \partial_x \beta = 0, \\ (\partial_t + \alpha \cdot \partial_x)\beta - \beta \partial_x \alpha = 0. \end{cases}$$

Without loss of generality, we will consider a uniform outflow $(\alpha, \beta) = (0, 1)$. Moreover, we consider (b_1, b_2) as a small perturbation around the uniform background magnetic field $e = (1, 0)$. Then using the notation that

$$(b_1, b_2) = (1 + f, g),$$

we may rewrite the initial-boundary problem (1.2)-(1.4) as

$$\begin{cases} \partial_t u + (u\partial_x + v\partial_y)u - \partial_y^2 u = (1 + f)\partial_x f + g\partial_y f, \\ \partial_t f + (u\partial_x + v\partial_y)f - \partial_y^2 f = (1 + f)\partial_x u + g\partial_y u, \\ \partial_x u + \partial_y v = 0, \quad \partial_x f + \partial_y g = 0, \\ (u, v, f, g)|_{y=0} = (0, 0, 0, 0), \quad \lim_{y \rightarrow +\infty} (u, f) = (0, 0), \\ (u, f)|_{t=0} = (u_{in}, f_{in}), \end{cases} \quad (1.5)$$

where $f_{in} = b_{in} - 1$. This paper aims to investigate the global-in-time existence and uniqueness of solutions to the MHD layer system (1.5) in the Sobolev space.

Notations. Before stating the main result, we first list some notations to be used throughout the paper. We will use $\|\cdot\|_{L^2}$ and $(\cdot, \cdot)_{L^2}$ to denote the norm and inner product of $L^2 = L^2(\mathbb{R}_+^2)$ and use the notation $\|\cdot\|_{L_y^2}$ and $(\cdot, \cdot)_{L_y^2}$ when the variable y is specified. Similar notation will be used for L^∞ . Moreover, denote by $L_x^p L_y^q = L^p(\mathbb{R}; L^q(\mathbb{R}_+))$ the classical Lebesgue space, and similarly for the Sobolev space $H_x^p H_y^q$. For any given norm $\|\cdot\|$ we use the convention that

$$\|A\| := \left(\sum_{1 \leq j \leq k} \|A_j\|^2 \right)^{\frac{1}{2}}$$

for vector-valued functions $A = (A_1, A_2, \dots, A_k)$.

We define the weighted Lebesgue space with a given non-negative function ω by setting

$$L_\omega^2 := \left\{ h : \mathbb{R}_+^2 \rightarrow \mathbb{R}; \|h\|_{L_\omega^2} := \left(\int_{\mathbb{R}_+^2} \omega h^2 dx dy \right)^{\frac{1}{2}} < +\infty \right\},$$

which is equipped with the norm $\|\cdot\|_{L_\omega^2}$ and the inner product $(\cdot, \cdot)_{L_\omega^2}$. More generally, define

$$H_\omega^{m,k} := \left\{ h : \mathbb{R}_+^2 \rightarrow \mathbb{R}; \|h\|_{H_\omega^{m,k}} := \left(\sum_{i=0}^m \sum_{j=0}^k \|\partial_x^i \partial_y^j h\|_{L_\omega^2}^2 \right)^{\frac{1}{2}} < +\infty \right\}. \quad (1.6)$$

In this paper we will work with the following time-dependent weight function:

$$\mu := \mu(t, y) = \exp\left(\frac{y^2}{4\langle t \rangle}\right), \quad (1.7)$$

where here and below

$$\langle t \rangle := 1 + t.$$

More generally, we define

$$\mu_\lambda := \mu_\lambda(t, y) = \exp\left(\frac{\lambda y^2}{4\langle t \rangle}\right), \quad 0 \leq \lambda \leq 1. \quad (1.8)$$

The parameter λ in (1.8) is allowed to be negative so that we have

$$\forall \lambda < 0, \forall t \geq 0, \quad \|\mu_\lambda\|_{L_y^2} \leq C_\lambda \langle t \rangle^{\frac{1}{4}} \quad (1.9)$$

with constant C_λ depending only on λ .

We will solve system (1.5) in the weighted Sobolev space $H_\mu^{8,0} \cap H_{\mu_\lambda}^{5,1}$ with any $0 \leq \lambda < 1$. So the initial data u_{in}, f_{in} in (1.5) satisfy the compatibility condition that

$$(u_{in}, f_{in})|_{y=0} = \lim_{y \rightarrow +\infty} (u_{in}, f_{in}) = (0, 0). \quad (1.10)$$

Theorem 1.1. *Assume the initial data in (1.5) satisfy that $u_{in}, f_{in} \in H_{\mu_{in}}^{8,0} \cap H_{\mu_{in}}^{5,1}$ with*

$$\mu_{in} := \mu(0, y) = \exp\left(\frac{y^2}{4}\right).$$

Moreover suppose that the compatibility condition (1.10) is fulfilled and that

$$\forall x \in \mathbb{R}, \quad \int_0^{+\infty} u_{in}(x, y) dy = 0 \text{ and } \int_0^{+\infty} f_{in}(x, y) dy = 0. \quad (1.11)$$

Let $0 < \delta \leq \frac{1}{25}$ be an arbitrarily given number. Then for any $0 \leq \lambda < 1$, there exists a small constant $\varepsilon_0 > 0$ and a constant $C > 0$, both depending on λ and δ , such that if

$$\|(u_{in}, f_{in})\|_{H_{\mu_{in}}^{8,0}} + \|(u_{in}, f_{in})\|_{H_{\mu_{in}}^{5,1}} \leq \varepsilon_0, \quad (1.12)$$

then the MHD boundary layer system (1.5) admits a unique global-in-time solution $u, f \in L^\infty([0, +\infty[; H_\mu^{8,0} \cap H_{\mu_\lambda}^{5,1})$, satisfying that

$$\sup_{t \geq 0} \langle t \rangle^{\frac{1-\delta}{4}} \|(u, f)\|_{H_\mu^{8,0}} + \left(\int_0^{+\infty} \langle t \rangle^{\frac{1-\delta}{2}} \|(\partial_y u, \partial_y f)\|_{H_\mu^{8,0}}^2 dt \right)^{\frac{1}{2}} \leq C\varepsilon_0,$$

and that

$$\begin{aligned} & \sup_{t \geq 0} \sum_{0 \leq k \leq 1} \langle t \rangle^{\frac{5-\delta}{4} + \frac{k}{2}} \|(\partial_y^k u, \partial_y^k f)\|_{H_{\mu_\lambda}^{5,0}} \\ & + \left(\sum_{0 \leq k \leq 1} \int_0^{+\infty} \langle t \rangle^{\frac{5-\delta}{2} + k} \|(\partial_y^{k+1} u, \partial_y^{k+1} f)\|_{H_{\mu_\lambda}^{5,0}}^2 dt \right)^{\frac{1}{2}} \leq C\varepsilon_0. \end{aligned}$$

Recall μ, μ_λ are defined in (1.7) and (1.8), respectively.

Remark 1.2. Without the smallness assumption (1.12), the MHD boundary layer system (1.5) is usually ill-posed in the Sobolev space, seeing [22, Theorem 1.2].

Remark 1.3. As to be seen below, the proof of Theorem 1.1 combines the cancellation mechanism in [20] and the idea of linearly-good unknowns in [27]. Precisely, we will apply the cancellation mechanism for the treatment of top tangential derivatives and meanwhile the linearly-good unknowns idea when dealing with lower-order derivatives. The former technique is used to overcome the loss of derivatives and the advantage of the latter one is that the Sobolev norm of the new good unknowns decays faster than that of the velocity and magnetic fields themselves. More details can be found in Subsection 2.2 below.

We briefly review some related works on the Prandtl and MHD boundary layer equations. In the absence of magnetic fields, the MHD layer system (1.5) is reduced to the classical Prandtl equation which was derived by L. Prandtl in 1904. The mathematical study on the Prandtl boundary layer has a long history, and the well/ill-posedness property is well-explored in various function spaces, seeing e.g. [1, 2, 5–7, 12–19, 31–33] and the references therein. To explore the well-posedness theory of the Prandtl and MHD boundary layer equations, the main difficulty lies in the loss of derivatives which occurs in the nonlocal terms

$$v(t, x, y) = - \int_0^y \partial_x u(t, x, \tilde{y}) d\tilde{y} \quad \text{and} \quad g(t, x, y) = - \int_0^y \partial_x f(t, x, \tilde{y}) d\tilde{y}.$$

Compared with the Prandtl equations, the MHD layer system (1.5) has additional degeneracy property caused by the normal component g of magnetic fields. In order to overcome the loss of derivatives, there are basically two main settings for the well-posedness theory of the Prandtl and MHD boundary layer equations, based on whether or not the structural assumption is imposed. One refers to the analytic or more general Gevrey spaces without any structural assumption on the initial data, and another one refers to function spaces with finite order of regularity under some structural conditions in two space dimensions.

For the 2D Prandtl equation, Oleinik's monotonicity condition plays a crucial role when investigating the well-posedness in spaces with finite order of regularity, seeing e.g. Oleinik-Samokhin's book [26] for the local existence and uniqueness of classical solutions by using the Crocco transformation. Using a new idea of cancellation mechanism, two groups Alexandre-Wang-Xu-Yang [1] and Masmoudi-Wong [25] independently proved the local Sobolev well-posedness through the energy method. Hence, the loss of tangential derivatives can be overcome by using either Crocco transformation or cancellation mechanism under the monotonicity condition. For the MHD boundary layer equations, thanks to the stabilizing effect of magnetic fields, the Sobolev well-posedness in the two-dimensional case is also available even when the velocity field does not satisfy Oleinik's monotonicity assumption; In fact, inspired by the idea of cancellation mechanism initiated by [1, 25], Liu-Xie-Yang [23] established the local Sobolev well-posedness under the hypothesis that the tangential component of magnetic fields does not degenerate.

Without any structural assumption on initial data, motivated by the abstract Cauchy-Kowalewski theorem it is natural to work with the analytic functions space to overcome the loss of one-order derivative. The local well-posedness theory in the analytic functions setting was proved by Sammartino-Caflisch [28]. Furthermore, by virtue of the intrinsic cancellation structure of the Prandtl equations, the analyticity regularity may be relaxed to general Gevrey class. In fact, based on a new understanding of the cancellation mechanism, Dietert and Gérard-Varet [5] and Li-Masmoudi-Yang [18] proved in the 2D and 3D cases, respectively, that the Prandtl equations are indeed local well-posed in Gevrey 2 space. Note that the Gevrey index 2 in [5, 18] is optimal in view of the ill-posedness theory in the work [7] of Gérard-Varet and Dormy. Hence the optimality of the well/ill-posedness theory for the 2D and 3D Prandtl equations without any structural assumption is well justified. However, for the counterpart of the MHD boundary layer system, the Gevrey well-posedness is more complicated because of the additional loss of derivatives in magnetic fields. For this, so far only Gevrey index up to $\frac{3}{2}$ rather than 2 is achieved for the local well-posedness for the 2D and 3D MHD boundary systems(cf. [21]). Compared with the Prandtl counterparts [5, 18], this Gevrey index $\frac{3}{2}$ is expected not to be critical for the well-posedness theory of the MHD boundary layer equations.

The aforementioned works are mainly about the local existence. Much less is known for the global well-posedness theory. Here we mention Xin-Zhang's work [31] on the global existence of weak solutions to the 2D Prandtl equations under an additional favorable pressure condition. On the other hand, without the monotonicity condition, boundary layer separation is well expected (see e.g. [4, 29] for the justification of the Goldstein singularity) and there are lots of works on the instability phenomena. Interested readers may refer to the works of E-Engquist [6] about the construction of blowup solutions for large data (see also Collot-Ghoul-Masmoudi [3] for the formulation of singularity), Grenier [10] on the unstable Euler shear flow that yields instability of the Prandtl equations, and [7, 8, 11, 12] about the instability around shear flows.

However, for small and analytic data we may expect the global solutions to the Prandtl and MHD boundary layer equations even when the structural assumptions are violated. In fact, inspired by the nonlinearly-good unknown idea in Masmoudi-Wong' work [25], Ignatova-Vicol [13] introduced a new linearly-good unknown with decay rate faster than that of the velocity itself and then take advantage of the better decay rate to construct the almost global analytic solutions for the 2D Prandtl equations, after the earlier work [33] of Zhang-Zhang on the long time behavior. This idea of linearly-good unknown is further developed by Paicu-Zhang [27], where the global analytic solution is established by introducing another new linearly-good unknown with the analytic norm decaying fast enough. We refer to Wang-Wang-Zhang [30] for the further generalization of [27] to the Gevrey setting, and [17, 24] for the application of the linearly-good unknown idea to the global well-posedness of the MHD boundary layer equations in the analytic setting. Note the works mentioned above are limited to the analytic or Gevrey setting and the global existence in Sobolev spaces remains open.

The purpose of this work is to study the global well-posedness of the MHD boundary layer system (1.5) in the Sobolev setting.

The organization of the paper is as follow. In Section 2 we state the key *a priori* estimate for proving Theorem 1.1 and discuss the idea in the proof of the *a priori* estimate. Sections 3-5 are devoted to proving the *a priori* estimate. In Appendices A and B we present the derivation of the evolution equations obeyed by the auxiliary functions introduced in this text.

2. *A priori* estimate

The key part for proving Theorem 1.1 is to derive *a priori* energy estimate for system (1.5). Then the global-in-time existence and uniqueness will follow from a standard regularization procedure. Hence, we only present in detail the proof of the *a priori* estimate stated in Subsection 2.2 and omit the regularization procedure for brevity.

2.1. Cancellation mechanism and good unknowns

Because of the loss of tangential derivatives in nonlocal terms, the Prandtl and MHD boundary layer equations are usually ill-posed in the Sobolev setting (cf. [7, 22] for instance). To overcome the degeneracy caused by the absence of the horizontal diffusion, two groups Alexandre-Wang-Xu-Yang [1] and Masmoudi-Wong [25] introduced independently the following nonlinearly-good unknown for the Prandtl equations:

$$\Phi := \partial_y u - \frac{\partial_y^2 u}{\partial_y u} = (\partial_y u) \partial_y \left(\frac{u}{\partial_y u} \right), \quad (2.1)$$

provided Oleinik's monotonicity condition is fulfilled (i.e., $\partial_y u \neq 0$). Note the quantity Φ above behaves better than u and $\partial_y u$, and in fact Φ satisfies an evolution equation where the terms involving the loss of tangential derivatives are cancelled out. This advantage, with the help of Hardy's inequality, enables to conclude the local-in-time well-posedness of the 2D Prandtl equation in Sobolev spaces (cf. [1, 25] for more details). Inspired by this cancellation idea, under the hypothesis that the tangential component of magnetic field does not degenerate, Liu-Xie-Yang [23] proved the local well-posedness for the 2D MHD boundary layer system, basing on the observation of a new cancellation mechanism. This justifies the stabilizing effect of magnetic fields.

The idea in [25] was further developed by Ignatova-Vicol [13], where the authors introduced a new linearly-good unknown of the following form:

$$\Psi := \partial_y u - \frac{\partial_y^2 \phi(t, y)}{\partial_y \phi(t, y)} u = (\partial_y \phi) \partial_y \left(\frac{u}{\partial_y \phi} \right), \quad (2.2)$$

with $\phi(t, y)$ well chosen function such that $-\frac{\partial_y^2 \phi(t, y)}{\partial_y \phi(t, y)}$ is the heat self-similar variable $\frac{y}{2\langle t \rangle}$ (e.g. $\partial_y \phi = \mu$ with μ defined in (1.7)). Note that Oleinik's monotonicity condition does not hold anymore in [13], so that in order to overcome the one-order loss of tangential derivative, the strong tangential analyticity is required for initial data. By a

new observation that the analytic norm of the good unknown Ψ in (2.2) decays faster than that of u itself, Ignatova-Vicol [13] established the almost global-in-time solution to the 2D Prandtl equation in the space of analytic functions. Motivated by Ψ in (2.2), Paicu-Zhang [27] worked with another new linearly-good unknown:

$$G = u + \frac{y}{2\langle t \rangle} \varphi \quad \text{with} \quad \partial_y \varphi = u. \quad (2.3)$$

Compared with the quantity Ψ in (2.2), the analytical norm of $\partial_y G$ decays almost like $\langle t \rangle^{-\frac{7}{4}}$, faster than the decay rate of the analytical norm of Ψ which is almost like $\langle t \rangle^{-\frac{5}{4}}$. Basing on this new decay rate, Paicu-Zhang [27] established the global-in-time existence of solutions to the 2D Prandtl equation with small analytic data.

2.2. Methodology and statement of *a priori* estimate

This work is strongly inspired by the idea of cancellation mechanism and good unknowns recalled in the previous part. On one side, the cancellation mechanism works well for the Sobolev well-posedness, but only the local-in-time existence may be expected. On the other side, one may get the global-in-time well-posedness using the idea of linearly-good unknowns, but ask strong analytic or Gevrey regularity for initial data. Rather than the sole use of the cancellation mechanism or the linearly-good unknown idea in the aforementioned works, we will combine the two techniques in this work.

To investigate the Sobolev well-posedness of system (1.5), the main difficulty arises from the loss of tangential derivatives in the terms v and g . To overcome this degeneracy, we adopt the cancellation mechanism observed in [20], which is inspired by the earlier work [23]. Precisely, define

$$u_m := \begin{cases} \partial_x^m u + \frac{\partial_y u}{1+f} \partial_x^{m-1} g, & \text{if } m \geq 1, \\ u, & \text{if } m = 0, \end{cases} \quad (2.4)$$

and

$$f_m := \begin{cases} \partial_x^m f + \frac{\partial_y f}{1+f} \partial_x^{m-1} g, & \text{if } m \geq 1, \\ f, & \text{if } m = 0. \end{cases} \quad (2.5)$$

Similar to the nonlinearly-good unknown Φ in (2.1), in the evolution equations of u_m and f_m above, the terms involving the loss of derivatives are cancelled out, that is (see Appendix B for more detail),

$$\begin{cases} \left(\partial_t + u \partial_x + v \partial_y - \partial_y^2 \right) u_m - ((1+f) \partial_x + g \partial_y) f_m = l.o.t., \\ \left(\partial_t + u \partial_x + v \partial_y - \partial_y^2 \right) f_m - ((1+f) \partial_x + g \partial_y) u_m = l.o.t., \end{cases} \quad (2.6)$$

where *l.o.t.* refers to lower-order terms. This enables to perform energy estimates in the Sobolev setting for the above system. On the other hand, the decay rate of the Sobolev norms of u_m and f_m is not fast enough to ensure the global-in-time existence. For this

we introduce the good unknown in (2.3) and define two linearly-good unknowns \mathcal{U} and \mathcal{F} by setting

$$\mathcal{U}(t, x, y) := u(t, x, y) + \frac{y}{2\langle t \rangle} \int_0^y u(t, x, \tilde{y}) d\tilde{y}, \quad \mathcal{F} := f + \frac{y}{2\langle t \rangle} \int_0^y f d\tilde{y}. \quad (2.7)$$

Applying ∂_y to the equations above yields that

$$\int_0^y u d\tilde{y} = 2\langle t \rangle (\partial_y \mathcal{U} - \partial_y u) - yu, \quad \int_0^y f d\tilde{y} = 2\langle t \rangle (\partial_y \mathcal{F} - \partial_y f) - yf.$$

Then if the terms on the right-hand side trend to 0 as y going to $+\infty$, we have

$$\int_0^{+\infty} u dy = 0 \quad \text{and} \quad \int_0^{+\infty} f dy = 0. \quad (2.8)$$

Thus we may rewrite \mathcal{U} and \mathcal{F} in (2.7) as

$$\mathcal{U} = u - \frac{y}{2\langle t \rangle} \int_y^{+\infty} u d\tilde{y}, \quad \mathcal{F} = f - \frac{y}{2\langle t \rangle} \int_y^{+\infty} f d\tilde{y}. \quad (2.9)$$

Different from system (2.6), there is no cancellation mechanism in the evolution equations solved by the top tangential derivatives of \mathcal{U} and \mathcal{F} , so that the loss of tangential derivatives may occur. However, we can perform Sobolev estimates for the lower order derivatives of \mathcal{U} and \mathcal{F} . When dealing with the quadratic nonlinear terms, we will control top tangential derivatives of u and f in terms of u_m and f_m (see Lemma 3.3 below), and the lower order derivatives of u and f by the $H_\mu^{5,1}$ -norms of \mathcal{U} and \mathcal{F} (see Lemma 3.1 below). Roughly speaking, u_m and f_m behave as the heat equation so that the decay rate in the L_μ^2 space is almost like $-\frac{1}{4}$ order with respect to time (see Lemma 3.4 below). On the other hand, \mathcal{U} and \mathcal{F} behave as a heat equation with an additional damping term $\frac{1}{\langle t \rangle}$, which yields that the decay rate of the L_μ^2 of \mathcal{U} and \mathcal{F} is almost like $-\frac{5}{4}$ order (see Remark 3.5 below). This motivates us to define the energy as follows.

Definition 2.1. Let δ be an arbitrarily fixed number, satisfying that

$$0 < \delta \leq \frac{1}{25}.$$

We define the energy $\mathcal{E}_\delta = \mathcal{E}_\delta(t)$ by

$$\mathcal{E}_\delta := \sum_{m \leq 8} \langle t \rangle^{\frac{1-\delta}{2}} \|(u_m, f_m)\|_{L_\mu^2}^2 + \sum_{0 \leq k \leq 1} \left(\frac{\delta}{2}\right)^k \langle t \rangle^{\frac{5-\delta}{2}+k} \|(\partial_y^k \mathcal{U}, \partial_y^k \mathcal{F})\|_{H_\mu^{5,0}}^2. \quad (2.10)$$

Accordingly, define $\mathcal{D}_\delta = \mathcal{D}_\delta(t)$ by

$$\begin{aligned} \mathcal{D}_\delta := & \sum_{m \leq 8} \delta \langle t \rangle^{\frac{1-\delta}{2}} \|(\partial_y u_m, \partial_y f_m)\|_{L_\mu^2}^2 \\ & + \sum_{0 \leq k \leq 1} \left(\frac{\delta}{2}\right)^{k+1} \langle t \rangle^{\frac{5-\delta}{2}+k} \|(\partial_y^{k+1} \mathcal{U}, \partial_y^{k+1} \mathcal{F})\|_{H_\mu^{5,0}}^2. \end{aligned} \quad (2.11)$$

Recall u_m, f_m and \mathcal{U}, \mathcal{F} are defined in (2.4), (2.5) and (2.7), and the weighted Sobolev spaces are defined in (1.6).

Theorem 2.2 (A priori estimate). *Let the initial data u_{in} and f_{in} in (1.5) satisfy assumption (1.11) and the compatibility condition (1.10). Then there exists a small constant $\varepsilon > 0$ such that if*

$$\mathcal{E}_\delta(0) \leq \varepsilon^2 \quad (2.12)$$

and if $u, f \in L^\infty([0, +\infty[; H_\mu^{8,0} \cap H_{\mu_\lambda}^{5,1})$, $0 \leq \lambda < 1$, is any global solution to system (1.5) satisfying that

$$\forall t > 0, \quad \mathcal{E}_\delta(t) + \frac{1}{2} \int_0^t \mathcal{D}_\delta(s) ds \leq 2\varepsilon^2,$$

then

$$\forall t > 0, \quad \mathcal{E}_\delta(t) + \frac{1}{2} \int_0^t \mathcal{D}_\delta(s) ds \leq \varepsilon^2. \quad (2.13)$$

The proof of Theorem 2.2 is postponed to Sections 3-5. By bootstrap principle, if $u, f \in L^\infty([0, +\infty[; H_\mu^{8,0} \cap H_{\mu_\lambda}^{5,1})$ solve the MHD boundary layer system (1.5), then assertion (2.13) holds true, provided the smallness assumption (2.12) is fulfilled.

Remark 2.3. As shown by Lemma 3.1 in the next section, the weighted Sobolev norm of lower order derivatives of (u, f) can be controlled by that of $(\mathcal{U}, \mathcal{F})$, that is, for any $0 \leq \lambda < 1$ it holds that

$$\sum_{0 \leq k \leq 1} \langle t \rangle^{\frac{s-\delta}{2}+k} \|(\partial_y^k u, \partial_y^k f)\|_{H_{\mu_\lambda}^{5,0}}^2 \leq C \sum_{0 \leq k \leq 1} \langle t \rangle^{\frac{s-\delta}{2}+k} \|(\partial_y^k \mathcal{U}, \partial_y^k \mathcal{F})\|_{H_\mu^{5,0}}^2 \leq C \mathcal{E}_\delta,$$

and similarly,

$$\sum_{0 \leq k \leq 1} \langle t \rangle^{\frac{s-\delta}{2}+k} \|(\partial_y^{k+1} u, \partial_y^{k+1} f)\|_{H_{\mu_\lambda}^{5,0}}^2 \leq C \mathcal{D}_\delta,$$

where the constant C depends on δ and λ . Moreover, if \mathcal{E}_δ is small enough, then it follows from Lemma 3.3 below that

$$\langle t \rangle^{\frac{1-\delta}{2}} \|(u, f)\|_{H_\mu^{8,0}}^2 \leq C \mathcal{E}_\delta \quad \text{and} \quad \langle t \rangle^{\frac{1-\delta}{2}} \|(\partial_y u, \partial_y f)\|_{H_\mu^{8,0}}^2 \leq C \mathcal{D}_\delta.$$

So the $H_\mu^{8,0} \cap H_{\mu_\lambda}^{5,1}$ -norm of (u, f) can be controlled by the energy \mathcal{E}_δ . On the other hand, if $\|f_{in}\|_{L^\infty} \leq \frac{1}{2}$, then we can verify directly that, by Lemma 2.5 below,

$$\mathcal{E}_\delta(0) \leq C \|(u_{in}, f_{in})\|_{H_{\mu_{in}}^{8,0}}^2 \left(1 + \|(u_{in}, f_{in})\|_{H_{\mu_{in}}^{5,1}}^2\right) + C \|(u_{in}, f_{in})\|_{H_{\mu_{in}}^{5,1}}^2.$$

So assumption (1.12) yields the smallness condition (2.12).

2.3. Inequalities with time-dependent weights

We list in this part some weighted inequalities to be used frequently in the proof of Theorem 2.2.

Lemma 2.4 (cf. Lemma 2.5 in [30]). *Let μ_λ be defined in (1.8), and let h be a function belonging to H^1 in y variable, which decays to zero sufficiently fast as $y \rightarrow +\infty$. Then for $0 \leq \lambda \leq 1$ we have*

$$\frac{\lambda^{\frac{1}{2}}}{(2\langle t \rangle)^{\frac{1}{2}}} \|h\|_{L_{\mu_\lambda}^2} \leq \|\partial_y h\|_{L_{\mu_\lambda}^2}, \quad (2.14)$$

and

$$\frac{\lambda^{\frac{1}{2}}}{2\langle t \rangle^{\frac{1}{2}}} \|h\|_{L^2_{\mu_\lambda}} + \frac{\lambda}{4} \left\| \frac{y}{\langle t \rangle} h \right\|_{L^2_{\mu_\lambda}} \leq 2 \|\partial_y h\|_{L^2_{\mu_\lambda}}. \quad (2.15)$$

Proof. For the sake of completeness we re-present the proof given in [30, Lemma 2.5]. It suffices to consider the case of $0 < \lambda \leq 1$. Using integration by parts yields

$$\begin{aligned} \int_0^{+\infty} h^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy &= - \int_0^{+\infty} y \partial_y \left(h^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} \right) dy \\ &= -2 \int_0^{+\infty} y h (\partial_y h) e^{\frac{\lambda y^2}{4\langle t \rangle}} dy - \frac{\lambda}{2\langle t \rangle} \int_0^{+\infty} y^2 h^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy. \end{aligned} \quad (2.16)$$

This, with the fact that

$$-2 \int_0^{+\infty} y h (\partial_y h) e^{\frac{\lambda y^2}{4\langle t \rangle}} dy \leq \frac{\lambda}{2\langle t \rangle} \int_0^{+\infty} y^2 h^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy + \frac{2\langle t \rangle}{\lambda} \int_0^{+\infty} (\partial_y h)^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy,$$

yields

$$\int_0^{+\infty} h^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy \leq \frac{2\langle t \rangle}{\lambda} \int_0^{+\infty} (\partial_y h)^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy.$$

Multiplying the above estimate by $\frac{\lambda}{2\langle t \rangle}$, we obtain assertion (2.14) in Lemma 2.4.

On the other hand, combining (2.16) with the fact that

$$-2 \int_0^{+\infty} y h (\partial_y h) e^{\frac{\lambda y^2}{4\langle t \rangle}} dy \leq \frac{\lambda}{4\langle t \rangle} \int_0^{+\infty} y^2 h^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy + \frac{4\langle t \rangle}{\lambda} \int_0^{+\infty} (\partial_y h)^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy,$$

yields

$$\int_0^{+\infty} h^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy + \frac{\lambda}{4\langle t \rangle} \int_0^{+\infty} y^2 h^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy \leq \frac{4\langle t \rangle}{\lambda} \int_0^{+\infty} (\partial_y h)^2 e^{\frac{\lambda y^2}{4\langle t \rangle}} dy.$$

Multiplying the above estimate by $\frac{\lambda}{4\langle t \rangle}$, we obtain the second assertion (2.15). The proof of Lemma 2.4 is thus completed. \square

Lemma 2.5. *Let μ_λ be defined in (1.8) and let h be a function belonging to H^1 in y variable, which decays to zero sufficiently fast as $y \rightarrow +\infty$. Then*

$$\forall 0 \leq \lambda < 1, \quad \|\mu_{\frac{\lambda}{2}} h\|_{L^\infty_y} \leq C_\lambda \langle t \rangle^{\frac{1}{4}} \|\mu_{\frac{\lambda+1}{4}} \partial_y h\|_{L^2_y},$$

where the constant C_λ depends only on λ .

Proof. It just follows from straightforward verification. In fact, observe μ_λ is an increasing function of y . Then for any $0 \leq \lambda < 1$ it holds that

$$\begin{aligned} \|\mu_{\frac{\lambda}{2}} h\|_{L^\infty_y} &= \left| \mu_{\frac{\lambda}{2}} \int_y^{+\infty} \partial_y h d\tilde{y} \right| \leq \left| \int_y^{+\infty} \mu_{\frac{\lambda+1}{4}}(\tilde{y}) \mu_{\frac{\lambda-1}{4}}(\tilde{y}) \partial_y h d\tilde{y} \right| \\ &\leq \|\mu_{\frac{\lambda-1}{4}}\|_{L^2_y} \|\mu_{\frac{\lambda+1}{4}} \partial_y h\|_{L^2_y} \leq C_\lambda \langle t \rangle^{\frac{1}{4}} \|\mu_{\frac{\lambda+1}{4}} \partial_y h\|_{L^2_y}, \end{aligned}$$

the last inequality using (1.9). This completes the proof of Lemma 2.5. \square

3. Preliminary lemmas

Before proving the *a priori* estimate, we first list some lemmas to be used frequently in the proof. We begin with the treatment of $\partial_x^m u$ and $\partial_x^m f$. Although these terms are not involved in the definition of \mathcal{E}_δ and \mathcal{D}_δ when $m \geq 1$, as to be seen in Lemmas 3.1 and 3.3 below, they can be controlled by the auxiliary functions u_m and f_m , and furthermore by $\partial_x^m \mathcal{U}$ and $\partial_x^m \mathcal{F}$ if $m \leq 5$.

To simplify the notations, throughout the proof we will use C_λ to denote some generic constants depending only on the number λ in (1.8). Moreover we use C to denote some generic constants which may vary from line to line and depend only on the Sobolev embedding constants and λ , but are independent of δ and ε .

Lemma 3.1. *Let μ_λ be defined in (1.8). For any $0 \leq \lambda < 1$ it holds that*

$$\|\partial_y^k u\|_{H_{\mu_\lambda}^{5,0}} \leq C_\lambda \|\partial_y^k \mathcal{U}\|_{H_{\mu_\lambda}^{5,0}}, \quad \|\partial_y^k f\|_{H_{\mu_\lambda}^{5,0}} \leq C_\lambda \|\partial_y^k \mathcal{F}\|_{H_{\mu_\lambda}^{5,0}}, \quad k = 0, 1, 2. \quad (3.1)$$

Proof. We proceed to prove (3.1) for $k = 0, 1$ and 2. It suffices to derive the estimate on u , since the treatment of f is similar.

Step 1. (Estimate on u). By definition (2.9) of \mathcal{U} and (2.8), we have

$$\begin{cases} \left(\partial_y + \frac{y}{2\langle t \rangle} \right) \int_y^{+\infty} u d\tilde{y} = -u + \frac{y}{2\langle t \rangle} \int_y^{+\infty} u d\tilde{y} = -\mathcal{U}, \\ \int_0^{+\infty} u d\tilde{y} = 0. \end{cases} \quad (3.2)$$

Solving this ODE system (3.2), we get

$$\int_y^{+\infty} u d\tilde{y} = -\exp\left(-\frac{y^2}{4\langle t \rangle}\right) \int_0^y \exp\left(\frac{\tilde{y}^2}{4\langle t \rangle}\right) \mathcal{U} d\tilde{y}.$$

Then we have, for any $0 \leq m \leq 5$ and any $0 \leq \lambda < 1$,

$$\begin{aligned} & \exp\left(\frac{\lambda y^2}{8\langle t \rangle}\right) \int_y^{+\infty} \partial_x^m u d\tilde{y} \\ &= -\exp\left(\frac{(\lambda-1)y^2}{8\langle t \rangle}\right) \int_0^y \exp\left(\frac{\tilde{y}^2 - y^2}{8\langle t \rangle}\right) \exp\left(\frac{\tilde{y}^2}{8\langle t \rangle}\right) \partial_x^m \mathcal{U} d\tilde{y}. \end{aligned} \quad (3.3)$$

On the other hand, observing

$$\forall \xi > 0, \quad e^{-\xi^2} \int_0^\xi e^{z^2} dz \leq 2,$$

we have

$$\int_0^y \exp\left(\frac{\tilde{y}^2 - y^2}{4\langle t \rangle}\right) d\tilde{y} = 2\langle t \rangle^{\frac{1}{2}} \exp\left(-\frac{y^2}{4\langle t \rangle}\right) \int_0^{\frac{y}{2\sqrt{\langle t \rangle}}} e^{z^2} dz \leq 4\langle t \rangle^{\frac{1}{2}}.$$

As a result, in view of (3.3),

$$\begin{aligned} \mu_\lambda(t, y) \left| \int_y^{+\infty} \partial_x^m u d\tilde{y} \right|^2 &= \left| \exp\left(\frac{\lambda y^2}{8\langle t \rangle}\right) \int_y^{+\infty} \partial_x^m u d\tilde{y} \right|^2 \\ &\leq \exp\left(\frac{(\lambda-1)y^2}{4\langle t \rangle}\right) \left[\int_0^y \exp\left(\frac{\tilde{y}^2 - y^2}{4\langle t \rangle}\right) d\tilde{y} \right] \|\partial_x^m \mathcal{U}\|_{L_\mu^2}^2 \\ &\leq 4\langle t \rangle^{\frac{1}{2}} \exp\left(\frac{(\lambda-1)y^2}{4\langle t \rangle}\right) \|\partial_x^m \mathcal{U}\|_{L_\mu^2}^2, \end{aligned}$$

which yields that, for any $0 \leq m \leq 5$,

$$\begin{aligned} \left\| \frac{y}{2\langle t \rangle} \int_y^{+\infty} \partial_x^m u d\tilde{y} \right\|_{L_{\mu_\lambda}^2}^2 &\leq \langle t \rangle^{\frac{1}{2}} \int_0^{+\infty} \frac{y^2}{\langle t \rangle^2} \exp\left(\frac{(\lambda-1)y^2}{4\langle t \rangle}\right) dy \|\partial_x^m \mathcal{U}\|_{L_\mu^2}^2 \leq C_\lambda \|\partial_x^m \mathcal{U}\|_{L_\mu^2}^2, \end{aligned}$$

where in the last inequality we used the fact that

$$\forall 0 \leq \lambda < 1, \quad \langle t \rangle^{\frac{1}{2}} \int_0^{+\infty} \frac{y^2}{\langle t \rangle^2} \exp\left(\frac{(\lambda-1)y^2}{4\langle t \rangle}\right) dy = \int_0^{+\infty} y^2 e^{\frac{(\lambda-1)}{4}y^2} dy \leq C_\lambda.$$

Thus, using (2.9) yields

$$\forall 0 \leq \lambda < 1, \quad \|u\|_{H_{\mu_\lambda}^{5,0}} \leq \left\| \frac{y}{2\langle t \rangle} \int_y^{+\infty} u d\tilde{y} \right\|_{H_{\mu_\lambda}^{5,0}} + \|\mathcal{U}\|_{H_{\mu_\lambda}^{5,0}} \leq C_\lambda \|\mathcal{U}\|_{H_\mu^{5,0}}. \quad (3.4)$$

Step 2). (Estimate on $\partial_y u$). We apply ∂_y to the representation of \mathcal{U} in (2.9) to get

$$\partial_y u = \partial_y \mathcal{U} + \frac{1}{2\langle t \rangle} \int_y^{+\infty} u d\tilde{y} - \frac{y}{2\langle t \rangle} u.$$

This yields

$$\|\partial_y u\|_{H_{\mu_\lambda}^{5,0}} \leq \|\partial_y \mathcal{U}\|_{H_{\mu_\lambda}^{5,0}} + \frac{1}{2\langle t \rangle} \left\| \int_y^{+\infty} u d\tilde{y} \right\|_{H_{\mu_\lambda}^{5,0}} + \frac{1}{2\langle t \rangle} \|yu\|_{H_{\mu_\lambda}^{5,0}}. \quad (3.5)$$

For the second term on the right-hand side, we use Lemma 2.4 and (3.4) to get

$$\frac{1}{2\langle t \rangle} \left\| \int_y^{+\infty} u d\tilde{y} \right\|_{H_{\mu_\lambda}^{5,0}} \leq \frac{C_\lambda}{\langle t \rangle^{\frac{1}{2}}} \|u\|_{H_{\mu_\lambda}^{5,0}} \leq \frac{C_\lambda}{\langle t \rangle^{\frac{1}{2}}} \|\mathcal{U}\|_{H_\mu^{5,0}}^2 \leq C_\lambda \|\partial_y \mathcal{U}\|_{H_\mu^{5,0}}^2.$$

Moreover, for any $0 \leq \lambda < 1$, we use the fact that $\|y^2 e^{-ay^2}\|_{L^\infty} \leq C_a$ for any constant $a > 0$, to compute

$$\frac{1}{2\langle t \rangle} \|yu\|_{H_{\mu_\lambda}^{5,0}} \leq \frac{C_\lambda}{\langle t \rangle^{\frac{1}{2}}} \|u\|_{H_{\mu_\lambda}^{5,0}} \leq \frac{C_\lambda}{\langle t \rangle^{\frac{1}{2}}} \|\mathcal{U}\|_{H_\mu^{5,0}} \leq C_\lambda \|\partial_y \mathcal{U}\|_{H_\mu^{5,0}}, \quad (3.6)$$

the second inequality using (3.4) and the last inequality following from Lemma 2.4.

Now we combine the above estimates with (3.5) to conclude that

$$\forall 0 \leq \lambda < 1, \quad \|\partial_y u\|_{H_{\mu_\lambda}^{5,0}} \leq C_\lambda \|\partial_y \mathcal{U}\|_{H_\mu^{5,0}}. \quad (3.7)$$

Step 3). (Estimate on $\partial_y^2 u$). We apply ∂_y^2 to the representation of \mathcal{U} in (2.9) to get

$$\partial_y^2 u = \partial_y^2 \mathcal{U} - \frac{1}{\langle t \rangle} u - \frac{y}{2 \langle t \rangle} \partial_y u.$$

Using Lemma 2.4 and (3.4) again yields that, for any $0 \leq \lambda < 1$,

$$\frac{1}{\langle t \rangle} \|u\|_{H_{\mu,\lambda}^{5,0}} \leq \frac{C_\lambda}{\langle t \rangle} \|\mathcal{U}\|_{H_\mu^{5,0}} \leq \frac{C_\lambda}{\langle t \rangle^{\frac{1}{2}}} \|\partial_y \mathcal{U}\|_{H_\mu^{5,0}} \leq C_\lambda \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}}.$$

Moreover, repeating the argument in (3.6) and then using (3.7) we have

$$\frac{1}{2 \langle t \rangle} \|y \partial_y u\|_{H_{\mu,\lambda}^{5,0}} \leq \frac{C_\lambda}{\langle t \rangle^{\frac{1}{2}}} \|\partial_y u\|_{H_{\mu,\frac{\lambda+1}{2}}^{5,0}} \leq \frac{C_\lambda}{\langle t \rangle^{\frac{1}{2}}} \|\partial_y \mathcal{U}\|_{H_\mu^{5,0}} \leq C_\lambda \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}},$$

the last inequality following from Lemma 2.4. As a result, we combine the above estimates to conclude that, for any $0 \leq \lambda < 1$,

$$\|\partial_y^2 u\|_{H_{\mu,\lambda}^{5,0}} \leq \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}} + \frac{1}{\langle t \rangle} \|u\|_{H_{\mu,\lambda}^{5,0}} + \frac{1}{2 \langle t \rangle} \|y \partial_y u\|_{H_{\mu,\lambda}^{5,0}} \leq C_\lambda \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}}.$$

This together with (3.4) and (3.7) yields the desired estimate on u in (3.1). The proof of Lemma 3.1 is thus completed. \square

As an immediate consequence of Lemma 3.1 and definitions (2.10) and (2.11) of \mathcal{E}_δ and \mathcal{D}_δ , we have the following

Corollary 3.2. *For any $0 \leq \lambda < 1$ and any $t \geq 0$ it holds that*

$$\sum_{0 \leq k \leq 1} \left(\frac{\delta}{2}\right)^{\frac{k}{2}} \langle t \rangle^{\frac{5-\delta}{4} + \frac{k}{2}} \|(\partial_y^k u, \partial_y^k f)\|_{H_{\mu,\lambda}^{5,0}} \leq C_\lambda (\mathcal{E}_\delta(t))^{\frac{1}{2}},$$

and

$$\sum_{0 \leq k \leq 1} \left(\frac{\delta}{2}\right)^{\frac{k+1}{2}} \langle t \rangle^{\frac{5-\delta}{4} + \frac{k}{2}} \|(\partial_y^{k+1} u, \partial_y^{k+1} f)\|_{H_{\mu,\lambda}^{5,0}} \leq C_\lambda (\mathcal{D}_\delta(t))^{\frac{1}{2}}.$$

Thus if

$$\sup_{t \geq 0} \mathcal{E}_\delta(t) \leq 2\varepsilon^2,$$

then

$$\forall 0 \leq \lambda < 1, \forall t \geq 0, \quad \sum_{0 \leq k \leq 1} \left(\frac{\delta}{2}\right)^{\frac{k}{2}} \langle t \rangle^{\frac{5-\delta}{4} + \frac{k}{2}} \|(\partial_y^k u, \partial_y^k f)\|_{H_{\mu,\lambda}^{5,0}} \leq \varepsilon C_\lambda. \quad (3.8)$$

The following lemma is devoted to controlling $(\partial_x^m u, \partial_x^m f)$ in terms of (u_m, f_m) under the smallness assumption.

Lemma 3.3. *There exists a small constant $\varepsilon > 0$ such that if*

$$\sup_{t \geq 0} \mathcal{E}_\delta(t) \leq 2\varepsilon^2,$$

then for each $1 \leq m \leq 8$ it holds that

$$\langle t \rangle^{\frac{1-\delta}{4}} \|(\partial_x^m u, \partial_x^m f)\|_{L_\mu^2} \leq C \langle t \rangle^{\frac{1-\delta}{4}} \|(u_m, f_m)\|_{L_\mu^2} \leq C (\mathcal{E}_\delta(t))^{\frac{1}{2}} \leq C\varepsilon, \quad (3.9)$$

and that

$$\delta^{\frac{1}{2}} \langle t \rangle^{\frac{1-\delta}{4}} \|(\partial_x^m \partial_y u, \partial_x^m \partial_y f)\|_{L_\mu^2} \leq C(\mathcal{D}_\delta(t))^{\frac{1}{2}}.$$

Proof. From Lemma 2.5 and assertion (3.8) in Corollary 3.2, it follows that

$$\|f\|_{L^\infty} \leq C \langle t \rangle^{\frac{1}{4}} \|\partial_y f\|_{H_{\mu^{\frac{1}{2}}}^{1,0}} \leq C\varepsilon \delta^{-\frac{1}{2}}.$$

Thus we can find a small $\varepsilon > 0$ such that $C\varepsilon \delta^{-\frac{1}{2}} \leq \frac{1}{2}$. This yields

$$\forall (t, x, y) \in [0, +\infty[\times \mathbb{R}_+^2, \quad \frac{1}{2} \leq 1 + f(t, x, y) \leq \frac{3}{2}. \quad (3.10)$$

Thus, by definition (2.4) of u_m ,

$$\forall 1 \leq m \leq 8, \quad \|\partial_x^m u\|_{L_\mu^2} \leq \|u_m\|_{L_\mu^2} + 2\|(\partial_x^{m-1} g) \partial_y u\|_{L_\mu^2}.$$

On the other hand, we use Lemma 2.5 to compute

$$\begin{aligned} \|(\partial_x^{m-1} g) \partial_y u\|_{L_\mu^2} &\leq \|\mu_{\frac{1}{4}} \partial_x^{m-1} g\|_{L_x^2 L_y^\infty} \|\mu_{\frac{1}{4}} \partial_y u\|_{L_x^\infty L_y^2} \\ &\leq C \langle t \rangle^{\frac{1}{4}} \|\partial_x^m f\|_{L_\mu^2} \|\partial_y u\|_{H_{\mu^{\frac{1}{2}}}^{1,0}} \leq C\varepsilon \delta^{-\frac{1}{2}} \|\partial_x^m f\|_{L_\mu^2}, \end{aligned} \quad (3.11)$$

the last inequality using (3.8) again. Combining the above estimates yields

$$\|\partial_x^m u\|_{L_\mu^2} \leq \|u_m\|_{L_\mu^2} + C\varepsilon \delta^{-\frac{1}{2}} \|\partial_x^m f\|_{L_\mu^2}.$$

Similarly, we have

$$\|\partial_x^m f\|_{L_\mu^2} \leq \|f_m\|_{L_\mu^2} + C\varepsilon \delta^{-\frac{1}{2}} \|\partial_x^m u\|_{L_\mu^2}.$$

As a result, by choosing ε such that $\varepsilon \delta^{-\frac{1}{2}}$ is small sufficiently, we have

$$\forall 1 \leq m \leq 8, \quad \|(\partial_x^m u, \partial_x^m f)\|_{L_\mu^2} \leq C\|(u_m, f_m)\|_{L_\mu^2},$$

which with definition (2.10) of \mathcal{E}_δ yields the first assertion in Lemma 3.3.

It remains to prove the second assertion. By definition (2.4) of u_m ,

$$\partial_x^m \partial_y u = \partial_y u_m + \frac{\partial_y u}{1+f} \partial_x^m f - \frac{\partial_y^2 u}{1+f} \partial_x^{m-1} g + \frac{(\partial_y u) \partial_y f}{(1+f)^2} \partial_x^{m-1} g. \quad (3.12)$$

Following the argument in (3.11) we can verify that

$$\left\| \frac{\partial_y^2 u}{1+f} \partial_x^{m-1} g \right\|_{L_\mu^2} \leq C \langle t \rangle^{\frac{1}{4}} \|\partial_x^m f\|_{L_\mu^2} \|\partial_y^2 u\|_{H_{\mu^{\frac{1}{2}}}^{1,0}} \leq C\varepsilon \langle t \rangle^{\frac{\delta}{4}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{1,0}},$$

the last inequality using Lemma 3.1 and the fact that

$$\langle t \rangle^{\frac{1}{4}} \|\partial_x^m f\|_{L_\mu^2} \leq \langle t \rangle^{\frac{\delta}{4}} \langle t \rangle^{\frac{1-\delta}{4}} \|\partial_x^m f\|_{L_\mu^2} \leq C\varepsilon \langle t \rangle^{\frac{\delta}{4}} \quad (3.13)$$

due to (3.9). Similarly,

$$\begin{aligned} \left\| \frac{\partial_y u}{1+f} \partial_x^m f \right\|_{L_\mu^2} &\leq 2\|\mu_{\frac{1}{4}} \partial_y u\|_{L^\infty} \|\partial_x^m f\|_{L_\mu^2} \leq C \langle t \rangle^{\frac{1}{4}} \|\partial_y^2 u\|_{H_{\mu^{\frac{3}{4}}}^{1,0}} \|\partial_x^m f\|_{L_\mu^2} \\ &\leq C \langle t \rangle^{\frac{1}{4}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{1,0}} \|\partial_x^m f\|_{L_\mu^2} \leq C\varepsilon \langle t \rangle^{\frac{\delta}{4}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{1,0}}. \end{aligned}$$

Moreover, using Lemma 2.5 and estimate (3.8) as well as (3.13),

$$\begin{aligned} \left\| \frac{(\partial_y u) \partial_y f}{(1+f)^2} \partial_x^{m-1} g \right\|_{L_\mu^2} &\leq 4 \|\mu_{\frac{1}{8}} \partial_y u\|_{L^\infty} \|\mu_{\frac{1}{8}} \partial_y f\|_{L_x^\infty L_y^2} \|\mu_{\frac{1}{4}} \partial_x^{m-1} g\|_{L_x^2 L_y^\infty} \\ &\leq C \langle t \rangle^{\frac{1}{4}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{1,0}} \|\partial_y f\|_{H_\mu^{1,0}} \left(\langle t \rangle^{\frac{1}{4}} \|\partial_x^m f\|_{L_\mu^2} \right) \leq C \varepsilon^2 \delta^{-\frac{1}{2}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{1,0}}. \end{aligned}$$

Combining the above estimates with (3.12) yields that, for any $1 \leq m \leq 8$,

$$\|\partial_x^m \partial_y u\|_{L_\mu^2} \leq \|\partial_y u_m\|_{L_\mu^2} + C \varepsilon \delta^{-\frac{1}{2}} \langle t \rangle^{\frac{\delta}{4}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{1,0}},$$

and thus, by definition (2.11) of \mathcal{D}_δ and the fact that $\varepsilon \delta^{-1}$ is small,

$$\delta^{\frac{1}{2}} \langle t \rangle^{\frac{1-\delta}{4}} \|\partial_x^m \partial_y u\|_{L_\mu^2} \leq (\mathcal{D}_\delta(t))^{\frac{1}{2}} + C \varepsilon \delta^{-1} (\delta \langle t \rangle^{\frac{7-\delta}{4}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{1,0}}) \leq C (\mathcal{D}_\delta(t))^{\frac{1}{2}}.$$

Similarly for $\|\partial_x^m \partial_y f\|_{L_\mu^2}$. Thus the second assertion in Lemma 3.3 follows. The proof is thus completed. \square

Lemma 3.4 (Technical lemma). *Let $\varphi \in H_\mu^{0,1}$ and $\psi \in L_\mu^2$ satisfy that*

$$(\partial_t - \partial_y^2) \varphi = \psi, \quad \varphi \partial_y \varphi|_{y=0} = 0. \quad (3.14)$$

Then we have

$$\frac{d}{dt} \|\varphi\|_{L_\mu^2}^2 + \frac{1-\delta}{2\langle t \rangle} \|\varphi\|_{L_\mu^2}^2 + \delta \|\partial_y \varphi\|_{L_\mu^2}^2 \leq 2(\psi, \varphi)_{L_\mu^2}, \quad (3.15)$$

and

$$\frac{d}{dt} \left(\langle t \rangle^{\frac{1-\delta}{2}} \|\varphi\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{1-\delta}{2}} \|\partial_y \varphi\|_{L_\mu^2}^2 \leq 2 \langle t \rangle^{\frac{1-\delta}{2}} (\psi, \varphi)_{L_\mu^2}. \quad (3.16)$$

Proof. It suffices to prove the first assertion (3.15), since the second one (3.16) will follow if we multiply (3.15) by $\langle t \rangle^{\frac{1-\delta}{2}}$ and then use the fact that

$$\frac{d}{dt} \langle t \rangle^{\frac{1-\delta}{2}} = \langle t \rangle^{\frac{1-\delta}{2}} \frac{1-\delta}{2\langle t \rangle}.$$

By virtue of the fact that

$$\partial_t \mu = -\frac{y^2}{4\langle t \rangle^2} \mu, \quad \partial_y^2 \mu = \frac{1}{2\langle t \rangle} \mu + \frac{y^2}{4\langle t \rangle^2} \mu,$$

we compute

$$2 \int_{\mathbb{R}_+^2} (\partial_t \varphi) \varphi \mu dx dy = \frac{d}{dt} \|\varphi\|_{L_\mu^2}^2 - \int_{\mathbb{R}_+^2} \varphi^2 (\partial_t \mu) dx dy = \frac{d}{dt} \|\varphi\|_{L_\mu^2}^2 + \frac{\|y\varphi\|_{L_\mu^2}^2}{4\langle t \rangle^2},$$

and

$$\begin{aligned} -2 \int_{\mathbb{R}_+^2} (\partial_y^2 \varphi) \varphi \mu dx dy &= 2 \|\partial_y \varphi\|_{L_\mu^2}^2 + 2 \int_{\mathbb{R}_+^2} (\partial_y \varphi) \varphi \partial_y \mu dx dy \\ &= 2 \|\partial_y \varphi\|_{L_\mu^2}^2 - \int_{\mathbb{R}_+^2} \varphi^2 \partial_y^2 \mu dx dy = 2 \|\partial_y \varphi\|_{L_\mu^2}^2 - \frac{1}{2\langle t \rangle} \|\varphi\|_{L_\mu^2}^2 - \frac{\|y\varphi\|_{L_\mu^2}^2}{4\langle t \rangle^2}. \end{aligned}$$

As a result, we take L_μ^2 -product with 2φ in (3.14) and then combine the two equations above to get

$$\frac{d}{dt} \|\varphi\|_{L_\mu^2}^2 + 2\|\partial_y \varphi\|_{L_\mu^2}^2 - \frac{1}{2\langle t \rangle} \|\varphi\|_{L_\mu^2}^2 = 2(\psi, \varphi)_{L_\mu^2}. \quad (3.17)$$

Moreover, it follows from (2.14) in Lemma 2.4 that

$$\frac{1}{2\langle t \rangle} \|\varphi\|_{L_\mu^2}^2 \leq \|\partial_y \varphi\|_{L_\mu^2}^2,$$

and thus

$$2\|\partial_y \varphi\|_{L_\mu^2}^2 = \delta \|\partial_y \varphi\|_{L_\mu^2}^2 + (2 - \delta) \|\partial_y \varphi\|_{L_\mu^2}^2 \geq \delta \|\partial_y \varphi\|_{L_\mu^2}^2 + \frac{2 - \delta}{2\langle t \rangle} \|\varphi\|_{L_\mu^2}^2.$$

Substituting the above estimate into (3.17) yields the first assertion in Lemma 3.4. The proof is thus completed. \square

Remark 3.5. If $\varphi \in H_\mu^{0,1}$ and $\psi \in L_\mu^2$ satisfy that

$$(\partial_t - \partial_y^2)\varphi + \frac{1}{\langle t \rangle}\varphi = \psi, \quad \varphi \partial_y \varphi|_{y=0} = 0,$$

then it follows from (3.15) in Lemma 3.4 that

$$\frac{d}{dt} \|\varphi\|_{L_\mu^2}^2 + \frac{5 - \delta}{2\langle t \rangle} \|\varphi\|_{L_\mu^2}^2 + \delta \|\partial_y \varphi\|_{L_\mu^2}^2 \leq 2(\psi, \varphi)_{L_\mu^2}.$$

Thus, similar to (3.16), it follows from the above estimate that

$$\frac{d}{dt} \left(\langle t \rangle^{\frac{5-\delta}{2}} \|\varphi\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{5-\delta}{2}} \|\partial_y \varphi\|_{L_\mu^2}^2 \leq 2 \langle t \rangle^{\frac{5-\delta}{2}} (\psi, \varphi)_{L_\mu^2}.$$

4. Estimate on the auxiliary functions u_m and f_m

This part is devoted to dealing with the terms (u_m, f_m) in definitions (2.10) and (2.11) of \mathcal{E}_δ and \mathcal{D}_δ . The main result can be stated as follows.

Proposition 4.1. *Let ε be the small constant given in Lemma 3.3. Suppose*

$$\forall t > 0, \quad \mathcal{E}_\delta(t) + \frac{1}{2} \int_0^t \mathcal{D}_\delta(s) ds \leq 2\varepsilon^2. \quad (4.1)$$

Then

$$\frac{d}{dt} \sum_{m \leq 8} \langle t \rangle^{\frac{1-\delta}{2}} \|(u_m, f_m)\|_{L_\mu^2}^2 + \delta \sum_{m \leq 8} \langle t \rangle^{\frac{1-\delta}{2}} \|(\partial_y u_m, \partial_y f_m)\|_{L_\mu^2}^2 \leq C\varepsilon\delta^{-2}\mathcal{D}_\delta. \quad (4.2)$$

Proof. It suffices to consider the case of $1 \leq m \leq 8$, since the energy estimate for (u_m, f_m) with $m = 0$ is straightforward. For each fixed $1 \leq m \leq 8$, the auxiliary

functions u_m, f_m solve the following equations (see Appendix B for the derivation in detail):

$$\begin{cases} \left(\partial_t + u\partial_x + v\partial_y - \partial_y^2 \right) u_m - ((1+f)\partial_x + g\partial_y) f_m = \sum_{1 \leq j \leq 3} S_{m,j}, \\ \left(\partial_t + u\partial_x + v\partial_y - \partial_y^2 \right) f_m - ((1+f)\partial_x + g\partial_y) u_m = \sum_{1 \leq j \leq 3} \tilde{S}_{m,j}, \end{cases} \quad (4.3)$$

where $S_{m,j}$ and $\tilde{S}_{m,j}$ are given in (B.5)-(B.10). It suffices to perform estimates for the first evolution equation in (4.3). To do so we recall the representations of $S_{m,j}$ given in (B.5)-(B.7), that is,

$$\begin{aligned} S_{m,1} = & \sum_{k=1}^m \binom{m}{k} [(\partial_x^k f) \partial_x^{m-k+1} f - (\partial_x^k u) \partial_x^{m-k+1} u] \\ & + \sum_{k=1}^{m-1} \binom{m}{k} [(\partial_x^k g) \partial_x^{m-k} \partial_y f - (\partial_x^k v) \partial_x^{m-k} \partial_y u], \end{aligned} \quad (4.4)$$

and

$$\begin{aligned} S_{m,2} = & \frac{\partial_y u}{1+f} \sum_{k=1}^{m-1} \binom{m-1}{k} [(\partial_x^k f) \partial_x^{m-k} v - (\partial_x^k g) \partial_x^{m-k} u] \\ & - \frac{\partial_y u}{1+f} \sum_{k=1}^{m-1} \binom{m-1}{k} [(\partial_x^k u) \partial_x^{m-k} g - (\partial_x^k v) \partial_x^{m-k} f], \end{aligned} \quad (4.5)$$

and

$$\begin{aligned} S_{m,3} = & \left[\frac{g\partial_y f}{1+f} + 2\partial_y \left(\frac{\partial_y u}{1+f} \right) \right] \partial_x^m f - \frac{g\partial_y u}{1+f} \partial_x^m u - 2 \frac{(\partial_y f)^2 \partial_y u}{(1+f)^3} \partial_x^{m-1} g \\ & + \left[\frac{(\partial_y f) \partial_x f - (\partial_y u) \partial_x u}{1+f} + \frac{g(\partial_y f)^2 - g(\partial_y u)^2 + 2(\partial_y f) \partial_y^2 u}{(1+f)^2} \right] \partial_x^{m-1} g. \end{aligned} \quad (4.6)$$

Applying Lemma 3.4 to the first equation in (4.3), we have

$$\begin{aligned} & \frac{d}{dt} \left(\langle t \rangle^{\frac{1-\delta}{2}} \|u_m\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{1-\delta}{2}} \|\partial_y u_m\|_{L_\mu^2}^2 \\ & \leq 2 \langle t \rangle^{\frac{1-\delta}{2}} \left(\left[(1+f)\partial_x + g\partial_y \right] f_m, u_m \right)_{L_\mu^2} + 2 \langle t \rangle^{\frac{1-\delta}{2}} \|u_m\|_{L_\mu^2} \sum_{1 \leq j \leq 3} \|S_{m,j}\|_{L_\mu^2} \\ & \quad - 2 \langle t \rangle^{\frac{1-\delta}{2}} \left((u\partial_x + v\partial_y) u_m, u_m \right)_{L_\mu^2}. \end{aligned} \quad (4.7)$$

We claim that

$$\sum_{1 \leq j \leq 3} \|S_{m,j}\|_{L_\mu^2} \leq C \langle t \rangle^{-\frac{1}{2}} \left(\varepsilon \delta^{-\frac{1}{2}} \|(\partial_y u, \partial_y f)\|_{H_\mu^{8,0}} + \varepsilon \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}} + \delta^{-2} \mathcal{D}_\delta \right), \quad (4.8)$$

with the proof postponed. Estimate (4.8) with Lemma 2.4 yields

$$\begin{aligned}
& \langle t \rangle^{\frac{1-\delta}{2}} \|u_m\|_{L_\mu^2} \sum_{1 \leq j \leq 3} \|S_{m,j}\|_{L_\mu^2} \\
& \leq C \langle t \rangle^{\frac{1-\delta}{2}} \|\partial_y u_m\|_{L_\mu^2} \left(\varepsilon \delta^{-\frac{1}{2}} \|(\partial_y u, \partial_y f)\|_{H_\mu^{8,0}} + \varepsilon \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}} \right) \\
& \quad + C \delta^{-2} \langle t \rangle^{-\frac{\delta}{2}} \|u_m\|_{L_\mu^2} \mathcal{D}_\delta \\
& \leq C \varepsilon \delta^{-2} \mathcal{D}_\delta,
\end{aligned}$$

the last inequality using definitions (2.10) and (2.11) of \mathcal{E}_δ and \mathcal{D}_δ , Lemma 3.3 and the smallness assumption (4.1). For the last term on the right-hand side of (4.7), integrating by parts and then using the divergence-free condition $\partial_x u + \partial_y v = 0$, we have

$$\begin{aligned}
& \left| \langle t \rangle^{\frac{1-\delta}{2}} \left((u \partial_x + v \partial_y) u_m, u_m \right)_{L_\mu^2} \right| = \frac{1}{2} \langle t \rangle^{\frac{1-\delta}{2}} \left| \int_{\mathbb{R}_+^2} v u_m^2 \partial_y \left(e^{\frac{y^2}{4\langle t \rangle}} \right) dx dy \right| \\
& \leq C \langle t \rangle^{\frac{1-\delta}{2}} \|v\|_{L^\infty} \|u_m\|_{L_\mu^2} \left\| \frac{y}{\langle t \rangle} u_m \right\|_{L_\mu^2} \leq C \langle t \rangle^{\frac{1-\delta}{2} + \frac{1}{4}} \|u\|_{H_\mu^{2,0}} \|u_m\|_{L_\mu^2} \left\| \frac{y}{\langle t \rangle} u_m \right\|_{L_\mu^2} \\
& \leq C \varepsilon \langle t \rangle^{\frac{1-\delta}{2} + \frac{1}{4}} \langle t \rangle^{-\frac{5-\delta}{4}} \|u_m\|_{L_\mu^2} \left\| \frac{y}{\langle t \rangle} u_m \right\|_{L_\mu^2} \leq C \varepsilon \langle t \rangle^{\frac{1-\delta}{2}} \|\partial_y u_m\|_{L_\mu^2}^2 \leq C \varepsilon \delta^{-1} \mathcal{D}_\delta,
\end{aligned} \tag{4.9}$$

where in the second inequality we used Lemma 2.5 and in the last line we used (3.8) and (2.15) as well as definition (2.11) of \mathcal{D}_δ . Substituting the above estimates into (4.7) yields

$$\begin{aligned}
& \frac{d}{dt} \left(\langle t \rangle^{\frac{1-\delta}{2}} \|u_m\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{1-\delta}{2}} \|\partial_y u_m\|_{L_\mu^2}^2 \\
& \leq 2 \langle t \rangle^{\frac{1-\delta}{2}} \left([(1+f)\partial_x + g\partial_y] f_m, u_m \right)_{L_\mu^2} + C \varepsilon \delta^{-2} \mathcal{D}_\delta.
\end{aligned}$$

Similarly, repeating the argument above for the second equation in (4.3) we have

$$\begin{aligned}
& \frac{d}{dt} \left(\langle t \rangle^{\frac{1-\delta}{2}} \|f_m\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{1-\delta}{2}} \|\partial_y f_m\|_{L_\mu^2}^2 \\
& \leq 2 \langle t \rangle^{\frac{1-\delta}{2}} \left([(1+f)\partial_x + g\partial_y] u_m, f_m \right)_{L_\mu^2} + C \varepsilon \delta^{-2} \mathcal{D}_\delta.
\end{aligned}$$

On the other hand, by a similar argument to that in (4.9), we have

$$\begin{aligned}
& \langle t \rangle^{\frac{1-\delta}{2}} \left([(1+f)\partial_x + g\partial_y] f_m, u_m \right)_{L_\mu^2} + \langle t \rangle^{\frac{1-\delta}{2}} \left([(1+f)\partial_x + g\partial_y] u_m, f_m \right)_{L_\mu^2} \\
& = - \langle t \rangle^{\frac{1-\delta}{2}} \int_{\mathbb{R}_+^2} g f_m u_m \partial_y \left(e^{\frac{y^2}{4\langle t \rangle}} \right) dx dy \leq C \varepsilon \delta^{-1} \mathcal{D}_\delta.
\end{aligned}$$

As a result, combining the above estimates yields that, for any $1 \leq m \leq 8$,

$$\frac{d}{dt} \left(\langle t \rangle^{\frac{1-\delta}{2}} \|(u_m, f_m)\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{1-\delta}{2}} \|(\partial_y u_m, \partial_y f_m)\|_{L_\mu^2}^2 \leq C \varepsilon \delta^{-2} \mathcal{D}_\delta.$$

Note the above estimate still holds true for $m = 0$ by performing straightforward energy estimate for system (1.5). Thus assertion (4.2) in Proposition 4.1 follows.

So to complete the proof of Proposition 4.1, it remains to prove assertion (4.8), and we proceed through the following three steps. In the following discussion, let $1 \leq m \leq 8$ be fixed.

Step 1 (Estimate on $S_{m,1}$). For any k with $1 \leq k \leq [\frac{m}{2}] \leq 4$, we use Sobolev inequality to compute

$$\begin{aligned} \|(\partial_x^k f) \partial_x^{m+1-k} f\|_{L_\mu^2} &\leq \|\mu_{\frac{1}{4}} \partial_x^k f\|_{L_x^\infty L_y^2} \|\mu_{\frac{1}{4}} \partial_x^{m+1-k} f\|_{L_x^2 L_y^\infty} \\ &\leq C \langle t \rangle^{\frac{1}{4}} \|f\|_{H_\mu^{5,0}} \|\partial_y f\|_{H_\mu^{8,0}} \leq C \varepsilon \langle t \rangle^{\frac{1}{4} - \frac{5-\delta}{4}} \|\partial_y f\|_{H_\mu^{8,0}} \leq C \varepsilon \langle t \rangle^{-\frac{1}{2}} \|\partial_y f\|_{H_\mu^{8,0}}, \end{aligned}$$

the last line using Lemma 2.5 as well as (3.8). Similarly, for any k with $[\frac{m}{2}] + 1 \leq k \leq m \leq 8$, we have $m+1-k \leq 4$ and thus

$$\begin{aligned} \|(\partial_x^k f) \partial_x^{m+1-k} f\|_{L_\mu^2} &\leq \|\mu_{\frac{1}{4}} \partial_x^k f\|_{L_x^2 L_y^\infty} \|\mu_{\frac{1}{4}} \partial_x^{m+1-k} f\|_{L_x^\infty L_y^2} \\ &\leq C \langle t \rangle^{\frac{1}{4}} \|\partial_y f\|_{H_\mu^{8,0}} \|f\|_{H_\mu^{5,0}} \leq C \varepsilon \langle t \rangle^{-\frac{1}{2}} \|\partial_y f\|_{H_\mu^{8,0}}. \end{aligned}$$

Combining the above estimates yields that

$$\sum_{k=1}^m \binom{m}{k} \|(\partial_x^k f) \partial_x^{m+1-k} f\|_{L_\mu^2} \leq C \varepsilon \langle t \rangle^{-\frac{1}{2}} \|\partial_y f\|_{H_\mu^{8,0}}. \quad (4.10)$$

In the same way we have that

$$\sum_{k=1}^m \binom{m}{k} \|(\partial_x^k u) \partial_x^{m+1-k} u\|_{L_\mu^2} \leq C \varepsilon \langle t \rangle^{-\frac{1}{2}} \|\partial_y u\|_{H_\mu^{8,0}}, \quad (4.11)$$

and that, observing $\partial_y g = -\partial_x f$ and $\|f\|_{H_\mu^{8,0}} \leq (2 \langle t \rangle)^{\frac{1}{2}} \|\partial_y f\|_{H_\mu^{8,0}}$ by Lemma 2.4,

$$\begin{aligned} \sum_{k=1}^{m-1} \binom{m-1}{k} \|(\partial_x^k g) \partial_x^{m-k} \partial_y f\|_{L_\mu^2} &\leq C \sum_{k=1}^{[\frac{m-1}{2}]} \|\mu_{\frac{1}{4}} \partial_x^k g\|_{L^\infty} \|\mu_{\frac{1}{4}} \partial_x^{m-k} \partial_y f\|_{L^2} \\ &\quad + C \sum_{k=[\frac{m-1}{2}]+1}^{m-1} \|\mu_{\frac{1}{4}} \partial_x^k g\|_{L_x^2 L_y^\infty} \|\mu_{\frac{1}{4}} \partial_x^{m-k} \partial_y f\|_{L_x^\infty L_y^2} \\ &\leq C \langle t \rangle^{\frac{1}{4}} \|f\|_{H_\mu^{5,0}} \|\partial_y f\|_{H_\mu^{8,0}} + C \langle t \rangle^{\frac{1}{4}} \|f\|_{H_\mu^{8,0}} \|\partial_y f\|_{H_\mu^{5,0}} \\ &\leq C \left(\langle t \rangle^{\frac{1}{4}} \|f\|_{H_\mu^{5,0}} + \langle t \rangle^{\frac{1}{4}} \langle t \rangle^{\frac{1}{2}} \|\partial_y f\|_{H_\mu^{5,0}} \right) \|\partial_y f\|_{H_\mu^{8,0}} \\ &\leq C \varepsilon \delta^{-\frac{1}{2}} \langle t \rangle^{-\frac{1}{2}} \|\partial_y f\|_{H_\mu^{8,0}}, \end{aligned} \quad (4.12)$$

the last inequality following from (3.8). Similarly,

$$\sum_{k=1}^{m-1} \binom{m-1}{k} \|(\partial_x^k v) \partial_x^{m-k} \partial_y u\|_{L_\mu^2} \leq C \varepsilon \delta^{-\frac{1}{2}} \langle t \rangle^{-\frac{1}{2}} \|\partial_y u\|_{H_\mu^{8,0}}.$$

Combining the two estimates above and estimates (4.10)-(4.11) with (4.4), we conclude that

$$S_{m,1} \leq C\varepsilon\delta^{-\frac{1}{2}} \langle t \rangle^{-\frac{1}{2}} \|(\partial_y u, \partial_y f)\|_{H_\mu^{8,0}}. \quad (4.13)$$

Step 2 (Estimate on $S_{m,2}$). In view of (3.10), we follow the argument in (4.12) to compute, using Lemmas 2.5 and 3.3 and assertion (3.8),

$$\begin{aligned} \sum_{k=1}^{m-1} \binom{m-1}{k} \left\| \frac{\partial_y u}{1+f} (\partial_x^k f) \partial_x^{m-k} v \right\|_{L_\mu^2} &\leq C \sum_{k=1}^{m-1} \|\mu_{\frac{1}{4}} \partial_y u\|_{L^\infty} \|(\partial_x^k f) \partial_x^{m-k} v\|_{L_{\mu^{\frac{1}{2}}}^2} \\ &\leq C \langle t \rangle^{\frac{1}{4}} \|\partial_y^2 u\|_{H_{\mu^{\frac{1}{2}}}^{1,0}} \left(\langle t \rangle^{\frac{1}{4}} \|f\|_{H_{\mu^{\frac{3}{4}}}^{5,0}} \|u\|_{H_\mu^{8,0}} + \langle t \rangle^{\frac{1}{4}} \|f\|_{H_\mu^{8,0}} \|u\|_{H_{\mu^{\frac{3}{4}}}^{5,0}} \right) \\ &\leq C\varepsilon^2 \langle t \rangle^{\frac{1}{2}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{1,0}} \langle t \rangle^{-\frac{5-\delta}{4}} \langle t \rangle^{-\frac{1-\delta}{4}} \leq C\varepsilon \langle t \rangle^{-\frac{1}{2}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}}. \end{aligned}$$

Similarly,

$$\begin{aligned} \sum_{k=1}^{m-1} \binom{m-1}{k} \left(\left\| \frac{\partial_y u}{1+f} (\partial_x^k g) \partial_x^{m-k} u \right\|_{L_\mu^2} + \left\| \frac{\partial_y u}{1+f} (\partial_x^k u) \partial_x^{m-k} g \right\|_{L_\mu^2} \right) \\ + \sum_{k=1}^{m-1} \binom{m-1}{k} \left\| \frac{\partial_y u}{1+f} (\partial_x^k v) \partial_x^{m-k} f \right\|_{L_\mu^2} \leq C\varepsilon \langle t \rangle^{-\frac{1}{2}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}}. \end{aligned}$$

Combining the estimates above with (4.5), we conclude that

$$\|S_{m,2}\|_{L_\mu^2} \leq C\varepsilon \langle t \rangle^{-\frac{1}{2}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}}. \quad (4.14)$$

Step 3 (Estimate on $S_{m,3}$). It just follows from straightforward computation. In fact, using (3.10) again yields

$$\begin{aligned} &\left\| \left[\frac{g \partial_y f}{1+f} + 2\partial_y \left(\frac{\partial_y u}{1+f} \right) \right] \partial_x^m f \right\|_{L_\mu^2} \\ &\leq C \|\mu_{\frac{1}{4}} \partial_x^m f\|_{L_x^2 L_y^\infty} \|\mu_{\frac{1}{4}} (g \partial_y f + \partial_y^2 u + (\partial_y u) \partial_y f)\|_{L_x^\infty L_y^2} \\ &\leq C \langle t \rangle^{\frac{1}{4}} \|\partial_y f\|_{H_\mu^{8,0}} \|\mu_{\frac{1}{4}} (g \partial_y f + \partial_y^2 u + (\partial_y u) \partial_y f)\|_{L_x^\infty L_y^2}. \end{aligned}$$

On the other hand, using Lemmas 3.1 and 2.5 we compute

$$\|\mu_{\frac{1}{4}} \partial_y^2 u\|_{L_x^\infty L_y^2} \leq C \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}},$$

and

$$\begin{aligned} \|\mu_{\frac{1}{4}} (\partial_y u) \partial_y f\|_{L_x^\infty L_y^2} &\leq \|\mu_{\frac{1}{8}} \partial_y u\|_{L^\infty} \|\mu_{\frac{1}{8}} \partial_y f\|_{L_x^\infty L_y^2} \leq C \langle t \rangle^{\frac{1}{4}} \|\partial_y^2 u\|_{H_{\mu^{\frac{1}{2}}}^{5,0}} \|\partial_y f\|_{H_{\mu^{\frac{1}{2}}}^{5,0}} \\ &\leq C \langle t \rangle^{\frac{1}{4}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}} \|\partial_y f\|_{H_{\mu^{\frac{1}{2}}}^{5,0}} \leq C\varepsilon\delta^{-\frac{1}{2}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}}, \end{aligned}$$

the last inequality using (3.8). Similarly, using Lemma 2.4 additionally,

$$\begin{aligned} \|\mu_{\frac{1}{4}} g \partial_y f\|_{L_x^\infty L_y^2} &\leq C \langle t \rangle^{\frac{1}{4}} \|f\|_{H_{\mu^{\frac{3}{4}}}^{5,0}} \|\partial_y f\|_{H_{\mu^{\frac{1}{2}}}^{5,0}} \\ &\leq C \langle t \rangle^{\frac{1}{4} + \frac{1}{2}} \|f\|_{H_{\mu^{\frac{3}{4}}}^{5,0}} \|\partial_y^2 f\|_{H_{\mu^{\frac{1}{2}}}^{5,0}} \leq C\varepsilon \|\partial_y^2 \mathcal{F}\|_{H_\mu^{5,0}}. \end{aligned}$$

As a result, combining the above estimates, we conclude that

$$\left\| \left[\frac{g \partial_y f}{1+f} + 2 \partial_y \left(\frac{\partial_y u}{1+f} \right) \right] \partial_x^m f \right\|_{L_\mu^2} \leq C \delta^{-\frac{1}{2}} \langle t \rangle^{\frac{1}{4}} \|\partial_y f\|_{H_\mu^{8,0}} \|(\partial_y^2 \mathcal{U}, \partial_y^2 \mathcal{F})\|_{H_\mu^{5,0}}.$$

Following the above argument with slight modification, we can verify that the L_μ^2 -norms of the remaining terms in (4.6) are bounded above from above by

$$C \delta^{-\frac{1}{2}} \langle t \rangle^{\frac{1}{4}} \left(\|\partial_y u\|_{H_\mu^{8,0}} + \|f\|_{H_\mu^{8,0}} \right) \|(\partial_y^2 \mathcal{U}, \partial_y^2 \mathcal{F})\|_{H_\mu^{5,0}},$$

and thus

$$\begin{aligned} S_{m,3} &\leq C \delta^{-\frac{1}{2}} \langle t \rangle^{\frac{1}{4}} \left(\|\partial_y u\|_{H_\mu^{8,0}} + \|f\|_{H_\mu^{8,0}} + \|\partial_y f\|_{H_\mu^{8,0}} \right) \|(\partial_y^2 \mathcal{U}, \partial_y^2 \mathcal{F})\|_{H_\mu^{5,0}} \\ &\leq C \delta^{-\frac{1}{2}} \langle t \rangle^{\frac{3}{4}} \|(\partial_y u, \partial_y f)\|_{H_\mu^{8,0}} \|(\partial_y^2 \mathcal{U}, \partial_y^2 \mathcal{F})\|_{H_\mu^{5,0}} \leq C \delta^{-2} \langle t \rangle^{-\frac{1}{2}} \mathcal{D}_\delta(t), \end{aligned}$$

the second inequality using Lemma 2.4 and the last inequality following from Lemma 3.3 as well as definition 2.11 of \mathcal{D}_δ . This with (4.13) and (4.14) yields (4.8). Thus the proof of Proposition 4.1 is completed. \square

5. Estimate on the auxiliary functions \mathcal{U} and \mathcal{F}

In this section, we first derive the estimate on $(\mathcal{U}, \mathcal{F})$ defined in (2.9), and then complete the proof of Theorem 2.2.

Proposition 5.1. *Let ε be the small constant given in Lemma 3.3. Suppose that the small assumption (4.1) holds. Then*

$$\frac{d}{dt} \left(\langle t \rangle^{\frac{5-\delta}{2}} \|(\mathcal{U}, \mathcal{F})\|_{H_\mu^{5,0}}^2 \right) + \delta \langle t \rangle^{\frac{5-\delta}{2}} \|(\partial_y \mathcal{U}, \partial_y \mathcal{F})\|_{H_\mu^{5,0}}^2 \leq C \varepsilon \delta^{-1} \mathcal{D}_\delta. \quad (5.1)$$

Proof. *Step 1).* Note that \mathcal{U} and \mathcal{F} satisfy the following equations (see Appendix A for the derivation):

$$\begin{aligned} (\partial_t - \partial_y^2) \mathcal{U} + \frac{1}{\langle t \rangle} \mathcal{U} - \partial_x \mathcal{F} &= -(u \partial_x + v \partial_y) u + (f \partial_x + g \partial_y) f \\ &\quad + \frac{y}{2 \langle t \rangle} \int_y^{+\infty} \left[(u \partial_x + v \partial_y) u - (f \partial_x + g \partial_y) f \right] d\tilde{y} \end{aligned} \quad (5.2)$$

and

$$\begin{aligned} (\partial_t - \partial_y^2) \mathcal{F} + \frac{1}{\langle t \rangle} \mathcal{F} - \partial_x \mathcal{U} &= -(u \partial_x + v \partial_y) f + (f \partial_x + g \partial_y) u \\ &\quad + \frac{y}{2 \langle t \rangle} \int_y^{+\infty} \left[(u \partial_x + v \partial_y) f - (f \partial_x + g \partial_y) u \right] d\tilde{y}. \end{aligned} \quad (5.3)$$

Observe the two equations above are supplemented with the boundary condition that

$$\mathcal{U}|_{y=0} = \mathcal{F}|_{y=0} = 0. \quad (5.4)$$

We first perform estimates for \mathcal{U} . For any $m \in \mathbb{Z}_+$ with $0 \leq m \leq 5$, applying ∂_x^m to the equation (5.2) yields

$$(\partial_t - \partial_y^2) \partial_x^m \mathcal{U} + \frac{1}{\langle t \rangle} \partial_x^m \mathcal{U} = \partial_x^{m+1} \mathcal{F} + R_{m,1} + R_{m,2} \quad (5.5)$$

with

$$\begin{cases} R_{m,1} = \partial_x^m \left[- (u \partial_x + v \partial_y) u + (f \partial_x + g \partial_y) f \right], \\ R_{m,2} = \frac{y}{2 \langle t \rangle} \int_y^{+\infty} \partial_x^m \left[(u \partial_x + v \partial_y) u - (f \partial_x + g \partial_y) f \right] d\tilde{y}. \end{cases} \quad (5.6)$$

We take the L_μ^2 -product with $\partial_x^m \mathcal{U}$ in (5.5) and then observe Remark 3.5, to obtain that

$$\begin{aligned} & \frac{d}{dt} \left(\langle t \rangle^{\frac{5-\delta}{2}} \|\partial_x^m \mathcal{U}\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{5-\delta}{2}} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 \\ & \leq 2 \langle t \rangle^{\frac{5-\delta}{2}} \left(\partial_x^{m+1} \mathcal{F}, \partial_x^m \mathcal{U} \right)_{L_\mu^2} + 2 \langle t \rangle^{\frac{5-\delta}{2}} \sum_{1 \leq j \leq 2} \|R_{m,j}\|_{L_\mu^2} \|\partial_x^m \mathcal{U}\|_{L_\mu^2}. \end{aligned} \quad (5.7)$$

Step 2). In this step we will deal with the last term on the right-hand side of (5.7), and prove that

$$\sum_{1 \leq j \leq 2} \|R_{m,j}\|_{L_\mu^2} \leq C \varepsilon \langle t \rangle^{-\frac{1}{2}} \|(\partial_y \mathcal{U}, \partial_y \mathcal{F})\|_{H_\mu^{5,0}}. \quad (5.8)$$

For each $0 \leq m \leq 5$ we have $\|\partial_x^m (u \partial_x u)\|_{L_\mu^2} \leq \|(\mu_{\frac{1}{4}} u) \mu_{\frac{1}{4}} \partial_x u\|_{H_x^5 L_y^2}$ and thus

$$\|\partial_x^m (u \partial_x u)\|_{L_\mu^2} \leq C \|\mu_{\frac{1}{4}} u\|_{H_x^5 L_y^\infty} \|\mu_{\frac{1}{4}} \partial_x u\|_{H_x^5 L_y^2} \leq C \langle t \rangle^{\frac{1}{4}} \|\partial_y u\|_{H_{\mu_{3/4}}^{5,0}} \|\partial_x u\|_{H_{\mu_{3/4}}^{5,0}}, \quad (5.9)$$

the last inequality using Lemma 2.5. For the last term on the right-hand side, it follows from the interpolation inequality that

$$\begin{aligned} \|\partial_x u\|_{H_{\mu_{3/4}}^{5,0}} & \leq \|u\|_{H_{\mu_{3/4}}^{6,0}} \leq C \|u\|_{H_{\mu_{3/4}}^{8,0}}^{\frac{1}{3}} \|u\|_{H_{\mu_{3/4}}^{5,0}}^{\frac{2}{3}} \\ & \leq C \varepsilon \langle t \rangle^{-\frac{1-\delta}{4} \times \frac{1}{3}} \langle t \rangle^{-\frac{5-\delta}{4} \times \frac{2}{3}} \leq C \varepsilon \langle t \rangle^{-\frac{11}{12} + \frac{\delta}{4}}, \end{aligned} \quad (5.10)$$

where in the last line we use Lemma 3.3 and assertion (3.8). As a result, we substitute (5.10) into (5.9) and then use Lemma 3.1 as well as the fact that $0 < \delta \leq \frac{1}{25}$, to conclude that

$$\|\partial_x^m (u \partial_x u)\|_{L_\mu^2} \leq C \varepsilon \langle t \rangle^{-\frac{11}{12} + \frac{\delta}{4}} \langle t \rangle^{\frac{1}{4}} \|\partial_y u\|_{H_{\mu_{3/4}}^{5,0}} \leq C \varepsilon \langle t \rangle^{-\frac{1}{2}} \|\partial_y \mathcal{U}\|_{H_\mu^{5,0}}. \quad (5.11)$$

Following the above argument with slight modification, we can verify that, for any $0 \leq m \leq 5$,

$$\begin{aligned} \|\partial_x^m (v \partial_y u)\|_{L_\mu^2} & \leq C \|\mu_{1/4} v\|_{H_x^5 L_y^\infty} \|\mu_{1/4} \partial_y u\|_{H_x^5 L_y^2} \\ & \leq C \langle t \rangle^{\frac{1}{4}} \|\partial_x u\|_{H_{\mu_{3/4}}^{5,0}} \|\partial_y u\|_{H_{\mu_{3/4}}^{5,0}} \leq C \varepsilon \langle t \rangle^{-\frac{1}{2}} \|\partial_y \mathcal{U}\|_{H_\mu^{5,0}}, \end{aligned}$$

where the first inequality in the last line follows from Lemma 2.5 and the last inequality holds because of (5.10). Similarly,

$$\|\partial_x^m(f\partial_x f)\|_{L_\mu^2} + \|\partial_x^m(g\partial_y f)\|_{L_\mu^2} \leq C\varepsilon \langle t \rangle^{-\frac{1}{2}} \|\partial_y \mathcal{F}\|_{H_\mu^{5,0}}.$$

Combining the above estimates and (5.11) with (5.6) yields

$$\|R_{m,1}\|_{L_\mu^2} \leq C\varepsilon \langle t \rangle^{-\frac{1}{2}} \|(\partial_y \mathcal{U}, \partial_y \mathcal{F})\|_{H_\mu^{5,0}}. \quad (5.12)$$

In view of (5.6), we apply (2.15) in Lemma 2.4 with

$$h = \int_y^{+\infty} \partial_x^m \left[(u\partial_x + v\partial_y)u - (f\partial_x + g\partial_y)f \right] d\tilde{y},$$

to obtain that, observing $\partial_y h = R_{m,1}$,

$$\|R_{m,2}\|_{L_\mu^2} = \left\| \frac{y}{2\langle t \rangle} h \right\|_{L_\mu^2} \leq C \|R_{m,1}\|_{L_\mu^2},$$

which with (5.12) yields (5.8).

Step 3). For each $m \leq 5$, using (5.8) and Lemma 2.4 yields

$$\begin{aligned} \langle t \rangle^{\frac{5-\delta}{2}} \sum_{1 \leq j \leq 2} \|R_{m,j}\|_{L_\mu^2} \|\partial_x^m \mathcal{U}\|_{L_\mu^2} &\leq C\varepsilon \langle t \rangle^{\frac{5-\delta}{2}} \langle t \rangle^{-\frac{1}{2}} \|(\partial_y \mathcal{U}, \partial_y \mathcal{F})\|_{H_\mu^{5,0}} \|\mathcal{U}\|_{H_\mu^{5,0}} \\ &\leq C\varepsilon \langle t \rangle^{\frac{5-\delta}{2}} \|(\partial_y \mathcal{U}, \partial_y \mathcal{F})\|_{H_\mu^{5,0}}^2 \leq C\varepsilon \delta^{-1} \mathcal{D}_\delta, \end{aligned}$$

the last inequality using definition (2.11) of \mathcal{D}_δ . Substituting the above estimate into (5.7), we conclude that

$$\begin{aligned} \frac{d}{dt} \left(\langle t \rangle^{\frac{5-\delta}{2}} \|\partial_x^m \mathcal{U}\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{5-\delta}{2}} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 \\ \leq 2 \langle t \rangle^{\frac{5-\delta}{2}} \left(\partial_x^{m+1} \mathcal{F}, \partial_x^m \mathcal{U} \right)_{L_\mu^2} + C\varepsilon \delta^{-1} \mathcal{D}_\delta. \end{aligned}$$

Similarly, we repeat the above argument for the evolution equation (5.3) of \mathcal{F} to conclude that

$$\begin{aligned} \frac{d}{dt} \left(\langle t \rangle^{\frac{5-\delta}{2}} \|\partial_x^m \mathcal{F}\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{5-\delta}{2}} \|\partial_x^m \partial_y \mathcal{F}\|_{L_\mu^2}^2 \\ \leq 2 \langle t \rangle^{\frac{5-\delta}{2}} \left(\partial_x^{m+1} \mathcal{U}, \partial_x^m \mathcal{F} \right)_{L_\mu^2} + C\varepsilon \delta^{-1} \mathcal{D}_\delta. \end{aligned}$$

Combining the two estimates above and observing the fact that

$$\left(\partial_x^{m+1} \mathcal{F}, \partial_x^m \mathcal{U} \right)_{L_\mu^2} + \left(\partial_x^{m+1} \mathcal{U}, \partial_x^m \mathcal{F} \right)_{L_\mu^2} = 0,$$

we obtain the assertion in Proposition 5.1. The proof of is thus completed. \square

Proposition 5.2. *Under the hypothesis of Proposition 5.1, we have*

$$\begin{aligned} \frac{d}{dt} \left(\frac{\delta}{2} \langle t \rangle^{\frac{7-\delta}{2}} \|(\partial_y \mathcal{U}, \partial_y \mathcal{F})\|_{H_\mu^{5,0}}^2 \right) + \frac{\delta^2}{2} \langle t \rangle^{\frac{7-\delta}{2}} \|(\partial_y^2 \mathcal{U}, \partial_y^2 \mathcal{F})\|_{H_\mu^{5,0}}^2 \\ \leq \frac{\delta}{2} \langle t \rangle^{\frac{5-\delta}{2}} \|(\partial_y \mathcal{U}, \partial_y \mathcal{F})\|_{H_\mu^{5,0}}^2 + C\varepsilon \delta^{-1} \mathcal{D}_\delta. \quad (5.13) \end{aligned}$$

Proof. For any $0 \leq m \leq 5$, applying ∂_y to equation (5.5) of $\partial_x^m \mathcal{U}$ yields

$$(\partial_t - \partial_y^2) \partial_x^m \partial_y \mathcal{U} + \frac{1}{\langle t \rangle} \partial_x^m \partial_y \mathcal{U} = \partial_x^{m+1} \partial_y \mathcal{F} + \partial_y R_{m,1} + \partial_y R_{m,2}, \quad (5.14)$$

recalling $R_{m,1}$ and $R_{m,2}$ are given in (5.6). By the boundary condition in (1.5), the trace at $y = 0$ of the source term in (5.2) is equal to 0. This with (5.4) yields that, taking $y = 0$ in equation (5.2),

$$\partial_y^2 \mathcal{U}|_{y=0} = 0 \quad \text{and} \quad R_{m,1}|_{y=0} = R_{m,2}|_{y=0} = 0.$$

Thus we take the L_μ^2 -product with $\partial_x^m \partial_y \mathcal{U}$ on both sides of (5.14) and then apply Remark 3.5 to get

$$\begin{aligned} & \frac{d}{dt} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 + \frac{5-\delta}{2\langle t \rangle} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 + \delta \|\partial_x^m \partial_y^2 \mathcal{U}\|_{L_\mu^2}^2 \\ & \leq 2 \left(\partial_x^{m+1} \partial_y \mathcal{F}, \partial_x^m \partial_y \mathcal{U} \right)_{L_\mu^2} + 2 \sum_{1 \leq j \leq 2} (\partial_y R_{m,j}, \partial_x^m \partial_y \mathcal{U})_{L_\mu^2}, \end{aligned}$$

or equivalently,

$$\begin{aligned} & \frac{d}{dt} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 + \frac{7-\delta}{2\langle t \rangle} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 + \delta \|\partial_x^m \partial_y^2 \mathcal{U}\|_{L_\mu^2}^2 \\ & \leq \frac{1}{\langle t \rangle} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 + 2 \left(\partial_x^{m+1} \partial_y \mathcal{F}, \partial_x^m \partial_y \mathcal{U} \right)_{L_\mu^2} + 2 \sum_{1 \leq j \leq 2} \|R_{m,j}\|_{L_\mu^2} \|\partial_x^m \partial_y^2 \mathcal{U}\|_{L_\mu^2}. \end{aligned}$$

Multiplying the above equation by $\langle t \rangle^{\frac{7-\delta}{2}}$ and observing

$$\frac{d}{dt} \langle t \rangle^{\frac{7-\delta}{2}} = \langle t \rangle^{\frac{7-\delta}{2}} \frac{7-\delta}{2\langle t \rangle},$$

we get

$$\begin{aligned} & \frac{d}{dt} \left(\langle t \rangle^{\frac{7-\delta}{2}} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{7-\delta}{2}} \|\partial_x^m \partial_y^2 \mathcal{U}\|_{L_\mu^2}^2 \leq \langle t \rangle^{\frac{5-\delta}{2}} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 \\ & + 2 \langle t \rangle^{\frac{7-\delta}{2}} \left(\partial_x^{m+1} \partial_y \mathcal{F}, \partial_x^m \partial_y \mathcal{U} \right)_{L_\mu^2} + 2 \langle t \rangle^{\frac{7-\delta}{2}} \sum_{1 \leq j \leq 2} \|R_{m,j}\|_{L_\mu^2} \|\partial_x^m \partial_y^2 \mathcal{U}\|_{L_\mu^2}. \end{aligned}$$

For the last term on the right-hand side, we use (5.8) and Lemma 2.4 as well as definition (2.11) of \mathcal{D}_δ , to conclude that, for each $m \leq 5$,

$$\begin{aligned} \langle t \rangle^{\frac{7-\delta}{2}} \sum_{1 \leq j \leq 2} \|R_{m,j}\|_{L_\mu^2} \|\partial_x^m \partial_y^2 \mathcal{U}\|_{L_\mu^2} & \leq C\varepsilon \langle t \rangle^{\frac{7-\delta}{2}} \langle t \rangle^{-\frac{1}{2}} \|(\partial_y \mathcal{U}, \partial_y \mathcal{F})\|_{H_\mu^{5,0}} \|\partial_y^2 \mathcal{U}\|_{H_\mu^{5,0}} \\ & \leq C\varepsilon \langle t \rangle^{\frac{7-\delta}{2}} \|(\partial_y^2 \mathcal{U}, \partial_y^2 \mathcal{F})\|_{H_\mu^{5,0}}^2 \leq C\varepsilon \delta^{-2} \mathcal{D}_\delta. \end{aligned}$$

Combining the two estimates above, we have, for any $0 \leq m \leq 5$,

$$\begin{aligned} & \frac{d}{dt} \left(\langle t \rangle^{\frac{7-\delta}{2}} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{7-\delta}{2}} \|\partial_x^m \partial_y^2 \mathcal{U}\|_{L_\mu^2}^2 \\ & \leq \langle t \rangle^{\frac{5-\delta}{2}} \|\partial_x^m \partial_y \mathcal{U}\|_{L_\mu^2}^2 + 2 \langle t \rangle^{\frac{7-\delta}{2}} \left(\partial_x^{m+1} \partial_y \mathcal{F}, \partial_x^m \partial_y \mathcal{U} \right)_{L_\mu^2} + C\varepsilon \delta^{-2} \mathcal{D}_\delta. \end{aligned}$$

Similarly, for any $0 \leq m \leq 5$ it holds that

$$\begin{aligned} \frac{d}{dt} \left(\langle t \rangle^{\frac{7-\delta}{2}} \|\partial_x^m \partial_y \mathcal{F}\|_{L_\mu^2}^2 \right) + \delta \langle t \rangle^{\frac{7-\delta}{2}} \|\partial_x^m \partial_y^2 \mathcal{F}\|_{L_\mu^2}^2 \\ \leq \langle t \rangle^{\frac{5-\delta}{2}} \|\partial_x^m \partial_y \mathcal{F}\|_{L_\mu^2}^2 + 2 \langle t \rangle^{\frac{7-\delta}{2}} \left(\partial_x^{m+1} \partial_y \mathcal{U}, \partial_x^m \partial_y \mathcal{F} \right)_{L_\mu^2} + C\varepsilon \delta^{-2} \mathcal{D}_\delta. \end{aligned}$$

As a result, we combine the above two estimates with the fact that

$$\left(\partial_x^{m+1} \partial_y \mathcal{F}, \partial_x^m \partial_y \mathcal{U} \right)_{L_\mu^2} + \left(\partial_x^{m+1} \partial_y \mathcal{U}, \partial_x^m \partial_y \mathcal{F} \right)_{L_\mu^2} = 0,$$

to obtain that

$$\begin{aligned} \frac{d}{dt} \left(\langle t \rangle^{\frac{7-\delta}{2}} \|(\partial_y \mathcal{U}, \partial_y \mathcal{F})\|_{H_\mu^{5,0}}^2 \right) + \delta \langle t \rangle^{\frac{7-\delta}{2}} \|(\partial_y^2 \mathcal{U}, \partial_y^2 \mathcal{F})\|_{H_\mu^{5,0}}^2 \\ \leq \langle t \rangle^{\frac{5-\delta}{2}} \|(\partial_y \mathcal{U}, \partial_y \mathcal{F})\|_{H_\mu^{5,0}}^2 + C\varepsilon \delta^{-2} \mathcal{D}_\delta. \end{aligned}$$

Multiplying the above estimate by $\frac{\delta}{2}$ yields the assertion in Proposition 5.2. The proof is thus completed. \square

Completing the proof of Theorem 2.2. We take summation of estimates (5.1) and (5.13) and observe the first term on the right-hand side of (5.13) can be absorbed by that on the left side of (5.1); this gives

$$\begin{aligned} \frac{d}{dt} \sum_{0 \leq k \leq 1} \left[\left(\frac{\delta}{2} \right)^k \langle t \rangle^{\frac{5-\delta}{2}+k} \|(\partial_y^k \mathcal{U}, \partial_y^k \mathcal{F})\|_{H_\mu^{5,0}}^2 \right] \\ + \sum_{0 \leq k \leq 1} \left(\frac{\delta}{2} \right)^{k+1} \langle t \rangle^{\frac{5-\delta}{2}+k} \|(\partial_y^{k+1} \mathcal{U}, \partial_y^{k+1} \mathcal{F})\|_{H_\mu^{5,0}}^2 \leq C\varepsilon \delta^{-1} \mathcal{D}_\delta. \end{aligned}$$

Moreover, by Proposition 4.1,

$$\frac{d}{dt} \sum_{m \leq 8} \langle t \rangle^{\frac{1-\delta}{2}} \|(u_m, f_m)\|_{L_\mu^2}^2 + \sum_{m \leq 8} \delta \langle t \rangle^{\frac{1-\delta}{2}} \|(\partial_y u_m, \partial_y f_m)\|_{L_\mu^2}^2 \leq C\varepsilon \delta^{-2} \mathcal{D}_\delta.$$

In view of definitions (2.10) and (2.11) of \mathcal{E}_δ and \mathcal{D}_δ , we combine the above estimates to conclude that

$$\forall t > 0, \quad \frac{d}{dt} \mathcal{E}_\delta(t) + \mathcal{D}_\delta(t) \leq C\varepsilon \delta^{-2} \mathcal{D}_\delta(t).$$

We choose ε small enough such that $C\varepsilon \delta^{-2} \leq \frac{1}{2}$. Then

$$\forall t > 0, \quad \mathcal{E}_\delta(t) + \frac{1}{2} \int_0^t \mathcal{D}_\delta(s) ds \leq \mathcal{E}_\delta(0) \leq \varepsilon^2.$$

The proof of Theorem 2.2 is thus completed. \square

Appendix A. Equations for \mathcal{U} and \mathcal{F}

This part is devoted to deriving the equations solved by \mathcal{U} and \mathcal{F} , recalling

$$\mathcal{U} = u - \frac{y}{2\langle t \rangle} \int_y^{+\infty} u d\tilde{y} \quad \text{and} \quad \mathcal{F} = f - \frac{y}{2\langle t \rangle} \int_y^{+\infty} f d\tilde{y}. \quad (\text{A.1})$$

To do so, integrating on $[y, +\infty)$ with respect to y for the first equation in (1.5), we get

$$(\partial_t - \partial_y^2) \int_y^{+\infty} u d\tilde{y} - \partial_x \int_y^{+\infty} f d\tilde{y} = \int_y^{+\infty} \left[(f\partial_x + g\partial_y)f - (u\partial_x + v\partial_y)u \right] d\tilde{y}.$$

Multiplying the above equation by $\frac{y}{2\langle t \rangle}$ and observing that

$$\begin{aligned} & \frac{y}{2\langle t \rangle} (\partial_t - \partial_y^2) \int_y^{+\infty} u d\tilde{y} \\ &= (\partial_t - \partial_y^2) \left(\frac{y}{2\langle t \rangle} \int_y^{+\infty} u d\tilde{y} \right) + \underbrace{\frac{y}{2\langle t \rangle^2} \int_y^{+\infty} u d\tilde{y} - \frac{1}{\langle t \rangle} u}_{=-\frac{1}{\langle t \rangle} u}, \end{aligned}$$

we get

$$\begin{aligned} & (\partial_t - \partial_y^2) \left(\frac{y}{2\langle t \rangle} \int_y^{+\infty} u d\tilde{y} \right) - \partial_x \left(\frac{y}{2\langle t \rangle} \int_y^{+\infty} f d\tilde{y} \right) - \frac{1}{\langle t \rangle} \mathcal{U} \\ &= \frac{y}{2\langle t \rangle} \int_y^{+\infty} \left[(f\partial_x + g\partial_y)f - (u\partial_x + v\partial_y)u \right] d\tilde{y}. \end{aligned}$$

Then in view of (A.1), we subtract the above equation by the first equation in (1.5) to get

$$\begin{aligned} (\partial_t - \partial_y^2) \mathcal{U} + \frac{1}{\langle t \rangle} \mathcal{U} - \partial_x \mathcal{F} &= -(u\partial_x + v\partial_y)u + (f\partial_x + g\partial_y)f \\ &+ \frac{y}{2\langle t \rangle} \int_y^{+\infty} \left[(u\partial_x + v\partial_y)u - (f\partial_x + g\partial_y)f \right] d\tilde{y}. \end{aligned}$$

In the same way, we have

$$\begin{aligned} (\partial_t - \partial_y^2) \mathcal{F} + \frac{1}{\langle t \rangle} \mathcal{F} - \partial_x \mathcal{U} &= -(u\partial_x + v\partial_y)f + (f\partial_x + g\partial_y)u \\ &+ \frac{y}{2\langle t \rangle} \int_y^{+\infty} \left[(u\partial_x + v\partial_y)f - (f\partial_x + g\partial_y)u \right] d\tilde{y}. \end{aligned}$$

Appendix B. Equations for u_m and f_m

We begin with the evolution equation solved by g . In view of (2.8), we may write

$$g = - \int_0^y \partial_x f d\tilde{y} = \int_y^\infty \partial_x f d\tilde{y} \quad \text{and} \quad v = - \int_0^y \partial_x u d\tilde{y} = \int_y^\infty \partial_x u d\tilde{y}.$$

Using integration by part yields

$$\int_y^{+\infty} v \partial_y f d\tilde{y} = \int_y^{+\infty} (\partial_x u) f d\tilde{y} - v f, \quad \int_y^{+\infty} g \partial_y u d\tilde{y} = \int_y^{+\infty} (\partial_x f) u d\tilde{y} - g u.$$

Then integrating the second equation in (1.5) over $(y, +\infty)$ with respect to the normal variable, we have

$$\partial_t \int_y^\infty f(\tilde{y}) d\tilde{y} - v f - \partial_y^2 \int_y^\infty f(\tilde{y}) d\tilde{y} = \partial_x \int_y^\infty u(\tilde{y}) d\tilde{y} - g u.$$

Applying ∂_x to the above equation yields

$$\partial_t g - (\partial_x v) f - v \partial_x f - \partial_y^2 g = \partial_x v - (\partial_x g) u - g \partial_x u.$$

This with the divergence-free condition yields

$$\partial_t g + u \partial_x g + v \partial_y g - \partial_y^2 g = (1 + f) \partial_x v - g \partial_x u. \quad (\text{B.1})$$

Applying ∂_x^{m-1} to equation (B.1) and using divergence conditions again, we have

$$\left(\partial_t + u \partial_x + v \partial_y - \partial_y^2 \right) \partial_x^{m-1} g = (1 + f) \partial_x^m v - g \partial_x^m u + I_m, \quad (\text{B.2})$$

where

$$\begin{aligned} I_m = & \sum_{k=1}^{m-1} \binom{m-1}{k} \left[(\partial_x^k f) \partial_x^{m-k} v - (\partial_x^k g) \partial_x^{m-k} u \right] \\ & - \sum_{k=1}^{m-1} \binom{m-1}{k} \left[(\partial_x^k u) \partial_x^{m-k} g - (\partial_x^k v) \partial_x^{m-k} f \right]. \end{aligned}$$

On the other hand, we apply ∂_x^m to first second equations in (1.5) to obtain

$$\begin{aligned} & \left(\partial_t + u \partial_x + v \partial_y - \partial_y^2 \right) \partial_x^m u - ((1 + f) \partial_x + g \partial_y) \partial_x^m f \\ & = -(\partial_x^m v) \partial_y u + (\partial_x^m g) \partial_y f + J_m \quad (\text{B.3}) \end{aligned}$$

with

$$\begin{aligned} J_m = & \sum_{k=1}^m \binom{m}{k} \left[(\partial_x^k f) \partial_x^{m-k+1} f - (\partial_x^k u) \partial_x^{m-k+1} u \right] \\ & + \sum_{k=1}^{m-1} \binom{m}{k} \left[(\partial_x^k g) \partial_x^{m-k} \partial_y f - (\partial_x^k v) \partial_x^{m-k} \partial_y u \right]. \end{aligned}$$

Multiplying equation (B.2) by $\frac{\partial_y u}{1+f}$ and taking summation with the equation (B.3) for $\partial_x^m u$, we obtain that, recalling $u_m = \partial_x^m u + \frac{\partial_y u}{1+f} \partial_x^{m-1} g$,

$$\begin{aligned} & \left(\partial_t + u \partial_x + v \partial_y - \partial_y^2 \right) u_m - ((1 + f) \partial_x + g \partial_y) \partial_x^m f \\ & = (\partial_x^m g) \partial_y f + J_m - \frac{g \partial_y u}{1+f} \partial_x^m u + \frac{\partial_y u}{1+f} I_m + K_m, \quad (\text{B.4}) \end{aligned}$$

where, denoting by $[P, Q]$ the commutator between two operators P and Q ,

$$\begin{aligned} K_m &= \left[\left(\partial_t + u\partial_x + v\partial_y - \partial_y^2 \right), \frac{\partial_y u}{1+f} \right] \partial_x^{m-1} g \\ &= \left[\partial_x \partial_y f + \frac{g\partial_y^2 f}{1+f} \right] \partial_x^{m-1} g - \left[\frac{(\partial_y u)\partial_x u}{1+f} + \frac{g(\partial_y u)^2}{(1+f)^2} \right] \partial_x^{m-1} g \\ &\quad + \left[2\frac{(\partial_y f)\partial_y^2 u}{(1+f)^2} - 2\frac{(\partial_y f)^2 \partial_y u}{(1+f)^3} \right] \partial_x^{m-1} g + 2\left(\partial_y \frac{\partial_y u}{1+f} \right) \partial_x^m f. \end{aligned}$$

Moreover, recalling $f_m = \partial_x^m f + \frac{\partial_y f}{1+f} \partial_x^{m-1} g$,

$$\begin{aligned} &-((1+f)\partial_x + g\partial_y)\partial_x^m f = -((1+f)\partial_x + g\partial_y)\left(f_m - \frac{\partial_y f}{1+f}\partial_x^{m-1} g\right) \\ &= -((1+f)\partial_x + g\partial_y)f_m + (\partial_y f)\partial_x^m g - \frac{g\partial_y f}{1+f}\partial_x^m f \\ &\quad + \left[\partial_x \partial_y f + \frac{g\partial_y^2 f}{1+f} \right] \partial_x^{m-1} g - \left[\frac{(\partial_y f)\partial_x f}{1+f} + \frac{g(\partial_y f)^2}{(1+f)^2} \right] \partial_x^{m-1} g. \end{aligned}$$

Substituting the above equation into (B.4) and observing the term $(\partial_x^m g)\partial_y f$ therein is cancelled out, we have

$$\begin{aligned} &\left(\partial_t + u\partial_x + v\partial_y - \partial_y^2 \right) u_m - ((1+f)\partial_x + g\partial_y) f_m \\ &= \frac{g\partial_y f}{1+f}\partial_x^m f - \frac{g\partial_y u}{1+f}\partial_x^m u + J_m + \frac{\partial_y u}{1+f}I_m + \tilde{K}_m, \end{aligned}$$

where

$$\begin{aligned} \tilde{K}_m &= \left[\frac{(\partial_y f)\partial_x f}{1+f} + \frac{g(\partial_y f)^2}{(1+f)^2} \right] \partial_x^{m-1} g - \left[\frac{(\partial_y u)\partial_x u}{1+f} + \frac{g(\partial_y u)^2}{(1+f)^2} \right] \partial_x^{m-1} g \\ &\quad + \left[2\frac{(\partial_y f)\partial_y^2 u}{(1+f)^2} - 2\frac{(\partial_y f)^2 \partial_y u}{(1+f)^3} \right] \partial_x^{m-1} g + 2\left(\partial_y \frac{\partial_y u}{1+f} \right) \partial_x^m f. \end{aligned}$$

Equivalently,

$$\left(\partial_t + u\partial_x + v\partial_y - \partial_y^2 \right) u_m - ((1+f)\partial_x + g\partial_y) f_m = \sum_{1 \leq j \leq 3} S_{m,j}$$

with

$$\begin{aligned} S_{m,1} = J_m &= \sum_{k=1}^m \binom{m}{k} \left[(\partial_x^k f)\partial_x^{m-k+1} f - (\partial_x^k u)\partial_x^{m-k+1} u \right] \\ &\quad + \sum_{k=1}^{m-1} \binom{m}{k} \left[(\partial_x^k g)\partial_x^{m-k}\partial_y f - (\partial_x^k v)\partial_x^{m-k}\partial_y u \right], \end{aligned} \tag{B.5}$$

and

$$S_{m,2} = \frac{\partial_y u}{1+f} I_m = \frac{\partial_y u}{1+f} \sum_{k=1}^{m-1} \binom{m-1}{k} [(\partial_x^k f) \partial_x^{m-k} v - (\partial_x^k g) \partial_x^{m-k} u] \\ - \frac{\partial_y u}{1+f} \sum_{k=1}^{m-1} \binom{m-1}{k} [(\partial_x^k u) \partial_x^{m-k} g - (\partial_x^k v) \partial_x^{m-k} f], \quad (\text{B.6})$$

and

$$S_{m,3} = \frac{g \partial_y f}{1+f} \partial_x^m f - \frac{g \partial_y u}{1+f} \partial_x^m u + \tilde{K}_m \\ = \left[\frac{g \partial_y f}{1+f} + 2 \partial_y \left(\frac{\partial_y u}{1+f} \right) \right] \partial_x^m f - \frac{g \partial_y u}{1+f} \partial_x^m u - 2 \frac{(\partial_y f)^2 \partial_y u}{(1+f)^3} \partial_x^{m-1} g \\ + \left[\frac{(\partial_y f) \partial_x f - (\partial_y u) \partial_x u}{1+f} + \frac{g (\partial_y f)^2 - g (\partial_y u)^2 + 2 (\partial_y f) \partial_y^2 u}{(1+f)^2} \right] \partial_x^{m-1} g. \quad (\text{B.7})$$

By similar computation to above, we have that

$$\left(\partial_t + u \partial_x + v \partial_y - \partial_y^2 \right) f_m - ((1+f) \partial_x + g \partial_y) u_m = \sum_{1 \leq j \leq 3} \tilde{S}_{m,j}$$

with

$$\tilde{S}_{m,1} = \sum_{k=1}^m \binom{m}{k} [(\partial_x^k f) \partial_x^{m-k+1} u - (\partial_x^k u) \partial_x^{m-k+1} f] \\ + \sum_{k=1}^{m-1} \binom{m}{k} [(\partial_x^k g) \partial_x^{m-k} \partial_y u - (\partial_x^k v) \partial_x^{m-k} \partial_y f], \quad (\text{B.8})$$

$$\tilde{S}_{m,2} = \sum_{k=1}^{m-1} \binom{m-1}{k} \frac{\partial_y f}{1+f} [(\partial_x^k f) \partial_x^{m-k} v - (\partial_x^k g) \partial_x^{m-k} u] \\ - \sum_{k=1}^{m-1} \binom{m-1}{k} \frac{\partial_y f}{1+f} [(\partial_x^k u) \partial_x^{m-k} g - (\partial_x^k v) \partial_x^{m-k} f], \quad (\text{B.9})$$

and

$$\tilde{S}_{m,3} = \left[\frac{g \partial_y u}{1+f} + 2 \partial_y \left(\frac{\partial_y f}{1+f} \right) \right] \partial_x^m f - \frac{g \partial_y f}{1+f} \partial_x^m u \\ + \left[\frac{(\partial_x u) \partial_y f - (\partial_y u) \partial_x f}{1+f} + \frac{2 (\partial_y f) \partial_y^2 f}{(1+f)^2} - \frac{2 (\partial_y f)^3}{(1+f)^3} \right] \partial_x^{m-1} g. \quad (\text{B.10})$$

Acknowledgements. The work was supported by NSF of China (Nos. 12325108, 12131017, 12221001), and the Natural Science Foundation of Hubei Province (No. 2019CFA007).

References

- [1] R. Alexandre, Y.-G. Wang, C.-J. Xu, and T. Yang. Well-posedness of the Prandtl equation in Sobolev spaces. *J. Amer. Math. Soc.*, 28(3):745–784, 2015.
- [2] D. Chen, Y. Wang, and Z. Zhang. Well-posedness of the linearized Prandtl equation around a non-monotonic shear flow. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 35(4):1119–1142, 2018.
- [3] C. Collot, T.-E. Ghoul, and N. Masmoudi. Singularities and unsteady separation for the inviscid two-dimensional Prandtl system. *Arch. Ration. Mech. Anal.*, 240(3):1349–1430, 2021.
- [4] A.-L. Dalibard and N. Masmoudi. Separation for the stationary Prandtl equation. *Publ. Math. Inst. Hautes Études Sci.*, 130:187–297, 2019.
- [5] H. Dietert and D. Gérard-Varet. Well-posedness of the Prandtl equations without any structural assumption. *Ann. PDE*, 5(1):Paper No. 8, 51, 2019.
- [6] W. E and B. Engquist. Blowup of solutions of the unsteady Prandtl’s equation. *Comm. Pure Appl. Math.*, 50(12):1287–1293, 1997.
- [7] D. Gérard-Varet and E. Dormy. On the ill-posedness of the Prandtl equation. *J. Amer. Math. Soc.*, 23(2):591–609, 2010.
- [8] D. Gérard-Varet and T. Nguyen. Remarks on the ill-posedness of the Prandtl equation. *Asymptot. Anal.*, 77(1-2):71–88, 2012.
- [9] D. Gérard-Varet and M. Prestipino. Formal derivation and stability analysis of boundary layer models in MHD. *Z. Angew. Math. Phys.*, 68(3):Paper No. 76, 16, 2017.
- [10] E. Grenier. On the nonlinear instability of Euler and Prandtl equations. *Comm. Pure Appl. Math.*, 53(9):1067–1091, 2000.
- [11] E. Grenier, Y. Guo, and T. T. Nguyen. Spectral instability of general symmetric shear flows in a two-dimensional channel. *Adv. Math.*, 292:52–110, 2016.
- [12] Y. Guo and T. Nguyen. A note on Prandtl boundary layers. *Comm. Pure Appl. Math.*, 64(10):1416–1438, 2011.
- [13] M. Ignatova and V. Vicol. Almost global existence for the Prandtl boundary layer equations. *Arch. Ration. Mech. Anal.*, 220(2):809–848, 2016.
- [14] I. Kukavica, N. Masmoudi, V. Vicol, and T. K. Wong. On the local well-posedness of the Prandtl and hydrostatic Euler equations with multiple monotonicity regions. *SIAM J. Math. Anal.*, 46(6):3865–3890, 2014.
- [15] I. Kukavica and V. Vicol. On the local existence of analytic solutions to the Prandtl boundary layer equations. *Commun. Math. Sci.*, 11(1):269–292, 2013.
- [16] I. Kukavica, V. Vicol, and F. Wang. The van Dommelen and Shen singularity in the Prandtl equations. *Adv. Math.*, 307:288–311, 2017.
- [17] S. Li and F. Xie. Global solvability of 2D MHD boundary layer equations in analytic function spaces. *J. Differential Equations*, 299:362–401, 2021.
- [18] W.-X. Li, N. Masmoudi, and T. Yang. Well-posedness in Gevrey function space for 3D Prandtl equations without structural assumption. *Comm. Pure Appl. Math.*,

- 75(8):1755–1797, 2022.
- [19] W.-X. Li, D. Wu, and C.-J. Xu. Gevrey class smoothing effect for the Prandtl equation. *SIAM J. Math. Anal.*, 48(3):1672–1726, 2016.
- [20] W.-X. Li and R. Xu. Well-posedness in Sobolev spaces of the two-dimensional MHD boundary layer equations without viscosity. *Electron. Res. Arch.*, 29(6):4243–4255, 2021.
- [21] W.-X. Li and T. Yang. Well-posedness of the MHD boundary layer system in Gevrey function space without structural assumption. *SIAM J. Math. Anal.*, 53(3):3236–3264, 2021.
- [22] C.-J. Liu, D. Wang, F. Xie, and T. Yang. Magnetic effects on the solvability of 2D MHD boundary layer equations without resistivity in Sobolev spaces. *J. Funct. Anal.*, 279(7):108637, 45, 2020.
- [23] C.-J. Liu, F. Xie, and T. Yang. MHD boundary layers theory in Sobolev spaces without monotonicity I: Well-posedness theory. *Comm. Pure Appl. Math.*, 72(1):63–121, 2019.
- [24] N. Liu and P. Zhang. Global small analytic solutions of MHD boundary layer equations. *J. Differential Equations*, 281:199–257, 2021.
- [25] N. Masmoudi and T. K. Wong. Local-in-time existence and uniqueness of solutions to the Prandtl equations by energy methods. *Comm. Pure Appl. Math.*, 68(10):1683–1741, 2015.
- [26] O. A. Oleinik and V. N. Samokhin. *Mathematical models in boundary layer theory*, volume 15 of *Applied Mathematics and Mathematical Computation*. Chapman & Hall/CRC, Boca Raton, FL, 1999.
- [27] M. Paicu and P. Zhang. Global existence and the decay of solutions to the Prandtl system with small analytic data. *Arch. Ration. Mech. Anal.*, 241(1):403–446, 2021.
- [28] M. Sammartino and R. E. Caflisch. Zero viscosity limit for analytic solutions, of the Navier-Stokes equation on a half-space. I. Existence for Euler and Prandtl equations. *Comm. Math. Phys.*, 192(2):433–461, 1998.
- [29] W. Shen, Y. Wang, and Z. Zhang. Boundary layer separation and local behavior for the steady Prandtl equation. *Adv. Math.*, 389:Paper No. 107896, 25, 2021.
- [30] C. Wang, Y. Wang, and P. Zhang. On the global small solution of 2-D Prandtl system with initial data in the optimal Gevrey class. *Adv. Math.*, 440:Paper No. 109517, 69, 2024.
- [31] Z. Xin and L. Zhang. On the global existence of solutions to the Prandtl’s system. *Adv. Math.*, 181(1):88–133, 2004.
- [32] C.-J. Xu and X. Zhang. Long time well-posedness of Prandtl equations in Sobolev space. *J. Differential Equations*, 263(12):8749–8803, 2017.
- [33] P. Zhang and Z. Zhang. Long time well-posedness of Prandtl system with small and analytic initial data. *J. Funct. Anal.*, 270(7):2591–2615, 2016.

(W.-X. Li) School of Mathematics and Statistics, & Hubei Key Laboratory of Computational Science, Wuhan University Wuhan 430072, China

E-mail address: wei-xi.li@whu.edu.cn

(Z. Xu) School of Mathematics and Statistics, Wuhan University, Wuhan 430072, China

E-mail address: xuzhan@whu.edu.cn

(A. Yang) Department of Mathematics, The Chinese University of Hong Kong, Shatin, Hong Kong

E-mail address: ayang@math.cuhk.edu.hk