

EXISTENCE OF AN EXTREMAL FUNCTION OF SOBOLEV CRITICAL EMBEDDING WITH AN α -HOMOGENEOUS WEIGHT

PETR GURKA AND DANIEL HAUER

ABSTRACT. In our previous publication [*Calc. Var. Partial Differential Equations*, 60(1):Paper No. 16, 27, 2021], we delved into examining a critical Sobolev-type embedding of a Sobolev weighted space into an exponential weighted Orlicz space. We specifically determined the optimal Moser-type constant for this embedding, utilizing the monomial weight introduced by Cabré and Ros-Oton [*J. Differential Equations*, 255(11):4312–4336, 2013]. Towards the conclusion of that paper, we pledged to explore the existence of an extremal function within this framework. In this current work, we not only provide a positive affirmation to this inquiry but extend it to a broader range of weights known as α -homogeneous weights.

1. INTRODUCTION

It is well-known that Trudinger-Moser type inequalities can be considered as the critical case of Sobolev embeddings. To be more precise, recall that the Sobolev space $W^{1,d}(\mathbb{R}^d)$, $d \in \mathbb{N}$, is embedded into the Lebesgue space $L^q(\mathbb{R}^d)$ for any $q \in [d, \infty)$, although $W^{1,d}(\mathbb{R}^d)$ is not embedded into $L^\infty(\mathbb{R}^d)$. It was proved in 1967 by Trudinger [14] that there is an embedding of this type of $W_0^{1,d}(\Omega)$ into the Orlicz space $L^\Phi(\Omega)$, with Ω being a bounded domain in \mathbb{R}^d , and the Young function $\Phi(x) = \exp(\alpha|x|^{d/(d-1)})$, $t > 0$, where α is a positive constant. However, this embedding was announced earlier by Yudovich [15] without a proof and, in a slightly weaker form, proved by Pokhozhaev [12]. In 1971 Moser [11] showed that if $\alpha \leq \alpha_d = d\omega_{d-1}^{1/(d-1)}$, where ω_{d-1} denotes the surface area of the unit sphere in \mathbb{R}^d , then

$$\sup_u \int_\Omega \exp(\alpha|x|^{d/(d-1)}) dx < \infty,$$

where the supremum is taken over all functions u from the unit ball of the space $W_0^{1,d}(\Omega)$. When $\alpha > \alpha_d$, this supremum becomes infinite.

A natural question arises as to whether there exists a function u such that the supremum is attained. The first step towards solving this problem was made in 1986 by Carleson and Chang [3], who showed that an extremal function u exists if Ω is a ball in \mathbb{R}^d . In 1988, Struwe [13] demonstrated that this result continues to hold for small perturbations of the ball in \mathbb{R}^2 , and in 1992, Flucher [4] extended the result to any bounded domain in \mathbb{R}^2 . Finally, in 1995, Lin [10] generalised the result to any dimension. Note that a key ingredient of his proof is the use of d -Green's functions, the singular solutions to the d -Laplacian.

In our previous paper, published in 2021 [5], we studied a weighted version of the Trudinger-Moser inequality and its relationship with the corresponding Sobolev-type embedding. This research was based on recent results by Cabré and Ros-Oton [1] and Lam [9], whose function spaces involved the so-called *monomial* weight. In our paper, we presented some weighted analogues of the classical results. Namely, we found the “optimal” Orlicz spaces (within the class of Orlicz spaces) of the corresponding critical Sobolev-type embedding and showed that this embedding is not compact. Moreover, we proved, that an embedding into any properly larger Orlicz space is compact, and we also derived a corresponding *concentrated compactness* principle of P.-L. Lions. However, we left the question of the existence of an extremal function unsolved. In this paper, we aim to address this gap, at least for the case where Ω is a ball in \mathbb{R}^d . We are working in a slightly more general context, as our weight is a *homogeneous* weight (more details about the weights can be found in Section 2).

2020 *Mathematics Subject Classification.* 46E35, 46E30, 35A23, 26D10.

Key words and phrases. Trudinger-Moser inequality, Moser constant, critical Sobolev embedding, Orlicz exponential space, α -homogeneous weight, monomial weight, extremal function.

The research of the first author was supported by grant No. GA23-04720S of the Czech Science Foundation; the second author's research was partially supported by the two Australian Research Council grant DP200101065 and DP220100067.

2. PRELIMINARIES

Weighted measure on an convex cone. We denote by Σ an open convex cone in \mathbb{R}^d , $d \in \mathbb{N}$, $d \geq 2$, with vertex at the origin. Following [2], $w: \overline{\Sigma} \rightarrow [0, \infty)$ is a nonnegative (not identically zero) continuous function that is α -homogeneous, $\alpha > 0$, and such that $w^{1/\alpha}$ is concave in Σ . Recall that w is α -homogeneous if

$$w(\kappa x) = \kappa^\alpha w(x) \quad \text{for any } x \in \overline{\Sigma} \text{ and all } \kappa > 0. \quad (2.1)$$

Throughout the paper μ denotes the weighted measure on Σ defined as

$$d\mu(x) = w(x) dx.$$

For future reference, we set

$$D = d + \alpha. \quad (2.2)$$

Important examples of α -homogeneous functions are the *monomial weights*. Namely, let $\Sigma = \Sigma_1 \cap \dots \cap \Sigma_k$, $k \in \{1, \dots, d\}$, where $\Sigma_j \neq \mathbb{R}^d$, $j = 1, \dots, k$, are open convex cones in \mathbb{R}^d with vertex at the origin. Given $A_1, \dots, A_k > 0$, the weight $w: \overline{\Sigma} \rightarrow [0, \infty)$ defined as

$$w(x) = \prod_{j=1}^k \text{dist}(x, \partial \Sigma_j)^{A_j}, \quad x \in \overline{\Sigma},$$

satisfies the α -homogeneity assumptions. For other interesting examples we refer the reader to [2, Section 2]. In particular, when $\Sigma_j = \{x \in \mathbb{R}^d; x_j > 0\}$, $j = 1, \dots, k$, the function (cf. also [1])

$$w(x) = x_1^{A_1} \dots x_k^{A_k}, \quad x \in \overline{\Sigma},$$

we considered in our previous paper [5].

Basic definitions.

Measure space. By the symbol (X, ν) we denote a measure space X with a nonnegative σ -finite measure ν . If Ω is a ν -measurable subset of X , we denote by $\nu(\Omega)$ the ν -measure of Ω , that is, $\nu(\Omega) = \int_{\Omega} d\nu(x)$. If $X = \mathbb{R}^d$ with its n -dimensional Lebesgue measure $d\nu(x) = dx$ and $M \subset \mathbb{R}^d$ is a Lebesgue measurable set in \mathbb{R}^d , we use the notation $|M| = \int_{\Omega} dx$.

Lebesgue space. If (X, ν) is a measure space, we denote by $L^p(\Omega, \nu)$, $p \in [1, \infty]$, the *Lebesgue space*, of all ν -measurable functions f on a ν -measurable set Ω in X , equipped with the norm

$$\|f\|_{p, \Omega, \nu} = \begin{cases} \left(\int_{\Omega} |f(x)|^p d\nu(x) \right)^{1/p} & \text{if } p < \infty, \\ \nu\text{-ess sup}_{x \in \Omega} |f(x)| & \text{if } p = \infty. \end{cases}$$

If $X = \mathbb{R}^d$ and ν is the d -dimensional Lebesgue measure, then we write $\|\cdot\|_{p, \Omega}$ instead of $\|\cdot\|_{p, \Omega, \nu}$. Moreover, we simply write $\|\cdot\|_{p, \nu}$ (or $\|\cdot\|_p$) when $\Omega = X$ (or $\Omega = \mathbb{R}^d$). For $p \in [1, \infty]$ we define the *Hölder conjugate* number $p' \in [1, \infty]$ by the equality $\frac{1}{p} + \frac{1}{p'} = 1$.

Sobolev space. Let Ω be a domain in a measure space (\mathbb{R}^d, ν) with a nonnegative Borel measure ν and let $p \in [1, \infty]$. The *Sobolev space* $W_0^{1,p}(\Omega, \nu)$ is defined as the closure of $C_c^\infty(\Omega)$ (the space of infinitely differentiable functions with compact supports in Ω) with respect to the norm

$$\|u\| = \|\nabla u\|_{p, \Omega, \nu},$$

where ∇u is the gradient of u and $|\nabla u|$ is its Euclidean length, that is,

$$\nabla u = \left(\frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n} \right), \quad |\nabla u| = \left(\sum_{j=1}^n (\partial u / \partial x_j)^2 \right)^{1/2}$$

(to simplify the notation we are writing $\|\nabla u\|_{p, \Omega, \nu}$ instead of $\| |\nabla u| \|_{p, \Omega, \nu}$).

Distribution function and rearrangements. For a measurable function f in Σ we define its *distribution function with respect to μ* as

$$f_{*\mu}(\tau) = \mu(\{x \in \Sigma : |f(x)| > \tau\}), \quad \tau > 0,$$

and its *nonincreasing rearrangement with respect to μ* as

$$f_{\mu}^*(t) = \inf\{\tau > 0 : f_{*\mu}(\tau) \leq t\}, \quad t > 0.$$

The function f_{μ}^{\star} defined as

$$f_{\mu}^{\star}(x) = f_{\mu}^*(C_D|x|^D), \quad x \in \mathbb{R}^d, \quad (2.3)$$

where

$$C_D = \mu(B_1 \cap \Sigma), \quad (2.4)$$

is the *radial rearrangement of f with respect to μ* . We use the notation

$$B_r = \{x \in \mathbb{R}^d : |x| < r\}, \quad r > 0.$$

Thanks to the α -homogeneity of w , we have

$$\mu(B_r \cap \Sigma) = C_D r^D, \quad r > 0.$$

Note that the function f_{μ}^{\star} is nonnegative and radially nonincreasing. Though it is defined on the whole \mathbb{R}^d , it depends only on function values of f in Σ . Furthermore, the functions f and f_{μ}^{\star} are equimeasurable with respect to μ , that is,

$$\mu(\{x \in \Sigma : |f(x)| > \tau\}) = \mu(\{x \in \Sigma : |f_{\mu}^{\star}(x)| > \tau\}), \quad \tau > 0.$$

Pólya-Szegő inequality. We need the following important result.

Theorem 2.1 ([6, Theorem 3.1]). *Let $1 \leq p < \infty$. For every function $u \in C_c^1(\mathbb{R}^d)$, its radial rearrangement u_{μ}^{\star} with respect to μ is locally absolutely continuous and*

$$\|\nabla u_{\mu}^{\star}\|_{p,\mu} \leq \|\nabla u\|_{p,\mu}. \quad (2.5)$$

In particular, $\text{supp } u_{\mu}^{\star} = \overline{B}_R$ where R is such that $\mu(B_R \cap \Sigma) = \mu(\text{supp } u \cap \Sigma)$.

We need another obvious result.

Lemma 2.2. *Let $\Psi : (0, \infty) \rightarrow (0, \infty)$ be an increasing and measurable function. Then*

$$\int_{\mathbb{R}^d} \Psi(u_{\mu}^{\star}(x)) \, d\mu(x) = \int_{\mathbb{R}^d} \Psi(|u(x)|) \, d\mu(x).$$

Reduction to the one-dimensional problem.

Proposition 2.3. *Let $u \in C_c^1(\mathbb{R}^d)$ be a nonnegative function with $\text{supp } u \subset \Omega$ such that $\mu(\Omega) < \infty$. Assume that $0 < a \leq c_D$ and $R > 0$ is the number satisfying $\mu(B_R \cap \Sigma) = \mu(\Omega \cap \Sigma)$. Let u_{μ}^{\star} be the radial rearrangement of u with respect to μ . Set*

$$U(|x|) = u_{\mu}^{\star}(x), \quad x \in \mathbb{R}^d \cap \Sigma \quad (2.6)$$

and

$$\varphi(t) = C_D^{1/D'} U(Re^{-t/D}), \quad t \in (0, \infty). \quad (2.7)$$

Then φ satisfies that $\varphi' \geq 0$ on $(0, \infty)$ and $\varphi(0) = 0$. Moreover,

$$\int_{\Omega \cap \Sigma} |\nabla u(x)|^D \, d\mu(x) \geq \int_{B_R \cap \Sigma} |\nabla u_{\mu}^{\star}(x)|^D \, d\mu(x) = \int_0^{\infty} (\varphi'(t))^D \, dt$$

and

$$\begin{aligned} & \frac{1}{\mu(\Omega \cap \Sigma)} \int_{\Omega \cap \Sigma} \exp(a(u(x))^{D'}) \, d\mu(x) \\ &= \frac{1}{\mu(B_R \cap \Sigma)} \int_{B_R \cap \Sigma} \exp(a(u_{\mu}^{\star}(x))^{D'}) \, d\mu(x) = \int_0^{\infty} \exp(\beta \varphi^{D'}(t) - t) \, dt, \end{aligned}$$

where $\beta = a/C_D \in (0, 1]$.

Proof. By Lemma 2.2 with $\Psi(t) = \exp(at^{D'})$ we have

$$\frac{1}{\mu(\Omega \cap \Sigma)} \int_{\Omega \cap \Sigma} \exp(a|u(x)|^{D'}) d\mu(x) = \frac{1}{\mu(B_R \cap \Sigma)} \int_{B_R \cap \Sigma} \exp(a(u^\star(x))^{D'}) d\mu(x). \quad (2.8)$$

Since u^\star is radially symmetric, we obtain

$$\begin{aligned} \int_{B_R \cap \Sigma} \exp(a(u^\star(x))^{D'}) d\mu(x) &= \int_0^R \left(\int_{x \in \Sigma, |x|=r} \exp(a(u^\star(x))^{D'}) w(x) d\sigma(x) \right) dr \\ &= \int_0^R \exp(a(U(r))^{D'}) \left(\int_{x \in \Sigma, |x|=r} w(x) d\sigma(x) \right) dr = P_w(B_1 \cap \Sigma) \int_0^R \exp(a(U(r))^{D'}) r^{D-1} dr, \end{aligned}$$

where $U(|x|) = u^\star(x)$ and $P_w(B_1 \cap \Sigma) = \int_{x \in \Sigma, |x|=1} w(x) d\sigma(x)$ (σ is the Hausdorff measure on the unit sphere centered at the origin). We make the change of variables

$$r = Re^{-t/D} \quad (2.9)$$

in the last integral to obtain

$$\int_0^R \exp(a(U(r))^{D'}) r^{D-1} dr = D^{-1} R^D \int_0^\infty \exp(a(U(Re^{-t/D}))^{D'} - t) dt.$$

Setting $\beta = a/C_D \in (0, 1]$ and using the fact, that

$$C_D = \mu(B_R \cap \Sigma) = \int_{B_R \cap \Sigma} w(x) dx = D^{-1} P_w(B_1 \cap \Sigma) R^D,$$

we arrive at

$$\frac{1}{\mu(B_R \cap \Sigma)} \int_{B_R \cap \Sigma} \exp(a(u^\star(x))^{D'}) d\mu(x) = \int_0^\infty \exp(\beta(\varphi(t))^{D'} - t) dt, \quad (2.10)$$

where the function φ is defined by (2.7). Similarly, since $|\nabla u^\star(x)| = |U'(|x|)|$, $x \in \mathbb{R}^d \cap \Sigma$, we have

$$\int_{B_R \cap \Sigma} |\nabla u^\star(x)|^D d\mu(x) = P_w(B_1 \cap \Sigma) \int_0^R |U'(r)|^D r^{D-1} dr.$$

Differentiating (2.7) we have, for all $t \in (0, \infty)$,

$$\varphi'(t) = -D^{-1} C_D^{1/D'} U'(Re^{-t/D}) Re^{-t/D} = -D^{-1} C_D^{1/D'} U'(r)r.$$

Observe that, since U is nonincreasing, the right hand side is nonnegative. Change of variables (2.9) gives

$$\begin{aligned} \int_0^R |U'(r)|^D r^{D-1} dr &= D^{-1} (D^{-1} C_D^{1/D'})^{-D} \int_0^\infty |D^{-1} C_D^{1/D'} U'(Re^{-t/D}) Re^{-t/D}|^D dt \\ &= D^{-1} (D^{-1} C_D^{1/D'})^{-D} \int_0^\infty |\varphi'(t)|^D dt. \end{aligned}$$

That is,

$$\int_{B_R \cap \Sigma} |\nabla u^\star(x)|^D d\mu(x) = P_w(B_1 \cap \Sigma) D^{-1} (D^{-1} C_D^{1/D'})^{-D} \int_0^\infty |\varphi'(t)|^D dt = \int_0^\infty |\varphi'(t)|^D dt.$$

The proof is completed. \square

Lemma 2.4. *Let $q \in (1, \infty)$. Consider nonnegative functions $\varphi \in C^1[0, \infty)$ such that $\varphi(0) = 0$, $\varphi' \geq 0$ on $(0, \infty)$ and $\int_0^\infty (\varphi'(t))^q dt \leq 1$. Then*

$$A(q) = \sup_{\varphi} \int_0^\infty \exp((\varphi(t))^{q'} - t) dt < \infty. \quad (2.11)$$

Proof. The proof for $q \geq 2$ was given by Moser [11], for $q \in (1, 2)$ it was extended by Jodeit [8]. \square

Lemma 2.5. *The supremum in (2.11) is attained for every $q > 1$.*

Proof. The result for $q \in \mathbb{N}$, $q \geq 2$ was proved first by Carleson and Chang [3], it was extended to real $q > 1$ by Hudson and Leckband [7]. \square

3. MAIN RESULT

Now we formulate the main theorem.

Theorem 3.1. *Assume that $p \in [1, \infty)$ and w is an α -homogeneous weight such that $\alpha > 0$. Then*

$$S = \sup_u \frac{1}{\mu(\Omega \cap \Sigma)} \int_{\Omega \cap \Sigma} \exp(a|u(x)|^{D'}) d\mu(x) < \infty,$$

where the supremum is taken over all $u \in C_c^1(\mathbb{R}^d)$ such that $\text{supp } u \subset \Omega$ with $\mu(\Omega) < \infty$ and $\|\nabla u\|_{p,\mu} \leq 1$. If $\Omega = B_R$ with some $R > 0$, then the supremum is attained.

Proof. To prove that S is finite we apply Proposition 2.3 to obtain

$$1 \geq \int_{\Omega \cap \Sigma} |\nabla u(x)|^D d\mu(x) \geq \int_0^\infty (\varphi'(t))^D dt,$$

and so, by Lemma 2.4 and Proposition 2.3,

$$\sup_u \frac{1}{\mu(\Omega \cap \Sigma)} \int_{\Omega \cap \Sigma} \exp(a|u(x)|^{D'}) d\mu(x) = \sup_\varphi \int_0^\infty \exp((\varphi(t))^{D'} - t) dt < \infty.$$

For proving the second part it is enough to take the function

$$u(x) = U(|x|), \quad x \in B_R \cap \Sigma,$$

where U is the function such that (2.7) holds and φ is the extremal function from Lemma 2.5 when $q = D$. \square

REFERENCES

- [1] X. Cabré and X. Ros-Oton. Sobolev and isoperimetric inequalities with monomial weights. *J. Differential Equations*, 255(11):4312–4336, 2013.
- [2] X. Cabré, X. Ros-Oton, and J. Serra. Sharp isoperimetric inequalities via the ABP method. *J. Eur. Math. Soc. (JEMS)*, 18(12):2971–2998, 2016.
- [3] L. Carleson and S.-Y. A. Chang. On the existence of an extremal function for an inequality of J. Moser. *Bull. Sci. Math. (2)*, 110(2):113–127, 1986.
- [4] M. Flucher. Extremal functions for the Trudinger-Moser inequality in 2 dimensions. *Comment. Math. Helv.*, 67(3):471–497, 1992.
- [5] P. Gurka and D. Hauer. More insights into the Trudinger-Moser inequality with monomial weight. *Calc. Var. Partial Differential Equations*, 60(1):Paper No. 16, 27, 2021.
- [6] P. Gurka, J. Lang, and Z. Mihula. Quantitative analysis of optimal sobolev-lorentz embeddings with α -homogeneous weights, 2023. preprint.
- [7] S. Hudson and M. Leckband. Extremals for a Moser-Jodeit exponential inequality. *Pacific J. Math.*, 206(1):113–128, 2002.
- [8] M. Jodeit, Jr. An inequality for the indefinite integral of a function in L^q . *Studia Math.*, 44:545–554, 1972. Collection of articles honoring the completion by Antoni Zygmund of 50 years of scientific activity, VI.
- [9] N. Lam. Sharp Trudinger-Moser inequalities with monomial weights. *NoDEA Nonlinear Differential Equations Appl.*, 24(4):Art. 39, 21, 2017.
- [10] K.-C. Lin. Extremal functions for Moser's inequality. *Trans. Amer. Math. Soc.*, 348(7):2663–2671, 1996.
- [11] J. Moser. A sharp form of an inequality by N. Trudinger. *Indiana Univ. Math. J.*, 20:1077–1092, 1970/71.
- [12] S. Pokhozhaev. Eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$. *Soviet Math. Doklady*, 6:1408–1411, 1965.
- [13] M. Struwe. Critical points of embeddings of $H_0^{1,n}$ into Orlicz spaces. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 5(5):425–464, 1988.
- [14] N. S. Trudinger. On imbeddings into Orlicz spaces and some applications. *J. Math. Mech.*, 17:473–483, 1967.
- [15] V. Yudovich. Some estimates connected with integral operators and with solutions of elliptic equations. *Soviet Math. Doklady*, 2:746–749, 1961.

P. GURKA, DEPARTMENT OF MATHEMATICS, CZECH UNIVERSITY OF LIFE SCIENCES PRAGUE, 165 21, PRAGUE 6, CZECH REPUBLIC;
DEPARTMENT OF MATHEMATICS, COLLEGE OF POLYTECHNICS JIHLAVA, TOLSTÉHO 16, 586 01, JIHLAVA, CZECH REPUBLIC; DEPARTMENT
OF MATHEMATICAL ANALYSIS, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83, 186 75 PRAHA 8,
CZECH REPUBLIC

Email address: gurka@tf.czu.cz

ORCID: 0000-0002-0995-4711

D. HAUER, BRANDENBURG UNIVERSITY OF TECHNOLOGY COTTBUS–SENFENBERG, PLATZ DER DEUTSCHEN EINHEIT 1, 03046
COTTBUS, GERMANY; SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF SYDNEY, SYDNEY, NSW, 2006, AUSTRALIA

Email address: daniel.hauer@b-tu.de

ORCID: 0000-0001-6210-7389