

# SECTION METHOD AND FRÉCHET POLYNOMIALS

Dan M. DĂIANU  
University "Politehnica" of Timisoara

September 18, 2024

*Tribute to Professor Borislav Crstici on the 100th anniversary of his birth*

## Abstract

Using the section method we characterize the solutions  $f : U \rightarrow Y$  of the following four equations

$$\sum_{i=0}^n (-1)^{n-i} \binom{n}{i} f \left( \sqrt[n]{u^m + iv^m} \right) = (n!) f(v),$$

$$f(u) + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} f \left( \sqrt[n]{u^m + iv^m} \right) = 0,$$

$$\sum_{i=0}^n (-1)^{n-i} \binom{n}{i} f \left( \arcsin |\sin u \sin^i v| \right) = (n!) f(v) \text{ and}$$

$$f(u) + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} f \left( \arcsin |\sin u \sin^i v| \right) = 0,$$

where  $m \geq 2$  and  $n$  are positive integers,  $U \subseteq \mathbb{R}$  is a maximally relevant real domain and  $(Y, +)$  is an  $(n!)$ -divisible Abelian group.<sup>1</sup>

## 1 Introduction

Educated at the Cluj school of functional equations, approximations and convexity founded by Academician Tiberiu Popoviciu, Professor Borislav Crstici had among his main concerns the functional characterization of polynomials and their generalizations. Without pretending exhaustiveness, we mention here Professor's thesis

<sup>1</sup>Mathematical Subject Classification(2010):39B52, 39A70

Keywords and phrases: *monomial, Fréchet polynomial, section method.*

dedicated to the functional equations that define polynomials [7] and the additions made in [18], [19] and [8]. A brief presentation of Professor's personality is given in [6]. In this context we mention the works [1], [2], [3], [4], [5], [9], [10], [11], [12], [13], [20] and [21], published only in the last decade and which contain some generalizations and analyses of different types of polynomials.

Let  $m, n$  be positive integers,  $m \geq 2$  and  $(Y, +)$  be an  $(n!)$ -divisible Abelian group - i.e. the group homomorphism  $Y \rightarrow Y, y \mapsto (n!)y$  is an isomorphism. This paper is dedicated to characterize the solutions  $f : \mathbb{R} \rightarrow Y$  of the equations

$$\sum_{i=0}^n (-1)^{n-i} \binom{n}{i} f \left( \sqrt[m]{u^m + iv^m} \right) = (n!) f(v) \text{ for all } u, v \in \mathbb{R}, \quad (1)$$

$$f(u) + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} f \left( \sqrt[m]{u^m + iv^m} \right) = 0 \text{ for all } u, v \in \mathbb{R}, \quad (2)$$

and the solutions  $f : U := \mathbb{R} \setminus \{k\pi \mid k \in \mathbb{Z}\} \rightarrow Y$  of the equations

$$\sum_{i=0}^n (-1)^{n-i} \binom{n}{i} f \left( \arcsin |\sin u \sin^i v| \right) = (n!) f(v) \text{ for all } u, v \in U, \quad (3)$$

$$f(u) + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} f \left( \arcsin |\sin u \sin^i v| \right) = 0 \text{ for all } u, v \in U. \quad (4)$$

The tool use for these characterizations is the section method [14], [15].

## 2 Framework

Everywhere in what follows  $(X, +)$  is a commutative semigroup,  $n$  is a positive integer and  $(Y, +)$  is an  $(n!)$ -divisible Abelian group. We denote by  $\mathcal{S}_i$  the set of the solutions of equation (i); for instance  $\mathcal{S}_1$  is the set of all functions  $f : \mathbb{R} \rightarrow Y$  that satisfy equation (1). Let  $j$  be a nonnegative integer; we will use the operator

$$\Delta_y^j : Y^X \rightarrow Y^X, \Delta_y^j \rho(x) := \sum_{i=0}^j (-1)^{j-i} \binom{j}{i} \rho(x + iy)$$

for  $y \in X$ ;  $\mathcal{M}_j(X, Y)$  denotes the  $j$ -monomials, i.e. the solutions  $\rho : X \rightarrow Y$  of the equation

$$\Delta_y^j \rho(x) = (j!) \rho(y) \text{ for all } x, y \in X$$

and  $\mathcal{P}_n(X, Y)$  denotes the (Fréchet)  $n$ -polynomials, i.e. the solutions  $\rho : X \rightarrow Y$  of the equation

$$\Delta_y^{n+1} \rho(x) = 0 \text{ for all } x, y \in X.$$

The first characterization of continuous real  $n$ -polynomials by this equation was realized by Fréchet in [17]. A detailed analysis of Fréchet polynomials in the present framework was given by Djoković in [16]; from this paper we will use only the following result.

**Lemma 2.1.** *Let  $\rho : X \rightarrow Y$ . Then  $\rho \in \mathcal{P}_n(X, Y)$  if and only if there exists  $\rho_j \in \mathcal{M}_j(X, Y)$  for all  $j \in \{0, 1, \dots, n\}$  such that  $\rho = \sum_{j=0}^n \rho_j$ .*

The section method [14], [15] provides, among other things, a technique for solving - partial or total - some equations whose solutions are composite functions. We will give below only a few rudiments of this method adapted to our goals.

Let  $g : U \rightarrow X$  be a surjection and  $g' : X \rightarrow U$  be a section of  $g$  (i.e.  $g \circ g' = id_X$ ). Let also the functions

$$G : Y^X \times X^2 \rightarrow Y, H : Y \times Y \rightarrow Y$$

and the equation

$$G(f \circ g', (g(u), g(v))) = H(f(u), f(v)) \text{ for all } u, v \in U, \quad (5)$$

where the unknown is  $f : U \rightarrow Y$ . The equation

$$G(\rho, (x, y)) = H(\rho(x), \rho(y)) \text{ for all } x, y \in X, \quad (6)$$

where  $\rho : X \rightarrow Y$  is the unknown, is *the characteristic* of equation (5).

We will use the following results extracted from Th. 2.4.1, Th 2.4.2 and Th. 2.6.6 in [14].

- Lemma 2.2.** 1.  $\{f \circ g' | f \in \mathcal{S}_5\} \subseteq \mathcal{S}_6$ .  
 2.  $\mathcal{S}_5^c := \{\rho \circ g | \rho \in \mathcal{S}_6\} \subseteq \mathcal{S}_5$ .  
 3. If  $f \in \mathcal{S}_5$  and  $u_0 \in U$  such that the function

$$f(U) \rightarrow Y, y \mapsto H(f(u_0), y) \text{ (or } y \mapsto H(y, f(u_0)))$$

is one-to-one, then  $f \in \mathcal{S}_5^c$ .

The functions in  $\mathcal{S}_5^c$  are named *canonical solutions* (of equation (5)). Thus Lemma 2.2.2 gives a partial solution for equation (5) and Lemma 2.2.3 provides sufficient conditions under which a solution of equation (5) is a canonical solution.

In the following we will use the notions and the conventions introduced above.

### 3 Radical-Fréchet equations

Let  $m \geq 2$  be an integer,  $U = \mathbb{R}$  and  $(X, +)$  be the additive semigroup defined by

$$(X, +) := \begin{cases} (\mathbb{R}, +) & \text{if } m \text{ is odd} \\ (\mathbb{R}_+, +) & \text{if } m \text{ is even} \end{cases},$$

where  $\mathbb{R}_+ := [0, \infty)$ .

First we characterize the solutions of the *radical-monomial* equation (1).

**Theorem 3.1.** *Let  $f : \mathbb{R} \rightarrow Y$  be a function. Then  $f$  is a solution of equation (1) if and only if there exists an  $n$ -monomial  $\rho \in \mathcal{M}_n(X, Y)$  such that*

$$f(u) = \rho(u^m) \text{ for all } u \in \mathbb{R}. \quad (7)$$

*Proof.* We apply the section method for the surjection

$$g : \mathbb{R} \rightarrow X, u \mapsto u^m,$$

its section

$$g' : X \rightarrow \mathbb{R}, x \mapsto \sqrt[m]{x},$$

and the functions

$$G : Y^X \times X^2 \rightarrow Y, G(\rho, (x, y)) := \Delta_y^n \rho(x),$$

$$H : Y \times Y \rightarrow Y, H(y_1, y_2) := (n!) y_2.$$

Then equation (1) becomes equation (5) and its characteristic is equation (6). Therefore the characteristic of equation (1) is exactly the  $n$ -monomial equation

$$\Delta_y^n \rho(x) = (n!) \rho(y) \text{ for all } x, y \in X$$

and all its solutions are in  $\mathcal{M}_n(X, Y)$ .

1. Let  $\rho \in \mathcal{M}_n(X, Y)$  and  $f : \mathbb{R} \rightarrow Y$  defined by (7). Then  $f = \rho \circ g$  and, according to Lemma 2.2.2,  $f$  is a solution of equation (1).

2. For proving the converse it suffices to show that  $\mathcal{S}_1 \subseteq \mathcal{S}_1^c$ . Let  $f$  be a solution of equation (1). Since  $(Y, +)$  is  $(n!)$ -divisible, the function

$$f(\mathbb{R}) \rightarrow Y, y \mapsto H(0, y) = (n!) y$$

is injective. According to Lemma 2.2.3,  $f$  is a canonical solution of equation (1), i.e. there exists  $\rho \in \mathcal{M}_n(X, Y)$  such that  $f(u) = \rho(u^m)$  for all  $u \in \mathbb{R}$ .  $\square$

Now we are in position to characterize the solutions of the *radical-Fréchet* equation (2).

**Theorem 3.2.** *Let  $f : \mathbb{R} \rightarrow Y$  be a function. Then  $f$  is a solution of equation (2) if and only if there exists  $\rho_i \in \mathcal{M}_i(X, Y)$  for all  $i \in \{0, 1, \dots, n\}$  such that*

$$f(u) = \rho_0(u^m) + \rho_1(u^m) + \dots + \rho_n(u^m) \text{ for all } u \in \mathbb{R}. \quad (8)$$

*Proof.* As in the proof of the previous theorem, let

$$g : \mathbb{R} \rightarrow X, g(u) := u^m \text{ and } g' : X \rightarrow \mathbb{R}, g'(x) := \sqrt[m]{x}.$$

Let also be the functions

$$G : Y^X \times X^2 \rightarrow Y, G(\rho, (x, y)) := \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} \rho(x + iy) \text{ and}$$

$$H : Y \times Y \rightarrow Y, H(y_1, y_2) := -y_1.$$

Then equation (2) can be written in the form (5) and its characteristic is (6) or, equivalent, the Fréchet polynomial equation

$$\Delta_y^{n+1} \rho(x) = 0 \text{ for all } x, y \in X.$$

We note that the solutions of the last equation are given by  $\mathcal{P}_n(X, Y)$  and their characterization is given by Lemma 2.1.

1. Let  $\rho_i \in \mathcal{M}_i(X, Y)$  for all  $i \in \{0, 1, \dots, n\}$  and  $f : \mathbb{R} \rightarrow Y$  defined by (8). Then  $\rho := \sum_{i=0}^n \rho_i \in \mathcal{P}_n(X, Y)$  (by Lemma 2.1) and  $f = \rho \circ g \in \mathcal{S}_2$  (by Lemma 2.2.2).

2. Let  $f \in \mathcal{S}_2$ . To show that  $f$  can be expressed by (8) with  $\rho_i \in \mathcal{M}_i(X, Y)$  - or, equivalent, that  $f = \rho \circ g$ , where  $\rho := \sum_{i=0}^n \rho_i \in \mathcal{P}_n(X, Y)$  -, it is sufficient to prove that  $f$  is a canonical solution of equation (2). But the function

$$f(\mathbb{R}) \rightarrow Y, y \mapsto H(y, 0) = -y$$

is injective; according to Lemma 2.2.3,  $f \in \mathcal{S}_2^c$ , and the theorem is completely proved.  $\square$

## 4 Arcsine-Fréchet equations

Before proceeding to the characterizations of the solutions of the *arcsine-Fréchet* equations (3) and (4), let us note that  $U := \mathbb{R} \setminus \{k\pi | k \in \mathbb{Z}\}$  is the maximal domain on which these equations has nontrivial solutions; indeed if there is  $k \in \mathbb{Z}$  such that  $k\pi$  is in the domain, for  $u = k\pi$  in (3) we get  $0 = (n!) f(v)$ , and, since  $(Y, +)$  is  $(n!)$ -divisible we immediately obtain  $f = 0$ , i.e.  $\mathcal{S}_3 = \{0\}$ ; analogously, for  $v = k\pi$  in (4) we get  $f = 0$  and  $\mathcal{S}_4 = \{0\}$ .

In the following lines, the set  $X := (-\infty, 0]$  is endowed with the addition of real numbers, hence  $(X, +)$  is an Abelian semigroup.

**Theorem 4.1.** *Let  $f : U := \mathbb{R} \setminus \{k\pi | k \in \mathbb{Z}\} \rightarrow Y$ . Then  $f \in \mathcal{S}_3$  if and only if there exists an  $n$ -monomial  $\rho \in \mathcal{M}_n(X, Y)$  and*

$$f(u) = \rho(\ln |\sin u|) \text{ for all } u \in U. \quad (9)$$

*Proof.* We apply the section method for

$$g : U \rightarrow X, g(u) := \ln |\sin u| \text{ for all } u \in U,$$

its section

$$g' : X \rightarrow U, g'(x) := \arcsin e^x \text{ for all } x \in X,$$

and the functions  $G, H$  defined by

$$G : Y^X \times X^2 \rightarrow Y, G(\rho, (x, y)) := \Delta_y^n \rho(x),$$

$$H : Y \times Y \rightarrow Y, H(y_1, y_2) := (n!) y_2.$$

We note that equation (3) becomes equation (5) and its characteristic is equation (6) or, the monomial equation

$$\Delta_y^n \rho(x) = (n!) \rho(y) \text{ for all } x, y \in X.$$

1. If  $\rho \in \mathcal{M}_n(X, Y)$  and  $f : U \rightarrow Y$  is defined by (9), then  $f = \rho \circ g$  and - from Lemma 2.2.2 -  $f$  is a solution of equation (3).

2. Let  $f \in \mathcal{S}_3$  and  $u_0 \in U$ . Since  $(Y, +)$  is  $(n!)$ -divisible, the function

$$f(U) \rightarrow Y, y \mapsto H(f(u_0), y) = (n!) y$$

is an injection and, from Lemma 2.2.3, there exists an  $n$ -monomial  $\rho \in \mathcal{M}_n(X, Y)$  such that  $f = \rho \circ g$ ; therefore  $f$  satisfies relation (9).  $\square$

Finally we characterize the solutions of equation (4).

**Theorem 4.2.** *Let  $f : U \rightarrow Y$  be a function. Then  $f$  is a solution of equation (4) if and only if there exists the monomials  $\rho_i \in \mathcal{M}_i(X, Y)$  for  $i \in \{0, 1, \dots, n\}$  such that*

$$f(u) = \sum_{i=0}^n \rho_i(\ln |\sin u|) \text{ for all } u \in U. \quad (10)$$

*Proof.* Let

$$g : U \rightarrow X, g(u) := \ln |\sin u| \text{ for all } u \in U,$$

$$g' : X \rightarrow U, g'(x) := \arcsin e^x \text{ for all } x \in X,$$

$$G : Y^X \times X^2 \rightarrow Y, G(\rho, (x, y)) := \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} \rho(x + iy) \text{ and}$$

$$H : Y \times Y \rightarrow Y, H(y_1, y_2) := -y_1.$$

We note that equation (4) can be rewritten in the form (5) and, consequently, its characteristic is equation (6) or, equivalent, the Fréchet polynomial equation

$$\Delta_y^{n+1} \rho(x) = 0 \text{ for all } x, y \in X.$$

Then, according to Lemma 2.2.2,  $\mathcal{S}_4^c \subseteq \mathcal{S}_4$ . Moreover, if  $f \in \mathcal{S}_4$  and  $u_0$  is an arbitrary number in  $U$ , the function

$$f(U) \rightarrow Y, y \mapsto H(y, f(u_0)) = -y$$

is bijective; from Lemma 2.2.3 we have  $f \in \mathcal{S}_4^c$ . Hence

$$\mathcal{S}_4 = \{\rho \circ g \mid \rho \in \mathcal{P}_n(X, Y)\},$$

and, by Lemma 2.1,

$$\mathcal{P}_n(X, Y) = \mathcal{M}_0(X, Y) + \mathcal{M}_1(X, Y) + \cdots + \mathcal{M}_n(X, Y).$$

Consequently, if  $f : U \rightarrow Y$ , then  $f \in \mathcal{S}_4$  if and only if there exist  $\rho_i \in \mathcal{M}_i(X, Y)$  for  $i \in \{0, 1, \dots, n\}$  such that  $f(u) = \sum_{i=0}^n \rho_i((\ln |\sin u|))$  for all  $u \in U$ , and the theorem is completely proved.  $\square$

## References

- [1] A.G. Aksoy, J.M. Almira, On Montel and Montel-Popoviciu theorems in several variables, *Aequat. Math.*, 89, (2015), 1335–1357.
- [2] J.M. Almira, Characterization of polynomials as solutions of certain functional equations, *J. Math. An. Appl.* 459(2), (2018), 1016–1028.
- [3] J.M. Almira, Using Aichinger’s equation to characterize polynomial functions, *Aequat. Math.*, 97, (2023), 899–907.
- [4] J.M. Almira, L. Székelyhidi, Local polynomials and the Montel theorem, *Aequat. Math.*, 89, (2015), 329–338.
- [5] J.M. Almira, L. Székelyhidi, Montel–Type Theorems for Exponential Polynomials, *Demonstratio Mathematica*, 49(2), (2016), 197–212.
- [6] G. Cristescu, D.M. Dăianu, Professor Borislav Crstici (1924–2014), the right man in the right place at the right time, *Ann. Tiberiu Popoviciu Semin. Funct. Equ. Approx. Convexity*, 12, (2014), 23–32.

- [7] B. Crstici, On the functional equations that define polynomials (in Romanian), thesis, "Babeş-Bolyai" University, Cluj-Napoca (1969).
- [8] B. Crstici, Sur certains equations fonctionnelle de T. Popoviciu pour la caracterisation des polynomes, *Ann. Tiberiu Popoviciu Semin. Funct. Equ. Approx. Convexity*, (1996), 41-47.
- [9] D.M. Dăianu, Recursive procedure in the stability of Fréchet polynomials, *Adv. Difference Equ.* Vol. 2014(16), (2014), 1-19.
- [10] D.M. Dăianu, A stability criterion for Fréchet's first polynomial equation, *Aequat. Math.* 88, (2014), 233–241.
- [11] D.M. Dăianu, Taylor type formula with Fréchet polynomials, *Aequat. Math.*, 92(3), (2018), 695–707.
- [12] D.M. Dăianu, Fixed Point Approach to the Stability of Generalized Polynomials, *Fixed Point Theory* 20(1), (2019), 135-156.
- [13] D.M. Dăianu, C. Mîndruța, Polynomials of Arithmetically Homogeneous Functions: Stability and Hyperstability, *Result Math.* 74(1), (2019), 1-31.
- [14] D.M. Dăianu, C. Mîndruța, Sections in functional equations, *Aequat. Math.*, 94(2), (2020), 931–943.
- [15] D.M. Dăianu, Sections in functional equations: stability and hyperstability, *Aequat. Math.* 96, (2022), 867–883.
- [16] D. Djoković, A representation theorem for  $(X_1 - 1)(X_2 - 1) \dots (X_n - 1)$  and its applications, *Ann. Pol. Math.* 22, (1969), 189-198.
- [17] M. Fréchet, Une definition fonctionnelle des polynômes, *Nouv. Ann. Math.* 9, (1909), 145-182.
- [18] P.L. Kannappan, B. Crstici, Two functional identities characterizing polynomials, *Ann. Tiberiu Popoviciu Semin. Funct. Equ. Approx. Convexity*, (1989), 175-180.
- [19] M. Neagu, B. Crstici, About a Functional Characterization of Polynomials, *Bul. Ştiinţ. Univ. Politeh. Timiş. Ser. Mat.Fiz.*, (1987), 59-62.
- [20] L. Székelyhidi, Characterization of exponential polynomials on commutative hypergroups, *Ann. Funct. Anal.* 5(2), (2014), 53-60.
- [21] I.V. Tikhonov, V.B. Sherstyukov, D.G. Tsvetkovich, Generalized Popoviciu Expansions for Bernstein Polynomials of a Rational Module, *J. Math. Sci.* 263, (2022), 532–578.



Dan M. Daianu – Department of Mathematics,  
'Politehnica' University of Timișoara,  
P-ta Victoriei 2, 300 006, Timișoara, ROMANIA  
E-mail: dan,daiianu.m@gmail.com