POSITIVE SOLUTIONS OF SINGULAR ELLIPTIC PROBLEMS WITH UNBOUNDED EXPONENTS AND UNBOUNDED CONVECTION TERM

ANDERSON L. A. DE ARAUJO, HAMILTON P. BUENO, AND KAMILA F. L. MADALENA

ABSTRACT. In this paper, we study the existence of a solution for a class of Dirichlet problems with a singularity and a convection term. Precisely, we consider the existence of a positive solution to the Dirichlet problem

$$-\Delta_p u = \frac{\lambda}{u^{\alpha}} + f(x, u, \nabla u)$$

in a bounded, smooth domain Ω . The convection term has exponents with no upper limitations neither in *u* nor in ∇u . This is somewhat unexpected and rare. So, we address a wide range of problems not yet contained in the literature.

The solution of the problem combines the definition of an auxiliary problem, the method of sub- and super-solution and Schauder's fixed point theorem.

1. INTRODUCTION

In this paper, we shall establish results on the existence of positive solutions to the following *p*-Laplacian problem with a singular nonlinearity and a nonlinear convection term.

(1)
$$\begin{cases} -\Delta_p u = \frac{\lambda}{u^{\alpha}} + f(x, u, \nabla u) & \text{in } \Omega \\ u > 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

where $-\Delta_p u = -\operatorname{div}(|\nabla u|^{p-2}\nabla u)$ is the *p*-Laplacian operator for 1 , $<math>\Omega \subset \mathbb{R}^N \ (N \ge 2)$ is a bounded, smooth domain, $0 < \alpha < 1$, and $f: \Omega \times \mathbb{R} \times \mathbb{R}^N \to \mathbb{R}$ is a continuous nonlinearity satisfying

(2)
$$0 \leq f(x,t,\xi) \leq a|t|^{r_1} + b|\xi|^{r_2} \quad a,b > 0.$$

where $\lambda > 0$ is a parameter and $r_1, r_2 \in (0, p-1) \cup (p-1, \infty)$.

Note that right-hand side of equation (1) presents two main difficulties. One is the presence of the singular term, which makes it unbounded, and therefore we cannot directly use the techniques applied in Bueno and Ercole [5], see a brief review of this paper in the sequel. Another difficulty is that the exponents r_1 of u and r_2 of ∇u are allowed to be supercritical and imposes severe restrictions on the use of techniques due to the lack of embedding of Sobolev spaces into suitable spaces.

In the past years, singular elliptic equations have been challenging mathematicians. An extensive literature is devoted to such problems with singularity, specially from the point of view of theoretical analysis. The following problem has

²⁰²⁰ Mathematics Subject Classification. 35J92, 35B09, 35B33, 35J25, 35J62.

Key words and phrases. quasilinear elliptic equation, positive solution, convection growth, supercritical growth.

First author was supported by FAPEMIG APQ-02375-21, RED-00133-21 and CNPq.

been studied in many papers

(3)
$$\begin{cases} -\Delta_p u = \eta(x)u^{-\alpha} & \text{in } \Omega\\ u > 0 & \text{in } \Omega\\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\eta(x) \ge 0$ in Ω and $0 < \alpha < 1$. In [6], Canino, Sciunzi and Trombetta showed the existence and uniqueness of solutions to (3) when $p \ne 2$ in many configurations.

In [15], Giacomoni, Schindler and Takáč studied the existence and multiplicity of weak solutions to the *p*-Laplacian problem

$$\begin{cases} -\Delta_p u &= \lambda u^{-\delta} + u^s & \text{in } \Omega \\ u &> 0 & \text{in } \Omega \\ u &= 0 & \text{on } \partial\Omega \end{cases}$$

where 1 0, and $0 < \delta < 1$. As usual, $p^* = \frac{Np}{N-p}$ if 1 is arbitrarily large if <math>p = N, and $p^* = \infty$ if p > N.

Also, singular problems driven by more general operators are also in the literature. For example,

$$\begin{cases} -\operatorname{div}\left(a(|\nabla u|^p)|\nabla u|^{p-2}\nabla u\right) &= u^{-\alpha} + u^{\beta} \quad \text{in } \Omega\\ u &> 0 \qquad \text{in } \Omega\\ u &= 0 \qquad \text{on } \partial\Omega, \end{cases}$$

was considered by Corrêa, Corrêa and Figueiredo in [8] for a bounded, smooth domain $\Omega \subset \mathbb{R}^N$ ($N \ge 3$), exponents $0 < \alpha, \beta < p - 1$ and a quite general function $a: \mathbb{R}^+ \to \mathbb{R}^+$, which allows the authors to consider a wide range of problems, including the *p*-Laplacian operator Δ_p and the (p, q)-Laplacian among others.

More details on topics related to singular problems without convection terms can be found in [3, 4, 8, 14, 21] and references therein.

Elliptic problems with convection terms also have been considered in various frameworks. In [5], using the method of sub- and super-solutions combined with a global estimate on the gradient, Bueno and Ercole prove the existence of at least one positive solution for the problem

$$\begin{cases} -\Delta_p u &= \beta f(x, u, \nabla u) + \lambda h(x, u) & \text{in } \Omega \\ u &= 0 & \text{on } \partial\Omega, \end{cases}$$

where h, f are continuous nonlinearities satisfying $0 \leq \omega_1(x)u^{q-1} \leq h(x, u) \leq \omega_2(x)u^{q-1}$ with 1 < q < p and $0 \leq f(x, u, \nabla u) \leq \omega_3(x)u^a |v|^b$ with a, b > 0 and the functions $\omega_i, 1 \leq i \leq 3$, are positive, continuous weights in $\overline{\Omega}$. It is important to note that the results presented in [5] are rare in the literature when the exponents of u and $|\nabla u|$ are greater than p - 1.

In [13], Faria, Miyagaki and Motreanu proved existence of a positive solution for the following quasi-linear elliptic problem involving the (p, q)-Laplacian and a convection term by aplying the Galerkin's method and a Schauder basis.

$$\begin{cases} -\Delta_p u - \mu \Delta_q u &= f(x, u, \nabla u) & \text{in } \Omega \\ u &> 0 & \text{in } \Omega \\ u &= 0 & \text{on } \partial\Omega, \end{cases}$$

In [19], Liu, Motreanu, and Zeng, using the method of sub- and super-solution, truncation techniques, nonlinear regularity theory, Leray-Schauder alternative principle, and set-valued analysis, established a result of existence of a positive solution to the problem

$$\begin{cases} -\Delta_p u &= f(x, u(x), \nabla u(x)) + g(x, u(x)) & \text{in } \Omega \\ u &> 0 & \text{in } \Omega \\ u &= 0 & \text{on } \partial\Omega, \end{cases}$$

where $f: \Omega \times \mathbb{R} \times \mathbb{R}^N \to \mathbb{R}$ satisfies a suitable growth condition and the semilinear function $g: \Omega \times (0, \infty) \to \mathbb{R}$ is singular at s = 0, that is, $\lim_{s \to 0^+} g(x, s) = +\infty$. It is noteworthy that the conditions (2) in this paper are significantly less restrictive than those on f in [19].

We state the main results of the paper. First we consider the case $0 < r_1, r_2 < p - 1$.

Theorem 1.1. Suppose f is a continuous function satisfying (2) such that $r_1, r_2 .$ $Then the problem (1) has at least one positive solution <math>u \in W_0^{1,p}(\Omega)$ for each $\lambda > 0$.

In order to handle the case $p-1 < r_1, r_2$ we need to assume a stronger assumption on the regularity of Ω : either $\partial \Omega \in C^{1,1}$ or Ω convex and $\partial \Omega \in C^{1,\tau}$, $0 < \tau < 1$. In either case we are allowed to apply Propositions 2.3 and 2.4 (stated in Section 2), which demand $\partial \Omega \in W^2 L^{N-1,1}$, where $W^2 L^{N-1,1}$ denotes the Lorentz-Sobolev space. As remarked by Cianchi and Maz'ya in [7], the assumption $\partial \Omega \in W^2 L^{N-1,1}$ means that Ω is locally the subgraph of a function of N-1 variables whose second-order weak derivatives belong to the Lorentz space $L^{N-1,1}$ (see Section 2 for basic results about this space) and $\partial \Omega \in C^{1,0}$.

So, with stronger hypotheses on $\partial \Omega$, we obtain the next result.

Theorem 1.2. Let $\Omega \subset \mathbb{R}^N$ be a bounded domain such that $\partial \Omega \in C^{1,1}$ and $q > \max\{N, p'\}$, where $p' = \frac{p}{p-1}$. Suppose that f is a continuous function satisfying (2), $0 < \alpha q < 1$ and $r_1, r_2 > p - 1$. Then there is a constant $A^* > 0$ such that, for each $\lambda \in (0, A^*)$, there is $M := M(\lambda)$ such that the problem (1) has at least one positive solution u_λ satisfying

$$\lambda u_0 \leq u_\lambda \leq M u_0$$
 and $\|\nabla u_\lambda\|_{\infty} \leq M$.

As a consequence of this result, we can obtain that, $u_{\lambda} \to 0$ and $|\nabla u_{\lambda}| \to 0$, uniformly in Ω , as $\lambda \to 0^+$, see the last part of the proof of Theorem 1.2.

We now briefly describe the plan of the paper. We collect a few results on the *p*-Laplacian operator in Section 2. In Section 3 we define an auxiliary problem, which will be useful in the proof of Theorem 1.1. The existence of a positive solution to the auxiliary problem is obtained in Lemma 3.2 by using the Galerkin method and a Schauder basis in $W_0^{1,p}(\Omega)$ combined with a consequence of Brouwer's fixed point theorem. The proof of Theorem 1.1 is then obtained by showing that the solution of the auxiliary problem converges to the solution of problem (1), using that $W_0^{1,p}(\Omega)$ is a reflexive space and the embedding $W_0^{1,p}(\Omega) \hookrightarrow L^p(\Omega)$. The comparison principle (see Lemma 2.2) and the solution u_0 of problem (8) (see below) guarantee the positivity of the solution of problem (1).

The proof of Theorem 1.2 is presented in Section 4 by applying Schauder's fixed point theorem. The main tools are Proposition 2.6, where we show the existence

of suitable λ and M to use the comparison principle combined with the solution u_0 of problem (8). To show the L^{∞} estimates on the gradient we apply Proposition 2.4, established in [12].

2. PRELIMINARIES

Let $W_0^{1,p}(\Omega)$ stand for the usual Sobolev space endowed with the norm $||u|| = ||\nabla u||_p$. We denote $p^* = \frac{Np}{N-p}$ if 1 . In the case <math>p = N, then $p^* \in (p, \infty)$ can be taken arbitrarily large and $p^* = \infty$ if p > N.

Definition 2.1. We say that $u \in W_0^{1,p}(\Omega)$ is a solution (respectively, sub-solution and super-solution) of (1) if $g_{\lambda}(\cdot, u, \nabla u) := \lambda u^{-\alpha} + f(x, u, \nabla u) \in L^p(\Omega)$,

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \varphi dx - \int_{\Omega} g_{\lambda}(x, u, \nabla u) \varphi dx = 0 \ (\leqslant 0, \ge 0)$$

for all $\varphi \in W_0^{1,p}(\Omega)$, $\varphi \ge 0$ and

$$u = 0$$
 (resp. $\leq 0, \geq 0$) on $\partial \Omega$.

The condition on $\partial \Omega$ is understood in $W^{1-1/p,p}(\partial \Omega)$, i.e, in the sense of traces.

By an ordered pair of sub- and super-solutions we mean a sub-solution α and a super-solution β such that $\alpha \leq \beta$ a.e.

It is well-known that the equation $-\Delta_p u = v$ in Ω with Dirichlet boundary condition has a unique weak solution $u \in C^{1,\sigma}(\overline{\Omega})$ for some $\sigma \in (0,1)$ if $v \in L^{\infty}(\Omega)$, see [10, 18, 24]. Moreover, if $v \ge 0$ and $v \ne 0$, then u > 0. Hence $\partial u/\partial \eta < 0$ on $\partial \Omega$, where η is the unit normal vector to $\partial \Omega$ pointing outward of Ω . Furthermore, u is bounded from above and from below by positive multiples of the distance function dist $(x, \partial \Omega)$.

Additionally, the associated solution operator $(-\Delta_p)^{-1} : L^{\infty}(\Omega) \to C^1(\overline{\Omega})$ is positive, continuous and compact. In synthesis, $(-\Delta_p)^{-1}$ can be viewed as a strongly positive operator on $C(\overline{\Omega})$, i.e., $v \in P$ implies $(-\Delta_p)^{-1}v \in int(P)$, where P denotes the cone of positive functions belonging $C(\overline{\Omega})$.

The following comparison result is true in the case of a bounded domain $\Omega \subset \mathbb{R}^N$ and $u, v \in W^{1,p}(\Omega) \cap C(\overline{\Omega})$, $1 . If <math>-\Delta_p u \leq -\Delta_p v$ on Ω in the weak sense and $u \leq v$ on $\partial\Omega$, then $u \leq v$ in Ω .

In order to consider problem (1), let us recall some results, starting with the Dirichlet problem

(4)
$$\begin{cases} -\Delta_p u = f(x) & \text{in } \Omega\\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

The following comparison result, proved by Cuesta and Takáč [9, Proposition 2.3], will be very important in the sequel.

Lemma 2.2. Let f be a non-negative continuous function such that f(u) is nonincreasing for $u \in (0, \infty)$, where 1 < p. Assume that, for p > 1, $u, v \in W^{1,p}(\Omega)$ are weak solutions such that

$$\begin{array}{rcl} -\Delta_p u & \geqslant & f(u), & u > 0 & in \ \Omega, \\ -\Delta_p v & \leqslant & f(v), & v > 0 & in \ \Omega, \end{array}$$

and $u \ge v$ in $\partial \Omega$. Then $u \ge v$ in Ω . We now recall the definition of the Lorentz spaces. For a measurable function $\phi: \Omega \to \mathbb{R}$ and t > 0, we denote by

$$\mu_{\phi}(t) = |\{x \in \Omega : |\phi(x)| > t\}|$$

its *distribution function*, $|\cdot|$ standing for the Lebesgue measure in \mathbb{R}^N . For $0 \le r \le |\Omega|$, the decreasing rearrangement ϕ^* of ϕ is defined by

$$\phi^*(r) = \sup \{t > 0 : \mu_\phi(t) > r\},\$$

and the Lorentz space $L^{p,q}$ is defined for $1 and <math>1 \leq q < \infty$ by

$$L^{p,q} = \left\{ \phi \colon \Omega \to \mathbb{R} \text{ measurable} : \left(\int_{0}^{|\Omega|} \frac{1}{t} [\phi^*(t)t^{1/p}]^q dt \right)^{1/q} < \infty \right\}$$

endowed with norm

$$\|\phi\|_{p,q} = \left(\int_{0}^{|\Omega|} \frac{1}{t} [\phi^*(t)t^{1/p}]^q dt\right)^{1/q}$$

Denoting the usual norm in the space $L^q(\Omega)$ by $\|\cdot\|_q$, for $1 < q < \infty$ we have the identity

$$L^{q,q} = L^{q}(\Omega)$$
 and $\|\phi\|_{q,q} = \|\phi\|_{q}$.

Furthermore, the following continuous embeddings are true for $1 < s < N < q < \infty$.

$$L^q \hookrightarrow L^{N,1} \hookrightarrow L^{N,s} \hookrightarrow L^{N,N} = L^N \hookrightarrow L^{N,q} \hookrightarrow L^s,$$

see [1] or [23] for details.

The next two results are important because they relate the norm $\|\nabla u_p\|_{\infty}^{p-1}$ of a solution $u_p \in W_0^{1,p}(\Omega)$ of problem (4) with the norm $\|f\|_q$ by means of an explicit dependence on p, although not necessarily optimal.

Proposition 2.3. [12, Theorem 1.3] Let Ω be a bounded domain of \mathbb{R}^2 such that $\partial \Omega \in W^2 L^{\theta,1}$ for some $\theta > 1$. Suppose that $f \in L^q(\Omega)$ for some q > 2, and let $u_p \in W^{1,p}_0(\Omega)$ be a solution of problem (4). Then,

$$\|\nabla u_p\|_{\infty}^{p-1} \leqslant C \begin{cases} 2^{\frac{p}{p-1}} (p-1)^{-\frac{2\theta}{\theta-1}} \|f\|_q & \text{if } 1$$

for some constant *C* depending at most on *q* and Ω . Moreover, if the assumption $\partial \Omega \in W^2 L^{\theta,1}$ is replaced with the assumption that is Ω convex, then

$$\|\nabla u_p\|_{\infty}^{p-1} \leqslant C \begin{cases} 2^{\frac{p}{p-1}} \|f\|_q & \text{if } 1$$

for some constant *C* depending at most on *q* and Ω .

Proposition 2.4. [12, Theorem 1.2] Let Ω be a bounded domain of \mathbb{R}^N , $N \ge 3$, such that $\partial \Omega \in W^2 L^{\theta,1}$ for some $\theta > N - 1$, and assume that $f \in L^{N,1}(\Omega)$. Let $u_p \in W_0^{1,p}(\Omega)$ be a weak solution of the Dirichlet problem (4). Then the estimate

$$\|\nabla u_p\|_{\infty}^{p-1} \leqslant C \begin{cases} 2^{\frac{p}{p-1}} (p-1)^{-\frac{\theta N}{\theta - (N-1)}} \|f\|_{N,1} & \text{if } 1$$

holds for a constant C, depending at most on N and Ω . Moreover, if the assumption $\partial \Omega \in W^2 L^{\theta,1}$ is replaced with the assumption that is Ω convex, then

$$\|\nabla u_p\|_{\infty}^{p-1} \leqslant C \begin{cases} 2^{\frac{p}{p-1}} \|f\|_{N,1} & \text{if } 1$$

holds for constant *C* depending at most on *N* and Ω .

The two last results were originally established by Chianchi and Maz'ya [7] in a more general context for general constants C. However, the explicit dependence of the constants on p was obtained by Ercole in [12].

Now, suppose that there exist constants $0 < c < \infty$ and $0 < \delta < 1$ such that

$$0 \leqslant f(x) \leqslant c \, \mathrm{d}_{\Omega}(x)^{-\delta}$$

holds for almost all $x \in \Omega$, where $d_{\Omega} \in W^{1,\infty}(\Omega) \cap C_0(\overline{\Omega})$ denotes the distance function to the boundary defined by

$$\mathbf{d}_{\Omega}(x) := \operatorname{dist}(x, \partial \Omega) = \inf_{y \in \partial \Omega} |x - y|, \ x \in \overline{\Omega}.$$

When $0 < \delta < 1$, the integrability requirement given by

$$\int_{\Omega} d_{\Omega}(x)^{-\delta} dx < \infty$$

is a consequence of [17, Lemma 1] (see also [22, Lemma B.1]). Hence, the condition $0 < \alpha q < 1$ in Theorem 1.2 implies that

(6)
$$\int_{\Omega} d_{\Omega}(x)^{-\alpha q} dx < \infty$$

The following result is a simplified version of the one obtained by Giacomoni, Schindler and Takáč. [15, Theorem B.1].

Proposition 2.5. Assume that f satisfies the growth hypothesis (5). Let $u \in W_0^{1,p}(\Omega)$ be the (unique) weak solution of problem (4). In addition, assume

(7)
$$0 \leq u(x) \leq Cd_{\Omega}(x)$$
 for almost all $x \in \Omega$,

where *C* is a constant, $0 \leq C < \infty$. Then there exist constants β and Γ , $0 < \beta < 1$ and $0 \leq \Gamma < \infty$, depending solely on Ω , *N*, *p*, on the constants *c*, δ in (5), and on the constant *C* in (7), such that *u* satisfies $u \in C^{1,\beta}(\overline{\Omega})$ and

$$\|u\|_{C^{1,\beta}(\bar{\Omega})} \leq \Gamma$$

To prove Theorems 1.1 and 1.2 in this paper, we consider the unique solution $u_0 \in W_0^{1,p}(\Omega) \cap C^{1,\beta}(\overline{\Omega})$ of the *p*-Laplacian problem

(8)
$$\begin{cases} -\Delta_p u_0 = u_0^{-\alpha} & \text{in } \Omega \\ u_0 > 0 & \text{in } \Omega \\ u_0 = 0 & \text{on } \partial \Omega \end{cases}$$

see [6, Theorem 1.3] for further results about this problem.

The next result is technical and will be crucial in the proof of our Theorem 1.2.

Proposition 2.6. Let u_0 be the solution of (8). There are positive constants M and λ such that

(i)
$$aM^{r_1} \|u_0\|_{\infty}^{r_1} + bM^{r_2} \leqslant \frac{\lambda^{1-\alpha}}{\|u_0\|_{\infty}^{\alpha}};$$

(5)

(*ii*)
$$2^{\frac{1}{p-1}}\lambda^{\frac{1-\alpha}{p-1}} \leq M;$$

(*iii*) $M \leq \lambda^{\frac{2-p}{\alpha}}.$

Proof. Of course,

$$aM^{r_1} \|u_0\|_{\infty}^{r_1} \leqslant \frac{\lambda^{1-\alpha}}{2\|u_0\|_{\infty}^{\alpha}} \quad \Leftrightarrow \quad M \leqslant \frac{\lambda^{\frac{1-\alpha}{r_1}}}{\left(2a\|u_0\|_{\infty}^{r_1+\alpha}\right)^{\frac{1}{r_1}}}$$

and

$$bM^{r_2} \leqslant \frac{\lambda^{1-\alpha}}{2\|u_0\|_{\infty}^{\alpha}} \quad \Leftrightarrow \quad M \leqslant \frac{\lambda^{\frac{1-\alpha}{r_2}}}{(2b\|u_0\|_{\infty}^{\alpha})^{\frac{1}{r_2}}}.$$

Comparing the inequality in (ii) with the first expression obtained for M, we observe that

$$2^{\frac{1}{p-1}}\lambda^{\frac{1-\alpha}{p-1}} < \frac{\lambda^{\frac{1-\alpha}{r_1}}}{\left(2a\|u_0\|_{\infty}^{r_1+\alpha}\right)^{\frac{1}{r_1}}}$$

is equivalent to

(9)

$$\lambda^{\frac{(1-\alpha)(r_1+1-p)}{r_1(p-1)}} < \frac{1}{2^{\frac{1}{p-1}} \left(2a \|u_0\|_{\infty}^{r_1+\alpha}\right)^{\frac{1}{r_1}}}$$

Solving (9) for λ we obtain

$$\lambda < \frac{1}{2^{\frac{r_1}{(1-\alpha)(r_1+1-p)}} \left(2a\|u_0\|_{\infty}^{r_1+\alpha}\right)^{\frac{p-1}{(1-\alpha)(r_1+1-p)}}}$$

Similarly, we derive

$$\lambda < \frac{1}{2^{\frac{r_2}{(1-\alpha)(r_2+1-p)}} \left(2b\|u_0\|_{\infty}^{\alpha}\right)^{\frac{p-1}{(1-\alpha)(r_2+1-p)}}}.$$

Now let us consider (*ii*) and (*iii*). We have

$$2^{\frac{1}{p-1}}\lambda^{\frac{1-\alpha}{p-1}} < \lambda^{\frac{2-p}{\alpha}} \Leftrightarrow \lambda^{\frac{\alpha(1-\alpha)+(p-1)(p-2)}{\alpha(p-1)}} < \frac{1}{2^{\frac{1}{p-1}}} \Leftrightarrow \lambda < \frac{1}{2^{\frac{1}{\alpha(1-\alpha)+(p-1)(p-2)}}}.$$

Therefore, in order to obtain (i) - (ii) - (iii) it is enough to define

$$A \coloneqq \min\left\{\frac{1}{2^{\frac{r_1}{(1-\alpha)(r_1+1-p)}} \left(2a\|u_0\|_{\infty}^{r_1+\alpha}\right)^{\frac{p-1}{(1-\alpha)(r_1+1-p)}}}, \frac{1}{2^{\frac{r_2}{(1-\alpha)(r_2+1-p)}} \left(2b\|u_0\|_{\infty}^{\alpha}\right)^{\frac{p-1}{(1-\alpha)(r_2+1-p)}}}, \frac{1}{2^{\frac{\alpha}{\alpha(1-\alpha)+(p-1)(p-2)}}}, 1\right\}$$

and consider

$$0 < \lambda < A,$$

since the last inequality implies

$$2^{\frac{1}{p-1}}\lambda^{\frac{1-\alpha}{p-1}} < \min\left\{\frac{\lambda^{\frac{1-\alpha}{r_1}}}{\left(2a\|u_0\|_{\infty}^{r_1+\alpha}\right)^{\frac{1}{r_1}}}, \frac{\lambda^{\frac{1-\alpha}{r_2}}}{\left(2b\|u_0\|_{\infty}^{\alpha}\right)^{\frac{1}{r_2}}}, \lambda^{\frac{2-p}{\alpha}}\right\}$$

With such a choice of λ , we can choose *M* satisfying

$$2^{\frac{1}{p-1}}\lambda^{\frac{1-\alpha}{p-1}} \leqslant M \leqslant \min\left\{\frac{\lambda^{\frac{1-\alpha}{r_1}}}{\left(2a\|u_0\|_{\infty}^{r_1+\alpha}\right)^{\frac{1}{r_1}}}, \frac{\lambda^{\frac{1-\alpha}{r_2}}}{\left(2b\|u_0\|_{\infty}^{\alpha}\right)^{\frac{1}{r_2}}}, \lambda^{\frac{2-p}{\alpha}}\right\},$$

and we are done.

We now turn our attention to the singular problem

(10)
$$\begin{cases} -\Delta_p u = \frac{\lambda}{u^\delta} + u^s & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega \end{cases}$$

where $1 , <math>p - 1 < s \le p^* - 1$, $\lambda > 0$, and $0 < \delta < 1$. The following result is established in [15, Lemma A.7].

Lemma 2.7. Each positive weak solution u of problem (10) satisfies

$$c_{\lambda} \mathbf{d}_{\Omega} \leq u \leq K_{\lambda} \mathbf{d}_{\Omega}$$
 a.e. in Ω ,

where $0 < c_{\lambda} \leq K_{\lambda} < \infty$ are some constants independent of u.

In the course of the proof of Theorems 1.1 and 1.2, we denote by λ_1 and ϕ_1 the first eigenpair of the *p*-Laplacian, that is,

(11)
$$\begin{cases} -\Delta_p \phi_1 = \lambda_1 \phi_1^{p-1} & \text{in } \Omega\\ \phi_1 = 0 & \text{on } \partial\Omega, \end{cases}$$

with $\phi_1 > 0$ satisfying $\|\phi_1\|_{\infty} = 1$. Note that we have $\phi_1 > 0$ in Ω and $\frac{\partial \phi_1}{\partial \nu} < 0$ on $\partial \Omega$, see [24, Theorem 5]. Hence, since $\phi_1 \in C^1(\overline{\Omega})$, there are constants ℓ and $L, 0 < \ell < L$, such that $\ell d_{\Omega}(x) \leq \phi_1(x) \leq L d_{\Omega}(x)$ for all $x \in \Omega$.

3. AN AUXILIARY PROBLEM

For each $\varepsilon > 0$, we consider the auxiliary problem

(12)
$$\begin{cases} -\Delta_p u = \frac{\lambda}{(|u|+\varepsilon)^{\alpha}} + f(x,u,\nabla u) & \text{in } \Omega\\ u > 0 & \text{in } \Omega\\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

To prove Theorem 1.1 we first show the existence of a solution to the auxiliary problem (12) by applying the Galerkin method.

In order to do so, we state a useful consequence of Brouwer's fixed point theorem, which was proved in [2]. The statement is a subtle generalization of the classical result, allowing us to work with an arbitrary norm $|\cdot|_m$ in \mathbb{R}^m . We denote by (\cdot, \cdot) the usual inner product in \mathbb{R}^m .

Lemma 3.1. Let $F: (\mathbb{R}^m, |\cdot|_m) \to (\mathbb{R}^m, |\cdot|_m)$ be a continuous function such that $(F(\xi), \xi) \ge 0$ for every $\xi \in \mathbb{R}^m$ with $|\xi|_m = R$ for some R > 0. Then, there exists z_0 in the closed ball $\bar{B}_r^m(0) = \{z \in \mathbb{R}^m : |z|_m \le R\}$ such that $F(z_0) = 0$.

Lemma 3.2. For any $\varepsilon > 0$ the problem (12) admits a positive weak solution $u_{\varepsilon} \in W_0^{1,p}(\Omega) \cap C^{1,\sigma}(\overline{\Omega})$ for each $\lambda > 0$ and some $0 < \sigma < 1$.

Proof. Fix $\varepsilon > 0$ and consider $\mathcal{B} = \{e_1, ..., e_m, ...\}$ a Schauder basis of $W_0^{1,p}(\Omega)$. For each $m \in \mathbb{N}$ and $\xi = (\xi_1, ..., \xi_m) \in \mathbb{R}^m$, define the *m*-dimensional subspace $V_m := [e_1, ..., e_m] \subset W_0^{1,p}(\Omega)$ and the norm

$$|\xi|_m := \left\| \sum_{j=1}^m \xi_j e_j \right\|_{W_0^{1,p}(\Omega)}$$

It follows that the map $T_m: (\mathbb{R}^m, |\cdot|_m) \to \left(V_m, \|\cdot\|_{W_0^{1,p}(\Omega)}\right)$ given by $T_m(\xi) = \frac{m}{2}$

$$\sum_{j=1}^{j} \xi_j e_j = u \text{ is an isometric isomorphism. Consider the function } F \colon \mathbb{R}^m \to \mathbb{R}^m,$$
$$F(\xi) = (F_1(\xi), F_2(\xi), \dots, F_m(\xi)), \text{ given by}$$

$$F_j(\xi) = \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla e_j - \lambda \int_{\Omega} \frac{e_j}{(\varepsilon + |u|)^{\alpha}} - \int_{\Omega} f(x, u, \nabla u) e_j, \ 1 \le j \le m.$$

Claim. $F \colon \mathbb{R}^m \to \mathbb{R}^m$ satisfies the hypotheses of Lemma 3.1.

We start showing that F is continuous. Indeed, for $k \in \mathbb{N}$, suppose that $\zeta^k \to \zeta^0$ in \mathbb{R}^m and consider $u_k = \sum_{i=1}^m \zeta_i^k e_i \in V_m$. For each fixed $j \in \{1, 2, ..., m\}$, note that

$$F_j(\zeta^k) = \int_{\Omega} |\nabla u_k|^{p-2} \nabla u_k \nabla e_j - \lambda \int_{\Omega} \frac{e_j}{(\varepsilon + |u_k|)^{\alpha}} - \int_{\Omega} f(x, u_k, \nabla u_k) e_j.$$

We will only give attention to the term $\int_{\Omega} f(x, u_k, \nabla u_k) e_j$. Since

$$u_k = \sum_{i=1}^m \zeta_i^k e_i \to u_0 \coloneqq \sum_{i=1}^m \zeta_i^0 e_i \quad \text{in} \quad W_0^{1,p}(\Omega),$$

we obtain $u_k \to u_0$ and $\nabla u_k \to \nabla u_0$, both in $L^p(\Omega)$.

Since $r_1, r_2 it follows from (2) that$

$$0 \leq f(x, t, \xi) \leq c_1 + c_2 |t|^{p-1} + c_3 |\xi|^{p-1},$$

in $\Omega \times \mathbb{R} \times \mathbb{R}^m$. The continuity of the Nemytskii operator yields

(13)
$$f(\cdot, u_k, \nabla u_k) \to f(\cdot, u_0, \nabla u_0)$$
 in $L^{\frac{p}{p-1}}(\Omega)$.

Hence,

$$\begin{aligned} \left| \int_{\Omega} f(x, u_k, \nabla u_k) e_j - \int_{\Omega} f(x, u_0, \nabla u_0) e_j \right| \\ (14) \qquad \leq \left\| f(\cdot, u_k, \nabla u_k) - f(\cdot, u_0, \nabla u_0) \right\|_{L^{\frac{p}{p-1}}(\Omega)} \|e_j\|_{L^p(\Omega)} \to 0 \text{ as } n \to \infty. \end{aligned}$$

Thus, we conclude that $F_j(\zeta^k) \to F_j(\zeta^0)$, proving the continuity of *F*.

It follows from (2), Holder's inequality and the pertinent Sobolev embedding that

$$\langle F(\xi),\xi\rangle \ge \|u\|_{W_0^{1,p}(\Omega)}^p - \lambda c_1 \|u\|_{W_0^{1,p}(\Omega)}^{1-\alpha} - c_2 \|u\|_{W_0^{1,p}(\Omega)}^{r_1+1} - c_3 \|u\|_{W_0^{1,p}(\Omega)}^{r_2+1},$$

where c_1, c_2 and c_3 are positive constants independent of m and u. Because $r_1, r_2 and <math>1 - \alpha < p$, we obtain that $\langle F(\xi), \xi \rangle$ is coercive, that is, for each $\lambda > 0$ there exists $R = R(\lambda) > 0$ large enough such that $|\xi|_m := ||u||_{W_0^{1,p}(\Omega)} = R$ implies $\langle F(\xi), \xi \rangle \ge 0$.

As a consequence of Lemma 3.1, there exists $z_0 \in \mathbb{R}^m$ with $|z_0| \leq R$ such that $F(z_0) = 0$. Taking into account the isometric isomorphism $T_m : \mathbb{R}^m \to V_m$, there exist $u_m \in V_m$ with $||u_m||_{W_0^{1,p}(\Omega)} \leq R$ such that, for all $v \in V_m$,

(15)
$$\int_{\Omega} |\nabla u_m|^{p-2} \nabla u_m \nabla v = \lambda \int_{\Omega} \frac{v}{(\varepsilon + |u_m|)^{\alpha}} + \int_{\Omega} f(x, u_m, \nabla u_m) v.$$

Since *R* does not depend on *m*, the sequence (u_m) is bounded in $W_0^{1,p}(\Omega)$. Thus, for a subsequence, there exists $u \in W_0^{1,p}(\Omega)$ such that

$$u_m \rightarrow u$$
 weakly in $W_0^{1,p}(\Omega)$.

Consequently, we conclude that $u_m \to u$ in $L^p(\Omega)$ and $u_m(x) \to u(x)$ a.e. in Ω . Observe also that

(16)
$$||u||_{W_0^{1,p}(\Omega)} \leq \liminf_{m \to \infty} ||u_m||_{W_0^{1,p}(\Omega)} \leq R.$$

We now claim that

(17)
$$u_m \to u \text{ in } W_0^{1,p}(\Omega).$$

Since \mathcal{B} is a Schauder basis of $W_0^{1,p}(\Omega)$, there exists a unique sequence $(\alpha_n)_{n \ge 1}$ in \mathbb{R} such that $u = \sum_{j=1}^{\infty} \alpha_j e_j$. Thus, as $m \to \infty$,

(18)
$$\phi_m = \sum_{j=1}^m \alpha_j e_j \to u \text{ in } W_0^{1,p}(\Omega).$$

Considering the test function $(u_m - \phi_m) \in V_m$ in (15) yields

$$\begin{split} \int_{\Omega} |\nabla u_m|^{p-2} \nabla u_m \nabla (u_m - \phi_m) \\ &= \lambda \int_{\Omega} \frac{(u_m - \phi_m)}{(\varepsilon + |u_m|)^{\alpha}} + \int_{\Omega} f(x, u_m, \nabla u_m) (u_m - \phi_m) \\ &\leqslant \frac{\lambda}{\varepsilon^{\alpha}} \|u_m - \phi_m\|_{L^1(\Omega)} + \tilde{c}_1 \|u_m\|_{W_0^{1,p}(\Omega)}^{r_2} \|u_m - \phi_m\|_{L^{r_2+1}(\Omega)} + \\ &+ \tilde{c}_2 \|u_m\|_{W_0^{1,p}(\Omega)}^{r_1} \|u_m - \phi_m\|_{L^{r_1+1}(\Omega)} \to 0 \quad \text{as} \quad m \to \infty. \end{split}$$

where we use (2), Holder's inequality, the boundness of $||u_m||_{W_0^{1,p}(\Omega)}$, (17) and (18). Thus, we obtain

$$\lim_{m \to \infty} \int_{\Omega} |\nabla u_m|^{p-2} \nabla u_m \nabla (u_m - u) = 0$$

Now it is sufficient to apply the (S_+) property of $-\Delta_p$ (see [20, Proposition 3.5]) to obtain (17).

If $k \in \mathbb{N}$, then for every $m \ge k$ and $v_k \in V_k$, we obtain

$$|\nabla u_m|^{p-2} \nabla u_m \nabla v_k = \lambda \int_{\Omega} \frac{v_k}{(\varepsilon + |u_m|)^{\alpha}} + \int_{\Omega} f(x, u_m \nabla u_m) v_k.$$

Since $[V_k]_{k\in\mathbb{N}}$ is dense in $W_0^{1,p}(\Omega)$ we conclude that, for all $v \in W_0^{1,p}(\Omega)$,

$$\int_{\Omega} |\nabla u_m|^{p-2} \nabla u_m \nabla v = \lambda \int_{\Omega} \frac{v}{(\varepsilon + |u_m|)^{\alpha}} + \int_{\Omega} f(x, u_m, \nabla u_m) v.$$

Proceeding as in (13) and (14) we obtain

$$f(\cdot, u_m, \nabla u_m) \to f(\cdot, u, \nabla u) \quad \text{in} \quad L^{\frac{p}{p-1}}(\Omega)$$

and

$$\begin{split} \left| \int_{\Omega} f(x, u_m, \nabla u_m) v - \int_{\Omega} f(x, u, \nabla u) v \right| \\ & \leq \left\| f(\cdot, u_m, \nabla u_m) - f(\cdot, u, \nabla u) \right\|_{L^{\frac{p}{p-1}}(\Omega)} \|v\|_{L^p(\Omega)} \to 0 \text{ as } m \to \infty. \end{split}$$

Therefore,

$$\int_{\Omega} f(x, u_m, \nabla u_m) v \to \int_{\Omega} f(x, u, \nabla u) v, \quad \forall v \in W_0^{1, p}(\Omega).$$

As before, we conclude that

$$\int_{\Omega} |\nabla u_m|^{p-2} \nabla u_m \nabla v \to \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla v,$$

what yields, for all $v \in W_0^{1,p}(\Omega)$,

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla v = \lambda \int_{\Omega} \frac{v}{(\varepsilon + |u|)^{\alpha}} + \int_{\Omega} f(x, u, \nabla u) v.$$

Furthermore, since $u^- \in W_0^{1,p}(\Omega)$, we have

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla u^{-} = \lambda \int_{\Omega} \frac{u^{-}}{(\varepsilon + |u|)^{\alpha}} + \int_{\Omega} f(x, u, \nabla u) u^{-},$$

from what follows

$$-\|u^-\|_{W^{1,p}_0(\Omega)} \ge \lambda \int_{\Omega/\{u(x)>0\}} \frac{u^-}{(\varepsilon+|u|)^{\alpha}} \ge 0.$$

Then $u_{-} \equiv 0$ a.e. in Ω . Therefore,

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla v = \lambda \int_{\Omega} \frac{v}{(\varepsilon + u)^{\alpha}} + \int_{\Omega} f(x, u, \nabla u) v, \quad \forall v \in W_0^{1, p}(\Omega).$$

The equation in (12) guarantees that $u \neq 0$. By [16, Theorem 7.1] we infer that $u \in L^{\infty}(\Omega)$. Furthermore, regularity up to the boundary ensures that $u \in C^{1,\sigma}(\overline{\Omega})$ with some $\sigma \in (0,1)$, see [18, Theorem 1]. It results from the strong maximum principle that u > 0 in Ω , completing the proof that u is a solution of problem (12). We are done.

4. PROOF OF THEOREM 1.1

Proof. We consider first the existence of a solution by taking successively $\varepsilon = 1/n$, $n \in \mathbb{N}$. According to Lemma 3.2, for each $\lambda > 0$, there exists a solution $u_n \in C^{1,\sigma}(\overline{\Omega})$ of the auxiliary problem (12), that is, for all $v \in W_0^{1,p}(\Omega)$ we have

(19)
$$\int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n \nabla v = \lambda \int_{\Omega} \frac{v}{(\frac{1}{n} + u_n)^{\alpha}} + \int_{\Omega} f(x, u_n, \nabla u_n) v,$$

and, in particular, (19) is valid for all $v \in C_c^1(\Omega)$.

Taking into account (16) and proceeding as in the proof of (17), we obtain a subsequence (u_n) such that

(20)
$$u_n \to u \text{ in } W_0^{1,p}(\Omega).$$

In order to handle the first integral on the right side of (19), let us consider $w = \lambda^{\frac{1}{p-1-\alpha}} u_0$, where u_0 is the unique solution of (8). Of course, w is a *positive* solution to the problem

(21)
$$\begin{cases} -\Delta_p w = \frac{\lambda}{w^{\alpha}} & \text{in } \Omega\\ w = 0 & \text{on } \partial\Omega. \end{cases}$$

Since u_n is a solution of (12), we have

$$-\Delta_p\left(u_n + \frac{1}{n}\right) = -\Delta_p u_n \ge \frac{\lambda}{\left(u_n + \frac{1}{n}\right)^{\alpha}}$$

and $(u_n + 1/n) > 0$ on $\partial\Omega$. In particular, $u_n + \frac{1}{n}$ is a super-solution of (8) and, as a consequence of Lemma 2.2, we obtain $u_n + \frac{1}{n} \ge w$.

Let ϕ_1 be the first eigenfunction of the p-Laplacian, as in (11). For each $\beta \in (0, \lambda_1^{-1/(p-1-\alpha)})$ we have that $\beta \phi_1$ satisfies

$$\begin{cases} -\Delta_p(\beta\phi_1) \leqslant \frac{1}{(\beta\phi_1)^{\alpha}} & \text{in } \Omega\\ \beta\phi_1 > 0 & \text{in } \Omega\\ \beta\phi_1 = 0 & \text{on } \partial\Omega \end{cases}$$

In particular, $\beta \phi_1$ is a sub-solution of (8). Another application of Lemma 2.2 yields $u_0 \ge \beta \phi_1$.

Since there is a constante $\ell > 0$ such that $\ell d_{\Omega} \leq \phi_1$, we obtain

$$(22) u_0 \ge k d_\Omega,$$

where $k = \ell \beta$. Therefore,

$$v = \lambda^{\frac{1}{p-1-\alpha}} u_0 \ge k_1 d_{\Omega_2}$$

where $k_1 = k\lambda^{\frac{1}{p-1-\alpha}}$. Consequently, $u_n + \frac{1}{n} \ge w \ge k_1 d_{\Omega}$, what implies

$$\frac{|v|}{\left(u_n + \frac{1}{n}\right)^{\alpha}} \leqslant \frac{|v|}{\left(k_1 d_{\Omega}\right)^{\alpha}}$$

for all $v \in C_0^1(\Omega)$.

It follows from (6) that $\frac{v}{(k_1 d_\Omega)^{\alpha}}$ is integrable. Since $u_n + \frac{1}{n} \to u$ a.e in Ω , we conclude that $u \ge k_1 d_\Omega > 0$ a.e. in Ω and

$$\frac{v}{(\frac{1}{n}+u_n)^{lpha}}
ightarrow \frac{v}{u^{lpha}}$$
 a.e. in Ω

as $n \to \infty$.

Now, an application of Lebesgue's dominated convergence theorem yields

(23)
$$\int_{\Omega} \frac{v}{(\frac{1}{n} + u_n)^{\alpha}} \to \int_{\Omega} \frac{v}{u^{\alpha}} \text{ as } n \to +\infty$$

Since $C_0^1(\Omega)$ is dense in $W_0^{1,p}(\Omega)$ and u_n solves (19), it follows from (20) and (23) as $n \to \infty$ that, for all $v \in W_0^{1,p}(\Omega)$,

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla v = \lambda \int_{\Omega} \frac{v}{u^{\alpha}} + \int_{\Omega} f(x, u, \nabla u) v, \ \forall v \in W_0^{1, p}(\Omega).$$

The proof is complete.

5. PROOF OF THEOREM 1.2

Proof. Let us consider λ and M satisfying Proposition 2.6. Define the set

$$\mathcal{A} := \left\{ v \in W_0^{1,p}(\Omega) : \lambda u_0 \leqslant v \leqslant M u_0, \ \|\nabla v\|_{\infty} \leqslant M \right\}.$$

We will prove that for each $v \in A$ there exists an unique positive solution u of the problem

(24)
$$\begin{cases} -\Delta_p u &= \lambda v^{-\alpha} + f(x, v, \nabla v) & \text{in } \Omega \\ u &> 0 & \text{in } \Omega \\ u &= 0 & \text{on } \partial\Omega. \end{cases}$$

We know that there are positive constants ℓ and C_2 such that

$$\ell d_{\Omega}(x) \leqslant \phi_1(x) \leqslant C_2 u_0(x),$$

thus yielding, for $c = \ell/C_2$,

(25)
$$cd_{\Omega}(x) \leq u_0(x)$$

If $p' = \frac{p}{p-1}$ and $q > \max\{N, p'\}$, by our hypothesis we have $0 < \alpha q < 1$. Then

$$\frac{1}{u_0(x)^{q\alpha}} \leqslant \frac{1}{[cd_\Omega(x)]^{q\alpha}}.$$

Thus, it follows from (6) that

$$\int_{\Omega} \left(\frac{1}{u_0(x)^{\alpha}} \right)^q = \int_{\Omega} \frac{1}{u_0(x)^{q\alpha}} \leqslant \int_{\Omega} \frac{1}{[cd_{\Omega}(x)]^{q\alpha}} < \infty,$$

and, since $\lambda u_0 \leq v$,

$$\begin{aligned} \left(\int_{\Omega} \left| \frac{\lambda}{v^{\alpha}} + f(x, v, \nabla v) \right|^{q} \right)^{\frac{1}{q}} &\leq \left(\int_{\Omega} \left(\left| \frac{\lambda}{v^{\alpha}} \right| + a|v|^{r_{1}} + b|\nabla v|^{r_{2}} \right)^{q} \right)^{\frac{1}{q}} \\ &\leq \left(\int_{\Omega} \left(\frac{\lambda^{1-\alpha}}{|u_{0}|^{\alpha}} + aM^{r_{1}} \|u_{0}\|_{\infty}^{r_{1}} + bM^{r_{2}} \right)^{q} \right)^{\frac{1}{q}} \\ &\leq \left(\int_{\Omega} \left(\frac{\lambda^{1-\alpha}}{|u_{0}|^{\alpha}} + \frac{\lambda^{1-\alpha}}{|u_{0}|^{\alpha}} \right)^{q} \right)^{\frac{1}{q}} = \left(\int_{\Omega} \left(\frac{2\lambda^{1-\alpha}}{|u_{0}|^{\alpha}} \right)^{q} \right)^{\frac{1}{q}} \\ &= 2\lambda^{1-\alpha} \left\| \frac{1}{u_{0}^{\alpha}} \right\|_{L^{q}(\Omega)} < \infty. \end{aligned}$$

(2

We conclude that $\frac{\lambda}{v^{\alpha}} + f(x, v, \nabla v) \in L^q(\Omega)$ and, since q > p', it follows that

$$\frac{\lambda}{v^{\alpha}} + f(x, v, \nabla v) \in L^{p'}(\Omega).$$

By applying [11, Theorem 8], we obtain the existence and uniqueness of the solution u.

We now show that $u \in A$, proving first that $\lambda u_0 \leq u \leq M u_0$. In fact, let $v \in A$. Proposition 2.6 guarantees that

$$-\Delta_p u \geqslant \frac{\lambda}{v^{\alpha}} \geqslant \frac{\lambda}{M^{\alpha} u_0^{\alpha}} = \frac{\lambda}{M^{\alpha}} \frac{1}{u_0^{\alpha}} \geqslant \lambda^{p-1} \frac{1}{u_0^{\alpha}} = \lambda^{p-1} (-\Delta_p u_0) = -\Delta_p (\lambda u_0).$$

The comparison principle imply that $u \ge \lambda u_0$.

On the other hand, it follows from Proposition 2.6 and (2) that

$$-\Delta_p u = \frac{\lambda}{v^{\alpha}} + f(x, v, \nabla v) \leqslant \frac{\lambda}{\lambda^{\alpha} u_0^{\alpha}} + a|v|^{r_1} + b|\nabla v|^{r_2}$$
$$\leqslant \frac{\lambda^{1-\alpha}}{u_0^{\alpha}} + aM^{r_1} \|u_0\|_{\infty} + bM^{r_2} \leqslant \frac{\lambda^{1-\alpha}}{u_0^{\alpha}} + \frac{\lambda^{1-\alpha}}{\|u_0\|_{\infty}^{\alpha}}$$
$$\leqslant \frac{2\lambda^{1-\alpha}}{u_0^{\alpha}} = 2\lambda^{1-\alpha} (-\Delta_p u_0) = -\Delta_p (2^{\frac{1}{p-1}} \lambda^{\frac{1-\alpha}{p-1}} u_0).$$

The comparison principle and Proposition 2.6 imply that $u \leq 2^{\frac{1}{p-1}} \lambda^{\frac{1-\alpha}{p-1}} u_0 \leq M u_0$, that is,

$$\lambda u_0 \leqslant u \leqslant M u_0.$$

We will now show $\|\nabla u\|_{\infty} \leq M$. If N = 2, since $\frac{\lambda}{v^{\alpha}} + f(x, v, \nabla v) \in L^{q}(\Omega)$, it follows from Proposition 2.3 that

$$\|\nabla u\|_{\infty}^{p-1} \leq c_p \left\| \frac{\lambda}{v^{\alpha}} + f(\,\cdot\,,v,\nabla v) \right\|_{L^q(\Omega)} \leq c_p 2\lambda^{1-\alpha} \left\| \frac{1}{u_0^{\alpha}} \right\|_{L^q(\Omega)},$$

and therefore

$$\|\nabla u\|_{\infty} \leqslant \tilde{c_p} 2^{\frac{1}{p-1}} \lambda^{\frac{1-\alpha}{p-1}},$$

where $\tilde{c_p} = c_p^{\frac{1}{p-1}}$. If $N \ge 3$, since q > N, we have $L^q(\Omega) \subset L^{N,1}(\Omega)$, allowing us to conclude that $rac{\lambda}{v^{lpha}}+f(x,v,
abla v)\in L^{N,1}(\Omega).$ So, (26) yields

$$\left\|\frac{\lambda}{v^{\alpha}} + f(.,v,\nabla v)\right\|_{L^{N,1}(\Omega)} \leq \overline{c} \left\|\frac{\lambda}{v^{\alpha}} + f(.,v,\nabla v)\right\|_{L^{q}(\Omega)} < \infty.$$

It follows from Proposition 2.4 that

$$\begin{aligned} \|\nabla u\|_{\infty}^{p-1} &\leq C_p \left\| \frac{\lambda}{v^{\alpha}} + f(\,\cdot\,,v,\nabla v) \right\|_{L^{N,1}(\Omega)} \\ &\leq C_p \,\bar{c} \left\| \frac{\lambda}{v^{\alpha}} + f(\,\cdot\,,v,\nabla v) \right\|_{L^q(\Omega)} \qquad \leq C_p \,\bar{c} \, 2\lambda^{1-\alpha} \left\| \frac{1}{u_0^{\alpha}} \right\|_{L^q(\Omega)} \end{aligned}$$

and we conclude that

$$\|\nabla u\|_{\infty} \leqslant \hat{C}_p 2^{\frac{1}{p-1}} \lambda^{\frac{1-\alpha}{p-1}},$$

where $\hat{C}_p = (C_p \overline{c})^{\frac{1}{p-1}}$. Thus, by considering $\tilde{C}_p = \max{\{\tilde{c}_p, \hat{C}_p\}}$, we have for all $N \ge 2$

$$\|\nabla u\|_{\infty} \leqslant \tilde{C}_p 2^{\frac{1}{p-1}} \lambda^{\frac{1-\alpha}{p-1}}.$$

14

Now, choosing λ so that $0 < \lambda < A^*$, where

$$\begin{aligned} A^* \coloneqq \min \left\{ \frac{1}{2^{\frac{r_1}{(1-\alpha)(r_1+1-p)}} \left(2a\|u_0\|_{\infty}^{r_1+\alpha}\right)^{\frac{p-1}{(1-\alpha)(r_1+1-p)}}}, \frac{1}{2^{\frac{r_2}{(1-\alpha)(r_2+1-p)}} \left(2b\|u_0\|_{\infty}^{\alpha}\right)^{\frac{p-1}{(1-\alpha)(r_2+1-p)}}}, \\ \frac{1}{2^{\frac{\alpha}{\alpha(1-\alpha)+(p-1)(p-2)}}}, \frac{1}{\left(\tilde{C}_p 2^{\frac{1}{p-1}}\right)^{\frac{r_1(p-1)}{(1-\alpha)(r_1+1-p)}} \left(2a\|u_0\|_{\infty}^{r_1+\alpha}\right)^{\frac{p-1}{(1-\alpha)(r_1+1-p)}}}}{\frac{1}{\left(\tilde{C}_p 2^{\frac{1}{p-1}}\right)^{\frac{r_2(p-1)}{(1-\alpha)(r_2+1-p)}}}, \frac{1}{\left(\tilde{C}_p 2^{\frac{1}{p-1}}\right)^{\frac{\alpha(p-1)}{\alpha(1-\alpha)+(p-1)(p-2)}}}, 1\right\}}, \end{aligned}$$

we obtain $M = M(\lambda)$ such that

$$\tilde{C}_{p}2^{\frac{1}{p-1}}\lambda^{\frac{1-\alpha}{p-1}} \leqslant M \leqslant \min\left\{\frac{\lambda^{\frac{1-\alpha}{r_{1}}}}{\left(2a\|u_{0}\|_{\infty}^{r_{1}+\alpha}\right)^{\frac{1}{r_{1}}}}, \frac{\lambda^{\frac{1-\alpha}{r_{2}}}}{\left(2b\|u_{0}\|_{\infty}^{\alpha}\right)^{\frac{1}{r_{2}}}}, \lambda^{\frac{2-p}{\alpha}}\right\}$$

Thus

$$\|\nabla u\|_{\infty} \leqslant \tilde{C}_p 2^{\frac{1}{p-1}} \lambda^{\frac{1-\alpha}{p-1}} \leqslant M.$$

Claim. We have $u \in C^{1,\beta}(\overline{\Omega})$ for some $0 < \beta < 1$, uniformly with respect to $v \in \mathcal{A}$.

In order to prove the Claim, we apply Proposition 2.5. Indeed, since each solution *U* of (10) is a super-solution of (21), the comparison principle (Lemma 2.2) yields $w = \lambda^{\frac{1}{p-1+\alpha}} u_0 \leq U$.

By Lemma 2.7 there is a positive constant K_{λ} such that $U \leq K_{\lambda}d_{\Omega}$, consequently, $u_0 \leq K_{\lambda}d_{\Omega}$. Moreover, by (22) there is a positive constante k such that $u_0 \geq kd_{\Omega}$. Furthermore, since

$$\begin{split} \frac{\lambda}{v^{\alpha}} + f(x, v, \nabla v) &\leqslant \frac{\lambda}{v^{\alpha}} + a|v|^{r_1} + b|\nabla v|^{r_2} \\ &\leqslant \frac{\lambda^{1-\alpha}}{|u_0|^{\alpha}} + aM^{r_1} \|u_0\|_{\infty}^{r_1} + bM^{r_2} \leqslant \frac{2\lambda^{1-\alpha}}{|u_0|^{\alpha}} \\ &\leqslant \frac{K_1}{d_{\Omega}^{\alpha}} = K_1 d_{\Omega}^{-\alpha} \text{ a.e in } \Omega, \end{split}$$

where $K_1 = \frac{2\lambda^{1-\alpha}}{k^{\alpha}}$, as a consequence of (25). Moreover, (27) implies that

$$u \leqslant M u_0 \leqslant M K_\lambda d_\Omega.$$

So, Proposition 2.5 guarantees that the solution of (24) satisfies $u \in C^{1,\beta}(\overline{\Omega})$ for some $0 < \beta < 1$.

As consequence of the previous arguments, the operator

$$\begin{array}{cccc} T: \mathcal{A} & \longrightarrow & W_0^{1,p}(\Omega) \\ v & \longmapsto & u, \end{array}$$

is well-defined, u being the unique positive solution of (24). Moreover, T is continuous and compact. In fact, let $(v_n)_{n\in\mathbb{N}} \subset \mathcal{A}$ be a bounded sequence in $W_0^{1,p}(\Omega)$ such that $T(v_n) = u_n$. Since

$$\frac{\lambda}{v_n^{\alpha}} + f(x, v_n, \nabla v_n) \leqslant \frac{K_1}{d_{\Omega}^{\alpha}} = K_1 d_{\Omega}^{-\alpha}, \text{ for all } n \in \mathbb{N},$$

arguments like the previous ones and Proposition 2.5 yield that

$$\|u_n\|_{C^{1,\beta}(\bar{\Omega})} \leq \Gamma$$

or, equivalently,

$$||T(v_n)||_{C^{1,\beta}(\bar{\Omega})} \leq \Gamma.$$

Hence, \mathcal{A} is a equicontinuous subset. Therefore, there is a convergent subsequence of $(T(v_n))_{n\in\mathbb{N}}$ in $C^1(\overline{\Omega})$ which, in particular, is convergent in $W_0^{1,p}(\Omega)$. Thus, since \mathcal{A} is a bounded, convex set invariant under T, it follows from Schauder's Fixed Point Theorem the existence of a fixed point u_{λ} for T. Of course, the fixed point u_lambda satisfies (1), since $-\Delta_p u = \lambda u^{-\alpha} + f(x, u, \nabla u)$.

Availability of data and materials. This declaration is not applicable.

Competing interests: The authors declare that they have no conflict of interests.

Authors' contributions. All authors contributed equally to the article.

References

- [1] R. A. Adams, Sobolev Spaces. Academic Press, 1975.
- [2] A. L. A. de Araujo, L. F. O. Faria, Existence, nonexistence, and asymptotic behavior of solutions for N-Laplacian equations involving critical exponential growth in the whole R^N. Math. Ann. 384 (2022), no. 3–4, 1469–1507.
- [3] Y. Bai, D. Motreanu and S. Zeng, Continuity results for parametric nonlinear singular Dirichlet problems. Adv. Nonlinear Anal. 9 (2020), no.1, 372–387.
- [4] Y. Bai, N. S. Papageorgiou and S. Zeng, A singular eigenvalue problem for the Dirichlet (p, q)-Laplacian. Math. Z. 300 (2022), no. 1, 325–345.
- [5] H. Bueno and G. Ercole, A quasilinear problem with fast growing gradient. Appl. Math. Letters 26 (2013), 520–523.
- [6] A. Canino, B. Sciunzi, and A. Trombetta, Existence and uniqueness for p-Laplace equations involving singular nonlinearities. NoDEA Nonlinear Differential Equations Appl. 23 (2016), no. 2, Art. 8, 18 pp.
- [7] A. Cianchi and V. G. Maz'ya, *Global Lipschitz regularity for a class of quasilinear elliptic equations*. Comm. Partial Differential Equations 36 (2011), no. 1, 100–133.
- [8] F.J. Corrêa, A.S. Corrêa and G. Figueiredo, Positive solution for a class of p&q-singular elliptic equation. Nonlinear Anal. Real World Appl. 16 (2014), 163–169.
- [9] M. Cuesta and P. Takáč, A strong comparison principle for positive solutions of degenerate elliptic equations. Differential Integral Equations 13 (2000), no. 4-6, 721–746.
- [10] E. DiBenedetto, C^{1+α} local regularity of weak solutions of degenerate elliptic equations. Nonlinear Anal. 7 (1983), no. 8, 827–850 (1998).
- [11] G. Dinca, P. Jebelean and J. Mawhin, Variational and topological methods for Dirichlet problems with p-Laplacian. Port. Math. (N.S.) 58 (2001), no.3, 340–378.
- [12] G. Ercole, On a global gradient estimate in p-Laplacian problems, to appear in Israel J. Math.
- [13] L. F. O. Faria, O. H. Miyagaki and D. Motreanu, Comparison and positive solutions for problems with (p, q)-Laplacian and convection term. Proc. Edinb. Math. Soc. 57 (2014), no. 2, 687–698 (2014).
- [14] L. Gasiński and N. S. Papageorgiou, Nonlinear elliptic equations with singular terms and combined nonlinearities. Ann. Henri Poincaré 13 (2012), no. 3, 481–512.
- [15] J. Giacomoni, I. Schindler and P. Takáč, Sobolev versus Hölder local minimizers and existence of multiple solutions for a singular quasilinear equation. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 6 (2007), 117–158.
- [16] O.A. Ladyzhenskaja, N.N. Ural'tseva, Linear and Quasilinear Elliptic Equations. Academic Press, New York-London, 1968, xviii+495 pp.
- [17] A. C. Lazer and P. J. Mckenna, On a singular nonlinear elliptic boundary-value problem, Proc. Amer. Math. Soc. 111 (1991), no. 3, 721–730.
- [18] G. M. Lieberman, Boundary regularity for solutions of degenerate elliptic equations. Nonlinear Anal. 12 (1988), no. 11, 1203–1219.

- [19] Z. Liu, D. Motreanu, and S. Zeng, Positive solutions for nonlinear singular elliptic equations of p-Laplacian type with dependence on the gradient. Calc. Var. Partial Differential Equations 58 (2019), no. 1, Paper no. 28, 22 pp.
- [20] D. Motreanu, V. V. Motreanu and N. S. Papageorgiou, Multiple constant sign and nodal solutions for Nonlinear Neumann eigenvalue problems. Ann. Sc. Norm. Sup. Pisa Cl. Sci. (5) 10 (2011), no. 3, 72–755.
- [21] N. S. Papageorgiou, V. D. Rădulescu and L. Wen, Strongly singular nonhomogeneous eigenvalue problems. Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM 117 (2023), no.1, Paper No. 32, 17 pp.
- [22] M. F. Stapenhorst, A class of singular elliptic equations. Ph. D. dissertation, Universidade Estadual de Campinas (Unicamp), Departamento de Matemática, Estatística e Computação Científica, Campinas, Brasil.
- [23] L. Tartar, Imbedding theorems of Sobolev spaces into Lorentz spaces. Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8) 1 (1998), no. 3, 479–500.
- [24] J. L. Vázquez, A strong maximum principle for some quasilinear elliptic equations. Appl. Math. Optim. 12 (1984), no. 3, 191–202.

(Anderson L.A. de Araujo) DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE VI-ÇOSA, 36570-900, VIÇOSA, MG, BRAZIL

Email address: anderson.araujo@ufv.br

(Hamilton P. Bueno) DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE MINAS GE-RAIS, 31270-901 - BELO HORIZONTE - MG, BRAZIL

Email address: hamilton@mat.ufmg.br

(Kamila F. L. Madalena) DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE MINAS GERAIS, 31270-901 - BELO HORIZONTE - MG, BRAZIL

Email address: kamilalobo@ufmg.br