

Complex $D(2, 1; \zeta)$ and spin chain solutions from Chern-Simons theory

El Hassan Saidi

¹ *LPHE-MS, Science Faculty, Mohammed V University in Rabat, Morocco*

² *Centre of Physics and Mathematics, CPM- Morocco*

E-mail: e.saidi@um5r.ac.ma

ABSTRACT: Using properties of $OSp(4|2)$ and $PSL(2|2)$, we investigate the super geometry of the parametric $D(2, 1; \zeta)$ labeled by variable ζ belonging to $\mathbb{C} \setminus \{-1, 0\}$ and we give applications in the study of integrable superspin chains. This 9|8 dimensional Lie supergroup has three orthogonal isospins in its even part $SL(2, \mathbb{C})^{\otimes 3}$ assembled by the tri-fundamental $2^{\otimes 3}$ with odd parity. It undergoes contractions at $\zeta = -1, 0$ where an $SL(2, \mathbb{C})$ gets decompactified into commutative \mathbb{C}^3 interpreted in terms of three central extensions. By help of the obtained characteristic features of $D(2, 1; \zeta)$ and their local structures at the special points $\zeta = \pm 1$, we calculate the Lax operator $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ solving the RLL equation describing the integrability of the superspin chain $\mathfrak{d}(2, 1; \zeta)$. We also complete missing results regarding the calculation of $\mathcal{L}_{psl(2|2)}^{(\mu)}$ and $\mathcal{L}_{osp(4|2)}^{(\mu)}$. Other features of the four super Dynkin diagrams $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ and weight graphs of $\mathfrak{d}(2, 1; \zeta)$ as well as discrete automorphisms are also given.

KEYWORDS: Super Dynkin diagrams and super graphs of $\mathfrak{d}(2,1;\zeta)$, supergeometry of $D(2, 1; \zeta)$, decompactification of $SL(2, \mathbb{C})$, integrable superspin chain, calculation of L-operators in 4D CS theory.

Contents

1	Introduction	1
2	Complex geometry of distinguished $SL(2 2)$	5
2.1	$SL(2 2)$ as a 15D hypersurface \mathfrak{h}_{15} in $\mathbb{C}^{8 8}$	6
2.1.1	Defining relations and properties	6
2.1.2	$GL(2)$ submanifolds inside $SL(2 2)$	8
2.1.3	Even and odd parts of $SL(2 2)$	14
2.2	Decompactifying the S^2 inside $SL(2, \mathbb{C})$	17
2.2.1	$SL(2, \mathbb{C})$ inside $D(2, 1; \zeta)_{\bar{0}}$	17
2.2.2	Decompactified S^2 and shrinking \tilde{S}^2	18
3	Graded 2-cycles in $D(2, 1; \zeta)$	19
3.1	Basis sets $\tilde{\Pi}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ and outer automorphisms	19
3.1.1	Gradation of super Dynkin $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$	21
3.1.2	Super 2-cycles in $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$	23
3.2	Realising the distinguished $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ and $\tilde{\mathcal{C}}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ systems	24
3.2.1	Roots $\tilde{\alpha}$ as function of ζ	26
3.2.2	Weight diagram of $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ from the $\Phi_{osp(4 2)}$ graph	27
4	Geometry of $D(2, 1; \zeta)$ as deformation of $O\mathcal{S}p(4 2)$	30
4.1	Bosonic subspace $D(2, 1; \zeta)_{\bar{0}}$	31
4.2	Odd part $D(2, 1; \zeta)_{\bar{1}}$	32
5	Integrable $D(2,1; \zeta)$ superspin chain	34
5.1	Integrable $sl(2 2)$ superspin chain	36
5.1.1	Distinguished $sl(2 2)$ spin chain	36
5.1.2	Calculating L-operator $\mathcal{L}_{sl_2 2}^{\mu_3}$	38
5.2	Integrable $osp(4 2)$ superspin chain	40
5.2.1	Distinguished orthosymplectic spin chain $osp(4 2)$	40
5.2.2	Computing the orthosymplectic $\mathcal{L}_{osp(4 2)}^{\mu_3}$	42
5.3	Lax operator of $D(2,1;\zeta)$ superspin chain	44
5.3.1	$\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$ as deformation of $\mathcal{L}_{osp}^{\mu_3}$	44
5.3.2	Calculation of $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$	45

6	Conclusion and comments	47
A	Characteristic data of $SL(2 2)$	49
B	Orthosymplectic $osp(4 2)$	56
C	Scheneurt realisation of $\mathfrak{d}(2,1;\zeta)$	59
D	Integrable superspin chains in CS theory	62

1 Introduction

The exceptional 17 dimensional Lie superalgebra $\mathfrak{d}(2,1;\zeta)$ has been subject to increasing interest during last decades; it has three orthogonal isospins $sl(2)^{\oplus 3}$ glued together by 8 fermionic charges making it somehow a special super symmetric system [1]-[9]. It is a parametric superalgebra labeled by a continuous parameter ζ belonging to the complex line \mathbb{C} deprived of the points $\{-1, 0\}$ where it contracts into two semi-simple sub- superalgebras. Generally speaking, the exceptional $\mathfrak{d}(2,1;\zeta)$ was introduced by V. G Kac [1] as a deformation of the orthosymplectic 17- dim Lie superalgebra $osp(4|2)$ located at $\zeta = 1$. This deviation is imagined in this paper as a characteristic datum of the exotic $\mathfrak{d}(2,1;\zeta)$ and will be used as a fundamental algorithm in the present study.

The even part $\mathfrak{d}(2,1;\zeta)_0$ of this superalgebra has several real forms including $su(1, 1)^{\oplus 3}$ and $su(1, 1) \oplus so(1, 3)$ as well as the remarkable $sl(2, \mathbb{R}) \oplus su(2, \mathbb{R})^{\oplus 2}$ [10] which found interesting applications in type II strings on $AdS_3 \times S^3 \times S^3$ [11]-[20]. It has been also considered in the study of exotic integrable superspin chains [21]-[26] going beyond the classical ABCD superchains [27-33] by using Bethe ansatz and algebraic methods [34-37]. Here, the superspin chain $\mathfrak{d}(2,1;\zeta)$ will be revisited from the view of 4D Chern-Simons (CS) theory [38-43] where the solution of the RLL equation (1.1) signing the integrability of the superspin chain is given by the computation of the Lax operator that we denote like

$$\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}$$

The CS investigation of the exceptional integrable superspin chain $\mathfrak{d}(2,1;\zeta)$ and the calculation of the parametric L-operator $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}$ constitute therefore one of the main results of the present work (see [section 5](#) and [Appendix D](#)).

Algebraically, the complexified super $\mathfrak{d}_{\mathbb{C}}(2,1;\zeta)$ —for short $\mathfrak{d}(2,1;\zeta)$ — is classified as a basic Lie superalgebra in the dictionary of Frappat and Sorba [44, 45]. It has four parametric super Dynkin diagrams with three nodes $(\alpha_1, \alpha_2, \alpha_3)$ [45–47] that we denote collectively as

$$S\mathfrak{D}\mathfrak{D}_{\zeta}^{(\eta)}$$

with label $\eta = 0, 1, 2, 3$. These super diagrams are effectively given by the graphs of the **Figure 9**; they can be pictured into a unified form as depicted by the **Figure 1**. However,




Figure 1: A unified representation of the four super Dynkin diagrams of the Lie superalgebra $\mathfrak{d}(2,1,\zeta)$ given by the **Figure 9**. The four super Dynkin diagrams $S\mathfrak{D}\mathfrak{D}_{\zeta}^{(\eta)}$ organise into a singlet $S\mathfrak{D}\mathfrak{D}_{\zeta}^{(0)}$ and a triplet $S\mathfrak{D}\mathfrak{D}_{\zeta}^{(i)}$ of the permutation group \mathcal{S}_3 . The closed $S\mathfrak{D}\mathfrak{D}_{\zeta}^{(0)}$ has three odd simple roots versus an odd simple root for the open $S\mathfrak{D}\mathfrak{D}_{\zeta}^{(i)}$; it is in the blue centre of the triangle.

the four super $\mathfrak{D}\mathfrak{D}_{\zeta}^{(\eta)}$ are not completely independent; they split in two blocks like $\mathbf{1} \oplus \mathbf{3}$ (blue singlet and red triplet in the **Figure 1**). The three $S\mathfrak{D}\mathfrak{D}_{\zeta}^{(1)}$, $S\mathfrak{D}\mathfrak{D}_{\zeta}^{(2)}$, $S\mathfrak{D}\mathfrak{D}_{\zeta}^{(3)}$ are exchanged by an outer automorphism symmetry given by the group \mathcal{Z}_3 . Because of this symmetry, one can then focus on one of them; say the $S\mathfrak{D}\mathfrak{D}_{\zeta}^{(2)}$ with odd simple root α_2 as it will be done in this paper; see the **Figure 2** for illustration. Notice that the discrete outer automorphism \mathcal{Z}_3 is in fact a subgroup of the non abelian symmetric group \mathcal{S}_3 that act by permuting the so-called *Kaplansky* parameters (s_1, s_2, s_3) used for the description of $\mathfrak{d}(2,1;\zeta)$ by the isomorphic *Scheunert* Lie superalgebra $\Gamma(s_1, s_2, s_3)$ [3, 48, 49]. Here, the complex parameters s_i , which are constrained like $s_1 + s_2 + s_3 = 0$ with $s_1 s_2 s_3 \neq 0$, are realised as $2s_1 = 1$, $2s_2 = -1 - \zeta$ and $2s_3 = \zeta$. This choice breaks \mathcal{S}_3 down to its




Figure 2: Even simple roots: $\tilde{\alpha}_1 = (\epsilon_1 - \epsilon_2)$ and $\tilde{\alpha}_3 = \sqrt{\zeta}(\epsilon_1 + \epsilon_2)$. Odd simple root $\tilde{\alpha}_2 = \frac{1}{2}\sqrt{2(1+\zeta)}\delta - \frac{1}{2}(\epsilon_1 - \epsilon_2) - \frac{1}{2}\sqrt{\zeta}(\epsilon_1 + \epsilon_2)$. These roots are functions of ζ .

subgroup \mathcal{Z}_2 which fixes s_1 and permutes s_2 and s_3 ; that is the \mathcal{Z}_2 generated by the transposition $\tau_{23} : (s_1, s_2, s_3) \rightarrow (s_1, s_3, s_2)$.

The parametric super $\mathfrak{D}\mathfrak{D}_\zeta^{(\eta)}$ will play an important role in this study; they share fundamental aspects with the super Dynkin diagrams of $\mathfrak{psl}(2|2)$ and $\mathfrak{osp}(4|2)$ sitting at the special points $\zeta = -1$ and $+1$; thus capturing important information on the exceptional super symmetry. For the singular limit $\zeta \rightarrow -1$, the exceptional $\mathfrak{d}(2,1;\zeta)$ contracts into $\mathfrak{psl}(2|2) \oplus \mathbb{C}^3$ [50–52] where one of the three $\mathfrak{sl}(2, \mathbb{C})$ s making the even part $\mathfrak{d}(2,1;\zeta)_0$ gets replaced by the abelian \mathbb{C}^3 whose translation generators (P_1, P_2, P_3) have been interpreted like three commuting central extensions (see Appendix A for details). For the regular limit $\zeta \rightarrow +1$, the $\mathfrak{d}(2,1;\zeta)$ coincides with the 17 dimensional orthosymplectic $\mathfrak{osp}(4|2)$ providing in turn an important tool for engineering models with $\mathfrak{d}(2,1;\zeta)$ symmetry characterised by a finite set of deformation parameters $\{\kappa_i\}$ which are functions of the complex ζ as it will be shown throughout this investigation.

In this paper, we aim for two main objectives motivated by (1) completing partial results on the super group $D(2,1;\zeta)$ and (2) by the application of the construction in integrable systems based on Chern-Simons potentials. For that, we first revisit the study of the complexified $\mathfrak{d}(2,1;\zeta)$ from algebraic geometry view; then use this description to get more insight into the parametric supergroup manifold $D(2,1;\zeta)$ in order to develop the study the integrable $\mathfrak{d}(2,1;\zeta)$ superspin chain by using 4D CS theory. Because of the complexity in directly solving the RLL integrability equation (D.6) of the $\mathfrak{d}(2,1;\zeta)$ superchain namely,

$$\mathbf{R}_{rs}^{ik}(z-w) \mathbf{L}_j^r(z) \mathbf{L}_l^s(w) = \mathbf{L}_r^i(w) \mathbf{L}_s^k(z) \mathbf{R}_{jl}^{rs}(z-w) \quad (1.1)$$

we get around this difficulty by taking advantage of its local structures at the special points $\zeta = \pm 1$ where live $\mathfrak{psl}(2|2)$ and $\mathfrak{osp}(4|2)$. Here, \mathbf{R}_{rs}^{ik} is the usual R-matrix of Yang Baxter equation and \mathbf{L}_j^i is the matrix realisation of the Lax operator. In this way, we aim to reach the parametric $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}$ by seeking to continuously join the two superchain solutions at $\zeta = \pm 1$ while respecting the constraints required by the exceptional symmetry.

However, an inspection of the literature on integrable superspin chains reveals that there are no results on the derivation of the Lax operators for the $\mathfrak{psl}(2|2)$ and $\mathfrak{osp}(4|2)$ superchains by using the CS method. Results in this regard are only known for the superalgebras $\mathfrak{psl}(m|n)$ with $m \neq n$ and $\mathfrak{osp}(2k|2l)$ with $k \geq 3$ and $l \geq 1$. To overcome this, we start by filling in this gap by first calculating the missing $\mathcal{L}_{\mathfrak{psl}(2|2)}$ and $\mathcal{L}_{\mathfrak{osp}(4|2)}$ Lax operators; and then proceed to calculate $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}$. As a result of this approach, we derive a family of $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}$ operators labeled by three complex parameters $(\kappa_0, \kappa_1, \kappa_2)$ which are functions of ζ . A particular matrix representation of the obtained solutions is given by the following 6×6 Lax matrix with $\kappa_1 = 1$ and $\kappa_2 = -\kappa_0$,

$$\begin{pmatrix} (z^{\frac{1}{2}} + \kappa_0^2 z^{-\frac{1}{2}} b' c') \delta_j^i + z^{-\frac{1}{2}} \beta^i \gamma_j & z^{-\frac{1}{2}} (c \beta^i - \kappa_0^2 b' \gamma_n \varepsilon^{ni}) & z^{-\frac{1}{2}} \beta^i & \kappa_0 z^{-\frac{1}{2}} b' \varepsilon^{ji} \\ z^{-\frac{1}{2}} (b \gamma_j - \kappa_0^2 c' \varepsilon_{jn} \beta^n) & z^{\frac{1}{2}} + z^{-\frac{1}{2}} (bc + \kappa_0^2 \beta^n \gamma_n) & z^{-\frac{1}{2}} b & -\kappa_0 z^{-\frac{1}{2}} \beta^j \\ z^{-\frac{1}{2}} \gamma_j & z^{-\frac{1}{2}} c & z^{-\frac{1}{2}} & 0 \\ \kappa_0 z^{-\frac{1}{2}} c' \varepsilon_{ji} & -\kappa_0 z^{-\frac{1}{2}} \gamma_i & 0 & z^{-\frac{1}{2}} \delta_i^j \end{pmatrix} \quad (1.2)$$

Here z is the usual complex variable (rapidity) used in the 4D CS theory whose field action $S_{4dCS}[A]$ is reported in the [Appendix D](#); see eq(D.2). The two pairs (β^i, γ_i) are fermionic oscillators satisfying $\{\beta^j, \gamma_i\} = \delta_i^j$; while the two singlets (b, c) and (b', c') are bosonic oscillators obeying $[b, c] = 1 = [b', c']$. Moreover, the parametric function $\kappa_0(\zeta)$ in (1.2) is equal to $\frac{1}{2}\zeta(\zeta + 1)$; it is given by the determinant $\det(\alpha_i \cdot \alpha_j)$ of the intersection matrix of the simple roots $(\alpha_1, \alpha_2, \alpha_3)$ generating the distinguished root system of $\mathfrak{d}(2, 1; \zeta)$ with odd α_2 and even (α_1, α_3) defining $S\mathfrak{D}\mathfrak{D}_\zeta^{(2)}$. The values of the L-matrix (1.2) at $\zeta = \pm 1$ recovers precisely the expressions the matrix representations of obtained $\mathcal{L}_{\mathfrak{psl}(2|2)}$ and $\mathcal{L}_{\mathfrak{osp}(4|2)}$.

The organisation of this paper is as follows: In [section 2](#), we study the algebraic geometry of the distinguished complex 14D supermanifold of $PSL(2|2)$ with emphasis on its 6D even $PSL(2|2)_0 \simeq SL(3)^{\otimes 2}$ part and the 8D odd $PSL(2|2)_1$ module $2 \times \mathbf{2}^{\otimes 2}$. The [section 3](#) and [section 4](#) are devoted to the exceptional $D(2, 1; \zeta)$. There, we investigate the geometry of the complex 17D supermanifold of the supergroup $D(2, 1; \zeta)$, its algebraic properties in link with the orthosymplectic $OSp(4|2)$ as well as with its decompactification into $PSL(2|2) \times \mathbb{C}^3$. In [section 5](#), we study the integrable superspin chain $\mathfrak{d}(2, 1; \zeta)$ by using the 4D Chern-Simons theory. Because of lack of results on integrable $\mathfrak{sl}(n|n)$ superchains in Chern-Simons formulation, we fill in this gap by calculating $\mathcal{L}_{\mathfrak{psl}(2|2)}$ in subsection 5.1. In subsection 5.2, we also determine $\mathcal{L}_{\mathfrak{osp}(4|2)}$ since it has not

been calculated before; and in subsection 5.3, we calculate the L-operator for the $\mathfrak{d}(2,1;\zeta)$ superchain. In [section 6](#), we give conclusion and additional comments. Auxiliary sections are devoted to *four* appendices. In [Appendix A](#), we give useful properties of the complex Lie superalgebras $\mathfrak{sl}(2|2)$ and its simple subalgebra $\mathfrak{psl}(2|2)$. There, we use the 4×4 super matrix realisation of $\mathfrak{sl}(2|2)$ to describe the algebraic and geometric properties of the super manifold $SL(2|2)$. In [Appendix B](#), we give some helpful ingredients of $\mathfrak{osp}(4|2)$ and study its oscillator realisation as well as its link with the exceptional super $D(2,1;\zeta)$ at $\zeta = 1$. In [Appendix C](#), we give complement tools regarding $D(2,1;\zeta)$ and the realisation of its four possible roots systems. In [Appendix D](#), we recall some basic tools on the 4D Chern-Simons theory for integrable super systems.

2 Complex geometry of distinguished $SL(2|2)$

In this section, we use the complex 4×4 super matrix realisation of $SL(2|2)$ with the 2×2 block matrix decomposition [\[53\]](#),

$$\mathbf{M}_{4 \times 4} = \begin{pmatrix} \mathbf{A}_{2 \times 2} & \mathbf{B}_{2 \times 2} \\ \mathbf{C}_{2 \times 2} & \mathbf{D}_{2 \times 2} \end{pmatrix} \quad (2.1)$$

as well as the distinguished root system $\Phi_{\mathfrak{sl}(2|2)}^{\Pi_1}$ to study useful algebraic and geometric properties of the $SL(2|2)$ supermanifold denoted here after like $\mathcal{X}_{\mathfrak{sl}(2|2)}^{\Pi_1}$. This 15D supermanifold is associated with the distinguished simple root basis $\Pi_1 = (\alpha_1, \alpha_2, \alpha_3)$ having odd α_2 and even α_1, α_3 [\[44\]](#); see the [Appendix A](#) for the four possible $SL(2|2)$ root systems $\Phi_{\mathfrak{sl}(2|2)}^{\Pi_\eta}$ labeled by $\eta = 0, 1, 2, 3$ and the associated graphs of the [Figure 15](#). Through this analysis, we also use the two possible real forms of $SL(2, \mathbb{C})$ to study other properties $\mathcal{X}_{\mathfrak{sl}(2|2)}^{\Pi_1}$ such as critical limits given by $SU(2)$ singularity of complex surfaces and its $SU(1,1)$ conical homologue thought of here in term of vanishing "gap energy" ideas as described by eqs [\(2.32\)](#)-[\(2.35\)](#).

The investigation given in this section should be also viewed as a front matter for addressing the $(9|8)$ -dimensional (17D) supergeometry of the parametric $D(2,1;\zeta)$ containing $PSL(2|2)$ as a complex $(6|8)$ -dimensional (14D) submanifold. The results obtained here as well as the tools given below will be used for the geometric interpretation of the contraction $D(2,1;\zeta) \rightarrow PSL(2|2)$ in terms of the vanishing of the (dual) 2-cycle

$$\mathcal{C}_{\tilde{\psi}} = \mathcal{C}_{\alpha_1 + 2\alpha_2 + \alpha_3} \quad (2.2)$$

sitting in the normal space $D(2,1;\zeta) \setminus PSL(2|2)$. In the above relation, the $\tilde{\psi} = \alpha_1 + 2\alpha_2 + \alpha_3$ is the long root of $D(2,1;\zeta)$ and $\mathcal{C}_{\tilde{\psi}}$ the associated homological 2-cycle in the

parametric $\mathcal{X}_{0(2,1;\zeta)}^{\Pi_1}$; its area $\mathcal{A}(\mathcal{C}_{\tilde{\psi}}) = \int_{\mathcal{C}_{\tilde{\psi}}} J_{Kahler}$ is a function of ζ ; it vanishes in the limit $\zeta \rightarrow -1$ as it will be shown later on.

2.1 SL(2|2) as a 15D hypersurface \mathfrak{h}_{15} in $\mathbb{C}^{8|8}$

We first study the algebraic relations describing the supermanifold $\mathcal{X}_{sl(2|2)}^{\Pi_1}$ of the distinguished Lie supergroup $SL(2|2)$. Then, we turn to give useful properties of this complex 15 dimensional- space \mathfrak{h}_{15} and its submanifolds; in particular $PSL(2|2)$, $SL(2)$ and $GL(1)$.

2.1.1 Defining relations and properties

The geometric description of the complex 15 dimensional $SL(2|2)$ is done in terms of graded complex 4×4 matrices (2.1) that can be also presented like $\mathbf{M}_{4 \times 4} = \exp(\mathbf{H}_{4 \times 4})$ with vanishing supertrace $str(\mathbf{H}_{4 \times 4}) = 0$. It is done by generalising usual constructions of bosonic symmetries $SU(2)$ and $SL(2, \mathbb{R}) \simeq SU(1, 1)$ describing the two real forms of $SL(2, \mathbb{C})$ [20, 54]. To that purpose, we consider a generic complex super matrix

$$\mathbf{M}_{4 \times 4} = \begin{pmatrix} \mathbf{X}_1 & \mathbf{Y}_2 \\ \mathbf{Y}_1 & \mathbf{X}_2 \end{pmatrix} \Leftrightarrow (\mathbf{M})_{\mathbf{B}}^{\mathbf{A}} = \begin{pmatrix} (\mathbf{X}_1)_b^a & (\mathbf{Y}_2)_\beta^a \\ (\mathbf{Y}_1)_b^\alpha & (\mathbf{X}_2)_\beta^\alpha \end{pmatrix} \quad (2.3)$$

belonging to $\text{End}(\mathbb{C}^{2|2})$ where the canonical vector basis $\{\mathbf{e}_1, \mathbf{e}_2; \mathbf{e}_3, \mathbf{e}_4\}$ of $\mathbb{C}^{2|2}$ are ordered as (even, even, odd, odd) that is (bbff) in the notation used in (A.26) termed as distinguished ordering; here the label (b) refers to bosonic and the (f) to fermionic parities. In eq(2.3), the complex $\mathbf{X}_{1,2}$ and $\mathbf{Y}_{1,2}$ are four 2×2 block matrices with entry variables as

$$\mathbf{X}_i = \begin{pmatrix} x_i & u_i \\ y_i & v_i \end{pmatrix}, \quad \mathbf{Y}_i = \begin{pmatrix} \xi_i & \phi_i \\ \zeta_i & \chi_i \end{pmatrix} \quad (2.4)$$

The 8+8 complex entries of the super matrix (2.3) reads in a condensed way like $(\mathbf{M})_{\mathbf{B}}^{\mathbf{A}}$ with labels $a, b = 1, 2$ for the first $SL(2)_1$ and $\alpha, \beta = 1, 2$ for the second $SL(2)_2$ making the even part $PSL(2|2)_0$. This labeling shows that \mathbf{Y}_1 and \mathbf{Y}_2 sit in the bi-fundamental of $SL(2)_1 \times SL(2)_2$ while \mathbf{X}_1 and \mathbf{X}_2 are in the adjoints of $GL(2)_1$ and $GL(2)_2$ respectively. By substituting (2.4), we have

$$\mathbf{M}_{\mathbf{B}}^{\mathbf{A}} = \begin{pmatrix} x_1 & u_1 & \xi_2 & \phi_2 \\ y_1 & v_1 & \zeta_2 & \chi_2 \\ \xi_1 & \phi_1 & x_2 & u_2 \\ \zeta_1 & \chi_1 & y_2 & v_2 \end{pmatrix} \quad (2.5)$$

with entries belonging to $\mathbb{C}^{8|8}$. In term of the super matrix \mathbf{M} , the defining equation of the complex $SL(2|2)$ manifold is given by the unimodular condition of the Berezinian (super determinant: sdet) $Ber(\mathbf{M}_{4 \times 4}) = 1$ turning into a constraint relation among the 16 complex variables in eq(2.5). This super determinant can be presented in a handy form using the 2×2 sub-block matrices $\mathbf{X}_{1,2}$ and $\mathbf{Y}_{1,2}$ like $\det[\mathbf{X}_1 - \mathbf{Y}_1 \mathbf{X}_2^{-1} \mathbf{Y}_2] \det(\mathbf{X}_2)^{-1} = 1$; thus defining a complex 15 dim hypersurface \mathfrak{h}_{15} in $\mathbb{C}^{8|8}$. By rewriting this constraint relation like

$$\mathfrak{h}_{15} : \quad \det(\mathbf{X}_1) \cdot \det[\mathbf{I} - \mathbf{X}_1^{-1} \mathbf{Y}_1 \mathbf{X}_2^{-1} \mathbf{Y}_2] \cdot \det(\mathbf{X}_2^{-1}) = 1 \quad (2.6)$$

one gets a correspondence between:

- On one hand, the triple factorisation of eq(2.6) given by the three blocks $\det(\mathbf{X}_1)$, $\det[\mathbf{I} - \mathbf{X}_1^{-1} \mathbf{Y}_1 \mathbf{X}_2^{-1} \mathbf{Y}_2]$ and $\det(\mathbf{X}_2^{-1})$; and
- On the other hand, the subalgebras $sl(2, \mathbb{C})_A$, $sl(1|1)_{AB}$ and $sl(2, \mathbb{C})_B$ in $sl(2|2)$. These subalgebras are indexed by the simple roots (α_1) , (α_2) and (α_3) where α_2 is fermionic and (α_1, α_3) bosonic.

One also derives other interesting properties on distinguished $SL(2|2)$ geometry; in particular the two useful ones:

(A) projective symmetry: Eq(2.6) has a remarkable invariance given by the projective transformation

$$\mathbb{C}^* : \mathbf{X}_i \rightarrow \lambda \mathbf{X}_i \quad , \quad \mathbf{Y}_i \rightarrow \lambda \mathbf{Y}_i \quad (2.7)$$

where λ is a non vanishing \mathbb{C} -number. This projective symmetry **(i)** allows to fix one of the complex degrees of freedom in $\mathbf{M}_{4 \times 4}$; for example $\det(\mathbf{X}_1) = 1$ expanding like $x_1 v_1 - y_1 u_1 = 1$, and **(ii)** permits to construct the defining relation of the 14 dimensional supermanifold $PSL(2|2)$ in terms of the coset

$$PSL(2|2) = SL(2|2)/\mathbb{C}^* \quad , \quad \mathbb{C}^* \simeq GL(1, \mathbb{C}) \quad (2.8)$$

(B) $sl(2)$ singularities: For the case where the two even submatrices \mathbf{X}_i take the diagonal values $z_i I_{2 \times 2}$ where the z_i 's are complex numbers [i.e: $\mathbf{X}_i \in GL(1, \mathbb{C})_i$], we have $\det(\mathbf{X}_i) = z_i^2$. If moreover, the two complex numbers are taken equal ($z_1 = z_2 = z$), that is

$$\mathbf{M}_{4 \times 4} = \begin{pmatrix} z I_{2 \times 2} & \mathbf{Y}_2 \\ \mathbf{Y}_1 & z I_{2 \times 2} \end{pmatrix} \quad (2.9)$$

we have the equality $\det(\mathbf{X}_1) = \det(\mathbf{X}_2)$; and then eq(2.6) becomes

$$\det \left[\mathbf{I}_{2 \times 2} - \frac{1}{z^2} \mathbf{Y}_1 \mathbf{Y}_2 \right] = 1 \quad (2.10)$$

Natural solutions of this particular relation are obtained by taking $\mathbf{Y}_1 = w_1 \mathbf{I}_{2 \times 2}$ and $\mathbf{Y}_2 = w_2 \mathbf{I}_{2 \times 2}$ where w_1 and w_2 are complex numbers; then putting into (2.10), we get

$$\left(1 - \frac{w_1 w_2}{z^2} \right)^2 = 1 \quad \Rightarrow \quad 1 - \frac{w_1 w_2}{z^2} = \pm 1 \quad \Rightarrow \quad w_1 w_2 = (1 \mp 1) z^2 \quad (2.11)$$

thus leading to two remarkable solutions corresponding to the \pm sectors:

(1) **First solution:** it is given by the vanishing condition $w_1 w_2 = 0$ corresponding to a triangular matrix $\mathbf{M}_{4 \times 4} = \mathbf{T}_{4 \times 4}$ where $w_1 = 0$ and $w_2 \neq 0$ or $w_1 \neq 0$ and $w_2 = 0$. For the case $w_1 = w_2 = 0$, it reduces simply to $\mathbf{D}_{4 \times 4} = z \mathbf{I}_{2 \times 2} \oplus z \mathbf{I}_{2 \times 2}$. This diagonal supermatrix describes a complex curve \mathcal{C} within $SL(2|2)$. Also, the $\mathbf{D}_{4 \times 4}$ is the central element of the super group $SL(2|2)$ showing that the hypersurface \mathfrak{h}_{15} describing this complex 15 dimensional supermanifold has within it a homological 2-cycle which turns out to be intimately related to the $\mathcal{C}_{\tilde{\psi}}$ given by (2.2).

(2) **Second solution:** it is given by the interesting relation

$$w_1 w_2 = 2z^2 \quad \Rightarrow \quad \mathbf{Y}_1 \mathbf{Y}_2 = 2z^2 \mathbf{I}_{2 \times 2} \quad (2.12)$$

leading to

$$(\mathbf{M})_{\mathbf{B}}^{\mathbf{A}} = \begin{pmatrix} z\delta_b^a & w_2\delta_\beta^a \\ w_1\delta_b^\alpha & z\delta_\beta^\alpha \end{pmatrix} \quad (2.13)$$

It describes a complex surface Σ inside $SL(2|2)$ given by the complex equation $w_1 w_2 = 2z^2$. This surface has an $\mathfrak{sl}(2)$ singularity at $z=0$ [55, 56] corresponding to the vanishing of the central 2-cycle $\mathcal{C}_{\tilde{\psi}}$. This feature shows that the complex 15 dimensional manifold \mathfrak{h}_{15} describing $SL(2|2)$ factorises like

$$\mathfrak{h}_{15} = PSL(2|2) \times \mathcal{C}_{\tilde{\psi}} \quad (2.14)$$

2.1.2 $GL(2)$ submanifolds inside $SL(2|2)$

The complex super $SL(2|2)$ has a 7-dim even part $SL(2|2)_{\bar{0}}$ and 8-dim odd one $SL(2|2)_{\bar{1}}$. The even part factorises like $SL(2)_1 \times SL(2)_2 \otimes \mathbb{C}^*$; it has two $GL(2, \mathbb{C})$ blocks reading in terms of the \mathbf{X}_i 's in eq(2.4) as follows

$$\mathbf{M}_{4 \times 4} |_{SL(2|2)_{\bar{0}}} = \begin{pmatrix} \mathbf{X}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{X}_2 \end{pmatrix} \quad (2.15)$$

with (i) super determinant $Ber(\mathbf{M}_{4 \times 4}) = 1$ requiring $(\det \mathbf{X}_1)(\det \mathbf{X}_2)^{-1} = 1$; and (ii) the projective symmetry $\mathbb{C}^* : (\mathbf{X}_1, \mathbf{X}_2) \rightarrow \lambda(\mathbf{X}_1, \mathbf{X}_2)$ permitting to set $\det \mathbf{X}_1 = 1$ and then $\det \mathbf{X}_2 = 1$; thus restricting the $GL(2, \mathbb{C})_i$ blocks to $SL(2, \mathbb{C})_i$. Below, we describe useful features of the geometry of the $GL(2, \mathbb{C})$ s while focussing on one of them; say $GL(2, \mathbb{C})_1$. To that purpose, we begin by the 2×2 matrix realisation of the Lie group $GL(2, \mathbb{C})_1$ given by

$$\mathbf{X}_1 = \begin{pmatrix} x_1 & u_1 \\ y_1 & v_1 \end{pmatrix}, \quad \mathbf{X}_1^{-1} = \frac{1}{\mu} \begin{pmatrix} v_1 & -u_1 \\ -y_1 & x_1 \end{pmatrix} := \begin{pmatrix} \tilde{v}_1 & -\tilde{u}_1 \\ -\tilde{y}_1 & \tilde{x}_1 \end{pmatrix} \quad (2.16)$$

with $\det \mathbf{X}_1 = \mu$ where μ is a fixed complex number ($\mu = \text{cte}$) parameterising the abelian factor $GL(1, \mathbb{C})$ in the group $GL(2, \mathbb{C})_1$ which factorises like $GL(2)_1 = SL(2)_1 \times GL(1)_1$. The determinant of the above even matrix describes a 3D complex space \mathcal{V}_1 (resp. $\tilde{\mathcal{V}}_1$) parameterised by the 4 coordinates (x_1, y_1, u_1, v_1) [resp. $(\tilde{x}_1, \tilde{y}_1, \tilde{u}_1, \tilde{v}_1)$] constrained as $x_1 v_1 - y_1 u_1 = \mu$ (resp. $\tilde{x}_1 \tilde{v}_1 - \tilde{y}_1 \tilde{u}_1 = 1/\mu$). For simplicity, we rewrite these constraints just like

$$\mathcal{V} : \quad xv - yu = \mu \quad , \quad \tilde{\mathcal{V}} : \quad \tilde{x}\tilde{v} - \tilde{y}\tilde{u} = \tilde{\mu} = \frac{1}{\mu} \quad (2.17)$$

by dropping out the subscript 1. As we will see below, it turns out that it is more interesting to work with the inverse matrix \mathbf{X}_1^{-1} than the \mathbf{X}_1 . We refer to the use of \mathbf{X}_1^{-1} as the dual parametrisation.

In the realisations (2.17), the complex threefold \mathcal{V}_1 (resp. $\tilde{\mathcal{V}}_1$) is given by a complex 3D hypersurface $\det \mathbf{X}_1 = \mu$ (resp. $\det \mathbf{X}_1^{-1} = 1/\mu$) in the ambient complex space $\mathbb{C}^{4|0}$ with coordinates (x, y, u, v) [resp. $(\tilde{x}, \tilde{y}, \tilde{u}, \tilde{v})$]. For later use, notice that when we study $D(2, 1; \zeta)$, we will think about the modulus μ as a function of the ζ parameter namely

$$\frac{1}{\mu} = \kappa_0(\zeta) \quad , \quad \kappa_0 = \frac{1}{2}\zeta(\zeta + 1) \quad (2.18)$$

with $\kappa_0(1) = 1$ and $\kappa_0(-1) = 0$ corresponding to $OSp(4|2)$ and $PSL(2|2)$ respectively. Notice moreover the following:

- (a) the complex 3D hypersurfaces \mathcal{V} and $\tilde{\mathcal{V}}$ are dual threefolds with duality transformation given by the mapping $(\tilde{x}, \tilde{y}, \tilde{u}, \tilde{v}) = \frac{1}{\mu}(v, -y, -u, x)$.
- (b) the complex 3D hypersurface $xv - yu = \mu$ contains two special real threefolds:
 - (i) a parametric compact 3-sphere $\mathbb{S}^3(\mu)$ that can roughly imagined in term of a group coset as

$$\mathbb{S}^3(\mu) \simeq SO(4)/SO(3) \simeq SO(3) \simeq SU(2) \quad (2.19)$$

This 3-sphere sits in \mathbb{C}^4 at the $SU(2)$ -locus given by

$$SU(2) : \quad (u, v) = (-\bar{y}, \bar{x}) \quad (2.20)$$

and is described by the equation

$$\mathbb{S}^3(\mu) : \quad x\bar{x} + y\bar{y} = |\mu| \geq 0 \quad (2.21)$$

(ii) a parametric non compact "pseudo 3-sphere" $\mathbb{S}^{1,2}(\mu)$ that can be also thought of in terms of groups as follows

$$\mathbb{S}^{1,2}(\mu) \simeq SO(2, 2)/SO(1, 2) \simeq SO(1, 2) \simeq SU(1, 1) \quad (2.22)$$

It sits in \mathbb{C}^4 at the $SU(1, 1)$ -locus

$$SU(1, 1) : \quad (u, v) = (\bar{y}, \bar{x}) \quad (2.23)$$

and is given by the algebraic equation

$$\mathbb{S}^{1,2}(\mu) : \quad x\bar{x} - y\bar{y} = |\mu| \quad (2.24)$$

Comparing (2.20) and (2.23), we learn that these two loci can be mapped into each other by a discrete Z_2 transformation acting on the variables as follows

$$\begin{array}{ccc} SU(2) & : & (u, v) = (-\bar{y}, \bar{x}) \\ \downarrow & & \downarrow \\ SU(1,1) & : & (u, v) = (+\bar{y}, \bar{x}) \end{array} \quad (2.25)$$

In the limit $|\mu| \rightarrow \infty$, we have the "decompactifications" to the flat spaces

$$\lim_{|\mu| \rightarrow \infty} \mathbb{S}^3(\mu) \simeq \mathbb{R}^3 \quad , \quad \lim_{|\mu| \rightarrow \infty} \mathbb{S}^{1,2}(\mu) \simeq \mathbb{R}^{1,2} \quad (2.26)$$

By setting $x = \xi_1 + i\xi_2$ and $y = \xi_3 + i\xi_4$ with real ξ_i , eqs(2.17) expand as

$$\begin{array}{l} \mathbb{S}^3(\mu) \quad : \quad \xi_1^2 + \xi_2^2 + \xi_3^2 + \xi_4^2 = |\mu| \\ \mathbb{S}^{1,2}(\mu) \quad : \quad \xi_1^2 + \xi_2^2 - \xi_3^2 - \xi_4^2 = |\mu| \end{array} \quad (2.27)$$

(c) the dual hypersurface $\tilde{\mathcal{V}}$ described by the relation $\tilde{x}\tilde{v} - \tilde{y}\tilde{u} = 1/\mu$ contains also two special real threefolds given by the real forms of $SL(2, \mathbb{C})$ namely:

(i) a 3-sphere $\tilde{\mathbb{S}}^3(\mu) \simeq SU(2)$ (dual to the real $\mathbb{S}^3(\mu)$) sitting at the locus

$$SU(2) : \quad (\tilde{u}, \tilde{v}) = (-\bar{\tilde{y}}, \bar{\tilde{x}}) \quad (2.28)$$

it is described by the equation $|\tilde{x}|^2 + |\tilde{y}|^2 = 1/|\mu|$. This 3-sphere shrinks to a point in the limit $|\mu| \rightarrow \infty$, i.e:

$$\lim_{|\mu| \rightarrow \infty} \tilde{\mathbb{S}}^3(\mu) \simeq \{(\tilde{x}, \tilde{y}) = (0, 0)\} \quad (2.29)$$

We refer to this degenerate limit as an $SU(2)$ *singularity*. For later use, notice that this singularity can be expressed in terms of vanishing "gap energy" used in the study of topological insulators and topological matter [59–63, 86, 87]. Indeed, expressing the equation of the 3-sphere in terms of $|\tilde{x}|_{\pm}$ and $|\tilde{y}|_{\pm}$ like

$$|\tilde{x}|_{\pm} = \pm \sqrt{\frac{1}{|\mu|} - |\tilde{y}|^2} \quad \Leftrightarrow \quad |\tilde{y}|_{\pm} = \pm \sqrt{\frac{1}{|\mu|} - |\tilde{x}|^2} \quad (2.30)$$

with $0 \leq |\tilde{x}|^2 \leq \frac{1}{|\mu|}$ and $0 \leq |\tilde{y}|^2 \leq \frac{1}{|\mu|}$; it follows that the "gap energy" defined as

$$\Delta |\tilde{x}|_g = \min(|\tilde{x}|_+ - |\tilde{x}|_-) \quad , \quad \Delta |\tilde{y}|_g = \min(|\tilde{y}|_+ - |\tilde{y}|_-) \quad (2.31)$$

vanishes identically for $|\mu| \rightarrow \infty$.

(ii) a "pseudo 3-sphere" $\tilde{\mathbb{S}}^{1,2}(\mu) \simeq SU(1, 1)$ (dual to the real $\mathbb{S}^{1,2}(\mu)$) sitting in \mathbb{C}^4 at the locus

$$SU(1, 1) : \quad (\tilde{u}, \tilde{v}) = (\tilde{y}, \tilde{x}) \quad (2.32)$$

it is described by the equation $|\tilde{x}|^2 - |\tilde{y}|^2 = 1/|\mu|$. This relation can be also presented in terms of branches like

$$|\tilde{x}|_{\pm} = \pm \sqrt{|\tilde{y}|^2 + \frac{1}{|\mu|}} \quad \Leftrightarrow \quad |\tilde{y}|_{\pm} = \pm \sqrt{|\tilde{x}|^2 - \frac{1}{|\mu|}} \quad (2.33)$$

with $|\tilde{x}|^2 \geq \frac{1}{|\mu|}$ and $|\tilde{y}|^2 \geq 0$. In this case, the "gap energy" is as follows

$$\Delta |\tilde{x}|_g = 2\sqrt{\frac{1}{|\mu|}} \quad , \quad \Delta |\tilde{y}|_g = 0 \quad , \quad |\tilde{x}|^2 = \frac{1}{|\mu|} \quad (2.34)$$

In the limit $|\mu| \rightarrow \infty$, this gap energy vanishes and the two hyperbols $|\tilde{x}|_{\pm} = \pm \sqrt{|\tilde{y}|^2 + 1/|\mu|}$ (resp. $|\tilde{y}|_{\pm} = \pm \sqrt{|\tilde{x}|^2 - 1/|\mu|}$) are given by two cones $|\tilde{y}| = +|\tilde{x}|$ and $|\tilde{y}| = -|\tilde{x}|$ with opposite apex. In this case, the pseudo 3-sphere shrinks to the cones

$$\lim_{|\mu| \rightarrow \infty} \tilde{\mathbb{S}}^{1,2}(\mu) \simeq \{|\tilde{y}| = \pm |\tilde{x}|\} \quad (2.35)$$

We refer to this conical limit as an " $SU(1, 1)$ singularity" in analogy with the $SU(2)$ singularity where the 3-sphere $\tilde{\mathbb{S}}^3(\mu) \rightarrow 0$.

Similar relations to eqs(2.27) can be also written down for the dual manifold; we have

$$\begin{aligned}\tilde{\mathbb{S}}^3(\mu) &: \tilde{\xi}_1^2 + \tilde{\xi}_2^2 + \tilde{\xi}_3^2 + \tilde{\xi}_4^2 = \frac{1}{|\mu|} \\ \tilde{\mathbb{S}}^{1,2}(\mu) &: \tilde{\xi}_1^2 + \tilde{\xi}_2^2 - \tilde{\xi}_3^2 - \tilde{\xi}_4^2 = \frac{1}{|\mu|}\end{aligned}\quad (2.36)$$

with

$$\tilde{\mathbb{S}}^{1,2}(\mu): \quad \tilde{\xi}_{1\pm} = \pm \sqrt{\frac{1}{|\mu|} + \tilde{\xi}_4^2 + \tilde{\xi}_3^2 - \tilde{\xi}_2^2} \quad (2.37)$$

In the limit $|\mu| \rightarrow \infty$, they contract to

$$\begin{aligned}\tilde{\mathbb{S}}^3(\infty) &: \tilde{\xi}_1^2 + \tilde{\xi}_2^2 + \tilde{\xi}_3^2 + \tilde{\xi}_4^2 = 0 \\ \tilde{\mathbb{S}}^{1,2}(\infty) &: \tilde{\xi}_1^2 + \tilde{\xi}_2^2 - \tilde{\xi}_3^2 - \tilde{\xi}_4^2 = 0\end{aligned}\quad (2.38)$$

with the two hyperboloids

$$\tilde{\mathbb{S}}^{1,2}(\infty): \quad \tilde{\xi}_{1\pm} = \pm \sqrt{\tilde{\xi}_4^2 + \tilde{\xi}_3^2 - \tilde{\xi}_2^2} \quad (2.39)$$

colliding at $\tilde{\xi}_4^2 + \tilde{\xi}_3^2 = \tilde{\xi}_2^2$.

In what follows, we will use the dual $\tilde{\mathcal{V}}$ containing $\tilde{\mathbb{S}}^3$ and $\tilde{\mathbb{S}}^{1,2}$ instead of the \mathcal{V} having \mathbb{S}^3 and the pseudo $\mathbb{S}^{1,2}$; this is because $\tilde{\mathbb{S}}^3$ and $\tilde{\mathbb{S}}^{1,2}$ have direct applications in our calculations; they are intimately linked with the simple root α of two real forms of $SL(2, \mathbb{C})$ and the associated homological 2-cycle \mathcal{C}_α of its algebraic geometry; i.e:

$$\alpha \in \Phi_{SL_2} \quad \leftrightarrow \quad \mathcal{C}_\alpha \in H_2(SL_2)$$

For convenience, we will refer to $\tilde{\mathcal{V}}$ simply like \mathcal{X} and often drops out the twilda label on the $\tilde{\mathbb{S}}^3$ and $\tilde{\mathbb{S}}^{1,2}$ by referring to them just as \mathbb{S}^3 and $\mathbb{S}^{1,2}$.

Given (2.26) as well as (2.29) and (2.35), an interesting question arises here: How much differs the non compact 3D hypersurface \mathcal{X} from the flat $\mathbb{C}^3 \simeq \mathbb{R}^6$?. We answer this question by considering two local representations:

A) \mathcal{X} as cotangent $\mathbf{T}^*\tilde{\mathbb{S}}^3(\mu)$ or $\mathbf{T}^*\tilde{\mathbb{S}}^{1,2}(\mu)$

Starting from (2.17) and taking the limit $\frac{1}{\mu} \rightarrow 0$, one learns that \mathcal{X} is given by a complex 3D conifold [64–66]

$$\mathcal{X}: \quad vx - yu = \frac{1}{\mu} \rightarrow 0 \quad (2.40)$$

By setting setting $x = a + b$ and $v = \bar{a} - \bar{b}$ as well as $y = c + d$ and $u = \bar{c} - \bar{d}$ with complex (a,b,c,d), we can expand $vx - yu = 1/\mu$ as follows (by using $\text{Im } \mu = 0$, and $\text{Re } \mu \gg 0$)

$$\begin{aligned}\mathfrak{B} &: |a|^2 + |d|^2 - |b|^2 - |c|^2 = +\frac{1}{\mu\bar{\mu}} \text{Re } \mu = \frac{1}{\text{Re } \mu} \\ \mathfrak{F} &: b\bar{a} - a\bar{b} + d\bar{c} - c\bar{d} = -\frac{1}{\mu\bar{\mu}} \text{Im } \mu = 0\end{aligned}\quad (2.41)$$

These two real equations can be solved in two interesting ways given by the two real forms SU(2) and SU(1,1) as follows

SU(2)	:	$b = c = 0$,	SU(1,1)	:	$b = d = 0$	(2.42)
$\mathfrak{B} = \mathbb{S}^3$:	$ a ^2 + d ^2 = \frac{1}{\text{Re}\mu}$		$\mathfrak{B} = \mathbb{S}^{1,2}$:	$ a ^2 - c ^2 = \frac{1}{\text{Re}\mu}$	
\mathcal{X}	=	$T^*\mathbb{S}^3$		\mathcal{X}	=	$T^*\mathbb{S}^{1,2}$	

with T^*E designating the cotangent space.

B) \mathcal{X} as complex $\mathbb{C} \times T^*\tilde{\mathbb{S}}^2(\mu)$ or $\mathbb{C} \times T^*\tilde{\mathbb{S}}^{1,1}(\mu)$:

These are local complex fibrations of \mathcal{X} that may be derived by using: (i) the cotangent $T^*\mathbb{S}^3$ of and the $T^*\mathbb{S}^{1,2}$ of eq(2.42); and (ii) substituting the Hopf fibration $\mathbb{S}^3 \simeq \mathbb{S}^2 \times \mathbb{S}^1$ and $\mathbb{S}^{1,2} \simeq \mathbb{S}^{1,1} \times \mathbb{S}^1$. We end up with

SU(2)/U(1)	:	$b = c = 0$,	$a \equiv e^{i\vartheta}a$	$d \equiv e^{-i\vartheta}d$	(2.43)
$\mathfrak{B} = \mathbb{S}^2$:	$ a ^2 + d ^2 = \frac{1}{\text{Re}\mu}$		$\mathbb{C} \times \Sigma_2 = \mathbb{C} \times T^*\mathbb{S}^2$		
\mathcal{X}	=	$\mathbb{C} \times \Sigma_2 = \mathbb{C} \times T^*\mathbb{S}^2$		$T^*\mathbb{S}^2$		

where we have used $\mathbb{C} \simeq T^*\mathbb{S}^1$ and

SU(1,1)/U(1)	:	$b = d = 0$,	$a \equiv e^{i\vartheta}a$	$c \equiv e^{-i\vartheta}d$	(2.44)
$\mathfrak{B} = \mathbb{S}^{1,1}$:	$ a ^2 - d ^2 = \frac{1}{\text{Re}\mu}$		$\mathbb{C} \times \Sigma_{1,1} = \mathbb{C} \times T^*\mathbb{S}^{1,1}$		
\mathcal{X}	=	$\mathbb{C} \times \Sigma_{1,1} = \mathbb{C} \times T^*\mathbb{S}^{1,1}$		$T^*\mathbb{S}^{1,1}$		

To establish these local fibrations explicitly, we (i) denote by (z_1, z_2, z_3) the three free complex variables parameterising the flat \mathbb{C}^3 , and (ii) look for expressing the relation $xv - yv = 1/\mu$ as a non linear function $v = f(z_1, z_2, z_3)$ with variable v as in (2.17). To that purpose, we set $(z_1, z_2, z_3) = (1/x, y, -u)$; this allows to factorise the constraint relation (2.17) like

$$\begin{aligned}
 v &= z_1 G(z) = z_1 \left[\frac{1}{\mu} - \sigma(z) \right] \\
 \sigma &= z_2 z_3
 \end{aligned}
 \tag{2.45}$$

where z_1 parameterises a complex line \mathbb{C} and where $G(z)$ describes a complex surface Σ given by the equation $G(z) = 1/\mu - \sigma(z)$ with zeros at $z_2 z_3 = 1/\mu$ describing a projective line. With the factorisation (2.45), we then have $\mathcal{X} = \mathbb{C} \times \Sigma$. Notice that by substituting $z_2 = y$ and $z_3 = -u$, the zeros equation $z_2 z_3 = 1/\mu$ reads as $yu = -1/\mu$.

Moreover, substituting $u = \mp \bar{y}$ giving the loci of the \mathbb{S}^3 and $\mathbb{S}^{1,2}$ as given by (2.25), we end up with

$$SU(2) \quad : \quad \bar{y}y = +\frac{1}{\operatorname{Re} \mu} \quad , \quad SU(1,1) \quad : \quad \bar{y}y = -\frac{1}{\operatorname{Re} \mu} \quad (2.46)$$

putting a constraint on the sign of $\operatorname{Re} \mu$; which is positive for $SU(2)$ and negative for $SU(1,1)$. Recall that for the case where $z_2 z_3 = z^2$, we have an ALE space with $SU(2)$ singularity describing a complex surface $\Sigma_2 \sim \mathbb{C} \times \mathbb{S}^2$ with shrinking 2-sphere to a point. This surface has a graphic representation in terms of toric diagrams as depicted by the **Figure 3**.




Figure 3: A toric representation of the complex surface $\Sigma_2 \sim \mathbb{C} \times \mathbb{C}\mathbb{P}^1$. The compact line is in red color. The non compact directions is given by the line \mathbb{C} . In real variables, we have $\mathcal{X} \simeq \mathbb{R}^2 \times \mathbb{S}_\alpha^2$; the 2-sphere (2-cycle \mathcal{C}_α) is associated with the simple root α of $\mathfrak{su}(2)$.

2.1.3 Even and odd parts of $SL(2|2)$

The even part of $SL(2|2)_\bar{0}$ is given by $SL(2)_1 \times SL(2)_2 \times \mathbb{C}^*$; it has two isospin copies; and as such it has a discrete \mathbb{Z}_2 outer automorphism acting by the transposition $SL(2)_1 \leftrightarrow SL(2)_2$. To implement this property, the geometry of $SL(2|2)_\bar{0}$ denoted like \mathcal{X}_{120} factorises as $\mathcal{X}_{12} \times \mathbb{C}^*$ with \mathcal{X}_{12} given by the product $\mathcal{X}_1 \times \mathcal{X}_2$. The complex manifold \mathcal{X}_{12} is a complex 6D manifold sitting in the $\mathbb{C}_{12}^{4|4} \sim \mathbb{C}_1^4 \times \mathbb{C}_2^4$ parameterised by (x_1, y_1, u_1, v_1) and (x_2, y_2, u_2, v_2) . Because $SL(2)$ has two real forms $SU(2)$ and $SU(1,1)$, one distinguishes the two particular real forms of $PSL(2|2)_\bar{0}$ given by

$$\begin{aligned} \text{(i)} \quad & : \quad PSU(2|2)_\bar{0} \quad = \quad SU(2)_1 \times SU(2)_2 \\ \text{(ii)} \quad & : \quad PSU(1,1|2)_\bar{0} \quad = \quad SU(1,1)_1 \times SU(2)_2 \end{aligned} \quad (2.47)$$

For local graphic representations, we use the $SU(2|2)$ having compact 2-spheres \mathbb{S}_1^2 and \mathbb{S}_2^2 . Using the toric graph of the **Figure 3**, the \mathcal{X}_{12} can be represented by two disconnected




Figure 4: Toric graph of the fibration $\mathfrak{B}_{12} \times \mathfrak{F}_{12}$ describing the $(\Sigma_2)_1 \times (\Sigma_2)_2$ geometry of the $SU(2)_1 \times SU(2)_2$ symmetry without bi-fundamentals. The red segments represent $\mathbb{C}\mathbb{P}^1$ curves.

toric diagrams [67–69] as depicted by the **Figure 4**. Notice that the two (red) 2-spheres in $\mathbb{S}_1^2 \times \mathbb{S}_2^2$ have an interpretation in terms of the even¹ roots β_1 and β_2 of the even $psu(2|2)_{\bar{0}} = su(2) \oplus su(2)$.

The complex 6D fibration \mathcal{X}_{12} is defined by the algebraic equations

$$\begin{aligned} \det \mathbf{X}_1 &: x_1 y_1 - u_1 v_1 = \frac{1}{\mu} \\ \det \mathbf{X}_2 &: x_2 y_2 - u_2 v_2 = \frac{1}{\mu} \end{aligned} \quad (2.48)$$

they can be merged into one constraint relation given by Berezinian $Ber(\mathbf{M}) = 1$ of the super matrix \mathbf{M} combining \mathbf{X}_1 and \mathbf{X}_2 as follows

$$\mathbf{M} = \begin{pmatrix} \mathbf{X}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{X}_2 \end{pmatrix}, \quad \mathbf{X}_i = \begin{pmatrix} x_i & u_i \\ y_i & v_i \end{pmatrix} \quad (2.49)$$

Recall that the $Ber(\mathbf{M})$ of the above super matrix is given by $(\det \mathbf{X}_1)(\det \mathbf{X}_2)^{-1}$; it reads explicitly as follows

$$\frac{x_1 y_1 - u_1 v_1}{x_2 y_2 - u_2 v_2} = 1 \quad (2.50)$$

it describes a 7D hypersurface in $\mathbb{C}^{4|4}$ and gives the complex 7D geometry of $SL(2|2)_{\bar{0}}$. In this regards, notice the two following:

- (i) the super determinant $Ber(\mathbf{M})$ is invariant under the scaling $\det \mathbf{X}_i \rightarrow \lambda^2 \det \mathbf{X}_i$ induced from the projective transformation $\mathbf{X}_i \rightarrow \lambda \mathbf{X}_i$.
- (ii) this scaling describes precisely the central element H_0 generating the $GL(1, \mathbb{C})$ subgroup of $SL(2|2)_{\bar{0}}$; thus permitting to define the 14D geometry of $PSL(2|2)_{\bar{0}}$.

¹ Sometimes these simple roots are denoted like $\beta_1 = \alpha_1$ and $\beta_2 = \alpha_3$.

For the eight dimensional odd part $\mathfrak{sl}(2|2)_{\bar{1}}$; it is given by the bi-fundamental representations $\mathbf{2}_1 \otimes \mathbf{2}_2^c$ and $\mathbf{2}_1^c \otimes \mathbf{2}_2$ of $SL(2, \mathbb{C})_1 \times SL(2, \mathbb{C})_2$. These bi-fundamentals are sketched by the **Figure 5**. The two nodes in this **Figure** corresponds to $\mathbf{2}_1 \otimes \mathbf{2}_1^c$ and




Figure 5: On left, the nodes in red and blue represent the two $\mathfrak{sl}(2, \mathbb{C})$ s making the even $\mathfrak{psl}(2|2)_{\bar{0}}$. The bridge between the two nodes given by the bi-fundamentals represent the odd generators of $\mathfrak{psl}(2|2)_{\bar{1}}$ relating the two $\mathfrak{sl}(2, \mathbb{C})$ s. A dual description of this graph is given by the super Dynkin diagrams of the **Figure 15**. On the right, the graphic representation in terms of the roots.

$\mathbf{2}_2 \otimes \mathbf{2}_2^c$; the links to $\mathbf{2}_i \otimes \mathbf{2}_j^c$ and $\mathbf{2}_i^c \otimes \mathbf{2}_j$. The bridge between the two (red) 2-spheres is given by (green) fermionic 2-cycles generated by α_2 and $\alpha_1 + \alpha_2 + \alpha_3$ as well as $\alpha_1 + \alpha_2$ and $\alpha_2 + \alpha_3$. They are represented by the green lines shown on the **Figure 6**. They permit to glue together the two toric copies of the **Figure 4**.




Figure 6: Toric representation of the gluing of the surfaces Σ_1 and Σ_2 in the super-geometry of $PSL(2|2)$. The red segments refer to the even cycles; the four green ones designate the odd cycles associated with the odd roots.

In summary, the geometric description of the 15 dimensional $\mathfrak{sl}(2|2)$ is given by the complex super matrix (2.3-2.5) with super determinant $Ber(\mathcal{M}) = 1$. This constraint

defines a complex 15 dim hypersurface in $\mathbb{C}^{8|8}$ reading explicitly as follows

$$\frac{\det(\mathbf{X}_1)}{\det(\mathbf{X}_2)} \det[\mathbf{I} - \mathbf{X}_1^{-1} \mathbf{Y}_1 \mathbf{X}_2^{-1} \mathbf{Y}_2] = 1 \quad (2.51)$$

It shows that (i) the \mathbf{X}_1 and \mathbf{X}_2 are coupled through the bi-fundamentals \mathbf{Y}_1 and \mathbf{Y}_2 like $\mathbf{X}_1^{-1} \mathbf{Y}_1 \mathbf{X}_2^{-1} \mathbf{Y}_2$ and (ii) the $Ber(\mathbf{M}) = 1$ is invariant under the scaling $\mathbf{M} \rightarrow \lambda \mathbf{M}$ generated by the projective transformations $\mathbf{X}_i \rightarrow \lambda \mathbf{X}_i$ and $\mathbf{Y}_i \rightarrow \lambda \mathbf{Y}_i$. By taking the coset with respect to this projective scaling, we get the complex 14D manifold describing $PSL(2|2)$.

2.2 Decompactifying the \mathbb{S}^2 inside $SL(2, \mathbb{C})$

Here, we consider the representation $\mathcal{X} \simeq \mathbb{C}^2 \times \mathbb{CP}^1$ to investigate the decompactification of projective \mathbb{CP}^1 inside $SL(2, \mathbb{C})$ towards the non compact line \mathbb{C} . This decompactification is done here with the idea of applying it when studying the contraction of the exceptional super group $D(2, 1; \zeta)$ down to $PSL(2|2)$. It is achieved by taking the infinite limit $|\mu| \rightarrow \infty$ of the area of the real 2-sphere (2.35); or equivalently the shrinking of its dual given by the limit $|\tilde{\mu}| \rightarrow 0$ as shown by the **Figure 7**.




Figure 7: Variation of the area of the real 2-sphere \mathbb{S}^2 (in blue) and its dual $\tilde{\mathbb{S}}^2$ (in red). On the left, the sphere is given by $|x^2| + |y^2| = |\mu|$ and on the right it is given by $|\tilde{x}^2| + |\tilde{y}^2| = |\tilde{\mu}|$. They are functions of the Kähler parameter ρ . The area of the real \mathbb{S}^2 explodes at the singular $\zeta = -1, 0$ in eq(2.59) while the dual $\tilde{\mathbb{S}}^2$ shrinks.

2.2.1 $SL(2, \mathbb{C})$ inside $D(2, 1; \zeta)_{\bar{0}}$

We start by recalling that the even part of the exceptional group $D(2, 1; \zeta)_{\bar{0}}$ is given by the product of three isospin groups as follows

$$\begin{aligned} D(2, 1; \zeta)_{\bar{0}} &= SL(2, \mathbb{C})_1 \times SL(2, \mathbb{C})_2 \times SL(2, \mathbb{C})_3 \\ &\simeq SO(4, \mathbb{C})_{12} \times SL(2, \mathbb{C})_3 \end{aligned} \quad (2.52)$$

The isospin group we want to decompactify is given by the third factor $SL(2, \mathbb{C})_3$ by using the analysis considered in deriving eq(2.35). Instead of using the complex 2×2 matrix \mathbf{X}_3 like

$$\mathbf{X}_3 = \begin{pmatrix} x_3 & u_3 \\ y_3 & v_3 \end{pmatrix} \quad \rightarrow \quad \mathbf{X}_3 = \begin{pmatrix} x & u \\ y & v \end{pmatrix} \quad (2.53)$$

with determinant $xv - yu = \mu$, we work with the dual parametrisation

$$\mathbf{X}_3^{-1} = \frac{1}{\mu} \begin{pmatrix} v_3 & -u_3 \\ -y_3 & x_3 \end{pmatrix} \quad \rightarrow \quad \mathbf{X}_3^{-1} = \frac{1}{\mu} \begin{pmatrix} v & -u \\ -y & x \end{pmatrix} \quad (2.54)$$

with determinant as

$$\mathcal{X}_3 : xv - yu = \frac{1}{\mu} \quad , \quad \mathcal{X}_3 := \tilde{\mathcal{V}}_3 \quad (2.55)$$

defining a complex 3D hypersurface \mathcal{X}_3 in \mathbb{C}^4 . To study the decompactification of the $SL(2, \mathbb{C})_3$, we focus on the locus $(v, u) = (\bar{x}, -\bar{y})$ where lives the (dual) 3-sphere $\tilde{\mathbb{S}}^3$,

$$|x|^2 + |y|^2 = \frac{1}{\varrho} \quad , \quad \varrho = |\mu| \quad (2.56)$$

On this locus, it lives also a 2-sphere $\tilde{\mathbb{S}}^2$ descending from the Hopf fibration of $\tilde{\mathbb{S}}^3 \simeq \tilde{\mathbb{S}}^2 \times \tilde{\mathbb{S}}^1$. Under the identification $(x, y) \simeq (e^{i\theta}x, e^{-i\theta}y)$, one gets a 2-sphere with area given by

$$\mathcal{A}(\mathbb{S}^2) = 4\pi\varrho \quad , \quad \mathcal{A}(\tilde{\mathbb{S}}^2) = \frac{4\pi}{\varrho} \quad (2.57)$$

2.2.2 Decompactified \mathbb{S}^2 and shrinking $\tilde{\mathbb{S}}^2$

From eq(2.57), we see that the decompactification of the 2-sphere towards \mathbb{C} can be realised by taking the limit $\varrho \rightarrow \infty$ in the sphere; because $\frac{1}{\varrho} \rightarrow 0$, the dual $\tilde{\mathbb{S}}^2$ shrinks then to a point. Below, we think about the real parameter ϱ as the absolute value of a complexified Kahler parameter $t(\zeta)$. This Kahler modulus is a function of the parameter ζ of the exceptional super group $D(2, 1; \zeta)$ as follows

$$t = \zeta(\zeta + 1) \quad , \quad t = \frac{1}{\varrho} e^{i\chi} \quad (2.58)$$

It vanishes for the values $\zeta = 0, -1$ where the area of the real 2-sphere

$$\mathcal{A}(\mathbb{S}^2) = \frac{4\pi}{|\zeta(\zeta + 1)|} \quad (2.59)$$

diverges towards the volume of \mathbb{R}^2 . For these values, the dual area $\mathcal{A}(\tilde{\mathbb{S}}^2) = 4\pi |\zeta(\zeta + 1)|$ shrinks. The variation of this area is given by the **Figure 7**.

3 Graded 2-cycles in $D(2, 1; \zeta)$

In this section, we first describe the four parametric super Dynkin graphs $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ of complex $\mathfrak{d}(2, 1; \zeta)$ and their discrete outer automorphisms by using (i) the complex parameter ζ and (ii) the *Kaplansky* variables (s_1, s_2, s_3) constrained as $\sum s_i = 0$ and $\prod s_i \neq 0$ [49, 50]. Then, we use the root/2-cycle correspondence

$$\begin{array}{ccc} D(2, 1; \zeta) & : & H_2 [D(2, 1; \zeta)] \\ \text{algebraic roots } \boldsymbol{\alpha} & \leftrightarrow & \text{geometric 2-cycles } \tilde{\mathcal{C}}_{\tilde{\boldsymbol{\alpha}}} \end{array} \quad (3.1)$$

and the ansatz [1, 70]

$$D(2, 1; \zeta) \quad \leftrightarrow \quad \text{continuous deformation of } \text{OSp}(4|2) = D(2, 1) \quad (3.2)$$

to extract useful information on the parametric super 2-cycles $\tilde{\mathcal{C}}_{\tilde{\boldsymbol{\alpha}}}[\zeta]$ in the geometry of complex $D(2, 1; \zeta)$. These parametric 2-cycles are functions of ζ ; they are thought of as a continuous deformation of the orthosymplectic 2-cycles $\mathcal{C}_{\boldsymbol{\alpha}}^{osp}$ of the super $\text{OSp}(4|2)$ further described in the [Appendix B](#). Also, they contract in the super 2-cycles of $\text{PSL}(2|2)$ for $\zeta = -1$.

Recall that $\mathfrak{d}(2, 1; \zeta)$ has four possible basis sets $\Pi_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ of simple roots that we denote like $\{\tilde{\boldsymbol{\alpha}}_1^{(\eta)}, \tilde{\boldsymbol{\alpha}}_2^{(\eta)}, \tilde{\boldsymbol{\alpha}}_3^{(\eta)}\}$ with index $\eta = 0, 1, 2, 3$; the twild label is used in order to distinguish the roots of $\mathfrak{d}(2, 1; \zeta)$ from their homologue² $\{\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2, \boldsymbol{\alpha}_3\}$ of $\mathfrak{psl}(2|2)$ and $\mathfrak{osp}(4|2)$. These simple roots are functions of the deformation parameters; i.e:

$$\tilde{\boldsymbol{\alpha}}_i^{(\eta)}(\boldsymbol{\alpha}^{osp}; \mathfrak{s}) \quad \Leftrightarrow \quad \tilde{\boldsymbol{\alpha}}_i^{(\eta)}(\boldsymbol{\alpha}^{osp}; \zeta) \quad (3.3)$$

they are constrained by consistency conditions including the $\mathfrak{osp}(4|2)$ and the $\mathfrak{sl}(2|2)$ symmetries sitting at $\zeta = \pm 1$; thus requiring $\tilde{\boldsymbol{\alpha}}_i|_{\zeta=1} = \boldsymbol{\alpha}_i^{osp(4|2)}$ and $\tilde{\boldsymbol{\alpha}}_i|_{\zeta=-1} = \boldsymbol{\alpha}_i^{sl(2|2)}$. Other useful properties on $\mathfrak{d}(2, 1; \zeta)$ such as the grading of the roots (homological 2-cycles) and outer automorphisms are reported in the [Appendix C](#).

3.1 Basis sets $\tilde{\Pi}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ and outer automorphisms

The Lie superalgebra $\mathfrak{d}(2, 1; \zeta)$ of the super group $D(2, 1; \zeta)$ is a parametric graded algebra with complex parameter ζ ; it has rank 3 and 17 dimensions and was interpreted as a deformation of the orthosymplectic $\mathfrak{osp}(4|2)$ sitting at $\zeta = 1$ [1, 71–75]. Also, the $\mathfrak{d}(2, 1; \zeta)$ has the same rank 3 as for $\mathfrak{sl}(2|2)$; but two dimensions bigger.

² To avoid confusion between the roots systems of $\mathfrak{sl}(2|2)$ and $\mathfrak{osp}(4|2)$, we sometimes use the convention notation $\boldsymbol{\alpha}_i^{sl(2|2)}$ and $\boldsymbol{\alpha}_i^{osp}$.

Its even part $\mathfrak{d}(2, 1; \zeta)_{\bar{0}}$ has 9 dimensions; it is given by

$$sl(2, \mathbb{C})_1 \oplus sl(2, \mathbb{C})_2 \oplus sl(2, \mathbb{C})_3 \quad (3.4)$$

exhibiting a manifest \mathcal{S}_3 automorphism symmetry that permutes the three isospins. Denoting the simple roots of these $sl(2)_i$ s by β_i (with diagonal intersection matrix $\beta_i \cdot \beta_j = 4s_i \delta_{ij}$), the group elements σ of \mathcal{S}_3 acts like $\beta_i \rightarrow \beta_{\sigma(i)}$.

The odd part $\mathfrak{d}(2, 1; \zeta)_{\bar{1}}$ is eight dimensional; it is given by the tri-fundamental representation $\mathbf{2}_1 \otimes \mathbf{2}_2 \otimes \mathbf{2}_3$ as depicted by the **Figure 8** extending the **Figure 5**. From this




Figure 8: The nodes in red, blue and green represent the three $sl(2, \mathbb{C})$ s making the even $D(2, 1; \zeta)_{\bar{0}}$. The bridges between the three nodes given by the tri-fundamentals represent the odd generators of $D(2, 1; \zeta)_{\bar{1}}$ relating the three $sl(2, \mathbb{C})$ s.

description, we see that the 14 dimensional $PSL(2|2)_{\bar{0}}$ sitting at $\zeta = -1$ descends from $D(2, 1; \zeta)_{\bar{0}}$ by decompactifying $SL(2, \mathbb{C})_3$ into \mathbb{C}^3 by taking the area (2.59) to infinity. In Lie algebra language we have for $\zeta \rightarrow -1$,

$$\mathfrak{d}(2, 1; \zeta)_{\bar{0}} \quad \rightarrow \quad \mathfrak{d}(2, 1; -1)_{\bar{0}} = psl(2|2)_{\bar{0}} \oplus \mathbb{C} \oplus \mathbb{C} \oplus \mathbb{C} \quad (3.5)$$

The exceptional $\mathfrak{d}(2, 1; \zeta)$ has four super Dynkin diagrams $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2, 1; \zeta)}^{(\eta)}$ given by the pictures (a), (b), (c) and (d) of the **Figure 9**. The second picture 9-(b) with odd $\tilde{\alpha}_2$ will be further discussed in **subsection 3.2** in connection with the correspondence (3.1). As for $sl(2|2)$, here also the sets $\Pi_{\mathfrak{d}(2, 1; \zeta)}^{(\eta)}$ (below $\tilde{\Pi}^{(\mu)}$) labeled by $\eta = 0, 1, 2, 3$ giving the four possible bases of simple roots spanning the root system $\Phi_{\mathfrak{d}(2, 1; \zeta)}^{(\eta)}$ (for short $\tilde{\Phi}^{(\eta)}$). Each basis set $\tilde{\Pi}^{(\eta)}$ is generated by three simple roots $\{\tilde{\alpha}_1, \tilde{\alpha}_2, \tilde{\alpha}_3\}$ characterised by the Z_2 -parity of the $\tilde{\alpha}_i$'s.




Figure 9: The four super Dynkin diagrams of the Lie superalgebra $D(2,1;\zeta)$. They are as in [44] where here we have used the S_i parameters constrained like $S_1+S_2+S_3 = 0$ and $S_1S_2S_3 \neq 0$. These diagrams will be imagined in terms of the intersection matrix \mathcal{I}_{ij} instead of \mathcal{K}_{ij} .

3.1.1 Gradation of super Dynkin $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$

Depending on the parity of the simple roots, we distinguish four graded roots systems

basis	$\tilde{\Pi}^{(1)}$	$\tilde{\Pi}^{(2)}$	$\tilde{\Pi}^{(3)}$	$\tilde{\Pi}^{(0)}$
root system	$\tilde{\Phi}^{(1)}$	$\tilde{\Phi}^{(2)}$	$\tilde{\Phi}^{(3)}$	$\tilde{\Phi}^{(0)}$
\mathcal{S}_3 repres.	triplet			singlet

(3.6)

The three $\tilde{\Pi}^{(i)}$ (resp. $\tilde{\Phi}^{(i)}$) form a triplet of the discrete permutation group \mathcal{S}_3 while $\tilde{\Phi}^{(0)}$ is a singlet. Their roots contents are as follows:

(1) **the root system $\tilde{\Phi}^{(1)}$** : (simple $\tilde{\alpha}_1$ odd)

It is generated by the simple roots $\tilde{\alpha}_1, \tilde{\alpha}_2, \tilde{\alpha}_3$ with $\tilde{\alpha}_1$ odd and $\tilde{\alpha}_2, \tilde{\alpha}_3$ even. The seven positive roots of $\tilde{\Phi}^{(1)}$ are collected in the following Table

root systems	fermionic subset	bosonic subset
$\tilde{\Phi}^{(1)}$	$\tilde{\alpha}_1,$ $\tilde{\alpha}_1 + \tilde{\alpha}_2, \quad \tilde{\alpha}_1 + \tilde{\alpha}_3,$ $\tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3$	$\tilde{\alpha}_2, \quad \tilde{\alpha}_3,$ $2\tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3$

(3.7)

The 6 even step operators $E_{\pm\beta_i}$ labeled by the even roots

$$\begin{aligned}
\beta_1 &= 2\tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3 & \leftrightarrow & \mathcal{C}_{\beta_1} := \mathbb{S}_\varrho^2 \text{ or } \mathbb{S}_\varrho^{1,1} \\
\beta_2 &= \tilde{\alpha}_2 & \leftrightarrow & \mathcal{C}_{\beta_2} \\
\beta_3 &= \tilde{\alpha}_3 & \leftrightarrow & \mathcal{C}_{\beta_3}
\end{aligned}$$
(3.8)

generate $\oplus_i sl(2, \mathbb{C})_i$. In (3.8), we have also given the 2-sphere \mathbb{S}_ℓ^2 or the pseudo $\mathbb{S}_\ell^{1,1}$ associated with the even roots β_i characterising the isospin $sl(2, \mathbb{C})_{\beta_i}$. Here β_1 is the long root.

(2) the root systems $\tilde{\Phi}^{(2)}$ (with odd $\tilde{\alpha}_2$) and $\tilde{\Phi}^{(3)}$ (with odd $\tilde{\alpha}_3$)

The root contents of these systems are obtained from $\tilde{\Phi}^{(1)}$ by performing cyclic permutations of the pictures (a), (b), (c) of the **Figure 9**. This discrete symmetry acts as

$$(\tilde{\alpha}_1, \tilde{\alpha}_2, \tilde{\alpha}_3) \quad \rightarrow \quad (\tilde{\alpha}_2, \tilde{\alpha}_3, \tilde{\alpha}_1) \quad \rightarrow \quad (\tilde{\alpha}_3, \tilde{\alpha}_1, \tilde{\alpha}_2) \quad (3.9)$$

The three basis sets $(\tilde{\Pi}_1, \tilde{\Pi}_2, \tilde{\Pi}_3)$ form then a triplet under the \mathcal{Z}_3 automorphism symmetry. Recall that this discrete symmetry \mathcal{Z}_3 is a subgroup of the symmetric group \mathcal{S}_3 permuting the three canonical weight vectors $(\epsilon_1, \epsilon_2, \epsilon_3)$ given by eq(C.1) reported in [Appendix C](#).

The contents of the sets $\tilde{\Phi}^{(2)}$ and $\tilde{\Phi}^{(3)}$ are collected in the following Table

root systems	fermionic subset	bosonic subset	
$\tilde{\Phi}^{(2)}$	$\tilde{\alpha}_2,$ $\tilde{\alpha}_2 + \tilde{\alpha}_1, \quad \tilde{\alpha}_2 + \tilde{\alpha}_3,$ $\tilde{\alpha}_2 + \tilde{\alpha}_1 + \tilde{\alpha}_3$	$\tilde{\alpha}_1, \quad \tilde{\alpha}_3,$ $\tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3$	(3.10)
$\tilde{\Phi}^{(3)}$	$\tilde{\alpha}_3,$ $\alpha_3 + \tilde{\alpha}_2, \quad \tilde{\alpha}_3 + \tilde{\alpha}_1,$ $\tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3$	$\tilde{\alpha}_1, \quad \tilde{\alpha}_2,$ $\tilde{\alpha}_1 + \tilde{\alpha}_2 + 2\tilde{\alpha}_3$	

Here the 6 even step operators $E_{\pm\beta_i}$ generating $\oplus_i sl(2, \mathbb{C})_i$ are labeled for the case of $\tilde{\Phi}_{\mathfrak{g}(2,1;\zeta)}^{(2)}$ by the even roots

$$\begin{aligned} \tilde{\Phi}^{(2)} : \quad \beta'_1 &= \tilde{\alpha}_1 & \leftrightarrow & \mathcal{C}_{\beta'_1} \\ \beta'_2 &= \tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3 & \leftrightarrow & \mathcal{C}_{\beta'_2} := \mathbb{S}_\ell^2 \text{ or } \mathbb{S}_\ell^{1,1} \\ \beta'_3 &= \tilde{\alpha}_3 & \leftrightarrow & \mathcal{C}_{\beta'_3} \end{aligned} \quad (3.11)$$

Because $\beta'_i \cdot \beta'_j = 0$ for $i < j$, these 3-spheres have no direct intersections; they couple through the odd roots. For the case of $\tilde{\Phi}^{(3)}$, we have

$$\begin{aligned} \tilde{\Phi}^{(3)} : \quad \beta''_1 &= \tilde{\alpha}_1 & \leftrightarrow & \mathcal{C}_{\beta''_1} \\ \beta''_2 &= \tilde{\alpha}_2 & \leftrightarrow & \mathcal{C}_{\beta''_2} \\ \beta''_3 &= \tilde{\alpha}_1 + \tilde{\alpha}_2 + 2\tilde{\alpha}_3 & \leftrightarrow & \mathcal{C}_{\beta''_3} := \mathbb{S}_\ell^2 \text{ or } \mathbb{S}_\ell^{1,1} \end{aligned} \quad (3.12)$$

(3) **fourth root system** $\tilde{\Phi}^{(0)}$:

It is generated by the simple roots $\tilde{\alpha}_1, \tilde{\alpha}_2, \tilde{\alpha}_3$ which all of them are odd roots.

Here, the seven positive roots are given by

root systems	fermionic subset	bosonic subset	
$\tilde{\Phi}^{(0)}$	$\tilde{\alpha}_1, \tilde{\alpha}_2, \tilde{\alpha}_3,$ $\tilde{\alpha}_3 + \tilde{\alpha}_1 + \tilde{\alpha}_2$	$\tilde{\alpha}_1 + \tilde{\alpha}_2, \tilde{\alpha}_2 + \tilde{\alpha}_3$ $\tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3$	(3.13)

For this special case, the 6 even step operators $E_{\pm\beta_i}$ generating $\oplus_i sl(2, \mathbb{C})_i$ are labeled by the even roots

$$\begin{aligned}
 & \beta_1 = \tilde{\alpha}_1 + \tilde{\alpha}_2 && \leftrightarrow && \mathcal{C}_{\beta_1} \\
 \text{case } \tilde{\Phi}^{(0)} : & \beta_2 = \tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3 && \leftrightarrow && \mathcal{C}_{\beta_2} \\
 & \beta_3 = \tilde{\alpha}_2 + \tilde{\alpha}_3 && \leftrightarrow && \mathcal{C}_{\beta_3}
 \end{aligned} \tag{3.14}$$

3.1.2 Super 2-cycles in $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$

Using (i) the correspondence (3.1) associating a homological 2-cycle $\tilde{\mathcal{C}}_{\tilde{\alpha}}$ to each root $\tilde{\alpha}$ in the set $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$, and (ii) knowing the root systems content (3.7-3.13), one ends up with four parametric systems of homological 2-cycles that we denote like

$$\tilde{\mathcal{C}}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)} \quad , \quad \eta = 0, 1, 2, 3$$

These 2-cycle systems are in 1:1 with the four $\tilde{\Phi}^{(\eta)}$ and then with the four parametric super Dynkin diagrams

$$S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)} \quad , \quad \eta = 0, 1, 2, 3$$

Below, we consider the 2-cycles in the triplet $\tilde{\Phi}^{(i)} \simeq \tilde{\mathcal{C}}_{\mathfrak{d}(2,1;\zeta)}^{(i)}$ while focussing on the system $\tilde{\Phi}^{(2)}$ due to the \mathcal{S}_3 symmetry. So, we have for the set $\tilde{\mathcal{C}}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ the following 2-cycle content

the set $\tilde{\mathcal{C}}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$:	<table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="padding: 5px;">odd 2-cycles</th> <th style="padding: 5px;">even 2-cycles</th> </tr> </thead> <tbody> <tr> <td style="text-align: center; padding: 5px;">$\tilde{\mathcal{C}}_{\tilde{\alpha}_2}$</td> <td style="text-align: center; padding: 5px;">$\tilde{\mathcal{C}}_{\tilde{\alpha}_1}$</td> </tr> <tr> <td style="padding: 5px;">$\tilde{\mathcal{C}}_{\tilde{\alpha}_2 + \tilde{\alpha}_1} \quad , \quad \tilde{\mathcal{C}}_{\tilde{\alpha}_2 + \tilde{\alpha}_3}$</td> <td style="text-align: center; padding: 5px;">$\tilde{\mathcal{C}}_{\tilde{\psi}}$</td> </tr> <tr> <td style="text-align: center; padding: 5px;">$\tilde{\mathcal{C}}_{\tilde{\alpha}_2 + \tilde{\alpha}_1 + \tilde{\alpha}_3}$</td> <td style="text-align: center; padding: 5px;">$\tilde{\mathcal{C}}_{\tilde{\alpha}_3}$</td> </tr> </tbody> </table>	odd 2-cycles	even 2-cycles	$\tilde{\mathcal{C}}_{\tilde{\alpha}_2}$	$\tilde{\mathcal{C}}_{\tilde{\alpha}_1}$	$\tilde{\mathcal{C}}_{\tilde{\alpha}_2 + \tilde{\alpha}_1} \quad , \quad \tilde{\mathcal{C}}_{\tilde{\alpha}_2 + \tilde{\alpha}_3}$	$\tilde{\mathcal{C}}_{\tilde{\psi}}$	$\tilde{\mathcal{C}}_{\tilde{\alpha}_2 + \tilde{\alpha}_1 + \tilde{\alpha}_3}$	$\tilde{\mathcal{C}}_{\tilde{\alpha}_3}$	(3.15)
odd 2-cycles	even 2-cycles									
$\tilde{\mathcal{C}}_{\tilde{\alpha}_2}$	$\tilde{\mathcal{C}}_{\tilde{\alpha}_1}$									
$\tilde{\mathcal{C}}_{\tilde{\alpha}_2 + \tilde{\alpha}_1} \quad , \quad \tilde{\mathcal{C}}_{\tilde{\alpha}_2 + \tilde{\alpha}_3}$	$\tilde{\mathcal{C}}_{\tilde{\psi}}$									
$\tilde{\mathcal{C}}_{\tilde{\alpha}_2 + \tilde{\alpha}_1 + \tilde{\alpha}_3}$	$\tilde{\mathcal{C}}_{\tilde{\alpha}_3}$									

where $\tilde{\psi}$ stands for the long root given by $\tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3$. This corresponds to the super Dynkin diagram $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ given by the **Figure 9**-(b). In this regard, notice the following: (i) the parametric $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ contains the distinguished super Dynkin diagram $S\mathcal{D}\mathcal{D}_{psl(2|2)}^{(1)}$ given by the **Figure 15**-(a) in appendix A;

$$S\mathcal{D}\mathcal{D}_{psl(2|2)}^{(1)} \subset S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$$

(ii) The parametric super 2-cycle system (3.15) contains the 2-cycles $\mathcal{C}_{psl(2|2)}^{(1)}$ illustrated by the **Figure 6** (green segments) namely

odd 2-cycles	even 2-cycles
\mathcal{C}_{α_2}	\mathcal{C}_{α_1}
$\mathcal{C}_{\alpha_1+\alpha_2} \quad , \quad \mathcal{C}_{\alpha_2+\alpha_3}$	—
$\mathcal{C}_{\alpha_1+\alpha_2+\alpha_3}$	\mathcal{C}_{α_3}

the set $\mathcal{C}_{psl(2|2)}^{(1)}$:

(3.16)

where the 2-cycle $\tilde{\mathcal{C}}_{\tilde{\psi}}$ in (3.15) associated with the even root $\tilde{\psi} = \tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3$ gets decoupled; i.e contracted to zero ($\mathcal{C}_{\tilde{\psi}} \simeq 0$). The other six 2-cycles $\tilde{\mathcal{C}}_{\tilde{\alpha}}$ in (3.15) get mapped to the six \mathcal{C}_{α} in the above (3.16).

3.2 Realising the distinguished $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ and $\tilde{\mathcal{C}}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ systems

To realise the positive roots of the system $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ of the super $\mathfrak{d}(2,1;\zeta)$ [resp. the associated $\tilde{\mathcal{C}}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$], we use the three following quantities: **(1)** the three orthogonal canonical weight vectors $\{\epsilon_1, \epsilon_2, \delta\}$ with $\epsilon_i \cdot \epsilon_j = \delta_{ij}$ and $\delta \cdot \delta = -1$; **(2)** the *Kaplansky* variables (s_1, s_2, s_3) , and **(3)** the roots (3.10) which for convenience we recall them here after

$$\begin{aligned}
 \text{odd :} \quad & \tilde{\alpha}_2, \quad \tilde{\alpha}_2 + \tilde{\alpha}_1 + \tilde{\alpha}_3 & ; \quad & \text{even :} \quad & \tilde{\alpha}_1, \quad \tilde{\alpha}_3, \\
 & \tilde{\alpha}_2 + \tilde{\alpha}_1, \quad \tilde{\alpha}_2 + \tilde{\alpha}_3 & & & \tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3
 \end{aligned}
 \tag{3.17}$$

We represent the three simple roots $(\tilde{\alpha}_1, \tilde{\alpha}_2, \tilde{\alpha}_3)$ generating $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ as follows

$$\begin{aligned}
 \tilde{\alpha}_1 &= \sqrt{2s_1} (\epsilon_1 - \epsilon_2) \\
 \tilde{\alpha}_2 &= \sqrt{s_1 + s_3} \delta - \sqrt{\frac{s_1}{2}} (\epsilon_1 - \epsilon_2) - \sqrt{\frac{s_3}{2}} (\epsilon_1 + \epsilon_2) \\
 \tilde{\alpha}_3 &= \sqrt{2s_3} (\epsilon_1 + \epsilon_2)
 \end{aligned}
 \tag{3.18}$$

showing that the roots (3.17) are indeed function of the parameters s_i . Notice that for $(s_1, s_2, s_3) = (1/2, -1, 1/2)$, the above relations reduce to

$$\begin{aligned}
 \tilde{\alpha}_1 &= \epsilon_1 - \epsilon_2 \equiv \alpha_1^{osp} \\
 \tilde{\alpha}_2 &= \delta - \epsilon_1 \equiv \alpha_2^{osp} \\
 \tilde{\alpha}_3 &= \epsilon_1 + \epsilon_2 \equiv \alpha_3^{osp}
 \end{aligned}
 \tag{3.19}$$

giving precisely the simple roots of the orthosymplectic Lie superalgebra $\mathfrak{osp}(4|2)$ further commented in the [Appendix B](#). These α_i^{osp} 's have well defined parities; α_1^{osp} and α_3^{osp} are even while α_2^{osp} is odd. From the parametric realisation (3.18), we also learn the following features:

(i) the realisation of the non simple roots (3.17) are given by

$$\begin{aligned}
\tilde{\alpha}_1 + \tilde{\alpha}_2 &= \sqrt{S_1 + S_3} \delta + \frac{1}{2}\sqrt{2S_1} (\epsilon_1 - \epsilon_2) - \frac{1}{2}\sqrt{2S_3} (\epsilon_1 + \epsilon_2) \\
\tilde{\alpha}_2 + \tilde{\alpha}_3 &= \sqrt{S_1 + S_3} \delta - \frac{1}{2}\sqrt{2S_1} (\epsilon_1 - \epsilon_2) + \frac{1}{2}\sqrt{2S_3} (\epsilon_1 + \epsilon_2) \\
\tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3 &= \sqrt{S_1 + S_3} \delta + \frac{1}{2}\sqrt{2S_1} (\epsilon_1 - \epsilon_2) + \frac{1}{2}\sqrt{2S_3} (\epsilon_1 + \epsilon_2)
\end{aligned} \tag{3.20}$$

together with the even long root $\tilde{\psi} = \tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3$ pointing in the δ -direction as follows³:

$$\tilde{\psi} = 2\sqrt{S_1 + S_3} \delta \tag{3.21}$$

For $S_1 + S_3 = 1$, this long root reduce to $\psi^{osp} = 2\delta$; this is the case of the orthosymplectic point $(S_1, S_2, S_3) = (1/2, -1, 1/2)$. For the case $S_1 + S_3 = 0$ as for the situation $(S_1, S_2, S_3) = (1/2, 0, -1/2)$, the long root vanishes identically; this corresponds to $\psi^{sl(2|2)} = 0$.

(ii) Using the property $S_1 + S_2 + S_3 = 0$, the intersection matrix of the simple roots like $\tilde{\mathcal{J}}_{ij}(s) = \tilde{\alpha}_i \cdot \tilde{\alpha}_j$ and the corresponding super Cartan matrix $\tilde{\mathcal{K}}_{ij}(s)$ read in the patch $S_1 \neq 0$ as follows

$$\tilde{\mathcal{J}}_{ij}(s) = \begin{pmatrix} 4S_1 & -2S_1 & 0 \\ -2S_1 & 0 & -2S_3 \\ 0 & -2S_3 & 4S_3 \end{pmatrix}, \quad \tilde{\mathcal{K}}_{ij}(s) = \begin{pmatrix} 2 & -1 & 0 \\ 1 & 0 & S_3/S_1 \\ 0 & -1 & 2 \end{pmatrix} \tag{3.22}$$

The determinant $\det \tilde{\mathcal{J}}_{ij}(s) = 16S_1S_2S_3$; it is singular for $\prod S_i = 0$; it the case of the Lie superalgebra $\mathfrak{psl}(2|2)$ sitting at $(S_1, S_2, S_3) = (1/2, 0, -1/2)$; i.e: $\det \mathcal{J}_{ij}^{(sl(2|2))} = 0$. For the Lie superalgebra $\mathfrak{osp}(4|2)$ sitting at $(S_1, S_2, S_3) = (1/2, -1, 1/2)$, we have $\det \mathcal{J}_{ij}^{(osp)} = -4$. The determinant of the super Cartan

$$\tilde{\mathcal{K}}_{ij}(s) = -\frac{2S_2}{S_1} \delta_{Dirac}(S_1 + S_2 + S_3) \tag{3.23}$$

(iii) In terms of the even roots $(\tilde{\alpha}_1, \tilde{\psi}, \tilde{\alpha}_3)$, the fermionic roots read as follows

$$\begin{aligned}
\tilde{\alpha}_2 &= +\frac{1}{2}\tilde{\psi} - \frac{1}{2}\tilde{\alpha}_1 - \frac{1}{2}\tilde{\alpha}_3 \\
\tilde{\alpha}_1 + \tilde{\alpha}_2 &= +\frac{1}{2}\tilde{\psi} + \frac{1}{2}\tilde{\alpha}_1 - \frac{1}{2}\tilde{\alpha}_3 \\
\tilde{\alpha}_2 + \tilde{\alpha}_3 &= +\frac{1}{2}\tilde{\psi} - \frac{1}{2}\tilde{\alpha}_1 + \frac{1}{2}\tilde{\alpha}_3 \\
\tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3 &= +\frac{1}{2}\tilde{\psi} + \frac{1}{2}\tilde{\alpha}_1 + \frac{1}{2}\tilde{\alpha}_3
\end{aligned} \tag{3.24}$$

By using the correspondence $\tilde{\alpha} \leftrightarrow \tilde{\mathcal{C}}_{\tilde{\alpha}}$ of eq(3.1), we get the system of 2-cycles (3.15) in the supermanifold describing $D(2|1; \zeta)$.

³ It would be interesting to explore the situation where the Kaplansky parameters (S_1, S_2, S_3) are solved like $S_1 + S_3 = n^2$ with integer n. For this case we have $S_2 = -n^2$ and the long root is even and is given by $\tilde{\psi} = 2n \delta$.

3.2.1 Roots $\tilde{\alpha}$ as function of ζ

Here, we solve the conditions $s_1+s_2+s_3 = 0$ and $s_1s_2s_3 \neq 0$ on the *Kaplansky* variables in terms of the complex parameter ζ labeling $D(2|1;\zeta)$ like

$$2s_1 = 1 \quad , \quad 2s_2 = -1 - \zeta \quad , \quad 2s_3 = \zeta \quad (3.25)$$

This solution breaks the permutation symmetry \mathcal{S}_3 down to its subgroup \mathcal{Z}_2 . This discrete \mathcal{Z}_2 fixes the axis s_1 and permutes s_2 and s_3 ; it has a fix point at $\zeta = 1/2$.

Putting (3.25) into (3.18), we get

$$\begin{aligned} \tilde{\alpha}_1 &= (\epsilon_1 - \epsilon_2) \\ \tilde{\alpha}_2 &= \frac{1}{2}\sqrt{2(1+\zeta)} \delta - \frac{1}{2}(\epsilon_1 - \epsilon_2) - \frac{1}{2}\sqrt{\zeta}(\epsilon_1 + \epsilon_2) \\ \tilde{\alpha}_3 &= \sqrt{\zeta}(\epsilon_1 + \epsilon_2) \end{aligned} \quad (3.26)$$

and

$$\begin{aligned} \tilde{\alpha}_1 + \tilde{\alpha}_2 &= \sqrt{\frac{1+\zeta}{2}} \delta + \frac{1}{2}(\epsilon_1 - \epsilon_2) - \frac{1}{2}\sqrt{\zeta}(\epsilon_1 + \epsilon_2) \\ \tilde{\alpha}_2 + \tilde{\alpha}_3 &= \sqrt{\frac{1+\zeta}{2}} \delta - \frac{1}{2}(\epsilon_1 - \epsilon_2) + \frac{1}{2}\sqrt{\zeta}(\epsilon_1 + \epsilon_2) \\ \tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3 &= \sqrt{\frac{1+\zeta}{2}} \delta + \frac{1}{2}(\epsilon_1 - \epsilon_2) + \frac{1}{2}\sqrt{\zeta}(\epsilon_1 + \epsilon_2) \end{aligned} \quad (3.27)$$

as well as the long root

$$\tilde{\psi} = \sqrt{2(1+\zeta)} \delta \quad , \quad \tilde{\psi}^2 = -2(1+\zeta) \quad (3.28)$$

The intersection matrix $\tilde{\mathcal{J}}_{ij} = \tilde{\alpha}_i \cdot \tilde{\alpha}_j$ and the corresponding super Cartan matrix $\tilde{\mathcal{K}}_{ij}(\zeta)$ read in terms of ζ as follows

$$\tilde{\mathcal{J}}_{ij}(\zeta) = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & -\zeta \\ 0 & -\zeta & 2\zeta \end{pmatrix} \quad , \quad \tilde{\mathcal{K}}_{ij}(\zeta) = \begin{pmatrix} 2 & -1 & 0 \\ 1 & 0 & \zeta \\ 0 & -1 & 2 \end{pmatrix} \quad (3.29)$$

The determinant $\det \tilde{\mathcal{J}}_{ij}(\zeta) = -2\zeta(\zeta+1)$ vanishes at $\zeta = -1, 0$ while the $\det \tilde{\mathcal{K}}_{ij}(\zeta) = 2(\zeta+1)$ vanishes at $\zeta = -1$. Below, we give the Lie superalgebras at the two special points $\zeta = \pm 1$:

A) Orthosymplectic $\mathfrak{osp}(4|2)$: $\zeta = 1 \bmod \mathcal{Z}_2$: This is a special point sitting on the real line of the plan $\mathbb{C} \setminus \{-1, 0\}$. The simple roots are given by

$$\begin{aligned} \tilde{\alpha}_1 &= \epsilon_1 - \epsilon_2 = \alpha_1^{osp} \\ \tilde{\alpha}_2 &= \delta - \epsilon_1 = \alpha_2^{osp} \\ \tilde{\alpha}_3 &= \epsilon_1 + \epsilon_2 = \alpha_3^{osp} \end{aligned} \quad (3.30)$$

with long root $\psi^{osp} = 2\delta$. The intersection matrix $\tilde{\mathcal{J}}(\zeta)|_{\zeta=1} = \mathcal{J}^{osp}$ and the associated super Cartan matrix \mathcal{K}^{osp} as follows

$$\mathcal{J}^{osp} = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & -1 \\ 0 & -1 & 2 \end{pmatrix}, \quad \mathcal{K}^{osp} = \begin{pmatrix} 2 & -1 & 0 \\ 1 & 0 & 1 \\ 0 & -1 & 2 \end{pmatrix} \quad (3.31)$$

The 4 other composite positive roots are given by

$$\begin{aligned} \alpha_1^{osp} + \alpha_2^{osp} &= \delta - \epsilon_2 & \alpha_1^{osp} + \alpha_2^{osp} + \alpha_3^{osp} &= \delta + \epsilon_1 \\ \alpha_2^{osp} + \alpha_3^{osp} &= \delta + \epsilon_2 & \alpha_1^{osp} + 2\alpha_2^{osp} + \alpha_3^{osp} &= 2\delta \end{aligned} \quad (3.32)$$

B) Roots at the point $\zeta = -1 \bmod \mathcal{Z}_2$: This special point corresponds to the Lie superalgebra $\mathfrak{sl}(2|2)$ with simple roots $\tilde{\alpha}_i|_{\zeta=-1} = \alpha_i^{sl_2|2}$ as follows

$$\begin{aligned} \tilde{\alpha}_1^{sl_2|2} &= (\epsilon_1 - \epsilon_2) \\ \tilde{\alpha}_2^{sl_2|2} &= \delta - \frac{1}{2}(\epsilon_1 - \epsilon_2) - \frac{i}{2}(\epsilon_1 + \epsilon_2) \\ \tilde{\alpha}_3^{sl_2|2} &= i(\epsilon_1 + \epsilon_2) \end{aligned} \quad (3.33)$$

where we used $\sqrt{-1} = i$. These roots are related to the roots of $\mathfrak{osp}(4|2)$ as follows

$$\begin{aligned} \alpha_1^{sl_2|2} &= \alpha_1^{osp} \\ 2\tilde{\alpha}_2^{sl_2|2} &= 2\delta - \alpha_1^{osp} - i\tilde{\alpha}_3^{sl_2|2} \\ \alpha_3^{sl_2|2} &= i\alpha_3^{osp} \end{aligned} \quad (3.34)$$

indicating that $\tilde{\alpha}_1^{sl_2|2} + 2\tilde{\alpha}_2^{sl_2|2} + \tilde{\alpha}_3^{sl_2|2} = 0 \bmod (2\delta)$. Here, the long root $\tilde{\psi}|_{\zeta=-1}$ vanishes [$\tilde{\psi}|_{\zeta=-1} = 0 \bmod (2\delta)$]. In these relations, the intersection matrix $\mathcal{J}^{sl(2|2)}$ of the simple roots and the corresponding super Cartan matrix $\mathcal{K}^{sl(2|2)}$ read as follows

$$\mathcal{J}^{sl(2|2)} = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & 1 \\ 0 & 1 & -2 \end{pmatrix}, \quad \mathcal{K}^{sl(2|2)} = \begin{pmatrix} 2 & -1 & 0 \\ 1 & 0 & -1 \\ 0 & -1 & 2 \end{pmatrix} \quad (3.35)$$

3.2.2 Weight diagram of $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ from the $\Phi_{\mathfrak{osp}(4|2)}$ graph

Because the Lie superalgebra $\mathfrak{d}(2,1;\zeta)$ has four parametric root systems $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ in 1:1 with the four possible $\mathcal{SD}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ and subsequently in 1:1 with the four parametric 2-cycles $\mathcal{C}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$, i.e:

$$\begin{aligned} \text{root systems} &: \text{super Dynkin diagram} &: \text{2-cycles} \\ \tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)} &\leftrightarrow \mathcal{SD}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)} &\leftrightarrow \mathcal{C}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)} \end{aligned} \quad (3.36)$$

one distinguishes 4 types of Root diagrams that we denote like

$$\mathfrak{RD}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$$

Using the root system (3.17), we can draw the diagram $\mathfrak{RD}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ of the super $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$. This is a parametric graph labeled by ζ

$$\mathfrak{RD}_{\mathfrak{d}(2,1;\zeta)}^{(2)} := \widetilde{\mathfrak{RD}}^{(2)}[\zeta] \quad , \quad \eta = 0, 1, 2, 3$$

As such, it can be thought of as a fibration over the complex line ζ of the root diagram $\mathfrak{RD}_{osp(4|2)}$ of the orthosymplectic root system $\Phi_{osp(4|2)}$ given by eqs(3.30,3.32). This fibration is depicted by the **Figure 10** where the $\mathfrak{RD}_{sl(2|2)}$ at $\zeta = -1$ is also shown. The




Figure 10: Graphic representation of $\widetilde{\mathfrak{RD}}^{(2)}[\zeta]$ as a fibration over the complex line ζ of the graph $\mathfrak{RD}_{osp(4|2)}$ sitting at $\zeta = 1$. At the point $\zeta = -1$, it contracts to the root diagram $\mathfrak{RD}_{sl(2|2)}$.

properties of the parametric $\widetilde{\mathfrak{RD}}^{(2)}[\zeta]$ descend from those of $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ among which we cite the following:

- (1) **values at $\zeta = \pm 1$:** The simple roots of $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ are $\{\tilde{\alpha}_1(\zeta), \tilde{\alpha}_2(\zeta), \tilde{\alpha}_3(\zeta)\}$ with odd $\tilde{\alpha}_2(\zeta)$ and even $\tilde{\alpha}_1(\zeta), \tilde{\alpha}_3(\zeta)$; they are functions of ζ as in (3.26). Then, the $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ is a parametric root system with values at $\zeta = \pm 1$ given by

$$\tilde{\Phi}_{\mathfrak{d}(2,1;+1)}^{(2)} = \Phi_{osp(4|2)} \quad , \quad \tilde{\Phi}_{\mathfrak{d}(2,1;-1)}^{(2)} = \Phi_{psl(2|2)} \quad (3.37)$$

From these features, we deduce:

$$\widetilde{\mathfrak{RD}}^{(2)}|_{\zeta=1} = \mathfrak{RD}_{osp(4|2)} \quad , \quad \widetilde{\mathfrak{RD}}^{(2)}|_{\zeta=-1} = \mathfrak{RD}_{psl(2|2)} \quad (3.38)$$

The Cartan-Weyl operators associated with the roots $\tilde{\alpha}$ include:

- (i) the three parametric Chevalley triplets $(\tilde{H}_{\tilde{\alpha}_i}, \tilde{E}_{\pm\tilde{\alpha}_i})$ with Cartan charges $\tilde{H}_{\tilde{\alpha}_i}$ given by

$$\tilde{E}_{+\tilde{\alpha}_i} \tilde{E}_{-\tilde{\alpha}_i} - (-)^{|\tilde{\alpha}_i||\tilde{\alpha}_i|} \tilde{E}_{-\tilde{\alpha}_i} \tilde{E}_{+\tilde{\alpha}_i}$$

with $|\tilde{\alpha}|$ standing for the degree of $\tilde{\alpha}$; and

(ii) the composite operators associated with non simple roots

$$\tilde{E}_{\pm(\tilde{\alpha}_1+\tilde{\alpha}_2)} \quad , \quad \tilde{E}_{\pm(\tilde{\alpha}_3+\tilde{\alpha}_2)} \quad , \quad \tilde{E}_{\pm(\tilde{\alpha}_1+\tilde{\alpha}_3+\tilde{\alpha}_2)} \quad , \quad \tilde{E}_{\pm(\tilde{\alpha}_1+\tilde{\alpha}_3+2\tilde{\alpha}_2)} \quad (3.39)$$

They are given by the graded Serre relations [52]; and are functions of ζ .

(2) **the diagram $\widetilde{\mathfrak{RD}}^{(2)}$** : The long root $\tilde{\psi} = \tilde{\psi}(\zeta)$ of the system $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2+)}$ and the long ψ^{osp} of the $\Phi_{osp(4|2)}$ are given by the even roots

$$\tilde{\psi} = \tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3 \quad , \quad \psi^{osp} = \alpha_1^{osp} + 2\alpha_2^{osp} + \alpha_3^{osp}$$

The ψ^{osp} and $\tilde{\psi}$ are also the highest weight vectors of the "adjoint" representation of $osp(4|2)$ and $\mathfrak{d}(2,1;\zeta)$ respectively. The root diagram $\mathfrak{RD}_{osp(4|2)}$ and subsequently the diagram $\widetilde{\mathfrak{RD}}^{(2)}$ are depicted by the **Figure 11**. The $\widetilde{\mathfrak{RD}}^{(2)}$ is imagined as the fibration of the root diagram of $osp(4|2)$ over the complex line $\mathbb{C} \setminus \{-1, 0\}$ with variable ζ . The parametric $\widetilde{\mathfrak{RD}}^{(2)}[\zeta]$ is built as follows:




Figure 11: The root diagram of $\widetilde{\mathfrak{RD}}^{(2)}|_{\zeta=1} = \mathfrak{RD}_{osp(4|2)}$. For generic ζ , we get the parametric root diagram $\widetilde{\mathfrak{RD}}^{(2)}$. In red the 3+3 even roots of $\mathfrak{d}(2,1;\zeta)|_{\zeta=+1}$ and in blue the 4+4 odd ones. From this diagram, we learn the weight vectors of the adjoint representation of $\mathfrak{d}(2,1;\zeta)|_{\zeta=+1}$ and its highest/lowest weight vectors.

We start from the highest weight state $|\tilde{\psi}\rangle$ which is: (i) an eigenvalue of the diagonal Cartan charge operators $\tilde{H}_{\tilde{\alpha}_i}|\tilde{\psi}\rangle = \tilde{q}_i|\tilde{\psi}\rangle$; and (ii) annihilated by $\tilde{E}_{-\tilde{\alpha}_i}$; that is $\tilde{E}_{-\tilde{\alpha}_i}|\tilde{\psi}\rangle = 0$.

Then, we act successively by appropriate monomials of the step operators $\tilde{E}_{\pm\tilde{\alpha}}$; thus generating the states of the adjoint representation. For example, we have

$$|\tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3\rangle \simeq E_{\tilde{\alpha}_2}|\tilde{\psi}\rangle \quad (3.40)$$

and

$$|\tilde{\alpha}_2\rangle \simeq E_{\tilde{\alpha}_1+\tilde{\alpha}_3}E_{\tilde{\alpha}_2}|\tilde{\psi}\rangle \quad (3.41)$$

as well as

$$|\tilde{\alpha}_1\rangle \simeq E_{\tilde{\alpha}_2+\tilde{\alpha}_3}E_{+\tilde{\alpha}_2}|\tilde{\psi}\rangle, \quad |\tilde{\alpha}_3\rangle \simeq E_{\tilde{\alpha}_1+\tilde{\alpha}_2}E_{\tilde{\alpha}_2}|\tilde{\psi}\rangle \quad (3.42)$$

4 Geometry of $D(2, 1; \zeta)$ as deformation of $OSp(4|2)$

In this section, we study geometric properties of the 9|8-dim supermanifold describing the exceptional Lie super group $D(2, 1; \zeta)$ by help of the following features:

- (1) the isometry $D(2, 1; 1) \simeq OSp(4|2)$ at the point $\zeta = 1$ of the complex line $\mathbb{C} \setminus \{-1, 0\}$.
The $D(2, 1; \zeta)$ has three parametric even roots (2-cycles) that we denote like $\tilde{\beta}_1(\zeta)$, $\tilde{\beta}_2(\zeta)$ and $\tilde{\beta}_3(\zeta)$. Their values at the special point $\zeta = 1$ give the roots (2-cycles) β_i^{osp} of $OSp(4|2)$. So, we have $\tilde{\beta}_i|_{\zeta=1} = \beta_i^{osp}$.
- (2) continuous deformation of the Kahler parameter $\tilde{\varrho}_3(\zeta)$ of the even 2-cycle $\tilde{\beta}_3(\zeta)$ of the 17D geometry of $D(2, 1; \zeta)$. We start from the special point

$$\beta_3^{osp} = \tilde{\beta}_3(\zeta)|_{\zeta=1} \quad (4.1)$$

of $OSp(4|2)$ and we move towards the critical point

$$\beta_3^{sl(2|2)} = \tilde{\beta}_3(\zeta)|_{\zeta=-1} \quad (4.2)$$

of $PSL(2|2) \times \mathbb{C}^3$. For this continuous deformation, the Kahler parameter $\tilde{\varrho}_3(\zeta)$ of the 2-cycle $\tilde{\beta}_3$ in $SL(2, \mathbb{C})_3$ is taken like $2/|\zeta(\zeta+1)|$; its value at $\zeta = 1$ is equal to $\varrho_3^{osp} = 1$ while at $\zeta = -1$, it is given by the diverging $\varrho_3^{sl(2|2)} \rightarrow \infty$.

To undertake this study, we first describe the geometry of the bosonic part given by

$$\begin{aligned} D(2, 1; \zeta)_{\bar{0}} &\simeq SL(2, \mathbb{C})^{\otimes 3} & \text{for } \zeta \in \mathbb{C} \setminus \{-1, 0\} \\ D(2, 1; \zeta)_{\bar{0}} &\simeq SL(2, \mathbb{C})^{\otimes 2} \otimes \mathbb{C}^3 & \text{for } \zeta = \{-1, 0\} \end{aligned} \quad (4.3)$$

Then, we consider the super extension by implementing the odd part $D(2, 1; \zeta)_{\bar{1}}$.

4.1 Bosonic subspace $D(2, 1; \zeta)_0$

The complex geometry of the bosonic $D(2, 1; \zeta)_0$ is given by a parametric 9 dimensional manifold $SL(2)^{\otimes 3}$ with parametric roots $\tilde{\beta}_i$ that are functions of ζ [70]. It is denoted like $\tilde{\mathcal{X}}_{123}[\zeta]$ in analogy with the \mathcal{X}_{120} geometry of $SL(2|2)_0 \simeq SL(2)^{\otimes 2} \times \mathbb{C}^*$ sitting at $\zeta = -1$.

By using (2.48), the $\tilde{\mathcal{X}}_{123}$ factorises like $\tilde{\mathcal{X}}_1 \times \tilde{\mathcal{X}}_2 \times \tilde{\mathcal{X}}_3$ where $\tilde{\mathcal{X}}_i \simeq \tilde{\mathcal{C}}_{\tilde{\beta}_i} \times \tilde{\mathfrak{F}}_i$ with 2-cycles $\tilde{\mathcal{C}}_{\tilde{\beta}_i}$ (1D complex curves) and complex surfaces $\tilde{\mathfrak{F}}_i$. The three 2-cycles $\tilde{\mathcal{C}}_{\tilde{\beta}_i}$ are associated with the even roots of $D(2, 1; \zeta)$; they are given by

$$\begin{aligned} \tilde{\beta}_1 &= \tilde{\alpha}_1 & \leftrightarrow & \tilde{\mathcal{C}}_{\tilde{\beta}_1} \\ \tilde{\beta}_2 &= \tilde{\alpha}_3 & \leftrightarrow & \tilde{\mathcal{C}}_{\tilde{\beta}_2} \\ \tilde{\beta}_3 &= \tilde{\psi} & \leftrightarrow & \tilde{\mathcal{C}}_{\tilde{\beta}_3} \end{aligned} \quad (4.4)$$

with $\tilde{\psi} = \tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3$ and vanishing intersections

$$\tilde{\beta}_1 \cdot \tilde{\beta}_2 = 0 \quad , \quad \tilde{\beta}_1 \cdot \tilde{\beta}_3 = 0 \quad , \quad \tilde{\beta}_2 \cdot \tilde{\beta}_3 = 0 \quad (4.5)$$

By substituting the factorisation $\tilde{\mathcal{C}}_{\tilde{\beta}_i} \times \tilde{\mathfrak{F}}_i$ into $\tilde{\mathcal{X}}_1 \times \tilde{\mathcal{X}}_2 \times \tilde{\mathcal{X}}_3$, we can express the complex $\tilde{\mathcal{X}}_{123}$ as the fibration $\tilde{\mathfrak{B}}_{123} \times \tilde{\mathfrak{F}}_{123}$ with 6-cycle

$$\tilde{\mathfrak{B}}_{123} = \tilde{\mathcal{C}}_{\tilde{\beta}_1} \times \tilde{\mathcal{C}}_{\tilde{\beta}_2} \times \tilde{\mathcal{C}}_{\tilde{\beta}_3} \quad (4.6)$$

Because each isospin group of $\otimes_{i=1}^3 SL(2)_{\tilde{\beta}_i}$ can be realised by complex 2×2 matrices \mathbf{X}_i , the geometric description of the super $D(2, 1; \zeta)_0$ can be imagined in terms of parametric 6×6 matrix $\mathbf{D} = \mathbf{D}(\zeta)$ as follows

$$\mathbf{D} = \begin{pmatrix} \mathbf{X}_1 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{X}_2 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{X}_3 \end{pmatrix} \quad \text{with} \quad \mathbf{X}_i = \begin{pmatrix} x_i & u_i \\ y_i & v_i \end{pmatrix} \quad (4.7)$$

Moreover, as each $\mathbf{X}_i(\zeta)$ describes an $SL(2)_{\tilde{\beta}_i}$, the complex variables $\{(x_i, y_i, u_i, v_i)_{i=1,2,3}\}$ parameterising $(\text{End}\mathbb{C}^2)^3$ are constrained by the conditions $\det \mathbf{X}_i = 1$ reading explicitly as

$$\begin{aligned} \mathbf{X}_1 &: x_1 v_1 - y_1 u_1 = 1 \\ \mathbf{X}_2 &: x_2 v_2 - y_2 u_2 = 1 \\ \mathbf{X}_3 &: x_3 v_3 - y_3 u_3 = 1 \end{aligned} \quad (4.8)$$

Using (4.7) and setting $x_i = a_i + b_i$ and $v_i = \bar{a}_i - \bar{b}_i$ as well as $y_i = c_i + d_i$ and $u_i = \bar{c}_i - \bar{d}_i$, we can expand these relationships (4.8) as follows

$$\begin{aligned} \tilde{\mathfrak{B}}_1 &: |a_1|^2 + |d_1|^2 - |b_1|^2 - |c_1|^2 = 1 \\ \tilde{\mathfrak{F}}_1 &: b_1 \bar{a}_1 - a_1 \bar{b}_1 + d_1 \bar{c}_1 - c_1 \bar{d}_1 = 0 \end{aligned} \quad (4.9)$$

and

$$\begin{aligned}\tilde{\mathfrak{B}}_2 & : |a_2|^2 + |d_2|^2 - |b_2|^2 - |c_2|^2 = 1 \\ \tilde{\mathfrak{F}}_2 & : b_2\bar{a}_2 - a_2\bar{b}_2 + d_2\bar{c}_2 - c_2\bar{d}_2 = 0\end{aligned}\tag{4.10}$$

as well as

$$\begin{aligned}\tilde{\mathfrak{B}}_3 & : |a_3|^2 + |d_3|^2 - |b_3|^2 - |c_3|^2 = 1 \\ \tilde{\mathfrak{F}}_3 & : b_3\bar{a}_3 - a_3\bar{b}_3 + d_3\bar{c}_3 - c_3\bar{d}_3 = 0\end{aligned}\tag{4.11}$$

On the locus $b_i = c_i = 0$, these relations become

$$\begin{aligned}\mathbb{S}_1^2 & : |a_1|^2 + |d_1|^2 = 1 \\ \mathbb{S}_2^2 & : |a_2|^2 + |d_2|^2 = 1 \\ \mathbb{S}_3^2 & : |a_3|^2 + |d_3|^2 = 1\end{aligned}\tag{4.12}$$

which by using the parametrisation $v_3 = \varrho_3\bar{x}_3$ and $u_3 = -\varrho_3\bar{y}_3$ [see also the discussion between eqs(2.53) and (2.59)], they take the form

$$\begin{aligned}\mathbb{S}_1^2 & : |a_1|^2 + |d_1|^2 = 1 \\ \mathbb{S}_2^2 & : |a_2|^2 + |d_2|^2 = 1 \\ \mathbb{S}_3^2 & : |x_3|^2 + |y_3|^2 = \frac{1}{\varrho_3}\end{aligned}\tag{4.13}$$

with positive $\varrho_3 = |\zeta(\zeta + 1)|/2$ and area

$$vol(\mathbb{S}_{\tilde{\beta}_3}^2) = \frac{8\pi}{|\zeta(\zeta + 1)|}\tag{4.14}$$

In this geometric description, the contraction $\zeta \rightarrow -1$ reduces the 6+3 dimensional $D(2, 1; \zeta)_{\bar{0}}$ down to the 6 dimensional $PSL(2|2)_{\bar{0}}$. It corresponds to the decompactification of the 2-cycle $\tilde{\mathcal{C}}_{\tilde{\beta}_3}|_{\zeta=-1}$ into $\mathbb{R}^2 \simeq \mathbb{C}$ for 2-sphere or $\mathbb{R}^{1,1}$ for non compact pseudo 2-sphere.

4.2 Odd part $D(2, 1; \zeta)_{\bar{1}}$

The contribution of the odd part in the geometric description of the exceptional supergroup $D(2, 1; \zeta)$ is given by the tri-fundamental $(1/2, 1/2, 1/2)$ of the even part $D(2, 1; \zeta)_{\bar{0}} = \Pi_{i=1}^3 SL(2)_{\tilde{\beta}_i}$. Because of the Z_3 -graded matrix description for $D(2, 1; \zeta)$ due to the tri-fundamental, we cannot apply the method used for $SL(2|2)$. However, we can get around this difficulty by using (i) the *isomorphism* $D(2, 1; 1)_{\bar{0}} \simeq OSp(4|2)_{\bar{0}}$ with

$$\begin{aligned}osp(4|2)_{\bar{0}} & \simeq so_{\mathbb{C}}(4) \oplus sp_{\mathbb{C}}(2) \\ so_{\mathbb{C}}(4) & \simeq sl(2, \mathbb{C}) \oplus sl(2, \mathbb{C})\end{aligned}\tag{4.15}$$

and (ii) the odd sector given by the module $(\mathbf{4}, \mathbf{2})$ of the group $SO_{\mathbb{C}}(4) \times SP_{\mathbb{C}}(2)$. In this orthosymplectic setting, the parametric geometry of $D(2, 1; 1)$ can be investigated by using 6×6 matrices of $OSp(4|2)$. In this regard, recall that the super $OSp(4|2)$ is a

subgroup of the linear super group $GL(4|2)$ given by super $4|2 \times 4|2$ matrices \mathbf{M} and super transpose \mathbf{M}^{st} ,

$$\mathbf{M} = \begin{pmatrix} \mathbf{A}_{4 \times 4} & \mathbf{B}_{4 \times 2} \\ \mathbf{C}_{2 \times 4} & \mathbf{D}_{2 \times 2} \end{pmatrix}, \quad \mathbf{M}^{st} = \begin{pmatrix} \mathbf{A}_{4 \times 4}^t & \mathbf{C}_{4 \times 2}^t \\ -\mathbf{B}_{2 \times 4}^t & \mathbf{D}_{2 \times 2}^t \end{pmatrix} \quad (4.16)$$

constrained as follows

$$\mathbf{M}^{st} \mathbf{J}_{4|2} \mathbf{M} = \mathbf{J}_{4|2} \quad (4.17)$$

with non vanishing super determinant $Ber(\mathbf{M})$ factorising like $\det(\mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C}) \det \mathbf{D}^{-1}$.

In this relation, the super orthosymplectic metric $\mathbf{J}_{4|2}$ is given by

$$\mathbf{J}_{4|2} = \begin{pmatrix} \mathbf{0}_{2 \times 2} & \mathbf{I}_{2 \times 2} & 0 & 0 \\ \mathbf{I}_{2 \times 2} & \mathbf{0}_{2 \times 2} & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix} \quad (4.18)$$

with $\mathbf{I}_{2 \times 2}$ being the identity 2×2 matrix. The graded \mathbf{M} and super transpose \mathbf{M}^{st} have even and odd submatrix blocks; the even block is given by $\mathbf{M}_{\bar{0}} = \mathbf{A}_{4 \times 4} \oplus \mathbf{D}_{2 \times 2}$; its super determinant

$$\det \mathbf{M}_{\bar{0}} = \det \mathbf{A}_{4 \times 4} \det \mathbf{D}_{2 \times 2}^{-1} \quad (4.19)$$

is invariant under the complex scaling

$$\mathbf{A}_{4 \times 4} \rightarrow \lambda \mathbf{A}_{4 \times 4}, \quad \mathbf{D}_{2 \times 2} \rightarrow \lambda^2 \mathbf{D}_{2 \times 2} \quad (4.20)$$

This is because $\det \mathbf{A}_{4 \times 4} \rightarrow \lambda^4 \det \mathbf{A}_{4 \times 4}$ and $\det \mathbf{D}_{2 \times 2} \rightarrow \lambda^4 \det \mathbf{D}_{2 \times 2}$. Extending this scaling to the odd sector $\mathbf{M}_{\bar{1}} = \mathbf{B}_{4 \times 2} \oplus \mathbf{C}_{2 \times 4}$, the invariance of the Berezinian $Ber(\mathbf{M})$ requires, in addition to the transformation of the even sector, the scalings

$$\mathbf{B}_{4 \times 2} \rightarrow \lambda^2 \mathbf{B}_{4 \times 2}, \quad \mathbf{C}_{2 \times 4} \rightarrow \lambda \mathbf{C}_{2 \times 4} \quad (4.21)$$

Furthermore, by putting (4.18) into the condensed form

$$\mathbf{J}_{4|2} = \begin{pmatrix} G_{4 \times 4} & 0 \\ 0 & \Omega_{2 \times 2} \end{pmatrix}, \quad G_{4 \times 4} = \begin{pmatrix} \mathbf{0}_{2 \times 2} & \mathbf{I}_{2 \times 2} \\ \mathbf{I}_{2 \times 2} & \mathbf{0}_{2 \times 2} \end{pmatrix}, \quad \Omega_{2 \times 2} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (4.22)$$

we can split the condition $\mathbf{M}^{st} \mathbf{J}_{4|2} \mathbf{M} = \mathbf{J}_{4|2}$ as follows

$$\begin{aligned} \mathbf{A}_{4 \times 4}^t G_{4 \times 4} \mathbf{A}_{4 \times 4} + \mathbf{C}_{4 \times 2}^t \Omega_{2 \times 2} \mathbf{C}_{2 \times 4} &= G_{4 \times 4} \\ \mathbf{D}_{2 \times 2}^t \Omega_{2 \times 2} \mathbf{D}_{2 \times 2} - \mathbf{B}_{2 \times 4}^t G_{4 \times 4} \mathbf{B}_{4 \times 2} &= \Omega_{2 \times 2} \\ \mathbf{A}_{4 \times 4}^t G_{4 \times 4} \mathbf{B}_{4 \times 2} + \mathbf{C}_{4 \times 2}^t \Omega_{2 \times 2} \mathbf{D}_{2 \times 2} &= 0 \\ \mathbf{D}_{2 \times 2}^t \Omega_{2 \times 2} \mathbf{C}_{2 \times 4} - \mathbf{B}_{2 \times 4}^t G_{4 \times 4} \mathbf{A}_{4 \times 4} &= 0 \end{aligned} \quad (4.23)$$

Notice that from (4.23), we learn that (i) the third and the fourth relations are related under transpose involution; and (ii) the odd blocks $\mathbf{B}_{4 \times 2}$ and $\mathbf{C}_{4 \times 2}$ are linked like

$$\mathbf{C}_{4 \times 2}^t = \mathbf{A}_{4 \times 4}^t G_{4 \times 4} \mathbf{B}_{4 \times 2} \mathbf{D}_{2 \times 2}^{-1} \Omega_{2 \times 2} \quad (4.24)$$

leaving free the block matrices $\mathbf{A}_{4 \times 4}$ and $\mathbf{B}_{4 \times 2}$ as well as $\mathbf{D}_{2 \times 2}$. The remaining relations give $10 + 1 + 8 = 19$ constraint relations amongst the $6 \times 6 = 36$ variables in eq(4.16); thus leaving 17 free degrees of freedom parameterising the complex variety of super $OSp(4|2)$.

In summary, the even sector has 9 degrees of freedom; 6 degrees coming from 4×4 (antisymmetric) matrix $\mathbf{A}_{4 \times 4}$ and 3 ones from the 2×2 (symmetric) matrix $\mathbf{D}_{2 \times 2}$. The odd sector has 8 degrees of freedom coming from the 4×2 matrix \mathbf{B} or equivalently from the 2×4 matrix $\mathbf{C}_{4 \times 2} \sim (\mathbf{B}_{2 \times 4})^t$ as in (4.24).

The link of the orthosymplectic description $OSp(4|2)$ to the 15 dimensional $SL(2|2)$ is obtained by imposing two conditions: First, by fixing the super determinant like

$$Ber(\mathbf{M}) = 1$$

this reduces the dimension from 17 down to 16. Second, by taking the coset by the scaling transformation

$$\begin{pmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{pmatrix} \rightarrow \begin{pmatrix} \lambda \mathbf{A} & \lambda^2 \mathbf{B} \\ \lambda \mathbf{C} & \lambda^2 \mathbf{D} \end{pmatrix} \quad (4.25)$$

this identification reduces further the dimension 16 down to 15. Eqs(4.23) are invariant under (4.25).

5 Integrable $D(2,1; \zeta)$ superspin chain

In this section, we give an application of the above analysis on the parametric $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ and the $S\mathfrak{D}\mathfrak{D}^{(2)}[\zeta]$ in the study of integrable superspin chains by using 4D Chern-Simons (CS) theory with $D(2,1; \zeta)$ symmetry. This will be done by calculating one of the basic operators in integrable spin systems namely the parametric Lax operator

$$\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\mu)} := \mathbf{L}(\zeta)$$

which, in the 4D CS gauge theory, is given by the Costello- Gaiotto- Yagi formula $e^X z^\mu e^Y$ [78]. Useful details on the Chern-Simons construction [38, 39, 76] and on the triplet (z^μ, X, Y) are reported in the Appendix D. Here, we focus on the calculation of

$\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\mu)}$ by using interpolation ideas motivated by algebraic geometry properties between the local data on $\mathcal{L}_{sl(2|2)}^{(\mu)}$ and $\mathcal{L}_{osp(4|2)}^{(\mu)}$ living at $\zeta = \pm 1$.

To undertake this study, we recall that integrable distinguished superspin chain $\mathfrak{d}(2,1;\zeta)$ is characterised by a root system $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ generated by three simple roots $(\tilde{\alpha}_1, \tilde{\alpha}_2, \tilde{\alpha}_3)$ with odd $\tilde{\alpha}_2$ and even $\tilde{\alpha}_1, \tilde{\alpha}_3$. This distinguished root basis is realised in terms of the canonical weights $(\epsilon_1, \epsilon_2, \delta)$ as in (3.18) namely,

$$\begin{aligned}\tilde{\alpha}_1 &= (\epsilon_1 - \epsilon_2) \\ \tilde{\alpha}_2 &= \frac{1}{2}\sqrt{2(1+\zeta)}\delta - \frac{1}{2}(\epsilon_1 - \epsilon_2) - \frac{1}{2}\sqrt{\zeta}(\epsilon_1 + \epsilon_2) \\ \tilde{\alpha}_3 &= \sqrt{\zeta}(\epsilon_1 + \epsilon_2)\end{aligned}\tag{5.1}$$

where ζ belongs to $\mathbb{C} \setminus \{-1, 0\}$. For these parametric simple roots, the intersection matrix $\tilde{\alpha}_i \cdot \tilde{\alpha}_j$ is a parametric 3×3 matrix $\tilde{\mathcal{J}}_{ij}(\zeta) = \tilde{\alpha}_i \cdot \tilde{\alpha}_j$ given by

$$\tilde{\mathcal{J}}_{ij}(\zeta) = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & -\zeta \\ 0 & -\zeta & 2\zeta \end{pmatrix}, \quad \det \tilde{\mathcal{J}}_{ij}(\zeta) = -2\zeta(1+\zeta)\tag{5.2}$$

with vanishing value ($\det \tilde{\mathcal{J}}_{ij}(\zeta) = 0$) at $\zeta = -1, 0$; and $\det \tilde{\mathcal{J}}_{ij}(\zeta)|_{\zeta=1} = -4$ for $osp(4|2)$ and super Cartan matrices as in (3.31) and (3.35). Eq(5.2) contains as subsystems two particular super symmetries located at $\zeta = \pm 1$ and then two integrable superspin sub-chains namely:

- the integrable $sl(2|2)$ superspin chain with Lax operator $\mathcal{L}_{sl(2|2)}^{(\mu)}$ at $\zeta = -1$; and
- the integrable $osp(4|2)$ spin chain with L-operator $\mathcal{L}_{osp(4|2)}^{(\mu)}$ at $\zeta = 1$.

To determine the $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\mu)}$, we use the following features:

- (i) the $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\mu)}$ must inherit properties of $\mathfrak{d}(2,1;\zeta)$; it should be a parametric operator with parameter ζ ; and must have critical values at $\zeta = -1, 0$ where $\det(\tilde{\alpha}_i \cdot \tilde{\alpha}_j) = 0$ and where $\mathfrak{d}(2,1;\zeta)$ contracts to semi-simple Lie sub-superalgebras.
- (ii) the expression of the parametric $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\mu)}$ must reproduce the values of $\mathcal{L}_{sl(2|2)}^{(\mu)}$ and $\mathcal{L}_{osp(4|2)}^{(\mu)}$ at $\zeta = \pm 1$; that is:

$$\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\mu)} \Big|_{\zeta=-1} = \mathcal{L}_{sl(2|2)}^{(\mu)} \quad , \quad \mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\mu)} \Big|_{\zeta=+1} = \mathcal{L}_{osp(4|2)}^{(\mu)}\tag{5.3}$$

Using these local features, we first calculate $\mathcal{L}_{sl(2|2)}^{(\mu)}$; thus completing results in [46, 77] concerning $\mathcal{L}_{sl(m|n)}^{(\mu)}$ for $m \neq n$. Then, we compute $\mathcal{L}_{osp(4|2)}^{(\mu)}$ which also completes results for $\mathcal{L}_{osp(2m|2n)}^{(\mu)}$ obtained in [46] for $m \geq 3$. After that, we give an approach to connect the $\mathcal{L}_{sl(2|2)}^{(\mu)}$ and $\mathcal{L}_{osp(4|2)}^{(\mu)}$ and then deduce a parametric realisation of $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\mu)}$.

5.1 Integrable $sl(2|2)$ superspin chain

We begin by recalling that the semi-simple Lie superalgebra $sl(2|2)$ [resp. the simple $psl(2|2)$] has rank $r = 3$ [resp. 2] and dimensions 15 [resp. 14]. It has two sectors: **(i)** an even sector $sl(2|2)_{\bar{0}}$ [resp. $psl(2|2)_{\bar{0}}$] describing bosons and splitting as

$$\begin{aligned} sl(2|2)_{\bar{0}} &= sl(2) \oplus sl(2) \oplus gl(1)_z \\ psl(2|2)_{\bar{0}} &= sl(2) \oplus sl(2) \end{aligned} \quad (5.4)$$

where $gl(1)_z$ is generated by the diagonal central element (identity operator). **(ii)** an odd sector $sl(2|2)_{\bar{1}}$ [resp. $psl(2|2)_{\bar{1}}$] given by the module $2|\bar{2} \oplus \bar{2}|2$ describing fermions.

The Levi- decomposition of the $sl(2|2)$ superalgebra reads in general as follows

$$\begin{aligned} sl(2|2) &\rightarrow \mathbf{N}_+^{(k,l)} \oplus \mathbf{l}_\mu^{(k,l)} \oplus \mathbf{N}_-^{(k,l)} \\ \mathbf{l}_\mu^{(k,l)} &= sl(k|l) \oplus sl(2-k|2-l) \oplus gl(1)_\mu \end{aligned} \quad (5.5)$$

where $\mathbf{l}_\mu^{(k,l)}$ is the Levi sub-superalgebra with $0 \leq k \leq 2$ and $0 \leq l \leq 2$ and where the nilpotent subalgebra $\mathbf{N}_\pm^{(k,l)}$ have the dimensions

$$\dim \mathbf{N}_+^{(k,l)} = \dim \mathbf{N}_-^{(k,l)} = (k+l)(4-k-l) \quad (5.6)$$

In graphical setting, the 3-grading (5.5) correspond to the cutting of any of the 3 nodes (bosonic or fermionic) of the four possible super Dynkin diagram reported in [Appendix A](#). This is a special property of $sl(m|n)$ where all nodes act like minuscule coweights; and as such the μ can be labeled like $\mu_{k,l}$ [46, 76, 77].

5.1.1 Distinguished $sl(2|2)$ spin chain

For our concern, we will be interested into the distinguished super Dynkin diagram of $sl(2|2)$ given by the **Figure 12**, and integers $(k, l) = (0, 1)$. Here, the Levi- decomposition




Figure 12: Distinguished super Dynkin diagram of the Lie superalgebra $sl(2|2)$. The odd root is given by the Green node α_2 with vanishing $\alpha_2 \cdot \alpha_2 = 0$.

of $sl(2|2)$ corresponds to the cutting of the (blue) node α_3 ; it is formulated like

$$\begin{aligned} sl(2|2) &: gl(1)_{\mu_3} \oplus sl(2|1) \oplus \mathbf{N}_+ \oplus \mathbf{N}_- \\ 15 \text{ dim} &: 1 + 8 + 3 + \bar{3} \\ 12 \text{ roots} &: 0 + 6 + 3 + \bar{3} \end{aligned} \quad (5.7)$$

with $sl(2|1)_0 = s[gl(2) \oplus gl(1)]$ and Levi-superalgebra $l_{\mu_3}^{sl(2|2)} = gl(1)_{\mu_3} \oplus sl(2|1)$. The interaction matrix $\alpha_i \cdot \alpha_j$ of the simple roots making the diagram of the **Figure 15** reads as

$$\alpha_i \cdot \alpha_j = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & 1 \\ 0 & 1 & -2 \end{pmatrix}, \quad \det(\alpha_i \cdot \alpha_j) = 0 \quad (5.8)$$

Under this decomposition, we have the following features:

(i) the positive root system $\Phi_{sl(2|2)}^+$ gets splitted like

$\Phi_{sl(2 2)}^+$	$\Phi_{sl(2 1)}^+$	\mathbf{N}_+
$\alpha_1, \alpha_2, \alpha_3,$	$\alpha_1, \alpha_2,$	α_3
$\alpha_2 + \alpha_1, \alpha_2 + \alpha_3$	$\alpha_2 + \alpha_1$	$\alpha_2 + \alpha_3$
$\alpha_1 + \alpha_2 + \alpha_3$	—	$\alpha_1 + \alpha_2 + \alpha_3$

(5.9)

and similarly for negative roots. For those 3+3 roots characterising the nilpotent subalgebras \mathbf{N}_{\pm} in (5.7), their realisations in terms of the canonical weight vectors $(\varepsilon_1, \varepsilon_2; \delta_1, \delta_2)$ read as

$$\alpha_3 = \delta_1 - \delta_2, \quad \alpha_2 + \alpha_3 = \varepsilon_2 - \delta_2, \quad \alpha_1 + \alpha_2 + \alpha_3 = \varepsilon_1 - \delta_2 \quad (5.10)$$

From this realisation, we deduce the 4×4 matrix representation of the step operators namely the even $E_{\pm\alpha_3}$ and the odd $F_{\pm(\alpha_2+\alpha_3)}, F_{\pm(\alpha_1+\alpha_2+\alpha_3)}$. For the example of the step operators associated with the positive roots $+\alpha$, we have the matrix representation

$$\begin{aligned} F_{\alpha_1+\alpha_2+\alpha_3} &= |\varepsilon_1\rangle \langle \delta_2| \equiv \tilde{\mathfrak{F}}_1 \\ F_{\alpha_2+\alpha_3} &= |\varepsilon_2\rangle \langle \delta_2| \equiv \tilde{\mathfrak{F}}_2 \\ E_{+\alpha_3} &= |\delta_1\rangle \langle \delta_2| \equiv \mathcal{E} \end{aligned} \quad (5.11)$$

from which we see that the odd $F_{\alpha_1+\alpha_2+\alpha_3}$ and $F_{\alpha_2+\alpha_3}$ form an isodoublet

$$\tilde{\mathfrak{F}}_i = (\tilde{\mathfrak{F}}_1, \tilde{\mathfrak{F}}_2)$$

while the even $E_{+\alpha_3}$ is an isosinglet \mathcal{E} . Similar features hold for the operators $F_{-\alpha_1-\alpha_2-\alpha_3}, F_{-\alpha_2-\alpha_3}$ and $E_{-\alpha_3}$ associated with negative roots. From this description, we also learn that the nilpotent representation $\mathbf{N}_+ = \mathbf{3}$ reduces as

$$\mathbf{3} = \mathbf{2} \oplus \mathbf{1}$$

describing a fermionic $sl(2)$ isodoublet $\tilde{\mathfrak{F}}_i \sim \mathbf{2}$ and a bosonic isosinglet $\mathcal{E} \sim \mathbf{1}$. The same thing is valid for $\mathbf{N}_- = \bar{\mathbf{3}}$ decomposing as $\bar{\mathbf{2}} \oplus \bar{\mathbf{1}}$.

(ii) the fundamental representation $\mathbf{2|2}$ of the super $sl(2|2)$ decomposes into irreps of $sl(2|1)$ as follows

$$\mathbf{2|2} \quad \rightarrow \quad \mathbf{2|} (\mathbf{1}_{1/2} \oplus \mathbf{1}_{-1/2}) \quad (5.12)$$

The subscripts $\pm 1/2$ designate the charges under the $gl(1)_{\mu_3}$ corresponding to the cutted node μ_3 dual to α_3 . These representations are labeled by the four basis states $|e_\lambda\rangle$ for $sl(2|2)$, the triplet $|e_a\rangle$ and the singlet $|e_\lambda\rangle$ for $sl(2|1)$. This convention notation is illustrated in the following table

algebra	$sl(2 2)$	$sl(2 1) \oplus gl(1)_{\mu_3}$	
reps	$\mathbf{2 2}$	$\mathbf{2 1}_{1/2} \oplus \mathbf{0 1}_{-1/2}$	(5.13)
basis states	$ e_\lambda\rangle$	$ e_a\rangle \oplus e_\lambda\rangle$	

with the gradings

$$|e_\lambda\rangle = |\varepsilon_i\rangle \oplus |\delta_j\rangle \quad , \quad |e_a\rangle = |\varepsilon_i\rangle \oplus |\delta_1\rangle \quad , \quad |e_\lambda\rangle = |\delta_2\rangle \quad (5.14)$$

where $i, j = 1, 2$. Using these quantum states ordered as $(|\varepsilon_1\rangle, |\varepsilon_2\rangle, |\delta_1\rangle, |\delta_2\rangle)$, we can write down the action of the coweight μ_3 ; it is given by help of the projectors $\Pi_1 = \sum_{a=1}^3 |e_a\rangle \langle e_a|$ and $\Pi_2 = |\delta_2\rangle \langle \delta_2|$ as follows

$$\begin{aligned} \mu_3 &= \frac{1}{2} (|\varepsilon_1\rangle \langle \varepsilon_1| + |\varepsilon_2\rangle \langle \varepsilon_2| + |\delta_1\rangle \langle \delta_1|) - \frac{1}{2} |\delta_2\rangle \langle \delta_2| \\ &= \frac{1}{2} \Pi_1 - \frac{1}{2} \Pi_2 \end{aligned} \quad (5.15)$$

The μ_3 has a vanishing super trace $str(\mu_3) = 0$.

5.1.2 Calculating L-operator $\mathcal{L}_{sl_{2|2}}^{\mu_3}$

Having the above tools at hand, we can now calculate the Lax operator of the integrable distinguished $sl(2|2)$ superspin chain by using the Costello- Gaiotto- Yagi (CGY) formula [76–78]

$$\mathcal{L}_{sl_{2|2}}^{\mu_3} = e^X z^{\mu_3} e^Y \quad (5.16)$$

The X and Y are nilpotent operators belonging to \mathbf{N}_+ and \mathbf{N}_- appearing in (5.7); their explicit expressions are needed for the computation of the Lax operator. The z^{μ_3} is diagonal given by (5.15) while the X and Y are determined by resolving the Levi-constraints

$$[\mu_3, X] = X \quad , \quad [\mu_3, Y] = -Y \quad , \quad [X, Y] = \mathbb{C}\mu_3 \quad (5.17)$$

These conditions are solved in terms of three pairs of harmonic oscillators: an even (b, c) —a singlet—; and two odd (β^i, γ_i) —a doublet— as follows

$$\begin{aligned} X &= b |\delta_1\rangle \langle \delta_2| + \sum_{i=1}^2 \beta^i |\varepsilon_i\rangle \langle \delta_2| \\ Y &= c |\delta_2\rangle \langle \delta_1| + \sum_{i=1}^2 \gamma_i |\delta_2\rangle \langle \varepsilon^i| \end{aligned} \quad (5.18)$$

which can be shortly expressed like

$$X = |\Psi_X\rangle \langle \delta_2| \quad , \quad Y = |\delta_2\rangle \langle \Psi_Y| \quad (5.19)$$

with $|\Psi_X\rangle = \beta^i |\varepsilon_i\rangle + b |\delta_1\rangle$ and $\langle \Psi_Y| = \gamma_j \langle \varepsilon^j| + c \langle \delta_1|$. In this regard, recall that in the 4D Chern-Simons description of integrable superspin chains, the pair (b, c) is a bosonic harmonic oscillator while the two pairs (β^i, γ_i) are two fermionic oscillators. Quantum mechanically, these pairs obey the graded commutation relations

$$[b, c] = 1 \quad , \quad \{\beta^i, \gamma_j\} = \delta_j^i$$

instead of the classical orthosymplectic Poisson brackets. For the explicit calculations given below, we treat them classically, quantum corrections can be implemented straightforwardly as done in [76]. Using the above relationships, we can compute the expressions of monomials of X and Y for the calculation of (5.16). We have

$$\begin{aligned} X^2 &= 0 \quad , \quad X\Pi_1 = \Pi_1 Y = 0 \\ Y^2 &= 0 \quad , \quad \Pi_2 X = Y\Pi_2 = 0 \end{aligned} \quad (5.20)$$

and $XY = |\Psi_X\rangle \langle \Psi_Y|$ expanding like

$$XY = \beta^i \gamma_j |\varepsilon_i\rangle \langle \varepsilon^j| + bc |\delta_1\rangle \langle \delta_1| + b\gamma_j |\delta_1\rangle \langle \varepsilon^j| + c\beta^i |\varepsilon_i\rangle \langle \delta_1| \quad (5.21)$$

Using these relations, the Lax operator $e^X z^{\mu_3} e^Y$ expands like $(1 + X) z^{\mu_3} (1 + Y)$. By substituting (5.15-5.18), we get

$$\mathcal{L}_{sl_{2|2}}^{\mu_3} = z^{\mu_3} + X z^{\mu_3} + z^{\mu_3} Y + X z^{\mu_3} Y \quad (5.22)$$

In terms of the matrix representation $\mathcal{L}_{sl_{2|2}}^{\mu_3} = \sum |e_A\rangle L_B^A \langle e^B|$ in the basis $(|\varepsilon_1\rangle, |\varepsilon_2\rangle, |\delta_1\rangle, |\delta_2\rangle)$, the L-operator reads in block submatrices as follows

$$L_B^A = \begin{pmatrix} z^{1/2} \delta_j^i + z^{-1/2} \beta^i \gamma_j & z^{-1/2} c \beta^i & z^{-1/2} \beta^i \\ z^{-1/2} b \gamma_j & z^{1/2} + z^{-1/2} bc & z^{-1/2} b \\ z^{-1/2} \gamma_j & z^{-1/2} c & z^{-1/2} \end{pmatrix} \quad (5.23)$$

This L-matrix agrees with the solution obtained in the superspin chain literature in [80] using algebraic methods to solve the RLL equation.

5.2 Integrable $\mathfrak{osp}(4|2)$ superspin chain

In this subsection, we calculate the Lax operator $\mathcal{L}_{osp}^{\mu_3}$ of distinguished integrable $\mathfrak{osp}(4|2)$ superchain by using the 4D Chern-Simons formalism; for an alternative algebraic method, see [81]. Here, the L-operator is given by the CGY formula $e^X z^{\mu_3} e^Y$ with (z^{μ_3}, X, Y) valued in $\mathfrak{osp}(4|2)$. Recall that the distinguished super Dynkin diagram of the orthosymplectic $\mathfrak{osp}(4|2)$ superalgebra is given by the **Figure 13**. This super diagram




Figure 13: Distinguished super Dynkin diagram of the Lie superalgebra $\mathfrak{osp}(4|2)$. The odd root is given by the Green node α_2 with vanishing $\alpha_2 \cdot \alpha_2 = 0$.

can be thought of in terms of the intersection matrix of the simple roots $(\alpha_1, \alpha_2, \alpha_3)$ labeling its three nodes with odd α_2 . This intersection matrix reads as follows

$$\alpha_i \cdot \alpha_j = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & -1 \\ 0 & -1 & 2 \end{pmatrix}, \quad \det(\alpha_i \cdot \alpha_j) = -4 \quad (5.24)$$

The super diagram **13** has an outer automorphism \mathcal{Z}_2 permuting the α_1 and the α_3 simple roots (red nodes in the diagram) with coweights μ_1 and μ_3 which are needed for the calculation of the associated Lax operators $\mathcal{L}_{osp}^{\mu} = e^X z^{\mu} e^Y$.

5.2.1 Distinguished orthosymplectic spin chain $\mathfrak{osp}(4|2)$

The Levi-decomposition (3-grading) for the distinguished orthosymplectic $\mathfrak{osp}(4|2)$ chain is obtained by the *cutting* the node α_3 in the super Dynkin diagram of the **Figure 13**. This decomposition is given by

$$\begin{aligned} \mathfrak{osp}(4|2) &: \mathfrak{gl}(1)_{\mu_3} \oplus \mathfrak{sl}(2|1) \oplus \mathbf{N}_+^{osp} \oplus \mathbf{N}_-^{osp} \\ 17 \text{ dim} &: 1 + 8 + 4 + \bar{4} \\ 14 \text{ roots} &: 0 + 6 + 4 + \bar{4} \end{aligned} \quad (5.25)$$

with $\mathfrak{sl}(2|1) \simeq \mathfrak{osp}(2|2)$ and $\mathfrak{gl}(1)_{\mu_3} \simeq \mathfrak{so}(2, \mathbb{C})$ and where $\mathfrak{l}_{\mu_3}^{osp} = \mathfrak{gl}(1)_{\mu_3} \oplus \mathfrak{sl}(2|1)$ is the Levi-subalgebra. Comparing the 3-grading in (5.25) with the 3-grading (5.7), we learn

that we have the isomorphism $\mathfrak{l}_{\mu_3}^{sl_2|2} \simeq \mathfrak{l}_{\mu_3}^{osp}$ while the nilpotent subalgebras \mathbf{N}_{\pm}^{osp} and $\mathbf{N}_{\pm}^{sl_2|2}$ are related like

$$\mathbf{N}_{\pm}^{osp} \simeq \mathbf{N}_{\pm}^{sl_2|2} \oplus \mathbf{1}_{\pm} \quad \Leftrightarrow \quad \mathbf{4}_{\pm}^{osp} \simeq \mathbf{3}_{\pm}^{sl_2|2} \oplus \mathbf{1}_{\pm} \quad (5.26)$$

showing that \mathbf{N}_{\pm}^{osp} has one dimension bigger with respect to $\mathbf{N}_{\pm}^{sl_2|2}$.

Below, we use the 6-dimensional vector representation of $osp(4|2)$ labeled by six canonical weight vectors $\{\mathbf{e}_1, \dots, \mathbf{e}_6\}$ thought of in terms of the $\pm\epsilon_i$ and $\pm\delta$ weight vectors ($\delta^2 = \epsilon_i^2 = -1$) as follows

$$\mathbf{e}_1 = \epsilon_1, \quad \mathbf{e}_2 = \epsilon_2, \quad \mathbf{e}_3 = \delta, \quad \mathbf{e}_4 = \bar{\delta}, \quad \mathbf{e}_5 = \bar{\epsilon}_2, \quad \mathbf{e}_6 = \bar{\epsilon}_1 \quad (5.27)$$

This set of quantum states can be combined into two subset like $|e_a\rangle \oplus |\bar{e}^a\rangle$ with $|e_a\rangle = (|\epsilon_1\rangle, |\epsilon_2\rangle, |\delta\rangle)$ and $|\bar{e}^a\rangle = (|\bar{\delta}\rangle, |\bar{\epsilon}^2\rangle, |\bar{\epsilon}^1\rangle)$. For the bras, we have $\langle e^a| = (\langle\epsilon^1|, \langle\epsilon^2|, \langle\delta|)$ and $|\bar{e}_{\bar{a}}\rangle = (\langle\bar{\delta}|, \langle\bar{\epsilon}_2|, \langle\bar{\epsilon}_1|)$. Under the decomposition (5.25), we have the following interesting features:

(a) the positive root system $\Phi_{osp(4|2)}^+$ gets splitted into $\Phi_{sl(2|1)}^+ \oplus \mathbf{N}_+^{osp}$ like

$\Phi_{osp(4 2)}^+$	$\Phi_{sl(2 1)}^+$	\mathbf{N}_+^{osp}	(5.28)
$\alpha_1, \quad \alpha_2, \quad \alpha_3,$	$\alpha_1, \quad \alpha_2,$	$\alpha_3, \quad \alpha_1 + 2\alpha_2 + \alpha_3$	
$\alpha_2 + \alpha_1, \quad \alpha_1 + \alpha_2 + \alpha_3$	$\alpha_2 + \alpha_1$	$\alpha_2 + \alpha_3,$	
$\alpha_2 + \alpha_3, \quad \alpha_1 + 2\alpha_2 + \alpha_3$	—	$\alpha_1 + \alpha_2 + \alpha_3$	

with the simple roots realised as $\alpha_1 = \epsilon_1 - \epsilon_2$, $\alpha_2 = \delta - \epsilon_1$ and $\alpha_3 = \epsilon_1 + \epsilon_2$.

For the roots characterising the nilpotent sector \mathbf{N}_+^{osp} , we have the roots

$$\begin{aligned} \alpha_3 &= \epsilon_1 + \epsilon_2 \\ \alpha_3 + \alpha_2 &= \delta + \epsilon_2 \\ \alpha_3 + \alpha_2 + \alpha_1 &= \delta + \epsilon_1 \\ \alpha_3 + 2\alpha_2 + \alpha_1 &= 2\delta \end{aligned} \quad (5.29)$$

from which we learn that the $\mathbf{N}_+ = \mathbf{4}$ reduces like

$$\mathbf{4} = \mathbf{2} \oplus \mathbf{1} \oplus \mathbf{1}$$

where the doublet $\mathbf{2}$ describes the two odd roots $\delta + \epsilon_i$ and the two singlets $\mathbf{1} \oplus \mathbf{1}$ corresponding to the even $\epsilon_1 + \epsilon_2$ and the even long root 2δ .

(b) the fundamental representation $\mathbf{4|2}$ of the orthosymplectic Lie algebra $osp(4|2)$ decomposes as the sum of two irreps of $sl(2|1)$ as follows

$$(\mathbf{4|2}) = (\mathbf{2|1})_{+\frac{1}{2}} \oplus (\bar{\mathbf{2}}|\bar{\mathbf{1}})_{-\frac{1}{2}} \quad (5.30)$$

The subscripts $\pm 1/2$ designate the charges under the Cartan subalgebra $gl(1)_{\mu_3}$. These representations are described by the basis states $|e_a\rangle \oplus |\bar{e}^{\bar{a}}\rangle$ like

algebra	osp(4 2)		$sl(2 1) \oplus gl(1)_{\mu_3}$		
reps	$\mathbf{4 2}$		$\mathbf{2 1}_{1/2}$	\oplus	$\bar{\mathbf{2}} \bar{\mathbf{1}}_{-1/2}$
basis states	$ e_a\rangle$		$ e_a\rangle$	\oplus	$ \bar{e}^{\bar{a}}\rangle$

(5.31)

5.2.2 Computing the orthosymplectic $\mathcal{L}_{osp(4|2)}^{\mu_3}$

Given the above quantum states ordered as in (5.27), we can:

(i) write down the action of the coweight μ_3^{osp} (dual to α_3^{osp}); it is given by using projectors Π and $\bar{\Pi}$ as follows

$$\mu_3^{osp} = \frac{1}{2}\Pi - \frac{1}{2}\bar{\Pi} \quad (5.32)$$

with

$$\begin{aligned} \Pi &= |\epsilon_1\rangle \langle \epsilon^1| + |\epsilon_2\rangle \langle \epsilon^2| + |\delta_1\rangle \langle \delta^1| \\ \bar{\Pi} &= |\bar{\delta}^1\rangle \langle \bar{\delta}_1| + |\bar{\epsilon}^2\rangle \langle \bar{\epsilon}_2| + |\bar{\epsilon}^1\rangle \langle \bar{\epsilon}_1| \end{aligned} \quad (5.33)$$

It has a vanishing super trace $str(\mu_3^{osp}) = 0$.

(ii) calculate the super L-operator $\mathcal{L}_{osp}^{\mu_3}$ (or equivalently $\mathcal{L}_{osp}^{\mu_1}$); it is given by a 6×6 matrix having the sub-blocks form

$$(L_{osp})_B^A = \begin{pmatrix} L_b^a & L_{\bar{b}}^{\bar{a}} \\ L_{\bar{b}}^a & L_b^{\bar{a}} \end{pmatrix} \quad (5.34)$$

with basis vectors $|e_a\rangle$ for the representation $(\mathbf{2|1})_{+1/2}$ ($|e_a\rangle = (|\epsilon_1\rangle, |\epsilon_2\rangle, |\delta\rangle)$); and $|\bar{e}^{\bar{a}}\rangle$ for the representation $(\bar{\mathbf{2}}|\bar{\mathbf{1}})_{-1/2}$ with $|\bar{e}^{\bar{a}}\rangle = (|\bar{\delta}\rangle, |\bar{\epsilon}^2\rangle, |\bar{\epsilon}^1\rangle)$.

Using eqs(5.29-5.31), we can determine the super Lax operator $\mathcal{L}_{osp}^{\mu_3}$ by help of the representation $e^{X_{osp}} z^{\mu_3} e^{Y_{osp}}$ with μ_3^{osp} as in (5.32). For the nilpotent X_{osp} and Y_{osp} , they expand like

$$\begin{aligned} X_{osp} &= \beta^i \mathcal{X}_i + bX_0 + b'X'_0 \\ Y_{osp} &= \gamma_i \mathcal{Y}^i + cY_0 + c'Y'_0 \end{aligned} \quad (5.35)$$

and obey the Levi- constraint relations (5.17) namely $[\mu_3, X] = X$ and $[\mu_3, Y] = -Y$. In these relations the pairs (b,c) and (b',c') are bosonic oscillators and the pairs (β^i, γ_i)

are fermionic ones. The above mentioned Levi- conditions can be solved as follows

$$\begin{aligned}
\mathcal{X}_i &= |\epsilon_i\rangle \langle \bar{\delta}| - |\delta\rangle \langle \bar{\epsilon}_i| & \mathcal{Y}^i &= |\bar{\delta}\rangle \langle \epsilon^i| - |\bar{\epsilon}^i\rangle \langle \delta| \\
X_0 &= |\delta\rangle \langle \bar{\delta}| & Y_0 &= |\bar{\delta}\rangle \langle \delta| \\
X'_0 &= |\epsilon_1\rangle \langle \bar{\epsilon}_2| - |\epsilon_2\rangle \langle \bar{\epsilon}_1| & Y'_0 &= |\bar{\epsilon}^2\rangle \langle \epsilon^1| - |\bar{\epsilon}^1\rangle \langle \epsilon^2|
\end{aligned} \tag{5.36}$$

where $|\epsilon_1\rangle \langle \bar{\epsilon}_2| - |\epsilon_2\rangle \langle \bar{\epsilon}_1|$ can be expressed shortly as $\epsilon^{kl}|\epsilon_k\rangle \langle \bar{\epsilon}_l|$ in terms of the usual antisymmetric tensor ϵ^{kl} . We also have $|\bar{\epsilon}^l\rangle \langle \epsilon^k| \epsilon_{kl}$ for the singlet $|\bar{\epsilon}^2\rangle \langle \epsilon^1| - |\bar{\epsilon}^1\rangle \langle \epsilon^2|$. Substituting these expressions into (5.35), we can present X_{osp} and Y_{osp} like

$$\begin{aligned}
X_{osp} &= \beta^k (|\epsilon_k\rangle \langle \bar{\delta}| - |\delta\rangle \langle \bar{\epsilon}_k|) + b |\delta\rangle \langle \bar{\delta}| + b' \epsilon^{kl} |\epsilon_k\rangle \langle \bar{\epsilon}_l| \\
Y_{osp} &= \gamma_n (|\bar{\delta}\rangle \langle \epsilon^n| - |\bar{\epsilon}^n\rangle \langle \delta|) + c |\bar{\delta}\rangle \langle \delta| + c' |\bar{\epsilon}^m\rangle \langle \epsilon^n| \epsilon_{nm}
\end{aligned} \tag{5.37}$$

From these relations, we learn the properties $X_{osp}^2 = 0 = Y_{osp}^2$ indicating that $e^{X_{osp}} = 1 + X_{osp}$ and $e^{Y_{osp}} = 1 + Y_{osp}$. We also learn the useful features $X_{osp}\Pi = 0 = \Pi Y_{osp}$ as well as $\Pi X_{osp} = X_{osp}$ and $\Pi Y_{osp} = Y_{osp}$. For convenience, we express the above X_{osp} and Y_{osp} as follows

$$\begin{aligned}
X_{osp} &= |\Phi_X\rangle \langle \bar{\delta}| - |\Psi_X^k\rangle \langle \bar{\epsilon}_k| \\
Y_{osp} &= |\bar{\delta}\rangle \langle \Phi_Y| - |\bar{\epsilon}^l\rangle \langle \Psi_{Yl}|
\end{aligned} \tag{5.38}$$

with

$$\begin{aligned}
|\Psi_X^k\rangle &= b' \epsilon^{kl} |\epsilon_l\rangle + \beta^k |\delta\rangle & |\Phi_X\rangle &= \beta^k |\epsilon_k\rangle + b |\delta\rangle \\
\langle \Psi_{Yn}| &= c' \langle \epsilon^m| \epsilon_{mn} + \gamma_n \langle \delta| & \langle \Phi_Y| &= \gamma_n \langle \epsilon^n| + c \langle \delta|
\end{aligned} \tag{5.39}$$

leading to

$$\begin{aligned}
X_{osp} Y_{osp} &= |\Phi_X\rangle \langle \Phi_Y| + |\Psi_X^k\rangle \langle \Psi_{Yk}| \\
|\Phi_X\rangle \langle \Phi_Y| &= \beta^i \gamma_j |\epsilon_i\rangle \langle \epsilon^j| + c \beta^i |\epsilon_i\rangle \langle \delta| + b \gamma_j |\delta\rangle \langle \epsilon^j| + bc |\delta\rangle \langle \delta| \\
|\Psi_X^k\rangle \langle \Psi_{Yk}| &= b' c' |\epsilon_l\rangle \langle \epsilon^l| + \beta^k c' \epsilon_{mk} |\delta\rangle \langle \epsilon^m| + b' \gamma_k \epsilon^{kl} |\epsilon_l\rangle \langle \delta| + \beta^k \gamma_k |\delta\rangle \langle \delta|
\end{aligned} \tag{5.40}$$

Using these relations, the Lax operator $e^X z^{\mu_3} e^Y$ expands like $(1 + X) z^{\mu_3} (1 + Y)$. By substituting (5.15-5.18), we get $\mathcal{L}_{osp}^{\mu_3} = z^{\mu_3} + z^{-1/2} (X + Y + XY)$. The matrix elements L_B^A in the basis $(|e_a\rangle, |\bar{e}^a\rangle)$ reads in block matrices as follows

$$\mathcal{L}_{osp}^{\mu_3} = \begin{pmatrix} z^{1/2} \delta_j^i + z^{-1/2} (b' c' \delta_j^i + \beta^i \gamma_j) & z^{-1/2} (c \beta^i + b' \gamma_k \epsilon^{ki}) & z^{-1/2} \beta^i & z^{-1/2} b' \epsilon^{ij} \\ z^{-1/2} (b \gamma_j + c' \epsilon_{jl} \beta^l) & z^{1/2} + z^{-1/2} (bc + \beta^l \gamma_l) & z^{-1/2} b & -z^{-1/2} \beta^j \\ z^{-1/2} \gamma_j & z^{-1/2} c & z^{-1/2} & 0 \\ z^{-1/2} c' \epsilon_{ij} & -z^{-1/2} \gamma_i & 0 & z^{-1/2} \delta_i^j \end{pmatrix} \tag{5.41}$$

This L-matrix completes the series of the expressions of the L-operators for $\text{osp}(2m|2n)$ given in [46] for $m \geq 3$ and $n \geq 1$. Notice that by setting $b' = c' = 0$ in (5.41), it reduces down to $\mathcal{L}_{osp}^{\mu_3}|_{b'=c'=0}$ reading as follows

$$\begin{pmatrix} z^{1/2}\delta_j^i + z^{-1/2}\beta^i\gamma_j & z^{-1/2}c\beta^i & z^{-1/2}\beta^i & 0 \\ z^{-1/2}b\gamma_j & z^{1/2} + z^{-1/2}(bc + \beta^l\gamma_l) & z^{-1/2}b & -z^{-1/2}\beta^j \\ z^{-1/2}\gamma_j & z^{-1/2}c & z^{-1/2} & 0 \\ 0 & -z^{-1/2}\gamma_i & 0 & z^{-1/2}\delta_i^j \end{pmatrix} \quad (5.42)$$

containing in turns $\mathcal{L}_{sl(2|2)}^{\mu_3}$ as a 4×4 submatrix as follows

$$\begin{pmatrix} & 0 \\ \left[\mathcal{L}_{sl(2|2)}^{\mu_3} \right]_{4 \times 4} & -z^{-1/2}\beta^j \\ & 0 \\ 0 & -z^{-1/2}\gamma_i & 0 & z^{-1/2}\delta_i^j \end{pmatrix} \quad (5.43)$$

5.3 Lax operator of $\mathfrak{D}(2,1;\zeta)$ superspin chain

Here, we use the obtained L-operators $\mathcal{L}_{sl(2|2)}^{\mu_3}$ and $\mathcal{L}_{osp}^{\mu_3}$ given by eqs(5.23) and (5.41) to determine $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$. We think about (5.23) as the value $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}|_{\zeta=-1}$ and about (5.41) like the value of $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}|_{\zeta=+1}$ as indicated by (5.3).

5.3.1 $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$ as deformation of $\mathcal{L}_{osp}^{\mu_3}$

Despite our knowledge of the values $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}|_{\zeta=\pm 1}$ given by (5.23) and (5.41), this is not enough to construct $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$ for generic values of ζ . To get around this difficulty, we use the fact that $\mathfrak{d}(2,1;\zeta)$ can be seen as a deformation of $\text{osp}(4|2)=\mathfrak{d}(2,1;1)$. In this vision, we propose to construct $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$ starting from the orthosymplectic solutions (5.37) that led to $\mathcal{L}_{osp}^{\mu_3}$ and consider their deformations towards $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$ with the conditions (5.3); i.e:

$$\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3} = \mathcal{L}_{osp}^{\mu_3} + \delta\mathcal{L}(\zeta) \quad (5.44)$$

So, to determine the expression of the L-operator of the integrable distinguishable superspin chain $\mathfrak{d}(2,1;\zeta)$, we start from the solution (5.37) and use the fact that $\mathfrak{d}(2,1;\zeta)$ and $\text{osp}(4|2)$ have the same dimensions ($\dim = 17$) and the same rank ($r = 3$) to think about $\mu_3^{\mathfrak{d}(2,1;\zeta)}$ like

$$\mu_3^{\mathfrak{d}(2,1;\zeta)} = \frac{1}{2}\Pi - \frac{1}{2}\bar{\Pi} \quad (5.45)$$

and the nilpotent operators $X_{\mathfrak{d}(2,1;\zeta)}$ and $Y_{\mathfrak{d}(2,1;\zeta)}$ as follows

$$\begin{aligned} X_{\mathfrak{d}(2,1;\zeta)} &= \beta^k (\kappa_1 |\epsilon_k\rangle \langle \bar{\delta}| + \kappa_2 |\delta\rangle \langle \bar{\epsilon}_k|) + b |\delta\rangle \langle \bar{\delta}| + b' \kappa_0 \epsilon^{kl} |\epsilon_k\rangle \langle \bar{\epsilon}_l| \\ Y_{\mathfrak{d}(2,1;\zeta)} &= \gamma_n (\kappa_1 |\bar{\delta}\rangle \langle \epsilon^n| + \kappa_1 |\bar{\epsilon}^n\rangle \langle \delta|) + c |\bar{\delta}\rangle \langle \delta| + c' \bar{\kappa}_0 |\bar{\epsilon}^m\rangle \langle \epsilon^n| \epsilon_{nm} \end{aligned} \quad (5.46)$$

In the above relations, the 3+3 moduli

$$\kappa_i(\zeta) \quad , \quad \bar{\kappa}_i(\zeta) \quad (5.47)$$

capture the deformation away from the $\mathfrak{osp}(4|2)$ solution (5.35-5.36). These are functions of ζ that we determine by using the two following conditions:

- (i) Eqs (5.45-5.46) must satisfy the Levi-constraint relations $[\mu_3, X] = X$ and $[\mu_3, Y] = -Y$. Actually, eqs(5.46) are particular deformations of (5.37) because a general deformation would involve more free parameters than the six $(\kappa_i, \bar{\kappa}_i)$ in eq(5.47). Eq(5.46) were chosen because they are the general ones effectively solving $[\mu_3, X] = X$ and $[\mu_3, Y] = -Y$. A reduced family of solutions is given by setting $\kappa_i(\zeta) = \bar{\kappa}_i(\zeta)$; thus restricting the number of free deformation functions to 3. Further simpler solutions are given by putting extra constraints on the three $\kappa_i(\zeta)$ as we will see below.
- (ii) the $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$ based on (5.46) is a function of $\kappa_i(\zeta)$, $\bar{\kappa}_i(\zeta)$; its value should reproduce the solutions obtained previously in deriving L-operators $\mathcal{L}_{sl(2|2)}^{\mu_3}$ and $\mathcal{L}_{osp}^{\mu_3}$ given by eqs(5.23) and (5.41). So, we must have

$\mathfrak{d}(2, 1; \zeta)$:	κ_1	κ_2	κ_0	$\bar{\kappa}_1$	$\bar{\kappa}_2$	$\bar{\kappa}_0$	(5.48)
$sl(2 2)$:	1	0	0	1	0	0	
$\mathfrak{osp}(4 2)$:	1	-1	1	1	-1	1	

indicating that κ_2 and κ_0 may be related like $\kappa_2 = -\kappa_0$; thus restricting the number of free functions.

5.3.2 Calculation of $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$

To determine the L-operator $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$ while using (5.45-5.46), we rewrite eq(5.46) as follows

$$\begin{aligned} X_{\mathfrak{d}(2,1;\zeta)} &= |\Phi_X\rangle \langle \bar{\delta}| + |\Psi_X^k\rangle \langle \bar{\epsilon}_k| \\ Y_{\mathfrak{d}(2,1;\zeta)} &= |\bar{\delta}\rangle \langle \Phi_Y| + |\bar{\epsilon}^l\rangle \langle \Psi_{Yl}| \end{aligned} \quad (5.49)$$

with

$$\begin{aligned} |\Phi_X\rangle &= \kappa_1 \beta^k |\epsilon_k\rangle + b |\delta\rangle \\ \langle \Phi_Y| &= \bar{\kappa}_1 \gamma_n \langle \epsilon^n| + c \langle \delta| \end{aligned} \quad (5.50)$$

and

$$\begin{aligned} |\Psi_X^k\rangle &= \kappa_0 b' \epsilon^{kl} |\epsilon_l\rangle + \kappa_2 \beta^k |\delta\rangle \\ \langle \Psi_{Yn}| &= \bar{\kappa}_0 c' \langle \epsilon^m| \epsilon_{mn} + \bar{\kappa}_2 \gamma_n \langle \delta| \end{aligned} \quad (5.51)$$

where (β^k, γ_k) are fermionic oscillators, and (b, c) and (b', c') are bosonic oscillator; they are the same oscillators as for $\text{osp}(4|2)$. From these relations, we calculate the product $XY = |\Phi_X\rangle \langle \Phi_Y| + |\Psi_X^n\rangle \langle \Psi_{Y_n}|$ expanding as follows

$$\begin{aligned} |\Phi_X\rangle \langle \Phi_Y| &= (\kappa_1 \bar{\kappa}_1) \beta^k \gamma_n |\epsilon_k\rangle \langle \epsilon^n| + bc |\delta\rangle \langle \delta| + \\ &\quad \kappa_1 c \beta^k |\epsilon_k\rangle \langle \delta| + \bar{\kappa}_1 b \gamma_n |\delta\rangle \langle \epsilon^n| \end{aligned} \quad (5.52)$$

and

$$\begin{aligned} |\Psi_X^n\rangle \langle \Psi_{Y_n}| &= (\kappa_0 \bar{\kappa}_0) b' c' |\epsilon_m\rangle \langle \epsilon^m| + (\kappa_2 \bar{\kappa}_2) \beta^n \gamma_n |\delta\rangle \langle \delta| + \\ &\quad (\kappa_0 \bar{\kappa}_2) b' \gamma_n \epsilon^{nl} |\epsilon_l\rangle \langle \delta| + (\kappa_2 \bar{\kappa}_0) c' \epsilon_{mn} \beta^n |\delta\rangle \langle \epsilon^m| \end{aligned} \quad (5.53)$$

Moreover, using the properties and $X^2 = Y^2 = 0$, the L-operator $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$ reduces to $(1+X)z^{\mu_3}(1+Y)$ with $z^{\mu_3} = z^{1/2}\Pi + z^{-1/2}\bar{\Pi}$ and X, Y as in (5.49). By help of the properties

$$\begin{aligned} \Pi X &= X & X \bar{\Pi} &= X & X \Pi &= \Pi Y &= 0 \\ \bar{\Pi} Y &= Y & Y \Pi &= Y & Y \bar{\Pi} &= \bar{\Pi} X &= 0 \end{aligned} \quad (5.54)$$

it can be further reduced down to the form $z^{\mu_3} + z^{-1/2}(X+Y+XY)$. By substituting, the $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{\mu_3}$ reads as follows

$$\begin{pmatrix} (z^{\frac{1}{2}} + \kappa_0 \bar{\kappa}_0 z^{-\frac{1}{2}} b' c') \delta_j^i + \kappa_1 \bar{\kappa}_1 z^{-\frac{1}{2}} \beta^i \gamma_j & z^{-\frac{1}{2}} (\kappa_1 c \beta^i + \kappa_0 \bar{\kappa}_2 b' \gamma_n \epsilon^{ni}) & \kappa_1 z^{-\frac{1}{2}} \beta^i & \kappa_0 z^{-\frac{1}{2}} b' \epsilon^{ji} \\ z^{-\frac{1}{2}} (\bar{\kappa}_1 b \gamma_j + \bar{\kappa}_0 \kappa_2 c' \epsilon_{jn} \beta^n) & z^{\frac{1}{2}} + z^{-\frac{1}{2}} (bc + \kappa_2 \bar{\kappa}_2 \beta^n \gamma_n) & z^{-\frac{1}{2}} b & \kappa_2 z^{-\frac{1}{2}} \beta^j \\ z^{-\frac{1}{2}} \bar{\kappa}_1 \gamma_j & z^{-\frac{1}{2}} c & z^{-\frac{1}{2}} & 0 \\ \bar{\kappa}_0 z^{-\frac{1}{2}} c' \epsilon_{ji} & \bar{\kappa}_2 z^{-\frac{1}{2}} \gamma_i & 0 & z^{-\frac{1}{2}} \delta_i^j \end{pmatrix} \quad (5.55)$$

By taking $\bar{\kappa}_i(\zeta) = \kappa_i(\zeta)$, the L-operator has three free functions constrained like

$sl(2 2)$	$osp(4 2)$	
$\kappa_1(\zeta) _{\zeta=-1} = 1$	$\kappa_1(\zeta) _{\zeta=+1} = +1$	(5.56)
$\kappa_2(\zeta) _{\zeta=-1} = 0$	$\kappa_2(\zeta) _{\zeta=+1} = -1$	
$\kappa_0(\zeta) _{\zeta=-1} = 0$	$\kappa_0(\zeta) _{\zeta=+1} = +1$	

A particular solution is given by taking $\kappa_1(\zeta) = 1$ and $\kappa_2(\zeta) = -\kappa_0(\zeta)$. Using the property $\det(\tilde{\alpha}_i, \tilde{\alpha}_j) = -2\zeta(1+\zeta)$, it is then natural to think about the deformation function $\kappa_0(\zeta)$ as given by $\frac{1}{2}\zeta(1+\zeta)$.

6 Conclusion and comments

To conclude this study, we summarise our main finds along with some additional comments. Motivated by getting more insight into the properties of $D(2, 1; \zeta)$ and its applications in (i) integrable exotic superspin chains; and (ii) in building type II string 3D models, we revisited useful algebraic and geometric aspects on this 9|8 dimensional exceptional parametric symmetry. In the present work, we focused on the algebraic geometry of complexified $D(2, 1; \zeta)$ and its application in the study of the integrable superspin chain with Lie superalgebra $\mathfrak{d}(2, 1; \zeta)$ within the framework of 4D Chern-Simons theory. The real forms of this parametric super symmetry will be considered in a forthcoming occasion focussing on applications aimed at the study of parametric AdS_3 strings along the lines of [14, 82–84].

To undertake the investigation on the algebraic geometry of the parametric $D(2, 1; \zeta)$ and its applications in integrable super systems à la Chern-Simons, we took advantage of its local structures at some special points of the complex variable ζ ; in particular:

- (1) at $\zeta = 1$ where $D(2, 1; \zeta)$ identifies with the 9|8-dim orthosymplectic $OSp(4|2)$.
- (2) at the singular point $\zeta = -1$ where $D(2, 1; \zeta)$ contracts to $PSL(2|2) \times \mathbb{C}^3$.

In order to describe geometrical 2-cycles of $D(2, 1; \zeta)$, we considered it useful to start by analysing $SL(2|2)$ since (i) it is a subspace of it; and (ii) it contains two $SL(2)$ submanifolds permitting to make use toric geometry to address $SL(2|2)$ and then $D(2, 1; \zeta)$. In section 2, we showed that $PSL(2|2)$ can be thought of in term of the toric diagram given by the **Figure 6**. In this regard, it would be interesting to extend this construction to $OSp(4|2)$ and $D(2, 1; \zeta)$ which have not been addressed in present study.

Using the correspondence (3.1) and the ansatz (3.2), we studied the four graded homological 2-cycles sets $\mathcal{C}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ in the supermanifold $D(2, 1; \zeta)$ with label $\eta = 0, 1, 2, 3$; see eqs(3.15–3.16). For that, we described the four parametric super Dynkin graphs $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ of complex $\mathfrak{d}(2, 1; \zeta)$ and their discrete outer automorphisms. These diagrams are of two kinds given by (i) the close graph $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(0)}$ with three odd simple roots; and (ii) the 3 open diagrams $S\mathcal{D}\mathcal{D}_{\mathfrak{d}(2,1;\zeta)}^{(i)}$ related by outer automorphisms as shown by the **Figure 9**. By extending to $\mathfrak{d}(2, 1; \zeta)$ the usual correspondence $\alpha \leftrightarrow \mathcal{C}_\alpha$ in ADE geometries linking roots α of ADE Lie algebras with 2-cycle \mathcal{C}_α in asymptotic locally Euclidean (ALE) spaces, we end up with the four parametric sets $\mathcal{C}_{\mathfrak{d}(2,1;\zeta)}^{(\eta)}$ and their relationships with $\mathcal{C}_{osp(4|2)}$ and $\mathcal{C}_{psl(2|2)}$.

By help of these specific features and thinking about $D(2, 1; \zeta)$ as a continuous deviation of $OSp(4|2)$ with deformation parameter ζ , we derived interesting properties on

exceptional symmetry. These specific features allowed us to shed more light on the relationships with $\text{OSp}(4|2)$ and with the 6|8-dim $\text{PSL}(2|2)$. They also made it possible to calculate the Lax operator for the superspin chain by using 4D Chern-Simons (CS) theory with $\text{D}(2,1; \zeta)$ symmetry. In this regard, we computed the Lax operator $\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\mu)}$ having 3 free parameters $\kappa_1(\zeta)$, $\kappa_2(\zeta)$ and $\kappa_0(\zeta)$ which are functions of ζ . It is given by eq(5.55) which recovers as particular values of the distinguished $\mathcal{L}_{sl(2|2)}^{(\mu)}$ and $\mathcal{L}_{osp(4|2)}^{(\mu)}$ by taking κ_i as in (5.56). By setting $\kappa_1 = 1$ and $\kappa_2 = -\kappa_0$, one is left with a simple expression of L-operator having one free function parameter κ_0 and reading as follows

$$z^{-1/2} \begin{pmatrix} (z + \kappa_0^2 b' c') \delta_j^i + \beta^i \gamma_j & c \beta^i - \kappa_0^2 b' \gamma_n \varepsilon^{ni} & \beta^i & \kappa_0 b' \varepsilon^{ji} \\ (b \gamma_j - \kappa_0^2 c' \varepsilon_{jn} \beta^n & z + bc + \kappa_0^2 \beta^n \gamma_n & b & -\kappa_0 \beta^j \\ \gamma_j & c & 1 & 0 \\ \kappa_0 c' \varepsilon_{ji} & -\kappa_0 \gamma_i & 0 & \delta_i^j \end{pmatrix} \quad (6.1)$$

where the pairs (β^i, γ_i) are fermionic oscillators and where (b, c) and (b', c') are bosonic oscillators; they satisfy the usual graded commutation relations. An interesting expression of $\kappa_0(\zeta)$ is given by $\frac{1}{2}\zeta(1 + \zeta)$ interpreted in terms of the determinant $\det(\tilde{\alpha}_i, \tilde{\alpha}_j)$ which is equal $-2\zeta(1 + \zeta)$; is also the area of the 2-cycle $\tilde{\mathcal{C}}_{\tilde{\psi}}$. In terms of the *Kaplansky* parameters S_i with the realisation $(2S_1, 2S_2, 2S_3) = (1, -1 - \zeta, \zeta)$, the value of κ_0 is given by $-4S_1 S_2 S_3$.

From the above Lax matrix, we learn that by setting $\kappa_0 = 1$ (corresponding to $\zeta = 1$), we obtain the L-operator $\mathcal{L}_{osp(4|2)}^{(\mu)}$ given by (5.41). Recall that at $\zeta = 1$, we have the orthosymplectic point $\mathfrak{d}(2, 1; +1) \simeq osp(4|2)$. Similarly, for the case $\kappa_0 = 0$ ($\zeta = -1$), eq(6.1) becomes

$$\mathcal{L}_{\mathfrak{d}(2,1;\zeta)}^{(\mu)} \Big|_{\zeta=-1} = \begin{pmatrix} \mathcal{L}_{sl_2|2}^{\mu} & 0 \\ 0 & z^{-\frac{1}{2}} \delta_i^j \end{pmatrix} \quad (6.2)$$

with $\mathcal{L}_{sl_2|2}^{\mu}$ given by eq(5.23).

In the end, we would like to add that along with the three super Dynkin diagrams $\text{SD}\mathfrak{D}_{\mathfrak{d}(2,1;\zeta)}^{(1)}$, $\text{SD}\mathfrak{D}_{\mathfrak{d}(2,1;\zeta)}^{(2)}$ and $\text{SD}\mathfrak{D}_{\mathfrak{d}(2,1;\zeta)}^{(3)}$ related by \mathcal{S}_3 outer automorphisms, one also has $\text{SD}\mathfrak{D}_{\mathfrak{d}(2,1;\zeta)}^{(0)}$ behaving as a singlet under \mathcal{S}_3 as shown by the **Figure 1** and the **Figure 9**. Here we have studied the integrable superspin chain associated with the open super Dynkin diagrams; it would be interesting to look for the remaining expression of Lax operator associated with the closed $\text{SD}\mathfrak{D}_{\mathfrak{d}(2,1;\zeta)}^{(0)}$.

Appendices

We give four appendices A, B, C and D where we report some technical details completing the analysis given in this paper. In [Appendix A](#), we give useful characteristic data on the algebraic geometry of $SL(2|2)$. In [Appendix B](#), we describe the orthosymplectic $osp(4|2)$ Lie algebra and its realisation using graded oscillators. In [Appendix C](#), we recall interesting aspects on the Scheneurt description of $\mathfrak{d}(2,1;\zeta)$; it uses a realisation of the simple roots $\tilde{\alpha}_i$ different from (3.18). In [Appendix D](#), we provide tools for the study of integrable superspin chains by the help of CS gauge potentials valued in $\mathfrak{d}(2,1;\zeta)$.

A Characteristic data of $SL(2|2)$

In this appendix, we give properties of the complex semi-simple $sl(2|2)$ and its simple subalgebra $psl(2|2)$ used in the main text in link with the parametric super algebra $\mathfrak{d}(2,1;\zeta)$ at $\zeta = -1$. The 14 dimensional $psl(2|2)$ is defined like $psl(2|2) = sl(2|2)/\langle H_0 \rangle$ where $H_0 = \mathbb{C}I_{2|2}$ is the central element of $sl(2|2)$.

Complex superalgebra and real forms

The even part $sl(2|2)_0$ of the Lie super algebra $sl(2|2)$ is 7 dimensional (sup dim 7|8); it given by the $sl(2)_1 \oplus sl(2)_2 \oplus \mathbb{C}$ with matrix generators $J_{b_i}^{a_i} = \sum_{x=1}^3 \mathbf{J}_x (\boldsymbol{\sigma}^x)_{b_i}^{a_i}$ as

symmetry	$sl(2, \mathbb{C})_1$	$sl(2, \mathbb{C})_3$	$\mathbb{C} := \mathbb{C}_3$	
generators	$J_{b_1}^{a_1} := J_b^a$	$R_{b_2}^{a_2} := R_\beta^\alpha$	H_0	(A.1)
action	rotation	rotation	central charge	

obeying the commutation relations

$$\begin{aligned}
 [J_b^a, J_d^c] &= \delta_b^c J_d^a - \delta_d^a J_b^c \\
 [R_\beta^\alpha, R_\delta^\gamma] &= \delta_\beta^\gamma R_\delta^\alpha - \delta_\delta^\alpha R_\beta^\gamma \\
 [H_0, J_b^a] &= [H_0, R_\beta^\alpha] = 0
 \end{aligned}
 \tag{A.2}$$

The odd part $sl(2|2)_1$ has 4+4 fermionic generators Q_a^α and G_α^a transforming in the $(2, 2^*)$ and $(2^*, 2)$ representations of $SL(2) \otimes SL(2)$; they obey the anticommutation relations

$$\begin{aligned}
 \{Q_a^\alpha, G_\beta^b\} &= \delta_\beta^\alpha J_a^b + \delta_a^b R_\beta^\alpha + \delta_a^b \delta_\beta^\alpha H_0 \\
 \{Q_a^\alpha, Q_b^\beta\} &= \varepsilon^{\alpha\beta} \varepsilon_{ab} H_+ \\
 \{G_\alpha^a, G_\beta^b\} &= \varepsilon^{ab} \varepsilon_{\alpha\beta} H_-
 \end{aligned}
 \tag{A.3}$$

where H_+ and H_- are two additional central charges generating two extra directions. Together with the original central charge, they form a triplet $P_i = (H_0, H_\pm)$ having an

interpretation in the 17 dimensional $\mathfrak{d}(2, 1; \zeta)$ at $\zeta = -1$. Because the $SL(2, \mathbb{C})$ group has two real forms given by the compact $SU(2)$ and the non compact $sl(2, \mathbb{R})$; the above relations should be adapted accordingly. For example, eq(A.1) can be interpreted either as

symmetry	$su(2)$	$su(2)$	\mathbb{R}
generators	J_b^a	R_β^α	H_0

(A.4)

with label $a, b = 1, 2, 3$ (first 3-sphere \mathbb{S}_1^3) and $\alpha, \beta = 4, 5, 6$ (second 3-sphere \mathbb{S}_2^3); or like

symmetry	$sl(2, \mathbb{R})$	$su(2)$	\mathbb{R}
generators	L_ν^μ	J_b^a	H_0

(A.5)

with spacetime label $\mu, \nu = 0, 1, 2$ (AdS_3) and internal $a, b = 3, 4, 5$ (3-sphere \mathbb{S}^3). Along with (A.4-A.5), one might also consider the situation with symmetry $sl(2, \mathbb{R}) \oplus sl(2, \mathbb{R}) \oplus \mathbb{R}$ (two pseudo-spheres $\mathbb{S}_i^{1,2} \simeq SU(1, 1)_i$).

Distinguished Cartan Weyl operators

The distinguished root system $\Phi_{sl(2|2)}^{(dis)}$ of the semi-simple Lie super algebra $sl(2|2)$ has 12 roots: 6 positive roots and 6 negative ones. They are generated by three simple roots $(\alpha_1, \alpha_2, \alpha_3)$ with parity as follows

even parity (bosonic)	odd parity (fermionic)
α_1, α_3	$\alpha_2, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$

(A.6)

In the Cartan Weyl basis, the generators of the even part of the Lie super algebra $sl(2|2)_0$ is labeled by two bosonic simple roots α_1 and α_3 as follows

symmetry	$sl(2, \mathbb{C})_A$	$sl(2, \mathbb{C})_B$	\mathbb{C}
generators	$E_{\pm\alpha_1}, H_{\alpha_1}$	$E_{\pm\alpha_3}, H_{\alpha_3}$	H_0

(A.7)

obeying the commutation relations

$$\begin{aligned}
 [E_{+\alpha_i}, E_{-\alpha_i}] &= H_{\alpha_i} \\
 [H_{\alpha_i}, E_{\pm\alpha_j}] &= \pm \mathcal{K}_{ij} E_{\pm\alpha_j}
 \end{aligned}
 \tag{A.8}$$

with super Cartan matrix \mathcal{K}_{ij} . The fermionic operators G_a^α and Q_a^α generating $sl(2|2)_1$ are labeled by the 4 fermionic roots (A.6) which for convenience we set as follows

$$\begin{aligned}
 \gamma_1 &= \alpha_2 & Q_{\pm\gamma_1} &= Q_{\pm\alpha_2} \\
 \gamma_2 &= \alpha_1 + \alpha_2 & Q_{\pm\gamma_2} &= Q_{\pm(\alpha_1 + \alpha_2)} \\
 \gamma_3 &= \alpha_2 + \alpha_3 & Q_{\pm\gamma_3} &= Q_{\pm(\alpha_2 + \alpha_3)} \\
 \gamma_4 &= \alpha_1 + \alpha_2 + \alpha_3 & Q_{\pm\gamma_4} &= Q_{\pm(\alpha_1 + \alpha_2 + \alpha_3)}
 \end{aligned}
 \tag{A.9}$$

The Chevalley generators are given by

symmetry	$sl(2, \mathbb{C})_A$	$sl(1 1)_{AB}$	$sl(2, \mathbb{C})_B$	(A.10)
generators	$E_{\pm\alpha_1}, H_{\alpha_1}$	$E_{\pm\alpha_2}, H_{\alpha_2}$	$E_{\pm\alpha_3}, H_{\alpha_3}$	

obeying the graded commutation relations $[H_{\alpha_i}, E_{\pm\alpha_j}] = \pm\mathcal{K}_{ij}E_{\pm\alpha_j}$ with super Cartan matrix $\mathcal{K} = (\mathcal{K}_{ij})$ factorising like $\mathcal{K} = \mathcal{J}^{diag} \mathcal{J}^{sym}$ with

$$\mathcal{J}^{diag} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad \mathcal{J}^{sym} = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & 1 \\ 0 & 1 & -2 \end{pmatrix}, \quad \mathcal{K} = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & 2 \end{pmatrix} \quad (\text{A.11})$$

where the symmetric part \mathcal{J}^{sym} is given by the intersection matrix $\alpha_i \cdot \alpha_j$. The other generators are given by the graded Serre relations $(Ad_{E_{\pm\alpha_j}})^{n_{jk}}(E_{\pm\alpha_k})$ [?].

Super oscillator realisation of $sl(2|2)$

A realisation of the Chevalley generators is given by 2+2 pairs of graded quantum oscillators (b_i, b_i^\dagger) and (f_i, f_i^\dagger) obeying commutation relations $[b_i, b_j^\dagger] = \delta_{ij}$ and anticommutations $\{f_i, f_j^\dagger\} = \delta_{ij}$. Depending on whether we are considering the complexified $sl(2|2)$ or its real forms $su(2|2)$ or $su(1,1|2)$, we distinguish different realisations. For the example $su(2|2)$, we have the following realisation of its subalgebras $su(2)_A$, $su(1|1)_{AB}$ and $su(2)_B$ respectively labelled by the simple roots α_1 , α_2 and α_3 :

- **Subalgebra $su(2)_A$:** It is contained in $su(2|2)$ and realised in terms of bosonic oscillators as follows

$$E_{+\alpha_1} = b_1^\dagger b_2 \quad , \quad E_{-\alpha_1} = b_2^\dagger b_1 \quad , \quad H_{\alpha_1} = b_1^\dagger b_1 - b_2^\dagger b_2 \quad (\text{A.12})$$

Notice that for the case $sl(2, \mathbb{C})$ contained into $sl(2|2)$, we have instead of (b_i, b_i^\dagger) the complexified oscillators (b_i, c_i) with $[b_i, c_j] = \delta_{ij}$. For this case, the above realisation extends like

$$E_{+\alpha_1} = c_1 b_2 \quad , \quad E_{-\alpha_1} = c_2 b_1 \quad , \quad H_{\alpha_1} = c_1 b_1 - c_2 b_2 \quad (\text{A.13})$$

- **Case of $su(1|1)_{AB}$:** We have

$$E_{+\alpha_2} = b_2^\dagger f_1 \quad , \quad E_{-\alpha_2} = f_1^\dagger b_2 \quad , \quad H_{\alpha_2} = b_2^\dagger b_2 + f_1^\dagger f_1 \quad (\text{A.14})$$

satisfying the cross commutators $[H_{\alpha_1}, E_{+\alpha_2}] = -E_{+\alpha_2}$ and $[H_{\alpha_2}, E_{+\alpha_1}] = -E_{+\alpha_1}$ giving the link with the $su(2)_A$ inside $su(2|2)$. Similar relationships can be written down for $sl(1|1)_{AB}$ by using the bosonic (b_i, c_i) and the fermionic (f_i, g_i) .

• **Subalgebra $\mathfrak{su}(2)_3$** : It is realised in terms of fermionic oscillators like

$$E_{+\alpha_3} = f_1^+ f_2 \quad , \quad E_{-\alpha_3} = f_2^+ f_1 \quad , \quad H_{\alpha_3} = f_1^+ f_1 - f_2^+ f_2 \quad (\text{A.15})$$

with cross commutations $[H_{\alpha_2}, E_{+\alpha_3}] = +E_{+\alpha_3}$ and $[H_{\alpha_3}, E_{+\alpha_2}] = -E_{+\alpha_2}$.

Notice that for $\mathfrak{su}(2|2)$ the total hermitian hamiltonian

$$H_0 = b_1^\dagger b_1 + b_2^\dagger b_2 + f_1^+ f_1 + f_2^+ f_2 \quad (\text{A.16})$$

has the properties: (i) it commutes with the three triplets of Chevalley generators $E_{\pm\alpha_i}$, H_{α_i} (central element). (ii) it is given by the linear combination of the Cartan generators

$$H_0 = H_{\alpha_1} + 2H_{\alpha_2} - H_{\alpha_3} \quad (\text{A.17})$$

Distinguished root diagram of $\mathfrak{psl}(2|2)$

Here, we construct the root diagram of $\mathfrak{psl}(2|2)$. It has 12 roots (6 positive and 6 negative). Its highest weight state $|hws\rangle$ is given by the long root $\boldsymbol{\psi} = \boldsymbol{\alpha}_1 + \boldsymbol{\alpha}_2 + \boldsymbol{\alpha}_3$. The weight diagram relating these roots is depicted by the **Figure 14**. It describes the adjoint representation of $\mathfrak{psl}(2|2)$ with weight vectors as

$$\begin{array}{ll} w_1 = (1, 1, 1) & w_8 = (0, 0, 0) \\ w_2 = (1, 1, 0) & w_9 = (0, 0, -1) \\ w_3 = (0, 1, 1) & w_{10} = (-1, 0, 0) \\ w_4 = (0, 1, 0) & w_{11} = (0, -1, 0) \\ w_5 = (1, 0, 0) & w_{12} = (0, -1, -1) \\ w_6 = (0, 0, 1) & w_{13} = (-1, -1, 0) \\ w_7 = (0, 0, 0) & w_{14} = (-1, -1, -1) \end{array} \quad , \quad (\text{A.18})$$

Starting from the highest weight vector $|\boldsymbol{\psi}\rangle$ satisfying $E_{-\alpha}|\boldsymbol{\psi}\rangle = 0$, one can generate the descendent states by applying the (creation/annihilation) step operators $E_{\pm\alpha}$. For example, the state $|\boldsymbol{\alpha}_1 + \boldsymbol{\alpha}_2\rangle$ is reached by the action

$$E_{\alpha_3}|\boldsymbol{\psi}\rangle \quad (\text{A.19})$$

and the state $|\boldsymbol{\alpha}_2 + \boldsymbol{\alpha}_3\rangle$ is reached as $E_{\alpha_1}|\boldsymbol{\psi}\rangle$ while there no direct way to go from $|\boldsymbol{\psi}\rangle$ towards $|\boldsymbol{\alpha}_2\rangle$ since we have $|\boldsymbol{\alpha}_2\rangle = E_{\alpha_1}E_{\alpha_3}|\boldsymbol{\psi}\rangle$. The full set of transitions can be read from the diagram of the **Figure ??**.




Figure 14: Root diagram of $\mathfrak{psl}(2|2)$. In red the 2+2 even roots of $\mathfrak{psl}(2|2)_0$ and in blue the 4+4 odd ones. From this diagram, we learn (i) its highest (lowest) weight vectors, and (ii) the list of the weight vectors of the adjoint representation of $\mathfrak{psl}(2|2)$.




Figure 15: Four graded Dynkin diagrams (super DD) of the Lie superalgebra $\mathfrak{sl}(2|2)$. The first (top left) has one odd simple root (black node α_2). The fourth graph (bottom right) has three (black) odd roots. The four graphs are associated with the orderings $bbff, fbbf, bffb, bfbf$. Here, we have drawn the intersections $I_{ij}^{(\mu)} = \alpha_i^{(\mu)} \cdot \alpha_j^{(\mu)}$.

Four super Dynkin diagrams

The super Lie algebra $\mathfrak{sl}(2|2)$ has four super Dynkin diagrams that we denote like $\text{SDD}_1, \text{SDD}_2, \text{SDD}_3$ and SDD_0 and which are given by the **Figure 15**. Each $\text{SDD}_\mu^{\mathfrak{sl}(2|2)}$

has three simple roots $\{\alpha_1^{(\mu)}, \alpha_2^{(\mu)}, \alpha_3^{(\mu)}\}$ whose parity depends on the grading of the canonical weight vectors $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4\}$. For convenience, we denote these canonical weight vectors like $\{\mathbf{b}_1, \mathbf{b}_2, \mathbf{f}_1, \mathbf{f}_2\}$; they generate the 2+2 graded space $\mathbb{C}^{2|2}$ with pairings as $\mathbf{b}_i \cdot \mathbf{b}_j = \delta_{ij}$ and $\mathbf{f}_i \cdot \mathbf{f}_j = -\delta_{ij}$ as well as $\mathbf{b}_i \cdot \mathbf{f}_j = 0$. Along with these canonical weight vectors, we can introduce the following "mean weight vector"

$$\epsilon = \frac{1}{4} (\mathbf{b}_1 + \mathbf{b}_2 - \mathbf{f}_1 - \mathbf{f}_2) \quad (\text{A.20})$$

that turns out to play an important role in $\mathfrak{sl}(2|2)$; it labels the central element H_0 .

Four basis sets of simple roots

Because of the intimate relationship between roots α of Lie algebras and 2-cycles \mathcal{C}_α of complex manifolds like the familiar ADE geometries of ALE surfaces [?], we think it interesting to describe explicitly the four sets of simple roots $\{\alpha_1^{(\mu)}, \alpha_2^{(\mu)}, \alpha_3^{(\mu)}\}$ and the associated 2-cycle basis sets of the $\text{SL}(2|2)$ geometry $\{\mathcal{C}_{\alpha_1^{(\mu)}}, \mathcal{C}_{\alpha_2^{(\mu)}}, \mathcal{C}_{\alpha_3^{(\mu)}}\}$. Due to the different orderings of the canonical weight vectors $\{\mathbf{b}_1, \mathbf{b}_2, \mathbf{f}_1, \mathbf{f}_2\}$, the superalgebra $\mathfrak{sl}(2|2)$ has four basis sets of simple roots systems Π_μ labelled by $\mu = 0, 1, 2, 3$;

$$\Pi_0 \quad , \quad \Pi_1 \quad , \quad \Pi_2 \quad , \quad \Pi_3 \quad (\text{A.21})$$

These four sets can be determined from the four possible realisations of the long roots. We denote these four realisations like ψ_1, ψ_2, ψ_3 and ψ_0 . According to the orderings (bbff), (bffb), (fbbf) and (bfbf), we have the following:

- **Ordering (bbff):** long root $\psi_1 = \mathbf{b}_1 - \mathbf{f}_2$

$$\begin{aligned} \mathbf{b}_1 - \mathbf{f}_2 &= (\mathbf{b}_1 - \mathbf{b}_2) + (\mathbf{b}_2 - \mathbf{f}_1) + (\mathbf{f}_1 - \mathbf{f}_2) \\ \psi_1 &= \alpha_1 + \alpha_2 + \alpha_3 \end{aligned} \quad (\text{A.22})$$

from which we read $\alpha_1 = \mathbf{b}_1 - \mathbf{b}_2$ (even) and $\alpha_2 = \mathbf{b}_2 - \mathbf{f}_1$ (odd) as well as $\alpha_3 = \mathbf{f}_1 - \mathbf{f}_2$ (even).

- **Ordering (bffb):** long root $\psi_2 = \mathbf{b}_1 - \mathbf{b}_2$

$$\begin{aligned} \mathbf{b}_1 - \mathbf{b}_2 &= (\mathbf{b}_1 - \mathbf{f}_1) + (\mathbf{f}_1 - \mathbf{f}_2) + (\mathbf{f}_2 + \mathbf{b}_2) \\ \psi_2 &= \alpha'_1 + \alpha'_2 + \alpha'_3 \end{aligned} \quad (\text{A.23})$$

with odd .

- **Ordering (fbbf):** long root $\psi_3 = \mathbf{f}_1 - \mathbf{f}_2$

$$\begin{aligned} \mathbf{f}_1 - \mathbf{f}_2 &= (\mathbf{f}_1 - \mathbf{b}_1) + (\mathbf{b}_1 - \mathbf{b}_2) + (\mathbf{b}_2 - \mathbf{f}_2) \\ \psi_3 &= \alpha''_1 + \alpha''_2 + \alpha''_3 \end{aligned} \quad (\text{A.24})$$

- **Ordering (bfbf)**: long root $\psi_0 = \mathbf{b}_1 - \mathbf{f}_2$

$$\begin{aligned}\mathbf{b}_1 - \mathbf{f}_2 &= (\mathbf{b}_1 - \mathbf{f}_1) + (\mathbf{f}_1 - \mathbf{b}_2) + (\mathbf{b}_2 - \mathbf{f}_2) \\ \psi_0 &= \alpha_1''' + \alpha_2''' + \alpha_3'''\end{aligned}\tag{A.25}$$

So, the four basis sets of the root system $\Phi_{sl(2|2)}$ are as follows

basis sets	α_1	α_2	α_3	odd roots	even roots
$\Pi_1(b, b, f, f)$	$\mathbf{b}_1 - \mathbf{b}_2$	$\mathbf{b}_2 - \mathbf{f}_1$	$\mathbf{f}_1 - \mathbf{f}_2$	α_2	α_1, α_3
$\Pi_2(b, f, f, b)$	$\mathbf{b}_1 - \mathbf{f}_1$	$\mathbf{f}_1 - \mathbf{f}_2$	$\mathbf{f}_2 - \mathbf{b}_2$	α_1, α_3	α_2
$\Pi_3(f, b, b, f)$	$\mathbf{f}_1 - \mathbf{b}_1$	$\mathbf{b}_1 - \mathbf{b}_2$	$\mathbf{b}_2 - \mathbf{f}_2$	α_1, α_3	α_2
$\Pi_0(b, f, b, f)$	$\mathbf{b}_1 - \mathbf{f}_1$	$\mathbf{f}_1 - \mathbf{b}_2$	$\mathbf{b}_2 - \mathbf{f}_2$	$\alpha_1, \alpha_2, \alpha_3$	-

where the Π_2 and Π_3 are distinguished by the different realisation of α_2 (red and blue colors in the **Figure 15**). By using the correspondence algebraic roots/ 2-cycles ($\alpha \leftrightarrow \mathcal{C}_\alpha$), we have the associated 2-cycles basis sets

2-cycles basis sets	odd roots	even roots
Π_1	\mathcal{C}_{α_2}	$\mathcal{C}_{\alpha_1}, \mathcal{C}_{\alpha_3}$
Π_2	$\mathcal{C}_{\alpha_1}, \mathcal{C}_{\alpha_3}$	\mathcal{C}_{α_2}
Π_3	$\mathcal{C}_{\alpha_1}, \mathcal{C}_{\alpha_3}$	\mathcal{C}_{α_2}
Π_0	$\mathcal{C}_{\alpha_1}, \mathcal{C}_{\alpha_2}, \mathcal{C}_{\alpha_3}$	-

Super Dynkin graphs

Because $\Phi_{sl(2|2)}$ is given by $\Phi_{sl(2|2)}^+ \cup \Phi_{sl(2|2)}^-$, the four root systems $\Phi_{sl(2|2)}^{\Pi_\mu}$ of $sl(2|2)$ have each 6 positive and 6 negative graded roots. For $\Phi_{sl(2|2)}^+$, we have

roots	fermionic subset $sl(2 2)_{\bar{1}}$	bosonic subset $sl(2 2)_0$
$\Phi_{sl(2 2)}^{\Pi_1}$	$\alpha_2, \alpha_1 + \alpha_2 + \alpha_3,$ $\alpha_2 + \alpha_1, \alpha_2 + \alpha_3$	α_1, α_3
$\Phi_{sl(2 2)}^{\Pi_2}$	$\alpha_1, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3$	$\alpha_2, \alpha_1 + \alpha_2 + \alpha_3$
$\Phi_{sl(2 2)}^{\Pi_3}$	$\alpha_1, \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3$	$\alpha_2, \alpha_1 + \alpha_2 + \alpha_3$
$\Phi_{sl(2 2)}^{\Pi_0}$	$\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2 + \alpha_3$	$\alpha_1 + \alpha_2, \alpha_2 + \alpha_3$

By using the relations (A.26), one can write down the realisation of the positive roots α of the system $\Phi_{sl(2|2)}^{\Pi_\mu}$ in terms of the canonical weight vectors. For the example of the

graded root system $\Phi_{sl(2|2)}^{\Pi_1}$ with weight vectors ordered like $(\mathbf{b}_1, \mathbf{b}_2, \mathbf{f}_1, \mathbf{f}_2)$, we have

$$\begin{aligned} \alpha_1 &= \mathbf{b}_1 - \mathbf{b}_2 \quad , & \alpha_1 + \alpha_2 &= \mathbf{b}_1 - \mathbf{f}_1 \\ \alpha_2 &= \mathbf{b}_2 - \mathbf{f}_1 \quad , & \alpha_2 + \alpha_3 &= \mathbf{b}_2 - \mathbf{f}_2 \\ \alpha_3 &= \mathbf{f}_1 - \mathbf{f}_2 \quad , & \alpha_1 + \alpha_2 + \alpha_3 &= \mathbf{b}_1 - \mathbf{f}_2 \end{aligned} \quad (\text{A.29})$$

Similar realisations can be written down for the other orderings; see eqs(A.23-A.25). The four super Dynkin diagrams $\text{DD}_\mu^{sl(2|2)}$ associated with the four orderings of the canonical weights are depicted by the **Figure 15**.

B Orthosymplectic $\text{osp}(4|2)$

In this appendix, we give some tools of $\text{osp}(4|2)$ and study its oscillator realisation as well as its link with the exceptional super $\mathfrak{D}(2, 1; \zeta)$ at $\zeta = 1$. Recall that the distinguished super Cartan matrix $\mathcal{K}_{D(2,1;\zeta)}$ and the intersection matrix $\tilde{\alpha}_i \cdot \tilde{\alpha}_j$ with odd $\tilde{\alpha}_2$ given by

$$\mathcal{K}_{D(2,1;\zeta)} = \begin{pmatrix} 2 & -1 & 0 \\ 1 & 0 & \zeta \\ 0 & -1 & 2 \end{pmatrix} \quad , \quad \tilde{\alpha}_i \cdot \tilde{\alpha}_j = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & -\zeta \\ 0 & -\zeta & 2\zeta \end{pmatrix} \quad (\text{B.1})$$

take the following $\text{osp}(4|2)$ values

$$\mathcal{K}_{D(2,1;\zeta)} = \begin{pmatrix} 2 & -1 & 0 \\ 1 & 0 & 1 \\ 0 & -1 & 2 \end{pmatrix} \quad , \quad \tilde{\alpha}_i \cdot \tilde{\alpha}_j = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & -1 \\ 0 & -1 & 2 \end{pmatrix} \quad (\text{B.2})$$

Root system of $\text{osp}(4|2)$

Safety

Here, we use the isomorphism between the triple-sum of isospins $sl(2)_{\beta_1} \oplus sl(2)_{\beta_2} \oplus sl(2)_{\beta_3}$ and the bi-sum $so(4)_{\beta_1, \beta_2} \oplus sp(2)_{\beta_3}$ to realise $\mathfrak{d}(2, 1; \zeta)$ at $\zeta = 1$. The root system $\Phi_{\text{osp}(4|2)}$ has 14 roots; 7 positive and 7 negative. The set of the 7 positive splits into an even subset $\{\beta_1, \beta_2, \beta_3\}$ of three bosonic roots; and an odd subset $\{\gamma_1, \gamma_2, \gamma_3, \gamma_4\}$ of fermionic roots.

- *Super Dynkin diagrams of $\text{osp}(4|2)$*

The structure of the super Dynkin diagrams of $\text{osp}(4|2)$ follow from the parametric super Dynkin diagrams of $\mathfrak{d}(2, 1; \zeta)$ at $\zeta = 1$. These super diagrams are given by the four graphs

of the **Figure 9** with Kaplansky parameters (s_1, s_2, s_3) realised for open graphs as follows

odd root	$2s_1$	$2s_2$	$2s_3$
α_1	-2	1	1
α_2	1	-2	1
α_3	1	1	-2

(B.3)

These super diagrams are given by **the Figure 16**. For the three open diagrams, the




Figure 16: Super Dynkin diagrams of the orthosymplectic Lie superalgebra $osp(4|2)$.

simple roots can be realised in terms of the graded canonical weights ϵ_i and δ as follow

simple root	α_1	α_2	α_3
Fig 9-(a)	$\delta - \epsilon_1$	$\epsilon_1 + \epsilon_2$	$\epsilon_1 - \epsilon_2$
Fig 9-(b)	$\epsilon_1 - \epsilon_2$	$\delta - \epsilon_1$	$\epsilon_1 + \epsilon_2$
Fig 9-(c)	$\epsilon_1 + \epsilon_2$	$\epsilon_1 - \epsilon_2$	$\delta - \epsilon_1$

(B.4)

For the other non simple root, see eq(B.9) below. Regarding the closed super Dynkin diagram, the three simple roots are $(\alpha_1, \alpha_2, \alpha_3)$ fermionic; they can be realised as follows

simple root	α_1	α_2	α_3
Fig 9-(d)	$\delta - \epsilon_2$	$\epsilon_1 - \delta$	$\delta + \epsilon_2$

(B.5)

together with

$$\begin{aligned}
 \alpha_1 + \alpha_2 &= \epsilon_1 - \epsilon_2 & \alpha_1 + \alpha_2 + \alpha_3 &= \delta + \epsilon_1 \\
 \alpha_2 + \alpha_3 &= \epsilon_1 + \epsilon_2 & \alpha_1 + 2\alpha_2 + \alpha_3 &= 2\epsilon_1
 \end{aligned}$$
(B.6)

- *Distinguished basis set $(\alpha_1, \alpha_2, \alpha_3)$ with α_2 odd.*

For the distinguished super Dynkin diagram with one odd root taken as α_2 (i.e: $\alpha_2^2 = 0$) and the other two (α_1, α_3) even, we have the following realisation of the roots of $\Phi_{osp(4|2)}$

in terms of three canonical weight vectors even ϵ_i ($\epsilon_1^2 = \epsilon_2^2 = 1$) and odd δ ($\delta^2 = -1$) [44, 45]. First, the simple roots are realised like $\alpha_1 = \epsilon_1 - \epsilon_2$ and $\alpha_2 = \delta - \epsilon_1$ as well as $\alpha_3 = \epsilon_1 + \epsilon_2$ with the automorphism symmetry \mathbb{Z}_2 acting like

$$\mathbb{Z}_2 : \begin{array}{ll} (\epsilon_1, \delta, \epsilon_2) & \rightarrow (\epsilon_1, \delta, -\epsilon_2) \\ (\alpha_1, \alpha_2, \alpha_3) & \rightarrow (\alpha_3, \alpha_2, \alpha_1) \end{array} \quad (\text{B.7})$$

and intersection $\alpha_i \cdot \alpha_j$ as follows

$$\alpha_i \cdot \alpha_j = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & -1 \\ 0 & -1 & 2 \end{pmatrix}, \quad \tilde{\alpha}_i \cdot \tilde{\alpha}_j|_{\zeta=1} = \alpha_i \cdot \alpha_j \quad (\text{B.8})$$

corresponding to setting $\zeta = 1$ in eq(B.1). The 4 other composite positive roots are given by

$$\begin{array}{ll} \alpha_1 + \alpha_2 = \delta - \epsilon_2 & \alpha_1 + \alpha_2 + \alpha_3 = \delta + \epsilon_1 \\ \alpha_2 + \alpha_3 = \delta + \epsilon_2 & \alpha_1 + 2\alpha_2 + \alpha_3 = 2\delta \end{array}, \quad (\text{B.9})$$

from which (i) we read the three even roots β_i of $sl(2)_{\beta_1} \oplus sl(2)_{\beta_2} \oplus sl(2)_{\beta_3}$ namely $\beta_1 = \alpha_1$ and $\beta_2 = \alpha_3$ and $\beta_3 = 2\delta$ (three even 2-cycles); and (ii) the four odd roots $\gamma_1 = \alpha_2$ and $\gamma_2 = \alpha_1 + \alpha_2$ as well as $\gamma_3 = \alpha_2 + \alpha_3$ and $\gamma_4 = \alpha_1 + \alpha_2 + \alpha_3$.

Three pairs of oscillators for $\text{osp}(4|2)$

The super $\text{osp}(4|2)$ splits as $\text{osp}(4|2)_0 \oplus \text{osp}(4|2)_1$ with $\text{osp}(4|2)_0 = so(4) \oplus sp(2)$. The oscillator realisation of the $so(4, \mathbb{C})$ generators is achieved by two pairs of fermionic oscillators (f_1, g_1) , (f_2, g_2) satisfying the anticommutation relations $\{f_i, g_j\} = \delta_{ij}$ and $\{f_i, f_j\} = \{g_i, g_j\} = 0$. For a unitary theory, the g_i 's are identified with f_i^\dagger . For simplicity, we consider the particular oscillator realisation with (f_1^\dagger, f_1) , (f_2^\dagger, f_2) satisfying the anticommutation relations $\{f_i, f_j^\dagger\} = \delta_{ij}$ in terms of which the 6 generators of $so(4, \mathbb{R}) \simeq su(2) \oplus su(2)$ are realised as follows

$$\begin{array}{ll} E_{+\beta_1} = f_1^\dagger f_2 & E_{+\beta_2} = f_1^\dagger f_2^\dagger \\ E_{-\beta_1} = f_2^\dagger f_1 & E_{-\beta_2} = f_2 f_1 \\ H_{\beta_1} = f_1^\dagger f_1 - f_2^\dagger f_2 & H_{\beta_2} = f_1^\dagger f_1 + f_2^\dagger f_2 - 1 \end{array}, \quad (\text{B.10})$$

Notice that this oscillator realisation can be motivated by the following expressions of the simple roots in terms of the canonical weight vectors

$$\pm\beta_1 = \pm(\epsilon_1 - \epsilon_2), \quad \pm\beta_2 = \pm(\epsilon_1 + \epsilon_2) \quad (\text{B.11})$$

Similarly, the realisation of the symplectic $sp(2, \mathbb{R})$ generators is achieved by a bosonic oscillator b, b^\dagger satisfying $[b, b^\dagger] = 1$ and $[b, b] = 0$. Here, the three generators are given by

$$\begin{aligned} E_{+\beta_3} &= -\frac{1}{2}b^\dagger b^\dagger \\ E_{-\beta_3} &= +\frac{1}{2}bb \\ H_{\beta_3} &= b^\dagger b + \frac{1}{2} \end{aligned} \tag{B.12}$$

Here also, this realisation can be motivated from the realisation $\beta_3 = 2\delta$.

The 4+4 odd generators of $osp(4|2, \mathbb{R})_{\bar{1}}$ are given by

$$\begin{aligned} E_{+\gamma_1} &= b^\dagger f_1 & E_{+\gamma_3} &= b^\dagger f_2^\dagger \\ E_{+\gamma_2} &= b^\dagger f_2 & E_{+\gamma_4} &= b^\dagger f_1^\dagger \\ E_{-\gamma_1} &= f_1^\dagger b & E_{-\gamma_3} &= f_2 b \\ E_{-\gamma_2} &= f_2^\dagger b & E_{-\gamma_4} &= f_1 b \end{aligned} \tag{B.13}$$

From these relations we calculate the anticommutator $\{E_{+\gamma_1}, E_{-\gamma_1}\}$ of the odd Chevalley generator giving $H_{\gamma_1} = b^\dagger b + f_1^\dagger f_1$.

C Scheneurt realisation of $\mathfrak{d}(2,1;\zeta)$

In this appendix, we give the Scheneurt method for realising the roots of the super Lie algebra $\mathfrak{d}(2,1;\zeta)$ in terms of weight vectors in \mathbb{C}^3 .

Scheneurt construction

Following [44, 51], the roots of the system $\tilde{\Phi}_{\mathfrak{d}(2,1;\zeta)}$ can be realised by using three complex canonical weight vectors $(\epsilon_1, \epsilon_2, \epsilon_3)$ generating \mathbb{C}^3 . These complex weight vectors have (non hermitian) pairing products as $\epsilon_i \cdot \epsilon_j = S_i \delta_{ij}$ in which the S_i 's are non vanishing complex numbers ($S_i \in \mathbb{C}^*$). The properties of these weight vectors go beyond the canonical values ± 1 used in $sl(2|2)$. So, non zero pairings are given by

$$\epsilon_1 \cdot \epsilon_1 = S_1 \quad , \quad \epsilon_2 \cdot \epsilon_2 = S_2 \quad , \quad \epsilon_3 \cdot \epsilon_3 = S_3 \tag{C.1}$$

In the relations the S_i 's are not completely free; these are three complex numbers $S_i = |S_i| e^{i\theta_i}$ such as $S_1 + S_2 + S_3 = 0$ and $S_1 \times S_2 \times S_3 \neq 0$. These conditions are required by the graded Jacobi identity in $\mathfrak{d}(2,1;\zeta)$ [49–51, 85]; a feature nicely exhibited on the **Figure 9**. In this regard, recall that $\mathfrak{d}(2,1;\zeta)$ is isomorphic to the simple Lie superalgebra $\Gamma(S_1, S_2, S_3)$ [49, 51]. Notice also that the above constraint relations can be solved in

several ways; an interesting way is given by restricting to $s_3/s_1 = \zeta$ and $s_2/s_1 = -1 - \zeta$ with complex $\zeta \neq 0, -1$ due to $s_1 s_2 s_3 \neq 0$. With this choice, we have

$$\frac{s_1 s_2 s_3}{s_1^3} = -\zeta(1 + \zeta) \quad (\text{C.2})$$

having zeros at $\zeta = 0, -1$. It breaks the \mathcal{S}_3 symmetry down to its subgroup \mathcal{Z}_2 generated by the transposition $\zeta \rightarrow -1 - \zeta$; it has a fix point at $\zeta = -1/2$ ($s_2 = s_3$) and exchanges the two singularities $\zeta = 0, -1$ as depicted by the **Figure ??**. Using the weight vectors $(\varepsilon_1, \varepsilon_2, \varepsilon_3)$, the simple roots $(\tilde{\alpha}_1, \tilde{\alpha}_2, \tilde{\alpha}_3)$ generating the basis set $\Pi_{\mathfrak{d}(2,1;\zeta)}^2$ are realised as follows

$$\begin{aligned} \tilde{\alpha}_1 &= 2\varepsilon_1 & \tilde{\alpha}_1 \cdot \tilde{\alpha}_1 &= 4s_1 \\ \tilde{\alpha}_2 &= \varepsilon_2 - \varepsilon_1 - \varepsilon_3 & \tilde{\alpha}_2 \cdot \tilde{\alpha}_2 &= 0 \\ \tilde{\alpha}_3 &= 2\varepsilon_3 & \tilde{\alpha}_3 \cdot \tilde{\alpha}_3 &= 4s_3 \end{aligned} \quad (\text{C.3})$$

with the properties

$$\begin{aligned} \tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3 &= \varepsilon & \varepsilon &= \varepsilon_1 + \varepsilon_2 + \varepsilon_3 \\ \tilde{\alpha}_1^2 + \tilde{\alpha}_2^2 + \tilde{\alpha}_3^2 &= -4s_2 & \varepsilon^2 &= 0 \end{aligned} \quad (\text{C.4})$$

The intersection matrix $\tilde{\mathcal{I}}_{ij}$ associated with the above realisation (C.3) reads as

$$\tilde{\mathcal{I}}_{ij}(s) = \begin{pmatrix} 4s_1 & -2s_1 & 0 \\ -2s_1 & 0 & -2s_3 \\ 0 & -2s_3 & 4s_3 \end{pmatrix}, \quad \tilde{\mathcal{I}}_{ij}(\zeta) = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & -\zeta \\ 0 & -\zeta & 2\zeta \end{pmatrix} \quad (\text{C.5})$$

with the property $\sum_{j=1}^3 \tilde{\mathcal{I}}_{ij}(s) = 2s_i$. By setting $2s_1 = 1$ and $2s_2 = -1 - \zeta$ as well as $2s_3 = \zeta$, the intersection $\tilde{\mathcal{I}}_{ij}(\zeta)$ with determinant $\det \tilde{\mathcal{I}}_{ij} = -2\zeta(\zeta + 1)$. The super Cartan $\tilde{\mathcal{K}}(\zeta)$ is given by $\tilde{\mathcal{J}} \cdot \tilde{\mathcal{I}}(\zeta)$ with [44, 45]

$$\tilde{\mathcal{J}} = \frac{-1}{1 + \zeta} \begin{pmatrix} \frac{4+\zeta}{2} & 3 & \frac{3}{2} \\ \frac{2+\zeta}{2} & 2 + \zeta & \frac{1}{2} \\ \frac{2+\zeta}{2} & 1 & -\frac{2+\zeta}{2\zeta} \end{pmatrix}, \quad \tilde{\mathcal{K}}_{ij} = \begin{pmatrix} 2 & -1 & 0 \\ 1 & 0 & \zeta \\ 0 & -1 & 2 \end{pmatrix} \quad (\text{C.6})$$

Comparing these matrices of $\mathfrak{d}(2, 1; \zeta)$ with their $\mathfrak{sl}(2|2)$ homologue given by the relations (??), we see that they can be recovered from the above (C.3) by setting $\zeta = -1$. For the two singular points $\zeta = -1$, we have critical matrices

$$\zeta = -1: \quad \tilde{\mathcal{I}}_{ij} = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & 1 \\ 0 & 1 & -2 \end{pmatrix} \quad (\text{C.7})$$

Roots system in terms of weight vectors

Here, we give the expression of the 7 positive roots $\tilde{\alpha}$ in terms of the canonical weight vectors

$$\begin{aligned}
\tilde{\alpha}_1 &= 2\varepsilon_1 \\
\tilde{\alpha}_2 &= \varepsilon_2 - \varepsilon_1 - \varepsilon_3 \\
\tilde{\alpha}_3 &= 2\varepsilon_3 \\
\tilde{\alpha}_1 + \tilde{\alpha}_2 &= \varepsilon_1 + \varepsilon_2 - \varepsilon_3 \\
\tilde{\alpha}_2 + \tilde{\alpha}_3 &= \varepsilon_2 - \varepsilon_1 + \varepsilon_3 \\
\tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3 &= \varepsilon_1 + \varepsilon_2 + \varepsilon_3 \\
\tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3 &= 2\varepsilon_2
\end{aligned} \tag{C.8}$$

Notice that using these weight vectors, the long root of $\mathfrak{d}(2,1;\zeta)$ reads as

$$\tilde{\psi} = \tilde{\alpha}_1 + 2\tilde{\alpha}_2 + \tilde{\alpha}_3 \tag{C.9}$$

it can be split in two ways like the sum of odd roots; either as $\tilde{\alpha}_2 + (\tilde{\alpha}_1 + \tilde{\alpha}_2 + \tilde{\alpha}_3)$; or like $(\tilde{\alpha}_1 + \tilde{\alpha}_2) + (\tilde{\alpha}_2 + \tilde{\alpha}_3)$. Using eq(C.9), we can express the odd root $\tilde{\alpha}_2$ in terms of halves of the bosonic $\tilde{\alpha}_1$, $\tilde{\alpha}_3$ and $\tilde{\psi}$ like

$$\tilde{\alpha}_2 = \frac{1}{2}\tilde{\psi} - \frac{1}{2}\tilde{\alpha}_1 - \frac{1}{2}\tilde{\alpha}_3 \tag{C.10}$$

So, instead of (C.8), we can present the roots into representations of \mathcal{S}_3 . For the 6 even roots, we have the triplet

$$\begin{aligned}
\pm\tilde{\alpha}_1 &= \pm 2\varepsilon_1 & \tilde{\alpha}_1^2 &= 4S_1 \\
\pm\tilde{\psi} &= \pm 2\varepsilon_2 & \tilde{\psi}^2 &= 4S_2 \\
\pm\tilde{\alpha}_3 &= \pm 2\varepsilon_3 & \tilde{\alpha}_3^2 &= 4S_3
\end{aligned} \tag{C.11}$$

For the 8 odd roots, they split into two singlets $\pm\tilde{\gamma}_0$ with

$$\begin{aligned}
\tilde{\gamma}_0 &= \varepsilon_2 + \varepsilon_1 + \varepsilon_3 \\
&= \frac{1}{2}\tilde{\alpha}_1 + \frac{1}{2}\tilde{\psi} + \frac{1}{2}\tilde{\alpha}_3
\end{aligned} \tag{C.12}$$

and two triplets $\pm\tilde{\gamma}_i$ with

$$\begin{aligned}
\tilde{\gamma}_1 &= +\frac{1}{2}\tilde{\alpha}_1 - \frac{1}{2}\tilde{\psi} - \frac{1}{2}\tilde{\alpha}_3 = +\varepsilon_1 - \varepsilon_2 - \varepsilon_3 \\
\tilde{\gamma}_2 &= -\frac{1}{2}\tilde{\alpha}_1 - \frac{1}{2}\tilde{\psi} + \frac{1}{2}\tilde{\alpha}_3 = -\varepsilon_1 - \varepsilon_2 + \varepsilon_3 \\
\tilde{\gamma}_3 &= -\frac{1}{2}\tilde{\alpha}_1 + \frac{1}{2}\tilde{\psi} - \frac{1}{2}\tilde{\alpha}_3 = -\varepsilon_1 + \varepsilon_2 - \varepsilon_3
\end{aligned} \tag{C.13}$$

D Integrable superspin chains in CS theory

In this appendix, we provide tools for the study of integrable superspin chains [24, 25] by the help of CS gauge potentials [38, 39]. For concreteness, we focus on the integrable $SL(2|2)$ superspin chain as it is one of the systems considered in this study. For more technical details and other applications, we report to [46] and refs therein.

$SL(2|2)$ potentials in 4D CS theory

Safety

The Lie superalgebra $sl(2|2)$ of the Lie supergroup $SL(2|2)$ over the complex field \mathbb{C} has rank 3 and 15 dimensions. It belongs to the sub-family $sl(n|m)$ with $m = n \geq 2$ having a central charge generator. Apart from the very special $sl(1|1)$ member, the super $sl(2|2)$ constitutes the leading symmetry of superspin chains described by 4D Chern-Simons 1-form potential $A(x)$ with expansion

$$A(x) = \mathfrak{A}_0(x) H_0 + \sum_{\text{even root } \beta} [A_\beta^0(x) H_\beta + A_\beta^\pm(x) E_{\pm\beta}] + \sum_{\text{odd root } \gamma} A_\gamma^\mp(x) F_{\pm\gamma} \quad (\text{D.1})$$

where the $H_0, H_\beta, E_{\pm\beta}$ and $F_{\pm\gamma}$ are matrix generators and where $\mathfrak{A}_0(x), A_\beta^0(x), A_\beta^\pm(x)$ and $A_\gamma^\mp(x)$ are (partial) 1-form potentials along the 15 directions of $sl(2|2)$. The above 4D CS gauge field $A(x)$ has a central potential

$$\mathfrak{A}_0(x) H_0$$

which is absent for the bigger family of $sl(n|m)$ superspin chains with $n \neq m$. Recall that $sl(n|m)$ contains two main sub-families labeled by $n \neq m$ and $n = m$. The dynamics of $A(x)$ is described by the gauge field action

$$S_{4dCS} = \int_{\Sigma \times \mathcal{C}} dz \wedge \text{tr}(A \wedge dA + \frac{2}{3} A \wedge A \wedge A) \quad (\text{D.2})$$

where Σ is a topological real plane (roughly \mathbb{R}^2) parametrised by real variables (x, y) ; and a holomorphic curve \mathcal{C} (say \mathbb{CP}^1) labeled by the complex coordinate z (rapidity) [38, 39, 41]. Notice that the gauge potential $A(x)$ is a partial 1-form potential; it is given by $A_x dx + A_y dy + A_{\bar{z}} d\bar{z}$ where we have dropped the component $A_z dz$ because of the dz factor in the holomorphic volume form measure in (D.2).

The field equation of motion following from the S_{4dCS} is given by the zero of the 2-form curvature; i.e:

$$F = dA + A \wedge A = 0 \quad (\text{D.3})$$

indicating that observables constructed out of it are trivial.

Line defects and RLL equation

Despite that the 2-form field strength $F = 0$, there are still observables describing physical entities. An interesting example is given by super Wilson loops $W_{\xi_z}^{2|2}$ defined by the supertrace of the holonomy of the gauge field A along a loop ξ_z in the 4D space $\mathbb{R}^2 \times \mathbb{CP}^1$ as follows [77, 86, 87]

$$W_{\xi_z}^{2|2} = \text{str}_{2|2} \left[P \exp \left(\oint_{\xi_z} A \right) \right] \quad (\text{D.4})$$

It plays a crucial role in dealing with (super) Yang Baxter equation of 2D (super) integrable models. Another interesting example is given by the super 't Hooft lines $\mathcal{L}^\mu(z)$ carrying a minuscule coweight charge μ ; it is defined as [88]-[91]

$$\mathcal{L}^\mu(z) = P \exp \left(\int_y A_y(z) \right) \quad (\text{D.5})$$

where the transport of the gauge fields $A_y(z)$ is measured from $y < 0$ to $y > 0$. The couplings of these topological super line defects is given by line's crossings and are effectively described by the Yang Baxter R-matrix and the so called Lax operator [41, 54, 92]. A typical example of such coupling is given by the $\mathfrak{sl}(2|2)$ superchain depicted by the **Figure 17**, where the 't Hooft line plays the role of a transfer matrix modeling the




Figure 17: Realization of an $\mathfrak{sl}(2|2)$ superspin chain of L nodes in the fundamental representation. In the 4D Chern-Simons theory, it is described by L vertical super Wilson lines (in blue) crossing a horizontal 't Hooft line (in red).

interactions between the electrically charged super atoms along the superchain. In this realisation, each intersection of an electric $W_{\xi_z}^{2|2}$, carrying a vector quantum space $2|2$, with the magnetic $\text{tH}_{\gamma_0}^\mu$ carrying an auxiliary (oscillator) space \mathbb{A} , yields the super Lax operator for the corresponding node of the superspin chain. This coupling operator acts on the tensor product of $\text{End}(2|2)$ of the Wilson and the algebra \mathbb{A} of functions in the phase space of the 't Hooft line. As for the Yang-Baxter equation, the crossing of Wilson and 't Hooft lines verifies the RLL equation of integrability namely

$$\mathbf{R}_{rs}^{ik}(z-w) \mathbf{L}_j^r(z) \mathbf{L}_l^s(w) = \mathbf{L}_r^i(w) \mathbf{L}_s^k(z) \mathbf{R}_{jl}^{rs}(z-w) \quad (\text{D.6})$$

In this relation, the $\mathbf{L}_n^m(z)$ is the matrix realization of the L-operator and $\mathbf{R}_{rs}^{ik}(z-w)$ is the usual R-matrix solving the Yang-Baxter equation. This RLL equation is represented by the graph of the **Figure 18**. For further details and applications, we refer to [38, 39]




Figure 18: Graphic representation of the RLL equation in terms of intersecting line defects in $SL(2|2)$ 4D CS theory.

for the general set up of the 4D Chern-Simons (CS) theory. Also, we refer to [34, 46] for explicit field realisations, examples of superspin chains and detailed calculations. For applications in strings, see for instance [34, 78] and [93–96].

Concerning the algebraic structure of the 1-form graded potential $A(x)$ given by eq(D.1), it has two blocs $A^{(even)}(x)$ and $A^{(odd)}(x)$. In this regard, recall that the even part $\mathfrak{sl}(2|2)_0$ of the superalgebra $\mathfrak{sl}(2|2)$ is given by $\mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus \mathbb{C}I_{id}$ where $\mathfrak{sl}(2)$ is the usual complex $\mathfrak{sl}(2, \mathbb{C})$ and the block $\mathbb{C}I_{id}$ is a central element spanned by the identity operator I_{id} commuting with all other generators; $[I_{id}, \mathfrak{sl}(2)] = 0$. The $\mathfrak{sl}(2|2)_0$ is 7 dimensional ($7=3+3+1$). The odd part $\mathfrak{sl}(2|2)_1$ is 8 dimensional generated by fermionic charges $F_{\pm\gamma}$. The CS potentials associated with these even and odd parts are given by

$$A^{(even)} = \mathfrak{a}_0(x) H_0 + \sum_{\text{even root } \beta} (A_{\beta}(x) H_{\beta} + A_{\mp\beta}(x) E_{\pm\beta}) \quad (\text{D.7})$$

and

$$A^{(odd)} = \sum_{\text{odd root } \gamma} A_{\mp\gamma}(x) F_{\pm\gamma} \quad (\text{D.8})$$

Their topological 2-form field strengths are

$$\begin{aligned} F^{(even)} &= dA^{(even)} + A^{(even)} \wedge A^{(even)} + A^{(odd)} \wedge A^{(odd)} \\ F^{(odd)} &= dA^{(odd)} + A^{(odd)} \wedge A^{(even)} + A^{(even)} \wedge A^{(odd)} \end{aligned} \quad (\text{D.9})$$

they vanish on shell. The explicit expression of the Lax operator $\mathbf{L}_j^r(z)$ for the open super Dynkin diagram of the superspin chain $\mathfrak{d}(2,1;\zeta)$ solving (D.6) is given by eq(5.55).

References

- [1] V.G. Kac, *Lie superalgebras*, Advances in Math. 26 (1977) 8-96 MR0486011.
- [2] I. Heckenberger, F. Spill, A. Torrielli, H. Yamane, *Drinfeld second realization of the quantum affine superalgebras of $D(2, 1; x)^{(1)}$ via the Weyl groupoid*, Publ.Res.Inst.Math.Sci.Kyoto B8:171,2008, arXiv:0705.1071 [math.QA].
- [3] E. Poletaeva, *Embedding of the Lie superalgebra $D(2, 1; \alpha)$ into the Lie superalgebra of pseudodifferential symbols on $S^{1|2}$* , J. Math. Phys. 48 (2007) 103504, 17 pp. e-print arXiv:0709.0083.
- [4] J. de Boer, A. Pasquinucci and K. Skenderis, “*AdS/CFT dualities involving large 2D $N=4$ superconformal symmetry*,” Adv. Theor. Math. Phys. 3 (1999) 577 [arXiv:hep-th/9904073].
- [5] S. Gukov, E. Martinec, G.W. Moore and A. Strominger, “*The Search for a holographic dual to $AdS_3 \times S^3 \times S^3 \times S^1$* ,” Adv. Theor. Math. Phys. 9 (2005) 435 [hep-th/0403090].
- [6] Lorenz Eberhardt, Matthias R. Gaberdiela and Wei Li, *A holographic dual for string theory on $AdS_3 \times S^3 \times S^3 \times S^1$* , J. High Energ. Phys. 2017, 111 (2017), arXiv:1707.02705 [hep-th].
- [7] Niklas Beisert, Reimar Hecht, Ben Hoare, *Maximally extended $sl(2|2)$, q -deformed $d(2, 1; \varepsilon)$ and 3D kappa-Poincaré*, J. Phys. A. 50, 314003 (2017), arXiv:1704.05093v2 [math-ph].
- [8] N. Beisert and M. de Leeuw, *The RTT realization for the deformed $gl(2|2)$ Yangian*, J. Phys.A: Math. Theor. 47, 305201 (2014), doi:10.1088/1751-8113/47/30/305201.
- [9] N. Beisert, M. de Leeuw and R. Hecht, *Maximally extended $sl(2|2)$ as a quantum double*, J. Phys. A: Math. Theor. 49, 434005 (2016), doi:10.1088/1751-8113/49/43/434005.
- [10] M. Parker, “*Classification of real simple Lie superalgebras of classical type*,” J. Math. Phys., vol. 21, no. 4, pp. 689-697, 1980.
- [11] Antonio Pittelli, *Yangian Symmetry of String Theory on $AdS_3 \times S^3 \times S^3 \times S^1$ with Mixed 3-form Flux*, Nuclear Physics B 935, 2018, 271-289, arXiv:1711.02468v2 [hep-th].
- [12] I. Bars, C. Deliduman, D. Minic, *String Theory on AdS_3 Revisited*, arXiv:hep-th/9907087v2 .
- [13] Constantin Bachas, Eric D’Hoker, John Estes, Darya Krym, *M-theory Solutions Invariant under $D(2, 1; \gamma) \oplus D(2, 1; \gamma)$* , <https://doi.org/10.1002/prop.201300039>, arXiv:1312.5477v2 [hep-th].

- [14] Lorenz Eberhardt, Matthias R. Gaberdiel, *Strings on $AdS_3 \times S^3 \times S^3 \times S^1$* . J. High Energ. Phys. 2019, 35 (2019)., arXiv:1904.01585v1 [hep-th].
- [15] G. Giribet, C. Hull, M. Kleban, M. Porrati, E. Rabinovici, *Superstrings on AdS_3 at $k=1$* , JHEP08(2018)204, arXiv:1803.04420v3 [hep-th].
- [16] Lorenz Eberhardt, Matthias R. Gaberdiel, *String theory on AdS_3 and the symmetric orbifold of Liouville theory*, Nuclear Physics B 948, 2019, 114774, arXiv:1903.00421v1 [hep-th].
- [17] Gaston Giribet, *String correlators in AdS_3 from FZZ duality*, JHEP12(2021)012, arXiv:2110.04197v2 [hep-th].
- [18] Lorenz Eberhardt, Matthias R. Gaberdiel, Rajesh Gopakumar, *Deriving the AdS_3/CFT_2 Correspondence*, J. High Energ. Phys. 2020, 136 (2020), arXiv:1911.00378v2 [hep-th].
- [19] Sammani, R., Boujakhrou, Y., Saidi, E. H., Laamara, R. A., & Drissi, L. B. (2023). *Higher spin AdS_3 gravity and Tits-Satake diagrams*. Physical Review D, 108(10), 106019.
- [20] R. Sammani, E. H Saidi, *Black Flowers and Real Forms of Higher Spin Symmetries*, JHEP (2024), arXiv:2406.01328v1 [hep-th].
Finiteness of 3D higher spin gravity Landscape, Classical and Quantum Gravity 2024, DOI 10.1088/1361-6382/ad7cba
- [21] Niklas Beisert, Egor Im, *Classical Lie Bialgebras for AdS/CFT Integrability by Contraction and Reduction*, SciPost Phys. 14, 157 (2023), arXiv:2210.11150v2 [hep-th].
- [22] Alessandro Torrielli, *Review of AdS/CFT Integrability, Chapter VI.2: Yangian Algebra*, Letters in Mathematical Physics 99(1), arXiv:1012.4005v5 [hep-th].
- [23] Andrea Prinsloo, Vidas Regelskis, Alessandro Torrielli, *Integrable open spin-chains in AdS_3/CFT_2 correspondences*, Phys. Rev. D 92, 106006 (2015), arXiv:1505.06767v3 [hep-th].
- [24] N.I. Stoilova and J. Van der Jeugt, *A classification of generalized quantum statistics associated with basic classical Lie superalgebras*, J. Math. Phys. 46 (2005) 113504[math-ph/0504013].
- [25] N.I. Stoilova and J. Van der Jeugt, *A classification of generalized quantum statistics associated with the exceptional Lie (super)algebras*, J. Math. Phys. 48 (2007) 043504 [math-ph/0611085].
- [26] Andrea Fontanella, Alessandro Torrielli, *Massless sector of AdS_3 superstrings: a geometric interpretation*, Phys. Rev. D 94, 066008 (2016), arXiv:1608.01631v3 [hep-th].

- [27] R. Frassek, A. Tsymbaliuk, Rational Lax matrices from antidominantly shifted extended Yangians: BCD types, arXiv :2104 .14518[math .RT].
- [28] V.V. Bazhanov, R. Frassek, T.L. ukowski, C. Meneghelli, M. Staudacher, Baxter Q-operators and representations of Yangians, Nucl. Phys. B 850 (2011) 148, arXiv :1010 .3699.
- [29] R. Frassek, Oscillator realisations associated to the D-type Yangian: towards the operatorial Q-system of orthogonal spin chains, Nuclear Phys. B 956 (2020) 115063, 22 [2001.06825].
- [30] G. Ferrando, R. Frassek, V. Kazakov, QQ-system and Weyl-type transfer matrices in integrable $SO(2r)$ spin chains, 2008.04336.
- [31] Y. Boujakhrou, E.H. Saidi, R.A. Laamara and L.B. Drissi, 't Hooft lines of ADE-type and topological quivers, SciPost Phys. 15 (2023) 078 [arXiv:2303.13879].
- [32] Youssra Boujakhrou, El Hassan Saidi, *Minuscul ABCDE Lax Operators from 4D Chern-Simons Theory*, Nucl.Phys.B 981 (2022) 115859, arXiv:2207.14777 [hep-th]
- [33] Katsushi Ito, Mingshuo Zhu, *ODE/IM correspondence and supersymmetric affine Toda field equations*, Nucl.Phys B985, December 2022, 116004, arXiv:2206.08024v2 [hep-th].
- [34] N. Ishtiaque, S.F. Moosavian, S. Raghavendran and J. Yagi, Superspin chains from superstring theory, SciPost Phys. 13 (2022) 083 [arXiv:2110.15112].
- [35] Niklas Beisert, Matthias Staudacher, *The $N=4$ SYM Integrable Super Spin Chain*, Nuclear Physics B670, 2003, p 439-463, arXiv:hep-th/0307042v3.
- [36] Daniel Arnaudon, Jean Avan, Nicolas Crampé, Anastasia Doikou, Luc Frappat, Eric Ragoucy, *Bethe Ansatz equations and exact S matrices for the $osp(M|2n)$ open super spin chain, bethe ansatz an algeb methods*, Nucl.Phys. B687 (2004) 257-278, arXiv:math-ph/0310042v2.
- [37] N. Beisert, The $su(2|2)$ dynamic S-matrix, Adv. Theor. Math. Phys. 12 (2008) 945–979. e-print arXiv:hep-th/0511082.
- [38] K. Costello, E. Witten and M. Yamazaki, *Gauge theory and integrability*, I, ICCM Not. 6 (2018) 46, arXiv:1709.09993 [hep-th].
- [39] K. Costello, E. Witten and M. Yamazaki, *Gauge theory and integrability*, II, ICCM Not. 6 (2018) 120 arXiv:1802.01579 [hep-th].
- [40] K. Costello, M. Yamazaki, *Gauge Theory And Integrability*, III, 1908.02289.
- [41] E.H. Saidi, *Quantum line operators from Lax pairs*, J. Math. Phys. 61 (2020) 063501, arXiv :1812 .06701[hep -th].

- [42] Kevin Costello, Bogdan Stefański jr, *The Chern-Simons Origin of Superstring Integrability*, Phys. Rev. Lett. 125, 121602 (2020), arXiv:2005.03064v3 [hep-th].
- [43] Edward Witten, Integrable Lattice Models From Gauge Theory,
- [44] L. Frappat, P. Sorba, *Dictionary on Lie Superalgebras*, ENSLAPP-AL-600/96 and DSF-T-30/96, arXiv:hep-th/9607161.
- [45] L. Frappat, A. Sciarrino, and P. Sorba, *Structure of Basic Lie Superalgebras and of their Affine Extensions*, Commun. Math. Phys. 121, 457-500 (1989).
- [46] Youssra Boujakhrou, El Hassan Saidi, Rachid Ahl Laamara, Lalla Btissam Drissi, *Superspin Chains Solutions from 4D Chern-Simons Theory*, J. High Energ. Phys. 2024, 43 (2024), arXiv:2309.04337v2 [hep-th].
- [47] Elena Poletaeva, *On matrix realizations of the Lie superalgebra $D(2, 1; \alpha)$* , Journal of Geometry and Physics 60 (2010) 1656–1664, arXiv:1008.2433 [math.RT].
- [48] M. Gunaydin, Modern Physics Letters A, 6, 3239 (1991)
- [49] M. Scheunert, *The Theory of Lie Superalgebras*, in: Lecture Notes in Mathematics vol. 716, Springer, Berlin, 1979.
- [50] K. Iohara , Y. Koga, *Central extensions of Lie superalgebras*, Comment. Math. Helv. 76 (2001) 110-154.
- [51] Leyu Han, *Centres of centralizers of nilpotent elements in exceptional lie superalgebras*, Journal of Algebra and Its Applications (2021): 2250053, arXiv:2203.04423v1 [math.RT].
- [52] V.K. Dobrev, *Note on Centrally Extended $su(2|2)$ and Serre Relations*, Fortsch.Phys.57:542-545,2009, arXiv:0903.0511v3 [hep-th].
- [53] N.I. Stoilova, J. Van der Jeugt, *Orthosymplectic $Z_2 \times Z_2$ -graded Lie superalgebras and parastatistics*, J. Phys. A: Math. Theor. 57 095202 (2024), arXiv:2402.11952 [math-ph].
- [54] Y Boujakhrou et al 2022, *Lax operator and superspin chains from 4D CS gauge theory*, J. Phys. A: Math. Theor. 55 415402,
- [55] K. Intriligator, N. Seiberg, *Mirror Symmetry in Three Dimensional Gauge Theories*, Phys.Lett.B387:513-519,1996, arXiv:hep-th/9607207.
- [56] Saidi, E.H. *Mutation symmetries in BPS quiver theories: building the BPS spectra*. J. High Energ. Phys. 2012, 18 (2012), arXiv:1204.0395v2 [hep-th].
- [57] Edward Witten, *Fermion Path Integrals And Topological Phases*, Rev. Mod. Phys. 88, 35001 (2016), arXiv:1508.04715.
- [58] Edward Witten, *Three Lectures On Topological Phases Of Matter*, La Rivista del Nuovo Cimento, 39 (2016) 313-370.

- [59] Liang Fu, Charles Kane, Eugene Mele, Topological Insulators in Three Dimensions, Phys. Rev. Lett. 98 (2007) 106803
- [60] Xiao-Liang Qi, Shou-Cheng Zhang, Topological insulators and superconductors, Rev. Mod. Phys. 83 (2011) 1057-1110.
- [61] L. B. Drissi, S. Lounis, E. H. Saidi, *Higher order topological matter and fractional chiral states*, Eur. Phys. J. Plus 137, 796 (2022), arXiv:2211.02362 [cond-mat.mes-hall].
- [62] L. B. Drissi, E. H. Saidi, *A signature index for third order topological insulators*, Journal Phys Condensed Matter, 32 (36) (2020), arXiv:2207.02901v1 [cond-mat.mes-hall]
- [63] Lalla Btissam Drissi, El Hassan Saidi, *Domain Walls in Topological Tri-hinge Matter*, European Physical Journal Plus 136, (68) (2021), arXiv:2206.11984 [cond-mat.mtrl-sci].
- [64] Rhiannon Gwyn, Anke Knauf, *Conifolds and Geometric Transitions*, Rev.Mod.Phys.8012:1419-1453,2008, arXiv:hep-th/0703289v3.
- [65] B.S. Acharya, L. Foscolo, M. Najjar and E.E. Svanes, *New G2-conifolds in M-theory and their Field Theory Interpretation*, arXiv:2011.06998 [hep-th].
- [66] El Hassan Saidi, *Topological $SL(2)$ Gauge Theory on Conifold*, arXiv:hep-th/0601020.
- [67] N.C. Leung, C. Vafa, *Branes and Toric Geometry*, Adv.Theor.Math.Phys.2:91-118,1998.
- [68] A. Belhaj, L. B. Drissi, J. Rasmussen, E. H. Saidi, A. Sebbar, *Toric Calabi-Yau supermanifolds and mirror symmetry*, J.Phys.A38:6405-6418,2005, arXiv:hep-th/0410291v2.
- [69] R.Abounasr, M.Ait Ben Haddou, A.El Rhalami, E.H.Saidi, *Algebraic Geometry Realization of Quantum Hall Soliton*, J.Math.Phys. 46 (2005) 022302, arXiv:hep-th/0406036.
- [70] Sigiswald Barbier, Sam Claerebout, *A superunitary Fock model of the exceptional Lie supergroup $D(2,1;\alpha)$* , arXiv:2301.07982v2 [math.RT].
- [71] Sergey Fedoruk, Evgeny Ivanov, Olaf Lechtenfeld, *New $D(2,1;\alpha)$ Mechanics with Spin Variables*, JHEP 1004:129,2010, arXiv:0912.3508v2 [hep-th].
- [72] Sergey Fedoruk, Evgeny Ivanov, Olaf Lechtenfeld, *$OSp(4|2)$ Superconformal Mechanics*, JHEP 0908:081,2009arXiv:0905.4951v2 [hep-th].
- [73] Constantin Bachas, Eric D'Hoker, John Estes, Darya Krym, *M-theory Solutions Invariant under $D(2, 1; \gamma) \oplus D(2, 1; \gamma)$* , arXiv:1312.5477v2 [hep-th].
- [74] L.B Drissi, E.H Saidi, *From orthosymplectic structure to super topological matter*, Nuclear Physics B, Volume 989, April 2023, 116128, arXiv:2303.12584v1 [hep-th].

- [75] E. H Saidi, L. B Drissi, *5D N=1 super QFT: symplectic quivers*, Nuclear Physics B 974, January 2022, 115632, arXiv:2112.04695 [hep-th].
- [76] Youssra Boujakhrou, El Hassan Saidi, *On Exceptional 't Hooft Lines in 4D-Chern-Simons Theory*, Nuclear Physics B 980, 2022, 115795, arXiv:2204.12424v1 [hep-th].
- [77] Y. Boujakhrou, E.H Saidi, R. Ahl Laamara, L.B Drissi, *'t Hooft lines of ADE-type and Topological Quivers*, SciPost Phys. 15, 078 (2023), arXiv:2303.13879v1 [hep-th].
- [78] K. Costello, D. Gaiotto and J. Yagi, *Q-operators are 't Hooft lines*, arXiv:2103.01835.
- [79] R. Frassek, *Algebraic Bethe ansatz for Q-operators: The Heisenberg spin chain*, J.Phys. A 48 (2015) 294002 [arXiv:1504.04501].
- [80] R. Frassek, T. Lukowski, C. Meneghelli, M. Staudacher, *Oscillator Construction of $su(n|m)$ Q-Operators*, <https://doi.org/10.48550/arXiv.1012.6021>.
- [81] Rouven Frassek, Alexander Tsybaliuk, *Orthosymplectic superoscillator Lax matrices*, arXiv:2309.14199 [math-ph].
- [82] Sebastian Gerigk, *String States on $AdS_3 \times S^3$ from the Supergroup*, J. High Energ. Phys. 2012, 84 (2012), arXiv:1208.0345v3 [hep-th].
- [83] Gerhard Gotz, Thomas Quella, Volker Schomerus, *The WZNW model on $PSU(1,1|2)$* , JHEP 0703:003,2007, arXiv:hep-th/0610070v2.
- [84] Matthias R. Gaberdiel, Bin Guo, and Samir D. Mathur, *Tensionless strings on AdS_3 orbifolds*, arXiv:2312.01348v1 [hep-th]
- [85] I.M. Musson, *Lie superalgebras and enveloping algebras*, Graduate Studies in Mathematics 131, Amer. Math. Soc. (2012).
- [86] K. Costello and M. Yamazaki, *Gauge Theory And Integrability, III*, arXiv:1908.02289.
- [87] T. Okuda, *Line operators in supersymmetric gauge theories and the 2d-4d relation*, in *New Dualities of Supersymmetric Gauge Theories*, J. Teschner ed., Springer, Cham (2016), p. 195-222 [DOI:10.1007/978-3-319-18769-3_7] [arXiv:1412.7126].
- [88] H. Hayashi, T. Okuda and Y. Yoshida, *ABCD of 't Hooft operators*, JHEP 04 (2021) 241 [arXiv:2012.12275].
- [89] K. Maruyoshi, T. Ota and J. Yagi, *Wilson-'t Hooft lines as transfer matrices*, JHEP 01 (2021) 072 [arXiv:2009.12391].
- [90] A. Kapustin and E. Witten, *Electric-Magnetic Duality And The Geometric Langlands Program*, Commun. Num. Theor. Phys. 1 (2007) 1 [hep-th/0604151].

- [91] A. Kapustin, Wilson-'t Hooft operators in four-dimensional gauge theories and S-duality, Phys. Rev. D 74 (2006) 025005 [hep-th/0501015].
- [92] V.V. Bazhanov, T. Lukowski, C. Meneghelli and M. Staudacher, A Shortcut to the Q-Operator, J. Stat. Mech. 1011 (2010) P11002 [arXiv:1005.3261].
- [93] H. Hayashi, T. Okuda, Y. Yoshida, ABCD of 't Hooft operators. J. High Energ. Phys. 2021, 241 (2021). [https://doi.org/10.1007/JHEP04\(2021\)241](https://doi.org/10.1007/JHEP04(2021)241).
- [94] A. Kapustin, E. Witten, Electric-Magnetic Duality And The Geometric Langlands Program, Communications in Number Theory and Physics Volume 1 (2007) Number 1 (arXiv:hep-th/0604151).
- [95] A. Kapustin, Wilson-'t Hooft operators in four-dimensional gauge theories and S-duality, Phys.Rev. D74 (2006) 025005 (arXiv:hep-th/0501015).
- [96] B. Charbonneau, J. Hurtubise, *Singular Hermitian-Einstein monopoles on the product of a circle and a Riemann surface*, <https://doi.org/10.48550/arXiv.0812.0221>.
- [97] K. Maruyoshi, T. Ota, J. Yagi, *Wilson- 't Hooft lines as transfer matrices*. J. High Energ. Phys. 2021, 72 (2021). [https://doi.org/10.1007/JHEP01\(2021\)072](https://doi.org/10.1007/JHEP01(2021)072).