

NEW MATRIX PERTURBATION BOUNDS VIA CONTOUR EXPANSION

PHUC TRAN, VAN VU

ABSTRACT. Matrix perturbation bounds (such as Weyl and Davis-Kahan) are frequently used in many branches of mathematics.

Most of the classical results in this area are optimal, in the worst case analysis. However, in modern applications, both the ground and the noise matrices frequently have extra structural properties. For instance, it is often assumed that the ground matrix is essentially low rank, and the noise matrix is random or pseudo-random.

We aim to rebuild a part of perturbation theory, adapting to these modern assumptions. We will do this using a contour expansion argument, which enables us to exploit the skewness among the leading eigenvectors of the ground and the noise matrix (which is significant when the two are uncorrelated) to our advantage. In the current paper, we focus on the perturbation of eigenspaces. This helps us to introduce the arguments in the cleanest way, avoiding the more technical consideration of the general case. In applications, this case is also one of the most useful. More general results appear in a subsequent paper.

Our method has led to several improvements, which have direct applications in central problems. Among others, we derive a sharp result for perturbation of a low rank matrix with random perturbation, answering an open question in this area. Next, we derive new results concerning the spike model, an important model in statistics, bridging two different directions of current research. Finally, we use our results on the perturbation of eigenspaces to derive new results concerning eigenvalues of deterministic and random matrices. In particular, we obtain new results concerning the outliers in the deformed Wigner model and the least singular value of random matrices with non-zero mean.

1. INTRODUCTION

1.1. Perturbation theory. Let A and E be symmetric real matrices of size n , and $\tilde{A} := A + E$. We view A as the truth (or data/ground) matrix, and E as noise.

Consider the spectral decomposition

$$A = \sum_{i=1}^n \lambda_i u_i u_i^T,$$

where λ_i are the eigenvalues of A and u_i the corresponding eigenvectors. We order the eigenvalues decreasingly, $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. We also order the singular values of A in similar manner $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n$. It is well known that $\{\sigma_1, \dots, \sigma_n\} = \{|\lambda_1|, \dots, |\lambda_n|\}$. Therefore, $\sigma_i = |\lambda_{\pi(i)}|$ for a permutation $\pi \in S_n$. We use notation $\tilde{\lambda}_i, \tilde{u}_i$, etc for \tilde{A} , with the same ordering.

Department of Mathematics, Yale University, 219 Prospect Ave, New Haven, Connecticut 06511, USA
Email: phuc.tran@yale.edu, van.vu@yale.edu

Remark 1.2. To simplify the presentation, we assume that the eigenvalues (singular values) are different, so the eigenvectors (singular vectors) are well-defined (up to signs). Our argument works for matrices with multiple eigenvalues, and the results will be stated in this generality. We will also consider rectangular matrices; see Section 2.13.

In applications involving large matrices, researchers often pay attention to the leading eigenvectors, eigenvalues, and eigenspaces of the matrix. The following matrix functionals are of fundamental interest, not only in mathematics, but also in a number of other branches of sciences. We denote the (general) functional by g .

- $g(A) = \lambda_1$, the largest eigenvalue of A
- $g(A) = \sigma_1 = \|A\|$, the largest singular value (or the spectral norm) of A .
- $g(A) = u_1$, the leading eigenvector of A .
- $g(A) = \Pi_p := \sum_{i=1}^p u_i u_i^T$, the orthogonal projection onto the eigenspace spanned by the leading p eigenvectors of A .
- $g(A) = \Pi_{(p)} = \sum_{i=1}^p u_{\pi(i)} u_{\pi(i)}^T$, the orthogonal projection onto the singular space spanned by the leading p singular vectors of A .
- $g(A) = \Pi_{(p)} A := A_p$, the best rank p approximation of A .

A matrix perturbation bound is a bound for the difference

$$\|g(A + E) - g(A)\| = \|g(\tilde{A}) - g(A)\|,$$

given a pair of functional g and an appropriate norm $\|\cdot\|$ (notice that g can be a scalar, a vector, or a matrix). In this paper, we use l_2 norm for a vector and spectral norm for a matrix.

The theory of matrix perturbation has a long history; see [14, 33, 48, 67, 84]. Let us discuss a few classical results in this area.

Theorem 1.3 (Weyl [14, 85]). *For any $1 \leq i \leq n$,*

$$(1) \quad \left| \tilde{\lambda}_i - \lambda_i \right| \leq \|E\|.$$

To state the next result, we consider a subset $S \subset \{1, \dots, n\}$ and denote by Π_S the orthogonal projection on to the subspace spanned by the eigenvectors of $u_i, i \in S$. Define $\delta_S = \min\{|\lambda_i - \lambda_j|, i \in S, j \notin S\}$.

Theorem 1.4 (Davis-Kahan [14, 28]). *We have*

$$(2) \quad \|\tilde{\Pi}_S - \Pi_S\| \leq \frac{2\|E\|}{\delta_S}.$$

The most important special case is when $S = \{1, \dots, p\}$. In this case, we obtain the following estimate for the perturbation of Π_p , the orthogonal projection onto the leading p -eigenspace of A .

Corollary 1.5. *Let $\delta_p := \lambda_p - \lambda_{p+1}$. Then*

$$(3) \quad \left\| \tilde{\Pi}_p - \Pi_p \right\| \leq 2 \frac{\|E\|}{\delta_p},$$

Remark 1.6. The reader can work out a similar result for $\Pi_{(p)}$ as an exercise. Notice that any meaningful application of Theorem 1.4 or Corollary 1.5 should require the gap-to-noise assumption

$$(4) \quad \delta_S \geq 2\|E\|.$$

As a matter of fact, one often needs to replace the constant 2 by a much larger quantity, depending on the strength of the bound required in applications.

1.7. Sharpness of the classical bounds. These classical bounds are sharp. For instance, consider Weyl's inequality. If the first eigenvalue of both A and E are realized by the same eigenvector, namely $Eu_1 = \mu_1 u_1$, where μ_1 , the largest eigenvalue of E is also its spectral norm, then $\mu_1 + \lambda_1$ is the first eigenvalue of \tilde{A} , with u_1 being the corresponding eigenvector.

Let us now consider the sharpness of Davis-Kahan theorem through the following example. Set

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 1-\epsilon & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 1-2\epsilon & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 1-3\epsilon & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 1-4\epsilon & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 0 & 1/2 & \cdots & 0 \\ \cdots & & & & & & & \\ 0 & 0 & 0 & 0 & 0 & 0 & \cdots & 1/2 \end{pmatrix},$$

$$E := \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 5\epsilon & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 5\epsilon & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ \cdots & & & & & & & \\ 0 & 0 & 0 & 0 & 0 & 0 & \cdots & 0 \end{pmatrix}, \quad \tilde{A} = A + E = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 1+4\epsilon & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 1+3\epsilon & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 1-3\epsilon & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 1-4\epsilon & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 0 & 1/2 & \cdots & 0 \\ \cdots & & & & & & & \\ 0 & 0 & 0 & 0 & 0 & 0 & \cdots & 1/2 \end{pmatrix}.$$

In this situation, we are interested in the perturbation of the first eigenvector of A , which is e_1 . The gap $\delta_1 = \lambda_1 - \lambda_2 = \epsilon$ and the norm of E is 5ϵ . The first eigenvector of \tilde{A} is e_2 , which is orthogonal to e_1 (so the perturbation is maximum). This shows that Davis-Kahan theorem is sharp up to the constant factor 2. (If one replaces 2 by .2, the theorem would fail.)

We notice here that in both constructions, the sharpness of the bounds is due to a very strong correlation between the leading eigenvectors of A and those of E . If we replace the diagonal 2×2 non-trivial block of E in the above construction by a random 2×2 matrix

with the same norm, then the first eigenvector of \tilde{A} will be very close to that of A . (The simple computation is left as an exercise.)

1.8. Matrices in modern applications. In modern applications, especially those from data science, both the ground matrix A and the noise matrix E usually have some spectral properties.

For instance, the following pattern often emerges: A has a small number of large leading eigenvalues (in absolute value), with the rest being relatively small compared to these. In statistics, these eigenvalues are referred to as “spikes”; see Section 5 for a discussion.

A special case is when A has low rank (thus most eigenvalues are zero). This case appears abundantly in applications. For instance, the very intense research on the matrix completion problem (Netflix prize type problem) in recent years, started with Candes-Tao paper [21], has been based on this assumption; the problem would have been unfeasible otherwise.

For the noise matrix E , the most common and natural assumption is that it is random, or semi-random in certain sense. This is just the nature of noise. What is worth mentioning here is that in many problems, the noise matrix is something we define out of the context of the problem, rather than caused by nature. Even in this cases, this “artificial” noise is still random. Let us illustrate this point by the following example:

Example. Statistical Block Model Statistical Block Model is a popular problem in statistics and combinatorics (in combinatorics it is often called the hidden partition problem). Given a vertex set $V = \{1, \dots, n\}$, we partition V into k subsets V_1, \dots, V_k , where k is small (typically a constant not depending on n). Between a vertex $u \in V_i$ and a vertex $v \in V_j$ ($1 \leq i \leq j \leq k$), we draw a random edge with probability p_{ij} , independently. The goal is to recover the partition under some conditions on the densities p_{ij} , given one instance of the resulting random graph. There is a huge literature on this problem; see [1, 17, 27, 37, 38, 62, 65] and the references therein.

Let \tilde{A} be the adjacency of the random graph and A its expectation. We can write $\tilde{A} = A + E$, where $E := \tilde{A} - A$ is a (block) random matrix with zero mean and independent entries. Notice that by construction A has at most k different rows, and thus it has rank at most k . The “noise” matrix E is not real noise, it is created by us.

Researchers frequently use the spectral method to attack this problem; see, for example [1, 30, 37, 56, 81]. The key observation here is that in many settings, the leading eigenvectors of A reveal the partition. Thus, it is natural to consider the leading eigenvectors of \tilde{A} , and use perturbation bounds to show that these vectors still reveal a good approximation of the truth. In many cases, one can further refine this approximation to obtain the exact partition. We will discuss a concrete example in Section 5.11.

It is clear that in this situation, the leading eigenvectors of A and E are uncorrelated.

1.9. Our goal and achievements. We goal is to rework and improve a part of perturbation theory, taking into account the modern assumptions discussed above. Our key idea is to exploit the skewness of the noise matrix E with respect to the *important* eigenvectors of A

(this skewness rules out most of the sharpness constructions of the classical results, such as those considered in subsection 1.7).

Our key quantity to measure this "skewness" is $x := \max_{u,v} |u^T E v|$, where u and v run over the set of (important) eigenvectors of A ; see Definition 2.2 for more details. In general, $\|E v\|$ can be as big as $\|E\|$. However, if there is no correlation between E and the eigenvectors of A , then the two vectors $E v$ and u must be in general position with respect to each other, and one can typically hope for a \sqrt{n} saving when considering their inner product.

Example: Random noise. It is a well-known fact from random matrix theory that if E is a (random) Wigner matrix, then with probability $1 - o(1)$, $\|E\| = (2 + o(1))\sqrt{n}$, but for any fix unit vectors u and v , $|u^T E v|$ is only $O(\log n)$. (A Wigner matrix is a random symmetric matrix whose upper diagonal entries are i.i.d subgaussian random variables with mean 0 and variance 1.) The estimate for $\|E\|$ is a well-known result of Füredi and Komlós [38]. The $O(\log n)$ bound is a simple consequence of Chernoff concentration inequality, see [78].

With our approach, we can replace $\|E\|$ by x in a number of situations, with relatively mild assumptions and some adjustments. Let us present a typical result. Let A be a low rank matrix, $\text{rank} A = r = O(1)$, and E be a Wigner matrix. We are interested in the perturbation of the leading eigenvector of A . In this setting, Davis-Kahan bound (Corollary 1.5) gives (with high probability)

$$\|\tilde{u}_1 - u_1\| = O\left(\frac{\|E\|}{\delta_1}\right) = O\left(\frac{\sqrt{n}}{\delta_1}\right),$$

where $\delta_1 = \lambda_1 - \lambda_2$ is the gap between the first and second eigenvalues of A . On the other hand, our new results yield, with high probability, that

$$\|\tilde{u}_1 - u_1\| = O\left(\frac{\log n}{\delta_1} + \frac{\|E\|}{\lambda_1}\right);$$

(see Corollary 5.8 and the discussion following it).

Notice that the bound $\frac{\|E\|}{\delta_1}$ is now replaced by $\frac{\log n}{\delta_1}$, which is basically $\frac{x}{\delta_1}$ (see the example above), resulting in a (basically) \sqrt{n} saving, as mentioned above. The reader may notice that we have a new error term $\frac{\|E\|}{\lambda_1}$. This term is necessary. Considering from the engineering viewpoint, if the signal of the ground matrix A (represented by λ_1) is significantly weaker than the intensity of the noise (represented by $\|E\|$), then the information of the ground matrix must be lost, and the perturbed matrix behaves like a random one. For a quantitative statement, we refer to the BBP transition phenomenon in random matrix theory [6], discovered by Ben Arous, Baik, and Peche in the asymmetric setting; see [12] for the symmetric version.

The above result is a very special case of a general theorem on perturbation of eigenspaces, which is the main object of our paper. This theorem is deterministic, but it contains statement involving random matrices, such as the above, as a special case; see Theorem 2.12.

In this paper, we consider matrix functionals $g(A)$ of the following form. Let $f(z)$ be an analytic function, and S a subset of the index set $\{1, 2, \dots, n\}$. Consider the spectral decomposition $A = U\Lambda U^T$ of A , where Λ is the diagonal matrix with the i th entry being λ_i . We consider

$$(5) \quad g(A) := f_S(A) = U f_S(\Lambda) U^T,$$

where $f_S(\Lambda)$ is the diagonal matrix with the i th entry being $f(\lambda_i)$ if $i \in S$ and zero otherwise. This is one of the most commonly matrix functionals considered in the literature; see [14]. For instance, it is easy to see that both the orthogonal projection onto an eigenvector subspace and the low rank approximation can be written using $f \equiv 1$ and $f(z) = z$ (together with a proper choice of S), respectively. These two operators alone are already of huge importance with respect to applications in numerical analysis and data science.

Our method works for any function f (such as $f \equiv 1$, $f(z) = z$, $f(z) = z^2$, $f(z) = e^z$, etc) in (5). In this paper, we focus on the simplest case when $f \equiv 1$. This is equivalent to considering the perturbation of eigenspaces (Davis-Kahan type estimates). The function $f_S(A)$ is exactly the orthogonal projection onto the eigenvectors spanned by the eigenvalues with indices in S . If $S = \{i\}$, for a particular index i , then we obtain a perturbation bound for the eigenvector u_i .

Focusing on this case will help us to present the main ideas in the cleanest manner, avoiding rather technical analytical details of the general case. Moreover, this special case is already useful in many applications, discussed throughout the paper. We will present the general case in a forthcoming paper [72].

It is clear that one cannot represent the eigenvalues and singular values (say λ_1 or σ_1) using (5), as $f_S(A)$ is always a matrix. On the other hand, somewhat surprisingly, we can still use our results to derive new perturbation results for eigenvalues and singular values, via an additional trick; see Section 4 for details. As a matter of fact, we can handle all functionals g listed after Remark 1.2.

Our investigation makes use of a technique called "contour expansion", which is of independent interest and we will present it in details in Section 3. In the next section, we describe our main results.

2. NEW PERTURBATION BOUNDS FOR EIGENSPACES

As already mentioned, in this paper, we solely focus on the case when the functional $f \equiv 1$. This is equivalent to measuring perturbation of eigensubspaces.

2.1. New eigenspace perturbation bounds. Let A be an $n \times n$ symmetric matrix with eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ and corresponding eigenvectors u_1, \dots, u_n . Let S be a proper subset of the index set $\{1, 2, \dots, n\}$ and denote

$$\Lambda_S := \{\lambda_i | i \in S\} \quad \text{and} \quad \Pi_S := \sum_{i \in S} u_i u_i^T.$$

Π_S is the orthogonal projection on to the space spanned by eigenvectors selected by S . We would like to bound its perturbation. In this general setting, we allow A to have multiple

eigenvalues, so in order to make Π_S well-defined, we assume that Λ_S and $\Lambda_{\bar{S}}$ are separated, and let

$$\delta_S := \min_{i \in S, j \notin S} |\lambda_i - \lambda_j| > 0.$$

Now we choose a quantity $\bar{\lambda} > 0$ and let $N_{\bar{\lambda}}(S)$ be the (spectral) $\bar{\lambda}$ -neighborhood of S :

$$(6) \quad N_{\bar{\lambda}}(S) := \{j \in [n] \mid |\lambda_i - \lambda_j| \leq \bar{\lambda} \text{ for some } i \in S\},$$

We call the eigenvectors corresponding to $N_{\bar{\lambda}}(S)$ the *important* eigenvectors. This definition is motivated by the fact that we do not need to pay attention to eigenvectors whose corresponding eigenvalues are far away. The vectors we need to pay special care to are those relatively close to those in S , as they can swap with them via the perturbation.

We set

$$(7) \quad p := |S|; r := |N_{\bar{\lambda}}(S)|.$$

Now we turn our attention to the noise matrix E . To measure the interaction of E with the eigenvectors, we define

Definition 2.2. Define

- $x := \max_{i,j} |u_i^T E u_j|$, where $i, j \in N_{\bar{\lambda}}(S)$.
- $y := \max_{i,j,k} |u_i^T E \left(\sum_{l \notin N_{\bar{\lambda}}(S)} \frac{u_l u_l^T}{\lambda_k - \lambda_l} \right) E u_j|$, where $i \neq j \in N_{\bar{\lambda}}(S)$ and $k \in S$.
- $w := \max_i \|u_i^T E\|$, where $i \in N_{\bar{\lambda}}(S)$.

As mentioned a few times before, the skewness quantity x plays an important role; it represents the skewness of the noise matrix E with respect to the important eigenvectors of A . The key assumption for our results is

$$(8) \quad (\mathbf{C0}) \quad \max \left\{ \frac{\sqrt{p} \|E\|}{\bar{\lambda}}, \frac{rx}{\delta_S}, \frac{\sqrt{r}w}{\sqrt{\bar{\lambda}}\delta_S} \right\} \leq \frac{1}{12}.$$

Now we are ready to state our results for two important cases of S .

2.2.1. Projection onto the leading eigenspace. Let $S := \{1, 2, \dots, p\}$, then $N_{\bar{\lambda}}(S) := \{1, 2, \dots, r\}$, in which r is the smallest index such that $|\lambda_p - \lambda_{r+1}| > \bar{\lambda}$.

Theorem 2.3. *Under Assumption C0*

$$(9) \quad \|\tilde{\Pi}_p - \Pi_p\| \leq 12\sqrt{p} \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{r}x}{\delta_p} + \frac{\sqrt{r}y}{\delta_p} \right).$$

Let us compare this theorem with Corollary 1.5 (Davis-Kahan bound). To simplify the analysis, we assume that $r = O(1)$. Thus, the (9) is essentially

$$\|\tilde{\Pi}_p - \Pi_p\| = O \left(\frac{\|E\|}{\bar{\lambda}} + \frac{x}{\delta_p} + \frac{y}{\delta_p} \right).$$

On the other hand, Corollary 1.5 (Davis-Kahan theorem) yields

$$\|\tilde{\Pi}_p - \Pi_p\| = O \left(\frac{\|E\|}{\delta_p} \right).$$

The key improvement here is that we can replace the spectral norm $\|E\|$ by the skewness x , resulting in the term $\frac{x}{\delta_p}$ instead of $\frac{\|E\|}{\delta_p}$. This is the message we like to emphasize throughout the paper. This improvement enables us to obtain a strong bound even when the gap δ_p is relatively small (say much smaller than $\|E\|$, which makes the original Davis-Kahan bound vacuous).

The two new errors are of smaller order of magnitude in many applications. Moreover, they are necessary (representing the truth), and not the artifact of our analysis; see Section 5 and Section 14 for more details. Our result is thus sharp up to a constant factor (in terms of both the bound and the assumption).

Assume that we need to show that $\|\tilde{\Pi}_p - \Pi_p\| \leq \epsilon$, for some given small quantity ϵ . We can use Theorem 2.3 as follows. First set $\bar{\lambda} = 24\epsilon^{-1}\|E\|$. This makes the contribution of the first term $12\frac{\sqrt{p}\|E\|}{\bar{\lambda}}$ in the bound $\epsilon/2$. Next, we bound r , the number of important vectors with respect to $\bar{\lambda}$. Here some information about the spectral distribution of A will help. For instance, if the spectrum of A decays fast, and p is small, then r is also relatively small. In general, to make r not too large, we need to assume that the spectrum of A does not cluster too heavily near λ_p . The hope is that the contribution from the last two terms on the RHS of (9) will be smaller than $\epsilon/2$, which would typically be the case if E interacts with the eigenvectors (especially the important ones) in a favorable manner. We will present a few examples in the discussion below.

2.4. Dealing with the y term. So far we have emphasized the role of the skewness x . On the other hand, the new error term involving y also needs some care, as the definition of y is slightly technical. In many cases, we already obtain good results using the following trivial (and often generous) estimate

$$(10) \quad y \leq \frac{\|E\|^2}{\lambda}.$$

Corollary 2.5. *Under Assumption C0*

$$(11) \quad \|\tilde{\Pi}_p - \Pi_p\| \leq 12\sqrt{p} \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{r}x}{\delta_p} + \frac{\sqrt{r}\|E\|^2}{\delta_p\bar{\lambda}} \right).$$

Let us make a quick comparison between Corollary 2.5 and Theorem 1.5, via a toy problem. For convenience, we ignore all constant terms and write $A \lesssim B$ if $A \leq cB$ for some positive constant c .

Assume that $p = O(1)$ and the eigenvalues of A have roughly equal spacing, say, $\lambda_i - \lambda_{i+1} \approx K\|E\| := \delta$ for some quantity $K > 0$, at least in some neighborhood of λ_p . In this case, Theorem 1.5 gives

$$(12) \quad \|\tilde{\Pi}_p - \Pi_p\| \lesssim \frac{\|E\|}{\delta} \approx \frac{1}{K}.$$

On the other hand, Corollary 2.5 yields

$$\|\tilde{\Pi}_p - \Pi_p\| \lesssim \frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{r}x}{\delta} + \frac{\sqrt{r}\|E\|^2}{\bar{\lambda}\delta}.$$

We can set $\bar{\lambda} = \epsilon^{-1}K\|E\|$, for some small quantity $\epsilon > 0$ to be chosen. Thus, $\sqrt{r} = \epsilon^{-1}$ and

$$\frac{\|E\|}{\bar{\lambda}} = \frac{\epsilon}{K}; \quad \frac{\sqrt{r}\|E\|^2}{\bar{\lambda}\delta} = \frac{1}{K^2}.$$

Clearly, both terms improve (12). The trade-off is the middle term

$$\frac{\sqrt{r}x}{\delta} = \frac{|u_i^T E u_j|}{\epsilon K \|E\|},$$

for some important unit vectors u_i, u_j (which realizes the max in the definition of x ; see Definition 2.2). As mentioned several times before, our point here is that if E interacts with u_i, u_j in a favorable manner, then $|u_i^T E u_j|$ is much less than $\|E\|$. We can then use this saving to offset the ϵ term in the denominator, leading to an improvement over (12).

If the critical gap $\lambda_p - \lambda_{p+1}$ is smaller (equals $\frac{1}{2}\|E\|$, say), then the original Davis-Kahan bound (Corollary 1.5) is vacuous, as the bound becomes bigger than 1. On the other hand, our above argument still works if the other gaps remain the same ($K\|E\|$ for some large K). The only difference is that the term $\frac{\sqrt{r}\|E\|^2}{\bar{\lambda}\delta_p}$ becomes $\frac{1}{K}$ (instead of $\frac{1}{K^2}$). Thus, we still get a strong bound. This illustrates the important fact that our theorems *do not* require the gap-to-noise assumption (4), which has been a critical and standard assumption for eigenspace perturbation bounds since Davis-Kahan's original paper.

For a finer analysis, let us point out that our above argument still works even if *all other gaps are also small* (equal $\frac{1}{2}\|E\|$, say), given that we can improve just a little bit on the

trivial bound (10) (by replacing $\|E\|^2$ by a smaller quantity); see, for instance, the setting of Corollary 2.7 below and many results in Section 5, where E is random.

Set

$$\bar{x} := \max_{i,j} |u_i^T E u_j|, \text{ where } 1 \leq i, j \leq n.$$

We can also bound y using \bar{x} . It is clear that in the current setting (with $S = \{1, \dots, p\}$)

$$(13) \quad y \leq \bar{x}^2 \sum_{l \geq r+1} \frac{1}{\lambda_p - \lambda_l}.$$

Corollary 2.6. *Under Assumption C0*

$$(14) \quad \|\tilde{\Pi}_p - \Pi_p\| \leq 12\sqrt{p} \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{r}x}{\delta_p} + \frac{\sqrt{r}\bar{x}^2}{\delta_p} \sum_{l \geq r+1} \frac{1}{\lambda_p - \lambda_l} \right).$$

The presence of the term $\sum_{l \geq r+1} \frac{1}{\lambda_p - \lambda_l}$ allows us to use information about the spectral distribution of A for our advantage. For instance, if the eigenvalues beyond λ_r has minimum gap $t := \min_{l \geq r} \lambda_l - \lambda_{l+1} > 0$, then

$$\sum_{l \geq r+1} \frac{1}{\lambda_p - \lambda_l} \leq \sum_{h=1}^n \frac{1}{\bar{\lambda} + ht} \approx t^{-1} \int_t^{nt} \frac{1}{\bar{\lambda} + x} dx \leq t^{-1} \log n.$$

In certain situations, there is a natural choice for the important vectors. For instance, if A has low rank ($\text{rank} A = r \ll n$) and the non-zero eigenvalues of A are positive and of the same order of magnitude, then it is tempting to let the important eigenvectors be the non-trivial ones. In this case, $\bar{\lambda} = |\lambda_p|$. Furthermore, the definition of y simplifies significantly, as all non-important eigenvalues are zero. In this case,

$$y = \frac{1}{\sigma} \max_{i \neq j \in I} u_i^T E \left(I_n - \sum_{l \in I} u_l u_l^T \right) E u_j,$$

where I denotes of the important vectors (a short hand for $N_{\bar{\lambda}}(S)$) and

$$(15) \quad \sigma := \min_{1 \leq i \leq p} |\lambda_i|.$$

By the triangle inequality, it is easy to see that

$$(16) \quad y \leq \frac{1}{\sigma} \left(\max_{i \neq j \in I} |u_i^T E^2 u_j| + r x^2 \right).$$

Set

$$(17) \quad \bar{y} := \max_{i \neq j \in I} |u_i^T E^2 u_j|,$$

we have

Corollary 2.7. *Assume that A has rank r with r positive eigenvalues. Under Assumption C0*

$$(18) \quad \|\tilde{\Pi}_p - \Pi_p\| \leq 12\sqrt{p} \left(\frac{\|E\|}{\lambda_p} + \frac{\sqrt{r}x}{\delta_p} + \frac{\sqrt{r}\bar{y}}{\delta_p\sigma} + \frac{r^{3/2}x^2}{\delta_p\sigma} \right).$$

The quantity \bar{y} is, in spirit, quite similar to the quantity x (see Definition 2.2), as it captures the *skewness* between E and the eigenvectors of A . Similar to the situation with x , if E interacts favorably with the important eigenvectors of A , then $\bar{y} \ll \|E\|^2$. Indeed, if the two vectors Eu_i and Eu_j (with length at most $\|E\|$) are in general position with respect to each other, then they would be near orthogonal and one can expect a saving. For instance if E is a (random) Wigner matrix then, with high probability, we can save a factor of order \sqrt{n} .

Remark 2.8. The assumption that all eigenvalues are positive is for convenience. We can easily modify the theorem for the general case; details are left as an exercise.

2.8.1. *Projection on to the leading singular space.* Consider $S = \{\pi(1), \dots, \pi(p)\}$; we can rewrite $S = S_1 \cup S_2$, $S_1 := \{\lambda_1, \dots, \lambda_{k-1}\}$, $S_2 := \{\lambda_n, \lambda_{n-1}, \dots, \lambda_{n-(p-k)}\}$, for some value $1 \leq k \leq p$. Thus, $N_{\bar{\lambda}}(S) = \{1, 2, \dots, r_+, n - r_- + 1, n - r_- + 2, \dots, n\}$ for properly chosen r_+ and r_- ; set $r := r_+ + r_-$. Denote $\bar{\delta}_p := \min\{\delta_{k-1}, \delta_{n-(p-k)-1}\}$.

Theorem 2.9. *Under assumption C0*

$$(19) \quad \|\tilde{\Pi}_{(p)} - \Pi_{(p)}\| \leq 12\sqrt{p} \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{r}x}{\bar{\delta}_p} + \frac{\sqrt{r}y}{\bar{\delta}_p} \right),$$

given that the RHS is less than 1.

Similarly, by the trivial bound $y \leq \frac{\|E\|^2}{\bar{\lambda}}$, we obtain the following corollaries.

Corollary 2.10. *Under Assumption C0*

$$(20) \quad \|\tilde{\Pi}_{(p)} - \Pi_{(p)}\| \leq 12\sqrt{p} \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{r}x}{\bar{\delta}_p} + \frac{\sqrt{r}\|E\|^2}{\bar{\delta}_p\bar{\lambda}} \right).$$

Corollary 2.11. *Let $r = \text{rank } A$. Under Assumption C0*

$$(21) \quad \|\tilde{\Pi}_{(p)} - \Pi_{(p)}\| \leq 12\sqrt{p} \left(\frac{\|E\|}{|\lambda_p|} + \frac{\sqrt{r}x}{\bar{\delta}_p} + \frac{\sqrt{r}\bar{y}}{\bar{\delta}_p\sigma} + \frac{r^{3/2}x^2}{\bar{\delta}_p\sigma} \right).$$

2.11.1. *Main Theorem: Projection with general S .* Both of the above two theorems are corollaries of our main result below, which treats an arbitrary set S . To state the theorem, let us first draw a simple closed curve (contour) Γ containing the eigenvalues with indices in S inside it (and no other eigenvalues of A) such that the distance from any eigenvalue of A to Γ is at least $\delta/2$ (for some quantity $\delta > 0$) and the distance from $\lambda_j, j \notin N_{\bar{\lambda}}(S)$ to Γ is at least $\bar{\lambda}/2$. The simplest way to do so is to construct Γ using horizontal and vertical segments, where the vertical ones bisect the intervals connecting an eigenvalue in S with its nearest neighbor (left or right) outside S . Let \tilde{S} be the set of eigenvalues of \tilde{A} inside Γ and set

$$\tilde{\Lambda}_{\tilde{S}} := \{\tilde{\lambda}_i | i \in \tilde{S}\} \quad \text{and} \quad \tilde{\Pi}_{\tilde{S}} = \sum_{i \in \tilde{S}} \tilde{u}_i \tilde{u}_i^T.$$

The following assumption replaces Assumption **C0**.

$$(22) \quad (\mathbf{D0}) \quad \max \left\{ \frac{\sqrt{p} \|E\|}{\bar{\lambda}}, \frac{rx}{\delta}, \frac{\sqrt{r}w}{\sqrt{\bar{\lambda}\delta}} \right\} \leq \frac{1}{12}.$$

Theorem 2.12. *Under Assumption **D0***

$$(23) \quad \|\tilde{\Pi}_{\tilde{S}} - \Pi_S\| \leq 12\sqrt{p} \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{r}x}{\delta_S} + \frac{\sqrt{r}y}{\delta_S} \right).$$

We would like to point out that we have two different parameters δ and δ_S , one in the condition and one in the conclusion of the theorem. Clearly, we can choose $\delta = \delta_S$ as in the special case when $S = \{1, \dots, p\}$ (by definition δ is always at most δ_S .) In this case, **D0** and **C0** are the same. However, we intentionally keep two parameters here, as we can use the flexibility of choosing δ to derive interesting consequences; see Section 4. In fact, under **D0**, **C0** is satisfied.

To derive Theorem 2.3 from Theorem 2.12, we notice that if the bound $12\sqrt{p} \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{r}x}{\delta_S} + \frac{\sqrt{r}y}{\delta_S} \right)$ is strictly less than 1, then the two subspaces spanned by S and \tilde{S} must have the same dimension. Thus, if in Theorem 2.3, we make the contour very long to the right, then $\tilde{\Pi}_{\tilde{S}}$ must be precisely $\tilde{\Pi}_p$. One can make a similar argument to prove Theorem 2.9.

2.13. Rectangular matrices. In this section, we extend our theorems to the non-symmetric case. Our object now is an $m \times n$ matrix A of rank $r_A \leq \min\{m, n\}$. Consider the singular decomposition of A

$$A := U\Sigma V^T,$$

where $\Sigma = \text{diag}(\sigma_1, \dots, \sigma_{r_A})$ is a diagonal matrix containing the non-zero singular values $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_{r_A}$. The columns of the matrices $U = (u_1, \dots, u_{r_A})$ and $V = (v_1, \dots, v_{r_A})$ are the left and right singular vectors of A , respectively. By definition

$$U^T U = V^T V = I_{r_A}.$$

Let S be a subset of $\{1, 2, \dots, r_A\}$. Using our notation, we set

$$(24) \quad \Pi_S^{left} = \sum_{i \in S} u_i u_i^T, \Pi_S^{right} = \sum_{i \in S} v_i v_i^T.$$

When $S = \{1, \dots, p\}$, we simply write Π_p^{left} (Π_p^{right}). Let E be an $m \times n$ “noise” matrix and set $\tilde{A} = A + E$. Define $\tilde{\Pi}_S^{left}, \tilde{\Pi}_S^{right}$ accordingly. Our goal is to find the upper bounds of left and right eigenspace perturbations, i.e.

$$\left\| \tilde{\Pi}_S^{left} - \Pi_S^{left} \right\|, \left\| \tilde{\Pi}_S^{right} - \Pi_S^{right} \right\|.$$

We will symmetrize A and \tilde{A} to use our main theorem. There are two natural ways to do this, with different advantages.

Additive symmetrization. Set $\mathcal{A} := \begin{pmatrix} 0 & A \\ A^T & 0 \end{pmatrix}, \mathcal{E} := \begin{pmatrix} 0 & E \\ E^T & 0 \end{pmatrix}$

It is well known that the eigenvalues of \mathcal{A} are $\sigma_1, \dots, \sigma_{r_A}, -\sigma_1, \dots, -\sigma_{r_A}$ and with corresponding eigenvectors $(u_1, v_1), \dots, (u_{r_A}, v_{r_A}), (u_1, -v_1), \dots, (u_{r_A}, -v_{r_A})$ (where (u, v) is the concatenation of u and v). The application of Theorem 2.12 in this case is simple and direct. On the other hand, because of the concatenation, one cannot separate the left singular space from the right one, and so one will obtain a bound on

$$\max \left\{ \left\| \tilde{\Pi}_S^{left} - \Pi_S^{left} \right\|, \left\| \tilde{\Pi}_S^{right} - \Pi_S^{right} \right\| \right\}.$$

Multiplicative symmetrization. Here we consider $M = AA^T, \tilde{M} := (A + E)(A^T + E^T)$. The eigenvalues of M are $\sigma_1^2, \dots, \sigma_{r_A}^2$ and its eigenvectors are the left singular vectors of A . Thus, we can focus solely on the perturbation of the left singular space. (Setting $M = A^T A$ of course will take care of the right one.) On the other hand, the new “noise” matrix has become $\tilde{E} = EA^T + A^T E + EE^T$, and its treatment requires more care, and also the assumption **C0** will become a bit more complicated.

In this paper, we consider only additive symmetrization. The consideration of the multiplicative one is a bit more technical and will be addressed in a later paper.

2.14. Results using additive symmetrization. Following Theorem 2.12, we choose a quantity $\bar{\sigma}$ and let $N_{\bar{\sigma}}(S)$ be the $\bar{\sigma}$ -neighborhood of S :

$$(25) \quad N_{\bar{\sigma}}(S) := \{i | \sigma_i - \sigma_j \leq \bar{\sigma} \text{ for some } j \in S\}.$$

$$(26) \quad p := |S|, r = |N_{\bar{\sigma}}(S)|, \text{ and } \delta_S = \min_{\sigma_i \in S, \sigma_j \notin S} |\sigma_i - \sigma_j|.$$

Now we have the analogue of Definition 2.2.

Definition 2.15.

- $x := \max_{i,j} |u_i^T E v_j|$ where $i, j \in N_{\bar{\sigma}}(S)$.
- $y := \max_{i,j,k} \left\{ \left| v_i^T E^T \left(\sum_{l \notin N_{\bar{\sigma}}(S)} \frac{u_l u_l^T}{\sigma_k - \sigma_l} \right) E v_j \right|, \left| u_i^T E \left(\sum_{l \notin N_{\bar{\sigma}}(S)} \frac{v_l v_l^T}{\sigma_k - \sigma_l} \right) E^T u_j \right| \right\}$ where $i < j \in N_{\bar{\sigma}}(S)$ and $k \in S$.
- $w := \max_{i,j} \{ \|E v_i\|, \|E^T u_j\| \}$ where $i, j \in N_{\bar{\sigma}}(S)$.

Next are the analogue of Assumptions **C0** and **D0**:

$$(27) \quad (\mathbf{C1}) : \max \left\{ \frac{\sqrt{p} \|E\|}{\bar{\sigma}}, \frac{rx}{\delta_S}, \frac{\sqrt{r}w}{\sqrt{\bar{\sigma}\delta_S}} \right\} \leq \frac{1}{12}.$$

$$(28) \quad (\mathbf{D1}) : \max \left\{ \frac{\sqrt{p} \|E\|}{\bar{\sigma}}, \frac{rx}{\delta}, \frac{\sqrt{r}w}{\sqrt{\bar{\sigma}\delta}} \right\} \leq \frac{1}{12}.$$

Similar to the previous section, we draw a contour Γ which encircles those singular values with indices in S (viewed as positive eigenvalues of \mathcal{A}) such that the distance from any eigenvalues of \mathcal{A} to Γ is at least $\delta/2$, and the distance from $\sigma_j, j \notin N_{\bar{\sigma}}(S)$ to Γ is at least $\bar{\sigma}/2$. Let \tilde{S} be the set of eigenvalues of \tilde{A} inside Γ . Denote

$$\tilde{\Pi}_{\tilde{S}}^{left} = \sum_{i \in \tilde{S}} \tilde{u}_i \tilde{u}_i^T \quad \text{and} \quad \tilde{\Pi}_{\tilde{S}}^{right} = \sum_{i \in \tilde{S}} v_i v_i^T.$$

Theorem 2.16. *Under Assumptions **D1**,*

$$(29) \quad \max \left\{ \left\| \tilde{\Pi}_{\tilde{S}}^{left} - \Pi_S^{left} \right\|, \left\| \tilde{\Pi}_{\tilde{S}}^{right} - \Pi_S^{right} \right\| \right\} \leq 12\sqrt{2p} \left(\frac{\|E\|}{\bar{\sigma}} + \frac{\sqrt{rx} + \sqrt{ry}}{\delta_S} \right).$$

We also obtain the following corollaries for $S = \{1, 2, \dots, p\}$ and $S = \{\pi(1), \dots, \pi(p)\}$.

Corollary 2.17. *Under Assumptions **C1**,*

$$(30) \quad \max \left\{ \left\| \tilde{\Pi}_p^{left} - \Pi_p^{left} \right\|, \left\| \tilde{\Pi}_p^{right} - \Pi_p^{right} \right\| \right\} \leq 12\sqrt{2p} \left(\frac{\|E\|}{\bar{\sigma}} + \frac{\sqrt{rx} + \sqrt{ry}}{\delta_p} \right).$$

Using the trivial bound $y \leq \frac{\|E\|^2}{\bar{\sigma}}$, we obtain

Corollary 2.18. *Under Assumptions **C1**,*

$$(31) \quad \max \left\{ \left\| \tilde{\Pi}_p^{left} - \Pi_p^{left} \right\|, \left\| \tilde{\Pi}_p^{right} - \Pi_p^{right} \right\| \right\} \leq 12\sqrt{2p} \left(\frac{\|E\|}{\bar{\sigma}} + \frac{\sqrt{rx}}{\delta_p} + \frac{\sqrt{r}\|E\|^2}{\delta_p \bar{\sigma}} \right).$$

The constant 12 gets replaced by $12\sqrt{2}$ due to the concatenation and usage of the Pythagorean theorem.

2.19. Structure of the rest of the paper. The rest of the paper has 8 main sections. In the next section, Section 3, we introduce our contour argument. We believe that this argument is of independent interest, and are currently applying it to few other topics beyond the one discussed here.

The next four sections contain applications in a number of different areas.

- In Section 4, we use Theorem 2.12 to derive perturbation bounds for the eigenvalues. It is somewhat surprising that we can use the contour for this purpose. We will be able to achieve new bounds not only for the leading eigenvalues but also for the smallest one (corresponding to the least singular value). We can apply these new bounds to obtain a more complete understanding of the behavior of outliers in deformed Wigner matrices, a popular topic in random matrix theory.
- In Section 5, we derive several new bounds for the case when the noise E is random. Among others, these partially settle an open problem raised in [59]. Here is also where we can see very concrete quantization of the assumption that E interacts favorably with the eigenvectors, leading to strong improvements over classical results. We will use the results derived in this section to treat the so-called “many hidden cliques” problem, rooted in random graph theory and theoretical computer science.
- In Section 6, we consider the spiked model in statistics. This is a model of fundamental importance in the study of covariance matrices and has been heavily studied; see [82]. On the other hand, this is also a model where our new results apply naturally, as the number of “spikes” (large eigenvalues) is usually small, making the spectrum of the matrix convenient for our method. In this section, we obtain new bounds for perturbation of the leading eigenvectors of the covariance matrix (Principal Component Analysis), which improves and complements many earlier results.
- In Section 7, we discuss applications in various areas of statistics and theoretical computer science, including matrix completion, differential privacy, and Gaussian mixture. Due to the length of the paper and the technicality involved in these applications, we will only discuss the nature of these applications; full details will appear in future papers.

The next 3 sections are devoted to the proofs

- In Section 8, we present the three main technical lemmas of the proof, and derive Theorem 2.12 from these lemmas.
- In Section 9.1, we first prove a combinatorial lemma, an essential tool used for the combinatorial profiling (mentioned earlier). Next, we prove the three main lemmas.
- In Section 10, we prove the theorems from Sections 5 and 6. Many technical parts of these proofs will be deferred to the Appendices.

We conclude the paper with Section 14, where we discuss the sharpness of our new results.

3. THE CONTOUR EXPANSION ARGUMENT

In this section, we introduce our key technical idea. Our strategy is to reduce the action of the noise matrix E to the subspace spanned by the important eigenvectors of the ground matrix A . The improvement here is two-fold. First, this makes it possible to replace $\|E\|$ with $x := \max |u_i^T E u_j|$, which represents the interaction of E with the eigenvectors of A . Second, there is a dimension reduction mechanism built in the proof, which allows us to consider only the important eigenvectors, thus reducing the dimension of the problem significantly. This reduction is important in a number of applications; see, for instance, Section 5.1.

We start with Cauchy theorem. Consider a contour Γ and a point $a \notin \Gamma$, together with a complex function f which is analytic on Γ and its interior. Cauchy theorem asserts that

$$(32) \quad \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} \frac{f(z)}{z-a} dz = f(a)$$

if a is inside Γ and zero otherwise. (Here and later \mathbf{i} denotes $\sqrt{-1}$.) Now consider the spectral decomposition of A , $A = U\Lambda U^T$, and define

$$f(A) := U f(\Lambda) U^T$$

where $f(\Lambda)$ is diagonal with entries $f(\lambda_i)$. Let Γ be a contour containing $\lambda_i, i \in S$ where S is a subset of $\{1, \dots, n\}$, and assume that all $\lambda_j, j \notin S$ are outside Γ . One obtains the classical contour formula, which is well known in numerical analysis and mathematical physics literature

$$(33) \quad \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} f(z)(z-A)^{-1} dz = \sum_{i \in S} f(\lambda_i) u_i u_i^T := f_S(A);$$

see [14, 48]. By choosing an appropriate f , we obtain many important matrix functionals

- $f \equiv 1$, $S = \{1, \dots, p\}$, then $\sum_{i \in S} f(\lambda_i) u_i u_i^T = \Pi_p$, the orthogonal projection on the subspace spanned by the first p eigenvectors of A .
- $f \equiv 1$, $S = \{\pi(1), \dots, \pi(p)\}$, then $\sum_{i \in S} f(\lambda_i) u_i u_i^T = \Pi_{(p)}$, the orthogonal projection on the subspace spanned by the first p singular vectors of A .
- $f(z) = z$, $S = \{\pi(1), \dots, \pi(p)\}$, then $\sum_{i \in S} f(\lambda_i) u_i u_i^T = \Pi_{(p)} A = A_p$, the best rank p approximation of A .
- $f(z) = z^2$, $S = \{\pi(1), \dots, \pi(p)\}$, then $\sum_{i \in S} f(\lambda_i) u_i u_i^T = \Pi_{(p)} A^2$, the best rank p approximation of A^2 .

For a moment, assume that the eigenvalues $\tilde{\lambda}_i, i \in S$ are inside Γ , and all $\tilde{\lambda}_j, j \notin S$ are outside, then

$$(34) \quad \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} f(z)(z - \tilde{A})^{-1} dz = \sum_{i \in S} f(\tilde{\lambda}_i) \tilde{u}_i \tilde{u}_i^T := f_S(\tilde{A}).$$

Thus, one obtain a contour identity for the perturbation

$$(35) \quad f_S(\tilde{A}) - f_S(A) = \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} f(z)[(z - \tilde{A})^{-1} - (z - A)^{-1}] dz.$$

Using the resolvent formula

$$M^{-1} - (M + N)^{-1} = (M + N)^{-1} N M^{-1}$$

and the fact that $z - A = (z - \tilde{A}) + E$, we obtain

$$(z - \tilde{A})^{-1} - (z - A)^{-1} = (z - A)^{-1} E (z - \tilde{A})^{-1}.$$

Repeating this argument, we obtain (formally at least)

$$(36) \quad (z - \tilde{A})^{-1} - (z - A)^{-1} = \sum_{s=1}^{\infty} (z - A)^{-1} [E(z - A)^{-1}]^s.$$

(To make this identity valid we need to make sure that the RHS converges, but let us skip this issue for a moment.) Now set

$$(37) \quad F_s := \int_{\Gamma} f(z)(z - A)^{-1} [E(z - A)^{-1}]^s dz.$$

We have

$$f_S(\tilde{A}) - f_S(A) = \frac{1}{2\pi\mathbf{i}} \sum_{s=1}^{\infty} F_s.$$

Therefore,

$$(38) \quad \left\| f_S(\tilde{A}) - f_S(A) \right\| \leq \frac{1}{2\pi} \sum_{s=1}^{\infty} \|F_s\|.$$

The Taylor expansion idea is well-known and has been used by many researchers; see for instance [48, Chapter 2]. The key matter is to bound $\|F_s\|$. In [48, Chapter 2], one simply used

$$\|F_s\| \leq \|E\|^s \max_{z \in \Gamma} |f(z)| \int_{\Gamma} \|(z - A)^{-1}\|^{s+1} |dz| = O \left[\max_{z \in \Gamma} |f(z)| \left(\frac{\|E\|}{\delta_S} \right)^s \right].$$

In the case when $f \equiv 1$, if we sum up the RHS, then we obtain exactly the Davis-Kahan bound, with no improvement.

Our approach to estimate $\|F_s\|$ is significantly different, involving two extra expansions and a some combinatorial ideas. Notice that $(z - A)^{-1} = \sum_{i=1}^n \frac{1}{z - \lambda_i} u_i u_i^T$. In order to control F_s ,

we split the RHS into two parts $P = \sum_{i \in I} \frac{1}{z - \lambda_i} u_i u_i^T$ and $Q = \sum_{j \in J} \frac{1}{z - \lambda_j} u_j u_j^T$, where I is the set of *important* indices, and J its complement. Then

$$f(z)(z - A)^{-1}[E(z - A)^{-1}]^s = f(z)(P + Q)[E(P + Q)]^s.$$

We now expand this product as the sum of 2^{s+1} terms, each is of the form $f(z)R_1ER_2 \dots ER_{s+1}$, where R_i is either P or Q . This is the second expansion in our proof, after the Taylor expansion.

At a first look, the situation looks worse, as the number of terms is enormous. However, we have two key observations which will help with the analysis. First, notice that the only term which does not involve any important direction is $f(z)QE \dots EQ$ (which involves only Q operator). Using the formula for Q , we further expand this product into the sum of $(n - r)^{s+1}$ terms, each of the form

$$f(z) \prod_{l_i \in J} \frac{1}{z - \lambda_{l_i}} u_{l_1} u_{l_1}^T E u_{l_2} u_{l_2}^T E \dots E u_{l_s} u_{l_s}^T.$$

The key point here is that all eigenvalues λ_{l_i} involving in this form are *outside* the contour, thus the integral of every product $f(z) \prod_{l_i \in J} \frac{1}{z - \lambda_{l_i}}$ is zero. Therefore, $\int_{\Gamma} f(z)QE \dots EQ dz$ vanishes completely.

For the remaining terms, we are not so lucky. They simply will not vanish. Our second key observation is that any of these terms contains many P 's, each of which adjacent to an E . We will rely crucially on this property to exploit the interaction of the important directions (contained in P) with the noise matrix E .

For a concrete example, let us consider the term $f(z)PEP$ ($s = 1$). Writing $P = \sum_{i \in I} \frac{1}{z - \lambda_i} u_i u_i^T$ and expanding the product (this is the third expansion in the analysis), we obtain

$$f(z)PEP = \sum_{i_1, i_2 \in I} \frac{f(z)}{(z - \lambda_{i_1})(z - \lambda_{i_2})} u_{i_1} u_{i_1}^T E u_{i_2} u_{i_2}^T = \sum_{i_1, i_2 \in I} \frac{f(z)}{(z - \lambda_{i_1})(z - \lambda_{i_2})} u_{i_1} u_{i_2}^T (u_{i_1}^T E u_{i_2}).$$

As u_{i_1}, u_{i_2} are unit vectors, a trivial bound for $|u_{i_1}^T E u_{i_2}|$ is $\|E\|$. However, as mentioned before, if E is in a general position with respect to both vectors u_{i_1} and u_{i_2} , then we expect a saving. The reader can see that this naturally leads to the definition of the critical ‘‘skewness’’ parameter x (see Definition 2.2).

The main technical part of the proof is to handle $f(z)(P + Q)[E(P + Q)]^s$. After the second and third expansions, this is an (enormous) sum of terms of the form

$$h(z)u_{l_1} u_{l_1}^T E u_{l_2} u_{l_2}^T E \dots E u_{l_{s+1}} u_{l_{s+1}}^T,$$

where

$$h(z) = \frac{f(z)}{\prod_{i \in I} (z - \lambda_i) \times \prod_{j' \in J} (z - \lambda_{j'})}.$$

This function $h(z)$ changes from term to term. The technicality of the argument comes from the fact that we will need to treat the complex part and the linear algebraic part of the formula together, as they compensate each other in some delicate way.

There are many ways to integrate $h(z)$. (For instance, after many trials, we have found out that using the residue theorem is a *wrong* way.) We end up designing an algorithm to execute this integration. The key point here is that we can use this algorithm to obtain a combinatorial profile of the term, in form of a graph. We next use this combinatorial profile to split the whole collection of terms into many subsums, which we will be able to estimate separately, by a combination of analytic, linear algebraic, and combinatorial tools.

In this paper, we focus on the special case when $f \equiv 1$, which corresponds to the orthogonal projection onto a selected eigensubspace. We do so in order to present the key ideas of the method in the simplest manner, avoiding some heavy analytic consideration of the general case. This special case already has plenty applications in many different areas. The treatment of f will be presented in a sequential paper [72].

Our argument leads to sharp results. Technically, we are going to estimate F_1 and F_2 almost precisely, and obtain sharp bounds, $\|F_1\| \leq h_1$ and $\|F_2\| \leq h_2$. The sum $h := h_1 + h_2$ will contain all the terms in our main theorem. We next show that the remaining F_s satisfies

$$(39) \quad \|F_s\| \leq 2^{3-s} h,$$


or some estimate of this sort (with 2^{3-s} replaced by another fast decaying function). Putting these together, we would obtain

$$\sum_{s=1}^{\infty} \|F_s\| \leq 3h.$$

Once this plan has been properly executed, we obtain the desired bound, which is sharp, as the computation concerning F_1 and F_2 are exact, and their contribution dominates the rest.

4. PERTURBATION OF EIGENVALUES

4.1. Eigenvalue perturbation bounds via contour optimization. It is somewhat surprising that we can use the contour argument to derive bounds for the eigenvalues. To present the idea, let us consider the perturbation of λ_1 . To start, we draw a contour around λ_1



We make sure that λ_1 is the only eigenvalue of A inside this contour. Now let us assume that using this contour, we have a non-trivial bound for $\tilde{\Pi}_1 - \Pi_1$, namely

$$(40) \quad \|\tilde{\Pi}_1 - \Pi_1\| < 1.$$

Since both $\tilde{\Pi}_1$ and Π_1 are orthogonal projections, the subspaces they are projecting onto must have the same dimension. Therefore, we conclude that the contour contains exactly one eigenvalue of \tilde{A} . But since we can push the right edge of the contour to (plus) infinity, this eigenvalue has to be $\tilde{\lambda}_1$. This way, we obtain

$$\lambda_1 - \tilde{\lambda}_1 \leq \bar{\delta},$$

where $\bar{\delta}$ is the distance from λ_1 to the left edge of the contour.

Using this method, we can have a bound on $\lambda_1 - \tilde{\lambda}_1$ by optimizing $\bar{\delta}$, under assumption **D0**. To make **D0** hold, we choose $\bar{\lambda} \geq 12\|E\|$, and define $r = |N_{\bar{\lambda}}(S)|$ with respect to this $\bar{\lambda}$. Set $\delta := \max\{12rx, 144rw^2/\bar{\lambda}\}$. This is the smallest δ for which **D0** holds; this value of δ is legitimate if $\delta \leq \delta_1 = \lambda_1 - \lambda_2$. Our $\bar{\delta}$ is exactly $\delta/2$. We obtain

Corollary 4.2. *Choose $\bar{\lambda} \geq 12\|E\|$ and set $r := |N_{\bar{\lambda}}(\{1\})|$. Assume that*

$$\delta_1 = \lambda_1 - \lambda_2 \geq \delta := \max\{12rx, 144rw^2/\bar{\lambda}\}.$$

Assume furthermore that

$$12 \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{rx}}{\delta_1} + \frac{\sqrt{ry}}{\delta_1} \right) < 1.$$

Then

$$(41) \quad \lambda_1 - \tilde{\lambda}_1 \leq \max\{6rx, 72rw^2/\bar{\lambda}\}.$$

If we choose $\bar{\lambda} = \frac{\lambda_1}{2}$, then r is simply the number of eigenvalues λ_i such that $\lambda_i \geq \frac{\lambda_1}{2}$. The bound becomes $O\left(\frac{rw^2}{\lambda_1} + rx\right) = O\left(r\left(\frac{\|E\|^2}{\lambda_1} + x\right)\right)$.

Corollary 4.3. *Assume that there are $O(1)$ eigenvalues larger than $\lambda_1/2$, and $\|E\| = o(\lambda_1)$, $\|E\| = o(\delta_1)$. Then*

$$\lambda_1 - \tilde{\lambda}_1 = O\left(\frac{\|E\|^2}{\lambda_1} + x\right).$$

Notice that in this setting $\frac{\|E\|^2}{\lambda_1} = o(\|E\|)$. Thus our bound is better than the Weyl bound $\lambda_1 - \tilde{\lambda}_1 \leq \|E\|$ if $x \ll \|E\|$. (Recall that Weyl bound is sharp A and E share the same leading eigenvector.)

We can also derive an upper bound for $\tilde{\lambda}_1 - \lambda_1$, via the “holing” trick, which we discuss in the next section. Theorem 4.2 is sharp, and sheds some lights on behavior of outliers in deformed Wigner matrices, a recent topic in random matrix theory; see Section 4.11.

It is clear that one can obtain similar bounds for other eigenvalues by placing the contour properly. For instance, by considering a contour separating $\lambda_1, \dots, \lambda_p$ from the rest of the spectrum, we can show

Corollary 4.4. *Choose $\bar{\lambda} \geq 12\|E\|$ and set $r := |N_{\bar{\lambda}}(\{1, \dots, p\})|$. Assume that*

$$\delta_p = \lambda_p - \lambda_{p+1} \geq \delta := \max\{12rx, 144rw^2/\bar{\lambda}\}.$$

Assume furthermore that

$$12 \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{r}x}{\delta_p} + \frac{\sqrt{r}y}{\delta_p} \right) < 1.$$

Then

$$(42) \quad \lambda_p - \tilde{\lambda}_p \leq \max\{6rx, 72rw^2/\bar{\lambda}\}.$$

4.5. The “holing” trick and perturbation of the least singular value. Assume that A has full rank and $\sigma_{\min} > 0$ be the least singular value of A . Without loss of generality, we can assume $\sigma_{\min} = \lambda_i$, where λ_i is the smallest positive eigenvalue of A (thus λ_{i+1} is negative and has absolute value at least σ_{\min}).

By Weyl inequality $\tilde{\sigma}_{\min} \geq \sigma_{\min} - \|E\|$, providing that the RHS is positive. If $\sigma_{\min} < \|E\|$, then in principle, \tilde{A} can be singular. We are going to show that if E interacts nicely with the eigenvectors of eigenvalues near zero, then $\tilde{\sigma}_{\min}$ is still comparable with σ_{\min} , even when it is much smaller than $\|E\|$. Consequently, in these cases, the noise does not make the matrix ill-conditioned; it hardly changes the condition number.

For a number $T > 0$, let $\Lambda(T)$ be the set of eigenvalues with absolute value at most T , and $x(T) := \max |u_i^T E u_j|$, where u_i, u_j are eigenvectors corresponding to two eigenvalues in $\Lambda(T)$. Let $r(T) = |\Lambda(T)|$.

Theorem 4.6. *If there is a choice of T such that*

$$(43) \quad 48 \left(\frac{\|E\|}{T} + \frac{r(T)x(T)}{\sigma_{\min}} + \frac{r(T)\|E\|^2}{T\sigma_{\min}} \right) < 1,$$

then $\tilde{\sigma}_{\min} \geq \frac{1}{2}\sigma_{\min}$.

To prove this theorem, we artificially add an all zero row and column to both A and E (say, the bottom row). This way, we have a new pair A_0 and E_0 , where A_0 has a zero eigenvalue with eigenvector $(0, 0, \dots, 0, 1)$. Now we consider a contour around zero with width $\sigma_{\min}/2$. We apply our argument with respect to A_0, E_0 and this contour, with T playing the role

of $\bar{\lambda}$. The set S has exactly one eigenvalue (zero) in it. The assumption (43) guarantees that $\|\tilde{\Pi}_S - \Pi_S\| < 1$ (the constant 48 is just a convenient choice, any large enough constant would do), so the contour must contain exactly one eigenvalue of $\tilde{A}_0 = A_0 + E_0$. But by the construction, \tilde{A}_0 also has a zero eigenvalue. This implies that $\tilde{A} = A + E$ does not have any eigenvalue in the contour, proving the theorem.

Let E be a Wigner matrix and A a symmetric matrix with $\sigma_{\min} \geq n^{1/3+\epsilon}$, for some small constant $\epsilon > 0$. It is clear that $\|E\| \approx 2\sqrt{n} \gg \sigma_{\min}$.

Corollary 4.7. *Consider A and E as above. Assume furthermore that A has at most $n^{1/3}$ eigenvalues with absolute value at most $n^{2/3+\epsilon}$. Then with probability $1 - o(1)$, $\tilde{\sigma}_{\min} \geq \frac{1}{2}\sigma_{\min}$. Consequently, the condition number of \tilde{A} is at most (asymptotically) twice that of A .*

Proof of Corollary 4.7. To prove the first part of the corollary, set $T = n^{2/3+\epsilon}$. We have $r(T) \leq n^{1/3}$. Standard results in random matrix theory show that with probability $1 - o(1)$, $\|E\| = (2 + o(1))\sqrt{n}$ and $x(T) = O(\log n)$. An easy computation shows that

$$\left(\frac{\|E\|}{T} + \frac{r(T)x(T)}{\sigma_{\min}} + \frac{r(T)\|E\|^2}{T\sigma_{\min}} \right) = o(1),$$


and thus Theorem 4.6 implies that $\tilde{\sigma}_{\min} \geq \frac{1}{2}\sigma_{\min}$.

To prove the second statement, recall that the condition number $\kappa(A) = \frac{\sigma_{\max}(A)}{\sigma_{\min}(A)}$. By the assumption of the corollary, $\sigma_{\max}(A) \geq n^{2/3+\epsilon} \gg \|E\|$, with probability $1 - o(1)$. Thus, with this probability, $\sigma_{\max}(\tilde{A}) = (1 + o(1))\sigma_{\max}(A)$, concluding the proof.

The constants $1/3, 2/3$ can be further improved; however, we choose not to do it here, to maintain the flow of the paper.

The holing trick. Notice what really happens is that we show that a *hole* (a contour with no eigenvalue inside) with respect to A remains a hole with respect to \tilde{A} .

4.8. Upper bounds for eigenvalue perturbation. We can use the holing trick to obtain an upper bound for $\tilde{\lambda}_1 - \lambda_1$, making Corollary 4.2 both-sided, as follows. We extend the matrix A to $A_\lambda = \begin{pmatrix} A & 0 \\ 0 & \lambda \end{pmatrix}$, and consider $\tilde{A}_\lambda = A_\lambda + E_0$. We set $\lambda > \lambda_1$, and consider a contour which goes sufficiently far to the right which contains only λ . This is a hole with respect to A , and if it remains so, with respect to \tilde{A} , then we obtain $\tilde{\lambda}_1 - \lambda_1 \leq \bar{\delta}$, where $\bar{\delta}$ is the distance from λ_1 to the left edge of the contour.



Corollary 4.9. *Choose $\bar{\lambda} \geq 24\|E\|$ and set $r := |N_{\bar{\lambda}}(\{1\})|$. Set $\delta := \max\{12rx, 144rw^2/\bar{\lambda}\}$. Then*

$$(44) \quad \tilde{\lambda}_1 - \lambda_1 \leq \bar{\delta} = \delta/2 = \max\{6rx, 72rw^2/\bar{\lambda}\}$$

Proof of Corollary 4.9. Set $\lambda = \lambda_1 + 16\delta$. The set S contains only λ and so $p = 1$, and $\delta_S = 16\delta$. Consider all eigenvalues of A of distance at most $\bar{\lambda}' := 16\delta + \bar{\lambda}$ from λ to be important. Notice that these eigenvalues (there are r of them) are of distance at most $\bar{\lambda} \geq 24\|E\|$ from λ_1 , and thus this set is well defined given A .

It is clear that with $\bar{\lambda}$ and δ as in the Corollary, Assumption **D0** holds. We set the contour so that the distance from its left edge to λ_1 is $\bar{\delta}$. The distance from the left edge to λ is 15δ . Notice that by construction $\delta_S = 16\delta$, and it is easy to check that the bound in Theorem 2.12 is less than 1. Applying the holing trick, we conclude that

$$\tilde{\lambda}_1 - \lambda_1 \leq \bar{\delta} = \delta/2 = \max\{6rx, 72rw^2/\bar{\lambda}\},$$

proving the Corollary.

Corollary 4.10. *Assume that $\lambda_1 \geq 48\|E\|$ and A has r eigenvalues $\geq \lambda_1/2$. Let $x := \max_{i,j \leq r} |u_i E u_j|$. Then*

$$(45) \quad \tilde{\lambda}_1 - \lambda_1 \leq \max \left\{ 6rx, 72r \frac{\|E\|^2}{\lambda_1} \right\}.$$

Proof of Corollary 4.10. We set $\bar{\lambda} := \lambda_1/2$ and apply the previous Corollary. It is clear that $\bar{\lambda} \geq 24\|E\|$. Since $w \leq \|E\|$ (see Definition 2.2), the bounds follows.

It is clear that one can obtain similar results for other eigenvalues, by the same argument.

4.11. Random matrices with low rank perturbation. The case when A has low rank and E is Wigner has been studied heavily in random matrix theory. In random matrix theory, E is viewed as the main object and A as the perturbation. The model $A + E$ is often referred to as deformed Wigner matrices; see [11, 12, 22, 50, 58] and the references therein.

In this setting, one assumes that A has rank $r = O(1)$, and the eigenvalues $\lambda_1 \geq \dots \geq \lambda_r$ of A are of order $\Theta(\sqrt{n}) = \Theta(\|E\|)$. The following result was obtained by Banaych-Georges and Nadakuditi.

Theorem 4.12. [12] *Let A be a diagonal matrix of rank r , where all eigenvalues are $O(\sqrt{n})$. If $\lambda_i > \sqrt{n}$, then with probability $1 - o(1)$*

$$(46) \quad \tilde{\lambda}_i - \lambda_i = (1 + o(1)) \frac{n}{\lambda_i}.$$

This theorem says that if $\lambda_i > \sqrt{n}$, then $\tilde{\lambda}_i$ will be an outlier of \tilde{A} (being outside the edges of the semi-circle distribution). This is the symmetric analogue of the BBP threshold phenomenon first observed by Ben-Arous, Baik, and Peche in [6].

In a later paper [50], Knowles and Jin removed the diagonal condition on A , and also gave a strong estimate for the $o(1)$ error term in (46).

It is an important assumption in these studies that at the beginning λ_i has to be on the same scale as $\|E\|$, namely $\lambda_i = \Theta(\sqrt{n})$. A natural question arises

Question 4.13. *What happens if $\lambda_i = \omega(\sqrt{n})$?*

In particular, one may wonder if $\tilde{\lambda}_i - \lambda_i$ is still of order $\frac{n}{\lambda_i}$. The following example shows that this is no longer true, if λ_i/\sqrt{n} tends to infinity sufficiently fast. Let A be a diagonal matrix with just one top entry equal $\lambda_1 = n^{1+\epsilon}$, and E is GOE. In this case,

$$\frac{n}{\lambda_1} = n^{-\epsilon} = o(1).$$

On the other hand, the top diagonal entry of $\tilde{A} = A + E$ is $\lambda_1 + N(0, 2)$. Since $\tilde{\lambda}_1$ dominates this entry, we have, with probability 1, that $\tilde{\lambda}_1 - \lambda_1 \geq N(0, 2)$, which is typically much larger than $\frac{n}{\lambda_1}$. (Here and later, $N(\mu, \sigma^2)$ denotes the gaussian random variable with mean μ and variance σ^2 .)

Corollary 4.10 provides an answer to this question, showing that besides the term $\frac{n}{\lambda_1} = \Theta(\frac{\|E\|^2}{\lambda_1})$, we need a correction term of order x . Notice that for E being Wigner, $u^T E v \approx N(0, 1)$ for any fixed unit vectors u and v . Thus, our result is best possible, up to a constant factor, given the above example.

Corollary 4.14. *Assume that E is Wigner. Let A be a diagonal matrix of rank r , where $r = O(1)$. If $\lambda_1 \geq 48\sqrt{n}$, then with probability $1 - o(1)$*

$$(47) \quad \tilde{\lambda}_1 - \lambda_1 = O\left(\frac{n}{\lambda_1} + x\right).$$

The proof is left as an exercise. The same result holds for other eigenvalues (under a mild assumption). Corollary 47 holds for very general model of random matrices (not necessarily Wigner), given that we are able to estimate $\|E\|$ (and then replace $48\sqrt{n}$ by $24\|E\|$). This result goes beyond cases considered in the random matrix theory literature.

5. EIGENSPACE PERTURBATION WITH RESPECT TO RANDOM NOISE

In this section, we consider the case when E is random, with independent, but not necessarily identically distributed entries. In particular, we allow the entries to have different variances, and the variance profile is going to play an important role in our analysis.

In subsection 5.1 below, we discuss several previous results in this topic, and a problem that has been a partial motivation of our study. Next, we present new results which give a partial solution to this problem and also lead to improvements in many situations.

5.1. Existing results for eigenspace perturbation. The case when A is low rank (say, r) and E is random was first studied by the second author in [80]. The leading intuition in [80] is that with respect to A , E should behave like an r dimensional random matrix, which would yield quantitative improvements.

In the ideal case when E is a Wigner matrix, its norm is $\Theta(\sqrt{n})$ (with high probability). Thus, Theorem 1.5 yields a bound of order $\frac{\sqrt{n}}{\delta_p}$. In the spirit of the above intuition, we would expect to replace \sqrt{n} by \sqrt{r} , or (perhaps more realistically) by some quantity depending polynomially on r and $\log n$. In [80], the second author proved

Theorem 5.2. *For any positive constants α_1, α_2 there is a positive constant C such that the following holds. Assume that $r := \text{rank}A \leq n^{1-\alpha_1}$ and $\lambda_1 \leq n^{\alpha_2}$. Let E be a random Bernoulli matrix. Then with high probability*

$$\left\| \tilde{\Pi}_1 - \Pi_1 \right\|^2 \leq C \left(\frac{\sqrt{r \log n}}{\delta_1} + \frac{\|E\|^2}{\delta_1 \lambda_1} + \frac{\|E\|}{\lambda_1} \right).$$

The first term in the bound is $\frac{\sqrt{r \log n}}{\delta_1}$, mirroring the above intuition. In many cases, this happens to be the leading term. However, we have an extra square on the LHS. Continuing in this direction, about 10 years ago, O'Rourke, Wang, and the second author [59] obtained the following stronger and more general theorem.

Definition 5.3. We say the $m \times n$ random matrix E is (C_1, c_1, γ) -concentrated if for any unit vectors $u \in \mathbb{R}^m, v \in \mathbb{R}^n$, and every $t > 0$,

$$\mathbf{P}(|u^T E v| > t) \leq C_1 \exp(-c_1 t^\gamma).$$

Theorem 5.4 ([59]). *Assume that E is (C_1, c_1, γ) -concentrated for a trio of constants $C_1, c_1, \gamma > 0$ and A has rank r . Let $1 \leq p \leq r$ be an integer. Then for any $t > 0$,*

$$\left\| \tilde{\Pi}_p - \Pi_p \right\| \leq 4\sqrt{2p} \left(\frac{\|E\|}{\lambda_p} + \frac{tr^{1/\gamma}}{\delta_p} + \frac{\|E\|^2}{\lambda_p \delta_p} \right),$$

with probability at least

$$1 - 6C_1 9^j \exp\left(-c_1 \frac{\delta_p^\gamma}{8^\gamma}\right) - 2C_1 9^{2r} \exp\left(-c_1 r \frac{t^\gamma}{4^\gamma}\right).$$

In the case when E is Wigner with subgaussian entries, $\gamma = 2$. If we further assume that $p = O(1)$, then Theorem 5.4 implies that with probability at least .99

$$\left\| \tilde{\Pi}_p - \Pi_p \right\| = O\left(\frac{\|E\|}{\lambda_p} + \frac{r^{1/2}}{\delta_p} + \frac{\|E\|^2}{\lambda_p \delta_p} \right).$$

We notice that in this bound, besides the term $\frac{r^{1/2}}{\delta_p}$, which corresponds to our dimension reduction intuition, there are two other terms. The term $\frac{\|E\|}{\lambda_p}$ is the noise-to-signal ratio, and is necessary. Studies in random matrix theory [6, 12] showed that when this ratio is large, the sum $A + E$ behaves like a totally random matrix, so all information from A is lost. More mysterious is the $\frac{\|E\|^2}{\lambda_p \delta_p}$ term, which also appeared in Theorem 5.2 (which had a different proof). Unlike the other two, there is no natural explanation for this term. Should it be there or is it just an artifact of the proof?

Question 5.5. *Is the third term $\frac{\|E\|^2}{\lambda_p \delta_p}$ necessary?*

Motivated by this question, very recently O'Rourke, Wang, and the second author [60] considered the case when E is GOE. In this case, they showed that only the first two terms are sufficient. (Their original setting is for rectangular matrices; the study of the symmetric case is actually simpler, using the same arguments.) In particular, they showed (under mild assumptions) that with high probability,

$$(48) \quad \left\| \tilde{\Pi}_p - \Pi_p \right\| = O \left(\frac{r^2 \log n}{\delta_p} + \frac{\|E\|}{\lambda_p} \right).$$

The same result holds for general Wigner matrix, but the proof requires new technical lemmas. This gives the impression that the answer to Question 5.5 is negative.

In [51], Koltchinskii and Xia also considered Gaussian noise and a general A (which can have arbitrary rank). Among others, they showed (under some assumption) that with probability $1 - o(1)$

$$\left\| \tilde{\Pi}_p - \Pi_p \right\| = O \left(\left(\frac{\|E\|}{\delta_{(p)}} \right)^2 + \frac{\sqrt{\log n}}{\delta_{(p)}} \right).$$

If we could ignore the term $\frac{\sqrt{\log n}}{\delta_{(p)}}$ on the RHS, then this is a quadratic improvement over the original Davis-Kahan theorem (Theorem 1.5). On the other hand, here we still need the gap-to-noise assumption (4). For many other related works, also see [25, 83, 86] and the references therein.

5.6. New results. Before stating the main results, we list a few ensembles of random matrices with increasing generality.

- **Wigner matrices:** This is the most well-known class of random matrices in the literature. Wigner matrices are Hermitian matrices with upper diagonal entries being independent (but not necessarily iid) random variables with mean 0 and variance 1. The diagonal entries are independent with mean 0 and variance 2. (This class contains GOE and GUE as a subcase).
- **Generalized Wigner matrices:** Wigner matrices are Hermitian matrices with upper diagonal and diagonal entries being independent (but not necessarily iid) random variables with mean 0 and variance $\sigma_{ij}^2, 1 \leq i \leq j \leq n$. The second moments σ_{ij}^2 do not need to be the same, but are comparable, namely there is a constant $C > 0$ such

that. $\sigma_{ij}^2/\sigma_{lk}^2 \leq C$ for all i, j, k, l . Finally $\sum_j \sigma_{ij}^2 = \sum_j \sigma_{lj}^2$ for any i and l . The last condition means the expectations of the square length of the rows are the same.

- Regular matrices: We further extend the class of Generalized Wigner matrices by dropping the condition that the moments are comparable. Thus a regular random matrix is a Hermitian matrix with upper diagonal and diagonal entries being independent (but not necessarily iid) random variables with mean 0 and variance $\sigma_{ij}^2, 1 \leq i \leq j \leq n$, where $\sum_j \sigma_{ij}^2 = \sum_j \sigma_{lj}^2$ for any i and l .

As already mentioned, in the case when E is random, the critical parameters x and y (see Definition 2.2) are relatively easy to estimate, and we can achieve significant saving compared to classical bounds.

Regarding Question 5.5, results in this section reveal a subtle fact. It has turned out that in the case when E is regular (which contains Wigner matrices as a subcase), the answer to Question 5.5 is indeed negative (which is consistent with the Gaussian case treated by O’rourke et al.; see (48)). On the other hand, the answer is positive in general; the error term in question *is not* an artifact of the proof, it has to be there, and originates from the third error term in Theorem 2.12. In the case when E is regular, this error term becomes negligible by a magic algebraic cancellation; see the discussion leading to (51).

Let $E = (\xi_{ij})_{i,j=1}^n$ be an $n \times n$ real symmetric random matrix with independent, 0-mean entries. We further assume that with probability 1,

$$|\xi_{ij}| \leq K, \sigma_{ij}^2 \leq \sigma^2, \forall 1 \leq i, j \leq n.$$

In particular, when the first assumption holds, we say that the matrix is K -bounded.

In the low-rank setting, it is convenient to choose the parameter r in Theorem 2.12 to be exactly the rank of A . Given this choice of r , we have that $\lambda_j = 0$, for all $j \geq r + 1$. Define

$$(49) \quad \epsilon_1 := \frac{\|E\|}{\lambda_p}, \epsilon_2 = \frac{1}{\delta_p}, \eta := \frac{\|E\|^2}{\lambda_p \delta_p}.$$

For a regular random matrices E with entries $\xi_{ij}, 1 \leq i, j \leq n$, let

$$m_2 := \frac{1}{n} \sum_j \sigma_{ij}^2.$$

By the definition of the regular matrices, m_2 does not depend on the choice of i . Furthermore, let

$$m_4 := \max_i \frac{1}{n} \sum_j \sigma_{ij}^4.$$

Particularly, when E is a K -bounded Wigner matrix, $m_2 := \frac{n+1}{n} \approx 1, m_4 \leq K^2 m_2$. Similar to previous sections, we now define our key assumption

$$(50) \quad (\mathbf{C2}) : \max \{ \sqrt{p} \epsilon_1, r t_1 \epsilon_2, \sqrt{r \eta} \} \leq \frac{1}{12}.$$

Here t_1 will be a parameter that is part of the theorems.

Theorem 5.7. *Assume that A is a symmetric matrix with rank r and E is a random symmetric matrix with independent entries. Assume that Assumption **C2** holds for some choice of t_1 . Then for any $t_2 > 0$, we have with probability at least*

$$1 - r^2 \exp\left(\frac{-t_1^2/2}{2\sigma^2 + Kt_1}\right) - \frac{5r(r-1)nm_4}{2t_2^2},$$

that

$$\left\| \tilde{\Pi}_p - \Pi_p \right\| \leq 12\sqrt{p} \left(\epsilon_1 + \sqrt{r}t_1\epsilon_2 + \frac{\sqrt{r}(t_2 + \mu)}{\|E\|} \epsilon_1\epsilon_2 \right),$$

where

$$\mu := \max_{1 \leq k < l \leq r} \left\{ \sum_i u_{ki}u_{li} \sum_j \sigma_{ji}^2 \right\}.$$

Notice that when E is regular, $\sum_j \sigma_{ji}^2$ does not depend on i . Thus, μ is precisely zero (as the vectors u_k and u_l are orthogonal). This is the “magic” that leads to the negative answer for Question 5.5 in the regular case. To see this, let us elaborate a bit further. Notice that with $\mu = 0$, the bound in the theorem reduces to

$$(51) \quad \left\| \tilde{\Pi}_p - \Pi_p \right\| \leq 12\sqrt{p} \left(\epsilon_1 + \sqrt{r}t_1\epsilon_2 + \frac{\sqrt{r}t_2}{\|E\|} \epsilon_1\epsilon_2 \right).$$

Assume that we want the bound to hold with probability at least $1 - \alpha$, we can work out the optimal value of t_1 and t_2 (up to constant factors) as follows

$$t_1 = 4(\sigma + K)\sqrt{\log(r/\alpha)}; t_2 = 5r\sqrt{nm_4}.$$

In particular, when E has subgaussian entries, we can (using a standard truncation argument) set $K = \log n$. Furthermore, $m_2 = 1$ and $m_4 \leq C$ for some absolute constant C (which can be bounded using the Φ_2 norm of the entries). In this case, we can set

$$t_1 = \sqrt{\log(r/\alpha)} \log n; t_2 = Cr\sqrt{n} = Cr\|E\|.$$

Plugging this back and using the definition of ϵ_1, ϵ_2 , we have

Corollary 5.8. *Assume that A is a symmetric matrix with rank r and E is a random symmetric matrix with independent entries. Let $\alpha > 0$ be a small quantity which may depend*

on n . Assume that Assumption **C2** for the above choice of t_1 . Then with probability at least $1 - \alpha$

$$\left\| \tilde{\Pi}_p - \Pi_p \right\| \leq 12\sqrt{\bar{p}} \left(\frac{\|E\|}{\lambda_p} + \frac{\sqrt{r \log(r/\alpha)} \log n}{\delta_p} + \frac{Cr^{3/2}\|E\|}{\lambda_p \delta_p} \right).$$

Notice that if $r^{3/2} = o(\delta_p)$, then the third term on the RHS is negligible compared to the first term, and we have the bound

$$\left\| \tilde{\Pi}_p - \Pi_p \right\| \leq 12\sqrt{\bar{p}} \left(\frac{\|E\|}{\lambda_p} + \frac{\sqrt{r \log(r/\alpha)} \log n}{\delta_p} \right),$$

which partially answers Question 5.5.

Remark 5.9. In the most of applications, p has order of $\Theta(1)$. The RHS is shortly written as $\tilde{O}\left(\frac{\|E\|}{\lambda_p} + \frac{\sqrt{r}}{\delta_p}\right)$. As discussed above, these two terms are inevitable. When $r = \Theta(n^\epsilon)$ for some $\epsilon > 0$, \sqrt{r} cannot be covered by \tilde{O} .

Finally, we address the general case when A has a high rank. In this case, we choose the parameter r (which is no longer the rank of A) following Theorem 2.12. Redefine

$$\epsilon_1 = \frac{\|E\|}{\lambda_p - \lambda_{r+1}}; \epsilon_2 = \frac{1}{\delta_p}; \eta := \frac{\|E\|^2}{(\lambda_p - \lambda_{r+1})\delta_p}.$$

$$s_{jj}^{\bar{p}} := \sum_{l>r} \frac{u_{lj}^2}{\lambda_{\bar{p}} - \lambda_l}, \text{ for each } 1 \leq j \leq r, 1 \leq \bar{p} \leq p.$$

Theorem 5.10. *Let A be a symmetric matrix and E be symmetric random noise with independent entries. Under Assumption **C2**, for any $t_1, t_2 > 0$, with probability at least $1 - r^2 \exp\left(\frac{-t_1^2/2}{2\sigma^2 + Kt_1}\right) - \frac{6r(r-1)pn\sigma^2(\sigma^2 + K^2)}{t_2^2}$,*

$$\left\| \tilde{\Pi}_p - \Pi_p \right\| \leq 12\sqrt{\bar{p}} \left(\epsilon_1 + \sqrt{r}t_1\epsilon_2 + \frac{\sqrt{r}(t_2 + \mu)}{\|E\|}\epsilon_1\epsilon_2 \right),$$

in which

$$\mu := \max_{\substack{1 \leq k < l \leq r \\ 1 \leq \bar{p} \leq p}} \left\{ \sum_i u_{ki}u_{li} \sum_j \sigma_{ji}^2 s_{jj}^{(\bar{p})} \right\}.$$

5.11. A toy application: Finding many hidden cliques. The hidden clique problem is a well-studied problem in random graph theory and theoretical computer science.

Problem. Hide a clique X of size k in the random graph $G(n, p)$. Find X in polynomial time.

Set $p = 1/2$, the largest clique in $G(n, 1/2)$, with overwhelming probability, has size approximately $2 \log n$. Thus for any $k > (2 + \epsilon) \log n$, with any constant $\epsilon > 0$, X would be abnormally large and therefore detectable, by brute-force at least. However, finding X in polynomial time is a different matter, and the best current bound for k is $k \geq c\sqrt{n}$ for any constant $c > 0$.

This result was first achieved by Alon, Krivelevich, and Sudakov [1]. They first find X when c is sufficiently large, then use a simple sampling trick to reduce the case of small c to this case. To find the clique for a large c , their algorithm essentially works as follows. Consider the adjacency matrix where we write 1 for an edge and -1 for a non-edge. (For simplicity we allow loops, and assume that the hidden clique has all the loops.) Compute the first eigenvector of this matrix and locate the first largest k coordinates in absolute value. Call this set Y . This is an approximation of the clique X . It is relatively simple to modify Y (by combinatorial means) to obtain the true set X . (For instance, one can define X to be those vertices which have at least $\frac{3}{4}k$ neighbors in Y .)

The critical first step relies on Davis-Kahan theorem. Let A be the matrix with 1 at an edge in the clique and 0 otherwise. Thus it has a $k \times k$ all-one block. It is clear that A has rank 1, with $\lambda_1 = k$, and $u_1 = \frac{1}{\sqrt{k}} I_{clq}$, where I_{clq} is the indicator vector of the clique. Thus, u_1 gives us the exact location of the vertices in the clique.

The adjacency matrix \tilde{A} (which is the input of the algorithm) is the sum of A and a random ± 1 matrix E . It is easy to show that E has norm $(2 + o(1))\sqrt{n}$, with probability $1 - o(1)$. By Theorem 1.5, we see that for $c = 200$, $\|\tilde{u}_1 \tilde{u}_1^T - u_1 u_1^T\| \leq \frac{1}{50}$. We can conclude, via a routine computation, that $|Y \Delta X| \leq \frac{k}{20}$ (with room to spare). This sufficiently small error will guarantee (via a simple combinatorial argument) that the cleaning step would identify X , with probability $1 - o(1)$. Thus, the heart of the matter here is to have $\|\tilde{u}_1 \tilde{u}_1^T - u_1 u_1^T\| \leq \frac{1}{50}$, or any sufficiently small constant ϵ , for that matter.

In the above setting, there is one clique of size $200\sqrt{n}$. Now we consider a more complex setting, where there are, say, 200 disjoint cliques of decreasing size $200\sqrt{n}, 199\sqrt{n}, \dots, 2\sqrt{n}, \sqrt{n}$, respectively. Again we would like to find the largest one. The matrix A now has 200 disjoint blocks and has rank 200, with $\lambda_1 = 200\sqrt{n}, \lambda_2 = 199\sqrt{n}, \dots, \lambda_{200} = \sqrt{n}$.

Let us point out that in this setting, Theorem 1.5 no longer applies, as the gap $\lambda_1 - \lambda_2 = \sqrt{n} \leq \|E\|$, so the bound in Theorem 1.5 becomes larger than 1, which is vacuous.

On the other hand, our new results guarantee that the previous algorithm still works, with no modification. Indeed, we can use Corollary 5.8. Choose $p = 1$ and $\alpha = \log^{-1} n = o(1)$ (any function tending to 0 with n). Notice that the second and last term on the RHS is only $o(1)$. Thus, the dominating term in the bound is $\frac{\|E\|}{\lambda_1}$, with the denominator being λ_1 (as in the case with only one clique discussed above), not $\lambda_1 - \lambda_2$ as Theorem 1.5 dictates. The rest of the analysis is similar to the argument presented above and omitted. We also refer to [13] for a new analysis showing that the cleaning step is not required, namely the large coordinates of \tilde{u}_1 are exactly the vertices of the clique.

6. SPIKED COVARIANCE MATRICES

6.1. The general covariance problem and spiked model. Let $X = [\xi_1 \ \xi_2 \ \dots \ \xi_d]^T$ be a random vector with covariance matrix $M = (m_{ij})_{1 \leq i, j \leq d}$, $m_{ij} = m_{ji} = \mathbf{Cov}(\xi_i, \xi_j)$. A fundamental problem in data science is to learn the unknown covariance matrix M using the sample covariance matrix $\tilde{M} := \frac{1}{n} \sum_{i=1}^n X_i X_i^T$, where X_i are iid samples of X .

Define $E = \tilde{M} - M$. It is clear that E is a mean zero random matrix (however its entries are in general not independent). In this section, we focus on the problem of learning the leading eigenvectors and eigenspaces of M , which are the core of Principal Component Analysis, a fundamental technique in many areas of sciences.

Formally, given the spectral decompositions of M and \tilde{M} ,

$$M = \sum_{i=1}^d \lambda_i u_i u_i^T = U \Lambda U^T, \quad \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d,$$

$$\text{and } \tilde{M} = \sum_{i=1}^d \tilde{\lambda}_i \tilde{u}_i \tilde{u}_i^T = \tilde{U} \tilde{\Lambda} \tilde{U}^T, \quad \tilde{\lambda}_1 \geq \dots \geq \tilde{\lambda}_d,$$

we want to estimate the differences

$$\|\tilde{u}_p \tilde{u}_p^T - u_p u_p^T\|, \quad \|\tilde{\Pi}_p - \Pi_p\|,$$

for some small p . For the sake of a simpler presentation, we will focus on eigenvector perturbation; the argument for eigenspace perturbation is similar.

Recently, researchers have paid great attention to spiked models; see, for instance, [7, 8, 9, 16, 35, 44, 45, 47, 54, 55] and the references therein. Informally speaking, in a spiked model, the matrix M has few important (large) eigenvalues (the "spikes"), while the remaining ones are small and comparable (see below for concrete examples). In this section, we are going to treat the following popular model

Spiked Population Model: $X = M^{1/2} Y$, where M is a spiked matrix and Y is d -dimensional random vector such that $\mathbf{E} Y Y^T = I_d$. Therefore,

$$\mathbf{E} X X^T = U \Lambda^{1/2} U^T (\mathbf{E} Y Y^T) U \Lambda^{1/2} U^T = U \Lambda U^T = M.$$

There have been two main approaches to analyze the behavior of these models.

The first approach is to compute the limiting behavior, inspired by works from random matrix theory [6, 12]. For this to be possible, one needs to assume that there is an infinite sequence of collections of samples $\{X_1^{(d)}, \dots, X_n^{(d)}\}_{n=1}^\infty$, where both d and n tends to infinity. Thus, there is a sequence of covariance matrices M_d and another sequence of sample covariance matrices \tilde{M}_d , with d tending to the infinity. Next, one computes the limit of the perturbations, for example: $\lim_{n, d \rightarrow \infty} \|\tilde{u}_p \tilde{u}_p^T - u_p u_p^T\|$. Under some conditions, researchers have computed this limit exactly, and so we have an idea about the asymptotic behavior of the perturbation. However, to make results of this kind useful in applications, one needs to estimate the rate of convergence. Furthermore, the assumptions required so techniques from random matrix theory (such as Stieltjes transform) can work are often too strict for applications.

In the second approach, one attacks the original problem with d, n fixed and large, and there is only one pair of M and \tilde{M} . The popular strategy here is to combine matrix perturbation results with Matrix Bernstein Inequality [74, 82]. These results are useful in applications and hold under rather mild assumptions. On the other hand, in many interesting cases, the bounds are weaker than the limiting results.

In what follows, we give a brief survey about these situations, and then present our new results, which can be viewed as a combination of “the best of two worlds”.

6.2. Existing bounds for Spiked Population Model. Let us consider a special case of the model, introduced by Johnstone, who initiated the study of spiked models [44]. In this model

- Y is d -dimensional random vector, whose entries are i.i.d with mean 0, variance 1 and finite 4^{th} moment.
- d and n are of the same order of magnitude: $\frac{d}{n} = \Theta(1)$.
- M has the “blocked” structure $\begin{pmatrix} \Lambda_r & 0 \\ 0 & I_{d-r} \end{pmatrix}$, in which Λ_r is an $r \times r$ matrix, whose eigenvalues $\Theta(1) = \lambda_1 > \lambda_2 > \dots > \lambda_r > 1$ are the spikes.

Thus he required that all eigenvalues are of the same order of magnitude (even the spiked ones), and also the gaps between the spikes have that same magnitude.

Approach 1: Limiting results

Assume that the ratio d/n tends to a positive constant γ as d, n tend to infinity. In [46], Johnstone proved

Theorem 6.3. *If $\lambda_p > 1 + \sqrt{\gamma}$, then almost surely*

$$(52) \quad \|\tilde{u}_p \tilde{u}_p^T - u_p u_p^T\| \rightarrow \sqrt{\frac{\lambda_p \gamma}{(\lambda_p - 1)^2 + (\lambda_p - 1)\gamma}}.$$

The RHS is a bit complicated. Let us consider the case where spike λ_p dominates γ (say the ratio γ/λ_p is sufficiently small), then the order of magnitude of the RHS is $\sqrt{\frac{\gamma}{\lambda_p}}$. So, at least on the heuristic level, we have

$$(53) \quad \|\tilde{u}_p \tilde{u}_p^T - u_p u_p^T\| \leq 2\sqrt{\frac{\gamma}{\lambda_p}}.$$

(As discussed earlier, to turn this into a rigorous bound, one needs to control the rate of convergence.)

Johnstone's spiked model is generalized by Bai, Yao, and Zhang, who replaced the identity matrix I_{d-r} by a general $(d-r) \times (d-r)$ matrix V_d , with two additional conditions

- (1) The ESD of V_d converges to a non-random limiting distribution H . Define $\varphi(\lambda) := \lambda + \gamma \int \frac{t\lambda}{\lambda-t} dH(t)$.
- (2) The eigenvalues of Λ_r ($\lambda_1 > \lambda_2 > \dots > \lambda_r$) are outside the support of H .

Under these conditions, they proved [10]

Theorem 6.4. *Assume that $\varphi'(\lambda_p) > 0$. Then almost surely,*

$$(54) \quad \|\tilde{u}_p \tilde{u}_p^T - u_p u_p^T\| \rightarrow \sqrt{1 - \frac{\lambda_p \varphi'(\lambda_p)}{\varphi(\lambda_p)}}.$$

Since $\varphi(\lambda_p) := \lambda_p + \gamma \int \frac{t\lambda_p}{\lambda_p-t} dH(t)$ and $\varphi'(\lambda_p) = 1 - \gamma \int \frac{t^2}{(\lambda_p-t)^2} dH(t)$, equivalently, the RHS is

$$(55) \quad \sqrt{\frac{\gamma}{\lambda_p} \times \frac{\int \frac{t\lambda_p}{\lambda_p-t} dH - \int \frac{t^2\lambda_p}{(\lambda_p-t)^2} dH}{1 + \gamma \int \frac{t}{\lambda_p-t} dH}}.$$

By a routine, but somewhat tedious computation, one can show that if λ_p dominates γ and $\max_{x \in \text{supp}H} |x|$, then similarly to (53), the LHS of (55) is also of order $\sqrt{\frac{\gamma}{\lambda_p}}$.

Remark 6.5. In Johnstone's spiked population model, $\varphi(\lambda_p) = \lambda_p + \frac{\gamma\lambda_p}{\lambda_p-1}$, $\varphi'(\lambda_p) = 1 - \frac{\gamma}{(\lambda_p-1)^2}$.

Approach 2: Direct Estimates. Here one starts with the classical Davis-Kahan theorem, and uses the Matrix Bernstein inequality to estimate $\|E\|$. We follow the analysis from Wainwright's book [82, Chapter 8].

In this approach, we do not need to make strong assumptions on the asymptotic of the spikes. In the limiting approach, the spikes all have to be order 1 (same as the non-spikes). In this approach, they are allowed to tend to infinity as functions of n (for instance $n^{1/4}$). This flexibility is important as spikes, by definition, are supposed to be considerably larger than then the average eigenvalues. Furthermore, in the limiting approach, the gaps between the spikes should also be of order 1 (so comparable to the spikes themselves). This condition can also be omitted, for instance, we can allow $\lambda_1 - \lambda_2 = O(n^{-1/4}\lambda_1)$.

Assume that there is a positive number L such that with probability 1, $\|XX^T - M\| \leq L$. Let $r_\lambda := \frac{\sum_{i=1}^d \lambda_i}{\lambda_1} = \frac{\text{Trace } M}{\lambda_1}$, be the *relative rank* of M . By Matrix Bernstein inequality

(Section 12.3), we have, with probability $1 - o(1)$

$$\begin{aligned}
 \|E\| &\leq \left(2 + \max_{1 \leq i \leq d} |\mathbf{E}y_i^4 - 3|\right) \left(\lambda_1 \sqrt{\frac{r_\lambda \log(n+d)}{n}} + L \log(n+d)\right) \\
 (56) \quad &= \tilde{\Omega} \left(\lambda_1 \sqrt{\frac{r_\lambda}{n}} + L\right) \\
 &= \tilde{\Omega} \left(\sqrt{\lambda_1 \gamma \frac{\text{Trace } M}{d}}\right).
 \end{aligned}$$

If $\text{Trace } M = \Theta(d)$, then

$$(57) \quad \|E\| = \tilde{\Omega}(\sqrt{\lambda_1 \gamma}).$$

Combining this with Davis-Kahan bound (Theorem 1.5), we obtain; for more details see [82, Chapter 8].

Theorem 6.6 (Eigenvector Perturbation). *Let $\delta_{(p)} = \min\{\delta_{p-1}, \delta_p\}$. Then with probability $1 - o(1)$*

$$(58) \quad \|\tilde{u}_p \tilde{u}_p^T - u_p u_p^T\| \leq 2 \frac{\|E\|}{\delta_{(p)}} = \tilde{\Omega} \left(\frac{\sqrt{\lambda_1 \gamma}}{\delta_{(p)}}\right).$$

It is a bit weird to have $\tilde{\Omega}$ on the RHS. We can turn it into (the more meaningful) $\tilde{\Theta}$ in the case the L term in (56) is negligible. Our point here that even when this term can be deleted, the bound is still off. Indeed, notice that we can rewrite the RHS of (58) as $\tilde{\Omega} \left(\frac{\sqrt{\lambda_1 \lambda_p}}{\delta_{(p)}} \sqrt{\frac{\gamma}{\lambda_p}}\right)$.

Thus, in comparison to (53), we are off by a factor $\frac{\sqrt{\lambda_1 \lambda_p}}{\delta_{(p)}}$, which can be large if the spikes and the gaps are of different order of magnitudes.

6.6.1. *New bounds for Spiked Population Model.* Following the above summary, the question that naturally arises is whether we can achieve the “best of two worlds”:

Question 6.7. *Can we obtain the optimal (up to a polylog term) bound $\tilde{\Omega} \left(\sqrt{\frac{\gamma}{\lambda_p}}\right)$ in the non-limit setting, without only modest assumptions on both the spikes and the gaps.*

Using Theorem 2.12, we give a positive answer to this question. We pay special attention to the case when the spikes tend to infinity as a function of n and d .

Theorem 6.8. *Assume that the model $X = M^{1/2}Y, Y = [y_1 \dots y_d]^T$ satisfies the following conditions:*

- *The entries $\{y_i\}_{i=1}^d$ are 8-wise independent random variables with mean 0, variance 1 and finite 4th moment.*

- $M = \begin{pmatrix} \Lambda_r & 0 \\ 0 & V_d \end{pmatrix}$, where r is a constant and the spikes $\lambda_r < \dots < \lambda_1 = O(n \wedge d)$ are eigenvalues of Λ_r . The non-spikes are eigenvalues of V_d and have of order 1.
- $p \leq r$ is a natural number and $\delta_{(p)} > 12 \max \left\{ \gamma, \frac{\lambda_1^{3/2}}{d^{1/2}} \right\}$.

Then, with probability $1 - o(1)$,

$$(59) \quad \|\tilde{u}_p \tilde{u}_p^T - u_p u_p^T\| = \tilde{O} \left(\sqrt{\frac{\gamma}{\lambda_p}} \right).$$

In this theorem, we allow the spikes to be as large as d (the dimension). The spikes need to be of the same magnitude, but the gap $\delta_{(p)}$ can be much smaller. For example, we can set $d = n^{1+\epsilon}$, $\delta = n^\epsilon$, $\lambda_i = n^{3\epsilon}$ for some constant $0 < \epsilon < \frac{1}{3}$. Our theorem gives a strong bound $n^{-\epsilon}$ in this case, while it seems the existing methods do not yield any useful information. Notice that in this setting, we can afford to have much less samples than dimension. Another new feature is that Theorem 6.8 does not require the coordinates of y be mutually independent.

Now we turn to relax the block structure of M .

Theorem 6.9. Assume that our spiked population model $X = M^{1/2}Y$, $Y = [y_1 \dots y_d]^T$ satisfies the following conditions:

- The entries $\{y_i\}_{i=1}^d$ are 8-wise independent random variables with mean 0, variance 1 and finite 4th moment.
- M is an arbitrary positive semi-definite matrix with r spikes - $O(n \wedge d) = \lambda_1 > \lambda_2 > \dots > \lambda_r$. All non-spikes eigenvalues have order 1.
- For some $p \leq r$, $\delta_{(p)} > 12 \max \left\{ \gamma, \frac{\lambda_1^{3/2}}{d^{1/2}} \right\}$.

Then, with probability $1 - o(1)$,

$$(60) \quad \|\tilde{u}_p \tilde{u}_p^T - u_p u_p^T\| = \tilde{O} \left(\sqrt{\frac{\gamma}{\lambda_p}} + \frac{c_r d}{\delta_{(p)} n} \right).$$

where

$$c_r := \max_{\substack{1 \leq i, j \leq r \\ l, l' > r}} \left| \sum_{k=1}^d (\mathbf{E}y_k^4 - 3) u_{ik} u_{jk} u_{l'k} u_{lk} \right|.$$

Remark 6.10. The extra term here (compared to the previous theorem) is $\frac{c_r d}{\delta_{(p)} n}$. Trivially $c_r \leq \max_k |\mathbf{E}y_k^4 - 3|$. Even with this trivial estimate, our bound is still meaningful if $d = o(\delta_{(p)} n)$. When the vectors $\{u_i\}_{1 \leq i \leq d}$ are delocalized, namely, their coordinates are of order $d^{-1/2+o(1)}$, then $c_r \leq \max_k |\mathbf{E}y_k^4 - 3| d^{-1+o(1)}$. In this case, the extra term becomes $\frac{\max_k |\mathbf{E}y_k^4 - 3| d^{o(1)}}{\delta_{(p)} n}$, which is often negligible.

Both theorems above are corollaries of the following general theorem. First, we need an analogue of Assumption **C0**

$$(61) \quad (\mathbf{C3}) : \max \left\{ \frac{\sqrt{p}\|E\|}{\lambda_p - \lambda_{r+1}}, \frac{r\lambda_1}{\delta_{(p)}\sqrt{n}}, \frac{\sqrt{r}\|E\|}{\sqrt{(\lambda_p - \lambda_{r+1})\delta_{(p)}}} \right\} < \frac{1}{12}.$$

The similar eigenspace perturbation result will be obtained under Assumption **C3'**:

$$(62) \quad (\mathbf{C3}') : \max \left\{ \frac{\sqrt{p}\|E\|}{\lambda_p - \lambda_{r+1}}, \frac{r\lambda_1}{\delta_p\sqrt{n}}, \frac{\sqrt{r}\|E\|}{\sqrt{(\lambda_p - \lambda_{r+1})\delta_p}} \right\} < \frac{1}{12}.$$

Theorem 6.11 (Eigenvectors perturbation). *For any given covariance matrix M , consider the spiked population model $X = M^{1/2}Y$, $Y = [y_1, \dots, y_d]^T$, in which y_i s are identity 8-wise independent with mean 0, variance 1 and finite 4th moment. There exists a universal constant $C > 0$ such that for each $1 \leq p \leq r$, under Assumption **C3**, with probability at least $1 - \frac{r^2}{t_1^2} - \frac{r^2 p}{t_2^2}$, we have*

(63)

$$\|\tilde{u}_p \tilde{u}_p^T - u_p u_p^T\| \leq C\sqrt{p} \left(\frac{\|E\|}{\lambda_p - \lambda_{r+1}} + \frac{\sqrt{r}\lambda_1 t_1}{\delta_{(p)}\sqrt{n}} + \frac{\sqrt{r}\sqrt{\lambda_1 \lambda_2} t_2}{\delta_{(p)}n} \frac{\sqrt{\sum_{l>r} \lambda_l^2}}{\lambda_p - \lambda_{r+1}} + \frac{\sqrt{r}(\sum_{l>r} \lambda_l)}{\delta_{(p)}} \frac{c_r \sqrt{\lambda_1 \lambda_2} t_2}{n} \right),$$

in which

$$c_r := \max_{\substack{1 \leq i, j \leq r \\ l, l' > r}} \left| \sum_{k=1}^d (\mathbf{E}y_k^4 - 3) u_{ik} u_{jk} u_{l'k} u_{lk} \right|.$$

Notice that in this theorem, we make no assumption on the spikes. The parameter r is chosen arbitrarily, M can have more than r spikes.

7. OTHER APPLICATIONS

7.1. Matrix completion. Let A be a large matrix, which we can only observe partially (say we can only see one per cent of its entries). A central problem in data science, the matrix completion problem, is to recover A , given the observed entries and some constrains on A . The Netflix prize problem is a famous example [41, 57], where the rows of A are viewers, the columns are movies, and the entries are ratings. Each viewer usually rates only a small number of movies, and we want to predict their ratings on the remaining ones. Any large recommendation system gives rise to a similar problem .

This problem is of huge importance with, clearly, very real applications; see, for example [2, 3, 4, 15, 20, 64, 70]. Of course, the task is only plausible if we have some constrains on A , and here a popular assumption is that A has low rank or approximately low rank. Another assumption, which is necessary, is that A has some kind of incoherence property; see the references above for definition.

If we assume that each entry could be observed with probability p (say, $p = .01$), independently, then we can form a matrix \tilde{A} as follows. Let write 0 for the entries that we cannot

observe, and scale up the observed entries by $1/p$ (for instance, 4 would become $4/p$). The resulting matrix \tilde{A} is a random matrix with independent entries and the expectation $\mathbf{E}\tilde{A} = A$. This way, we can write $\tilde{A} = A + E$, where A is approximately low rank and E is random with zero mean.

Researchers have been using spectral algorithms to attack the problem. The basic strategy is similar to one used for the hidden clique problem. One first computes a low rank approximation of \tilde{A} , and uses perturbation bounds to show that it approximates A in some norm. Next, one cleans up this approximation to obtain the truth; see [13, 19, 21, 23, 49].

To our best knowledge, most results obtained for the matrix completion problem have a dependence on κ , the ratio between the largest and smallest non-trivial singular values of A . The quality of the results decreases if κ gets larger; see the main results of [21, 49] for instance. In a new paper [13], this dependence is removed, but at the cost of the assumption that the non-trivial singular values of A are well separated. For the case when the entries of A are integer (such as in the Netflix problem), the algorithm in [13] works as follows

- Step 1. Compute the singular values of \tilde{A} . Threshold at some properly chosen parameter σ
- Step 2. Compute the low rank approximation A' of \tilde{A} with respect to the singular values larger than σ .
- Step 3. Round off all entries of A' to the nearest integer.

This is perhaps the simplest and fastest algorithm for the matrix completion problem. With the new tools presented in this paper, we can show that the performance of this algorithm does not depend on κ , and at the same time, the separation assumption on the singular value is not required either.

The analysis of the above algorithm shows that with high probability, $\|A' - A\|_\infty < 1/2$, which immediately leads to the desired result. The same idea was used in [13], but in their proof, the separation assumption is essential.

Using the contour expansion method, combined with some new ideas, we came up with a general scheme to obtain perturbation bound in the infinity norm. In particular, we can prove the above claim under much milder assumptions, compared to [13]. Details will appear in a future paper [73].

7.2. Differential Privacy. Computing with privacy is an important area in computer science. Let A be a data matrix, whose rows are indexed by individuals and may contain their sensitive data. We want to compute some functional $g(A)$ of A , but do not want to compromise the privacy of those individuals.

The theory of differential privacy has been developed to resolve this problem. We would not repeat a detailed introduction here and refer the readers to [31]. Very roughly speaking, the idea is to add random noise to A in order to protect sensitive data. (How much noise is added depends on the level of privacy we want to guarantee; again see [31] for a precise setting.)

The trade-off here is, of course, that the added random noise will alter the output $g(A)$. Estimating the output's accuracy is usually referred to as the "utility" problem, and is, obviously of critical importance.

It is clear that this problem is an idea application for our new method, as our new bounds are the most effective when the noise matrix E is random. As it requires an in-depth discussion in order to present the new results in this direction in a meaningful manner, we will present the details in a future paper [71].

8. PROOF OF THEOREM 2.12

We will use $N_{\tilde{\lambda}}(S)^c, S^c$ to denote the sets $\{1, 2, \dots, n\} \setminus N_{\tilde{\lambda}}(S)$ and $\{1, 2, \dots, n\} \setminus S$ respectively.

8.1. The simple spectrum assumption. It is more convenient for us to work with matrices having a simple spectrum (namely, all eigenvalues are different). We can make this assumption by the following simple trick. Instead of our matrix A (which may have multiple eigenvalues), we consider $A_\epsilon := A + \epsilon G$, where G is GOE (as a matter of fact, any random matrix with a continuous distribution would do). For any non-zero ϵ , A_ϵ has a simple spectrum with probability 1. Next, we bound the difference $\Pi_S \tilde{A}_\epsilon - \Pi_S A_\epsilon$ under the simple spectrum assumption. We then finish the proof by noticing that $\Pi_S A_\epsilon - \Pi_S A$ tends to zero with ϵ . (It is important here that there is no index $i \in S$ and $j \in S^c$ so that $\lambda_i = \lambda_j$.)

8.2. Contour Estimates. We will often use contour estimate for functions of the form $\frac{1}{(z-a)^l}$ for some positive integer l . As a corollary of (32), we have

Lemma 8.3. *Let a, b be two points not on the boundary of Γ . Then*

$$\frac{1}{2\pi i} \int_{\Gamma} \frac{1}{(z-a)(z-b)} dz = \frac{1}{b-a}$$

if exactly b lies inside Γ and a lies outside Γ . In all other cases, the integral equals zero.

To prove this lemma, notice that

$$(64) \quad \int_{\Gamma} \frac{1}{(z-a)(z-b)} dz = \frac{1}{b-a} \left(\int_{\Gamma} \frac{1}{z-a} dz - \int_{\Gamma} \frac{1}{z-b} dz \right).$$

The claim follows from (32).

8.4. Estimating the first term $\|F_1\|$. We follow the ideas leading up to (38). In this section, we estimate the first term $\|F_1\|$. To start, we split $(z-A)^{-1}$ into two parts, as follows

$$(z-A)^{-1} = \sum_{i=1}^n \frac{1}{z-\lambda_i} u_i u_i^T = P + Q,$$

where

$$(65) \quad P := \sum_{i \in N_{\tilde{\lambda}}(S)} \frac{u_i u_i^T}{z-\lambda_i},$$

$$(66) \quad Q := \sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{z - \lambda_j}.$$

For all $s \geq 1$, we can rewrite F_s as

$$(67) \quad F_s(E) = \int_{\Gamma} (P + Q)[E(P + Q)]^s dz.$$

In particular

$$F_1 = \int_{\Gamma} (P + Q)E(P + Q)dz = \int_{\Gamma} (PEP + QEP + PEQ + QEQ) dz.$$

Since Γ does not enclose any point from the set $\{\lambda_j\}_{j \in N_{\bar{\lambda}}(S)^c}$, $\int_{\Gamma} \frac{1}{(z - \lambda_j)(z - \lambda_{j'})} dz = 0$ for any pair $j, j' \in N_{\bar{\lambda}}(S)^c$. Therefore

$$(68) \quad \int_{\Gamma} QEQ dz = \sum_{j, j' \in N_{\bar{\lambda}}(S)^c} \int_{\Gamma} \frac{u_j u_j^T E u_{j'} u_{j'}^T}{(z - \lambda_j)(z - \lambda_{j'})} dz = 0.$$

Next, we have

$$\begin{aligned} \frac{1}{2\pi i} \int_{\Gamma} PEQ dz &= \frac{1}{2\pi i} \sum_{\substack{i \in N_{\bar{\lambda}}(S) \\ j \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)} u_i u_i^T E u_j u_j^T dz \\ &= \sum_{\substack{i \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{1}{\lambda_i - \lambda_j} u_i u_i^T E u_j u_j^T \text{ (by Lemma 8.3)} \\ &= \sum_{i \in S} u_i u_i^T E \left(\sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{\lambda_i - \lambda_j} \right). \end{aligned}$$

Therefore,

$$\begin{aligned}
(69) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} PEQ dz \right\| &= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i \in S} (\mathbf{v}^T u_i) \left(u_i^T E \left(\sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{\lambda_i - \lambda_j} \right) \mathbf{w} \right) \\
&\leq \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i \in S} |\mathbf{v}^T u_i| \left| u_i^T E \left(\sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{\lambda_i - \lambda_j} \right) \mathbf{w} \right| \\
&\leq \left(\max_{\|\mathbf{v}\|=1} \sum_{i \in S} |\mathbf{v}^T u_i| \right) \times \left\| E \left(\sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{\lambda_i - \lambda_j} \right) \right\| \\
&\leq \left(\max_{\|\mathbf{v}\|=1} \sum_{i \in S} |\mathbf{v}^T u_i| \right) \times \frac{\|E\|}{\bar{\lambda}} \\
&\leq \frac{\sqrt{p} \|E\|}{\bar{\lambda}} \text{ (by Cauchy-Schwarz and the fact that } |S| = p \text{)}.
\end{aligned}$$

The same estimate holds for $\| \int_{\Gamma} QEP dz \|$. Now we consider the last term $\int_{\Gamma} PEP dz$, which can be expressed as

$$\int_{\Gamma} \sum_{i,j \in N_{\bar{\lambda}}(S)} \frac{1}{(z - \lambda_i)(z - \lambda_j)} u_j u_j^T E u_i u_i^T dz = \int_{\Gamma} \sum_{i,j \in N_{\bar{\lambda}}(S)} \frac{1}{(z - \lambda_i)(z - \lambda_j)} u_j (u_j^T E u_i) u_i^T dz.$$

By Lemma 8.3 $|\int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)} dz| = \frac{2\pi}{|\lambda_j - \lambda_i|}$ if $\lambda_i \neq \lambda_j$ and exactly one of them lies inside Γ . In all other cases, the integral is zero. Furthermore, by Definition 2.2, $|u_j^T E u_i| \leq x$. We

obtain

$$\begin{aligned}
(70) \quad & \frac{1}{2\pi} \left\| \int_{\Gamma} PEP dz \right\| \\
&= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \left(\sum_{\substack{i \in S \\ j \in N_{\bar{\lambda}}(S) \setminus S}} \mathbf{v}^T u_i \frac{(u_i^T E u_j)}{\lambda_i - \lambda_j} u_j^T \mathbf{w} + \sum_{\substack{i \in S \\ j \in N_{\bar{\lambda}}(S) \setminus S}} \mathbf{v}^T u_j \frac{(u_j^T E u_i)}{\lambda_i - \lambda_j} u_i^T \mathbf{w} \right) \\
&\leq \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \left(\sum_{\substack{i \in S \\ j \in N_{\bar{\lambda}}(S) \setminus S}} |\mathbf{v}^T u_i| \left| \frac{(u_i^T E u_j)}{\lambda_i - \lambda_j} \right| |u_j^T \mathbf{w}| + \sum_{\substack{i \in S \\ j \in N_{\bar{\lambda}}(S) \setminus S}} |\mathbf{v}^T u_j| \left| \frac{(u_j^T E u_i)}{\lambda_i - \lambda_j} \right| |u_i^T \mathbf{w}| \right) \\
&\leq 2 \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{\substack{i \in S \\ j \in N_{\bar{\lambda}}(S) \setminus S}} |\mathbf{v}^T u_i| \left| \frac{(u_i^T E u_j)}{\lambda_i - \lambda_j} \right| |u_j^T \mathbf{w}| \\
&\leq \frac{2x}{\delta_S} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i \in S, j \in N_{\bar{\lambda}}(S) \setminus S} |\mathbf{v}^T u_i| |u_j^T \mathbf{w}| \\
&= \frac{2x}{\delta_S} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \left(\sum_{i \in S} |\mathbf{v}^T u_i| \right) \left(\sum_{j \in N_{\bar{\lambda}}(S) \setminus S} |u_j^T \mathbf{w}| \right) \\
&\leq \frac{2\sqrt{p(r-p)}x}{\delta_S} \text{ (by Cauchy-Schwarz).}
\end{aligned}$$

By (68), (69) (and the line following that), and (70), we obtain

$$(71) \quad \frac{1}{2\pi} \|F_1\| \leq \frac{2\sqrt{p}\|E\|}{\bar{\lambda}} + \frac{2\sqrt{pr}x}{\delta_S}.$$

8.5. Estimating the second term $\|F_2\|$. In this subsection, we bound $\|F_2\|$. The bounds here and from previous sections will give us all the terms in the bound of Theorem 2.12.

Similar to the last section, we consider the expansion

$$\begin{aligned}
F_2 &= \int_{\Gamma} (P+Q)E(P+Q)E(P+Q)dz \\
&= \left(\int_{\Gamma} QEQEQdz + \int_{\Gamma} QEPEQdz \right) + \left(\int_{\Gamma} PEQEPdz + \int_{\Gamma} PEPEPdz \right) + \\
&\quad + \int_{\Gamma} PEQEQdz + \int_{\Gamma} PEPEQdz + \int_{\Gamma} QEQEPdz + \int_{\Gamma} QEPEPdz.
\end{aligned}$$

Since Γ does not enclose any point from the set $\{\lambda_j\}_{j \in N_{\bar{\lambda}}(S)}^c$, $\int_{\Gamma} \frac{1}{(z-\lambda_{j_1})(z-\lambda_{j_2})(z-\lambda_{j_3})} dz = 0$ for any $j_1, j_2, j_3 \in N_{\bar{\lambda}}(S)^c$. Therefore

$$(72) \quad \int_{\Gamma} QEQEQdz = \sum_{j_1, j_2, j_3 \in N_{\bar{\lambda}}(S)^c} \int_{\Gamma} \frac{u_{j_1} u_{j_1}^T E u_{j_2} u_{j_2}^T E u_{j_3} u_{j_3}^T}{(z - \lambda_{j_1})(z - \lambda_{j_2})(z - \lambda_{j_3})} dz = 0.$$

Next, we have

$$\begin{aligned} \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} QEPEQdz &= \frac{1}{2\pi\mathbf{i}} \sum_{\substack{i \in N_{\bar{\lambda}}(S) \\ j, j' \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)(z - \lambda_{j'})} u_{j'} u_{j'}^T E u_i u_i^T E u_j u_j^T dz \\ &= \sum_{\substack{i \in S \\ j, j' \in N_{\bar{\lambda}}(S)^c}} \frac{1}{(\lambda_i - \lambda_j)(\lambda_i - \lambda_{j'})} u_{j'} u_{j'}^T E u_i u_i^T E u_j u_j^T \quad (\text{by Lemma 9.3}) \\ &= \sum_{i \in S} \left(\sum_{j' \in N_{\bar{\lambda}}(S)^c} \frac{u_{j'} u_{j'}^T}{\lambda_i - \lambda_{j'}} \right) E u_i u_i^T E \left(\sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{\lambda_i - \lambda_j} \right). \end{aligned}$$

Therefore,

$$(73) \quad \begin{aligned} \frac{1}{2\pi} \left\| \int_{\Gamma} QEPEQdz \right\| &\leq \sum_{i \in S} \left| \sum_{j' \in N_{\bar{\lambda}}(S)^c} \frac{u_{j'} u_{j'}^T}{\lambda_i - \lambda_{j'}} \right| \|E u_i\| \|u_i^T E\| \left| \sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{\lambda_i - \lambda_j} \right| \\ &\leq \frac{p \|E\|^2}{\bar{\lambda}^2} \\ &\leq \frac{1}{12} \frac{\sqrt{p} \|E\|}{\bar{\lambda}} \quad (\text{by Assumption D0}). \end{aligned}$$

The same estimates hold for $\frac{1}{2\pi} \left\| \int_{\Gamma} PEQEQdz \right\|$, $\frac{1}{2\pi} \left\| \int_{\Gamma} QEQEPdz \right\|$. Now, we consider the term $\int_{\Gamma} PEQEPdz$, which can be expressed as

$$\begin{aligned} \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} PEQEPdz &= \frac{1}{2\pi\mathbf{i}} \sum_{\substack{i, i' \in N_{\bar{\lambda}}(S) \\ j \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)(z - \lambda_{i'})} u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T dz \\ &= \frac{1}{2\pi\mathbf{i}} \sum_{\substack{i, i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)(z - \lambda_{i'})} u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T dz \\ &\quad + \frac{1}{2\pi\mathbf{i}} \sum_{\substack{i \in S \\ i' \in N_{\bar{\lambda}}(S) \setminus S \\ j \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)(z - \lambda_{i'})} u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T dz \\ &\quad + \frac{1}{2\pi\mathbf{i}} \sum_{\substack{i \in N_{\bar{\lambda}}(S) \setminus S \\ i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)(z - \lambda_{i'})} u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T dz \\ &\quad (\text{for the case } i, i' \in N_{\bar{\lambda}}(S) \setminus S, \text{ the integral } \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)(z - \lambda_{i'})} dz = 0). \end{aligned}$$

By Lemma 9.3, we can compute exactly the integral $\int_{\Gamma} \frac{1}{(z-\lambda_i)(z-\lambda_j)(z-\lambda_{i'})} dz$ for each group of (i, i', j) and obtain

$$\begin{aligned}
(74) \quad \frac{1}{2\pi i} \int_{\Gamma} PEQEP dz &= \sum_{\substack{i, i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{-1}{(\lambda_i - \lambda_j)(\lambda_{i'} - \lambda_j)} u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T \\
&+ \sum_{\substack{i \in S \\ i' \in N_{\bar{\lambda}}(S) \setminus S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{1}{(\lambda_i - \lambda_{i'}) (\lambda_i - \lambda_j)} u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T \\
&+ \sum_{\substack{i \in N_{\bar{\lambda}}(S) \setminus S \\ i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{1}{(\lambda_{i'} - \lambda_i) (\lambda_{i'} - \lambda_j)} u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T.
\end{aligned}$$

We bound the norm of the first sum as follows

$$\begin{aligned}
(75) \quad & \left\| \sum_{\substack{i, i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{-1}{(\lambda_i - \lambda_j)(\lambda_{i'} - \lambda_j)} u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T \right\| \\
&= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{\substack{i, i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{\mathbf{v}^T u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T \mathbf{w}}{(\lambda_i - \lambda_j)(\lambda_{i'} - \lambda_j)} \\
&= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i, i' \in S} \mathbf{v}^T u_{i'} u_{i'}^T E \left(\sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{(\lambda_i - \lambda_j)(\lambda_{i'} - \lambda_j)} \right) E u_i u_i^T \mathbf{w} \\
&\leq \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i, i' \in S} |\mathbf{v}^T u_{i'}| \left| u_{i'}^T E \left(\sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{(\lambda_i - \lambda_j)(\lambda_{i'} - \lambda_j)} \right) E u_i \right| |u_i^T \mathbf{w}| \\
&\leq \frac{p \|E\|^2}{\lambda^2} \text{(by the definition of } \bar{\lambda} \text{ and Cauchy-Schwarz)}.
\end{aligned}$$

Next, we bound the norm of the second sum

$$\begin{aligned}
& \left\| \sum_{\substack{i \in S \\ i' \in N_{\bar{\lambda}}(S) \setminus S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{1}{(\lambda_i - \lambda_{i'}) (\lambda_i - \lambda_j)} u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T \right\| \\
&= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{\substack{i \in S \\ i' \in N_{\bar{\lambda}}(S) \setminus S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{\mathbf{v}^T u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T \mathbf{w}}{(\lambda_i - \lambda_{i'}) (\lambda_i - \lambda_j)} \\
(76) \quad &= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{\substack{i \in S \\ i' \in N_{\bar{\lambda}}(S) \setminus S}} \frac{\mathbf{v}^T u_{i'} u_{i'}^T E}{\lambda_i - \lambda_{i'}} \left(\sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{\lambda_i - \lambda_j} \right) E u_i u_i^T \mathbf{w} \\
&\leq \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{\substack{i \in S \\ i' \in N_{\bar{\lambda}}(S) \setminus S}} \frac{|\mathbf{v}^T u_{i'}|}{\lambda_i - \lambda_{i'}} \left| u_{i'}^T E \left(\sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{\lambda_i - \lambda_j} \right) E u_i \right| |u_i^T \mathbf{w}| \\
&\leq \frac{y}{\delta_S} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{\substack{i \in S \\ i' \in N_{\bar{\lambda}}(S) \setminus S}} |\mathbf{v}^T u_{i'}| \cdot |u_i^T \mathbf{w}| \\
&\leq \frac{\sqrt{p r} y}{\delta_S} \text{ (by Cauchy-Schwartz)}.
\end{aligned}$$

A similar estimate holds for the norm of the third sum. Putting the bounds together, we obtain

$$(77) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} P E Q E P dz \right\| \leq \frac{p \|E\|^2}{\bar{\lambda}^2} + \frac{2\sqrt{p r} y}{\delta_S}.$$

Now we consider the fourth term $\int_{\Gamma} P E P E P dz$. By expansion, we have

$$\int_{\Gamma} P E P E P dz = \sum_{i_1, i_2, i_3 \in N_{\bar{\lambda}}(S)} \int_{\Gamma} \frac{u_{i_1} u_{i_1}^T E u_{i_2} u_{i_2}^T E u_{i_3} u_{i_3}^T}{(z - \lambda_{i_1})(z - \lambda_{i_2})(z - \lambda_{i_3})} dz.$$

Notice that the integral is non-zero when there is one or two of $\{i_1, i_2, i_3\}$ inside the contour Γ . Clearly, there are 6 such scenarios, and we split $\int_{\Gamma} P E P E P dz$ into 6 corresponding subsums. We will bound the norm of each subsum by $\frac{\sqrt{p r} x^2}{\delta_S^2}$. We present here the treatment for the case $i_1 \in S$ and $i_2, i_3 \in N_{\bar{\lambda}}(S) \setminus S$; the treatment of other cases are similar. Consider

$$\begin{aligned}
& \left\| \sum_{\substack{i_1 \in S \\ i_2, i_3 \in N_{\bar{\lambda}}(S) \setminus S}} \int_{\Gamma} \frac{u_{i_1} u_{i_1}^T E u_{i_2} u_{i_2}^T E u_{i_3} u_{i_3}^T}{(z - \lambda_{i_1})(z - \lambda_{i_2})(z - \lambda_{i_3})} dz \right\| \\
&= \left\| \sum_{\substack{i_1 \in S \\ i_2, i_3 \in N_{\bar{\lambda}}(S) \setminus S}} \frac{u_{i_1} u_{i_1}^T E u_{i_2} u_{i_2}^T E u_{i_3} u_{i_3}^T}{(\lambda_{i_1} - \lambda_{i_2})(\lambda_{i_1} - \lambda_{i_3})} \right\| \\
&= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{\substack{i_1 \in S \\ i_2, i_3 \in N_{\bar{\lambda}}(S) \setminus S}} \frac{\mathbf{v}^T u_{i_1} u_{i_1}^T E u_{i_2} u_{i_2}^T E u_{i_3} u_{i_3}^T \mathbf{w}}{(\lambda_{i_1} - \lambda_{i_2})(\lambda_{i_1} - \lambda_{i_3})} \\
&\leq \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{\substack{i_1 \in S \\ i_2, i_3 \in N_{\bar{\lambda}}(S) \setminus S}} \frac{|\mathbf{v}^T u_{i_1}| |u_{i_1}^T E u_{i_2}| |u_{i_2}^T E u_{i_3}| |u_{i_3}^T \mathbf{w}|}{(\lambda_{i_1} - \lambda_{i_2})(\lambda_{i_1} - \lambda_{i_3})} \\
&\leq \frac{rx^2}{\delta_S^2} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{\substack{i_1 \in S \\ i_3 \in N_{\bar{\lambda}}(S) \setminus S}} |\mathbf{v}^T u_{i_1}| |u_{i_3}^T \mathbf{w}| \\
&\leq \frac{\sqrt{pr}rx^2}{\delta_S^2} \text{(by Cauchy-Schwarz)}.
\end{aligned}$$

Therefore,

$$(78) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} PEPEP dz \right\| \leq \frac{6\sqrt{pr}rx^2}{\delta_S^2}.$$

Finally, we bound $\int_{\Gamma} PEPEQ dz$. Again, we expand the product and obtain

$$\begin{aligned}
\frac{1}{2\pi\mathbf{i}} \int_{\Gamma} PEPEQ dz &= \frac{1}{2\pi\mathbf{i}} \sum_{\substack{i, i' \in N_{\bar{\lambda}}(S) \\ j \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)(z - \lambda_{i'})} u_{i'} u_{i'}^T E u_i u_i^T E u_j u_j^T dz \\
&= \frac{1}{2\pi\mathbf{i}} \sum_{\substack{i, i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)(z - \lambda_{i'})} u_{i'} u_{i'}^T E u_j u_j^T E u_i u_i^T dz \\
&\quad + \frac{1}{2\pi\mathbf{i}} \sum_{\substack{i \in S \\ i' \in N_{\bar{\lambda}}(S) \setminus S \\ j \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)(z - \lambda_{i'})} u_{i'} u_{i'}^T E u_i u_i^T E u_j u_j^T dz \\
&\quad + \frac{1}{2\pi\mathbf{i}} \sum_{\substack{i \in N_{\bar{\lambda}}(S) \setminus S \\ i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{1}{(z - \lambda_i)(z - \lambda_j)(z - \lambda_{i'})} u_{i'} u_{i'}^T E u_i u_i^T E u_j u_j^T dz.
\end{aligned}$$

Applying Lemma 9.3 to compute the integral, we obtain

$$\begin{aligned}
(79) \quad \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} PEPEQdz &= \sum_{\substack{i,i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{-1}{(\lambda_i - \lambda_j)(\lambda_{i'} - \lambda_j)} u_{i'} u_{i'}^T E u_i u_i^T E u_j u_j^T \\
&+ \sum_{\substack{i \in S \\ i' \in N_{\bar{\lambda}}(S) \setminus S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{1}{(\lambda_i - \lambda_{i'})(\lambda_i - \lambda_j)} u_{i'} u_{i'}^T E u_i u_i^T E u_j u_j^T \\
&+ \sum_{\substack{i \in N_{\bar{\lambda}}(S) \setminus S \\ i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{1}{(\lambda_{i'} - \lambda_i)(\lambda_{i'} - \lambda_j)} u_{i'} u_{i'}^T E u_i u_i^T E u_j u_j^T.
\end{aligned}$$

We bound the norm of the first sum as follows

$$\begin{aligned}
(80) \quad &\left\| \sum_{\substack{i,i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{-1}{(\lambda_i - \lambda_j)(\lambda_{i'} - \lambda_j)} u_{i'} u_{i'}^T E u_i u_i^T E u_j u_j^T \right\| \\
&= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{\substack{i,i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{1}{(\lambda_i - \lambda_j)(\lambda_{i'} - \lambda_j)} \mathbf{v}^T u_{i'} u_{i'}^T E u_i u_i^T E u_j u_j^T \mathbf{w} \\
&= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i,i' \in S} \mathbf{v}^T u_{i'} u_{i'}^T E u_i u_i^T E \left(\sum_{j \in N_{\bar{\lambda}}^c} \frac{u_j u_j^T}{(\lambda_i - \lambda_j)(\lambda_{i'} - \lambda_j)} \right) \mathbf{w} \\
&\leq \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i,i' \in S} |\mathbf{v}^T u_{i'}| |u_{i'}^T E u_i| |u_i^T E| \left| \sum_{j \in N_{\bar{\lambda}}^c} \frac{u_j u_j^T}{(\lambda_i - \lambda_j)(\lambda_{i'} - \lambda_j)} \mathbf{w} \right| \\
&\leq \frac{px\|E\|}{\bar{\lambda}^2} \max_{\|\mathbf{v}\|=1} \sum_{i' \in S} |\mathbf{v}^T u_{i'}| \\
&\leq \frac{px}{\bar{\lambda}} \times \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \text{(by Cauchy-Schwarz)}.
\end{aligned}$$

By a similar argument, we also obtain

$$\left\| \sum_{\substack{i \in S \\ i' \in N_{\bar{\lambda}}(S) \setminus S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{1}{(\lambda_i - \lambda_{i'})(\lambda_i - \lambda_j)} u_{i'} u_{i'}^T E u_i u_i^T E u_j u_j^T \right\| \leq \frac{\sqrt{p}rx}{\delta_S} \times \frac{\sqrt{p}\|E\|}{\bar{\lambda}},$$

and

$$\left\| \sum_{\substack{i \in N_{\bar{\lambda}}(S) \setminus S \\ i' \in S \\ j \in N_{\bar{\lambda}}(S)^c}} \frac{1}{(\lambda_{i'} - \lambda_i)(\lambda_{i'} - \lambda_j)} u_{i'} u_{i'}^T E u_i u_i^T E u_j u_j^T \right\| \leq \frac{rx}{\delta_S} \times \frac{\sqrt{p} \|E\|}{\lambda}.$$

Therefore, by Assumption **D0** and the fact that $\delta \leq \delta_S$, we have

$$(81) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} PEPEQ dz \right\| \leq 3 \times \frac{1}{12} \times \frac{\sqrt{p} \|E\|}{\lambda} = \frac{1}{4} \frac{\sqrt{p} \|E\|}{\lambda}.$$

The same estimate holds for $\frac{1}{2\pi} \left\| \int_{\Gamma} QEPEP dz \right\|$.

Combining (72), (73) (and the line following that), (77), (78), and (81) (and the line following that), we obtain a bound for $\|F_2\|$

$$(82) \quad \begin{aligned} \frac{1}{2\pi} \|F_2\| &\leq \frac{3}{12} \frac{\sqrt{p} \|E\|}{\bar{\lambda}} + \frac{p \|E\|^2}{\bar{\lambda}^2} + \frac{2\sqrt{pr}y}{\delta_S} + \frac{6\sqrt{pr}rx^2}{\delta_S^2} + \frac{2}{4} \frac{\sqrt{p} \|E\|}{\bar{\lambda}} \\ &\leq \left(\frac{3}{12} + \frac{\sqrt{p} \|E\|}{\bar{\lambda}} + \frac{1}{2} \right) \frac{\sqrt{p} \|E\|}{\bar{\lambda}} + \frac{6rx}{\delta_S} \times \frac{\sqrt{pr}x}{\delta_S} + \frac{2\sqrt{pr}y}{\delta_S} \\ &\leq \frac{\sqrt{p} \|E\|}{\bar{\lambda}} + \frac{\sqrt{pr}x}{2\delta_S} + \frac{2\sqrt{pr}y}{\delta_S} \quad (\text{since } \frac{\sqrt{p} \|E\|}{\bar{\lambda}}, \frac{rx}{\delta_S} \leq \frac{1}{12} \text{ by Assumption } \mathbf{D0}) \\ &\leq 2\sqrt{p} \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{r}x}{\delta_S} + \frac{\sqrt{r}y}{\delta_S} \right). \end{aligned}$$

8.6. Estimating $\|F_s\|$ for a general s . In this subsection, we present three technical lemmas for estimating $\|F_s\|$, with a general s . These lemmas are the core of the proof. Combining these lemmas with (38), we obtain a proof of Theorem 2.12, which we present in the next subsection. The proofs of the lemmas are involved and will be presented in the next section.

Following (67), we expand $(P+Q)[E(P+Q)]^s$ into the sum of 2^{s+1} operators, each of which is a product of alternating Q -blocks and P -blocks

$$(QEQE \dots QE)(PEPE \dots PE) \dots (PEPE \dots PE)(QEQE \dots Q),$$

where we allow the first and last blocks to be empty.

We code each operator like this by the numbers of Q 's and the numbers of P 's in each blocks. If there are $(k+1)$ Q -blocks and k P -blocks, for some integer k , we let $\alpha_1, \dots, \alpha_{k+1}$ and β_1, \dots, β_k be these numbers. These numbers satisfy the following conditions

$$\begin{aligned} \alpha_1, \alpha_{k+1} &\geq 0, \\ \alpha_i, \beta_j &\geq 1, \quad \forall 1 < i < k+1, \quad \forall 1 \leq j \leq k, \\ \alpha_1 + \dots + \alpha_{k+1} + \beta_1 + \dots + \beta_k &= s+1. \end{aligned}$$

In what follows, we set $\alpha = (\alpha_1, \dots, \alpha_{k+1}) \in \mathbb{Z}^{k+1}$ and $\beta = (\beta_1, \dots, \beta_k) \in \mathbb{Z}^k$, and use $M(\alpha; \beta)$ to denote the corresponding operator.

Example: given $s = 10, k = 2, \alpha_1 = 3, \alpha_2 = 2, \alpha_3 = 1, \beta_1 = 2, \beta_2 = 3$, we have

$$\int_{\Gamma} M(3, 2, 1; 2, 3) dz = \int_{\Gamma} \underbrace{(QEQQEQE)}_{\substack{\alpha_1=3 \\ \text{numbers of } Q}} \underbrace{(PEPE)}_{\substack{\beta_1=2 \\ \text{numbers of } P}} \underbrace{(QEQQE)}_{\substack{\alpha_2=2 \\ \text{numbers of } Q}} \underbrace{(PEPEPE)}_{\substack{\beta_2=3 \\ \text{numbers of } P}} \underbrace{Q}_{\substack{\alpha_3=1 \\ \text{numbers of } Q}} dz.$$

We are going to bound $\left\| \int_{\Gamma} M(\alpha, \beta) dz \right\|$ for each pair α, β separately, and use the triangle inequality to add up the bounds. For s and k fixed, we split the collection of pairs $(\alpha; \beta)$ into the following types, according to the values of α_1 and α_{k+1}

- Type I: $\alpha_1, \alpha_{k+1} > 0$.
- Type II: $\alpha_1, \alpha_{k+1} = 0$.
- Type III: either $(\alpha_1 = 0, \alpha_{k+1} > 0)$ or $(\alpha_1 > 0, \alpha_{k+1} = 0)$.

Let $s_1 := \sum_{i=1}^{k+1} \alpha_i$ and $s_2 := \sum_{j=1}^k \beta_j$. We prove three lemmas bounding $\left\| \int_{\Gamma} M(\alpha, \beta) dz \right\|$ with respect to the above 3 types. We assume that Assumption **D0** holds in all of three lemmas.

Lemma 8.7. *For a pair (α, β) of Type I,*

$$(83) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| \leq \left(\frac{\sqrt{p} \|E\|}{\bar{\lambda}} + \frac{px}{\bar{\lambda}} \right) \frac{2^{s_2+s-1}}{12^{s-1}}.$$

Lemma 8.8. *For a pair (α, β) of Type II,*

- If $M(\alpha; \beta) \neq PEQEPE\dots QEP$, i.e., $(s_1, s_2) \neq (k-1, k)$, then

$$(84) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| \leq \left(\frac{\sqrt{p} \|E\|}{\bar{\lambda}} + \frac{\sqrt{pr}x}{\delta_S} \right) \frac{2^{s_2+s-1}}{12^{s-1}},$$

- If $M(\alpha; \beta) = PEQEPEQE\dots EQEP$, i.e., $(s_1, s_2) = (k-1, k)$, then

$$(85) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} PEQEPEQE\dots EQEP dz \right\| \leq \left(\frac{px}{\bar{\lambda}} + \frac{\sqrt{p} \|E\|}{\bar{\lambda}} \right) \frac{2^{3s/2}}{12^{s-1}} + \frac{(s+2)\sqrt{pr}y}{\delta_S} \left(\frac{2\sqrt{rw}}{\sqrt{\bar{\lambda}\delta_S}} \right)^{s-2}.$$

Lemma 8.9. *For a pair (α, β) of Type III,*

$$(86) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| \leq \left(\frac{\sqrt{p} \|E\|}{\bar{\lambda}} + \frac{px}{\bar{\lambda}} \right) \frac{2^{s_2+s-1}}{12^{s-1}}.$$

8.10. Proof of Theorem 2.12 using the lemmas. For any integer $1 \leq s_2 \leq s$, the expansion of F_s contains $\binom{s+1}{s_2}$ operators $M(\alpha; \beta)$, each of which has exactly s_2 P operators. A simple consideration reveals that among these, $\binom{s-1}{s_2}$ are of Type I, $\binom{s-1}{s_2-2}$ are of Type II, and $2\binom{s-1}{s_2-1}$ are of Type III. As shown in Section 3, the only term in the expansion with $s_2 = 0$ vanishes by Cauchy's theorem.

Putting (83), (84), (85), (86) together and summing over s_2 , we show that $\frac{1}{2\pi}\|F_s\|$ is at most

$$\left[\sum_{s_2=1}^{s+1} \binom{s-1}{s_2} \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \frac{2^{s_2+s-1}}{12^{s-1}} \right] + \left[\left(\frac{px}{\lambda} + \frac{\sqrt{p}\|E\|}{\lambda} \right) \frac{2^{3s/2}}{12^{s-1}} + \frac{(s+2)\sqrt{pr}y}{\delta_S} \left(\frac{2\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right)^{s-2} \right] + \left[\sum_{s_2=1}^{s+1} \binom{s-1}{s_2-2} \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{\sqrt{pr}x}{\delta_S} \right) \frac{2^{s_2+s-1}}{12^{s-1}} \right] + \left[\sum_{s_2=1}^{s+1} 2 \binom{s-1}{s_2-1} \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \times \frac{2^{s_2+s-1}}{12^{s-1}} \right],$$

here $\binom{s-1}{l} = 0$ for either $l > s-1$ or $l < 0$. Consider the first term

$$\begin{aligned} \sum_{s_2=1}^{s+1} \binom{s-1}{s_2} \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \frac{2^{s_2+s-1}}{12^{s-1}} &= \sum_{s_2=1}^{s+1} \binom{s-1}{s_2} 2^{s_2} \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \frac{2^{s-1}}{12^{s-1}} \\ &= \left(\sum_{s_2=1}^{s+1} \binom{s-1}{s_2} 2^{s_2} 1^{s-1-s_2} \right) \times \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \frac{2^{s-1}}{12^{s-1}} \\ &= 3^{s-1} \times \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \frac{2^{s-1}}{12^{s-1}} \\ &= \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \frac{1}{2^{s-1}}. \end{aligned}$$

Using a similar argument, we have

$$\sum_{s_2=1}^{s+1} \binom{s-1}{s_2-2} \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{\sqrt{pr}x}{\delta_S} \right) \frac{2^{s_2+s-1}}{12^{s-1}} = 4 \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{\sqrt{pr}x}{\delta_S} \right) \frac{1}{2^{s-1}},$$

(here we pair up 2^{s_2-2} with $\binom{s-1}{s_2-2}$, the remaining factor is 4), and

$$\sum_{s_2=1}^{s+1} 2 \binom{s-1}{s_2-1} \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \times \frac{2^{s_2+s-1}}{12^{s-1}} = 4 \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \times \frac{1}{2^{s-1}},$$

(we pair up 2^{s_2-1} with $\binom{s-1}{s_2-1}$, the remaining factor is $2 \times 2 = 4$). Finally, we have

$$\left[\left(\frac{px}{\lambda} + \frac{\sqrt{p}\|E\|}{\lambda} \right) \frac{2^{3s/2}}{12^{s-1}} + \frac{(s+2)\sqrt{pr}y}{\delta_S} \left(\frac{2\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right)^{s-2} \right] \leq \left[\frac{2}{3} \left(\frac{px}{\lambda} + \frac{\sqrt{p}\|E\|}{\lambda} \right) \frac{1}{4^{s-2}} + \frac{\sqrt{pr}y}{\delta_S} \frac{s+2}{6^{s-2}} \right]$$

where we use Assumption **D0** (and the fact that $\delta \leq \delta_S$) to bound $\frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S}$ and the trivial estimate that $\frac{2^{3/2}}{12} \leq \frac{3}{12} = \frac{1}{4}$. It follows that

$$(87) \quad \begin{aligned} \frac{1}{2\pi} \|F_s\| &\leq \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \frac{1}{2^{s-1}} + \left[\frac{2}{3} \left(\frac{px}{\lambda} + \frac{\sqrt{p}\|E\|}{\lambda} \right) \frac{1}{4^{s-2}} + \frac{\sqrt{pr}y}{\delta_S} \frac{s+2}{6^{s-2}} \right] + \\ &\quad + 4 \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{\sqrt{pr}x}{\delta_S} \right) \frac{1}{2^{s-1}} + 4 \left(\frac{\sqrt{p}\|E\|}{\lambda} + \frac{px}{\lambda} \right) \frac{1}{2^{s-1}}. \end{aligned}$$

The RHS converges. By summing (over s), we can bound $\frac{1}{2\pi} \sum_{s \geq 2} \|F_s\|$ by

$$(88) \quad 10 \frac{\sqrt{p} \|E\|}{\bar{\lambda}} + 4 \frac{\sqrt{prx}}{\delta_S} + 6 \frac{px}{\bar{\lambda}} + \frac{26}{5} \frac{\sqrt{pry}}{\delta_S}.$$

(Here we omit the routine calculation and do not try to optimize the constants.) Combining (38), (71), (88), we obtain

$$(89) \quad \|\Pi_S - \tilde{\Pi}_S\| \leq \frac{2\sqrt{p} \|E\|}{\bar{\lambda}} + \frac{2\sqrt{prx}}{\delta_S} + \left(10 \frac{\sqrt{p} \|E\|}{\bar{\lambda}} + 6 \frac{px}{\bar{\lambda}} + 4 \frac{\sqrt{prx}}{\delta_S} + \frac{26}{5} \frac{\sqrt{pry}}{\delta_S} \right).$$

Since $\frac{px}{\bar{\lambda}} \leq \frac{\sqrt{prx}}{\delta_S}$, (89) implies

$$\|\Pi_S - \tilde{\Pi}_S\| \leq \frac{12\sqrt{p} \|E\|}{\bar{\lambda}} + \frac{12\sqrt{prx}}{\delta_S} + 6 \frac{\sqrt{pry}}{\delta_S} \leq 12\sqrt{p} \left(\frac{\|E\|}{\bar{\lambda}} + \frac{\sqrt{rx}}{\delta_S} + \frac{\sqrt{ry}}{\delta_S} \right),$$

completing the proof.

9. PROOFS OF THE MAIN LEMMAS

In this section, we prove the three main lemmas, Lemma 8.7, Lemma 8.8, and Lemma 8.9.

9.1. A combinatorial lemma. An important part of our analysis is to compute the contour integral

$$\int_{\Gamma} \prod_{j=1}^l \frac{1}{z - \lambda_{i_j}},$$

for a sequence of indices i_1, \dots, i_l . There are many ways to do this, which could lead to different looking (but equivalent) formulae.

The result in this section provides a combinatorial way to compute the integral. This will be essential in regrouping the terms in after the expansion.

We start with a few definitions. Fix a natural number s . Given a parameter $0 \leq T \leq s + 1$, consider two sequences of (not necessarily different) indices: $X = [i_1, \dots, i_T]$ and $Y = [i_{T+1}, \dots, i_{s+1}]$.

Let $\mathcal{L}(T, s)$ denote the set of all sequences of natural numbers m_1, m_2, \dots, m_T such that $m_1 + \dots + m_T = s$.

For each sequence $L = m_1, \dots, m_T \in \mathcal{L}(T, s)$, we define a directed bipartite graph $G(X, Y|L)$, with X and Y being the vertex sets, as follow:


- The edges go from X to Y : $\forall e \in \mathcal{E}(X, Y|L)$, $e = e^+ \rightarrow e^- : e^+ \in X$ and $e^- \in Y$.

- For each $1 \leq j \leq T$, i_j in X is connected to $i_{j'}$ in Y , for all j' in the interval

$$[(s+1) - ((m_1 - 1) + \dots + (m_j - 1)), (s+1) - ((m_1 - 1) + \dots + (m_{j-1} - 1))].$$

In particular, i_1 is connected to all $i_{j'}$ with j' in $[(s+1) - (m_1 - 1), s+1]$. Thus, if $m_1 = 1$, then i_1 is connected to only i_{s+1} . It is clear that given T and L , the graph is defined up to isomorphism. Denote the set of (directed) edges of the graph $G(X, Y|L)$ by $\mathcal{E}(X, Y|L)$.

Example: Given two sequences of indices: $X = [1, 1, 2]$ and $Y = [3, 3, 9]$. We have $T = 3$, $s = 5$, $i_1 = 1, i_2 = 1, i_3 = 2, i_4 = 3, i_5 = 3, i_6 = 9$ and a sequence $(m_1, m_2, m_3) := (1, 1, 3)$, we obtain:



Remark 9.2. $G(X, Y|L)$ has exactly s edges and for every vertex j in Y , there is at least one edge ending at j .

In what follows, we fix two sequences X, Y of indices and the parameter T . Thus, for each sequence $L \in \mathcal{L}(T, s)$, we obtain a graph $G(X, Y|L)$. Let $\lambda_{i_1}, \dots, \lambda_{i_{s+1}}$ be the sequence of eigenvalues corresponding to the sequence of indices i_1, \dots, i_{s+1} . Define

$$w(X, Y|L) := \prod_{e \in \mathcal{E}(X, Y|L)} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})}.$$

If either X or Y is the empty set, we simply set $w(X, Y|L) = 0$ for any $L \in \mathcal{L}(T, s)$.

Example: For $X = [1, 1, 2]$ and $Y = [3, 3, 9]$ and $L = (m_1, m_2, m_3) = (1, 1, 3)$

$$w(X, Y|L) = \frac{1}{(\lambda_1 - \lambda_9)(\lambda_1 - \lambda_9)(\lambda_2 - \lambda_9)(\lambda_2 - \lambda_3)(\lambda_2 - \lambda_3)}.$$

Lemma 9.3. Consider a contour Γ , a parameter $0 \leq T \leq s+1$, and two sequences $X = [i_1, \dots, i_T], Y = [i_{T+1}, \dots, i_{s+1}]$ of indices. Assume that $\lambda_{i_1}, \dots, \lambda_{i_T}$ are inside the contour Γ and $\lambda_{i_{T+1}}, \dots, \lambda_{i_{s+1}}$ are outside. Then,

$$(90) \quad \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} \frac{1}{\prod_{j=1}^{s+1} (z - \lambda_{i_j})} dz = (-1)^{T+1} \sum_{L \in \mathcal{L}(T,s)} w(X, Y|L).$$

Proof of Lemma 9.3. We use induction on s . In the base case $s = 1$, T is either 0, 1 or 2, and the claim follows from Lemma 8.3.

Now assume that for all $1 \leq r \leq s$, the induction hypothesis holds. This means that given any two sequences $X_1 = [i'_1, i'_2, \dots, i'_{T_1}]$, $Y_1 = [i'_{T_1+1}, \dots, i'_r]$ of indices (their total length is at most s) such that $\lambda_{i'_1}, \dots, \lambda_{i'_{T_1}}$ are inside the contour Γ and $\lambda_{i'_{T_1+1}}, \dots, \lambda_{i'_r}$ are outside,

$$\frac{1}{2\pi\mathbf{i}} \int_{\Gamma} \frac{1}{\prod_{j=1}^r (z - \lambda_{i'_j})} dz = (-1)^{T_1+1} \sum_{L \in \mathcal{L}(T_1, r-1)} w(X_1, Y_1|L).$$

We will prove for two sequences $X = [i_1, \dots, i_T]$, $Y = [i_{T+1}, \dots, i_{s+1}]$ (which have $s+1$ indices in total) and a parameter $0 \leq T \leq s+1$,

$$\frac{1}{2\pi\mathbf{i}} \int_{\Gamma} \frac{1}{\prod_{j=1}^{s+1} (z - \lambda_{i_j})} dz = (-1)^{T+1} \sum_{L \in \mathcal{L}(T,s)} w(X, Y|L).$$

For each $l < k$, let $F(l, k) = \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} \frac{1}{\prod_{j=l}^k (z - \lambda_{i_j})} dz$. The LHS is $F(1, s+1)$. We have

$$(91) \quad \begin{aligned} F(1, s+1) &= \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} \frac{1}{(z - \lambda_{i_1})(z - \lambda_{i_{s+1}})} \frac{1}{\prod_{j=2}^s (z - \lambda_{i_j})} dz \\ &= \frac{1}{2\pi\mathbf{i}} \times \frac{1}{(\lambda_{i_1} - \lambda_{i_{s+1}})} \int_{\Gamma} \left(\frac{1}{z - \lambda_{i_1}} - \frac{1}{z - \lambda_{i_{s+1}}} \right) \frac{1}{\prod_{j=2}^s (z - \lambda_{i_j})} dz \\ &= \frac{1}{(\lambda_{i_1} - \lambda_{i_{s+1}})} \left(\frac{1}{2\pi\mathbf{i}} \int_{\Gamma} \frac{1}{\prod_{j=1}^s (z - \lambda_{i_j})} dz - \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} \frac{1}{\prod_{j=2}^{s+1} (z - \lambda_{i_j})} dz \right) \\ &= \frac{1}{(\lambda_{i_1} - \lambda_{i_{s+1}})} [F(1, s) - F(2, s+1)]. \end{aligned}$$

Now, consider two sequences: $X' := \{i_2, \dots, i_T\}$ and $Y := \{i_{T+1}, \dots, i_{s+1}\}$, which have s indices in total. Notice that for any sequence of natural numbers m_2, \dots, m_T such that $m_2 + \dots + m_T = s-1$, we have

$$(92) \quad G(X, Y|1, m_2, \dots, m_T) = G(X', Y|m_2, \dots, m_T) \cup (\vec{e}_{i_1, i_{s+1}}),$$

where $\vec{e}_{i_1, i_{s+1}}$ is the directed edge from i_1 to i_{s+1} . Therefore,

$$w(X, Y|1, m_2, \dots, m_T) = \frac{1}{(\lambda_{i_1} - \lambda_{i_{s+1}})} \times w(X', Y|m_2, \dots, m_T).$$

Applying the induction hypothesis on $\lambda_{i_2}, \dots, \lambda_{i_{s+1}}$, we obtain

$$(93) \quad \begin{aligned} \frac{1}{(\lambda_{i_1} - \lambda_{i_{s+1}})} F(2, s+1) &= \sum_{m_2 + \dots + m_T = s-1} \frac{1}{(\lambda_{i_1} - \lambda_{i_{s+1}})} \times (-1)^{(T-1)+1} w(X', Y | m_2, \dots, m_T) \\ &= \sum_{1+m_2+\dots+m_T=s} (-1)^T w(X, Y | 1, m_2, \dots, m_T). \end{aligned}$$

Similarly, considering two other sequences $X := \{i_1, \dots, i_T\}$ and $Y' := \{i_{T+1}, \dots, i_s\}$ (which have s indices in total), and applying the induction hypothesis, we obtain

$$(94) \quad \begin{aligned} \frac{1}{(\lambda_{i_1} - \lambda_{i_{s+1}})} F(1, s) &= \sum_{m_1 + \dots + m_T = s-1} \frac{1}{(\lambda_{i_1} - \lambda_{i_{s+1}})} \times (-1)^{T+1} w(X, Y' | m_1, \dots, m_T) \\ &= \sum_{\substack{m_1 + \dots + m_T = s \\ m_1 > 1}} (-1)^{T+1} w(X, Y | m_1, \dots, m_T). \end{aligned}$$

Combining (91), (93) and (94), we obtain

$$(95) \quad \begin{aligned} F(1, s+1) &= \frac{1}{\lambda_{i_1} - \lambda_{i_{s+1}}} [F(1, s) - F(2, s+1)] \\ &= \sum_{\substack{m_1 + \dots + m_T = s \\ m_1 > 1}} (-1)^{T+1} w(X, Y | m_1, \dots, m_T) - \sum_{\substack{m_1 + m_2 + \dots + m_T = s \\ m_1 = 1}} (-1)^T w(X, Y | m_1, m_2, \dots, m_T) \\ &= \sum_{L \in \mathcal{L}(T, s)} (-1)^{T+1} w(X, Y | L). \end{aligned}$$

This verifies the claim for $s+1$, completing the proof. \square

We will also use the following simple lemma several times in the upcoming subsections.

Lemma 9.4. *Let a_1, a_2, a_3 be positive numbers and l_1, l_2, l_3 be non negative integers. One has*

$$a_1^{l_1} a_2^{l_2} a_3^{l_3} \leq \max\{a_1, a_2, a_3\}^{l_1+l_2+l_3}.$$

If $\max\{l_1, l_2\} \geq 1$, then

$$a_1^{l_1} a_2^{l_2} a_3^{l_3} \leq (a_1 + a_2) \times \max\{a_1, a_2, a_3\}^{l_1+l_2+l_3-1}.$$

9.5. Proof of Lemma 8.7. Fix (α, β) of Type I, define

$$(96) \quad \begin{aligned} A_l &:= \alpha_1 + \alpha_2 + \dots + \alpha_l \quad \text{for each } 1 \leq l \leq k, \\ B_\ell &:= \beta_1 + \beta_2 + \dots + \beta_\ell \quad \text{for each } 1 \leq \ell \leq k-1, \\ A_0 &= B_0 = 0. \end{aligned}$$

Note that for a pair (α, β) of Type I, $s_1 \geq k+1$. Multiplying out all the terms in P (there are s_2 of P 's) and all terms in Q (there are s_1 of Q 's), we obtain the expansion

$$\int_{\Gamma} M(\alpha; \beta) dz = \sum_{\substack{i_1, i_2, \dots, i_{s_2} \in N_{\bar{\lambda}}(S) \\ j_1, \dots, j_{s_1} \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{dz}{\prod_{\ell=1}^{s_1} (z - \lambda_{j_{\ell}}) \prod_{l=1}^{s_2} (z - \lambda_{i_l})} \times$$

$$\left\{ \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} \left(u_{j_{l+A_h}} u_{j_{l+A_h}}^T E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} \left(u_{i_{l+B_h}} u_{i_{l+B_h}}^T E \right) \right] \right\} \times \left[\prod_{l=1}^{\alpha_{k+1}-1} \left(u_{j_{l+A_k}} u_{j_{l+A_k}}^T E \right) \right] u_{j_{s_1}} u_{j_{s_1}}^T,$$

where the indices j_{ℓ} 's represent the Q -operators, and the indices i_l 's represent the P -operators. To make the grouping of the indices more clear, let us present a small example.

Example. Consider $s = 10, k = 2$, and $\alpha = (3, 2, 1); \beta = (2, 3)$. In this case

$$s_1 = 3 + 2 + 1 = 6, s_2 = 2 + 3 = 5,$$

and we have

$$\int_{\Gamma} M(3, 2, 1; 2, 3) dz = \int_{\Gamma} \underbrace{(QEQQE)}_{\substack{\alpha_1=3 \\ \text{numbers of } Q}} \underbrace{(PEPE)}_{\substack{\beta_1=2 \\ \text{numbers of } P}} \underbrace{(QEQE)}_{\substack{\alpha_2=2 \\ \text{numbers of } Q}} \underbrace{(PEPEPE)}_{\substack{\beta_2=3 \\ \text{numbers of } P}} \underbrace{Q}_{\substack{\alpha_3=1 \\ \text{numbers of } Q}} dz$$

into

$$\sum_{\substack{i_1, i_2, \dots, i_5 \in N_{\bar{\lambda}}(S) \\ j_1, \dots, j_6 \in N_{\bar{\lambda}}(S)^c}} \int_{\Gamma} \frac{dz}{\prod_{\ell=1}^6 (z - \lambda_{j_{\ell}}) \prod_{l=1}^5 (z - \lambda_{i_l})} [(u_{j_1} u_{j_1}^T E)(u_{j_2} u_{j_2}^T E)(u_{j_3} u_{j_3}^T E)(u_{i_1} u_{i_1}^T E)(u_{i_2} u_{i_2}^T E)]$$

$$\times [(u_{j_4} u_{j_4}^T E)(u_{j_5} u_{j_5}^T E)(u_{i_3} u_{i_3}^T E)(u_{i_4} u_{i_4}^T E)(u_{i_5} u_{i_5}^T E)] \times [u_{j_6} u_{j_6}^T].$$

From now on, to reduce unnecessary indices, we will write $\sum_{i_1, \dots, i_{s_2}}$ and $\sum_{j_1, \dots, j_{s_1}}$ instead of $\sum_{i_1, \dots, i_{s_2} \in N_{\bar{\lambda}}(S)}$ and $\sum_{j_1, \dots, j_{s_1} \in N_{\bar{\lambda}}(S)^c}$. The next idea is to use Lemma 9.3 to treat $\int_{\Gamma} \frac{dz}{\prod_{\ell=1}^{s_1} (z - \lambda_{j_{\ell}}) \prod_{l=1}^{s_2} (z - \lambda_{i_l})}$. In order to do that, we need to know which eigenvalues are inside/outside the contour Γ .

We have two sequences of indices: $[i_1, \dots, i_{s_2}]$, whose elements are in $N_{\bar{\lambda}}(S)$ and $[j_1, \dots, j_{s_1}]$, whose elements are in $N_{\bar{\lambda}}(S)^c$. The order of the elements in the sequence $[i_1, \dots, i_{s_2}]$ ($[j_1, \dots, j_{s_1}]$) is inherited from the positions of P 's (Q 's resp.), counting from left to right. Now, we fix a sequence i_1, \dots, i_{s_2} and split it exactly into two following sub-sequences. The first one (denoted by X) is the subsequence of indices i_l in S , i.e. λ_{i_l} is inside contour Γ . The second one (denoted by $I_2 = [i_{l_1}, i_{l_2}, \dots, i_{l_{|I_2|}}]$) is obtained from i_1, \dots, i_{s_2} by removing X . Define the sequence Y as

$$Y = [i_{l_1}, i_{l_2}, \dots, i_{l_{|I_2|}}, j_1, j_2, \dots, j_{s_1}].$$

Example: Continue with the above example ($s = 10, k = 2$ and so on), we further set $N_{\bar{\lambda}}(S) = \{1, 2, 3, 4, 5\}$ and $S = \{1, 2, 3\}$. Consider the sequence $[i_1, i_2, i_3, i_4, i_5] = [1, 4, 2, 2, 3]$. In this case,

$$X = [1, 2, 2, 3] \text{ and } Y = [4, j_1, j_2, j_3, j_4, j_5, j_6].$$

Let $T = |X|$ and apply the Lemma 9.3, we have

$$(97) \quad \begin{aligned} \frac{1}{2\pi i} \int_{\Gamma} \frac{dz}{\prod_{\ell=1}^{s_1} (z - \lambda_{j_\ell}) \prod_{l=1}^{s_2} (z - \lambda_{i_l})} &= \sum_{L \in \mathcal{L}(T, s)} (-1)^{T+1} w(X, Y|L) \\ &= \sum_{L \in \mathcal{L}(T, s)} (-1)^{T+1} \prod_{e \in \mathcal{E}(X, Y|L)} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})}. \end{aligned}$$

Notice that once the sequence $[i_1, i_2, \dots, i_{s_2}]$ and L are fixed, then the set X is well-defined, and the shape of the graph $G(X, Y|L)$ is also determined, regardless the particular values of j_l . In this case, we can write \mathcal{E}_L instead of $\mathcal{E}(X, Y|L)$ (for the edge set of the graph). Let us illustrate with a concrete example.


Example: Continue with the previous example ($s = 10, k = 2$ and so on). We further set $N_\lambda(S) = \{1, 2, 3, 4, 5\}$ and $S = \{1, 2, 3\}$. Consider the sequence $[i_1, i_2, i_3, i_4, i_5] = [1, 4, 2, 2, 3]$. In this case,

$$X = [1, 2, 2, 3] \quad \text{and} \quad Y = [4, j_1, j_2, j_3, j_4, j_5, j_6].$$

Thus, $T = 4$. We set

$$L = (m_1, m_2, m_3, m_4) = (4, 3, 2, 1).$$

The graph $G(1, 2, 2, 3; 4, j_1, j_2, j_3, j_4, j_5, j_6 | 4, 3, 2, 1)$ is



The set of edges $\mathcal{E}(X, Y|L)$ is

$\{(1 \rightarrow j_6), (1 \rightarrow j_5), (1 \rightarrow j_4), (1 \rightarrow j_3), (2 \rightarrow j_3), (2 \rightarrow j_2), (2 \rightarrow j_1), (2 \rightarrow j_1), (2 \rightarrow 4), (3 \rightarrow 4)\}$, and $w(X, Y|L)$ equals

$$\frac{1}{(\lambda_1 - \lambda_{j_6})(\lambda_1 - \lambda_{j_5})(\lambda_1 - \lambda_{j_4})(\lambda_1 - \lambda_{j_3})(\lambda_2 - \lambda_{j_3})(\lambda_2 - \lambda_{j_2})(\lambda_2 - \lambda_{j_1})(\lambda_2 - \lambda_{j_1})(\lambda_2 - \lambda_4)(\lambda_3 - \lambda_4)}.$$

Therefore, we can write

$$(98) \quad \begin{aligned} &\frac{1}{2\pi i} \int_{\Gamma} M(\alpha; \beta) dz \\ &= \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} \sum_{j_1, \dots, j_{s_1}} (-1)^{T+1} \prod_{e \in \mathcal{E}_L} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \times \\ &\quad \left\{ \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} \left(u_{j_{l+A_h}} u_{j_{l+A_h}}^T E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} \left(u_{i_{l+B_h}} u_{i_{l+B_h}}^T E \right) \right] \right\} \times \left[\prod_{l=1}^{\alpha_{k+1}-1} \left(u_{j_{l+A_k}} u_{j_{l+A_k}}^T E \right) \right] u_{j_{s_1}} u_{j_{s_1}}^T. \end{aligned}$$

Now, we "redistribute" the terms in $\prod_{e \in \mathcal{E}_L} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})}$. By the definition of Y , e^- either is in I_2 or equals j_l for some $1 \leq l \leq s_1$. Therefore,

$$\prod_{e \in \mathcal{E}_L} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} = \left[\prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \right] \times \prod_{l=1}^{s_1} \left[\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_l}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \right].$$

Example: Continue with the above example, we can rewrite $w(X, Y|L) = \prod_{e \in \mathcal{E}_L} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})}$ as

$$\frac{1}{(\lambda_2 - \lambda_4)(\lambda_3 - \lambda_4)} \times \frac{1}{(\lambda_2 - \lambda_{j_1})(\lambda_2 - \lambda_{j_1})} \times \frac{1}{(\lambda_2 - \lambda_{j_2})} \times \\ \frac{1}{(\lambda_1 - \lambda_{j_3})(\lambda_2 - \lambda_{j_3})} \times \frac{1}{(\lambda_1 - \lambda_{j_4})} \times \frac{1}{(\lambda_1 - \lambda_{j_5})} \times \frac{1}{(\lambda_1 - \lambda_{j_6})}.$$

For each $1 \leq l \leq s_1$, we group the factor $\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_l}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})}$ with $u_{j_l} u_{j_l}^T$. This way, we have

$$\frac{1}{2\pi i} \int_{\Gamma} M(\alpha; \beta) dz \\ = \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} \sum_{j_1, \dots, j_{s_1}} (-1)^{T+1} \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \times \\ \left\{ \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_h}} (\lambda_{e^+} - \lambda_{e^-})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_{l+B_h}} u_{i_{l+B_h}}^T E) \right] \right\} \times \\ \times \left[\prod_{l=1}^{\alpha_{k+1}-1} \left(\frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_k}} (\lambda_{e^+} - \lambda_{e^-})} E \right) \right] \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{s_1}} (\lambda_{e^+} - \lambda_{e^-})}.$$

We can distribute the sum over j_1, j_2, \dots, j_{s_1} into each position of $u_{j_l} u_{j_l}^T$ in the product and obtain

$$\begin{aligned}
& \frac{1}{2\pi i} \int_{\Gamma} M(\alpha; \beta) dz \\
&= \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} (-1)^{T+1} \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \times \\
& \left\{ \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\sum_{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{\substack{e \in \mathcal{E}_L; e^- = j_{l+A_h}}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \times \left[\left(\prod_{l=1}^{\beta_{h+1}} u_{i_{l+B_h}} u_{i_{l+B_h}}^T E \right) \right] \right\} \times \\
& \times \left[\prod_{l=1}^{\alpha_{k+1}-1} \left(\sum_{j_{l+A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_k}}} (\lambda_{e^+} - \lambda_{j_{l+A_k}})} E \right) \right] \times \left(\sum_{j_{s_1} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{s_1}}} (\lambda_{e^+} - \lambda_{j_{s_1}})} \right) \\
&= \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} (-1)^{T+1} \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \times W \times \\
& \times \left[\prod_{l=1}^{\alpha_{k+1}-1} \left(\sum_{j_{l+A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_k}}} (\lambda_{e^+} - \lambda_{j_{l+A_k}})} E \right) \right] \times \left(\sum_{j_{s_1} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{s_1}}} (\lambda_{e^+} - \lambda_{j_{s_1}})} \right),
\end{aligned}$$

where

$$(99) \quad W = \left\{ \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\sum_{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{\substack{e \in \mathcal{E}_L; e^- = j_{l+A_h}}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \times \left[\left(\prod_{l=1}^{\beta_{h+1}} u_{i_{l+B_h}} u_{i_{l+B_h}}^T E \right) \right] \right\}.$$

Therefore, by triangle inequality,

$$\begin{aligned}
(100) \quad & \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| \\
& \leq \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} \left| \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \right| \times \|W\| \times \\
& \left[\prod_{l=1}^{\alpha_{k+1}-1} \left(\left\| \sum_{j_{l+A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_k}}} (\lambda_{e^+} - \lambda_{j_{l+A_k}})} \|E\| \right\| \right) \right] \times \left\| \sum_{j_{s_1} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{s_1}}} (\lambda_{e^+} - \lambda_{j_{s_1}})} \right\|.
\end{aligned}$$

Now, we bound the terms from left to right. First,

$$(101) \quad \left| \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \right| \leq \frac{1}{\delta_S^{|E(I_2)|}},$$

where $E(I_2)$ is the set of edges ending in I_2 in $G(X, Y|L)$. Next, we can split W into

$$\begin{aligned}
W &= \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}-1} \left(\sum_{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_h}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \\
&\times \left[\left(\sum_{j_{\alpha_{h+1}+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{\alpha_{h+1}+A_h}} u_{j_{\alpha_{h+1}+A_h}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{\alpha_{h+1}+A_h}} (\lambda_{e^+} - \lambda_{j_{\alpha_{h+1}+A_h}})} \right) E \left(\prod_{l=1}^{\beta_{h+1}} u_{i_{l+B_h}} u_{i_{l+B_h}}^T E \right) \right] \\
&= \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}-1} \left(\sum_{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_h}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \\
&\times \left[\left(\sum_{j_{A_{h+1}} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{A_{h+1}}} u_{j_{A_{h+1}}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{A_{h+1}}} (\lambda_{e^+} - \lambda_{j_{A_{h+1}}})} \right) E \left(\prod_{l=1}^{\beta_{h+1}} u_{i_{l+B_h}} u_{i_{l+B_h}}^T E \right) \right],
\end{aligned}$$

(by Definition 96, we have $\alpha_{h+1} + A_h = A_{h+1}$).

Therefore,

$$(102) \quad \begin{aligned} \|W\| &\leq \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}-1} \left(\left\| \sum_{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_h}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} \right\| \cdot \|E\| \right) \right] \\ &\times \left\| \sum_{j_{A_{h+1}} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{A_{h+1}}} u_{j_{A_{h+1}}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{A_{h+1}}} (\lambda_{e^+} - \lambda_{j_{A_{h+1}}})} \right\| \cdot \left\| E \left(\prod_{l=1}^{\beta_{h+1}} u_{i_{l+B_h}} u_{i_{l+B_h}}^T E \right) \right\|. \end{aligned}$$

Notice that

$$(103) \quad \begin{aligned} \left\| E \left(\prod_{l=1}^{\beta_{h+1}} u_{i_{l+B_h}} u_{i_{l+B_h}}^T E \right) \right\| &= \left\| E u_{i_{1+B_h}} \left(\prod_{l=1}^{\beta_{h+1}-1} (u_{i_{l+B_h}}^T E u_{i_{l+1+B_h}}) \right) u_{i_{B_{h+1}}}^T E \right\| \\ &\leq \|E u_{i_{1+B_h}}\| \cdot \|u_{i_{B_{h+1}}}^T E\| \times \prod_{l=1}^{\beta_{h+1}-1} \|u_{i_{l+B_h}}^T E u_{i_{l+1+B_h}}\| \\ &\leq w^2 \cdot x^{\beta_{h+1}-1} \text{(by Definition 2.2)}. \end{aligned}$$

Moreover, for any fixed $1 \leq l \leq s_1$,

$$(104) \quad \left\| \sum_{j_l \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_l} u_{j_l}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_l} (\lambda_{e^+} - \lambda_{j_l})} \right\| \leq \frac{1}{\bar{\lambda}^{d(j_l)}},$$

where $d(j_l)$ is the degree of j_l in $G(X, Y|L)$ (this degree does not depend on the particular value of j_l). Applying (103) and (104) on the RHS of (102), we obtain

$$(105) \quad \begin{aligned} \|W\| &\leq \prod_{h=0}^{k-1} \left[w^2 x^{(\beta_{h+1}-1)} \|E\|^{(\alpha_{h+1}-1)} \left(\prod_{1 \leq l \leq \alpha_{h+1}} \frac{1}{\bar{\lambda}^{d(j_{l+A_h})}} \right) \right] \\ &= \prod_{h=0}^{k-1} \left[w^2 x^{(\beta_{h+1}-1)} \|E\|^{(\alpha_{h+1}-1)} \left(\prod_{1+A_h \leq l \leq A_{h+1}} \frac{1}{\bar{\lambda}^{d(j_l)}} \right) \right] \\ &= w^{2k} x^{\sum_{h=0}^{k-1} (\beta_{h+1}-1)} \|E\|^{\sum_{h=0}^{k-1} (\alpha_{h+1}-1)} \prod_{h=0}^{k-1} \left(\prod_{1+A_h \leq l \leq A_{h+1}} \frac{1}{\bar{\lambda}^{d(j_l)}} \right) \\ &= w^{2k} x^{\sum_{l=1}^k (\beta_l-1)} \|E\|^{\sum_{l=1}^k (\alpha_l-1)} \prod_{1 \leq l \leq A_k} \frac{1}{\bar{\lambda}^{d(j_l)}} \\ &= \frac{w^{2k} x^{\sum_{l=1}^k (\beta_l-1)} \|E\|^{\sum_{l=1}^k (\alpha_l-1)}}{\bar{\lambda}^{\sum_{l=1}^k d(j_l)}}. \end{aligned}$$

Arguing as in (104), we also have

$$(106) \quad \left\| \sum_{j_{l+A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_k}} (\lambda_{e^+} - \lambda_{j_{l+A_k}})} \right\| \leq \frac{1}{\bar{\lambda}^{d(j_{l+A_k})}} \text{ for all } 1 \leq l \leq \alpha_{k+1} - 1,$$

$$\left\| \sum_{j_{s_1} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{s_1}} (\lambda_{e^+} - \lambda_{j_{s_1}})} \right\| \leq \frac{1}{\bar{\lambda}^{d(j_{s_1})}},$$

and then,

$$\left[\prod_{l=1}^{\alpha_{k+1}-1} \left(\left\| \sum_{j_{l+A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_k}} (\lambda_{e^+} - \lambda_{j_{l+A_k}})} \right\| \|E\| \right) \right] \times \left\| \sum_{j_{s_1} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{s_1}} (\lambda_{e^+} - \lambda_{j_{s_1}})} \right\|$$

$$\leq \|E\|^{\alpha_{k+1}-1} \prod_{1+A_k \leq l \leq s_1} \frac{1}{\bar{\lambda}^{d(j_l)}} = \frac{\|E\|^{\alpha_{k+1}-1}}{\bar{\lambda}^{\sum_{l=1+A_k}^{s_1} d(j_l)}}.$$

Combining (100), (101), (105), and (106), we finally obtain

$$(107) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| \leq \sum_{i_1, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} \frac{w^{2k} x^{\sum_{i=1}^k (\beta_i - 1)} \|E\|^{\sum_{i=1}^{k+1} (\alpha_i - 1)}}{\delta_S^{|E(I_2)|} \bar{\lambda}^{\sum_{l=1}^{s_1} d(j_l)}}.$$

Set $r_c(T, L) := \sum_{l=1}^{s_1} (d(j_l) - 1) \geq 0$. Then, by Remark 9.2,

$$(108) \quad \sum_{l=1}^{s_1} d(j_l) = r_c(T, L) + s_1 \text{ and } |E(I_2)| = s - (r_c(T, L) + s_1) = s_2 - 1 - r_c(T, L).$$

We can rewrite above inequality as

$$(109) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| \leq \sum_{i_1, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} \frac{\|E\|^{s_1 - (k+1)} w^{2k} x^{s_2 - k}}{\delta_S^{s_2 - 1 - r_c(T, L)} \bar{\lambda}^{s_1 + r_c(T, L)}}.$$

Notice that if we fix T and a sequence $L \in \mathcal{L}(T, s)$, the shape of the graph $G(X, Y|L)$ and then $r_c(T, L)$ are uniquely determined. Hence, if two different sequences $[i_1, \dots, i_{s_2}]$ and $[i'_1, \dots, i'_{s_2}]$ has the same number T of elements in S , then their corresponding $r_c(T, L)$'s are the same for each fixed $L \in \mathcal{L}(T, s)$. Moreover, when we fix T , the choices for i_1, \dots, i_{s_2} are at most

$\binom{s_2}{T} p^T (r-p)^{s_2-T} < \binom{s_2}{T} p^T r^{s_2-T}$. Thus, (107) leads to

$$\begin{aligned}
\frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} p^T r^{s_2-T} \sum_{L \in \mathcal{L}(T, s)} \frac{\|E\|^{s_1-(k+1)} w^{2k} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}} \\
&= M_1(\alpha; \beta) + M_2(\alpha; \beta), \text{ where} \\
(110) \quad M_1(\alpha; \beta) &:= \sum_{T=1}^{s_2} \binom{s_2}{T} p^T r^{s_2-T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) \leq k-1} \frac{\|E\|^{s_1-(k+1)} w^{2k} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}}, \\
M_2(\alpha; \beta) &:= \sum_{T=1}^{s_2} \binom{s_2}{T} p^T r^{s_2-T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) > k-1} \frac{\|E\|^{s_1-(k+1)} w^{2k} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}}.
\end{aligned}$$

In order to bound $M_1(\alpha; \beta)$ and $M_2(\alpha; \beta)$ from above, we need the following property of $r_c(T, L)$.

Lemma 9.6.

$$r_c(T, L) \leq T - 1.$$

Proof of Lemma 9.6. By Remark 9.2, we have $|E(I_2)| \geq |I_2|$. Since $|E(I_2)| = s_2 - (r_c(T, L) + 1)$ and $|I_2| = s_2 - T$, we obtain

$$r_c(T, L) \leq T - 1. \quad \square$$

Back to our estimation, we can bound $M_1(\alpha; \beta)$ from above as follows. Since $w \leq \|E\|$, $w^{2k} \leq w^{2(k-1-r_c(T, L))} \|E\|^{2(r_c(T, L)+1)}$. Thus,

$$\begin{aligned}
(111) \quad M_1(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} p^T r^{s_2-T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) \leq k-1} \frac{\|E\|^{s_1-(k+1)+2(r_c(T, L)+1)} w^{2(k-1-r_c(T, L))} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}} \\
&= \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) \leq k-1} p^T r^{s_2-T} \left(\frac{\|E\|}{\bar{\lambda}} \right)^{T_1} \left(\frac{x}{\delta_S} \right)^{T_2} \left(\frac{w}{\sqrt{\lambda} \delta_S} \right)^{T_3},
\end{aligned}$$

where

$$T_1 = s_1 - (k-1) + 2r_c(T, L) \geq 2,$$

$$T_2 = s_2 - k,$$

$$T_3 = 2(k-1-r_c(T, L)).$$

Redistributing the factors \sqrt{p}, r, \sqrt{r} to $\frac{\|E\|}{\bar{\lambda}}, \frac{x}{\delta_S}, \frac{w}{\sqrt{\lambda} \delta_S}$, respectively, we can rewrite the RHS of (111) as

$$\sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) \leq k-1} \left(\frac{p}{r} \right)^{T-(r_c(T, L)+1)} p^{\frac{k+1-s_1}{2}} \left(\frac{\sqrt{p} \|E\|}{\bar{\lambda}} \right)^{T_1} \left(\frac{rx}{\delta_S} \right)^{T_2} \left(\frac{\sqrt{r} w}{\sqrt{\lambda} \delta_S} \right)^{T_3}.$$

Since $T \geq r_c(T, L) + 1$ (Lemma (9.6)) and $s_1 - (k + 1) \geq 0$ ((α, β) is Type I),

$$0 \leq \left(\frac{p}{r}\right)^{T-(r_c(T,L)+1)} \leq 1 \text{ and } 0 \leq p^{\frac{k+1-s_1}{2}} \leq 1.$$

We can omit these factors and obtain

$$\begin{aligned} M_1(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{T_1} \left(\frac{rx}{\delta_S}\right)^{T_2} \left(\frac{\sqrt{r}w}{\sqrt{\lambda}\delta_S}\right)^{T_3} \\ &= \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \times \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{T_1-1} \left(\frac{rx}{\delta_S}\right)^{T_2} \left(\frac{\sqrt{r}w}{\sqrt{\lambda}\delta_S}\right)^{T_3}. \end{aligned}$$

By the setting of $M_1(\alpha; \beta)$, $r_c(T, L) \leq k - 1$, and then $T_3 \geq 0$. We also have $T_2 \geq 0$ and $T_1 - 1 \geq 1$ by the definition of Type I (α, β) . Therefore, by Lemma 9.4,

(112)

$$\begin{aligned} M_1(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right) \times \max \left\{ \frac{\sqrt{p}\|E\|}{\bar{\lambda}}, \frac{rx}{\delta_S}, \frac{\sqrt{r}\|E\|}{\sqrt{\lambda}\delta_S} \right\}^{T_1+T_2+T_3-1} \\ &= \left(\sum_{T=1}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s): \\ r_c(T,L) \leq k-1}} \binom{s_2}{T} \right) \times \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right) \times \max \left\{ \frac{\sqrt{p}\|E\|}{\bar{\lambda}}, \frac{rx}{\delta_S}, \frac{\sqrt{r}\|E\|}{\sqrt{\lambda}\delta_S} \right\}^{T_1+T_2+T_3-1} \\ &\leq \left(\sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T,s)} \binom{s_2}{T} \right) \times \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right) \times \max \left\{ \frac{\sqrt{p}\|E\|}{\bar{\lambda}}, \frac{rx}{\delta_S}, \frac{\sqrt{r}\|E\|}{\sqrt{\lambda}\delta_S} \right\}^{T_1+T_2+T_3-1}. \end{aligned}$$

Since $|\mathcal{L}(T, s)| = \binom{s-1}{T-1} \leq 2^{s-1}$ for all $T \leq s$, we have

$$(113) \quad \sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T,s)} \binom{s_2}{T} \leq 2^{s-1} \sum_{T=1}^{s_2} \binom{s_2}{T} \leq 2^{s_2+s-1}.$$

Thus, (112) becomes

$$\begin{aligned} M_1(\alpha; \beta) &\leq 2^{s_2+s-1} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right) \times \max \left\{ \frac{\sqrt{p}\|E\|}{\bar{\lambda}}, \frac{rx}{\delta_S}, \frac{\sqrt{r}\|E\|}{\sqrt{\lambda}\delta_S} \right\}^{T_1+T_2+T_3-1} \\ (114) \quad &\leq \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right) \frac{2^{s_2+s-1}}{12^{T_1+T_2+T_3-1}} \text{ (by Assumption } \mathbf{D0} \text{ and the fact that } \delta \leq \delta_S) \\ &= \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right) \frac{2^{s_2+s-1}}{12^{s-1}} \text{ (since } T_1 + T_2 + T_3 = s). \end{aligned}$$

Similarly, we can bound $M_2(\alpha; \beta)$ as follows. Since $w \leq \|E\|$, $w^{2k} \leq \|E\|^{2k}$ and hence

$$(115)$$

$$\begin{aligned} M_2(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} p^T r^{s_2-T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) > k-1} \frac{\|E\|^{s_1+k-1} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}} \\ &= \sum_{T=1}^{s_2} \binom{s_2}{T} p^T r^{s_2-T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) > k-1} \left(\frac{x}{\delta_S}\right)^{s_2-1-r_c(T, L)} \left(\frac{x}{\bar{\lambda}}\right)^{r_c(T, L)-(k-1)} \left(\frac{\|E\|}{\bar{\lambda}}\right)^{k-1+s_1}. \end{aligned}$$

Distributing each factors r, p, \sqrt{p} to $\frac{x}{\delta_S}, \frac{x}{\bar{\lambda}}, \frac{\|E\|}{\bar{\lambda}}$ respectively, we can rewrite the RHS as

$$\sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \left(\frac{p}{r}\right)^{T-(r_c(T, L)+1)} p^{\frac{k+1-s_1}{2}} \left(\frac{rx}{\delta_S}\right)^{s_2-1-r_c(T, L)} \left(\frac{px}{\bar{\lambda}}\right)^{r_c(T, L)-(k-1)} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{k-1+s_1}.$$

Since $T \geq r_c(T, L) + 1$ (Lemma (9.6)) and $s_1 - (k + 1) \geq 0$ ((α, β) is Type I),

$$0 \leq \left(\frac{p}{r}\right)^{T-(r_c(T, L)+1)} \leq 1 \quad \text{and} \quad 0 \leq p^{\frac{k+1-s_1}{2}} \leq 1.$$

We can omit these factors and obtain

$$\begin{aligned} M_2(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \left(\frac{rx}{\delta_S}\right)^{s_2-1-r_c(T, L)} \left(\frac{px}{\bar{\lambda}}\right)^{r_c(T, L)-(k-1)} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{k-1+s_1} \\ &= \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \frac{px}{\bar{\lambda}} \times \left(\frac{rx}{\delta_S}\right)^{s_2-1-r_c(T, L)} \left(\frac{px}{\bar{\lambda}}\right)^{r_c(T, L)-(k-1)-1} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{k-1+s_1}. \end{aligned}$$

By the setting of $M_2(\alpha; \beta)$, $r_c(T, L) - (k - 1) - 1 \geq 0$. By definitions of $s_1, s_2, r_c(T, L)$, we also have $s_2 - 1 - r_c(T, L), k - 1 + s_1 \geq 0$. Applying Lemma 9.4, we finally obtain

(116)

$$\begin{aligned}
M_2(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \frac{px}{\lambda} \times \max \left\{ \frac{rx}{\delta_S}, \frac{px}{\lambda}, \frac{\sqrt{p} \|E\|}{\lambda} \right\}^{s_2-1-r_c(T, L)+r_c(T, L)-(k-1)-1+k-1+s_1} \\
&= \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \frac{px}{\lambda} \times \max \left\{ \frac{rx}{\delta_S}, \frac{px}{\lambda}, \frac{\sqrt{p} \|E\|}{\lambda} \right\}^{s-1} \quad (\text{since } s_1 + s_2 - 2 = s - 1) \\
&= \left(\sum_{T=1}^{s_2} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \binom{s_2}{T} \right) \times \frac{px}{\lambda} \times \max \left\{ \frac{rx}{\delta_S}, \frac{px}{\lambda}, \frac{\sqrt{p} \|E\|}{\lambda} \right\}^{s-1} \\
&\leq \left(\sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T, s)} \binom{s_2}{T} \right) \times \frac{px}{\lambda} \times \max \left\{ \frac{rx}{\delta_S}, \frac{px}{\lambda}, \frac{\sqrt{p} \|E\|}{\lambda} \right\}^{s-1} \\
&\leq 2^{s_2+s-1} \frac{px}{\lambda} \max \left\{ \frac{rx}{\delta_S}, \frac{\sqrt{p} \|E\|}{\lambda} \right\}^{s-1} \quad (\text{by (113) and } \frac{px}{\lambda} \leq \frac{rx}{\delta_S}) \\
&\leq \frac{px}{\lambda} \frac{2^{s_2+s-1}}{12^{s-1}} \quad (\text{by Assumption D0 and the fact that } \delta \leq \delta_S).
\end{aligned}$$

Combining (110), (112) and (116), we obtain that

$$(117) \quad \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| \leq \left(\frac{\sqrt{p} \|E\|}{\lambda} + \frac{px}{\lambda} \right) \frac{2^{s_2+s-1}}{12^{s-1}}.$$

It proves Lemma 8.7.

9.7. Proof of Lemma 8.8. Similar to (98), we obtain

$$\begin{aligned}
&\frac{1}{2\pi i} \int_{\Gamma} M(\alpha; \beta) dz \\
&= \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} \sum_{j_1, \dots, j_{s_1}} (-1)^{T+1} \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \times \left[\prod_{l=1}^{\beta_1} (u_{i_l} u_{i_l}^T E) \right] \times \\
&\quad \left\{ \prod_{h=1}^{k-2} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\frac{u_{j_l+A_h} u_{j_l+A_h}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_l+A_h} (\lambda_{e^+} - \lambda_{j_l+A_h})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_l+B_h} u_{i_l+B_h}^T E) \right] \right\} \times \\
&\quad \times \left[\prod_{l=1}^{\alpha_k} \left(\frac{u_{j_l+A_{k-1}} u_{j_l+A_{k-1}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_l+A_{k-1}} (\lambda_{e^+} - \lambda_{j_l+A_{k-1}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{k-1}} (u_{i_l+B_{k-1}} u_{i_l+B_{k-1}}^T E) \right] u_{i_{s_2}} u_{i_{s_2}}^T.
\end{aligned}$$

Similar to the previous subsection, we distribute the sum over j_1, j_2, \dots, j_{s_2} according to each position of $u_{j_l} u_{j_l}^T$ in the product, and obtain

$$\begin{aligned}
& \frac{1}{2\pi\mathbf{i}} \int_{\Gamma} M(\alpha; \beta) dz \\
&= \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} (-1)^{T+1} \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \times \left[\prod_{l=1}^{\beta_1} (u_{i_l} u_{i_l}^T E) \right] \times \\
& \left\{ \prod_{h=1}^{k-2} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\sum_{\substack{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c \\ e \in \mathcal{E}_L; e^- = j_{l+A_h}}} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_{l+B_h}} u_{i_{l+B_h}}^T E) \right] \right\} \times \\
& \times \left[\prod_{l=1}^{\alpha_k} \left(\sum_{\substack{j_{l+A_{k-1}} \in N_{\bar{\lambda}}(S)^c \\ e \in \mathcal{E}_L; e^- = j_{l+A_{k-1}}} \frac{u_{j_{l+A_{k-1}}} u_{j_{l+A_{k-1}}}^T}{\prod (\lambda_{e^+} - \lambda_{j_{l+A_{k-1}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{k-1}} (u_{i_{l+B_{k-1}}} u_{i_{l+B_{k-1}}}^T E) \right] u_{i_{s_2}} u_{i_{s_2}}^T.
\end{aligned}$$

Notice that once we fix the sequence $[i_1, \dots, i_{s_2}]$ and $L \in \mathcal{L}(T, s)$, the graph $G(X, Y|L)$ and $r_c(T, L)$ are determined. We can split the double sum $\sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} (\dots)$ above into two sub-sums:

$$\begin{aligned}
(118) \quad \Sigma_1 &:= \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) = 0}} (\dots), \\
\Sigma_2 &:= \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > 0}} (\dots),
\end{aligned}$$

and then,

$$\frac{1}{2\pi\mathbf{i}} \int_{\Gamma} M(\alpha; \beta) dz = \Sigma_1 + \Sigma_2.$$

We prove two lemmas bounding $\|\Sigma_1\|$ and $\|\Sigma_2\|$. We assume that Assumption D0 holds in all of two lemmas.

Lemma 9.8. *If $(s_2, s_1) \neq (k, k-1)$, then*

$$(119) \quad \|\Sigma_1\| \leq \left(\frac{\sqrt{p}rx}{\delta_S} + \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right) \frac{\sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) = 0} \binom{s_2}{T}}{12^{s-1}}.$$

If $(s_2, s_1) = (k, k-1)$, then

$$(120) \quad \|\Sigma_1\| \leq \frac{(s+2)\sqrt{rpy}}{\delta_S} \left(\frac{2\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right)^{s-2}.$$

Lemma 9.9.

$$(121) \quad \|\Sigma_2\| \leq \frac{px}{\bar{\lambda}} \frac{\sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) > k-1}} \binom{s_2}{T}}{12^{s-1}} + \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right) \frac{\sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ 1 \leq r_c(T,L) \leq k-1}} \binom{s_2}{T}}{12^{s-1}}.$$

Together Lemma 9.9 and (119) imply that if $(s_2, s_1) \neq (k, k-1)$, then

$$\begin{aligned} \|\Sigma_1\| + \|\Sigma_2\| &\leq \left(\frac{\sqrt{pr}x}{\delta_S} + \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right) \frac{\sum_{T=1}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L)=0}} \binom{s_2}{T}}{12^{s-1}} \\ &\quad + \frac{px}{\bar{\lambda}} \frac{\sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) > k-1}} \binom{s_2}{T}}{12^{s-1}} + \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right) \frac{\sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ 1 \leq r_c(T,L) \leq k-1}} \binom{s_2}{T}}{12^{s-1}}. \end{aligned}$$

Replacing $\frac{px}{\bar{\lambda}}$ by $\frac{\sqrt{pr}x}{\delta_S}$ (bigger one) and rearranging the RHS, we further obtain

$$(122) \quad \begin{aligned} \|\Sigma_1\| + \|\Sigma_2\| &\leq \frac{\sqrt{pr}x}{\delta_S} \frac{\sum_{T=1}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L)=0}} \binom{s_2}{T} + \sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s); \\ r_c(T,L) > k-1}} \binom{s_2}{T}}{12^{s-1}} + \\ &\quad + \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \frac{\sum_{T=1}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L)=0}} \binom{s_2}{T} + \sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s); \\ 1 \leq r_c(T,L) \leq k-1}} \binom{s_2}{T}}{12^{s-1}} \\ &\leq \frac{\sqrt{pr}x}{\delta_S} \frac{\sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T,s)} \binom{s_2}{T}}{12^{s-1}} + \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \frac{\sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T,s)} \binom{s_2}{T}}{12^{s-1}} \\ &\leq \left(\frac{\sqrt{pr}x}{\delta_S} + \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right) \frac{2^{s_2+s-1}}{12^{s-1}} \text{ (by (113)).} \end{aligned}$$

Therefore, if $(s_2, s_1) \neq (k, k-1)$, we have

$$\frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) \right\| \leq \|\Sigma_1\| + \|\Sigma_2\| \leq \left(\frac{\sqrt{pr}x}{\delta_S} + \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right) \frac{2^{s_2+s-1}}{12^{s-1}}.$$

This proves the first part of Lemma 8.8.

On the other hand, together Lemma 9.9 and (120) imply that if $(s_2, s_1) = (k, k-1)$, then

$$(123) \quad \begin{aligned} \|\Sigma_1\| + \|\Sigma_2\| &\leq \\ &\frac{(s+2)\sqrt{rpy}}{\delta_S} \left(\frac{2\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right)^{s-2} + \frac{px}{\bar{\lambda}} \frac{\sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) > k-1}} \binom{s_2}{T}}{12^{s-1}} + \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right) \frac{\sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ 1 \leq r_c(T,L) \leq k-1}} \binom{s_2}{T}}{12^{s-1}}. \end{aligned}$$

Moreover, by (113),

$$\max \left\{ \frac{\sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) > k-1}} \binom{s_2}{T}}{12^{s-1}}, \frac{\sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ 1 \leq r_c(T,L) \leq k-1}} \binom{s_2}{T}}{12^{s-1}} \right\} \leq \frac{\sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T,s)} \binom{s_2}{T}}{12^{s-1}} \leq \frac{2^{s_2+s-1}}{12^{s-1}}.$$

We can rewrite (123) into

$$(124) \quad \|\Sigma_1\| + \|\Sigma_2\| \leq \frac{(s+2)\sqrt{rpy}}{\delta_S} \left(\frac{2\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right)^{s-2} + \left(\frac{px}{\bar{\lambda}} + \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right) \frac{2^{s_2+s-1}}{12^{s-1}}.$$

This implies that if $(s_2, s_1) = (k, k-1)$,

$$\frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) \right\| \leq \|\Sigma_1\| + \|\Sigma_2\| \leq \left(\frac{px}{\bar{\lambda}} + \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right) \frac{2^{s_2+s-1}}{12^{s-1}} + \frac{(s+2)\sqrt{rpy}}{\delta_S} \left(\frac{2\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right)^{s-2}.$$

In this case, $s_2 = k, s_1 = k-1, s = s_1 + s_2 - 1 = 2(k-1)$ and hence $s_2 = s/2 + 1, s_2 + s - 1 = 3s/2$. The RHS can be rewritten as

$$\left(\frac{px}{\bar{\lambda}} + \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right) \frac{2^{3s/2}}{12^{s-1}} + \frac{(s+2)\sqrt{rpy}}{\delta_S} \left(\frac{2\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right)^{s-2}.$$

This proves the second part of Lemma 8.8.

9.9.1. *Proof of Lemma 9.9.* Recall that for any square matrix M ,

$$(125) \quad \|M\| := \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \mathbf{v}^T M \mathbf{w}.$$

Therefore,

$$\begin{aligned} & \|\Sigma_2\| \\ &= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) > 0} (-1)^{T+1} \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \times \mathbf{v}^T \left[\prod_{l=1}^{\beta_1} (u_{i_l} u_{i_l}^T E) \right] \times \\ & \left\{ \prod_{h=1}^{k-2} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\sum_{\substack{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c \\ e \in \mathcal{E}_L; e^- = j_{l+A_h}}} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_{l+B_h}} u_{i_{l+B_h}}^T E) \right] \right\} \times \\ & \times \left[\prod_{l=1}^{\alpha_k} \left(\sum_{\substack{j_{l+A_{k-1}} \in N_{\bar{\lambda}}(S)^c \\ e \in \mathcal{E}_L; e^- = j_{l+A_{k-1}}} \frac{u_{j_{l+A_{k-1}}} u_{j_{l+A_{k-1}}}^T}{\prod (\lambda_{e^+} - \lambda_{j_{l+A_{k-1}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{k-1}} (u_{i_{l+B_{k-1}}} u_{i_{l+B_{k-1}}}^T E) \right] u_{i_{s_2}} u_{i_{s_2}}^T \mathbf{w}. \end{aligned}$$

By triangle inequality,

(126)

$$\begin{aligned} \|\Sigma_2\| &\leq \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) > 0} \left| \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \right| \times \left\| \mathbf{v}^T \prod_{l=1}^{\beta_1} (u_{i_l} u_{i_l}^T E) \right\| \times \\ &\left\| \prod_{h=1}^{k-2} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\sum_{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_h}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_{l+B_h}} u_{i_{l+B_h}}^T E) \right] \right] \right\| \times \\ &\times \left\| \left[\prod_{l=1}^{\alpha_k} \left(\sum_{j_{l+A_{k-1}} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_{k-1}}} u_{j_{l+A_{k-1}}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_{k-1}}} (\lambda_{e^+} - \lambda_{j_{l+A_{k-1}})} E \right) \times \left[\prod_{l=1}^{\beta_{k-1}} (u_{i_{l+B_{k-1}}} u_{i_{l+B_{k-1}}}^T E) \right] u_{i_{s_2}} u_{i_{s_2}}^T \mathbf{w} \right] \right\|. \end{aligned}$$

We proceed in a fashion similar to the previous subsection. First,

$$(127) \quad \left| \frac{1}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} (\lambda_{e^+} - \lambda_{e^-})} \right| \leq \frac{1}{\delta_S^{|E(I_2)|}},$$

in which $E(I_2)$ is the set of edges ending in I_2 in $G(X, Y|L)$. Second,

$$(128) \quad \begin{aligned} \left\| \mathbf{v}^T \prod_{l=1}^{\beta_1} (u_{i_l} u_{i_l}^T E) \right\| &\leq |\mathbf{v}^T u_{i_1}| \times \left[\prod_{l=1}^{\beta_1-1} |u_{i_l}^T E u_{i_{l+1}}| \right] \times \|u_{i_{\beta_1}}^T E\| \\ &\leq |\mathbf{v}^T u_{i_1}| x^{\beta_1-1} w \text{ (by Definition 2.2)}. \end{aligned}$$

Third, by repeating the procedure of bounding $\|W\|$ from the previous subsection, we obtain

(129)

$$\begin{aligned} &\left\| \prod_{h=1}^{k-2} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\sum_{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_h}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_{l+B_h}} u_{i_{l+B_h}}^T E) \right] \right] \right\| \\ &\leq \prod_{h=1}^{k-2} \left[w^2 x^{(\beta_{h+1}-1)} \|E\|^{(\alpha_{h+1}-1)} \left(\prod_{1+A_h \leq l \leq A_{h+1}} \frac{1}{\bar{\lambda}^{d(j_l)}} \right) \right] \\ &= \frac{w^{2(k-2)} x^{\sum_{l=2}^{k-1} (\beta_l-1)} \|E\|^{\sum_{l=2}^{k-1} (\alpha_l-1)}}{\bar{\lambda}^{\sum_{l=1}^{A_{k-1}} d(j_l)}} \text{ (since } \alpha_1 = 0 \text{ and then } 1 + A_1 = 1 + 0 = 1). \end{aligned}$$

Finally, since $\alpha_k + A_{k-1} = A_k$, we have

$$\begin{aligned}
& \left\| \left[\prod_{l=1}^{\alpha_k} \left(\sum_{j_{l+A_{k-1}} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_{k-1}}} u_{j_{l+A_{k-1}}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_{k-1}}} (\lambda_{e^+} - \lambda_{j_{l+A_{k-1}}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{k-1}} \left(u_{i_{l+B_{k-1}}} u_{i_{l+B_{k-1}}}^T E \right) \right] u_{i_{s_2}} u_{i_{s_2}}^T \mathbf{w} \right\| \\
&= \left\| \left[\prod_{l=1}^{\alpha_{k-1}} \left(\sum_{j_{l+A_{k-1}} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_{k-1}}} u_{j_{l+A_{k-1}}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_{k-1}}} (\lambda_{e^+} - \lambda_{j_{l+A_{k-1}}})} E \right) \right] \times \left(\sum_{j_{A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{A_k}} u_{j_{A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{A_k}}} (\lambda_{e^+} - \lambda_{j_{A_k}})} E \right) \right. \\
&\quad \left. \times \left[\prod_{l=1}^{\beta_{k-1}} \left(u_{i_{l+B_{k-1}}} u_{i_{l+B_{k-1}}}^T E \right) \right] \times u_{i_{s_2}} u_{i_{s_2}}^T \mathbf{w} \right\|.
\end{aligned}$$

We can rewrite the RHS into

$$\begin{aligned}
& \left\| \left[\prod_{l=1}^{\alpha_{k-1}} \left(\sum_{j_{l+A_{k-1}} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_{k-1}}} u_{j_{l+A_{k-1}}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_{k-1}}} (\lambda_{e^+} - \lambda_{j_{l+A_{k-1}}})} E \right) \right] \times \left(\sum_{j_{A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{A_k}} u_{j_{A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{A_k}}} (\lambda_{e^+} - \lambda_{j_{A_k}})} \right) \right. \\
&\quad \left. \times \left\{ E \left[\prod_{l=1}^{\beta_{k-1}} \left(u_{i_{l+B_{k-1}}} u_{i_{l+B_{k-1}}}^T E \right) \right] u_{i_{s_2}} \right\} \times (u_{i_{s_2}}^T \mathbf{w}) \right\| \\
&= \left\| \left[\prod_{l=1}^{\alpha_{k-1}} \left(\sum_{j_{l+A_{k-1}} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_{k-1}}} u_{j_{l+A_{k-1}}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_{k-1}}} (\lambda_{e^+} - \lambda_{j_{l+A_{k-1}}})} E \right) \right] \times \left(\sum_{j_{A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{A_k}} u_{j_{A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{A_k}}} (\lambda_{e^+} - \lambda_{j_{A_k}})} \right) \right. \\
&\quad \left. \times (E u_{i_{1+B_{k-1}}}) \times \left[\prod_{l=1}^{\beta_{k-1}} \left(u_{i_{l+B_{k-1}}}^T E u_{i_{l+1+B_{k-1}}} \right) \right] \times (u_{i_{s_2}}^T \mathbf{w}) \right\| \\
&\leq \left[\prod_{l=1}^{\alpha_{k-1}} \left\| \sum_{j_{l+A_{k-1}} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_{k-1}}} u_{j_{l+A_{k-1}}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_{k-1}}} (\lambda_{e^+} - \lambda_{j_{l+A_{k-1}}})} \right\| \|E\| \right] \times \left\| \sum_{j_{A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{A_k}} u_{j_{A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{A_k}}} (\lambda_{e^+} - \lambda_{j_{A_k}})} \right\| \\
&\quad \times \|E u_{i_{1+B_{k-1}}}\| \times \left[\prod_{l=1}^{\beta_{k-1}} \left\| u_{i_{l+B_{k-1}}}^T E u_{i_{l+1+B_{k-1}}} \right\| \right] \times |u_{i_{s_2}}^T \mathbf{w}|.
\end{aligned}$$

By (104) and Definition 2.2, the RHS above is at most

$$\left[\|E\|^{(\alpha_k-1)} \left(\prod_{1+A_{k-1} \leq l \leq A_k} \frac{1}{\bar{\lambda}^{d(j_l)}} \right) w x^{(\beta_k-1)} \right] \times \left| u_{i_{s_2}}^T \mathbf{w} \right| = \frac{w x^{\beta_k-1} \|E\|^{\alpha_k-1} \times \left| u_{i_{s_2}}^T \mathbf{w} \right|}{\bar{\lambda}^{\sum_{l=1+A_{k-1}}^{A_k} d(j_l)}}.$$

It implies

$$(130) \quad \left\| \left[\prod_{l=1}^{\alpha_k} \left(\sum_{j_{l+A_{k-1}} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_{k-1}}} u_{j_{l+A_{k-1}}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_{k-1}}} (\lambda_{e^+} - \lambda_{j_{l+A_{k-1}}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_k-1} \left(u_{i_{l+B_{k-1}}} u_{i_{l+B_{k-1}}}^T E \right) \right] u_{i_{s_2}} u_{i_{s_2}}^T \mathbf{w} \right\| \\ \leq \frac{w x^{\beta_k-1} \|E\|^{\alpha_k-1} \times \left| u_{i_{s_2}}^T \mathbf{w} \right|}{\bar{\lambda}^{\sum_{l=1+A_{k-1}}^{A_k} d(j_l)}}.$$

Combining (126), (127), (128), (129), and (130), we obtain

$$(131) \quad \|\Sigma_2\| \leq \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i_1, \dots, i_{s_2}} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > 0}} |\mathbf{v}^T u_{i_1}| \left| u_{i_{s_2}}^T \mathbf{w} \right| \frac{w^{2(k-1)} x^{\sum_{l=1}^k (\beta_l-1)} \|E\|^{\sum_{l=2}^k (\alpha_l-1)}}{\delta_S^{|E(I_2)|} \bar{\lambda}^{\sum_{l=1}^{s_1} d(j_l)}} \\ = \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i_1, \dots, i_{s_2}} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > 0}} |\mathbf{v}^T u_{i_1}| \left| u_{i_{s_2}}^T \mathbf{w} \right| \frac{\|E\|^{s_1-(k-1)} w^{2(k-1)} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}} \text{ (by (108))} \\ = \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > 0}} \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| \left| u_{i_{s_2}}^T \mathbf{w} \right| \right] \frac{\|E\|^{s_1-(k-1)} w^{2(k-1)} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}},$$

here recall that $X(i_1, \dots, i_{s_2})$ is the sub-sequence of $[i_1, \dots, i_{s_2}]$ whose elements are in S . Note that in this case $r_c(T, L) \geq 1$, by Lemma 9.6, $T \geq 2$ (that is why our sum runs from $T = 2$ to $T = s_2$).

Moreover,

$$\begin{aligned} \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \right] &= \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ i_1, i_{s_2} \in S \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \right] + \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ i_1 \in S, i_{s_2} \notin S \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \right] \\ &+ \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ i_1 \notin S, i_{s_2} \in S \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \right] + \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ i_1, i_{s_2} \notin S \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \right]. \end{aligned}$$

When $i_1, i_{s_2} \in S$ and $|X(i_1, \dots, i_{s_2})| = T \geq 2$, the choices for $[i_2, \dots, i_{s_2-1}]$ is at most $\binom{s_2-2}{T-2} p^{T-2} r^{s_2-T}$. Thus, by Cauchy-Schwarz,

$$\begin{aligned} (132) \quad \sum_{\substack{i_1, \dots, i_{s_2} \\ i_1, i_{s_2} \in S \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| &\leq \binom{s_2-2}{T-2} p^{T-2} r^{s_2-T} \times \left(\sum_{i_1 \in S} |\mathbf{v}^T u_{i_1}| \right) \left(\sum_{i_{s_2} \in S} |u_{i_{s_2}}^T \mathbf{w}| \right) \\ &\leq \binom{s_2-2}{T-2} p^{T-2} r^{s_2-T} \times \sqrt{p} \sqrt{p} \\ &= \binom{s_2-2}{T-2} p^{T-1} r^{s_2-T}. \end{aligned}$$

By a similar argument, we also have

$$(133) \quad \sum_{\substack{i_1, \dots, i_{s_2} \\ i_1 \in S, i_{s_2} \notin S \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \leq \binom{s_2-2}{T-1} p^{T-1} r^{s_2-T-1} \sqrt{pr}.$$

$$(134) \quad \sum_{\substack{i_1, \dots, i_{s_2} \\ i_1 \notin S, i_{s_2} \in S \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \leq \binom{s_2-2}{T-1} p^{T-1} r^{s_2-T-1} \sqrt{pr}.$$

$$(135) \quad \sum_{\substack{i_1, \dots, i_{s_2} \\ i_1, i_{s_2} \notin S \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \leq \binom{s-2}{T} p^T r^{s-2-T} \sqrt{r} \sqrt{r} = \binom{s-2}{T} p^T r^{s-1-T}.$$

Combining (132), (133), (134), and (135), we obtain

$$\begin{aligned}
(136) \quad \sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| &\leq \binom{s_2-2}{T-2} p^{T-1} r^{s_2-T} + 2 \binom{s_2-2}{T-1} \sqrt{pr} p^{T-1} r^{s_2-T-1} + \\
&+ \binom{s_2-2}{T} p^T r^{s_2-T-1} \\
&\leq \binom{s_2}{T} p^{T-1} r^{s_2-T}.
\end{aligned}$$

Combining (131) and (136), we have

$$\begin{aligned}
(137) \quad \|\Sigma_2\| &\leq \sum_{T=2}^{s_2} \binom{s_2}{T} p^{T-1} r^{s_2-T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) > 0} \frac{\|E\|^{s_1-(k-1)} w^{2(k-1)} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}} \\
&= M_1(\alpha; \beta) + M_2(\alpha; \beta), \text{ where} \\
M_1(\alpha; \beta) &:= \sum_{T=2}^{s_2} \binom{s_2}{T} p^{T-1} r^{s_2-T} \sum_{L \in \mathcal{L}(T, s); 1 \leq r_c(T, L) \leq k-1} \frac{\|E\|^{s_1-(k-1)} w^{2(k-1)} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}}, \\
M_2(\alpha; \beta) &:= \sum_{T=2}^{s_2} \binom{s_2}{T} p^{T-1} r^{s_2-T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L) > k-1} \frac{\|E\|^{s_1-(k-1)} w^{2(k-1)} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}}.
\end{aligned}$$

Similar to the arguments of bounding $M_1(\alpha; \beta)$, $M_2(\alpha; \beta)$ for $(\alpha; \beta)$ of Type I from the previous subsection, we obtain the following lemmas.

Lemma 9.10. *For any pair $(\alpha; \beta)$ of Type II,*

$$(138) \quad M_1(\alpha; \beta) \leq \left(\frac{\sqrt{p} \|E\|}{\bar{\lambda}} \right) \frac{\sum_{T=2}^{s_2} \sum_{L \in \mathcal{L}(T, s); 1 \leq r_c(T, L) \leq k-1} \binom{s_2}{T}}{12^{s-1}}.$$

Lemma 9.11. *For any pair $(\alpha; \beta)$ of Type II,*

$$(139) \quad M_2(\alpha; \beta) \leq \frac{px}{\bar{\lambda}} \frac{\sum_{T=2}^{s_2} \sum_{L \in \mathcal{L}(T, s); r_c > k-1} \binom{s_2}{T}}{12^{s-1}}.$$

The proofs of these lemmas can be found in Section 13. Combining (137), Lemma 9.10 and Lemma 9.11, we finally obtain

$$\begin{aligned}
\|\Sigma_2\| &\leq M_1(\alpha; \beta) + M_2(\alpha; \beta) \\
&\leq \frac{px}{\bar{\lambda}} \frac{\sum_{T=2}^{s_2} \sum_{L \in \mathcal{L}(T, s); r_c > k-1} \binom{s_2}{T}}{12^{s-1}} + \left(\frac{\sqrt{p} \|E\|}{\bar{\lambda}} \right) \frac{\sum_{T=2}^{s_2} \sum_{L \in \mathcal{L}(T, s); 1 \leq r_c(T, L) \leq k-1} \binom{s_2}{T}}{12^{s-1}}.
\end{aligned}$$

This proves Lemma 9.9.

9.11.1. *Proof of Lemma 9.8 - first part.* Recall that in this case, $(s_2, s_1) \neq (k, k-1)$. Together with definition of (s_2, s_1) , we have

$$(140) \quad \max\{s_2 - k, s_1 - (k-1)\} \geq 1.$$

Back to our estimation on $\|\Sigma_1\|$. First, repeating the argument on bounding $\|\Sigma_2\|$ up to (131), we also obtain

$$(141) \quad \|\Sigma_1\| \leq \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{T=1}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L)=0}} \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \right] \frac{\|E\|^{s_1-(k-1)} w^{2(k-1)} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T,L)} \bar{\lambda}^{s_1+r_c(T,L)}}.$$

Similar to (136), if $T \geq 2$,

$$\sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \leq \binom{s_2}{T} p^{T-1} r^{s_2-T}.$$

If $T = 1$, the first case $i_1, i_{s_2} \in S$ doesn't exist, combining (133), (134), and (135), we obtain

$$(142) \quad \sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=1}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| = \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ i_1 \in S, i_{s_2} \notin S \\ |X(i_1, \dots, i_{s_2})|=1}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \right] + \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ i_1 \notin S, i_{s_2} \in S \\ |X(i_1, \dots, i_{s_2})|=1}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \right] + \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ i_1, i_{s_2} \notin S \\ |X(i_1, \dots, i_{s_2})|=1}} |\mathbf{v}^T u_{i_1}| |u_{i_{s_2}}^T \mathbf{w}| \right] \\ \leq 2 \binom{s_2-2}{0} \sqrt{pr} r^{s_2-2} + \binom{s_2-2}{1} pr^{s_2-2} \\ \leq s_2 \sqrt{pr} r^{s_2-2}.$$

Combining (141), (136), and (142), we obtain

$$\|\Sigma_1\| \leq M_3(\alpha; \beta) + M_4(\alpha; \beta), \text{ where}$$

$$(143) \quad M_3(\alpha; \beta) := \sum_{L \in \mathcal{L}(1,s)} s_2 \sqrt{pr} r^{s_2-2} \frac{\|E\|^{s_1-(k-1)} w^{2(k-1)} x^{s_2-k}}{\delta_S^{s_2-1} \bar{\lambda}^{s_1}} \quad (T=1, r_c(T,L)=0),$$

$$M_4(\alpha; \beta) := \sum_{T=2}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L)=0}} \binom{s_2}{T} p^{T-1} r^{s_2-T} \frac{\|E\|^{s_1-(k-1)} w^{2(k-1)} x^{s_2-k}}{\delta_S^{s_2-1} \bar{\lambda}^{s_1}}.$$

We can rewrite $M_3(\alpha; \beta)$ as

$$M_3(\alpha; \beta) = s_2 \sum_{L \in \mathcal{L}(1, s)} \sqrt{\frac{p}{r}} \left(\frac{rx}{\delta_S} \right)^{s_2 - k} \left(\frac{\|E\|}{\lambda} \right)^{s_1 - (k-1)} \left(\frac{\sqrt{rw}}{\sqrt{\lambda \delta_S}} \right)^{2(k-1)}.$$

Notice that by definition of (s_1, s_2) , $s_2 \geq k$, $s_1 \geq k - 1$, all the exponents are non negative. Applying Lemma 9.4 under condition (140), we have

(144)

$$\begin{aligned} M_3(\alpha; \beta) &\leq s_2 \sum_{L \in \mathcal{L}(1, s)} \sqrt{\frac{p}{r}} \left(\frac{rx}{\delta_S} + \frac{\|E\|}{\lambda} \right) \max \left\{ \frac{\|E\|}{\lambda}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda \delta_S}} \right\}^{s_1 + s_2 - 2} \\ &= s_2 \sum_{L \in \mathcal{L}(1, s)} \sqrt{\frac{p}{r}} \left(\frac{rx}{\delta_S} + \frac{\|E\|}{\lambda} \right) \max \left\{ \frac{\|E\|}{\lambda}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda \delta_S}} \right\}^{s-1} \quad (\text{since } s_1 + s_2 = s + 1) \\ &= s_2 \sum_{L \in \mathcal{L}(1, s)} \left(\frac{\sqrt{pr}x}{\delta_S} + \frac{\sqrt{p}\|E\|}{\sqrt{r\lambda}} \right) \max \left\{ \frac{\|E\|}{\lambda}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda \delta_S}} \right\}^{s-1} \\ &\leq s_2 \sum_{L \in \mathcal{L}(1, s)} \left(\frac{\sqrt{pr}x}{\delta_S} + \frac{\sqrt{p}\|E\|}{\sqrt{r\lambda}} \right) \frac{1}{12^{s-1}} \quad (\text{by Assumption } \mathbf{D0} \text{ and the fact that } \delta \leq \delta_S) \\ &\leq s_2 \sum_{L \in \mathcal{L}(1, s)} \left(\frac{\sqrt{pr}x}{\delta_S} + \frac{\sqrt{p}\|E\|}{\lambda} \right) \frac{1}{12^{s-1}}. \end{aligned}$$

Next, $M_4(\alpha; \beta)$ can be rewritten as

$$\sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L)=0} \left(\frac{p}{r} \right)^{T-1} \left(\frac{rx}{\delta_S} \right)^{s_2 - k} \left(\frac{\|E\|}{\lambda} \right)^{s_1 - (k-1)} \left(\frac{\sqrt{rw}}{\sqrt{\lambda \delta_S}} \right)^{2(k-1)}.$$

By the definition of $M_4(\alpha; \beta)$, $T \geq 2$ and hence $\left(\frac{p}{r}\right)^{T-1} \leq \frac{p}{r}$. Hence,

$$(145) \quad M_4(\alpha; \beta) \leq \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L)=0} \frac{p}{r} \left(\frac{rx}{\delta_S} \right)^{s_2 - k} \left(\frac{\|E\|}{\lambda} \right)^{s_1 - (k-1)} \left(\frac{\sqrt{rw}}{\sqrt{\lambda \delta_S}} \right)^{2(k-1)}.$$

Note that by the definition of (s_1, s_2) , all exponents are non-negative. Applying Lemma 9.4 under condition (140), we have

(146)

$$\begin{aligned} \left(\frac{rx}{\delta_S} \right)^{s_2 - k} \left(\frac{\|E\|}{\lambda} \right)^{s_1 - (k-1)} \left(\frac{\sqrt{rw}}{\sqrt{\lambda \delta_S}} \right)^{2(k-1)} &\leq \left(\frac{\|E\|}{\lambda} + \frac{rx}{\delta_S} \right) \max \left\{ \frac{\|E\|}{\lambda}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda \delta_S}} \right\}^{s_1 + s_2 - 2} \\ &= \left(\frac{\|E\|}{\lambda} + \frac{rx}{\delta_S} \right) \max \left\{ \frac{\|E\|}{\lambda}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda \delta_S}} \right\}^{s-1}. \end{aligned}$$

Combining (145) and (146), we have

$$\begin{aligned}
M_4(\alpha; \beta) &\leq \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L)=0} \frac{p}{r} \left(\frac{\|E\|}{\lambda} + \frac{rx}{\delta_S} \right) \max \left\{ \frac{\|E\|}{\lambda}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right\}^{s-1} \\
(147) \quad &= \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T, s); r_c(T, L)=0} \left(\frac{p\|E\|}{r\lambda} + \frac{px}{\delta_S} \right) \max \left\{ \frac{\|E\|}{\lambda}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right\}^{s-1} \\
&\leq \left(\frac{px}{\delta_S} + \frac{p\|E\|}{r\lambda} \right) \frac{\sum_{T=2}^{s_2} \sum_{L \in \mathcal{L}(T, s); r_c(T, L)=0} \binom{s_2}{T}}{12^{s-1}}, \\
&\quad \text{(by Assumption **D0** and the fact that } \delta \leq \delta_S \text{).}
\end{aligned}$$

Together (143), (144) and (147) imply that if $(s_2, s_1) \neq (k, k-1)$, then

$$\begin{aligned}
(148) \quad \|\Sigma_1\| &\leq M_3(\alpha; \beta) + M_4(\alpha; \beta) \\
&\leq s_2 \sum_{L \in \mathcal{L}(1, s)} \left(\frac{\sqrt{pr}x}{\delta_S} + \frac{\sqrt{p}\|E\|}{\lambda} \right) \frac{1}{12^{s-1}} + \left(\frac{px}{\delta_S} + \frac{p\|E\|}{r\lambda} \right) \frac{\sum_{T=2}^{s_2} \sum_{L \in \mathcal{L}(T, s); r_c(T, L)=0} \binom{s_2}{T}}{12^{s-1}} \\
&\leq s_2 \sum_{L \in \mathcal{L}(1, s)} \left(\frac{\sqrt{pr}x}{\delta_S} + \frac{\sqrt{p}\|E\|}{\lambda} \right) \frac{1}{12^{s-1}} + \left(\frac{\sqrt{pr}x}{\delta_S} + \frac{\sqrt{p}\|E\|}{\lambda} \right) \frac{\sum_{T=2}^{s_2} \sum_{L \in \mathcal{L}(T, s); r_c(T, L)=0} \binom{s_2}{T}}{12^{s-1}} \\
&= \left(\frac{\sqrt{pr}x}{\delta_S} + \frac{\sqrt{p}\|E\|}{\lambda} \right) \frac{\sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T, s); r_c(T, L)=0} \binom{s_2}{T}}{12^{s-1}}.
\end{aligned}$$

This proves the first part of Lemma 9.8.

9.11.2. *Proof of Lemma 9.8 - second part.* Now we consider the very special case when $(s_2, s_1) = (k, k-1)$. In this case,

$$\begin{aligned}
(149) \quad M(\alpha, \beta) &= PEQEPEQEPEQ \dots PEQEP, \\
\alpha_1 &= \alpha_{k+1} = 0, \alpha_2 = \dots = \alpha_k = \beta_1 = \dots = \beta_k = 1, \\
s &= s_1 + s_2 - 1 = 2k - 2.
\end{aligned}$$

Again, write $P = \sum_{i \in N_{\bar{\lambda}}(S)} \frac{1}{z - \lambda_i} u_i u_i^T$ and $Q = \sum_{j \in N_{\bar{\lambda}}(S)^c} \frac{u_j u_j^T}{z - \lambda_j}$. Multiplying out all the terms, we can express $PEQEPEQE \dots PEQEP$ as

$$\sum_{i_1, i_2, \dots, i_k \in N_{\bar{\lambda}}(S)} \sum_{j_1, j_2, \dots, j_{k-1} \in N_{\bar{\lambda}}(S)^c} \frac{1}{(z - \lambda_{i_k}) \prod_{l=1}^{k-1} [(z - \lambda_{i_l})(z - \lambda_{j_l})]} u_{i_1} u_{i_k}^T \prod_{l=1}^{k-1} (u_{i_l}^T E u_{j_l} u_{j_l}^T E u_{i_{l+1}}).$$

Following the procedure above, we can expand $\frac{1}{2\pi i} \int_{\Gamma} PEQEPEQ\dots PEQEPEdz$ as

$$(150) \quad \sum_{i_1, i_2, \dots, i_k} \sum_{j_1, j_2, \dots, j_{k-1}} \sum_{L \in \mathcal{L}(T, 2k-2)} (-1)^{T+1} w(X, Y|L) u_{i_1} \left[\prod_{l=1}^{k-1} (u_{i_l}^T E u_{j_l} u_{j_l}^T E u_{i_{l+1}}) \right] u_{i_k}^T.$$

Splitting $w(X, Y|L) = \prod_{e \in \mathcal{E}_L} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} = \left[\prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \right] \times \prod_{l=1}^{s_1} \left[\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_l}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \right]$ and distributing the sum over j_1, \dots, j_{k-1} according to each position of $u_{j_l} u_{j_l}^T$ in the product, (150) becomes

$$\sum_{i_1, i_2, \dots, i_k} \sum_{L \in \mathcal{L}(T, s)} \frac{(-1)^{T+1}}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} (\lambda_{e^+} - \lambda_{e^-})} u_{i_1} \prod_{l=1}^{k-1} \left[u_{i_l}^T E \left(\sum_{j_l \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_l} u_{j_l}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_l}} (\lambda_{e^+} - \lambda_{j_l})} \right) E u_{i_{l+1}} \right] u_{i_k}^T.$$

Therefore, in this particular case, Σ_1 (defined in (118)) becomes

(151)

$$\sum_{i_1, i_2, \dots, i_k} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) = 0}} \frac{(-1)^{T+1}}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} (\lambda_{e^+} - \lambda_{e^-})} u_{i_1} \prod_{l=1}^{k-1} \left[u_{i_l}^T E \left(\sum_{j_l \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_l} u_{j_l}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_l}} (\lambda_{e^+} - \lambda_{j_l})} \right) E u_{i_{l+1}} \right] u_{i_k}^T.$$

Recall that $r_c(T, L) := \sum_{l=1}^{s_1} (d(j_l) - 1)$. So the condition $r_c(T, L) = 0$ implies that $d(j_l) = 1$ for all $1 \leq l \leq k-1$. It means: j_l is connected to a unique vertex e_l^+ in the graph $G(X, Y|L)$. Therefore, Σ_1 equals

$$\sum_{i_1, i_2, \dots, i_k} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) = 0}} \frac{(-1)^{T+1}}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} (\lambda_{e^+} - \lambda_{e^-})} u_{i_1} \prod_{l=1}^{k-1} \left[u_{i_l}^T E \left(\sum_{j_l \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_l} u_{j_l}^T}{(\lambda_{e_l^+} - \lambda_{j_l})} \right) E u_{i_{l+1}} \right] u_{i_k}^T.$$

Similar to the argument of bounding $\|\Sigma_1\|$ from the previous case, we have

(152)

$$\begin{aligned} \|\Sigma_1\| &\leq \sum_{\substack{i_1, i_2, \dots, i_k \\ L \in \mathcal{L}(T, 2k-1) \\ r_c(T, L) = 0}} \max_{\|\mathbf{v}\| = \|\mathbf{w}\| = 1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \frac{1}{\delta_S^{|E(I_2)|}} \prod_{l=1}^{k-1} \left| u_{i_l}^T E \left(\sum_{j_l \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_l} u_{j_l}^T}{\lambda_{e_l^+} - \lambda_{j_l}} \right) E u_{i_{l+1}} \right| \\ &\leq \sum_{\substack{i_1, i_2, \dots, i_k \\ L \in \mathcal{L}(T, 2k-1) \\ r_c(T, L) = 0}} \max_{\|\mathbf{v}\| = \|\mathbf{w}\| = 1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \frac{1}{\delta_S^{k-1}} \prod_{l=1}^{k-1} \left| u_{i_l}^T E \left(\sum_{j_l \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_l} u_{j_l}^T}{\lambda_{e_l^+} - \lambda_{j_l}} \right) E u_{i_{l+1}} \right|, \end{aligned}$$

since $|E(I_2)| = s_2 - 1 - r_c(T, L) = k - 1$.

Consider the special case when all the elements of X are the same, namely, $X = [i, i, \dots, i]$, with k repetitions, $Y = [j_1, \dots, j_{k-1}]$, and $L \in \mathcal{L}(T, 2k-2)$. In this case, the graph $G(X, Y|L)$ is non-trivial if $I \in S$ and $T = k$. Since $|Y| = k-1 = \frac{|\mathcal{E}(X, Y|L)|}{2}$, there exists at least one vertex j_l such that $d(j_l) \geq 2$, which implies that $r_c(X, Y|L) > 0$. Therefore, the RHS of (152) does not contain any sequence $[i_1, i_2, \dots, i_k] = [i, i, \dots, i]$. We call a sequence $[i_1, i_2, \dots, i_k] \neq [i, i, \dots, i]$ "non-uniform", and rewrite (152) as

$$\|\Sigma_1\| \leq \sum_{\substack{\text{non-uniform} \\ i_1, i_2, \dots, i_k \\ L \in \mathcal{L}(T, 2k-1); r_c(k, L)=0}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \frac{1}{\delta_S^{k-1}} \prod_{l=1}^{k-1} \left| u_{i_l}^T E \left(\sum_{j_l \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_l} u_{j_l}^T}{\lambda_{e_l^+} - \lambda_{j_l}} \right) E u_{i_{l+1}} \right|.$$

Observe that for any $1 \leq l \leq k-1$, by Definition 2.2,

$$\left| u_{i_l}^T E \left(\sum_{j_l \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_l} u_{j_l}^T}{\lambda_{e_l^+} - \lambda_{j_l}} \right) E u_{i_{l+1}} \right| \leq \|u_{i_l}^T E\| \left\| \sum_{j_l \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_l} u_{j_l}^T}{\lambda_{e_l^+} - \lambda_{j_l}} \right\| \|E u_{i_{l+1}}\| \leq \frac{w^2}{\bar{\lambda}}.$$

Furthermore, since $[i_1, \dots, i_k] \neq [i, \dots, i]$, there exists at least one pair $i_{l'} \neq i_{l'+1}$, and hence

$$\left| u_{i_{l'}}^T E \left(\sum_{j_{l'} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l'}} u_{j_{l'}}^T}{\lambda_{e_{l'}^+} - \lambda_{j_{l'}}} \right) E u_{i_{l'+1}} \right| \leq y \text{ (by Definition 2.2).}$$

Together these observations imply that

$$(153) \quad \prod_{l=1}^{k-1} \left| u_{i_l}^T E \left(\sum_{j_l \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_l} u_{j_l}^T}{\lambda_{e_l^+} - \lambda_{j_l}} \right) E u_{i_{l+1}} \right| \leq \frac{y w^{2(k-2)}}{\bar{\lambda}^{k-2}}.$$

Therefore, (152) implies

$$\|\Sigma_1\| \leq \sum_{\substack{\text{non-uniform} \\ i_1, i_2, \dots, i_k}} \sum_{\substack{L \in \mathcal{L}(T, 2k-1) \\ r_c(k, L)=0}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \frac{1}{\delta_S^{k-1}} \frac{y w^{2(k-2)}}{\bar{\lambda}^{k-2}}.$$

By a well-known combinatorial fact,

$$|\mathcal{L}(T, 2k-1)| = \binom{(2k-1)-1}{T-1} = \binom{2k-2}{T-1} \leq 2^{2k-2}$$

for $T \leq 2k-1$. Thus, the last bound on $\|\Sigma_1\|$ implies

$$(154) \quad \begin{aligned} \|\Sigma_1\| &\leq \sum_{\substack{\text{non-uniform} \\ i_1, i_2, \dots, i_k}} 2^{2k-2} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \frac{y w^{2(k-2)}}{\bar{\lambda}^{k-2} \delta_S^{k-1}} \\ &= \left(\sum_{\substack{\text{non-uniform} \\ i_1, i_2, \dots, i_k}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \right) 2^{2k-2} \frac{y w^{2(k-2)}}{\bar{\lambda}^{k-2} \delta_S^{k-1}}. \end{aligned}$$

As there is at least one index i_l such that $i_l \in S$, we have

$$\begin{aligned}
(155) \quad \sum_{\substack{\text{non-uniform} \\ i_1, i_2, \dots, i_k}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| &\leq \sum_{l=1}^k \sum_{\substack{i_1, i_2, \dots, i_k \\ i_l \in S}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \\
&= \sum_{l=2}^{k-1} \sum_{\substack{i_1, i_2, \dots, i_k \\ i_l \in S}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \\
&\quad + \sum_{\substack{i_1, i_2, \dots, i_k \\ i_1 \in S}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \\
&\quad + \sum_{\substack{i_1, i_2, \dots, i_k \\ i_k \in S}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}|.
\end{aligned}$$

Similar to the argument in (132) and (142), we have

$$(156) \quad \sum_{\substack{i_1, i_2, \dots, i_k \\ i_l \in S}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \leq \sqrt{r} \sqrt{r} p r^{k-3} \leq \sqrt{p r} r^{k-2}, \text{ for any } 2 \leq l \leq k-1,$$

$$(157) \quad \sum_{\substack{i_1, i_2, \dots, i_k \\ i_1 \in S}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \leq \sqrt{p} \sqrt{r} r^{k-2} = \sqrt{p r} r^{k-2},$$

$$(158) \quad \sum_{\substack{i_1, i_2, \dots, i_k \\ i_k \in S}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \leq \sqrt{r} \sqrt{p r} r^{k-2} = \sqrt{p r} r^{k-2}.$$

Together (155), (156), (157), and (158) imply that

$$(159) \quad \sum_{\substack{\text{non-uniform} \\ i_1, i_2, \dots, i_k}} \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} |\mathbf{v}^T u_{i_1}| |u_{i_k}^T \mathbf{w}| \leq k \sqrt{p r} r^{k-2} = k \sqrt{p r} r^{k-2}.$$

Combining (154) and (159), we obtain

$$\|\Sigma_1\| \leq k \sqrt{p r} r^{k-2} \times 2^{2k-2} \frac{y w^{2(k-2)}}{\bar{\lambda}^{k-2} \delta_S^{k-1}}.$$

As $k = s/2 + 1$, we can write this as

$$\|\Sigma_1\| \leq \frac{(s+2) \sqrt{p r} y}{\delta_S} \left(\frac{2 \sqrt{r} w}{\sqrt{\bar{\lambda} \delta_S}} \right)^{s-2}.$$

This proves the second part of Lemma 9.8.

9.12. **Proof of Lemma 8.9.** In the setting of this lemma, $s_1 \geq k$. Without loss of generality, it is sufficient to handle the case that $\alpha_1 = 0, \alpha_{k+1} > 0$. Similar to previous subsections, we have

$$\begin{aligned}
& \frac{1}{2\pi i} \int_{\Gamma} M(\alpha; \beta) dz \\
&= \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} \sum_{j_1, \dots, j_{s_1}} (-1)^{T+1} \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \times \left[\prod_{l=1}^{\beta_1} (u_{i_l} u_{i_l}^T E) \right] \times \\
& \left\{ \prod_{h=1}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_h}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_{l+B_h}} u_{i_{l+B_h}}^T E) \right] \right\} \times \\
& \times \left[\prod_{l=1}^{\alpha_{k+1}-1} \left(\frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_k}}} (\lambda_{e^+} - \lambda_{j_{l+A_k}})} E \right) \right] \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{s_1}}} (\lambda_{e^+} - \lambda_{j_{s_1}})}.
\end{aligned}$$

Once again, we distribute the sum over j_1, j_2, \dots, j_{s_2} according to each position of $u_{j_l} u_{j_l}^T$ in the product, and obtain

$$\begin{aligned}
& \frac{1}{2\pi i} \int_{\Gamma} M(\alpha; \beta) dz \\
&= \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} (-1)^{T+1} \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \times \left[\prod_{l=1}^{\beta_1} (u_{i_l} u_{i_l}^T E) \right] \times \\
& \left\{ \prod_{h=1}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\sum_{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{e \in \mathcal{E}_L; e^- = j_{l+A_h}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_{l+B_h}} u_{i_{l+B_h}}^T E) \right] \right\} \times \\
& \times \left[\prod_{l=1}^{\alpha_{k+1}-1} \left(\sum_{j_{l+A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_k}}} (\lambda_{e^+} - \lambda_{j_{l+A_k}})} E \right) \right] \times \left(\sum_{j_{s_1} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{s_1}}} (\lambda_{e^+} - \lambda_{j_{s_1}})} \right).
\end{aligned}$$

Next, by (125),

$$\begin{aligned}
& \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| \\
&= \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} (-1)^{T+1} \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \times \left[\mathbf{v}^T \prod_{l=1}^{\beta_1} (u_{i_l} u_{i_l}^T E) \right] \times \\
& \left\{ \prod_{h=1}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\sum_{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{\substack{e \in \mathcal{E}_L; e^- = j_{l+A_h}}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_{l+B_h}} u_{i_{l+B_h}}^T E) \right] \right\} \times \\
& \times \left[\prod_{l=1}^{\alpha_{k+1}-1} \left(\sum_{j_{l+A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_k}}} (\lambda_{e^+} - \lambda_{j_{l+A_k}})} E \right) \right] \times \left(\sum_{j_{s_1} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{s_1}}} (\lambda_{e^+} - \lambda_{j_{s_1}})} \right) \mathbf{w}.
\end{aligned}$$

By the triangle inequality, the RHS is at most

$$\begin{aligned}
& \max_{\|\mathbf{v}\|=\|\mathbf{w}\|=1} \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} \left| \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \right| \times \left\| \mathbf{v}^T \prod_{l=1}^{\beta_1} (u_{i_l} u_{i_l}^T E) \right\| \times \\
& \left\| \prod_{h=1}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\sum_{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{\prod_{\substack{e \in \mathcal{E}_L; e^- = j_{l+A_h}}} (\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_{l+B_h}} u_{i_{l+B_h}}^T E) \right] \right\| \times \\
& \times \left\| \prod_{l=1}^{\alpha_{k+1}-1} \left(\sum_{j_{l+A_k} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{l+A_k}}} (\lambda_{e^+} - \lambda_{j_{l+A_k}})} E \right) \right\| \times \left\| \sum_{j_{s_1} \in N_{\bar{\lambda}}(S)^c} \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{\prod_{\substack{e \in \mathcal{E}_L \\ e^- = j_{s_1}}} (\lambda_{e^+} - \lambda_{j_{s_1}})} \mathbf{w} \right\|.
\end{aligned}$$

Since $\|w\| = 1$, this is at most

$$\begin{aligned} & \max_{\|\mathbf{v}\|=1} \sum_{i_1, i_2, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} \left| \prod_{\substack{e \in \mathcal{E}_L \\ e^- \in I_2}} \frac{1}{(\lambda_{e^+} - \lambda_{e^-})} \right| \times \left\| \mathbf{v}^T \prod_{l=1}^{\beta_1} (u_{i_l} u_{i_l}^T E) \right\| \times \\ & \left\| \prod_{h=1}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} \left(\sum_{\substack{j_{l+A_h} \in N_{\bar{\lambda}}(S)^c \\ e \in \mathcal{E}_L; e^- = j_{l+A_h}}} \frac{u_{j_{l+A_h}} u_{j_{l+A_h}}^T}{(\lambda_{e^+} - \lambda_{j_{l+A_h}})} E \right) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} (u_{i_{l+B_h}} u_{i_{l+B_h}}^T E) \right] \right\| \times \\ & \times \left\| \prod_{l=1}^{\alpha_{k+1}-1} \left(\sum_{\substack{j_{l+A_k} \in N_{\bar{\lambda}}(S)^c \\ e \in \mathcal{E}_L \\ e^- = j_{l+A_k}}} \frac{u_{j_{l+A_k}} u_{j_{l+A_k}}^T}{(\lambda_{e^+} - \lambda_{j_{l+A_k}})} E \right) \right\| \times \left\| \sum_{\substack{j_{s_1} \in N_{\bar{\lambda}}(S)^c \\ e \in \mathcal{E}_L \\ e^- = j_{s_1}}} \frac{u_{j_{s_1}} u_{j_{s_1}}^T}{(\lambda_{e^+} - \lambda_{j_{s_1}})} \right\|. \end{aligned}$$

Using the estimates in (127), (128), (129), and (106), we obtain

$$\begin{aligned} (160) \quad & \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| \leq \max_{\|\mathbf{v}\|=1} \sum_{i_1, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} |\mathbf{v}^T u_{i_1}| \frac{w^{2k-1} x^{\sum_{l=1}^k (\beta_l - 1)} \|E\|^{\sum_{l=2}^{k+1} (\alpha_l - 1)}}{\delta_S^{|E(I_2)|} \bar{\lambda}^{\sum_{l=1}^{s_1} d(j_l)}} \\ & = \max_{\|\mathbf{v}\|=1} \sum_{i_1, \dots, i_{s_2}} \sum_{L \in \mathcal{L}(T, s)} |\mathbf{v}^T u_{i_1}| \frac{\|E\|^{s_1-k} w^{2k-1} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}} \\ & = \max_{\|\mathbf{v}\|=1} \sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T, s)} \left[\sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| \right] \frac{\|E\|^{s_1-k} w^{2k-1} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T, L)} \bar{\lambda}^{s_1+r_c(T, L)}}. \end{aligned}$$

Moreover, similar to the argument in (132), by Cauchy-Schwarz, we have

$$\begin{aligned} \sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| &= \sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=T \\ i_1 \in S}} |\mathbf{v}^T u_{i_1}| + \sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=T \\ i_1 \in N_{\bar{\lambda}}(S) \setminus S}} |\mathbf{v}^T u_{i_1}| \\ &\leq \binom{s_2-1}{T-1} \sqrt{p} p^{T-1} r^{s_2-T} + \binom{s_2-1}{T} \sqrt{r} p^T r^{s_2-T-1}. \end{aligned}$$

Since $\sqrt{r} p^T r^{s_2-T-1} = p^T r^{s_2-T-1/2} \leq p^{T-1/2} r^{s_2-T}$ and $\binom{s_2-1}{T-1} + \binom{s_2-1}{T} = \binom{s_2}{T}$, we have

$$(161) \quad \sum_{\substack{i_1, \dots, i_{s_2} \\ |X(i_1, \dots, i_{s_2})|=T}} |\mathbf{v}^T u_{i_1}| \leq \binom{s_2}{T} p^{T-1/2} r^{s_2-T}.$$

Combining (160) and (161), we obtain

(162)

$$\begin{aligned} \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} p^{T-1/2} r^{s_2-T} \sum_{L \in \mathcal{L}(T,s)} \frac{\|E\|^{s_1-k} w^{2k} x^{s_2-k}}{\delta_S^{s_2-1-r_c} \bar{\lambda}^{s_1+r_c}} \\ &= M_1(\alpha; \beta) + M_2(\alpha; \beta), \text{ where} \\ M_1(\alpha; \beta) &:= \sum_{T=1}^{s_2} \binom{s_2}{T} p^{T-1/2} r^{s_2-T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} \frac{\|E\|^{s_1-k} w^{2k-1} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T,L)} \bar{\lambda}^{s_1+r_c(T,L)}}, \\ M_2(\alpha; \beta) &:= \sum_{T=1}^{s_2} \binom{s_2}{T} p^{T-1/2} r^{s_2-T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) > k-1} \frac{\|E\|^{s_1-k} w^{2k-1} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T,L)} \bar{\lambda}^{s_1+r_c(T,L)}}. \end{aligned}$$

By a similar argument of bounding $M_1(\alpha; \beta)$, $M_2(\alpha; \beta)$ from previous subsections, we obtain the following lemmas.

Lemma 9.13. *For any pair $(\alpha; \beta)$ of Type III,*

$$(163) \quad M_1(\alpha; \beta) \leq \frac{\sqrt{p} \|E\|}{\bar{\lambda}} \times \frac{2^{s_2+s-1}}{12^{s-1}}.$$

Lemma 9.14. *For any pair $(\alpha; \beta)$ of Type III,*

$$(164) \quad M_2(\alpha; \beta) \leq \frac{px}{\bar{\lambda}} \frac{2^{s_2+s-1}}{12^{s-1}}.$$

The proofs of these lemmas can be found in Section 13. Combining (162), Lemma 9.14 and Lemma 9.13, we finally have

$$(165) \quad \begin{aligned} \frac{1}{2\pi} \left\| \int_{\Gamma} M(\alpha; \beta) dz \right\| &\leq M_1(\alpha; \beta) + M_2(\alpha; \beta) \\ &\leq \left(\frac{\sqrt{p} \|E\|}{\bar{\lambda}} + \frac{px}{\bar{\lambda}} \right) \times \frac{2^{s_2+s-1}}{12^{s-1}}. \end{aligned}$$

This proves Lemma 8.9.

9.15. Proof of Expansion (36). To conclude the proof of Theorem 2.12, we still need to prove that (under Assumption **D0**) the series $\sum_{s=1}^{\infty} \|(z-A)^{-1}[E(z-A)^{-1}]^s\|$ converges.

Define $G_s := \|(z-A)^{-1}[E(z-A)^{-1}]^s\|$. By the triangle inequality, we have

$$(166) \quad G_s \leq \sum_{(\alpha; \beta)} \|M(\alpha; \beta)\|,$$

in which $\alpha, \beta, M(\alpha; \beta)$ are as defined in Subsection 8.6.

Lemma 9.16. *For any pair (α, β) ,*

$$(167) \quad \|M(\alpha, \beta)\| \leq \frac{2r}{6^s \delta}.$$

Combining (166), Lemma 9.16, and the fact that there are exactly 2^{s+1} pairs (α, β) in the expansion of G_s , we obtain

$$G_s \leq 2^{s+1} \times \frac{2r}{6^s \delta} = \frac{4r}{3^s \delta}.$$

Therefore,

$$\sum_{s=1}^{\infty} \|(z - A)^{-1} [E(z - A)^{-1}]^s\| \leq \sum_{s=1}^{\infty} \frac{4r}{3^s \delta} = \frac{2r}{\delta}.$$

So, our last duty is to prove Lemma 9.16. We first bound $\|M(\alpha; \beta)\|$ for $(\alpha; \beta)$ of Type I; the proofs for the other types are similar. Multiplying out all the terms in the operators P that appear in $M(\alpha; \beta)$ (there are s_2 such operators), we obtain

$$\begin{aligned} & M(\alpha; \beta) \\ &= \sum_{i_1, i_2, \dots, i_{s_2}} \frac{1}{\prod_{l=1}^{s_2} (z - \lambda_{i_l})} \left\{ \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}} (QE) \right] \times \left[\prod_{l=1}^{\beta_{h+1}} \left(u_{i_l+B_h} u_{i_l+B_h}^T E \right) \right] \right\} \times \left[\prod_{l=1}^{\alpha_{k+1}-1} (QE) \right] Q \\ &= \sum_{i_1, i_2, \dots, i_{s_2}} \frac{1}{\prod_{l=1}^{s_2} (z - \lambda_{i_l})} \left\{ \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}-1} (QE) \right] QE \left[\prod_{l=1}^{\beta_{h+1}} \left(u_{i_l+B_h} u_{i_l+B_h}^T E \right) \right] \right\} \times \left[\prod_{l=1}^{\alpha_{k+1}-1} (QE) \right] Q \\ &= \sum_{i_1, i_2, \dots, i_{s_2}} \frac{1}{\prod_{l=1}^{s_2} (z - \lambda_{i_l})} \left\{ \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}-1} (QE) \right] Q \left[E \prod_{l=1}^{\beta_{h+1}} \left(u_{i_l+B_h} u_{i_l+B_h}^T E \right) \right] \right\} \times \left[\prod_{l=1}^{\alpha_{k+1}-1} (QE) \right] Q. \end{aligned}$$

By the triangle inequality,

$$\begin{aligned} & (168) \\ & \|M(\alpha; \beta)\| \leq \\ & \sum_{i_1, i_2, \dots, i_{s_2}} \left| \frac{1}{\prod_{l=1}^{s_2} (z - \lambda_{i_l})} \right| \times \prod_{h=0}^{k-1} \left[\prod_{l=1}^{\alpha_{h+1}-1} \|QE\| \right] \|Q\| \left\| E \prod_{l=1}^{\beta_{h+1}} \left(u_{i_l+B_h} u_{i_l+B_h}^T E \right) \right\| \times \left[\prod_{l=1}^{\alpha_{k+1}-1} \|QE\| \right] \|Q\|. \end{aligned}$$

On the other hand, Assumption **D0** implies that $|z - \lambda_{j_l}| \geq \bar{\lambda}/2$ for all z on the contour and $j_l \in N_{\bar{\lambda}}(S)^c$ and that $|z - \lambda_i| \geq \delta/2$ for all z on the contour and eigenvalue $\lambda_i, i \in N_{\bar{\lambda}}(S)$. Thus, we have

$$(169) \quad \left| \frac{1}{\prod_{l=1}^{s_2} (z - \lambda_{i_l})} \right| \leq \frac{2^{s_2}}{\delta^{s_2}}$$

and

$$(170) \quad \|Q\| \leq \frac{2}{\bar{\lambda}}.$$

Moreover, the estimate in (103) shows that

$$(171) \quad \left| E \prod_{l=1}^{\beta_{h+1}} \left(u_{i_l+B_h} u_{i_l+B_h}^T E \right) \right| \leq x^{\beta_{h+1}-1} w^2.$$

Combining (168), (169), (171), and the fact that there are exactly $s_1 - (k + 1)$ factors $\|QE\|$, we obtain

$$\begin{aligned} \|M(\alpha; \beta)\| &\leq \sum_{i_1, \dots, i_{s_2}} \frac{x^{s_2-k} w^{2k} \|QE\|^{s_1-1-k} \|Q\|^{k+1} 2^{s_2}}{\delta^{s_2}} \\ &\leq \sum_{i_1, \dots, i_{s_2}} \frac{x^{s_2-k} (\|Q\| \cdot \|E\|)^{s_1-1-k} w^{2k} \|Q\|^{k+1} 2^{s_2}}{\delta^{s_2}} \\ &= \sum_{i_1, \dots, i_{s_2}} \frac{x^{s_2-k} \|E\|^{s_1-1-k} w^{2k} \|Q\|^{s_1} 2^{s_2}}{\delta^{s_2}}. \end{aligned}$$

Since $w \leq \|E\|$, we further obtain

$$\begin{aligned} (172) \quad \|M(\alpha; \beta)\| &\leq \sum_{i_1, \dots, i_{s_2}} \frac{x^{s_2-k} \|E\|^{s_1+1-k} w^{2(k-1)} \|Q\|^{s_1} 2^{s_2}}{\delta^{s_2}} \\ &\leq \sum_{i_1, \dots, i_{s_2}} \frac{x^{s_2-k} \|E\|^{s_1+1-k} w^{2(k-1)} 2^{s_2+s_1}}{\bar{\lambda}^{s_1} \delta^{s_2}} \text{ (by (170)).} \end{aligned}$$

Note that there are r^{s_2} choices for i_1, \dots, i_{s_2} and $s_1 + s_2 = s + 1$. The last formula in (172) becomes

$$2^{s+1} r^{s_2} \frac{x^{s_2-k} \|E\|^{s_1+1-k} w^{2(k-1)}}{\bar{\lambda}^{s_1} \delta^{s_2}} = 2^{s+1} \left(\frac{rx}{\delta}\right)^{s_2-k} \left(\frac{\sqrt{rw}}{\sqrt{\lambda\delta}}\right)^{2(k-1)} \left(\frac{\|E\|}{\lambda}\right)^{s_1+1-k} \frac{r}{\delta}.$$

Applying Lemma 9.4, we finally have

$$\begin{aligned} \|M(\alpha; \beta)\| &\leq \frac{2^{s+1} r}{\delta} \max \left\{ \frac{rx}{\delta}, \frac{\|E\|}{\lambda}, \frac{\sqrt{rw}}{\sqrt{\lambda\delta}} \right\}^{s_2-k+2(k-1)+s_1-(k-1)} \\ &= \frac{2^{s+1} r}{\delta} \max \left\{ \frac{rx}{\delta}, \frac{\|E\|}{\lambda}, \frac{\sqrt{rw}}{\sqrt{\lambda\delta}} \right\}^{s_2+s_1-1} \\ &\leq \frac{2^{s+1} r}{\delta} \left(\frac{1}{12}\right)^{s_2+s_1-1} \text{ (by Assumption D0)} \\ &= \frac{2^{s+1} r}{12^s \delta} \text{ (since } s_1 + s_2 - 1 = s) \\ &= \frac{2r}{6^s \delta}. \end{aligned}$$

We complete our proof for Lemma 9.16 and then the proof of Theorem 2.12.

10. PROOFS OF THE MAIN THEOREMS FROM SECTIONS 5 AND 6

10.1. Proof of Theorems 5.7 and 5.10. In order to prove Theorem 5.10, we need to bound the quantities x, y for the random noise E and then apply Theorem 2.12. First, we use the following lemma to estimate the x -upper bound.

Lemma 10.2. *Let $E = (\xi_{kl})_{k,l=1}^n$ be a $n \times n$ real symmetric random matrix where*

$$\{\xi_{kl} : 1 \leq k \leq l \leq n\}$$

is a collection of independent random variables each with mean zero. Further assume

$$\sup_{1 \leq k \leq l \leq n} |\xi_{kl}| \leq K, \quad \sup_{1 \leq k \leq l \leq n} \left(\mathbf{E}(\xi_{kl}^2) = \sigma_{kl}^2 \right) \leq \sigma^2.$$

with probability 1. Then for any $1 \leq i \leq j \leq r$ and every $t > 0$

$$\mathbf{P}(|u_i^T E u_j| \geq t) \leq \exp\left(\frac{-t^2/2}{2\sigma^2 + Kt}\right).$$

Proof. Note that for any $1 \leq i, j \leq r$,

$$u_i^T E u_j = \sum_{k,l} u_{ik} \xi_{kl} u_{jl} = \sum_{k < l} (u_{ik} u_{jl} + u_{il} u_{jk}) \xi_{kl} + \sum_k u_{ik} u_{jk} \xi_{kk}.$$

Therefore, $\mathbf{E}(u_i^T E u_j) = 0$. Furthermore,

$$\begin{aligned} \mathbf{Var}(u_i^T E u_j) &= \sum_{k < l} (u_{ik} u_{jl} + u_{il} u_{jk})^2 \sigma_{kl}^2 + \sum_k u_{ik}^2 u_{jk}^2 \sigma_{kk}^2 \\ &\leq \sum_{k < l} 2(u_{ik}^2 u_{jl}^2 + u_{il}^2 u_{jk}^2) \sigma_{kl}^2 + \sum_k u_{ik}^2 u_{jk}^2 \sigma_{kk}^2 \leq 2 \sum_{k,l} u_{ik}^2 u_{jl}^2 \sigma_{kl}^2. \end{aligned}$$

By assumption about upper bound of σ_{kl}^2 , we obtain

$$\mathbf{Var}(u_i^T E u_j) \leq 2\sigma^2 \sum_{k,l} u_{ik}^2 u_{jl}^2 = 2\sigma^2 \sum_k u_{ik}^2 \sum_l u_{jl}^2 = 2\sigma^2.$$

By the fact that $|u_{ik} u_{jl} \xi_{kl}| \leq K, \forall k, l, i, j$, we can apply the Bernstein inequality and get

$$(173) \quad \mathbf{P}(|u^T E v| \geq t) \leq \exp\left(\frac{-t^2/2}{2\sigma^2 + Kt}\right), \quad \forall t > 0.$$

□

If all ξ_{ij} additionally are sub-Gaussian random variables, we have

$$\mathbf{P}(|u^T E v| \geq t) \leq \exp\left(\frac{-t^2}{4\sigma^2}\right).$$

To obtain the y -upper bound, we will use moment method to bound $G_{ij} := u_i^T E \left(\sum_{l > r} \frac{u_l u_l^T}{\lambda_k - \lambda_j} \right) E u_j$ for a fixed triple (k, i, j) : $1 \leq i < j \leq r, 1 \leq k \leq p$ and then apply the union bound on all feasible triples (i, j, k) . Indeed, we have the following lemma.

Lemma 10.3. *Let $E = (\xi_{ij})_{1 \leq i, j \leq n}$ be a random $n \times n$ matrix with independent, 0-mean entries. We further assume that with probability 1,*

$$|\xi_{ij}| \leq K, \quad \sigma_{ij}^2 \leq \sigma^2, \quad \forall 1 \leq i, j \leq n.$$

One has

$$(174) \quad \mathbf{E}G_{12}^2 \leq \mu^2 + \frac{n\sigma^2(6\sigma^2 + 6K^2)}{(\lambda_p - \lambda_{r+1})^2},$$

here $\mu := \mathbf{E}G_{12}$. In particular, if E 's entries are i.i.d, then $\mu = 0$.

Therefore, by Chebyshey's inequality, we obtain:

Corollary 10.4. *For any $t > 0$*

$$\mathbf{P}(|G_{12} - \mu| \geq \frac{t}{\lambda - \lambda_{r+1}}) \leq \frac{6n\sigma^2(\sigma^2 + K^2)}{t^2}.$$

In particular, if E is i.i.d, then $\mu = 0$ and

$$\mathbf{P}(|G_{12}| \geq \frac{t}{\lambda - \lambda_{r+1}}) \leq \frac{6n\sigma^2 K^2}{t^2}.$$

The computations of G_{12} 's moments will be presented later in Appendix A - Section 11. Now, we are ready to prove Theorem 5.10 . Let Ω be the compound event defined by

$$\Omega = \left\{ |u_i^T E u_j| \leq t_1, \forall 1 \leq i, j \leq r \right\} \cap \left\{ \left| u_i^T E \left(\sum_{l>r} \frac{u_l u_l^T}{\lambda_k - \lambda_l} \right) E u_j \right| \leq \frac{t_2}{\lambda_p - \lambda_{r+1}}, \forall 1 \leq i < j \leq r, 1 \leq k \leq p \right\}.$$

By Lemma 10.2 and Corollary 10.4, Ω happens with probability at least

$$1 - r^2 \exp\left(\frac{-t_1^2/2}{2\sigma^2 + K t_1}\right) - \frac{6r(r-1)pn\sigma^2(\sigma^2 + K^2)}{t_2^2}.$$

Now, applying Theorem 2.12 with Assumption **C0** and on when event Ω happens, we have

$$\left\| \tilde{\Pi}_p - \Pi_p \right\| \leq 12\sqrt{p} \left(\epsilon_1 + \sqrt{r} t_1 \epsilon_2 + \frac{\sqrt{r}(t_2 + \mu)}{\|E\|} \epsilon_1 \epsilon_2 \right).$$

Remark 10.5. The proof of Theorem 5.7 follows the same procedure above except that one needs to use Lemma 10.6 and Corollary 10.7 instead of Lemma 10.3, Corollary 10.4

Lemma 10.6. *Let $E = (\xi_{ij})_{1 \leq i, j \leq n}$ be a random $n \times n$ matrix with independent, 0-mean entries. One has*

$$(175) \quad \mathbf{E}(E u_1 \cdot E u_2)^2 \leq 5nm_4 + \mu^2,$$

here, $\mu := \mathbf{E}E u_1 \cdot E u_2$.

In particular, if E is regular, then $\mu = 0$.

Corollary 10.7. *For any $t > 0$*

$$\mathbf{P}(|E u_1 \cdot E u_2 - \mu| \geq t) \leq \frac{5nm_4}{t^2}.$$

In particular, if E is regular, then

$$\mathbf{P}(|E u_1 \cdot E u_2| \geq t) \leq \frac{5nm_4}{t^2}.$$

10.8. **Proof of Theorem 2.16.** Given Theorem 2.12, we are going to prove Theorem 2.16. First, we apply Linearization method, considering the $(m+n) \times (m+n)$ matrices

$$\mathcal{A} := \begin{pmatrix} 0 & A \\ A^T & 0 \end{pmatrix}, \quad \mathcal{E} := \begin{pmatrix} 0 & E \\ E^T & 0 \end{pmatrix}, \quad \text{and } \tilde{\mathcal{A}} := \mathcal{A} + \mathcal{E}.$$

The non-zero eigenvalues of \mathcal{A} are $\pm\sigma_1, \dots, \pm\sigma_{r_A}$, with corresponding eigenvectors

$$\mathbf{u}_j := \frac{1}{\sqrt{2}}(u_j^T, v_j^T)^T, \quad \text{and } \mathbf{u}_{j+r} := \frac{1}{\sqrt{2}}(u_j^T, -v_j^T)^T.$$

Thus \mathcal{A} admits the spectral decomposition

$$\mathcal{A} = \mathcal{U}\mathcal{D}\mathcal{U}^T,$$

where $\mathcal{U} := (\mathbf{u}_1, \dots, \mathbf{u}_{2r_A})$ is the matrix with columns $\mathbf{u}_1, \dots, \mathbf{u}_{2r_A}$. We expand Γ into Γ' which contains exactly $S' = S \cup -S$ and define $\mathcal{U}_{S'} := (\mathbf{u}_i)_{i \in S'}$. By the decomposition of $\tilde{\mathcal{A}} = \tilde{\mathcal{U}}\tilde{\mathcal{D}}\tilde{\mathcal{U}}^T$, we also define $\tilde{\mathcal{U}}, \tilde{S}' = \tilde{S} \cup -\tilde{S}$, and $\tilde{\mathcal{U}}_{\tilde{S}'}$ respectively.

Note that for all $i, j \in N_{\bar{\sigma}}(S)$,

$$\begin{aligned} 2\mathbf{u}_i \mathcal{E} \mathbf{u}_j &= u_i^T E v_j + v_i^T E^T u_j, \\ 2\mathbf{u}_i \mathcal{E} \mathbf{u}_{j+r_A} &= -u_i^T E v_j + v_i^T E^T u_j, \\ 2\mathbf{u}_{i+r_A} \mathcal{E} \mathbf{u}_{j+r_A} &= -(u_i^T E v_j + v_i^T E^T u_j). \end{aligned}$$

For each $1 \leq i \leq 2r_A$, we define

$$(176a) \quad \bar{i} := \begin{cases} i, & \text{if } i \leq r_A \\ i - r_A, & \text{if } i > r_A. \end{cases}$$

Therefore, for any $1 \leq i, j \leq 2r_A$ and $\bar{i}, \bar{j} \in N_{\bar{\sigma}}(S)$,

$$(177) \quad |\mathbf{u}_i^T \mathcal{E} \mathbf{u}_j| \leq \frac{1}{2} (|u_i^T E v_j| + |v_i^T E^T u_j|) \leq x.$$

Similarly, we also obtain that for all $1 \leq i < j \leq 2r_A$ and $\bar{i}, \bar{j} \in N_{\bar{\sigma}}(S)$,

$$(178) \quad \begin{aligned} 4\mathbf{u}_i^T \mathcal{E} \left(\sum_{\substack{l \\ \bar{l} \notin N_{\bar{\sigma}}(S)}} \frac{\mathbf{u}_l \mathbf{u}_l^T}{\sigma_k - \sigma_l} \right) \mathcal{E} \mathbf{u}_j &= \pm v_i^T E^T \sum_{\substack{l \\ \bar{l} \notin N_{\bar{\sigma}}(S)}} \frac{u_l u_l^T}{\sigma_k - \sigma_l} E v_j \pm u_i^T E \sum_{\substack{l \\ \bar{l} \notin N_{\bar{\sigma}}(S)}} \frac{v_l v_l^T}{\sigma_k - \sigma_l} E^T u_j \\ &= 2 \left(\pm v_i^T E^T \sum_{\substack{l \\ \bar{l} \notin N_{\bar{\sigma}}(S)}} \frac{u_l u_l^T}{\sigma_k - \sigma_l} E v_j \pm u_i^T E \sum_{\substack{l \\ \bar{l} \notin N_{\bar{\sigma}}(S)}} \frac{v_l v_l^T}{\sigma_k - \sigma_l} E^T u_j \right). \end{aligned}$$

Thus, for all $1 \leq \bar{i} < \bar{j} \leq r$ and $\sigma_k \in S$,

$$(179) \quad \left| \mathbf{u}_i^T \mathcal{E} \left(\sum_{\substack{l \\ \bar{l} \notin N_{\bar{\sigma}}(S)}} \frac{\mathbf{u}_l \mathbf{u}_l^T}{\sigma_k - \sigma_l} \right) \mathcal{E} \mathbf{u}_j \right| \leq \frac{1}{2} \left(\left| v_i^T E^T \sum_{\substack{l \\ \bar{l} \notin N_{\bar{\sigma}}(S)}} \frac{u_l u_l^T}{\sigma_k - \sigma_l} E v_j \right| + \left| u_i^T E \sum_{\substack{l \\ \bar{l} \notin N_{\bar{\sigma}}(S)}} \frac{v_l v_l^T}{\sigma_k - \sigma_l} E^T u_j \right| \right) \leq y.$$

In fact, we still obtain Lemma 9.8 - second part (Section 9.1) if we consider the particular case $(\tilde{i}_1, \dots, \tilde{i}_k) = (i, \dots, i)$, $1 \leq i \leq r$ instead of the case $(i_1, \dots, i_k) = (i, \dots, i)$ from the original proof. Hence, (177), (179), and the fact that $\|\mathcal{E}\| = \|E\|$ are enough for us to apply Theorem 2.12 on \mathcal{A} and \mathcal{E} . It means that we do not need to require $\left| \mathbf{u}_i^T \mathcal{E} \left(\sum_{l \notin N_{\bar{\sigma}}(S)} \frac{\mathbf{u}_l \mathbf{u}_l^T}{\sigma_k - \sigma_l} \right) \mathcal{E} \mathbf{u}_{i+r_A} \right| \leq y$.

Obtain

$$(180) \quad \left\| \mathcal{U}_{S'} \mathcal{U}_{S'}^T - \tilde{\mathcal{U}}_{\tilde{S}} \tilde{\mathcal{U}}_{\tilde{S}'}^T \right\| \leq 12\sqrt{p} \left(\frac{\|E\|}{\bar{\sigma}} + \frac{\sqrt{r}x}{\delta} + \frac{\sqrt{r}y}{\delta} \right).$$

Furthermore by [59, Lemma 26],

$$(181) \quad \max \left\{ \left\| \tilde{\Pi}_{\tilde{S}}^{left} - \Pi_S^{left} \right\|, \left\| \tilde{\Pi}_{\tilde{S}}^{right} - \Pi_S^{right} \right\| \right\} \leq \sqrt{2} \left\| \mathcal{U}_{S'} \mathcal{U}_{S'}^T - \tilde{\mathcal{U}}_{\tilde{S}} \tilde{\mathcal{U}}_{\tilde{S}'}^T \right\|.$$

Combining (180) and (181), we complete the proof.

10.9. Proof of Theorem 6.11. Recall the setting and notations from Section 6. Our proof is the combination of Theorem 2.12 and two following lemmas - the estimates of the corresponding parameters x and y for the spiked population model. Note that our assumption is $X = M^{1/2}Y$, $Y = [y_1, \dots, y_d]^T$, in which y_i s are identity 8-wise independent with mean 0, variance 1 and finite 4th moment.

Lemma 10.10 (x -bound). *There is a universal constant C such that for any $t > 0$, with probability at least $1 - \frac{r^2}{t^2}$,*

$$(182) \quad x \leq C \frac{\lambda_1 t}{\sqrt{n}}.$$

Lemma 10.11 (y -bound). *There is a universal constant C such that, for any $t > 0$, with probability at least $1 - \frac{pr^2 t^2}{t^2}$,*

$$(183) \quad y \leq C \left(\frac{c_r \sqrt{\lambda_1 \lambda_2} t}{n} \frac{\sum_{l>r} \lambda_l}{\lambda_p - \lambda_{r+1}} + \frac{\sqrt{\lambda_1 \lambda_2} t}{n} \sqrt{\sum_{l>r} \frac{\lambda_l^2}{(\lambda_p - \lambda_l)^2}} \right),$$

in which

$$c_r := \max_{\substack{1 \leq i < j \leq r \\ l, l' > r}} \left| \sum_{\alpha} (\mathbf{E} y_{\alpha}^4 - 3) u_{i\alpha} u_{j\alpha} u_{l\alpha} u_{l'\alpha} \right| + \frac{1}{\sqrt{n}}.$$

These lemmas are proven via moment method. These computations will be presented in Appendix B.

11. APPENDIX A: MOMENT COMPUTATIONS

11.1. Moment computation of $|Eu_i \cdot Eu_j|$. To obtain a more friendly version (for applications), we will bound $|Eu_i \cdot Eu_j|$ (the quantity \bar{y} for low-rank matrix perturbation) in the case that E is a matrix with independent entries. Without loss of generality, take $i = 1$ and $j = 2$. We have

$$\bar{G}_{12} := Eu_1 \cdot Eu_2 = \sum_l \sum_i \xi_{li} u_{1i} \sum_j \xi_{lj} u_{2j}.$$

As the ξ_{li}, ξ_{lj} are independent with zero mean, the expectation comes from the diagonal term

$$\mathbf{E}\bar{G}_{12} = \sum_l \sum_i \mathbf{E}\xi_{li}^2 u_{1i} u_{2i} = \sum_i u_{1i} u_{2i} \sum_l \mathbf{E}\xi_{li}^2.$$

Let $S_i = \sum_l \mathbf{E}\xi_{li}^2$. If the random matrix E is regular, each $S_i = nm_2$, and the expectation of \bar{G}_{12} is zero as u_1 and u_2 are orthogonal. If the matrix is not generalized Wigner, then

$$\mu := \mathbf{E}\bar{G}_{12} = \sum_i S_i u_{1i} u_{2i},$$

with S_i defined as above (in geometrical term, S_i is the expectation of the square of the length of the i th row of E). We now turn to the second moment

$$\bar{G}_{12}^2 = \sum_{\substack{i,j,l \\ i',j',l'}} \xi_{il} \xi_{lj} \xi_{i'l'} \xi_{l'j'} u_{1i} u_{2j} u_{1i'} u_{1j'}.$$

Thus,

$$\mathbf{E}\bar{G}_{12}^2 = \sum_{\substack{i,j,l \\ i',j',l'}} u_{1i} u_{2j} u_{1i'} u_{1j'} \mathbf{E}\xi_{il} \xi_{lj} \xi_{i'l'} \xi_{l'j'}.$$

Since $\mathbf{E}(\xi_{ij}) = 0$, most of the expectations on the RHS are zero. The non-trivial ones can be divided into four cases.

Case 1. (Fourth moment case) $\{i, l\} = \{l, j\} = \{i', l'\} = \{l', j'\}$. This can only happens if $i = j = i' = j', l = l'$ or $i = j = l', i' = j' = l$.

In the first scenario, we receive

$$\sum_i (u_{1i} u_{2i})^2 \sum_l \mathbf{E}\xi_{il}^4 \leq h_{12} n m_4.$$

In the second scenario, we receive

$$\sum_{i,l} u_{1i} u_{2i} u_{1l} u_{2l} \mathbf{E}\xi_{il}^4.$$

As $u_{1i}u_{2i}u_{1l}u_{2l} \leq \frac{1}{2}((u_{1i}u_{2i})^2 + (u_{1l}u_{2l})^2)$, the last quantity can also be bounded by $h_{12}nm_4$. Thus, the contribution in Case 1 is at most $2h_{12}nm_4$.

Case 2. $\{i, l\} = \{l, j\} \neq \{i', l'\} = \{l', j'\}$. This could only happen if $i = j, i' = j'$ and $\{i, l\} \neq \{i', l'\}$. The contribution from this case is

$$\sum_{i, i', l, l'} u_{1i}u_{2i}u_{1i'}u_{2i'} \mathbf{E}\xi_{il}^2 \mathbf{E}\xi_{i'l'}^2 - D,$$

where D collect all diagonal terms with $\{i, l\} = \{i'l'\}$. Without any restriction on the indices, the sum $\sum_{i, i', l, l'} u_{1i}u_{2i}u_{1i'}u_{2i'} \mathbf{E}\xi_{ij}^2 \mathbf{E}\xi_{i'l'}^2$ equals

$$\left(\sum_{i, l} u_{1i}u_{2i} \mathbf{E}\xi_{il}^2 \right)^2 = \left(\sum_i u_{1i}u_{2i} \sum_l \mathbf{E}\xi_{il}^2 \right)^2 = \mu^2.$$

The term D can be bounded (in absolute value) by $h_{12}nm_4$ similarly to Case 1. Thus, the contribution of this case is at most

$$h_{12}nm_4 + \mu^2.$$

Case 3. $\{i, l\} = \{i', l'\} \neq \{l, j\} = \{l', j'\}$. It happens if $(i = i', l = l', j = j', i \neq j)$. The contribution from this case is

$$\sum_{i, j, l} \mathbf{E}\xi_{il}^2 \xi_{jl}^2 u_{1i}^2 u_{2j}^2 = \sum_{i, j} u_{1i}^2 u_{2j}^2 \sum_l \mathbf{E}\xi_{il}^2 \xi_{jl}^2.$$

By Cauchy-Schwartz

$$\sum_l \mathbf{E}\xi_{il}^2 \xi_{jl}^2 \leq \frac{1}{2} \sum_l \mathbf{E}(\xi_{il}^4 \xi_{jl}^4) \leq nm_4.$$

Furthermore, as u_1, u_2 are unit vectors $\sum_{i, j} u_{1i}^2 u_{2j}^2 = 1$. Thus, the contribution in this case is bounded from above by nm_4 .

Case 4. $\{i, l\} = \{l', j'\} \neq \{l, j\} = \{i', l'\}$. It happens if $(i = j', i' = j, l = l', i \neq j)$. The contribution in this case is

$$\left| \sum_{i, j, l} u_{1i}u_{2i}u_{1j}u_{2j} \mathbf{E}\xi_{il}^2 \xi_{jl}^2 \right| \leq \sum_{i, j} |u_{1i}u_{2i}u_{1j}u_{2j}| \sum_l \mathbf{E}\xi_{il}^2 \xi_{jl}^2.$$

By the argument in the previous case,

$$\sum_l \mathbf{E} \xi_{il}^2 \xi_{jl}^2 \leq nm_4.$$

Furthermore, by Cauchy-Schwartz

$$\sum_{i,j} |u_{1i} u_{2i} u_{1j} u_{2j}| = \sum_i |u_{1i} u_{2i}| \sum_j |u_{1j} u_{2j}| \leq 1.$$

Thus, the contribution from this case is bounded by nm_4 .

Putting all the cases together,

$$\mathbf{E} \bar{G}_{12}^2 \leq 3h_{12}nm_4 + 2nm_4 + \mu^2 \leq 5nm_4 + \mu^2.$$

Recall that $\mathbf{E} \bar{G}_{12} = \mu$, so the variance of \bar{G}_{12} is at most $5nm_4$. By Chebyshev's inequality, we obtain

Corollary 11.2. *For any $t > 0$*

$$\mathbf{P}(|\bar{G}_{12} - \mu| \geq t) \leq \frac{5nm_4}{t^2}.$$

In particular, if E is regular, then $\mu = 0$ and

$$\mathbf{P}(|\bar{G}_{12}| \geq t) \leq \frac{5nm_4}{t^2}.$$

11.3. Moment computation of $\left| u_i^T E \left(\sum_{l>r} \frac{u_l u_l^T}{\lambda_k - \lambda_j} \right) E u_j \right|$. In this subsection, via moment method, we are going to bound $\left| u_i^T E \left(\sum_{l>r} \frac{u_l u_l^T}{\lambda_k - \lambda_j} \right) E u_j \right|$ for a fixed triple (k, i, j) : $1 \leq i < j \leq r, 1 \leq k \leq p$, i.e. $S := \{1, 2, \dots, p\}, N_{\bar{\lambda}}(S) = \{1, 2, \dots, r\}$. Here, we will work with only the assumption that E is a matrix with independent entries. A similar computation also applies for a general subset S and its corresponding $N_{\bar{\lambda}}(S)$.

Without loss of generality, take $i = 1$ and $j = 2$. We further denote

$$G_{12} = u_1^T E \left(\sum_{l>r} \frac{u_l u_l^T}{\bar{\sigma}_l} \right) E u_2, \quad \bar{\sigma}_l = \lambda_k - \lambda_l, \quad \text{for some } 1 \leq k \leq p.$$

We have

$$\begin{aligned} (184) \quad G_{12} &= \sum_{l>r} \sum_{k,i,j,m} u_{1k} \xi_{ki} u_{li} u_{lj} \xi_{jm} u_{2m} \frac{1}{\bar{\sigma}_l} \\ &= \sum_{l>r} \sum_{k,i,j,m} u_{1k} u_{2m} \frac{u_{li} u_{lj}}{\bar{\sigma}_l} \xi_{ki} \xi_{jm}. \end{aligned}$$

Therefore,

$$\begin{aligned}
\mathbf{E}G_{12} &= \sum_{l>r} \sum_{k,i,j,m} u_{1k}u_{2m} \frac{u_{li}u_{lj}}{\bar{\sigma}_l} \mathbf{E}\xi_{ki}\xi_{jm} \\
(185) \quad &= \sum_{l>r} \sum_{k,i} u_{1k}u_{2i} \frac{u_{li}u_{lk}}{\bar{\sigma}_l} \mathbf{E}\xi_{ki}^2 + \sum_{l>r} \sum_{k,i} u_{1k}u_{2k} \frac{u_{li}u_{li}}{\bar{\sigma}_l} \mathbf{E}\xi_{ki}^2 - \sum_{l>r} \sum_k u_{1k}u_{2k} \frac{u_{lk}^2}{\bar{\sigma}_l} \mathbf{E}\xi_{kk}^2 \\
&= \sum_{l>r} \sum_{k,i} u_{1k}u_{2i} \frac{u_{li}u_{lk}}{\bar{\sigma}_l} \mathbf{E}\xi_{ki}^2 + \mu - \sum_{l>r} \sum_k u_{1k}u_{2k} \frac{u_{lk}^2}{\bar{\sigma}_l} \mathbf{E}\xi_{kk}^2.
\end{aligned}$$

For the second moment, we have

$$\begin{aligned}
\mathbf{E}G_{12}^2 &= \sum_{l,l'>r} \sum_{\substack{k,i,j,m \\ k',i',j',m'}} u_{1k}u_{2m} \frac{u_{li}u_{lj}}{\bar{\sigma}_l} u_{1k'}u_{2m'} \frac{u_{l'i'}u_{l'j'}}{\bar{\sigma}_{l'}} \mathbf{E}\xi_{ki}\xi_{jm}\xi_{k'i'}\xi_{j'm'} \\
(186) \quad &= M_1 + M_2 + M_3 + M_4,
\end{aligned}$$

in which

$$\begin{aligned}
M_1 &:= \sum_{l,l'>r} \sum_{\substack{(k,i)=(j,m) \neq \\ (k',i')=(j',m')}} (u_{1k}u_{2m} \frac{u_{li}u_{lj}}{\bar{\sigma}_l} u_{1k'}u_{2m'} \frac{u_{l'i'}u_{l'j'}}{\bar{\sigma}_{l'}} \mathbf{E}\xi_{ki}^2 \mathbf{E}\xi_{k'i'}^2) \\
M_2 &:= \sum_{l,l'>r} \sum_{\substack{(k,i)=(k',i') \neq \\ (j,m)=(j',m')}} (u_{1k}u_{2m} \frac{u_{li}u_{lj}}{\bar{\sigma}_l} u_{1k'}u_{2m'} \frac{u_{l'i'}u_{l'j'}}{\bar{\sigma}_{l'}} \mathbf{E}\xi_{ki}^2 \mathbf{E}\xi_{jm}^2) \\
(187) \quad M_3 &:= \sum_{l,l'>r} \sum_{\substack{(k,i)=(j',m') \neq \\ (k',i')=(j,m)}} (u_{1k}u_{2m} \frac{u_{li}u_{lj}}{\bar{\sigma}_l} u_{1k'}u_{2m'} \frac{u_{l'i'}u_{l'j'}}{\bar{\sigma}_{l'}} \mathbf{E}\xi_{ki}^2 \mathbf{E}\xi_{jm}^2) \\
M_4 &:= \sum_{l,l'>r} \sum_{\substack{(k,i)=(j,m) = \\ (k',i')=(j',m')}} (u_{1k}u_{2m} \frac{u_{li}u_{lj}}{\bar{\sigma}_l} u_{1k'}u_{2m'} \frac{u_{l'i'}u_{l'j'}}{\bar{\sigma}_{l'}} \mathbf{E}\xi_{ki}^4).
\end{aligned}$$

In fact,

$$M_5 := M_1 + M_4 - (\mathbf{E}G_{12})^2 = \sum_{l,l'} \sum_{\substack{(k,i)=(j,m) = \\ (k',i')=(j',m')}} (u_{1k}u_{2m} \frac{u_{li}u_{lj}}{\bar{\sigma}_l} u_{1k'}u_{2m'} \frac{u_{l'i'}u_{l'j'}}{\bar{\sigma}_{l'}}) (\mathbf{E}\xi_{ki}^4 - (\mathbf{E}\xi_{ki}^2)^2).$$

To estimate $\mathbf{E}G_{12}^2$, we only need to bound M_2 and M_5 , since M_3 is M_2 with indices permutation.

11.3.1. *Bounding M_2 .* We have

$$\begin{aligned}
M_2 &= \sum_{l,l'>r} \sum_{\substack{(k,i) \neq \\ (j,m)}} \sigma_{ki}^2 \sigma_{jm}^2 \left(\frac{(u_{1k})^2 (u_{2m})^2 u_{li}u_{lj}u_{l'i}u_{l'j}}{\bar{\sigma}_l \bar{\sigma}_{l'}} + \frac{(u_{1k})^2 u_{2m} u_{2j} u_{li}u_{lj}u_{l'i}u_{l'm}}{\bar{\sigma}_l \bar{\sigma}_{l'}} \right) \\
(188) \quad &+ \sum_{l,l'>r} \sum_{\substack{(k,i) \neq \\ (j,m)}} \sigma_{ki}^2 \sigma_{jm}^2 \left(\frac{u_{1k}u_{1i} (u_{2m})^2 u_{li}u_{lj}u_{l'k}u_{l'j}}{\bar{\sigma}_l \bar{\sigma}_{l'}} + \frac{u_{1k}u_{1i}u_{2m}u_{2j}u_{li}u_{lj}u_{l'k}u_{l'm}}{\bar{\sigma}_l \bar{\sigma}_{l'}} \right).
\end{aligned}$$

The first sub-sum can be bounded as what follows. First, we recall the notation that

$$s_{ij} = \sum_{l>r} \frac{u_{li}u_{lj}}{\bar{\sigma}_l}.$$

(189)

$$\begin{aligned} \sum_{l,l'>r} \sum_{\substack{(k,i) \neq \\ (j,m)}} \sigma_{ki}^2 \sigma_{jm}^2 \frac{(u_{1k})^2 (u_{2m})^2 u_{li} u_{lj} u_{l'i} u_{l'j}}{\bar{\sigma}_l \bar{\sigma}_{l'}} &= \sum_{k,i,j,m} \sigma_{ki}^2 \sigma_{jm}^2 u_{1k}^2 u_{2m}^2 s_{ij}^2 \\ &\leq \sigma^4 \sum_{k,i,j,m} u_{1k}^2 u_{2m}^2 s_{ij}^2 \\ &= \sigma^4 \sum_{l,l'>r} \sum_{k,i,j,m} \frac{(u_{1k})^2 (u_{2m})^2 u_{li} u_{lj} u_{l'i} u_{l'j}}{\bar{\sigma}_l \bar{\sigma}_{l'}} \\ &= \sigma^4 \sum_{l>r,k,m} \frac{u_{1k}^2 u_{2m}^2}{\bar{\sigma}_l^2} \left(\text{since } \sum_i u_{li} u_{l'i} = \mathbf{1}_{l=l'} \right) \\ &= \sigma^4 \sum_{l>r} \frac{1}{\bar{\sigma}_l^2} \leq \frac{\sigma^4 (n-r)}{(\lambda - \lambda_{r+1})^2}. \end{aligned}$$

We bound the second sub-sum as follows

(190)

$$\begin{aligned} \sum_{l,l'>r} \sum_{\substack{(k,i) \neq \\ (j,m)}} \sigma_{ki}^2 \sigma_{jm}^2 \frac{(u_{1k})^2 u_{2m} u_{2j} u_{li} u_{lj} u_{l'i} u_{l'm}}{\bar{\sigma}_l \bar{\sigma}_{l'}} &= \sum_k u_{1k}^2 \sum_i \sigma_{ki}^2 \sum_{j,m} (u_{2j} s_{ij}) \sigma_{jm}^2 (u_{2m} s_{im}) \\ &= \sum_k u_{1k}^2 \sum_i \sum_{j,m} (\sigma_{ki} u_{2j} s_{ij}) \sigma_{jm}^2 (\sigma_{ki} u_{2m} s_{im}). \end{aligned}$$

For each k , consider the $n \times 1$ vectors $U_{i,k} = [\sigma_{ki} u_{2j} s_{ij}]_{1 \leq j \leq n}^T$ and the $n \times n$ matrix $\Sigma = (\sigma_{j,m}^2)_{1 \leq j,m \leq n}$. The absolute value of above RHS can be rewritten as

$$\begin{aligned} \left| \sum_k u_{1k}^2 \sum_i \sum_{j,m} (\sigma_{ki} u_{2j} s_{ij}) \sigma_{jm}^2 (\sigma_{ki} u_{2m} s_{im}) \right| &= \sum_k u_{1k}^2 \sum_i |U_{i,k}^T \Sigma U_{i,k}| \\ &\leq \sum_k u_{1k}^2 \|\Sigma\| \sum_i \|U_{i,k}\|^2 \\ &\leq n \sigma^2 \sum_k u_{1k}^2 \sum_i \sum_j \sigma_{ki}^2 u_{2j}^2 s_{ij}^2 \\ (191) \quad &\leq n \sigma^4 \sum_k u_{1k}^2 \sum_i \sum_j \sum_{l,l'>r} u_{2j}^2 \frac{u_{li} u_{lj} u_{l'i} u_{l'j}}{\bar{\sigma}_l \bar{\sigma}_{l'}} \\ &= n \sigma^4 \sum_j u_{2j}^2 \sum_l \frac{u_{lj}^2}{\bar{\sigma}_l^2} \\ &\leq n \sigma^4 \sum_j u_{2j}^2 \frac{\sum_l u_{lj}^2}{(\lambda - \lambda_{r+1})^2} \leq \frac{n \sigma^4}{(\lambda - \lambda_{r+1})^2}. \end{aligned}$$

The third sub-sum is estimated by the similar argument. Finally, the fourth sub-sum can be bounded as follows.

(192)

$$\begin{aligned}
\left| \sum_{l,l'>r} \sum_{\substack{(k,i) \neq \\ (j,m)}} \sigma_{ki}^2 \sigma_{jm}^2 \frac{u_{1k} u_{1i} u_{2m} u_{2j} u_{li} u_{lj} u_{l'k} u_{l'm}}{\bar{\sigma}_l \bar{\sigma}_{l'}} \right| &= \left| \sum_{k,i,j,m} u_{1k} \sigma_{ki}^2 u_{1i} u_{2m} \sigma_{jm}^2 u_{2j} s_{ij} s_{km} \right| \\
&= \left| \sum_{k,i} (u_{1k} \sigma_{ki}^2 u_{1i}) \sum_{j,m} (u_{2m} s_{km} \sigma_{jm}^2 (u_{2j} s_{ij})) \right| \\
&= \left| \sum_{k,i} u_{1k} (\sigma_{ki}^2 r_{ki}) u_{1i} \right|, \quad r_{ki} := \sum_{j,m} (u_{2m} s_{km} \sigma_{jm}^2 (u_{2j} s_{ij})) \\
&= |u_1^T \Sigma' u_1|, \quad \Sigma' := (\sigma_{ki}^2 r_{ki})_{1 \leq k,i \leq n} \\
&\leq \|\Sigma'\| \leq \|\Sigma'\|_F
\end{aligned}$$

Moreover,

$$\begin{aligned}
\|\Sigma'\|_F^2 &\leq \sigma^4 \sum_{1 \leq k,i \leq n} r_{ki}^2 \\
(193) \quad &= \sigma^4 \sum_{1 \leq k,i \leq n} |V_{2,k}^T \Sigma V_{2,i}|^2 \quad (V_{2,k} := [u_{2m} s_{km}]_{1 \leq m \leq n}^T, V_{2,i} := [u_{2m} s_{im}]_{1 \leq m \leq n}^T) \\
&\leq \sigma^4 \|\Sigma\|^2 \sum_{k,i} \|V_{2,k}\|^2 \cdot \|V_{2,i}\|^2.
\end{aligned}$$

Similarly, $\|\Sigma\|^2 \leq n^2 \sigma^4$ and

$$\sum_{k,i} \|V_{2,k}\|^2 \cdot \|V_{2,i}\|^2 = \left(\sum_k \sum_m u_{2m}^2 s_{km}^2 \right)^2.$$

And,

$$\begin{aligned}
\sum_k \sum_m u_{2m}^2 s_{km}^2 &= \sum_m u_{2m}^2 \sum_k s_{km}^2 \\
(194) \quad &= \sum_m u_{2m}^2 \sum_{l,l'>r} \frac{(\sum_k u_{lk} u_{l'k}) u_{lm} u_{l'm}}{\bar{\sigma}_l \bar{\sigma}_{l'}} \\
&= \sum_m u_{2m}^2 \sum_{l>r} \frac{u_{lm}^2}{\bar{\sigma}_l^2} \\
&\leq \frac{\sum_m u_{2m}^2}{(\lambda - \lambda_{r+1})^2}.
\end{aligned}$$

Therefore,

$$(195) \quad \|\Sigma'\| \leq \frac{n\sigma^4}{(\lambda - \lambda_{r+1})^2}$$

Combining (189), (191), (195), we obtain

$$(196) \quad |M_2| \leq \frac{3n\sigma^4}{(\lambda - \lambda_{r+1})^2}$$

11.3.2. *Bounding M_5 .* Denote $\sigma_{4,ki} = \mathbf{E}\xi_{ki}^4 - (\mathbf{E}\xi_{ki}^2)^2$ and $\bar{\sigma}_4 = \max_{1 \leq k, i \leq n} \sigma_{4,ki}$. We have $M_5 = \sum_{l, l' > r} \sum_{\substack{(k, i) = (j, m) \\ (k', i') = (j', m')}} (\dots) (\mathbf{E}\xi_{ki}^4 - (\mathbf{E}\xi_{ki}^2)^2)$, which equals

$$(197) \quad \sum_{l, l' > r} \frac{1}{\bar{\sigma}_l \bar{\sigma}_{l'}} \sum_{k, i} \sigma_{4,ki} \times (u_{1k} u_{2i} u_{li} u_{lk} u_{1k} u_{2i} u_{l'i} u_{l'k} +$$

$$u_{1k} u_{2i} u_{li} u_{lk} u_{1i} u_{2i} u_{l'k} u_{l'k} +$$

$$u_{1k} u_{2k} u_{li} u_{li} u_{1k} u_{2i} u_{l'i} u_{l'k} +$$

$$u_{1k} u_{2i} u_{li} u_{lk} u_{1k} u_{2k} u_{l'i} u_{l'i} +$$

$$u_{1k} u_{2i} u_{li} u_{lk} u_{1i} u_{2k} u_{l'k} u_{l'i} +$$

$$u_{1k} u_{2k} u_{li} u_{li} u_{1k} u_{2k} u_{l'i} u_{l'i} +$$

$$u_{1k} u_{2k} u_{li} u_{li} u_{1i} u_{2i} u_{l'k} u_{l'k} +$$

$$u_{1k} u_{2k} u_{li} u_{li} u_{1i} u_{2k} u_{l'k} u_{l'i}).$$

It equals

$$(198) \quad \sum_{l, l' > r} \frac{1}{\bar{\sigma}_l \bar{\sigma}_{l'}} \sum_{k, i} \sigma_{4,ki} \times ((u_{1k})^2 u_{lk} u_{l'k} (u_{2i})^2 u_{li} u_{l'i} +$$

$$u_{1k} u_{lk} (u_{l'k})^2 (u_{2i})^2 u_{li} u_{1i} +$$

$$(u_{1k})^2 u_{2k} u_{l'k} (u_{li})^2 u_{2i} u_{l'i} +$$

$$(u_{1k})^2 u_{lk} u_{2k} u_{2i} u_{li} (u_{l'i})^2 +$$

$$(u_{1k} u_{lk} u_{2k} u_{l'k}) (u_{2i} u_{li} u_{1i} u_{l'i}) +$$

$$(u_{1k})^2 (u_{2k})^2 (u_{li})^2 (u_{l'i})^2 +$$

$$u_{1k} u_{2k} (u_{l'k})^2 (u_{li})^2 u_{1i} u_{2i} +$$

$$u_{1k} (u_{2k})^2 u_{l'k} (u_{li})^2 u_{1i} u_{l'i}).$$

Here, we have 8 sub-sums. Without losing generality, we only present how to work with two first ones, others are estimated by a similar argument. We bound the first one as what

follows.

$$\begin{aligned}
(199) \quad & \sum_{l,l'>r} \frac{1}{\bar{\sigma}_l \bar{\sigma}_{l'}} \sum_{k,i} \sigma_{4,ki} (u_{1k})^2 u_{lk} u_{l'k} (u_{2i})^2 u_{li} u_{l'i} \leq \sum_{l,l'>r} \frac{1}{\bar{\sigma}_l \bar{\sigma}_{l'}} \sum_{k,i} \sigma_{4,ki} |u_{1k}|^2 |u_{lk}| |u_{l'k}| |u_{2i}|^2 |u_{li}| |u_{l'i}| \\
& \leq \sum_{k,i} \sigma_{4,ki} |u_{1k}|^2 |u_{2i}|^2 \left(\frac{|u_{lk}| |u_{li}|}{\bar{\sigma}_l} \right)^2 \\
& \leq \sum_{k,i} \sigma_{4,ki} |u_{1k}|^2 |u_{2i}|^2 \frac{1}{(\lambda - \lambda_{r+1})^2} \left(\sum_l u_{lk}^2 \right) \left(\sum_l u_{li}^2 \right) \\
& \leq \sum_{k,i} \sigma_{4,ki} |u_{1k}|^2 |u_{2i}|^2 \frac{1}{(\lambda - \lambda_{r+1})^2} \\
& \leq \frac{\bar{\sigma}_4 (\sum_k u_{1k}^2) (\sum_i u_{2i}^2)}{(\lambda - \lambda_{r+1})^2} \\
& \leq \frac{\bar{\sigma}_4}{(\lambda - \lambda_{r+1})^2}.
\end{aligned}$$

The second sub-sum can be bounded as follows.

$$\begin{aligned}
(200) \quad & \sum_{l,l'>r} \frac{1}{\bar{\sigma}_l \bar{\sigma}_{l'}} \sum_{k,i} \sigma_{4,ki} u_{1k} u_{lk} (u_{l'k})^2 (u_{2i})^2 u_{li} u_{l'i} \leq \sum_{l,l'>r} \sum_{k,i} \sigma_{4,ki} |u_{1k}| |u_{li}| (u_{2i})^2 \frac{|u_{lk}| |u_{li}|}{\bar{\sigma}_l} \frac{(u_{l'k})^2}{\bar{\sigma}_{l'}} \\
& \leq \frac{\bar{\sigma}_4}{(\lambda - \lambda_{r+1})^2} \left(\sum_k |u_{1k}| \right) \left(\sum_i |u_{li}| u_{2i}^2 \right) \\
& \leq \frac{\bar{\sigma}_4}{(\lambda - \lambda_{r+1})^2} \left(\sum_k |u_{1k}| \right) \left(\sum_i |u_{li}| \right) \\
& \leq \frac{n \bar{\sigma}_4}{(\lambda - \lambda_{r+1})^2}.
\end{aligned}$$

All leads to

$$(201) \quad |M_5| \leq \frac{6n\bar{\sigma}_4}{(\lambda - \lambda_{r+1})^2} \leq \frac{6n\sigma^2 K^2}{(\lambda - \lambda_{r+1})^2}$$

Remark 11.4.

- (1) If E is random matrix with i.i.d entries, $\mathbf{E}G_{12} = 0$.
- (2) In the general case, with the same procedure above, we can bound $\mathbf{E}G_{12}$ by $\frac{(\sqrt{n}+1)\sigma^2}{\lambda - \lambda_{r+1}} + \mu$.

Putting all above estimates together, we finally obtain

$$\mathbf{E}G_{12}^2 \leq \mu^2 + \frac{n\sigma^2(6\sigma^2 + 6K^2)}{(\lambda - \lambda_{r+1})^2}.$$

12. APPENDIX B: ESTIMATING COVARIANCE PARAMETERS x AND y

Recall that $X = [\xi_1 \dots \xi_d]^T$. For each $1 \leq i \leq n$, denote $X_i := [\xi_1^{(i)} \dots \xi_d^{(i)}]^T$.

We will use the following notations

$$\begin{aligned}\xi_{i_1, i_2, \dots, i_r}^{(i)} &:= \xi_{i_1}^{(i)} \xi_{i_2}^{(i)} \dots \xi_{i_r}^{(i)}, \\ \xi_{i_1, i_2, \dots, i_r} &:= \xi_{i_1} \dots \xi_{i_r}.\end{aligned}$$

Denote $E = (e_{lk})_{1 \leq l, k \leq d}$, and then by construction,

$$e_{kl} = \frac{1}{n} \sum_{i=1}^n (\xi_k^{(i)} \xi_l^{(i)} - m_{kl}).$$

12.1. General Spiked Model.

12.1.1. *Bounding x .* Without the loss of generality, we now find the upper bound of $u_1^T E u_2$. We have

$$\begin{aligned}(202) \quad u_1^T E u_2 &= \sum_{k, l} u_{1k} u_{2l} e_{kl} \\ &= \frac{1}{n} \sum_{1 \leq k, l \leq d} \sum_{1 \leq i \leq n} u_{1k} u_{2l} (\xi_k^{(i)} \xi_l^{(i)} - m_{kl}).\end{aligned}$$

Therefore, $\mathbf{E} u_1^T E u_2 = 0$. Furthermore,

$$\begin{aligned}(203) \quad \mathbf{E}(u_1^T E u_2)^2 &= \frac{1}{n^2} \sum_{\substack{k, l \\ k', l'}} \sum_{1 \leq i, i' \leq n} u_{1k} u_{2l} u_{1k'} u_{2l'} \mathbf{E} \left[(\xi_k^{(i)} \xi_l^{(i)} - m_{kl}) (\xi_{k'}^{(i')} \xi_{l'}^{(i')} - m_{k'l'}) \right] \\ &= \frac{1}{n^2} \sum_{1 \leq i \leq n} \sum_{\substack{k, l \\ k', l'}} u_{1k} u_{2l} u_{1k'} u_{2l'} \mathbf{E} \left[(\xi_k^{(i)} \xi_l^{(i)} - m_{kl}) (\xi_{k'}^{(i)} \xi_{l'}^{(i)} - m_{k'l'}) \right] \\ &= \frac{1}{n^2} \sum_{1 \leq i \leq n} \sum_{\substack{k, l \\ k', l'}} u_{1k} u_{2l} u_{1k'} u_{2l'} \left[\mathbf{E} \left(\xi_{k, l, k', l'}^{(i)} \right) - m_{kl} m_{k'l'} \right] \\ &= \frac{1}{n} \sum_{\substack{k, l \leq d \\ k', l' \leq d}} u_{1k} u_{2l} u_{1k'} u_{2l'} \left(\mathbf{E} \xi_{k, l, k', l'} - m_{kl} m_{k'l'} \right).\end{aligned}$$

So (if) we can consider an $d^2 \times 1$ random vector $X' = [\xi_{k, l}]_{1 \leq k, l \leq d}^T$ and the $d^2 \times 1$ deterministic unit vector $U_{12} = [u_{k, l}^{12}]_{1 \leq k, l \leq d}^T$, $u_{kl}^{12} = u_{1k} u_{2l}$. And then,

$$(204) \quad \mathbf{E}(u_1^T E u_2)^2 = \frac{1}{n} U_{12}^T \text{Cov}(X') U_{12},$$

in which $\text{Cov}(X')$ is the covariance matrix of X' (size $d^2 \times d^2$). It looks like we can bound it by $\frac{\|\text{Cov}(X')\|}{n}$.

12.1.2. *Bounding y .* For the general setting of covariance matrix X , we need to compute the first moment of $u_1^T E \sum_{l>r} \frac{u_l u_l^T}{\lambda - \lambda_l} E u_2$:

(205)

$$\begin{aligned} \mathbf{E} u_1^T E \sum_{l>r} \frac{u_l u_l^T}{\lambda - \lambda_l} E u_2 &= \frac{1}{n^2} \sum_{k,j,m,i} \sum_{l>r} u_{1k} \frac{u_{lj} u_{lm}}{\lambda - \lambda_l} u_{2i} \mathbf{E} \left(\sum_I \xi_k^{(I)} \xi_j^{(I)} - m_{kj} \right) \left(\sum_J \xi_m^{(J)} \xi_i^{(J)} - m_{mi} \right) \\ &= \frac{1}{n} \sum_{k,j,m,i} \sum_{l>r} u_{1k} \frac{u_{lj} u_{lm}}{\lambda - \lambda_l} u_{2i} \mathbf{E} (\xi_k \xi_j - m_{kj}) (\xi_m \xi_i - m_{mi}) \\ &= \frac{1}{n} \sum_{l>r} \frac{1}{\lambda - \lambda_l} \mathbf{E} \left(\left(\sum_k u_{1k} \xi_k \right) \left(\sum_j \xi_j u_{lj} \right) - \mathbf{E} \left(\sum_k u_{1k} \xi_k \right) \left(\sum_j \xi_j u_{lj} \right) \right) \times \\ &\quad \left(\left(\sum_i u_{2i} \xi_i \right) \left(\sum_m \xi_m u_{lm} \right) - \mathbf{E} \left(\sum_i u_{2i} \xi_i \right) \left(\sum_m \xi_m u_{lm} \right) \right). \end{aligned}$$

When M is low-rank, the above equation can be reduced into:

$$\begin{aligned} \mathbf{E} u_1^T E^2 u_2 &= \sum_{1 \leq j,k,l \leq d} u_{1k} u_{2l} e_{kj} e_{jl} \\ &= \mathbf{E} \frac{1}{n^2} \sum_{1 \leq j,k,l \leq d} \sum_{1 \leq i,i' \leq n} u_{1k} u_{2l} (\xi_k^{(i)} \xi_j^{(i)} - m_{kj}) (\xi_l^{(i')} \xi_j^{(i')} - m_{jl}) \\ (206) \quad &= \frac{1}{n^2} \sum_{1 \leq j,k,l \leq d} \sum_{1 \leq i \leq n} u_{1k} u_{2l} \mathbf{E} \left[(\xi_k^{(i)} \xi_j^{(i)} - m_{kj}) (\xi_l^{(i)} \xi_j^{(i)} - m_{jl}) \right] \\ &= \frac{1}{n^2} \sum_{1 \leq j,k,l \leq d} \sum_{1 \leq i \leq n} u_{1k} u_{2l} \left(\mathbf{E} \xi_{j,j,k,l}^{(i)} - m_{kj} m_{lj} \right) \\ &= \frac{1}{n} \sum_{1 \leq j,k,l \leq d} u_{1k} u_{2l} (\mathbf{E} \xi_{j,j,k,l} - m_{kj} m_{lj}). \end{aligned}$$

For each particular model, we will need to replace $\xi_k, \xi_k^{(i)}, \xi_{k,l,k',l'}$ by the exact formulas depending on $\{y_1, \dots, y_d\}$.

12.2. Spiked Population Model.

12.2.1. *Bounding x .* Recall $X = M^{1/2} Y$ and we need to compute

$$\mathbf{E} (u_1^T E u_2)^2 = \frac{1}{n} \sum_{\substack{k,l \leq d \\ k',l' \leq d}} u_{1k} u_{2l} u_{1k'} u_{2l'} (\mathbf{E} \xi_{k,l,k',l'} - m_{kl} m_{k'l'}).$$

Now, we replace $m_{kl} m_{k'l'}$ and $\mathbf{E} \xi_{k,l,k',l'}$ by the exact formulas depending on $\{y_1, \dots, y_d\}$. Since

$$\begin{aligned} X &= M^{1/2} Y = \sum_{1 \leq i \leq d} \sqrt{\lambda_i} u_i u_i^T Y \\ (207) \quad &= \sum_{1 \leq i \leq d} \sqrt{\lambda_i} (u_i^T Y) u_i, \end{aligned}$$

then,

$$\begin{aligned}
(208) \quad \xi_k &:= \sum_{i=1}^d \sqrt{\lambda_i} (u_i^T Y) u_{ik} \\
&= \sum_{i=1}^d \sqrt{\lambda_i} u_{ik} \sum_j u_{ij} y_j \\
&= \sum_{j=1}^d y_j \left(\sum_{i=1}^d u_{ik} u_{ij} \sqrt{\lambda_i} \right) = \sum_{j=1}^d y_j s_{jk},
\end{aligned}$$

in which

$$s_{jk} := \sum_{i=1}^d u_{ik} u_{ij} \sqrt{\lambda_i}.$$

Then,

$$m_{kl} = \mathbf{E}(\xi_k \xi_l) = \sum_{j, j'=1}^d (\mathbf{E} y_j y_{j'}) s_{jk} s_{j'l} = \sum_{\alpha=1}^d s_{\alpha k} s_{\alpha l}.$$

And,

$$\begin{aligned}
(209) \quad \mathbf{E}(\xi_k \xi_l \xi_{k'} \xi_{l'}) &= \sum_{j_1, j_2, j_3, j_4=1}^d \mathbf{E}(y_{j_1} y_{j_2} y_{j_3} y_{j_4}) s_{j_1 k} s_{j_2 l} s_{j_3 k'} s_{j_4 l'} \\
&= \sum_{1 \leq \alpha \neq \beta \leq d} (s_{\alpha k} s_{\alpha l} s_{\beta k'} s_{\beta l'} + s_{\alpha k} s_{\beta l} s_{\alpha k'} s_{\beta l'} + s_{\alpha k} s_{\beta l} s_{\beta k'} s_{\alpha l'}) + \sum_{\alpha=1}^d \mathbf{E} y_{\alpha}^4 s_{\alpha k} s_{\alpha l} s_{\alpha k'} s_{\alpha l'} \\
&= \sum_{\alpha, \beta=1}^d (s_{\alpha k} s_{\alpha l} s_{\beta k'} s_{\beta l'} + s_{\alpha k} s_{\beta l} s_{\alpha k'} s_{\beta l'} + s_{\alpha k} s_{\beta l} s_{\beta k'} s_{\alpha l'}) + \sum_{\alpha=1}^d (\mathbf{E} y_{\alpha}^4 - 3) s_{\alpha k} s_{\alpha l} s_{\alpha k'} s_{\alpha l'}.
\end{aligned}$$

Therefore,

$$(210) \quad \mathbf{E}(\xi_k \xi_l \xi_{k'} \xi_{l'}) - m_{kl} m_{k'l'} = \sum_{\alpha, \beta=1}^d (s_{\alpha k} s_{\beta l} s_{\alpha k'} s_{\beta l'} + s_{\alpha k} s_{\beta l} s_{\beta k'} s_{\alpha l'}) + \sum_{\alpha=1}^d (\mathbf{E} y_{\alpha}^4 - 3) s_{\alpha k} s_{\alpha l} s_{\alpha k'} s_{\alpha l'}.$$

On the other hand,

$$\begin{aligned}
(211) \quad \sum_{k, l, k', l'=1}^d \sum_{\alpha, \beta=1}^d (u_{1k} s_{\alpha k}) (u_{1k'} s_{\alpha k'}) (u_{2l} s_{\beta l}) (u_{2l'} s_{\beta l'}) &= \sum_{\alpha, \beta=1}^d \left(\sum_{k=1}^d u_{1k} s_{\alpha k} \right) \left(\sum_{k'=1}^d u_{1k'} s_{\alpha k'} \right) \left(\sum_{l=1}^d u_{2l} s_{\beta l} \right) \left(\sum_{l'=1}^d u_{2l'} s_{\beta l'} \right) \\
&= \sum_{\alpha, \beta=1}^d \lambda_1 \lambda_2 u_{1\alpha}^2 u_{2\beta}^2 \\
&= \lambda_1 \lambda_2.
\end{aligned}$$

By (210), (211), and the similar computations for $\sum_{k,k',l,l'=1}^d u_{1k}u_{1k'}u_{2l}u_{2l'}$ $\sum_{\alpha,\beta=1}^d (s_{\alpha k}s_{\beta l}s_{\beta k'}s_{\alpha l'})$ and $\sum_{k,k',l,l'=1}^d u_{1k}u_{1k'}u_{2l}u_{2l'}$ $\sum_{\alpha=1}^d (\mathbf{E}y_{\alpha}^4 - 3)s_{\alpha k}s_{\alpha l}s_{\alpha k'}s_{\alpha l'}$, we obtain

$$(212) \quad \mathbf{E}(u_1^T E u_2)^2 = \frac{\lambda_1 \lambda_2}{n} \left(1 + \sum_{\alpha=1}^d (\mathbf{E}y_{\alpha}^4 - 3) u_{1\alpha}^2 u_{2\alpha}^2 \right).$$

By the same procedure, we have

$$\mathbf{E}(u_1^T E u_1)^2 = \frac{\lambda_1^2}{n} \left(1 + \sum_{\alpha=1}^d (\mathbf{E}y_{\alpha}^4 - 3) u_{1\alpha}^4 \right).$$

Therefore, by Markov inequality and union bound, if all y_i s have finite 4th moment, then with probability at least $1 - r^2 t_1^{-2}$,

$$x = O\left(\frac{\lambda_1 t_1}{\sqrt{n}}\right),$$

for any $t_1 > 0$.

12.2.2. *Bounding y .* In this subsection, to reduce the unnecessary complexity of indices, we will shortly write $\sum_j(\dots)$ for $\sum_{j=1}^d(\dots)$ and all other summations will be written in full detail.

First, we compute the first moment of $u_1^T E \sum_{l>r} \frac{u_l u_l^T}{\lambda - \lambda_l} E u_2$, which is

$$(213) \quad \mathbf{E} u_1^T E \sum_{l>r} \frac{u_l u_l^T}{\lambda - \lambda_l} E u_2 = \frac{1}{n} \sum_{l>r} \frac{1}{\lambda - \lambda_l} \mathbf{E} \left(\left(\sum_k u_{1k} \xi_k \right) \left(\sum_j \xi_j u_{lj} \right) - \mathbf{E} \left(\sum_k u_{1k} \xi_k \right) \left(\sum_j \xi_j u_{lj} \right) \right) \times \\ \left(\left(\sum_i u_{2i} \xi_i \right) \left(\sum_m \xi_m u_{lm} \right) - \mathbf{E} \left(\sum_i u_{2i} \xi_i \right) \left(\sum_m \xi_m u_{lm} \right) \right).$$

Following above computation, note that $\sum_k u_{1k} \xi_k = \sum_j y_j (\sum_i u_{ij} \sqrt{\lambda_i} \sum_k u_{ik} u_{1k}) = \sqrt{\lambda_1} \sum_j y_j u_{1j}$, we can rewrite above sum as

$$(214) \quad \frac{1}{n} \sum_{l>r} \frac{\lambda_l \sqrt{\lambda_1 \lambda_2}}{\lambda - \lambda_l} \mathbf{E} \left(\left(\sum_{j_1} y_{j_1} u_{1j_1} \right) \left(\sum_{j_2} y_{j_2} u_{lj_2} \right) - \mathbf{E} \left(\sum_{j_1} y_{j_1} u_{1j_1} \right) \left(\sum_{j_2} y_{j_2} u_{lj_2} \right) \right) \times \\ \left(\left(\sum_{j_3} y_{j_3} u_{2j_3} \right) \left(\sum_{j_4} y_{j_4} u_{lj_4} \right) - \mathbf{E} \sum_{j_3} y_{j_3} u_{2j_3} \left(\sum_{j_4} y_{j_4} u_{lj_4} \right) \right) \\ = \frac{1}{n} \sum_{l>r} \frac{\lambda_l \sqrt{\lambda_1 \lambda_2}}{\lambda - \lambda_l} \mathbf{E} \left(\left(\sum_{j_1} y_{j_1} u_{1j_1} \right) \left(\sum_{j_2} y_{j_2} u_{lj_2} \right) - \sum_{\alpha} u_{1\alpha} u_{l\alpha} \right) \left(\left(\sum_{j_3} y_{j_3} u_{2j_3} \right) \left(\sum_{j_4} y_{j_4} u_{lj_4} \right) - \sum_{\beta} u_{2\beta} u_{l\beta} \right) \\ = \frac{\sqrt{\lambda_1 \lambda_2}}{n} \sum_{l>r} \frac{\lambda_l}{\lambda - \lambda_l} \left[\sum_{\alpha} (\mathbf{E}y_{\alpha}^4 - 3) u_{1\alpha} u_{2\alpha} u_{l\alpha}^2 \right].$$

Now, we compute the second moment of $u_1^T E \sum_{l>r} \frac{u_l u_l^T}{\lambda - \lambda_l} E u_2$, which is

$$\frac{1}{n^4} \sum_{\substack{k,j,m,i, \\ k',j',m',i' \\ l>r \\ l'>r}} u_{1k} u_{1k'} u_{2i} u_{2i'} \frac{u_{lj} u_{lm} u_{l'j'} u_{l'm'}}{(\lambda - \lambda_l)(\lambda - \lambda_{l'})} \mathbf{E} \left(\sum_{I_1=1}^n \cdots \right) \left(\sum_{I_2=1}^n \cdots \right) \left(\sum_{I_3=1}^n \cdots \right) \left(\sum_{I_4=1}^n \cdots \right).$$

Here

$$\begin{aligned} \sum_{I_1=1}^n \cdots &= \sum_{I_1=1}^n \left(\sum_k u_{1k} \xi_k^{(I_1)} \right) \left(\sum_j \xi_j^{(I_1)} u_{lj} \right) - \mathbf{E} \left(\sum_k u_{1k} \xi_k^{(I_1)} \right) \left(\sum_j \xi_j^{(I_1)} u_{lj} \right), \\ \sum_{I_2=1}^n \cdots &= \sum_{I_2=1}^n \left(\sum_i u_{2i} \xi_i^{(I_2)} \right) \left(\sum_m \xi_m^{(I_2)} u_{lm} \right) - \mathbf{E} \left(\sum_i u_{2i} \xi_i^{(I_2)} \right) \left(\sum_m \xi_m^{(I_2)} u_{lm} \right), \\ \sum_{I_3=1}^n \cdots &= \sum_{I_3=1}^n \left(\sum_{k'} u_{1k'} \xi_{k'}^{(I_3)} \right) \left(\sum_{j'} \xi_{j'}^{(I_3)} u_{l'j'} \right) - \mathbf{E} \left(\sum_{k'} u_{1k'} \xi_{k'}^{(I_3)} \right) \left(\sum_{j'} \xi_{j'}^{(I_3)} u_{l'j'} \right), \\ \sum_{I_4=1}^n \cdots &= \sum_{I_4=1}^n \left(\sum_{i'} u_{2i'} \xi_{i'}^{(I_4)} \right) \left(\sum_{m'} \xi_{m'}^{(I_4)} u_{l'm'} \right) - \mathbf{E} \left(\sum_{i'} u_{2i'} \xi_{i'}^{(I_4)} \right) \left(\sum_{m'} \xi_{m'}^{(I_4)} u_{l'm'} \right). \end{aligned}$$

Note that the expectations of the term $(\sum_k u_{1k} \xi_k^{(I_1)}) (\sum_j \xi_j^{(I_1)} u_{lj}) - \mathbf{E}(\sum_k u_{1k} \xi_k^{(I_1)}) (\sum_j \xi_j^{(I_1)} u_{lj})$ and the similar ones are zero. Therefore, after totally expanding above sum, the non trivial terms can be grouped up into the following cases:

- $I_1 = I_2 = I_3 = I_4 = I$, we can drop the upper index I and pay a factor n . The following sum is obtained:

$$M_2 = \frac{1}{n^3} \sum_{\substack{k,j,m,i, \\ k',j',m',i' \\ l>r \\ l'>r}} u_{1k} u_{1k'} u_{2i} u_{2i'} \frac{u_{lj} u_{lm} u_{l'j'} u_{l'm'}}{(\lambda - \lambda_l)(\lambda - \lambda_{l'})} \mathbf{E}(\xi_k \xi_j - m_{kj}) (\xi_m \xi_i - m_{mi}) (\xi_{k'} \xi_{j'} - m_{k'j'}) (\xi_{m'} \xi_{i'} - m_{m'i'}).$$

- $I_1 = I_3 \neq I_2 = I_4$, we can drop the upper indices I_1, I_2 and pay a factor of $n(n-1)$. The following sum is obtained

$$\begin{aligned} M_1(I_1 = I_3) &= \frac{n-1}{n^3} \sum_{\substack{k,j,m,i, \\ k',j',m',i' \\ l>r \\ l'>r}} u_{1k} u_{1k'} u_{2i} u_{2i'} \frac{u_{lj} u_{lm} u_{l'j'} u_{l'm'}}{(\lambda - \lambda_l)(\lambda - \lambda_{l'})} \times \\ &\quad \left(\mathbf{E}(\xi_k \xi_j - m_{kj}) (\xi_{k'} \xi_{j'} - m_{k'j'}) \right) \left(\mathbf{E}(\xi_m \xi_i - m_{mi}) (\xi_{m'} \xi_{i'} - m_{m'i'}) \right), \end{aligned}$$

- $I_1 = I_2 \neq I_3 = I_4$, we follow the same procedure and obtain the sum of $M_1(I_1 = I_2)$.
- $I_1 = I_4 \neq I_2 = I_3$, we follow the same procedure and obtain the sum of $M_1(I_1 = I_4)$.

We have,

$$\mathbf{E} \left(u_1^T E \sum_{l>r} \frac{u_l u_l^T}{\lambda - \lambda_l} E u_2 \right)^2 = M_2 + \sum_{j=2}^4 M_1(I_1 = I_j).$$

Similar to the first moment, we obtain

(215)

$$\begin{aligned}
M_1(I_1 = I_3) &= \frac{\lambda_1 \lambda_2}{n^2} \sum_{\substack{l > r \\ l' > r}} \frac{\lambda_l \lambda_{l'}}{(\lambda - \lambda_{l'}) (\lambda - \lambda_l)} \times \\
&\quad \left(\mathbf{E} \left(\left(\sum_{j_1} y_{j_1} u_{1j_1} \right) \left(\sum_{j_2} y_{j_2} u_{lj_2} \right) - \sum_{\alpha} u_{1\alpha} u_{l\alpha} \right) \left(\left(\sum_{j_3} y_{j_3} u_{1j_3} \right) \left(\sum_{j_4} y_{j_4} u_{l'j_4} \right) - \sum_{\beta} u_{1\beta} u_{l'\beta} \right) \right) \times \\
&\quad \left(\mathbf{E} \left(\left(\sum_{j_1} y_{j_1} u_{2j_1} \right) \left(\sum_{j_2} y_{j_2} u_{lj_2} \right) - \sum_{\alpha} u_{2\alpha} u_{l\alpha} \right) \left(\left(\sum_{j_3} y_{j_3} u_{2j_3} \right) \left(\sum_{j_4} y_{j_4} u_{l'j_4} \right) - \sum_{\beta} u_{2\beta} u_{l'\beta} \right) \right) \\
&= \frac{\lambda_1 \lambda_2}{n^2} \sum_{\substack{l > r \\ l' > r}} \frac{\lambda_l \lambda_{l'}}{(\lambda - \lambda_{l'}) (\lambda - \lambda_l)} \times \\
&\quad \left(\sum_{\beta} u_{l\beta} u_{l'\beta} + \sum_{\alpha} (\mathbf{E} y_{\alpha}^4 - 3) u_{1\alpha}^2 u_{l\alpha} u_{l'\alpha} \right) \left(\sum_{\beta} u_{l\beta} u_{l'\beta} + \sum_{\alpha} (\mathbf{E} y_{\alpha}^4 - 3) u_{2\alpha}^2 u_{l\alpha} u_{l'\alpha} \right) \\
&= \frac{\lambda_1 \lambda_2}{n^2} \sum_{l > r} \frac{\lambda_l^2}{(\lambda - \lambda_l)^2} \left(1 + \sum_{\alpha} (\mathbf{E} y_{\alpha}^4 - 3) u_{1\alpha}^2 u_{l\alpha}^2 \right) \left(1 + \sum_{\alpha} (\mathbf{E} y_{\alpha}^4 - 3) u_{2\alpha}^2 u_{l\alpha}^2 \right) + \\
&\quad \frac{\lambda_1 \lambda_2}{n^2} \sum_{l \neq l' > r} \frac{\lambda_l \lambda_{l'}}{(\lambda - \lambda_l) (\lambda - \lambda_{l'})} \left(\sum_{\alpha} (\mathbf{E} y_{\alpha}^4 - 3) u_{1\alpha}^2 u_{l\alpha} u_{l'\alpha} \right) \left(\sum_{\beta} (\mathbf{E} y_{\beta}^4 - 3) u_{2\beta}^2 u_{l\beta} u_{l'\beta} \right)
\end{aligned}$$

Similar computation on $M_1(I_1 = I_2)$:

(216)

$$\begin{aligned}
M_1(I_1 = I_2) &= \frac{\lambda_1 \lambda_2}{n^2} \sum_{\substack{l > r \\ l' > r}} \frac{\lambda_l \lambda_{l'}}{(\lambda - \lambda_{l'}) (\lambda - \lambda_l)} \times \\
&\quad \left(\mathbf{E} \left(\left(\sum_{j_1} y_{j_1} u_{1j_1} \right) \left(\sum_{j_2} y_{j_2} u_{lj_2} \right) - \sum_{\alpha} u_{1\alpha} u_{l\alpha} \right) \left(\left(\sum_{j_3} y_{j_3} u_{2j_3} \right) \left(\sum_{j_4} y_{j_4} u_{l'j_4} \right) - \sum_{\beta} u_{2\beta} u_{l'\beta} \right) \right) \times \\
&\quad \left(\mathbf{E} \left(\left(\sum_{j_1} y_{j_1} u_{1j_1} \right) \left(\sum_{j_2} y_{j_2} u_{l'j_2} \right) - \sum_{\alpha} u_{1\alpha} u_{l'\alpha} \right) \left(\left(\sum_{j_3} y_{j_3} u_{2j_3} \right) \left(\sum_{j_4} y_{j_4} u_{lj_4} \right) - \sum_{\beta} u_{2\beta} u_{l\beta} \right) \right) \\
&= \frac{\lambda_1 \lambda_2}{n} \sum_{\substack{l > r \\ l' > r}} \frac{\lambda_l \lambda_{l'}}{(\lambda - \lambda_{l'}) (\lambda - \lambda_l)} \left[\sum_{\alpha} (\mathbf{E} y_{\alpha}^4 - 3) u_{1\alpha} u_{2\alpha} u_{l\alpha} u_{l'\alpha} \right]^2.
\end{aligned}$$

And,

$$(217) \quad M_1(I_1 = I_4) = \frac{\lambda_1 \lambda_2}{n} \left[\sum_{l > r} \frac{\lambda_l}{\lambda - \lambda_l} \sum_{\alpha} (\mathbf{E} y_{\alpha}^4 - 3) u_{1\alpha} u_{2\alpha} u_{l\alpha}^2 \right]^2.$$

$$\begin{aligned}
(218) \quad M_2 &= \frac{\lambda_1 \lambda_2}{n^3} \sum_{l, l' > r} \frac{\lambda_l \lambda_{l'}}{(\lambda - \lambda_l)(\lambda - \lambda_{l'})} \mathbf{E} \left(\sum_{j_1} y_{j_1} u_{l j_1} \right)^2 \left(\sum_{j_2} y_{j_2} u_{l j_2} \right)^2 \left(\sum_{j_3} y_{j_3} u_{l j_3} \right)^2 \left(\sum_{j_4} y_{j_4} u_{l j_4} \right)^2 \\
&\leq C \frac{\lambda_1 \lambda_2}{n^3} \left(\sum_{l > r} \frac{\lambda_l}{\lambda - \lambda_l} \right)^2, \text{ for some universal constant } C < 8!.
\end{aligned}$$

Combining (215), (216), (218), we have, with probability at least $1 - t^{-2}$,

$$y_{12} \leq C_r \frac{\sqrt{\lambda_1 \lambda_2} t}{n} \frac{\sum_{l > r} \lambda_l}{\lambda_p - \lambda_{r+1}} + \frac{\sqrt{\lambda_1 \lambda_2} t}{n} \sqrt{\sum_{l > r} \frac{\lambda_l^2}{(\lambda_p - \lambda_l)^2}},$$

in which

$$C_r := C \left(\max_{\substack{1 \leq i < j \leq r \\ l, l' > r}} \left| \sum_{\alpha} (\mathbf{E} y_{\alpha}^4 - 3) u_{i\alpha} u_{j\alpha} u_{l\alpha} u_{l'\alpha} \right| + \frac{1}{\sqrt{n}} \right).$$

12.3. Estimate Covariance Noise $\|E\|$. In this section, we will estimate the covariance noise for each spiked models. The main tool is Matrix Bernstein inequality. Assume that

$$\|XX^T - M\| \leq L \text{ and } \|\mathbf{E}(XX^T - M)^2\| = V.$$

Theorem 12.4 (Matrix Bernstein inequality). *Consider a finite sequences $\{S_k\}$ of independent, random matrices with common dimension $d_1 \times d_2$. Assume that*

$$\mathbf{E}S_k = 0 \text{ and } \|S_j\| \leq L \text{ for all index } k.$$

Introduce the random matrix

$$Z := \sum_k S_k.$$

Let $v(Z)$ be the matrix variance statistic of the sum:

$$\begin{aligned}
(219) \quad v(Z) &:= \max \{ \|\mathbf{E}(ZZ^*)\|, \|\mathbf{E}Z^*Z\| \} \\
&= \max \left\{ \left\| \sum_k \mathbf{E}(S_k S_k^*) \right\|, \left\| \sum_k \mathbf{E}S_k^* S_k \right\| \right\}
\end{aligned}$$

Then,

$$(220) \quad \mathbf{E}\|Z\| \leq \sqrt{2v(Z) \log(d_1 + d_2)} + \frac{1}{3}L \log(d_1 + d_2).$$

Furthermore for all $t \geq 0$,

$$(221) \quad \mathbf{P} \{ \|Z\| \geq t \} \leq (d_1 + d_2) \exp \left(\frac{-t^2/2}{v(Z) + Lt/3} \right).$$

Applying Matrix Bernstein Inequality on $E = \sum_{i=1}^n E_i$, $E_i = \frac{1}{n} (X_i X_i^T - M)$, without further information about X , we obtain:

Lemma 12.5.

$$(222) \quad \mathbf{P} \{ \|E\| \geq t \} \leq (d+n) \exp \left(\frac{-t^2 n/2}{V + 2Lt/3} \right).$$

And hence, with high probability,

$$(223) \quad \|E\| \leq \sqrt{\frac{V \log(n+d)}{n}} + CL \log(n+d).$$

Proof. We need to bound

$$\|E_i\| = \left\| \frac{1}{n} (XX^T - M) \right\| = \frac{1}{n} \|XX^T - M\|,$$

and

$$\left\| \sum_i \frac{1}{n^2} \mathbf{E}(X_i X_i^T - M)^2 \right\| = \left\| \frac{1}{n} \mathbf{E}(XX^T - M)^2 \right\|.$$

All together implies that

$$(224) \quad \mathbf{P} \{ \|E\| \geq t \} \leq (d+n) \exp \left(\frac{-t^2/2}{\frac{\mathbf{E}(XX^T - M)^2}{n} + \frac{2Lt}{3n}} \right) = (d+n) \exp \left(\frac{-t^2 n/2}{V + 2Lt/3} \right).$$

We complete the proof. \square

We are going to show the more concrete bounds for the considered models above.

Lemma 12.6. *If X is spiked population model, then with high probability,*

$$(225) \quad \|E\| \leq \left(2 + \max_{1 \leq i \leq d} |\mathbf{E}y_i^4 - 3| \right) \left(\lambda_1 \sqrt{\frac{r_\lambda \log(n+d)}{n}} + L \log(n+d) \right).$$

Proof. Similar to the proof of Lemma 12.5, we need to bound

$$\left\| \frac{1}{n} (\mathbf{E}XX^T XX^T - M^2) \right\|.$$

Recall $X := M^{1/2}Y$, therefore,

$$(226) \quad \begin{aligned} \mathbf{E}XX^T XX^T &= \mathbf{E}M^{1/2}YY^T MYY^T M^{1/2} \\ &= M^{1/2} \mathbf{E}YY^T MYY^T M^{1/2} \\ &= M^{1/2} \mathbf{E}(Y^T M Y) Y Y^T M^{1/2} \\ &= M^{1/2} \left(\mathbf{E} \left(\sum_{i,j} m_{ij} y_i y_j \right) (y_k y_l) \right)_{1 \leq k, l \leq d} M^{1/2}. \end{aligned}$$

Since $\mathbf{E}y_i = 0, \mathbf{E}y_i^2 = 1 \forall 1 \leq i \leq d$, we have

$$(227) \quad \begin{aligned} \mathbf{E}\left(\sum_{i,j} m_{ij}y_iy_j\right)(y_ky_l) &= 2m_{kl} \text{ for all } k \neq l, \\ \mathbf{E}\left(\sum_{i,j} m_{ij}y_iy_j\right)(y_ky_l) &= 2m_{kk} + \left(\sum_i m_{ii}\right) + (\mathbf{E}y_k^4 - 3)m_{kk}. \end{aligned}$$

Thus,

$$(228) \quad \begin{aligned} \mathbf{E}X X^T X X^T &= \mathbf{E}M^{1/2}Y Y^T M Y Y^T M^{1/2} \\ &= 2M^2 + \left(\sum_i m_{ii}\right)M + M^{1/2}DM^{1/2}, \end{aligned}$$

D is a diagonal matrix, whose diagonal entries are $(\mathbf{E}y_k^4 - 3)m_{kk}, \forall 1 \leq k \leq d$. \square

Thus

$$(229) \quad \begin{aligned} \|\mathbf{E}X X^T X X^T - M^2\| &= \|M^2 + \left(\sum_i m_{ii}\right)M + M^{1/2}DM^{1/2}\| \\ &\leq \left(2 + \max_{1 \leq i \leq d} |\mathbf{E}y_i^4 - 3|\right) \left(\sum_i \lambda_i\right)\lambda_1. \end{aligned}$$

13. APPENDIX C: PROOFS OF LEMMAS 9.10, 9.11, 9.13, AND 9.14

13.1. Proof of Lemma 9.10. We can bound $M_1(\alpha; \beta)$ in this case as follows. We replace $w^{2(k-1)}$ by $w^{2(k-1-r_c(T,L))}\|E\|^{2r_c(T,L)}$ and obtain

$$M_1(\alpha; \beta) \leq \sum_{T=2}^{s_2} \binom{s_2}{T} p^{T-1} r^{s_2-T} \sum_{L \in \mathcal{L}(T,s); 1 \leq r_c(T,L) \leq k-1} \frac{\|E\|^{s_1-(k-1)+2r_c(T,L)} w^{2(k-1-r_c(T,L))} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T,L)} \bar{\lambda}^{s_1+r_c(T,L)}}.$$

Similar to the argument in (111), we can arrange the factors in the RHS and get

$$(230) \quad \begin{aligned} M_1(\alpha; \beta) &\leq \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); 1 \leq r_c(T,L) \leq k-1} p^{T-1} r^{s_2-T} \left(\frac{\|E\|}{\bar{\lambda}}\right)^{T_1} \left(\frac{x}{\delta_S}\right)^{T_2} \left(\frac{w}{\sqrt{\bar{\lambda}\delta_S}}\right)^{T_3} \\ &= \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); 1 \leq r_c(T,L) \leq k-1} \left(\frac{p}{r}\right)^{T-(r_c(T,L)+1)} p^{\frac{k-1-s_1}{2}} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{T_1} \left(\frac{rx}{\delta_S}\right)^{T_2} \left(\frac{\sqrt{rw}}{\sqrt{\bar{\lambda}\delta_S}}\right)^{T_3}, \end{aligned}$$

where

$$\begin{aligned} T_1 &= s_1 - (k-1) + 2r_c(T,L), \\ T_2 &= s_2 - k, \\ T_3 &= 2(k-1-r_c(T,L)). \end{aligned}$$

Since (α, β) is type II, $s_1 \geq k-1$ and hence $0 \leq p^{\frac{k-1-s_1}{2}} \leq 1$. By Lemma 9.6, we also have $0 \leq \left(\frac{p}{r}\right)^{T-(r_c(T,L)+1)} \leq 1$. We can omit these factors and obtain

(231)

$$\begin{aligned} M_1(\alpha; \beta) &\leq \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); 1 \leq r_c(T,L) \leq k-1} \left(\frac{\sqrt{p}\|E\|}{\lambda}\right)^{T_1} \left(\frac{rx}{\delta_S}\right)^{T_2} \left(\frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S}\right)^{T_3} \\ &= \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); 1 \leq r_c(T,L) \leq k-1} \frac{\sqrt{p}\|E\|}{\lambda} \times \left(\frac{\sqrt{p}\|E\|}{\lambda}\right)^{T_1-1} \left(\frac{rx}{\delta_S}\right)^{T_2} \left(\frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S}\right)^{T_3}. \end{aligned}$$

By the setting of $M_1(\alpha; \beta)$ for (α, β) of type II, we have $T_2, T_3 \geq 0$. Particularly, $T_1 \geq 2r_c(T, L) \geq 2$. Therefore, by Lemma 9.4, $M_1(\alpha; \beta)$ is less than or equals

$$\begin{aligned} &\sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T,s) \\ 1 \leq r_c(T,L) \leq k-1}} \left(\frac{\sqrt{p}\|E\|}{\lambda}\right) \max\left\{\frac{\|E\|}{\lambda}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S}\right\}^{T_1+T_2+T_3-1} \\ &= \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T,s) \\ 1 \leq r_c(T,L) \leq k-1}} \left(\frac{\sqrt{p}\|E\|}{\lambda}\right) \max\left\{\frac{\|E\|}{\lambda}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S}\right\}^{s-1}, \end{aligned}$$

since $T_1 + T_2 + T_3 = s$. Moreover, by Assumption **D0** and the fact that $\delta \leq \delta_S$, this RHS is at most

$$(232) \quad \left(\frac{\sqrt{p}\|E\|}{\lambda}\right) \frac{\sum_{T=2}^{s_2} \sum_{L \in \mathcal{L}(T,s); 1 \leq r_c(T,L) \leq k-1} \binom{s_2}{T}}{12^{s-1}}.$$

This proves Lemma 9.10.

13.2. Proof of Lemma 9.11. We bound $M_2(\alpha; \beta)$ in this case as follows. Replacing $w^{2(k-1)}$ by $\|E\|^{2(k-1)}$, we obtain

$$M_2(\alpha; \beta) \leq \sum_{T=2}^{s_2} \binom{s_2}{T} p^{T-1} r^{s_2-T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) > k-1} \frac{\|E\|^{s_1+(k-1)} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T,L)} \lambda^{s_1+r_c(T,L)}},$$

whose RHS can be rewritten into

$$\sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) > k-1} p^{T-1} r^{s_2-T} \left(\frac{x}{\delta_S}\right)^{s_2-1-r_c(T,L)} \left(\frac{x}{\lambda}\right)^{r_c(T,L)-(k-1)} \left(\frac{\|E\|}{\lambda}\right)^{k-1+s_1}.$$

Distributing each factors r, p, \sqrt{p} to $\frac{x}{\delta_S}, \frac{x}{\lambda}, \frac{\|E\|}{\lambda}$, respectively, we can rewrite the RHS as

$$\sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) > k-1}} \left(\frac{p}{r}\right)^{T-(r_c(T,L)+1)} p^{\frac{k-1-s_1}{2}} \left(\frac{rx}{\delta_S}\right)^{s_2-1-r_c(T,L)} \left(\frac{px}{\lambda}\right)^{r_c(T,L)-(k-1)} \left(\frac{\sqrt{p}\|E\|}{\lambda}\right)^{k-1+s_1}.$$

Since $T \geq r_c(T, L) + 1$ (Lemma 9.6) and $s_1 - (k-1) \geq 0$ ((α, β) is Type II),

$$0 \leq \left(\frac{p}{r}\right)^{T-(r_c(T,L)+1)} \leq 1 \text{ and } 0 \leq p^{\frac{k-1-s_1}{2}} \leq 1.$$

We can omit these factors and obtain

$$\begin{aligned} M_2(\alpha; \beta) &\leq \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) > k-1}} \left(\frac{rx}{\delta_S}\right)^{s_2-1-r_c(T,L)} \left(\frac{px}{\bar{\lambda}}\right)^{r_c(T,L)-(k-1)} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{k-1+s_1} \\ &= \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) > k-1}} \frac{px}{\bar{\lambda}} \times \left(\frac{rx}{\delta_S}\right)^{s_2-1-r_c(T,L)} \left(\frac{px}{\bar{\lambda}}\right)^{r_c(T,L)-(k-1)-1} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{k-1+s_1}. \end{aligned}$$

By the setting of $M_2(\alpha; \beta)$, $r_c(T, L) - (k - 1) - 1 \geq 0$. By definitions of $s_1, s_2, r_c(T, L)$, we also have $s_2 - 1 - r_c(T, L), k - 1 + s_1 \geq 0$. Applying Lemma 9.4, we finally obtain

(233)

$$\begin{aligned} M_2(\alpha; \beta) &\leq \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) > k-1}} \frac{px}{\bar{\lambda}} \times \max \left\{ \frac{rx}{\delta_S}, \frac{px}{\bar{\lambda}}, \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right\}^{s_2-1-r_c(T,L)+r_c(T,L)-(k-1)-1+k-1+s_1} \\ &= \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) > k-1}} \frac{px}{\bar{\lambda}} \times \max \left\{ \frac{rx}{\delta_S}, \frac{px}{\bar{\lambda}}, \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right\}^{s_2+s_1-2} \\ &\leq \sum_{T=2}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) > k-1} \frac{px}{\bar{\lambda}} \max \left\{ \frac{rx}{\delta_S}, \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \right\}^{s-1} \\ &\quad (\text{since } s_1 + s_2 - 2 = s - 1 \text{ and } \frac{px}{\bar{\lambda}} \leq \frac{rx}{\delta_S}) \\ &\leq \frac{px}{\bar{\lambda}} \frac{\sum_{T=2}^{s_2} \sum_{L \in \mathcal{L}(T,s); r_c > k-1} \binom{s_2}{T}}{12^{s-1}} \quad (\text{by Assumption D0 and the fact that } \delta \leq \delta_S). \end{aligned}$$

This proves Lemma 9.11.

13.3. Proof of Lemma 9.13. We bound $M_1(\alpha; \beta)$ as follows. Similarly, replacing w^{2k-1} by $w^{2k-2-2r_c(T,L)}\|E\|^{2r_c(T,L)+1}$, we have

$$M_1(\alpha; \beta) \leq \sum_{T=1}^{s_2} \binom{s_2}{T} p^{T-1/2} r^{s_2-T} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) \leq k-1}} \frac{\|E\|^{s_1-(k-1)+2r_c(T,L)} w^{2(k-1-r_c(T,L))} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T,L)} \bar{\lambda}^{s_1+r_c(T,L)}}.$$

Rearranging the factors in the RHS, we further obtain

$$\begin{aligned} M_1(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} p^{T-1/2} r^{s_2-T} \left(\frac{\|E\|}{\bar{\lambda}}\right)^{T_1} \left(\frac{x}{\delta_S}\right)^{T_2} \left(\frac{w}{\sqrt{\lambda}\delta_S}\right)^{T_3} \\ &= \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} \left(\frac{p}{r}\right)^{T-(r_c(T,L)+1)} p^{\frac{k-s_1}{2}} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{T_1} \left(\frac{rx}{\delta_S}\right)^{T_2} \left(\frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S}\right)^{T_3}, \end{aligned}$$

where

$$\begin{aligned} T_1 &= s_1 - (k - 1) + 2r_c(T, L), \\ T_2 &= s_2 - k, \\ T_3 &= 2(k - 1 - r_c(T, L)). \end{aligned}$$

Since $T \geq r_c(T, L) + 1$, $s_1 \geq k$, and then $0 \leq \left(\frac{p}{r}\right)^{T-(r_c(T,L)+1)} p^{\frac{k-s_1}{2}} \leq 1$, we have

$$\begin{aligned} M_1(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{T_1} \left(\frac{rx}{\delta_S}\right)^{T_2} \left(\frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S}\right)^{T_3} \\ &= \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \times \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{T_1-1} \left(\frac{rx}{\delta_S}\right)^{T_2} \left(\frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S}\right)^{T_3} \end{aligned}$$

By the setting of $M_1(\alpha; \beta)$, $r_c(T, L) \leq k - 1$ and hence $T_3 \geq 0$. We also have $T_2 \geq 0$ by definition of s_2 . Since (α, β) is type III, $s_1 \geq k$ and hence $T_1 - 1 \geq 0$. Therefore, by Lemma 9.4, we finally obtain

(234)

$$\begin{aligned} M_1(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \times \left\{ \frac{\sqrt{p}\|E\|}{\bar{\lambda}}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right\}^{T_1+T_2+T_3-1} \\ &= \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \times \left\{ \frac{\sqrt{p}\|E\|}{\bar{\lambda}}, \frac{rx}{\delta_S}, \frac{\sqrt{rw}}{\sqrt{\lambda}\delta_S} \right\}^{s-1} \quad (\text{since } T_1 + T_2 + T_3 = s) \\ &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) \leq k-1} \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \times \frac{1}{12^{s-1}} \quad (\text{by Assumption D0 and the fact that } \delta \leq \delta_S) \\ &= \left(\sum_{T=1}^{s_2} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) \leq k-1}} \binom{s_2}{T} \right) \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \times \frac{1}{12^{s-1}} \\ &\leq \left(\sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T,s)} \binom{s_2}{T} \right) \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \times \frac{1}{12^{s-1}} \\ &\leq \frac{\sqrt{p}\|E\|}{\bar{\lambda}} \times \frac{2^{s_2+s-1}}{12^{s-1}} \quad (\text{by (113)}). \end{aligned}$$

This proves Lemma 9.13

13.4. Proof of Lemma 9.14. We bound $M_2(\alpha; \beta)$ in this case as follows. Replacing w^{2k-1} by $\|E\|^{2k-1}$, we get

$$M_2(\alpha; \beta) \leq \sum_{T=1}^{s_2} \binom{s_2}{T} p^{T-1/2} r^{s_2-T} \sum_{L \in \mathcal{L}(T,s); r_c(T,L) > k-1} \frac{\|E\|^{s_1+k-1} x^{s_2-k}}{\delta_S^{s_2-1-r_c(T,L)} \bar{\lambda}^{s_1+r_c(T,L)}}.$$

Similar to the argument in (115), the RHS can be rewritten as

(235)

$$\sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T,s) \\ r_c(T,L) > k-1}} \left(\frac{p}{r}\right)^{T-(r_c(T,L)+1)} p^{\frac{k-s_1}{2}} \left(\frac{rx}{\delta_S}\right)^{s_2-1-r_c(T,L)} \left(\frac{px}{\bar{\lambda}}\right)^{r_c(T,L)-(k-1)} \left(\frac{\sqrt{p}\|E\|}{\bar{\lambda}}\right)^{k-1+s_1}.$$

By Lemma 9.6, we have $T - (r_c(T, L) + 1) \geq 0$ and hence $0 \leq \left(\frac{p}{r}\right)^{T-(r_c(T, L)+1)} \leq 1$. Since (α, β) is of Type III, $s_1 \geq k$ and then $0 \leq p^{\frac{k-s_1}{2}} \leq 1$. Omitting these factors, we obtain

$$\begin{aligned} M_2(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \left(\frac{rx}{\delta_S}\right)^{s_2-1-r_c(T, L)} \left(\frac{px}{\lambda}\right)^{r_c(T, L)-(k-1)} \left(\frac{\sqrt{p}\|E\|}{\lambda}\right)^{k-1+s_1} \\ &= \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \frac{px}{\lambda} \times \left(\frac{rx}{\delta_S}\right)^{s_2-1-r_c(T, L)} \left(\frac{px}{\lambda}\right)^{r_c(T, L)-(k-1)-1} \left(\frac{\sqrt{p}\|E\|}{\lambda}\right)^{k-1+s_1}. \end{aligned}$$

By the setting of $M_2(\alpha; \beta)$, $r_c(T, L) - (k-1) - 1 \geq 0$. We also have $s_2 - 1 - r_c(T, L) \geq s_2 - T \geq 0$ and $k - 1 + s_1 \geq 0$. By Lemma 9.4, we further obtain

$$\begin{aligned} M_2(\alpha; \beta) &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \frac{px}{\lambda} \max \left\{ \frac{rx}{\delta_S}, \frac{px}{\lambda}, \frac{\sqrt{p}\|E\|}{\lambda} \right\}^{s_1+s_2-2} \\ &= \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \frac{px}{\lambda} \max \left\{ \frac{rx}{\delta_S}, \frac{px}{\lambda}, \frac{\sqrt{p}\|E\|}{\lambda} \right\}^{s-1} \\ &\leq \sum_{T=1}^{s_2} \binom{s_2}{T} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \frac{px}{\lambda} \max \left\{ \frac{rx}{\delta_S}, \frac{\sqrt{p}\|E\|}{\lambda} \right\}^{s-1} \quad (\text{since } \frac{px}{\lambda} \leq \frac{rx}{\delta_S}) \\ (236) \quad &= \left(\sum_{T=1}^{s_2} \sum_{\substack{L \in \mathcal{L}(T, s) \\ r_c(T, L) > k-1}} \binom{s_2}{T} \right) \times \frac{px}{\lambda} \max \left\{ \frac{rx}{\delta_S}, \frac{\sqrt{p}\|E\|}{\lambda} \right\}^{s-1} \\ &\leq \left(\sum_{T=1}^{s_2} \sum_{L \in \mathcal{L}(T, s)} \binom{s_2}{T} \right) \times \frac{px}{\lambda} \max \left\{ \frac{rx}{\delta_S}, \frac{\sqrt{p}\|E\|}{\lambda} \right\}^{s-1} \\ &\leq 2^{s+s-1} \frac{px}{\lambda} \max \left\{ \frac{rx}{\delta_S}, \frac{\sqrt{p}\|E\|}{\lambda} \right\}^{s-1} \quad (\text{by (113)}) \\ &\leq \frac{px}{\lambda} \frac{2^{s_2+s-1}}{12^{s-1}} \quad (\text{by Assumption D0 and the fact that } \delta \leq \delta_S). \end{aligned}$$

This proves Lemma 9.14.

14. APPENDIX D: SHARPNESS OF THEOREM 2.12

In this section, we fix the $n \times n$ matrix

$$A := \begin{pmatrix} \lambda & 0 & 0 & \cdots & 0 \\ 0 & \lambda - \delta & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ \cdots & & & & \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix},$$

for some $\lambda > \delta > 0$. Clearly, $\lambda_1 = \lambda, \lambda_2 = \lambda - \delta$ are the first and second eigenvalues of A with the corresponding eigenvectors e_1, e_2 , respectively. We will show that our bound for $\|\tilde{u}_1 \tilde{u}_1^T - u_1^T u_1^T\| = \|\tilde{\Pi}_1 - \Pi_1\|$ is sharp by setting up three different noise matrices E , in which each term from Theorem 2.12 dominates and represents the true value of the perturbation. We set $\bar{\lambda} = \lambda$. Hence, $r = 2$ and $N_{\bar{\lambda}} = \{1, 2\}$.

In the following subsections, we will compute $\|\tilde{u}_1 \tilde{u}_1^T - u_1 u_1^T\| = \|\tilde{\Pi}_1 - \Pi_1\|$ by the sine formula:

$$(237) \quad \|\tilde{u}_1 \tilde{u}_1^T - u_1^T u_1^T\| = |\sin \angle(\tilde{u}_1, u_1)| = \sqrt{1 - \cos^2 \angle(\tilde{u}_1, u_1)} = \sqrt{1 - \langle \tilde{u}_1, u_1 \rangle^2}.$$

Therefore, given our setting $u_1 := e_1 = [1\ 0\ 0 \cdots 0]^T$ and that $\tilde{u}_1 = [\tilde{u}_{11} \ \tilde{u}_{12} \ \tilde{u}_{13} \ \cdots \ \tilde{u}_{1n}]^T$, we have

$$(238) \quad \|\tilde{u}_1 \tilde{u}_1^T - u_1^T u_1^T\| = \sqrt{\sum_{j>1} \tilde{u}_{1j}^2}.$$

14.1. Necessity of the first term. Consider

$$E = \begin{pmatrix} 0 & 0 & \mu & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 \\ \mu & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 \\ \cdots & & & & & \\ 0 & 0 & 0 & 0 & \cdots & 0 \end{pmatrix}, \text{ for some } 0 < \mu < \lambda.$$

Thus, $\tilde{A} = \begin{pmatrix} \lambda & 0 & \mu & \cdots & 0 \\ 0 & \lambda - \delta & 0 & \cdots & 0 \\ \mu & 0 & 0 & \cdots & 0 \\ \cdots & & & & \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix}$. By a routine computation, we can show that the first

eigenvalue $\tilde{\lambda}_1 = \frac{\lambda + \sqrt{\lambda^2 + 4\mu^2}}{2}$, with corresponding eigenvector

$$\tilde{u}_1 = [\tilde{u}_{11} \ 0 \ \tilde{u}_{13} \ 0 \ 0 \dots 0]^T,$$

where

$$\tilde{u}_{11} := \frac{\lambda + \sqrt{\lambda^2 + 4\mu^2}}{2\sqrt{\tilde{\lambda}_1^2 + \mu^2}} \text{ and } \tilde{u}_{13} := \frac{\mu}{\sqrt{\tilde{\lambda}_1^2 + \mu^2}}.$$

By 238, we have

$$(239) \quad \|\tilde{u}_1 \tilde{u}_1^T - u_1^T u_1^T\| = |\tilde{u}_{13}| = \frac{\mu}{\sqrt{\tilde{\lambda}_1^2 + \mu^2}} = \Theta\left(\frac{\mu}{\lambda}\right) = \Theta\left(\frac{\|E\|}{\lambda}\right).$$

Moreover, in this setting $x = y = 0$ (see Definition 2.2). The first term $\frac{\|E\|}{\lambda}$ obviously dominates and presents the true value of $\|\tilde{u}_1 \tilde{u}_1^T - u_1 u_1^T\|$.

14.2. **Necessity of the second term.** Consider

$$E = \begin{pmatrix} 0 & \mu & 0 & 0 & \cdots & 0 \\ \mu & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & \sqrt{\lambda} & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 \\ \cdots & & & & & \\ 0 & 0 & 0 & 0 & \cdots & 0 \end{pmatrix}, \text{ for some } 0 < \mu \ll \delta < \lambda.$$

Thus, $\tilde{A} = \begin{pmatrix} \lambda & \mu & 0 & \cdots & 0 \\ \mu & \lambda - \delta & 0 & \cdots & 0 \\ 0 & 0 & \sqrt{\lambda} & \cdots & 0 \\ \cdots & & & & \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix}$. By a routine computation, one can show that first eigenvalue $\tilde{\lambda}_1 = (\lambda - \frac{\delta}{2}) + \frac{\sqrt{\delta^2 + 4\mu^2}}{2}$ with the corresponding eigenvector

$$\tilde{u}_1 = [\tilde{u}_{11} \ \tilde{u}_{12} \ 0 \ 0 \ 0 \dots 0]^T,$$

where

$$\tilde{u}_{11} := \frac{\delta + \sqrt{\delta^2 + 4\mu^2}}{2\sqrt{(\tilde{\lambda}_1 - \lambda + \delta)^2 + \mu^2}} \text{ and } \tilde{u}_{12} := \frac{\mu}{\sqrt{(\tilde{\lambda}_1 - \lambda + \delta)^2 + \mu^2}}.$$

By 238,

$$(240) \quad \|\tilde{u}_1 \tilde{u}_1^T - u_1^T u_1^T\| = |\tilde{u}_{12}| = \frac{\mu}{\sqrt{(\tilde{\lambda}_1 - \lambda + \delta)^2 + \mu^2}} = \Theta\left(\frac{\mu}{\delta}\right) = \Theta\left(\frac{x}{\delta}\right).$$

On the other hand, in this case, it is easy to check that $y = 0$ and $\|E\| = \Theta(\sqrt{\lambda})$. Therefore,

$$\max\left\{\frac{\|E\|}{\lambda}, \frac{y}{\delta}\right\} = \Theta\left(\frac{\|E\|}{\lambda}\right) = \Theta\left(\frac{1}{\sqrt{\lambda}}\right).$$

We choose $\lambda = n^2, \delta = n^{1/2}, \mu = n^{1/4}$. In this setting, the second term $\frac{x}{\delta}$ obviously dominates and presents the true value of $\|\tilde{u}_1 \tilde{u}_1^T - u_1 u_1^T\|$.

14.3. **Necessity of the third term.** Consider

$$E = \begin{pmatrix} 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ \cdots & & & & & & & \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & \cdots & 0 & 0 & 0 \end{pmatrix}.$$

In this case, $x = 0, y = \Theta(\frac{1}{\lambda})$, and $\|E\| = \Theta(1)$. Choose $\lambda = n^{\epsilon_1}, \delta = n^{-\epsilon}$ for some $\epsilon_1 > \epsilon > 0$. Thus,

$$\frac{\|E\|}{\lambda} = \Theta\left(\frac{1}{\lambda}\right), \frac{x}{\delta} = 0, \frac{y}{\delta} = \Theta\left(\frac{1}{\lambda\delta}\right).$$

As $\delta \ll 1$, it is clear that the term $\frac{y}{\delta}$ is the dominating term.

Let $\tilde{\lambda}_1$ be the first eigenvalue of \tilde{A} . Since $\tilde{A} := \begin{pmatrix} \lambda & 0 & 0 & 0 & \cdots & 0 & 0 & 1 \\ 0 & \lambda - \delta & 0 & 0 & \cdots & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ \dots & & & & & & & \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & \cdots & 0 & 0 & 0 \end{pmatrix}$, the corre-

sponding first eigenvector has the form

$$\tilde{u}_1 = [\tilde{u}_{11} \ \tilde{u}_{12} \ 0 \ \cdots \ 0 \ \tilde{u}_{1n}],$$

in which $\tilde{u}_{11}, \tilde{u}_{12}, \tilde{u}_{1n}$ satisfy the following system of equations:

$$(241) \quad \begin{aligned} \lambda \tilde{u}_{11} + \tilde{u}_{1n} &= \tilde{\lambda}_1 \tilde{u}_{11}, \\ (\lambda - \delta) \tilde{u}_{12} + \tilde{u}_{1n} &= \tilde{\lambda}_1 \tilde{u}_{12}, \\ \tilde{u}_{11} + \tilde{u}_{12} &= \tilde{\lambda}_1 \tilde{u}_{1n}, \\ \tilde{u}_{11}^2 + \tilde{u}_{12}^2 + \tilde{u}_{1n}^2 &= 1. \end{aligned}$$

These equations imply that

$$(\tilde{\lambda}_1 - \lambda) \tilde{u}_{11} = \tilde{u}_{1n}, \quad (\tilde{\lambda}_1 - \lambda + \delta) \tilde{u}_{12} = \tilde{u}_{1n},$$

and then,

$$1 = \tilde{u}_{11}^2 + \tilde{u}_{12}^2 + \tilde{u}_{1n}^2 = \tilde{u}_{1n}^2 \left(\frac{1}{(\tilde{\lambda}_1 - \lambda)^2} + \frac{1}{(\tilde{\lambda}_1 - \lambda + \delta)^2} + 1 \right).$$

Thus,

$$(242) \quad \tilde{u}_{1n}^2 = \frac{(\tilde{\lambda}_1 - \lambda)^2}{1 + (\tilde{\lambda}_1 - \lambda)^2 + \frac{(\tilde{\lambda}_1 - \lambda)^2}{(\tilde{\lambda}_1 - \lambda + \delta)^2}} \quad \text{and} \quad \tilde{u}_{12}^2 = \frac{\frac{(\tilde{\lambda}_1 - \lambda)^2}{(\tilde{\lambda}_1 - \lambda + \delta)^2}}{1 + (\tilde{\lambda}_1 - \lambda)^2 + \frac{(\tilde{\lambda}_1 - \lambda)^2}{(\tilde{\lambda}_1 - \lambda + \delta)^2}}.$$

Together (238) and (242) imply that

$$(243) \quad \|\tilde{u}_1 \tilde{u}_1^T - u_1^T u_1^T\|^2 = \frac{(\tilde{\lambda}_1 - \lambda)^2 + \frac{(\tilde{\lambda}_1 - \lambda)^2}{(\tilde{\lambda}_1 - \lambda + \delta)^2}}{1 + (\tilde{\lambda}_1 - \lambda)^2 + \frac{(\tilde{\lambda}_1 - \lambda)^2}{(\tilde{\lambda}_1 - \lambda + \delta)^2}}.$$

On the other hand, since $\tilde{\lambda}_1$ is the largest root of the equation $\det(xI_n - \tilde{A})$, by Schur's formula for determinant, $\tilde{\lambda}_1 > \lambda$ satisfies the following equation:

$$(244) \quad \tilde{\lambda}_1 = \frac{1}{\tilde{\lambda}_1 - \lambda} + \frac{1}{\tilde{\lambda}_1 - \lambda + \delta}.$$

Since $\tilde{\lambda}_1 > \lambda = n^{\epsilon_1}$, Equation (244) requires that $\tilde{\lambda}_1 = \lambda + o(1)$, and that

$$\frac{1}{\tilde{\lambda}_1 - \lambda} < \tilde{\lambda}_1 < \frac{2}{\tilde{\lambda}_1 - \lambda}.$$

These requirements imply that

$$\tilde{\lambda}_1 = \Theta(n^{\epsilon_1}) \quad \text{and} \quad \tilde{\lambda}_1 - \lambda = \Theta(n^{-\epsilon_1}).$$

Therefore, we can simplify (243) as

$$\|\tilde{u}_1 \tilde{u}_1^T - u_1^T u_1^T\|^2 = \Theta \left(\frac{1}{\left(\frac{\tilde{\lambda}_1 - \lambda + \delta}{\lambda_1 - \lambda} \right)^2 + 1} \right) = \Theta \left(\frac{1}{(1 + n^{\epsilon_1 - \epsilon})^2 + 1} \right) = \Theta \left(\frac{1}{(\lambda \delta)^2} \right).$$

It means

$$\|\tilde{u}_1 \tilde{u}_1^T - u_1^T u_1^T\| = \Theta \left(\frac{y}{\delta} \right).$$

We complete our claim.

14.4. Necessity of the assumptions. In this section, we show that our assumptions (such as **C0**, **D0** etc) are also sharp (up to a constant factor).

We focus on **C0**. In this assumption, the term $\frac{\sqrt{p}\|E\|}{\lambda}$ and $\frac{\sqrt{pr}x}{\delta}$ appears in the bound. Thus, even to make the bound meaningful, we need to assume that this two terms are bounded by a small constant.

For the last term, let us consider matrix $A := \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \cdots & & & \\ 0 & 0 & \cdots & 0 \end{pmatrix}$, with $\lambda_1 = c_1 \sqrt{n}$, $\lambda_2 =$

$c_2 \sqrt{n}$ for some positive constants $c_1 > c_2 > 0$.

Let $E := \begin{pmatrix} 0 & 0 \\ 0 & W_{n-1} \end{pmatrix}$, in which W_{n-1} is $(n-1) \times (n-1)$ Wigner matrix.

Results from random matrix theory [12, 50] show that $\tilde{\lambda}_2 \approx \lambda_2 + \frac{\|E\|^2}{4\lambda_2}$ when n tends to infinity; see also Section 4.

If $\frac{\|E\|^2}{4\lambda_2} > \lambda_1 - \lambda_2 = \delta$, then $\tilde{\lambda}_2$ becomes the leading eigenvalue. By our construction, u_1 and \tilde{u}_1 are perpendicular. This means that in order to obtain any meaningful bound on $\sin(u_1, \tilde{u}_1)$, we must assume

$$\frac{\|E\|^2}{4\lambda_2} < \delta,$$

which is equivalent to

$$\frac{\|E\|^2}{\lambda_2 \delta} \leq 4.$$

Acknowledgment. We would like to thank T. Tao and R. Vershynin for valuable remarks. The research is partially supported by Simon Foundation award SFI-MPS-SFM-00006506 and NSF grant AWD 0010308.

REFERENCES

- [1] N. Alon, M. Krivelevich, and B. Sudakov, *Finding a Large Hidden Clique in a Random Graph*, Proceedings of the Ninth Annual ACM-SIAM Symposium on Discrete Algorithm. **13** (1998), 594-598.
- [2] Y. Amit, M. Fink, N. Srebro, and S. Ullman, *Uncovering shared structures in multiclass classification*, Proceedings of the Twenty-fourth International Conference on Machine Learning. (2007), 17-24, <https://dl.acm.org/doi/10.1145/1273496.1273499>.
- [3] A. Argyriou, T. Evgeniou, and M. Pontil, *Multitask feature learning*, Neural Information Processing Systems. **19** (2007).
- [4] Y. Azar, A. Flat, A. Karlin, F. McSherry, and J. Sala, *Spectral analysis of data*, Proceedings of the Thirty-third Annual ACM Symposium on Theory of Computing. (2001), 619-626.
- [5] K. Azuma, *Weighted sums of certain dependent random variables*, Tokohu Mathematical Journal. **19** (1967), 357-367.
- [6] J. Baik, G. Ben Arous, and S. Peche, *Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices*, Ann. Probab. **33** (2005), 1643-1697. MR2165575
- [7] J. Baik and J.W. Silverstein, *Eigenvalues of large sample covariance matrices of spiked population models*, J. Multivariate Anal. **97** (2006), no. 6, 1382-1408. <https://doi.org/10.1016/j.jmva.2005.08.003>.
- [8] Z. Bai and J. Yao, *Central limit theorems for eigenvalues in a spiked population model*, Annales de l'institut Henri Poincaré - Probabilités et Statistiques. **44** (2008), 447-474.
- [9] Z. Bai and J. Yao, *On sample eigenvalues in a generalized spiked population model*, Journal of Multivariate Analysis. **106** (2012), 167-177.
- [10] Z. Bai, J. Yao, and S. Zheng, *Large sample covariance matrices and high-dimensional data analysis*, Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press, 2015.
- [11] F. Benaych-Georges, A. Guionnet, and M. Maida, *Fluctuations of the extreme eigenvalues of finite rank deformations of random matrices*, Electron. J. Probab. **16** (2011), 1621 - 1662. <https://doi.org/10.1214/EJP.v16-929>.
- [12] F. Benaych-Georges and R.R. Nadakuditi, *The eigenvalues and eigenvectors of finite, low rank perturbations of large random matrices*, Adv. Math. **227** (2011), no. 1, 494-521.
- [13] A. Bhardwaj, and V. Vu, *Matrix Perturbation: Davis-Kahan in the Infinity Norm*, Proceedings of the 2024 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA), Society for Industrial and Applied Mathematics. (2024), 880-934.
- [14] R. Bhatia. *Matrix Analysis*, Springer: New York, NY, 2013.
- [15] P. Biswas, T-C Lian, T-C Wang, and Y. Ye, *Semidefinite programming based algorithm for sensor network localization*, ACM Trans. Sen. Netw. **2** (2006), no. 2, 188-220.
- [16] A. Bloemendal, A. Knowles, H-T Yau, and J. Yin, *On the principal components of sample covariance matrices*, Probability Theory and Related Fields. **164** (2016), 459-552.
- [17] A. Blum and J. Spencer, *Coloring random and semi-random k-colorable graphs*, J. Algorithm. **19** (1995), no. 2, 204-234.
- [18] T.T. Cai and A. Zhang, *Rate-optimal perturbation bounds for singular subspaces with applications to high-dimensional statistics*. (2018), 60-89.
- [19] E.J. Candes and B. Recht. *Exact Matrix Completion via Convex Optimization*, Communications of the ACM. **55** (2008), no. 6, 111-119.
- [20] E.J. Candes, J. Romberg, and T. Tao. *Robust uncertainly principles: exact signal reconstruction from highly incomplete frequency information*, IEEE Trans. Inform. Theory. **52** (2006), no. 2, 489-509.
- [21] E.J. Candes and T. Tao. *The power of convex relaxatin: Near-optimal matrix completion*, IEEE Trans. Inform. Theory. **56** (2010), 2053-2080.
- [22] M. Capitaine, C. Donati-Martin, and D. Feral, *The largest eigenvalues of finite rank deformation of large Wigner matrices: convergence and nonuniversality of the fluctuations*, Ann. Prob. **37** (2009), 1-47.
- [23] S. Chatterjee, *Matrix estimation by universal singular value thresholding*, Ann. Stat. **43** (2015): 177-214.
- [24] K. Chaudhuri, A. Sarwate, and K. Sinha, *Near-optimal differentially private principal components*, Advances in neural information processing systems. **25** (2012), 989-997.
- [25] Y. Chen, C. Chen, and J. Fan, *Asymmetry helps: Eigenvalue and Eigenvector analyses of asymmetrically perturbed low-rank matrices*, Ann. Stat. **49** (2021), no. 1, 435.
- [26] Y. Chen, C. Cheng, and Y. Wei, *Tackling small eigen-gaps: Fine-grained eigenvector estimation and inference under heteroscedastic noise*, IEEE Transactions on Information Theory **67** (2021), no. 11, 7380-7419.

- [27] A. Condon and R. M. Karp, *Algorithm for graph partitioning on the planted partition model*, Random Struct. Algorithms. **18** (2001), no.2, 116-140.
- [28] C. Davis and W.M. Kahan, *The rotation of eigenvectors by a perturbation, III*. SIAM J. Numer. Anal. **7** (1970), 1-46.
- [29] S. Deerwester, S. T. Dumais, and R. Harshman, *Indexing by Latent Semantic Analysis*, Journal of the American Society for Information Science. **41** (1990), no. 6, 391-407.
- [30] P. Drineas, A. Frieze, R. Kannan, S. Vempala, and V. Vinay. *Clustering large graphs via the singular value decomposition: theoretical advances in data clustering*, Machine Learning. **56** (2004), 9-33.
- [31] C. Dwork, F. McSherry, K. Nissim, and A. Smith, *Calibrating noise to sensitivity in private data analysis*, Theory of cryptography conference. Springer (2006), 265-284.
- [32] C. Dwork, K. Talwar, A. Thakurta, and L. Zhang. *Analyze Gauss: Optimal bounds for privacy-preserving principal component analysis*, Proceedings of the forty-sixth annual ACM symposium on Theory of computing. (2014), 11-20.
- [33] F.J. Dyson and M. Lal Mehta. *Random matrices and the statistical theory of energy levels IV*, J. math. Phys. **4** (1963), 701-12.
- [34] J. Eldridge, M. Belkin and Y. Wang, *Unperturbed: spectral analysis beyond Davis-Kahan*, Algorithmic learning theory. PMLR (2018), 321-358.
- [35] J. Fan and W. Wang, *Asymptotics of empirical eigenstructure for high dimensional spiked covariance*, Ann. Stat. **45** (2017), no. 3, 1342-1374.
- [36] U. Feige and R. Krauthgamer, *Finding and certifying a large hidden clique in a semirandom graph*, Random Struct. Algorithms. **16** (2000), no. 2, 195-208.
- [37] U. Feige and D. Ron, *Finding hidden cliques in linear time*, Discrete Mathematics and Theoretical Computer Science DMTCS Proceedings, 21st International Meeting on Probabilistic, Combinatorial, and Asymptotic Methods in the Analysis of Algorithms, 2010.
- [38] Z. Füredi and J. Komlós, *The eigenvalues of random symmetric matrices*, Combinatorica, **1**, 233-241, 1981.
- [39] J. Ginibre, *Statistical ensembles of complex, quaternion, and real matrices*, Journal of Mathematical Physics. **6** (1965), no. 3, 440-449.
- [40] V.L. Girko, *Circular law, Theory of Probability & Its Applications*, **29** (1985), no. 4, 694-706.
- [41] D. Goldberg, D. Nichols, B.M. Oki, and D. Terry, *Using collaborative filtering to weave an information tapestry*, Communications of the ACM. **35** (1992), 61-70.
- [42] F. Götze, H. Sambale, and A. Sinulis, *Concentration inequalities for polynomials in α -sub-exponential random variables*. (2021), 1-22. arXiv:1903.05964v1.
- [43] W. Hoeffding, *Probability inequalities for sums of bounded random variable*, Journal of the American Statistical Association. **58** (1963), no. 301, 13-30.
- [44] I.M. Johnstone, *On the distribution of the largest eigenvalue in principal components analysis*, Ann. Statist. **29** (2001), 295-327. MR1863961.
- [45] I.M. Johnstone and A.Y. Lu. *Sparse principal component analysis*. (2009). arXiv: 0901.4392.
- [46] I.M. Johnstone and D. Paul, *PCA in high dimensions: An orientation*, Proceedings of the IEEE. **106** (2018), no. 8, 1277-1292.
- [47] N. El Karoui, *On the largest eigenvalue of Wishart matrices with identity covariance when n, p and $p/n \rightarrow \infty$* . (2003). ArXiv: 0309355v1.
- [48] T. Kato, *Perturbation Theory for Linear Operators*, Classics in Mathematics, Springer: New York, NY, 1980.
- [49] R.H. Keshavan, A. Montanari, and S. Oh, *Matrix completion from a few entries*, IEEE International Symposium on Information Theory. **56** (2010), no. 6, 2980-2998. doi: 10.1109/ISIT.2009.5205567.
- [50] A. Knowles and J. Yin, *The isotropic semicircle law and deformation of Wigner matrices*, Communications on Pure and Applied Mathematics. **66** (2013), no. 11, 1663-1749.
- [51] V. Koltchinskii and D. Xia, *Perturbation of linear forms of singular vectors under Gaussian noise*, High dimensional probability VII, Progr. Probab. **71** (2016), Springer, 397-423.
- [52] O. Mangoubi and N.K. Vishnoi, *Re-analyze Gauss: Bounds for private matrix approximation via Dyson Brownian motion*, Advances in Neural Information Processing System. **35** (2022), 38585-38599.
- [53] O. Mangoubi and N.K. Vishnoi, *Private Covariance Approximation and Eigenvalue-Gap Bounds for Complex Gaussian Perturbations*, The Thirty Sixth Annual Conference on Learning Theory. (2023), 1522-1587.
- [54] JS. Marron and S. Jung, *PCA consistency in high dimension, low sample size context*, Ann. Statist. **37** (2009), no. 6B, 4104-4130.

- [55] JS. Marron, D. Shen, H. Shen, and H. Zhu, *The statistics and mathematics of high dimension low sample size asymptotics*, *Statist. Sinica.* **26** (2016), no.4, 1747-1770.
- [56] F. McSherry. *Spectral partitioning of random graphs*, *Proceedings 42nd IEEE Symposium on Foundations of Computer Science.* (2001), 529-537.
- [57] *Netflix prize*, <http://www.netflixprize.com/>.
- [58] A. Pizzo, D. Renfrew, and A. Soshnikov, *On finite rank deformations of Wigner matrices*, *Annales de l'IHP Probabilités et statistiques.* **49** (2013), no. 1, 64-94.
- [59] S. O'Rourke, V. Vu, and K. Wang, *Random Perturbation of low rank matrices: improving classical bounds*, *Linear Algebra and its Applications.* **540**, 2018, 26-59.
- [60] S. O'Rourke, V. Vu, and K. Wang, *Matrices with Gaussian noise: Optimal estimates for singular subspace perturbation*, *IEEE Transactions on Information Theory.* (2023).
- [61] D. Paul, *Asymptotics of sample eigenstructure for a large dimensional spiked covariance model*, *Statistica Sinica.* **17** (2007), 1617-1642.
- [62] K. Rohe, S. Chatterjee, and B. Yu, *Spectral clustering and the high-dimensional stochastic blockmodel*, *Ann. Statist.* **39** (2011), no. 4, 1878-1915.
- [63] W. Schudy and M. Sviridenko, *Concentration and Moment Inequalities for Polynomials of Independence Random Variables*, *Proceedings of the twenty-third annual ACM-SIAM symposium on Discrete Algorithms.* (2012), 437-446.
- [64] A. Singer, *A remark on global positioning from local distances*, *Proc. Natl. Acad. Sci. USA.* **105** (2008), no. 28, 9507-9511.
- [65] T.A.B. Snijders and K. Nowicki, *Estimation and prediction for stochastic blockmodels for graphs with latent block structure*, *Journal of Classification.* **14** (1997), no. 1, 75-100.
- [66] W.L. Steiger, *Some Kolmogoroff-type inequalities for bounded random variables*, *Biometrika.* **54** (1967), 641-647.
- [67] G.W. Stewart and J. Sun. *Matrix Perturbation Theory.* Academic Press, 1990.
- [68] T. Tao and V. Vu, *Random Matrices: the Distribution of the Smallest Singular Values*, *Geom. Funct. Anal.* **20** (2010), 260–297. <https://doi.org/10.1007/s00039-010-0057-8>.
- [69] T. Tao and V. Vu., *Random matrices: A general approach for the least singular value problem.* (2008). [Arxiv.org/pdf/0805.3167v1](https://arxiv.org/pdf/0805.3167v1).
- [70] C. Tomasi and T. Kanada, *Shape and motion from image streams under orthography: a factorization method*, *International Journal of Computer Vision.* **9** (1992), no. 2, 137-154.
- [71] P. Tran, N.K. Vishnoi, and V. Vu. (2024) [Manuscript in preparation].
- [72] P. Tran, and V. Vu. (2024) [Manuscript in preparation].
- [73] L. Tran and V. Vu. (2024) [Manuscript in preparation].
- [74] J.A. Tropp, *An Introduction to Matrix Concentration Inequalities*, *Foundations and Trends in Machine Learning.* **8** (2015), no. 1-2, 1-230.
- [75] M. Udell and A. Townsend, *Why are big data matrices approximately low rank?* *SIAM Journal on Mathematics of Data Science.* **1.1** (2019), 144-160.
- [76] N. Vaswani and H. Gou, *Correlated-PCA: Principal Component's Analysis when Data and Noise are correlated*, *Advances in Neural Information Processing System.* (2016) pp. 1768-1776.
- [77] N. Vaswani and P. Narayanamurthy, *Finite sample guarantees for PCA in non-isotropic and data-dependent noise.* (2017). [ArXiv: 1709.06255](https://arxiv.org/abs/1709.06255).
- [78] R. Vershynin, *High Dimensional Probability*, Cambridge Univ. Press, 2019.
- [79] V. Vu, *Spectral norm of random matrices*, *Combinatorica*, **27**, 721-736, 2007.
- [80] V. Vu, *Singular vectors under random perturbation*, *Random Structures and Algorithm.* **39** (2011), no. 4, 526-538.
- [81] V. Vu, *A simple SVD Algorithm for finding hidden partitions*, *Combinatorics, Probability and Computing.* **27** (2018), no. 1, 124-140.
- [82] M.J. Wainwright. *High-Dimensional Statistics: A Non-Asymptotic view point*, *Cambridge Series in Statistical and Probabilistic Mathematics*, NY (2019).
- [83] R. Wang, *Singular vector perturbation under Gaussian noise*, *SIAM J. Matrix Anal. Appl.* **36** (2015), no. 1, 158-177.
- [84] P.-A. Wedin, *Perturbation bounds in connection with singular value decomposition*, *Nordisk Tidskr. Informationsbehandling (BIT).* **12** (1972), 99-111.
- [85] Hermann. Weyl, *Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen*, *Mathematische Annalen.* **71** (1912), no. 4, 441-479.

- [86] D. Xia and F. Zhou, *The sup-norm perturbation of HOSVD and low-rank tensor denoising*, J. Mach. Learn. Res. **20** (2019), no. 61, 42.
- [87] A. Y. Zhang, H.Y. Zhou, *Leave-one-out singular subspace perturbation analysis for spectral clustering*. (2022). arXiv:2205.14855.