

S -FP-injective modules

Driss Bennis and Ayoub Bouziri

Abstract

Let R be a commutative ring, and let S be a multiplicative subset of R . In this paper, we introduce and investigate the notion of S -FP-injective modules. Among other results, we show that, under certain conditions, a ring R is S -Noetherian if and only if every S -FP-injective R -module is S -injective. Moreover, we establish, under certain conditions, counterparts of Matlis, Stenström and Cheatham-Stone's characterizations of S -coherent rings.

Mathematics Subject Classification (2020): 13C13, 13E15, 13E99.

Key Words: S -Noetherian rings, S -coherent rings, S -FP-injective modules.

1 Introduction

Throughout this paper, R is a commutative ring with identity, all modules are unitary and S is a multiplicative subset of R ; that is, $1 \in S$ and $s_1 s_2 \in S$ for any $s_1 \in S, s_2 \in S$. Unless explicitly stated otherwise, when we refer to a multiplicative subset S of R , we implicitly assume that $0 \notin S$. This assumption will be used in the sequel without explicit mention. Let M be an R -module. As usual, we use M^+ and M_S to denote, respectively, the character module $\text{Hom}_{\mathbb{Z}}(M, \mathbb{Q}/\mathbb{Z})$ and the localization of M at S . Recall that $M_S \cong M \otimes_R R_S$.

In 1940, Baer initially introduced the notion of injective modules, greatly enriching the study of ring theory. An R -module M is said to be injective if $\text{Ext}_R^1(N, M) = 0$ for any R -module N . In 1967, Maddox [11] extended the notion of injective modules to that of absolutely pure modules. Recall that an R -module M is said to be absolutely pure if it is a pure submodule of every module containing it. A well-known characterization states that an R -module M is absolutely pure if and only if $\text{Ext}_R^1(P, M) = 0$ for any finitely presented R -module P [20, Proposition 2.6]. This characterization led to the terminology of FP-injective modules instead of absolutely pure modules (FP for finitely presented), which was first used by Stenström [20].

Like injective modules, FP-injective modules are an important tool for characterizing some classical rings. Megibben [12] showed that a ring R is semihereditary if and only if any homomorphic image of an FP-injective R -module is FP-injective, and it is Noetherian if and only if any FP-injective R -module is injective. He also proved that a ring R is von Neumann regular if and only if

any R -module is FP-injective [12, Theorem 5]. In 1970, Stenström showed that a ring R is coherent if and only if any direct limit of FP-injective modules is FP-injective [20, Theorem 3.2]. In 1981, Cheatham and Stone, in their work [7, Theorem 1], established characterization for coherent rings using the notion of character modules as bellow:

Theorem 1.1 ([7], Theorem 1) *The following statements are equivalent for a ring R :*

1. R is coherent.
2. M is FP-injective if and only if M^+ is a flat R -module.
3. M is FP-injective if and only if M^{++} is an injective R -module.
4. M is flat if and only if M^{++} is a flat R -module.

In 1982, Matlis [13] characterized coherent rings as those rings for which the duality homomorphisms are isomorphisms for certain modules:

Theorem 1.2 ([13], Theorem 1) *Let R be a commutative ring. The following statements are equivalent:*

1. R is coherent.
2. The induced morphisms $\text{Tor}_R^n(\text{Hom}_R(N, E), M) \rightarrow \text{Hom}_R(\text{Ext}_R^n(M, N), E)$ are isomorphisms for all $n \geq 0$, whenever E is injective and M is finitely presented.
3. $\text{Hom}_R(N, E)$ is a flat R -module for all injective R -modules N and E .

In the last years, the notion of the S -property has drawn the attention of several authors. This notion was introduced in 2002 by D. D. Anderson and Dumitrescu, where they defined the concepts of S -finite modules and S -Noetherian rings [1]. Namely, an R -module M is said to be S -finite module if there exist a finitely generated submodule N of M and $s \in S$ such that $sM \subseteq N$. A commutative ring R is said to be S -Noetherian if every ideal of R is S -finite. To extend coherent rings by multiplicative sets, Bennis and El Hajoui [6] introduced the notions of S -finitely presented modules and S -coherent rings. An R -module M is said to be S -finitely presented if there exists an exact sequence of R -modules $0 \rightarrow K \rightarrow L \rightarrow M \rightarrow 0$, where L is a finitely generated free R -module and K is an S -finite R -module. Moreover, a commutative ring R is called S -coherent, if every finitely generated ideal of R is S -finitely presented. Among other results, they showed that S -coherent rings have a similar characterization to the classical one given by Chase for coherent rings [6, Theorem 3.8].

There are many notions and results of a homological nature that can be generalized from Noetherian (resp., coherent) rings to S -Noetherian rings (resp.,

S -coherent rings). In this context, Qi et al. [18], inspired by the classical case, introduced the notion of S -flat modules and provided a new characterization of S -coherent rings in terms of these modules. An R -module M is said to be S -flat if, for any finitely generated ideal I of R , the natural homomorphism $(I \otimes_R M)_S \rightarrow (R \otimes_R M)_S$ is a monomorphism [18, Definition 2.5]; equivalently, M_S is a flat R_S -module [18, Proposition 2.6]. Notably, any flat R -module is S -flat. A general framework for S -flat modules was developed in the paper [3].

Recently, in [4], the notion of S -injective modules was introduced, and it was shown that many results can be extended from Noetherian rings to S -Noetherian rings. For example, it was demonstrated that, under certain conditions on S , a ring R is S -Noetherian if and only if every direct limit of S -injective R -modules is S -injective, if and only if the class of S -injective modules is closed under direct sums [5, Proposition 2.13 and Corollary 2.17]. Additionally, a counterpart of Cheatham-Stone's characterizations of S -Noetherian rings was presented using the notion of character modules [5, Theorem 2.22].

The primary motivation of this work is to introduce and study an S -version of FP-injective modules, extending some results on coherent and Noetherian rings to S -coherent rings and S -Noetherian rings, respectively.

The organization of the paper is as follows: In Section 2, we introduce the notion of S -FP-injective modules, which serves as a weak version of S -injective modules and an S -version of FP-injective modules (see Definition 2.1). In Theorem 2.6, we provide some characterizations of S -FP-injective modules. We then proceed to explore some properties of these modules. For example, we demonstrate that the class of S -FP-injective modules is closed under direct summands, direct products, and, under certain conditions, finite direct sums. Furthermore, we establish that, under certain conditions, the class of all S -injective modules coincides with the class of S -FP-injective modules if and only if R is S -Noetherian (see Proposition 2.15). This extends the well-known result that a ring is Noetherian if and only if every FP-injective module is injective [12, Theorem 3]. In Section 3, we provide new characterizations for S -coherent rings in the case where R_S is finitely presented as an R -module. Firstly, we introduce an S -version of Matlis' Theorem in Theorem 3.3. Then, we present an S -version of Stenström's characterization of coherent rings in Theorem 3.4. Finally, we conclude this section with a counterpart of Cheatham and Stone's characterizations for S -coherent rings in Theorem 3.6.

2 S -FP-injective modules

Recall that an exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ is pure if and only if, for every finitely presented R -module F , the induced sequence $0 \rightarrow \text{Hom}_R(F, L) \rightarrow \text{Hom}_R(F, M) \rightarrow \text{Hom}_R(F, N) \rightarrow 0$ is exact. An R -module M is FP-injective if and only if every monomorphism $M \rightarrow N$ is pure; that is, the induced exact sequence $0 \rightarrow M \rightarrow N \rightarrow N/M \rightarrow 0$ is pure. Now, we introduce S -versions of these notions:

Definition 2.1 1. An exact sequence of R -modules $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is said to be S -pure if the induced sequence $0 \rightarrow \text{Hom}_R(P_S, A) \rightarrow \text{Hom}_R(P_S, B) \rightarrow \text{Hom}_R(P_S, C) \rightarrow 0$ is exact for any finitely presented R -module P .

2. A monomorphism $f : M \rightarrow N$ is said to be S -pure if the induced exact sequence $0 \rightarrow M \rightarrow N \rightarrow \text{coker}(f) \rightarrow 0$ is S -pure.

3. A submodule M of an R -module N is said to be an S -pure submodule if the inclusion $M \rightarrow N$ is an S -pure monomorphism.

4. An R -module M is said to be S -FP-injective (or absolutely S -pure) if every monomorphism $M \rightarrow N$ is S -pure.

Recall from [17] that an R -module C is said to be S -weakly cotorsion if $\text{Ext}_R^1(R_S, C) = 0$. An R -module F is said to be S -strongly flat if $\text{Ext}_R^1(F, C) = 0$ for every S -weakly cotorsion R -module C . We will see that any S -FP-injective R is S -weakly cotorsion. However, the converse does not hold in general (see the discussion just after Theorem 2.6).

According to [5, Definition 2.1], an R -module E is said to be S -injective if, whenever $i : A \rightarrow B$ is a monomorphism and $h : A_S \rightarrow E$ is any morphism of R -modules, there exists a morphism of R -modules g making the following diagram commutes:

$$\begin{array}{ccc}
 & E & \\
 & \uparrow h & \swarrow g \\
 0 & \longrightarrow A_S & \xrightarrow{i_S} B_S
 \end{array}$$

Next, in Corollary 2.9, we will see that any S -weakly cotorsion S -injective R -module is S -FP-injective. However, an S -FP-injective R -module need not be S -injective (see Example 2.16). In Proposition 2.15, we will show that if R is S -Noetherian, then any S -FP-injective module is S -injective. Moreover, the converse holds (at least) when R_S is finitely presented and the S -torsion in R is bounded; that is, there exists an element $s_0 \in S$ such that $sr = 0$ for $s \in S$ and $r \in R$ implies $s_0r = 0$. This definition can be found in [17].

Remarks 2.2 1. Let P be an S -finitely presented R -module. Then P_S is a finitely presented R_S -module by [6, Remark 3.4 (3)]. By [14, Proposition 3.19], there exists a finitely presented R -module P' such that $P_S \cong P'_S$. Thus, an exact sequence $0 \rightarrow K \rightarrow L \rightarrow M \rightarrow 0$ is S -pure if and only if, for every S -finitely presented R -module P , the induced sequence $0 \rightarrow \text{Hom}_R(P_S, K) \rightarrow \text{Hom}_R(P_S, L) \rightarrow \text{Hom}_R(P_S, M) \rightarrow 0$ is exact.

2. Let \mathcal{X} be a class of modules. Recall from [22, Section 1] that an exact sequence $0 \rightarrow K \rightarrow L \rightarrow M \rightarrow 0$ of R -modules is called \mathcal{X} -pure exact if, for any $X \in \mathcal{X}$, the induced morphism $\text{Hom}_R(X, L) \rightarrow \text{Hom}_R(X, M)$ is surjective. Thus, an exact sequence $0 \rightarrow K \rightarrow L \rightarrow M \rightarrow 0$ is S -pure if and only if it is FP_S -pure exact, where $\text{FP}_S = \{P_S | P \text{ is finitely presented}\}$.

3. The $U(R)$ -pure exact sequences (resp., monomorphisms, submodules) are exactly the pure exact sequences (resp., monomorphisms, submodules), where $U(R)$ denotes the set of units in R .
4. The $U(R)$ -FP-injective modules are exactly the FP-injective modules, where $U(R)$ denotes the set of units in R .

Proposition 2.3 1. Let $0 \rightarrow K \rightarrow L \rightarrow M \rightarrow 0$ be an exact sequence of R -modules. Then $0 \rightarrow K \rightarrow L \rightarrow M \rightarrow 0$ is S -pure if and only if the induced sequence $0 \rightarrow \text{Hom}_R(R_S, K) \rightarrow \text{Hom}_R(R_S, L) \rightarrow \text{Hom}_R(R_S, M) \rightarrow 0$ is a pure exact sequence.

2. A monomorphism $K \rightarrow L$ is S -pure if and only if $0 \rightarrow \text{Hom}_R(R_S, N) \rightarrow \text{Hom}_R(R_S, M)$ is a pure monomorphism.

Proof. 1. This follows immediately from the following natural isomorphism:

$$\text{Hom}_R(A, \text{Hom}_R(B, C)) \cong \text{Hom}_R(A \otimes_R B, C),$$

for any R -modules A , B , and C [19, Theorem 2.75].

2. This follows from (1). ■

The following lemma will be needed:

Lemma 2.4 Let M be an S -weakly cotorsion R -module, then

$$\text{Ext}_R^1(N_S, M) \cong \text{Ext}_R^1(N, \text{Hom}_R(R_S, M)).$$

for any R -module N .

Proof. Let $0 \rightarrow M \rightarrow E \rightarrow E/M \rightarrow 0$ be an exact sequence with E is an injective R -module. Since M is S -weakly cotorsion, the induced sequence

$$0 \rightarrow \text{Hom}_R(R_S, M) \rightarrow \text{Hom}_R(R_S, E) \rightarrow \text{Hom}_R(R_S, E/M) \rightarrow 0$$

is exact. Consider the following commutative diagram:

$$\begin{array}{ccccccc} \text{Hom}_R(N_S, E) & \longrightarrow & \text{Hom}_R(N_S, E/M) & \longrightarrow & C & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ \text{Hom}_R(N, \text{Hom}_R(R_S, E)) & \longrightarrow & \text{Hom}_R(N, \text{Hom}_R(R_S, E/M)) & \longrightarrow & D & \longrightarrow & 0 \end{array}$$

where $C = \text{Ext}_R^1(N_S, M)$ and $D = \text{Ext}_R^1(N, \text{Hom}_R(R_S, M))$. The upper row is exact because E is injective. The lower row is exact because $\text{Hom}_R(R_S, E)$ is injective. Since, by [19, Theorem 2.75], the first two vertical arrows are isomorphisms, $\text{Ext}_R^1(N_S, M) \cong \text{Ext}_R^1(N, \text{Hom}_R(R_S, M))$. ■

Recall that an R -module M is S -injective if and only if $\text{Hom}_R(R_S, M)$ is injective [5, Proposition 2.7]. Therefore, the next result follows immediately from Lemma 2.4.

Corollary 2.5 *The following are equivalent for an S -weakly cotorsion R -module M :*

1. M is S -injective,
2. $\text{Ext}_R^1(N_S, M) = 0$ for any R -module N ,
3. $\text{Ext}_R^1(N, M) = 0$ for any R_S -module N .

An S -weakly cotorsion R -module M need not be S -injective (see Corollary 2.7 and Example 2.16). Recall that an R -module C is said to be S - h -reduced if $\text{Hom}_R(R_S, C) = 0$. Thus, every S - h -reduced module is S -injective, but an S - h -reduced module is not necessarily S -weakly cotorsion. In fact, the S - h -reduced S -weakly cotorsion modules are the so-called S -contramodule R -modules, which are extensively studied in papers [15, 16]. Whether the assumption in Corollary 2.5 can be removed remains an interesting question.

Recall that an R -module M is FP-injective if and only if it is a pure submodule of an injective R -module. In the next result, we provide a corresponding result for S -FP-injective modules in terms of S -pureness and S -injective modules.

Theorem 2.6 *The following assertions are equivalent for an R -module M :*

1. M is S -FP-injective,
2. M is injective with respect to the exact sequences $0 \rightarrow A \rightarrow B \rightarrow C_S \rightarrow 0$ with C is S -finitely presented,
3. M is injective with respect to the exact sequences $0 \rightarrow A \rightarrow B \rightarrow C_S \rightarrow 0$ with C is finitely presented,
4. $\text{Ext}_R^1(C_S, M) = 0$ for every S -finitely presented R -module C ,
5. $\text{Ext}_R^1(C_S, M) = 0$ for every finitely presented R -module C ,
6. M is an S -pure submodule of an injective module,
7. M is an S -pure submodule of an S -weakly cotorsion S -injective module.

Proof. The equivalences 2. \Leftrightarrow 3. and 4. \Leftrightarrow 5. follow from the fact that every finitely presented R_S -module come from a finitely presented R -module [14, Proposition 3.19], and the localization P_S of any finitely presented (resp., S -finitely presented) R -module P is a finitely presented R_S -module.

1. \Rightarrow 4. Consider an exact sequence $\mathcal{E} = 0 \rightarrow M \rightarrow N \xrightarrow{g} C_S \rightarrow 0$, where C is an S -finitely presented R -module. According to (1) and Remarks 2.2, there exists $h : C_S \rightarrow N$ such that $g \circ h = 1_{C_S}$. Therefore, \mathcal{E} splits. Hence, $\text{Ext}_R^1(C_S, M) = 0$ by [19, Theorem 7.31].

2. \Rightarrow 5. Similar to 1. \Rightarrow 4.

4. \Rightarrow 1. and 5. \Rightarrow 2. follow immediately by using the long exact sequences.

1. \Rightarrow 6. This follows from the fact that every R -module M can be embedded as a submodule of an injective R -module.

6. \Rightarrow 7. This follows from the fact that every injective R -module is both S -weakly cotorsion and S -injective .

7. \Rightarrow 5. If $\mathcal{E} = 0 \rightarrow M \rightarrow E \xrightarrow{g} M' \rightarrow 0$ is an S -pure monomorphism with E being both S -weakly cotorsion and S -injective, then, by Corollary 2.5, for any finitely presented R -module C , the induced sequence

$$\mathrm{Hom}_R(C_S, E) \xrightarrow{g^*} \mathrm{Hom}_R(C_S, M') \rightarrow \mathrm{Ext}_R^1(C_S, M) \rightarrow 0$$

is exact. Since \mathcal{E} is S -pure, g^* is an epimorphism. Consequently, $\mathrm{Ext}_R^1(C_S, M) = 0$, for any finitely presented R -module C . \blacksquare

It follows from Theorem 2.6(5) that every S -FP-injective R -module is S -weakly cotorsion. However, an S -weakly cotorsion R -module need not be S -FP-injective. Indeed, let S be a multiplicative subset of a Noetherian ring R such that R_S is not a semisimple ring. If every S -weakly cotorsion R -module is S -FP-injective, then for any ideal I of R , we have

$$\mathrm{Ext}_R^1(R_S/I_S, C) = 0$$

for any S -weakly cotorsion R -module C . Hence, R_S/I_S is an S -strongly flat R -module for any ideal I of R . Hence, by [2, Lemma 3.1], R_S/I_S is a projective R_S -module for any ideal I of R . Then, R_S is semisimple, which contradicts our assumption. Thus, there exists an S -weakly cotorsion R -module that is not S -FP-injective.

Corollary 2.7 *The following assertion are equivalent for an R -module M :*

1. M is S -FP-injective,
2. M is S -weakly cotorsion and, $\mathrm{Hom}_R(R_S, M)$ is an FP-injective R -module.

Proof. This follows from Lemma 2.4 and Theorem 2.6 (5). \blacksquare

The following lemma will be needed:

Lemma 2.8 *If R_S is finitely presented as an R -module, then for every S -finitely presented R -module N , N_S is finitely presented as an R -module.*

Proof. Let $0 \rightarrow K \rightarrow F \rightarrow N \rightarrow 0$ be an exact sequence such that F is a finitely generated free R -module and K is S -finite; that is, there exists a finitely generated submodule K' of K such that $sK \subseteq K'$ for some $s \in S$. Notice that $K'_S = K_S$. Since R_S is a flat R -module, the induced sequence $0 \rightarrow K_S = K'_S \rightarrow F_S \rightarrow N_S \rightarrow 0$ is exact. Since R_S is finitely generated as R -module and $K_S = K'_S$ is finitely generated as an R_S -modules, K_S is finitely generated as an R -module. Since R_S is finitely presented, then F_S , being a direct sum of finitely many copies of R_S , is finitely presented as an R -module. Therefore, N_S is finitely presented as an R -module by [10, Theorem 2.1.2]. \blacksquare

Corollary 2.9 1. Assume that R_S is finitely presented as R -module, then every FP-injective R -module is S -FP-injective.

2. Let M be an S -weakly cotorsion R -module. If M is S -injective, then it is S -FP-injective.

Proof. 1. Let M be an FP-injective R -module and N a finitely presented R -module. Then, N_S is a finitely presented R_S -module. Since R_S is finitely presented, by Lemma 2.8, N_S is finitely presented as an R -module. Then, $\text{Ext}_R^1(N_S, M) = 0$. Therefore, M is S -FP-injective by Theorem 2.6.

2. This is clear by Theorem 2.6 (5) and Corollary 2.5. ■

In [8, Proposition 1.2], Couchot proved that, if R is a coherent ring, then each localization of any FP-injective R -module is an FP-injective R_S -module. Here, we prove that the localization M_S of any S -FP-injective R -module M is FP-injective as an R_S -module.

Proposition 2.10 Let M be an R -module. If M is S -FP-injective, then M_S is an FP-injective R_S -module.

Proof. Consider a monomorphism $M_S \rightarrow E$ where E is an injective R_S -module. Let P be a finitely presented R_S -module. By [14, Proposition 3.19], there exists a finitely presented R -module P' such that $P'_S \cong P$. Then, by using the fact that $\text{Hom}_R(A, B) = \text{Hom}_{R_S}(A, B)$ whenever A and B are R_S -modules, one can see that

$$\text{Hom}_{R_S}(P, E) \rightarrow \text{Hom}_{R_S}(P, E/M_S) \rightarrow 0$$

is an exact sequence. Thus, M_S is an FP-injective R_S -module. ■

It follows from Proposition 2.10 that, for a multiplicative subset S of R , if every FP-injective R -module is S -FP-injective, then the localization at S of any FP-injective R -module is an FP-injective R_S -module. Hence, by Corollary 2.9, if R_S is finitely presented as an R -module, then the localization at S of any FP-injective R -module is an FP-injective R_S -module. Moreover, in Proposition 2.11, we prove that these concepts are equivalent for R_S -modules.

Proposition 2.11 The following assertions are equivalent for an R_S -module M :

1. M is S -FP-injective as an R -module.
2. M is FP-injective as an R_S -module.
3. M is FP-injective as an R -module.

Proof. 1. \Rightarrow 2. By [19, Corollary 4.79], M is naturally isomorphic to its localization M_S as R_S -modules. Thus, M is FP-injective as an R_S -module by Proposition 2.10.

2. \Rightarrow 3. This is [14, Theorem 3.20].

3. \Rightarrow 1. Let $M \rightarrow E$ be a monomorphism where E is an injective R_S -module. Then, by (3), for every finitely presented R -module P , the morphism $\text{Hom}_R(P, E) \rightarrow \text{Hom}_R(P, E/M)$ is an epimorphism. The induced morphism

$$\text{Hom}_R(P, E)_S \rightarrow \text{Hom}_R(P, E/M)_S$$

is also an epimorphism. By [19, Lemma 4.87 and Corollary 4.79 (ii)], $\text{Hom}_R(P, M)_S \cong \text{Hom}_{R_S}(P_S, M)$. Moreover, $\text{Hom}_R(A, B) = \text{Hom}_{R_S}(A, B)$ whenever A and B are R_S -modules. Thus, $\text{Hom}_R(P_S, E) \rightarrow \text{Hom}_R(P_S, E/M)$ is an epimorphism, and therefore, M is S -FP-injective by Theorem 2.6 (6). \blacksquare

Megibben [12] gave an equivalent way to define FP-injective modules: An R -module M is FP-injective if and only if every diagram

$$\begin{array}{ccc} & M & \\ & \uparrow & \swarrow \text{---} \\ 0 & \longrightarrow P' & \longrightarrow P \end{array}$$

with P' finitely generated and P projective can be completed to a commutative diagram. Here, we extend this result to S -FP-injective modules.

Theorem 2.12 *The following statements are equivalent for an R -module M :*

1. M is S -FP-injective.
2. M is S -weakly cotorsion, and every diagram

$$\begin{array}{ccc} & M & \\ & \uparrow & \swarrow \text{---} \\ 0 & \longrightarrow P'_S & \longrightarrow P_S \end{array}$$

where P is a finitely generated projective R -module and P' is an S -finitely generated submodule of P , can be completed to a commutative diagram.

3. M is S -weakly cotorsion, and every diagram

$$\begin{array}{ccc} & M & \\ & \uparrow & \swarrow \text{---} \\ 0 & \longrightarrow P'_S & \longrightarrow P_S \end{array}$$

where P is a finitely generated projective (free) R -module and P' is a finitely generated submodule of P , can be completed to a commutative diagram.

Proof. 1. \Rightarrow 2. Assume that M is S -FP-injective. The induced exact sequence:

$$0 \rightarrow P'_S \rightarrow P_S \rightarrow (P'/P)_S \rightarrow 0.$$

gives rise to the exact sequence:

$$\text{Hom}(P_S, M) \rightarrow \text{Hom}(P'_S, M) \rightarrow \text{Ext}_R^1((P'/P)_S, M).$$

Since M is S -FP-injective and P/P' is S -finitely presented, $\text{Ext}_R^1((P/P)_S, M) = 0$. Thus the sequence

$$\text{Hom}(P_S, M) \rightarrow \text{Hom}(P'_S, M) \rightarrow 0$$

is exact, and therefore the appropriate diagram can be completed.

2. \Rightarrow 3. Trivial.

3. \Rightarrow 1. We show that $\text{Hom}_R(R_S, M)$ is FP-injective. By Corollary 2.7, it follows that M is S -FP-injective. Let P' be a finitely generated submodule of a finitely generated free R -module F . Then we have the exact sequence $0 \rightarrow P' \rightarrow F \rightarrow F/P' \rightarrow 0$ which gives rise to the commutative diagram with exact rows:

$$\begin{array}{ccccc} \text{Hom}_R(F_S, M) & \longrightarrow & \text{Hom}_R(P'_S, M) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \\ \text{Hom}_R(F, \text{Hom}_R(R_S, M)) & \longrightarrow & \text{Hom}_R(P', \text{Hom}_R(R_S, M)) & & \end{array}$$

The upper row is exact by assumption. Moreover, all the vertical maps are isomorphisms. Hence, $\text{Hom}_R(F, \text{Hom}_R(R_S, M)) \rightarrow \text{Hom}_R(P', \text{Hom}_R(R_S, M))$ is an epimorphism. Therefore, $\text{Hom}_R(R_S, M)$ is FP-injective as desired. \blacksquare

According to [12, Theorem 4 and Remark on page 565], if R is a coherent ring, then an R -module M is FP-injective if and only if $\text{Ext}_R^1(R/I, M) = 0$ for any finitely generated ideal I of R . Replacing "coherent" with " S -coherent," we prove the following theorem, which will be useful in the next Section.

Recall from [6, Definition 3.3] that a ring R is called S -coherent, if every finitely generated ideal of R is S -finitely presented.

Theorem 2.13 *Suppose that R is S -coherent. Then the following assertions are equivalent for an R -module M :*

1. M is S -FP-injective.
2. M is S -weakly cotorsion, and $\text{Ext}_R^1(R_S/I_S, M) = 0$ for all S -finitely generated ideals I of R .
3. M is S -weakly cotorsion, and $\text{Ext}_R^1(R_S/I_S, M) = 0$ for all finitely generated ideals I of R .

Proof. 1. \Rightarrow 2. and 2. \Rightarrow 3. are straightforward.

3. \Rightarrow 1. Assume that $\text{Ext}_R^1(R_S/I_S, M) = 0$ for all S -finitely generated ideals I . Consider an arbitrary homomorphism $f : N_S \rightarrow M$, where N is a finitely

generated submodule of a finitely generated free R -module F . We need to show that f can be extended to F_S . By Theorem 2.13, it follows that M is S -FP-injective. We prove this by induction on the number n of generators of F . The case $n = 1$ holds by assumption. For $n \geq 2$, we write $F = Rx \oplus V$, where V is free with $(n - 1)$ generators.

Let $I = \{r \in R/rx \in N + V\}$. The morphism $rx + g \in N \rightarrow r \in I$, induces an isomorphism $I \cong N/(N \cap V)$. Consequently, I is finitely generated. Since R is S -coherent, I is S -finitely presented. Hence, by [6, Theorem 2.4], $(N \cap V)$ is S -finite; that is, there exists a finitely generated submodule K of $N \cap V$ such that $s(N \cap V) \subseteq K$ for some $s \in S$. In particular, $(N \cap V)_S = K_S$. Thus, we can assume that $N \cap V$ is finitely generated.

By induction, there exists a homomorphism $g : V_S \rightarrow M$ such that $g|_{(N \cap V)_S} = f|_{(N \cap V)_S}$. Hence, there is a unique homomorphism $h : (N + V)_S \rightarrow M$ extending both g and f . Define $\theta : I_S \rightarrow M$ by $\theta(\frac{r}{s}) = h(\frac{rx}{s})$ for all $r \in I$. Since $\text{Ext}_R^1(R_S/I_S, M) = 0$, there is a homomorphism $\psi : R_S \rightarrow M$ that extends θ . The mapping $k : F_S \rightarrow M$ defined by $k(\frac{rx+v}{s}) = \psi(\frac{r}{s}) + g(\frac{v}{s})$ for all $r \in R$, $s \in S$, and $v \in V$ is a homomorphism extending f . ■

There are questions that arise naturally about classes of modules. Two of these are whether the direct sum of elements of the class remains in the class and whether the direct limit of elements in the class also remains in the class. The direct sum of FP-injective modules is always FP-injective, but it is not easy to see whether this holds for S -FP-injective modules. Here, we examine specific cases, but we leave the general case as an open question.

Proposition 2.14 *Let $(M_i)_{i \in I}$ be a family of R -modules. Then*

1. $\prod_{i \in I} M_i$ is S -FP-injective if and only if each M_i is S -FP-injective. In particular, every direct summand of an S -FP-injective module is S -FP-injective.
2. Assume that R_S is finitely presented as an R -module. Then $\bigoplus_{i \in I} M_i$ is S -FP-injective if and only if each M_i is S -FP-injective.

Proof. 1. By Theorem 2.6, $\prod_{i \in I} M_i$ is S -FP-injective if and only if, for any finitely presented R -module P , $\text{Ext}_R^1(P_S, \prod_{i \in I} M_i) = 0$. By [19, Proposition 7.22], we have $\text{Ext}_R^1(P_S, \prod_{i \in I} M_i) \cong \prod_{i \in I} \text{Ext}_R^1(P_S, M_i)$. Thus, $\text{Ext}_R^1(P_S, \prod_{i \in I} M_i) = 0$ if and only if, for each $i \in I$, $\text{Ext}_R^1(P_S, M_i) = 0$. Again, by Theorem 2.6, this holds if and only if, for each $i \in I$, M_i is S -FP-injective.

2. Since R_S is flat, it is projective [21, Theorem 2.6.15]. Hence, every R -module is S -weakly cotorsion. Thus, by Corollary 2.7, $\bigoplus_{i \in I} M_i$ is S -FP-injective if and only if $\text{Hom}_R(R_S, \bigoplus_{i \in I} M_i)$ is FP-injective. By [21, Theorem 2.6.10], we have

$$\text{Hom}_R(R_S, \bigoplus_{i \in I} M_i) \cong \bigoplus_{i \in I} \text{Hom}_R(R_S, M_i).$$

Since the class of FP-injective modules is closed under direct sums, $\bigoplus_{i \in I} \text{Hom}_R(R_S, M_i)$ is FP-injective if and only if, for each $i \in I$, $\text{Hom}_R(R_S, M_i)$ is FP-injective. Again, by Corollary 2.7, this happens if and only if, for each $i \in I$, M_i is S -FP-injective. ■

Megibben [12, Theorem 3] showed that a ring R is Noetherian if and only if every FP-injective R -module is injective. Here, we extend this result in some particular cases. The general case is left as an open question.

Proposition 2.15 *If R is S -Noetherian, then every S -FP-injective module is S -injective. Moreover, if the S -torsion in R is bounded and R_S is finitely presented as an R -module, then the converse holds true.*

Proof. Assume that R is S -Noetherian. Let M be an S -FP-injective R -module. Let I be an ideal of R . Since R is S -Noetherian, I is S -finite. Hence, $\text{Ext}_R^1(R_S/I_S, M) = 0$. Therefore, M is S -injective by [5, Proposition 2.4].

Conversely, suppose that every S -FP-injective module is S -injective. If R is not S -Noetherian, then by [5, Corollary 2.16], there exists a family $(M_i)_{i \in I}$ of S -injective modules such that $\bigoplus_{i \in I} M_i$ is not S -injective. However, by Proposition 2.14, $\bigoplus_{i \in I} M_i$ is S -FP-injective, a contradiction. Thus, R is S -Noetherian. ■

Now, utilizing Proposition 2.15, we can demonstrate that the classes of S -FP-injective modules and S -injective modules are not coincident in general.

Example 2.16 *Let R' be a commutative ring and S' be a multiplicative subset of R' such that the S' -torsion in R' is bounded, and $R'_{S'}$ is finitely presented (for an example of such a ring, please see [5, Example 2.23]). Let R'' be a non-Noetherian commutative ring. Consider the ring $R = R' \times R''$ with the multiplicative subset $S = S' \times U(R'')$, where $U(R'')$ denotes the set of units in R'' . Then, there exists an S -FP-injective R -module which is not S -injective.*

Proof. Firstly, notice that the S -torsion in R is bounded, and $R_S = (R')_{S'} \times R''$ is a finitely presented R -module. Moreover, since R'' is not $(U(R''))$ -Noetherian, R is not S -Noetherian. Hence, according to Proposition 2.15, there exists an S -FP-injective module which is not S -injective. ■

3 S -coherent rings

In this section, we provide some characterizations of S -coherence. Specifically, considering particular multiplicative subsets S , we present counterparts of Matlis, Stenström, and Cheatham-Stone's characterizations of S -coherent rings.

Recall that an R -module M is said to be S -flat if for any finitely generated ideal I of R , the natural homomorphism $I \otimes_R M \rightarrow R \otimes_R M$ is an S -monomorphism; equivalently, M_S is a flat R_S -module [18, Proposition 2.6].

We need the following lemmas:

Lemma 3.1 *Let $(F_i)_{i \in I}$ be a family of S -flat modules. For each $i \in I$, let K_i be a submodule of F_i such that F_i/K_i is S -flat. Suppose that R_S is finitely presented. Then, the following assertions are equivalent:*

1. $\prod_{i \in I} F_i$ is S -flat.
2. $\prod_{i \in I} K_i$ and $\prod_{i \in I} F_i/K_i$ are S -flat.

Proof. 2. \Rightarrow 1. Let N be a finitely presented R -module. Consider the induced exact sequences:

$$\mathrm{Tor}_R^1\left(\prod_{i \in I} K_i, N_S\right) \rightarrow \mathrm{Tor}_R^1\left(\prod_{i \in I} F_i, N_S\right) \rightarrow \mathrm{Tor}_R^1\left(\prod_{i \in I} F_i/K_i, N_S\right)$$

By [4, Proposition 2.5], the first and last terms are zero, which implies that $\mathrm{Tor}_R^1\left(\prod_{i \in I} F_i, N_S\right) = 0$. Hence, $\prod_{i \in I} F_i$ is S -flat, again by [4, Proposition 2.5].

1. \Rightarrow 2. Let N be a finitely presented R -module. Consider the following commutative diagram:

$$\begin{array}{ccccc} 0 \rightarrow & \mathrm{Tor}_R^1\left(\prod_{i \in I} F_i/K_i, N_S\right) & \rightarrow & \left(\prod_{i \in I} K_i\right) \otimes_R N_S & \rightarrow & \left(\prod_{i \in I} F_i\right) \otimes_R N_S \\ & \downarrow & & \downarrow \cong & & \downarrow \cong \\ & \prod_{i \in I} \mathrm{Tor}_R^1(F_i/K_i, N_S) & \rightarrow & \prod_{i \in I} (K_i \otimes_R N_S) & \rightarrow & \prod_{i \in I} (F_i \otimes_R N_S) \end{array}$$

Since R_S is finitely presented, N_S is a finitely presented R -module by Lemma 2.8. Hence, by [9, Theorem 3.2.22], the right two vertical homomorphisms are isomorphisms. Since, for each $i \in I$, F_i/K_i is S -flat, we have $\mathrm{Tor}_R^1(F_i/K_i, N_S) = 0$. Hence, $\prod_{i \in I} \mathrm{Tor}_R^1(F_i/K_i, N_S) = 0$, and so $\mathrm{Tor}_R^1\left(\prod_{i \in I} F_i/K_i, N_S\right) = 0$. Therefore, $\prod_{i \in I} F_i/K_i$ is S -flat by [4, Proposition 2.5].

Now, using [4, Proposition 2.5] and the induced exact sequence

$$\mathrm{Tor}_R^2\left(\prod_{i \in I} F_i/K_i, N_S\right) \rightarrow \mathrm{Tor}_R^1\left(\prod_{i \in I} K_i, N_S\right) \rightarrow \mathrm{Tor}_R^1\left(\prod_{i \in I} F_i, N_S\right)$$

we deduce that $\prod_{i \in I} K_i$ is S -flat too. \blacksquare

Recall that if E is an injective cogenerator R -module (universal injective in the terminology of [12]), then $R \subseteq \mathrm{Hom}_R(E, E)$ and $\mathrm{Hom}_R(E, E)$ is flat if and only if $\mathrm{Hom}_R(E, E)/R$ is flat. To provide an S -version of Matlis's characterization for coherent rings, we need an S -version of this result:

Lemma 3.2 *Let E be an injective cogenerator R -module and let $H = \mathrm{Hom}_R(E, E)$ then, H is S -flat if and only if H/R is S -flat.*

Proof. The "if" part follows from the fact that the class of S -flat modules is closed under extension [4, Lemma 2.1]. For the "only if" part, let I_S be an ideal of R_S , where I is an ideal of R . By [10, Lemma 2.6.5], $HI \cap R = I$. Hence, $(HI \cap R)_S = H_S I_S \cap R_S = I_S$. Then, by [10, Theorem 1.2.3], H_S is a flat R_S -module. ■

Theorem 3.3 *Let R be a commutative ring and $S \subseteq R$ be a multiplicative subset such that R_S is finitely presented as an R -module. Then the following assertions are equivalent:*

1. R is S -coherent.
2. R_S is a coherent ring.
3. The induced morphisms $\text{Tor}_R^n(\text{Hom}_R(N, E), M_S) \rightarrow \text{Hom}_R(\text{Ext}_R^n(M_S, N), E)$ are isomorphisms for all $n \geq 0$ whenever E is injective and M is S -finitely presented.
4. The induced morphisms $\text{Tor}_R^n(\text{Hom}_R(N, E), M_S) \rightarrow \text{Hom}_R(\text{Ext}_R^n(M_S, N), E)$ are isomorphisms for all $n \geq 0$ whenever E is injective and M is finitely presented.
5. $\text{Hom}_R(N, E)$ is an S -flat R -module for all S -FP-injective R -modules N and any injective R -module E .
6. $\text{Hom}_R(N, E)$ is an S -flat R -module for all S -injective R -modules N and any injective module E .
7. $\text{Hom}_R(N, E)$ is an S -flat R -module for all injective R -modules N and any injective module E .

Proof. 1. \Rightarrow 2. This is [6, Remarks 3.4(3)].

2. \Rightarrow 3. Let M be an S -finitely presented R -module. By [6, Remark 3.4], M_S is a finitely presented R_S -module. Since R_S is a coherent ring, M_S has a projective resolution composed of finitely generated R_S -modules [10, Corollary 2.5.2].

On the other hand, as R_S is a finitely generated projective R -module [19, Theorem 3.56], every finitely generated projective R_S -module is also a finitely generated projective R -module. Therefore, M_S has a projective resolution composed of finitely generated R -modules. Consequently, (3) follows from [10, Theorem 1.1.8].

3. \Rightarrow 4. and 6. \Rightarrow 7. are obvious.

4. \Rightarrow 5. This follows from Theorem 2.6 and [4, Proposition 2.5].

5. \Rightarrow 6. This follows from Corollary 2.9.

7. \Rightarrow 1. Let E be an injective cogenerator. Then, $H := \text{Hom}_R(E, E)$ is S -flat by (7). Hence, by Lemma 3.2, H/R is S -flat.

Let I be an index set. Denote, for each $i \in I$, $R_i = R$, $H_i = H$, and $E_i = E$. We have, by (4), $\prod_{i \in I} H_i = \text{Hom}_R(E, \prod_{i \in I} E_i)$ is S -flat because $\prod_{i \in I} E_i$ is

injective. Then $\prod_{i \in I} R_i$ is S -flat by Lemma 3.1 and therefore, R is S -coherent by [18, Theorem 4.4]. \blacksquare

Next, we present an S -version of Stenström's characterizations of coherent rings.

Theorem 3.4 *Suppose that R_S is finitely presented. Then the following assertions are equivalent:*

1. R is S -coherent.
2. Every direct limit of S -FP-injective R -modules is an S -FP-injective R -module.
3. Every direct limit of FP-injective R_S -modules is an FP-injective R_S -module.

Proof. 1. \Rightarrow 2. Let $(M_i)_{i \in J}$ be a direct system of S -FP-injective modules over a directed set J . Let I be a finitely generated ideal of R . We construct the following commutative diagram:

$$\begin{array}{ccccccc}
\varinjlim \operatorname{Hom}_R(R_S, M_i) & \rightarrow & \varinjlim \operatorname{Hom}_R(I_S, M_i) & \rightarrow & \varinjlim \operatorname{Ext}_R^1(R_S/I_S, M_i) & \rightarrow & 0 \\
\downarrow & & \downarrow & & \downarrow & & \\
\operatorname{Hom}_R(R_S, \varinjlim M_i) & \rightarrow & \operatorname{Hom}_R(I_S, \varinjlim M_i) & \rightarrow & \operatorname{Ext}_R^1(R_S/I_S, \varinjlim M_i) & \rightarrow & 0
\end{array}$$

Since R is S -coherent, I is S -finitely presented. Hence, I_S is finitely presented R_S -module. Given that R_S is finitely presented, by Lemma 2.8, I_S is also a finitely presented R -module. Therefore, the first two vertical morphisms in the above diagram are isomorphisms. Consequently, by the Five Lemma, $\operatorname{Ext}_R^1(R_S/I_S, \varinjlim M_i) \cong \varinjlim \operatorname{Ext}_R^1(R_S/I_S, M_i) = 0$. Thus, $\varinjlim M_i$ is S -FP-injective by Theorem 2.13.

2. \Rightarrow 3. This follows from Proposition 2.11.

3. \Rightarrow 1. By [20, Theorem 3.2], R_S is a coherent ring. Since R_S is finitely presented, R is S -coherent by Theorem 3.3. \blacksquare

Finally, we present an S -counterpart to the classical result by Cheatham and Stone [7, Theorem 1]. To do so, recall the following lemma:

Lemma 3.5 ([5], Lemma 2.5.2) *An R -module M is S -flat if and only if its character $\operatorname{Hom}_{\mathbb{Z}}(M, \mathbb{Q}/\mathbb{Z})$ is S -injective.*

Theorem 3.6 *Let R be a ring and S a multiplicative subset of R . Consider the following assertions:*

1. R is S -coherent,
2. M is S -FP-injective if and only if M^+ is S -flat,
3. M is S -FP-injective if and only if M^{++} is S -injective,

4. M is S -flat if and only if M^{++} is S -flat,

The implications $2. \Rightarrow 3. \Rightarrow 4. \Rightarrow 1.$ hold true. Assuming that R_S is finitely presented as an R -module, then all the four assertions are equivalent.

Proof. $2. \Rightarrow 3.$ This follows from Lemma 3.5.

$3. \Rightarrow 4.$ Let M be an S -flat R -module. Then, by Lemma 3.5, M^+ is S -injective; so, by (3) M^{+++} is S -injective and hence, again by Lemma 3.5, M^{++} is S -flat.

Conversely, if M^{++} is S -flat, then M , being a pure submodule of M^{++} [23, Proposition 2.3.5], is S -flat by [3, Lemma 2.1].

$4. \Rightarrow 1.$ Let $(M_i)_{i \in I}$ be a family of flat R -modules. We aim to show that $\prod_{i \in I} M_i$ is S -flat. It follows then from [18, Theorem 4.4] that R is S -coherent. Since every flat R -module is S -flat, $\bigoplus_{i \in I} M_i$ is S -flat, and hence

$$\left(\bigoplus_{i \in I} M_i\right)^{++} \cong \left(\prod_{i \in I} M_i^+\right)^+$$

is S -flat by (4). By [7, Lemma 1], $\bigoplus_{i \in I} M_i^+$ is a pure submodule of $\prod M_i^+$. Then the induced sequence

$$\left(\prod M_i^+\right)^+ \rightarrow \left(\bigoplus M_i^+\right)^+ \rightarrow 0$$

splits. Thus, $\prod M_i^{++} \cong \left(\bigoplus M_i^+\right)^+$ is S -flat. Since, for each $i \in I$, M_i is a pure submodule of M^{++} , $\prod M_i$ is a pure submodule of $\prod M_i^{++}$ by [7, Lemma 1]. Hence, $\prod_{i \in I} M_i$ is S -flat by [3, Lemma 2.1].

$1. \Rightarrow 2.$ Assume R_S is finitely presented. Since R is S -coherent, R_S is a coherent ring. Then, by [5, Lemma 2.21], we have the following isomorphism:

$$\mathrm{Tor}_R^1(M^+, (R/I)_S) \cong \mathrm{Ext}_R^1((R/I)_S, M)^+$$

for any finitely generated ideal I of R . Thus, (2) follows from Theorem 2.13 and [4, Proposition 2.5]. \blacksquare

Let R be an S -perfect ring; that is, every S -flat R -module is projective [3, Definition 4.1]. Then, R_S is a finitely presented R -module. Indeed, R_S is projective and cyclic as an R -module [3, Theorem 4.9]. Therefore, by [19, Proposition 3.11], R_S is finitely presented. Thus, Theorems 3.3, 3.4 and 3.6 characterize, in particular, when an S -perfect ring R is S -coherent. We conclude with the following example:

Example 3.7 Let R_1 be a semisimple ring and let S_1 be a multiplicative subset of R_1 . Let R_2 be a non-coherent commutative ring. Consider the ring $R = R_1 \times R_2$ with the multiplicative subset $S = S_1 \times 0$. Then,

1. R_S is finitely presented.

2. R is S -coherent, but it not is coherent.

Proof. 1. Since R_1 is S_1 -perfect, $(R_1)_{S_1}$ is a finitely generated projective R_1 -module by [3, Theorem 4.9]. Then $R_S \cong (R_1)_{S_1} \times 0$ is a finitely generated projective R -module, hence it is finitely presented.

2. R is S -coherent by [6, Proposition 3.5]. ■

References

- [1] D. D. Anderson and T. Dumitrescu, *S-Noetherian rings*, Commun. Algebra. **30**(2002) 4407-4416.
- [2] S. Bazzoni and L. Positselski, *S-almost perfect commutative rings*, J. Algebra. **532**(2019) 323-356.
- [3] D. Bennis and A. Bouziri, *When every S-flat module is (flat) projective*, Commun. in Algebra. **52**(10)(2024) 4480-4491.
- [4] D. Bennis and A. Bouziri, *S-flat cotorsion pair*, Bull. Korean Math. Soc. to appear. <https://doi.org/10.48550/arXiv.2403.09242>
- [5] D. Bennis and A. Bouziri, *On S-injective modules*, Available from: <https://arxiv.org/abs/2406.02041>
- [6] D. Bennis and M. El Hajoui, *On S-coherence*, J. Korean Math. Soc. **55**(6)(2018) 1499-1512.
- [7] T. J. Cheatham and D. R. Stone, *Flat and projective character modules*, Proc. Amer. Math. Soc. **81**(1981) 175-177.
- [8] F. Couchot, *Exemples d'anneaux auto-fp-injectifs*, Commun. Algebra. **10**(4)(1982) 339-360.
- [9] E. E. Enochs and O. M. G. Jenda, *Relative Homological Algebra*, Berlin-New York: Walter de Gruyter, 2000.
- [10] S. Glaz, *Commutative Coherent Rings*. Lecture Notes in Mathematics, vol. **1371**, Spring-Verlag, Berlin, 1989.
- [11] B. Maddox, *Absolutely pure modules*, Proc. Amer. Math. Soc. **18**(1967) 155-158.
- [12] C. Megibben, *Absolutely pure modules*, Proc. Amer. Math. Soc. **26**(1970) 561-566.
- [13] E. Matlis, *Commutative coherent rings*, Can. J. Math. **6**(1982) 1240-1244

- [14] K. R. Pinzon, *Absolutely Pure Modules*, Thesis, University of Kentucky, 2005.
- [15] L. Positselski, *Contraadjusted modules, contramodules, and reduced cotorsion modules*, Mosc. Math. J. **17**(3)(2017) 385-455.
- [16] L. Positselski, *Triangulated Matlis equivalence*, J. Algebra Appl. **17**(4)(2018) 1850067.
- [17] L. Positselski and A. Slávik, *On strongly flat and weakly cotorsion modules*, Math. Z. **291**(3-4)(2019) 831-875.
- [18] W. Qi, X. Zhang and W. Zhao, *New Characterizations of S -coherent rings*, J. Algebra Appl. **22**(2023) 2350078.
- [19] J. Rotman, *An Introduction to Homological Algebra*, Academic Press, New York, 2009.
- [20] B. Stenström, *Coherent rings and FP-injective modules*, J. London Math. Soc. **2**(2)(1970) 323-329.
- [21] F. G. Wang and H. Kim, *Foundations of Commutative Rings and Their Modules*, Singapore, Springer, 2016.
- [22] R.B. Warfield, *Purity and algebraic compactness for modules*, Pac. J. Math. **28**(3)(1969) 699-719.
- [23] J. Xu, *Flat covers of modules*, Lecture notes in mathematics, vol **1634**, 1996.

Driss Bennis: Faculty of Sciences, Mohammed V University in Rabat, Rabat, Morocco.

e-mail address: driss.bennis@fsr.um5.ac.ma; driss_bennis@hotmail.com

Ayoub Bouziri: Faculty of Sciences, Mohammed V University in Rabat, Rabat, Morocco.

e-mail address: ayoubbouziri66@gmail.com; ayoub_bouziri@um5.ac.ma