

THE EULER CHARACTERISTIC OF \mathcal{A}_g VIA HODGE INTEGRALS

AITOR IRIBAR LOPEZ

ABSTRACT. We prove the Harder-Siegel formula for the Euler characteristic of \mathcal{A}_g via the intersection theory of $\overline{\mathcal{M}}_g$ and a vanishing result for lambda classes on the boundary of the toroidal compactifications of \mathcal{A}_g , recently proven by Canning, Molcho, Oprea and Pandharipande.

INTRODUCTION

The moduli space of principally polarized abelian varieties of dimension g is a smooth Deligne-Mumford stack of dimension $\binom{g+1}{2}$, usually denoted by \mathcal{A}_g . Over the complex numbers, this is the same as saying that \mathcal{A}_g is an orbifold, and it can be identified with the stack quotient of the Siegel upper half space by the group of symplectic matrices with integer coefficients

$$(1) \quad \mathcal{A}_g \cong [\mathcal{H}_g / \mathrm{Sp}(2g, \mathbb{Z})].$$

The Gauss-Bonnet formula [Har71] of Harder shows that the Euler characteristic of \mathcal{A}_g can be obtained from the volume of a fundamental domain for the action of $\mathrm{Sp}(2g, \mathbb{Z})$ on \mathcal{H}_g , which was known to Siegel [Sie36]:

Theorem 1. *The Euler characteristic of \mathcal{A}_g is*

$$(2) \quad \chi(\mathcal{A}_g) = \zeta(-1) \cdots \zeta(1-2g).$$

We employ the *logarithmic Gauss-Bonnet formula* to reduce the calculation of the Euler characteristic of \mathcal{A}_g to an intersection theoretic problem on the toroidal compactifications of \mathcal{A}_g , which is solved via Hodge integrals and the Torelli map

On every toroidal compactification of \mathcal{A}_g , there is a universal semiabelian scheme $\pi : \mathcal{G}_g \rightarrow \overline{\mathcal{A}}_g$ with a zero section s . The *Hodge bundle* is a vector bundle of rank g defined by

$$\mathbb{E}_g = s^* \Omega_\pi,$$

and its Chern classes are denoted by $\lambda_i = c_i(\mathbb{E}_g)$. Let $\overline{\mathcal{M}}_{g,n}$ denote the moduli space of stable curves of genus g with n markings, and let

$$\pi : \overline{\mathcal{C}}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$$

be the universal curve, with sections s_1, \dots, s_n corresponding to the markings. It is a smooth DM stack of dimension $3g - 3 + n$ and it has a *Hodge bundle*, defined by

$$\mathbb{E}_g = \pi_* \omega_\pi.$$

We denote its Chern classes by λ_i also. On some toroidal compactifications there is a Torelli morphism

$$\mathrm{Tor} : \overline{\mathcal{M}}_{g,n} \rightarrow \overline{\mathcal{A}}_g,$$

as shown in [Ale04, Nam76], and $\mathrm{Tor}^* \mathbb{E}_g = \mathbb{E}_g$. Esnault and Viehweg showed in [EV02] that the λ classes satisfy *Mumford's relation*:

$$(3) \quad (1 + \lambda_1 + \dots + \lambda_g)(1 - \lambda_1 + \dots + (-1)^g \lambda_g) = c(\mathbb{E}_g \oplus \mathbb{E}_g^\vee) = 1 \text{ in } \mathrm{CH}^*(\overline{\mathcal{A}}_g).$$

On $\overline{\mathcal{M}}_{g,n}$ there are also line bundles $\mathbb{L}_i = s_i^* \omega_\pi$ representing the cotangent space at the i -th marking, and its first Chern class is ψ_i . The ψ classes and the λ classes are part of the *tautological ring* of $\overline{\mathcal{M}}_{g,n}$ (see [FP13, Pan18] for the definition). Integrals of the form

$$\int_{\overline{\mathcal{M}}_{g,n}} \lambda_1^{a_1} \dots \lambda_g^{a_g} \psi_1^{b_1} \dots \psi_n^{b_n}$$

are *Hodge integrals*, and they appear naturally in Gromov-Witten theory.

We will consider the locus \mathcal{B}_g of semiabelian varieties which are trivial extension of an abelian variety by a torus. λ_g is proportional to the class $[\mathcal{B}_g]$ when we restrict ourselves to an open subset of $\overline{\mathcal{A}}_g$ given by degenerations of torus rank at most 1 [vdG99, EvdG05]. Then we compute the proportionality factor in two ways. The first is through the isomorphism $\mathcal{B}_g \cong \mathcal{A}_{g-1} \times \mathrm{B}\mathbb{Z}/2\mathbb{Z}$, and the second one is pulling back via the Torelli map. We will see that the formula for the Euler characteristic (2) follows from the evaluation of the Hodge integrals

$$\int_{\overline{\mathcal{M}}_g} \lambda_g \lambda_{g-1} \lambda_{g-2} \quad \text{and} \quad \int_{\overline{\mathcal{M}}_{g,1}} \frac{\lambda_g \lambda_{g-1} c(\mathbb{E}_g^\vee)}{1 - \psi},$$

which had been computed by Faber and Pandharipande [FP00a, FP00b].

To a polarization on an abelian variety of dimension g we can associate a list of numbers $\delta = (d_1, \dots, d_g)$, where $d_i \mid d_{i+1}$, and there is a moduli stack of abelian varieties of dimension g together with a polarization of type δ , denoted by $\mathcal{A}_{g,\delta}$. When $d_i = 1$ for all i , we recover \mathcal{A}_g . These moduli spaces are part of a tower of étale maps. By the degree calculations of [IL24], the Euler characteristics of all the moduli spaces $\mathcal{A}_{g,\delta}$ is determined:

Theorem 2. *For a list of positive integers $\delta = (d_1, \dots, d_g)$ such that $d_i \mid d_{i+1}$ for all i ,*

$$\chi(\mathcal{A}_{g,\delta}) = \left(d_g^{2g-2} d_{g-1}^{2g-6} \dots d_1^{-2g+2} \prod_{1 \leq i < j \leq g} \prod_{p \mid d_j/d_i} \frac{(1 - p^{-2(j-i+1)})}{(1 - p^{-2(j-i)})} \right) \chi(\mathcal{A}_g),$$

where the product is over primes p .

We will prove in Corollary 7 that the non-vanishing of $\chi(\mathcal{A}_g)$ implies the non-vanishing of $\lambda_{g-1} \dots \lambda_1$ on $\mathrm{CH}^*(\mathcal{A}_g)$. This result was first established in [vdG99, Corollary 1.3] using the geometry of \mathcal{A}_g over fields of positive characteristic.

Further directions. The main geometric input for this formula is that λ_g is proportional to $[\mathcal{B}_g]$ on $\mathcal{A}_g^{\leq 1}$, which follows from a residue calculation in [EvdG05] to express λ_g in terms of boundary strata of the toroidal compactifications of \mathcal{A}_g . Improvements of this result for the locus given by degenerations of abelian varieties of torus rank at most k for small k would lead to more connections to the intersection theory of $\overline{\mathcal{M}}_g$. Johannes Schmitt has checked that $[\mathcal{B}_2]$ is *not* proportional to λ_2 on $\overline{\mathcal{A}}_2$, so a deeper understanding is needed for torus rank at least 2.

Note that \mathcal{B}_g is one of the two components of the closure of the *product locus* $\mathcal{A}_1 \times \mathcal{A}_{g-1}$ in $\mathcal{A}_g^{\leq 1}$. In [COP24], the authors compute $\mathrm{Tor}^*([\mathcal{A}_1 \times \mathcal{A}_{g-1}])$ and prove that it is a tautological class on $\mathcal{M}_g^{\mathrm{ct}}$. Given our presentation of the fibered product $\mathcal{M}_g^{\leq 1} \times_{\mathcal{A}_g} \mathcal{B}_g$ in Lemma 4, we think it is reasonable to expect the following:

Conjecture 1. $\mathrm{Tor}^*([\overline{\mathcal{A}_1 \times \mathcal{A}_{g-1}}])$ lies in the tautological ring $R^*(\mathcal{M}_g^{\leq 1})$.

Acknowledgements. This article is the result of a series of conversations between the author and Rahul Pandharipande during the Alpine Algebraic Geometry Workshop in Obergurgl, Austria in September 2024. The author is also grateful to Sam Molcho and Johannes Schmitt for conversations about \mathcal{A}_g and its compactifications, to Gerard van der Geer for his comments about the history of the formula for $\chi(\mathcal{A}_g)$ and to Dragos Oprea for pointing out the article [EvdG05]. This project has received support from SNF-200020-219369.

LOGARITHMIC EULER CHARACTERISTIC

If \mathcal{Y} is a smooth DM-stack of dimension n with a smooth compactification $\overline{\mathcal{Y}}$ such that the complement D of \mathcal{Y} is a normal crossing divisor then the sheaf of meromorphic differentials having at most log poles along D is a vector bundle of rank n , denoted by $\Omega_{\overline{\mathcal{Y}}}(\log D)$ and it is well-known that

$$\chi(\mathcal{Y}) = (-1)^n \int_{\overline{\mathcal{Y}}} c_n(\Omega_{\overline{\mathcal{Y}}}(\log D)),$$

where χ is the Euler characteristic (see [CMZ20, Section 2] for a proof). When \mathcal{Y} is \mathcal{A}_g and $\overline{\mathcal{Y}}$ is one of its toroidal compactifications (constructed in [AMRT10] over the complex numbers, or [CF91] over the integers), $\Omega_{\overline{\mathcal{A}_g}}(\log D)$ is the *canonical extension* of $\Omega_{\mathcal{A}_g}$ (see [CF91, VI.4.1] for details) so we have the following:

$$\Omega_{\overline{\mathcal{A}_g}}(\log D) = \text{Sym}^2 \mathbb{E}_g.$$

The top Chern class of $\text{Sym}^2 \mathbb{E}_g$ can be computed by the Giambelli formula ([Ful84, Example 14.5.1]), and it equals

$$2^g \begin{vmatrix} \lambda_g & 0 & 0 & \dots & 0 \\ \lambda_{g-2} & \lambda_{g-1} & \lambda_g & \dots & 0 \\ \lambda_{g-4} & \lambda_{g-3} & \lambda_{g-2} & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & \lambda_1 \end{vmatrix} = 2^g \left(\lambda_g \lambda_{g-1} \dots \lambda_1 + \sum_{k=0}^{g-1} \underbrace{\lambda_g \dots \lambda_{g-k+1} \lambda_{g-k}^2}_{=0 \text{ by Mumford's relation (3)}} p_k(\lambda_1, \dots, \lambda_g) \right),$$

so

$$(4) \quad \chi(\mathcal{A}_g) = (-1)^{\binom{g+1}{2}} 2^g \int_{\overline{\mathcal{A}_g}} \lambda_1 \dots \lambda_g.$$

In particular, the integral on the right hand side does not depend on the toroidal compactification, and more generally any integral of λ classes is independent of the toroidal compactification, by choosing a common roof, and it makes sense on non-smooth toroidal compactifications considering Chern classes as operational Chow classes.

An abelian variety of dimension 1 is an elliptic curve, so

$$(5) \quad \chi(\mathcal{A}_1) = -2 \int_{\overline{\mathcal{A}_1}} \lambda_1 = -2 \int_{\overline{\mathcal{M}}_{1,1}} \psi = \zeta(-1).$$

λ_{g-k} -EVALUATIONS

Any toroidal compactification of \mathcal{A}_g has a canonical map to the Satake compactification

$$\beta : \overline{\mathcal{A}_g} \longrightarrow \overline{\mathcal{A}_g}^{\text{Sat}} = \mathcal{A}_g \sqcup \mathcal{A}_{g-1} \sqcup \dots \sqcup \mathcal{A}_0,$$

and we can obtain partial compactifications $\mathcal{A}_g^{\leq k} = \beta^{-1}(\mathcal{A}_g \sqcup \dots \sqcup \mathcal{A}_{g-k})$ of semiabelian varieties of torus rank at most k . It is shown in [CMOP24] that

$$\lambda_{g-k}|_{\overline{\mathcal{A}}_g \setminus \mathcal{A}_g^{\leq k}} = 0.$$

This result guarantees that the natural integration maps

$$\epsilon_{\mathcal{A}_g^{\leq k}}: \mathrm{CH}^{\binom{g+1}{2}+k-g}(\mathcal{A}_g^{\leq k}) \longrightarrow \mathbb{Q}$$

given by

$$\alpha \mapsto \int_{\overline{\mathcal{A}}_g} \bar{\alpha} \lambda_{g-k},$$

where $\bar{\alpha}$ is any extension of α to $\overline{\mathcal{A}}_g$ are well-defined. On the moduli space of curves we can define

$$\epsilon_{\mathcal{M}_g^{\leq k}}: \mathrm{CH}^{2g-3+k}(\mathcal{M}_g^{\leq k}) \longrightarrow \mathbb{Q}$$

analogously, where $\mathcal{M}_g^{\leq k}$ is the moduli space of stable curves whose dual graph has Betti number at most k .

If the Torelli morphism extends to $\overline{\mathcal{A}}_g$, and $[\mathcal{J}_g^{\leq k}]$ is the pushforward of 1 under this Torelli morphism to $\mathcal{A}_g^{\leq k}$, then the integration maps are related by the identity

$$(6) \quad \epsilon_{\mathcal{M}_g^{\leq k}}(\mathrm{Tor}^*(\alpha)) = \epsilon_{\mathcal{A}_g^{\leq k}}(\alpha \cdot [\mathcal{J}_g^{\leq k}]).$$

THE LOCUS \mathcal{B}_g

There is a natural map

$$j: \mathcal{A}_{g-1} \rightarrow \partial \mathcal{A}_g^{\leq 1}$$

sending A to the semiabelian variety $A \times \mathbb{G}_m$, whose image we denote by \mathcal{B}_g . It is isomorphic to $\mathcal{A}_{g-1} \times \mathrm{B}\mathbb{Z}/2\mathbb{Z}$ because of the extra automorphism of \mathbb{G}_m . The map j extends to toroidal compactifications of \mathcal{A}_{g-1} and satisfies $j^* \mathbb{E}_g = \mathbb{E}_{g-1} \oplus \mathcal{O}$. Therefore, the normal bundle to j is

$$N_j = \mathrm{Sym}^2(\mathbb{E}_{g-1}^\vee \oplus \mathcal{O}_{\mathcal{A}_{g-1}}) - \mathrm{Sym}^2(\mathbb{E}_{g-1}^\vee) = \mathbb{E}_{g-1}^\vee \oplus \mathcal{O}_{\mathcal{A}_{g-1}}.$$

We have the following result, first proven in cohomology¹ [vdG99, Proposition 1.10], and then in the Chow groups [EvdG05, Theorem 1.1]:

Theorem 3. *In the Chow ring of $\mathcal{A}_g^{\leq 1}$, the following holds:*

$$\lambda_g = \frac{|B_{2g}|}{2g} [\mathcal{B}_g],$$

where B_{2g} is the g -th even Bernoulli number.

We will give a new proof of the proportionality factor under the assumption that the two cycles are proportional. Let $\tau(g) \in \mathbb{Q}$ be such that

$$\lambda_g = \tau(g) [\mathcal{B}_g].$$

¹In fact, the result in cohomology is enough for our calculation, since the λ_{g-k} -evaluation maps factor through the cycle class map.

PULLBACK OF \mathcal{B}_g TO $\mathcal{M}_g^{\leq 1}$

Consider the Cartesian diagram

$$(7) \quad \begin{array}{ccc} \mathcal{Z} & \longrightarrow & \mathcal{M}_g^{\leq 1} \\ \downarrow & & \downarrow \text{Tor} \\ \mathcal{B}_g & \xrightarrow{j} & \mathcal{A}_g^{\leq 1} \end{array}$$

where j is a regular embedding.

Lemma 4. *For a partition $\mu = (g_1, \dots, g_l)$ of $g-1$ with $g_1 \leq \dots \leq g_l$, let $\mathcal{M}_{1,l}^{cycle}$ be the substack of $\overline{\mathcal{M}}_{1,l}$ given by curves without rational tails and whose normalization is a union of rational curves, and consider the gluing map*

$$\xi_\mu : \mathcal{M}_{g_1,1}^{ct} \times \dots \times \mathcal{M}_{g_l,1}^{ct} \times \mathcal{M}_{1,l}^{cycle} \longrightarrow \mathcal{M}_g^{\leq 1}$$

that attaches the marked point of the i -th moduli space of compact type to the i -th market point of $\mathcal{M}_{1,l}^{cycle}$. The images of the morphisms ξ_μ when μ runs through all the partitions of $g-1$ are disjoint and cover \mathcal{Z} .

Proof. Consider a prestable curve D of genus h whose dual graph is a cycle of length k . Denote by D_1, \dots, D_k the irreducible components of \tilde{D} , and consider points $p_i, q_i \in D_i$ such that q_i and p_{i+1} are identified in C . Then, $\text{Jac}(D)$ is the \mathbb{G}_m -extension of $\text{Jac}(\tilde{D})$ that corresponds to

$$(\mathcal{O}_{D_1}(p_1 - q_1), \dots, \mathcal{O}_{D_k}(p_k - q_k)) \in \text{Pic}^{0, \dots, 0}(\tilde{D}) \cong \text{Jac}(\tilde{D})^\vee.$$

Therefore, the semiabelian variety $\text{Jac}(D)$ lies in \mathcal{B}_h if and only if $\mathcal{O}_{D_i}(p_i - q_i) = \mathcal{O}_{D_i}$; this is, if and only if all the D_i are rational.

Now consider a general stable curve C of genus g whose dual graph Γ_C has Betti number 1. There is a subgraph of Γ_C which is minimal among all the subgraphs that have Betti number 1; let $D \subset C$ be the curve that corresponds to such a graph. Then $\overline{C \setminus D}$ is a disjoint union of l curves C_i of compact type, and by stability $g(C_i) > 0$. Moreover,

$$\text{Jac}(C) = \text{Jac}(D) \times \prod \text{Jac}(C_i),$$

so $\text{Jac}(C)$ lies in \mathcal{B}_g if and only if $\text{Jac}(D)$ lies in \mathcal{B}_g , which happens if and only if \tilde{D} is a union of rational curves (and in particular $g(D) = 1$), so C lies in the image of ξ_μ for the partition $\mu = (g(C_1), \dots, g(C_l))$. □

Remark 1. Note that

$$\dim(\mathcal{M}_{g_1,1}^{ct} \times \dots \times \mathcal{M}_{g_l,1}^{ct} \times \mathcal{M}_{1,l}^{cycle}) = \sum_{i=1}^l (3(g_i - 1) + 1) + (l - 1) = \dim(\mathcal{M}_g^{\leq 1}) - (l + 1),$$

and that $\mathcal{M}_{1,l}^{cycle}$ is not proper except for the case $l = 1$, where it is isomorphic to $\text{B}\mathbb{Z}/2\mathbb{Z}$. In particular,

$$\int_{\mathcal{M}_{1,1}^{cycle}} 1 = \frac{1}{2}.$$

When $\mu = (g-1)$, the image of ξ_μ is defined to be the *principal locus*.




FIGURE 1. Two examples of the dual graph of a curve of genus 11 in \mathcal{Z} . They correspond to the partitions $(1, 2, 3, 4)$ and (10)

Lemma 5. *The principal locus is closed in \mathcal{Z} .*

Proof. If $\mu = (g_1, \dots, g_l)$ then the dual graph of a curve in the image of ξ_μ is a stable cycle with l markings and a collection of rooted trees together with an assignment of a tree to each marking on the cycle. There are five types of edges:

- edges belonging to the cycle, when it has length at least 2,
- edges belonging to the tree,
- edges between a vertex of the cycle and the root of the associated tree when the genus of the root is not 0,
- edges between a vertex of the cycle and the root of the tree when the genus of the root is 0, or
- the edge of the cycle, when it has length 1.

If we contract an edge of the first two types, the curve stays in ξ_μ . If we contract an edge of the third type, the new curve lies in the image of $\xi_{\mu'}$ where μ' is a refinement of μ , and the last two contractions take the curve outside of \mathcal{Z} . The partition $(g-1)$ is not the refinement of any other partition, so the principal locus is closed under specialization. \square

In particular, $\xi_{(g-1)}$ is proper. Let \mathcal{Z}' be the closure of the complement of the principal locus in \mathcal{Z} .

Lemma 6. $\lambda_{g-2} \in \text{CH}^{g-2}(\mathcal{M}_g^{\leq 1})$ vanishes when restricted to \mathcal{Z}' .

Proof. From the discussion in Lemma 5, the points in $\xi_{(g-1)}(\mathcal{M}_{g,1} \times \mathcal{M}_{1,1}^{\text{cycle}})$ cannot be in \mathcal{Z}' , so the normalization of any curve in \mathcal{Z}' has to be a union of rational curves and curves of genus g_1, \dots, g_l with $\sum g_i = g-1$ and $l \geq 2$, but note that

$$c_{g-2} \left(\bigoplus \mathbb{E}_{g_i} \right) = \sum_{i=1}^l c_{g_i-1}(\mathbb{E}_{g_i}) \cdot \prod_{j \neq i} c_{g_j}(\mathbb{E}_{g_j}),$$

which vanishes because the domain of all the morphisms ξ_μ is a space of curves of compact type and $\lambda_h|_{\mathcal{M}_h^{\text{ct}}} = 0$. \square

$\xi_{(g-1)}$ is a regular embedding with normal bundle $\mathbb{L}^\vee \boxplus \mathcal{O}_{\mathcal{M}_{1,1}^{cyc}}$, where \mathbb{L} is the cotangent line, and therefore the excess class for the principal component in the diagram (7) is

$$c_{top} \left(N_j - N_{\xi_{(g-1)}} \right) = c_{g-2}(\mathbb{E}_{g-1}^\vee - \mathbb{L}^\vee).$$

By the residual intersection formula [Ful84, Corollary 9.2.3],

$$j^!([\mathcal{B}_g]) = \xi_{(g-1),*} \left(\left[\frac{c(\mathbb{E}_{g-1}^\vee)}{1-\psi} \right]_{g-2} \right) + \mathbf{R}$$

where \mathbf{R} is the residual class supported on \mathcal{Z}' . By Lemma 6, $\lambda_{g-2}j^!([\mathcal{B}_g])$ has a natural extension to $\overline{\mathcal{M}}_g$, namely,

$$\lambda_{g-2} \left[\overline{\xi}_{(g-1),*} \left(\frac{c(\mathbb{E}_{g-1}^\vee)}{1-\psi} \right) \right]_g \in \text{CH}^{2g-2}(\overline{\mathcal{M}}_g),$$

where $\overline{\xi}_{(g-1)} : \overline{\mathcal{M}}_{g-1,1} \times \mathcal{M}_{1,1}^{cyc} \rightarrow \overline{\mathcal{M}}_g$ is the gluing map.

Proof of Theorem 3. Using formula (6),

$$\epsilon_{\mathcal{A}_g^{\leq 1}}(\lambda_{g-2}[\mathcal{B}_g] \cdot [\mathcal{J}_g^{\leq 1}]) = \left(\int_{\mathcal{M}_{1,1}^{cyc}} 1 \right) \left(\int_{\overline{\mathcal{M}}_{g-1,1}} \frac{\lambda_{g-1}\lambda_{g-2}c(\mathbb{E}_{g-1}^\vee)}{1-\psi} \right) = \frac{1}{2} \frac{1}{(2g-2)!} \frac{|B_{2g-2}|}{2g-2},$$

by the work in [FP00b]. In [FP00a], the authors also showed that

$$\epsilon_{\mathcal{A}_g^{\leq 1}}(\lambda_{g-2}\lambda_g \cdot [\mathcal{J}_g^{\leq 1}]) = \int_{\overline{\mathcal{M}}_g} \lambda_g \lambda_{g-1} \lambda_{g-2} = \frac{1}{2(2g-2)!} \frac{|B_{2g-2}|}{2g-2} \frac{|B_{2g}|}{2g}.$$

Dividing the last two equations, we obtain the value of $\tau(g)$. □

THE EULER CHARACTERISTIC OF $\mathcal{A}_{g,\delta}$

First, we see that the evaluation of $\tau(g)$ for all g is equivalent to the knowledge of the Euler characteristic of \mathcal{A}_g :

Proof of Theorem 1. Since $j^*\mathbb{E}_g = \mathbb{E}_{g-1} \oplus \mathcal{O}$, where $j : \mathcal{A}_{g-1} \rightarrow \mathcal{A}_g^{\leq 1}$, we see that

$$\int_{\overline{\mathcal{A}}_g} \lambda_g \dots \lambda_1 = \epsilon_{\mathcal{A}_g^{\leq 1}}([\tau(g)[\mathcal{B}_g]\lambda_{g-1} \dots \lambda_1]) = \frac{\tau(g)}{2} \int_{\overline{\mathcal{A}}_{g-1}} \lambda_{g-1} \dots \lambda_1,$$

and so, by the logarithmic Gauss-Bonnet (4), we see that

$$\chi(\mathcal{A}_g) = (-1)^g \tau(g) \chi(\mathcal{A}_{g-1})$$

and note that $(-1)^g \tau(g) = (-1)^g \frac{|B_{2g}|}{2g} = \zeta(1-2g)$. □

In order to compare \mathcal{A}_g and $\mathcal{A}_{g,\delta}$, we introduce a level structure. Let θ be a polarization on an abelian variety. Then

$$\ker(\theta) \cong (\mathbb{Z}/d_1\mathbb{Z} \times \dots \times \mathbb{Z}/d_g\mathbb{Z})^2$$

as symplectic groups, where (d_1, \dots, d_g) is the type of θ . The moduli space of triplets (A, θ, F) , where $(A, \theta) \in \mathcal{A}_{g,\delta}$ and F is a symplectic basis of $\ker(\theta)$ is denoted by $\mathcal{A}_{g,\delta}^{\text{lev}}$, and has étale

morphisms

$$\begin{array}{ccc} & \mathcal{A}_{g,\delta}^{\text{lev}} & \\ \pi_\delta \swarrow & & \searrow \varphi_\delta \\ \mathcal{A}_{g,\delta} & & \mathcal{A}_g \end{array},$$

where π_δ forgets the symplectic basis and φ_δ sends an abelian variety X to its quotient by a Lagrangian subgroup of $\ker(\theta)$, see [IL24, Section 2] for details.

It follows that

$$\chi(\mathcal{A}_{g,\delta}) = \frac{\deg(\varphi_\delta)}{\deg(\pi_\delta)} \chi(\mathcal{A}_g),$$

and

$$\frac{\deg(\varphi_\delta)}{\deg(\pi_\delta)} = d_g^{2g-2} d_{g-1}^{2g-6} \dots d_1^{-2g+2} \prod_{1 \leq i < j \leq g} \prod_{p|d_j/d_i} \frac{(1-p^{-2(j-i+1)})}{(1-p^{-2(j-i)})}$$

was computed in [IL24, Proposition 26]. This proves Theorem 2.

THE HIRZEBRUCH-MUMFORD PROPORTIONALITY THEOREM

Consider the Lagrangian Grassmanian LG_g , that parametrizes Lagrangian subspaces of dimension g inside a symplectic vector space of dimension $2g$. It is a smooth projective variety of dimension $\binom{g+1}{2}$. The universal Lagrangian subspace

$$\mathbb{S}_g \rightarrow \text{LG}_g$$

defines a vector bundle of rank g , with Chern classes $x_i = c_i(\mathbb{S}_g)$, and $\text{Sym}^2(\mathbb{S}_g)$ is the cotangent bundle to LG_g . Mumford shows in [Mum77] that there is a constant $K(g)$ such that for any $a_i \in \mathbb{N}$,

$$\int_{\overline{\mathcal{A}}_g} \lambda_1^{a_1} \dots \lambda_g^{a_g} = K(g) \int_{\text{LG}_g} x_1^{a_1} \dots x_n^{a_n}.$$

This constant has been determined in [vdG99, Theorem 1.13] by the Gauss-Bonnet formula:

$$\chi(\mathcal{A}_g) = K(g) \chi(\text{LG}_g).$$

An approach to prove the formula for $\chi(\mathcal{A}_g)$ could be to determine $K(g)$ in an alternative way.

Van der Geer also shows that the assignment $\lambda_i \mapsto x_i$ defines an isomorphism between the subring of $\text{CH}^*(\mathcal{A}_g)$ generated by the classes $\lambda_1, \dots, \lambda_{g-1}$ (also known as the *tautological ring*) and the cohomology ring of LG_{g-1} . One of the steps of the proof given there relied on a characteristic p argument and the ampleness of λ_1 . We give a different proof:

Corollary 7 ([vdG99, Corollary 1.4]). $\lambda_1 \dots \lambda_{g-1} \neq 0$ in $\text{CH}_g(\mathcal{A}_g)$.

Proof. If it were 0 then

$$0 = \epsilon_{\mathcal{A}_g}(\lambda_1 \dots \lambda_{g-1}) = \int_{\overline{\mathcal{A}}_g} \lambda_1 \dots \lambda_g = (-1)^{\binom{g+1}{2}} 2^{-g} \chi(\mathcal{A}_g),$$

a contradiction. □

REFERENCES

- [Ale04] Valery Alexeev. Compactified jacobians and the Torelli map. *Publ. RIMS*, 4:1241–1265, 2004.
- [AMRT10] Avner Ash, David Mumford, Michael Rapoport, and Yung-sheng Tai. *Smooth Compactifications of Locally Symmetric Varieties*. Cambridge Mathematical Library. Cambridge University Press, 2 edition, 2010.
- [CF91] Ching-Li Chai and Gerd Faltings. *Degeneration of Abelian Varieties*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics. Springer Berlin Heidelberg, 1991.
- [CMOP24] Samir Canning, Sam Molcho, Dragos Oprea, and Rahul Pandharipande. Tautological projection for cycles on the moduli space of abelian varieties, 2024. [arXiv:2401.15768](#).
- [CMZ20] Matteo Costantini, Martin Möller, and Jonathan Zachhuber. The Chern classes and the Euler characteristic of the moduli spaces of abelian differentials, 2020. [arXiv:2006.12803](#).
- [COP24] Samir Canning, Dragos Oprea, and Rahul Pandharipande. Tautological and non-tautological cycles on the moduli space of abelian varieties, 2024. [arXiv:2408.08718](#).
- [EV02] Hélène Esnault and Eckart Viehweg. Chern classes of Gauss–Manin bundles of weight 1 vanish. *K-Theory*, 26:287–305, 07 2002.
- [EvdG05] Torsten Ekedahl and Gerard van der Geer. Cycles representing the top Chern class of the Hodge bundle on the moduli space of abelian varieties. *Duke Math. J.*, 129(1):187–199, 2005.
- [FP00a] Carel Faber and Rahul Pandharipande. Hodge integrals and Gromov–Witten theory. *Inventiones Mathematicae*, 139:173–199, 2000.
- [FP00b] Carel Faber and Rahul Pandharipande. Logarithmic series and Hodge integrals in the tautological ring. With an appendix by Don Zagier. *Michigan Mathematical Journal*, 48(1):215 – 252, 2000.
- [FP13] Carel Faber and Rahul Pandharipande. Tautological and nontautological cohomology of the moduli space of curves. In Gavril Farkas and Ian Morrison, editors, *Handbook of moduli*, volume 24 of *Adv. Lect. Math.*, pages 293–330. International Press, 2013.
- [Ful84] William Fulton. *Intersection Theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete : a series of modern surveys in mathematics. Folge 3. Springer-Verlag, 1984.
- [Har71] G. Harder. A Gauss–Bonnet formula for discrete arithmetically defined groups. *Annales scientifiques de l’École Normale Supérieure*, Ser. 4, 4(3):409–455, 1971.
- [IL24] Aitor Iribar López. Twisted products on moduli spaces of abelian varieties. *in preparation*, 2024.
- [Mum77] David Mumford. Hirzebruch’s proportionality theorem in the non-compact case. *Inventiones mathematicae*, 42:239–272, 1977.
- [Nam76] Yukio Namikawa. A new compactification of the Siegel space and degenerations of abelian varieties i–ii. *Math. Ann.*, 221(201):91–141 201–241, 1976.
- [Pan18] Rahul Pandharipande. A calculus for the moduli space of curves. In Tommaso de Fernex, Brendan Hassett, Mircea Mustata, Martin Olsson, Mihnea Popa, and Richard Thomas, editors, *Algebraic geometry: Salt Lake City 2015*, volume 97.1 of *Pure Math.*, pages 459–487. American Mathematical Society, 2018.
- [Sie36] Carl Ludwig Siegel. The volume of the fundamental domain for some infinite groups. *Trans. Amer. Math. Soc.*, 39(2):209–218, 1936.
- [vdG99] Gerard van der Geer. *Cycles on the Moduli Space of Abelian Varieties*, pages 65–89. Vieweg+Teubner Verlag, Wiesbaden, 1999.