INTERIOR OF CERTAIN SUMS AND CONTINUOUS IMAGES OF VERY THIN CANTOR SETS

YEONWOOK JUNG AND CHUN-KIT LAI

ABSTRACT. We show that for all Cantor set K_1 on \mathbb{R}^d , it is always possible to find another Cantor set K_2 so that the sum $g(K_1) + K_2$ (where g is a C^1 local diffeomorphism) has non-empty interior, and the existence of the interior is robust under small perturbation of the mapping. More generally, we can also show that the image set $H(\alpha, K_1, K_2)$, where H is some C^1 function on $\mathbb{R}^N \times \mathbb{R}^d \times \mathbb{R}^d$ with nonvanishing Jacobian, have non-empty interior for α all in an open ball of \mathbb{R}^N . This result allows us to show that all Cantor sets are not topologically universal using C^1 local diffeomorphism, proving a stronger version of the topological Erdős similarity conjecture. Moreover, we are also able to construct a Cantor set of dimension d on \mathbb{R}^{2d} , whose distance set has an interior.

1. INTRODUCTION

1.1. **Background.** In this paper, **a Cantor set** on \mathbb{R}^d always refers to a compact, totally disconnected and perfect set. We are interested in determining if the image set of Cantor sets under continuously differentiable functions possessed an interior. The simplest case will be about the Minkowski sum of two Cantor sets. Let (K_1, K_2) be a pair of Cantor sets on \mathbb{R}^d . One of the basic questions studied by many mathematicians mainly on \mathbb{R}^1 is that

(Qu): For which pair of Cantor sets (K_1, K_2) , $K_1 + \alpha K_2$ has non-empty interior for all α in an interval of \mathbb{R} ?

The study of the topological and the dimensional properties for the algebraic sum of Cantor sets $K_1 + K_2$ arises naturally in dynamical system, number theory and harmonic analysis [1, 3, 12, 13]. In dynamical system, the famous Palis-Taken's conjecture (see e.g. [21]) asserted that for generic dynamically defined Cantor sets, the algebraic sum either has empty interior or contains an interval . Substantial progress was made by Moreria and Yoccoz for non-linear Cantor sets [4]. However, there exists dynamically defined Cantor set K such that K - K has positive measure but no interior [23] (See also [24]).

Note that $K_1 + \alpha K_2$ is the orthogonal projection of $K_1 \times K_2$ onto the line with slope $\tan(\alpha)$. Therefore, $\dim_H(K_1 + \alpha K_2) \leq \dim_H(K_1) + \dim_B(K_2)$ (by the product

²⁰²⁰ Mathematics Subject Classification. Primary: 28A80, 28A75.

formula for dimension [7]). If $K_1 + \alpha K_2$ has interior, it is necessary that

$$\dim_H(K_1) + \dim_B(K_2) \ge 1.$$

On the other hand, on \mathbb{R}^1 , using the notion of Newhouse thickness $\tau(K)$ via the Newhouse gap lemma, one knows that if

$$\tau(K_1)\tau(K_2) \ge 1,$$

then the $K_1 + \alpha K_2$ always contains an interior for all $\alpha \neq 0$. Nonetheless, there is a gap between the sum of the dimensions being larger than 1 and the product of the thickness larger than 1, so the question is undetermined for many pairs of Cantor sets. In the case of self-similar sets K_1 and K_2 defined by two maps whose contraction ratios are a and b respectively, there is a mysterious region \mathcal{R} for which question has remained open ([27]). We also refer to [22, 28, 20] for some recent results about this direction. The arithmetic sum of Cantor sets is largely open on higher dimensions because there is no obvious choice for defining the Newhouse thickness. Yavicoli [30] recently defined a new version of thickness on \mathbb{R}^d , which allows us to determine certain (K_1, K_2) fulfills the question stated. In another direction, Feng and Wu introduced another type of thickness τ_{FW} , on \mathbb{R}^d in which adding enough number of Cantor sets with $\tau_{FW} \geq c > 0$ will ensure the sum has an non-empty interior [9].

The question about the algebraic sum of Cantor sets can be generalized to the following non-linear setting. As far as we know, this is first due to Simon and Taylor [26]. Let $H : \Lambda \times \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}^d$, where $\Lambda \subset \mathbb{R}^N$ (can be thought as a parameter set) be a continuously differentiable function, denoted by $C^1(\Lambda \times \mathbb{R}^d \times \mathbb{R}^d)$. We can ask under what condition on the pair of Cantor sets (K_1, K_2) , we have the image set

$$H(\alpha, K_1, K_2) = \{H(\alpha, k_1, k_2) : k_1 \in K_1, k_2, \in K_2\}$$

possesses an non-empty interior for all α inside an open ball of Λ . When $H(\alpha, x, y) = x + \alpha y$, this reduces back to (**Qu**). Simon and Taylor showed that if $\tau(K_1)\tau(K_2) > 1$ (with d = 1) and H has non-vanishing partial derivatives, then $(H(\alpha, K_1, K_2))^{\circ} \neq \emptyset$ around an interval of α . This setting allows us to gain extra flexibility to study different problems. In [26], they considered $H(x, y) = x^2 + y^2$, which represents the pinned distance set at the origin of $K_1 \times K_2$. They showed

$$\Delta_0(K_1 \times K_2) = \{\sqrt{x^2 + y^2} : (x, y) \in K_1 \times K_2\}$$

has non-empty interior if $\tau(K_1)\tau(K_2) > 1$. This offers us a connection to the celebrated Falconer's distance set conjecture. McDonald and Taylor further generalized this result to distance sets determined by trees [19]. Jiang also studied the result under similar settings, but with more Cantor sets and he obtained conditions for which $H(K_1, K_2)$ is entirely an interval [15]. 1.2. Main Contribution of this paper. In the aforementioned results, they are mostly on \mathbb{R}^1 and a handful of them touches on \mathbb{R}^d . Moreover, there is no conclusion about the arithmetic sum or continuous C^1 image of Cantor sets if one of the Cantor sets has Newhouse thickness zero or other notions of thickness is zero. In this paper, we will prove a version of these types of theorems that will work for all Cantor sets regardless of dimensions and thickness and we believe that this is the first type of such theorems that work for all Cantor sets. Adopting the setting by Simon and Taylor [26], we have the following theorem:

Theorem 1.1. Let $N \geq 1$ and $\Lambda \subset \mathbb{R}^N$ be a set with interior and let $\alpha_0 \in \Lambda^\circ$. Let $Q_1 \subset Q_2 \subset \mathbb{R}^d$ and $Q_1^\circ \neq \emptyset$. Let H be a C^1 function on $\Lambda \times Q_1 \times Q_2$ such that the Jacobian on Q_2 is invertible (see (5.11) for the precise definition). Then for all Cantor set $K_1 \subset Q_1$, there exists a Cantor set $K_2 \subset Q_2$ and a cube centered at α_0 , $\mathbb{Q}_{\epsilon}(\alpha_0)$ ($\epsilon > 0$) such that

$$\left(\bigcap_{\alpha\in\mathbf{Q}_{\epsilon}(\alpha_{0})}H(\alpha,K_{1},K_{2})\right)^{\circ}\neq\varnothing.$$

As one direct application, we let $\mathsf{GL}_d(\mathbb{R})$ to be the set of all $d \times d$ invertible matrices and we embed it naturally on \mathbb{R}^{d^2} as an open subset. In other words, $\mathsf{GL}_d(\mathbb{R})$ is endowed with the Frobenius norm: $||T||_F = \sqrt{\sum_{i,j=1}^d |T_{ij}|^2}$ and $T = (T_{ij})_{1 \le i,j \le d}$. Consider the map $H : \mathsf{GL}_d(\mathbb{R}) \times \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}^d$ defined by

$$H(T, x, y) = Tx + y.$$

Then the Jacobians on x and y are respectively T and the identity transformation, which are invertible. Hence, we obtain the following corollary immediately from Theorem 1.1.

Corollary 1.2. Let K_1 be a Cantor set on \mathbb{R}^d and let $T_0 \in GL_d(\mathbb{R})$. Then there a exists another Cantor set K_2 and $\epsilon > 0$ such that

$$\left(\bigcap_{\{T:\|T-T_0\|_F<\epsilon\}} (TK_1+K_2)\right)^{\circ}\neq\emptyset.$$

We can prove this theorem more generally by considering invertible C^1 functions, not just invertible linear transformations. To this end, we endow the space of all continuously differentiable mapping from \mathbb{R}^d to \mathbb{R}^d , $C^1(\mathbb{R}^d)$, by the topology of local uniform convergence, which turns $C^1(\mathbb{R}^d)$ into a separable metric space, denoted by $(C^1(\mathbb{R}^d), \mathbf{d})$. Inside, we let $C^1_{inv}(\mathbb{R}^d)$ be all the local diffeomorphisms on \mathbb{R}^d i.e. the functions whose Jacobian is invertible at all points (See Section 4 for more details). This is our main theorem under this general setting. **Theorem 1.3.** Let K_1 be a Cantor set in \mathbb{R}^d and let $g_0 \in C^1_{inv}(\mathbb{R}^d)$. Then there exists $\varepsilon > 0$ and a Cantor set K_2 such that

(1.1)
$$\left(\bigcap_{g\in\mathsf{B}^{\mathsf{d}}_{\varepsilon}(g_0)}(g(K_1)+K_2)\right)^{\circ}\neq\varnothing.$$

Corollary 1.2 and Theorem 1.3 show that for all Cantor sets K_1 , we can always find another K_2 such that (**Qu**) is true. We notice that if K_1 has positive Newhouse thickness, then we can find another K_2 so the product of the thickness is larger than 1, then (**Qu**) is holds for the pair (K_1, K_2). If K_1 has positive Hausdorff dimension, we were informed by Shmerkin that we can use fractal percolation to construct the Cantor set K_2 so that (**Qu**) holds for the pair (K_1, K_2) ([25, Theorem 13.1]). Therefore, the novelty of this theorem lies on the Cantor sets that are both Newhouse thickness and Hausdorff dimension zero, which we refer to these Cantor sets as *very thin*, as our title suggested.

The idea of the proof will rely on what we call the *containment lemma*, namely, given a certain separation data of the first Cantor set, we can construct the other Cantor set whose convex hull contains the first one and two Cantor sets will intersect. Moreover, the intersection is robust under small C^1 -perturbation. This can be easily achieved on \mathbb{R}^1 using the gap data (Lemma 2.1). In high dimension, the data will yield useful information only if K is *non-degenerate* or *uniformly non-degenerate* (Theorem 3.4). However, we can show that these are minor assumptions, since we can apply an orthogonal transformation to the Cantor set so that the resulting Cantor is uniformly non-degenerate (See Section 6 and 7). Therefore, upon orthogonal transformations, we still have our containment lemma.

These results yield many applications about continuous image of Cantor sets. We will discuss two main examples in Section 8. Concerning the sum set in Theorem 1.3, we will see this is related to the Erdős similarity problem. Indeed, we will be able to solve a very general topological version of the Erdős similarity problem (Theorem 8.1), generalizing a previous result in [10]. Similar to the results by Simon and Taylor [26], we are also able to say something about the pinned distance set of Cantor sets of Hausdorff dimension 1. We will show that there exists a Cantor set on \mathbb{R}^2 of dimension 1, whose pinned distance set, hence its distance set, possesses an interior. The same also holds true for \mathbb{R}^d where d is even. Indeed, such sets trivially exist if it is connected such as a line segment. However, the fact that our set is totally disconnected seems to be new. Further applications can also be made if we consider other C^1 functions similar to [15], but we are not going to write in full details in this paper.

We will organize our paper as follows: In Section 2, we will prove our results on \mathbb{R}^1 , which is much simpler. We will then prove the containment lemma on \mathbb{R}^d in Section

3 for non-degenerate Cantor sets. In Section 4, we will introduce the uniformly nondegenerate Cantor sets and the topology of C^1 function. In Section 5, we will prove our main theorem, Theorem 1.1 and Theorem 1.3. In Sections 6 and 7, we will study those non-degenerate and uniformly non-degenerate Cantor sets respectively. We will show that all will become uniformly non-degenerate if we apply an orthogonal transformation. Finally, in Section 8, we will discuss the applications of our theory to topological Erdős similarity problem and the distance set problems.

2. Containment lemma on \mathbb{R}^1

We first define our notation. We let $\Sigma^0 = \{\emptyset\}$, $\Sigma^n = \{0, 1\}^n$ and $\Sigma^* = \bigcup_{n=1}^{\infty} \Sigma^n$. We will naturally concatenate σ and σ' in Σ' and denote it by $\sigma\sigma'$. We say that σ is a **descendant** of σ' (or σ' is an **ancestor** of σ) if $\sigma = \sigma'\tilde{\sigma}$ for some $\tilde{\sigma}$. If $\tilde{\sigma}$ has length 1, σ is a **child** of σ' .

We describe a Cantor set in \mathbb{R} as a binary Cantor set. Let K be a Cantor set and let $I_{\emptyset}(K)$ be the convex hull of K. Then the complement of the K in $I_{\emptyset}(K)$ is a countable union of disjoint bounded open intervals, which we call (**bounded**) gaps. We first write them as G_n , $n = 1, 2, \cdots$ with $|G_1| \ge |G_2| \ge \cdots$ and if they are of the same length, we will enumerate them from the leftmost one. We now define

$$I_{\varnothing}(K) = I_0(K) \cup U_{\emptyset}(K) \cup I_1(K), \text{ and } U_{\emptyset}(K) := G_1$$

where we also define by convention that $I_0(K)$ is on the left hand side of G_1 and $I_1(K)$ is on the right of G_1 . Suppose that $I_{\sigma}(K)$, $\sigma \in \Sigma^*$ has been defined. We will let $U_{\sigma}(K)$ to be the largest open interval from the collection of $\{G_n\}_{n\in\mathbb{N}}$ that is in the I_{σ} . Then we define

$$I_{\sigma}(K) = I_{\sigma 0}(K) \cup U_{\sigma}(K) \cup I_{\sigma 1}(K).$$

In this way, the Cantor set K can be represented as

$$K = \bigcup_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma^n} I_{\sigma}(K).$$

A Cantor set is called a **centrally symmetric Cantor set** if for all $n \in \mathbb{N}$, $|U_{\sigma}(K)|$ are equal for all $\sigma \in \Sigma^n$. If we let ℓ_n be the length of the gap at the n^{th} stage, i.e. $\ell_n = |U_{\sigma}(K)|$ then

(2.1)
$$\ell_n < |I_{\sigma}| = \frac{|I_{\emptyset}(K)| - \ell_0 - 2\ell_1 - \dots - 2^{n-1}\ell_{n-1}}{2^n} \quad \forall \sigma \in \Sigma^n.$$

Conversely, if we are given a sequence $0 < \ell_n < 1$ that satisfies (2.1), then we can always construct a centrally symmetric Cantor set whose *n*-th stage gap are of length ℓ_n . We also recall that K is a **fat Cantor set** if K has positive Lebesgue measure. It happens if and only if

$$|I_{\emptyset}(K)| - \sum_{n=1}^{\infty} \sum_{\sigma \in \Sigma^n} |I_{\sigma}(K)| > 0.$$

Lemma 2.1. (Containment lemma) Let K be a Cantor set in \mathbb{R} . Suppose that \widetilde{K} is a Cantor set in \mathbb{R} with $I_{\emptyset}(K) \subset I_{\emptyset}(\widetilde{K})$ and

(2.2)
$$\max\{|U_{\sigma}(\tilde{K})|: \sigma \in \Sigma^n\} < \min\{|U_{\sigma}(K)|: \sigma \in \Sigma^n\} \quad \forall n \in \mathbb{N}.$$

Then, $K \cap \widetilde{K} \neq \emptyset$.

Proof. Since K and \widetilde{K} are compact, it suffices to show that there exists a sequence $(\alpha_n, \widetilde{\alpha}_n) \in K \times \widetilde{K}$ such that $\lim_{n \to \infty} |\alpha_n - \widetilde{\alpha}_n| = 0$. To show this, we first prove the following claim.

Claim. For each $n \in \mathbb{N}$, there exists $\sigma_n, \sigma'_n \in \Sigma^n$ where σ_n is a child of σ_{n-1} and σ'_n is a child of σ'_{n-1} , such that $I_{\sigma_n}(K) \subset I_{\sigma'_n}(\widetilde{K})$.

Assuming the claim, we can just take $\alpha_n \in K \cap I_{\sigma_n}(K)$ and $\alpha'_n \in I_{\sigma'_n}(\tilde{K})$. Since $|I_{\sigma'_n}(K)| \to 0$ as $n \to \infty$ and $|\alpha_n - \alpha'_n| \leq |I_{\sigma'_n}(K)|$, this finishes the proof.

We now prove the above claim by induction. Note that the base case is true since $I_{\emptyset}(K) \subset I_{\emptyset}(\tilde{K})$ by assumption. For the induction hypothesis, suppose that there exist σ_n and σ'_n such that $I_{\sigma_n}(K) \subset I_{\sigma'_n}(\tilde{K})$ where σ_n and σ'_n form a chain of children up to n. We now proceed to the induction step of n + 1.

Let us write $I_{\sigma_n}(K) = [a_0, b_0]$ and $I_{\sigma'_n}(\widetilde{K}) = [c_0, d_0]$. Then let us denote the next children $I_{\sigma_n 0}(K)$, $I_{\sigma_n 1}(K)$, $I_{\sigma'_n 0}(\widetilde{K})$, $I_{\sigma'_n 1}(\widetilde{K})$ respectively by $[a_0, a], [b, b_0], [c_0, c]$ and $[d, d_0]$, so that

$$I_{\sigma_n}(K) = [a_0, a] \cup U_{\sigma_n}(K) \cup [b, b_0], \ I_{\sigma'_n}(\tilde{K}) = [c_0, c] \cup U_{\sigma'_n}(\tilde{K}) \cup [d, d_0].$$

By induction hypothesis $c_0 \leq a_0$ and $b_0 \leq d_0$. From our assumption,

$$d - c = |U_{\sigma_n}(\widetilde{K})| < \min\{|U_{\sigma_n}(K)| : \sigma_n \in \Sigma^n\} \le |U_{\sigma_n}(K)| = b - a.$$

This means that we must have a < c or d < b. In the first case, $[c_0, c] \supset [a_0, a]$ meaning that $I_{\sigma'_n 0}(\widetilde{K}) \supset I_{\sigma_n 0}(K)$, while in the second case, we have $I_{\sigma'_n 1}(\widetilde{K}) \supset I_{\sigma_n 1}(K)$. This shows that the claim holds true for n + 1 and hence completes the whole proof. \Box

- Remark 2.2. (1) We note that there always exists \tilde{K} satisfying (2.2), since by taking $\ell_n > 0$ small enough, we can always construct a symmetric Cantor set whose n^{th} -stage gap length satisfies both (2.2) and (2.1).
 - (2) For convenience, we arrange the gaps in decreasing lengths. Indeed, the proof did not use this fact. If we successively remove a gap, not necessarily the largest gap, from $I_{\sigma}(K)$, we obtain another representation of the Cantor set and we can still find \tilde{K} for the containment lemma to hold in this situation.

(3) Unfortunately, the resulting Cantor set \widetilde{K} is always a fat Cantor set. It is because the Lebesgue measure of \widetilde{K} is given by

$$|I_{\emptyset}(\widetilde{K})| - \sum_{n=1}^{\infty} \sum_{\sigma \in \Sigma^n} |I_{\sigma}(\widetilde{K})| > |I_{\emptyset}(K)| - \sum_{n=1}^{\infty} \sum_{\sigma \in \Sigma^n} |I_{\sigma}(K)| \ge 0$$

where we used the fact that $I_{\emptyset}(K) \subset I_{\emptyset}(\widetilde{K})$ and (2.2) in the inequality above.

The following corollary gives a stable perturbation under small scaling for the containment lemma.

Corollary 2.3. If K is a Cantor set in \mathbb{R} , then there exists a Cantor set \widetilde{K} in \mathbb{R} and $\delta > 0$ such that $K \cap (\lambda \widetilde{K} + t) \neq \emptyset$ for all $\lambda \in (1 - \delta, 1 + \delta)$ and for all $t \in (-\delta, \delta)$. In particular, $K + \lambda \widetilde{K}$ has a non-empty interior.

Proof. We take the Cantor set \widetilde{K} such that $I_{\emptyset}(\widetilde{K})$ strictly contain $I_{\emptyset}(K)$ and satisfying (2.2) with right hand side replaced by $1/2 \cdot \min\{|U_{\sigma}(K)| : \sigma \in \Sigma^n\}$. Then, $K \cap \widetilde{K} \neq \emptyset$. Note that $I_{\sigma}(\lambda \widetilde{K} + t) = \lambda I_{\sigma}(\widetilde{K}) + t$. By choosing δ sufficiently small, the conditions $I_{\emptyset}(K) \subset I_{\emptyset}(\lambda \widetilde{K} + t)$ and (2.2) holds true with \widetilde{K} replaced by $\lambda \widetilde{K} + t$ for all $\lambda \in (1 - \delta, 1 + \delta)$ and $t \in (-\delta, \delta)$. The containment lemma (Lemma 2.1) gives our desired conclusion.

2.1. Interior for general C^1 functions. We are going to follow the setup obtained by Simon and Taylor. We will denote by $I_{\delta}(a)$ the closed interval of length δ and centered at a. Let Λ be a compact parameter interval with center α_0 and let J_1 and J_2 be two compact intervals. We are also given $H \in C^1(\Lambda \times J_1 \times J_2)$ such that $H(\alpha, x, y)$ has non-vanishing partial derivatives H_x and H_y on $\Lambda \times J_1 \times J_2$.

Fix
$$(u_1, u_2) \in (J_1 \times J_2)^\circ$$
. Set $c_0 = H(\alpha_0, u_1, u_2)$ and let $\delta_0 > 0$ so that

(2.3) $\mathbf{I}_{\delta_0}(\alpha_0) \times \mathbf{I}_{\delta_0}(u_1) \times \mathbf{I}_{\delta_0}(u_2) \subset (\Lambda \times J_1 \times J_2)^{\circ}.$

Let also

$$S = \mathbf{I}_{\delta_0}(c_0) \times \mathbf{I}_{\delta_0}(\alpha_0) \times \mathbf{I}_{\delta_0}(u_1) \times \mathbf{I}_{\delta_0}(u_2).$$

and $F: S \to \mathbb{R}$ defined by

$$F(c, \alpha, x, y) = H(\alpha, x, y) - c.$$

Note that $F_y = H_y \neq 0$ by assumption and we have $F(c_0, \alpha_0, u_1, u_2) = 0$. By the implicit function theorem, we can find a $\delta_1 \in (0, \delta_0)$, a closed set

(2.4) $Z = \mathbf{I}_{\delta_1}(c_0) \times \mathbf{I}_{\delta_1}(\alpha_0) \times \mathbf{I}_{\delta_1}(u_1)$

and a function $G: Z \to I_{\delta_0}(u_2)$ such that $G(c_0, \alpha_0, u_1) = u_2$ and

(2.5)
$$H(\alpha, x, G(c, \alpha, x)) = c$$

We define

(2.6)
$$g_{c,\alpha}(x) = G(c,\alpha,x), \ (c,\alpha) \in \mathbf{I}_{\delta_1}(c_0) \times \mathbf{I}_{\delta_1}(\alpha_0)$$

The following lemma was proved by Taylor and Simon.

Lemma 2.4. For each i = 1, 2, let $K_i \subset J_i$ be two Cantor sets. Suppose that

$$g_{c,\alpha}(K_1) \cap K_2 \neq \emptyset$$
 for all $(c,\alpha) \in I_{\delta_1}(c_0) \times I_{\delta_1}(\alpha_0)$.

Then

$$\mathbf{I}_{\delta_1}(c_0) \subset \bigcap_{\alpha \in \mathbf{I}_{\delta_1}(\alpha_0)} H(\alpha, K_1, K_2).$$

Proof. We prove it here for the sake of completeness. For each $(c, \alpha) \in I_{\delta_1}(c_0) \times I_{\delta_1}(\alpha_0)$, our assumption implies that there exist $k_1 \in K_1$ and $k_2 \in K_2$ such that $g_{c,\alpha}(k_1) = k_2$. Hence, $H(\alpha, k_1, k_2) = H(\alpha, k_1, g_{c,\alpha}(k_1)) = c$. This shows that $c \in H(\alpha, K_1, K_2)$, and we obtain our desired conclusion.

We also need a derivative estimate.

Lemma 2.5. There exists $\eta > 0$ such that

$$|g'_{c,\alpha}(x)| \ge \eta > 0, \ \forall (c,\alpha,x) \in N.$$

Proof. From (2.5),

$$g'_{c,\alpha}(x) = -\frac{H_x(\alpha, x, G(c, \alpha, x))}{H_u(\alpha, x, G(c, \alpha, x))}.$$

From our assumption, H_x and H_y are continuous functions that never vanish. Hence, they never change sign. This implies that $g'_{c,\alpha}$ never changes sign and $g_{c,\alpha}$ is strictly monotone. Furthermore, since H_x, H_y are continuous functions on the compact set Z, there exists a $\eta > 0$ such that our desired conclusion holds.

Using the containment lemma, our main theorem is the following:

Theorem 2.6. Let Λ , J_1 , J_2 be compact intervals, $H \in C^1(\Lambda \times J_1 \times J_2)$ with non vanishing partial derivatives H_x and H_y , and $J_1 \subset J_2$ and α_0 be the center of Λ . Then for all Cantor sets $K_1 \subset J_1$, there exists a Cantor set $K_2 \subset J_2$ and a nondegenerate interval $I_{\tilde{\epsilon}}(\alpha_0)$ such that

$$\left(\bigcap_{\alpha\in I_{\tilde{\epsilon}}(\alpha_0)} H(\alpha, K_1, K_2)\right)^{\circ} \neq \emptyset.$$

Proof. Let u_1 be a right-endpoint of a gap of K_1 and choose δ_0 such that $\mathbf{I}_{\delta_0}(u_1) \cap K_1$ is a non-degenerate Cantor set inside J_1° and (2.3) is fulfilled. Let $u_2 = u_1 \in J_2$. We also set $c_0 = H(\alpha_0, u_1, u_2)$. Using our setting of H, we can find a function $G : Z \to \mathbb{R}$ satisfying (2.5) and $G(c_0, \alpha_0, u_1) = u_1$. Recall also that Z is of the form (2.4) for some $\delta_1 \in (0, \delta_0)$. By choosing δ_1 smaller if necessary, we can assume that $\widetilde{K_1} = \mathbf{I}_{\delta_1}(u_1) \cap K_1$ is a Cantor set inside $K_1 \cap J_1^{\circ}$. Let $g_{c,\alpha}(x) = G(c, \alpha, x)$. Our goal is to construct $K_2 \subset J_2$ such that the condition in Lemma 2.4 holds for $\widetilde{K_1}$. We now represent $\widetilde{K_1}$ as

$$\widetilde{K}_1 = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma^n} I_{\sigma}(\widetilde{K}_1)$$

and $U_{\sigma}(C)$ are the gaps of $\widetilde{K_1}$. For $n = 0, 1, 2, \cdots$, the n^{th} stage gaps are the collection of gaps $U_{\sigma}(\widetilde{K_1})$ where $\sigma \in \Sigma^n$. For all $(c, \alpha) \in I_{\delta_1}(c_0) \times I_{\delta_1}(\alpha_0)$, the collection of all the n^{th} gaps of $g_{c,\alpha}(\widetilde{K})$ are

$$\{g_{c,\alpha}(U_{\sigma}(\widetilde{K}_1)): \sigma \in \Sigma^n\}$$

We now claim that there exists $\eta_n > 0$ such that for all $(c, \alpha) \in I_{\delta_1}(c_0) \times I_{\delta_1}(\alpha_0)$.

(2.7)
$$\min\{|g_{c,\alpha}(U_{\sigma}(K_1))|: \sigma \in \Sigma^n\} \ge \eta_n$$

To see this claim, we let for a given $\sigma \in \Sigma^n$, $U_{\sigma}(\widetilde{K}_1) = (t_1, t_2)$. Then, by the mean value theorem,

$$|g_{c,\alpha}(U_{\sigma}(K_1))| = |g_{c,\alpha}(t_2) - g_{c,\alpha}(t_1)| = |g'_{c,\alpha}(\zeta)||t_2 - t_1|.$$

for some $\zeta \in (t_1, t_2)$. Using Lemma 2.5,

$$|g_{c,\alpha}(U_{\sigma}(K_1))| \ge \eta \cdot |U_{\sigma}(K_1)|.$$

As η is independent of c and α and there are only finitely many n^{th} stage gaps, we can take $\eta_n = \eta \cdot \min\{|U_{\sigma}(\widetilde{K}_1)| : \sigma \in \Sigma^n\}$. This justifies (2.7).

Next, we note that all $g_{c,\alpha}(\widetilde{K_1})$ are inside $I_{\delta_0}(u_1)$. We construct a centrally symmetric Cantor set K_2 whose convex hull is $I_{\delta_0}(u_1)$ and the length of the n^{th} stage gaps are less than η_n . By the containment lemma (Lemma 2.1), $g_{c,\alpha}(\widetilde{K_1}) \cap K_2 \neq \emptyset$ for all $(c,\alpha) \in I_{\delta_1}(c_0) \times I_{\delta_1}(\alpha_0)$. In particular, the condition for Lemma 2.4 is fulfilled. This completes the proof.

3. Containment lemma in \mathbb{R}^d

3.1. Nested representation of Cantor sets. In this section, we are going to generalize the containment lemma to higher dimensions. Since there is no canonical way to represent a Cantor set on \mathbb{R}^d , we will consider the following way to represent our Cantor sets.

Definition 3.1. A nested representation of a Cantor set *C* is a countable family of compact connected sets $\mathcal{R} = \{R_{\sigma} : \sigma \in \bigcup_{k=1}^{\infty} \Sigma^k\}$ defined inductively as follows:

(1) $C \subset R_{\emptyset}$ and

 $R_{\emptyset} \cap C = (R_1 \cap C) \cup \cdots (R_{m_{\emptyset}} \cap C)$

for some integer $m_{\emptyset} \geq 2$ where $R_1 \cdots, R_{m_{\emptyset}}$ are disjoint sets contained in R_{\emptyset} . We will denote by $\Sigma^1 = \{1, \cdots, m_{\emptyset}\}$ (2) Suppose that Σ^k has been defined and R_{σ} has been defined for all $\sigma \in \Sigma^k$. We define $R_{\sigma 1} \cdots R_{\sigma m_{\sigma}}$ be disjoint compact connected sets contained in R_{σ} such that

$$R_{\sigma} \cap C = (R_{\sigma 1} \cap C) \cup \dots \cup (R_{\sigma m_{\sigma}} \cap C)$$

- and we define $\Sigma^{k+1} = \bigcup_{\sigma \in \Sigma^k} \{\sigma\} \times \{1, \cdots, m_\sigma\}.$
- (3) $\lim_{k \to \infty} \max_{\sigma \in \Sigma^k} (\operatorname{diam}(R_{\sigma})) = 0.$

In this nested representation,

(3.1)
$$C = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma^n} R_{\sigma}$$

Moreover, for all $x \in C$, there exists a unique sequence of compact connected sets $R_{\sigma_1}, R_{\sigma_1\sigma_2}, \cdots$ such that

$$\{x\} = \bigcap_{n=1}^{\infty} R_{\sigma_1 \cdots \sigma_n}$$

and $\sigma_{i+1} \in \{1, \cdots, m_{\sigma_1 \cdots \sigma_i}\}$ for all $i \ge 1$.

We will adopt the following multi-index notations: Let $\Sigma^* = \bigcup_{k=1}^{\infty} \Sigma^k$ and Σ^{∞} be the collection of all infinite paths $\sigma_1 \sigma_2 \cdots$ where $\sigma_1 \cdots \sigma_n \in \Sigma^n$ for all $n \in \mathbb{N}$. For each $\sigma \in \Sigma^n$, we define

$$\Sigma^k(\sigma) = \{j_1 \cdots j_k : 1 \le j_1 \le m_\sigma, \ 1 \le j_r \le m_{\sigma j_1 \cdots j_{r-1}}\}.$$

We can naturally think Σ^* as a tree starting from the vertex \emptyset that branches out to m_{\emptyset} many edges, where we label $\{1, \dots, m_{\sigma}\}$ as the vertices and each vertex σ further branches out m_{σ} many edges with $\sigma 1, \dots, \sigma m_{\sigma}$ as the vertices. Σ^{∞} are the infinite paths in the tree.

Remark 3.2. (1) For most of our study of Cantor sets, R_{σ} are chosen to be cubes or balls. However, we only require our R_{σ} to be a compact connected set.

(2) Let

 $\mathcal{Q}_n = \{ [0, 2^{-n}]^d + 2^{-n}t : t \in \mathbb{Z}^d \}$

be the set of all axis-parallel dyadic cubes of side length 2^{-n} . For a given Cantor set $C \subset \mathbb{R}^d$, define

$$C_n = \bigcup \{ Q \in \mathcal{Q}_n : Q \cap C \neq \emptyset \},\$$

Note that C_n is a nested decreasing sequence of sets and $C = \bigcap_{n=1}^{\infty} C_n$. For each C_n , we can decompose it into its connected components. Then these connected components provide a natural nested representation for the Cantor set C.

We will denote by $\pi_j : \mathbb{R}^d \to \mathbb{R}$ the orthogonal projection onto the x_j -axis. For compact subsets $A, B \subset \mathbb{R}^d$, define

$$d_{\min}(A,B) = \min_{j=1,\cdots,d} d\left(\pi_j(A), \pi_j(B)\right).$$

where $d(E, F) = \min\{|x - y| : x \in E, y \in F\}$. $d_{\min}(A, B) > 0$ means that the two sets cannot overlap in any axis directions.

Definition 3.3. Let *C* be a Cantor set and $C = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma^n} R_{\sigma}$ be a nested representation of *C*. Let also $\Sigma_d^n = \{1, \dots, d+1\} \times \dots \times \{1, \dots, d+1\}$ (*n* times) and $\Sigma_d^* = \bigcup_{n=1}^{\infty} \Sigma_d^n$.

(1) We say that R_{σ} is **non-degenerate** if there exists $k \geq 1$ and there exists $\{A_1, \dots, A_{d+1}\} \subset \{R_{\sigma\sigma'} : \sigma' \in \Sigma^k(\sigma)\}$ such that

$$(3.2) d_{\min}(A_i, A_j) > 0$$

for all $i \neq j \in \{1, \dots, d+1\}$. Otherwise, R_{σ} will be called **degenerate**. We will call $\{A_i\}$ the **non-degenerate connected components** in R_{σ}

(2) C is **non-degenerate** if there exists a sub-Cantor set \widehat{C} generated by a sub-collection

$$\{\mathbf{R}_{\sigma}: \sigma \in \Sigma_d^*\}$$

in \mathcal{R} and it can be nest-represented as

(3.3)
$$\widehat{C} = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma_d^n} \mathbf{R}_{\sigma}$$

and $\{\mathbf{R}_{\sigma 1}, \cdots, \mathbf{R}_{\sigma (d+1)}\}\$ are the non-degenerate components of \mathbf{R}_{σ} for all $\sigma \in \Sigma_d^n$ and $n \geq 1$.

3.2. Containment lemma for non-degenerate Cantor sets. In this section, we will prove the containment lemma under the assumption that C is non-degenerate.

Theorem 3.4. (Containment Lemma on \mathbb{R}^d) Let C be a non-degenerate Cantor set in \mathbb{R}^d that contains a sub-Cantor set \widehat{C} as in (3.3). Let

(3.4)
$$d_k = \min_{\sigma \in \Sigma_d^k} \left\{ d_{\min} \left(\mathbf{R}_{\sigma p}, \mathbf{R}_{\sigma q} \right) : 1 \le p < q \le d+1 \right\}$$

Define a Cantor set $K = \overbrace{K_0 \times \cdots \times K_0}^{d \text{-times}} \subset \mathbb{R}^d$ such that

- (K0). I is any closed interval such that $conv(K) = I \times \cdots \times I \supset conv(C)$.
- (K1). K_0 is a centrally symmetric Cantor set on \mathbb{R}^1 obtained from the closed interval I (which we determine in (K0)) and removing an open interval of length $g_k < d_k$, in each remaining closed interval from the center at the k-th stage.

Then $C \cap K \neq \emptyset$.

Proof. For the centrally symmetric Cantor set $K_0 \subset \mathbb{R}^1$, as in the previous section, we can write it as

$$K_0 = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma_2^n} I_{\sigma}(K_0)$$

where $\Sigma_2 = \{0, 1\}$ and $I_{\sigma}(K_0)$ has the same length for all $\sigma \in \Sigma_2^n$. We write K as

$$K = \overbrace{K_0 \times \cdots \times K_0}^{d \text{-times}} = \bigcap_{n=1}^{\infty} \bigcup_{j=1}^{2^{nd}} S_{n,j}$$

where $S_{n,j}$ are cubes of the form $I_{\sigma_1}(K_0) \times \cdots \times I_{\sigma_d}(K_0)$ with $(\sigma_1, \cdots, \sigma_d) \in \Sigma_2^n \times \cdots \times \Sigma_2^n$, so there are 2^{nd} such cubes. To complete the proof, it suffices to establish the following claim which allows us to find a sequence of points from C and K converging to the intersections:

Claim. For all $n \in \mathbb{N}$, there exists $\sigma \in \Sigma_d^n$ and $S_{n,j}$ such that $\mathbb{R}_{\sigma} \subset S_{n,j}$.

Proof of claim. We proceed the proof by induction. The base case holds automatically by (K0) since $\mathbb{R}_{\emptyset} \subset S_{\emptyset} = I \times ... \times I$. Suppose that we have already constructed $\mathbb{R}_{\sigma} \subset S_{n,j}$ for some $\sigma \in \Sigma_d^n$ and $n \ge 1$. Let us write $\mathbb{S} = S_{n,j} = \prod_{k=1}^d [a_k, b_k]$ where $b_k - a_k$ are all the same length. By the requirement of centrally symmetric Cantor sets, \mathbb{S} is split up into 2^d many cubes $\mathbb{S}_1, \cdots \mathbb{S}_{2^d}$ on the corner so that

$$\mathbf{S} \setminus (\mathbf{S}_1 \cup \cdots \cup \mathbf{S}_{2^d}) = \bigcup_{k=1}^d G_k,$$

where

$$G_k = [a_1, b_1] \times \dots \times \left[\frac{a_k + b_k}{2} - \frac{g_{n+1}}{2}, \frac{a_k + b_k}{2} + \frac{g_{n+1}}{2}\right] \times \dots \times [a_d, b_d].$$

Note that G_k is a slit of thickness g_{n+1} centered at $x_k = \frac{a_k + b_k}{2}$ and is parallel to the x_k coordinate plane.

We claim that there exists $\mathbb{R}_{\sigma p}$ and \mathbb{S}_q such that $\mathbb{R}_{\sigma p} \subset \mathbb{S}_q$. Suppose the claim is false. Then we have for all $i \in \{1, \dots, d+1\}$, $\mathbb{R}_{\sigma i} \cap \mathbb{S} \setminus \mathbb{S}_j \neq \emptyset$ for all $j = 1, \dots 2^d$. Hence, $\mathbb{R}_{\sigma i}$ intersects $\mathbb{S} \setminus (\mathbb{S}_1 \cup \dots \cup \mathbb{S}_{2^d}) = \bigcup_{k=1}^d G_k$. Therefore, there exists k = k(i)such that $\mathbb{R}_{\sigma i} \cap G_k \neq \emptyset$. By pigeonhole principle, there exists $i \neq i'$ such that $\mathbb{R}_{\sigma i}$ and $\mathbb{R}_{\sigma i'}$ both intersects at the same G_k . However, this implies that

$$d_{\min}(\mathtt{R}_{\sigma i}, \mathtt{R}_{\sigma i'}) < g_{n+1} < d_{n+1}$$

by (K1). This is a contradiction to the definition of d_{n+1} in (3.4). This justifies the claim and hence completes the proof.

Remark 3.5. (1) We remark that the Cantor set K that we constructed will have positive Lebesgue measure since we can choose g_k arbitrarily small. However, due to the centrally symmetric Cantor set construction, it still has no interior. (2) Not all Cantor sets are non-degenerate. For example, if we embed the middlethird Cantor set into an axis parallel line on \mathbb{R}^2 , then no matter how we generate a nested representation, we would not be able to find a set of three connected components that satisfy (3.2), since these components eventually converges to the line. On the other hand, if we rotate the Cantor set so that it is no longer parallel to the axes, it will be non-degenerate. We will prove in Section 6 that all Cantor sets are non-degenerate after some orthogonal transformations (See Theorem 6.5). Indeed, we can show that all degenerate Cantor sets must lie in countably many axe-parallel hyperplanes.

4. Uniformly non-degeneracy and topology of C^1 functions

We are interested in determining if the intersection of Cantor sets in the containment lemma is stable under small perturbation of C^1 functions. Indeed, the containment lemma is stable if C is perturbed only by scaling and translation using the same argument in Corollary 2.3. To make it work in all C^1 perturbation, we need some stronger assumption on the non-degeneracy and we need to endow C^1 functions with topology of local uniform convergence. We will define it in this section.

4.1. Uniformly non-degenerate Cantor sets.

Definition 4.1. We say that a Cantor set K in \mathbb{R}^d is **uniformly non-degenerate** (**u.n.d.** in short) if K contains a sub-Cantor set \widehat{K} whose nested representation $\widehat{K} = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma_d^n} \mathbb{R}_{\sigma}$ such that there exist constants $C_1, C_2 > 0$ satisfying the following property: for all $\sigma \in \Sigma_d^*$ and $(p, q, j) \in \{1, \dots, d+1\} \times \{1, \dots, d+1\} \times \{1, \dots, d\}$,

(4.1)
$$C_1 \|x - x'\| \le |\pi_j(x - x')| \le C_2 \|x - x'\| \quad \forall x \in \mathbb{R}_{\sigma p} \text{ and } x' \in \mathbb{R}_{\sigma q}.$$

Notice that (4.1) is also equivalent to saying that $|\pi_j(x - x')|$ and $|\pi_k(x - x')|$ are comparable to each other for all $j \neq k$ and for all $\sigma \in \Sigma_d^*$. Intuitively, a Cantor set cannot be lying up arbitrarily close to some axes-parallel hyperplanes. Therefore, after a suitable orthogonal transformation, the Cantor set should be uniformly nondegenerate. In the next theorem, we will see that this is the case.

Theorem 4.2. Let $K \subset \mathbb{R}^d$ be a Cantor set. Then there exists an orthogonal linear transformation O on \mathbb{R}^d such that the image $\mathsf{O}(K)$ is **u.n.d.** on \mathbb{R}^d .

The proof however requires us some careful writing, we will postpone it in Section 7 to ensure the continuity of proving our main theorems.

4.2. Topology of $C^1(\mathbb{R}^d)$. We let $\mathsf{GL}_d(\mathbb{R})$ be the set of all $d \times d$ invertible matrices A. If $T \in \mathsf{GL}_d(\mathbb{R})$, we write $(T_{ij})_{i,j=1}^d$ to be the matrix representation of T in the standard basis of \mathbb{R}^d . the Recall that the **operator norm** and the **Frobenius norm** are defined by

$$||T|| = ||T||_{op} = \sup_{x \neq 0} \frac{||Tx||}{||x||}, \ ||T||_F = \left(\sum_{i,j=1}^d |T_{ij}|^2\right)^{1/2}.$$

Using the singular value decomposition, ||T|| is the maximum of the singular value of T, while $||T||_F$ is the square root of the sum of the square of all singular values (see e.g. [14]). Hence, we know that

$$\frac{1}{\sqrt{d}} \|T\|_F \le \|T\| \le \|T\|_F.$$

We will write

$$B_{\delta}(A) = \{T \in \mathsf{GL}_d(\mathbb{R}) : \|T - A\|_F < \delta\}.$$

The following lemma ensures the invertibility in a neighborhood of Frobenius norm.

Lemma 4.3. Let $T_0 \in \mathsf{GL}_d(\mathbb{R})$ and $T : \mathbb{R}^d \to \mathbb{R}^d$ be a linear map. Then there exists $\delta > 0$ such that $||T - T_0||_F < \delta$ implies $T \in \mathsf{GL}_d(\mathbb{R})$.

Proof. We know that $T_0 \in \mathsf{GL}_d(\mathbb{R})$ implies that we can always find c > 0 such that $||T_0x|| \ge c||x||$ for all $x \in \mathbb{R}^d$. Note that

$$||Tx|| \ge ||T_0x|| - ||(T - T_0)x|| \ge c||x|| - ||T - T_0||_F ||x|| \ge (c - \delta)||x||.$$

Hence, if we choose $\delta < c$, then T is invertible.

Let $g : \mathbb{R}^d \to \mathbb{R}^d$ and let J_g be the Jacobian matrix of g. In other words, if we write $g(x) = (g_1(x), \cdots, g_d(x))$, where g_j are its component functions, we have

$$J_g(x) = \begin{bmatrix} \frac{\partial g_1}{\partial x_1} & \cdots & \frac{\partial g_1}{\partial x_d} \\ & \cdots & & \\ \frac{\partial g_d}{\partial x_1} & \cdots & \frac{\partial g_d}{\partial x_d} \end{bmatrix}.$$

We aim at proving the following theorem for uniformly non-degenerate Cantor sets. With a slight abuse of notation, we define

$$C^{1}(\mathbb{R}^{d}) = \left\{ g : \mathbb{R}^{d} \to \mathbb{R}^{d} : \frac{\partial g_{i}}{\partial x_{j}} \in C(\mathbb{R}^{d}) \ \forall i, j \in \{1, 2 \cdots, d\} \right\}.$$

We define also

$$C^{1}_{inv}(\mathbb{R}^{d}) = \{ g \in C^{1}(\mathbb{R}^{d}) : J_{g}(x) \in \mathsf{GL}_{d}(\mathbb{R}) \ \forall x \in \mathbb{R}^{d} \}.$$

 $C^1_{\mathsf{inv}}(\mathbb{R}^d)$ are usually referred as local diffeomorphisms. Clearly, if g(x) = Tx + t for some $T \in \mathsf{GL}_d(\mathbb{R})$ is an affine transformation, then $J_g(x) = T$ and it belongs to $C^1_{\mathsf{inv}}(\mathbb{R}^d)$. Before we begin the proof, we need to endow $C^1(\mathbb{R}^d)$ with the metric of local

uniform convergence. To do this, we write $\mathbb{R}^d = \bigcup_{n=1}^{\infty} R_n$ where R_n is an increasing union of compact subsets. Define

(4.2)
$$\|g\|_{n,\infty} = \max_{x \in R_n} |g(x)| + \left(\sum_{i,j=1}^d \left(\max_{x \in R_n} \left|\frac{\partial g_j}{\partial x_i}\right|\right)^2\right)^{1/2} \\ = \max_{x \in R_n} |g(x)| + \left\|\left(\max_{x \in R_N} \left|\frac{\partial g_j}{\partial x_i}\right|\right)_{1 \le i,j \le d}\right\|_F$$

and for each $f, g \in C^1(\mathbb{R}^d)$, define

$$\mathsf{d}(f,g) = \sum_{n=1}^{\infty} 2^{-n} \frac{\|f-g\|_{n,\infty}}{1+\|f-g\|_{n,\infty}}.$$

It is a routine check that $(C^1(\mathbb{R}^d), \mathsf{d})$ is a metric space. The ball under this metric will be denoted by $\mathsf{B}^{\mathsf{d}}_f(\delta) := \{g \in C^1(\mathbb{R}^d) : \mathsf{d}(f,g) < \delta\}$. Apart from the balls in the topology, we need the following sets: For each $f \in C^1(\mathbb{R}^d)$, $x_0 \in \mathbb{R}^d$ and $\delta \in (0, 1)$, we define

$$\mathcal{G}_{f,x_0,\delta} := \{ g \in C^1(\mathbb{R}^d) : \|J_g(x_0) - J_f(x_0)\|_F < \delta \text{ and } |g(x_0) - f(x_0)| < \delta \}$$

and for each $\eta > 0$,

$$\mathcal{G}_{f,x_0,\delta,\eta} = \mathcal{G}_{f,x_0,\delta} \cap \{g \in C^1(\mathbb{R}^d) : |x - x_0| < \eta \implies \|J_g(x) - J_g(x_0)\|_F < \delta\}.$$

We also let

$$\Delta(f, x_0, \delta) = \sup\{\eta > 0 : |x - x_0| < \eta \implies ||J_f(x) - J_f(x_0)||_F < \delta\}.$$

Note that since the map $x \to ||J_f(x) - J_f(x_0)||_F$ is continuous, $\Delta(f, x_0, \delta) > 0$.

Lemma 4.4. Let $K \subset \mathbb{R}^d$ be a compact set and let $x_0 \in K$. Then there exists $N_K \in \mathbb{N}$ such that for all $f \in C^1(\mathbb{R}^d)$ and $0 < \delta < 2^{-N_K-1}$,

 $\mathsf{B}^{\mathsf{d}}_{\delta}(f) \subset \mathcal{G}_{f,x_0,c_K\cdot\delta},$

where $c_K = 2^{N_K+1}$. Moreover, for all $\eta \leq \Delta(f, x_0, \delta)$,

$$\mathsf{B}^{\mathsf{a}}_{\delta}(f) \subset \mathcal{G}_{f,x_0,(2c_K+1)\cdot\delta,\eta}$$

Proof. Let N_K be such that $K \subset R_{N_K}$ and let $c_K = 2^{N_K+1}$. If $g \in \mathsf{B}^{\mathsf{d}}_{\delta}(f)$, then $d(f,g) < \delta$, which implies that

$$\|f - g\|_{N_{K,\infty}} < 2^{N_{K}} \delta \cdot (1 + \|f - g\|_{N_{K,\infty}}).$$

Hence, $||f - g||_{N_{K,\infty}} < \frac{2^{N_K}\delta}{1-2^{N_K}\delta}$. As $\delta < 2^{-N_K-1}$, this implies that

$$||f - g||_{N_K,\infty} < 2^{N_K + 1} \delta.$$

But since $K \subset R_{N_K}$, from (4.2), it follows immediately that $|f(x) - g(x)| < 2^{N_K + 1}\delta$ and $||J_g(x) - J_f(x)||_F < 2^{N_K + 1}\delta$ for all $x \in K$, completing the proof of the first part. For the second part, we notice that if $\eta < \Delta(f, x_0, \delta)$, then $|x - x_0| < \eta$ implies $||J_f(x) - J_f(x_0)||_F < \delta$. Since we already proved that $||J_g(x) - J_f(x)||_F < 2^{N_K+1}\delta$ for all $x \in K$, by a triangle inequality, if $|x - x_0| < \eta$, then

$$\begin{aligned} \|J_g(x) - J_g(x_0)\|_F &\leq \|J_g(x) - J_f(x)\|_F + \|J_f(x) - J_f(x_0)\|_F + \|J_f(x_0) - J_g(x_0)\|_F \\ &\leq (2c_K + 1)\delta. \end{aligned}$$

This shows the last inclusion.

Lemma 4.5. $(C^1(\mathbb{R}^d), \mathsf{d})$ is a separable metric space and is a Lindelöf space. *i.e.* every open cover has a countable sub-cover.

Proof. The separability is a routine exercise and it is known that a separable metric space must be a Lindelöf space [29, Theorem 16.9]. \Box

5. Proof of the main theorems

We will prove Theorem 1.1 and Theorem 1.3 in this section. Let us begin with the robustness of intersection around the identity map (Theorem 5.1) and then around the general maps (Theorem 5.2).

Theorem 5.1. Let K be a **u.n.d.** Cantor set in \mathbb{R}^d with the sub-Cantor set $\widehat{K} = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma_d^n} \mathbb{R}_{\sigma}$ satisfying (4.1) with constant $C_1, C_2 > 0$ and let $x_0 \in \widehat{K}$. Then, there exist $\delta > 0$ such that for all $\eta > 0$, there exists a Cantor set $\widetilde{K} = \widetilde{K}(\delta, \eta) \subset \mathbb{R}^d$ such that for all $g \in \mathcal{G}_{I,x_0,\delta,\eta}$ (I denotes the identity mapping),

$$g(K) \cap K \neq \emptyset.$$

Proof. By Lemma 4.3, there exists $\delta_0 > 0$ such that for all linear maps $T : \mathbb{R}^d \to \mathbb{R}^d$, $||T - I||_F < \delta_0$ implies $T \in \mathsf{GL}_d(\mathbb{R})$. Then, choose $\delta = \delta(C_1, C_2, d)$ such that

(5.1)
$$0 < \delta < \min\left\{\frac{1}{2(1+C_1C_2^{-1}(d-1))}, \delta_0\right\}$$

and choose $\lambda = \lambda(C_1, C_2, d, \delta)$ such that

(5.2)
$$0 < \lambda < 1 - 2(1 + C_1 C_2^{-1} (d - 1))\delta$$

Let $g \in \mathcal{G}_{I,x_0,\delta,\eta}$. Then $\|J_g(x_0) - I\|_F < \delta$, $|g(x_0) - x_0| < \delta$ and for all $|x - x_0| < \eta$, we have $\|J_g(x) - J_g(x_0)\|_F < \delta$. Therefore,

$$||J_g(x) - I||_F \le ||J_g(x) - J_g(x_0)||_F + ||J_g(x_0) - I||_F < 2\delta.$$

Let $T_{ij}(x) = \frac{\partial g_i(x)}{\partial x_j}$. From the definition of Frobenius norm, for each j and $i \neq j$ in $\{1, 2, \dots, d\},\$

(5.3)
$$T_{jj}(x) \in (1 - 2\delta, 1 + 2\delta) \text{ and } T_{ji}(x) \in (-2\delta, 2\delta)$$

whenever $|x - x_0| < \eta$.

16

We now focus on $x_0 \in \widehat{K}$. We can find a $\sigma_1 \sigma_2 \cdots \in \Sigma_d^\infty$ such that $x_0 \in \bigcap_{n=1}^\infty \mathbb{R}_{\sigma_1 \cdots \sigma_n}$. Hence, there exists \mathbb{R}_{σ} such that $\mathbb{R}_{\sigma} \subset B_{\eta}(x_0)$. The sub-Cantor set $\mathbb{R}_{\sigma} \cap \widehat{K}$ is also uniformly non-degenerate satisfying (4.1) with constants C_1, C_2 . For simplicity of notation, we may just assume $\widehat{K} \subset B_{\eta}(x_0)$. If not, we will replace $\mathbb{R}_{\sigma} \cap \widehat{K}$ by \widehat{K} .

Note that for each $g \in \mathcal{G}_{I,x_0,\delta}$, $g(\widehat{K}) = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma_d^n} g(\mathbb{R}_{\sigma})$ is a nested representation of the Cantor set $g(\widehat{K})$. Note that $g(\widehat{K})$ is a Cantor set since $g \in \mathsf{GL}_d(\mathbb{R})$. For each $n \in \mathbb{N}$, define

$$a_n = \min_{\sigma \in \Sigma_d^n} \left\{ d_{\min} \left(\mathbf{R}_{\sigma p}, \mathbf{R}_{\sigma q} \right) : 1 \le p < q \le d+1 \right\},$$

$$b_n(g) = \min_{\sigma \in \Sigma_d^n} \left\{ d_{\min} \left(g(\mathbf{R}_{\sigma p}), g(\mathbf{R}_{\sigma q}) \right) : 1 \le p < q \le d+1 \right\}.$$

Let $n \in \mathbb{N}$, $\sigma \in \Sigma_d^n$, and $(p, q, j) \in \{1, \dots, d+1\} \times \{1, \dots, d+1\} \times \{1, \dots, d\}$ Then, for all $x = (x_1, \dots, x_d) \in \mathbb{R}_{\sigma p}$, $x' = (x'_1, \dots, x'_d) \in \mathbb{R}_{\sigma q}$, using mean value theorem, we can find ζ_k such that

(5.4)
$$|g_j(x) - g_j(x')| = \left| \sum_{k=1}^d T_{jk}(\zeta_k) \cdot (x_k - x'_k) \right| \\ \geq |T_{jj}(\zeta_j)| |x_j - x'_j| - \sum_{k \neq j} |T_{jk}(\zeta_k)| |x_k - x'_k|$$

(5.5)

$$\geq (1 - 2\delta)|x_j - x'_j| - 2(d - 1)\delta C_1 C_2^{-1} |x_j - x'_j|$$

$$= (1 - 2(1 + C_1 C_2^{-1} (d - 1))\delta) |x_j - x'_j|$$

$$> \lambda |x_j - x'_j| \qquad (by (5.2)),$$

where in (5.4), we used the reverse triangle inequality, and in (5.5), we used (5.3) and the fact that for all $k \neq j$

$$|\pi_k(x - x')| \ge C_1 ||x - x'|| \ge \frac{C_1}{C_2} |\pi_j(x - x')|,$$

which follows from our assumption (4.1). Taking infimum over all $x \in \mathbb{R}_{\sigma p}, x' \in \mathbb{R}_{\sigma q}$, minimum over $j \in \{1, 2, \dots, d\}$, minimum over integers p, q such that $1 \leq p < q \leq d+1$, and minimum over $\sigma \in \Sigma_d^n$ in order, we have

$$b_n(g) \ge \lambda a_n, \quad \forall n \in \mathbb{N} \quad \forall g \in \mathcal{G}_{I,x_0,\delta}$$

Next, we claim that there exists C = C(d) such that for all $g \in \mathcal{G}_{I,x_0,\delta,\eta}$,

(5.6)
$$|g(x) - y| \le C, \ \forall x, y \in \widehat{K}.$$

Indeed, using the mean value theorem, there exists $\zeta_{ij} \in B_{\eta}(x_0)$ such that

$$g(x) - g(x_0) = A(x - x_0)$$

where A is the $d \times d$ matrix $(T_{ij}(\zeta_{ij}))_{1 \le i,j \le d}$. Hence,

$$|g(x) - g(x_0)| \le ||A||_F \cdot \eta.$$

Using (5.3), $||A||_F \leq (d(1+\delta)^2 + d(d-1)\delta)^{1/2} < \sqrt{4d+d^2} := C'$. Therefore, recalling that $\widehat{K} \subset B_\eta(x_0)$, we have

$$|g(x) - y| \le |g(x) - g(x_0)| + |x_0 - y| \le (C' + 1)\eta < C' + 1$$

for all $x, y \in \widehat{K}$. The proof of (5.6) is complete. Consequently, for all $g \in \mathcal{G}_{I,x_0,\delta,\eta}$, $g(\widehat{K}) \subset \{x : d(x,\widehat{K}) < C\}$, the *C*-neighborhood of \widehat{K} .

To finish the proof, we construct \widetilde{K} to be a centrally symmetric Cantor set in \mathbb{R}^d whose convex hull strictly contains the *C*-neighborhood of \widehat{K} . Then this convex hull contains all $g(\widehat{K})$ with $J_g(x_0) \in B_{\delta}(I)$ and $g(x_0) \in B_{\delta}(x_0)$. Then we require the n^{th} stage gaps of \widetilde{K} to be given by $\frac{\lambda a_n}{2}$. Then, the condition of the containment lemma in \mathbb{R}^d (Theorem 3.4) is satisfied by the pairs $(g(\widehat{K}), \widetilde{K})$ for all g under our condition, implying that

$$g(K) \cap \widetilde{K} \neq \emptyset.$$

We notice that there is no need to be in a neighborhood of the identity transformation, we can actually begin with any $g_0 \in C^1(\mathbb{R}^d)$.

Theorem 5.2. Let $g_0 \in C^1(\mathbb{R}^d)$ and K be a Cantor set in \mathbb{R}^d such that $g_0(K)$ is a Cantor set that is **u.n.d.** with the sub-Cantor set $g_0(\widehat{K})$ and let $x_0 \in \widehat{K}$ such that $J_{g_0}(x_0) \in \mathsf{GL}_d(\mathbb{R})$. Then, there exist $\delta > 0$ such that for all sufficiently small $\eta > 0$, one can find a Cantor set $\widetilde{K} \subset \mathbb{R}^d$ such that for all $g \in \mathcal{G}_{g_0,x_0,\delta,\eta}$,

$$g(K) \cap K \neq \emptyset.$$

Proof. Before we begin our proof, we first notice a linear algebra fact. Let $A, B \in GL_d(\mathbb{R})$. Note that

$$\frac{1}{\sqrt{d}} \|\mathsf{A}\mathsf{B}^{-1} - I\|_F \le \|\mathsf{A}\mathsf{B}^{-1} - I\| \le \|\mathsf{A} - \mathsf{B}\| \|\mathsf{B}^{-1}\| \le \|\mathsf{A} - \mathsf{B}\|_F \|\mathsf{B}^{-1}\|_F.$$

Consequently,

(5.7)
$$\mathsf{A} \in B_{\delta}(\mathsf{B}) \implies \mathsf{A}\mathsf{B}^{-1} \in B_{\sqrt{d}||\mathsf{B}^{-1}||_{F}\delta}(I).$$

We now begin the proof. By the inverse function theorem, there exists an open set W

containing x_0 and an open set V containing $g_0(x_0)$ such that $g_0: W \to V$ is invertible. We now apply Theorem 5.1 to $g_0(\widehat{K})$. There exists $\delta_0 > 0$ such that for all $\eta_0 > 0$, there exists a Cantor set \widetilde{K} such that for all $g \in \mathcal{G}_{I,g_0(x_0),\delta_0,\eta_0}$, we have

(5.8)
$$g(g_0(K)) \cap K \neq \emptyset$$

We take $\delta = \min\left\{\frac{\delta_0}{3\sqrt{d}(\|J_{g_0}(x_0)^{-1}\|_F+1)}, \delta_0\right\}$ and sufficiently small $\eta > 0$ so that

- (i) $B_{\eta}(x_0) \subset W$,
- (ii) if $|x x_0| < \eta$, then $||J_{g_0}(x)^{-1}||_F \le ||J_{g_0}(x_0)^{-1}||_F + 1$, and
- (iii) $||J_{g_0}(x) J_{g_0}(x_0)||_F < \delta.$

Such η exists since $||J_{g_0}(x)^{-1}||_F$ and $||J_{g_0}(x) - J_{g_0}(x_0)||_F$ are continuous functions and W is open. Finally, as $g_0(B_\eta(x_0))$ is open, we can take $\eta' < \eta_0$ so that

 $|y - g_0(x_0)| < \eta' \Longrightarrow y = g_0(x)$ for some x where $|x - x_0| < \eta$.

For all $g \in \mathcal{G}_{g_0,x_0,\delta,\eta}$, we have

- (1) $J_g(x_0) \in B_{\delta}(J_{g_0}(x_0)),$
- (2) $|g(x_0) g_0(x_0)| < \delta$,
- (3) $|x-x_0| < \eta \Longrightarrow ||J_g(x) J_g(x_0)||_F < \delta.$

We claim that $g \circ g_0^{-1} \in \mathcal{G}_{I,g_0(x_0),\delta_0,\eta'}$. As $\mathcal{G}_{I,g_0(x_0),\delta_0,\eta'} \subset \mathcal{G}_{I,g_0(x_0),\delta_0,\eta_0}$. This combined with (5.8) implies that $g(K) \cap \widetilde{K} \neq \emptyset$ for all $g \in G_{g_0,x_0,\delta,\eta}$, completing the proof.

It remains to justify the claim, which requires us to show that

(a) $J_{g \circ g_0^{-1}}(g_0(x_0)) \in B_{\delta_0}(I)),$ (b) $|g \circ g_0^{-1}(g_0(x_0)) - g_0(x_0)| < \delta_0,$ (c) $|y - g_0(x_0)| < \eta' \Longrightarrow ||J_{g \circ g_0^{-1}}(y) - I||_F < \delta_0.$

Proof of (a). (1) and (5.7) implies that we have $J_g(x_0)J_{g_0}(x_0)^{-1} \in B_{\delta_0}(I)$. Thus, $J_{gg_0^{-1}}(g_0(x_0)) = J_g(x_0)J_{g_0}(x_0)^{-1} \in B_{\delta_0}(I)$, where we used the chain rule and the inverse function theorem for Jacobians.

Proof of (b). It follows from $|gg_0^{-1}(g_0(x_0)) - g_0(x_0)| = |g(x_0) - g_0(x_0)| < \delta < \delta_0$.

Proof of (c). For all $|y - g_0(x_0)| < \eta'$, $y = g_0(x)$ for some x such that $|x - x_0| < \eta$. Now,

$$\begin{aligned} \|J_{g \circ g_0^{-1}}(y) - I\|_F &= \|J_g(x)J_{g_0}(x)^{-1} - I\|_F \le \sqrt{d}\|J_{g_0}(x)^{-1}\|_F \|J_g(x) - J_{g_0}(x)\|_F \\ &\le \sqrt{d}\|J_{g_0}(x)^{-1}\|_F \left(\|J_g(x) - J_g(x_0)\|_F + \|J_g(x_0) - J_{g_0}(x_0)\|_F + \|J_{g_0}(x_0) - J_{g_0}(x)\|_F\right) \\ &\le 3\sqrt{d}(\|J_{g_0}(x_0)^{-1}\|_F + 1)\delta \le \delta_0. \end{aligned}$$

Note that $||J_g(x) - J_g(x_0)||_F < \delta$ follows from (3), $||J_g(x_0) - J_{g_0}(x_0)||_F < \delta$ follows from (1) and $||J_{g_0}(x_0) - J_{g_0}(x_0)||_F < \delta$ follows from (iii) and finally $||J_{g_0}(x)^{-1}||_F \le ||J_{g_0}(x_0)^{-1}||_F + 1$ follows from (ii).

Theorem 5.3. (=**Theorem 1.3**) Let K be a Cantor set in \mathbb{R}^d and let $g_0 \in C^1_{inv}(\mathbb{R}^d)$. Then, there exists $x_0 \in K$ and $\delta > 0$ such that for all sufficiently small $\eta > 0$, we can find a Cantor set $\widetilde{K} \subset \mathbb{R}^d$ such that for all $g \in \mathcal{G}_{g_0, x_0, \delta, \eta}$,

(5.9)
$$g(K) \cap K \neq \emptyset.$$

Moreover, there exists $\varepsilon > 0$ and a Cantor set \widehat{K} such that

(5.10)
$$\left(\bigcap_{g\in\mathsf{B}^{\mathsf{d}}_{\varepsilon}(g_0)}\left(g(K)+\widehat{K}\right)\right)^{\circ}\neq\emptyset.$$

Proof. By Theorem 4.2, we can find an orthogonal transformation O such that $\mathsf{O}(g_0(K))$ is a Cantor set that is **u.n.d.**. Let \widehat{K} be the sub-Cantor set of K such that $\mathsf{O}(g_0(\widehat{K}))$ is uniformly non-degenerate satisfying (4.1). We now choose $x_0 \in \widehat{K}$. By Theorem 5.2, we can find $\delta > 0$ such that for all η sufficiently small, there exists a Cantor set $\widetilde{K}_0 \subset \mathbb{R}^d$ such that for all $h \in \mathcal{G}_{\mathsf{O} \circ g_0, x_0, \delta, \eta}$,

$$h(K) \cap \widetilde{K}_0 \neq \emptyset.$$

To finish the proof, we take $g \in \mathcal{G}_{g_0,x_0,\delta/||\mathbf{O}||_F,\eta}$, then a routine calculation shows that $h = \mathbf{O} \circ g$ will satisfy $J_h(x_0) \in B_{\delta}(J_{\mathbf{O} \circ g_0}(x_0)), |h(x_0) - (\mathbf{O} \circ g_0)(x_0)| < \delta$ and $|x - x_0| < \eta$ $\Rightarrow |h(x) - h(x_0)| < \delta$. Thus, $h \in \mathcal{G}_{\mathbf{O},x_0,\delta,\eta}$ and hence $(\mathbf{O} \circ g)(K) \cap \widetilde{K}_0 \neq \emptyset$. The proof is complete by taking $\widetilde{K} = \mathbf{O}^{-1}(\widetilde{K}_0)$.

We now prove the last statement. Let $g \in \mathcal{G}_{g_0,x_0,\delta/2,\eta}$. Then, we notice that if we let $|t| < \delta/2$ and consider the map $\tilde{g}(x) = g(x) + t$, then $J_{\tilde{g}}(x_0) = J_g(x_0)$, so $\|J_{\tilde{g}}(x_0) - J_g(x_0)\| < \delta$. Also,

$$|\tilde{g}(x_0) - g_0(x_0)| \le |g(x_0) - g_0(x_0)| + |t| < \delta.$$

Finally, if $|x-x_0| < \eta$, then $||J_{\tilde{g}}(x) - J_{\tilde{g}}(x_0)|| = ||J_g(x) - J_g(x_0)|| < \delta/2$ by the definition of $g \in \mathcal{G}_{g_0,x_0,\delta/2,\eta}$. Hence, by (5.9), $(g(K) + t) \cap \widetilde{K} \neq \emptyset$. Therefore, $t \in g(K) - \widetilde{K}$. Defining $\widehat{K} = -\widetilde{K}$, we find that the Euclidean ball $\{t : |t| < \delta/2\}$ is a subset of $\left(\bigcap_{g \in \mathcal{G}_{g_0,x_0,\delta/2,\eta}} \left(g(K) + \widehat{K}\right)\right)^{\circ}$. By Lemma 4.4, the ball $\mathsf{B}^{\mathsf{d}}_{\delta/2(2c_K+1)}(g_0) \subset \mathcal{G}_{g_0,x_0,\delta/2,\eta}$. This shows (5.10) holds with $\varepsilon = \delta/2(2c_K+1)$ provided we chose δ and η small enough so that Lemma 4.4 can be applied.

5.1. Interior of general C^1 functions. In this subsection, we aim at proving an analogous theorem for Theorem 2.6 on \mathbb{R}^d under the uniformly non-degenerate assumption of Cantor sets. We denote by $\mathbb{Q}_{\delta}(a)$ the closed cube in \mathbb{R}^d with side length δ and centered at a. Let Λ be a compact set on \mathbb{R}^N (the parameter set) with nonempty interior and let α_0 be an interior point of Λ , and Q_1 and Q_2 be two sets in \mathbb{R}^d with nonempty interior. Also, let $H \in C^1(\Lambda \times Q_1 \times Q_2) \to \mathbb{R}^d$ and let the $d \times d$ matrices of Jacobians $J_{H,1}$ of H by x and $J_{H,2}$ of H by y be given by

(5.11)
$$J_{H,1} = \left(\frac{\partial H_i}{\partial x_j}\right)_{i,j=1}^d \text{ and } J_{H,2} = \left(\frac{\partial H_i}{\partial y_j}\right)_{i,j=1}^d$$

We will assume that $J_{H,2}$ is invertible on Q_2 .

Fix $(u_1, u_2) \in (Q_1 \times Q_2)^\circ$ and let $c_0 = H(\alpha_0, u_1, u_2)$. Choose $\delta_0 > 0$ such that (5.12) $\mathbb{Q}_{\delta_0}(\alpha_0) \times \mathbb{Q}_{\delta_0}(u_1) \times \mathbb{Q}_{\delta_0}(u_2) \subset (\Lambda \times Q_1 \times Q_2)^\circ$.

Let

$$S = \mathbf{Q}_{\delta_0}(c_0) \times \mathbf{Q}_{\delta_0}(\alpha_0) \times \mathbf{Q}_{\delta_0}(u_1) \times \mathbf{Q}_{\delta_0}(u_2)$$

and define $F: S \to \mathbb{R}^d$ by

$$F(c, \alpha, x, y) = H(\alpha, x, y) - c$$

By the implicit function theorem, there exist $\delta_1 \in (0, \delta_0)$ and a function $G : Z \to \mathbf{Q}_{\delta_0}(u_2)$, where Z is the closed set given by

$$Z = \mathbf{Q}_{\delta_1}(c_0) \times \mathbf{Q}_{\delta_1}(\alpha_0) \times \mathbf{Q}_{\delta_1}(u_1),$$

such that $G(c_0, \alpha_0, u_1) = u_2$ and

(5.13)
$$H(\alpha, x, G(c, \alpha, x)) = c$$

Define

(5.14)
$$g_{c,\alpha}(x) = G(c,\alpha,x), \quad (c,\alpha) \in \mathbf{Q}_{\delta_1}(c_0) \times \mathbf{Q}_{\delta_1}(\alpha_0).$$

Notice that the Jacobian of $g_{c,\alpha}$,

$$J_{g_{c,\alpha}}(x) = J_{H,2}^{-1}(c,\alpha,x) \cdot J_{H,1}(c,\alpha,x),$$

is invertible by our assumption. With this setup for \mathbb{R}^d , the following lemma can be proved using the same line of proof as in Lemma 2.4. We will omit the details of the proof.

Lemma 5.4. Let $K_1 \subset Q_1$ and $K_2 \subset Q_2$ be Cantor sets and $\delta_2 \leq \delta_1$. If

$$g_{c,\alpha}(K_1) \cap K_2 \neq \emptyset$$
 for all $(c,\alpha) \in \mathbb{Q}_{\delta_2}(c_0) \times \mathbb{Q}_{\delta_2}(\alpha_0)$,

then

$$\mathbf{Q}_{\delta_2}(c_0) \subset \bigcap_{\alpha \in \mathbf{Q}_{\delta_2}(\alpha_0)} H(\alpha, K_1, K_2).$$

We now conclude our main theorem.

Theorem 5.5 (= Theorem 1.1). Let $N \ge 1$ and $\Lambda \subset \mathbb{R}^N$ be a set with interior and let $\alpha_0 \in \Lambda^\circ$. Let $Q_1 \subset Q_2 \subset \mathbb{R}^d$ and $Q_1^\circ \ne \emptyset$. Let H be a C^1 function on $\Lambda \times Q_1 \times Q_2$ such that $J_{H,2}$ in (5.11) is invertible on Q_2 .

Then for all Cantor sets $K_1 \subset Q_1$, there exists a Cantor set $K_2 \subset Q_2$ and a cube $\mathbb{Q}_{\epsilon}(\alpha_0)$ ($\varepsilon > 0$) such that

$$\left(\bigcap_{\alpha\in\mathbf{Q}_{\varepsilon}(\alpha_0)}H(\alpha,K_1,K_2)\right)^{\circ}\neq\emptyset.$$

Proof. Let $u_1 \in K_1 \subset Q_1 \subset Q_2$ and let $c_0 = H(\alpha_0, u_1, u_1)$. Then the maps $g_{c,\alpha}$ are defined according to (5.13) and (5.14). By restricting to a sub-Cantor set, we may assume $K_1 \subset \mathsf{Q}_{\delta_1}(u_1)$. Using Theorem 4.2, there exists an orthogonal transformation O such that $\mathsf{O}(g_{c_0,\alpha_0}(K_1))$ is uniformly non-degenerate. Let us write $g_0 = \mathsf{O} \circ g_{c_0,\alpha_0}$

and let $g_0(\widehat{K}_1)$ be the sub-Cantor set inside $g_0(K_1)$ that realizes the uniform nondegeneracy of $g_0(K_1)$. If $x_0 \in \widehat{K}_1 \subset \mathbb{Q}_{\delta_1}(u_1)$, then

$$J_{g_0}(x_0) = J_{\mathsf{O}\circ g_{c_0,\alpha_0}}(x_0) = \left(\mathsf{O}\circ J_{g_{c_0,\alpha_0}}\right)(x_0) \in \mathsf{GL}_d(\mathbb{R}).$$

By Theorem 5.2, there exists $\delta > 0$ such that for all sufficiently small $\eta > 0$, one can find a Cantor set \widetilde{K} such that for all $g \in \mathcal{G}_{g_0,x_0,\delta,\eta}$, $g(K_1) \cap \widetilde{K} \neq \emptyset$. \widetilde{K} is chosen from the containment lemma by requiring the convex hull to strictly contain the convex hull of $g(K_1)$ for all $g \in \mathcal{G}_{g_0,x_0,\delta,\eta}$. In particular, we are only interested in the case that $g = \mathbf{O} \circ g_{c,\alpha}$ and that the convex hulls of $\mathbf{O}(g_{c,\alpha}(K_1))$ are contained in $\mathbf{O}(Q_2)$. Note that we may choose \widetilde{K} contained in $\mathbf{O}(Q_2)$. We claim that there exists $\gamma > 0$ such that if $|(c,\alpha) - (c_0,\alpha_0)| < \gamma$, then $\mathbf{O} \circ g_{c,\alpha} \in \mathcal{G}_{g_0,x_0,\delta,\gamma}$. Clearly, we can make γ sufficiently small so that we can apply Theorem 5.2. Then, we have $|(c,\alpha) - (c_0,\alpha_0)| < \gamma$ implies

$$\mathsf{O}(g_{c,\alpha}(K_1)) \cap K \neq \emptyset.$$

Taking the inverse of O, the proof is complete by defining $K_2 = O^{-1}(\tilde{K})$ and applying Lemma 5.4.

To justify the claim, we need to show that there exists $\gamma > 0$ such that if $|(c, \alpha) - (c_0, \alpha_0)| < \gamma$, then

(1)
$$||J_{O \circ g_{c,\alpha}}(x_0) - J_{O \circ g_{c_0,\alpha_0}}(x_0)||_F < \delta.$$

(2) $|O(g_{c,\alpha}(x_0)) - O(g_{c_0,\alpha_0}(x_0))| < \delta.$
(3) $|x - x_0| < \gamma \implies ||J_{O \circ g_{c,\alpha}}(x) - J_{O \circ g_{c,\alpha}}(x_0)||_F < \delta.$

We now consider the following three functions,

$$F_{1}(c, \alpha, x) = \|J_{g_{c,\alpha}}(x) - J_{g_{c_{0},\alpha_{0}}}(x_{0})\|_{F} = \sqrt{\sum_{i,j=1}^{d} \left|\frac{\partial g_{j,c,\alpha}}{\partial x_{i}}(x) - \frac{\partial g_{j,c_{0},\alpha_{0}}}{\partial x_{i}}(x_{0})\right|^{2}},$$

$$F_{2}(c, \alpha, x) = |g_{c,\alpha}(x_{0}) - g_{c_{0},\alpha_{0}}(x_{0})|,$$

$$F_{3}(c, \alpha, x) = \|J_{g_{c,\alpha}}(x) - J_{g_{c,\alpha}}(x_{0})\|_{F},$$

where $g_{j,c,\alpha}$ are the component functions of $g_{c,\alpha}$ and they all have continuous derivatives in $(c, \alpha, x) \in Z$. Note that all F_i are uniformly continuous on a compact subset Z' of Z containing (c_0, α_0, x_0) . Hence, there exists $\gamma > 0$, such that if $|(c, \alpha, x) - (c', \alpha', x')| < \gamma$, then $|F_i(c, \alpha, x) - F_i(c', \alpha', x')| < \delta/||\mathbf{O}||_F$ for all i = 1, 2, 3.

For (1), if $|(c, \alpha) - (c_0, \alpha_0)| < \gamma$, then $|(c, \alpha, x_0) - (c_0, \alpha_0, x_0)| < \gamma$. Hence, notice that $F_1(c_0, \alpha_0, x_0) = 0$ and that

$$\|J_{O \circ g_{c,\alpha}}(x_0) - J_{O \circ g_{c_0,\alpha_0}}(x_0)\|_F \le \|\mathsf{O}\|_F \cdot F_1(c,\alpha,x_0) < \delta$$

(2) can be done using the same argument. For (3), we take $\eta = \gamma$, then if $|x - x_0| < \eta$, then $|(c, \alpha, x) - (c, \alpha, x_0)| < \gamma$ and hence

$$||J_{O \circ g_{c,\alpha}}(x) - J_{O \circ g_{c,\alpha}}(x_0)||_F \le ||\mathsf{O}||_F \cdot F_3(c,\alpha,x) < \delta,$$

where we also used that $F_3(c, \alpha, x_0) = 0$. We have thus verified that there exists $\gamma > 0$ such that $\mathsf{O} \circ g_{c,\alpha} \in \mathcal{G}_{g_0,x_0,\delta,\gamma}$ for all $|(c,\alpha) - (c_0,\alpha_0)| < \gamma$. The claim follows and the proof is complete.

6. When is a Cantor set non-degenerate?

We will now show that being embedded into axis parallel hyperplanes is the only possibility for the Cantor sets failing to be non-degenerate. Let us begin the setup of the proof. We let

$$K = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma^n} R_{\sigma}$$

be a nested representation of K as defined in Definition 3.1 and (3.1).

Lemma 6.1. Suppose that R_{σ} is degenerate. Then $R_{\sigma\sigma'}$ is degenerate for all $\sigma' \in \Sigma^k(\sigma)$ and $k \ge 1$.

Proof. If there exists $R_{\sigma\sigma'}$ such that it is non-degenerate, then one can find d+1 non-degenerate connected components inside $R_{\sigma\sigma'}$. But then, these connected components are also contained in R_{σ} , meaning that R_{σ} is non-degenerate, a contradiction. So the lemma holds.

Lemma 6.2. Let $K \subset \mathbb{R}^d$ be a Cantor set with a nested representation

$$K = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma^n} R_{\sigma}.$$

Suppose that all R_{σ} are degenerate. Then, K is contained in at most d^2 axis parallel hyperplane.

Proof. We first prove the following claim.

Claim. Suppose that R_{\emptyset} is degenerate. Then, for each set of d + 1 points in C, there exist 2 distinct points that lie in an axis parallel hyperplane.

Proof of Claim. Let $x_1, x_2, \dots, x_{d+1} \in K$. Then, for each $i \in \{1, \dots, d+1\}$, there exist $\sigma_{i,1}\sigma_{i,2}\sigma_{i,3} \dots \in \Sigma^{\infty}$ such that

(6.1)
$$\{x_i\} = \bigcap_{n=1}^{\infty} R_{\sigma_{i,1}\cdots\sigma_{i,n}}.$$

Since R_{\emptyset} is degenerate by assumption, for all $n \in \mathbb{N}$, the compact connected sets $\{R_{\sigma_{1,1}\cdots\sigma_{1,n}}, \cdots, R_{\sigma_{d+1,1}\cdots\sigma_{d+1,n}}\}$ satisfies

$$d_{\min}\left(R_{\sigma_{i,1}\cdots\sigma_{i,n}}, R_{\sigma_{j,1}\cdots\sigma_{j,n}}\right) = 0$$

for all $1 \leq i < j \leq d+1$. Hence, for each $n \in \mathbb{N}$, there exists $(i_n, j_n) \in \{1, \dots, d+1\} \times \{1, \dots, d+1\}$ and $k_n \in \{1, \dots, d\}$ such that

$$d(\pi_{k_n}(R_{\sigma_{i_n,1}\cdots\sigma_{i_n,n}}),\pi_{k_n}(R_{\sigma_{j_n,1}\cdots\sigma_{j_n,n}}))=0.$$

Since for each $n \in \mathbb{N}$, (i_n, j_n, k_n) are taken from a finite set, there exists (i_*, j_*, k_*) such that

$$d\left(\pi_{k_{*}}(R_{\sigma_{i_{*},1}\cdots\sigma_{i_{*},n}}),\pi_{k_{*}}(R_{\sigma_{j_{*},1}\cdots\sigma_{j_{*},n}})\right) = 0$$

holds for infinitely many n, which, for simplicity of notation, we may assume it holds for all n. In particular, this means that $\pi_{k_*}(R_{\sigma_{i_*,1}\cdots\sigma_{i_*,n}})$ and $\pi_{k_*}(R_{\sigma_{j_*,1}\cdots\sigma_{j_*,n}})$ overlaps, and from (6.1), we conclude that

$$|\pi_{k^*}(x_{i^*}) - \pi_{k^*}(x_{j^*})| \le |\pi_{k_*}(R_{\sigma_{i_*,1}\cdots\sigma_{i_*,n}})| + |\pi_{k_*}(R_{\sigma_{j_*,1}\cdots\sigma_{j_*,n}})| \to 0$$

as $n \to \infty$ since the diameter of R_{σ} tends to zero as σ gets longer. Hence, x_{i^*} and x_{j^*} lies in the same axis parallel plane.

With this claim, we let

$$k_{\min} = \min\{k \ge 2 : \forall k \text{ distinct points in } C, \exists 2 \}$$

distinct points that lie in an axis parallel hyperplane.}.

The claim implies that k_{\min} exists and is at most d+1. We now take $x_1, \cdots x_{k_{\min}-1}$ distinct points in K such that none of the two points lies in the same plane, where we denote $x_i = (x_{i,1}, \cdots x_{i,d})$. Then for all $x \in K - \{x_1, \cdots x_{k_{\min}-1}\}$, x must be in lying in one of the planes $x = x_{i,j}$ for some $i \in \{1, \cdots k_{\min} - 1\}$ and $j \in \{1, \cdots d\}$. Hence, K must be inside $(k_{\min} - 1) \cdot d \leq d^2$ many axis parallel hyperplane. This completes the proof.

Lemma 6.3. Let $K \subset \mathbb{R}^d$ be a degenerate Cantor set. Then, there exists $\sigma \in \Sigma^*$ such that R_{σ} is degenerate.

Proof. Suppose that for all $\sigma \in \Sigma^*$, R_{σ} is non-degenerate. Then, starting from R_{\emptyset} , we can find d+1 many \mathbb{R}_i at certain level k such that they have a positive d_{\min} . From our assumption, all \mathbb{R}_i are non-degenerate, so each \mathbb{R} has d+1 descendants that are non-degenerate. Continuing inductively, we are able to construct a sub-Cantor set that satisfies (3.3), meaning that K is non-degenerate.

Lemma 6.4. Let $K \subset \mathbb{R}^{d-1}$ be a non-degenerate Cantor set. Then there exists an orthogonal linear transformation T on \mathbb{R}^d such that the image $\mathsf{T}(K \times \{0\} + \mathbf{x})$ is non-degenerate on \mathbb{R}^d for all $\mathbf{x} \in \mathbb{R}^d$.

Proof. Note that it suffices to find an orthogonal linear transformation T such that $T(K \times \{0\})$ is non-degenerate, since $T(K \times \{0\} + \mathbf{x})$ just differs from $T(K \times \{0\})$ by a translation. Let $\mathbf{e}_1, \cdots, \mathbf{e}_d$ be the standard basis of \mathbb{R}^d . Consider the subspace $W = \{(x_1, \cdots, x_d) \in \mathbb{R}^d : x_1 + x_2 = 0\}$. Then, using elementary linear algebra, $\{\mathbf{v}_1 = (-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0 \cdots, 0), \mathbf{e}_3, \cdots, \mathbf{e}_d\}$ is an orthonormal basis for W. We define the linear map T on \mathbb{R}^d by sending

$$T(e_1) = v_1, T(e_i) = e_{i+1}, i = 2, \dots d - 1, T(e_d) = f_d$$

where \mathbf{f}_d spans the orthogonal complement W^{\perp} of W, i.e., $W^{\perp} = \text{span}(\mathbf{f}_d)$. The linear map is clearly an orthogonal transformation and is therefore invertible and it maps the x_d -plane onto the subspace W.

We now let \widehat{K} be the non-degenerate Cantor set inside $K \subset \mathbb{R}^{d-1}$. Let us represent it as in (3.1) with $\mathbb{R}_{\sigma} \subset \mathbb{R}^{d-1}$. Now, we can write

$$\widehat{K} \times \{0\} = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma_{d-1}^{n}} \mathbb{R}_{\sigma} \times \{0\}, \quad \mathsf{T}(\widehat{K} \times \{0\}) = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma_{d-1}^{n}} \mathsf{T}(\mathbb{R}_{\sigma} \times \{0\})$$

We now claim that for all $\sigma \in \Sigma_{d-1}^n$,

(6.2)
$$d_{\min}(\mathsf{T}(\mathsf{R}_{\sigma p} \times \{0\}), \mathsf{T}(\mathsf{R}_{\sigma q} \times \{0\})) > 0$$

for all $1 \leq p < q \leq d$. Take $\mathbf{x} = (x_1, \cdots, x_{d-1}, 0) \in \mathbb{R}_{\sigma p} \times \{0\}, \mathbf{y} = (y_1, \cdots, y_{d-1}, 0) \in \mathbb{R}_{\sigma q} \times \{0\}$. From the definition of d_{\min} ,

$$|x_i - y_i| \ge d_{\min}(\mathbf{R}_{\sigma p}, \mathbf{R}_{\sigma q}) := k_0.$$

By the definition of T,

$$\mathsf{T}(\mathbf{x}) = \left(-\frac{1}{\sqrt{2}}x_1, \frac{1}{\sqrt{2}}x_1, x_2, \cdots, x_{d-1}\right), \ \mathsf{T}(\mathbf{y}) = \left(-\frac{1}{\sqrt{2}}y_1, \frac{1}{\sqrt{2}}y_1, y_2, \cdots, y_{d-1}\right).$$

Hence, all coordinates of $\mathsf{T}(\mathbf{x})$ and $\mathsf{T}(\mathbf{y})$ are distinct by a distance of $\frac{1}{\sqrt{2}}k_0$. This shows that (6.2) holds. From this claim, we can construct a non-degenerate sub-Cantor set from $T(\widehat{K} \times \{0\})$ by inductively taking d + 1 non-degenerate components of \mathbb{R}_{σ} from $\{\mathbb{R}_{\sigma st} : s, t \in \{1, \dots d\}\}$.

We are now ready to conclude our main theorem in this section.

Theorem 6.5. Let $K \subset \mathbb{R}^d$ be a Cantor set. Then there exists an invertible linear transformation T on \mathbb{R}^d such that the image $\mathsf{T}(K)$ is non-degenerate on \mathbb{R}^d .

Proof. We prove by induction on the dimension d that all Cantor set K on \mathbb{R}^d satisfy the property in the statement. When d = 2 and if K is non-degenerate, we are done. Suppose that K is degenerate. By Lemma 6.3, there exists R_{σ} in a nest representation of C such that R_{σ} is degenerate. By Lemma 6.1 and 6.2, $K \cap R_{\sigma}$ must be contained in a finite union of axis-parallel planes. Hence, we can take a sub-Cantor set $\widehat{K} \subset K$ lying in an axis parallel line. Note that \widehat{K} , as a Cantor set on \mathbb{R}^1 , is always nondegenerate. By Lemma 6.4, there exists an invertible linear transformation T such that $\mathsf{T}(\widehat{K})$ is non-degenerate.

Suppose that the lemma is true for d-1. If $K \subset \mathbb{R}^d$ is non-degenerate, then we are done by taking simply the identity transformation. If $K \subset \mathbb{R}^d$ is degenerate, we can apply Lemma 6.1, 6.2 and 6.3 to obtain a sub-Cantor set lying in an axis parallel plane. Without loss of generality, we can assume that the sub-Cantor set lies on the subspace is $\{x_d = 0\}$, so that it can be represented as $\widehat{K} \times \{0\}$. From the induction hypothesis, there exists a linear map $\mathsf{T}_0 : \mathbb{R}^{d-1} \to \mathbb{R}^{d-1}$ such that $\mathsf{T}_0(\widehat{K})$ is non-degenerate on \mathbb{R}^{d-1} . By lemma 6.4, there exists a linear map T_1 such that $\mathsf{T}_1(\mathsf{T}_0(\widehat{K}) \times \{0\})$ is non-degenerate on \mathbb{R}^d .

Hence, the linear map $\mathsf{T}_2 : \mathbb{R}^d \to \mathbb{R}^d$ defined by $\mathsf{T}_2((\mathbf{x}, 0) + \mathbf{e}_d) = \mathsf{T}_0(\mathbf{x}, 0) + \mathbf{e}_d$ is an invertible linear map and $\mathsf{T}_2(\widehat{K} \times \{0\}) = \mathsf{T}_0(\widehat{K}) \times \{0\}$. Therefore, $\mathsf{T}_1 \circ \mathsf{T}_2(\widehat{K} \times \{0\})$ is non-degenerate. As $\widehat{K} \times \{0\} \subset K$, $(\mathsf{T}_1 \circ \mathsf{T}_2)(K)$ is therefore, non-degenerate. This completes the proof.

7. When is a Cantor set uniformly non-degenerate?

Definition 7.1. Let $K = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma^n} R_{\sigma}$ be a nested representation of K. Let $\kappa > 0$ be a constant. For each $k \in \mathbb{N}$, let $\{A_1, \dots, A_{d+1}\} \subset \{R_{\sigma\sigma'} : \sigma' \in \Sigma^k(\sigma)\}$. We define the following ratios:

$$\mathcal{A}^{k}(R_{\sigma}, A_{1}, \cdots, A_{d+1}) = \min_{p \neq q} \min_{i \neq j} \left\{ \frac{|\pi_{i}(x) - \pi_{i}(x')|}{|\pi_{j}(x) - \pi_{j}(x')|} : x \in A_{p}, x' \in A_{q}, \right\};$$

$$\mathcal{B}^{k}(R_{\sigma}, A_{1}, \cdots, A_{d+1}) = \max_{p \neq q} \max_{i \neq j} \left\{ \frac{|\pi_{i}(x) - \pi_{i}(x')|}{|\pi_{j}(x) - \pi_{j}(x')|} : x \in A_{p}, x' \in A_{q} \right\}.$$

Also, we define $\mathcal{A}^k(R_{\sigma}, A_1, \cdots, A_{d+1}) = 0$ and $\mathcal{B}^k(R_{\sigma}, A_1, \cdots, A_{d+1}) = \infty$ if $d_{\min}(A_p, A_q) = 0$ for some $p \neq q$. We say that R_{σ} is κ -non-degenerate if there exists $k \geq 1$ and there exists $\{A_1, \cdots, A_{d+1}\} \subset \{R_{\sigma\sigma'} : \sigma' \in \Sigma^k(\sigma)\}$ such that

(7.1)
$$\kappa^{-1} \leq \mathcal{A}^k(R_{\sigma}, A_1, \cdots, A_{d+1}) \leq \mathcal{B}^k(R_{\sigma}, A_1, \cdots, A_{d+1}) \leq \kappa.$$

Otherwise, we say that R_{σ} is κ -degenerate. We say that K is κ -uniformly nondegenerate if there exists a sub-Cantor set $\widehat{K} = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma_{n}^{n}} \mathbb{R}_{\sigma}$ and

(7.2)
$$\kappa^{-1} \leq \mathcal{A}^{1}(\mathbb{R}_{\sigma}, \mathbb{R}_{\sigma 1}, \cdots, \mathbb{R}_{\sigma (d+1)}) \leq \mathcal{B}^{1}(\mathbb{R}_{\sigma}, \mathbb{R}_{\sigma 1}, \cdots, \mathbb{R}_{\sigma (d+1)}) \leq \kappa.$$

for all $\sigma \in \Sigma_d^*$.

It is clear that K is uniformly non-degenerate if and only if there exists $\kappa > 0$ such that K is κ -uniformly non-degenerate. We will assume throughout this section that $K \subset \mathbb{R}^d$ is a Cantor set with a nested representation

$$K = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma^n} R_{\sigma}.$$

Lemma 7.2. Suppose that K is not **u.n.d.**. Then for all $\kappa > 0$, there exists R_{σ} such that R_{σ} is κ -degenerate. Moreover, for all $\{x_1, \dots, x_{d+1}\} \subset R_{\sigma} \cap K$, there exists $x_p \neq x_q$ and there exists $i \neq j \in \{1, \dots, d+1\}$ such that

(7.3)
$$|\pi_i(x_p) - \pi_i(x_q)| \le \kappa^{-1} |\pi_j(x_p) - \pi_j(x_q)|.$$

Proof. For the first statement, we prove the contrapositive. Suppose that there exists $\kappa > 0$ such that for all $\sigma \in \Sigma^*$, R_{σ} is κ -non-degenerate. Then starting from R_{\emptyset} , we can find d+1 many R_i at certain level k such that (7.2) holds. From our assumption, each R_i is κ -non-degenerate, so there exist d+1 κ -non-degenerate descendants of R_i . Continuing inductively, we can construct a sub-Cantor set \hat{K} of K such that

all $\sigma \in \Sigma_d^*$ satisfies (7.2). Therefore, K is κ -non-degenerate, which contradicts our assumption that K is not **u.n.d.**.

For the second statement, we suppose for the sake of contradiction that there exists $\{x_1, \dots, x_{d+1}\} \subset R_{\sigma} \cap K$ such that for all distinct points $x_p \neq x_q$, for all $i \neq j$

(7.4)
$$|\pi_i(x_p) - \pi_i(x'_p)| > \kappa^{-1} |\pi_j(x_q) - \pi_j(x'_q)|.$$

For each $p = 1, \dots, d+1$, let us write $\{x_p\} = \bigcap_{n=1}^{\infty} R_{\sigma\sigma_{p,1}\cdots\sigma_{p,n}}$. Since K is totally disconnected, by taking sufficiently large n, $R_{\sigma\sigma_{i,1}\cdots\sigma_{i,n}}$ has a very small diameter. Since π_i are continuous functions, (7.4) continues to hold for all $x \in R_{\sigma\sigma_{p,1}\cdots\sigma_{p,n}}$ and $x' \in R_{\sigma\sigma_{q,1}\cdots\sigma_{q,n}}$. This implies that

$$\kappa^{-1} < \frac{|\pi_i(x) - \pi_i(x')|}{|x_j(x) - \pi_j(x')|} < \kappa$$

for all $x \in R_{\sigma\sigma_{p,1}\cdots\sigma_{p,n}}$ and $x' \in R_{\sigma\sigma_{q,1}\cdots\sigma_{q,n}}$ and for all $i \neq j \in \{1, \cdots, d+1\}$. This implies that R_{σ} is κ -non-degenerate, which is a contradiction.

Lemma 7.3. Suppose that K is not **u.n.d.**. Then, for all $\kappa > 0$, there exists a sub-Cantor set \widehat{K} of K such that

$$\widehat{K} = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma_d^n} R_{\sigma}$$

and the following condition holds:

(7.5)
$$\forall \sigma \in \Sigma_d^n, \exists p \neq q \in \{1, \cdots, d\} \text{ and } \exists i \neq j \in \{1, \cdots, d+1\}$$

such that for all $x \in \mathbb{R}_{\sigma p}$ and $x' \in \mathbb{R}_{\sigma q}, |\pi_i(x) - \pi_i(x')| \leq \kappa^{-1} |\pi_j(x) - \pi_j(x')|.$

Proof. Using Lemma 7.2, we can find 2κ -degenerate R_{σ} and d+1 points $x_1, \dots x_{d+1} \in R_{\sigma} \cap K$ such that (7.3) holds for some $x_p \neq x_q \in \{x_1, \dots, x_{d+1}\}$ and some $i \neq j \in \{1, \dots, d+1\}$ with constant $(2\kappa)^{-1}$. Similar to the proof in Lemma 7.2, by considering R_{σ} around each x_p , $p = 1, \dots d+1$, we can find $R_1, \dots R_{d+1}$ with $x_i \in R_i$ such that (7.3) holds for all points in R_p and R_q with constant κ^{-1} , which means (7.5) holds.

We now notice a simple fact that if K is not **u.n.d.**, then $K \cap \mathbb{R}_i$ is also not **u.n.d.** for all $i = 1, \dots, d+1$. Hence, we can apply Lemma 7.2 and the argument we just did on $K \cap \mathbb{R}_i$. Then inductively to obtain \widehat{K} that fulfills this lemma.

Before we proceed to the main theorem, we make the following simple but important observation. With slight abuse of notation, we say that x, x' are κ -degenerate if there exists $i \neq j$ such that $|\pi_i(x) - \pi_i(x')| < \kappa^{-1} |\pi_j(x) - \pi_j(x')|$. In this case, x - x' must be lying in a cone around some axis-parallel hyperplane. More precisely,

(7.6)
$$x - x' \in \bigcup_{i=1}^{d} \mathsf{P}_i \left(\{ (x_1, \cdots x_d) : x_1^2 + \cdots + x_{d-1}^2 < \alpha x_d^2 \} \right) := \mathcal{C}$$

where $\alpha = \tan^{-1}(\kappa^{-1})$ and P_i are the orthogonal transformation mapping the hyperplane $x_d = 0$ onto the hyperplane $x_i = 0$. If K is a compact set inside \mathcal{C} , then we can find an orthogonal transformation O so that O(x) and O(x') are κ -non-degenerate for all $x \neq x' \in K$.

Theorem 7.4. Let $K \subset \mathbb{R}^d$ be a Cantor set. Then there exists an orthogonal linear transformation O on \mathbb{R}^d such that the image O(K) is **u.n.d.** on \mathbb{R}^d .

Proof. Let R be the diameter of K and let $N = \binom{d+1}{2}$ and let $O_1, \cdots O_N, O_{N+1}$ be orthogonal transformations such that for each $i = 1, \cdots, N, N+1$, O_i maps the standard basis onto an orthonormal basis $\mathcal{B}_i = \{\mathbf{e}_{i,1}, \cdots, \mathbf{e}_{i,d}\}$. We will assume that all $\mathbf{e}_{i,j}$ are distinct unit vectors. Let also H_i be the union of the linear spans of all possible d-1 vectors chosen from \mathcal{B}_i , i.e. the union of all (d-1)-dimensional hyperplanes generated by taking d-1 vectors from $\{\mathbf{e}_{i,1}, \cdots, \mathbf{e}_{i,d}\}$. The cone generated by H_i as in (7.6) will denoted by $\mathcal{C}_{i,\kappa}$. We will choose κ so that all $\mathcal{C}_i = \mathcal{C}_{i,\kappa} \cap B(0,R)$ intersect only at the origin.

With all these parameters fixed, we consider the given Cantor set K. If K is uniformly non-degenerate, there is nothing prove. Otherwise, by Lemma 7.2, we can find R_{σ} so that R_{σ} is 2κ -degenerate and Lemma 7.3 implies that we can find a sub-Cantor set $\widehat{K} = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \Sigma_d^n} \mathbb{R}_{\sigma}$ such that for all $\sigma \in \Sigma_d^n$, there exists $p \neq q$ and $i \neq j$ such that

$$|\pi_i(x) - \pi_i(x')| \le \kappa^{-1} |\pi_j(x) - \pi_j(x')|$$

for all $x \in \mathbf{R}_{\sigma p}$ and $x' \in \mathbf{R}_{\sigma q}$.

Note that if there exists an orthogonal transformation O and there exists $\kappa > 0$ such that $O(\widehat{K})$ is κ -non-degenerate, then O(K) is uniformly non-degenerate, so the theorem is proved. Therefore, assume that for all orthogonal transformation O and for all $\kappa > 0$, $O(\widehat{K})$ is κ -degenerate.

We then take $O = O_1$ and κ in the first paragraph. By Lemma 7.3, we can further find a sub-Cantor set \widehat{K}_1 inside \widehat{K} such that \widehat{K}_1 satisfies (7.5) (if \widehat{K}_1 fails (7.5), then it will violate (7.3)) and $O_1(\widehat{K}_1)$ satisfies (7.5).

For each σ , \widehat{K}_1 already satisfies (7.5) for some $\mathbb{R}_{\sigma p}$ and $\mathbb{R}_{\sigma q}$. We notice that for the same pair, $O_1(\mathbb{R}_{\sigma p})$ and $O_1(\mathbb{R}_{\sigma q})$ cannot satisfy (7.5) due to choice of our κ , in which \mathcal{C}_i only intersects at origin. Hence, they must be coming from another pair.

Again, if there exists an orthogonal transformation O and there exists $\kappa > 0$ such that $O(\widehat{K}_1)$ is κ -non-degenerate, then the proof is done. Otherwise, we can further find a sub-Cantor set $\widehat{K}_2 \subset \widehat{K}_1$ such that

- (1) $O_2(\widehat{K}_2)$ satisfies (7.5)
- (2) $O_1(\hat{K}_2)$ satisfies (7.5),
- (3) \widehat{K}_2 satisfies (7.5).

Note that all the above three conditions must be from distinct pairs of $\mathbb{R}_{\sigma p}$ and $\mathbb{R}_{\sigma q}$ for all σ . We proceed with the proof inductively until all $N = \binom{d+1}{2}$ possible edges are

exhausted. Then a sub-Cantor set \widehat{K}_{N+1} with orthogonal transformation $O_{N+1}(\widehat{K}_{N+1})$ must be uniformly non-degenerate. The proof is complete.

8. Applications

8.1. Topological Erdős similarity problem. The Erdős similarity conjecture asserts that for all infinite sets P, there always exists a set E of positive Lebesgue measure such that E does not contain an affine copy of P. It has been studied by many authors (see e.g. [6, 2, 17]). One can also refer to [8] and the references therein for some recent progress. The conjecture remains open for all fast decaying sequences and even for Cantor sets with Newhouse thickness zero. In [10], Gallagher, Lai, and Weber proposed a topological version of the conjecture.

Conjecture: For all uncountable sets P, there is a dense G_{δ} set G that does not contain an affine copy of P.

In another paper of the authors [16], we showed that this conjecture is equivalent to the Borel conjecture and is actually independent of the ZFC axiomatic set theory. Despite its independence, the conjecture can be verified to be true for Cantor sets [10] on \mathbb{R}^d . In this paper, we offer another proof of this result using Theorem 1.1 and provide an even more general statement.

Theorem 8.1. Let K be a Cantor set on \mathbb{R}^d . Then there exists a dense G_{δ} set G such that G does not contain g(K) for all $g \in C^1_{inv}(\mathbb{R}^d)$.

Proof. We note that a dense G_{δ} set G does not contain any C^1_{inv} image of K if and only if $g(K) \cap F \neq \emptyset$ for all $g \in C^1_{inv}(\mathbb{R}^d)$ where $F = \mathbb{R}^d \setminus G$ is an nowhere dense F_{σ} set. Hence, we need to construct the set F.

For all $f \in C^1_{inv}(\mathbb{R}^d)$, by Theorem 5.3, there exists $\varepsilon = \varepsilon_f > 0$ and a Cantor set \widehat{K}_f such that

$$\left(\bigcap_{g\in\mathsf{B}^{\mathsf{d}}_{\varepsilon_{f}}(f)}(g(K)+\widehat{K})\right)^{\circ}\neq\emptyset.$$

Let $\delta_f > 0$ be such that the Euclidean ball $B_{\delta_f}(0)$ is inside the interior, so that

(8.1)
$$g(K) + \widehat{K}_f + \delta_f \mathbb{Z} = \mathbb{R}^d \quad \forall g \in \mathsf{B}^{\mathsf{d}}_{\varepsilon_f}(f).$$

Note that $C^1_{inv}(\mathbb{R}^d)$ is now covered by the open balls $\mathsf{B}^{\mathsf{d}}_{\varepsilon_f}(f)$. By Lemma 4.5, $C^1_{inv}(\mathbb{R}^d)$ is covered by countably many such balls. We call them $\mathsf{B}^{\mathsf{d}}_{\varepsilon_{fn}}(f_n)$, $n = 1, 2, 3, \cdots$. Let

$$F = \bigcup_{n=1}^{\infty} (\widehat{K}_{f_n} + \delta_{f_n} \mathbb{Z}).$$

From (8.1) and the covering property,

$$g(K) + F = \mathbb{R}^d \ \forall g \in C^1_{inv}(\mathbb{R}^d).$$

This means that $0 \in g(K) + F$ and thus $g(K) \cap (-F) \neq \emptyset$ for all $g \in C^1_{inv}(\mathbb{R}^d)$. It is clear that -F is an F_{σ} set. Since $\widehat{K}_{f_n} + \delta_{f_n}\mathbb{Z}$ is closed and nowhere dense, F is also nowhere dense by the Baire category theorem. This completes the proof. \Box

It is clear that the theorem will not be true if we consider only $C^1(\mathbb{R}^d)$ as constant functions lie inside $C^1(\mathbb{R}^d)$, which will map Cantor sets to one point.

As another remark, we do not know the Lebesgue measure of F in the theorem. If one can show that F has finite Lebesgue measure or even Lebesgue measure zero, then we will be able to show that the original Erdős similarity conjecture is true for all Cantor sets.

8.2. Pinned distance sets in even dimensions. We first recall the definition of pinned distance sets. Let $A \subset \mathbb{R}^d$, $t \in \mathbb{R}^d$, and $\alpha > 1$. Then, the pinned distance set at t with respect to norm $||x||_{\alpha} := (\sum_{i=1}^d |x_i|^{\alpha})^{1/\alpha}$ is the set

$$\Delta_t^{(\alpha)}(K_1 \times K_2) = \{ \|x - t\|_\alpha : x \in K_1 \times K_2 \}.$$

The famous Falconer distance set conjecture asserted that if a Borel set $E \subset \mathbb{R}^d$ has $\dim_H(E) > d/2$ (dim_H denotes Hausdorff dimension), then the distance set $\Delta(E) = \{ ||x - y||_2 : x, y \in E \}$ has positive Lebesgue measure. One can refer to ([18, Chapter 4]) for more details. For the current best known world record towards this conjecture, one can refer to [11, 5].

At the threshold value d/2, Falconer constructed an example for which the distance set has zero Lebesgue measure. From Simon and Taylor [26], one can construct a Cantor set on \mathbb{R}^2 whose dimension is arbitraily close to 1 (but never equal 1), such that its pinned distance set has an interior. Indeed, they show that if $\tau(K_1)\tau(K_2) > 1$, then $K = K_1 \times K_2$ is the desired set. By requiring K_1 and K_2 to be the symmetric Cantor sets with dissection ratio α and β , one can find some K has dimension arbitrarily close to 1 and $\tau(K_1)\tau(K_2) > 1$ is maintained. Using the result in our paper, we can construct a Cantor set K of dimension d/2 on \mathbb{R}^d with even d such that the pinned distance set has non-empty interior.

Theorem 8.2. For every even d and $\alpha > 1$, there exists a Cantor set K in \mathbb{R}^d with $\dim_H(K) = \frac{d}{2}$ such that the pinned distance set $\left(\Delta_t^{(\alpha)}(K)\right)^{\circ} \neq \emptyset$ for some $t \in K$.

Proof. Let Λ be any closed interval containing α as an interior point and Q_1, Q_2 be cubes in $\mathbb{R}^{d/2}$ such that $Q_2 \subset \{(y_1, \cdots, y_{d/2}) \in \mathbb{R}^{d/2} : y_i > 0, \forall i = 1, \cdots d/2\}$ and $Q_1 \subset Q_2^{\circ}$. For simplicity, we take t = 0. Define a C^1 function $H = H_t$ on $\Lambda \times Q_1 \times Q_2$ by

$$H_t\left(\alpha, (x_1, \cdots, x_{d/2}), (y_1, \cdots, y_{d/2})\right) \\= \left(\|(x_1, \cdots, x_{d/2}, y_1, \cdots, y_{d/2})\|_{\alpha}^{\alpha}, y_2, y_3, \cdots, y_{d/2}\right).$$

Then, the Jacobian matrix $J_H(y)$ of H is given by

| 10 | $\alpha y_1^{\alpha-1}$ | $\alpha y_2^{\alpha-1}$ | ••• | • • • | $\alpha y_{d/2}^{\alpha-1}$ | |
|----|-------------------------|-------------------------|-----|-------|-----------------------------|--|
| | 0 | 1 | 0 | ••• | 0 | |
| | 0 | 0 | 1 | ••• | 0 | |
| | ÷ | ÷ | ÷ | ••• | ÷ | |
| | 0 | 0 | 0 | • • • | 1 / | |

Note that $J_H(y)$ has determinant $\alpha y_1^{\alpha-1}$, which is nonzero if and only if $y_1 \neq 0$. By our assumption on Q_2 , the determinant is nonzero for all y. Let $K_1 \subset Q_1$ be any Cantor set with Hausdorff dimension zero. Then, Theorem 1.1 implies that there exists a Cantor set $K_2 \subset Q_2$ such that $H(\alpha, K_1, K_2)$ has a nonempty interior. Projecting onto the first coordinate of $\mathbb{R}^{d/2}$, we have that $\{||x||_{\alpha}^{\alpha} : x \in K_1 \times K_2\}$ has nonempty interior. Thus, $\Delta_0^{(\alpha)}(K_1 \times K_2)$ has nonempty interior. We notice that K_2 has a positive Lebesgue measure (See Remark 3.5 (1)), so $\dim_H(K_2) = \dim_B(K_2) = d/2$. By the well-known dimension estimate formula ([7]),

$$\dim_H(K_1) + \dim_H(K_2) \le \dim_H(K_1 \times K_2) \le \dim_H(K_1) + \dim_B(K_2),$$

we have $\dim_H(K_1 \times K_2) = \frac{d}{2}$. Finally, we can put a Cantor set K_0 of Hausdorff dimension 0 and it contains the origin. Let $K = K_0 \cup (K_1 \times K_2)$. Then K has our desired property stated in the theorem.

References

- Steve Astels, Cantor sets and numbers with restricted partial quotients, Trans. Amer. Math. Soc. 352 (2000), no. 1, 133–170. MR 1491854
- Jean Bourgain, Construction of sets of positive measure not containing an affine image of a given infinite structures, Israel J. Math. 60 (1987), no. 3, 333–344. MR 937795
- Karma Dajani, Vilmos Komornik, Derong Kong, and Wenxia Li, Algebraic sums and products of univoque bases, Indag. Math. (N.S.) 29 (2018), no. 4, 1087–1104. MR 3826515
- 4. Carlos Gustavo T. de A. Moreira and Jean-Christophe Yoccoz, Stable intersections of regular Cantor sets with large Hausdorff dimensions, Ann. of Math. (2) 154 (2001), no. 1, 45–96. MR 1847588
- Xiumin Du, Alex Iosevich, Yumeng Ou, Hong Wang, and Ruixiang Zhang, An improved result for Falconer's distance set problem in even dimensions, Math. Ann. 380 (2021), no. 3-4, 1215– 1231. MR 4297185
- Kenneth Falconer, On a problem of Erdős on sequences and measurable sets, Proc. Amer. Math. Soc. 90 (1984), no. 1, 77–78. MR 722418
- Fractal geometry, second ed., John Wiley & Sons, Inc., Hoboken, NJ, 2003, Mathematical foundations and applications. MR 2118797
- De-Jun Feng, Chun-Kit Lai, and Ying Xiong, Erdős similarity problem via bi-lipschitz embedding, To appear in IMRN (2024).
- De-Jun Feng and Yu-Feng Wu, On arithmetic sums of fractal sets in ℝ^d, Journal of the London Mathematical Society 104 (2021), no. 1, 35–65.
- John Gallagher, Chun-Kit Lai, and Eric Weber, On a topological Erdős similarity problem, Bull. Lond. Math. Soc. 55 (2023), no. 3, 1104–1119. MR 4599101

- Larry Guth, Alex Iosevich, Yumeng Ou, and Hong Wang, On Falconer's distance set problem in the plane, Invent. Math. 219 (2020), no. 3, 779–830. MR 4055179
- Marshall Hall, Jr., On the sum and product of continued fractions, Ann. of Math. (2) 48 (1947), 966–993. MR 22568
- Michael Hochman and Pablo Shmerkin, Local entropy averages and projections of fractal measures, Ann. of Math. (2) 175 (2012), no. 3, 1001–1059. MR 2912701
- Roger A. Horn and Charles R. Johnson, *Matrix analysis*, Cambridge University Press, Cambridge, 1985. MR 832183
- Kan Jiang, Obtaining an explicit interval for a nonlinear Newhouse thickness theorem, Math. Z. 301 (2022), no. 1, 1011–1037. MR 4405675
- 16. Yeonwook Jung and Chun-Kit Lai, Topological Erdős similarity conjecture and strong measure zero sets, preprint (2024).
- Mihalis Kolountzakis, Infinite patterns that can be avoided by measure, Bull. London Math. Soc. 29 (1997), no. 4, 415–424. MR 1446560
- Pertti Mattila, Fourier analysis and Hausdorff dimension, Cambridge Studies in Advanced Mathematics, vol. 150, Cambridge University Press, Cambridge, 2015. MR 3617376
- Alex McDonald and Krystal Taylor, Finite point configurations in products of thick Cantor sets and a robust nonlinear Newhouse gap lemma, Math. Proc. Cambridge Philos. Soc. 175 (2023), no. 2, 285–301. MR 4623515
- Piotr Nowakowski, When the algebraic difference of two central Cantor sets is an interval?, Ann. Fenn. Math. 48 (2023), no. 1, 163–185. MR 4535243
- Jacob Palis and Floris Takens, Hyperbolicity and sensitive chaotic dynamics at homoclinic bifurcations, Cambridge Studies in Advanced Mathematics, vol. 35, Cambridge University Press, Cambridge, 1993, Fractal dimensions and infinitely many attractors. MR 1237641
- Mehdi Pourbarat, On the arithmetic difference of middle Cantor sets, Discrete Contin. Dyn. Syst. 38 (2018), no. 9, 4259–4278. MR 3824702
- Atsuro Sannami, An example of a regular Cantor set whose difference set is a Cantor set with positive measure, Hokkaido Math. J. 21 (1992), no. 1, 7–24. MR 1153749
- Jörg Schmeling and Pablo Shmerkin, On the dimension of iterated sumsets, Recent developments in fractals and related fields, Appl. Numer. Harmon. Anal., Birkhäuser Boston, Boston, MA, 2010, pp. 55–72. MR 2742987
- Pablo Shmerkin and Ville Suomala, Spatially independent martingales, intersections, and applications, Mem. Amer. Math. Soc. 251 (2018), no. 1195, v+102. MR 3756896
- Károly Simon and Krystal Taylor, Interior of sums of planar sets and curves, Math. Proc. Cambridge Philos. Soc. 168 (2020), no. 1, 119–148. MR 4043823
- Boris Solomyak, On the measure of arithmetic sums of Cantor sets, Indag. Math. (N.S.) 8 (1997), no. 1, 133–141. MR 1617830
- Yuki Takahashi, Sums of two homogeneous Cantor sets, Trans. Amer. Math. Soc. 372 (2019), no. 3, 1817–1832. MR 3976578
- 29. Stephen Willard, General topology, Courier Corporation, 2012.
- Alexia Yavicoli, Thickness and a gap lemma in ℝ^d, Int. Math. Res. Not. IMRN (2023), no. 19, 16453–16477. MR 4651893

San Francisco State University Department of Mathematics, 1600 Holloway Ave, San Francisco, CA 94132

Email address: yjung4@sfsu.edu

San Francisco State University Department of Mathematics, 1600 Holloway Ave, San Francisco, CA 94132

 $Email \ address: \ \texttt{cklai@sfsu.edu}$