

LAGRANGIAN EXTENSIONS AND LEFT SYMMETRIC STRUCTURES ON THE FOUR-DIMENSIONAL REAL LIE SUPERALGEBRAS

SOFIANE BOUARROUDJ AND ANA-MARIA RADU

ABSTRACT. Over real numbers, Backhouse classified all four-dimensional Lie superalgebras. From this list, we will investigate those Lie superalgebras that can be obtained as Lagrangian extensions. Moreover, we investigate left-symmetric structures on these Lie superalgebras. Furthermore, except for two of them, they are all Novikov superalgebras.

1. INTRODUCTION

1.1. **Lagrangian extensions of Lie (super)algebras.** The notion of T^* -extension of Lie algebras was initially introduced by Bordemann [Bor]. A short explanation can be understood as the following. Given a Lie algebra \mathfrak{h} equipped with an \mathfrak{h}^* -valued bilinear map α , a T^* -extension of \mathfrak{h} is the Lie algebra structure on the vector space $\mathfrak{g} := \mathfrak{h} \oplus \mathfrak{h}^*$ constructed by means of the form α that satisfies a certain cohomological requirement. The Lie algebra \mathfrak{g} naturally admits an invariant non-degenerate *symmetric* bilinear form. Conversely, Bordemann showed in [Bor] that every finite-dimensional nilpotent Lie algebra \mathfrak{h} with an invariant symmetric non-degenerate bilinear form “is” a suitable T^* -extension. In particular, if \mathfrak{h} is a real finite-dimensional Lie algebra belonging to a Lie group G , then the T^* -extension of \mathfrak{h} is the Lie algebra of the cotangent bundle T^*G of G , considered as a Lie group. The terminology and notation are justified by this differential geometric fact.

A superization of this construction was given in [BBB]. In particular, it was shown that every solvable Lie superalgebra with an *even* symmetric non-degenerate bilinear form is either isomorphic to a T^* -extension of a certain Lie superalgebra or to an ideal of codimension one of a certain T^* -extension.

The notion of T^* -extension was also studied by Baues and Cortés in [BC] in the context of Lie algebras admitting a flat connection, and called *Lagrangian extensions* since the Lie sub-algebra \mathfrak{h}^* is a Lagrangian ideal in $\mathfrak{h} \oplus \mathfrak{h}^*$. Unlike the case studied in [Bor],

2020 *Mathematics Subject Classification.* 17B05; 17B55; 17D25.

Key words and phrases. Quasi-Frobenius Lie (super)algebra, Lagrangian extensions, Left symmetric structures, Novikov structures, Balinsky-Novikov structures.

SB is supported by the grant NYUAD-065.

the non-degenerate form in this context is anti-symmetric and defines a 2-cocycle on \mathfrak{h} . In addition, the Lie algebra \mathfrak{h} must admit a flat connection so that the Lagrangian extension is possible. Lie algebras with a non-degenerate anti-symmetric form satisfying the 2-cocycle conditions are called quasi-Frobenius, and are studied intensively in the literature, see, e.g., [E, O].

Lie algebras possessing both an invariant non-degenerate symmetric bilinear form and a non-degenerate 2-cocycle of its scalar cohomology, were studied in [BBM] and were called *quadratic symplectic* Lie algebras. If the ground field is algebraically closed, the authors showed that every quadratic symplectic Lie algebra is a T^* -extension of a Lie algebra which admits an invertible derivation. Necessary and sufficient conditions on \mathfrak{h} and on the cocycle used in the construction of the T^* -extension are investigated to ensure that the extended algebra admits a skew-symmetric derivation and, hence, a symplectic structure. As a result, complex symplectic quadratic Lie algebras with dimensions less than or equal to 8 are classified. Some of these Lie algebras correspond to some Lie groups that admit a bi-invariant pseudo-Riemannian metric as well as a left-invariant symplectic form. These Lie groups are nilpotent, and their geometry is very rich, see [BBM, MR] for more details.

The notion of Lagrangian extension was superized in [BM]. The construction is valid for arbitrary Lie superalgebras defined over a field of characteristic not two provided the Lie superalgebra admit a torsion-free flat connection. Moreover, they have observed that there are two ways to perform this extension: they consider either $\mathfrak{g} := \mathfrak{h} \oplus \mathfrak{h}^*$ or $\mathfrak{g} := \mathfrak{h} \oplus \Pi(\mathfrak{h}^*)$, where Π is the change of parity functor. They call them T^* -extension and ΠT^* -extensions of \mathfrak{h} , respectively. In the first case, the non-degenerate form is even, and in the second case, it is odd.

Several examples of Lagrangian extensions are given in [BM], including filiform Lie superalgebras and some Lie superalgebras over reals. As a source of examples, they used the list of 4-dimensional real Lie superalgebras classified by Backhouse in [B].

One of the main objectives of this paper is to complete the list of examples among the 4-dimensional Lie superalgebras and explain how they can be obtained as Lagrangian extensions, see Section 4. According to the Backhouse classification, this investigation will classify all four-dimensional real Lie superalgebras that admit (or don't admit) Lagrangian extensions.

1.2. Left-symmetric structures. Left-symmetric algebras (also known as Vinberg-Koszul algebras) play an important role in various fields of mathematics and mathematical physics; for instance, rooted tree algebras, operad theory, vertex algebras, convex homogeneous cones, affine manifolds, left-invariant affine structures on Lie groups, to name just a few. For a thorough review, see [B3].

Over the real and complex numbers left-symmetric algebras are closely related to the theory of left-invariant affine structures over Lie groups, see for example [M, GS] (and references therein).

The investigation and classification of left-symmetric structure on Lie (super)algebras is an important subject by its own, as initiated in [S]. The problem, however, seems to be difficult to handle.

For reductive Lie algebras, Baues [Ba] proved that $\mathfrak{gl}(n)$ is the only reductive Lie algebra with one-dimensional center and a simple semisimple ideal which admits left-symmetric algebras over an algebraically closed ground field (see also [B1]). Moreover, all left-symmetric algebras on $\mathfrak{gl}(n)$ were classified.

Simple Lie algebras over complex numbers do not admit left-symmetric structures, see [H]. The existence of such structures is not always guaranteed, even for nilpotent Lie algebras. There are filiform nilpotent Lie groups of dimension $10 \leq n \leq 13$ which do not admit any left-invariant affine structure, see [B, B3, BG]. Any filiform nilpotent Lie group of dimension $n \leq 9$ admits a left-invariant affine structure. Nevertheless, left-symmetric algebra structures on both finite-dimensional and infinite-dimensional Lie algebras have been studied extensively, see [Bai, B2, YZ] as well as [CL, KCB, TB], respectively. Left symmetric structures on the Virasoro superalgebra were studied in [KB].

In the super setting, it was proved in [DZ] that $\mathfrak{sl}(m+1|m)$ admits a left-symmetric superalgebra, and a full classification is offered in the case of $\mathfrak{sl}(2|1)$. Moreover, it was shown that $\mathfrak{sl}(m|1)$ does not admit left-symmetric superalgebras for $m \geq 3$.

There is also the notion of Novikov algebra. It was introduced in connection with the Poisson brackets of hydrodynamic type, see [BN].

A second aim of this paper is to provide explicitly a left-symmetric structure on the 4-dimensional real Lie superalgebras. Surprisingly enough, all these Lie superalgebras admit such a structure, given explicitly in Section 5. Furthermore, we show that these structures are Novikov, except for the two Lie superalgebras $(D_0^{10})^1$ and $(D_0^{10})^2$. In addition, all these Lie superalgebras admit a Balinsky Novikov structure. Classifying all left-symmetric structures on each of these Lie superalgebras remains an open problem.

2. MAIN CONCEPTS AND DEFINITIONS

The purpose of this section is to review the main definitions and theorems that will be used in this paper.

Throughout the text \mathbb{K} is a field of characteristic $p \neq 2$.

A Lie superalgebra is a \mathbf{Z}_2 -graded space $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ over a field of characteristic $p \neq 2$ together with a bilinear binary operator $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$, called the Lie superbracket that has the following properties: (for any homogeneous $x, y, z \in \mathfrak{g}$)

- (i) Super anti-commutativity: $[x, y] = -(-1)^{|x||y|}[y, x]$;

- (ii) Super Jacobi identity: $[x, [y, z]] = [[x, y], z] + (-1)^{|x||y|}[y, [x, z]]$ where $|x| = \bar{0}$ if $x \in \mathfrak{g}_{\bar{0}}$ and $|x| = \bar{1}$ if $x \in \mathfrak{g}_{\bar{1}}$;
- (iii) If $p = 3$, one must add $[x, [x, x]] = 0$ if $x \in \mathfrak{g}_{\bar{1}}$.

There is also the notion of Lie superalgebra in characteristic 2 but we are not studying them here.

Let us introduce the notion of bilinear forms over a superspace. For more details, see [LSoS].

Let V and W be two superspaces defined over an arbitrary field \mathbb{K} . Let $\omega \in \text{Bil}(V, W)$ be a *homogeneous* bilinear form. The Gram matrix $B = (B_{ij})$ associated to ω is given by the formula,

$$(1) \quad B_{ij} = (-1)^{|\omega||v_i|} \omega(v_i, v_j) \quad \text{for the basis vectors } v_i \in V.$$

This definition can be extended by linearity to non-homogeneous forms. Moreover, it allows us to identify a bilinear form $B(V, W)$ with an element of $\text{Hom}(V, W^*)$. Consider the *upsetting* of bilinear forms $u: \text{Bil}(V, W) \rightarrow \text{Bil}(W, V)$ given by the formula

$$(2) \quad u(\omega)(u, v) = (-1)^{|v||u|} \omega(v, u) \quad \text{for any homogeneous } v \in V \text{ and } u \in W.$$

In terms of the Gram matrix B of ω : the form ω is *symmetric* if and only if

$$(3) \quad u(B) = B, \quad \text{where } u(B) = \begin{pmatrix} R^t & (-1)^{|\omega|} T^t \\ (-1)^{|\omega|} S^t & -U^t \end{pmatrix}, \quad \text{for } B = \begin{pmatrix} R & S \\ T & U \end{pmatrix}.$$

Similarly, *anti-symmetry* of ω means that $u(B) = -B$.

An even (resp. odd) non-degenerate bilinear form is called *orthosymplectic* (resp. *periplectic*).

Proposition 2.1 (Superdimension constraints). *If $V = V_{\bar{0}} \oplus V_{\bar{1}}$ is a finite-dimensional superspace such that $V_{\bar{1}} \neq 0$, equipped with an anti-symmetric non-degenerate bilinear form ω , then:*

(i) *If $p(\omega) = \bar{0}$, then $\dim(V_{\bar{0}})$ is even.*

(ii) *If $p(\omega) = \bar{1}$, then $\dim(V_{\bar{0}}) = \dim(V_{\bar{1}})$.*

Proof. See, [BM] □

We will denote by Π the functor that assigns to V the superspace $\Pi(V)$ defined as the other copy of V with the opposite parity of its homogeneous components. We have:

$$(\Pi(V))_{\bar{0}} = V_{\bar{1}}, \quad (\Pi(V))_{\bar{1}} = V_{\bar{0}}.$$

The space $\Pi(V)$ consists of the linear combinations of elements that we denote by $\Pi(f)$ for every homogeneous $f \in V$. For every superspace V , denote the canonical odd homomorphism induced by the functor Π by

$$\Pi : V \rightarrow \Pi(V) \quad f \mapsto \Pi(f).$$

We will identify V and $\Pi(\Pi(V))$, naturally.

Let $\omega \in \text{Bil}(\mathfrak{g}, \mathfrak{g})$ be an anti-symmetric bilinear form on a Lie superalgebra \mathfrak{g} . The form ω is called **closed** if it satisfies:

$$(-1)^{|x||z|}\omega(x, [y, z]) + (-1)^{|z||y|}\omega(z, [x, y]) + (-1)^{|y||x|}\omega(y, [z, x]) = 0,$$

for all $x, y, z \in \mathfrak{g}_{\bar{i}}$. This means that ω is 2-cocycle with scalar values.

A Lie superalgebra \mathfrak{g} is called *quasi-Frobenius* if it is equipped with a closed anti-symmetric non-degenerate bilinear form ω . If ω is even (resp. odd) we call it *orthosymplectic* (resp. *perisymplectic*). We denote such a Lie superalgebra by (\mathfrak{g}, ω) . Recently, these Lie superalgebras were studied in [BE, BEM, BM] in the context of Lagrangian and double extensions.

A Lie superalgebra is called *Frobenius* if ω is exact, which means that there is $f \in \mathfrak{g}^*$ such that $\omega(x, y) = f([x, y])$ for all $x, y \in \mathfrak{g}$.

Let (\mathfrak{g}, ω) be a quasi-Frobenius Lie superalgebra. An ideal $I \subseteq \mathfrak{g}$ is called *Lagrangian* if and only if $I^\perp = I$.

2.1. Connections on Lie (super)algebras. To write down a Lagrangian extension of a Lie (super)algebra, we will need the notion of a connection on a Lie (super)algebra. In our context, connections are defined as non-associative products on a superalgebra with purely algebraic definitions. See [BM, MKSV] for the non-super case.

An **even** bilinear map $\nabla : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$, written as $(x, y) \rightarrow \nabla_x y$ is called a **connection** on \mathfrak{g} .

We will introduce the notion of **torsion** and **curvature** of a connection.

Let $\nabla : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ be a connection on \mathfrak{g} . For all $x, y, z \in \mathfrak{g}_{\bar{i}}$, the *torsion* T^∇ of ∇ is defined as

$$(4) \quad T(x, y) := \nabla_x y - (-1)^{|x||y|}\nabla_y x - [x, y].$$

The *curvature* R^∇ of ∇ is defined as

$$(5) \quad R^\nabla(x, y)h := \nabla_x \nabla_y z - (-1)^{|x||y|}\nabla_y \nabla_x z - \nabla_{[x, y]}z.$$

By bilinearity, the definition of T and R^∇ can be extended to the whole \mathfrak{g} .

If $T = 0$, then the connection ∇ is called *torsion-free*. The curvature R^∇ vanishes if and only if the map $\sigma^\nabla : \mathfrak{g} \rightarrow \text{End}(\mathfrak{g})$, $x \mapsto \nabla_x$ is a representation of \mathfrak{g} on itself. In this case, the connection ∇ will be called *flat*.

2.2. Left-Symmetric Lie (super)algebras. Let us review a few notions regarding left-symmetric structures.

A superalgebra $L = L_{\bar{0}} \oplus L_{\bar{1}}$ over a field \mathbb{K} equipped with an even bilinear operation \cdot is called a left-symmetric superalgebra (or an LSSA for short) if the associator

$$(x, y, z) := (x \cdot y) \cdot z - x \cdot (y \cdot z)$$

is supersymmetric in x and y , i.e., $(x, y, z) = (-1)^{|x||y|}(y, x, z)$; or, equivalently,

$$(x \cdot y) \cdot z - x \cdot (y \cdot z) = (-1)^{|x||y|}((y \cdot x) \cdot z - y \cdot (x \cdot z)), \text{ for all } x, y, z \in L_{\bar{i}}.$$

These superalgebras are Lie-admissible. The supercommutator $[x, y] := x \cdot y - (-1)^{|x||y|}y \cdot x$ defines then a Lie superalgebra structure on L and we denote it by $\mathfrak{g}(L)$. The resulting Lie superalgebra is called the associated Lie superalgebra of the LSSA. Note that if L admits an LSSA then the even part $L_{\bar{0}}$ also admits one. Associative superalgebras are all LSSAs.

In [H], Helmstetter showed that it is necessary to have $[L, L] \subsetneq L$ if L is a left symmetric algebra defined over the field of complex numbers. Over a field of characteristic p , this is not true anymore. Burde showed in [B2] that classical Lie algebras over a field $p > 3$ admit an LSS only in case p divides $\dim(L)$. In the super setting and over a field of prime characteristic, there has not been much research done so far.

The existence of a flat connection ∇ on a Lie superalgebra \mathfrak{g} will induce a left-symmetric structure given by

$$x \cdot y := \nabla_x(y) \text{ for all } x, y \in \mathfrak{g}.$$

Additionally, every quasi-Frobenius Lie superalgebra (\mathfrak{g}, ω) admits a left symmetric structure given by

$$\omega(x \cdot y, z) = (-1)^{|x||y|}\omega(y, [x, z]) \text{ for all } x, y \in \mathfrak{g}_{\bar{i}}, \text{ and } z \in \mathfrak{g}.$$

The converse is not always true. There will be several examples that illustrate that in Section 5.

Some subclasses of LSSAs are important as well. An LSSA with supercommutative right multiplication is called a *Novikov superalgebra* (cf. [BN, Xu]). Explicitly, a *Novikov superalgebra* (L, \cdot) is an LSSA with an additional condition:

$$(z \cdot x) \cdot y = (-1)^{|x||y|}(z \cdot y) \cdot x, \text{ for all } x, y \in L_{\bar{i}}, \text{ and } z \in L.$$

It turns out that there are two superizations of the notion of Novikov algebras. The second superization is due to [Bal] (see also [PB]) and will be referred to as *Balinsky Novikov superalgebras*, or *BN superalgebras* for short. They are defined as follows:

(1) Left symmetry and commutativity:

$$(x \cdot y) \cdot z - x \cdot (y \cdot z) = (y \cdot x) \cdot z - y \cdot (x \cdot z), \quad (x \cdot y) \cdot z = (x \cdot z) \cdot y,$$

for any homogeneous elements

$$x, y, z \in L_{\bar{0}}; \quad x, y \in L_{\bar{0}}, z \in L_{\bar{1}}; \quad x, z \in L_{\bar{0}}, y \in L_{\bar{1}}; \quad y, z \in L_{\bar{0}}, x \in L_{\bar{1}};$$

(2) Commutativity: $x \cdot y = y \cdot x$ for all $x, y \in L_{\bar{1}}$;

(3) Compatibility conditions:

$$(x \cdot y) \cdot z = (x \cdot z) \cdot y \quad \text{for all } x, y \in L_{\bar{1}} \text{ and } z \in L_{\bar{0}};$$

$$x \cdot (y \cdot z) = (x \cdot y) \cdot z + y \cdot (x \cdot z) \quad \text{for all } x \in L_{\bar{0}} \text{ and } y, z \in L_{\bar{1}};$$

$$x \cdot (y \cdot z) = (x \cdot y) \cdot z + (x \cdot z) \cdot y \quad \text{for all } x, y, z \in L_{\bar{1}}.$$

Balinsky Novikov superalgebras are also admissible Lie superalgebras. The Lie structure is given as follows:

$$[x, y] := x \cdot y - y \cdot x, \quad \text{for all } x, y \in L_{\bar{0}};$$

$$[x, y] := x \cdot y - \frac{1}{2}y \cdot x = -[y, x], \quad \text{for all } x \in L_{\bar{0}}, \text{ and } y \in L_{\bar{1}}$$

$$[x, y] := x \cdot y \quad \text{for all } x, y \in L_{\bar{1}}.$$

2.3. The 4-dimensional real Lie superalgebras. The ground field \mathbb{K} is assumed to be the field of real numbers \mathbb{R} . The classification of four-dimensional real Lie superalgebras has been carried out by Backhouse, see [B]. The Lie superalgebras with anti-symmetric bilinear forms were examined in [BM].

As in the [BM] paper, the tables in the Appendix represent the classification of 4-dimensional real Lie superalgebras as follows:

- (1) **Table 1** having the *trivial algebras*, i.e. $[\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] = \{0\}$ and with $\text{sdim} = 2|2$;
- (2) **Table 2** having the *non-trivial algebras*, i.e. $[\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] \neq \{0\}$ and with $\text{sdim} = 2|2$;
- (3) **Table 3** having the *trivial algebras*, i.e. $[\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] = \{0\}$ and with $\text{sdim} = 3|1$;
- (4) **Table 4** having the *trivial algebras*, i.e. $[\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] = \{0\}$ and with $\text{sdim} = 1|3$;
- (5) **Table 5** having the *non-trivial algebras*, i.e. $[\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] \neq \{0\}$ and with $\text{sdim} = 3|1$;
- (6) **Table 6** having the *non-trivial algebras*, i.e. $[\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] \neq \{0\}$ and with $\text{sdim} = 1|3$.

In this paper, we present only the forms as in Table 1-6 which are suitable for Lagrangian extensions, as opposed to [BM], where the forms are given in the form of parametric families. The classification of these structures up to a symplectomorphism is still an open problem. We also correct a claim in [BM] stating that only non-homogenous forms exist for the Lie superalgebras $(D_0^{10})^1$ and $(D_0^{10})^2$. It turns out that these two superalgebras both admit an even and an odd non-degenerate closed form, and both of them can be obtained as Lagrangian extensions.

3. THE NOTION OF T^* -EXTENSIONS AND ΠT^* - (LAGRANGIAN EXTENSIONS)

We will introduce the notion of T^* -extension and ΠT^* -extension of Lie superalgebras (Lagrangian extensions). Originally, the notion of T^* -extension for Lie algebras was first

introduced by Bordemann in [Bor]. Recently, it was studied in [BC] in the context of quasi-Frobenius Lie algebras, and then in [BM] in the context of quasi-Frobenius Lie superalgebras.

3.1. Flat connections and representations. Let (\mathfrak{h}, ∇) be a Lie superalgebra endowed with a flat connection ∇ . Since the connection ∇ is flat, the map

$$\mathfrak{h} \rightarrow \text{End}(\mathfrak{h}) \quad u \mapsto \nabla_u,$$

defines a representation. We define a map $\rho : \mathfrak{h} \rightarrow \text{End}(\mathfrak{h}^*)$, the dual representation, as follows (for all $u \in \mathfrak{h}$ and $\xi \in \mathfrak{h}^*$):

$$(6) \quad \rho : \mathfrak{h} \rightarrow \text{End}(\mathfrak{h}^*) \quad u \mapsto \rho(u), \text{ where } \rho(u) \cdot \xi = -(-1)^{|u||\xi|} \xi \circ \nabla_u.$$

Similarly, we have a representation

$$(7) \quad \chi : \mathfrak{h} \rightarrow \text{End}(\Pi(\mathfrak{h}^*)) \quad u \mapsto \chi(u) := (-1)^{|u|} \Pi \circ \rho(u) \circ \Pi.$$

Lemma 3.1 (Dual representation). *The maps (6) and (7) are indeed representations.*

Proof. See [BM]. □

3.2. Polarization and Lagrangian extensions. Following [BC] (for the supercase, see [BM]), a *polarization* for a quasi-Frobenius Lie superalgebra (\mathfrak{g}, ω) is a choice of a Lagrangian subalgebra \mathfrak{l} of (\mathfrak{g}, ω) . A *strong polarization* of a quasi-Frobenius Lie superalgebra (\mathfrak{g}, ω) is a pair (\mathfrak{a}, N) consisting of a Lagrangian *ideal* $\mathfrak{a} \subset \mathfrak{g}$ and a complementary Lagrangian *subspace* $N \subset \mathfrak{g}$. The quadruple $(\mathfrak{g}, \omega, \mathfrak{a}, N)$ is referred to as a *strongly polarized* quasi-Frobenius Lie superalgebra.

By means of a 2-cocycle $\alpha \in Z^2(\mathfrak{h}, \mathfrak{h}^*)$ (resp. $\beta \in Z^2(\mathfrak{h}, \Pi(\mathfrak{h}^*))$) we will construct a Lie superalgebra structure on $\mathfrak{g} := \mathfrak{h} \oplus \mathfrak{h}^*$ (resp. $\mathfrak{g} := \mathfrak{h} \oplus \Pi(\mathfrak{h}^*)$), called T^* -extension (resp. ΠT^* -extension) (Lagrangian extension). By construction, we will see that these Lie superalgebras are strongly polarized quasi-Frobenius.

Recall that the space \mathfrak{h}^* and $\Pi(\mathfrak{h}^*)$ are \mathfrak{h} -modules by means of the representations (6) and (7).

On the space $\mathfrak{g} := \mathfrak{h} \oplus \mathfrak{h}^*$: The brackets are defined as follows: (for all $u, v \in \mathfrak{h}$ and $\xi \in \mathfrak{h}^*$)

$$[u, v]_{\mathfrak{g}} := [u, v]_{\mathfrak{h}} + \alpha(u, v), \quad [u, \xi]_{\mathfrak{g}} := \rho(u) \cdot \xi.$$

An **even** form ω is defined on \mathfrak{g} as follows: (for all $u + \xi, v + \zeta \in \mathfrak{h} \oplus \mathfrak{h}^*$)

$$(8) \quad \omega(u + \xi, v + \zeta) = \xi(v) - (-1)^{|\zeta||u|} \zeta(u).$$

This construction will be referred to as the T^* -extension.

On the space $\mathfrak{k} := \mathfrak{h} \oplus \Pi(\mathfrak{h}^*)$: The brackets are defined as follows: (for any $u, v \in \mathfrak{a}$)

$$[u, v]_{\mathfrak{g}} := [u, v]_{\mathfrak{h}} + \beta(u, v), \quad [u, \Pi(\xi)]_{\mathfrak{k}} := \chi(u) \cdot \Pi(\xi).$$

We define an **odd** form κ on \mathfrak{k} as follows: (for all $u + \Pi(\xi), v + \Pi(\zeta) \in \mathfrak{h} \oplus \Pi(\mathfrak{h}^*)$)

$$(9) \quad \kappa(u + \Pi(\xi), v + \Pi(\zeta)) = \xi(v) - (-1)^{(|\zeta|+1)|u|} \zeta(u).$$

This construction will be referred to as the ΠT^* -*extension*.

Lemma 3.2 (Conditions on α and β). (i) *The form ω is closed on \mathfrak{g} if and only if*

$$(10) \quad (-1)^{|u||w|} \alpha(u, v)(w) + \circlearrowleft(u, v, w) = 0 \text{ for all } u, v, w \in \mathfrak{h}.$$

(ii) *The form κ is closed on \mathfrak{g} if and only if*

$$(11) \quad (-1)^{|u||w|} \Pi \circ \beta(u, v)(w) + \circlearrowleft(u, v, w) = 0 \text{ for all } u, v, w \in \mathfrak{h}.$$

Proof. See, [BM]. □

We arrive to the following theorem proved in [BM].

Theorem 3.3 (Lagrangian or T^* - and ΠT^* -extensions). *Let (\mathfrak{h}, ∇) be a flat Lie superalgebra. To every 2-cocycle $\alpha \in Z^2(\mathfrak{h}, \mathfrak{h}^*)$ (resp. $\beta \in Z^2(\mathfrak{h}, \Pi(\mathfrak{h}^*))$) which satisfies (10) (resp. (11)) one can canonically associate a strongly polarized quasi-Frobenius Lie superalgebra $(\mathfrak{g}, \omega, \mathfrak{h}^*, \mathfrak{h})$ (resp. $(\mathfrak{k}, \kappa, \Pi(\mathfrak{h}^*), \mathfrak{h})$), where $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{h}^*$ (resp. $\mathfrak{k} = \mathfrak{h} \oplus \Pi(\mathfrak{h}^*)$) and ω (resp. κ) is given as in (8) (resp. (9)).*

The converse of this theorem says that every orthosymplectic or periplectic quasi-Frobenius Lie superalgebra (\mathfrak{g}, ω) that has a Lagrangian ideal \mathfrak{a} can be obtained as a T^* -extension or ΠT^* -extension of a flat Lie superalgebra.

Let $(\mathfrak{g}, \omega, \mathfrak{a}, N)$ be a strongly polarized quasi-Frobenius Lie superalgebra. Here the form ω is either orthosymplectic or periplectic. Consider the quotient space $\mathfrak{h} = \mathfrak{g}/\mathfrak{a}$ (recall that $\mathfrak{a}^\perp = \mathfrak{a}$). We will construct a flat connection ∇ on \mathfrak{h} (see [BM]), superizing the construction of [BC]. For every $u, v \in \mathfrak{h}$, we denote by \tilde{u} and \tilde{v} their lift to \mathfrak{g} . We then write

$$(12) \quad \omega_{\mathfrak{h}}(\nabla_u v, a) = -(-1)^{|u||v|} \omega_{\mathfrak{g}}(\tilde{u}, [\tilde{v}, a]).$$

The pair (\mathfrak{h}, ∇) is called *the quotient flat Lie superalgebra associated with the Lagrangian ideal \mathfrak{a} of (\mathfrak{g}, ω)* .

Theorem 3.4 (Converse of Theorem 3.3). *Let $(\mathfrak{g}, \omega, \mathfrak{a}, N)$ be a strongly polarized orthosymplectic or periplectic quasi-Frobenius Lie superalgebra and (\mathfrak{h}, ∇) be its associated quotient flat Lie superalgebra.*

(i) *If the form ω is orthosymplectic, then there exists $\alpha \in Z^2(\mathfrak{h}, \mathfrak{h}^*)$ satisfying Eq. (10), such that $(\mathfrak{g}, \omega, \mathfrak{a}, N)$ is isomorphic to the T^* -extension of (\mathfrak{h}, ∇) by means of α .*

(ii) *If the form ω is periplectic, then there exists $\beta \in Z^2(\mathfrak{h}, \Pi(\mathfrak{h}^*))$ satisfying Eq. (11), such that $(\mathfrak{g}, \omega, \mathfrak{a}, N)$ is isomorphic to the ΠT^* -extension of (\mathfrak{h}, ∇) by means of β .*

Proof. See [BM]. □

4. THE 4-DIMENSIONAL LIE SUPERALGEBRAS AS LAGRANGIAN EXTENSIONS

Using Theorem 3.3 and Theorem 3.4 above, we will determine whether the 4-dimensional real Lie superalgebras classified by Backhouse (see the Appendix) are isomorphic to a T^* -extension or to a ΠT^* -extension of a smaller Lie superalgebra.

We will outline the proof and computation for the Lie superalgebra D^6 (row 2 in the table); however, we will only exhibit the connection ∇ for the other Lie superalgebras.

- (1) The Lie superalgebra D^5 : We correct here the description of the Lie superalgebra D_5 in [BM]. It is ΠT^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1|e_4\}$ with the bracket $[e_1, e_4]_{\mathfrak{h}} = e_4$, and where $\Pi\mathfrak{h}^*$ is spanned by $e_2 = -\Pi(e_4^*)|e_3 = \Pi(e_1^*)$. The 2-form $\beta \equiv 0$ and the connection ∇ on \mathfrak{h} is given by:

$$\nabla_{e_4}(e_1) = -e_4, \quad \nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(e_4) = 0, \quad \nabla_{e_4}(e_4) = 0.$$

- (2) The Lie superalgebra D^6 : We will show first that this Lie superalgebra does have a Lagrangian ideal. Indeed, consider $\mathfrak{a} = \text{Span}\{e_3, e_4\}$. This is obviously an ideal because of the multiplication table. Let us show it is Lagrangian. We have

$$\mathfrak{a}^\perp = \{v \in D^6 \mid \omega(v, x) = 0, \forall x \in \mathfrak{a}\},$$

where the form we are considering is given by $\omega = e_3^* \wedge e_1^* + e_4^* \wedge e_2^*$. Let $v = l_1 e_1 + l_2 e_2 + l_3 e_3 + l_4 e_4$. We have

$$0 = \omega(v, e_3) = -l_1 \implies l_1 = 0,$$

$$0 = \omega(v, e_4) = -l_2 \implies l_2 = 0.$$

This implies that $\mathfrak{a}^\perp = \mathfrak{a}$. The Lie superalgebra \mathfrak{h} is given by $\text{Span}\{e_1, e_2\}$ with the bracket $[e_1, e_2]_{\mathfrak{h}} = 0$. The connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(e_2) = -e_2, \quad \nabla_{e_2}(e_1) = -e_2, \quad \nabla_{e_2}(e_2) = e_1.$$

Now, we compute the torsion and curvature of ∇ , using Eqns. (4, 5). For the torsion, we have:

$$T(e_1, e_2) := \nabla_{e_1}(e_2) - (-1)^{|e_1||e_2|}\nabla_{e_2}(e_1) - [e_1, e_2]_{\mathfrak{h}} = -e_2 - (-e_2) - 0 = 0.$$

For the curvature, we have

$$\begin{aligned} R^\nabla(e_1, e_2)(e_1) &:= \nabla_{e_1}\nabla_{e_2}(e_1) - (-1)^{|e_1||e_2|}\nabla_{e_2}\nabla_{e_1}(e_1) - \nabla_{[e_1, e_2]_{\mathfrak{h}}}(e_1) \\ &= \nabla_{e_1}(-e_2) - \nabla_{e_2}(-e_1) = e_2 - e_2 = 0. \end{aligned}$$

Similarly,

$$\begin{aligned} R^\nabla(e_1, e_2)(e_2) &:= \nabla_{e_1}\nabla_{e_2}(e_2) - (-1)^{|e_1||e_2|}\nabla_{e_2}\nabla_{e_1}(e_2) - \nabla_{[e_1, e_2]_{\mathfrak{h}}}(e_2) \\ &= \nabla_{e_1}(e_1) - \nabla_{e_2}(-e_2) = -e_1 + e_1 = 0. \end{aligned}$$

Therefore, the connection is torsion-free and flat.

Thus, The Lie superalgebra D^6 is a ΠT^* -extension of the Lie superalgebra \mathfrak{h} , where $\Pi\mathfrak{h}^*$ is spanned by $e_3 = \Pi(e_1^*)|e_4 = \Pi(e_2^*)$. The 2-form $\beta \equiv 0$.

(3) The Lie superalgebra D_{pq}^7 : This Lie superalgebra is not quasi-Frobenius.

(4) The Lie superalgebra D_{-1q}^7 , with $q \leq 1$: This Lie superalgebra is a ΠT^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1, e_2\}$ with the bracket $[e_1, e_2]_{\mathfrak{h}} = e_2$, and where $\Pi\mathfrak{h}^*$ is spanned by $e_4 = \Pi(e_1^*), e_3 = \Pi(e_2^*)$. The 2-form $\beta \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -qe_1, \quad \nabla_{e_1}(e_2) = e_2, \quad \nabla_{e_2}(e_1) = 0, \quad \nabla_{e_2}(e_2) = 0.$$

(5) The Lie superalgebra D_{pq}^7 , with $p = -q, q \neq 0, -1$: This Lie superalgebra is a T^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1|e_4\}$ with the bracket $[e_1, e_4]_{\mathfrak{h}} = -pe_4$, and where \mathfrak{h}^* is spanned by $e_2 = e_1^*|e_3 = e_4^*$. The 2-form $\alpha \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(e_4) = -pe_4, \quad \nabla_{e_4}(e_1) = 0, \quad \nabla_{e_4}(e_4) = 0.$$

(6) The Lie superalgebra D_{pq}^7 , with $p = -q, q = -1$: In the case where the form is odd, then this Lie superalgebra is a ΠT^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1, e_2\}$ with the bracket $[e_1, e_2]_{\mathfrak{h}} = e_2$, and where $\Pi\mathfrak{h}^*$ is spanned by $e_3 = \Pi(e_1^*), e_4 = \Pi(e_2^*)$. The 2-form $\beta \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(e_2) = e_2, \quad \nabla_{e_2}(e_1) = 0, \quad \nabla_{e_2}(e_2) = 0.$$

In the case where the form is even, then this Lie superalgebra is a T^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1|e_4\}$ with the bracket $[e_1, e_4]_{\mathfrak{h}} = -e_4$, and where \mathfrak{h}^* is spanned by $e_2 = e_1^*|e_3 = e_4^*$. The 2-form $\alpha \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(e_4) = -e_4, \quad \nabla_{e_4}(e_1) = 0, \quad \nabla_{e_4}(e_4) = 0.$$

(7) The Lie superalgebra D_p^8 , with $p \neq 0$: This Lie superalgebra is not quasi-Frobenius.

(8) The Lie superalgebra D_{-1}^8 : This Lie superalgebra is a ΠT^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1, e_2\}$ with the bracket $[e_1, e_2]_{\mathfrak{h}} = e_2$, and where $\Pi\mathfrak{h}^*$ is spanned by $e_3 = \Pi(e_1^*), e_4 = \Pi(e_2^*)$. The 2-form $\beta \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = e_1 - e_2, \quad \nabla_{e_1}(e_2) = e_2, \quad \nabla_{e_2}(e_1) = 0, \quad \nabla_{e_2}(e_2) = 0.$$

- (9) The Lie superalgebra D_{pq}^9 , with $q > 0$: This Lie superalgebra is not quasi-Frobenius.
- (10) The Lie superalgebra D_q^{10} with $q \neq -1$: This Lie superalgebra is a ΠT^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1|e_4\}$ with the bracket $[e_1, e_4]_{\mathfrak{h}} = qe_4$, and where $\Pi\mathfrak{h}^*$ is spanned by $e_2 = \Pi(e_4^*)|e_3 = -(q+1)\Pi(e_1^*)$. The 2-form $\beta \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -(q+1)e_1, \quad \nabla_{e_1}(e_4) = -e_4, \quad \nabla_{e_4}(e_1) = -(q+1)e_4, \quad \nabla_{e_4}(e_4) = 0.$$

- (11) The Lie superalgebra $(D_{1/2, 1/2}^7)^1$: This Lie superalgebra does not admit a 2-dimensional Lagrangian ideal. Indeed, let us suppose the contrary, and let us denote by $\mathfrak{a} = \text{Span}\{X, Y\}$ such an ideal. Let us write $X = x_1e_1 + x_2e_2 + x_3e_3 + x_4e_4$ and $Y = y_1e_1 + y_2e_2 + y_3e_3 + y_4e_4$. We compute the brackets $[e_i, X]$ for all $i \in \{1, 2, 3, 4\}$. we get

$$\begin{aligned} [e_1, X] &= x_2e_2 + \frac{1}{2}x_3e_3 + \frac{1}{2}x_4e_4, & [e_2, X] &= -x_1e_2, \\ [e_3, X] &= -\frac{1}{2}x_1e_3 + x_3e_2, & [e_4, X] &= -\frac{1}{2}x_1e_4 + x_4e_2. \end{aligned}$$

If $x_1 \neq 0$, from the second bracket we get that $e_2 \in \mathfrak{a}$, followed by $e_3 \in \mathfrak{a}$ and $e_4 \in \mathfrak{a}$, which implies that the ideal is not of dimension 2. We are in the case where $x_1 = 0$ (and similarly $y_1 = 0$).

$$\begin{aligned} [e_1, X] &= x_2e_2 + \frac{1}{2}x_3e_3 + \frac{1}{2}x_4e_4, & [e_2, X] &= 0, \\ [e_3, X] &= x_3e_2, & [e_4, X] &= x_4e_2. \end{aligned}$$

We consider the following cases:

- (a) If $x_3 = x_4 = 0$. In this case, $e_2 \in \mathfrak{a}$ and we can write $X = e_2$ and $Y = y_2e_2 + y_3e_3 + y_4e_4$. We can simplify and write $Y = z_3e_3 + z_4e_4$. Since¹ $\omega(Y, Y) = z_3^2 + z_4^2 \neq 0$, it follows that \mathfrak{a} is not Lagrangian.
- (b) If $x_3 = 0$, and $x_4 \neq 0$. Looking at brackets again we get $[e_1, X] = x_2e_2 + \frac{1}{2}x_4e_4$, and $[e_4, X] = x_4e_2$, which implies $e_2, e_4 \in \mathfrak{a}$, so $\mathfrak{a} = \text{Span}\{e_2, e_4\}$. We have to check that \mathfrak{a} is not Lagrangian. Indeed, $\omega(e_4, e_4) = 1$.
- (c) If $x_3 \neq 0$, and $x_4 = 0$. This case will be similar to the previous one since the equations are symmetrical in x_3 and x_4 , which implies that no Lagrangian ideal exists.
- (d) If $x_3 \neq 0$, and $x_4 \neq 0$. Since all the previous cases showed that we can't find a Lagrangian ideal, and since the proof is symmetrical in X and Y without loss

¹As in [BM, BEM, BE], we adopt the following convention: $\langle e_i^*, e_j \rangle = \delta_{ij}$ and $\langle e_i^* \otimes e_j^*, e_k \otimes e_l \rangle = (-1)^{|e_k||e_j|} \langle e_i^*, e_k \rangle \langle e_j^*, e_l \rangle$.

of generality we can state that $y_3 \neq 0, y_4 \neq 0$. This implies that $e_2 \in \mathfrak{a}$ and that $x_2e_2 + \frac{1}{2}x_3e_3 + \frac{1}{2}x_4e_4 \in \mathfrak{a}$. We can write $X = e_2$ and $Y = z_3e_3 + z_4e_4$.

Once again, $\omega(Y, Y) = z_3^2 + z_4^2 \neq 0$ which implies that \mathfrak{a} is not Lagrangian.

- (12) The Lie superalgebra $(D_{1/2, 1/2}^7)^2$: This Lie superalgebra is a T^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1|x := e_3 - e_4\}$ with the bracket $[e_1, x]_{\mathfrak{h}} = \frac{1}{2}x$, and where \mathfrak{h}^* is spanned by $e_2 = e_1^*|e_3 + e_4 = -2x^*$. The 2-form $\alpha \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(x) = -\frac{1}{2}x, \quad \nabla_x(e_1) = -x, \quad \nabla_x(x) = 0.$$

- (13) The Lie superalgebra $(D_{1/2, 1/2}^7)^3$: This Lie superalgebra is not quasi-Frobenius.

- (14) The Lie superalgebra $(D_{1-p, p}^7)$, with $p \leq \frac{1}{2}$: This Lie superalgebra is a T^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1|e_4\}$ with the bracket $[e_1, e_4]_{\mathfrak{h}} = (1-p)e_4$, and where \mathfrak{h}^* is spanned by $e_2 = -e_1^*|e_3 = e_4^*$. The 2-form $\alpha \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(e_4) = -pe_4, \quad \nabla_{e_4}(e_1) = -e_4, \quad \nabla_{e_4}(e_4) = 0.$$

- (15) The Lie superalgebra $(D_{1/2}^8)$: This Lie superalgebra is not quasi-Frobenius.

- (16) The Lie superalgebra $(D_{1/2, p}^9)$, with $p > 0$: This Lie superalgebra does not admit a 2-dimensional Lagrangian ideal. Indeed, let us suppose there exists one, denoted by $\mathfrak{a} = \text{Span}\{X, Y\}$. Let us write $X = x_1e_1 + x_2e_2 + x_3e_3 + x_4e_4$ and $Y = y_1e_1 + y_2e_2 + y_3e_3 + y_4e_4$. We compute the brackets $[e_i, X]$ for all $i \in \{1, 2, 3, 4\}$. We get

$$\begin{aligned} [e_1, X] &= x_2e_2 + \frac{1}{2}x_3e_3 - x_3pe_4 + x_4pe_3 + \frac{1}{2}x_4e_4, \\ [e_2, X] &= -x_1e_2, \\ [e_3, X] &= -\frac{1}{2}x_1e_3 + px_1e_4 + x_3e_2, \\ [e_4, X] &= -\frac{1}{2}x_1e_4 - px_1e_3 + x_4e_2. \end{aligned}$$

We will analyze a few cases. If $x_1 \neq 0$, given the second bracket we get that $e_2 \in \mathfrak{a}$ and according to the last brackets $e_3 \in \mathfrak{a}$ and $e_4 \in \mathfrak{a}$. This case is not possible

since \mathfrak{a} has dimension 2. This implies that $x_1 = 0$. We get then:

$$\begin{aligned} [e_1, X] &= x_2 e_2 + \frac{1}{2} x_3 e_3 - x_3 p e_4 + x_4 p e_3 + \frac{1}{2} x_4 e_4, & [e_2, X] &= 0, \\ [e_3, X] &= x_3 e_2, & [e_4, X] &= x_4 e_2. \end{aligned}$$

For all the cases $x_3, x_4 = 0$ or $\neq 0$ we get that $e_2 \in \mathfrak{a}$. We can write $X = e_2$ and $Y = z_3 e_3 + z_4 e_4$. As $\omega(Y, Y) \neq 0$, it follows that \mathfrak{a} is not Lagrangian.

Thus, this Lie superalgebra does not admit a 2-dimensional Lagrangian ideal.

- (17) The Lie superalgebra $(D_0^{10})^1$: In the case where the form is odd, this Lie superalgebra is a ΠT^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1|e_4\}$ with the bracket $[e_4, e_4]_{\mathfrak{h}} = e_1$, and where \mathfrak{h}^* is spanned by $e_2 = -\Pi(e_4^*)|e_3 = \Pi(e_1^*)$. The 2-form $\beta \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(e_4) = -e_4, \quad \nabla_{e_4}(e_1) = -e_4, \quad \nabla_{e_4}(e_4) = \frac{1}{2}e_1.$$

In the case where the form is even, this Lie superalgebra is a T^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1|e_4\}$ with the bracket $[e_4, e_4]_{\mathfrak{h}} = e_1$, and where \mathfrak{h}^* is spanned by $e_2 = 2e_1^*|e_3 = e_4^*$. The 2-form $\alpha \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(e_4) = -e_4, \quad \nabla_{e_4}(e_1) = -e_4, \quad \nabla_{e_4}(e_4) = \frac{1}{2}e_1.$$

- (18) The Lie superalgebra $(D_0^{10})^2$: In the case where the form is odd, this Lie superalgebra is a ΠT^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1|e_4\}$ with the bracket $[e_4, e_4]_{\mathfrak{h}} = -e_1$, and where \mathfrak{h}^* is spanned by $e_2 = -\Pi(e_4^*)|e_3 = \Pi(e_1^*)$. The 2-form $\beta \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(e_4) = -e_4, \quad \nabla_{e_4}(e_1) = -e_4, \quad \nabla_{e_4}(e_4) = -\frac{1}{2}e_1.$$

In the case where the form is even, this Lie superalgebra is a T^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1|e_4\}$ with the bracket $[e_4, e_4]_{\mathfrak{h}} = -e_1$, and where \mathfrak{h}^* is spanned by $e_2 = 2e_1^*|e_3 = e_4^*$. The 2-form $\alpha \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(e_4) = -e_4, \quad \nabla_{e_4}(e_1) = -e_4, \quad \nabla_{e_4}(e_4) = -\frac{1}{2}e_1.$$

- (19) The Lie superalgebra $(2A_{1,1} + 2A)^1$: This Lie superalgebra is not quasi-Frobenius.

- (20) The Lie superalgebra $(2A_{1,1} + 2A)^2$: This Lie superalgebra is not quasi-Frobenius.

- (21) The Lie superalgebra $(2A_{1,1} + 2A)_p^3$, for $p = \frac{1}{2}$: This Lie superalgebra is a ΠT^* -extension of the abelian Lie superalgebra $\mathfrak{h} = \text{Span}\{e_3, e_4\}$, and where $\Pi\mathfrak{h}^*$ is spanned by $e_1 = -\Pi(e_4^*)$, $e_2 = \Pi(e_3^*)$. The connection ∇ is trivial, and the 2-form β is given by

$$\beta = \frac{1}{2}\Pi(e_4^*) \otimes e_3^* \wedge e_3^* - \frac{1}{2}\Pi(e_3^*) \otimes e_4^* \wedge e_4^* - \frac{1}{2}(\Pi(e_3^*) - \Pi(e_4^*)) \otimes e_3^* \wedge e_4^*.$$

- (22) The Lie superalgebra $(2A_{1,1} + 2A)_p^4$: This Lie superalgebra is not quasi-Frobenius.
 (23) The Lie superalgebra $(C_1^1 + A)$: This Lie superalgebra is a T^* -extension of the abelian Lie superalgebra $\mathfrak{h} = \text{Span}\{e_1 | x := e_3 - 2e_4\}$, and where \mathfrak{h}^* is spanned by $e_2 = \frac{1}{2}e_1^* | e_3 = x^*$. The 2-form α is given by:

$$\alpha = e_1^* \otimes (x^* \wedge x^*) + x^* \otimes (e_1^* \wedge x^*)$$

The connection ∇ is given by:

$$\nabla_{e_1}(e_1) = -e_1, \quad \nabla_{e_1}(x) = -x, \quad \nabla_x(e_1) = -x, \quad \nabla_x(x) = 0.$$

- (24) The Lie superalgebra $(C_{1/2}^1 + A)$: This Lie superalgebra does not admit a 2-dimensional Lagrangian ideal. Indeed, let us suppose there is one, denoted by $\mathfrak{a} = \text{Span}\{X, Y\}$. Let us write $X = x_1e_1 + x_2e_2 + x_3e_3 + x_4e_4$ and $Y = y_1e_1 + y_2e_2 + y_3e_3 + y_4e_4$. We compute the brackets $[e_i, X]$ for all $i \in \{1, 2, 3, 4\}$. We get:

$$[e_1, X] = x_2e_2 + \frac{1}{2}x_3e_3, \quad [e_2, X] = -x_1e_2, \quad [e_3, X] = -\frac{1}{2}x_1e_3 + x_3e_2, \quad [e_4, X] = 0.$$

We will analyze a few cases.

If $x_1 \neq 0$, given the second bracket implies that $e_2 \in \mathfrak{a}$ and according to the third bracket $e_3 \in \mathfrak{a}$, which implies that $\mathfrak{a} = \text{Span}\{e_2, e_3\}$. Since $\omega(e_2, e_3) \neq 0$, it follows that \mathfrak{a} is not Lagrangian.

If $x_1 = 0$, and $x_3 \neq 0$, we get that $e_2 \in \mathfrak{a}$, followed by $e_3 \in \mathfrak{a}$, so $\mathfrak{a} = \text{Span}\{e_2, e_3\}$, which we showed is not possible. It follows that $x_3 = 0$, and using the same reasoning as above, we also conclude that for Y , the scalars $y_1 = y_3 = 0$. Looking at the brackets, we either have $x_2 = 0$ or $e_2 \in \mathfrak{a}$. If $x_2 = 0$, then $X = x_4e_4$, and $Y = y_2e_2 + y_4e_4$. We can simplify and write $X = e_4$, and $Y = e_2$. Since $\omega(e_4, e_4) \neq 0$, it follows that \mathfrak{a} is not Lagrangian. On the other hand, if $x_2 \neq 0$, then $X = x_2e_2 + x_4e_4$, and $Y = y_2e_2 + y_4e_4$. We can simplify and write $X = e_2 + z_4e_4$, and $Y = e_4$. Again, \mathfrak{a} is not Lagrangian.

Thus, this Lie superalgebra does not admit a 2-dimensional Lagrangian ideal.

- (25) The Lie superalgebra $(C_{-1}^2 + A)$: This Lie superalgebra is not quasi-Frobenius.
 (26) The Lie superalgebra $(C^3 + A)$: This Lie superalgebra is a T^* -extension of the Lie superalgebra $\mathfrak{h} = \text{Span}\{e_2 | e_4\}$ with the bracket $[e_4, e_4]_{\mathfrak{h}} = e_2$, and where \mathfrak{h}^* is

spanned by $e_1 = 2e_2^* | e_3 = e_4^*$. The 2-form $\alpha \equiv 0$ and the connection ∇ is given by:

$$\nabla_{e_2}(e_2) = 0, \quad \nabla_{e_2}(e_4) = 0, \quad \nabla_{e_4}(e_2) = 0, \quad \nabla_{e_4}(e_4) = \frac{1}{2}e_2.$$

- (27) The Lie superalgebra $(C_0^5 + A)$: This Lie superalgebra is not quasi-Frobenius.
- (28) The Lie superalgebra D^1 : This Lie superalgebra doesn't admit a non-degenerate homogeneous closed form.
- (29) The Lie superalgebra D_q^2 , with $q \neq -1, 0$: This Lie superalgebra is not quasi-Frobenius.
- (30) The Lie superalgebra D_{-1}^2 : This Lie superalgebra doesn't admit a non-degenerate homogeneous closed form.
- (31) The Lie superalgebra D_{pq}^3 : This Lie superalgebra is not quasi-Frobenius.
- (32) The Lie superalgebra D_{pq}^{11} with $0 < |p| \leq |q| \leq 1$: This Lie superalgebra is not quasi-Frobenius.
- (33) The Lie superalgebra D_{pq}^{11} with $(p, q) = (-1, -1)$: This Lie superalgebra doesn't admit a non-degenerate homogeneous closed form.
- (34) The Lie superalgebra D_{pq}^{11} with $0 < |p| \leq 1, p = -q$: This Lie superalgebra doesn't admit a non-degenerate homogeneous closed form.
- (35) The Lie superalgebra D_{pq}^{11} with $0 < |p| < 1, q = -1$: This Lie superalgebra doesn't admit a non-degenerate homogeneous closed form.
- (36) The Lie superalgebra D^{12} : This Lie superalgebra doesn't admit a non-degenerate homogeneous closed form.
- (37) The Lie superalgebra D_p^{13} , with $p \neq 0$: This Lie superalgebra is not quasi-Frobenius.
- (38) The Lie superalgebra D_{-1}^{13} : This Lie superalgebra doesn't admit a non-degenerate homogeneous closed form.
- (39) The Lie superalgebra D_{pq}^{14} , with $p \neq 0, q \geq 0$: This Lie superalgebra is not quasi-Frobenius.
- (40) The Lie superalgebra D^{15} : This Lie superalgebra doesn't admit a non-degenerate homogeneous closed form.
- (41) The Lie superalgebra D^{16} : This Lie superalgebra is in fact not quasi-Frobenius.
- (42) The Lie superalgebra $(A_{3,1} + A)$: This Lie superalgebra is not quasi-Frobenius.
- (43) The Lie superalgebra $(D_{p,-1/2}^3)$, with $p \neq 0$: This Lie superalgebra is not quasi-Frobenius.
- (44) The Lie superalgebra $(D_{-1/2,-1/2}^3)$: This Lie superalgebra doesn't admit a non-degenerate homogeneous closed form.
- (45) The Lie superalgebra $(D_{-1/2}^2)^1$: This Lie superalgebra is not quasi-Frobenius.
- (46) The Lie superalgebra $(D_{-1/2}^2)^2$: This Lie superalgebra is not quasi-Frobenius.

- (47) The Lie superalgebra $(A_{1,1} + 3A)^1$: This Lie superalgebra is not quasi-Frobenius.
 (48) The Lie superalgebra $(A_{1,1} + 3A)^2$: This Lie superalgebra is not quasi-Frobenius.

5. LEFT-SYMMETRIC STRUCTURES ON THE 4-DIMENSIONAL REAL LIE SUPERALGEBRAS

In the list of 4-dimensional real Lie superalgebras, we will list left-symmetric structures that are compatible with the Lie structure. With the exception of $(D_0^{10})^1$ and $(D_0^{10})^2$, all of these structures are Novikov. Moreover, we show that each of these Lie superalgebras admit a Balinsky-Novikov structure. In line with standard practice, only non-zero products are displayed.

- (1) The Lie superalgebra D^5 : A Novikov structure is given by:

$$e_1 \cdot e_3 = e_3, \quad e_1 \cdot e_4 = e_4, \quad e_2 \cdot e_4 = e_3.$$

Additionally, this structure is also Balinsky-Novikov. As a matter of fact, the non-zero associators are given by:

$$(e_1, e_1, e_4) = -e_4, \quad (e_2, e_1, e_4) = -e_3, \quad (e_1, e_1, e_3) = -e_3, \quad (e_1, e_2, e_4) = -e_3.$$

- (2) The Lie superalgebra D^6 : A Novikov structure is given by:

$$e_1 \cdot e_3 = e_3, \quad e_1 \cdot e_4 = e_4, \quad e_2 \cdot e_3 = -e_4, \quad e_2 \cdot e_4 = e_3.$$

Additionally, this structure is also Balinsky-Novikov.

- (3) The Lie superalgebra D_{pq}^7 with $(p \geq q, pq \neq 0)$: A Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$e_1 \cdot e_1 = \gamma e_2, \quad e_1 \cdot e_2 = e_2, \quad e_1 \cdot e_3 = p e_3, \quad e_1 \cdot e_4 = q e_4.$$

Additionally, this structure is also Balinsky-Novikov.

- (4) The Lie superalgebra D_p^8 with $p \neq 0$: A Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$e_1 \cdot e_1 = \gamma e_2, \quad e_1 \cdot e_2 = e_2, \quad e_1 \cdot e_3 = p e_3, \quad e_1 \cdot e_4 = e_3 + p e_4.$$

Additionally, this structure is also Balinsky-Novikov.

- (5) The Lie superalgebra D_{pq}^9 with $q > 0$: A Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$e_1 \cdot e_1 = \gamma e_1, \quad e_1 \cdot e_2 = (1 + \gamma)e_2, \quad e_1 \cdot e_3 = (p + \gamma)e_3 - q e_4, \quad e_1 \cdot e_4 = q e_3 + (p + \gamma)e_4, \\ e_2 \cdot e_1 = \gamma e_2, \quad e_3 \cdot e_1 = \gamma e_3, \quad e_4 \cdot e_1 = \gamma e_4.$$

A Balinsky-Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$e_1 \cdot e_1 = \gamma e_2, \quad e_1 \cdot e_2 = e_2, \quad e_1 \cdot e_3 = p e_3 - q e_4, \quad e_1 \cdot e_4 = q e_3 + p e_4.$$

(6) The Lie superalgebra D_q^{10} : A Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -qe_1 + \gamma(q-1)e_2, & e_1 \cdot e_2 &= (1-q)e_2, & e_1 \cdot e_3 &= e_3, & e_1 \cdot e_4 &= \gamma e_3, \\ e_2 \cdot e_1 &= -qe_2, & e_3 \cdot e_1 &= -qe_3, & e_4 \cdot e_1 &= \gamma e_3 - qe_4, & e_4 \cdot e_2 &= -e_3. \end{aligned}$$

A Balinsky-Novikov structure is given by

$$\begin{aligned} e_1 \cdot e_1 &= -e_1, & e_1 \cdot e_3 &= \frac{1}{2}(1+2q)e_3, & e_1 \cdot e_4 &= \frac{1}{2}(2q-1)e_4, & e_2 \cdot e_1 &= -e_2, \\ e_2 \cdot e_4 &= e_3, & e_3 \cdot e_1 &= -e_3, & e_4 \cdot e_1 &= -e_4. \end{aligned}$$

(7) The Lie superalgebra $(D_{1/2, 1/2}^7)^1$: A Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -\frac{1}{2}e_1 + \lambda e_2, & e_1 \cdot e_2 &= \frac{1}{2}e_2, & e_2 \cdot e_1 &= -\frac{1}{2}e_2, & e_3 \cdot e_1 &= -\frac{1}{2}e_3, & e_3 \cdot e_3 &= \frac{1}{2}e_2, \\ e_3 \cdot e_4 &= \gamma e_2, & e_4 \cdot e_1 &= -\frac{1}{2}e_4, & e_4 \cdot e_3 &= -\gamma e_2, & e_4 \cdot e_4 &= \frac{1}{2}e_2. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= (\gamma-1)e_1 + \lambda e_2, & e_1 \cdot e_2 &= \gamma e_2, & e_1 \cdot e_3 &= \frac{\gamma}{2}e_3, & e_1 \cdot e_4 &= \frac{\gamma}{2}e_4, \\ e_2 \cdot e_1 &= (\gamma-1)e_2, & e_3 \cdot e_1 &= (\gamma-1)e_3, & e_3 \cdot e_3 &= e_2, & e_4 \cdot e_1 &= (\gamma-1)e_4, \\ e_4 \cdot e_4 &= e_2. \end{aligned}$$

(8) The Lie superalgebra $(D_{1/2, 1/2}^7)^2$: A Novikov structure is given by (where $\lambda \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -\frac{1}{2}e_1 + \lambda e_2, & e_1 \cdot e_2 &= \frac{1}{2}e_2, & e_2 \cdot e_1 &= -\frac{1}{2}e_2, & e_3 \cdot e_1 &= -\frac{1}{2}e_3, \\ e_3 \cdot e_3 &= \frac{1}{2}e_2, & e_4 \cdot e_1 &= -\frac{1}{2}e_4, & e_4 \cdot e_4 &= -\frac{1}{2}e_2. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= \gamma e_1 + \lambda e_2, & e_1 \cdot e_2 &= (1+\gamma)e_2, & e_1 \cdot e_3 &= \frac{1}{2}(1+\gamma)e_3, & e_1 \cdot e_4 &= \frac{1}{2}(1+\gamma)e_4, \\ e_2 \cdot e_1 &= \gamma e_2, & e_3 \cdot e_1 &= \gamma e_3, & e_3 \cdot e_3 &= e_2, & e_4 \cdot e_1 &= \gamma e_4, \\ e_4 \cdot e_4 &= -e_2. \end{aligned}$$

(9) The Lie superalgebra $(D_{1/2, 1/2}^7)^3$: A Novikov structure is given by (where $\lambda \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -\frac{1}{2}e_1 + \lambda e_2, & e_1 \cdot e_2 &= \frac{1}{2}e_2, & e_2 \cdot e_1 &= -\frac{1}{2}e_2, & e_3 \cdot e_1 &= -\frac{1}{2}e_3, \\ e_3 \cdot e_3 &= \frac{1}{2}e_2, & e_4 \cdot e_1 &= -\frac{1}{2}e_4. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= \gamma e_1 + \lambda e_2, & e_1 \cdot e_2 &= (1+\gamma)e_2, & e_1 \cdot e_3 &= \frac{1}{2}(1+\gamma)e_3, & e_1 \cdot e_4 &= \frac{1}{2}(1+\gamma)e_4, \\ e_2 \cdot e_1 &= \gamma e_2, & e_3 \cdot e_1 &= \gamma e_3, & e_3 \cdot e_3 &= e_2, & e_4 \cdot e_1 &= \gamma e_4. \end{aligned}$$

(10) The Lie superalgebra $(D_{1-p,p}^7)$ with $p \leq \frac{1}{2}$: A Novikov structure is given by (where $\lambda \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -pe_1 + \lambda e_2, & e_1 \cdot e_2 &= (1-p)e_2, & e_1 \cdot e_4 &= (1-2p)e_4, & e_2 \cdot e_1 &= -pe_2, \\ e_3 \cdot e_1 &= -pe_3, & e_3 \cdot e_4 &= e_2, & e_4 \cdot e_1 &= -pe_4. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= (\gamma-1)e_1 + \lambda e_2, & e_1 \cdot e_2 &= \gamma e_2, & e_1 \cdot e_3 &= \frac{1}{2}(2p + \gamma - 1)e_3, & e_1 \cdot e_4 &= \frac{1}{2}(1 - 2p + \gamma)e_4, \\ e_3 \cdot e_1 &= (\gamma-1)e_3, & e_3 \cdot e_4 &= e_2, & e_2 \cdot e_1 &= (\gamma-1)e_2, & e_4 \cdot e_1 &= (\gamma-1)e_4 \\ e_4 \cdot e_3 &= e_2. \end{aligned}$$

(11) The Lie superalgebra $(D_{1/2}^8)$: A Novikov structure is given by (where $\lambda \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -\frac{1}{2}e_1 + \lambda e_2, & e_1 \cdot e_2 &= \frac{1}{2}e_2, & e_1 \cdot e_4 &= e_3, & e_2 \cdot e_1 &= -\frac{1}{2}e_2, \\ e_3 \cdot e_1 &= -\frac{1}{2}e_3, & e_4 \cdot e_1 &= -\frac{1}{2}e_4, & e_4 \cdot e_4 &= \frac{1}{2}e_2. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= \gamma e_1 + \lambda e_2, & e_1 \cdot e_2 &= (1+\gamma)e_2, & e_1 \cdot e_3 &= \frac{1}{2}(1+\gamma)e_3, & e_1 \cdot e_4 &= e_3 + \frac{1}{2}(1+\gamma)e_4, \\ e_2 \cdot e_1 &= \gamma e_2, & e_3 \cdot e_1 &= \gamma e_3, & e_4 \cdot e_1 &= \gamma e_4, & e_4 \cdot e_4 &= e_2. \end{aligned}$$

(12) The Lie superalgebra $(D_{1/2,p}^9)$ with $p > 0$: A Novikov structure is given by (where $\lambda \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -\frac{1}{2}e_1 + \lambda e_2, & e_1 \cdot e_2 &= \frac{1}{2}e_2, & e_1 \cdot e_3 &= -pe_4, & e_1 \cdot e_4 &= pe_3, & e_2 \cdot e_1 &= -\frac{1}{2}e_2, \\ e_3 \cdot e_1 &= -\frac{1}{2}e_3, & e_3 \cdot e_3 &= \frac{1}{2}e_2, & e_4 \cdot e_1 &= -\frac{1}{2}e_4, & e_4 \cdot e_4 &= \frac{1}{2}e_2. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= (\gamma-1)e_1 + \lambda e_2, & e_1 \cdot e_2 &= \gamma e_2, & e_1 \cdot e_3 &= \frac{\gamma}{2}e_3 - pe_4, & e_1 \cdot e_4 &= pe_3 + \frac{\gamma}{2}e_4, \\ e_2 \cdot e_1 &= (\gamma-1)e_2, & e_3 \cdot e_1 &= (\gamma-1)e_3, & e_3 \cdot e_3 &= e_2, & e_4 \cdot e_1 &= (\gamma-1)e_4, \\ e_4 \cdot e_4 &= e_2. \end{aligned}$$

(13) The Lie superalgebra $(D_0^{10})^1$: An LSSA structure is given by (where $\gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -e_1, & e_1 \cdot e_4 &= -e_4, & e_2 \cdot e_1 &= -e_2, & e_2 \cdot e_4 &= (1+\gamma)e_3, & e_3 \cdot e_1 &= -e_3, \\ e_3 \cdot e_4 &= -\frac{1}{2}e_2, & e_4 \cdot e_1 &= -e_4, & e_4 \cdot e_2 &= \gamma e_3, & e_4 \cdot e_4 &= \frac{1}{2}e_1. \end{aligned}$$

Claim. There is no Novikov structure on $(D_0^{10})^1$ compatible with the Lie structure.

Proof. Let us write:

$$\begin{aligned} e_i \cdot e_j &= \sum_{i,j} \lambda_{ij}^1 e_1 + \sum_{i,j} \lambda_{ij}^2 e_2 & \text{if } |e_i| + |e_j| = \bar{0}, \text{ and} \\ e_i \cdot e_j &= \sum_{i,j} \lambda_{ij}^3 e_3 + \sum_{i,j} \lambda_{ij}^4 e_4 & \text{if } |e_i| + |e_j| = \bar{1}. \end{aligned}$$

The compatibility condition implies that

$$\begin{aligned} \lambda_{13}^3 &= \lambda_{31}^3 + 1, & \lambda_{13}^4 &= \lambda_{31}^4, & \lambda_{12}^2 &= \lambda_{21}^2 + 1, & \lambda_{21}^1 &= \lambda_{12}^1, & \lambda_{41}^3 &= \lambda_{14}^3, & \lambda_{41}^4 &= \lambda_{14}^4, \\ \lambda_{24}^3 &= \lambda_{42}^3 + 1, & \lambda_{42}^4 &= \lambda_{24}^4, & \lambda_{32}^3 &= \lambda_{23}^3, & \lambda_{32}^4 &= \lambda_{23}^4, & \lambda_{33}^1 &= 0, & \lambda_{33}^2 &= 0, & \lambda_{44}^1 &= \frac{1}{2}, \\ \lambda_{44}^2 &= 0 & \lambda_{34}^2 &= -\lambda_{43}^2 - \frac{1}{2}, & \lambda_{43}^1 &= -\lambda_{34}^1. \end{aligned}$$

Let us put

$$\begin{aligned} N(x, y, z) &: = (z \cdot x) \cdot y - (-1)^{|x||y|} (z \cdot y) \cdot x, \\ T(x, y, z) &: = (x, y, z) - (-1)^{|x||y|} (y, x, z). \end{aligned}$$

A direct calculation gives

$$N(e_4, e_4, e_4) = 0 \implies \lambda_{14}^3 = \lambda_{14}^4 = 0, \text{ also } N(e_4, e_1, e_4) = 0 \implies \lambda_{11}^1 = \lambda_{11}^2 = 0.$$

Similarly,

$$N(e_1, e_3, e_1) = 0 \implies \lambda_{31}^3 = \lambda_{13}^4 = 0, \text{ also } T(e_2, e_1, e_2) = 0 \implies \lambda_{21}^2 = \lambda_{12}^1 = \lambda_{22}^1 = \lambda_{22}^2 = 0.$$

Additionally,

$$N(e_4, e_2, e_1) = 0 \implies \lambda_{24}^4 = \lambda_{42}^3 = -1 \text{ also } N(e_4, e_2, e_4) = 0 \implies \lambda_{34}^1 = \lambda_{43}^2 = 0.$$

Now, $N(e_3, e_4, e_4) = \frac{1}{2}e_3$ which is a not zero. Therefore, it cannot be Novikov. \square

However, there is a Balinsky-Novikov structure given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= \gamma e_1 + \lambda e_2, & e_1 \cdot e_2 &= (1 + \gamma)e_2, & e_1 \cdot e_3 &= \frac{1}{2}(2 + \gamma)e_3, & e_1 \cdot e_4 &= -\lambda e_3 + \frac{\gamma}{2}e_4, \\ e_2 \cdot e_1 &= \gamma e_2, & e_2 \cdot e_4 &= -\gamma e_3, & e_3 \cdot e_1 &= \gamma e_3, & e_3 \cdot e_4 &= -\frac{1}{2}e_2, \\ e_4 \cdot e_1 &= -2\lambda e_3 + \gamma e_4, & e_4 \cdot e_2 &= -2(1 + \gamma)e_3 & e_4 \cdot e_3 &= -\frac{1}{2}e_2, & e_4 \cdot e_4 &= e_1. \end{aligned}$$

(14) The Lie superalgebra $(D_0^{10})^2$: AAn LSSA structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -e_1 - 2\lambda\gamma e_2, & e_1 \cdot e_4 &= \gamma e_3 - e_4, & e_2 \cdot e_1 &= -e_2, & e_2 \cdot e_4 &= e_3, & e_3 \cdot e_1 &= -e_3, \\ e_3 \cdot e_4 &= \frac{1}{2}(1 - 2\lambda)e_2, & e_4 \cdot e_1 &= \gamma e_3 - e_4, & e_4 \cdot e_3 &= \lambda e_2, & e_4 \cdot e_4 &= -\frac{1}{2}e_1. \end{aligned}$$

Claim. There is no Novikov structure on $(D_0^{10})^2$ compatible with the Lie structure.

However, there is a Balinsky-Novikov structure given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= (\gamma - 1)e_1 + \lambda e_2, & e_1 \cdot e_2 &= \gamma e_2, & e_1 \cdot e_3 &= \frac{1}{2}(1 + \gamma)e_3, & e_1 \cdot e_4 &= -\lambda e_3 + \frac{1}{2}(\gamma - 1)e_4, \\ e_2 \cdot e_1 &= (\gamma - 1)e_2, & e_2 \cdot e_4 &= (1 - \gamma)e_3, & e_3 \cdot e_1 &= (\gamma - 1)e_3, & e_3 \cdot e_4 &= \frac{1}{2}e_2, \\ e_4 \cdot e_1 &= -2\lambda e_3 + (\gamma - 1)e_4, & e_4 \cdot e_2 &= -2\gamma e_3 & e_4 \cdot e_3 &= \frac{1}{2}e_2, & e_4 \cdot e_4 &= -e_1. \end{aligned}$$

(15) The Lie superalgebra $(2A_{1,1} + 2A)^1$: A Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$e_3 \cdot e_3 = \frac{1}{2}e_1, \quad e_3 \cdot e_4 = \lambda e_1 + \gamma e_2, \quad e_4 \cdot e_3 = -\lambda e_1 - \gamma e_2, \quad e_4 \cdot e_4 = \frac{1}{2}e_2.$$

A Balinsky-Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= \gamma e_1, & e_1 \cdot e_3 &= \frac{\gamma}{2}e_3, & e_2 \cdot e_2 &= 2\lambda e_2, & e_2 \cdot e_4 &= \lambda e_4, \\ e_3 \cdot e_1 &= \gamma e_3, & e_3 \cdot e_3 &= e_1, & e_4 \cdot e_2 &= 2\lambda e_4, & e_4 \cdot e_4 &= e_2. \end{aligned}$$

(16) The Lie superalgebra $(2A_{1,1} + 2A)^2$: A Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$e_3 \cdot e_3 = \frac{1}{2}e_1, \quad e_3 \cdot e_4 = \gamma e_1, \quad e_4 \cdot e_3 = (1 - \gamma)e_1, \quad e_4 \cdot e_4 = \frac{1}{2}e_2.$$

A Balinsky-Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= 2\gamma e_1, & e_1 \cdot e_2 &= 2\gamma e_1, & e_1 \cdot e_3 &= \gamma e_3, & e_1 \cdot e_4 &= \gamma e_3, \\ e_2 \cdot e_1 &= 2\gamma e_1, & e_2 \cdot e_2 &= -2(\lambda - \gamma)e_1 + 2\lambda e_2, & e_2 \cdot e_3 &= \gamma e_3, & e_2 \cdot e_4 &= (\gamma - \lambda)e_3 + \lambda e_4, \\ e_3 \cdot e_1 &= 2\gamma e_3, & e_3 \cdot e_2 &= 2\gamma e_3, & e_3 \cdot e_3 &= e_1, & e_3 \cdot e_4 &= e_1, \\ e_4 \cdot e_1 &= 2\gamma e_3, & e_4 \cdot e_2 &= -2(\lambda - \gamma)e_3 + 2\lambda e_4, & e_4 \cdot e_3 &= e_1, & e_4 \cdot e_4 &= e_2. \end{aligned}$$

(17) The Lie superalgebra $(2A_{1,1} + 2A)_p^3$: A Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$e_3 \cdot e_3 = \frac{1}{2}e_1, \quad e_3 \cdot e_4 = \gamma e_1, \quad e_4 \cdot e_3 = (p - \gamma)e_1 + p e_2, \quad e_4 \cdot e_4 = \frac{1}{2}e_2.$$

A Balinsky-Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= \frac{2}{p}(p^2\gamma + \lambda(2p^2 - 1))e_1 + 2p\lambda e_2, & e_1 \cdot e_2 &= -2p\lambda e_1 + 2p\gamma e_2, \\ e_1 \cdot e_3 &= \frac{1}{p}(p^2\gamma + \lambda(p^2 - 1))e_3 + \lambda e_4, & e_1 \cdot e_4 &= -\lambda e_3 + p(\gamma + \lambda)e_4, \\ e_2 \cdot e_1 &= -2p\lambda e_1 + 2p\gamma e_2, & e_2 \cdot e_2 &= -2p\gamma e_1 - \frac{2}{p}(p^2\lambda + (2p^2 - 1)\gamma)e_2, \\ e_2 \cdot e_3 &= -p(\lambda + \gamma)e_3 + \gamma e_4, & e_2 \cdot e_4 &= -\gamma e_3 + \frac{1}{p}((1 - p^2)\gamma - p^2\lambda)e_4, \\ e_3 \cdot e_1 &= \frac{2}{p}(p^2\gamma + (p^2 - 1)\lambda)e_3 + 2\lambda e_4, & e_3 \cdot e_2 &= -2p(\gamma + \lambda)e_3 + 2\gamma e_4, \\ e_3 \cdot e_3 &= e_1, & e_3 \cdot e_4 &= p e_1 + p e_2, \\ e_4 \cdot e_1 &= -2\lambda e_3 + 2p(\gamma + \lambda)e_4, & e_4 \cdot e_2 &= -2\gamma e_3 - \frac{2}{p}((p^2 - 1)\gamma + p^2\lambda)e_4, \\ e_4 \cdot e_3 &= p e_1 + p e_2, & e_4 \cdot e_4 &= e_2. \end{aligned}$$

(18) The Lie superalgebra $(2A_{1,1} + 2A)_p^4$: A Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$e_3 \cdot e_3 = \frac{1}{2}e_1, \quad e_3 \cdot e_4 = \gamma e_1, \quad e_4 \cdot e_3 = (p - \gamma)e_1 - p e_2, \quad e_4 \cdot e_4 = \frac{1}{2}e_2.$$

A Balinsky-Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned}
e_1 \cdot e_1 &= -\frac{2}{p}(p^2\gamma - \lambda(2p^2 + 1))e_1 - 2p\lambda e_2, & e_1 \cdot e_2 &= 2p\lambda e_1 - 2p\gamma e_2, \\
e_1 \cdot e_3 &= \frac{1}{p}(-p^2\gamma + \lambda(p^2 + 1))e_3 + \lambda e_4, & e_1 \cdot e_4 &= \lambda e_3 + p(-\gamma + \lambda)e_4, \\
e_2 \cdot e_1 &= 2p\lambda e_1 - 2p\gamma e_2, & e_2 \cdot e_2 &= 2p\gamma e_1 - \frac{2}{p}(-p^2\lambda + (2p^2 + 1)\gamma)e_2, \\
e_2 \cdot e_3 &= p(-\gamma + \lambda)e_3 + \gamma e_4, & e_2 \cdot e_4 &= \gamma e_3 + \frac{1}{p}((-1 - p^2)\gamma + p^2\lambda)e_4, \\
e_3 \cdot e_1 &= \frac{2}{p}(-p^2\gamma + (p^2 + 1)\lambda)e_3 + 2\lambda e_4, & e_3 \cdot e_2 &= -2p(\gamma - \lambda)e_3 + 2\gamma e_4, \\
e_3 \cdot e_3 &= e_1, & e_3 \cdot e_4 &= pe_1 - pe_2, \\
e_4 \cdot e_1 &= 2\lambda e_3 - 2p(\gamma - \lambda)e_4, & e_4 \cdot e_2 &= 2\gamma e_3 - \frac{2}{p}((p^2 + 1)\gamma - p^2\lambda)e_4, \\
e_4 \cdot e_3 &= pe_1 - pe_2, & e_4 \cdot e_4 &= e_2.
\end{aligned}$$

(19) The Lie superalgebra $(C_1^1 + A)$: A Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$\begin{aligned}
e_1 \cdot e_1 &= \lambda e_1 + \gamma e_2, & e_1 \cdot e_2 &= (1 + \lambda)e_2, & e_1 \cdot e_3 &= (1 + \lambda)e_3, & e_1 \cdot e_4 &= \lambda e_4, \\
e_2 \cdot e_1 &= \lambda e_2, & e_3 \cdot e_1 &= \lambda e_3, & e_3 \cdot e_4 &= -\lambda e_2, & e_4 \cdot e_1 &= \lambda e_4, \\
e_4 \cdot e_3 &= (1 + \lambda)e_2.
\end{aligned}$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$\begin{aligned}
e_1 \cdot e_1 &= (\gamma - 1)e_1 + \lambda e_2, & e_1 \cdot e_2 &= \gamma e_2, & e_1 \cdot e_3 &= \frac{1}{2}(1 + \gamma)e_3, & e_1 \cdot e_4 &= \frac{1}{2}(\gamma - 1)e_4, \\
e_2 \cdot e_1 &= (\gamma - 1)e_2, & e_3 \cdot e_1 &= (\gamma - 1)e_3, & e_3 \cdot e_4 &= e_2, & e_4 \cdot e_1 &= (\gamma - 1)e_4, \\
e_4 \cdot e_3 &= e_2.
\end{aligned}$$

(20) The Lie superalgebra $(C_{1/2}^1 + A)$: A Novikov structure is given by:

$$e_1 \cdot e_1 = -\frac{1}{2}e_1 + \gamma e_2, \quad e_1 \cdot e_2 = \frac{1}{2}e_2, \quad e_2 \cdot e_1 = -\frac{1}{2}e_2, \quad e_3 \cdot e_1 = -\frac{1}{2}e_3, \quad e_3 \cdot e_3 = \frac{1}{2}e_2.$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$\begin{aligned}
e_1 \cdot e_1 &= \gamma e_1 + \lambda e_2, & e_1 \cdot e_2 &= (1 + \gamma)e_2, & e_1 \cdot e_3 &= \frac{1}{2}(1 + \gamma)e_3, & e_1 \cdot e_4 &= \frac{\gamma}{2}e_4, \\
e_2 \cdot e_1 &= \gamma e_2, & e_3 \cdot e_1 &= \gamma e_3, & e_3 \cdot e_3 &= e_2, & e_4 \cdot e_1 &= \gamma e_4.
\end{aligned}$$

(21) The Lie superalgebra $(C_{-1}^2 + A)$: A Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned}
e_1 \cdot e_1 &= (2\lambda - 1)e_1 + \gamma e_2, & e_1 \cdot e_2 &= (2\lambda - 1)e_2, & e_1 \cdot e_3 &= 2\lambda e_3, & e_1 \cdot e_4 &= 2(\lambda - 1)e_4, \\
e_2 \cdot e_1 &= (2\lambda - 1)e_2, & e_3 \cdot e_1 &= (2\lambda - 1)e_3, & e_3 \cdot e_4 &= (1 - \lambda)e_2, & e_4 \cdot e_1 &= (2\lambda - 1)e_4, \\
e_4 \cdot e_3 &= \lambda e_2.
\end{aligned}$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= \gamma e_1 + \lambda e_2, & e_1 \cdot e_2 &= \gamma e_2, & e_1 \cdot e_3 &= \frac{1}{2}(2 + \gamma)e_3, & e_1 \cdot e_4 &= \frac{1}{2}(\gamma - 2)e_4, \\ e_2 \cdot e_1 &= \gamma e_2, & e_3 \cdot e_1 &= \gamma e_3, & e_3 \cdot e_4 &= e_2, & e_4 \cdot e_1 &= \gamma e_4, \\ e_4 \cdot e_3 &= e_2. \end{aligned}$$

(22) The Lie superalgebra $(C^3 + A)$: A Novikov structure is given by:

$$e_4 \cdot e_1 = -e_3, \quad e_4 \cdot e_4 = \frac{1}{2}e_2.$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= \gamma e_1 + \lambda e_2, & e_1 \cdot e_2 &= \gamma e_2, & e_1 \cdot e_3 &= \frac{\gamma}{2}e_3, & e_1 \cdot e_4 &= e_3 + \frac{\gamma}{2}e_4, \\ e_2 \cdot e_1 &= \gamma e_2, & e_3 \cdot e_1 &= \gamma e_3, & e_4 \cdot e_1 &= \gamma e_4, & e_4 \cdot e_4 &= e_2. \end{aligned}$$

(23) The Lie superalgebra $(C_0^5 + A)$: A Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= 2\lambda e_1 + \gamma e_2, & e_1 \cdot e_2 &= 2\lambda e_2, & e_1 \cdot e_3 &= 2\lambda e_3 - e_4, & e_1 \cdot e_4 &= e_3 + 2\lambda e_4, \\ e_2 \cdot e_1 &= 2\lambda e_2, & e_3 \cdot e_1 &= 2\lambda e_3, & e_3 \cdot e_3 &= \frac{1}{2}e_2, & e_3 \cdot e_4 &= -\lambda e_2, & e_4 \cdot e_1 &= 2\lambda e_4, \\ e_4 \cdot e_3 &= \lambda e_2, & e_4 \cdot e_4 &= \frac{1}{2}e_2. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= \gamma e_1 + \lambda e_2, & e_1 \cdot e_2 &= \gamma e_2, & e_1 \cdot e_3 &= \frac{\gamma}{2}e_3 - e_4, & e_1 \cdot e_4 &= e_3 + \frac{\gamma}{2}e_4, \\ e_2 \cdot e_1 &= \gamma e_2, & e_3 \cdot e_1 &= \gamma e_3, & e_3 \cdot e_3 &= e_2, & e_4 \cdot e_1 &= \gamma e_4, \\ e_4 \cdot e_4 &= e_2. \end{aligned}$$

(24) The Lie superalgebra D^1 : A Novikov structure is given by (where $\lambda, \mu, \gamma \in \mathbb{R}$):

$$e_2 \cdot e_2 = \lambda e_1 + \gamma e_3, \quad e_2 \cdot e_3 = (1 + \mu)e_1, \quad e_2 \cdot e_4 = e_4, \quad e_3 \cdot e_2 = \mu e_1.$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$e_2 \cdot e_2 = \gamma e_1 + \lambda e_3, \quad e_2 \cdot e_4 = e_4, \quad e_3 \cdot e_2 = -e_1.$$

(25) The Lie superalgebra D_q^2 with $q \neq 0$: A Novikov structure is given by:

$$\begin{aligned} e_1 \cdot e_3 &= -qe_1, & e_2 \cdot e_3 &= -qe_2, & e_3 \cdot e_1 &= -(1 + q)e_1, & e_3 \cdot e_2 &= -e_1 - (q + 1)e_2, \\ e_3 \cdot e_3 &= -qe_3, & e_4 \cdot e_3 &= -qe_4. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$e_3 \cdot e_1 = -e_1, \quad e_3 \cdot e_2 = -e_1 - e_2, \quad e_3 \cdot e_3 = \lambda e_1 + \gamma e_2, \quad e_3 \cdot e_4 = qe_4.$$

(26) The Lie superalgebra D_{pq}^3 with $pq \neq 0$: A Novikov structure is given by (where $\lambda, \mu, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_3 &= (p + \gamma)e_1, & e_2 \cdot e_3 &= (p + \gamma)e_2, & e_3 \cdot e_1 &= \gamma e_1 + e_2, & e_3 \cdot e_2 &= -e_1 + \gamma e_2, \\ e_3 \cdot e_3 &= \lambda e_1 + \mu e_2 + (p + \gamma)e_3, & e_3 \cdot e_4 &= (p + q + \gamma)e_4, & e_4 \cdot e_3 &= (p + \gamma)e_4. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$e_3 \cdot e_1 = -pe_1 + e_2, \quad e_3 \cdot e_2 = -e_1 - pe_2, \quad e_3 \cdot e_3 = \lambda e_1 + \gamma e_2, \quad e_3 \cdot e_4 = qe_4.$$

(27) The Lie superalgebra D_{pq}^{11} with $0 < |p| \leq |q| \leq 1$: A Novikov structure is given by:

$$e_1 \cdot e_2 = e_2, \quad e_1 \cdot e_3 = pe_3, \quad e_1 \cdot e_4 = qe_4.$$

A Balinsky-Novikov structure is given by:

$$e_1 \cdot e_2 = e_2, \quad e_1 \cdot e_3 = pe_3, \quad e_1 \cdot e_4 = qe_4.$$

(28) The Lie superalgebra D^{12} : A Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$e_1 \cdot e_2 = e_2, \quad e_1 \cdot e_4 = (1 + \gamma)e_3, \quad e_4 \cdot e_1 = \gamma e_3.$$

A Balinsky-Novikov structure is given by:

$$\begin{aligned} e_1 \cdot e_1 &= -2e_1, & e_1 \cdot e_3 &= -e_3, & e_1 \cdot e_4 &= e_3 - e_4, & e_2 \cdot e_1 &= -2e_2, \\ e_3 \cdot e_1 &= -2e_3, & e_4 \cdot e_1 &= -2e_4. \end{aligned}$$

(29) The Lie superalgebra D_p^{13} with $p \neq 0$: A Novikov structure is given by:

$$e_1 \cdot e_2 = pe_2, \quad e_1 \cdot e_3 = e_3, \quad e_1 \cdot e_4 = e_3 + e_4.$$

A Balinsky-Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= 2(\gamma - 1)e_1, & e_1 \cdot e_2 &= (p - 1 + \gamma)e_2, & e_1 \cdot e_3 &= \gamma e_3, & e_1 \cdot e_4 &= e_3 + \gamma e_4, \\ e_2 \cdot e_1 &= 2(\gamma - 1)e_2, & e_3 \cdot e_1 &= 2(\gamma - 1)e_3, & e_4 \cdot e_1 &= 2(\gamma - 1)e_4. \end{aligned}$$

(30) The Lie superalgebra D_{pq}^{14} with $p \neq 0, q \geq 0$: A Novikov structure is given by:

$$e_1 \cdot e_2 = pe_2, \quad e_1 \cdot e_3 = qe_3 - e_4, \quad e_1 \cdot e_4 = e_3 + qe_4.$$

A Balinsky-Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -2qe_1, & e_1 \cdot e_2 &= (p - q)e_2, & e_1 \cdot e_3 &= -e_4, & e_1 \cdot e_4 &= e_3, \\ e_2 \cdot e_1 &= -2qe_2, & e_3 \cdot e_1 &= -2qe_3, & e_4 \cdot e_1 &= -2qe_4. \end{aligned}$$

(31) The Lie superalgebra D^{15} : A Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$e_1 \cdot e_3 = e_2, \quad e_1 \cdot e_4 = \gamma e_2 + e_3, \quad e_4 \cdot e_1 = \gamma e_2.$$

A Balinsky-Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= 2\gamma e_1, & e_1 \cdot e_2 &= \gamma e_2, & e_1 \cdot e_3 &= e_2 + \gamma e_3, & e_1 \cdot e_4 &= e_3 + \gamma e_4, \\ e_2 \cdot e_1 &= 2\gamma e_2, & e_3 \cdot e_1 &= 2\gamma e_3, & e_4 \cdot e_1 &= 2\gamma e_4. \end{aligned}$$

(32) The Lie superalgebra D^{16} : A Novikov structure is given by:

$$e_1 \cdot e_2 = e_2, \quad e_1 \cdot e_3 = e_2 + e_3, \quad e_1 \cdot e_4 = e_3 + e_4.$$

A Balinsky-Novikov structure is given by (where $\gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= 2(\gamma - 1)e_1, & e_1 \cdot e_2 &= \gamma e_2, & e_1 \cdot e_3 &= e_2 + \gamma e_3, & e_1 \cdot e_4 &= e_3 + \gamma e_4, \\ e_2 \cdot e_1 &= 2(\gamma - 1)e_2, & e_3 \cdot e_1 &= 2(\gamma - 1)e_3, & e_4 \cdot e_1 &= 2(\gamma - 1)e_4. \end{aligned}$$

(33) The Lie superalgebra $(A_{3,1} + A)$: A Novikov structure is given by (where $\lambda, \gamma, \mu \in \mathbb{R}$):

$$e_2 \cdot e_2 = \gamma e_1, \quad e_2 \cdot e_3 = (1 + \lambda)e_1, \quad e_3 \cdot e_2 = \lambda e_1, \quad e_3 \cdot e_3 = \mu e_1, \quad e_4 \cdot e_4 = \frac{1}{2}e_1.$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda, \mu \in \mathbb{R}$):

$$e_2 \cdot e_2 = \gamma e_1, \quad e_2 \cdot e_3 = \lambda e_1, \quad e_3 \cdot e_2 = (\lambda - 1)e_1, \quad e_3 \cdot e_3 = \mu e_1, \quad e_4 \cdot e_4 = e_1.$$

(34) The Lie superalgebra $(D_{p,-1/2}^3)$ with $p \neq 0$: A Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -\frac{1}{2}e_1 + \lambda e_2 + \gamma e_3, & e_1 \cdot e_2 &= \frac{1}{2}e_2, & e_1 \cdot e_3 &= \frac{1}{2}(2p - 1)e_3, & e_2 \cdot e_1 &= -\frac{1}{2}e_2, \\ e_3 \cdot e_1 &= -\frac{1}{2}e_3, & e_4 \cdot e_1 &= -\frac{1}{2}e_4, & e_4 \cdot e_4 &= \frac{1}{2}e_2. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$e_1 \cdot e_1 = \gamma e_2 + \lambda e_3, \quad e_1 \cdot e_2 = e_2, \quad e_1 \cdot e_3 = p e_3, \quad e_1 \cdot e_4 = \frac{1}{2}e_4, \quad e_4 \cdot e_4 = e_2.$$

(35) The Lie superalgebra $(D_{-1/2}^2)^1$: A Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -\frac{1}{2}e_1 + \lambda e_2 + \gamma e_3, & e_1 \cdot e_2 &= \frac{1}{2}e_2, & e_1 \cdot e_3 &= e_2 + \frac{1}{2}e_3, & e_2 \cdot e_1 &= -\frac{1}{2}e_2, \\ e_3 \cdot e_1 &= -\frac{1}{2}e_3, & e_4 \cdot e_1 &= -\frac{1}{2}e_4, & e_4 \cdot e_4 &= \frac{1}{2}e_2. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$e_1 \cdot e_1 = \gamma e_2 + \lambda e_3, \quad e_1 \cdot e_2 = e_2, \quad e_1 \cdot e_3 = e_2 + e_3, \quad e_1 \cdot e_4 = \frac{1}{2}e_4, \quad e_4 \cdot e_4 = e_2.$$

(36) The Lie superalgebra $(D_{-1/2}^2)^2$: A Novikov structure is given by (where $\lambda, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_1 \cdot e_1 &= -\frac{1}{2}e_1 + \lambda e_2 + \gamma e_3, & e_1 \cdot e_2 &= \frac{1}{2}e_2, & e_1 \cdot e_3 &= -e_2 + \frac{1}{2}e_3, & e_2 \cdot e_1 &= -\frac{1}{2}e_2, \\ e_3 \cdot e_1 &= -\frac{1}{2}e_3, & e_4 \cdot e_1 &= -\frac{1}{2}e_4, & e_4 \cdot e_4 &= \frac{1}{2}e_2. \end{aligned}$$

A Balinsky-Novikov structure is given by (where $\gamma, \lambda \in \mathbb{R}$):

$$e_1 \cdot e_1 = \gamma e_2 + \lambda e_3, \quad e_1 \cdot e_2 = e_2, \quad e_1 \cdot e_3 = -e_2 + e_3, \quad e_1 \cdot e_4 = \frac{1}{2}e_4, \quad e_4 \cdot e_4 = e_2.$$

(37) The Lie superalgebra $(A_{1,1} + 3A)^1$: A Novikov structure is given by (where $\lambda, \mu, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_2 \cdot e_2 &= \frac{1}{2}e_1, & e_2 \cdot e_3 &= \lambda e_1, & e_2 \cdot e_4 &= \gamma e_1, & e_3 \cdot e_2 &= -\lambda e_1, & e_3 \cdot e_3 &= \frac{1}{2}e_1, \\ e_3 \cdot e_4 &= \mu e_1, & e_4 \cdot e_2 &= -\gamma e_1, & e_4 \cdot e_3 &= -\mu e_1, & e_4 \cdot e_4 &= \frac{1}{2}e_1. \end{aligned}$$

A Balinsky-Novikov structure is given by:

$$e_2 \cdot e_2 = e_1, \quad e_3 \cdot e_3 = e_1, \quad e_4 \cdot e_4 = e_1.$$

(38) The Lie superalgebra $(A_{1,1} + 3A)^2$: A Novikov structure is given by (where $\lambda, \mu, \gamma \in \mathbb{R}$):

$$\begin{aligned} e_2 \cdot e_2 &= \frac{1}{2}e_1, & e_2 \cdot e_3 &= \lambda e_1, & e_2 \cdot e_4 &= \gamma e_1, & e_3 \cdot e_2 &= -\lambda e_1, & e_3 \cdot e_3 &= \frac{1}{2}e_1, \\ e_3 \cdot e_4 &= \mu e_1, & e_4 \cdot e_2 &= -\gamma e_1, & e_4 \cdot e_3 &= -\mu e_1, & e_4 \cdot e_4 &= -\frac{1}{2}e_1. \end{aligned}$$

A Balinsky-Novikov structure is given by:

$$e_2 \cdot e_2 = e_1, \quad e_3 \cdot e_3 = e_1, \quad e_4 \cdot e_4 = -e_1.$$

6. THE LIST OF FOUR-DIMENSIONAL REAL LIE SUPERALGEBRAS

The LSA	Relations in the basis: $e_1, e_2 e_3, e_4$	Symplectic structure ω	$p(\omega)$
D^5	$[e_1, e_3] = e_3,$ $[e_1, e_4] = e_4,$ $[e_2, e_4] = e_3$	$e_3^* \wedge e_1^* + e_4^* \wedge e_2^*$	$\bar{1}$
D^6	$[e_1, e_3] = e_3,$ $[e_1, e_4] = e_4,$ $[e_2, e_3] = -e_4,$ $[e_2, e_4] = e_3$	$e_3^* \wedge e_1^* + e_4^* \wedge e_2^*$	$\bar{1}$
$D_{pq}^7,$ $pq \neq 0,$ $p \geq q$	$[e_1, e_2] = e_2,$ $[e_1, e_3] = pe_3,$ $[e_1, e_4] = qe_4$	None	-
$D_{-1q}^7,$ $q \leq -1$	$[e_1, e_2] = e_2,$ $[e_1, e_3] = -e_3,$ $[e_1, e_4] = qe_4$	$e_4^* \wedge e_1^* + e_3^* \wedge e_2^*$	$\bar{1}$
$D_{pp}^7,$ $p = -q$ $q \neq 0, -1$	$[e_1, e_2] = e_2,$ $[e_1, e_3] = pe_3$ $[e_1, e_4] = -pe_4$	$e_2^* \wedge e_1^* - e_3^* \wedge e_4^*$	$\bar{0}$
$D_{pq}^7,$ $p = -q,$ $q = -1$	$[e_1, e_2] = e_2,$ $[e_1, e_3] = e_3,$ $[e_1, e_4] = -e_4$	$e_2^* \wedge e_1^* - e_3^* \wedge e_4^*$ $e_3^* \wedge e_1^* + e_4^* \wedge e_2^*$	$\bar{0}$ $\bar{1}$
$D_p^8,$ $p \neq 0$	$[e_1, e_2] = e_2,$ $[e_1, e_3] = pe_3,$ $[e_1, e_4] = e_3 + pe_4$	None	-
D_{-1}^8	$[e_1, e_2] = e_2,$ $[e_1, e_3] = -e_3,$ $[e_1, e_4] = e_3 - e_4$	$e_3^* \wedge e_1^* + e_4^* \wedge e_2^*$	$\bar{1}$
$D_{pq}^9,$ $q > 0$	$[e_1, e_3] = pe_3 - qe_4,$ $[e_1, e_2] = e_2,$ $[e_1, e_4] = qe_3 + pe_4$	None	-
D_q^{10}	$[e_1, e_2] = e_2,$ $[e_1, e_3] = (q+1)e_3,$ $[e_1, e_4] = qe_4,$ $[e_2, e_4] = e_3$	$(1+q)e_1^* \wedge e_3^* + e_2^* \wedge e_4^*,$ where $q \neq -1$	$\bar{1}$

TABLE 1. Trivial algebras (i.e. $[\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] = \{0\}$) with $\text{sdim} = 2|2$

The LSA	Relations in the basis: $e_1, e_2 \mid e_3, e_4$	Symplectic structure	$p(\omega)$
$(D_{1/2, 1/2}^7)^1$	$[e_1, e_2] = e_2, [e_1, e_3] = \frac{1}{2}e_3$ $[e_1, e_4] = \frac{1}{2}e_4, [e_3, e_3] = e_2$ $[e_4, e_4] = e_2$	$e_1^* \wedge e_2^* - \frac{1}{2}e_3^* \wedge e_3^* - \frac{1}{2}e_4^* \wedge e_4^*$	$\bar{0}$
$(D_{1/2, 1/2}^7)^2$	$[e_1, e_2] = e_2, [e_1, e_3] = \frac{1}{2}e_3$ $[e_1, e_4] = \frac{1}{2}e_4, [e_3, e_3] = e_2$ $[e_4, e_4] = -e_2$	$-e_1^* \wedge e_2^* + \frac{1}{2}e_3^* \wedge e_3^* - \frac{1}{2}e_4^* \wedge e_4^*$	$\bar{0}$
$(D_{1/2, 1/2}^7)^3$	$[e_1, e_2] = e_2, [e_1, e_3] = \frac{1}{2}e_3$ $[e_1, e_4] = \frac{1}{2}e_4, [e_3, e_3] = e_2$	None	—
$(D_{1-p, p}^7)$ $p \leq \frac{1}{2}$	$[e_1, e_2] = e_2, [e_1, e_3] = pe_3$ $[e_1, e_4] = (1-p)e_4,$ $[e_3, e_4] = e_2$	$e_1^* \wedge e_2^* - e_3^* \wedge e_4^*$	$\bar{0}$
$(D_{1/2}^8)$	$[e_1, e_2] = e_2, [e_1, e_3] = \frac{1}{2}e_3$ $[e_1, e_4] = e_3 + \frac{1}{2}e_4,$ $[e_4, e_4] = e_2$	None	—
$(D_{1/2, p}^9)$ $p > 0$	$[e_1, e_2] = e_2, [e_3, e_3] = e_2$ $[e_1, e_4] = pe_3 + \frac{1}{2}e_4, [e_4, e_4] = e_2$ $[e_1, e_3] = \frac{1}{2}e_3 - pe_4$	$e_1^* \wedge e_2^* - \frac{1}{2}e_3^* \wedge e_3^* - \frac{1}{2}e_4^* \wedge e_4^*$	$\bar{0}$
$(D_0^{10})^1$	$[e_1, e_2] = e_2, [e_1, e_3] = e_3$ $[e_2, e_4] = e_3, [e_4, e_4] = e_1$ $[e_3, e_4] = -\frac{1}{2}e_2$	$2e_2^* \wedge e_1^* - e_3^* \wedge e_4^*$ $e_1^* \wedge e_3^* + e_2^* \wedge e_4^*$	$\bar{0}$ $\bar{1}$
$(D_0^{10})^2$	$[e_1, e_2] = e_2, [e_1, e_3] = e_3$ $[e_2, e_4] = e_3, [e_4, e_4] = -e_1$ $[e_3, e_4] = \frac{1}{2}e_2$	$2e_2^* \wedge e_1^* - e_3^* \wedge e_4^*$ $e_1^* \wedge e_3^* + e_2^* \wedge e_4^*$	$\bar{0}$ $\bar{1}$
$(2A_{1,1} + 2A)^1$	$[e_3, e_3] = e_1, [e_4, e_4] = e_2$	None	—
$(2A_{1,1} + 2A)^2$	$[e_3, e_3] = e_1, [e_4, e_4] = e_2$ $[e_3, e_4] = e_1$	None	—
$(2A_{1,1} + 2A)_p^3$ $p > 0$	$[e_3, e_3] = e_1, [e_4, e_4] = e_2$ $[e_3, e_4] = p(e_1 + e_2)$	For $p \neq \frac{1}{2}$: None For $p = \frac{1}{2}$: $e_2^* \wedge e_3^* - e_1^* \wedge e_4^*$	— $\bar{1}$
$(2A_{1,1} + 2A)_p^4$ $p > 0$	$[e_3, e_3] = e_1, [e_4, e_4] = e_2$ $[e_3, e_4] = p(e_1 - e_2)$	None	—
$(C_1^1 + A)$	$[e_1, e_2] = e_2, [e_1, e_3] = e_3$ $[e_3, e_4] = e_2$	$\frac{1}{2}e_2^* \wedge e_1^* + \frac{1}{2}e_3^* \wedge e_4^* + \frac{1}{4}e_4^* \wedge e_4^*$	$\bar{0}$
$(C_{1/2}^1 + A)$	$[e_1, e_2] = e_2, [e_1, e_3] = \frac{1}{2}e_3$ $[e_3, e_3] = e_2$	$\lambda e_1^* \wedge e_2^* - \frac{\lambda}{2}e_3^* \wedge e_3^* - \frac{\mu}{2}e_4^* \wedge e_4^*,$ where $\lambda\mu \neq 0$	$\bar{0}$
$(C_{-1}^2 + A)$	$[e_1, e_3] = e_3, [e_1, e_4] = -e_4$ $[e_3, e_4] = e_2$	None	—
$(C^3 + A)$	$[e_1, e_4] = e_3, [e_4, e_4] = e_2$	$2e_1^* \wedge e_2^* - e_3^* \wedge e_4^*$	0
$(C_0^5 + A)$	$[e_1, e_3] = -e_4, [e_1, e_4] = e_3$ $[e_3, e_3] = e_2, [e_4, e_4] = e_2$	None	—

TABLE 2. Non-trivial algebras (i.e. $[\mathfrak{g}_{\bar{1}}, \mathfrak{g}_{\bar{1}}] \neq \{0\}$) with $\text{sdim} = 2 \mid 2$

The LSA	Relations in the basis: $e_1, e_2, e_3 e_4$	Symplectic structure	$p(\omega)$
D^1	$[e_2, e_3] = e_1, [e_2, e_4] = e_4$	$\lambda e_1^* \wedge e_3^* + \mu e_2 \wedge e_4^*$ $+ \nu e_1^* \wedge e_2^* + \gamma e_2 \wedge e_3^*$ where $\lambda\mu \neq 0$	NH
D_q^2 $q \neq -1, 0$	$[e_1, e_3] = e_1, [e_2, e_3] = e_1 + e_2,$ $[e_3, e_4] = qe_4$	None	-
D_{-1}^2	$[e_1, e_2] = e_1, [e_2, e_3] = e_1 + e_2,$ $[e_3, e_4] = -e_4$	$\lambda e_1^* \wedge e_3^* + \mu e_2^* \wedge e_3^* +$ $\nu e_2^* \wedge e_4^* + \gamma e_3^* \wedge e_4^*$ where $\lambda\nu \neq 0$	NH
D_{pq}^3 $pq \neq 0$	$[e_1, e_3] = pe_1 - e_2, [e_2, e_3] = e_1 + pe_2,$ $[e_3, e_4] = qe_4$	None	-

TABLE 3. Trivial algebras (i.e. $[\mathfrak{g}_1, \mathfrak{g}_1] = \{0\}$) with $\text{sdim} = 3|1$

The LSA	Relations in the basis: $e_1 e_2, e_3, e_4$	Symplectic structure	$p(\omega)$
D_{pq}^{11} $0 < p \leq q \leq 1$	$[e_1, e_2] = e_2, [e_1, e_3] = pe_3,$ $[e_1, e_4] = qe_4$	None	-
D_{pq}^{11} $(p, q) = (-1, -1)$	$[e_1, e_2] = e_2, [e_1, e_3] = pe_3,$ $[e_1, e_4] = qe_4$	$\lambda e_1^* \wedge e_2^* + \mu e_1^* \wedge e_3^* + \nu e_1^* \wedge e_4^*$ $+ \gamma e_2^* \wedge e_3^* + \delta e_2^* \wedge e_4^*,$ where $\nu\gamma - \mu\delta \neq 0$	NH
D_{pq}^{11} $0 < p \leq 1$ $p = -q$	$[e_1, e_2] = e_2, [e_1, e_3] = pe_3,$ $[e_1, e_4] = qe_4$	$\lambda e_1^* \wedge e_2^* + \mu e_1^* \wedge e_3^* + \nu e_1^* \wedge e_4^*$ $+ \gamma e_3^* \wedge e_4^*,$ where $\lambda\gamma \neq 0$	NH
D_{pq}^{11} $0 < p < 1$ $q = -1$	$[e_1, e_2] = e_2, [e_1, e_3] = pe_3,$ $[e_1, e_4] = qe_4$	$\lambda e_1^* \wedge e_2^* + \mu e_1^* \wedge e_3^* + \nu e_1^* \wedge e_4^*$ $+ \gamma e_2^* \wedge e_4^*,$ where $\mu\gamma \neq 0$	NH
D^{12}	$[e_1, e_2] = e_2, [e_1, e_4] = e_3,$	None	-
D_p^{13} $p \neq 0$ (generic)	$[e_1, e_2] = pe_2, [e_1, e_3] = e_3,$ $[e_1, e_4] = e_3 + e_4$	None	-
D_{-1}^{13}	$[e_1, e_2] = -e_2, [e_1, e_3] = e_3,$ $[e_1, e_4] = e_3 + e_4$	$\lambda e_1^* \wedge e_2^* + \mu e_1^* \wedge e_3^* + \gamma e_1^* \wedge e_4^*$ $+ \nu e_2^* \wedge e_4^*,$ where $\mu\nu \neq 0$	NH
D_{pq}^{14} $p \neq 0, q \geq 0$	$[e_1, e_2] = pe_2,$ $[e_1, e_3] = qe_3 - e_4,$ $[e_1, e_4] = qe_4 + e_3$	None	-
D^{15}	$[e_1, e_3] = e_2, [e_1, e_4] = e_3,$	$\lambda e_1^* \wedge e_2^* + \mu e_1^* \wedge e_3^* + \nu e_1^* \wedge e_4^*$ $+ \delta e_2^* \wedge e_4^* - \frac{1}{2}\delta e_3^* \wedge e_3^*,$ where $\delta^2(\mu^2 - 2\lambda\nu) \neq 0.$	NH
D^{16}	$[e_1, e_2] = e_2,$ $[e_1, e_3] = e_2 + e_3,$ $[e_1, e_4] = e_3 + e_4$	None	-

TABLE 4. Trivial algebras (i.e. $[\mathfrak{g}_1, \mathfrak{g}_1] = \{0\}$) with $\text{sdim} = 1|3$

The LSA	Relations in the basis: $e_1, e_2, e_3 \mid e_4$	Symplectic structure	$p(\omega)$
$(A_{3,1} + A)$	$[e_2, e_3] = e_1, [e_4, e_4] = e_1$	None	–
$(D_{p,-1/2}^3)$ $p \neq 0$	$[e_1, e_2] = e_2, [e_1, e_3] = pe_3,$ $[e_1, e_4] = \frac{1}{2}e_4, [e_4, e_4] = e_2$	None	–
$(D_{-1/2,-1/2}^3)$	$[e_1, e_2] = e_2, [e_1, e_3] = -\frac{1}{2}e_3,$ $[e_1, e_4] = \frac{1}{2}e_4, [e_4, e_4] = e_2$	$\frac{1}{2}\lambda e_1^* \wedge e_2^* + \mu e_1^* \wedge e_3^*$ $+\nu e_1^* \wedge e_4^* + \gamma e_3^* \wedge e_4^*$ $+\frac{1}{2}\lambda e_4^* \wedge e_4^*,$ where $\gamma\lambda \neq 0$	NH
$(D_{-1/2}^2)^1$	$[e_1, e_2] = e_2, [e_1, e_3] = e_2 + e_3,$ $[e_1, e_4] = \frac{1}{2}e_4, [e_4, e_4] = e_2$	None	–
$(D_{-1/2}^2)^2$	$[e_1, e_2] = e_2, [e_1, e_3] = -e_2 + e_3,$ $[e_1, e_4] = \frac{1}{2}e_4, [e_4, e_4] = e_2$	None	–

TABLE 5. Trivial algebras (i.e. $[\mathfrak{g}_1, \mathfrak{g}_1] \neq \{0\}$) with $\text{sdim} = 3 \mid 1$

The LSA	Relations in the basis: $e_1 \mid e_2, e_3, e_4$	Symplectic structure	$p(\omega)$
$(A_{1,1} + 3A)^1$	$[e_2, e_2] = e_1, [e_3, e_3] = e_1,$ $[e_4, e_4] = e_1$	None	–
$(A_{1,1} + 3A)^2$	$[e_2, e_2] = e_1, [e_3, e_3] = e_1,$ $[e_4, e_4] = -e_1$	None	–

TABLE 6. Trivial algebras (i.e. $[\mathfrak{g}_1, \mathfrak{h}_1] \neq \{0\}$) with $\text{sdim} = 1 \mid 3$

Acknowledgment. We would like to acknowledge Chengming Bai for pointing out that Novikov algebras have two superizations. Also, we would like to thank Saïd Benayadi, Isabel Cunha, and Quentin Ehret for reading the manuscript and making pertinent comments.

REFERENCES

- [B] Backhouse N., A classification of four-dimensional Lie superalgebras, *Journal of Mathematics Physics*, **19**, 2400, (1987).
- [Bai] Bai C., Bijective 1-cocycles and classification of 3-dimensional left-symmetric algebras, *Commun. Algebra* **37** (2009) 1016–1057.
- [BBB] Bajo I., Benayadi S and Bordemann M., Generalized double extension and descriptions of quadratic Lie superalgebras, arXiv: 0712.0228v1.
- [BBM] Bajo I., Benayadi S and Medina A., Symplectic structures on quadratic Lie algebras, *Journal of Algebra* **316** (2007) 174–188.
- [Bal] Balinsky A. A., Classification of the Virasoro, the Neveu-Schwarz and the Ramond type simple Lie superalgebras, *Funct. Anal. Appl.* **21** (1987) 308–309.
- [BN] Balinskii A. A., Novikov S. P., Poisson brackets of hydrodynamic type, Frobenius algebras and Lie algebras, *Sov. Math. Dokl.* **32**, 228–231 (1985).
- [Ba] Baues O., Left-symmetric algebras for $\mathfrak{gl}(n)$, *Trans. Am. Math. Soc.* **351** (1999) 2979–2996.

- [BC] Baues O. and Cortés V., Symplectic Lie groups I-III, Symplectic Reduction, Lagrangian extensions, and existence of Lagrangian normal subgroups. *Astérisque* 79, Société mathématique de France, 2016.
- [Be] Benoist Y., Une nilvariété non affine, *J. Diff. Geom.* 41 (1995), 21–52.
- [Bor] Bordemann M., Nondegenerate invariant bilinear forms on nonassociative algebras. *Acta Math. Univ. Comenianae*, Vol. LXVI, 2 (1997), 151–201.
- [BE] Bouarroudj S., Ehret Q., Double extensions of quasi-Frobenius Lie superalgebras with degenerate center, arXiv:2405.12128
- [BEM] Bouarroudj S., Ehret Q. and Maeda Y., Symplectic double extensions for restricted quasi-Frobenius Lie (super)algebras, *SIGMA. Symmetry Integrability Geom. Methods Appl.* **19**, (2023), 070, 29 pages.
- [BM] Bouarroudj S. and Maeda Y., Double and Lagrangian extensions for quasi-Frobenius Lie superalgebras, *J. Algebra Appl.* **22** (2023), no.12, Paper No. 2450001, 46 pp.
- [B1] Burde D., Left-invariant affine structures on reductive Lie groups, *J. Algebra* **181** (1996) 884–902.
- [B2] Burde D., Left-symmetric structures on simple modular Lie algebras, *J. Algebra* **169** (1994) 112–138.
- [B3] Burde D., Left-symmetric algebras, or pre-Lie algebras in geometry and physics. *Cent. Eur. J. Math.* **4** (2006) 323–357.
- [B4] Burde D., Affine structures on nilmanifolds. *Int. J. of Math.* 7 (1996), 599–616.
- [BG] Burde D., Grunewald F., Modules for certain Lie algebras of maximal class. *J. Pure Appl. Algebra* **99** (1995), 239–254.
- [CL] Chen H., Li J., Left-symmetric algebra structures on the W -algebra $W(2, 2)$, *Linear Algebra Appl.* **437** (2012) 1821–1834.
- [C] Chu B., Symplectic homogeneous spaces. *Trans. Amer. Math. Soc.* 197 (1974) 145–159.
- [DZ] Dimitrova I. and Zhang R., Left-symmetric superalgebras on special linear Lie superalgebras, *Journal of Algebra* **635** (2023) 384–410.
- [GS] Grunewald F., Segal D., On affine crystallographic groups, *J. Diff. Geom.* **40** (1994), 563–594.
- [E] Elashvili A. G. , Frobenius Lie algebras, *Funct. Anal. Appl.* **16**(4) (1982) 326–328.
- [H] Helmstetter J., Radical d’une algèbre symétrique à gauche, *Ann. Inst. Fourier Grenoble.* **29** (1979), 17–35.
- [KB] Kong X. and Bai C., Left-symmetric superalgebraic structures on the super-Virasoro algebras, *Pac. J. Math.* **235** (2008) 43–55.
- [KCB] Kong X., Chen H., Bai C., Classification of graded left-symmetric algebraic structures on Witt and Virasoro algebras, *Int. J. Math.* 22 (2011) 201–222.
- [LSoS] Leites D. (ed.) *Seminar on supersymmetry v. 1. Algebra and Calculus: Main chapters*, (J. Bernstein, D. Leites, V. Molotkov, V. Shander), MCCME, Moscow, 2012, 410 pp (in Russian; a version in English is in preparation but available for perusal).
- [M] Medina A., Flat left-invariant connections adapted to the automorphism structure of a Lie group, *J. Differ. Geom.* **16** (1981) 445–474.
- [MR] Medina R., Revoy Ph., Groupes de Lie à structure symplectique invariante, in: P. Dazord, A. Weinstein (Eds.), *Symplectic Geometry, Groupoids, and Intégrable Systems*, Séminaire Sud-Rhodanien de Géométrie, in: *Math. Sci. Res. Inst. Publ.*, Springer-Verlag, 1991, pp. 247–266.
- [MKSV] Munthe-Kaas H. Z., Stern A., Verdier O., Invariant connections, Lie algebra actions, and foundations of numerical integration on manifolds. *SIAM J. Appl. Algebra Geom.* 4 (2020), no. 1, 49–68.
- [O] Ooms I. O., On Frobenius Lie algebras, *Commun. Algebra* **8**(1) (1980) 13–52.

- [PB] Pei Y., Bai C., Balinsky-Novikov superalgebras and some infinite-dimensional Lie superalgebras, *Journal of Algebra and Its Applications*, Vol. 11, No. 6 (2012) 1250119 (29 pages).
- [S] Segal D., Free left-symmetric algebras and an analogue of the Poincaré-Birkhoff-Witt theorem, *J. Algebra* **164** (1994) 750–772.
- [TB] Tang X., Bai C., A class of non-graded left-symmetric algebraic structures on the Witt algebra, *Math. Nachr.* 285 (2012) 922–935.
- [V] Vinberg E. B., Theory of convex homogeneous cones, *Tr. Moscow Mat. Obshch.* 12 (1963) 303–358, *Transl. Moscow Math. Soc.* **12** (1963) 340–403.
- [YZ] Yang X., Zhu F., Cuspidal prehomogeneous vector spaces for reductive Lie groups and related algebraic and geometric structures, arXiv :1801 .02995.
- [Xu] Xu X. P., Variational calculus of supervariables and related algebraic structures, *J. Algebra* **223** (2000) 396–437.

DIVISION OF SCIENCE AND MATHEMATICS, NEW YORK UNIVERSITY ABU DHABI, P.O. BOX 129188,
ABU DHABI, UNITED ARAB EMIRATES.

Email address: sb3922@nyu.edu; amr1190@nyu.edu