

Two dark components of de Sitter Universe

G.E. Volovik¹

¹*Landau Institute for Theoretical Physics, acad. Semyonov av., 1a, 142432, Chernogolovka, Russia*
(Dated: October 11, 2024)

The de Sitter vacuum serves as the thermal bath for matter with local temperature, which is twice the Gibbons-Hawking temperature related to the cosmological horizon. This local temperature leads to the heating of matter. Here do not consider the real matter, but only the component which is related to the gravitational degrees of freedom and participates the thermodynamics of the de Sitter state. This gravitational component behaves as Zel'dovich stiff matter. The vacuum energy and the analog of the Zel'dovich stiff matter represent correspondingly the dark energy and dark matter. In equilibrium the positive partial pressure of dark matter compensates the negative partial pressure of quantum vacuum. That is why in the full equilibrium the total pressure is zero. This is rather similar to the correspondingly superfluid and normal components of superfluid liquid, which together produce the zero pressure of the liquid in the absence of environment. We extend this two dark components approach and the suggest the phenomenological theory, which describes the dynamics of the dark energy and the dark matter. For that we assume that the in dynamics, the gravitational dark matter also behaves as the real Zel'dovich stiff matter. In this phenomenological model, both components experience the power law decay due to the energy exchange between these components. Then it follows that their values at present time have the correct order of magnitude. How this phenomenology is supported by the microscopic physics is an open question.

PACS numbers:

Contents

| | |
|--|---|
| I. Introduction | 1 |
| II. Two component thermodynamics of the de Sitter state | 2 |
| A. Local temperature and local entropy | 2 |
| B. Modified Gibbs-Duhem relation | 2 |
| C. Two dark components of de Sitter | 3 |
| III. Two component dynamics of the de Sitter state | 4 |
| A. Phenomenological theory of two component dynamics | 4 |
| B. Dynamics without energy exchange | 4 |
| C. Dynamics with the energy exchange | 5 |
| IV. Discussion | 5 |
| References | 6 |

I. INTRODUCTION

The vacuum of the de Sitter spacetime is characterized by the local temperature $T = H/\pi$, where H is the Hubble parameter, see Ref.^{1,2} and references therein. This temperature describes the thermal processes of decay of the composite particles and the other activation processes, which are energetically forbidden in the Minkowski spacetime, but are allowed in the de Sitter background, see also Refs.³⁻⁵. In particular, this temperature determines the probability of the ionization of an atom in the de Sitter environment, $\exp(-E/T)$, where E is the ionization potential. This activation temperature is twice the Gibbons-Hawking⁶ temperature $T_{\text{GH}} = H/2\pi$ of the cosmological horizon, $T = 2T_{\text{GH}}$. As distinct from the T_{GH} , the activation temperature has no relation to the cosmological horizon. It describes the local processes, which take place well inside the cosmological horizon.

The factor of 2 difference between the local temperature and the temperature of the Hawking radiation from the cosmological horizon measured by the same observer has simple explanation. In case of the Hawking radiation the observer detects particles radiated from the horizon but cannot detect their partners, that are simultaneously (coherently) created on the other side of the horizon. This is similar to the doubling of the Hawking temperature of

black holes suggested by 't Hooft.^{7,8} In case of the black hole horizon, the suggested partner is in the mirror image of the black hole spacetime – "quantum clone inside black hole".

The local temperature $T = H/\pi$ determines the local temperature in de Sitter spacetime, giving rise to the local entropy and the modified Gibbs-Duhem relation of the de Sitter state. The latter is modified due to the gravitational degrees of freedom: the gravitational coupling $K = 1/16\pi G$ and the scalar Riemann curvature \mathcal{R} , which are the thermodynamically conjugate variables. As a result, the de Sitter can be considered as the mixture of two components: the vacuum energy (dark energy) and the artificial Zel'dovich stiff matter which plays the role of dark matter. It is the dark matter component, which is responsible for the thermodynamics of the de Sitter state with equilibrium temperature $T = H/\pi$. The two-component thermodynamics of the de Sitter equilibrium is summarized in equations (14)-(17).

In Section III we consider the dynamics of de Sitter which may follow from its thermodynamics. In the suggested phenomenological approach, it is assumed that the equations are modified in such a way, that away from equilibrium the gravitational dark matter behaves as the real stiff matter. The latter experiences the cool-down due to expansion, which is partially compensated by the energy exchange with the dark energy component. The energy exchange leads to the loss of the dark energy, and as a result both dark components have the power-law decay. Both components reach at present time the correct order of magnitude. How this phenomenology is supported by microscopic theory is an open question.

II. TWO COMPONENT THERMODYNAMICS OF THE DE SITTER STATE

A. Local temperature and local entropy

In the Painlevé-Gullstrand (PG) form^{9,10} the metric of the de Sitter expansion is:

$$ds^2 = -dt^2 + (dr - v(r)dt)^2 + r^2 d\Omega^2, \quad (1)$$

where the shift velocity is $v(r) = Hr$, and H is the Hubble parameter, see also Ref.¹¹.

From Friedmann equations of general relativity it follows that the vacuum energy density (which is the cosmological constant Λ) expressed in terms of the activation temperature $T = H/\pi$ is:²

$$\epsilon_{\text{vac}} = \Lambda = \frac{3}{8\pi G} H^2 = \frac{3\pi}{8G} T^2. \quad (2)$$

This determines the free energy density of the de Sitter state, $F = \epsilon_{\text{vac}} - T d\epsilon_{\text{vac}}/dT$, and thus the entropy density s_{dS} in this state:

$$s_{\text{dS}} = -\frac{\partial F}{\partial T} = \frac{3\pi}{4G} T = \frac{3}{4G} H. \quad (3)$$

The total entropy S of the Hubble volume surrounded by the cosmological horizon is $S = \frac{A}{4G}$, where A is the horizon area. This coincides with the entropy of the cosmological horizon suggested by Gibbons and Hawking, but it comes from the local entropy of the de Sitter state, rather than from the horizon degrees of freedom. Such agreement with the holographic bulk-horizon correspondence supports the local thermodynamics of the de Sitter state with $T = H/\pi$.

B. Modified Gibbs-Duhem relation

The quadratic dependence of vacuum energy on temperature is important for consideration of the thermodynamic Gibbs-Duhem relation for quantum vacuum. It leads to the reformulation of the vacuum pressure. The conventional vacuum pressure P_{vac} obeys the equation of state $w = -1$ and enters the energy momentum tensor of the vacuum medium in the form:

$$T^{\mu\nu} = \Lambda g^{\mu\nu} = \text{diag}(\epsilon_{\text{vac}}, P_{\text{vac}}, P_{\text{vac}}, P_{\text{vac}}), \quad P_{\text{vac}} = -\epsilon_{\text{vac}}. \quad (4)$$

In the de Sitter state the vacuum pressure is negative, $P_{\text{vac}} = -\epsilon_{\text{vac}} < 0$. This pressure P_{vac} does not satisfy the thermodynamic Gibbs-Duhem relation, $T s_{\text{dS}} = \epsilon_{\text{vac}} + P_{\text{vac}}$, because the right hand side of this equation is zero. The reason for that is that in this equation we did not take into account the gravitational degrees of freedom. Earlier it was shown, that gravity contributes to thermodynamics with the pair of the thermodynamically conjugate variables: the gravitational coupling $K = \frac{1}{16\pi G}$ and the Riemann curvature \mathcal{R} , see Refs.¹²⁻¹⁴. This is because the Einstein-Hilbert

action contains the gravitational term $K\mathcal{R}$, and its contribution to thermodynamics is somewhat similar to the work density.^{15–18} The gravitational thermodynamic variables allow us to write the modified Gibbs-Duhem relation:

$$Ts_{\text{dS}} = \epsilon_{\text{vac}} + P_{\text{vac}} - K\mathcal{R}. \quad (5)$$

This equation is obeyed, since $\epsilon_{\text{vac}} + P_{\text{vac}} = 0$, $\mathcal{R} = -12H^2$ and:

$$Ts_{\text{dS}} = -K\mathcal{R} = \frac{3}{4\pi G}H^2 = 2\epsilon_{\text{vac}}. \quad (6)$$

Eq.(5) suggests that one may introduce the pressure, which is modified by gravitational degrees of freedom:

$$P_{DM} = P_{\text{vac}} - K\mathcal{R} = \epsilon_{\text{vac}}. \quad (7)$$

We call it the pressure of the dark matter component. Then the Gibbs-Duhem relation becomes:

$$Ts_{\text{dS}} = \epsilon_{\text{vac}} + P_{DM}. \quad (8)$$

The dark matter pressure P_{DM} in Eq.(7) is positive, $P_{DM} = \epsilon_{\text{vac}} > 0$, and satisfies equation of state $w = 1$. This corresponds to stiff matter introduced by Zel'dovich,¹⁹ where the speed of sound is equal to the speed of light, $c_s^2 = c^2 dP/d\epsilon_{\text{vac}} = c^2$.

C. Two dark components of de Sitter

Let us show that the thermodynamics of the de Sitter expansion can be described in terms of two components of the de Sitter state: the vacuum component (dark energy, DE) and the stiff matter (dark matter, DM). These components have the following forms:

$$w = -1, \quad P_{\text{DE}}(H) = -\epsilon_{\text{DE}}(H) = -\frac{3}{8\pi G}H^2, \quad (9)$$

and

$$w = 1, \quad P_{\text{DM}}(T) = \epsilon_{\text{DM}}(T) = \frac{3\pi}{8G}T^2, \quad (10)$$

Then from equations (5)-(8) it follows that in equilibrium, i.e. at $T = H/\pi$, the total (thermodynamic) pressure is zero, as it should be in the absence of the external environment when the partial pressure of dark matter compensates the partial pressure of quantum vacuum:

$$P = P_{\text{DE}}(H = \pi T) + P_{\text{DM}}(H = \pi T) = 0. \quad (11)$$

In this equilibrium state, the energy density of dark matter is equal to the dark energy density:

$$\epsilon_{\text{DM}}(H = \pi T) = \epsilon_{\text{DE}}(H = \pi T). \quad (12)$$

Then the modified Gibbs-Duhem relation in Eq.(8) gets the physical meaning as the Gibbs-Duhem relation for dark matter:

$$Ts_{\text{DM}} = \epsilon_{\text{DM}} + P_{DM}, \quad s_{\text{DM}} \equiv s_{\text{dS}}. \quad (13)$$

Summarising, we have the following set of equations for the equilibrium de Sitter state, which is expressed in term of the two dark components:

$$P_{\text{DE}} = -\epsilon_{\text{DE}}, \quad (14)$$

$$P_{\text{DM}} = \epsilon_{\text{DM}}, \quad (15)$$

$$P = P_{\text{DE}} + P_{\text{DM}} = 0, \quad (16)$$

$$Ts_{\text{DM}} = \epsilon_{\text{DM}} + P_{DM}. \quad (17)$$

In this representation it is the the dark matter component, which is responsible for thermodynamics of the de Sitter state in Eq.(17). While the dark energy component represents the vacuum, and it acquires the nonzero value only

for compensation of the dark matter pressure in Eq.(16) in equilibrium. In the absence of environment, the external pressure is zero, $P = 0$. Because of that the energies of the two components in equilibrium have equal values in Eq. (12).

The similar situation takes place for the closed static Universe with positive curvature introduced by Einstein. In this case there are three contributions to the pressure: from matter, from vacuum energy and from the gravitational degrees of freedom (spatial curvature), see Sec. 29.4.5 in the book²⁰. Since the static Universe has no environment, in the full equilibrium the total pressure is zero, $P = P_{\text{vac}} + P_{\text{matter}} + P_{\text{gravity}} = 0$. The equilibrium conditions provide the connection between matter, dark energy and curvature of the Einstein Universe.

The same takes place in the absence of gravity, i.e. in Minkowski spacetime, where the Universe contains the non-gravitating quantum vacuum and the non-gravitating matter.²¹ In equilibrium and in the absence of environment, the partial pressure of matter is compensated by the partial pressure of the vacuum, $P = P_{\text{vac}} + P_{\text{matter}} = 0$. The analogous situation is in the two-fluid model of the superfluid liquid, where the role of matter is played by the system of quasiparticles on the background of the "quantum vacuum" – the ground state.²⁰

The equilibrium in Eqs. (14)-(17) is valid only for the artificial Zel'dovich matter, which comes from the gravitational degrees of freedom. For the real Zel'dovich matter, equilibrium is impossible, since due to expansion of the Universe the density of real matter must decrease with time. In the next Section we shall try to construct the dynamics of de Sitter using the possible connection between these two kinds of Zel'dovich stiff matter which originate from the matter fields and from gravity.

III. TWO COMPONENT DYNAMICS OF THE DE SITTER STATE

A. Phenomenological theory of two component dynamics

The two-fluid thermodynamics of the de Sitter state discussed in Sec. II is valid in thermal equilibrium, when $T = H/\pi$. Let us now speculate on what happens in the non-equilibrium state when the temperature of stiff matter is not equal to its equilibrium value H/π . In this case the energy exchange between two components may lead the temperature of stiff dark matter towards the equilibrium temperature $T = H/\pi$. The question arises how to describe the non-equilibrium state, in which $T \neq H/\pi$. Here we shall use the following phenomenological approach suggested in Ref.².

In Sec. II we obtained that in the de Sitter thermodynamics, the gravitational Zel'dovich stiff matter behaves as the real Zel'dovich matter. Now we assume that the Einstein equations are modified in such a way that the gravitational dark matter behaves as the real Zel'dovich matter also in dynamics. Then the dark matter decays with time due to expansion, but it also gets energy from the energy exchange with the dark energy component. On the other hand this energy exchange leads to decrease of the vacuum energy density, and finally the Universe approaches the equilibrium Minkowski vacuum state.

So, let us consider how the de Sitter state is modified if one takes into account that the temperature of dark matter changes both due to the expansion and due to the energy exchange with the dark energy. The only assumption in this phenomenological approach is that due to the energy exchange, the temperature of dark matter tends to approach the de Sitter local temperature, $T \rightarrow H/\pi$.

B. Dynamics without energy exchange

The Einstein equations for standard de Sitter expansion of the Universe give the equation for the stiff matter with $w = 1$, which density decreases with expansion of space:

$$\partial_t \epsilon_{\text{DM}} = -3H(w + 1)\epsilon_{\text{DM}} = -6H\epsilon_{\text{DM}}, \quad (18)$$

while the equation for the vacuum energy density with $w = -1$ is:

$$\partial_t \epsilon_{\text{DE}} = 0. \quad (19)$$

Of course, for the original gravitational stiff matter one has $\partial_t \epsilon_{\text{DM}} = 0$ instead of Eq.(18). But here we consider the phenomenological model, in which the real Zel'dovich matter has exactly the same energy density as the gravitational Zel'dovich matter in Eq.(10).

C. Dynamics with the energy exchange

Due to the energy exchange with the dark energy, the energy density of dark matter tends to approach the dark energy density in Eq.(12), $\epsilon_{\text{DM}}(T) \rightarrow \epsilon_{\text{DE}}(H)$. This suggests the following modification of Eq.(18), which now includes the term describing the energy exchange between DE and DM:

$$\partial_t \epsilon_{\text{DM}}(T) = -6H(\epsilon_{\text{DM}}(T) - \epsilon_{\text{DE}}(H)). \quad (20)$$

This term allows the thermalization of dark matter by the energy exchange with the dark energy.

The gain $6H\epsilon_{\text{DE}}(H)$ of the dark matter energy density in Eq.(20) must be compensated by loss of the dark energy. This modifies Eq.(19):

$$\partial_t \epsilon_{\text{DE}}(H) = -6H\epsilon_{\text{DE}}(H). \quad (21)$$

The equation (21) gives the following time dependence of dark energy at large time:

$$\epsilon_{\text{DE}}(t) = \frac{1}{6\pi G t^2}. \quad (22)$$

This gives the reasonable order of magnitude of the vacuum energy density in the present time. The same behaviour of the vacuum energy density was obtained using the Hawking 4-form field,²² see Refs.^{14,23}. In this case the role of dark matter is played by the oscillations of the vacuum energy during its decay.²⁴

However, in this simple model the dark matter, which obeys Eq.(20), does not approach the dark energy behaviour in Eq.(22), and instead it contains the logarithm:

$$\epsilon_{\text{DM}}(t) = \frac{\ln(t/t_0)}{3\pi G t^2}. \quad (23)$$

In this model the dark matter dominates, but anyway both components have the correct order of magnitude at present time, $\epsilon_{\text{DME}}(t = t_{\text{present}}) \sim \epsilon_{\text{DM}}(t = t_{\text{present}}) \sim H^2/G$. Finally they decay to zero value at $t \rightarrow \infty$, when the system approaches the fully equilibrium state with zero temperature – the Minkowski vacuum.

IV. DISCUSSION

It is shown that the de Sitter Universe can be represented in terms of two dark components in equations (14)-(17). These are the dark energy (the energy of the vacuum) and the dark matter, which comes from the gravitational degrees of freedom. Their partial pressures compensate each other in the equilibrium de Sitter state. In this representation, the dark matter is responsible for the thermodynamics of the de Sitter state, which is characterized by the local temperature $T = H/\pi$ and is similar to the thermodynamics of the Zel'dovich stiff matter. Let us recall that the local temperature of the de Sitter state is twice the Gibbons-Hawking temperature related to the cosmological horizon, and the stiff matter behaviour of the dark matter component is the consequence of this value of the local temperature.

We tried to extend this two-fluid approach to the dynamics of the Universe, assuming that in dynamics the dark matter component is also equivalent to the Zel'dovich stiff matter. Then we obtained the power-law decay of both dark the matter component in Eq.(23) and the dark energy component in Eq.(22). Both components acquire the correct order of magnitude at present time, and both finally decay to zero value at $t \rightarrow \infty$.

The reasonable decay law (22) for dark energy takes place only for the stiff matter. The question then arises: what might be the physical origin of this stiff matter and how is it related to the gravitational hard matter discussed in the Section II. In principle, it is possible, that the stiff matter comes just from the gravitational dark matter, and the modified equations reflect the microscopic physics, which supports both the thermodynamics and dynamics of the de Sitter. The question then remains open as to how to describe the corresponding degrees of freedom that lead to the considered thermodynamics and dynamics.

One possibility is to introduce the proper vacuum variable, which describes the quantum vacuum and its vacuum energy density. The working example of such variable is the Hawking 4-form field.²² This variable was used in particular in the so-called q -theory.^{14,21} However, the suggested form of the q -theory gives rise to the cold dark matter instead of the stiff matter.²⁴ The analogue of the cold matter is produced by the fast oscillations of the q -field. This q -theory mechanism also gives the required power-law decay of the dark energy similar to that in Eq.(22). However, this decay to the Minkowski vacuum takes place only in case of the proper choice of the value of corresponding analog of the chemical potential. This represents some form of fine tuning, while the two fluid dynamics does not require

the fine tuning. That is why one should try the other variables. For example, the stiff era can be generated by an appropriate $F(R)$ gravity term.²⁵

-
- ¹ G.E. Volovik, On the global temperature of the Schwarzschild-de Sitter spacetime, *Pis'ma v ZhETF* **118**, (2023), *JETP Lett.* **118**, (2023), <https://doi.org/10.1134/S0021364023601173>, arXiv:2304.09847.
- ² G.E. Volovik, Thermodynamics and decay of de Sitter vacuum, *Symmetry* **16**, 763 (2024), <https://doi.org/10.3390/sym16060763>
- ³ N. Arkani-Hamed and J. Maldacena, *Cosmological Collider Physics*, arXiv:1503.08043.
- ⁴ Matthew Reece, Lian-Tao Wang and Zhong-Zhi Xianyu Large-field inflation and the cosmological collider, *Phys. Rev. D* **107**, L101304 (2023).
- ⁵ Henry Maxfield and Zahra Zahraee, Holographic solar systems and hydrogen atoms: non-relativistic physics in AdS and its CFT dual, *JHEP* **11** (2022) 093.
- ⁶ G. W. Gibbons and S. W. Hawking, Cosmological event horizons, thermodynamics, and particle creation, *Phys. Rev. D* **15**, 2738 (1977).
- ⁷ G. 't Hooft, Quantum clones inside black holes, *Universe* **8**, 537 (2022).
- ⁸ Gerard 't Hooft, How an exact discrete symmetry can preserve black hole information or Turning a black hole inside out, *J. Phys.: Conf. Ser.* **2533**, 012015 (2023).
- ⁹ P. Painlevé, La mécanique classique et la théorie de la relativité, *C. R. Acad. Sci. (Paris)* **173**, 677 (1921).
- ¹⁰ A. Gullstrand, Allgemeine Lösung des statischen Einkörper-problems in der Einsteinschen Gravitations-theorie, *Arkiv. Mat. Astron. Fys.* **16**, 1-15 (1922).
- ¹¹ Rudeep Gaur and Matt Visser, Cosmology in Painlevé–Gullstrand coordinates, *JCAP*09(2022)030, arXiv:2207.08375.
- ¹² G.E. Volovik, Macroscopic quantum tunneling: from quantum vortices to black holes and Universe, *ZhETF* **162**, 449–454 (2022), *JETP* **135**, 388–408 (2022), arXiv:2108.00419.
- ¹³ G.E. Volovik, Varying Newton constant and black hole to white hole quantum tunneling, *MDPI, Universe* **6**, 133 (2020), arXiv:2003.10331.
- ¹⁴ F.R. Klinkhamer and G.E. Volovik, Dynamic vacuum variable and equilibrium approach in cosmology, *Phys. Rev. D* **78**, 063528 (2008); arXiv:0806.2805 [gr-qc].
- ¹⁵ S.A. Hayward, Unified first law of black-hole dynamics and relativistic thermodynamics, *Class. Quantum Grav.* **15**, 3147–3162 (1998).
- ¹⁶ Sean A. Hayward, Shinji Mukohyama and M.C. Ashworth, Dynamic black-hole entropy, *Phys. Lett. A* **256**, 347–350 (1999), arXiv:gr-qc/9810006.
- ¹⁷ T. Jacobson, Thermodynamics of Spacetime: The Einstein Equation of State, *Phys. Rev. Lett.* **75**, 1260–1263 (1995).
- ¹⁸ Shin'ichi Nojiri, Sergei D. Odintsov, Tanmoy Paul and Soumitra SenGupta, Horizon entropy consistent with FLRW equations for general modified theories of gravity and for all EoS of the matter field, arXiv:2307.05011.
- ¹⁹ Ya. B. Zel'dovich, The equation of state at ultrahigh densities and its relativistic limitations, *JETP* **14**, 1143 (1962).
- ²⁰ G.E. Volovik, *The Universe in a Helium Droplet*, Clarendon Press, Oxford (2003),
- ²¹ F.R. Klinkhamer and G.E. Volovik, Self-tuning vacuum variable and cosmological constant, *Phys. Rev. D* **77**, 085015 (2008); arXiv:0711.3170.
- ²² S.W. Hawking, The cosmological constant is probably zero, *Phys. Lett.* **134B**, 403 (1984).
- ²³ G.E. Volovik, From analogue models to gravitating vacuum, in: *Analogie spacetimes: The first thirty years*, eds. V. Cardoso, L.C.B. Crispino, S. Liberati, E.S. de Oliveira, M. Visser, Editoria Livraria da Fisica, Sao Paulo 2013, pp. 263–290; arXiv:1111.1155.
- ²⁴ F.R. Klinkhamer and G.E. Volovik, Dark matter from dark energy in q -theory, *Pis'ma ZhETF* **105**, 62–63 (2017), *JETP Lett.* **105**, 74–77 (2017), arXiv:1612.02326.
- ²⁵ S.D. Odintsov and V.K. Oikonomou, Early-time cosmology with stiff era from modified gravity *Phys. Rev. D* **96**, 104059 (2017).