Genus distribution polynomials for bicellular bicolored maps all with real zeros

Zi-Wei Bai, Ricky Xiaofeng Chen *

School of Mathematics, Hefei University of Technology 485 Danxia Road, Hefei, Anhui 230601, P. R. China 2021111458@mail.hfut.edu.cn, xiaofengchen@hfut.edu.cn

Abstract. Since 1960s, enumerating bicolored maps (thus maps) according to the numbers (and possibly types) of edges, faces, white vertices, black vertices and genus has been an important topic arising in many fields of mathematics and physics. To the best of our knowledge, no explicit formulas are known for the number of bicolored maps given the first four parameters unless the given number of faces is one. There is actually a vast literature for the one-face case. In particular, Jackson (1987), Zagier (1995) and Stanley (2011) respectively obtained some expressions for the generating polynomial of the numbers of one-face bicolored maps with given number of edges and white vertex degree distribution while tracking the number of black vertices. In this paper, we obtain an explicit formula for the corresponding generating polynomial for that of bicolored maps with two faces, i.e., bicellular, one face of length p and the other of length n - p. We also prove that the polynomial essentially has only real zeros and thus the log-concavity of the counting sequence.

Keywords: Bicolored maps, Permutation product, Genus distribution, Dessin d'enfant, Group character, Log-concavity Mathematics Subject Classifications 2020: 05A15, 05E10, 14H57

1 Introduction

A map is a 2-cell embedding of a connected graph in an orientable surface [23]. A bicolored map is a map where the vertices are colored black and white such that every edge is incident to two vertices of different colors. If every white vertex is of degree two in a bicolored map, then we may replace every white vertex and its incident edges with a single edge to obtain a map (with only black vertices). Thus, maps may be viewed as special bicolored maps. Bicolored maps

^{*}Corresponding author (ORCID: 0000-0003-1061-3049)



Figure 1: A labeled bicolored map of 8 edges and genus 2.

are also known as Grothendieck's dessins d'enfants in studying Riemann surfaces [16,23,24,34]. These objects lie at the crossroad of a number of research fields, e.g., topology, mathematical physics and representation theory, and we will be mainly interested in their enumerative and combinatorial properties.

An edge-end of an edge is sometimes called a half-edge (or a dart). A rooted map is a map where one half-edge is distinguished and called the root. As for rooted bicolored maps, the root is assumed to be a half-edge around a black vertex by convention. The considered maps and bicolored maps are always rooted in this paper. A labeled bicolored map of n edges is a rooted bicolored map with edges respectively having a label from the label set $[n] = \{1, 2, \ldots, n\}$ where in particular the edge containing the root is labeled 1. It is well-known (e.g. [23]) that a labeled bicolored map of n edges can be encoded into a triple in the set \mathfrak{S}_n of permutations on [n]. Specifically, a triple (α, β, γ) of permutations such that $\gamma = \alpha\beta$ (compose from right to left) and the group generated by α and β acts transitively on [n] determines a labeled bicolored map in the following manner: the cycles of γ encode the faces, the cycles of α encode the white vertices, and the cycles of β encode the black vertices, and an edge connects the same label in α and β . See Figure 1 for an example where the black vertices give the permutation (in cycle notation) $\beta = (1836)(2754)$, and the white vertices give the permutation $\alpha = (135)(26478)$. A face is a region of the embedding surface bounded by a cyclic sequence of edges (\ldots, i, i', \ldots) , where i' is the edge obtained by starting with the edge i, going counterclockwisely to the neighboring edge e around a black vertex, and then going counterclockwisely to the neighboring edge around a white vertex. It is not hard to see that a face corresponds to a cycle in $\gamma = (12857)(346)$. The requirement of transitive action forces the underlying graph to be connected. If the requirement is not satisfied, the triple may be viewed as a disconnected bicolored map. In the following, bicolored maps may be disconnected unless explicitly stated otherwise.

According to the Euler characteristic formula, the genus g of a bicolored map (defined as the genus of the underlying embedding surface) satisfies:

$$v - e + f = 2 - 2g, (1)$$

where v, e, f are respectively the numbers of vertices, edges and faces of the map. This translates to the following relation:

$$\kappa(\alpha) + \kappa(\beta) - n + \kappa(\gamma) = 2 - 2g,\tag{2}$$

where $\kappa(\pi)$ denotes the number of cycles in the permutation π .

Since 1960s, the enumeration of bicolored maps (thus maps) has been being centered around studying the enumerative and combinatorial properties of sets of bicolored maps filtered by different combinations of these parameters, i.e., the number of edges, the genus, the number of black and/or white vertices (and vertex degree distribution), and the number of faces (and face length distribution). Many of these studies are based on the permutation triple representation.

Up to our knowledge, if the set for enumeration contains bicolored maps with multiple faces, then usually the bicolored maps in the set have the same number of edges and genus, while both the numbers of faces and vertices may vary. See for instance [2,3,14,15,17,31] counting maps of n edges and genus g. If the set contains bicolored maps with the same number of faces (and possibly face length distribution), the same number of edges and genus, then it seems much harder to obtain the size of the set, even asymptotically. A few results exist (e.g., certain asymptotics in [4], recurrences in [15, 25], and functional relations in [9, 24]) and no explicit formulas are known unless the number of faces is one, i.e., one-face maps or unicellular maps. In fact, there are vast literature on studying the "easier" one-face (equivalently, one white vertex or one black vertex) bicolored maps. See [5, 7, 8, 10, 11, 19–21, 26, 27, 30, 32, 33] and the references therein. In particular, formulas for the genus distribution polynomial for one-face bicolored maps with any prescribed white vertex degree distribution have been obtain in Stanley [30], and somewhat more implicit in Jackson [21] and Zagier [33], all relying on the group character approach. Our main contributions here are explicit formulas for the genus distribution polynomial for bicellular (i.e., two-face) bicolored maps and its consequences.

To state our results more precisely, we introduce some notation first. The number of disjoint cycles of a permutation $\pi \in \mathfrak{S}_n$ is denoted by $\kappa(\pi)$, and the multiset consisting of the lengths of these disjoint cycles is called the cycle-type of π , and denoted by $\rho(\pi)$. We write $\rho(\pi)$ as a partition of n, i.e., a nonincreasing positive integer sequence $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_k)$ such that $\lambda_1 + \cdots + \lambda_k = n$. If in λ , there are m_i parts equal to i, we also write λ as $[1^{m_1}, 2^{m_2}, \ldots, n^{m_n}]$. We often omit i^{m_i} when $m_i = 0$ and simply write i^1 as i. These two representations will be used interchangeably whichever is more convenient. An n-cycle on [n] is a permutation of cycle-type (n), and a permutation on the set [2n] of cycle-type $[2^n]$ is called a fixed-point free involution.

Stanley [30] studied the following cycle distribution polynomial:

$$P_{\mu}(x) = \frac{1}{|C_{\mu}|} \sum_{\alpha \in \mathfrak{S}_{n}, \, \rho(\alpha) = \mu} x^{\kappa(\alpha\gamma)},\tag{3}$$

where γ is a fixed *n*-cycle and C_{μ} is the set of permutations on [n] with cycle-type μ . This can be interpreted as the distribution of the number of black vertices (thus genus) over all one-face bicolored maps with *n* edges and white vertex degree distribution given by $\mu \vdash n$. Note that the celebrated Harer-Zagier formula [20] is for the case $\mu = [2^n] \vdash 2n$. In [30], an expression for $P_{\mu}(x)$ was given in terms of the backward shift operator, and it was shown that $P_{\mu}(x)$ has purely imaginary zeros. In order for studying bicolored maps with multiple faces, one may study the above polynomial with γ being a permutation with multiple cycles. As discussed above, no formulas have been obtained for these polynomials.

Our first main result is a nice expression for the genus distribution polynomial of bicellular bicolored maps of face-type [p, n-p] (i.e., one face of length p and one face of length n-p) and white vertex degree distribution (abbr. WDD) μ :

$$P_{[p,n-p],\mu}(x) = \frac{1}{|\mathcal{C}_{\mu}|} \sum_{\alpha \in \mathfrak{S}_n, \, \rho(\alpha) = \mu} x^{\kappa(\alpha\gamma)},$$

where γ is a fixed permutation of cycle-type [p, n - p], and μ could be any fixed partition of n except some cases. We also obtain the genus distribution polynomials $P^{\bullet}_{[p,n-p],\mu}(x)$ for the corresponding connected bicolored maps. Based on these, we next prove that every root of the polynomial $P^{\bullet}_{[p,n-p],\mu}(x)$ is of purely imaginary. The latter implies the log-concavity of the coefficients which is closely related to the wellknown, three-decade old log-concavity conjecture of Gross, Robbins and Tucker [18] for topological embeddings of graphs.

2 Preliminaries

Let C_{λ} denote the conjugacy class of \mathfrak{S}_n containing permutations of the same cycle-type λ . For $\lambda = [1^{m_1}, 2^{m_2}, \ldots, n^{m_n}] \vdash n$, the number of elements contained in C_{λ} is well known to be

$$|\mathcal{C}_{\lambda}| = \frac{n!}{z_{\lambda}}, \text{ where } z_{\lambda} = \prod_{i=1}^{n} i^{m_i} m_i!.$$

The permutations on [n] having exactly k cycles are counted by the signless Stirling number of the first kind $\begin{bmatrix} n \\ k \end{bmatrix}$ which satisfies

$$(x)_n := x(x-1)\cdots(x-n+1) = \sum_{k=1}^n (-1)^{n-k} {n \brack k} x^k.$$
(4)

Let χ^{λ} denote the character associated to the irreducible representation indexed by λ , and f^{λ} denote its dimension. The readers are invited to consult Stanley [29], and Sagan [28] for the character theory of symmetric groups.

Lemma 2.1 (The hook length formula [28, 29]).

$$f^{\lambda} = \frac{n!}{\prod_{u \in \lambda} h_u},\tag{5}$$

where for $u = (i, j) \in \lambda$, i.e., the (i, j) cell in the Young diagram of λ , h(u) is the hook length of the cell u.

Suppose C is a conjugacy class of \mathfrak{S}_n indexed by $\alpha \vdash n$. We view $\chi^{\lambda}(\alpha)$ and $\chi^{\lambda}(C)$ as the same. Let

$$\mathfrak{m}_{\lambda,m} = \prod_{u \in \lambda} \frac{m + c(u)}{h(u)}, \qquad \mathfrak{c}_{\lambda,m} = \sum_{d=0}^m (-1)^d \binom{m}{d} \mathfrak{m}_{\lambda,m-d}.$$

where for $u = (i, j) \in \lambda$, c(u) = j - i. The following theorem was proved in [12].

Theorem 2.2 (Chen [12]). Let $\xi_{n,m}(C_1, \ldots, C_t)$ be the number of tuples $(\sigma_1, \sigma_2, \ldots, \sigma_t)$ such that the permutation $\pi = \sigma_1 \sigma_2 \cdots \sigma_t$ has m cycles, where σ_i belongs to a conjugacy class C_i of \mathfrak{S}_n . Then we have

$$\xi_{n,m}(C_1,\ldots,C_t) = \left(\prod_{i=1}^t |C_i|\right) \sum_{k=0}^{n-m} {m+k \brack m} \frac{(-1)^k}{(m+k)!} W_{n,m+k}(C_1,\ldots,C_t),$$
(6)

where

$$W_{n,m+k}(C_1,\ldots,C_t) = \sum_{\lambda \vdash n} \frac{\mathfrak{c}_{\lambda,m+k}}{(f^{\lambda})^{t-1}} \prod_{i=1}^t \chi^{\lambda}(C_i).$$
(7)

The case m = n of eq. (6) reduces to the famous Frobenius identity (e.g. [23]). See Chen [12] for discussion. The second author [13] has applied Theorem 2.2 to studying one-face hypermaps, and a plethora of results, e.g., the Harer-Zagier formula [20] and the Jackson formula [22, 27], have been obtained in a unified way. In the following, we will further explore Theorem 2.2 in order for studying bicellular maps.

3 Enumeration of bicellular bicolored maps

Obviously, the genus distribution polynomial

$$P_{[p,n-p],\mu}(x) = \frac{1}{|C_{\mu}| \cdot |C_{[p,n-p]}|} \sum_{m} \xi_{n,m}(C_{\mu}, C_{[p,n-p]}) x^{m}$$
$$= \sum_{m} \sum_{k=0}^{n-m} {m+k \brack m} \frac{(-1)^{k}}{(m+k)!} W_{n,m+k}(C_{\mu}, C_{[p,n-p]}) x^{m},$$

where the corresponding W-number is as follows:

$$W_{n,m+k} = \sum_{\lambda \vdash n} \frac{\mathfrak{c}_{\lambda,m+k}}{f^{\lambda}} \chi^{\lambda}([p,n-p]) \,\chi^{\lambda}(\mu). \tag{8}$$

Here we first assume p < n - p and $n - 2p \ge 2$. Throughout the section, we also assume

$$\mu = (\mu_1, \mu_2, \dots, \mu_d) = [l^{n_l}, \dots, q^{n_q}] \vdash n,$$

where $l \ge p+1$ and $q \ge p+1$ are respectively the minimum size and maximum size of a part in μ . Sometimes the number of parts contained in λ is written as $\ell(\lambda)$.

In the following, the evaluation of the involved characters are based on the Murnaghan-Nakayama rule (see, e.g., Stanley [29], and Sagan [28]).

Lemma 3.1. For $\lambda \vdash n \geq 2p + 2$, we have

٠

$$\chi^{\lambda}([p, n-p]) = \begin{cases} (-1)^j, & \lambda = [1^j, n-j], \ 0 \le j \le p-1, \\ (-1)^{j+1}, & \lambda = [1^j, n-j], \ n-p \le j \le n-1. \end{cases}$$
(9)

- For $\lambda = [1^j, 2^k, p-k+1, n-j-k-p-1]$ where $0 \le k \le p-1$ and $0 \le j \le n-2p-2$, $\chi^{\lambda}([p, n-p]) = (-1)^{j+1};$
- For $\lambda = [1^j, 2^k, p-k-j, n-k-p]$ where $0 \le k, j$ and $k+j \le p-2, \ \chi^{\lambda}([p, n-p]) = (-1)^j;$
- For $\lambda = [1^j, 2^k, p k j, n k p]$ where $0 \le k$ and $j \ge n 2p, \ \chi^{\lambda}([p, n p]) = (-1)^j;$
- For any other $\lambda \vdash n$, $\chi^{\lambda}([p, n-p]) = 0$.

The calculation of $\chi^{\lambda}(\mu)$ is crucial and generally hard. However, we observe that only the values for λ that give a non-zero value in Lemma 3.1 are needed.



Figure 2: Two scenarios that $\chi^{\lambda}(\mu)$ may not be zero, scenario (i) (left) and scenario (ii) (right), where the rim hook with slash is of size l > p.

Lemma 3.2. There holds

$$\chi^{\lambda}(\mu) = \begin{cases} (-1)^j, & \lambda = [1^j, n-j], \ 0 \le j \le p-1, \\ (-1)^{j+1-\ell(\mu)}, & \lambda = [1^j, n-j], \ n-p \le j \le n-1. \end{cases}$$
(10)

For $\lambda = [1^j, 2^k, p - k + 1, n - j - k - p - 1]$ where $0 \le k \le p - 1$ and $0 \le j \le n - 2p - 2$,

$$\chi^{\lambda}(\mu) = \sum_{j_{l}=0}^{n_{l}-1} \sum_{j_{l+1}=0}^{n_{l+1}} \cdots \sum_{j_{q}=0}^{n_{q}} \binom{n_{l}-1}{j_{l}} \binom{n_{l+1}}{j_{l+1}} \cdots \binom{n_{q}}{j_{q}} \times \left\{ (-1)^{j-\sum_{i\geq l} j_{i}} \delta_{j+p+1-l,\sum_{i\geq l} ij_{i}} + (-1)^{j+1-\sum_{i\geq l} j_{i}} \delta_{j+p+1,\sum_{i\geq l} ij_{i}} \right\}, \quad (11)$$

where $\delta_{x,y} = 1$ if x = y, and 0 otherwise, and $\chi^{\lambda}(\mu) = 0$ for any other λ that $\chi^{\lambda}([p, n-p])$ may not be zero.

Proof. Eq. (10) is easy to verify. We only prove the case $\lambda = [1^j, 2^k, p-k+1, n-j-k-p-1]$. There are only two scenarios that $\chi^{\lambda}(\mu)$ may not be zero for $\min(\mu) \ge p+1$ according to the Murnaghan-Nakayama rule:

- (i) The rim hook (or border strip) containing the cell (1,1)) is of size $l \ge p+1$ and also contains the cells $(1,2), \ldots, (1,p-k)$ but not the cell (1,p-k+1). See an illustration in Figure 2 left;
- (ii) The rim hook containing the cell (1,1) is of size $l \ge p+1$ and also contains the cells $(2,1),\ldots,(k+1,1)$ but not the cell (k+2,1). See an illustration in Figure 2 right.

For (i), the rim hook containing the cell (1, 1) necessarily contains the cell (k + 2, 1) since the size of the rim hook is at least p + 1. Then, this rim hook separates the remaining diagram into two independent pieces: the lower-left piece of size j + p + 1 - l and the upper-right piece of size n - j - p - 1. Next, we just need to fill each piece with the remaining $n_l - 1$ rim hooks of size l

and n_i rim hooks of size *i* for $l < i \leq q$. Suppose j_i rim hooks of size *i* go to the lower-left piece and the rest goes to the upper-right piece. It is not hard to see that the total height of these rim hooks is

$$j + k + 1 - \sum_{i \ge l} j_i + k + 1.$$

Thus, the contribution of such a filling to $\chi^{\lambda}(\mu)$ is given by $(-1)^{j-\sum_{i\geq l}j_i}\delta_{j+p+1-l,\sum_{i\geq l}ij_i}$. Therefore, the contribution of the first scenario is

$$\sum_{j_l=0}^{n_l-1} \sum_{j_{l+1}=0}^{n_{l+1}} \cdots \sum_{j_q=0}^{n_q} \binom{n_l-1}{j_l} \binom{n_{l+1}}{j_{l+1}} \cdots \binom{n_q}{j_q} \{(-1)^{j-\sum_{i\geq l} j_i} \delta_{j+p+1-l,\sum_{i\geq l} ij_i} \}$$

Analogously, the contribution of the second scenario is

$$\sum_{j_l=0}^{n_l-1} \sum_{j_{l+1}=0}^{n_{l+1}} \cdots \sum_{j_q=0}^{n_q} \binom{n_l-1}{j_l} \binom{n_{l+1}}{j_{l+1}} \cdots \binom{n_q}{j_q} \{(-1)^{j+1-\sum_{i\geq l} j_i} \delta_{j+p+1,\sum_{i\geq l} ij_i}\},$$

and the proof follows.

Next, we compute the factor $\frac{c_{\lambda,m+k}}{f^{\lambda}}$ in eq. (8), only for these λ 's which return a non-zero value in Lemma 3.2. First of all, we have

$$\frac{\mathfrak{c}_{\lambda,m}}{f^{\lambda}} = \frac{\sum_{d=0}^{m} (-1)^d \binom{m}{d} \prod_{u \in \lambda} \frac{m-d+c(u)}{h(u)}}{\frac{n!}{\prod_{u \in \lambda} h_u}} = \sum_{d=0}^{m} \frac{(-1)^d}{n!} \binom{m}{d} \prod_{u \in \lambda} \{m-d+c(u)\}.$$
 (12)

With this, the following two lemmas are obvious and their proofs are omitted.

Lemma 3.3. Suppose $\lambda = [1^j, n - j]$. Then, we have

$$\frac{\mathfrak{c}_{\lambda,r}}{f^{\lambda}} = \sum_{d=0}^{r} (-1)^d \binom{r}{d} \binom{r-d+n-j-1}{n}.$$

Lemma 3.4. For $\lambda = [1^j, 2^k, p-k+1, n-j-k-p-1]$ where $0 \le k \le p-1$ and $0 \le j \le n-2p-2$,

$$\frac{\mathfrak{c}_{\lambda,r}}{f^{\lambda}} = \frac{p!(n-p)!}{n!} \sum_{d=0}^{r} (-1)^d \binom{r}{d} \binom{r-d+p-k-1}{p} \binom{r-d+n-j-k-p-2}{n-p}.$$

In the rest of the paper, some further notation will be used: as usual, the notation $[y^n]f(y)$ means taking the coefficient of the term y^n in the power series expansion of f(y), and

$$\mathcal{V}_{\mu}(y) := \prod_{i=1}^{d} \{ (1+y)^{\mu_i} - 1 \}.$$

Theorem 3.5. The following holds:

$$P_{[p,n-p],\mu}(x) = \frac{p!(n-p)!}{n!} [y^{n-p}] \mathcal{V}_{\mu}(y) \sum_{i=0}^{p-1} \binom{x+i}{p} (1+y)^{x+i-p}.$$
 (13)

Sketch of the proof. In the light of Lemma 3.1 and Lemma 3.2, we can break the quantity $W_{n,r}$ into the following form:

$$W_{n,r} = \sum_{\substack{\lambda = [1^j, n-j]\\0 \le j \le p-1}} \frac{\mathfrak{c}_{\lambda,r}}{f^{\lambda}} \chi^{\lambda}([p, n-p]) \chi^{\lambda}(\mu) + \sum_{\substack{\lambda = [1^j, n-j],\\n-p \le j \le n-1}} \frac{\mathfrak{c}_{\lambda,r}}{f^{\lambda}} \chi^{\lambda}([p, n-p]) \chi^{\lambda}(\mu) + \sum_{\substack{\lambda = [1^j, 2^k, p-k+1, n-j-k-p-1],\\0 \le k \le p-1, \ 0 \le j \le n-2p-2]}} \frac{\mathfrak{c}_{\lambda,r}}{f^{\lambda}} \chi^{\lambda}([p, n-p]) \chi^{\lambda}(\mu).$$

We compute the third sum first.

$$A_{3} := \sum_{\lambda = [1^{j}, 2^{k}, p-k+1, n-j-k-p-1]} \frac{\mathfrak{c}_{\lambda, r}}{f^{\lambda}} \chi^{\lambda}([p, n-p]))\chi^{\lambda}(\mu)$$

$$= \sum_{j, k} \left\{ \sum_{d} (-1)^{d} \binom{r}{d} \binom{r-d+n-j-k-p-2}{n-p} \binom{r-d+p-k-1}{p} \frac{(n-p)!p!}{n!} \right\}$$

$$\times (-1)^{j+1} \left\{ \sum_{j_{l}}^{n_{l}-1} \sum_{j_{l+1}}^{n_{l+1}} \cdots \sum_{j_{q}}^{n_{q}} \binom{n_{l}-1}{j_{l}} \binom{n_{l+1}}{j_{l+1}} \cdots \binom{n_{q}}{j_{q}} \times \left\{ (-1)^{j+\sum_{i\geq l} j_{i}} \delta_{j+p+1-l,\sum_{i\geq l} ij_{i}} + (-1)^{j+1+\sum_{i\geq l} j_{i}} \delta_{j+p+1,\sum_{i\geq l} ij_{i}} \right\} \right\}.$$

We next compute

$$B_{1} := \sum_{j,k} \left\{ \sum_{d=0}^{r} (-1)^{d} {\binom{r}{d}} {\binom{r-d+n-j-k-p-2}{n-p}} {\binom{r-d+p-k-1}{p}} \right\} (-1)^{j+1} \\ \times \left\{ \sum_{j_{l}}^{n_{l}-1} \sum_{j_{l+1}}^{n_{l+1}} \cdots \sum_{j_{q}}^{n_{q}} {\binom{n_{l}-1}{j_{l}}} {\binom{n_{l+1}}{j_{l+1}}} \cdots {\binom{n_{q}}{j_{q}}} (-1)^{j+\sum_{i\geq l} j_{i}} \delta_{j+p+1-l,\sum_{i\geq l} ij_{i}} \right\} \\ = \sum_{k} \sum_{d} (-1)^{d} {\binom{r}{d}} {\binom{r-d+p-k-1}{p}} \left\{ \sum_{j_{l}}^{n_{l}-1} \sum_{j_{l+1}}^{n_{l+1}} \cdots \sum_{j_{q}}^{n_{q}} {\binom{n_{l}-1}{j_{l}}} {\binom{n_{l+1}}{j_{l+1}}} \cdots {\binom{n_{q}}{j_{q}}} \right\} \\ \times (-1)^{1+\sum_{i\geq l} j_{i}} {\binom{r-d+n-(\sum_{i\geq l} ij_{i}+l-p-1)-k-p-2}{n-p}} - U_{1},$$

where

$$U_1 = \sum_{k=0}^{p-1} \sum_d (-1)^d \binom{r}{d} \binom{r-d+p-k-1}{p} (-1)^{\ell(\mu)} \binom{r-d-k-1}{n-p}.$$

In the above computation, we are supposed to sum over all $0 \le j \le n - 2p - 2$ and j_i 's such that $j = \sum_{i\ge l} ij_i + l - p - 1$ (i.e., the Kronecker delta factor). However, with a careful analysis, we find that the only case that needs to be excluded is the case $j_l = n_l - 1$ and $j_i = n_i$ for i > l whence the subtraction of U_1 .

Then, we have

$$B_{1} + U_{1} = [y^{n-p}] \sum_{k} \sum_{d} (-1)^{d} \binom{r}{d} \binom{r-d+p-k-1}{p}$$

$$\times \sum_{j_{l}}^{n_{l}-1} \sum_{j_{l+1}}^{n_{l+1}} \cdots \sum_{j_{q}}^{n_{q}} \binom{n_{l}-1}{j_{l}} \binom{n_{l+1}}{j_{l+1}} \cdots \binom{n_{q}}{j_{q}} (-1)^{1+\sum_{i\geq l} j_{i}} (1+y)^{r-d+n-(\sum_{i\geq l} ij_{i}+l)-k-1}$$

$$= [y^{n-p}] \sum_{k=0}^{p-1} \sum_{d} (1+y)^{r-d+n-l-k-1} (-1)^{d} \binom{r}{d} \binom{r-d+p-k-1}{p}$$

$$\times \Big\{ -\{1-(1+y)^{-l}\}^{n_{l}-1} \prod_{i\geq l} \{1-(1+y)^{-i}\}^{n_{l}} \Big\}.$$

Analogously, we compute

$$B_{2} := \sum_{j,k} \left\{ \sum_{d} (-1)^{d} \binom{r}{d} \binom{r-d+n-j-k-p-2}{n-p} \binom{r-d+p-k-1}{p} \right\} (-1)^{j+1} \\ \times \left\{ \sum_{j_{l}}^{n_{l}-1} \sum_{j_{l+1}}^{n_{l+1}} \cdots \sum_{j_{q}}^{n_{q}} \binom{n_{l}-1}{j_{l}} \binom{n_{l+1}}{j_{l+1}} \cdots \binom{n_{q}}{j_{q}} (-1)^{j+1+\sum_{i\geq l} j_{i}} \delta_{j+p+1,\sum_{i\geq l} ij_{i}} \right\} \\ = \sum_{k} \sum_{d} (-1)^{d} \binom{r}{d} \binom{r-d+p-k-1}{p} \left\{ \sum_{j_{l}}^{n_{l}-1} \sum_{j_{l+1}}^{n_{l+1}} \cdots \sum_{j_{q}}^{n_{q}} \binom{n_{l}-1}{j_{l}} \binom{n_{l+1}}{j_{l+1}} \cdots \binom{n_{q}}{j_{q}} \right\} \\ \times (-1)^{\sum_{i\geq l} j_{i}} \binom{r-d+n-(\sum_{i\geq l} ij_{i}-p-1)-k-p-2}{n-p} \right\} - U_{2},$$

where

$$U_{2} = \sum_{k} \sum_{d} (-1)^{d} \binom{r}{d} \binom{r-d+p-k-1}{p} \binom{r-d+n-k-1}{n-p}$$

is dealing with the case $j_i = 0$ for all i. Then, we have

$$B_2 + U_2 = [y^{n-p}] \sum_k \sum_d (1+y)^{r-d+n-k-1} (-1)^d \binom{r}{d} \binom{r-d+p-k-1}{p} \times \{1 - (1+y)^{-l}\}^{n_l-1} \prod_{i>l} \{1 - (1+y)^{-i}\}^{n_i}.$$

With a careful inspection, we realize that the quantity $\frac{p!(n-p)!}{n!}U_1$ equals the second sum for $W_{n,r}$ while $\frac{p!(n-p)!}{n!}U_2$ equals the first sum. Therefore, we now have

$$W_{n,r} = \frac{p!(n-p)!}{n!} (B_1 + U_1 + B_2 + U_2)$$

= $[y^{n-p}] \frac{p!(n-p)!}{n!} \mathcal{V}_{\mu}(y) \sum_{k=0}^{p-1} \sum_{d \ge 0} (1+y)^{r-d-k-1} (-1)^d \binom{r}{d} \binom{r-d+p-k-1}{p}.$

Finally, we compute the generating function:

$$\sum_{m\geq 1} \sum_{k=0}^{n-m} (-1)^k \frac{\binom{m+k}{m}}{(m+k)!} W_{n,m+k} x^m$$

$$= \sum_{m\geq 1} \sum_{k=0}^{n-m} (-1)^k \frac{\binom{m+k}{m}}{(m+k)!} \frac{p!(n-p)!}{n!} [y^{n-p}] \mathcal{V}_{\mu}(y) \sum_{a\geq m+k-p} \binom{p+a}{a} y^a \binom{p}{p+a+1-m-k} x^m$$

$$= \frac{p!(n-p)!}{n!} [y^{n-p}] \mathcal{V}_{\mu}(y) \sum_{i=0}^{p-1} \binom{x+i}{p} (1+y)^{x+i-p}$$

which gives the desired polynomial $P_{[p,n-p],\mu}(x)$.

Note that if n - p = p or n - p = p + 1, then the only μ satisfying $\min(\mu) \ge p + 1$ is $\mu = [n]$. For these two exceptional cases, $P_{[p,n-p],[n]}(x)$ can be equivalently viewed as the genus distribution polynomial for certain one-face bicolored maps. By comparing our formula eq. (13) with Stanley's formula [30], we find that they agree with each other. Hence, the formulas in Theorem 3.5 holds for these two cases as well.

Moreover, it is easy to show bicolored maps with face-type (equivalently black vertex degree distribution) [p, n - p] and WDD μ with min $(\mu) \ge p + 1$ are always connected. Consequently, we have the following corollary.

Corollary 3.6. For $n - p \ge p$ and $\min(\mu) \ge p + 1$, we have

$$P^{\bullet}_{[p,n-p],\mu}(x) = \frac{p!(n-p)!}{n!} [y^{n-p}] \mathcal{V}_{\mu}(y) \sum_{i=0}^{p-1} \binom{x+i}{p} (1+y)^{x+i-p}.$$
 (14)

As mentioned, the Harer-Zagier formula [20] is the genus distribution polynomial for one-face bicolored maps with WDD being regular, i.e., every white vertex is of degree exactly two. From this point of view, the following corollary provides analogues of the Harer-Zagier formula, i.e., the genus distribution polynomials for two-face bicolored maps with WDD being regular.

Corollary 3.7. For p > 0 and $\mu = [k^d]$ where k > p, we have

$$\binom{dk}{p}P^{\bullet}_{[p,dk-p],[k^d]}(x) = \sum_{i=0}^{p-1} \sum_{j=0}^d (-1)^{d-j} \binom{d}{j} \binom{x+i}{p} \binom{x+i-p+jk}{dk-p}.$$
(15)

Furthermore, thanks to the expression in Theorem 3.5, we can prove the following results.

Theorem 3.8. Every zero of the polynomial $P_{[p,n-p],\mu}(x)$ has a real part 0.

Corollary 3.9. The sequence of the non-zero coefficients of $P_{[p,n-p],\mu}(x)$ is log-concave.

The cases for other μ 's and more related results will be presented later in the full version.

References

- N. M. Adrianov, An analog of the Harer–Zagier formula for unicellular bicolored maps, Func. Anal. Appl. 31 (1997), 149–155.
- [2] E. A. Bender, E. R. Canfield, The asymptotic number of rooted maps on a surface, J. Combin. Theory Ser. A 43 (1986), 244–257.
- [3] E. A. Bender, E. R. Canfield, The number of rooted maps on an orientable surface, J. Combin. Theory Ser. B 53 (1991), 293–299.
- [4] E. A. Bender, E. R. Canfield, L. B. Richmond, The asymptotic number of rooted maps on a surface II. Enumeration by vertices and faces, J. Combin. Theory Ser. A 63 (1993), 318–329.
- [5] O. Bernardi, A. H. Morales, Bijections and symmetries for the factorizations of the long cycle, Adv. Appl. Math. 50 (2013), 702–722.
- [6] Z.-W. Bai, R. X. F. Chen, Explicit formulas for a family of hypermaps beyond the one-face case, J. Combin. Theory Ser. A 206 (2024), Article 105905.
- [7] G. Chapuy, A new combinatorial identity for unicellular maps, via a direct bijective approach, Adv. Appl. Math. 47 (2011), 874–893.
- [8] G. Chapuy, V. Féray, É. Fusy, A simple model of trees for unicellular maps, J. Combin. Theory Ser. A 120 (2013), 2064–2092.
- G. Chapuy, W. Fang, Generating functions of bipartite maps on orientable surfaces, Electron. J. Combin. 23(3) (2016), #P3.31.
- [10] R. X. F. Chen, A versatile combinatorial approach of studying products of long cycles in symmetric groups, Adv. Appl. Math. 133 (2022), 102283.
- [11] R. X. F. Chen, C. M. Reidys, Plane permutations and applications to a result of Zagier-Stanley and distances of permutations, SIAM J. Discrete Math. 30(3) (2016), 1660–1684.
- [12] R. X. F. Chen, Towards studying the structure of triple Hurwitz numbers, submitted.
- [13] R. X. F. Chen, The Harer-Zagier and Jackson formulas and new results for one-face bipartite maps, Proc. Amer. Math. Soc. 152(10) (2024), 4245–4259.
- [14] R. Cori, B. Vauquelin, Planar maps are well labeled trees, Canad. J. Math. 33(5) (1981), 1023–1042.
- [15] S. R. Carrell, G. Chapuy, Simple recurrence formulas to count maps on orientable surfaces, J. Combin. Theory Ser. A 133 (2015), 58–75.
- [16] A. Grothendieck, Esquisse d'un programme, In Geometric Galois Action, vol.1, London Math. Soc. Lecture Note Ser., vol. 242, Cambridge University Press, Cambridge, 1997, 5–48 (English translation, 243–284).

- [17] Z. Gao, The number of degree restricted maps on general surfaces, Discrete Math. 123 (1993), 47–63.
- [18] J. L. Gross, D. P. Robbins, T. W. Tucker, Genus distributions for bouquets of circles, J. Combin. Theory Ser. B 47 (1989), 292–306.
- [19] A. Goupil, G. Schaeffer, Factoring n-cycles and counting maps of given genus, European J. Combin. 19(7) (1998), 819–834.
- [20] J. Harer, D. Zagier, The Euler characteristics of the moduli space of curves, Invent. Math. 85 (1986), 457–485.
- [21] D. M. Jackson, Counting cycles in permutations by group characters, with an application to a topological problem, Trans. Amer. Math. Soc. 299(2) (1987), 785–801.
- [22] D. M. Jackson, Some combinatorial problems associated with products of conjugacy classes of the symmetric group, J. Combin. Theory Ser. A 49 (1988), 363–369.
- [23] S. K. Lando, A. K. Zvonkin, Graphs on surfaces and their applications, volume 141 of Encyclopaedia of Mathematical Sciences, Springer-Verlag, 2004. Appendix by D. B. Zagier.
- [24] M. Kazarian, P. Zograf, Virasoro constraints and topological recursion for Grothendieck's dessin counting, Lett. Math. Phys. 105 (2015), 1057–1084.
- [25] B. Louf, Simple formulas for constellations and bipartite maps with prescribed degrees, Canad. J. Math. 73(1) (2021), 160–176.
- [26] A. H. Morales, E. A. Vassilieva, Bijective enumeration of bicolored maps of given vertex degree distribution, Discrete Math. Theoret. Comput. Sci. (2009), 661–672.
- [27] G. Schaeffer, E. Vassilieva, A bijective proof of Jackson's formula for the number of factorizations of a cycle, J. Combin. Theory Ser. A 115 (2008), 903–924.
- [28] B. E. Sagan, The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions, 2nd Edition, Springer, New York, 2001.
- [29] R. P. Stanley, Enumerative Combinatorics, vol. 2, Cambridge University Press, Cambridge, 1999.
- [30] R. P. Stanley, Two enumerative results on cycles of permutations, European J. Combin. 32 (2011), 937–943.
- [31] W. T. Tutte. A census of planar maps. Canad. J. Math., 15:249–271, 1963.
- [32] T. R. S. Walsh, A. B. Lehman, Counting rooted maps by genus. I, J. Combin. Theory Ser. B 13 (1972), 192–218.
- [33] D. Zagier, On the distribution of the number of cycles of elements in symmetric groups, Nieuw Arch. Wisk. (4) 13 (1995), 489–495.
- [34] L. Zapponi, What is a Dessin d'Enfant? Notices Amer. Math. Soc. 50(7) (2003), 788–789.