

A SEGMENT OF EULER PRODUCT ASSOCIATED TO A CERTAIN DIRICHLET SERIES

RAJAT GUPTA AND ADITI SAVALIA

ABSTRACT. In the spirit of the work of Hardy-Littlewood and Lavrik, we study the Dirichlet series associated to the generalized divisor function $\sigma_\alpha(n) := \sum_{d|n} d^\alpha$. We obtain an exact identity relating the Dirichlet series $\zeta(s)\zeta(s-\alpha)$ and a segment of the Euler product attached to it. Specifically, our main theorems are valid in the critical strip.

1. INTRODUCTION

In the early 20th century, Hardy and Littlewood [5] studied the approximation of the Riemann zeta function $\zeta(s) := \sum_{n=1}^{\infty} 1/n^s$ in the region $0 \leq \operatorname{Re}(s) \leq 1$. They expressed it as a sum of two finite series with two error terms, called an approximate functional equation, namely for $0 \leq \sigma \leq 1$,

$$\zeta(s) = \sum_{n \leq x} \frac{1}{n^s} + \chi(s) \sum_{n \leq y} \frac{1}{n^{1-s}} + \mathcal{O}\left(\frac{1}{x^\sigma}\right) + \mathcal{O}\left(\frac{|t|^{\frac{1}{2}-\sigma}}{y^{1-\sigma}}\right). \quad (1.1)$$

Here, $\chi(s) := 2(2\pi)^{s-1} \sin(\pi s/2) \Gamma(1-s)$, $s = \sigma + it$ with $|t| = 2\pi xy$, and $x, y > A$ for any $A > 0$. They called this identity a “compromise” between the series expression of $\zeta(s)$ and its functional equation $\zeta(s) = \chi(s)\zeta(1-s)$. In the same paper they also derived the approximation for $\zeta^2(s)$, namely that

$$\zeta^2(s) = \sum_{n \leq x} \frac{d(n)}{n^s} + \chi^2(s) \sum_{n \leq y} \frac{d(n)}{n^{1-s}} + \mathcal{O}\left(x^{1/2-\sigma} \log t\right), \quad (1.2)$$

for $0 \leq \sigma \leq 1$, where $x, y > A$ and $xy = (t/2\pi)^2$. Note that error terms in both (1.1) and (1.2) are best possible (see [8, Chapter 4]) in the sense that for some values of x, y , and t , the corresponding error terms are of the exact order of magnitude. Later, Heath-Brown [6, Lemma 1], [7, Lemma 1], and Lavrik [9] studied the weighted approximate functional equations. For example, they derived them for any positive integer power of $\zeta(s)$ similar to what we have in (1.1) and (1.2). In [16], Simonič calculated implied constants in the error terms in (1.1) and (1.2).

Lavrik [12] made further investigations to approximate $\zeta(s)$. Interestingly, rather than truncating the series representation of $\zeta(s)$, he considered the truncation of its Euler product. Indeed, he studied a segment of the Euler product which, in turn, extends the Euler product formula

$$\zeta(s) = \prod_p \left(1 - \frac{1}{p^s}\right)^{-1} \quad (1.3)$$

into the critical strip. We record his result [12, Equation (1)] here. Throughout the paper we are fixing the notation that \mathbb{P} and \mathbb{P} denote a finite set of primes and their product, respectively.

Theorem 1.1 (Lavrik, [12, Equation (1)]). *Let $\mathbb{P} \neq \emptyset$, $\Gamma(s)$ be the gamma function and denote $(n, \bar{\mathbb{P}})$ to be the greatest common divisor of n and $\bar{\mathbb{P}}$. Then for complex numbers s and τ , where $\operatorname{Re}(s) > 0$ ¹ with $s \neq 1$ and*

2010 *Mathematics Subject Classification.* Primary 11M41; Secondary, 11M06.

Keywords and phrases. Euler product, Dirichlet series, approximate functional equation, and divisor function.

¹Here we have corrected the condition from any complex s to $\operatorname{Re}(s) > 0$ which Lavrik missed while evaluating integration of $n = 1$ term coming from (6) in [12].

$\operatorname{Re}(\tau) > 0$, we have

$$\begin{aligned} \zeta(s) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^s}\right) &= 1 + \frac{(\pi\tau)^{s/2}}{\Gamma(s/2)} \left(\frac{\tau^{-1/2}}{s-1} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \right. \\ &\quad \left. + \int_1^\infty t^{s/2-1} \left(\sum_{\substack{n=2 \\ (n, \mathbb{P})=1}}^\infty e^{-\pi\tau n^2 t} + \frac{1}{t^s} \left(\sum_{\substack{n=2 \\ (n, \mathbb{P})=1}}^\infty e^{-\pi\tau n^2/t} - \frac{1}{2} \sqrt{\frac{t}{\tau}} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \right) \right) dt \right). \end{aligned} \quad (1.4)$$

As an immediate application of (1.4), in a follow-up paper, he gave an approximation [13, Theorem 1] of $\zeta(s)$ in terms of a finite segment of the Euler product and then used it to give a new method of deriving a zero free region for $\zeta(s)$. We refer readers to [13, Section 1.2] for further discussion on the connection between the approximate functional equation and the non-trivial zeros of the Riemann zeta function. Approximate functional equations are one of the very effective tools in analytic number theory through which we can express any L -function, in a region where its Dirichlet series does not converge, by a summation of finite Dirichlet polynomials up to some error term.

Our main goal in this paper is to express a Dirichlet series associated to a generalized divisor function $\sigma_\alpha(n) := \sum_{d|n} d^\alpha$ in terms of a segment of an Euler product in the critical strip where the Dirichlet series and its Euler product are not convergent, see our main result given in Theorem 3.2. Indeed, we derive as a special case of Theorem 3.2 at $\alpha \rightarrow 0$, the following identity for $\zeta^2(s)$ in terms of a segment of an Euler product and a series involving the modified Bessel function of the second kind of order zero. We record the identity here.

Theorem 1.2. *Let $\mathbb{P} \neq \emptyset$, $\operatorname{Re}(\tau) > 0$, and $\operatorname{Re}(s) > 0$ with $s \neq 1$. Then*

$$\begin{aligned} \zeta^2(s) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^s}\right)^2 &= 1 + \frac{\pi^s \tau^{s/2}}{\Gamma^2\left(\frac{s}{2}\right)} \left[\frac{1}{(s-1)\sqrt{\tau}} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \left(-2 \sum_{n \in \mathbb{P}} \frac{\mu(n)}{n} \log n + \left(\frac{1}{s-1} + \gamma - \log(4\pi\sqrt{\tau}) \right) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \right) \right. \\ &\quad + 2 \int_1^\infty t^{s/2-1} \left(\sum_{\substack{n=2 \\ (n, \mathbb{P})=1}}^\infty d(n) K_0(2n\pi\sqrt{\tau t}) + t^{-s} \left(\sum_{\substack{n=2 \\ (n, \mathbb{P})=1}}^\infty d(n) K_0\left(2n\pi\sqrt{\frac{\tau}{t}}\right) \right. \right. \\ &\quad \left. \left. - \frac{1}{4} \sqrt{\frac{t}{\tau}} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \left(-2 \sum_{n \in \mathbb{P}} \frac{\mu(n)}{n} \log n + \left(\gamma - \log\left(4\pi\sqrt{\frac{\tau}{t}}\right) \right) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \right) \right) \right) dt \right]. \end{aligned} \quad (1.5)$$

We adapt Lavrik's approach to achieve our goal. In particular, we derive a formula of type (1.4) for $\zeta(s)\zeta(s-\alpha)$ and its special cases, one of which is given in Theorem 1.2. To structure the proof of our main theorem in this paper, we will be employing Ramanujan's renowned version of the Guinand formula [2, p. 253], which he had discovered several years before Guinand [4] himself. The formula reads as follows:

Theorem 1.3 (B.C. Berndt, Y. Lee, and J. Sohn [2, p. 253]). *Let $K_\nu(z)$ denote the modified Bessel function of the second kind of order ν , and x, y are positive numbers such that $xy = \pi^2$. For z being any complex number, we have*

$$\begin{aligned} \sqrt{x} \sum_{n=1}^\infty \sigma_{-z}(n) n^{z/2} K_{z/2}(2nx) - \sqrt{y} \sum_{n=1}^\infty \sigma_{-z}(n) n^{z/2} K_{z/2}(2ny) \\ = \frac{1}{4} \Gamma\left(\frac{z}{2}\right) \zeta(z) \left(y^{(1-z)/2} - x^{(1-z)/2} \right) + \frac{1}{4} \Gamma\left(-\frac{z}{2}\right) \zeta(-z) \left(y^{(1+z)/2} - x^{(1+z)/2} \right). \end{aligned} \quad (1.6)$$

2. PRELIMINARIES

We refer readers to G. N. Watson's classical treatise for the definition and properties of the modified Bessel function of the second kind $K_\nu(z)$, see [18, pp. 15, 78]. The following lemmas will be used in the sequel.

Lemma 2.1. [18, pp. 202] Let $K_\nu(x)$ denote the modified Bessel function of order ν . As $z \rightarrow \infty$,

$$K_\nu(z) = \left(\frac{2}{\pi z}\right)^{1/2} e^{-z} \left(1 + O\left(\frac{1}{z}\right)\right). \quad (2.1)$$

Next, the Mellin transform [15] of a function is a multiplicative analog of the Fourier transform. The Mellin transform of a locally integrable function $f(x)$ on $(0, \infty)$ is given by

$$F(s) = \int_0^\infty f(x) x^{s-1} dx, \quad (2.2)$$

when the integral converges. Let the region of convergence of the above integral (2.2) be $a < \operatorname{Re}(s) = c < b$. Then the inversion formula for Mellin transformation gives

$$f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} F(s) x^{-s} ds, \quad (a < c < b).$$

We will be using a well-known result called Parseval's formula [15, Equation (3.1.13)] noted below.

Lemma 2.2. Let $F(s)$ and $G(s)$ be the Mellin transform of f and g respectively. Then for c lying in the common strip of analyticity of F and G ,

$$\int_0^\infty f(u/x) g(x) \frac{dx}{x} = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} F(s) G(s) u^{-s} ds.$$

We conclude this section by stating three lemmas which are quite useful to prove our main theorems.

Lemma 2.3. For $c > \max\{0, -\operatorname{Re}(s), -\operatorname{Re}(s) + \operatorname{Re}(\alpha)\}$ and $\operatorname{Re}(u^2) > 0$, we have

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma\left(\frac{z+s}{2}\right) \Gamma\left(\frac{z+s-\alpha}{2}\right) u^{-z} \frac{dz}{z} = 2u^{s-\alpha/2} \int_1^\infty t^{-\alpha/4+s/2-1} K_{\alpha/2}(2u\sqrt{t}) dt.$$

We prove the above Lemma 2.3 in Section 4.

Let the Dirichlet convolution of two arithmetic functions f, g be defined as

$$f * g(n) = \sum_{d|n} f(d) g(n/d).$$

Also, we consider μ to be the well-known Möbius function defined as $\mu(1) = 1$ and

$$\mu(n) = \begin{cases} (-1)^k & \text{if } n \text{ is a product of } k \text{ distinct primes,} \\ 0 & \text{otherwise.} \end{cases}$$

Moreover, we let I be the multiplicative identity function, namely $I(1) = 1$ and $I(n) = 0$ for $n > 1$. We also remind the reader that $(\mu N^\alpha)(n) = \mu(n) n^\alpha$ and that $\sigma_\alpha^{-1}(n) = (\mu * \mu N^\alpha)(n)$ in the sense of the Dirichlet inverse (refer [1, Theorem 2.20]). We state the following lemma without its proof as it trivially follows using elementary methods of number theory.

Lemma 2.4. For $z, \alpha \in \mathbb{C}$, we have

$$\sum_{k|\mathbb{P}^2} \frac{(\mu * \mu N^\alpha)(k)}{k^z} = \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^z}\right) \left(1 - \frac{1}{p^{z-\alpha}}\right).$$

Lemma 2.5. For $\mathbb{P} \neq \emptyset$ and $(n, \mathbb{P}^2) \neq 1$,

$$\mathcal{G}_\alpha(n) := \sum_{k|(n, \mathbb{P}^2)} \sigma_\alpha\left(\frac{n}{k}\right) (\mu * \mu N^\alpha)(k) = 0. \quad (2.3)$$

Proof. This can be seen as follows. Let σ_α^{-1} denote the Dirichlet inverse of σ_α , i.e., $\sigma_\alpha * \sigma_\alpha^{-1} = I$. Let n be such that $p^3|n$ for some prime p . Then

$$\sigma_\alpha^{-1}(n) = (\mu * \mu N^\alpha)(n) = \sum_{d|n} \mu\left(\frac{n}{d}\right) \mu(d) d^\alpha = 0,$$

as at least one of the d or n/d is not square free. Thus $\sigma_\alpha^{-1}(n) = 0$ for such n . In the other case, i.e., when there is no p with $p^3|n$, we can write $n = \bar{n} \cdot t$, such that $p|\bar{n} \Rightarrow p \in \mathbb{P}$, and $(t, \mathbb{P}) = 1$, indeed, $\bar{n} = (n, \mathbb{P}^2)$. Moreover we are in the case where $(n, \mathbb{P}^2) > 1$, thus we have

$$\sum_{k|(n, \mathbb{P}^2)} \sigma_\alpha \left(\frac{n}{k} \right) \sigma_\alpha^{-1}(k) = \sigma_\alpha(t) \sum_{k|\bar{n}} \sigma_\alpha \left(\frac{\bar{n}}{k} \right) \sigma_\alpha^{-1}(k) = \sigma_\alpha(t) (\sigma_\alpha * \sigma_\alpha^{-1})(\bar{n}) = 0.$$

□

3. MAIN THEOREMS AND INTERMEDIATE RESULTS

We begin this section by recording an interesting modular type relation. Recall the notation that \mathbb{P} and $\bar{\mathbb{P}}$ denote a finite collection of primes and their product, respectively.

Theorem 3.1. *For $\mathbb{P} \neq \emptyset$, $\operatorname{Re}(t/\tau) > 0$, and $\alpha \in \mathbb{C}$, we have*

$$\begin{aligned} & \sum_{k|\bar{\mathbb{P}}^2} \frac{(\mu * \mu N^\alpha)(k)}{k^{1+\alpha/2}} \sum_{n=1}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(\frac{2n\pi}{k} \sqrt{\frac{t}{\tau}} \right) \\ &= \sqrt{\frac{\tau}{t}} \sum_{\substack{n=1 \\ (n, \bar{\mathbb{P}})=1}}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(2n\pi \sqrt{\frac{\tau}{t}} \right) - (\mathcal{F}_\alpha(t) + \mathcal{F}_{-\alpha}(t)), \end{aligned} \quad (3.1)$$

where,

$$\mathcal{F}_\alpha(t) = \frac{1}{4} \Gamma \left(\frac{\alpha}{2} \right) \zeta(\alpha) \left(\frac{1}{\pi} \sqrt{\frac{\tau}{t}} \right)^{\alpha/2} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \left(1 - \frac{1}{p^{1-\alpha}} \right). \quad (3.2)$$

Using the above modular relation we will derive our main theorem stated below.

Theorem 3.2. *For $\mathbb{P} \neq \emptyset$, $\operatorname{Re}(\tau) > 0$, $\alpha \in \mathbb{C} \setminus \{0\}$, and $\operatorname{Re}(s) > \max\{0, \operatorname{Re}(\alpha)\}$ with $s \neq 1, 1 + \alpha$, we have*

$$\begin{aligned} & \zeta(s) \zeta(s - \alpha) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^s} \right) \left(1 - \frac{1}{p^{s-\alpha}} \right) = 1 + \frac{\pi^{s-\alpha/2} \tau^{s/2}}{\Gamma \left(\frac{s}{2} \right) \Gamma \left(\frac{s-\alpha}{2} \right)} \left(\frac{\tau^{-1/2}}{s-1} \mathcal{R}_{1-\alpha} + \frac{\tau^{-(1+\alpha)/2}}{s-1-\alpha} \mathcal{R}_{1+\alpha} \right. \\ & \quad + 2\tau^{-\alpha/4} \int_1^\infty t^{-\alpha/4+s/2-1} \left(\sum_{\substack{n=2 \\ (n, \bar{\mathbb{P}})=1}}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(2n\pi \sqrt{\tau t} \right) \right. \\ & \quad \left. \left. + \frac{t^{\alpha/2}}{t^s} \left(\sum_{\substack{n=2 \\ (n, \bar{\mathbb{P}})=1}}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(2n\pi \sqrt{\frac{\tau}{t}} \right) - \sqrt{\frac{t}{\tau}} (\mathcal{F}_\alpha(t) + \mathcal{F}_{-\alpha}(t)) \right) \right) dt \right), \end{aligned} \quad (3.3)$$

where

$$\mathcal{R}_\alpha = \pi^{-\alpha/2} \Gamma \left(\frac{\alpha}{2} \right) \zeta(\alpha) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \left(1 - \frac{1}{p^\alpha} \right),$$

and $\mathcal{F}_\alpha(t)$ is defined in (3.2).

It can be easily seen from Lemma 2.1 that both the series in the integrand on the right-hand side are absolutely convergent. Also the absolute convergence of the integral on the right-hand side above can be justified using the modular relation given in Theorem 3.1. Indeed, with the help of (3.1), the integrand can be written as multiplication of the factors $t^{-\alpha/4+s/2-1}$ and

$$\sum_{\substack{n=2 \\ (n, \bar{\mathbb{P}})=1}}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(2n\pi \sqrt{\tau t} \right) + \frac{t^{\alpha/2-s+1/2}}{\sqrt{\tau}} \sum_{k|\bar{\mathbb{P}}^2} \frac{(\mu * \mu N^\alpha)(k)}{k^{1+\alpha/2}} \sum_{n=1}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(\frac{2n\pi}{k} \sqrt{\frac{t}{\tau}} \right).$$

Therefore, again using Lemma 2.1, we can show that the integrand has an exponential decay.

We will show that Theorem 1.2 is a limiting case $\alpha \rightarrow 0$ of Theorem 3.2. Further restricting ourselves to the region $\operatorname{Re}(s) > 1$, Theorem 1.2 reduces to:

Corollary 3.3. For $\mathbb{P} \neq \emptyset$, $\operatorname{Re}(s) > 1$, and $\operatorname{Re}(\tau) > 0$, we obtain

$$\begin{aligned} \zeta^2(s) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^s}\right)^2 &= 1 + \frac{2\pi^s \tau^{s/2}}{\Gamma^2\left(\frac{s}{2}\right)} \int_1^\infty t^{s/2-1} \left(\sum_{\substack{n=2 \\ (n, \mathbb{P})=1}}^\infty d(n) K_0\left(2n\pi\sqrt{\tau t}\right) \right. \\ &\quad \left. + t^{-s} \sum_{\substack{n=2 \\ (n, \mathbb{P})=1}}^\infty d(n) K_0\left(2n\pi\sqrt{\frac{\tau}{t}}\right) \right) dt. \end{aligned} \quad (3.4)$$

Remark 3.4. Using [3, Formula 6.561.16, p. 676], we can further simplify the integration in (3.4) and derive the following elementary identity for $\operatorname{Re}(s) > 1$,

$$\sum_{\substack{n=1 \\ (n, \mathbb{P})=1}}^\infty \frac{d(n)}{n^s} = \zeta^2(s) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^s}\right)^2. \quad (3.5)$$

If we let \mathbb{P} be the collection of all primes in the above corollary, then we recover the well-known Euler identity, namely, we get for $\operatorname{Re}(s) > 1$,

$$\zeta^2(s) \prod_p \left(1 - \frac{1}{p^s}\right)^2 = 1.$$

4. PROOFS OF LEMMAS, MAIN THEOREM AND THEIR COROLLARIES

4.1. Proof of Lemma 2.3.

Proof of Lemma 2.3. For $\operatorname{Re}(z) > 0$, we have [14, p. 347, Equation (1)]

$$\int_0^\infty x^{z-1} e^{-x} dx = \Gamma(z). \quad (4.1)$$

Replacing z with $\frac{z+s}{2}$, and then applying change of variable $x \rightarrow x^2$ in (4.1), we get

$$2 \int_0^\infty x^{s+z-1} e^{-x^2} dx = \Gamma\left(\frac{z+s}{2}\right), \quad \operatorname{Re}(z) > -\operatorname{Re}(s). \quad (4.2)$$

Also, for $\operatorname{Re}(\lambda + \beta) > 0$, $\operatorname{Re}(\mu) > 0$, and $\operatorname{Re}(\mu + \nu) > 0$, we have [3, p. 657, Formula 6.455(1)]

$$\int_0^\infty x^{\mu-1} e^{-\beta x} \Gamma(\nu, \lambda x) dx = \frac{\lambda^\nu \Gamma(\mu + \nu)}{\mu(\lambda + \beta)^{\mu+\nu}} {}_2F_1\left(1, \mu + \nu; \mu + 1; \frac{\beta}{\lambda + \beta}\right). \quad (4.3)$$

Letting $\beta \rightarrow 0$, $x \rightarrow x^2$, $\mu = z/2$, $\lambda = 1$, and $\nu = \frac{s-\alpha}{2}$ in (4.3), we obtain for $\operatorname{Re}(z) > \max\{0, -\operatorname{Re}(s) + \operatorname{Re}(\alpha)\}$

$$\int_0^\infty x^{z-1} \Gamma\left(\frac{s-\alpha}{2}, x^2\right) dx = \frac{1}{z} \Gamma\left(\frac{z+s-\alpha}{2}\right), \quad (4.4)$$

where $\Gamma(a, x)$ is the incomplete gamma function defined as

$$\Gamma(a, x) := \int_x^\infty t^{a-1} e^{-t} dt.$$

Employing Parseval's formula stated in Lemma 2.2, for c in the intersection of $\operatorname{Re}(z) > 0$, $\operatorname{Re}(z) > -\operatorname{Re}(s)$ and $\operatorname{Re}(z) > -\operatorname{Re}(s) + \operatorname{Re}(\alpha)$, the identities (4.2) and (4.4) gives

$$\begin{aligned} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma\left(\frac{z+s}{2}\right) \Gamma\left(\frac{z+s-\alpha}{2}\right) \frac{u^{-z}}{z} dz &= 2 \int_0^\infty \Gamma\left(\frac{s-\alpha}{2}, \frac{u^2}{x^2}\right) e^{-x^2} x^s \frac{dx}{x} \\ &= 2 \int_0^\infty \left(\int_{u^2/x^2}^\infty t^{\frac{s-\alpha}{2}-1} e^{-t} dt \right) e^{-x^2} x^{s-1} dx. \end{aligned}$$

We make the change of variable $t \rightarrow \frac{u^2}{x^2}t$, to obtain

$$\begin{aligned} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma\left(\frac{z+s}{2}\right) \Gamma\left(\frac{z+s-\alpha}{2}\right) \frac{u^{-z}}{z} dz &= 2 \int_0^\infty \left(\int_1^\infty \left(\frac{u^2}{x^2}t\right)^{\frac{s-\alpha}{2}-1} e^{-u^2 t/x^2} dt \right) \frac{u^2}{x^2} e^{-x^2} x^{s-1} dx \\ &= 2u^{s-\alpha} \int_0^\infty \left(\int_1^\infty t^{\frac{s-\alpha}{2}-1} e^{-u^2 t/x^2} dt \right) e^{-x^2} x^{\alpha-1} dx \\ &= 2u^{s-\alpha} \int_1^\infty \left(\int_0^\infty e^{-u^2 t/x^2 - x^2} x^{\alpha-1} dx \right) t^{\frac{s-\alpha}{2}-1} dt, \end{aligned} \quad (4.5)$$

where in the last step, change of order of integration can be justified by invoking Watson's lemma [17, Theorem 2.4]. We have from [3, p. 368, 3.471, Formula (9)],

$$\int_0^\infty x^{\nu-1} e^{-\gamma x - \beta/x} dx = 2 \left(\frac{\beta}{\gamma}\right)^{\nu/2} K_\nu(2\gamma\sqrt{\beta}),$$

for $\text{Re}(\beta) > 0$ and $\text{Re}(\gamma) > 0$. We now use the above integral with change of variable $x \rightarrow x^2$ followed by taking $\beta = u^2 t$, $\gamma = 1$, and $\nu = \frac{\alpha}{2}$ in (4.5) to derive our lemma, namely, for $\text{Re}(z) > \max\{0, -\text{Re}(s), -\text{Re}(s) + \text{Re}(\alpha)\}$ and $\text{Re}(u^2) > 0$,

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma\left(\frac{z+s}{2}\right) \Gamma\left(\frac{z+s-\alpha}{2}\right) \frac{u^{-z}}{z} dz = 2u^{s-\alpha/2} \int_1^\infty t^{-\alpha/4+s/2-1} K_{\alpha/2}(2u\sqrt{t}) dt.$$

□

4.2. Proofs of Theorems 3.1 and 3.2.

Proof of Theorem 3.1. In Ramanujan-Guinand's formula take $z = \alpha$, $x = \frac{\pi}{k\sqrt{(\tau/t)}}$, and $y = \frac{\pi^2}{x} = \pi k\sqrt{(\tau/t)}$, and use the relation $\sigma_{-\alpha}(n) = n^{-\alpha}\sigma_\alpha(n)$, to get

$$\begin{aligned} \left(\frac{\pi}{k}\sqrt{\frac{t}{\tau}}\right)^{1/2} \sum_{n=1}^\infty \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2}\left(\frac{2n\pi}{k}\sqrt{\frac{t}{\tau}}\right) - \left(\pi k\sqrt{\frac{\tau}{t}}\right)^{1/2} \sum_{n=1}^\infty \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2}\left(2\pi n k\sqrt{\frac{\tau}{t}}\right) \\ = \frac{1}{4}\Gamma\left(\frac{\alpha}{2}\right)\zeta(\alpha) \left(\left(\pi k\sqrt{\frac{\tau}{t}}\right)^{(1-\alpha)/2} - \left(\frac{\pi}{k}\sqrt{\frac{t}{\tau}}\right)^{(1-\alpha)/2} \right) \\ + \frac{1}{4}\Gamma\left(-\frac{\alpha}{2}\right)\zeta(-\alpha) \left(\left(\pi k\sqrt{\frac{\tau}{t}}\right)^{(1+\alpha)/2} - \left(\frac{\pi}{k}\sqrt{\frac{t}{\tau}}\right)^{(1+\alpha)/2} \right). \end{aligned}$$

After considerable simplification we obtain

$$\sum_{n=1}^\infty \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2}\left(\frac{2n\pi}{k}\sqrt{\frac{t}{\tau}}\right) = k\sqrt{\frac{\tau}{t}} \sum_{n=1}^\infty \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2}\left(2\pi n k\sqrt{\frac{\tau}{t}}\right) - A_\alpha - A_{-\alpha}, \quad (4.6)$$

where

$$A_\alpha = A_\alpha(t) := -\frac{1}{4}\Gamma\left(\frac{\alpha}{2}\right)\zeta(\alpha) \left(k\sqrt{\frac{\tau}{t}} \left(\pi k\sqrt{\frac{\tau}{t}}\right)^{-\alpha/2} - \left(\frac{k}{\pi}\sqrt{\frac{\tau}{t}}\right)^{\alpha/2} \right).$$

We multiply both sides by $\frac{(\mu * \mu N^\alpha)(k)}{k^{1+\alpha/2}}$ and then take the sum over divisors of \mathbb{P}^2 . Thus, (4.6) changes to

$$\sum_{k|\mathbb{P}^2} \frac{(\mu * \mu N^\alpha)(k)}{k^{1+\alpha/2}} \sum_{n=1}^\infty \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2}\left(\frac{2n\pi}{k}\sqrt{\frac{t}{\tau}}\right) = \sqrt{\frac{\tau}{t}} \sum_{n=1}^\infty \frac{K_{\alpha/2}(2n\pi\sqrt{\frac{\tau}{t}})}{n^{\alpha/2}} \sum_{k|(\mathbb{P}^2, n)} (\mu * \mu N^\alpha)(k) \sigma_\alpha\left(\frac{n}{k}\right)$$

$$-\sum_{k|\mathbb{P}^2} \frac{(\mu * \mu N^\alpha)(k)}{k^{1+\alpha/2}} (A_\alpha + A_{-\alpha}). \quad (4.7)$$

The first sum on the right-hand side of the above equation can be split into two sums, namely,

$$\sum_{\substack{n=1 \\ (n, \mathbb{P}^2)=1}}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(2n\pi \sqrt{\frac{\tau}{t}} \right) + \sum_{\substack{n=1 \\ (n, \mathbb{P}^2) \neq 1}}^{\infty} \frac{\mathcal{G}_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(2n\pi \sqrt{\frac{\tau}{t}} \right), \quad (4.8)$$

where \mathcal{G}_α is defined in (2.3). Surprisingly, by employing Lemma 2.5, we see that the second sum in (4.8) amounts to zero.

Also note that

$$\begin{aligned} \sum_{k|\mathbb{P}^2} \frac{(\mu * \mu N^\alpha)(k)}{k^{1+\alpha/2}} A_\alpha &= -\frac{1}{4} \Gamma\left(\frac{\alpha}{2}\right) \zeta(\alpha) \left(\sqrt{\frac{\tau}{t}} \sum_{k|\mathbb{P}^2} \frac{(\mu * \mu N^\alpha)(k)}{k^\alpha} \left(\pi \sqrt{\frac{\tau}{t}} \right)^{-\alpha/2} \right. \\ &\quad \left. - \sum_{k|\mathbb{P}^2} \frac{(\mu * \mu N^\alpha)(k)}{k} \left(\frac{1}{\pi} \sqrt{\frac{\tau}{t}} \right)^{\alpha/2} \right) \\ &= -\frac{1}{4} \Gamma\left(\frac{\alpha}{2}\right) \zeta(\alpha) \left(\pi^{-\alpha/2} \left(\sqrt{\frac{\tau}{t}} \right)^{1-\alpha/2} \sum_{n|\mathbb{P}} \frac{\mu(n)}{n^\alpha} \sum_{m|\mathbb{P}} \mu(m) \right. \\ &\quad \left. - \left(\frac{1}{\pi} \sqrt{\frac{\tau}{t}} \right)^{\alpha/2} \sum_{n|\mathbb{P}} \frac{\mu(n)}{n} \sum_{m|\mathbb{P}} \frac{\mu(m)m^\alpha}{m} \right) \\ &= \frac{1}{4} \Gamma\left(\frac{\alpha}{2}\right) \zeta(\alpha) \left(\frac{1}{\pi} \sqrt{\frac{\tau}{t}} \right)^{\alpha/2} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \left(1 - \frac{1}{p^{1-\alpha}} \right) = \mathcal{F}(\alpha), \end{aligned} \quad (4.9)$$

where in the last step we use that for $\mathbb{P} \neq \emptyset$,

$$\sum_{m|\mathbb{P}} \mu(m) = 0, \quad \text{and} \quad \sum_{n|\mathbb{P}} \frac{\mu(n)}{n} \sum_{m|\mathbb{P}} \frac{\mu(m)m^\alpha}{m} = \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \left(1 - \frac{1}{p^{1-\alpha}} \right).$$

Similarly we obtain,

$$\sum_{k|\mathbb{P}^2} \frac{(\mu * \mu N^\alpha)(k)}{k^{1+\alpha/2}} A_{-\alpha} = \frac{1}{4} \Gamma\left(-\frac{\alpha}{2}\right) \zeta(-\alpha) \left(\frac{1}{\pi} \sqrt{\frac{\tau}{t}} \right)^{-\alpha/2} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \left(1 - \frac{1}{p^{1+\alpha}} \right) = \mathcal{F}(-\alpha) \quad (4.10)$$

using the facts that for $\mathbb{P} \neq \emptyset$,

$$\sum_{m|\mathbb{P}} \mu(m) = 0, \quad \text{and} \quad \sum_{n|\mathbb{P}} \frac{\mu(n)}{n^{1+\alpha}} \sum_{m|\mathbb{P}} \frac{\mu(m)}{m} = \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \left(1 - \frac{1}{p^{1+\alpha}} \right).$$

Hence, by plugging (4.8), (4.9), and (4.10) into (4.7), we derive our result. \square

Proof of Theorem 3.2. For $s = \sigma + it$, and $\alpha = \sigma_0 + it_0$, let us define $F(z)$ as

$$F(z) := \pi^{-z+\alpha/2} \Gamma\left(\frac{z}{2}\right) \Gamma\left(\frac{z-\alpha}{2}\right) \zeta(z) \zeta(z-\alpha) \frac{\tau^{(s-z)/2}}{z-s} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \left(1 - \frac{1}{p^z} \right).$$

Let \mathcal{C} be the contour determined by the line segments $[\sigma_1 - iT, \sigma_1 + iT]$, $[\sigma_1 + iT, \sigma_2 + iT]$, $[\sigma_2 + iT, \sigma_2 - iT]$, $[\sigma_2 - iT, \sigma_1 - iT]$, where $\sigma_1 < \min\{0, \sigma, \sigma_0\}$, $\sigma_2 > \max\{1, \sigma, 1 + \sigma_0\}$ and $T > |t| + |t_0|$.

Note that poles of $\Gamma\left(\frac{z}{2}\right)$ and $\Gamma\left(\frac{z-\alpha}{2}\right)$ at 0 and α are canceled by the zeros of the product over finite collection of primes, and the poles at $-2m$ and $\alpha - 2m$ for $m \in \mathbb{N}$ are canceled by the trivial zeros of $\zeta(z)$ and $\zeta(z-\alpha)$. For $\text{Re}(\tau) > 0$, the factor $\frac{\tau^{(s-1)/2}}{z-s}$ also contributes a simple pole at $z = s$. Thus we have poles of

$F(z)$ only at $z = 1, 1 + \alpha$ and s . The choice of σ_1, σ_2 , and T is such that all poles lie inside the contour. Let R_a denote residue of $F(z)$ at the pole $z = a$. Then we have,

(1) at $z = 1$, $F(z)$ has a simple pole with residue

$$R_1 = \lim_{z \rightarrow 1} (z-1)F(z) = \pi^{(\alpha-1)/2} \Gamma\left(\frac{1-\alpha}{2}\right) \zeta(1-\alpha) \frac{\tau^{(s-1)/2}}{1-s} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \left(1 - \frac{1}{p^{1-\alpha}}\right),$$

(2) at $z = 1 + \alpha$, $F(z)$ has a simple pole with residue

$$R_{1+\alpha} = \lim_{z \rightarrow 1+\alpha} (z-1-\alpha)F(z) = \pi^{-(1+\alpha)/2} \Gamma\left(\frac{1+\alpha}{2}\right) \zeta(1+\alpha) \frac{\tau^{(s-1-\alpha)/2}}{1+\alpha-s} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \left(1 - \frac{1}{p^{1+\alpha}}\right),$$

(3) at $z = s$, $F(z)$ has a simple pole with residue

$$R_s = \lim_{z \rightarrow s} (z-s)F(z) = \pi^{-s+\alpha/2} \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s-\alpha}{2}\right) \zeta(s) \zeta(s-\alpha) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^s}\right) \left(1 - \frac{1}{p^{s-\alpha}}\right).$$

Now applying Cauchy's residue theorem for $F(z)$ in the contour \mathcal{C} , we have

$$\frac{1}{2\pi i} \left(\int_{\sigma_2-iT}^{\sigma_2+iT} + \int_{\sigma_2+iT}^{\sigma_1+iT} + \int_{\sigma_1+iT}^{\sigma_1-iT} + \int_{\sigma_1-iT}^{\sigma_2-iT} \right) F(z) dz = R_1 + R_{1+\alpha} + R_s. \quad (4.11)$$

Using Stirling bound for $\Gamma\left(\frac{z}{2}\right)$, and $\Gamma\left(\frac{z-\alpha}{2}\right)$ with $|\arg(\tau)| < \pi/2$, which states that for $z' = \sigma' + it'$ with $a \leq \sigma' \leq b$ and $|t'| > 1$,

$$|\Gamma(z')| \ll |t'|^{\sigma' - \frac{1}{2}} e^{-\frac{\pi}{2}|t'|}, \quad (4.12)$$

and property of the Riemann Zeta function that $\zeta(z') \ll |t'|^{\frac{3}{2}+\delta}$ for $\sigma' > -\delta$, the integrals along the horizontal lines $[\sigma_1 + iT, \sigma_2 + iT]$ and $[\sigma_1 - iT, \sigma_2 - iT]$ tend to zero as $T \rightarrow \infty$. Hence, letting $T \rightarrow \infty$ in (4.11), we get

$$\frac{1}{2\pi i} \left(\int_{\sigma_2-i\infty}^{\sigma_2+i\infty} + \int_{\sigma_1+i\infty}^{\sigma_1-i\infty} \right) F(z) dz = R_1 + R_{1+\alpha} + R_s. \quad (4.13)$$

We first evaluate

$$f_1 := \frac{1}{2\pi i} \int_{\sigma_2-i\infty}^{\sigma_2+i\infty} \pi^{-z+\alpha/2} \Gamma\left(\frac{z}{2}\right) \Gamma\left(\frac{z-\alpha}{2}\right) \zeta(z) \zeta(z-\alpha) \frac{\tau^{(s-z)/2}}{z-s} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^z}\right) \left(1 - \frac{1}{p^{z-\alpha}}\right) dz. \quad (4.14)$$

By our choice of σ_2 , $\zeta(z)$ and $\zeta(z-\alpha)$ have the Euler product representations on the line segment $[\sigma_2 + iT, \sigma_2 - iT]$. Thus the factor

$$Z(z, \alpha) := \zeta(z) \zeta(z-\alpha) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^z}\right) \left(1 - \frac{1}{p^{z-\alpha}}\right), \quad (4.15)$$

for $\text{Re}(z) = \sigma_2$, reduces to

$$Z(z, \alpha) = \sum_{\substack{n=1 \\ (n, \mathbb{P})=1}}^{\infty} \frac{\sigma_\alpha(n)}{n^z}. \quad (4.16)$$

Upon using (4.16) in (4.14), and then changing the order of summation and integration we obtain,

$$f_1 = \frac{1}{2\pi i} \sum_{\substack{n=1 \\ (n, \mathbb{P})=1}}^{\infty} \frac{\sigma_\alpha(n)}{n^z} \int_{\sigma_2-i\infty}^{\sigma_2+i\infty} \pi^{-z+\alpha/2} \Gamma\left(\frac{z}{2}\right) \Gamma\left(\frac{z-\alpha}{2}\right) \frac{\tau^{(s-z)/2}}{z-s} dz.$$

By change of variable z by $z + s$, we derive,

$$f_1 = \pi^{-s+\alpha/2} \sum_{(n, \mathbb{P})=1} \frac{\sigma_\alpha(n)}{n^s} \left(\frac{1}{2\pi i} \int_{a-i\infty}^{a+i\infty} \Gamma\left(\frac{z+s}{2}\right) \Gamma\left(\frac{z+s-\alpha}{2}\right) (n\pi\sqrt{\tau})^{-z} \frac{dz}{z} \right), \quad (4.17)$$

for $a = \sigma_2 - \sigma > 0$. It is easy to verify that for $\operatorname{Re}(z) = a$, $\sigma_2 > \max\{1, \sigma, 1 + \sigma_0\}$ and $\operatorname{Re}(\tau) > 0$, all the conditions are satisfied to employ Lemma 2.3 in (4.17) with $u = n\pi\sqrt{\tau}$. This gives

$$f_1 = 2\tau^{s/2-\alpha/4} \sum_{\substack{n=1 \\ (n, \mathbb{P})=1}}^{\infty} \frac{\sigma_{\alpha}(n)}{n^{\alpha/2}} \int_1^{\infty} t^{-\alpha/4+s/2-1} K_{\alpha/2} \left(2n\pi\sqrt{\tau t} \right) dt.$$

The asymptotic bound on $K_{\nu}(z)$ from Lemma 2.1 allows us to interchange the order of summation and integration, and derive

$$f_1 = 2\tau^{s/2-\alpha/4} \int_1^{\infty} t^{-\alpha/4+s/2-1} \sum_{\substack{n=1 \\ (n, \mathbb{P})=1}}^{\infty} \frac{\sigma_{\alpha}(n)}{n^{\alpha/2}} K_{\alpha/2} \left(2n\pi\sqrt{\tau t} \right) dt. \quad (4.18)$$

Now we turn our attention towards,

$$f_2 := \frac{1}{2\pi i} \int_{\sigma_1+i\infty}^{\sigma_1-i\infty} \pi^{-z+\alpha/2} \Gamma\left(\frac{z}{2}\right) \Gamma\left(\frac{z-\alpha}{2}\right) \zeta(z) \zeta(z-\alpha) \frac{\tau^{(s-z)/2}}{z-s} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^z}\right) \left(1 - \frac{1}{p^{z-\alpha}}\right) dz. \quad (4.19)$$

Note that the functional equation for $\zeta(z)\zeta(z-\alpha)$ in variable z can be easily obtained by multiplying the functional equations for $\zeta(z)$ and $\zeta(z-\alpha)$ as

$$\pi^{-z+\alpha/2} \Gamma\left(\frac{z}{2}\right) \Gamma\left(\frac{z-\alpha}{2}\right) \zeta(z) \zeta(z-\alpha) = \pi^{-(1-z)-\alpha/2} \Gamma\left(\frac{1-z}{2}\right) \Gamma\left(\frac{1-z+\alpha}{2}\right) \zeta(1-z) \zeta(1-z+\alpha).$$

Employing the above functional relation in (4.19), we find that

$$f_2 = \frac{\pi^{-1-\alpha/2}}{2\pi i} \int_{\sigma_1+i\infty}^{\sigma_1-i\infty} \pi^z \Gamma\left(\frac{1-z}{2}\right) \Gamma\left(\frac{1-z+\alpha}{2}\right) \times \\ \zeta(1-z) \zeta(1-z+\alpha) \frac{\tau^{(s-z)/2}}{z-s} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^z}\right) \left(1 - \frac{1}{p^{z-\alpha}}\right) dz. \quad (4.20)$$

Our choice of σ_1 helps us to change the factor $\zeta(1-z)\zeta(1-z+\alpha)$ into a Dirichlet series,

$$\zeta(1-z) \zeta(1-z+\alpha) = \sum_{n=1}^{\infty} \frac{\sigma_{\alpha}(n)}{n^{1-z+\alpha}}. \quad (4.21)$$

Thus using (4.21), and then interchanging the order of summation and integration in (4.20), we get

$$f_2 = \frac{\pi^{-1-\alpha/2}}{2\pi i} \sum_{n=1}^{\infty} \frac{\sigma_{\alpha}(n)}{n^{\alpha+1}} \times \\ \int_{\sigma_1+i\infty}^{\sigma_1-i\infty} \pi^z \Gamma\left(\frac{1-z}{2}\right) \Gamma\left(\frac{1-z+\alpha}{2}\right) \frac{\tau^{(s-z)/2} n^z}{z-s} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^z}\right) \left(1 - \frac{1}{p^{z-\alpha}}\right) dz.$$

By applying the change of variable z by $s-z$, one obtains

$$f_2 = \frac{\pi^{s-1-\alpha/2}}{2\pi i} \sum_{n=1}^{\infty} \frac{\sigma_{\alpha}(n)}{n^{\alpha+1-s}} \int_{\sigma-\sigma_1-i\infty}^{\sigma-\sigma_1+i\infty} \Gamma\left(\frac{1+z-s}{2}\right) \Gamma\left(\frac{1+z-s+\alpha}{2}\right) \\ \times \left(\frac{\sqrt{\tau}}{n\pi}\right)^z \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^{s-z}}\right) \left(1 - \frac{1}{p^{s-z-\alpha}}\right) \frac{dz}{z}.$$

Invoking Lemma 2.4, with z replaced by $s-z$,

$$f_2 = \frac{\pi^{s-1-\alpha/2}}{2\pi i} \sum_{k|\mathbb{P}^2} \frac{(\mu * \mu N^{\alpha})(k)}{k^s} \sum_{n=1}^{\infty} \frac{\sigma_{\alpha}(n)}{n^{\alpha+1-s}} \times \\ \int_{\sigma-\sigma_1-i\infty}^{\sigma-\sigma_1+i\infty} \Gamma\left(\frac{1+z-s}{2}\right) \Gamma\left(\frac{1+z-s+\alpha}{2}\right) \left(\frac{n\pi}{\sqrt{\tau}k}\right)^{-z} \frac{dz}{z}. \quad (4.22)$$

Now we want to apply Lemma 2.3, with $u = \frac{n\pi}{k\sqrt{\tau}}$, $s \rightarrow 1-s$ and $\alpha \rightarrow -\alpha$. See that for $\operatorname{Re}(z) = \sigma - \sigma_1$, all the required conditions to employ Lemma 2.3 in (4.22) are satisfied. Thus we derive, for $\operatorname{Re}(\tau) > 0$,

$$f_2 = \frac{2}{\tau^{\alpha/4+(1-s)/2}} \sum_{k|\mathbb{P}^2} \frac{(\mu * \mu N^\alpha)(k)}{k^{1+\alpha/2}} \sum_{n=1}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} \int_1^\infty t^{\alpha/4-(s+1)/2} K_{\alpha/2} \left(\frac{2n\pi}{k} \sqrt{\frac{t}{\tau}} \right) dt. \quad (4.23)$$

The asymptotic bound on $K_\nu(z)$ from Lemma 2.1 will help us to interchange the summation and integration in (4.23), to obtain,

$$f_2 = 2\tau^{-\alpha/4-(1-s)/2} \int_1^\infty t^{\alpha/4-(s+1)/2} \sum_{k|\mathbb{P}^2} \frac{(\mu * \mu N^\alpha)(k)}{k^{1+\alpha/2}} \sum_{n=1}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(\frac{2n\pi}{k} \sqrt{\frac{t}{\tau}} \right) dt. \quad (4.24)$$

Invoking Theorem 3.1 in the above equation, we have

$$f_2 = 2\tau^{-\alpha/4-(1-s)/2} \int_1^\infty t^{\alpha/4-(s+1)/2} \left(\sqrt{\frac{\tau}{t}} \sum_{\substack{n=1 \\ (n, \mathbb{P}^2)=1}}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(2n\pi \sqrt{\frac{\tau}{t}} \right) - (\mathcal{F}(\alpha) + \mathcal{F}(-\alpha)) \right) dt, \quad (4.25)$$

where $\mathcal{F}(\alpha)$ is defined in (3.2).

Upon using (4.13), and (4.18), in (4.25), and noting that $(n, \mathbb{P}^2) = 1$ if and only if $(n, \mathbb{P}) = 1$, we derive

$$\begin{aligned} & 2\tau^{s/2-\alpha/4} \int_1^\infty t^{-\alpha/4+s/2-1} \left(\sum_{\substack{n=1 \\ (n, \mathbb{P})=1}}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(2n\pi \sqrt{\tau t} \right) \right. \\ & \quad \left. + \frac{t^{\alpha/2}}{t^s} \left(\sum_{\substack{n=1 \\ (n, \mathbb{P})=1}}^{\infty} \frac{\sigma_\alpha(n)}{n^{\alpha/2}} K_{\alpha/2} \left(2n\pi \sqrt{\frac{\tau}{t}} \right) - \sqrt{\frac{t}{\tau}} (\mathcal{F}_\alpha(t) + \mathcal{F}_{-\alpha}(t)) \right) \right) dt \\ & = \pi^{(\alpha-1)/2} \Gamma \left(\frac{1-\alpha}{2} \right) \zeta(1-\alpha) \frac{\tau^{(s-1)/2}}{1-s} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \left(1 - \frac{1}{p^{1-\alpha}} \right) \\ & \quad + \pi^{-(1+\alpha)/2} \Gamma \left(\frac{1+\alpha}{2} \right) \zeta(1+\alpha) \frac{\tau^{(s-1-\alpha)/2}}{1+\alpha-s} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \left(1 - \frac{1}{p^{1+\alpha}} \right) \\ & \quad + \pi^{-s+\alpha/2} \Gamma \left(\frac{s}{2} \right) \Gamma \left(\frac{s-\alpha}{2} \right) \zeta(s) \zeta(s-\alpha) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^s} \right) \left(1 - \frac{1}{p^{s-\alpha}} \right). \end{aligned} \quad (4.26)$$

From the left-hand side of the above equation (4.26), we will evaluate the integration of the $n = 1$ terms of both the series separately as:

$$\begin{aligned} & \int_1^\infty t^{-\alpha/4+s/2-1} K_{\alpha/2} \left(2\pi \sqrt{\tau t} \right) dt + \int_1^\infty t^{\alpha/4-s/2-1} K_{\alpha/2} \left(2\pi \sqrt{\frac{\tau}{t}} \right) dt \\ & = \int_0^\infty t^{-\alpha/4+s/2-1} K_{\alpha/2} \left(2\pi \sqrt{\tau t} \right) dt \\ & = \frac{1}{2} (\pi \sqrt{\tau})^{-s+\alpha/2} \Gamma \left(\frac{s}{2} \right) \Gamma \left(\frac{s-\alpha}{2} \right), \end{aligned} \quad (4.27)$$

provided $\operatorname{Re}(s) > \max\{0, \operatorname{Re}(\alpha)\}$ and $\operatorname{Re}(\tau) > 0$. In the last step we have used [3, Formula 6.561.16, p. 676], namely for $\operatorname{Re}(\nu + 1 \pm \mu) > 0$ and $\operatorname{Re}(a) > 0$,

$$\int_0^\infty x^\mu K_\nu(ax) dx = \frac{2^{\mu-1}}{a^{\mu+1}} \Gamma \left(\frac{1+\mu+\nu}{2} \right) \Gamma \left(\frac{1+\mu-\nu}{2} \right).$$

Now, dividing (4.26) by the factor $\pi^{-s+\alpha/2}\Gamma\left(\frac{s}{2}\right)\Gamma\left(\frac{s-\alpha}{2}\right)$ and simplifying the terms corresponding to $n = 1$ by employing (4.27), we derive (3.3) for $\text{Re}(s) > \max\{0, \text{Re}(\alpha)\}$. $s \neq 1, 1 + \alpha$, and $\text{Re}(\tau) > 0$. This completes the proof. \square

Next, we will provide proof of Theorem 1.2.

4.3. Proofs of Theorem 1.2 and Corollary 3.3.

Proof of Theorem 1.2. Letting $\alpha \rightarrow 0$ in (3.3), we get

$$\begin{aligned} \zeta^2(s) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^s}\right)^2 &= 1 + \frac{\pi^s \tau^{s/2}}{\Gamma^2\left(\frac{s}{2}\right)} \left[\lim_{\alpha \rightarrow 0} \left(\frac{\tau^{-1/2}}{s-1} \mathcal{R}_{1-\alpha} + \frac{\tau^{-(1+\alpha)/2}}{s-1-\alpha} \right) \right. \\ &\quad + 2 \int_1^\infty t^{s/2-1} \left\{ \sum_{\substack{n=2 \\ (n, \mathbb{P})=1}}^\infty d(n) K_0(2n\pi\sqrt{\tau t}) \right. \\ &\quad \left. \left. + t^{-s} \left(\sum_{\substack{n=2 \\ (n, \mathbb{P})=1}}^\infty d(n) K_0\left(2n\pi\sqrt{\frac{\tau}{t}}\right) - \sqrt{\frac{t}{\tau}} \lim_{\alpha \rightarrow 0} (\mathcal{F}_\alpha(t) + \mathcal{F}_{-\alpha}(t)) \right) \right\} dt \right]. \end{aligned} \quad (4.28)$$

We first compute the limit,

$$\lim_{\alpha \rightarrow 0} \left(\frac{\tau^{-1/2}}{s-1} \mathcal{R}_{1-\alpha} + \frac{\tau^{-(1+\alpha)/2}}{s-1-\alpha} \mathcal{R}_{1+\alpha} \right).$$

To do so, we expand each factors involving α around $\alpha = 0$ to get,

$$\begin{aligned} \mathcal{R}_{1-\alpha} &= \pi^{-(1-\alpha)/2} \Gamma\left(\frac{1-\alpha}{2}\right) \zeta(1-\alpha) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \left(1 - \frac{1}{p^{1-\alpha}}\right) \\ &= \left(1 + \left(\frac{1}{2} \log \pi + \log 2 + \frac{\gamma}{2}\right) \alpha + O(|\alpha|^2)\right) \left(-1/\alpha + \gamma + O(|\alpha|)\right) \\ &\quad \times \left(\sum_{n|\mathbb{P}} \frac{\mu(n)}{n} + \sum_{n|\mathbb{P}} \frac{\mu(n) \log n}{n} \alpha + O(|\alpha|^2)\right) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \\ &= -\frac{1}{\alpha} \sum_{n|\mathbb{P}} \frac{\mu(n)}{n} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) + \left(-\sum_{n|\mathbb{P}} \frac{\mu(n) \log n}{n} + \left(\frac{\gamma}{2} - \log(\sqrt{4\pi})\right) \sum_{n|\mathbb{P}} \frac{\mu(n)}{n}\right) \\ &\quad \times \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) + O_{\mathbb{P}}(|\alpha|). \end{aligned} \quad (4.29)$$

In the first step above, we have used the Taylor series expansion

$$\pi^{-(1-\alpha)/2} \Gamma\left(\frac{1-\alpha}{2}\right) = 1 + \left(\frac{1}{2} \log \pi + \log 2 + \frac{\gamma}{2}\right) \alpha + O(|\alpha|^2),$$

and the Laurent series expansion $\zeta(1-\alpha) = -1/\alpha + \gamma + O(|\alpha|)$ around $\alpha = 0$. In order to expand the product over primes involving α , we first write it in a different form as

$$\prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^{1-\alpha}}\right) = \sum_{n|\mathbb{P}} \frac{\mu(n)}{n^{1-\alpha}},$$

and then using the Laurent series expansion of n^α around $\alpha = 0$, we obtain

$$\prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^{1-\alpha}}\right) = \left(\sum_{n|\mathbb{P}} \frac{\mu(n)}{n} + \sum_{n|\mathbb{P}} \frac{\mu(n) \log n}{n} \alpha + O(|\alpha|^2)\right).$$

Similarly in $\mathcal{R}_{1+\alpha}$, using the Taylor series of $\pi^{-(1+\alpha)/2}\Gamma\left(\frac{1+\alpha}{2}\right)$, $\tau^{-\alpha/2}$, $(s-1-\alpha)^{-1}$, and the Laurent series of $\zeta(1+\alpha)$ around $\alpha = 0$, we have

$$\begin{aligned} \frac{\tau^{-\alpha/2}}{s-1-\alpha}\mathcal{R}_{1+\alpha} &= \left(1 - \frac{\log \tau}{2}\alpha + \mathcal{O}(|\alpha|^2)\right) \left(\frac{1}{s-1} + \frac{\alpha}{(s-1)^2} + \mathcal{O}(|\alpha|^2)\right) \\ &\quad \times \left(1 - \left(\frac{1}{2}\log \pi + \log 2 + \frac{\gamma}{2}\right)\alpha + \mathcal{O}(|\alpha|^2)\right) \left(\frac{1}{\alpha} + \gamma + \mathcal{O}(|\alpha|)\right) \\ &\quad \times \left(\sum_{n|\mathbb{P}} \frac{\mu(n)}{n} - \sum_{n|\mathbb{P}} \frac{\mu(n)\log n}{n}\alpha + \mathcal{O}(|\alpha|^2)\right) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \\ &= \frac{1}{\alpha(s-1)} \sum_{n|\mathbb{P}} \frac{\mu(n)}{n} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) + \frac{1}{(s-1)} \left(-\sum_{n|\mathbb{P}} \frac{\mu(n)\log n}{n}\right. \\ &\quad \left.+ \left(\frac{1}{s-1} + \frac{\gamma}{2} - \log(\sqrt{4\pi\tau})\right) \sum_{n|\mathbb{P}} \frac{\mu(n)}{n}\right) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) + \mathcal{O}_{\mathbb{P}}(|\alpha|). \end{aligned} \quad (4.30)$$

Therefore, around $\alpha = 0$, addition of (4.29) and (4.30) gives

$$\begin{aligned} \frac{\tau^{-1/2}}{s-1}\mathcal{R}_{1-\alpha} + \frac{\tau^{-(1+\alpha)/2}}{s-1-\alpha}\mathcal{R}_{1+\alpha} \\ = \frac{\tau^{-1/2}}{s-1} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \left(-2 \sum_{n|\mathbb{P}} \frac{\mu(n)\log n}{n} + \left(\frac{1}{s-1} + \gamma - \log(4\pi\sqrt{\tau})\right) \sum_{n|\mathbb{P}} \frac{\mu(n)}{n}\right) + \mathcal{O}_{\mathbb{P}}(\alpha). \end{aligned}$$

Thus,

$$\begin{aligned} \lim_{\alpha \rightarrow 0} \left(\frac{\tau^{-1/2}}{s-1}\mathcal{R}_{1-\alpha} + \frac{\tau^{-(1+\alpha)/2}}{s-1-\alpha}\mathcal{R}_{1+\alpha}\right) \\ = \frac{\tau^{-1/2}}{s-1} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \left(-2 \sum_{n|\mathbb{P}} \frac{\mu(n)\log n}{n} + \left(\frac{1}{s-1} + \gamma - \log(4\pi\sqrt{\tau})\right) \sum_{n|\mathbb{P}} \frac{\mu(n)}{n}\right). \end{aligned} \quad (4.31)$$

Similarly, we can find that,

$$\lim_{\alpha \rightarrow 0} (\mathcal{F}_{\alpha}(t) + \mathcal{F}_{-\alpha}(t)) = \frac{1}{4} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right) \left(-2 \sum_{n|\mathbb{P}} \frac{\mu(n)\log n}{n} + \left(\gamma - \log\left(4\pi\sqrt{\frac{\tau}{t}}\right)\right) \sum_{n|\mathbb{P}} \frac{\mu(n)}{n}\right). \quad (4.32)$$

We conclude by employing (4.31), (4.32) into (4.28), and writing $\sum_{n|\mathbb{P}} \frac{\mu(n)}{n}$ in the form of Dirichlet series $\prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right)$. \square

Now, we turn our attention to prove Corollary 3.3.

Proof of Corollary 3.3. In Theorem 1.2, we split the integration over t on the right-hand side, and consider the integral,

$$\int_1^\infty t^{-s/2-1/2} \left(-2 \sum_{n|\mathbb{P}} \frac{\mu(n)\log n}{n} + \left(\gamma - \log\left(4\pi\sqrt{\frac{\tau}{t}}\right)\right) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p}\right)\right) dt. \quad (4.33)$$

Using integral identities, for $\operatorname{Re}(s) > 1$,

$$\int_1^\infty t^{-s/2-1/2} dt = \frac{2}{s-1}, \quad (4.34)$$

$$\text{and } \int_1^\infty t^{-s/2-1/2} \log\left(4\pi\sqrt{\frac{\tau}{t}}\right) dt = -\frac{2}{(s-1)^2} + \frac{2}{(s-1)} \log(4\pi\sqrt{\tau}), \quad (4.35)$$

in (4.33), we obtain,

$$\begin{aligned} \int_1^\infty t^{-s/2-1/2} \left(-2 \sum_{n|\mathbb{P}} \frac{\mu(n) \log n}{n} + \left(\gamma - \log \left(4\pi \sqrt{\frac{\tau}{t}} \right) \right) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \right) dt \\ = -\frac{4}{(s-1)} \sum_{n|\mathbb{P}} \frac{\mu(n) \log n}{n} + 2 \left(\frac{\gamma}{(s-1)} + \frac{1}{(s-1)^2} - \frac{1}{(s-1)} \log(4\pi\sqrt{\tau}) \right) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right). \end{aligned}$$

Thus, for $\text{Re}(s) > 1$, and $\text{Re}(\tau) > 0$,

$$\begin{aligned} -\frac{1}{2\sqrt{\tau}} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \int_1^\infty t^{-s/2-1/2} \left(-2 \sum_{n|\mathbb{P}} \frac{\mu(n) \log n}{n} + \left(\gamma - \log \left(4\pi \sqrt{\frac{\tau}{t}} \right) \right) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \right) dt \\ = -\frac{1}{(s-1)\sqrt{\tau}} \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \left(-2 \sum_{n|\mathbb{P}} \frac{\mu(n)}{n} \log n + \left(\frac{1}{s-1} + \gamma - \log(4\pi\sqrt{\tau}) \right) \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p} \right) \right). \end{aligned} \quad (4.36)$$

From (4.36) we observe that the second and the last term on the right-hand side of (1.5) beautifully cancels out each other. Finally, the expression in (1.5) vastly simplifies to the required equation (3.4), for $\text{Re}(s) > 1$, and $\text{Re}(\tau) > 0$. \square

5. ACKNOWLEDGMENT

We are grateful to the reviewer who gave essential comments and feedback to improve our article. The authors would also like to thank Prof. Atul Dixit and Prof. Akshaa Vatwani for the insightful discussions and the critical first review of this manuscript.

REFERENCES

- [1] Apostol, T. M., *Introduction to Analytic Number Theory*, Undergraduate Texts in Mathematics, p. xii+338. Springer, New York-Heidelberg, 1976.
- [2] B.C. Berndt, Y. Lee, and J. Sohn, *Koshliakov's formula and Guinand's formula in Ramanujan's lost notebook*, Surveys in Number Theory, Series: Developments in Mathematics, vol. **17**, K. Alladi, ed., Springer, New York, 2008, pp. 21–42.
- [3] I. S. Gradshteyn and I. M. Ryzhik, eds., *Table of Integrals, Series, and Products*, 8th ed., Edited by D. Zwillinger, V. H. Moll, Academic Press, New York, 2015.
- [4] A. P. Guinand, *Some rapidly convergent series for the Riemann ξ -function*, Quart. J. Math. (Oxford) **6** (1955), 156–160.
- [5] G. H. Hardy and J. E. Littlewood, *The approximate functional equations for $\zeta(s)$ and $\zeta^2(s)$* , Proc. London Math. Soc., S2–29 (**1**), 81–97, 1929.
- [6] D. R. Heath-Brown, *Hybrid bounds for Dirichlet L-functions*, Invent. Math. **47** (1978), 149–170.
- [7] D. R. Heath-Brown, *The fourth power moment of the Riemann Zeta-function on the critical line*, Proc. London Math. Soc. (3), **38** (1979), 385–422.
- [8] A. Ivic, *The Riemann Zeta-Function*, John Wiley & Sons, New York, 1985.
- [9] A. F. Lavrik, *Functional equation for Dirichlet L-functions and the problem of divisors in arithmetic progressions. (Russian)*, Izv. Akad. Nauk SSSR Ser. Mat., **30:2** (1966), 433–448.
- [10] A. F. Lavrik, *Functional equations with parameter of zeta-functions, (Russian)*, Izv. Akad. Nauk SSSR Ser. Mat., **54:3** (1990), 501–521.
- [11] A. F. Lavrik, *A functional equation of the Riemann zeta function and an interval of the Euler product. (Russian)*, Dokl. Akad. Nauk SSSR 320 (1991), No. **1**, 29–33.
- [12] A. F. Lavrik, *Functional equation of the zeta function and a segment of the Euler product over prime numbers*, Soviet Math. Dokl., Vol 4, No. **2** (1992), 386–390.
- [13] A. F. Lavrik, *An approximate functional equation for the zeta function and a segment of the Euler product*, Russian Acad. Sci. Dokl. Math., Vol 46, No. **2** (1993), 201–204.
- [14] F. Oberhettinger, *Tables of Mellin Transforms*, Springer-Verlag, New York, 1974.
- [15] R. B. Paris and D. Kaminski, *Asymptotics and Mellin-Barnes Integrals*, Encyclopedia of Mathematics and its Applications, **85**. Cambridge University Press, Cambridge, 2001.
- [16] A. Simonič, *Explicit zero density estimate for the Riemann zeta-function near the critical line*, J. Math. Anal. Appl. 491(2020), no. **1**, 124303.
- [17] N. M. Temme, *Special functions: An introduction to the classical functions of mathematical physics*, Wiley-Interscience Publication, New York, 1996.

- [18] G. N. Watson, *Theory of Bessel Functions*, Cambridge University Press, second ed., London, 1966.

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, 6F, ASTRONOMY-MATHEMATICS BUILDING, NO. 1, SEC. 4, ROOSEVELT ROAD, TAIPEI 106319, TAIWAN,

CURRENT ADDRESS: THE UNIVERSITY OF TEXAS AT TYLER, TEXAS 75701, USA.

Email address: rajatgpt972@gmail.com, rgupta@uttyler.edu

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, MUMBAI 400076, MAHARASHTRA, INDIA

Email address, Corresponding author: aditiben.s@iitgn.ac.in, aditisav1995@gmail.com