

# Fully smooth one shot multipartite soft covering of quantum states without pairwise independence

Pranab Sen\*

## Abstract

We provide a powerful machinery to prove fully smooth one shot multipartite covering, aka convex split, type results for quantum states. In the important case of smooth multipartite convex split for classical quantum states, aka smooth multipartite soft covering, our machinery works even when certain marginals of these states do not satisfy pairwise independence. The paper [Sen24] gave the first proof of fully smooth multipartite convex split by simplifying and extending a technique called telescoping, developed originally for convex split by [CGB23]. However, that work as well as all earlier works on convex split assumed pairwise or even more independence amongst suitable marginals of the quantum states.

We develop our machinery by leveraging known results from [Sen21b] involving tilting and augmentation smoothing of quantum states, combined with a novel observation that a natural quantum operation ‘flattening’ quantum states actually preserves the fidelity. This machinery is powerful enough to lead to non pairwise independent results as mentioned above.

As an application of our soft covering lemma without pairwise independence, we prove the ‘natural’ one shot inner bounds for sending private classical information over a quantum wiretap interference channel, even when the classical encoders at the input lose pairwise independence in their encoding strategies to a certain extent. This result was unknown earlier even in the classical setting.

## 1 Introduction

The simultaneous smoothing bottleneck is a famous open problem in network quantum information theory [DF13]. A positive result for this problem would imply that many union and intersection type arguments carried out, sometimes implicitly, in network classical information theory extend similarly to the quantum setting. Most tasks in information theory can be decomposed into simpler tasks of one of two types: *packing* and *covering*. The earlier work of [Sen21b] gave a machinery for implementing union and intersection for packing tasks in one shot network quantum information theory, bypassing simultaneous smoothing. However it left open the question of implementing union and intersection for covering tasks, a lacuna which was explicitly pointed out in [DGHW20].

Recent exciting works of Cheng, Gao and Berta [CGB23], and Colomer and Winter [CW23], have introduced a telescoping cum mean-zero decomposition technique that bypasses the simultaneous smoothing bottleneck for intersection arguments for two fundamental problems in quantum

---

\*School of Technology and Computer Science, Tata Institute of Fundamental Research, Mumbai, India. Part of this work was done while the author was on sabbatical leave at the Centre for Quantum Technologies, National University of Singapore. Email: pranab.sen.73@gmail.com.

information theory viz. multipartite covering aka multipartite convex split, and multipartite decoupling respectively. However Cheng et al. did not state their multipartite convex split results in terms of smooth one shot quantities, leaving several basic one shot and finite blocklength achievability questions in network quantum information theory, e.g. inner bounds for the generalised one shot quantum Slepian Wolf [AJW18] problem, unanswered. Very recently, [Sen24] showed that the telescoping technique can in fact be simplified and further extended in order to prove fully smooth multipartite convex split and decoupling results for the first time. That work also provided several applications of the fully smooth convex split and decoupling theorems to important problems in network quantum information theory.

All the works referenced above as well as all earlier works on convex split, whether smooth or non smooth, assumed at least pairwise, often full independence or tensor product structure of certain marginals of quantum states. However, all the earlier proof techniques including telescoping fail when the pairwise independence requirement is dropped. Dropping pairwise independence becomes important for classical quantum soft covering lemmas. In multipartite soft covering, there are  $k$  classical parties,  $k \geq 1$ . Each party independently samples a set of classical symbols from a probability distribution. A  $k$ -tuple of symbols from the samples of the  $k$  parties is fed to a box whose output is a quantum state depending on the input  $k$ -tuple. The uniform average of the resulting input dependent quantum state is taken over all  $k$  tuples that can be constructed from the  $k$  sets sampled by the  $k$  parties. The aim is to ensure that the resulting sample averaged input dependent quantum state is close in trace distance to an ‘ideal’ fixed quantum state, in expectation over the random choices of the  $k$  parties. Dropping pairwise independence for soft covering means that a classical party chooses its set of symbols in a pairwise dependent fashion.

Soft covering lemmas are used e.g. to prove the privacy requirement for classical messages for various types of wiretap channels [RSW17, Wil17]. Relaxing pairwise independence in such settings can make computational sense as it will require senders to use less true random bits, an expensive resource, in their codebook construction step.

In this paper, we provide a new proof for fully smooth multipartite convex split of quantum states. Our proof does not use telescoping; instead it uses a completely different machinery. The advantage of using this more sophisticated machinery is that in the soft covering setting, we can show for the first time that our fully smooth inner bounds continue to hold even when the classical covering parties lose pairwise independence in a restricted way while choosing their set of samples. We believe this is a major conceptual contribution of our work.

In order to prove such powerful and general multipartite fully quantum convex split and soft classical quantum covering results, we develop new machinery by leveraging known results from [Sen21b] involving tilting and augmentation smoothing of quantum states, combined with a novel observation that a natural quantum operation ‘flattening’ quantum states actually preserves the fidelity. As an application of the power of this machinery, we prove the achievability of a natural rate region for sending private classical information over a wiretap quantum interference channel extending Chong, Motani, Garg and ElGamal inner bound [CMGE08] for the non-wiretap classical interference channel. Our scheme continues to work even when the classical encodings at the inputs to the channel lose pairwise independence to a certain extent. This was not known before, not even in the classical setting.

## 1.1 Organisation of the paper

Section 2 serves as a warmup, where we first recall the CMG covering problem introduced in [Sen24]. That work provided a smooth inner bound for CMG covering using telescoping. In Section 2, we will see how the telescoping proof of smooth CMG covering fails if the pairwise independence assumption between certain marginals of classical quantum states involved in the telescoping proof is removed. Some novel results about flattening quantum states, Schatten- $\ell_2$  norm and inner product of matrices, and tilting and augmentation smoothing of quantum states are stated and proved in Section 3. They will be required to prove fully smooth multipartite convex split in Section 4. The style of proof of convex split in Section 4 serves as a warmup to the proof of the smooth multipartite soft covering lemma without pairwise independence in Section 5. Section 5 also proves a smooth inner bound for the CMG covering problem that works even when certain pairwise independence assumptions are relaxed, addressing the drawback of telescoping pointed out earlier in Section 2. We use this inner bound for CMG covering in Section 6, where we finally obtain smooth inner bounds for sending private classical information over a quantum wiretap interference channel, even when certain pairwise independence requirements at the classical encoders are relaxed. Section 7 contains an alternate simpler proof of fully smooth multipartite convex split using our flattening technique. It serves as a warmup to Section 8, where we prove a fully smooth multipartite decoupling theorem using similar methods. The fully smooth multipartite decoupling theorem of Section 8 uses one extra ancilla qubit per sender unlike the fully smooth multipartite decoupling theorem in [Sen24]; however this extra qubit per sender does not affect any application of decoupling to the best of our knowledge. We finally make some concluding remarks and list directions for further research in Section 9.

## 2 Warmup: Telescoping fails without pairwise independence

The CMG covering problem defined in [Sen24] lies at the heart of obtaining inner bounds for private classical communication over a quantum wiretap interference channel. The name CMG is an acronym for Chong, Motani and Garg, the discoverers of one of the most well known inner bounds for the classical non-wiretap interference channel in the asymptotic independent and identically distributed (asymptotic iid) setting [CMGE08]. The paper [Sen12] first showed the achievability of the same inner bound for sending classical information over a quantum non-wiretap interference channel in the asymptotic iid setting. Subsequently, the achievability in the non-wiretap quantum case in the general one shot setting was proved in [Sen21a]. That paper used the machinery of [Sen21b] to bypass simultaneous smoothing for intersection problems arising in packing tasks. Later on in this paper, we will be able to extend the result of [Sen21a] to the one shot wiretap quantum case by combining the packing arguments with our fully smooth CMG covering lemma without pairwise independence (Lemma 19 below).

Let  $\mathcal{N}$  be a quantum channel with two inputs called Alice and Bob, and one output called Eve. The intuition here is that Eve is an eavesdropper trying to get information on the classical messages that Alice and Bob are sending to some other outputs of  $\mathcal{N}$ , not described here explicitly as they are not relevant for CMG covering. Though  $\mathcal{N}$  has quantum inputs and quantum outputs, we will assume without loss of generality that the inputs are classical, because in the inner bound below, there is a standard optimisation step over a choice of all ensembles of input states to Alice and Bob. Thus, we will henceforth think of  $\mathcal{N}$  as a classical-quantum (cq) channel with its two classical input alphabets being denoted by  $\mathcal{X}$ ,  $\mathcal{Y}$ , and the quantum output alphabet by  $\mathcal{E}$ .

Let  $0 \leq \epsilon \leq 1$ . On input  $(x, y) \in \mathcal{X} \times \mathcal{Y}$ , the channel outputs a quantum state

$$\sigma_{xy}^E := \mathcal{N}^{AB \rightarrow E}(\rho_x^A \otimes \rho_y^B)$$

on  $E$ , where  $\{\rho_x^A\}_x$ ,  $\{\rho_y^B\}_y$  are the ensembles of the so-called encoding quantum states at the quantum inputs  $A, B$  of the channel which is modelled by a completely positive trace non-increasing superoperator  $\mathcal{N} : AB \rightarrow E$ . To define the CMG covering problem, we need to define new alphabets  $\mathcal{X}'$ ,  $\mathcal{Y}'$ , following the scheme of the original paper [CMGE08]. We then pick a ‘control’ probability distribution

$$p(x', x, y', y) := p(x', x)p(y', y)$$

on the classical alphabet  $\mathcal{X}' \times \mathcal{X} \times \mathcal{Y}' \times \mathcal{Y}$ . The ‘control’ cq state for the CMG covering problem is now defined as

$$\sigma^{X'XY'YE} := \sum_{x', x, y', y} p(x', x, y', y) |x', x, y', y\rangle \langle x', x, y', y|^{X'XY'Y} \otimes \sigma_{xy}^E.$$

Though the state  $\sigma_{xy}^E$  depends only on  $x$  and  $y$ , for later convenience we will write it as  $\sigma_{x'xy'y}^E := \sigma_{xy}^E$ . Though this notation seems heavier now, it will be extremely useful when we take various marginals of  $\sigma^{X'XY'YE}$  in the entropic quantities and the proofs later on. Thus,

$$\sigma^{X'XY'YE} := \sum_{x', x, y', y} p(x', x)p(y', y) |x', x, y', y\rangle \langle x', x, y', y|^{X'XY'Y} \otimes \sigma_{x'xy'y}^E.$$

Let us say Alice and Bob are trying to send a pair of classical messages  $a$  and  $b$  respectively to Charlie. To obfuscate them from Eve, Alice and Bob independently do the following strategy. Alice chooses iid samples  $x'_1, \dots, x'_{L'}$  from the marginal distribution  $p^{X'}$ . Conditioned on each sample  $x'_{l'} \in \mathcal{X}'$ , Alice chooses iid samples  $x_{l',1}, \dots, x_{l',L}$  from the conditioned marginal distribution  $p^X | (X' = x'_{l'})$ . Thus, a total of  $L'L$  samples are chosen by Alice. Similarly, Bob chooses iid samples  $y'_1, \dots, y'_{M'}$  from the marginal distribution  $p^{Y'}$ . Conditioned on each sample  $y'_{m'} \in \mathcal{Y}'$ , Bob chooses iid samples  $y_{m',1}, \dots, y_{m',M}$  from the conditioned marginal distribution  $p^Y | (Y' = y'_{m'})$ . Thus, a total of  $M'M$  samples are chosen by Bob. Alice inputs a uniformly random sample from her chosen set into the channel  $\mathcal{N}$ . Similarly, Bob inputs a uniformly random sample from his chosen set into the channel  $\mathcal{N}$ . The hope is that this strategy obfuscates Eve’s received state so much that she is unable to figure out the actual input message pair  $(a, b)$ .

Let  $\sigma^E$  denote the marginal on  $E$  of the control state  $\sigma^{X'XY'YE}$ . Define

$$\sigma_{\vec{x}', \vec{x}, \vec{y}', \vec{y}}^E := (L'(L+1)M'(M+1))^{-1} \sum_{a'=1}^{L'} \sum_{a=1}^L \sum_{b'=1}^{M'} \sum_{b=1}^M \sigma_{x'_{a'} x_a y'_{b'} y_b}^E. \quad (1)$$

We would like to show that the so-called CMG convex split quantity defined below is small, when  $L', L, M', M$  are suitably large.

$$\text{CMG} := \mathbb{E}_{\vec{x}', \vec{x}, \vec{y}', \vec{y}} [\|\sigma_{\vec{x}', \vec{x}, \vec{y}', \vec{y}}^E - \sigma^E\|_1], \quad (2)$$

the expectation being over the choice of the samples  $\vec{x}' := (x'_1, \dots, x'_{L'})$ ,  $\vec{x} := (x_{1,1}, \dots, x'_{L',L})$ ,  $\vec{y}' := (y'_1, \dots, y'_{M'})$ ,  $\vec{y} := (y_{1,1}, \dots, y'_{M',M})$  according to the procedure described above. Recall that

for  $p > 0$ , the *Schatten- $\ell_p$  norm* of a matrix  $M$  is defined as  $\|M\|_p := \text{Tr} [(A^\dagger A)^{p/2}]$ . The *Schatten- $\ell_\infty$  norm*, aka operator norm induced from the Hilbert space norm, is defined by taking  $p \rightarrow +\infty$  in the above expression, resulting in  $\|M\|_\infty$  being the largest singular value of  $M$ . For normal matrices  $M$ , it equals the largest absolute value of an eigenvalue of  $M$ .

The paper [Sen24] gave a fully smooth inner bound for the CMG covering problem using telescoping. The telescoping based proof in that paper uses expressions like

$$\mathbb{E}_{x', x, \hat{x}, y', y} \left[ \text{Tr} \left[ (\tilde{\sigma}_{x'xy'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'xy'}^E(8) + \tilde{\sigma}_{x'y'}^E(8)) \right. \right. \\ \left. \left. (\tilde{\sigma}_{x'\hat{x}y'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'\hat{x}y'}^E(8) + \tilde{\sigma}_{x'y'}^E(8)) \right] \right], \quad (3)$$

in order to bound the CMG quantity in Equation 2 above. Above, a matrix like  $\tilde{\sigma}_{x'xy'y}^E(8)$  is a perturbed version of the matrix

$$\tilde{\sigma}_{x'xy'y}^E := (\sigma^E)^{-1/2} \circ \sigma_{x'xy'y}^E := (\sigma^E)^{-1/2} \sigma_{x'xy'y}^E (\sigma^E)^{-1/2},$$

where  $\sigma^{-1}$  is the so-called Moore-Penrose pseudoinverse of  $\sigma$  i.e. the inverse on the support of  $\sigma$  and the zero operator orthogonal to the support. Eight different perturbed versions of  $\tilde{\sigma}_{x'xy'y}^E$  called  $\tilde{\sigma}_{x'xy'y}^E(i)$ ,  $i \in [8]$ , are used at different places of the telescoping proof [Sen24] in order to prove fully smooth inner bounds for CMG covering.

The telescoping technique crucially requires that a telescoping term satisfy the so-called *mean zero property* [Sen24]. For a term like in Equation 3 above, various conditions need to be met in order for the mean zero requirement to be satisfied. One of those conditions is as follows:

$$\begin{aligned} & \mathbb{E}_{x', x, \hat{x}, y', y} \left[ \text{Tr} \left[ (\tilde{\sigma}_{x'xy'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'xy'}^E(8) + \tilde{\sigma}_{x'y'}^E(8)) \right. \right. \\ & \quad \left. \left. (\tilde{\sigma}_{x'\hat{x}y'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'\hat{x}y'}^E(8) + \tilde{\sigma}_{x'y'}^E(8)) \right] \right] \\ &= \mathbb{E}_{x', y', y} \left[ \text{Tr} \left[ \left( \mathbb{E}_{x|x'} [\tilde{\sigma}_{x'xy'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'xy'}^E(8) + \tilde{\sigma}_{x'y'}^E(8)] \right) \right. \right. \\ & \quad \left. \left. \left( \mathbb{E}_{\hat{x}|x'} [\tilde{\sigma}_{x'\hat{x}y'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'\hat{x}y'}^E(8) + \tilde{\sigma}_{x'y'}^E(8)] \right) \right] \right] \\ &= \mathbb{E}_{x', y', y} \left[ \text{Tr} \left[ (\tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'y'}^E(8) + \tilde{\sigma}_{x'y'}^E(8)) \right. \right. \\ & \quad \left. \left. (\tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'y'}^E(8) + \tilde{\sigma}_{x'y'}^E(8)) \right] \right] \\ &= 0. \end{aligned} \quad (4)$$

The equalities in the above equation only hold when in the obfuscating strategy of Alice described earlier, the distribution of  $x_{a'a}$  conditioned on  $x'_{a'}$  is pairwise independent from the distribution of  $x_{a'\hat{a}}$  conditioned on  $x'_{a'}$  for any pair  $a \neq \hat{a}$ .

We can now appreciate the minimum requirement of pairwise independence in the proofs of both non-smooth as well as smooth covering lemmas. Suppose the pairwise independence condition in Alice's obfuscating strategy is 'slightly broken'. The mean zero property in Equation 4 fails. By *slightly broken* we mean that conditioned on a particular choice of  $x'_{a'}$ ,  $x_{a'a}$ ,  $y'_{b'}$ ,  $y_{b'b}$ ,

$$\mathbb{E}_{x_{a'\hat{a}} | (x'_{a'}, x_{a'a}, y'_{b'}, y_{b'b})} [\sigma_{x'_{a'}, x_{a'\hat{a}}, y'_{b'}, y_{b'b}}^E] =: \dot{\sigma}_{x'_{a'}, x_{a'a}, y'_{b'}, y_{b'b}}^E \neq \sigma_{x'_{a'}, y'_{b'}, y_{b'b}}^E, \quad \dot{\sigma}_{x'_{a'}, x_{a'a}, y'_{b'}, y_{b'b}}^E \leq \epsilon^{-1} \sigma_{x'_{a'}, y'_{b'}, y_{b'b}}^E.$$

We require that the expectation above depend only on the actual symbols  $x'_{a'}$ ,  $x_{a'a}$ ,  $y'_{b'}$ ,  $y_{b'b}$  and not on their locations  $a'$ ,  $a$ ,  $\hat{a}$ ,  $b'$  or  $b$ .

This creates immense problems for telescoping. Suppose for simplicity that only the pairwise independence condition involving  $X'$  and  $X$  is slightly broken as described above i.e. conditioned on  $x'_{a'}$ , the distribution of  $x_{a'a}$  is not independent from the distribution of  $x_{a'\hat{a}}$  for  $a \neq \hat{a}$ . Other pairs like  $(x'_{a'}, x_{a'a})$  and  $(x'_{\hat{a}'}, x_{\hat{a}'\hat{a}})$  for  $a' \neq \hat{a}'$ , or  $y_{b'b}$  and  $y_{b'\hat{b}}$  conditioned on  $y'_{b'}$  for  $b \neq \hat{b}$  etc. continue to be independent. Now writing a term like

$$(\tilde{\sigma}_{x'_{a'}x_{a'a}y'_{b'}y_{b'b}}^E(8) - \tilde{\sigma}_{x'_{a'}y'_{b'}y_{b'b}}^E(8) - \tilde{\sigma}_{x'_{a'}x_{a'a}y'_{b'}}^E(8) + \tilde{\sigma}_{x'_{a'}y'_{b'}}^E(8))$$

in the telescoping inequalities, requires for example the following condition (and several other conditions also) to be met in order to satisfy the mean zero property. The following condition occurs in the telescoping proof of [Sen24] when that proof considers the mean zero requirement for the case  $a' = \hat{a}'$ ,  $a \neq \hat{a}$ ,  $b' = \hat{b}'$ ,  $b = \hat{b}$ ,

$$\begin{aligned} & \mathbb{E}_{x', x, \hat{x}, y', y} [\text{Tr} [(\tilde{\sigma}_{x'xy'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'xy'}^E(8) + \tilde{\sigma}_{x'y'}^E(8))(\tilde{\sigma}_{x'\hat{x}y'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'\hat{x}y'}^E(8) + \tilde{\sigma}_{x'y'}^E(8))]] \\ &= \mathbb{E}_{x', x, y', y} [\text{Tr} [(\tilde{\sigma}_{x'xy'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'xy'}^E(8) + \tilde{\sigma}_{x'y'}^E(8)) \\ &\quad \left( \mathbb{E}_{\hat{x}|(x'xy'y)} [\tilde{\sigma}_{x'\hat{x}y'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'\hat{x}y'}^E(8) + \tilde{\sigma}_{x'y'}^E(8)] \right)]] \\ &= \mathbb{E}_{x', x, y', y} [\text{Tr} [(\tilde{\sigma}_{x'xy'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'xy'}^E(8) + \tilde{\sigma}_{x'y'}^E(8))(\dot{\sigma}_{x'xy'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \dot{\sigma}_{x'xy'}^E(8) + \tilde{\sigma}_{x'y'}^E(8))]] \\ &\neq 0, \end{aligned}$$

which destroys the mean zero property. On the other hand, writing a term like

$$(\tilde{\sigma}_{x'_{a'}x_{a'a}y'_{b'}y_{b'b}}^E(8) - \dot{\sigma}_{x'_{a'}y'_{b'}y_{b'b}}^E(8) - \tilde{\sigma}_{x'_{a'}x_{a'a}y'_{b'}}^E(8) + \dot{\sigma}_{x'_{a'}y'_{b'}}^E(8))$$

in the telescoping inequalities gives an expression like the following to meet the mean zero requirement in the proof of [Sen24], when  $a' = \hat{a}'$ ,  $a \neq \hat{a}$ ,  $b' = \hat{b}'$ ,  $b = \hat{b}$ ,

$$\begin{aligned} & \mathbb{E}_{x', x, \hat{x}, y', y} [\text{Tr} [(\tilde{\sigma}_{x'xy'y}^E(8) - \dot{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'xy'}^E(8) + \dot{\sigma}_{x'y'}^E(8))(\tilde{\sigma}_{x'\hat{x}y'y}^E(8) - \dot{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'\hat{x}y'}^E(8) + \dot{\sigma}_{x'y'}^E(8))]] \\ &= \mathbb{E}_{x', x, y', y} [\text{Tr} [(\tilde{\sigma}_{x'xy'y}^E(8) - \dot{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'xy'}^E(8) + \dot{\sigma}_{x'y'}^E(8)) \\ &\quad \left( \mathbb{E}_{\hat{x}|(x'xy'y)} [\tilde{\sigma}_{x'\hat{x}y'y}^E(8) - \dot{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'\hat{x}y'}^E(8) + \dot{\sigma}_{x'y'}^E(8)] \right)]] \\ &= \mathbb{E}_{x', x, y', y} [\text{Tr} [(\tilde{\sigma}_{x'xy'y}^E(8) - \dot{\sigma}_{x'y'y}^E(8) - \tilde{\sigma}_{x'xy'}^E(8) + \dot{\sigma}_{x'y'}^E(8))(\dot{\sigma}_{x'xy'y}^E(8) - \tilde{\sigma}_{x'y'y}^E(8) - \dot{\sigma}_{x'xy'}^E(8) + \tilde{\sigma}_{x'y'}^E(8))]] \\ &\neq 0, \end{aligned}$$

which again destroys the mean zero property. In fact, there is no choice of term one could write down for a successful telescoping strategy when pairwise independence is slightly broken.

The proof of the non-smooth inner bound for CMG covering in [Sen24] handles the loss of pairwise independence without much trouble giving rise to an inequality like

$$\log L' + \log M' + \log M > D_\infty(\dot{\sigma}^{X'XY'YE} \parallel \sigma^{X'XY'Y} \otimes \sigma^E) + \log \epsilon^{-2}, \quad (5)$$

in the inner bound. Above, we have used the *non-smooth Rényi- $\infty$  divergence*, aka non-smooth max divergence. This and the *non-smooth Rényi-2 divergence* are defined below.

### Definition 1

$$\begin{aligned}
I_2(X' : E)_\sigma &:= D_2(\sigma^{X'E} \| \sigma^{X'} \otimes \sigma^E), \\
I_\infty(X' : E)_\sigma &:= D_\infty(\sigma^{X'E} \| \sigma^{X'} \otimes \sigma^E), \\
D_2(\alpha \| \beta) &:= 2 \log \|\beta^{-1/4} \alpha \beta^{-1/4}\|_2, \quad \text{if } \text{supp}(\alpha) \leq \text{supp}(\beta), +\infty \text{ otherwise}, \\
D_\infty(\alpha \| \beta) &:= \log \|\beta^{-1/2} \alpha \beta^{-1/2}\|_\infty, \quad \text{if } \text{supp}(\alpha) \leq \text{supp}(\beta), +\infty \text{ otherwise}.
\end{aligned}$$

where  $\alpha$  is a *subnormalised density matrix* and  $\beta$  is a positive semidefinite matrix acting on the same Hilbert space. The quantity  $I_\infty(X' : E)_\sigma$  is known as *non-smooth Rényi- $\infty$  mutual information*, aka non-smooth max mutual information under the joint state  $\sigma^{X'E}$ . The quantity  $I_2(X' : E)_\sigma$  is known as the *non-smooth Rényi-2 mutual information* under the joint state  $\sigma^{X'E}$ . In this paper, a subnormalised density matrix means a positive semidefinite matrix with trace at most one; a *normalised density matrix* means the trace is exactly one. Note that  $D_2(\alpha \| \beta) \leq D_\infty(\alpha \| \beta)$ , and that for a cq state  $\sigma^{X'E}$ ,

$$I_2(X' : E)_\sigma = \log \mathbb{E}_{x'} \text{Tr} [((\sigma^E)^{-1/4} \sigma_{x'}^E (\sigma^E)^{-1/4})^2] = \log \mathbb{E}_{x'} \text{Tr} [(\tilde{\sigma}_{x'}^E)^2],$$

the expectation over  $x'$  being taken according to the classical probability distribution  $\sigma^{X'}$ . For later use, we remark that for a cq state  $\sigma^{X'E}$ ,

$$I_\infty(X' : E)_\sigma = \max_{x'} D_\infty(\sigma_{x'}^E \| \sigma^E).$$

The rate inequality in Equation 5 above arises from a term like the one below, when the proof of the non-smooth inner bound for CMG covering in [Sen24] considers a situation where  $a' = \hat{a}'$ ,  $b' = \hat{b}'$ ,  $b = \hat{b}$  and  $a \neq \hat{a}$ :

$$\begin{aligned}
& \mathbb{E}_{\vec{x}, \vec{x}', \vec{y}, \vec{y}'} \left[ \text{Tr} [\tilde{\sigma}_{x'_a x_{a'} y'_b y_b}^E \tilde{\sigma}_{x'_a x_{\hat{a}'} y'_b y_{\hat{b}'}}^E] \right] \\
&= \mathbb{E}_{x' x \hat{x} y' y} [\text{Tr} [\tilde{\sigma}_{x' x y' y}^E \tilde{\sigma}_{x' \hat{x} y' y}^E]] = \mathbb{E}_{x' x \hat{x} y' y} [\text{Tr} [\sigma_{x' x y' y}^E (\sigma^E)^{-1/2} \sigma_{x' \hat{x} y' y}^E (\sigma^E)^{-1/2}]] \\
&= \mathbb{E}_{x' x y' y} \left[ \text{Tr} \left[ \sigma_{x' x y' y}^E (\sigma^E)^{-1/2} \mathbb{E}_{\hat{x} | (x' x y' y)} [\sigma_{x' \hat{x} y' y}^E] (\sigma^E)^{-1/2} \right] \right] \\
&= \mathbb{E}_{x' x y' y} [\text{Tr} [\sigma_{x' x y' y}^E (\sigma^E)^{-1/2} \dot{\sigma}_{x' x y' y}^E (\sigma^E)^{-1/2}]] \\
&\leq 2^{D_\infty(\dot{\sigma}^{X'XY'YE} \| \sigma^{X'XY'Y} \otimes \sigma^E)} \cdot \mathbb{E}_{x' x y' y} [\text{Tr} [\sigma_{x' x y' y}^E (\sigma^E)^{-1/2} \sigma^E (\sigma^E)^{-1/2}]] \\
&= 2^{D_\infty(\dot{\sigma}^{X'XY'YE} \| \sigma^{X'XY'Y} \otimes \sigma^E)} \cdot \mathbb{E}_{x' x y' y} [\text{Tr} [\sigma_{x' x y' y}^E]] = 2^{D_\infty(\dot{\sigma}^{X'XY'YE} \| \sigma^{X'XY'Y} \otimes \sigma^E)},
\end{aligned}$$

where the cq state  $\dot{\sigma}^{X'XY'YE}$  is defined in the natural fashion as

$$\dot{\sigma}^{X'XY'YE} := \sum_{x' x y' y} p(x', x) p(y', y) |x', x, y', y\rangle \langle x', x, y', y|^{X'XY'Y} \otimes \dot{\sigma}_{x' x y' y}^E.$$

Observe that if

$$\dot{\sigma}_{x' x y' y}^E = \sigma_{x' x y' y}^E,$$

which is what we have in the pairwise independent case, then,  $D_\infty(\dot{\sigma}^{X'XY'YE} \| \sigma^{X'XY'Y} \otimes \sigma^E) = D_\infty(\sigma^{X'XY'YE} \| \sigma^{X'XY'Y} \otimes \sigma^E)$ . Our non-smooth rate region then becomes exactly like the non-smooth

rate region for CMG covering in [Sen24] in the pairwise independent case, except for  $D_2(\|\cdot\|)$  in [Sen24]'s rate region being replaced by  $D_\infty(\|\cdot\|)$ . Now suppose

$$\forall x : \dot{\sigma}_{x'xy'y}^E < \epsilon^{-1} \sigma_{x'y'y}^E,$$

i.e. the ‘consequences’ of pairwise independence hold up to a ‘scale factor’ of  $\epsilon^{-1}$ . Then,

$$D_\infty(\dot{\sigma}^{X'XY'YE} \| \sigma^{X'XY'Y} \otimes \sigma^E) \leq D_\infty(\sigma^{X'Y'YE} \| \sigma^{X'Y'Y} \otimes \sigma^E) + \log \epsilon^{-1}.$$

Under an appropriate simultaneous smoothing conjecture, such a term should give rise to an inequality like

$$\log L' + \log M' + \log M > D_\infty^\epsilon(\sigma^{X'Y'YE} \| \sigma^{X'Y'Y} \otimes \sigma^E) + \log \epsilon^{-1} + \text{polylog}(\epsilon^{-1}),$$

where  $0 < \epsilon < 1$  is the so-called *smoothing parameter*. Above we used the smooth Rényi- $\infty$  divergence  $D_\infty^\epsilon(\|\cdot\|)$ . The  $\epsilon$ -smooth Rényi divergences are defined as follows:

$$D_2^\epsilon(\alpha \| \beta) := \min_{\alpha' \approx_\epsilon \alpha} D_2(\alpha' \| \beta), \quad D_\infty^\epsilon(\alpha \| \beta) := \min_{\alpha' \approx_\epsilon \alpha} D_\infty(\alpha' \| \beta),$$

where the minimisation is over all subnormalised density matrices  $\alpha'$  satisfying  $\|\alpha' - \alpha\|_1 \leq \epsilon(\text{Tr } \alpha)$ . The smooth mutual informations are defined accordingly.

$$I_2^\epsilon(X' : E)_\sigma := D_2^\epsilon(\sigma^{X'E} \| \sigma^{X'} \otimes \sigma^E), \quad I_\infty^\epsilon(X' : E)_\sigma := D_\infty^\epsilon(\sigma^{X'E} \| \sigma^{X'} \otimes \sigma^E).$$

We remark that several alternate definitions of non-smooth and smooth Rényi divergences have been provided in the literature. However the work of [ABJT19] shows that all the smooth Rényi- $p$  divergences for  $p > 1$  are essentially equivalent to  $D_\infty^\epsilon$  up to tweaks in the smoothing parameter  $\epsilon$  and dimension independent additive corrections that are polynomial in  $\log \epsilon^{-1}$ . So henceforth, our fully smooth convex split and soft covering results shall be stated in terms of  $D_\infty^\epsilon$  or quantities derived from  $D_\infty^\epsilon$  like  $I_\infty^\epsilon$ .

Another important entropic quantity derived from Rényi divergence is conditional entropy. The smooth conditional Rényi entropy is defined by

$$H_2^\epsilon(A|R)_\rho := -D_2^\epsilon(\rho^{AR} \| \mathbb{1}^A \otimes \rho^R), \quad H_{\min}^\epsilon(A|R)_\rho := -D_\infty^\epsilon(\rho^{AR} \| \mathbb{1}^A \otimes \rho^R).$$

Various alternate definitions of the smooth conditional Rényi-2 and Rényi min entropy have been given in the literature, but there all equivalent up to minor tweaks in the smoothing parameter  $\epsilon$  and small dimension independent additive terms polynomial in  $\log \epsilon^{-1}$ . Also,  $H_2^\epsilon(A|R)_\rho \approx H_{\min}^\epsilon(A|R)_\rho$  [ABJT19].

Now contrast the above expression with the following viz.

$$\begin{aligned} \log L' + \log M' + \log M &> I_2^\epsilon(X'Y'Y : E)_\sigma + \log \epsilon^{-2} = D_2^\epsilon(\sigma^{X'Y'YE} \| \sigma^{X'Y'Y} \otimes \sigma^E) + \log \epsilon^{-2} \\ &\approx D_\infty^\epsilon(\sigma^{X'Y'YE} \| \sigma^{X'Y'Y} \otimes \sigma^E) + \Theta(\log \epsilon^{-2}) \end{aligned}$$

obtained by telescoping in the pairwise independent case [Sen24]. Thus, simultaneous smoothing is still potentially much more powerful than telescoping when pairwise independence is slightly broken in the sense of  $\dot{\sigma}_{x'xy'y}^E$  being bounded by  $\sigma_{x'y'y}^E$  by at most a scale factor of  $\text{poly}(\epsilon^{-1})$ .

As we will see in Section 5, our machinery allows us to handle the slight breaking of pairwise independence in the sense described above, achieving the bound in terms of  $D_\infty^\epsilon$  that one would expect from simultaneous smoothing. We remark that sacrificing pairwise independence may be computationally beneficial for Alice’s obfuscation strategy, as it will require her to use a lesser number of pure random bits.



### 3 Preliminaries

We now state some basic definitions and facts for completeness. First is the well known *gentle measurement lemma*.

**Fact 1** *Let  $\rho$  be a normalised density matrix on a Hilbert space. Let  $\Pi$  be a POVM element the same space (i.e.  $0 \leq \Pi \leq \mathbb{1}$ ) satisfying  $\text{Tr} [\Pi\rho] \geq 1 - \epsilon$ . Then,*

$$\|\rho - \sqrt{\Pi}\rho\sqrt{\Pi}\|_1 < 2\sqrt{\epsilon}.$$

*If  $\rho$  is not normalised, we can still obtain a good gentle measurement upper bound by rescaling by  $\text{Tr} \rho$ .*

The *Löwner partial order* on Hermitian matrices  $A, B$  on the same Hilbert space is defined as  $A \leq B$  iff  $B - A$  is positive semidefinite. With this notation, we sometimes write  $A \geq 0$  to indicate that  $A$  is positive semidefinite.

Every Hermitian positive semidefinite matrix  $\rho^A$  on a Hilbert space  $A$ , with complex entries, has a *length preserving purification*  $\rho^{AR}$  on an extended Hilbert space  $A \otimes R$ , where  $R$  is another Hilbert space, often called the *reference*, of sufficiently large dimension. By a length preserving purification, we mean that  $\rho^{AR}$  is a rank one orthogonal projection, i.e. a *pure state*, whose of same Schatten- $\ell_1$  norm as that of  $\rho^A$ . Note that the Schatten- $\ell_1$  norm of a pure state  $\rho^{AR} = |\rho\rangle\langle\rho|^{AR}$  is the Euclidean  $\ell_2$ -length of the vector  $|\rho^A\rangle$ .

**Definition 2** *The fidelity between two positive semidefinite matrices  $\rho^A, \sigma^A$  on the same Hilbert space  $A$  is defined as*

$$F(\rho^A, \sigma^A) := \|\sqrt{\rho^A}\sqrt{\sigma^A}\|_1 = \max_{|\rho\rangle^{AR}, |\sigma\rangle^{AR}} |\langle\rho|\sigma\rangle|,$$

*where the maximisation is over all length preserving purifications  $|\rho\rangle^{AR}, |\sigma\rangle^{AR}$  of  $\rho^A, \sigma^A$ .*

Given the above definition of fidelity, a straightforward analogue of Uhlmann's theorem holds for pairs of positive semidefinite matrices. Also, we have the standard inequalities relating fidelity and trace distance for normalised density matrices.

**Fact 2** *Let  $\rho, \sigma$  be normalised density matrices in the same Hilbert space. Then,*

$$1 - F(\rho, \sigma) \leq \frac{1}{2}\|\rho - \sigma\|_1 \leq \sqrt{1 - F(\rho, \sigma)^2}.$$

In this paper, all superoperators will be *Hermitian preserving* i.e. they are linear operators on vector spaces of matrices that send Hermitian matrices to Hermitian matrices. We will only apply superoperators to Hermitian matrices inside our proofs.

We will need the notion of ‘intersection’ of two probability subdistributions  $p^X, q^X$  on the same alphabet  $X$  from [Sen21b].

$$(p \cap q)^X(x) := \min\{p(x), q(x)\}. \tag{6}$$

It is easy to see that

$$\|(p \cap q)^X - p^X\|_1 \leq \|p^X - q^X\|_1, \quad \|(p \cap q)^X - q^X\|_1 \leq \|p^X - q^X\|_1.$$

We state the following two folklore facts for completeness, which have been commonly used in earlier literature on decoupling.

**Fact 3** Let  $\rho^{AB}$  be a positive semidefinite matrix. Then  $\rho^{AB} \leq |A|(\mathbb{1}^A \otimes \rho^B)$ .

**Fact 4** Let  $\rho^{AB}$  be a subnormalised density matrix. Then  $H_{\min}^\epsilon(A|B)_\rho \leq \log |A|$ .

We now recall the so-called *matrix weighted Cauchy-Schwarz inequality*. This inequality has been extensively used in earlier literature on decoupling.

**Fact 5** Let  $M$  be Hermitian matrix on a Hilbert space. Let  $\sigma$  be a normalised density matrix on the same space such that  $\text{supp}(M) \subseteq \text{supp}(\sigma)$ . Then,

$$\|M\|_1 \leq \|\sigma^{-1/4} M \sigma^{-1/4}\|_2,$$

where  $\sigma^{-1}$  is the Moore-Penrose pseudoinverse of  $\sigma$  as defined earlier. If  $\text{Tr } \sigma \neq 1$ , still an appropriate inequality can be easily obtained by rescaling by  $\text{Tr } \sigma$ .

### 3.1 Shaved Cauchy-Schwarz inequality for matrix norms

Though the matrix weighted Cauchy Schwarz inequality is very useful, it does have a notable drawback in certain applications. Suppose we want to show an upper bound for  $\|M_1 - M_2\|_1$ , where  $M_1, M_2$  are Hermitian matrices. In several applications, including during the proof of the smooth multipartite convex split lemma in the next section,  $M_1, M_2$  have incomparable supports, and it is not clear what normalised density matrix  $\sigma$  should be used as the weighting matrix in Fact 5 so as to get a good upper bound on  $\|M_1 - M_2\|_1$ . For several of those applications, including for our proof in the next section, it turns out that one can apply the following *shaved Cauchy-Schwarz inequality* of Proposition 1 in order to get very good upper bounds on  $\|M_1 - M_2\|_1$ . The term ‘shaved’ refers to the use of the POVM element  $\Pi$  in Proposition 1 which shaves off the ‘bad parts’ of  $\rho_1, \rho_2$  and applies Cauchy Schwarz only on the good part  $\sqrt{\Pi}(\rho_1 - \rho_2)\sqrt{\Pi}$ .

**Proposition 1** Suppose  $\rho_1, \rho_2$  are two positive semidefinite matrices on the same Hilbert space. Let  $\Pi$  be a POVM element on the same space. Suppose  $\text{Tr } [\Pi \rho_i] \geq (1 - \epsilon_i)(\text{Tr } \rho_i)$ . Then,

$$\|\rho_1 - \rho_2\|_1 \leq \sqrt{\text{Tr } \Pi} \cdot \|\rho_1 - \rho_2\|_2 + (2 \text{Tr } \rho_1) \sqrt{\epsilon_1} + (2 \text{Tr } \rho_2) \sqrt{\epsilon_2}.$$

**Proof:** We just apply Fact 1 and the standard Cauchy-Schwarz inequality, together with the triangle inequality.

$$\begin{aligned} \|\rho_1 - \rho_2\|_1 &\leq \|\sqrt{\Pi}(\rho_1 - \rho_2)\sqrt{\Pi}\|_1 + \|\rho_1 - \sqrt{\Pi} \rho_1 \sqrt{\Pi}\|_1 + \|\rho_2 - \sqrt{\Pi} \rho_2 \sqrt{\Pi}\|_1 \\ &\leq \sqrt{\text{Tr } \Pi} \cdot \|\sqrt{\Pi}(\rho_1 - \rho_2)\sqrt{\Pi}\|_2 + (2 \text{Tr } \rho_1) \sqrt{\epsilon_1} + (2 \text{Tr } \rho_2) \sqrt{\epsilon_2} \\ &\leq \sqrt{\text{Tr } \Pi} \cdot \|\rho_1 - \rho_2\|_2 + (2 \text{Tr } \rho_1) \sqrt{\epsilon_1} + (2 \text{Tr } \rho_2) \sqrt{\epsilon_2}. \end{aligned}$$

□

### 3.2 Projector smoothing of matrix $\|\cdot\|_\infty$ norm

As we will see below, our novel flattening lemmas reduce considerations about  $D_\infty(\alpha|\beta)$  to considerations about  $\|\alpha'\|_\infty$ , where  $\alpha'$  is a matrix related to  $\alpha, \beta$ . Accordingly, considerations about  $D_\infty^\epsilon(\alpha|\beta)$  reduce to considerations about a smoothed version of  $\|\alpha'\|_\infty$ . Naively speaking, the smoothed version of  $\|\alpha'\|_\infty$  that arises from this reduction replaces positive semidefinite matrix  $\alpha'$

by a smoothed positive semidefinite matrix  $\hat{\alpha}'$  close to it leading to the *state smoothed Schatten- $\ell_\infty$  norm*.

$$\|\alpha'\|_\infty^\epsilon := \min_{\hat{\alpha}'} \|\hat{\alpha}'\|_\infty,$$

where the minimisation is over all positive semidefinite matrices  $\hat{\alpha}'$  satisfying  $\|\alpha' - \hat{\alpha}'\|_1 \leq \epsilon(\text{Tr } \rho)$ . However for our purposes, as will become clear later, it is better to smooth  $\alpha'$  by applying a suitable orthogonal projector to it. We call this approach *projector smoothing* as opposed to *state smoothing*. The following proposition relates the two types of smoothing in the context of the Schatten- $\ell_\infty$  norm.

**Proposition 2** *Let  $0 < \epsilon < 1/4$ . Let  $\rho$  be a positive semidefinite matrix on a Hilbert space. Let  $\rho'$  be a positive semidefinite matrix in the same space achieving the optimum in the definition of  $\|\rho\|_\infty^\epsilon$ . Then there exists an orthogonal projector in the same space such that*

$$\Pi\rho = \rho\Pi, \quad \text{Tr } [\Pi\rho] \geq (1 - \sqrt{\epsilon})(\text{Tr } \rho), \quad \|\Pi\rho\Pi\|_\infty \leq (1 + 2\sqrt{\epsilon})\|\rho'\|_\infty = (1 + 2\sqrt{\epsilon})\|\rho\|_\infty^\epsilon.$$

**Proof:** Consider the matrices  $\rho, \rho'$  expressed in the eigenbasis  $\{|i\rangle\}_i$  of  $\rho$ . Define the subset

$$\text{Bad} := \{i : \rho(i, i) > (1 + 2\sqrt{\epsilon})\rho'(i, i)\}.$$

Then,

$$\begin{aligned} \epsilon(\text{Tr } \rho) &\geq \|\rho - \rho'\|_1 \geq \sum_{i \in \text{Bad}} (\rho(i, i) - \rho'(i, i)) \geq \sum_{i \in \text{Bad}} \rho(i, i) \left(1 - \frac{1}{1 + 2\sqrt{\epsilon}}\right) \\ &= \frac{2\sqrt{\epsilon}}{1 + 2\sqrt{\epsilon}} \sum_{i \in \text{Bad}} \rho(i, i), \\ \implies \sum_{i \in \text{Bad}} \rho(i, i) &\leq \frac{(1 + 2\sqrt{\epsilon})\sqrt{\epsilon}}{2} (\text{Tr } \rho) < \sqrt{\epsilon}(\text{Tr } \rho). \end{aligned}$$

Let  $\Pi$  be the projector in the eigenbasis of  $\rho$  that exactly skips  $i \in \text{Bad}$ . Then,  $\Pi\rho = \rho\Pi$ ,  $\text{Tr } [\Pi\rho] > (1 - \sqrt{\epsilon})(\text{Tr } \rho)$ , and

$$\|\Pi\rho\Pi\|_\infty = \max_{i \notin \text{Bad}} \rho(i, i) \leq \max_{i \notin \text{Bad}} (1 + 2\sqrt{\epsilon})\rho'(i, i) \leq (1 + 2\sqrt{\epsilon})\|\rho'\|_\infty.$$

This completes the proof of the proposition.  $\square$

### 3.3 From matrix $\|\cdot\|_2$ norm to inner product

A crucial technique required in the proof of our smooth multipartite convex split lemma is a technique to convert the Schatten- $\ell_2$  norm of a positive semidefinite matrix into a Hilbert-Schmidt inner product of two related positive semidefinite matrices.

**Lemma 1** *Let  $0 < \epsilon < 1/4$ . Suppose  $\rho_1, \rho_2$  are positive semidefinite operators on the same Hilbert space. Let  $\|\rho_1 - \rho_2\|_1 \leq \epsilon$ . Then there exists a positive semidefinite matrix  $\rho'_1$  on the same space such that*

$$\rho'_1 \leq \rho_1, \quad \|\rho'_1 - \rho_1\|_1 < \sqrt{\epsilon}, \quad \|\rho'_1\|_2^2 \leq (1 + 2\sqrt{\epsilon}) \text{Tr } [\rho_1 \rho_2].$$

**Proof:** We follow the lines of the proof of Proposition 2, and work in the eigenbasis  $\{|i\rangle\}_i$  of  $\rho_1$ . Define the subset

$$\text{Bad} := \{i : \rho_1(i, i) > (1 + 2\sqrt{\epsilon})\rho_2(i, i)\} \implies \sum_{i \in \text{Bad}} \rho_1(i, i) < \sqrt{\epsilon}(\text{Tr } \rho).$$

Define

$$\rho'_1 := \sum_{i \notin \text{Bad}} \rho_1(i, i) |i\rangle\langle i| \implies \rho'_1 \leq \rho_1, \quad \|\rho_1 - \rho'_1\|_1 < \sqrt{\epsilon}(\text{Tr } \rho).$$

Now,

$$\begin{aligned} \|\rho'_1\|_2^2 &= \text{Tr}[(\rho'_1)^2] = \sum_{i \notin \text{Bad}} \rho_1(i, i)^2 \\ &\leq (1 + 2\sqrt{\epsilon}) \sum_{i \notin \text{Bad}} \rho_1(i, i)\rho_2(i, i) \leq (1 + 2\sqrt{\epsilon}) \sum_i \rho_1(i, i)\rho_2(i, i) \\ &= (1 + 2\sqrt{\epsilon}) \text{Tr}[\rho_1\rho_2]. \end{aligned}$$

This completes the proof of the lemma.  $\square$

### 3.4 Tilting and augmentation smoothing

In this subsection, we state some results about *tilting* and *augmentation smoothing* of quantum states from [Sen21b], adapted to the specific requirements of this paper.

Consider a normalised density matrix  $\rho^{XYM}$ . Introduce new Hilbert spaces  $L_X$ ,  $L_Y$  of sufficiently large (how large will be clarified below) and equal dimension  $L$ . The *augmented state* is defined as

$$\rho^{(L_X X)(L_Y Y)M} := \frac{\mathbb{1}^{L_X}}{L} \otimes \frac{\mathbb{1}^{L_Y}}{L} \otimes \rho^{XYM}.$$

As will become clear in Lemma 4 below, the aim of augmentation is to reduce the Schatten- $\ell_\infty$  norm of  $\rho^{L_X X}$ ,  $\rho^{L_Y Y}$ , which will play a crucial role in a *smoothing property* to be stated below.

Introduce orthogonal copies of the Hilbert space  $M$  which we will label as  $\{M(l_x)\}_{l_x \in [L]}$  and  $\{M(l_y)\}_{l_y \in [L]}$ . The orthogonal direct sum of  $M$  together with all its copies is denoted by  $\hat{M}$  i.e.

$$\hat{M} := M \oplus \left( \bigoplus_{l_x \in [L]} M(l_x) \right) \oplus \left( \bigoplus_{l_y \in [L]} M(l_y) \right) \cong M \oplus (M \otimes L_X) \oplus (M \otimes L_Y).$$

We interchangeably think of the augmented state as residing in  $L_X X L_Y Y M$  when we will write  $\rho^{(L_X X)(L_Y Y)M}$ , or as residing in  $L_X X L_Y Y \hat{M}$  when we will write  $\rho^{(L_X X)(L_Y Y)\hat{M}}$ .

Let  $0 < \epsilon < 1$ . Let  $|m\rangle^M$  denote a computational basis state of  $M$ . For any fixed  $l_x \in [L]$ ,  $l_y \in [L]$ , the corresponding *tilting isometry* embedding  $M$  into  $\hat{M}$  is defined as

$$T_{l_x, l_y, \epsilon}^{M \rightarrow \hat{M}}(|m\rangle^M) := \sqrt{1 - 2\epsilon}|m\rangle^M + \sqrt{\epsilon}|m(l_x)\rangle^{M(l_x)} + \sqrt{\epsilon}|m(l_y)\rangle^{M(l_y)},$$

with the extension to the entire vector space  $M$  by linearity. Intuitively, tilting a vector means that we rotate it towards an orthogonal direction by a small precise amount.

We next define some other tilting maps which are subnormally scaled isometric embeddings of  $M$  into  $\hat{M}$ . For example,

$$T_{l_x, \epsilon}^{M \rightarrow \hat{M}}(|m\rangle^M) := \sqrt{1 - 2\epsilon}|m\rangle^M + \sqrt{\epsilon}|m(l_x)\rangle^{M(l_x)}.$$

Similarly, we define  $T_{l_y, \epsilon}^{M \rightarrow \hat{M}}$ . The last two of them are isometric embeddings of  $M$  into  $\hat{M}$  with scale factor  $\sqrt{1 - \epsilon}$ .

Define another tilting map in the analogous fashion:

$$T_{\epsilon}^{L_X L_Y M \rightarrow L_X L_Y \hat{M}}(|l_x\rangle^{L_X} |l_y\rangle^{L_Y} |m\rangle^M) := |l_x\rangle^{L_X} |l_y\rangle^{L_Y} \otimes T_{l_x, l_y, \epsilon}^{M \rightarrow \hat{M}}(|m\rangle^M).$$

The tilting maps  $T_{\epsilon}^{L_X M \rightarrow L_X \hat{M}}$ ,  $T_{\epsilon}^{L_Y M \rightarrow L_Y \hat{M}}$  are defined similarly.

Finally, we define the tilting map:

$$T_{\epsilon}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} := \mathbb{I}^X \otimes \mathbb{I}^Y \otimes T_{\epsilon}^{L_X L_Y M \rightarrow L_X L_Y \hat{M}}.$$

The tilting maps  $T_{\epsilon}^{L_X X M \rightarrow L_X X \hat{M}}$ ,  $T_{\epsilon}^{L_Y Y M \rightarrow L_Y Y \hat{M}}$  are then defined similarly. We remark that all the above tilting maps are isometric embeddings with scale factors  $\sqrt{1 - \epsilon}$  or 1. Also, all the tilting maps defined above can be extended from spaces of vectors to spaces of Hermitian matrices in the natural fashion, becoming superoperators if need be.

We remark that the definition of tilting given above is slightly simpler than the definition in [Sen21b]. This simplification makes the proofs of subsequent statements much shorter; nevertheless it is still strong enough for all known applications.

Let  $\Pi^{XYM}(1)$ ,  $\Pi^{XYM}(2)$ ,  $\Pi^{XYM}(3)$  be certain orthogonal projectors in  $XYM$  satisfying

$$\text{Tr} [\Pi^{XYM}(i) \rho^{XYM}] \geq 1 - \sqrt{\epsilon}, \quad \forall i \in [3]. \quad (7)$$

In the context of smooth multipartite convex split, these projectors will be related to optimising states in the definitions of certain smooth entropic quantities. We note that, in general, there is no relation amongst them.

Define the  $\epsilon$ -tilted approximate intersection of the projectors  $\Pi^{XYM}(1)$ ,  $\Pi^{XYM}(2)$ ,  $\Pi^{XYM}(3)$  as follows:

$$\begin{aligned} \hat{\Pi}^{L_X X L_Y Y \hat{M}} &:= \mathbb{1}^{L_X X L_Y Y \hat{M}} - \\ &\quad \text{span} \{ (T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}} \otimes \mathbb{I}^{L_Y Y}) (\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{XYM} - \Pi^{XYM}(1))), \\ &\quad (T_{\epsilon^{1/4}}^{L_Y Y M \rightarrow L_Y Y \hat{M}} \otimes \mathbb{I}^{L_X X}) (\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{XYM} - \Pi^{XYM}(2))), \\ &\quad T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{XYM} - \Pi^{XYM}(3))) \}, \end{aligned} \quad (8)$$

where the notation ‘span’ above means that we take the orthogonal projection onto the vector space span of the supports of the operators in the expression. Intuitively, we are just defining intersection via de Morgan’s rule i.e. intersection is the complement of the union of complements. Complement of a subspace, or rather of the orthogonal projection onto that subspace, has a precise meaning in terms of matrices. However, there is a problem defining the union of subspaces as explained in more detail in [Sen21b]; naively taking the span of subspaces often results in a catastrophe as it can easily be the entire Hilbert space! So, we need the more sophisticated approach of [Sen21b] viz. tilt the subspaces into orthogonal directions, then take their span, and then take the complement.

Following [Sen21b] again, we now define a normalised followed by a subnormalised density matrix. Both of them are close to the augmented normalised density matrix  $\rho^{L_X X L_Y Y \hat{M}}$ .

$$\begin{aligned}\hat{\rho}^{L_X X L_Y Y \hat{M}} &:= T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}}(\rho^{L_X X L_Y Y M}), \\ \hat{\hat{\rho}}^{L_X X L_Y Y \hat{M}} &:= \hat{\Pi}^{L_X X L_Y Y \hat{M}} \circ \rho^{L_X X L_Y Y \hat{M}}.\end{aligned}\tag{9}$$

These two states satisfy the following properties.

**Lemma 2**  $\|\hat{\rho}^{L_X X L_Y Y \hat{M}} - \rho^{L_X X L_Y Y \hat{M}}\|_1 = 2\sqrt{2} \cdot \epsilon^{1/8}$ .

**Proof:** Easily follows from the definition of the tilting isometry.  $\square$

**Lemma 3**  $\text{Tr} [\hat{\rho}^{L_X X L_Y Y \hat{M}}] \geq 1 - 121\epsilon^{1/4}$ .

**Proof:(Sketch)** Easily follows from Equation 7 and Proposition 3 of [Sen21b] regarding the ‘union’ properties of the so-called *matrix tilted span*.  $\square$

**Corollary 1**  $\|\hat{\rho}^{L_X X L_Y Y \hat{M}} - \rho^{L_X X L_Y Y \hat{M}}\|_1 < 22 \cdot \epsilon^{1/8}$ .

**Proof:** Follows by applying Fact 1 to Lemma 3.  $\square$

**Corollary 2**  $\|\hat{\hat{\rho}}^{L_X X L_Y Y \hat{M}} - \hat{\rho}^{L_X X L_Y Y \hat{M}}\|_1 < 25 \cdot \epsilon^{1/8}$ .

**Proof:** Triangle inequality and Lemma 2 and Corollary 1.  $\square$

We now state an important lemma that follows from the smoothing properties of a tilted augmented state shown in [Sen21b]. It essentially states that tracing out  $L_X$  removes the tilt in  $\hat{M}$  along  $l_x$  in terms of the Schatten  $\|\cdot\|_\infty$  norm, and similarly for tracing out  $L_Y$ .

**Lemma 4** *For any  $l_x, l_y \in [L]$  and positive semidefinite matrix  $\sigma^M$ ,*

$$\begin{aligned}\|L^{-1} \sum_{l'_y \in [L]} T_{l_x, l'_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\sigma^M) - T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\sigma^M)\|_\infty &< 4\epsilon^{1/8} L^{-1/2} (\text{Tr } \sigma), \\ \|L^{-1} \sum_{l'_x \in [L]} T_{l'_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\sigma^M) - T_{l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\sigma^M)\|_\infty &< 4\epsilon^{1/8} L^{-1/2} (\text{Tr } \sigma), \\ \|L^{-2} \sum_{l'_x, l'_y \in [L]} T_{l'_x, l'_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\sigma^M) - \sigma^M\|_\infty &< 8\epsilon^{1/8} L^{-1/2} (\text{Tr } \sigma).\end{aligned}$$

*As a consequence, for any augmented normalised density matrix  $\rho^{L_X X L_Y Y M}$ ,*

$$\begin{aligned}\|\hat{\rho}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})\|_\infty &< 4\epsilon^{1/8} L^{-3/2}, \\ \|\hat{\rho}^{L_Y Y \hat{M}} - T_{\epsilon^{1/4}}^{L_Y Y M \rightarrow L_Y Y \hat{M}}(\rho^{L_Y Y M})\|_\infty &< 4\epsilon^{1/8} L^{-3/2}, \\ \|\hat{\hat{\rho}}^{\hat{M}} - \rho^M\|_\infty &< 8\epsilon^{1/8} L^{-1/2}.\end{aligned}$$

**Proof:** Follows from Proposition 4 of [Sen21b].  $\square$

### 3.5 Flattening quantum states

In this subsection, we shall show that a natural quantum operation called *flattening*, aka the ‘brother construction’, actually preserves the fidelity between quantum states. The *brother construction* is a well known argument in classical probability theory that is sometimes used to flatten a specific probability distribution by enlarging its support. It works as follows: Suppose  $p^X$  is a probability distribution on the alphabet  $X$ . Assume for simplicity that probability of each symbol in  $X$  is a rational number with denominator  $F_p$ . We note that taking  $F_p$  to be large enough approximates  $p^X$  quite well; this point will be made more precise (and also quantum!) when we actually state our technical lemmas below. So for now we assume that  $p(x) = \frac{a_x}{F_p}$  for all  $x \in X$ . We take a new alphabet  $L$  large enough, and define a probability distribution  $p^{XL}$  on  $XL$  as follows:

$$p^{XL}(x, l) := \begin{cases} \frac{1}{F_p} & l \leq a_x, \\ 0 & l > a_x. \end{cases}$$

It is now easy to see that if  $p^{XL}(x, l) \neq 0$  for some  $(x, l)$ , then  $p^{XL}(x, l) = \frac{1}{F_p}$ . In other words,  $p^{XL}$  is *flat on its support*. For a fixed  $x$ , the set of pairs  $\{(x, l) : l \leq a_x\}$  are called the ‘brothers’ of  $x$ .

Note that the above brothers construction was tailored for a specific probability distribution  $p^X$ . It will flatten only  $p^X$ . A lesser known fact about the construction is that flattening according to  $p^X$  nevertheless preserves the  $\ell_1$ -distance between any two probability distributions  $q_1^X, q_2^X$ . Flattening sometimes leads to cleaner proofs in classical probability theory. As we will see later in our proof of our smooth multipartite convex split lemma, the quantum analogue of flattening plays a crucial role in overcoming bottleneck due to non-commutativity of certain operators involved in the proof.

Flattening ideas have been used in quantum information theory before. Anshu and Jain [AJ22] take inspiration from the classical brothers construction and defines a unitary quantum flattening operation. However in order to define their unitary, they need to add a small ancilla in addition to the ‘brother’ Hilbert space. That ancilla is initialised with a suitable *embezzling pure state*, which is almost unchanged at the end of the unitary operation. Anshu and Jain use this unitary in the context of one of their more economical convex split lemmas. The extra ancilla containing the embezzling state does not hurt their application. However, such an ancilla creates serious problems in our attempt to flatten a quantum state in order overcome the non-commutativity bottleneck amongst certain operators in our smooth multipartite convex split proof. Hence, we cannot use the unitary flattening operation of [AJ22] in this paper.

It turns out that our final solution for this problem is even simpler than the unitary flattening operation of [AJ22]. We take the vanilla brothers construction outlined above and convert it in the most naive fashion into a CPTP superoperator acting on quantum states. Our flattening superoperator is not a unitary operation. It does not use embezzling states. As expected it is tailored to flatten a specific density matrix  $\sigma^A$  on the Hilbert space  $A$ . Serendipitously, it preserves the fidelity between any two quantum states  $\rho_1^A, \rho_2^A$ , though not the trace distance. Nevertheless, this novel observation suffices to meet all our requirements. We consider this simple but powerful operation to be a major conceptual contribution of this work.

We now describe our quantum flattening operation. Let  $\sigma^A$  be a positive semidefinite operator on the Hilbert space  $A$ . Let its eigenvalues and eigenvectors be  $\{(\sigma(a, a), |a\rangle^A)\}_a$ . Let  $0 < \delta < 1$ . Let  $F_\sigma$  be the smallest positive integer such that there exist integers  $\{f(a)\}_a$  with the property

that

$$\forall a : \sigma(a, a) \leq \frac{f(a)}{F_\sigma} \leq \sigma(a, a)(1 + \delta).$$

Then the *flattening superoperator with respect to  $\sigma^A$*  is defined as

$$\mathcal{F}_\sigma^{A \rightarrow AL}(\rho^A) := \sum_{l=1}^L A_l^{A \rightarrow AL}(\rho^A)(A_l^\dagger)^{AL \rightarrow A},$$

where  $\rho^A$  is any Hermitian matrix on  $A$ ,  $L := \max_a f(a)$  and for each  $l \in [L]$ ,  $A_l$  is a linear map from the Hilbert space  $A$  to the Hilbert space  $AL := A \otimes \mathbb{C}^L$  defined as:

$$A_l^{A \rightarrow AL}(|a\rangle^A) := \begin{cases} \frac{1}{\sqrt{f(a)}} |a\rangle^A |l\rangle^L & l \leq f(a), \\ 0 & l > f(a), \end{cases}$$

with the extension to the full vector space  $A$  by linearity.

To get a better geometric picture of  $\mathcal{F}_\sigma^{A \rightarrow AL}$ , it helps to order the eigenvectors of  $\sigma^A$  as  $\{|a_1\rangle, |a_2\rangle, \dots\}$  in non-decreasing order of  $f(a)$ . Introduce new notation  $A(1) \equiv A$ . Consider fresh copies  $A(l)$ ,  $2 \leq l \leq L$  of the Hilbert space  $A$ . We can then write

$$AL = A \otimes \mathbb{C}^L = \bigoplus_{l=1}^L A(l).$$

Define  $f(a_0) \equiv 0$ . For  $i \in [|A|]$ , define  $A_{(i)} := \text{span} \{|a_i\rangle, |a_{i+1}\rangle, \dots, |a_{|A|}\rangle\}$ . For  $i \in [A]$  and  $l \in [L]$ , define  $T_{i,l}$  to be the identity isometric embedding of  $A_{(i)}$  into  $A(l)$ . Depending on the context,  $T_{i,l}$  can also be thought of as the identity superoperator mapping matrices on  $A_{(i)}$  to matrices on  $A(l)$ . Then,

$$\mathcal{F}_\sigma^{A \rightarrow AL} = \bigoplus_{i=1}^{|A|} \bigoplus_{l=f(a_{i-1})+1}^{f(a_i)} T_{i,l}^{A_{(i)} \rightarrow A(l)},$$

i.e.,  $\mathcal{F}_\sigma^{A \rightarrow AL}$  is an orthogonal direct sum of projection superoperators.

We now prove the following lemma.

**Lemma 5** *The superoperator  $\mathcal{F}_\sigma^{A \rightarrow AL}$  is completely positive and trace preserving.*

**Proof:** The Kraus representation of  $\mathcal{F}_\sigma^{A \rightarrow AL}$  given above immediately implies it is completely positive. To check the trace preserving property, we work in the basis of  $A$  given by the eigenvectors of  $\sigma^A$  and obtain,

$$\begin{aligned} \sum_{l=1}^L A_l^\dagger A_l(a, a') &= \langle a | \left( \sum_{l=1}^L A_l^\dagger A_l \right) | a' \rangle = \sum_{l=1}^{\min\{f(a'), f(a)\}} \frac{1}{\sqrt{f(a')f(a)}} \langle a, l | a', l \rangle \\ &= \delta_{aa'} \sum_{l=1}^{f(a)} \frac{1}{f(a)} = \delta_{aa'}, \\ \implies \sum_{l=1}^L A_l^\dagger A_l(a, a') &= \mathbb{1}^A. \end{aligned}$$



This completes the proof of the lemma.  $\square$

Define the subspace  $F'_\sigma$  of  $AL$  by

$$F'_\sigma := \text{span} \{ |a\rangle^A |l\rangle^L : l \leq f(a) \}.$$

In other words, the image of  $\mathcal{F}_\sigma^{A \rightarrow AL}$  is contained in the span of Hermitian operators with support in  $F'_\sigma$ . The next lemma shows that the dimension of  $F'_\sigma$  is approximately  $F_\sigma(\text{Tr } \sigma)$ .

**Lemma 6**  $(\text{Tr } \sigma)F_\sigma \leq |F'_\sigma| \leq (1 + \delta)(\text{Tr } \sigma)F_\sigma$ .

**Proof:**

$$\begin{aligned} |F'_\sigma| &= \sum_a f(a) = F_\sigma \sum_a \frac{f(a)}{F_\sigma}, \\ F_\sigma \sum_a \sigma(a, a) &\leq F_\sigma \sum_a \frac{f(a)}{F_\sigma} \leq F_\sigma \sum_a \sigma(a, a)(1 + \delta) \\ \implies F_\sigma(\text{Tr } \sigma) &\leq |F'_\sigma| \leq (1 + \delta)F_\sigma(\text{Tr } \sigma). \end{aligned}$$

This completes the proof of the lemma.  $\square$

We are now finally in a position to prove that  $\mathcal{F}_\sigma^{A \rightarrow AL}$  flattens  $\sigma^A$ .

**Proposition 3** *Let  $\sigma^A$  be a positive semidefinite operator on  $A$ . Then,*

$$(1 + \delta)^{-1} \frac{\mathbb{1}^{F'_\sigma}}{F_\sigma} \leq \mathcal{F}_\sigma^{A \rightarrow AL}(\sigma) \leq \frac{\mathbb{1}^{F'_\sigma}}{F_\sigma}.$$

**Proof:**

$$\mathcal{F}_\sigma^{A \rightarrow AL}(\sigma) = \sum_a \sigma(a, a) \mathcal{F}_\sigma^{A \rightarrow AL}(|a\rangle\langle a|) = \sum_a \sigma(a, a) \sum_{l=1}^L A_l |a\rangle\langle a| A_l^\dagger = \sum_a \frac{\sigma(a, a)}{f(a)} |a\rangle\langle a|^A \sum_{l=1}^{f(a)} |l\rangle\langle l|^L.$$

But,

$$\begin{aligned} \sigma(a, a) &\leq \frac{f(a)}{F_\sigma} \leq (1 + \delta)\sigma(a, a) \\ \implies \frac{(1 + \delta)^{-1}}{F_\sigma} &\leq \frac{\sigma(a, a)}{f(a)} \leq \frac{1}{F_\sigma}. \end{aligned}$$

This implies

$$\frac{(1 + \delta)^{-1}}{F_\sigma} \sum_a |a\rangle\langle a|^L \sum_{l=1}^{f(a)} |l\rangle\langle l|^L \leq \mathcal{F}_\sigma^{A \rightarrow AL}(\sigma^A) \leq \frac{1}{F_\sigma} \sum_a |a\rangle\langle a|^L \sum_{l=1}^{f(a)} |l\rangle\langle l|^L.$$

But,  $\sum_a |a\rangle\langle a|^L \sum_{l=1}^{f(a)} |l\rangle\langle l|^L = \mathbb{1}^{F'_\sigma}$ . This completes the proof of the proposition.  $\square$

We now show the important property that flattening does not change the length preserving fidelity.

**Proposition 4** Let  $\sigma^A$ ,  $\rho_1^{AB}$ ,  $\rho_2^{AB}$  be positive semidefinite operators on their respective Hilbert spaces. Then,

$$F((\mathcal{F}_\sigma^{A \rightarrow LA} \otimes \mathbb{I}^B)(\rho_1^{AB}), (\mathcal{F}_\sigma^{A \rightarrow LA} \otimes \mathbb{I}^B)(\rho_2^{AB})) = F(\rho_1^{AB}, \rho_2^{AB}).$$

**Proof:** Consider the following unitary Stinespring dilation  $U^{A \rightarrow ALL'}$  of  $\mathcal{F}_\sigma^{A \rightarrow AL}$ , where  $ALL' := A \otimes \mathbb{C}^L \otimes \mathbb{C}^L$ .

$$U : |a\rangle^A \mapsto |a\rangle^A \otimes \left( \frac{1}{\sqrt{f(a)}} \sum_{l=1}^{f(a)} |l\rangle^L |l\rangle^{L'} \right).$$

Define for  $i = 1, 2$ ,  $\sigma_i^{L'LAB} := (U^{A \rightarrow ALL'} \otimes \mathbb{I}^B) \circ \rho_i^{AB}$ . Since unitary superoperators preserve fidelity, we have

$$F(\rho_1^{AB}, \rho_2^{AB}) = F(\sigma_1^{L'LAB}, \sigma_2^{L'LAB}).$$

Let  $|\sigma_1\rangle^{L'LABR}$ ,  $|\sigma_2\rangle^{L'LABR}$  be length preserving purifications of  $\sigma_1^{L'LAB}$ ,  $\sigma_2^{L'LAB}$  achieving their fidelity i.e.

$$F(\sigma_1^{L'LAB}, \sigma_2^{L'LAB}) = F(\sigma_1^{L'LABR}, \sigma_2^{L'LABR}) = \langle \sigma_1 | \sigma_2 \rangle.$$

Then,

$$|\sigma_1\rangle^{L'LABR} = \sum_{l=1}^L |l\rangle^{L'} |l\rangle^L |v_1(l)\rangle^{ABR}, \quad |\sigma_2\rangle^{L'LABR} = \sum_{l=1}^L |l\rangle^{L'} |l\rangle^L |v_2(l)\rangle^{ABR},$$

where  $\{|v_1(l)\rangle^{ABR}\}_l$ ,  $\{|v_2(l)\rangle^{ABR}\}_l$  are in general unnormalised, non-orthogonal pure states.

By the definition of the flattening operator  $\mathcal{F}_\sigma$ ,  $\sigma_i^{LAB} := (\mathcal{F}_\sigma^{A \rightarrow AL} \otimes \mathbb{I}^B)(\rho_i^{AB})$ ,  $i = 1, 2$  have an orthogonal direct sum structure indexed by  $l \in L$  i.e.

$$\sigma_i^{LAB} = \bigoplus_{l=1}^L |l\rangle\langle l|^L \otimes \sigma_i(l)^{AB},$$

for some positive semidefinite operators  $\sigma_i(l)^{AB}$ . In other words,  $\sigma_1^{LAB}$ ,  $\sigma_2^{LAB}$  are classical on  $L$ . As a result,  $\| |v_i(l)\rangle^{ABR} \|_2^2 = \text{Tr} [\sigma_i(l)^{AB}]$ , and  $|v_i(l)\rangle^{ABR}$  is a length preserving purification of  $\sigma_i(l)^{AB}$ . Moreover,

$$F(\sigma_1(l)^{AB}, \sigma_2(l)^{AB}) = \langle v_1(l) | v_2(l) \rangle.$$

Otherwise, the maximum overlap of length preserving purifications required by the fidelity of positive semidefinite operators  $\sigma_1^{LAB}$ ,  $\sigma_2^{LAB}$  will not be achieved.

We now investigate the fidelity of positive operators  $\sigma_1^{LAB}$  and  $\sigma_2^{LAB}$ . Take  $|\sigma_1\rangle^{L'LABR}$  as the purification of  $\sigma_1^{LAB}$ . Let  $|\sigma'_2\rangle^{L'LABR}$  be a fidelity achieving length preserving purification of  $\sigma_2^{LAB}$ . Since the flattening operation is trace preserving, we have  $\text{Tr} [\sigma_i^{LAB}] = \text{Tr} [\rho_i^{AB}]$ ,  $i = 1, 2$ . Let

$$|\sigma'_2\rangle^{L'LABR} = \sum_{l', l=1}^L |l'\rangle^{L'} |l\rangle^L |v_2(l', l)\rangle^{ABR},$$

Then the unnormalised pure state

$$|\sigma'_2(l)\rangle^{L'ABR} := \sum_{l'=1}^L |l'\rangle^{L'} |v_2(l', l)\rangle^{ABR}$$

is a length preserving purification of  $\sigma_2(l)^{AB}$  i.e.  $\|\sigma'_2(l)\rangle^{L'ABR}\|_2^2 = \text{Tr}[\sigma_2(l)^{AB}]$ . One way to maximise the overlap between  $|\sigma_1\rangle^{L'LABR}$  and  $|\sigma'_2\rangle^{L'LABR}$  is to set  $|v'_2(l', l)\rangle = 0$  if  $l' \neq l$  and  $|v'_2(l, l)\rangle = |v_2(l)\rangle$ . In other words, we can take  $|\sigma'_2\rangle^{L'LABR} = |\sigma_2\rangle^{L'LABR}$ .

Thus,

$$F(\sigma_1^{LAB}, \sigma_2^{LAB}) = \langle \sigma_1 | \sigma_2 \rangle = F(\sigma_1^{L'LAB}, \sigma_2^{L'LAB}) = F(\rho_1^{AB}, \rho_2^{AB}),$$

completing the proof of the proposition.  $\square$

## 4 Fully smooth multipartite convex split via flattening

In this section we prove a smooth multipartite convex split lemma, Lemma 7, using flattening, tilting and augmentation smoothing. At first glance, Lemma 7 looks like another version of the smooth multipartite convex split lemma proved in [Sen24], with the main difference being that Lemma 7 states its inner bounds stated in terms of  $D_\infty^\epsilon$  instead of  $D_2^\epsilon$  in [Sen24]. One may wonder about the need of giving another proof of smooth multipartite convex split, and that too a much longer and more complicated proof. There seems to be no advantage in doing so; the  $D_\infty^\epsilon$  formulation is not any stronger. In fact it is likely slightly worse than the  $D_2^\epsilon$  formulation. However, this exercise paves the way to prove a non pairwise independent smooth multipartite soft covering lemma for classical quantum states in the next section. This is where our more sophisticated machinery wins out as proving such a result is beyond the reach of telescoping. We will assume in this section, without loss of generality, that the density matrix  $\rho^{XYM}$  in the statement of convex split is normalised, by dividing by  $\text{Tr } \rho$ .

We first state the version of the smooth multipartite convex split lemma proved via flattening, tilting and augmentation smoothing.

**Lemma 7** *Let  $\rho^{XYM}$  be a subnormalised density matrix. Let  $\alpha^X, \beta^Y$  be normalised density matrices such that  $\text{supp}(\rho^X) \leq \text{supp}(\alpha^X)$  and  $\text{supp}(\rho^Y) \leq \text{supp}(\beta^Y)$ . Define the  $A$ -fold tensor product states  $\alpha^{X^A} := (\alpha^X)^{\otimes A}, \beta^{Y^B} := (\beta^Y)^{\otimes B}$ , and the  $(A-1)$ -fold tensor product states  $\alpha^{X^{-a}}, \beta^{Y^{-b}}$  for any  $a \in [A], b \in [B]$ . Define the convex split state*

$$\sigma^{X^A Y^B M} := (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \rho^{X_a Y_b M} \otimes \alpha^{X^{-a}} \otimes \beta^{Y^{-b}},$$

and the fully decoupled state

$$\tau^{X^A Y^B M} := \alpha^{X^A} \otimes \beta^{Y^B} \otimes \rho^M.$$

Suppose

$$\begin{aligned} \log A &> D_\infty^\epsilon(\rho^{XM} \| \alpha^X \otimes \rho^M) + \log \epsilon^{-1/20}, \\ \log B &> D_\infty^\epsilon(\rho^{YM} \| \beta^Y \otimes \rho^M) + \log \epsilon^{-1/20}, \\ \log A + \log B &> D_\infty^\epsilon(\rho^{XYM} \| \alpha^X \otimes \beta^Y \otimes \rho^M) + \log \epsilon^{-1/20}. \end{aligned}$$

Then,  $\|\sigma^{X^A Y^B M} - \tau^{X^A Y^B M}\|_1 < 10\epsilon^{1/64}(\text{Tr } \rho)$ .

We now proceed with the proof of Lemma 7. From the definitions of smooth  $D_\infty^\epsilon(\cdot \| \cdot)$ , we have subnormalised density matrices  $\rho^{XYM}(3)$ ,  $\rho^{XM}(1)$  and  $\rho^{YM}(2)$  satisfying

$$\|\rho^{XYM} - \rho^{XYM}(3)\|_1 \leq \epsilon, \quad \|\rho^{XM} - \rho^{XM}(1)\|_1 \leq \epsilon, \quad \|\rho^{YM} - \rho^{YM}(2)\|_1 \leq \epsilon, \quad (10)$$

such that

$$\begin{aligned}\rho^{XYM}(3) &\leq 2^{D_\infty^\epsilon(\rho^{XYM}\|\alpha^X\otimes\beta^Y\otimes\rho^M)}(\alpha^X\otimes\beta^Y\otimes\rho^M), \\ \rho^{XM}(1) &\leq 2^{D_\infty^\epsilon(\rho^{XM}\|\alpha^X\otimes\rho^M)}(\alpha^X\otimes\rho^M), \\ \rho^{YM}(2) &\leq 2^{D_\infty^\epsilon(\rho^{YM}\|\beta^Y\otimes\rho^M)}(\beta^Y\otimes\rho^M).\end{aligned}\tag{11}$$

Apply the flattening CPTP superoperator to get the normalised density matrix

$$\rho^{L_X X L_Y Y L_M M} := (\mathcal{F}_\alpha^{X \rightarrow L_X X} \otimes \mathcal{F}_\beta^{Y \rightarrow L_Y Y} \otimes \mathcal{F}_\rho^{M \rightarrow L_M M})(\rho^{XYM}).$$

Observe that

$$\rho^{L_X X L_M M} = (\mathcal{F}_\alpha^{X \rightarrow L_X X} \otimes \mathcal{F}_\rho^{M \rightarrow L_M M})(\rho^{XM}), \quad \rho^{L_Y Y L_M M} = (\mathcal{F}_\beta^{Y \rightarrow L_Y Y} \otimes \mathcal{F}_\rho^{M \rightarrow L_M M})(\rho^{YM}).$$

Define  $\rho^{L_X X L_Y Y L_M M}(3)$ ,  $\rho^{L_X X L_M M}(1)$ ,  $\rho^{L_Y Y L_M M}(2)$  via flattening in the natural fashion. Then we use Equation 10 and conclude,

$$\begin{aligned}\|\rho^{L_X X L_Y Y L_M M} - \rho^{L_X X L_Y Y L_M M}(3)\|_1 &< \epsilon, \\ \|\rho^{L_X X L_M M} - \rho^{L_X X L_M M}(1)\|_1 &< \epsilon, \quad \|\rho^{L_Y Y L_M M} - \rho^{L_Y Y L_M M}(2)\|_1 < \epsilon,\end{aligned}\tag{12}$$

since a CPTP superoperator cannot increase the trace distance.

By Lemma 6 and Proposition 3, we have

$$\begin{aligned}(1+\delta)^{-2} \frac{\mathbb{1}_{F'_\alpha}}{|F'_\alpha|} &\leq \mathcal{F}_\alpha^{X \rightarrow L_X X}(\alpha^X) \leq (1+\delta) \frac{\mathbb{1}_{F'_\alpha}}{|F'_\alpha|}, \\ (1+\delta)^{-2} \frac{\mathbb{1}_{F'_\beta}}{|F'_\beta|} &\leq \mathcal{F}_\beta^{Y \rightarrow L_Y Y}(\beta^Y) \leq (1+\delta) \frac{\mathbb{1}_{F'_\beta}}{|F'_\beta|}, \\ (1+\delta)^{-2} \frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|} \cdot \mathcal{F}_\rho^{M \rightarrow L_M M}(\rho^M) &\leq (1+\delta) \frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|}.\end{aligned}\tag{13}$$

Because a CPTP superoperator respects the Löwner partial order on Hermitian matrices, combining Equations 11 and 13 gives us

$$\begin{aligned}\rho^{L_X X L_Y Y L_M M}(3) &\leq (1+\delta)^3 2^{D_\infty^\epsilon(\rho^{XYM}\|\alpha^X\otimes\beta^Y\otimes\rho^M)} \frac{\mathbb{1}_{F'_\alpha} \otimes \mathbb{1}_{F'_\beta} \otimes \mathbb{1}_{F'_\rho}}{|F'_\alpha| |F'_\beta| |F'_\rho|}, \\ \rho^{L_X X L_M M}(1) &\leq (1+\delta)^2 2^{D_\infty^\epsilon(\rho^{XM}\|\alpha^X\otimes\rho^M)} \frac{\mathbb{1}_{F'_\alpha} \otimes \mathbb{1}_{F'_\rho}}{|F'_\alpha| |F'_\rho|}, \\ \rho^{L_Y Y L_M M}(2) &\leq (1+\delta)^2 2^{D_\infty^\epsilon(\rho^{YM}\|\beta^Y\otimes\rho^M)} \frac{\mathbb{1}_{F'_\beta} \otimes \mathbb{1}_{F'_\rho}}{|F'_\beta| |F'_\rho|}.\end{aligned}\tag{14}$$

In particular,

$$\text{supp}(\rho^{L_X X L_Y Y L_M M}(3)) \leq F'_\alpha \otimes F'_\beta \otimes F'_\rho, \quad \text{supp}(\rho^{L_X X L_M M}(1)) \leq F'_\alpha \otimes F'_\rho, \quad \text{supp}(\rho^{L_Y Y L_M M}(2)) \leq F'_\beta \otimes F'_\rho.$$

Applying Proposition 2 to Equations 12, 14 tells us that there exists orthogonal projectors  $\Pi^{L_X X L_Y Y L_M M}(3)$ ,  $\Pi^{L_X X L_M M}(1)$ ,  $\Pi^{L_Y Y L_M M}(2)$  such that

$$\begin{aligned}\text{Tr} [\Pi^{L_X X L_Y Y L_M M}(3) \rho^{L_X X L_Y Y L_M M}] &\geq 1 - \sqrt{\epsilon}, \\ \text{Tr} [\Pi^{L_X X L_M M}(1) \rho^{L_X X L_M M}] &\geq 1 - \sqrt{\epsilon}, \\ \text{Tr} [\Pi^{L_Y Y L_M M}(2) \rho^{L_Y Y L_M M}] &\geq 1 - \sqrt{\epsilon}, \\ \Pi^{L_X X L_Y Y L_M M}(3) \rho^{L_X X L_Y Y L_M M} &= \rho^{L_X X L_Y Y L_M M} \Pi^{L_X X L_Y Y L_M M}(3), \\ \Pi^{L_X X L_M M}(1) \rho^{L_X X L_M M} &= \rho^{L_X X L_M M} \Pi^{L_X X L_M M}(1), \\ \Pi^{L_Y Y L_M M}(2) \rho^{L_Y Y L_M M} &= \rho^{L_Y Y L_M M} \Pi^{L_Y Y L_M M}(2), \\ \|\Pi^{L_X X L_Y Y L_M M}(3) \circ \rho^{L_X X L_Y Y L_M M}\|_\infty &\leq \frac{(1+\delta)^3 (1+2\sqrt{\epsilon}) 2^{D_\infty^\epsilon(\rho^{XYM}\|\alpha^X\otimes\beta^Y\otimes\rho^M)}}{|F'_\alpha| |F'_\beta| |F'_\rho|}, \\ \|\Pi^{L_X X L_M M}(1) \circ \rho^{L_X X L_M M}\|_\infty &\leq \frac{(1+\delta)^2 (1+2\sqrt{\epsilon}) 2^{D_\infty^\epsilon(\rho^{XM}\|\alpha^X\otimes\rho^M)}}{|F'_\alpha| |F'_\rho|}, \\ \|\Pi^{L_Y Y L_M M}(2) \circ \rho^{L_Y Y L_M M}\|_\infty &\leq \frac{(1+\delta)^2 (1+2\sqrt{\epsilon}) 2^{D_\infty^\epsilon(\rho^{YM}\|\beta^Y\otimes\rho^M)}}{|F'_\beta| |F'_\rho|}.\end{aligned}\tag{15}$$

Propositions 3 and 4, and Lemma 6 taken together allow us to prove Lemma 8 below. Lemma 8 will be used as the first step in the proof of Lemma 7. We note that  $\delta$  can be taken as small as we please at the expense of increasing  $|F'_\sigma|$ . Essentially,  $\delta$  can be treated as a free parameter.

**Lemma 8** *Let  $A, B$  be positive integers. Let  $\alpha^X, \beta^Y, \rho^M, \sigma^{X^A Y^B M}$  be normalised density matrices. Define the normalised density matrix  $\tau^{X^A Y^B M} := (\alpha^X)^{\otimes A} \otimes (\beta^Y)^{\otimes B} \otimes \rho^M$ . Then,*

$$\begin{aligned}
& F(((\mathcal{F}_\alpha^{X \rightarrow L_X X})^{\otimes A} \otimes (\mathcal{F}_\beta^{Y \rightarrow L_Y Y})^{\otimes B} \otimes \mathcal{F}_\rho^{M \rightarrow L_M M})(\sigma^{X^A Y^B M}), \\
& \quad ((\mathcal{F}_\alpha^{X \rightarrow L_X X})^{\otimes A} \otimes (\mathcal{F}_\beta^{Y \rightarrow L_Y Y})^{\otimes B} \otimes \mathcal{F}_\rho^{M \rightarrow L_M M})(\tau^{X^A Y^B M})) \\
& = F(\sigma^{X^A Y^B M}, \tau^{X^A Y^B M}), \\
& (1 + \delta)^{-(A+B+1)} \left( \left( \frac{\mathbb{1}^{F'_\alpha}}{|F'_\alpha|} \right)^{\otimes A} \otimes \left( \frac{\mathbb{1}^{F'_\beta}}{|F'_\beta|} \right)^{\otimes B} \otimes \frac{\mathbb{1}^{F'_\rho}}{|F'_\rho|} \right) \\
& \leq ((\mathcal{F}_\alpha^{X \rightarrow L_X X})^{\otimes A} \otimes (\mathcal{F}_\beta^{Y \rightarrow L_Y Y})^{\otimes B} \otimes \mathcal{F}_\rho^{M \rightarrow L_M M})(\tau^{X^A Y^B M}) \\
& \leq (1 + \delta)^{A+B+1} \left( \left( \frac{\mathbb{1}^{F'_\alpha}}{|F'_\alpha|} \right)^{\otimes A} \otimes \left( \frac{\mathbb{1}^{F'_\beta}}{|F'_\beta|} \right)^{\otimes B} \otimes \frac{\mathbb{1}^{F'_\rho}}{|F'_\rho|} \right),
\end{aligned}$$

where  $L_X, L_Y, L_M$  are the additional Hilbert spaces required for the respective flattening superoperators.

In order to prepare for the proof of Lemma 7, we will assume that the states  $\sigma^{X^A Y^B M}$  and  $\tau^{X^A Y^B M}$  have been flattened as in Lemma 8, where the positive integers  $A, B$  come from the statement of Lemma 7. We will call the flattened states  $\sigma^{(L_X X)^A (L_Y Y)^B L_M M}$  and  $\tau^{(L_X X)^A (L_Y Y)^B L_M M}$ . We will choose the positive integers  $F_\alpha, F_\beta, F_\rho$  large enough, and correspondingly the dimensions of  $L_X, L_Y, L_M$  large enough so that  $\delta \leq \frac{\sqrt{\epsilon}}{2(A+B+1)}$ . Also to make the notation lighter, henceforth we will always work with the flattened states and the Hilbert spaces  $L_X X, L_Y Y$  and  $L_M M$ . Hence from now on we redefine  $X \equiv L_X X, Y \equiv L_Y Y$  and  $M \equiv L_M M$ . Under this redefinition, we restate Equation 15 as Equation 16 below. That is, we can assume that there are projectors  $\Pi^{XYM}(3)$ ,

$\Pi^{XM}(1)$  and  $\Pi^{YM}(2)$  such that

$$\begin{aligned}
\text{Tr} [\Pi^{XYM}(3)\rho^{XYM}] &\geq 1 - \sqrt{\epsilon}, \\
\text{Tr} [\Pi^{XM}(1)\rho^{XM}] &\geq 1 - \sqrt{\epsilon}, \\
\text{Tr} [\Pi^{YM}(2)\rho^{YM}] &\geq 1 - \sqrt{\epsilon}, \\
\Pi^{XYM}(3)\rho^{XYM} &= \rho^{XYM}\Pi^{XYM}(3), \\
\Pi^{XM}(1)\rho^{XM} &= \rho^{XM}\Pi^{XM}(1), \\
\Pi^{YM}(2)\rho^{YM} &= \rho^{YM}\Pi^{YM}(2), \\
\|\Pi^{XYM}(3)\rho^{XYM}\|_\infty &\leq \frac{(1+3\sqrt{\epsilon})2^{D_\infty^\epsilon(\rho^{XYM}\|\alpha^X\otimes\beta^Y\otimes\rho^M)}}{|F'_\alpha||F'_\beta||F'_\rho|}, \\
\|\Pi^{XM}(1)\rho^{XM}\|_\infty &\leq \frac{(1+3\sqrt{\epsilon})2^{D_\infty^\epsilon(\rho^{XM}\|\alpha^X\otimes\rho^M)}}{|F'_\alpha||F'_\rho|}, \\
\|\Pi^{YM}(2)\rho^{YM}\|_\infty &\leq \frac{(1+3\sqrt{\epsilon})2^{D_\infty^\epsilon(\rho^{YM}\|\beta^Y\otimes\rho^M)}}{|F'_\beta||F'_\rho|}, \\
F'_\alpha &\leq X, \quad F'_\beta \leq Y, \quad F'_\rho \leq M, \\
(1 - \sqrt{\epsilon})\frac{\mathbb{1}_{F'_\alpha}}{|F'_\alpha|} &\leq \alpha^X \leq (1 + \sqrt{\epsilon})\frac{\mathbb{1}_{F'_\alpha}}{|F'_\alpha|}, \\
(1 - \sqrt{\epsilon})\frac{\mathbb{1}_{F'_\beta}}{|F'_\beta|} &\leq \beta^Y \leq (1 + \sqrt{\epsilon})\frac{\mathbb{1}_{F'_\beta}}{|F'_\beta|}, \\
(1 - \sqrt{\epsilon})\frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|} &\leq \rho^M \leq (1 + \sqrt{\epsilon})\frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|}, \\
(1 - \sqrt{\epsilon})\frac{(\mathbb{1}_{F'_\alpha})^{\otimes A} \otimes (\mathbb{1}_{F'_\beta})^{\otimes B} \otimes (\mathbb{1}_{F'_\rho})}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|} &\leq \tau^{X^A Y^B M} \leq (1 + \sqrt{\epsilon})\frac{(\mathbb{1}_{F'_\alpha})^{\otimes A} \otimes (\mathbb{1}_{F'_\beta})^{\otimes B} \otimes (\mathbb{1}_{F'_\rho})}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|}.
\end{aligned} \tag{16}$$

In particular,

$$\text{supp}(\rho^{XYM}) \leq F'_\alpha \otimes F'_\beta \otimes F'_\rho, \quad \text{supp}(\sigma^{X^A Y^B M}) \leq (F'_\alpha)^{\otimes A} \otimes (F'_\beta)^{\otimes B} \otimes F'_\rho = \text{supp}(\tau^{X^A Y^B M}).$$

To finish the proof of Lemma 7, we only need to show

$$\|\sigma^{X^A Y^B M} - \tau^{X^A Y^B M}\|_1 \leq 20\epsilon^{1/32}, \tag{17}$$

because of Lemma 8 and Fact 2.

In order to prove the above inequality, we augment  $X$  to  $L_X X$ ,  $Y$  to  $L_Y Y$  as in Section 3.4. Note that  $L_X, L_Y$  are the additional Hilbert spaces required for augmentation and have nothing to do the  $L_X, L_Y$  used for flattening earlier. In fact, the current definition of the Hilbert space  $X$  is actually the old  $X$  tensored with the old flattening  $L_X$ ; same for the current definition of  $Y$ . Recall that the new spaces  $L_X, L_Y$  required for augmentation have the same dimension  $L$ . The value of  $L$  will be chosen later, and it will be sufficiently large for our later purposes. Keeping augmentation in mind, we define

$$\alpha^{L_X X} := \frac{\mathbb{1}^{L_X}}{L} \otimes \alpha^X, \quad \beta^{L_Y Y} := \frac{\mathbb{1}^{L_Y}}{L} \otimes \beta^Y, \quad \rho^{L_X X L_Y Y M} := \frac{\mathbb{1}^{L_X}}{L} \otimes \frac{\mathbb{1}^{L_Y}}{L} \otimes \rho^{XYM}.$$

Then, the  $A$ -fold tensor product state  $\alpha^{(L_X X)^A}$  and the  $(|A| - 1)$ -fold tensor product state  $\alpha^{(L_X X)^{-a}}$  for any  $a \in [A]$  can be defined as before in the natural fashion. A similar comment holds for  $\beta^{(L_Y Y)^B}$  and  $\beta^{(L_Y Y)^{-b}}$  for any  $b \in [B]$ .

As in Section 3.4, we enlarge the Hilbert space  $M$  into a larger space  $\hat{M}$ . Because of this containment, the state  $\rho^{L_X X L_Y Y \hat{M}}$  is identical to  $\rho^{L_X X L_Y Y M}$ ; which notation we use depends on the application. Define the normalised state  $\hat{\rho}^{L_X X L_Y Y \hat{M}}$  and subnormalised state  $\hat{\hat{\rho}}^{L_X X L_Y Y \hat{M}}$  as in

Equation 9 via the projectors  $\Pi^{XYM}(3)$ ,  $\Pi^{XYM}(2) := \mathbb{1}^X \otimes \Pi^{YM}(2)$ ,  $\Pi^{XYM}(1) := \mathbb{1}^Y \otimes \Pi^{XM}(1)$ . Define

$$\begin{aligned}
\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}} &:= \sigma^{(L_X X)^A (L_Y Y)^B M} \\
&= (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \rho^{(L_X X)_a (L_Y Y)_b M} \otimes \alpha^{(L_X X)^{-a}} \otimes \beta^{(L_Y Y)^{-b}}, \\
\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} &:= (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \alpha^{(L_X X)^{-a}} \otimes \beta^{(L_Y Y)^{-b}}, \\
\hat{\hat{\sigma}}^{(L_X X)^A (L_Y Y)^B \hat{M}} &:= (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \hat{\hat{\rho}}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \alpha^{(L_X X)^{-a}} \otimes \beta^{(L_Y Y)^{-b}}, \\
\tau^{(L_X X)^A (L_Y Y)^B \hat{M}} &:= \tau^{(L_X X)^A (L_Y Y)^B M} = \alpha^{(L_X X)^A} \otimes \beta^{(L_Y Y)^B} \otimes \rho^{L_X M}.
\end{aligned} \tag{18}$$

Observe that

$$\begin{aligned}
\|\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}} - \tau^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 &\equiv \|\sigma^{(L_X X)^A (L_Y Y)^B M} - \tau^{(L_X X)^A (L_Y Y)^B M}\|_1 \\
&= \|\sigma^{X^A Y^B M} - \tau^{X^A Y^B M}\|_1.
\end{aligned}$$

Note that  $\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}}$ ,  $\tau^{(L_X X)^A (L_Y Y)^B \hat{M}}$ ,  $\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}$  are normalised states and  $\hat{\hat{\sigma}}^{(L_X X)^A (L_Y Y)^B \hat{M}}$  is subnormalised. So to prove Lemma 7, we just need to show that

$$\|\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}} - \tau^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 \leq 17\epsilon^{1/32}, \tag{19}$$

because of Equations 17, 18.

Define the tensor product spaces

$$L_X \otimes F'_\alpha =: L_X F'_\alpha \leq L_X X, \quad L_Y \otimes F'_\beta =: L_Y F'_\beta \leq L_Y Y.$$

Observe that

$$\begin{aligned}
(1 - \sqrt{\epsilon}) \frac{(\mathbb{1}^{L_X F'_\alpha})^{-a} \otimes (\mathbb{1}^{L_Y F'_\beta})^{-b}}{(L|F'_\alpha|)^{A-1} (L|F'_\beta|)^{B-1}} &\leq \alpha^{X^{-a}} \otimes \beta^{Y^{-b}} \\
&\leq (1 + \sqrt{\epsilon}) \frac{(\mathbb{1}^{L_X F'_\alpha})^{-a} \otimes (\mathbb{1}^{L_Y F'_\beta})^{-b}}{(L|F'_\alpha|)^{A-1} (L|F'_\beta|)^{B-1}}, \\
(1 - \sqrt{\epsilon}) \frac{(\mathbb{1}^{L_X F'_\alpha})^{\otimes A} \otimes (\mathbb{1}^{L_Y F'_\beta})^{\otimes B} \otimes (\mathbb{1}^{F'_\rho})}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} &\leq \tau^{(L_X X)^A (L_Y Y)^B \hat{M}} \\
&\leq (1 + \sqrt{\epsilon}) \frac{(\mathbb{1}^{L_X F'_\alpha})^{\otimes A} \otimes (\mathbb{1}^{L_Y F'_\beta})^{\otimes B} (\mathbb{1}^{F'_\rho})}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|},
\end{aligned} \tag{20}$$

for any  $a \in [A]$ ,  $b \in [B]$ .

We now prove four lemmas which are crucially required in the following arguments, leading to the proof of Lemma 7.

**Lemma 9**

$$\text{Tr} [\hat{\rho}^{L_X X L_Y Y \hat{M}} \hat{\rho}^{L_X X L_Y Y \hat{M}}] \leq \frac{(1 + 3\sqrt{\epsilon}) 2^{D_\infty^\epsilon} (\rho^{XYM} \|\alpha^X \otimes \beta^Y \otimes \rho^M\|)}{(L|F'_\alpha|)(L|F'_\beta|)|F'_\rho|}.$$

**Proof:** Using Equations 9, 8, 16, we get

$$\begin{aligned}
&\text{Tr} [\hat{\rho}^{L_X X L_Y Y \hat{M}} \hat{\rho}^{L_X X L_Y Y \hat{M}}] \\
&= \text{Tr} [\hat{\rho}^{L_X X L_Y Y \hat{M}} (\hat{\Pi}^{L_X X L_Y Y \hat{M}} \circ \rho^{L_X X L_Y Y \hat{M}})] \\
&= \text{Tr} [T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\rho^{L_X X L_Y Y M})]
\end{aligned}$$

$$\begin{aligned}
& ((\mathbb{1}^{L_X X L_Y Y \hat{M}} - T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{XYM} - \Pi^{XYM}(3)))) \\
& \quad \circ (\hat{\Pi}^{L_X X L_Y Y \hat{M}} \circ \rho^{L_X X L_Y Y \hat{M}})) \\
= & \text{Tr} [(T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X X L_Y Y M}) \circ T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\rho^{L_X X L_Y Y M})) \\
& ((\mathbb{1}^{L_X X L_Y Y \hat{M}} - T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{XYM} - \Pi^{XYM}(3)))) \\
& \quad \circ \hat{\rho}^{L_X X L_Y Y \hat{M}})] \\
= & \text{Tr} [T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\rho^{L_X X L_Y Y M}) \\
& (T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X X L_Y Y M}) - \\
& \quad T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{XYM} - \Pi^{XYM}(3)))) \circ \hat{\rho}^{L_X X L_Y Y \hat{M}}] \\
= & \text{Tr} [T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} ((\frac{\mathbb{1}^{L_X L_Y}}{L^2}) \otimes \rho^{XYM}) \\
& (T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X L_Y} \otimes \Pi^{XYM}(3)) \circ \hat{\rho}^{L_X X L_Y Y \hat{M}})] \\
= & \text{Tr} [T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} ((\frac{\mathbb{1}^{L_X L_Y}}{L^2}) \otimes (\Pi^{XYM}(3) \circ \rho^{XYM})) \hat{\rho}^{L_X X L_Y Y \hat{M}}] \\
\leq & \|T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} ((\frac{\mathbb{1}^{L_X L_Y}}{L^2}) \otimes (\Pi^{XYM}(3) \circ \rho^{XYM}))\|_{\infty} \cdot \|\hat{\rho}^{L_X X L_Y Y \hat{M}}\|_1 \\
= & \|(\frac{\mathbb{1}^{L_X L_Y}}{L^2}) \otimes (\Pi^{XYM}(3) \circ \rho^{XYM})\|_{\infty} \cdot \|\hat{\rho}^{L_X X L_Y Y \hat{M}}\|_1 \\
= & L^{-2} \cdot \|\Pi^{XYM}(3) \circ \rho^{XYM}\|_{\infty} \cdot \|\hat{\rho}^{L_X X L_Y Y \hat{M}}\|_1 \leq \frac{(1 + 3\sqrt{\epsilon})2^{D_{\infty}^{\epsilon}(\rho^{XYM} \|\alpha^X \otimes \beta^Y \otimes \rho^M)}}{(L|F'_{\alpha}|)(L|F'_{\beta}|)|F'_{\rho}|}.
\end{aligned}$$

Above, we used the fact that

$$\mathbb{1}^{L_X X L_Y Y \hat{M}} - T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{XYM} - \Pi^{XYM}(3))) \geq \hat{\Pi}^{L_X X L_Y Y \hat{M}}$$

in the second equality,

$$\begin{aligned}
& (\mathbb{1}^{L_X X L_Y Y \hat{M}} - T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{XYM} - \Pi^{XYM}(3)))) \\
& (T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X X L_Y Y M})) \\
= & T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X X L_Y Y M}) \\
& - T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{XYM} - \Pi^{XYM}(3)))
\end{aligned}$$

in the fourth equality,  $T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}}$  is an isometry in the sixth and seventh equalities. The proof of the lemma is now complete.  $\square$

**Lemma 10**

$$\text{Tr} [\hat{\rho}^{L_X X \hat{M}} \hat{\rho}^{L_X X \hat{M}}] \leq \frac{4\epsilon^{1/8}}{L^{3/2}} + \frac{(1 + 3\sqrt{\epsilon})2^{D_{\infty}^{\epsilon}(\rho^{XM} \|\alpha^X \otimes \rho^M)}}{(L|F'_{\alpha}|)|F'_{\rho}|}.$$

**Proof:** Using Equations 9, 8, 16 and Lemma 4, we get

$$\text{Tr} [\hat{\rho}^{L_X X \hat{M}} \hat{\rho}^{L_X X \hat{M}}]$$



$$\begin{aligned}
&= \text{Tr} [(\hat{\rho}^{L_X X \hat{M}} \otimes \mathbb{1}^{L_Y Y}) \hat{\rho}^{L_X X L_Y Y \hat{M}}] \\
&= \text{Tr} [((T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \otimes \mathbb{1}^{L_Y Y}) \hat{\rho}^{L_X X L_Y Y \hat{M}}] \\
&\quad + \text{Tr} [((\hat{\rho}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \otimes \mathbb{1}^{L_Y Y}) \hat{\rho}^{L_X X L_Y Y \hat{M}}] \\
&= \text{Tr} [((T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \otimes \mathbb{1}^{L_Y Y}) \\
&\quad ((\mathbb{1}^{L_X X L_Y Y \hat{M}} - (T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}} \otimes \mathbb{1}^{L_Y Y})(\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{X Y M} - \mathbb{1}^Y \otimes \Pi^{X M}(1))))) \\
&\quad \circ (\hat{\Pi}^{L_X X L_Y Y \hat{M}} \circ \rho^{L_X X L_Y Y \hat{M}}))] \\
&\quad + \text{Tr} [(\hat{\rho}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \hat{\rho}^{L_X X \hat{M}}] \\
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \\
&\quad (\mathbb{1}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\mathbb{1}^{L_X} \otimes (\mathbb{1}^{X M} - \Pi^{X M}(1))))) \otimes \mathbb{1}^{L_Y Y}) \circ \hat{\rho}^{L_X X L_Y Y \hat{M}}] \\
&\quad + \text{Tr} [(\hat{\rho}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \hat{\rho}^{L_X X \hat{M}}] \\
&= \text{Tr} [((T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \\
&\quad (\mathbb{1}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\mathbb{1}^{L_X} \otimes (\mathbb{1}^{X M} - \Pi^{X M}(1))))) \circ \hat{\rho}^{L_X X \hat{M}}] \\
&\quad + \text{Tr} [(\hat{\rho}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \hat{\rho}^{L_X X \hat{M}}] \\
&\leq \text{Tr} [((T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \\
&\quad (\mathbb{1}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\mathbb{1}^{L_X} \otimes (\mathbb{1}^{X M} - \Pi^{X M}(1))))) \circ \hat{\rho}^{L_X X \hat{M}}] \\
&\quad + \|\hat{\rho}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})\|_{\infty} \cdot \|\hat{\rho}^{L_X X \hat{M}}\|_1 \\
&\leq \text{Tr} [((T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \\
&\quad (\mathbb{1}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\mathbb{1}^{L_X} \otimes (\mathbb{1}^{X M} - \Pi^{X M}(1))))) \circ \hat{\rho}^{L_X X \hat{M}}] + 4\epsilon^{1/8} L^{-3/2} \\
&= \text{Tr} [((T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\mathbb{1}^{L_X X M}) \circ T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M})) \\
&\quad (\mathbb{1}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\mathbb{1}^{L_X} \otimes (\mathbb{1}^{X M} - \Pi^{X M}(1))))) \circ \hat{\rho}^{L_X X \hat{M}}] + 4\epsilon^{1/8} L^{-3/2} \\
&= \text{Tr} [T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\rho^{L_X X M}) \\
&\quad ((T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\mathbb{1}^{L_X X M}) - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\mathbb{1}^{L_X} \otimes (\mathbb{1}^{X M} - \Pi^{X M}(1))))) \circ \hat{\rho}^{L_X X \hat{M}}] \\
&\quad + 4\epsilon^{1/8} L^{-3/2} \\
&= \text{Tr} [T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}((\frac{\mathbb{1}^{L_X}}{L}) \otimes \rho^{X M}) \\
&\quad ((T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}(\mathbb{1}^{L_X} \otimes \Pi^{X M}(1))) \circ \hat{\rho}^{L_X X \hat{M}})] + 4\epsilon^{1/8} L^{-3/2} \\
&= \text{Tr} [(T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}((\frac{\mathbb{1}^{L_X}}{L}) \otimes (\Pi^{X M}(1) \circ \rho^{X M})) \hat{\rho}^{L_X X \hat{M}})] + 4\epsilon^{1/8} L^{-3/2} \\
&\leq \|T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}((\frac{\mathbb{1}^{L_X}}{L}) \otimes (\Pi^{X M}(1) \circ \rho^{X M}))\|_{\infty} \cdot \|\hat{\rho}^{L_X X \hat{M}}\|_1 + 4\epsilon^{1/8} L^{-3/2} \\
&= \|(\frac{\mathbb{1}^{L_X}}{L}) \otimes (\Pi^{X M}(1) \circ \rho^{X M})\|_{\infty} \cdot \|\hat{\rho}^{L_X X \hat{M}}\|_1 + 4\epsilon^{1/8} L^{-3/2}
\end{aligned}$$

$$\begin{aligned}
&= L^{-1} \cdot \|\Pi^{XM}(1) \circ \rho^{XM}\|_\infty \cdot \|\hat{\rho}^{L_X X \hat{M}}\|_1 + 4\epsilon^{1/8} L^{-3/2} \\
&\leq \frac{(1 + 3\sqrt{\epsilon}) 2^{D_\infty^\epsilon(\rho^{XM} \|\alpha^X \otimes \rho^M)} (L|F'_\alpha|)|F'_\rho|}{(L|F'_\alpha|)|F'_\rho|} + 4\epsilon^{1/8} L^{-3/2}.
\end{aligned}$$

Above, we used the fact that

$$\mathbb{1}^{L_X X L_Y Y \hat{M}} - T_{\epsilon^{1/4}}^{L_X X L_Y Y M \rightarrow L_X X L_Y Y \hat{M}} (\mathbb{1}^{L_X L_Y} \otimes (\mathbb{1}^{XYM} - \mathbb{1}^Y \Pi^{XM}(1))) \geq \hat{\Pi}^{L_X X L_Y Y \hat{M}}$$

in the third equality,

$$\begin{aligned}
&(\mathbb{1}^{L_X X \hat{M}} - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}} (\mathbb{1}^{L_X} \otimes (\mathbb{1}^{XM} - \Pi^{XM}(1)))) (T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}} (\mathbb{1}^{L_X X M})) \\
&= T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}} (\mathbb{1}^{L_X X M}) - T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}} (\mathbb{1}^{L_X} \otimes (\mathbb{1}^{XM} - \Pi^{XM}(1)))
\end{aligned}$$

in the seventh equality,  $T_{\epsilon^{1/4}}^{L_X X M \rightarrow L_X X \hat{M}}$  is an isometry in the ninth and tenth equalities. The proof of the lemma is now complete.  $\square$

**Lemma 11**

$$\text{Tr} [\hat{\rho}^{L_Y Y \hat{M}} \hat{\rho}^{L_Y Y \hat{M}}] \leq \frac{4\epsilon^{1/8}}{L^{3/2}} + \frac{(1 + 3\sqrt{\epsilon}) 2^{D_\infty^\epsilon(\rho^{YM} \|\beta^Y \otimes \rho^M)} (L|F'_\beta|)|F'_\rho|}{(L|F'_\beta|)|F'_\rho|}.$$

**Proof:** Similar to proof of Lemma 10 above.  $\square$

**Lemma 12**

$$\text{Tr} [\hat{\rho}^{\hat{M}} \hat{\rho}^{\hat{M}}] \leq \frac{8\epsilon^{1/8}}{L^{1/2}} + \frac{1 + \sqrt{\epsilon}}{|F'_\rho|}.$$

**Proof:** Using Equation 16 and Lemma 4, we get

$$\begin{aligned}
&\text{Tr} [\hat{\rho}^{\hat{M}} \hat{\rho}^{\hat{M}}] \\
&= \text{Tr} [\rho^{\hat{M}} \hat{\rho}^{\hat{M}}] + \text{Tr} [(\hat{\rho}^{\hat{M}} - \rho^{\hat{M}}) \hat{\rho}^{\hat{M}}] \leq \frac{1 + \sqrt{\epsilon}}{|F'_\rho|} \text{Tr} [\mathbb{1}^{F'_\rho} \hat{\rho}^{\hat{M}}] + \|\hat{\rho}^{\hat{M}} - \rho^{\hat{M}}\|_\infty \cdot \|\hat{\rho}^{\hat{M}}\|_1 \\
&\leq \frac{1 + \sqrt{\epsilon}}{|F'_\rho|} \text{Tr} [\mathbb{1}^{\hat{M}} \hat{\rho}^{\hat{M}}] + 8\epsilon^{1/8} L^{-1/2} \leq \frac{1 + \sqrt{\epsilon}}{|F'_\rho|} + 8\epsilon^{1/8} L^{-1/2}.
\end{aligned}$$

This completes the proof of the lemma.  $\square$

By Lemma 2 and Equation 18, we have

$$\|\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 \leq \|\rho^{L_X X L_Y Y \hat{M}} - \hat{\rho}^{L_X X L_Y Y \hat{M}}\|_1 = 2\sqrt{2} \cdot \epsilon^{1/8}. \quad (21)$$

Hence by triangle inequality and Equation 19, in order to prove Lemma 7, it suffices to show that

$$\|\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} - \tau^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 \leq 14\epsilon^{1/32}. \quad (22)$$

By Corollary 2 and Equation 18, we have

$$\|\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 \leq \|\rho^{L_X X L_Y Y \hat{M}} - \hat{\rho}^{L_X X L_Y Y \hat{M}}\|_1 < 25 \cdot \epsilon^{1/8}.$$

By Lemma 1, there is a subnormalised state  $\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}}$  such that

$$\begin{aligned} \hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}} &\leq \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}, \\ \|\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 &< 5\epsilon^{1/16}, \\ \|\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_2^2 &\leq (1 + 10\epsilon^{1/16}) \text{Tr} [\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}]. \end{aligned} \quad (23)$$

Hence by triangle inequality and Equation 22, in order to prove Lemma 7, it suffices to show that

$$\|\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}} - \tau^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 \leq 9\epsilon^{1/32}. \quad (24)$$

Note that  $\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}}$  is subnormalised.

Define  $\Pi^{(L_X X)^A (L_Y Y)^B \hat{M}}$  to be the orthogonal projector from the ambient space  $(L_X X)^A (L_Y Y)^B \hat{M}$  onto

$$\text{supp}(\tau^{(L_X X)^A (L_Y Y)^B \hat{M}}) = (L_X \otimes F'_\alpha)^A \otimes (L_Y \otimes F'_\beta)^B \otimes F'_\rho \geq \text{supp}(\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}}).$$

By Equations 23, 21, we get

$$\begin{aligned} &\text{Tr} [\Pi^{(L_X X)^A (L_Y Y)^B \hat{M}} \tau^{(L_X X)^A (L_Y Y)^B \hat{M}}] \\ &= \text{Tr} [\tau^{(L_X X)^A (L_Y Y)^B \hat{M}}] = 1, \\ &\text{Tr} [\Pi^{(L_X X)^A (L_Y Y)^B \hat{M}} \hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}}] \\ &\geq \text{Tr} [\Pi^{(L_X X)^A (L_Y Y)^B \hat{M}} \sigma^{(L_X X)^A (L_Y Y)^B \hat{M}}] - \|\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 \\ &\quad - \|\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 \\ &= \text{Tr} [\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}}] - \|\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 \\ &\quad - \|\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1 \\ &\geq 1 - 2\sqrt{2} \cdot \epsilon^{1/8} - 5\epsilon^{1/16} \geq 1 - 8\epsilon^{1/16}. \end{aligned} \quad (25)$$

By Proposition 1 and Equations 24, 25, in order to prove Lemma 7, it suffices to show that

$$\sqrt{\text{Tr} [\Pi^{(L_X X)^A (L_Y Y)^B \hat{M}}]} \cdot \|\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}} - \tau^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_2 < 6\epsilon^{1/32}.$$

Observe that  $\text{Tr} [\Pi^{(L_X X)^A (L_Y Y)^B \hat{M}}] = L^{A+B} |F'_\alpha|^A |F'_\beta|^B |F'_\rho|$ . Hence it suffices to show that

$$\|\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}} - \tau^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_2^2 < \frac{35\epsilon^{1/16}}{L^{A+B} |F'_\alpha|^A |F'_\beta|^B |F'_\rho|}. \quad (26)$$

The left hand side of the above inequality is

$$\begin{aligned}
& \|\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}} - \tau^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_2^2 \\
&= \|\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_2^2 + \|\tau^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_2^2 \\
&\quad - 2 \operatorname{Tr} [\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}} \tau^{(L_X X)^A (L_Y Y)^B \hat{M}}] \\
&\leq (1 + 10\epsilon^{1/16}) \operatorname{Tr} [\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}] \\
&\quad + (1 + \sqrt{\epsilon})^2 \left\| \frac{(\mathbb{1}^{L_X F'_\alpha})^{\otimes A} \otimes (\mathbb{1}^{L_Y F'_\beta})^{\otimes B} \otimes (\mathbb{1}^{F'_\rho})}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} \right\|_2^2 \\
&\quad - 2(1 - \sqrt{\epsilon}) \operatorname{Tr} \left[ \hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}} \left( \frac{(\mathbb{1}^{L_X F'_\alpha})^{\otimes A} \otimes (\mathbb{1}^{L_Y F'_\beta})^{\otimes B} \otimes (\mathbb{1}^{F'_\rho})}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} \right) \right] \\
&= (1 + 10\epsilon^{1/16}) \operatorname{Tr} [\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}] + \frac{(1 + \sqrt{\epsilon})^2}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} \\
&\quad - \frac{2(1 - \sqrt{\epsilon})}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} \operatorname{Tr} [\hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}} \Pi^{(L_X X)^A (L_Y Y)^B \hat{M}}] \\
&\leq (1 + 10\epsilon^{1/16}) \operatorname{Tr} [\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}] + \frac{(1 + \sqrt{\epsilon})^2}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} \\
&\quad - \frac{2(1 - \sqrt{\epsilon})(1 - 8\epsilon^{1/16})}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} \\
&\leq (1 + 10\epsilon^{1/16}) \operatorname{Tr} [\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}] + \frac{12\epsilon^{1/16} - 1}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|},
\end{aligned} \tag{27}$$

where the first inequality follows from Equations 23, 20, and the second inequality follows from Equation 25.

From Equations 26 and 27, in order to prove Lemma 7, it suffices to show the following lemma.

**Lemma 13** *Suppose*

$$\sqrt{L} > \max \left\{ \frac{4|F'_\beta||F'_\rho|}{\epsilon^{3/8} 2D_\infty^\epsilon(\rho^{YM} \|\beta^Y \otimes \rho^M)}, \frac{4|F'_\alpha||F'_\rho|}{\epsilon^{3/8} 2D_\infty^\epsilon(\rho^{XM} \|\alpha^X \otimes \rho^M)}, \frac{8|F'_\rho|}{\epsilon^{3/8}} \right\},$$

and

$$\begin{aligned}
\log A &> D_\infty^\epsilon(\rho^{XM} \|\alpha^X \otimes \rho^M) + \log \epsilon^{-1/16}, \\
\log B &> D_\infty^\epsilon(\rho^{YM} \|\beta^Y \otimes \rho^M) + \log \epsilon^{-1/16}, \\
\log A + \log B &> D_\infty^\epsilon(\rho^{XYM} \|\alpha^X \otimes \beta^Y \otimes \rho^M) + \log \epsilon^{-1/16}.
\end{aligned}$$

Then,

$$\operatorname{Tr} [\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}] < \frac{1 + 11\epsilon^{1/16}}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|}.$$

**Proof:** Given the lower bound on  $\sqrt{L}$ , we can write down the following consequences of Lemmas 12, 10, 11, 9.

$$\begin{aligned}
\operatorname{Tr} [\hat{\rho}^{\hat{M}} \hat{\rho}^{\hat{M}}] &\leq \frac{1+2\sqrt{\epsilon}}{|F'_\rho|}, \\
\operatorname{Tr} [\hat{\rho}^{L_X X \hat{M}} \hat{\rho}^{L_X X \hat{M}}] &\leq \frac{(1+4\sqrt{\epsilon})2^{D_\infty^\epsilon(\rho^{XM} \|\alpha^X \otimes \rho^M)}}{(L|F'_\alpha|)|F'_\rho|}, \\
\operatorname{Tr} [\hat{\rho}^{L_Y Y \hat{M}} \hat{\rho}^{L_Y Y \hat{M}}] &\leq \frac{(1+4\sqrt{\epsilon})2^{D_\infty^\epsilon(\rho^{YM} \|\beta^Y \otimes \rho^M)}}{(L|F'_\beta|)|F'_\rho|}, \\
\operatorname{Tr} [\hat{\rho}^{L_X X L_Y Y \hat{M}} \hat{\rho}^{L_X X L_Y Y \hat{M}}] &\leq \frac{(1+3\sqrt{\epsilon})2^{D_\infty^\epsilon(\rho^{XYM} \|\alpha^X \otimes \beta^Y \otimes \rho^M)}}{(L|F'_\alpha|)(L|F'_\beta|)|F'_\rho|},
\end{aligned} \tag{28}$$

We have,

$$\begin{aligned}
& \text{Tr} [\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}] \\
&= (AB)^{-2} \cdot \text{Tr} \left[ \sum_{a, \hat{a}=1}^A \sum_{b, \hat{b}=1}^B (\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \alpha^{(L_X X)^{-a}} \otimes \beta^{(L_Y Y)^{-b}} \right. \\
&\quad \left. (\hat{\rho}^{(L_X X)_{\hat{a}} (L_Y Y)_{\hat{b}} \hat{M}} \otimes \alpha^{(L_X X)^{-\hat{a}}} \otimes \beta^{(L_Y Y)^{-\hat{b}}}) \right] \\
&\leq \frac{(1+\sqrt{\epsilon})^2}{(AB)^2 (L|F'_\alpha|)^{2(A-1)} (L|F'_\beta|)^{2(B-1)}} \\
&\quad \cdot \sum_{a, \hat{a}=1}^A \sum_{b, \hat{b}=1}^B \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b}}) \\
&\quad (\hat{\rho}^{(L_X X)_{\hat{a}} (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-\hat{a}}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-\hat{b}}})]
\end{aligned} \tag{29}$$

We analyse the above summation by considering several cases. Consider the following term for a fixed choice of  $a \neq \hat{a}$ ,  $b \neq \hat{b}$ .

$$\begin{aligned}
& \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b}}) (\hat{\rho}^{(L_X X)_{\hat{a}} (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-\hat{a}}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-\hat{b}}})] \\
&= \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{\hat{a}}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{\hat{b}}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a, \hat{a}}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b, \hat{b}}}) \\
&\quad (\hat{\rho}^{(L_X X)_{\hat{a}} (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)_a} \otimes \mathbb{1}^{(L_Y F'_\beta)_b} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a, \hat{a}}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b, \hat{b}}})] \\
&= \text{Tr} [((\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{\hat{a}}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{\hat{b}}}) \\
&\quad (\hat{\rho}^{(L_X X)_{\hat{a}} (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)_a} \otimes \mathbb{1}^{(L_Y F'_\beta)_b})) \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a, \hat{a}}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b, \hat{b}}}] \\
&= (L|F'_\alpha|)^{A-2} (L|F'_\beta|)^{B-2} \\
&\quad \cdot \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{\hat{a}}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{\hat{b}}}) \\
&\quad (\hat{\rho}^{(L_X X)_{\hat{a}} (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)_a} \otimes \mathbb{1}^{(L_Y F'_\beta)_b})] \\
&= (L|F'_\alpha|)^{A-2} (L|F'_\beta|)^{B-2} \\
&\quad \cdot \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X X)_{\hat{a}}} \otimes \mathbb{1}^{(L_Y Y)_{\hat{b}}}) \\
&\quad (\hat{\rho}^{(L_X X)_{\hat{a}} (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_X X)_a} \otimes \mathbb{1}^{(L_Y Y)_b})] \\
&= (L|F'_\alpha|)^{A-2} (L|F'_\beta|)^{B-2} \text{Tr} [\hat{\rho}^{\hat{M}} \hat{\rho}^{\hat{M}}] \leq \frac{(1+2\sqrt{\epsilon})(L|F'_\alpha|)^{A-2} (L|F'_\beta|)^{B-2}}{|F'_\rho|},
\end{aligned} \tag{30}$$

where we used Equation 28 in the inequality above. There are  $A(A-1)B(B-1)$  such terms.

Consider the following term for a fixed choice of  $a = \hat{a}$ ,  $b \neq \hat{b}$ .

$$\begin{aligned}
& \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b}}) (\hat{\rho}^{(L_X X)_a (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-\hat{b}}})] \\
&= \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{\hat{b}}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b, \hat{b}}}) \\
&\quad (\hat{\rho}^{(L_X X)_a (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_Y F'_\beta)_b} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b, \hat{b}}})] \\
&= \text{Tr} [((\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{\hat{b}}}) \\
&\quad (\hat{\rho}^{(L_X X)_a (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_Y F'_\beta)_b})) \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b, \hat{b}}}] \\
&= (L|F'_\alpha|)^{A-1} (L|F'_\beta|)^{B-2} \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{\hat{b}}}) \\
&\quad (\hat{\rho}^{(L_X X)_a (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_Y F'_\beta)_b})] \\
&= (L|F'_\alpha|)^{A-1} (L|F'_\beta|)^{B-2} \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_Y Y)_{\hat{b}}}) (\hat{\rho}^{(L_X X)_{\hat{a}} (L_Y Y)_{\hat{b}} \hat{M}} \otimes \mathbb{1}^{(L_Y Y)_b})] \\
&= (L|F'_\alpha|)^{A-1} (L|F'_\beta|)^{B-2} \text{Tr} [\hat{\rho}^{(L_X X) \hat{M}} \hat{\rho}^{(L_X X) \hat{M}}] \\
&\leq \frac{(1+4\sqrt{\epsilon})(L|F'_\alpha|)^{A-1} (L|F'_\beta|)^{B-2} 2^{D_\infty(\rho^{XM} \| \alpha^X \otimes \rho^M)}}{L|F'_\alpha| |F'_\beta|},
\end{aligned} \tag{31}$$

where we used Equation 28 in the inequality above. There are  $AB(B-1)$  such terms.

Consider the following term for a fixed choice of  $a \neq \hat{a}$ ,  $b = \hat{b}$ . We have similarly,

$$\begin{aligned} & \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b}}) \\ & \quad (\hat{\rho}^{(L_X X)_{\hat{a}} (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-\hat{a}}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b}})] \\ & \leq \frac{(1+4\sqrt{\epsilon})(L|F'_\alpha|)^{A-2}(L|F'_\beta|)^{B-1}2^{D_\infty^\epsilon}(\rho^{YM} \otimes \rho^M)}{L|F'_\beta||F'_\rho|}. \end{aligned} \quad (32)$$

There are  $A(A-1)B$  such terms.

Finally, consider the following term for a fixed choice of  $a = \hat{a}$ ,  $b = \hat{b}$ .

$$\begin{aligned} & \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b}}) \\ & \quad (\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b}})] \\ & = \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}}) \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b}}] \\ & = (L|F'_\alpha|)^{A-1}(L|F'_\beta|)^{B-1} \text{Tr} [\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}}] \\ & \leq \frac{(1+3\sqrt{\epsilon})(L|F'_\alpha|)^{A-1}(L|F'_\beta|)^{B-1}2^{D_\infty^\epsilon}(\rho^{XYM} \otimes \alpha^X \otimes \beta^Y \otimes \rho^M)}{(L|F'_\alpha|)(L|F'_\beta|)|F'_\rho|}, \end{aligned} \quad (33)$$

where we used Equation 28 in the inequality above. There are  $AB$  such terms.

From Equations 30, 31, 32, 33 we get

$$\begin{aligned} & \sum_{a, \hat{a}=1}^A \sum_{b, \hat{b}=1}^B \text{Tr} [(\hat{\rho}^{(L_X X)_a (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-a}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b}}) \\ & \quad (\hat{\rho}^{(L_X X)_{\hat{a}} (L_Y Y)_b \hat{M}} \otimes \mathbb{1}^{(L_X F'_\alpha)^{-\hat{a}}} \otimes \mathbb{1}^{(L_Y F'_\beta)^{-b}})] \\ & = \frac{A(A-1)B(B-1)(1+2\sqrt{\epsilon})(L|F'_\alpha|)^{A-2}(L|F'_\beta|)^{B-2}}{|F'_\rho|} \\ & \quad + \frac{AB(B-1)(1+4\sqrt{\epsilon})(L|F'_\alpha|)^{A-1}(L|F'_\beta|)^{B-2}2^{D_\infty^\epsilon}(\rho^{XM} \otimes \alpha^X \otimes \rho^M)}{L|F'_\alpha||F'_\rho|} \\ & \quad + \frac{A(A-1)B(1+4\sqrt{\epsilon})(L|F'_\alpha|)^{A-2}(L|F'_\beta|)^{B-1}2^{D_\infty^\epsilon}(\rho^{YM} \otimes \beta^Y \otimes \rho^M)}{L|F'_\beta||F'_\rho|} \\ & \quad + \frac{AB(1+3\sqrt{\epsilon})(L|F'_\alpha|)^{A-1}(L|F'_\beta|)^{B-1}2^{D_\infty^\epsilon}(\rho^{XYM} \otimes \alpha^X \otimes \beta^Y \otimes \rho^M)}{(L|F'_\alpha|)(L|F'_\beta|)|F'_\rho|}. \end{aligned} \quad (34)$$

From Equations 29, 34 we get,

$$\begin{aligned} & \text{Tr} [\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}] \\ & \leq \frac{(1+\sqrt{\epsilon})^2}{(AB)^2(L|F'_\alpha|)^{2(A-1)}(L|F'_\beta|)^{2(B-1)}} \\ & \quad \cdot \left[ \frac{A(A-1)B(B-1)(1+2\sqrt{\epsilon})(L|F'_\alpha|)^{A-2}(L|F'_\beta|)^{B-2}}{|F'_\rho|} \right. \\ & \quad + \frac{AB(B-1)(1+4\sqrt{\epsilon})(L|F'_\alpha|)^{A-1}(L|F'_\beta|)^{B-2}2^{D_\infty^\epsilon}(\rho^{XM} \otimes \alpha^X \otimes \rho^M)}{L|F'_\alpha||F'_\rho|} \\ & \quad + \frac{A(A-1)B(1+4\sqrt{\epsilon})(L|F'_\alpha|)^{A-2}(L|F'_\beta|)^{B-1}2^{D_\infty^\epsilon}(\rho^{YM} \otimes \beta^Y \otimes \rho^M)}{L|F'_\beta||F'_\rho|} \\ & \quad \left. + \frac{AB(1+3\sqrt{\epsilon})(L|F'_\alpha|)^{A-1}(L|F'_\beta|)^{B-1}2^{D_\infty^\epsilon}(\rho^{XYM} \otimes \alpha^X \otimes \beta^Y \otimes \rho^M)}{(L|F'_\alpha|)(L|F'_\beta|)|F'_\rho|} \right] \end{aligned}$$

$$\begin{aligned}
&\leq (1 + 7\sqrt{\epsilon}) \left[ \frac{1}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} + A^{-1} \cdot \frac{2^{D_\infty^\epsilon(\rho^{XM} \| \alpha^X \otimes \rho^M)}}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} \right. \\
&\quad \left. + B^{-1} \cdot \frac{2^{D_\infty^\epsilon(\rho^{YM} \| \beta^Y \otimes \rho^M)}}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} + (AB)^{-1} \cdot \frac{2^{D_\infty^\epsilon(\rho^{XYM} \| \alpha^X \otimes \beta^Y \otimes \rho^M)}}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} \right] \\
&\leq \frac{(1 + 7\sqrt{\epsilon})(1 + 3\epsilon^{1/16})}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|} \leq \frac{1 + 11\epsilon^{1/16}}{(L|F'_\alpha|)^A (L|F'_\beta|)^B |F'_\rho|}.
\end{aligned}$$

This finishes the proof of the lemma.  $\square$

Since the dimension  $L$  of the augmenting Hilbert spaces  $L_X, L_Y$  is a free parameter in our proof, it can be chosen large enough as in the requirement of Lemma 13. Thus Lemma 13 holds, which completes the proof of Lemma 7.

## 5 Smooth soft covering without pairwise independence

We will now prove a smooth multipartite soft covering classical quantum version of Lemma 7 where pairwise independence amongst the classical random variables can be relaxed slightly. For simplicity, we will prove the case with two parties only. However, analogous results hold for greater than two parties also, as the machinery will make amply clear below.

Let  $\mathcal{X}, \mathcal{Y}$  be classical alphabets. Let  $p^{XY}$  be a normalised probability distribution on  $XY$ , whose marginals are denoted by  $p^X, p^Y$ . Let  $q^X, q^Y$  be normalised probability distributions. For each  $(x, y) \in \mathcal{X} \times \mathcal{Y}$ , let  $\rho_{xy}^M$  be a subnormalised density matrix on  $M$ . We define the following classical quantum *control state*:

$$\rho^{XYM} := \sum_{(x,y) \in \mathcal{X} \times \mathcal{Y}} p^{XY}(x, y) |x, y\rangle \langle x, y|^{XY} \otimes \rho_{xy}^M.$$

Define the density matrices

$$\rho_x^M := \sum_y p^{Y|X=x}(y) \rho_{xy}^M, \quad \rho_y^M := \sum_x p^{X|Y=y}(x) \rho_{xy}^M,$$

and the classical quantum states

$$\rho^{XM, Y|X} := \sum_{x,y} p^{XY}(x, y) |x, y\rangle \langle x, y|^{XY} \otimes \rho_x^M, \quad \rho^{YM, X|Y} := \sum_{x,y} p^{XY}(x, y) |x, y\rangle \langle x, y|^{XY} \otimes \rho_y^M.$$

Intuitively,  $\rho_x^M$  is the quantum state induced on  $M$  when the  $X$  register of the control state  $\rho^{XYM}$  is measured and found to be equal to  $x$ . Similarly, the classical quantum state  $\rho^{XM, Y|X}$  is what one gets if one ‘decouples’ the register  $M$  from  $Y$ , making it depend only on the contents of  $X$ , without affecting the probability distribution on  $XY$ . Analogous remarks can be made for  $\rho_y^M$  and  $\rho^{YM, X|Y}$ .

Suppose  $\text{supp}(p^X) \leq \text{supp}(q^X)$  and  $\text{supp}(p^Y) \leq \text{supp}(q^Y)$ . Let  $A, B$  be positive integers. Let  $x^{(A)} := (x(1), \dots, x(A)) \in \mathcal{X}^A$  denote a  $|A|$ -tuple of elements from  $\mathcal{X}$ ;  $y^{(B)}$  has a similar meaning. For any pair of tuples  $x^{(A)}, y^{(B)}$ , define the *sample average covering state*

$$\sigma_{x^{(A)}, y^{(B)}}^M := (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \frac{p^{XY}(x(a), y(b))}{q^X(x(a)) q^Y(y(b))} \rho_{x(a), y(b)}^M, \quad (35)$$

where the fraction term above represents the ‘change of measure’ from the product probability distribution  $q^X \times q^Y$  to the joint probability distribution  $p^{XY}$ .

Let  $\bar{q}^{X^A}, \bar{q}^{Y^B}$  be two probability distributions on  $\mathcal{X}^A, \mathcal{Y}^B$  with the following properties:

1. The marginal on any coordinate  $\bar{q}^{X_a}$  for any  $a \in [A]$  equals  $q^X$ ;
2. The marginal on any two coordinates  $\bar{q}^{X_a X_{a'}}$  for any  $a, a' \in [A], a \neq a'$  is independent of  $a, a'$ ; hence it can be modelled as a distribution  $\bar{q}^{X X'}$ ;
3. Similar properties as the two above hold for  $\bar{q}^{Y^B}$ .

For any  $(x, y) \in \mathcal{X} \times \mathcal{Y}$ , define the positive semidefinite matrices

$$\begin{aligned} \rho(X)_{xy}^M &:= \sum_{y'} \frac{\bar{q}^{Y'|Y=y}(y') p^{Y|X=x}(y')}{q^Y(y')} \rho_{xy'}^M, \quad \rho(Y)_{xy}^M := \sum_{x'} \frac{\bar{q}^{X'|X=x}(x') p^{X|Y=y}(x')}{q^X(x')} \rho_{x'y}^M, \\ \rho()_{xy}^M &:= \sum_{x'y'} \frac{\bar{q}^{Y'|Y=y}(y') \bar{q}^{X'|X=x}(x') p(x', y')}{q^X(x') q^Y(y')} \rho_{x'y'}^M. \end{aligned}$$

**Definition 3** Let  $0 < \epsilon < 1$ . For the two party classical quantum soft covering problem, the consequences of loss of pairwise independence are said to hold up to scale factors of  $(\epsilon, f(\epsilon), g(\epsilon))$  if there exist probability distributions  $p(X)^{XY}, p(Y)^{XY}, p()^{XY}$  satisfying

$$\|p^{XY} - p(X)^{XY}\|_1 < \epsilon, \quad \|p^{XY} - p(Y)^{XY}\|_1 < \epsilon, \quad \|p^{XY} - p()^{XY}\|_1 < \epsilon,$$

such that

$$\rho(X)^{XYM} \leq f(\epsilon) \rho^{XM, Y|X}, \quad \rho(Y)^{XYM} \leq f(\epsilon) \rho^{YM, X|Y}, \quad \rho()^{XYM} \leq (1 + g(\epsilon)) (p^{XY} \otimes \rho^M),$$

where

$$\begin{aligned} \rho(X)^{XYM} &:= \sum_{(x,y) \in \mathcal{X} \times \mathcal{Y}} p(X)^{XY}(x, y) |x, y\rangle \langle x, y|^{XY} \otimes \rho(X)_{xy}^M, \\ \rho(Y)^{XYM} &:= \sum_{(x,y) \in \mathcal{X} \times \mathcal{Y}} p(Y)^{XY}(x, y) |x, y\rangle \langle x, y|^{XY} \otimes \rho(Y)_{xy}^M, \\ \rho()^{XYM} &:= \sum_{(x,y) \in \mathcal{X} \times \mathcal{Y}} p()^{XY}(x, y) |x, y\rangle \langle x, y|^{XY} \otimes \rho()_{xy}^M. \end{aligned}$$

In this paper, we shall take  $f(\epsilon) \leq 2^{\text{polylog}(\epsilon^{-1})}$  and  $g(\epsilon) \leq \epsilon^{\Theta(1)}$ .

We are now ready to give the statement of our smooth two party classical quantum covering lemma under slight relaxation of pairwise independence.

**Proposition 5** Consider the setting of the two party classical quantum soft covering problem. Suppose,

$$\begin{aligned} \log A &> D_\infty^\epsilon(\rho^{XM} \| q^X \otimes \rho^M) + \log f(e) + 3 + \log \epsilon^{-1/2}, \\ \log B &> D_\infty^\epsilon(\rho^{YM} \| q^Y \otimes \rho^M) + \log f(e) + 3 + \log \epsilon^{-1/2}, \\ \log A + \log B &> D_\infty^\epsilon(\rho^{XYM} \| q^X \otimes q^Y \otimes \rho^M) + 1 + \log \epsilon^{-1/2}. \end{aligned}$$

Then,

$$\mathbb{E}_{x^{(A)}, y^{(B)}} [\|\sigma_{x^{(A)}, y^{(B)}}^M - \rho^M\|_1] < (2\sqrt{15\epsilon^{1/64} + \sqrt{40\epsilon^{1/32} + 8e + g(e) + 4e}})(\text{Tr } \rho).$$

where the expectation is taken over independent choices of tuples  $x^{(A)}, y^{(B)}$  from the distributions  $\bar{q}^{X^A}, \bar{q}^{Y^B}$  which satisfy the consequences of loss of pairwise independence up to scale factors of  $(e, f(e), g(e))$ . Note that for the usual setting of  $e = \epsilon$ ,  $f(e) \leq 2^{\text{polylog}(e^{-1})}$  and  $g(e) \leq e^{\Theta(1)}$ , the right hand sides of the inequalities in the above rate region will have additive terms of  $\text{polylog}(\epsilon^{-1})$  and the covering error will be bounded by  $\epsilon^{\Theta(1)}$ .



**Remarks:**

1. In the pairwise independent setting,  $\bar{q}^{XX'} = q^X \times q^{X'}$ ,  $\bar{q}^{YY'} = q^Y \times q^{Y'}$ . Hence,  $\rho(X)^{XYM} = \rho^{XM,Y|X}$ ,  $\rho(Y)^{XYM} = \rho^{YM,X|Y}$  and  $\rho()^{XYM} = p^{XY} \otimes \rho^M$ . Thus,  $\bar{q}^{XX'}$ ,  $\bar{q}^{YY'}$  satisfy the consequences of loss of pairwise independence up to scale factors of  $(0, 1, 0)$  i.e. perfectly. The rate region and covering error promised by Proposition 5 is essentially the same as that given by the smooth soft covering lemma of [Sen24]. The main difference is that the rate region of Proposition 5 is given in terms of  $D_\infty^\epsilon(\cdot\|\cdot)$  versus  $D_2^\epsilon(\cdot\|\cdot)$  in [Sen24].

2. Proposition 5 cannot be proved by the telescoping technique. Consider a simplified setting where  $p^{XY} = p^X \otimes p^Y$ ,  $q^X = p^X$ ,  $q^Y = p^Y$ ,  $p^{XY} = p(X)^{XY} = p(Y)^{XY} = p()^{XY}$ , but  $q^{XX'} \neq q^X \otimes q^{X'}$ ,  $q^{YY'} \neq q^Y \otimes q^{Y'}$ . In other words,  $p^{XY}$  is a tensor product distribution, the sampling distributions  $q^X, q^Y$  are the same as the marginals  $p^X, p^Y$  but there is a loss of pairwise independence in sampling  $(x(a))_a$  as well as in sampling  $(y(b))_b$ ; nevertheless  $p^{XY}$  does not require perturbation in order to satisfy the operator inequalities involving  $f(e), g(e)$ . Telescoping starts off by using the mapping  $(x, y) \mapsto \rho'_{xy}{}^M := (\rho_{xy}^M - \rho_x^M - \rho_y^M + \rho^M)$  instead of using the original mapping  $(x, y) \mapsto \rho_{xy}^M$ . The mean zero requirement of telescoping is the fact that

$$\forall(x, y) : \quad \mathbb{E}_{(x, x') \sim p^X \times p^{X'}}[\rho'_{xy}{}^M \rho'_{x'y'}{}^M] = \mathbb{E}_{(y, y') \sim p^Y \times p^{Y'}}[\rho'_{xy}{}^M \rho'_{xy'}{}^M] = 0.$$

However under the ‘slight breaking of pairwise independence’, we have to evaluate the expectation under  $(x, x') \sim q^{XX'}$ , and similarly for  $(y, y')$ . Since  $q^{XX'} \neq p^X \times p^{X'}$ , the mean zero requirement fails. Replacing  $\rho'_{xy}{}^M$  with other candidates like  $(\rho_{xy}^M - \rho(X)_{xy}^M - \rho(Y)_{xy}^M + \rho()_{xy}^M)$  does not help either.

To prove Proposition 5, we generally follow the lines of the proof of Lemma 7. Several steps are simpler because  $X, Y$  are classical. In particular, we do not need to flatten  $q^X, q^Y$ . Only the flattening superoperator  $\mathcal{F}_\rho^{M \rightarrow L_M M}$  is needed. Again, without loss of generality, we assume that  $\rho^{XYM}$  is a normalised density matrix.

From the notion of intersection of classical probability distributions defined in Equation 6, we get

$$\|(p \cap p(X) \cap p(Y) \cap p())^{XY} - p^{XY}\|_1 < 3e.$$

Arguing along the lines of the initial part of the proof of Proposition 2 and defining a set

$$\text{Bad} := \{(x, y) : p^{XY}(x, y) > 4(p \cap p(X) \cap p(Y) \cap p())^{XY}(x, y)\},$$

we see that there is a subnormalised probability distribution  $\hat{p}^{XY}$  such that

$$\|\hat{p}^{XY} - p^{XY}\|_1 < 4e, \hat{p}^{XY} \leq p^{XY} < 4\hat{p}^{XY}, \hat{p}^{XY} \leq p(X)^{XY}, \hat{p}^{XY} \leq p(Y)^{XY}, \hat{p}^{XY} \leq p()^{XY}, \quad (36)$$

where the inequality  $p^{XY} < 4\hat{p}^{XY}$  above holds only over  $\text{supp}(\hat{p}^{XY})$ . We can define the sample average covering state  $\hat{\sigma}_{x^{(A)}, y^{(B)}}^M$  using  $\hat{p}^{XY}$  instead of  $p^{XY}$  as in Equation 35 in the natural fashion. Similarly, we can define the new classical quantum control state  $\hat{\rho}^{XYM}$ . Then for any  $x^{(A)}, y^{(B)}$ , using Equations 35, 36, we can easily see that  $\|\hat{\sigma}_{x^{(A)}, y^{(B)}}^M - \sigma_{x^{(A)}, y^{(B)}}^M\|_1 < 4e$ . So in order to prove Proposition 5, it suffices to prove

$$\mathbb{E}_{x^{(A)}, y^{(B)}} [\|\hat{\sigma}_{x^{(A)}, y^{(B)}}^M - \rho^M\|_1] < 2\sqrt{15\epsilon^{1/64} + \sqrt{40\epsilon^{1/32} + 8e + g(e)}}. \quad (37)$$

From the definitions of smooth  $D_\infty^\epsilon(\cdot|\cdot)$ , we have subnormalised classical quantum density matrices  $\rho^{XYM}(3)$ ,  $\rho^{XM}(1)$  and  $\rho^{YM}(2)$  satisfying

$$\|\rho^{XYM} - \rho^{XYM}(3)\|_1 \leq \epsilon, \quad \|\rho^{XM} - \rho^{XM}(1)\|_1 \leq \epsilon, \quad \|\rho^{YM} - \rho^{YM}(2)\|_1 \leq \epsilon,$$

such that

$$\begin{aligned} \rho^{XYM}(3) &\leq 2^{D_\infty^\epsilon(\rho^{XYM}\|q^X \otimes q^Y \otimes \rho^M)}(q^X \otimes q^Y \otimes \rho^M), \\ \rho^{XM}(1) &\leq 2^{D_\infty^\epsilon(\rho^{XM}\|q^X \otimes \rho^M)}(q^X \otimes \rho^M), \\ \rho^{YM}(2) &\leq 2^{D_\infty^\epsilon(\rho^{YM}\|q^Y \otimes \rho^M)}. \end{aligned} \quad (38)$$

Note that in general  $\rho^{XM}(1)$ ,  $\rho^{YM}(2)$  have nothing to do with the marginals of  $\rho^{XYM}(3)$ . From Equation 36 and the triangle inequality, we can write

$$\|\hat{\rho}^{XYM} - \rho^{XYM}(3)\|_1 \leq 4e + \epsilon, \quad \|\hat{\rho}^{XM} - \rho^{XM}(1)\|_1 \leq 4e + \epsilon, \quad \|\hat{\rho}^{YM} - \rho^{YM}(2)\|_1 \leq 4e + \epsilon. \quad (39)$$

We can extend the classical quantum states  $\rho^{XM}(1)$ ,  $\rho^{YM}(2)$  to classical quantum states  $\rho^{XM,Y|X}(1)$ ,  $\rho^{YM,X|Y}(2)$  in the natural fashion. Henceforth we will be working with the control state  $\hat{\rho}^{XYM}$  and probability distribution  $\hat{p}^{XY}$  only. So to ease the notation below, we rename  $\hat{p}^{XY}$  to  $p^{XY}$ ,  $\hat{\rho}^{XYM}$  to  $\rho^{XYM}$ ,  $\hat{\sigma}_{x(A),y(B)}^M$  to  $\sigma_{x(A),y(B)}^M$ . Under this simplified notation, in order to prove Proposition 5, Equation 37 rewritten says that it suffices to show

$$\mathbb{E}_{x(A),y(B)} [\|\sigma_{x(A),y(B)}^M - \rho^M\|_1] < 2\sqrt{15\epsilon^{1/64} + \sqrt{40\epsilon^{1/32} + 8e + g(e)}}. \quad (40)$$

Also, Equation 39 is rewritten as

$$\|\rho^{XYM} - \rho^{XYM}(3)\|_1 \leq 4e + \epsilon, \quad \|\rho^{XM} - \rho^{XM}(1)\|_1 \leq 4e + \epsilon, \quad \|\rho^{YM} - \rho^{YM}(2)\|_1 \leq 4e + \epsilon. \quad (41)$$

Apply the flattening CPTP superoperator to get the normalised density matrix

$$\rho^{XYL_MM} := (\mathbb{I}^{XY} \otimes \mathcal{F}_\rho^{M \rightarrow L_MM})(\rho^{XYM}).$$

Define  $\rho^{XYL_MM}(3)$ ,  $\rho^{XL_MM}(1)$ ,  $\rho^{YL_MM}(2)$ ,  $\rho(X)^{XYL_MM}$ ,  $\rho(Y)^{XYL_MM}$ ,  $\rho(\cdot)^{XYL_MM}$  via flattening in the natural fashion. Extend  $\rho^{XL_MM}(1)$ ,  $\rho^{YL_MM}(2)$  to classical quantum states  $\rho^{XL_MM,Y|X}(1)$ ,  $\rho^{YL_MM,X|Y}(2)$  in the natural fashion. Then we use Equation 41 and conclude,

$$\begin{aligned} \|\rho^{XYL_MM} - \rho^{XYL_MM}(3)\|_1 &< 4e + \epsilon, \\ \|\rho^{XL_MM} - \rho^{XL_MM}(1)\|_1 &< 4e + \epsilon, \quad \|\rho^{YL_MM} - \rho^{YL_MM}(2)\|_1 < 4e + \epsilon, \end{aligned} \quad (42)$$

since a CPTP superoperator cannot increase the trace distance.

By Lemma 6 and Proposition 3, we have

$$(1 + \delta)^{-2} \frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|} \leq \mathcal{F}_\rho^{M \rightarrow L_MM}(\rho^M) \leq (1 + \delta) \frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|}. \quad (43)$$

Because a CPTP superoperator respects the Löwner partial order on Hermitian matrices, combining Equations 38 and 43 gives us

$$\begin{aligned} \rho^{XYL_MM}(3) &\leq (1 + \delta) 2^{D_\infty^\epsilon(\rho^{XYM}\|q^X \otimes q^Y \otimes \rho^M)}(q^X \otimes q^Y \otimes (\frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|})), \\ \rho^{XL_MM}(1) &\leq (1 + \delta) 2^{D_\infty^\epsilon(\rho^{XM}\|q^X \otimes \rho^M)}(q^X \otimes (\frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|})), \\ \rho^{YL_MM}(2) &\leq (1 + \delta) 2^{D_\infty^\epsilon(\rho^{YM}\|q^Y \otimes \rho^M)}(q^Y \otimes (\frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|})). \end{aligned} \quad (44)$$

Applying Proposition 2 to Equations 42, 44 tells us that there exists classical quantum orthogonal projectors

$$\begin{aligned}\Pi^{XY L_M M}(3) &:= \sum_{xy} |x, y\rangle\langle x, y|^{XY} \otimes \Pi^{L_M M}(3)_{xy}, \\ \Pi^{X L_M M}(1) &:= \sum_x |x\rangle\langle x|^X \otimes \Pi^{L_M M}(1)_x, \\ \Pi^{Y L_M M}(2) &:= \sum_y |y\rangle\langle y|^Y \otimes \Pi^{L_M M}(2)_y,\end{aligned}$$

such that

$$\begin{aligned}\text{Tr} [\Pi^{XY L_M M}(3) \rho^{XY L_M M}] &\geq 1 - \sqrt{4e + \epsilon}, \\ \text{Tr} [\Pi^{X L_M M}(1) \rho^{X L_M M}] &\geq 1 - \sqrt{4e + \epsilon}, \\ \text{Tr} [\Pi^{Y L_M M}(2) \rho^{Y L_M M}] &\geq 1 - \sqrt{4e + \epsilon}, \\ \Pi^{XY L_M M}(3) \rho^{XY L_M M} &= \rho^{XY L_M M} \Pi^{XY L_M M}(3), \\ \Pi^{X L_M M}(1) \rho^{X L_M M} &= \rho^{X L_M M} \Pi^{X L_M M}(1), \\ \Pi^{Y L_M M}(2) \rho^{Y L_M M} &= \rho^{Y L_M M} \Pi^{Y L_M M}(2), \\ \forall(x, y) : \frac{p(x, y)}{q(x)q(y)} \|\Pi_{xy}^{L_M M}(3) \rho_{xy}^{L_M M}\|_\infty &\leq \frac{(1+\delta)(1+2\sqrt{4e+\epsilon})2^{D_\infty^\epsilon(\rho^{XY M} \|_{q^X \otimes q^Y \otimes \rho^M})}}{|F'_\rho|}, \\ \forall x : \frac{p(x)}{q(x)} \|\Pi_x^{L_M M}(1) \rho_x^{L_M M}\|_\infty &\leq \frac{(1+\delta)(1+2\sqrt{4e+\epsilon})2^{D_\infty^\epsilon(\rho^{X M} \|_{q^X \otimes \rho^M})}}{|F'_\rho|}, \\ \forall y : \frac{p(y)}{q(y)} \|\Pi_y^{L_M M}(2) \rho_y^{L_M M}\|_\infty &\leq \frac{(1+\delta)(1+2\sqrt{4e+\epsilon})2^{D_\infty^\epsilon(\rho^{Y M} \|_{q^Y \otimes \rho^M})}}{|F'_\rho|}.\end{aligned}\tag{45}$$

Define the *flattened sample average covering state*

$$\sigma_{x^{(A)}, y^{(B)}}^{L_M M} := (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \frac{p^{XY}(x(a), y(b))}{q^X(x(a))q^Y(y(b))} \rho_{x(a), y(b)}^{L_M M},$$

Proposition 4 and Fact 2 together with concavity of square root imply that

$$\begin{aligned}\mathbb{E}_{x^{(A)}, y^{(B)}} [\|\sigma_{x^{(A)}, y^{(B)}}^{L_M M} - \rho^{L_M M}\|_1] &\leq 2 \mathbb{E}_{x^{(A)}, y^{(B)}} [\sqrt{\|\sigma_{x^{(A)}, y^{(B)}}^{L_M M} - \rho^{L_M M}\|_1}] \\ &\leq 2 \sqrt{\mathbb{E}_{x^{(A)}, y^{(B)}} [\|\sigma_{x^{(A)}, y^{(B)}}^{L_M M} - \rho^{L_M M}\|_1]},\end{aligned}\tag{46}$$

where the expectation is taken over independent choices of tuples  $x^{(A)}, y^{(B)}$  from the distributions  $\bar{q}^{X^A}, \bar{q}^{Y^B}$ . We note that  $\delta$  can be taken as small as we please at the expense of increasing  $|F'_\rho|$ . Essentially,  $\delta$  can be treated as a free parameter. We will set  $\delta := \sqrt{\epsilon}/3$ .

To make the notation lighter, henceforth we will always work with the flattened states and the Hilbert space  $L_M M$ . Hence from now on we redefine  $M \equiv L_M M$ . Under this redefinition, we restate Equations 45, 43, and the consequences of Equation 36 as Equation 47 below. That is, we can assume that there are classical quantum orthogonal projectors  $\Pi^{XY M}(3)$ ,  $\Pi^{X M}(1)$  and  $\Pi^{Y M}(2)$

such that

$$\begin{aligned}
\text{Tr} [\Pi^{XYM}(3)\rho^{XYM}] &\geq 1 - \sqrt{\epsilon} - 2\sqrt{e}, \\
\text{Tr} [\Pi^{XM}(1)\rho^{XM}] &\geq 1 - \sqrt{\epsilon} - 2\sqrt{e}, \\
\text{Tr} [\Pi^{YM}(2)\rho^{YM}] &\geq 1 - \sqrt{\epsilon} - 2\sqrt{e}, \\
\Pi^{XYM}(3)\rho^{XYM} &= \rho^{XYM}\Pi^{XYM}(3), \\
\Pi^{XM}(1)\rho^{XM} &= \rho^{XM}\Pi^{XM}(1), \\
\Pi^{YM}(2)\rho^{YM} &= \rho^{YM}\Pi^{YM}(2), \\
\forall (x, y) \in \text{supp}(p^{XY}) : \rho(X)_{xy}^M &\leq 4f(e)\rho_x^M, \\
\forall (x, y) \in \text{supp}(p^{XY}) : \rho(Y)_{xy}^M &\leq 4f(e)\rho_y^M, \\
\forall (x, y) \in \text{supp}(p^{XY}) : \rho()_{xy}^M &\leq (1 + g(e))\rho^M, \\
\forall (x, y) : \frac{p(x, y)}{q(x)q(y)} \|\Pi_{xy}^M(3)\rho_{xy}^M\|_\infty &\leq \frac{(1+4\sqrt{e}+2\sqrt{\epsilon})2^{D_\infty^\epsilon(\rho^{XYM}\|_{q^X \otimes q^Y \otimes \rho^M})}}{|F'_\rho|}, \\
\forall x : \frac{p(x)}{q(x)} \|\Pi_x^M(1)\rho_x^M\|_\infty &\leq \frac{(1+4\sqrt{e}+2\sqrt{\epsilon})2^{D_\infty^\epsilon(\rho^{XM}\|_{q^X \otimes \rho^M})}}{|F'_\rho|}, \\
\forall y : \frac{p(y)}{q(y)} \|\Pi_y^M(2)\rho_y^M\|_\infty &\leq \frac{(1+4\sqrt{e}+2\sqrt{\epsilon})2^{D_\infty^\epsilon(\rho^{YM}\|_{q^Y \otimes \rho^M})}}{|F'_\rho|}, \\
F'_\rho &\leq M, \\
(1 - \sqrt{\epsilon}) \frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|} &\leq \rho^M \leq (1 + \sqrt{\epsilon}) \frac{\mathbb{1}_{F'_\rho}}{|F'_\rho|}.
\end{aligned} \tag{47}$$

To finish the proof of Proposition 5, we only need to show

$$\mathbb{E}_{x^{(A)}, y^{(B)}} [\|\sigma_{x^{(A)}, y^{(B)}}^M - \rho^M\|_1] \leq 15\epsilon^{1/64} + \sqrt{40\epsilon^{1/32} + 8e + g(e)}. \tag{48}$$

because of Equation 46.

In order to prove the above inequality, we augment  $X$  to  $L_X X$ ,  $Y$  to  $L_Y Y$  as in Section 3.4. Recall that the new spaces  $L_X$ ,  $L_Y$  required for augmentation have the same dimension  $L$ . The value of  $L$  will be chosen later, and it will be sufficiently large for our later purposes. Keeping augmentation in mind, we define

$$q^{L_X X} := \frac{\mathbb{1}^{L_X}}{L} \otimes q^X, \quad q^{L_Y Y} := \frac{\mathbb{1}^{L_Y}}{L} \otimes q^Y, \quad \rho^{L_X X L_Y Y M} := \frac{\mathbb{1}^{L_X}}{L} \otimes \frac{\mathbb{1}^{L_Y}}{L} \otimes \rho^{XYM}.$$

As in Section 3.4, we enlarge the Hilbert space  $M$  into a larger space  $\hat{M}$ . Because of this containment, the state  $\rho^{L_X X L_Y Y \hat{M}}$  is identical to  $\rho^{L_X X L_Y Y M}$ ; which notation we use depends on the application. Define tuples  $(l_x x)^{(A)}$ ,  $(l_y y)^{(B)}$  in the natural fashion. Define the normalised state  $\hat{\rho}^{L_X X L_Y Y \hat{M}}$  and subnormalised state  $\hat{\hat{\rho}}^{L_X X L_Y Y \hat{M}}$  as in Equation 9 via the classical quantum projectors  $\Pi^{XYM}(3)$ ,  $\Pi^{XYM}(2) := \Pi^{YM}(2) \otimes \mathbb{1}^X$ ,  $\Pi^{XYM}(1) := \Pi^{XM}(1) \otimes \mathbb{1}^Y$ . Observe that the states  $\hat{\rho}^{L_X X L_Y Y \hat{M}}$ ,  $\hat{\hat{\rho}}^{L_X X L_Y Y \hat{M}}$ , and the approximate intersection projector  $\hat{\Pi}^{L_X X L_Y Y \hat{M}}$  of

Equation 8 are now classical quantum. Thus we can write

$$\begin{aligned}
\hat{\rho}^{L_X X L_Y Y \hat{M}} &:= L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) |l_x, x, l_y, y\rangle \langle l_x, x, l_y, y|^{L_X X L_Y Y} \otimes \hat{\rho}_{l_x x l_y y}^{\hat{M}}, \\
\hat{\rho}_{l_x x l_y y}^{\hat{M}} &= T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho_{xy}^M), \\
\hat{\rho}^{L_X X L_Y Y \hat{M}} &:= L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) |l_x, x, l_y, y\rangle \langle l_x, x, l_y, y|^{L_X X L_Y Y} \otimes \hat{\rho}_{l_x x l_y y}^{\hat{M}}, \\
\forall i \in [3] : \hat{\Pi}^{XYM}(i) &:= \sum_{x, y} |x, y\rangle \langle x, y|^{XY} \otimes \Pi_{xy}^M(i), \\
\hat{\Pi}^{L_X X L_Y Y \hat{M}} &:= \sum_{l_x, x, l_y, y} |l_x, x, l_y, y\rangle \langle l_x, x, l_y, y|^{L_X X L_Y Y} \otimes \hat{\Pi}_{l_x x l_y y}^{\hat{M}}, \\
\hat{\Pi}_{l_x x l_y y}^{\hat{M}} &= \mathbb{1}^{\hat{M}} - \text{span} \{T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}((\mathbb{1}^M - \Pi_x^M(1))), T_{l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}((\mathbb{1}^M - \Pi_y^M(2))), \\
&\quad T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}((\mathbb{1}^M - \Pi_{xy}^M(3)))\}, \\
\hat{\rho}_{l_x x l_y y}^{\hat{M}} &= \hat{\Pi}_{l_x x l_y y}^{\hat{M}} \circ \rho_{xy}^M.
\end{aligned} \tag{49}$$

Define

$$\begin{aligned}
\sigma_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} &:= \sigma_{(l_x x)^{(A)}, (l_y y)^{(B)}}^M = \sigma_{x^{(A)}, y^{(B)}}^M \\
&= (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \frac{p^{XY}(x(a), y(b))}{q^X(x(a)) q^Y(y(b))} \rho_{x(a), y(b)}^M, \\
\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} &:= (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \frac{p^{XY}(x(a), y(b))}{q^X(x(a)) q^Y(y(b))} \hat{\rho}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}, \\
\hat{\hat{\sigma}}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} &:= (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \frac{p^{XY}(x(a), y(b))}{q^X(x(a)) q^Y(y(b))} \hat{\hat{\rho}}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}.
\end{aligned} \tag{50}$$

Observe that

$$\begin{aligned}
\forall ((l_x x)^{(A)}, (l_y y)^{(B)}) : \text{supp}(\sigma_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}) &\leq F'_\rho \leq M, \\
\|\rho^{XYM}\|_1 = \|\rho^M\|_1 = \|p^{XY}\|_1 &\geq 1 - 4e.
\end{aligned} \tag{51}$$

Define the probability distributions

$$\bar{q}^{(L_X X)^A} := \left(\frac{\mathbb{1}^{L_X}}{L}\right)^{\otimes A} \otimes \bar{q}^{X^A}, \quad \bar{q}^{(L_Y Y)^B} := \left(\frac{\mathbb{1}^{L_Y}}{L}\right)^{\otimes B} \otimes \bar{q}^{Y^B}.$$

Observe that

$$\begin{aligned}
\mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} [\|\sigma_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} - \rho^{\hat{M}}\|_1] &\equiv \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} [\|\sigma_{(l_x x)^{(A)}, (l_y y)^{(B)}}^M - \rho^M\|_1] \\
&= \mathbb{E}_{x^{(A)}, y^{(B)}} [\|\sigma_{x^{(A)}, y^{(B)}}^M - \rho^M\|_1],
\end{aligned}$$

where the expectation is taken over independent choices of tuples  $(l_x x)^{(A)}, (l_y y)^{(B)}$  from the distributions  $\bar{q}^{(L_X X)^A}, \bar{q}^{(L_Y Y)^B}$ . So to prove Proposition 5, we just need to show that

$$\mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} [\|\sigma_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} - \rho^{\hat{M}}\|_1] \leq 15\epsilon^{1/64} + \sqrt{40\epsilon^{1/32} + 8e + g(e)}. \tag{52}$$

because of Equations 48, 50.

We now prove four lemmas which are crucially required in the following arguments, leading to the proof of Proposition 5.

**Lemma 14**

$$\begin{aligned}
L^{-2} \sum_{l_x, x, l_y, y} q^X(x) q^Y(y) \text{Tr} \left[ \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
\leq \frac{(1 + 4\sqrt{e} + 2\sqrt{e}) 2^{D_\infty^\epsilon(\rho^{XYM} \| q^X \otimes q^Y \otimes \rho^M)}}{|F'_\rho|}.
\end{aligned}$$

**Proof:** Using Equations 49, 47 and arguing as in the proof of Lemma 9, we get

$$\begin{aligned}
& L^{-2} \sum_{l_x, x, l_y, y} q^X(x) q^Y(y) \operatorname{Tr} \left[ \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
&= L^{-2} \sum_{l_x, x, l_y, y} \operatorname{Tr} \left[ \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) (p^{XY}(x, y) \hat{\Pi}_{l_x x l_y y}^{\hat{M}} \circ \rho_{xy}^{\hat{M}}) \right] \\
&= L^{-2} \sum_{l_x, x, l_y, y} \operatorname{Tr} \left[ \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho_{xy}^M) \right) \right. \\
&\quad \left. (p^{XY}(x, y) ((\mathbf{1}^{\hat{M}} - T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}((\mathbf{1}^M - \Pi_{xy}^M(3)))) \circ (\hat{\Pi}_{l_x x l_y y}^{\hat{M}} \circ \rho_{xy}^{\hat{M}}))) \right] \\
&= L^{-2} \sum_{l_x, x, l_y, y} \operatorname{Tr} \left[ \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} (T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\mathbf{1}^M)) \circ (T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho_{xy}^M)) \right) \right. \\
&\quad \left. (p^{XY}(x, y) ((\mathbf{1}^{\hat{M}} - T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}((\mathbf{1}^M - \Pi_{xy}^M(3)))) \circ \hat{\rho}_{l_x x l_y y}^{\hat{M}})) \right] \\
&= L^{-2} \sum_{l_x, x, l_y, y} \operatorname{Tr} \left[ \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho_{xy}^M) \right) \right. \\
&\quad \left. (p^{XY}(x, y) ((T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\mathbf{1}^M) - T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}((\mathbf{1}^M - \Pi_{xy}^M(3)))) \circ \hat{\rho}_{l_x x l_y y}^{\hat{M}})) \right] \\
&= L^{-2} \sum_{l_x, x, l_y, y} \operatorname{Tr} \left[ \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho_{xy}^M) \right) (p^{XY}(x, y) ((T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\Pi_{xy}^M(3))) \circ \hat{\rho}_{l_x x l_y y}^{\hat{M}})) \right] \\
&= L^{-2} \sum_{l_x, x, l_y, y} \operatorname{Tr} \left[ \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\Pi_{xy}^M(3) \circ \rho_{xy}^M) \right) (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\
&\leq L^{-2} \sum_{l_x, x, l_y, y} \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \|T_{l_x, l_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\Pi_{xy}^M(3) \circ \rho_{xy}^M)\|_{\infty} \cdot (p^{XY}(x, y) \|\hat{\rho}_{l_x x l_y y}^{\hat{M}}\|_1) \\
&= L^{-2} \sum_{l_x, x, l_y, y} \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \|\Pi_{xy}^M(3) \circ \rho_{xy}^M\|_{\infty} \cdot (p^{XY}(x, y) \|\hat{\rho}_{l_x x l_y y}^{\hat{M}}\|_1) \\
&\leq L^{-2} \sum_{l_x, x, l_y, y} \frac{(1 + 4\sqrt{e} + 2\sqrt{e}) 2^{D_{\infty}^{\epsilon}}(\rho^{XYM} \|q^X \otimes q^Y \otimes \rho^M\|)}{|F'_{\rho}|} p^{XY}(x, y) \\
&\leq \frac{(1 + 4\sqrt{e} + 2\sqrt{e}) 2^{D_{\infty}^{\epsilon}}(\rho^{XYM} \|q^X \otimes q^Y \otimes \rho^M\|)}{|F'_{\rho}|}.
\end{aligned}$$

The proof of the lemma is now complete.  $\square$

**Lemma 15**

$$\begin{aligned}
& L^{-3} \sum_{l_x, x, l_y, y, l'_y, y'} q^X(x) \bar{q}^{Y Y'}(y, y') \operatorname{Tr} \left[ \left( \frac{p^{XY}(x, y')}{q^X(x) q^Y(y')} \hat{\rho}_{l_x x l'_y y'}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
&\leq 4\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 + \frac{4(1 + 4\sqrt{e} + 2\sqrt{e}) f(e) 2^{D_{\infty}^{\epsilon}}(\rho^{XM} \|q^X \otimes \rho^M\|)}{|F'_{\rho}|}.
\end{aligned}$$

**Proof:** Using Equations 49, 47 and Lemma 4, and arguing as in the proof of Lemma 10, we get

$$\begin{aligned}
& L^{-3} \sum_{l_x, x, l_y, y, l'_y, y'} q^X(x) \bar{q}^{Y Y'}(y, y') \operatorname{Tr} \left[ \left( \frac{p^{XY}(x, y')}{q^X(x) q^Y(y')} \hat{\rho}_{l_x x l'_y y'}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
&= L^{-3} \sum_{l_x, x, l_y, y, l'_y, y'} \frac{p^X(x)}{q^X(x)} \cdot \frac{p^{Y|X=x}(y') \bar{q}^{Y'|Y=y}(y')}{q^Y(y')} \operatorname{Tr} \left[ \hat{\rho}_{l_x x l'_y y'}^{\hat{M}} (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\
&= L^{-3} \sum_{l_x, x, l_y, y, l'_y, y'} \frac{p^X(x)}{q^X(x)} \cdot \frac{p^{Y|X=x}(y') \bar{q}^{Y'|Y=y}(y')}{q^Y(y')} \operatorname{Tr} \left[ (T_{l_x, l'_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho_{xy'}^M)) (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\
&= L^{-2} \sum_{l_x, x, l_y, y} \frac{p^X(x)}{q^X(x)} \operatorname{Tr} \left[ \left( L^{-1} \sum_{l'_y y'} \frac{p^{Y|X=x}(y') \bar{q}^{Y'|Y=y}(y')}{q^Y(y')} T_{l_x, l'_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho_{xy'}^M) \right) (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\
&= L^{-2} \sum_{l_x, x, l_y, y} \frac{p^X(x)}{q^X(x)} \operatorname{Tr} \left[ \left( \left( L^{-1} \sum_{l'_y} T_{l_x, l'_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}} \left( \sum_{y'} \frac{p^{Y|X=x}(y') \bar{q}^{Y'|Y=y}(y')}{q^Y(y')} \rho_{xy'}^M \right) \right) \right. \right. \\
&\quad \left. \left. - T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}} \left( \sum_{y'} \frac{p^{Y|X=x}(y') \bar{q}^{Y'|Y=y}(y')}{q^Y(y')} \rho_{xy'}^M \right) \right) (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\
&\quad + L^{-2} \sum_{l_x, x, l_y, y} \frac{p^X(x)}{q^X(x)} \operatorname{Tr} \left[ \left( T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}} \left( \sum_{y'} \frac{p^{Y|X=x}(y') \bar{q}^{Y'|Y=y}(y')}{q^Y(y')} \rho_{xy'}^M \right) \right) (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\
&\leq L^{-2} \sum_{l_x, x, l_y, y} \frac{p^X(x)}{q^X(x)} \left\| \left( L^{-1} \sum_{l'_y} T_{l_x, l'_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho(X)_{xy}^M) \right) - T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho(X)_{xy}^M) \right\|_{\infty} \cdot (p^{XY}(x, y) \|\hat{\rho}_{l_x x l_y y}^{\hat{M}}\|_1) \\
&\quad + L^{-2} \sum_{l_x, x, l_y, y} \frac{p^X(x)}{q^X(x)} \operatorname{Tr} \left[ (T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho(X)_{xy}^M)) (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\
&\leq 4\epsilon^{1/8} L^{-5/2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 \\
&\quad + L^{-2} \sum_{l_x, x, l_y, y} \frac{p^X(x)}{q^X(x)} \operatorname{Tr} \left[ (T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho(X)_{xy}^M)) (p^{XY}(x, y) (\hat{\Pi}_{l_x x l_y y}^{\hat{M}} \circ \rho_{xy}^{\hat{M}})) \right] \\
&= 4\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 \\
&\quad + L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \cdot \frac{p^X(x)}{q^X(x)} \\
&\quad \operatorname{Tr} \left[ T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho(X)_{xy}^M) ((\mathbb{1}^{\hat{M}} - T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}((\mathbb{1}^M - \Pi_x^M(1)))) \circ (\hat{\Pi}_{l_x x l_y y}^{\hat{M}} \circ \rho_{xy}^{\hat{M}})) \right] \\
&= 4\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1
\end{aligned}$$

$$\begin{aligned}
& + L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \cdot \frac{p^X(x)}{q^X(x)} \\
& \quad \text{Tr} [((T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\mathbb{1}^M)) \circ (T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho(X)_{xy}^M))) \\
& \quad \quad ((\mathbb{1}^{\hat{M}} - T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}((\mathbb{1}^M - \Pi_x^M(1)))) \circ \hat{\rho}_{l_x x l_y y}^{\hat{M}})] \\
= & \quad 4\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 \\
& + L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \cdot \frac{p^X(x)}{q^X(x)} \\
& \quad \text{Tr} [(T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho(X)_{xy}^M))((T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\mathbb{1}^M) - T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}((\mathbb{1}^M - \Pi_x^M(1)))) \circ \hat{\rho}_{l_x x l_y y}^{\hat{M}})] \\
= & \quad 4\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 \\
& + L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \cdot \frac{p^X(x)}{q^X(x)} \text{Tr} [(T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho(X)_{xy}^M))((T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\Pi_x^M(1))) \circ \hat{\rho}_{l_x x l_y y}^{\hat{M}})] \\
= & \quad 4\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 \\
& + L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \cdot \frac{p^X(x)}{q^X(x)} \text{Tr} [(T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\Pi_x^M(1) \circ \rho(X)_{xy}^M)) \hat{\rho}_{l_x x l_y y}^{\hat{M}}] \\
\leq & \quad 4\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 \\
& + 4f(e)L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \cdot \frac{p^X(x)}{q^X(x)} \text{Tr} [(T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\Pi_x^M(1) \circ \rho_x^M)) \hat{\rho}_{l_x x l_y y}^{\hat{M}}] \\
\leq & \quad 4\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 \\
& + 4f(e)L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \cdot \frac{p^X(x)}{q^X(x)} \|T_{l_x, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\Pi_x^M(1) \circ \rho_x^M)\|_\infty \cdot \|\hat{\rho}_{l_x x l_y y}^{\hat{M}}\|_1 \\
= & \quad 4\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 \\
& + 4f(e)L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \cdot \frac{p^X(x)}{q^X(x)} \|\Pi_x^M(1) \circ \rho_x^M\|_\infty \cdot \|\hat{\rho}_{l_x x l_y y}^{\hat{M}}\|_1 \\
\leq & \quad 4\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 \\
& + 4f(e)L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \cdot \frac{(1 + 4\sqrt{e} + 2\sqrt{\epsilon})2^{D_\infty^\epsilon}(\rho^{XM} \|q^X \otimes \rho^M\|)}{|F'_\rho|}
\end{aligned}$$



$$\leq 4\epsilon^{1/8} L^{-1/2} \sum_{x,y} p^{XY}(x,y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1 + \frac{4(1+4\sqrt{\epsilon}+2\sqrt{\epsilon})f(e)2^{D_\infty^\epsilon(\rho^{XM}\|q^X\otimes\rho^M)}}{|F'_\rho|}.$$

The proof of the lemma is now complete.  $\square$

**Lemma 16**

$$\begin{aligned} & L^{-3} \sum_{l_x, x, l_y, y, l'_x, x'} q^Y(y) \bar{q}^{XX'}(x, x') \text{Tr} \left[ \left( \frac{p^{XY}(x', y)}{q^X(x') q^Y(y)} \hat{\rho}_{l'_x x' l_y y}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\ & \leq 4\epsilon^{1/8} L^{-1/2} \sum_{x,y} p^{XY}(x,y) \frac{p^Y(y)}{q^Y(y)} \|\rho(Y)_{xy}^M\|_1 + \frac{4(1+4\sqrt{\epsilon}+2\sqrt{\epsilon})f(e)2^{D_\infty^\epsilon(\rho^{YM}\|q^Y\otimes\rho^M)}}{|F'_\rho|}. \end{aligned}$$

**Proof:** Similar to proof of Lemma 15 above.  $\square$

**Lemma 17**

$$\begin{aligned} & L^{-4} \sum_{l_x, x, l_y, y, l'_x, x', l'_y, y'} \bar{q}^{XX'}(x, x') \bar{q}^{YY'}(y, y') \text{Tr} \left[ \left( \frac{p^{XY}(x', y')}{q^X(x') q^Y(y')} \hat{\rho}_{l'_x x' l'_y y'}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\ & \leq 8\epsilon^{1/8} L^{-1/2} \sum_{x,y} p^{XY}(x,y) \|\rho()_{xy}^M\|_1 + \frac{(1+g(e)+2\sqrt{\epsilon})}{|F'_\rho|}. \end{aligned}$$

**Proof:** Using Equations 49, 47 and Lemma 4, and arguing as in the proof of Lemma 12, we get

$$\begin{aligned} & L^{-4} \sum_{l_x, x, l_y, y, l'_x, x', l'_y, y'} \bar{q}^{XX'}(x, x') \bar{q}^{YY'}(y, y') \text{Tr} \left[ \left( \frac{p^{XY}(x', y')}{q^X(x') q^Y(y')} \hat{\rho}_{l'_x x' l'_y y'}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\ & = L^{-4} \sum_{l_x, x, l_y, y, l'_x, x', l'_y, y'} \frac{p^{XY}(x', y') \bar{q}^{X'|X=x}(x') \bar{q}^{Y'|Y=y}(y')}{q^X(x') q^Y(y')} \text{Tr} \left[ \hat{\rho}_{l'_x x' l'_y y'}^{\hat{M}} (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\ & = L^{-4} \sum_{l_x, x, l_y, y, l'_x, x', l'_y, y'} \frac{p^{XY}(x', y') \bar{q}^{X'|X=x}(x') \bar{q}^{Y'|Y=y}(y')}{q^X(x') q^Y(y')} \text{Tr} \left[ (T_{l'_x, l'_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho_{x' y'}^M)) (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\ & = L^{-2} \sum_{l_x, x, l_y, y} \text{Tr} \left[ \left( L^{-2} \sum_{l'_x x' l'_y y'} \frac{p^{XY}(x', y') \bar{q}^{X'|X=x}(x') \bar{q}^{Y'|Y=y}(y')}{q^X(x') q^Y(y')} T_{l'_x, l'_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}}(\rho_{x' y'}^M) \right) \right. \\ & \quad \left. (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\ & = L^{-2} \sum_{l_x, x, l_y, y} \text{Tr} \left[ \left( \left( L^{-2} \sum_{l'_x l'_y} T_{l'_x, l'_y, \epsilon^{1/4}}^{M \rightarrow \hat{M}} \left( \sum_{x' y'} \frac{p^{XY}(x', y') \bar{q}^{X'|X=x}(x') \bar{q}^{Y'|Y=y}(y')}{q^X(x') q^Y(y')} \rho_{x' y'}^M \right) \right) \right. \right. \\ & \quad \left. \left. - \left( \sum_{x' y'} \frac{p^{XY}(x', y') \bar{q}^{X'|X=x}(x') \bar{q}^{Y'|Y=y}(y')}{q^X(x') q^Y(y')} \rho_{x' y'}^M \right) \right) (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \\ & \quad + L^{-2} \sum_{l_x, x, l_y, y} \text{Tr} \left[ \left( \sum_{x' y'} \frac{p^{XY}(x', y') \bar{q}^{X'|X=x}(x') \bar{q}^{Y'|Y=y}(y')}{q^X(x') q^Y(y')} \rho_{x' y'}^M \right) (p^{XY}(x, y) \hat{\rho}_{l_x x l_y y}^{\hat{M}}) \right] \end{aligned}$$

$$\begin{aligned}
&\leq L^{-2} \sum_{l_x, x, l_y, y} \left\| \left( L^{-2} \sum_{l'_x, l'_y} T_{l'_x, l'_y}^{M \rightarrow \hat{M}}(\rho()_{xy}^M) \right) - \rho()_{xy}^M \right\|_{\infty} \cdot (p^{XY}(x, y) \|\hat{\rho}_{l_x x l_y y}^{\hat{M}}\|_1) \\
&\quad + L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \text{Tr} [\rho()_{xy}^M \hat{\rho}_{l_x x l_y y}^{\hat{M}}] \\
&\leq 8\epsilon^{1/8} L^{-5/2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \|\rho()_{xy}^M\|_1 + (1 + g(e)) L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \text{Tr} [\rho^M \hat{\rho}_{l_x x l_y y}^{\hat{M}}] \\
&\leq 8\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \|\rho()_{xy}^M\|_1 + (1 + g(e))(1 + \sqrt{\epsilon}) L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \frac{\text{Tr} [\mathbb{1}^M \hat{\rho}_{l_x x l_y y}^{\hat{M}}]}{|F'_{\rho}|} \\
&\leq 8\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \|\rho()_{xy}^M\|_1 + (1 + g(e) + 2\sqrt{\epsilon}) L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \frac{\|\hat{\rho}_{l_x x l_y y}^{\hat{M}}\|_1}{|F'_{\rho}|} \\
&\leq 8\epsilon^{1/8} L^{-1/2} \sum_{x, y} p^{XY}(x, y) \|\rho()_{xy}^M\|_1 + \frac{(1 + g(e) + 2\sqrt{\epsilon})}{|F'_{\rho}|}.
\end{aligned}$$

The proof of the lemma is now complete.  $\square$

The reader may have noticed some differences between the statements of Lemma 17 and Lemma 12, Lemma 15 and Lemma 10, Lemma 16 and Lemma 11. The notable looking differences arise from corrections due to augmentation smoothing viz. from the application of Lemma 4. Part of the reason for these differences is the presence of terms like  $\rho(X)_{xy}$ ,  $\rho(Y)_{xy}$  etc. in the non-pairwise independent case, which do not exist in the pairwise independent case. The other part of the reason for the differences is cosmetic: the proof of the fully quantum smooth convex split lemma Lemma 7 uses flattening for  $\alpha^X$  and  $\beta^Y$ , whereas in the non pairwise independent classical quantum covering lemma Proposition 5, we do not need to flatten their classical analogues  $q^X$  and  $q^Y$ . Nevertheless, an effect similar to flattening  $\alpha^X$ ,  $\beta^Y$  persists in the statements of Lemmas 15, 16. Terms like  $\frac{1}{q^X(x)}$ ,  $\frac{1}{q^Y(y)}$  in the statements of Lemmas 15, 16 are related to flattening terms like  $\frac{1}{F'_{\alpha}}$ ,  $\frac{1}{F'_{\beta}}$  in the statements of Lemmas 10, 11.

By Lemma 2 and Equation 50, we have

$$\begin{aligned}
&\mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \left\| \sigma_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \right\|_1 \right] \\
&= (AB)^{-1} \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \left\| \sum_{a=1}^A \sum_{b=1}^B \frac{p^{XY}(x(a), y(b))}{q^X(x(a)) q^Y(y(b))} \rho_{(l_x x)^{(a)}, (l_y y)^{(b)}}^{\hat{M}} - \sum_{a=1}^A \sum_{b=1}^B \frac{p^{XY}(x(a), y(b))}{q^X(x(a)) q^Y(y(b))} \hat{\rho}_{(l_x x)^{(a)}, (l_y y)^{(b)}}^{\hat{M}} \right\|_1 \right] \\
&\leq (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \left\| \frac{p^{XY}(x(a), y(b))}{q^X(x(a)) q^Y(y(b))} \left( \rho_{(l_x x)^{(a)}, (l_y y)^{(b)}}^{\hat{M}} - \hat{\rho}_{(l_x x)^{(a)}, (l_y y)^{(b)}}^{\hat{M}} \right) \right\|_1 \right] \\
&= L^{-2} \sum_{l_x, x, l_y, y} q^X(x) q^Y(y) \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \left\| \rho_{l_x, x, l_y, y}^{\hat{M}} - \hat{\rho}_{l_x, x, l_y, y}^{\hat{M}} \right\|_1 \\
&= L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \left\| \rho_{l_x, x, l_y, y}^{\hat{M}} - \hat{\rho}_{l_x, x, l_y, y}^{\hat{M}} \right\|_1 \\
&= \left\| \rho^{L_X X L_Y Y \hat{M}} - \hat{\rho}^{L_X X L_Y Y \hat{M}} \right\|_1 \leq 2\sqrt{2} \cdot \epsilon^{1/8}.
\end{aligned} \tag{53}$$

Hence by triangle inequality and Equation 52, in order to prove Proposition 5, it suffices to show

that

$$\mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} [\|\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} - \rho^{\hat{M}}\|_1] \leq 12\epsilon^{1/64} + \sqrt{40\epsilon^{1/32} + 8e + g(e)}. \quad (54)$$

By Corollary 2 and Equation 50, we have

$$\begin{aligned} & \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \left\| \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} - \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \right\|_1 \right] \\ &= (AB)^{-1} \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \left\| \sum_{a=1}^A \sum_{b=1}^B \frac{p^{XY}(x(a), y(b))}{q^X(x(a))q^Y(y(b))} \hat{\rho}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} - \sum_{a=1}^A \sum_{b=1}^B \frac{p^{XY}(x(a), y(b))}{q^X(x(a))q^Y(y(b))} \hat{\rho}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \right\|_1 \right] \\ &\leq (AB)^{-1} \sum_{a=1}^A \sum_{b=1}^B \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \left\| \frac{p^{XY}(x(a), y(b))}{q^X(x(a))q^Y(y(b))} \left\| \hat{\rho}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} - \hat{\rho}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \right\|_1 \right\|_1 \right] \\ &= L^{-2} \sum_{l_x, x, l_y, y} q^X(x)q^Y(y) \frac{p^{XY}(x, y)}{q^X(x)q^Y(y)} \left\| \hat{\rho}_{l_x, x, l_y, y}^{\hat{M}} - \hat{\rho}_{l_x, x, l_y, y}^{\hat{M}} \right\|_1 \\ &= L^{-2} \sum_{l_x, x, l_y, y} p^{XY}(x, y) \left\| \hat{\rho}_{l_x, x, l_y, y}^{\hat{M}} - \hat{\rho}_{l_x, x, l_y, y}^{\hat{M}} \right\|_1 = \|\hat{\rho}^{L_X X L_Y Y \hat{M}} - \hat{\rho}^{L_X X L_Y Y \hat{M}}\|_1 < 25 \cdot \epsilon^{1/8}. \end{aligned}$$

Define

$$\text{Bad} := \{((l_x x)^{(A)}, (l_y y)^{(B)}) : \|\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} - \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}\|_1 > 25\epsilon^{1/16}\}.$$

By Markov's inequality,  $\Pr_{(l_x x)^{(A)}, (l_y y)^{(B)}} [\text{Bad}] < \epsilon^{1/16}$ . For  $((l_x x)^{(A)}, (l_y y)^{(B)}) \in \text{Bad}$ , we define  $\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} := \mathbf{0}$ . For  $((l_x x)^{(A)}, (l_y y)^{(B)}) \notin \text{Bad}$ , we apply Lemma 1 to the states  $\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}$ ,  $\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}$  to get a state  $\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}$  satisfying the following properties:

$$\begin{aligned} & \forall ((l_x x)^{(A)}, (l_y y)^{(B)}) : \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \leq \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}, \\ & \forall ((l_x x)^{(A)}, (l_y y)^{(B)}) \notin \text{Bad} : \|\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} - \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}\|_1 < 5\epsilon^{1/32}, \\ & \implies \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \left\| \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} - \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \right\|_1 \right] < 5\epsilon^{1/32} + \epsilon^{1/16} < 6\epsilon^{1/32}, \\ & \forall ((l_x x)^{(A)}, (l_y y)^{(B)}) : \|\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}\|_2^2 \leq (1 + 10\epsilon^{1/32}) \text{Tr} [\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}]. \end{aligned} \quad (55)$$

Hence by triangle inequality and Equations 54, 55, in order to prove Proposition 5, it suffices to show that

$$\mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} [\|\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} - \rho^{\hat{M}}\|_1] \leq 6\epsilon^{1/64} + \sqrt{40\epsilon^{1/32} + 8e + g(e)}. \quad (56)$$

Define  $\mathbb{1}^{F'}$  to be the orthogonal projector from the ambient space  $\hat{M}$  onto  $F'_\rho$ . Using Equations 51, 53, 55 (some of these equations were stated earlier for  $\text{Tr } \rho = 1$  but they continue to hold

in the natural fashion for general  $\text{Tr } \rho$  too), we get

$$\begin{aligned}
& \text{Tr } [\mathbb{1}^{F'_\rho} \rho^{\hat{M}}] \\
&= \text{Tr } [\rho^{\hat{M}}], \\
& \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\text{Tr } [\mathbb{1}^{F'_\rho} \hat{\sigma}'^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}}]] \\
&\geq \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} \left[ \text{Tr } [\mathbb{1}^{F'_\rho} \sigma^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}}] - \|\sigma^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} - \hat{\sigma}^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}}\|_1 \right. \\
&\quad \left. - \|\hat{\sigma}^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} - \hat{\sigma}'^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}}\|_1 \right] \\
&= \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\text{Tr } [\sigma^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}}]] \\
&\quad - \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\|\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1] \\
&\quad - \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\|\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1] \\
&= \text{Tr } \rho^{\hat{M}} - \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\|\sigma^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1] \\
&\quad - \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\|\hat{\sigma}^{(L_X X)^A (L_Y Y)^B \hat{M}} - \hat{\sigma}'^{(L_X X)^A (L_Y Y)^B \hat{M}}\|_1] \\
&\geq (\text{Tr } \rho^{\hat{M}})(1 - 2\sqrt{2} \cdot \epsilon^{1/8} - 6\epsilon^{1/32}) \geq (\text{Tr } \rho^{\hat{M}})(1 - 9\epsilon^{1/32}).
\end{aligned} \tag{57}$$

Using Proposition 1 and concavity of square root, we see that

$$\begin{aligned}
& \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\|\hat{\sigma}'^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} - \rho^{\hat{M}}\|_1] \\
&\leq \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} \left[ 2\sqrt{\text{Tr } [\rho^{\hat{M}}]} \sqrt{\text{Tr } [\rho^{\hat{M}}] - \text{Tr } [\mathbb{1}^{F'_\rho} \rho^{\hat{M}}]} \right. \\
&\quad \left. + 2\sqrt{\text{Tr } [\rho^{\hat{M}}]} \sqrt{\text{Tr } [\rho^{\hat{M}}] - \text{Tr } [\mathbb{1}^{F'_\rho} \hat{\sigma}'^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}}]} \right. \\
&\quad \left. + \sqrt{|F'_\rho|} \cdot \|\hat{\sigma}'^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} - \rho^{\hat{M}}\|_2 \right] \\
&\leq 2\sqrt{\text{Tr } [\rho^{\hat{M}}]} - \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\text{Tr } [\mathbb{1}^{F'_\rho} \hat{\sigma}'^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}}]] \\
&\quad + \sqrt{|F'_\rho|} \cdot \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\|\hat{\sigma}'^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} - \rho^{\hat{M}}\|_2] \\
&\leq 6\epsilon^{1/64} + \sqrt{|F'_\rho|} \cdot \mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\|\hat{\sigma}'^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} - \rho^{\hat{M}}\|_2]
\end{aligned}$$

From Equation 56 and convexity of the square function, in order to prove Proposition 5, it suffices to show that

$$\mathbb{E}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} [\|\hat{\sigma}'^{\hat{M}}_{(l_{xx})^{(A)}, (l_{yy})^{(B)}} - \rho^{\hat{M}}\|_2^2] < \frac{40\epsilon^{1/32} + 8e + g(e)}{|F'_\rho|}. \tag{58}$$

The left hand side of the above inequality is

$$\begin{aligned}
& \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} [\|\hat{\sigma}'_{(l_x x)^{(A)}, (l_y y)^{(B)}} - \rho^{\hat{M}}\|_2^2] \\
&= \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} [\|\hat{\sigma}'_{(l_x x)^{(A)}, (l_y y)^{(B)}}\|_2^2] + \|\rho^{\hat{M}}\|_2^2 \\
&\quad - 2 \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} [\text{Tr} [\hat{\sigma}'_{(l_x x)^{(A)}, (l_y y)^{(B)}} \rho^{\hat{M}}]] \\
&\leq (1 + 10\epsilon^{1/32}) \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \text{Tr} [\hat{\sigma}'_{(l_x x)^{(A)}, (l_y y)^{(B)}} \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}] \right] \\
&\quad + (1 + \sqrt{\epsilon})^2 \left\| \frac{\mathbf{1}_{F'_\rho}}{|F'_\rho|} \right\|_2^2 \\
&\quad - 2(1 - \sqrt{\epsilon}) \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \text{Tr} \left[ \hat{\sigma}'_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left( \frac{\mathbf{1}_{F'_\rho}}{|F'_\rho|} \right) \right] \right] \\
&= (1 + 10\epsilon^{1/32}) \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \text{Tr} [\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}] \right] + \frac{(1 + \sqrt{\epsilon})^2}{|F'_\rho|} \\
&\quad - \frac{2(1 - \sqrt{\epsilon})}{|F'_\rho|} \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \text{Tr} [\hat{\sigma}'_{(l_x x)^{(A)}, (l_y y)^{(B)}} \mathbf{1}_{F'_\rho}] \right] \\
&\leq (1 + 10\epsilon^{1/32}) \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \text{Tr} [\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}] \right] + \frac{(1 + \sqrt{\epsilon})^2}{|F'_\rho|} \\
&\quad - \frac{2(1 - \sqrt{\epsilon})(1 - 4e - 9\epsilon^{1/32})}{|F'_\rho|} \\
&\leq (1 + 10\epsilon^{1/32}) \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \text{Tr} [\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}] \right] \\
&\quad + \frac{8e + 23\epsilon^{1/32} - 1}{|F'_\rho|},
\end{aligned} \tag{59}$$

where the first inequality follows from Equations 55, 47, and the second inequality follows from Equations 57, 51.

From Equations 58 and 59, in order to prove Proposition 5, it suffices to show the following lemma.

**Lemma 18** *Suppose*

$$\begin{aligned}
\sqrt{L} &> \frac{8|F'_\rho|}{\epsilon^{3/8}} \cdot \max \left\{ \sum_{x,y} p^{XY}(x,y) \frac{p^X(x)}{q^X(x)} \|\rho(X)_{xy}^M\|_1, \right. \\
&\quad \left. \sum_{x,y} p^{XY}(x,y) \frac{p^Y(y)}{q^Y(y)} \|\rho(Y)_{xy}^M\|_1, \sum_{x,y} p^{XY}(x,y) \|\rho()_{xy}^M\|_1 \right\},
\end{aligned}$$

and

$$\begin{aligned}
\log A &> D_\infty^\epsilon(\rho^{XM} \| q^X \otimes \rho^M) + \log f(e) + 3 + \log \epsilon^{-1/2}, \\
\log B &> D_\infty^\epsilon(\rho^{YM} \| q^Y \otimes \rho^M) + \log f(e) + 3 + \log \epsilon^{-1/2}, \\
\log A + \log B &> D_\infty^\epsilon(\rho^{XYM} \| q^X \otimes q^Y \otimes \rho^M) + 1 + \log \epsilon^{-1/2}.
\end{aligned}$$

Then,

$$\mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \text{Tr} [\hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}}] \right] < \frac{1 + 6\epsilon^{1/2} + g(e)}{|F'_\rho|}.$$

**Proof:** Given the lower bound on  $\sqrt{L}$ , we can write down the following consequences of Lemmas 17,

15, 16, 14.

$$\begin{aligned}
& L^{-2} \sum_{l_x, x, l_y, y} q^X(x) q^Y(y) \text{Tr} \left[ \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
& \leq \frac{(1+4\sqrt{e}+2\sqrt{e})2^{D_\infty^\epsilon(\rho^{XYM} \| q^X \otimes q^Y \otimes \rho^M)}}{|F'_\rho|}, \\
& L^{-3} \sum_{l_x, x, l_y, y, l'_x, y'} q^X(x) \bar{q}^{YY'}(y, y') \text{Tr} \left[ \left( \frac{p^{XY}(x, y')}{q^X(x) q^Y(y')} \hat{\rho}_{l_x x l'_y y'}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
& \leq \frac{4(1+4\sqrt{e}+3\sqrt{e})f(e)2^{D_\infty^\epsilon(\rho^{XM} \| q^X \otimes \rho^M)}}{|F'_\rho|}, \\
& L^{-3} \sum_{l_x, x, l_y, y, l'_x, y'} q^Y(y) \bar{q}^{XX'}(x, x') \text{Tr} \left[ \left( \frac{p^{XY}(x', y)}{q^X(x') q^Y(y)} \hat{\rho}_{l'_x x' l_y y}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
& \leq \frac{4(1+4\sqrt{e}+3\sqrt{e})f(e)2^{D_\infty^\epsilon(\rho^{YM} \| q^Y \otimes \rho^M)}}{|F'_\rho|}, \\
& L^{-4} \sum_{l_x, x, l_y, y, l'_x, x', l'_y, y'} \bar{q}^{XX'}(x, x') \bar{q}^{YY'}(y, y') \text{Tr} \left[ \left( \frac{p^{XY}(x', y')}{q^X(x') q^Y(y')} \hat{\rho}_{l'_x x' l'_y y'}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
& \leq \frac{(1+g(e)+3\sqrt{e})}{|F'_\rho|}.
\end{aligned} \tag{60}$$

We have,

$$\begin{aligned}
& \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \text{Tr} \left[ \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \right] \right] \\
& = (AB)^{-2} \cdot \sum_{a, \hat{a}=1}^A \sum_{b, \hat{b}=1}^B \mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \frac{p^{XY}(x(\hat{a}), y(\hat{b}))}{q^X(x(\hat{a})) q^Y(y(\hat{b}))} \cdot \frac{p^{XY}(x(a), y(b))}{q^X(x(a)) q^Y(y(b))} \right. \\
& \quad \left. \text{Tr} \left[ \hat{\rho}_{(l_x x)(\hat{a}), (l_y y)(\hat{b})}^{\hat{M}} \hat{\rho}_{(l_x x)(a), (l_y y)(b)}^{\hat{M}} \right] \right] \\
& = (AB)^{-2} \cdot \sum_{a, \hat{a}=1}^A \sum_{b, \hat{b}=1}^B \mathbb{E}_{((l_x x)(\hat{a}), (l_x x)(a)), ((l_y y)(\hat{b}), (l_y y)(b))} \left[ \frac{p^{XY}(x(\hat{a}), y(\hat{b}))}{q^X(x(\hat{a})) q^Y(y(\hat{b}))} \cdot \frac{p^{XY}(x(a), y(b))}{q^X(x(a)) q^Y(y(b))} \right. \\
& \quad \left. \text{Tr} \left[ \hat{\rho}_{(l_x x)(\hat{a}), (l_y y)(\hat{b})}^{\hat{M}} \hat{\rho}_{(l_x x)(a), (l_y y)(b)}^{\hat{M}} \right] \right].
\end{aligned} \tag{61}$$

We analyse the above summation by considering several cases. Consider the following term for a fixed choice of  $a \neq \hat{a}$ ,  $b \neq \hat{b}$ .

$$\begin{aligned}
& \mathbb{E}_{((l_x x)(\hat{a}), (l_x x)(a)), ((l_y y)(\hat{b}), (l_y y)(b))} \left[ \frac{p^{XY}(x(\hat{a}), y(\hat{b}))}{q^X(x(\hat{a})) q^Y(y(\hat{b}))} \cdot \frac{p^{XY}(x(a), y(b))}{q^X(x(a)) q^Y(y(b))} \right. \\
& \quad \left. \text{Tr} \left[ \hat{\rho}_{(l_x x)(\hat{a}), (l_y y)(\hat{b})}^{\hat{M}} \hat{\rho}_{(l_x x)(a), (l_y y)(b)}^{\hat{M}} \right] \right] \\
& = L^{-4} \sum_{l_x, x, l_y, y, l'_x, x', l'_y, y'} \bar{q}^{XX'}(x, x') \bar{q}^{YY'}(y, y') \\
& \quad \text{Tr} \left[ \left( \frac{p^{XY}(x', y')}{q^X(x') q^Y(y')} \hat{\rho}_{l'_x x' l'_y y'}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x) q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
& \leq \frac{(1+g(e)+3\sqrt{e})}{|F'_\rho|},
\end{aligned} \tag{62}$$

where we used Equation 60 in the inequality above. There are  $A(A-1)B(B-1)$  such terms.

Consider the following term for a fixed choice of  $a = \hat{a}$ ,  $b \neq \hat{b}$ .

$$\begin{aligned}
& \mathbb{E}_{(l_x x)(a), (l_y y)(\hat{b}), (l_y y)(b)} \left[ \frac{p^{XY}(x(a), y(\hat{b}))}{q^X(x(a))q^Y(y(\hat{b}))} \cdot \frac{p^{XY}(x(a), y(b))}{q^X(x(a))q^Y(y(b))} \right. \\
& \quad \left. \text{Tr} [\hat{\rho}_{(l_x x)(a), (l_y y)(\hat{b})}^{\hat{M}} \hat{\rho}_{(l_x x)(a), (l_y y)(b)}^{\hat{M}}] \right] \\
&= L^{-3} \sum_{l_x, x, l_y, y, l'_y, y'} q^X(x) \bar{q}^{Y Y'}(y, y') \text{Tr} \left[ \left( \frac{p^{XY}(x, y')}{q^X(x)q^{Y'}(y')} \hat{\rho}_{l_x x l'_y y'}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x)q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
&\leq \frac{4(1+4\sqrt{e}+3\sqrt{e})f(e)2^{D_\infty^\epsilon(\rho^{XM} \| q^X \otimes \rho^M)}}{|F'_\rho|},
\end{aligned} \tag{63}$$

where we used Equation 60 in the inequality above. There are  $AB(B-1)$  such terms.

Consider the following term for a fixed choice of  $a \neq \hat{a}$ ,  $b = \hat{b}$ . We have similarly,

$$\begin{aligned}
& \mathbb{E}_{((l_x x)(\hat{a}), (l_x x)(a)), (l_y y)(b)} \left[ \frac{p^{XY}(x(\hat{a}), y(b))}{q^X(x(\hat{a}))q^Y(y(b))} \cdot \frac{p^{XY}(x(a), y(b))}{q^X(x(a))q^Y(y(b))} \right. \\
& \quad \left. \text{Tr} [\hat{\rho}_{(l_x x)(\hat{a}), (l_y y)(b)}^{\hat{M}} \hat{\rho}_{(l_x x)(a), (l_y y)(b)}^{\hat{M}}] \right] \\
&\leq \frac{4(1+4\sqrt{e}+3\sqrt{e})f(e)2^{D_\infty^\epsilon(\rho^{YM} \| q^Y \otimes \rho^M)}}{|F'_\rho|}.
\end{aligned} \tag{64}$$

There are  $A(A-1)B$  such terms.

Finally, consider the following term for a fixed choice of  $a = \hat{a}$ ,  $b = \hat{b}$ .

$$\begin{aligned}
& \mathbb{E}_{(l_x x)(a), (l_y y)(b)} \left[ \frac{p^{XY}(x(a), y(b))}{q^X(x(a))q^Y(y(b))} \cdot \frac{p^{XY}(x(a), y(b))}{q^X(x(a))q^Y(y(b))} \right. \\
& \quad \left. \text{Tr} [\hat{\rho}_{(l_x x)(a), (l_y y)(b)}^{\hat{M}} \hat{\rho}_{(l_x x)(a), (l_y y)(b)}^{\hat{M}}] \right] \\
&= L^{-2} \sum_{l_x, x, l_y, y} q^X(x) q^Y(y) \text{Tr} \left[ \left( \frac{p^{XY}(x, y)}{q^X(x)q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \left( \frac{p^{XY}(x, y)}{q^X(x)q^Y(y)} \hat{\rho}_{l_x x l_y y}^{\hat{M}} \right) \right] \\
&\leq \frac{(1+4\sqrt{e}+2\sqrt{e})2^{D_\infty^\epsilon(\rho^{XYM} \| q^X \otimes q^Y \otimes \rho^M)}}{|F'_\rho|},
\end{aligned} \tag{65}$$

where we used Equation 60 in the inequality above. There are  $AB$  such terms.

From Equations 61, 62, 63, 64, 65 we get

$$\begin{aligned}
& \mathbb{E}_{(l_x x)(A), (l_y y)(B)} \left[ \text{Tr} [\hat{\sigma}_{(l_x x)(A), (l_y y)(B)}^{\hat{M}} \hat{\sigma}_{(l_x x)(A), (l_y y)(B)}^{\hat{M}}] \right] \\
&= (AB)^{-2} \cdot \sum_{a, \hat{a}=1}^A \sum_{b, \hat{b}=1}^B \\
& \quad \mathbb{E}_{((l_x x)(\hat{a}), (l_x x)(a)), ((l_y y)(\hat{b}), (l_y y)(b))} \left[ \frac{p^{XY}(x(\hat{a}), y(\hat{b}))}{q^X(x(\hat{a}))q^Y(y(\hat{b}))} \cdot \frac{p^{XY}(x(a), y(b))}{q^X(x(a))q^Y(y(b))} \right. \\
& \quad \left. \text{Tr} [\hat{\rho}_{(l_x x)(\hat{a}), (l_y y)(\hat{b})}^{\hat{M}} \hat{\rho}_{(l_x x)(a), (l_y y)(b)}^{\hat{M}}] \right] \\
&\leq (AB)^{-2} \left( \frac{(A(A-1)B(B-1)(1+g(e)+3\sqrt{e}))}{|F'_\rho|} + \frac{4AB(B-1)(1+4\sqrt{e}+3\sqrt{e})f(e)2^{D_\infty^\epsilon(\rho^{XM} \| q^X \otimes \rho^M)}}{|F'_\rho|} \right. \\
& \quad + \frac{4A(A-1)B(1+4\sqrt{e}+3\sqrt{e})f(e)2^{D_\infty^\epsilon(\rho^{YM} \| q^Y \otimes \rho^M)}}{|F'_\rho|} \\
& \quad \left. + \frac{AB(1+4\sqrt{e}+2\sqrt{e})2^{D_\infty^\epsilon(\rho^{XYM} \| q^X \otimes q^Y \otimes \rho^M)}}{|F'_\rho|} \right) \\
&\leq \frac{(1+g(e)+3\sqrt{e})}{|F'_\rho|} + \frac{4(1+4\sqrt{e}+3\sqrt{e})f(e)2^{D_\infty^\epsilon(\rho^{XM} \| q^X \otimes \rho^M)}}{A|F'_\rho|} \\
& \quad + \frac{4(1+4\sqrt{e}+3\sqrt{e})f(e)2^{D_\infty^\epsilon(\rho^{YM} \| q^Y \otimes \rho^M)}}{B|F'_\rho|} + \frac{(1+4\sqrt{e}+2\sqrt{e})2^{D_\infty^\epsilon(\rho^{XYM} \| q^X \otimes q^Y \otimes \rho^M)}}{AB|F'_\rho|}.
\end{aligned} \tag{66}$$

From the constraints on  $\log A$ ,  $\log B$ , we get

$$\mathbb{E}_{(l_x x)^{(A)}, (l_y y)^{(B)}} \left[ \text{Tr} \left[ \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \hat{\sigma}_{(l_x x)^{(A)}, (l_y y)^{(B)}}^{\hat{M}} \right] \right] \leq \frac{(1 + g(e) + 6\sqrt{\epsilon})}{|F'_\rho|}.$$

This completes the proof of the lemma.  $\square$

Note that the entropic quantities in the statement of Lemma 18 are in terms of the original normalised probability distribution  $p^{XY}$ , and not in terms of  $\hat{p}^{XY}$  which was later renamed to  $p^{XY}$  for notational convenience. Since the dimension  $L$  of the augmenting Hilbert spaces  $L_X$ ,  $L_Y$  is a free parameter in our proof, it can be chosen large enough as in the requirement of Lemma 18. Thus Lemma 18 holds, which completes the proof of Proposition 5.

Arguing along similar lines, we can now prove a smooth multipartite soft covering lemma without pairwise independence. This is the main theorem of this paper.

**Theorem 1** *Let  $k$  be a positive integer. Let  $X_1, \dots, X_k$  be  $k$  classical alphabets. For any subset  $S \subseteq [k]$ , let  $X_S := (X_s)_{s \in S}$ . Let  $p^{X_{[k]}}$  be a normalised probability distribution on  $X_{[k]}$ . The notation  $p^{X_S}$  denotes the marginal distribution on  $X_S$ . Let  $q^{X_1}, \dots, q^{X_k}$  be normalised probability distributions on the respective alphabets. For each  $(x_1, \dots, x_k) \in X_{[k]}$ , let  $\rho_{x_1, \dots, x_k}^M$  be a subnormalised density matrix on  $M$ . The classical quantum control state is now defined as*

$$\rho^{X_{[k]}M} := \sum_{(x_1, \dots, x_k) \in X_{[k]}} p^{X_{[k]}}(x_1, \dots, x_k) |x_1, \dots, x_k\rangle \langle x_1, \dots, x_k|^{X_{[k]}} \otimes \rho_{x_1, \dots, x_k}^M.$$

Suppose  $\text{supp}(p^{X_i}) \leq \text{supp}(q^{X_i})$ . For any subset  $S \subseteq [k]$ , let  $q^{X_S} := \times_{s \in S} q^{X_s}$ . Let  $A_1, \dots, A_k$  be positive integers. For each  $i \in [k]$ , let  $x_i^{(A_i)} := (x_i(1), \dots, x_i(A_i))$  denote a  $|A_i|$ -tuple of elements from  $X_i$ . Denote the  $A_i$ -fold product alphabet  $X_i^{A_i} := X_i^{\times A_i}$ , and the product probability distribution  $q^{X_i^{A_i}} := (q^{X_i})^{\times A_i}$ . For any collection of tuples  $x_i^{(A_i)} \in X_i^{A_i}$ ,  $i \in [k]$ , we define the sample average covering state

$$\sigma_{x_1^{(A_1)}, \dots, x_k^{(A_k)}}^M := (A_1 \cdots A_k)^{-1} \sum_{a_1=1}^{A_1} \cdots \sum_{a_k=1}^{A_k} \frac{p^{X_{[k]}}(x_1(a_1), \dots, x_k(a_k))}{q^{X_1}(x_1(a_1)) \cdots q^{X_k}(x_k(a_k))} \rho_{x_1(a_1), \dots, x_k(a_k)}^M,$$

where the fraction term above represents the ‘change of measure’ from the product probability distribution  $q^{X_{[k]}}$  to the joint probability distribution  $p^{X_{[k]}}$ .

Let the covering be done using independent choices of tuples  $x_1^{(A_1)}, \dots, x_k^{(A_k)}$  from the distributions  $\bar{q}^{X_1^{A_1}}, \dots, \bar{q}^{X_k^{A_k}}$ , which satisfy the consequences of loss of pairwise independence, akin to the situation in Proposition 5, up to scale factors of  $(e, f(e), g(e))$ . Suppose for each non-empty subset  $\{i\} \neq S \subseteq [k]$ ,

$$\sum_{s \in S} \log A_s > D_\infty^\epsilon(\rho^{X_S M} \| q^{X_S} \otimes \rho^M) + \log f(e) + 3 + \log \epsilon^{-1/2}.$$

Then,

$$\mathbb{E}_{x_1^{(A_1)}, \dots, x_k^{(A_k)}} \left[ \left\| \sigma_{x_1^{(A_1)}, \dots, x_k^{(A_k)}}^M - \rho^M \right\|_1 \right] < (2\sqrt{10 \cdot 2^k \epsilon^{1/64}} + \sqrt{30 \cdot 2^{2k} \epsilon^{1/32} + 2^{k+1} e + g(e) + 2^k e}) (\text{Tr } \rho).$$



## Remarks:

1. The scale factor of  $f(e)$  in the consequences of loss of pairwise independence in the statement of Theorem 1 means that for any sequence of coordinates  $(a_1, \dots, a_k)$ ,  $a_i \in [A_i]$ , for any sequence of symbols  $(x_{a_1}, \dots, x_{a_k})$ , for any non-empty subset  $\{\} \neq S \subset [k]$ , for any sequence of coordinates  $(a'_s)_{s \in S}$  with  $a_s \neq a'_s$  for all  $s \in S$ ,

$$\mathbb{E}_{(x'_{a'_s})_{s \in S} | (x_{a_s})_{s \in S}} [\rho^M_{(x_{a_t})_{t \in \bar{S}}, (x'_{a'_s})_{s \in S}}] =: \rho(\bar{S})^M_{x_{a_1}, \dots, x_{a_k}} \leq f(e) \rho^M_{(x_{a_t})_{t \in \bar{S}}},$$

where the expectation is taken over independent choices from the distributions  $\bar{q}^{X'_s | X_s = x_{a_s}}$ ,  $s \in S$ , and  $\rho^M_{(x_{a_t})_{t \in \bar{S}}}$  is the quantum state of register  $M$  when the symbols  $(x_{a_t})_{t \in \bar{S}}$  are fixed and the symbols in  $(a_s)_{s \in S}$  are averaged over according to independent choices from the distribution  $q^{X_S}$ .

2. A factor of  $k$  comes in the error bound on the right hand side of Theorem 1 due to using Lemma 2 for  $k$  orthogonal tilting directions that arise from the  $k$  covering parties. However it is subsumed by factors of  $2^k$  that arise from the  $k$  party analogue of Lemma 3. The factor of  $2^{k+1}e$  in the error bound arises from the  $k$  party analogue of Equation 36, where we now have to do  $2^k - 1$  intersections of classical probability distributions.

Similarly, we can also show the following smooth version of CMG covering without pairwise independence.

**Lemma 19** *Consider the CMG covering problem of Section 2, where there is a loss of pairwise independence in Alice's obfuscation strategy when she chooses  $L$  symbols  $x_{\nu,1}, \dots, x_{\nu,L} \in \mathcal{X}$  conditioned on a sample  $x'_{\nu} \in \mathcal{X}'$ , and a similar loss of pairwise independence in Bob's obfuscation strategy. Suppose these are the only two places of loss of pairwise independence, with the consequences of the loss holding up to scale factors of  $(e, f(e), 0)$ . Suppose*

$$\begin{aligned} \log L' &> I_{\infty}^e(X' : E)_{\sigma} + 3 + \log \epsilon^{-1/2}, \\ \log M' &> I_{\infty}^e(Y' : E)_{\sigma} + 3 + \log \epsilon^{-1/2}, \\ \log L' + \log L &> I_{\infty}^e(X'X : E)_{\sigma} + 3 + \log \epsilon^{-1/2}, \\ \log M' + \log M &> I_{\infty}^e(Y'Y : E)_{\sigma} + 3 + \log \epsilon^{-1/2}, \\ \log L' + \log M' &> I_{\infty}^e(X'Y' : E)_{\sigma} + 3 + \log f(e) + \log \epsilon^{-1/2}, \\ \log L' + \log M' + \log M &> I_{\infty}^e(X'Y'Y : E)_{\sigma} + 3 + \log f(e) + \log \epsilon^{-1/2}, \\ \log L' + \log M' + \log L &> I_{\infty}^e(X'XY' : E)_{\sigma} + 3 + \log f(e) + \log \epsilon^{-1/2}, \\ \log L' + \log M' + \log L + \log M &> I_{\infty}^e(X'XY'Y : E)_{\sigma} + 3 + \log \epsilon^{-1/2}. \end{aligned}$$

Then,  $\text{CMG} < 2\sqrt{160 \cdot \epsilon^{1/64} + \sqrt{750\epsilon^{1/32} + 8e} + 4e}$

## 6 Inner bound for the wiretap interference channel

In a wiretap quantum interference channel, there are two senders Alice and Bob and their corresponding intended receivers Charlie and Damru. There is a third receiver Eve, corresponding to an eavesdropper. Alice would like to send a classical message  $m_1 \in [2^{R_1}]$  to Charlie. Similarly, Bob would like to send  $m_2 \in [2^{R_2}]$  to Damru. Charlie and Damru should be able to decode correctly with large probability at their respective channel outputs and recover  $m_1, m_2$ . It is okay if Damru

gets some information about Alice's message  $m_1$  in addition to the message  $m_2$  intended for him; similarly for Charlie. However, Eve should have at most negligible information about any message tuple  $(m_1, m_2)$ .

The parties have at their disposal a quantum channel  $\mathcal{N} : AB \rightarrow CDE$  with input Hilbert spaces  $\mathcal{A}, \mathcal{B}$  and output Hilbert spaces  $\mathcal{C}, \mathcal{D}, \mathcal{E}$ . Though  $\mathcal{N}$  has quantum inputs and quantum outputs, we will assume without loss of generality that the inputs are classical, because in the inner bound below, there is a standard optimisation step over a choice of all ensembles of input states to Alice and Bob. In other words, we fix classical to quantum encodings  $x \mapsto \rho_x^A, y \mapsto \rho_y^B$ . Thus, we will henceforth think of  $\mathcal{N}$  as a classical quantum (cq) channel with its two classical input alphabets being denoted by  $\mathcal{X}, \mathcal{Y}$ , and the quantum output alphabets by  $\mathcal{C}, \mathcal{D}, \mathcal{E}$ .

We now describe the strategy of Chong, Motani, Garg, El Gamal [CMGE08], adapted to the one shot quantum case by [Sen21a], to achieve a rate inner bound for sending  $(m_1, m_2)$  over  $\mathcal{N}$  when there is no eavesdropper Eve. As shown below, we will suitably modify that strategy to ensure privacy against Eve. For brevity, we will call the overall coding strategy as the CMG wiretap strategy. For this, we need to define additional classical alphabets  $\mathcal{X}', \mathcal{Y}'$ , another 'timesharing' classical alphabet  $\mathcal{Q}$ , a 'control probability distribution'

$$p(q, x', x, y', y) = p(q)p(x', x)p(y', y), \quad (67)$$

and a 'control classical quantum state'

$$\sigma^{QX'XY'YCDE} := \sum_{q, x', x, y', y} p(q)p(x', x)p(y', y) |q, x', x, y', y\rangle \langle q, x', x, y', y|^{QX'XY'Y} \otimes \sigma_{x'xy'y}^{CDE}, \quad (68)$$

where  $\sigma_{x'xy'y}^{CDE} := \sigma_{xy}^{CDE}$  similar to the definitions in Section 2.

The CMG wiretap strategy uses *rate splitting* for both Alice and Bob. By this we mean that Alice 'splits' her message  $m_1 \in [2^{R_1}]$  into a pair  $(m'_1, m''_1) \in [2^{R'_1}] \times [2^{R_1 - R'_1}]$ . Similarly Bob 'splits' his message  $m_2 \in [2^{R_2}]$  into a pair  $(m'_2, m''_2) \in [2^{R'_2}] \times [2^{R_2 - R'_2}]$ . The intention is that  $m'_1, m'_2$  should be decodable by both Charlie and Damru. The message  $m'_1$ , which should of course be decodable by Charlie, need not satisfy any requirement with respect to Damru; similarly for message  $m'_2$ . Of course, the state of Eve for any quadruple  $(m'_1, m''_1, m'_2, m''_2)$  should be almost independent of the quadruple for the privacy requirement to be met. The intuition, though not the actual decoding strategy, is that the  $m'_1, m'_2$  parts of the encoded messages are 'easy' to decode at both Charlie and Damru. So intuitively Charlie can decode  $m'_2$  and use it as side information in order to decode  $m''_1$ ; similarly for Damru. It is known that this strategy of rate splitting enhances the achievable inner bound in many important examples of interference channels. The quantities  $R'_1, R'_2$  can be optimised in a later step in order to get the largest inner bound possible.

We now describe the codebook, Alice and Bob's encoding and obfuscation, and Charlie and Damru's decoding strategies.

### Codebook:

Let  $0 \leq \epsilon \leq 1$ . Let  $L', L, M', M$  be positive integers. As will become clear very soon, these four integers play the obfuscation role as described for the CMG covering problem in Section 2. First generate a sample  $q$  from the distribution  $p^Q$ . For each message  $m'_1 \in [2^{R'_1}]$  independently generate samples  $x'(m'_1, 1), \dots, x'(m'_1, L')$  from the distribution  $p^{X'|Q=q}$ . Similarly, for each message  $m'_2 \in [2^{R'_2}]$  independently generate samples  $y'(m'_2, 1), \dots, y'(m'_2, M')$  from the distribution  $p^{Y'|Q=q}$ . Now for each codeword  $x'(m'_1, l')$ ,  $l' \in [L']$  independently generate samples  $x(m'_1, l', m''_1, l)$  from

the distribution  $p^{X|X'=x'(m'_1, l'), Q=q}$  for all  $(m''_1, l) \in [2^{R_1-R'_1}] \times [L]$ . Similarly for each codeword  $y'(m'_2, m')$ ,  $m' \in [M']$ , independently generate samples  $y(m'_2, m', m''_2, m)$  from the probability distribution  $p^{Y|Y'=y'(m'_2, m'), Q=q}$  for all  $(m''_2, m) \in [2^{R_2-R'_2}] \times [M]$ . These samples together constitute the random codebook. We augment the random codebook as in [Sen21a] to get a random augmented codebook. This random augmented codebook is revealed to Alice, Bob, Charlie, Damru and Eve.

### Encoding and obfuscation:

To send message  $m_1 = (m'_1, m''_1)$ , Alice picks up a symbol  $x(m'_1, l', m''_1, l)$  for a uniformly random pair  $(l', l) \in [L'] \times [L]$  from the codebook and inputs the state  $\rho_{x(m'_1, l', m''_1, l)}^A$  into the input  $\mathcal{A}$  of the channel  $\mathcal{N}$ . Similarly, to send message  $m_2 = (m'_2, m''_2)$ , Bob picks up the symbol  $y(m'_2, m', m''_2, m)$  for a uniformly random pair  $(m', m) \in [M'] \times [M]$  from the codebook and inputs the state  $\rho_{y(m'_2, m', m''_2, m)}^B$  into the input  $\mathcal{B}$  of  $\mathcal{N}$ .

### Decoding:

Charlie decodes the tuple  $m_1 = (m'_1, m''_1)$  using *simultaneous non-unique decoding* as explained in [Sen21a], obtaining his guess  $\hat{m}_1$  for  $m_1$ . Damru decodes the tuple  $m_2 = (m'_2, m''_2)$  using *simultaneous non-unique decoding*, obtaining his guess  $\hat{m}_2$  for  $m_2$ . By simultaneous decoding, we mean that Charlie's decoded pair  $(\hat{m}'_1, \hat{m}''_1)$  is obtained by applying a single POVM element to the state at the channel output  $\mathcal{C}$ ; this POVM element is not composed from 'simpler' POVM elements like e.g. the successive cancellation decoder which first decodes, say  $\hat{m}'_1$ , and then uses it as side information to decode  $\hat{m}''_1$ . By non-unique decoding, we mean that Charlie's decoding POVM element for a fixed tuple  $(\hat{m}'_1, \hat{m}''_1)$  can be thought of as a 'union', ranging over  $(l', l, m'_2) \in [L'] \times [L] \times [2^{R'_2}]$ , of projectors for each possible tuple  $(\hat{m}'_1, l', \hat{m}''_1, l, m'_2)$ . The 'actual or unique identity' of  $(l', l, m'_2)$  in the 'union' is immaterial as long as the correct  $(\hat{m}'_1, \hat{m}''_1) = (m'_1, m''_1)$  is decoded.

### Correctness:

We want that  $\Pr[(\hat{m}_1, \hat{m}_2) \neq (m_1, m_2)] \leq \epsilon$ , where the probability is over the uniform choice of the message pair  $(m_1, m_2) \in [2^{R_1}] \times [2^{R_2}]$ , and actions of the encoder, channel and decoders.

### Privacy:

We want that  $\mathbb{E}_{(m_1, m_2)}[\|\sigma_{m_1, m_2}^E - \sigma^E\|_1] \leq \epsilon$ , where the expectation is over the uniform choice of the message pair  $(m_1, m_2) \in [2^{R_1}] \times [2^{R_2}]$ ,  $\sigma_{m_1, m_2}^E$  denotes the quantum state of Eve when message pair  $(m_1, m_2)$  is transmitted, and  $\sigma^E$  is the marginal of the control state on  $E$  from Equation 68. Note that  $\sigma_{m_1, m_2}^E$  can be defined from the control state  $\sigma^{QX'XY'YE}$  given the codebook.

If there exist such coding schemes for a particular channel  $\mathcal{N}$ , we say that there exists an  $(R_1, R_2, \epsilon)$ -wiretap quantum interference channel code for sending private classical information through  $\mathcal{N}$ .

We are interested in the variant of the codebook construction where pairwise independence is lost in the following step of constructing Alice's part of the codebook.

Now for each codeword  $x'(m'_1, l')$ ,  $l' \in [L']$  independently generate samples  $x(m'_1, l', m''_1, l)$  from the distribution  $p^{X|X'=x'(m'_1, l'), Q=q}$  for all  $(m''_1, l) \in [2^{R_1-R'_1}] \times [L]$ .

A similar loss of pairwise independence occurs in the corresponding step of constructing Bob's part of the codebook. There is no further loss of pairwise independence. The consequences of the loss of pairwise independence hold up to scale factors of  $(e, f(e), 0)$ . Loss of pairwise independence in these parts of codebook construction makes computational sense as these parts require the largest amount of random bits because they contain most of the codeword samples.

It turns out that the simultaneous non-unique decoding strategy for Charlie and Damru without any privacy considerations, already described in [Sen21a], continues to work with a natural slight degradation of parameters when pairwise independence is lost up to scale factors of  $(e, f(e), 0)$ . This is because the strategy is based on the same machinery of tilting and augmentation smoothing. The requirement of privacy against Eve comes from the CMG covering problem. We have already shown a smooth inner bound for this problem under loss of pairwise independence in Lemma 19. Combining both simultaneous non-unique decoding for Charlie, Damru together with the CMG covering obfuscation against Eve, we get the following inner bound for sending private classical information over a quantum interference channel.

**Theorem 2** *Consider the quantum interference channel with classical quantum control state  $\sigma^{QX'XY'YCDE}$  defined in Equation 68. Let  $0 < \epsilon < 1$ . Consider a rate octuple  $(R'_1, L', R_1 - R'_1, L, R'_2, M', R_2 - R'_2, M)$  satisfying the following inequalities:*

$$\begin{aligned}
R_1 - R'_1 + L &< I_H^\epsilon(X : C|X'Y'Q)_\sigma - \log f(e) - 2 - \log \epsilon^{-1} \\
R_1 + L' + L &< I_H^\epsilon(X : C|Y'Q)_\sigma - 2 - \log \epsilon^{-1} \\
R_1 - R'_1 + L + R'_2 + M' &< I_H^\epsilon(XY' : C|X'Q)_\sigma - \log f(e) - 2 - \log \epsilon^{-1} \\
R_1 + L' + L + R'_2 + M' &< I_H^\epsilon(XY' : C|Q)_\sigma - 2 - \log \epsilon^{-1} \\
R_2 - R'_2 + M &< I_H^\epsilon(Y : D|X'Y'Q)_\sigma - \log f(e) - 2 - \log \epsilon^{-1} \\
R_2 + M' + M &< I_H^\epsilon(Y : D|X'Q)_\sigma - 2 - \log \epsilon^{-1} \\
R_2 - R'_2 + M + R'_1 + L' &< I_H^\epsilon(YX' : D|Y'Q)_\sigma - \log f(e) - 2 - \log \epsilon^{-1} \\
R_2 + M' + M + R'_1 + L' &< I_H^\epsilon(YX' : D|Q)_\sigma - 2 - \log \epsilon^{-1} \\
\\ 
L' &> I_\infty^\epsilon(X' : E)_\sigma + 3 + \log \epsilon^{-1}, \\
L' + L &> I_\infty^\epsilon(X'X : E)_\sigma + 3 + \log \epsilon^{-1}, \\
M' &> I_\infty^\epsilon(Y' : E)_\sigma + 3 + \log \epsilon^{-1}, \\
M' + M &> I_\infty^\epsilon(Y'Y : E)_\sigma + 3 + \log \epsilon^{-1}, \\
L' + M' &> I_\infty^\epsilon(X'Y' : E)_\sigma + 3 + \log f(e) + \log \epsilon^{-1}, \\
L' + M' + M &> I_\infty^\epsilon(X'Y'Y : E)_\sigma + 3 + \log f(e) + \log \epsilon^{-1}, \\
L' + M' + L &> I_\infty^\epsilon(X'XY' : E)_\sigma + 3 + \log f(e) + \log \epsilon^{-1}, \\
L' + M' + L + M &> I_\infty^\epsilon(X'XY'Y : E)_\sigma + 3 + \log \epsilon^{-1}.
\end{aligned}$$

Then there exists an  $(R_1, R_2, \alpha)$ -wiretap quantum interference channel code for sending private classical information through  $\mathcal{N}$  where

$$\alpha < 2^{2^{14}} \epsilon^{1/6} + 2\sqrt{160 \cdot \epsilon^{1/64} + \sqrt{750\epsilon^{1/32} + 8e} + 4e}.$$

Standard Fourier-Motzkin elimination of  $R'_1, R'_2, L', L, M', M$  can now be used to obtain an inner bound in terms of only  $R_1, R_2$ .

The above one shot theorem automatically gives us the following corollary in the asymptotic iid limit.

**Corollary 3** *Consider the setting of Theorem 2. Consider an octuple of rate per channel use  $(R'_1, L', R_1 - R'_1, L, R'_2, M', R_2 - R'_2, M)$  satisfying the following inequalities:*

$$\begin{aligned}
R_1 - R'_1 + L &< I(X : C|X'Y'Q)_\sigma - \log f(e) \\
R_1 + L' + L &< I(X : C|Y'Q)_\sigma \\
R_1 - R'_1 + L + R'_2 + M' &< I(XY' : C|X'Q)_\sigma - \log f(e) \\
R_1 + L' + L + R'_2 + M' &< I(XY' : C|Q)_\sigma \\
R_2 - R'_2 + M &< I(Y : D|X'Y'Q)_\sigma - \log f(e) \\
R_2 + M' + M &< I(Y : D|X'Q)_\sigma \\
R_2 - R'_2 + M + R'_1 + L' &< I(YX' : D|Y'Q)_\sigma - \log f(e) \\
R_2 + M' + M + R'_1 + L' &< I(YX' : D|Q)_\sigma
\end{aligned}$$

$$\begin{aligned}
L' &> I(X' : E)_\sigma \\
L' + L &> I(X'X : E)_\sigma \\
M' &> I(Y' : E)_\sigma \\
M' + M &> I(Y'Y : E)_\sigma \\
L' + M' &> I(X'Y' : E)_\sigma + \log f(e) \\
L' + M' + M &> I(X'Y'Y : E)_\sigma + \log f(e) \\
L' + M' + L &> I(X'XY' : E)_\sigma + \log f(e) \\
L' + M' + L + M &> I(X'XY'Y : E)_\sigma.
\end{aligned}$$

Then there exists a per channel use rate  $(R_1, R_2)$ -wiretap quantum interference channel code for sending private classical information through  $\mathcal{N}$  with asymptotically vanishing decoding and privacy error. Standard Fourier-Motzkin elimination of  $R'_1, R'_2, L', L, M', M$  can now be used to obtain an inner bound in terms of only  $R_1, R_2$ .

We note that the above corollary with loss of pairwise independence was unknown earlier even in the classical asymptotic iid setting.

## 7 Simpler proof of smooth multipartite convex split

In this section, we give a simpler proof of Lemma 7 which only tilts and augments the registers  $X, Y$  in the state  $\sigma^{X^A Y^B M}$ . The augmentation only doubles the dimensions of  $X, Y$ . The register  $M$  is not affected. The advantage of the simpler style of proof is that it can be extended to prove our variant of the fully smooth decoupling theorem in the next section. The simpler proof style is somewhat dual to the more involved proof in Section 4 which tilted and augmented register  $M$  but left registers  $X, Y$  unchanged. The advantage of the more involved proof was that it could be extended to prove the smooth multipartite soft covering lemma without pairwise independence in Section 5.

As in Section 4, we assume without loss of generality that the density matrix  $\rho^{XYM}$  is normalised. We flatten by applying the superoperators  $\mathcal{F}_\alpha^{X \rightarrow L_X X}, \mathcal{F}_\beta^{Y \rightarrow L_Y Y}, \mathcal{F}_\rho^{M \rightarrow L_M M}$  to  $\rho^{XYM}$ , followed by renaming  $X \equiv L_X X, Y \equiv L_Y Y$  and  $M \equiv L_M M$  as in Section 4. Thus, we get orthog-

onal projectors  $\Pi^{XYM}(3)$ ,  $\Pi^{XM}(1)$  and  $\Pi^{YM}(2)$  satisfying Equation 16 as in Section 4. To finish the proof of Lemma 7, we only need to show the validity of Equation 17 of Section 4.

We will now need to augment and tilt the Hilbert spaces  $X$ ,  $Y$ . We will not touch  $M$ . This is the place from where the present proof starts differing from the proof in Section 4. Define new Hilbert spaces  $\tilde{X}$ ,  $\tilde{Y}$  isomorphic to  $X$ ,  $Y$ . Define the natural isometric bijections  $|x\rangle^X \mapsto |x\rangle^{\tilde{X}}$ ,  $|y\rangle^Y \mapsto |y\rangle^{\tilde{Y}}$ . Define the augmented Hilbert spaces  $\hat{X} := X \oplus \tilde{X}$ ,  $\hat{Y} := Y \oplus \tilde{Y}$ . Let  $0 < \epsilon < 1$ . Define the tilting map

$$T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} : |x\rangle^X \mapsto \sqrt{1 - \epsilon^{1/4}} |x\rangle^X + \epsilon^{1/8} |x\rangle^{\tilde{X}}.$$

Define  $T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}}$  similarly. Define

$$\begin{aligned} \rho^{\hat{X}\hat{Y}M} &\equiv \rho^{XYM}, \\ \hat{\rho}^{\hat{X}\hat{Y}M} &:= (T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\rho^{XYM}), \\ \hat{\Pi}^{\hat{X}\hat{Y}M} &:= \mathbb{1}^{\hat{X}\hat{Y}M} \\ &\quad - \text{span} \{ ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM} - \Pi(1)^{XM})) \otimes \mathbb{1}^{\hat{Y}}, \\ &\quad ((T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{YM} - \Pi(2)^{YM})) \otimes \mathbb{1}^{\hat{X}}, \\ &\quad (T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM} - \Pi(3)^{XYM}) \}, \\ \hat{\rho}^{\hat{X}\hat{Y}M} &:= \hat{\Pi}^{\hat{X}\hat{Y}M} \circ \rho^{\hat{X}\hat{Y}M}, \\ \alpha^{\hat{X}} &:= \alpha^X, \quad \beta^{\hat{Y}} := \beta^Y, \\ \tau^{\hat{X}^A \hat{Y}^B M} &\equiv \tau^{X^A Y^B M} := \alpha^{X^A} \otimes \beta^{Y^B} \otimes \rho^M, \\ \sigma^{\hat{X}^A \hat{Y}^B M} &\equiv \sigma^{X^A Y^B M} := \frac{1}{AB} \sum_{a=1}^A \sum_{b=1}^B \rho^{X_a Y_b M} \otimes \alpha^{X^{-a}} \otimes \beta^{Y^{-b}} \\ &\equiv \frac{1}{AB} \sum_{a=1}^A \sum_{b=1}^B \rho^{\hat{X}_a \hat{Y}_b M} \otimes \alpha^{\hat{X}^{-a}} \otimes \beta^{\hat{Y}^{-b}}, \\ \hat{\sigma}^{\hat{X}^A \hat{Y}^B M} &:= \frac{1}{AB} \sum_{a=1}^A \sum_{b=1}^B \hat{\rho}^{\hat{X}_a \hat{Y}_b M} \otimes \alpha^{\hat{X}^{-a}} \otimes \beta^{\hat{Y}^{-b}}, \\ \hat{\hat{\sigma}}^{\hat{X}^A \hat{Y}^B M} &:= \frac{1}{AB} \sum_{a=1}^A \sum_{b=1}^B \hat{\hat{\rho}}^{\hat{X}_a \hat{Y}_b M} \otimes \alpha^{\hat{X}^{-a}} \otimes \beta^{\hat{Y}^{-b}}. \end{aligned} \tag{69}$$

The following properties can be proved easily as in Section 4.

$$\begin{aligned} \hat{\rho}^{\hat{X}M} &= (T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\rho^{XM}), \quad \hat{\rho}^{\hat{Y}M} = (T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\rho^{YM}), \\ \|\hat{\rho}^{\hat{X}\hat{Y}M} - \rho^{\hat{X}\hat{Y}M}\|_1 &\leq 4\sqrt{2}\epsilon^{1/8}, \\ \text{Tr} [\hat{\rho}^{\hat{X}\hat{Y}M}] &\geq 1 - 121\epsilon^{1/4}, \\ \implies \|\hat{\hat{\rho}}^{\hat{X}\hat{Y}M} - \hat{\rho}^{\hat{X}\hat{Y}M}\|_1 &\leq 22\epsilon^{1/8}, \quad \|\hat{\hat{\rho}}^{\hat{X}\hat{Y}M} - \hat{\rho}^{\hat{X}\hat{Y}M}\|_1 \leq 28\epsilon^{1/8}, \\ \implies \|\hat{\hat{\sigma}}^{\hat{X}^A \hat{Y}^B M} - \sigma^{\hat{X}^A \hat{Y}^B M}\|_1 &\leq 4\sqrt{2}\epsilon^{1/8}, \quad \|\hat{\hat{\sigma}}^{\hat{X}^A \hat{Y}^B M} - \hat{\sigma}^{\hat{X}^A \hat{Y}^B M}\|_1 \leq 28\epsilon^{1/8}. \end{aligned} \tag{70}$$

By Lemma 1, there exists a subnormalised density matrix  $\sigma'^{\hat{X}^A \hat{Y}^B M}$  such that

$$\begin{aligned} \|\sigma'^{\hat{X}^A \hat{Y}^B M} - \hat{\sigma}^{\hat{X}^A \hat{Y}^B M}\|_1 &\leq \sqrt{28}\epsilon^{1/16}, \\ \sigma'^{\hat{X}^A \hat{Y}^B M} &\leq \hat{\sigma}^{\hat{X}^A \hat{Y}^B M}, \\ \|\sigma'^{\hat{X}^A \hat{Y}^B M}\|_2^2 &\leq (1 + 11\epsilon^{1/16}) \text{Tr} [\hat{\sigma}^{\hat{X}^A \hat{Y}^B M} \hat{\hat{\sigma}}^{\hat{X}^A \hat{Y}^B M}], \\ \implies \|\sigma'^{\hat{X}^A \hat{Y}^B M} - \sigma^{\hat{X}^A \hat{Y}^B M}\|_1 &< 9\epsilon^{1/16}. \end{aligned} \tag{71}$$

From Equation 17 and the above calculation, it suffices to show

$$\|\sigma'^{\hat{X}^A \hat{Y}^B M} - \tau^{\hat{X}^A \hat{Y}^B M}\|_1 < 11\epsilon^{1/32}.$$

Define  $\Pi^{\hat{X}^A \hat{Y}^B M}$  to be the projector from  $\hat{X}^A \hat{Y}^B M$  onto  $\text{supp}(\tau^{\hat{X}^A \hat{Y}^B M}) = F'_\alpha{}^{\otimes A} \otimes F'_\beta{}^{\otimes B} \otimes F'_\rho$ . Then,

$$\text{Tr} [\sigma'^{\hat{X}^A \hat{Y}^B M} \Pi^{\hat{X}^A \hat{Y}^B M}] \geq \text{Tr} [\sigma^{\hat{X}^A \hat{Y}^B M} \Pi^{\hat{X}^A \hat{Y}^B M}] - \frac{1}{2} \|\sigma'^{\hat{X}^A \hat{Y}^B M} - \sigma^{\hat{X}^A \hat{Y}^B M}\|_1 \geq 1 - 5\epsilon^{1/16}, \quad (72)$$

as  $\text{Tr} [\sigma^{\hat{X}^A \hat{Y}^B M} \Pi^{\hat{X}^A \hat{Y}^B M}] = 1$  because  $\text{supp}(\sigma^{\hat{X}^A \hat{Y}^B M}) \leq \Pi^{\hat{X}^A \hat{Y}^B M}$ . By Proposition 1, it thus suffices to show

$$\sqrt{\text{Tr} [\Pi^{\hat{X}^A \hat{Y}^B M}]} \cdot \|\sigma'^{\hat{X}^A \hat{Y}^B M} - \tau^{\hat{X}^A \hat{Y}^B M}\|_2 < 11\epsilon^{1/32} - 5\epsilon^{1/32} = 6\epsilon^{1/32},$$

which implies that it suffices to show,

$$\|\sigma'^{\hat{X}^A \hat{Y}^B M} - \tau^{\hat{X}^A \hat{Y}^B M}\|_2^2 < \frac{36\epsilon^{1/16}}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|}. \quad (73)$$

Using Equations 71, 72, 16, the left hand side of the above equation can be simplified as

$$\begin{aligned} & \|\sigma'^{\hat{X}^A \hat{Y}^B M} - \tau^{\hat{X}^A \hat{Y}^B M}\|_2^2 \\ &= \|\sigma'^{\hat{X}^A \hat{Y}^B M}\|_2^2 - 2 \text{Tr} [\sigma'^{\hat{X}^A \hat{Y}^B M} \tau^{\hat{X}^A \hat{Y}^B M}] + \|\tau^{\hat{X}^A \hat{Y}^B M}\|_2^2 \\ &\leq (1 + 11\epsilon^{1/16}) \text{Tr} [\hat{\sigma}^{\hat{X}^A \hat{Y}^B M} \hat{\sigma}^{\hat{X}^A \hat{Y}^B M}] - \frac{2(1 - \sqrt{\epsilon}) \text{Tr} [\sigma'^{\hat{X}^A \hat{Y}^B M} \Pi^{\hat{X}^A \hat{Y}^B M}]}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|} + \frac{1 + \sqrt{\epsilon}}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|} \\ &< (1 + 11\epsilon^{1/16}) \text{Tr} [\hat{\sigma}^{\hat{X}^A \hat{Y}^B M} \hat{\sigma}^{\hat{X}^A \hat{Y}^B M}] - \frac{2(1 - \sqrt{\epsilon})(1 - 5\epsilon^{1/16})}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|} + \frac{1 + \sqrt{\epsilon}}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|} \\ &< (1 + 11\epsilon^{1/16}) \text{Tr} [\hat{\sigma}^{\hat{X}^A \hat{Y}^B M} \hat{\sigma}^{\hat{X}^A \hat{Y}^B M}] - \frac{1 - 13\epsilon^{1/16}}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|}. \end{aligned}$$

From the above and Equation 73, it thus suffices to show

$$\text{Tr} [\hat{\sigma}^{\hat{X}^A \hat{Y}^B M} \hat{\sigma}^{\hat{X}^A \hat{Y}^B M}] < \frac{1 + \epsilon^{1/16}}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|}. \quad (74)$$

We now prove four lemmas which are crucially required to prove Equation 74. They are the analogues of Lemmas 9, 10, 11, 12 in Section 4.

**Lemma 20**

$$\text{Tr} [\hat{\rho}^{\hat{X}^A \hat{Y}^B M} \hat{\rho}^{\hat{X}^A \hat{Y}^B M}] \leq \frac{(1 + 3\sqrt{\epsilon}) 2^{D_\infty^\epsilon(\rho^{XYM} \|\alpha^X \otimes \beta^Y \otimes \rho^M)}}{|F'_\alpha| |F'_\beta| |F'_\rho|}.$$

**Proof:** Using Equations 16, 69, 70, we get

$$\begin{aligned} & \text{Tr} [\hat{\rho}^{\hat{X}^A \hat{Y}^B M} \hat{\rho}^{\hat{X}^A \hat{Y}^B M}] \\ &= \text{Tr} [((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\rho^{XYM}))(\hat{\Pi}^{\hat{X}^A \hat{Y}^B M} \circ \rho^{\hat{X}^A \hat{Y}^B M})] \\ &= \text{Tr} [((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\rho^{XYM})) \\ &\quad ((\mathbb{1}^{\hat{X}^A \hat{Y}^B M} - ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM} - \Pi(3)^{XYM}))) \circ (\hat{\Pi}^{\hat{X}^A \hat{Y}^B M} \circ \rho^{\hat{X}^A \hat{Y}^B M})))] \end{aligned}$$

$$\begin{aligned}
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM})) \circ ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\rho^{XYM}))) \\
&\quad ((\mathbb{1}^{\hat{X}\hat{Y}M} - ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM} - \Pi(3)^{XYM}))) \circ \hat{\rho}^{\hat{X}\hat{Y}M})] \\
&= \text{Tr} [((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\rho^{XYM})) \\
&\quad (((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM})) \\
&\quad \circ (\mathbb{1}^{\hat{X}\hat{Y}M} - ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM} - \Pi(3)^{XYM})))) \\
&\quad \circ \hat{\rho}^{\hat{X}\hat{Y}M})] \\
&= \text{Tr} [((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\rho^{XYM})) \\
&\quad (((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM})) \\
&\quad - ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM} - \Pi(3)^{XYM}))) \circ \hat{\rho}^{\hat{X}\hat{Y}M})] \\
&= \text{Tr} [((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\rho^{XYM})) \\
&\quad (((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\Pi(3)^{XYM})) \circ \hat{\rho}^{\hat{X}\hat{Y}M})] \\
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\Pi(3)^{XYM})) \\
&\quad \circ ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\rho^{XYM}))) \hat{\rho}^{\hat{X}\hat{Y}M}] \\
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\Pi(3)^{XYM} \circ \rho^{XYM})) \hat{\rho}^{\hat{X}\hat{Y}M})] \\
&\leq \| (T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\Pi(3)^{XYM} \circ \rho^{XYM}) \|_{\infty} \cdot \| \hat{\rho}^{\hat{X}\hat{Y}M} \|_1 \\
&= \| \Pi(3)^{XYM} \circ \rho^{XYM} \|_{\infty} \cdot \text{Tr} [\hat{\rho}^{\hat{X}\hat{Y}M}] \leq \frac{(1 + 3\sqrt{\epsilon})2^{D_{\infty}^{\epsilon}}(\rho^{XYM} \| \alpha^X \otimes \beta^Y \otimes \rho^M)}{|F'_{\alpha}| |F'_{\beta}| |F'_{\rho}|}.
\end{aligned}$$

Above, we used the fact that

$$\mathbb{1}^{\hat{X}\hat{Y}M} - (T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM} - \Pi(3)^{XYM}) \geq \hat{\Pi}^{\hat{X}\hat{Y}M}$$

in the third equality,

$$\begin{aligned}
&((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM})) \\
&\quad \circ (\mathbb{1}^{\hat{X}\hat{Y}M} - (T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM} - \Pi(3)^{XYM})) \\
&= (T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM}) - (T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M)(\mathbb{1}^{XYM} - \Pi(3)^{XYM})
\end{aligned}$$

in the sixth equality,  $T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes T_{\epsilon^{1/4}}^{Y \rightarrow \hat{Y}} \otimes \mathbb{I}^M$  is an isometry in the tenth equality. The proof of the lemma is now complete.  $\square$

**Lemma 21**

$$\text{Tr} [\hat{\rho}^{\hat{X}M} \hat{\rho}^{\hat{X}M}] \leq \frac{(1 + 3\sqrt{\epsilon})2^{D_{\infty}^{\epsilon}}(\rho^{XM} \| \alpha^X \otimes \rho^M)}{|F'_{\alpha}| |F'_{\rho}|}.$$

**Proof:** Using Equations 16, 69, 70, we get

$$\text{Tr} [\hat{\rho}^{\hat{X}M} \hat{\rho}^{\hat{X}M}]$$



$$\begin{aligned}
&= \text{Tr} [(\hat{\rho}^{\hat{X}M} \otimes \mathbb{1}^{\hat{Y}}) \hat{\rho}^{\hat{X}\hat{Y}M}] = \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\rho^{XM})) \otimes \mathbb{1}^{\hat{Y}})(\hat{\Pi}^{\hat{X}\hat{Y}M} \circ \rho^{\hat{X}\hat{Y}M})] \\
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\rho^{XM})) \otimes \mathbb{1}^{\hat{Y}}) \\
&\quad ((\mathbb{1}^{\hat{X}\hat{Y}M} - ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM} - \Pi(1)^{XM})) \otimes \mathbb{1}^{\hat{Y}}) \circ (\hat{\Pi}^{\hat{X}\hat{Y}M} \circ \rho^{\hat{X}\hat{Y}M})))] \\
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM})) \circ ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\rho^{XM}))) \otimes \mathbb{1}^{\hat{Y}}) \\
&\quad (((\mathbb{1}^{\hat{X}M} - ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM} - \Pi(1)^{XM}))) \otimes \mathbb{1}^{\hat{Y}}) \circ \hat{\rho}^{\hat{X}\hat{Y}M})] \\
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\rho^{XM})) \otimes \mathbb{1}^{\hat{Y}}) \\
&\quad ((((((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM})) \circ (\mathbb{1}^{\hat{X}M} - ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM} - \Pi(1)^{XM})))) \otimes \mathbb{1}^{\hat{Y}}) \\
&\quad \circ \hat{\rho}^{\hat{X}\hat{Y}M})))] \\
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\rho^{XM})) \otimes \mathbb{1}^{\hat{Y}}) \\
&\quad ((((((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM})) - ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM} - \Pi(1)^{XM}))) \otimes \mathbb{1}^{\hat{Y}}) \circ \hat{\rho}^{\hat{X}\hat{Y}M})))] \\
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\rho^{XM})) \otimes \mathbb{1}^{\hat{Y}}) \\
&\quad ((((((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\Pi(1)^{XM})) \otimes \mathbb{1}^{\hat{Y}}) \circ \hat{\rho}^{\hat{X}\hat{Y}M})))] \\
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\Pi(1)^{XM})) \circ ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\rho^{XM}))) \otimes \mathbb{1}^{\hat{Y}}) \hat{\rho}^{\hat{X}\hat{Y}M}] \\
&= \text{Tr} [(((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\Pi(1)^{XM} \circ \rho^{XM})) \otimes \mathbb{1}^{\hat{Y}}) \hat{\rho}^{\hat{X}\hat{Y}M}] \\
&\leq \|((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\Pi(1)^{XM} \circ \rho^{XM})) \otimes \mathbb{1}^{\hat{Y}}\|_{\infty} \cdot \|\hat{\rho}^{\hat{X}\hat{Y}M}\|_1 \\
&= \|\Pi(1)^{XM} \circ \rho^{XM}\|_{\infty} \cdot \text{Tr} [\hat{\rho}^{\hat{X}\hat{Y}M}] \leq \frac{(1 + 3\sqrt{\epsilon})2^{D_{\infty}^{\epsilon}(\rho^{XM} \| \alpha^X \otimes \rho^M)}}{|F'_{\alpha}| |F'_{\rho}|}.
\end{aligned}$$

Above, we used the fact that

$$\mathbb{1}^{\hat{X}\hat{Y}M} - ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM} - \Pi(1)^{XM})) \otimes \mathbb{1}^{\hat{Y}} \geq \hat{\Pi}^{\hat{X}\hat{Y}M}$$

in the third equality,

$$\begin{aligned}
&((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM})) \circ (\mathbb{1}^{\hat{X}M} - ((T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM} - \Pi(1)^{XM}))) \\
&= (T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM}) - (T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M)(\mathbb{1}^{XM} - \Pi(1)^{XM})
\end{aligned}$$

in the sixth equality,  $T_{\epsilon^{1/4}}^{X \rightarrow \hat{X}} \otimes \mathbb{I}^M$  is an isometry in the tenth equality. The proof of the lemma is now complete.  $\square$

## Lemma 22

$$\text{Tr} [\hat{\rho}^{\hat{Y}M} \hat{\rho}^{\hat{X}\hat{Y}M}] \leq \frac{(1 + 3\sqrt{\epsilon})2^{D_{\infty}^{\epsilon}(\rho^{YM} \| \beta^Y \otimes \rho^M)}}{|F'_{\beta}| |F'_{\rho}|}.$$

**Proof:** Similar to proof of Lemma 21 above.  $\square$

## Lemma 23

$$\text{Tr} [\hat{\rho}^M \hat{\rho}^M] \leq \frac{1 + \sqrt{\epsilon}}{|F'_{\rho}|}.$$

**Proof:** Using Equations 69 and 16, we get

$$\mathrm{Tr} [\hat{\rho}^M \hat{\rho}^M] = \mathrm{Tr} [\rho^M \hat{\rho}^M] \leq \frac{1 + \sqrt{\epsilon}}{|F'_\rho|} \mathrm{Tr} [\mathbb{1}^{F'_\rho} \hat{\rho}^M] \leq \frac{1 + \sqrt{\epsilon}}{|F'_\rho|} \mathrm{Tr} [\hat{\rho}^M] \leq \frac{1 + \sqrt{\epsilon}}{|F'_\rho|}.$$

This completes the proof of the lemma.  $\square$

We now prove the following lemma which completes the simpler proof of Lemma 7 by finally showing the validity of Equation 74.

**Lemma 24** *Suppose*

$$\begin{aligned} \log A &> D_\infty^\epsilon(\rho^{XM} \| \alpha^X \otimes \rho^M) + \log \epsilon^{-1/20}, \\ \log B &> D_\infty^\epsilon(\rho^{YM} \| \beta^Y \otimes \rho^M) + \log \epsilon^{-1/20}, \\ \log A + \log B &> D_\infty^\epsilon(\rho^{XYM} \| \alpha^X \otimes \beta^Y \otimes \rho^M) + \log \epsilon^{-1/20}. \end{aligned}$$

*Then for sufficiently small  $\epsilon$ ,*

$$\mathrm{Tr} [\hat{\sigma}^{\hat{X}^A \hat{Y}^B M} \hat{\sigma}^{\hat{X}^A \hat{Y}^B M}] < \frac{1 + \epsilon^{1/16}}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|}.$$

**Proof:** We follow the proof method of Lemma 13 above, but use Lemmas 20, 21, 22, 23 instead of Lemmas 9, 10, 11, 12. We get

$$\mathrm{Tr} [\hat{\sigma}^{\hat{X}^A \hat{Y}^B M} \hat{\sigma}^{\hat{X}^A \hat{Y}^B M}] < \frac{1 + 40\epsilon^{1/20}}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|} < \frac{1 + \epsilon^{1/16}}{|F'_\alpha|^A |F'_\beta|^B |F'_\rho|}$$

for sufficiently small  $\epsilon$ . This finishes the proof of the lemma.  $\square$

The proof of Lemma 7 is now complete.

## 8 Fully smooth multipartite decoupling

In this section, we prove a fully smooth multipartite decoupling theorem where we have to enlarge the Hilbert space dimensions of the senders by a factor of 2. Our proof is inspired by the simpler proof of the fully smooth multipartite convex split lemma in Section 7.

The EPR state is defined as

$$\Phi^{A'A} := |A|^{-1} \sum_{a_1, a_2=1}^{|A|} |a_1, a_1\rangle^{A'A} \langle a_2, a_2|,$$

where  $A'$  is another Hilbert space of the same dimension as  $A$ . We remark that though Theorem 3 is stated in terms of expectation over the Haar measure on unitaries as is the convention, it continues to hold without change for expectation over a perfect 2-design of unitaries, and with minor additive terms for expectation over an approximate 2-design of unitaries.

**Theorem 3** *Let  $\mathcal{T}^{A_1 A_2 \rightarrow E}$  be a completely positive trace non increasing superoperator with Choi state given by  $\tau^{A'_1 A'_2 E} := (\mathcal{T}^{A_1 A_2 \rightarrow E} \otimes \mathbb{I}^{A'_1 A'_2})(\Phi^{A'_1 A'_1} \otimes \Phi^{A'_2 A'_2})$ . Let  $\rho^{A_1 A_2 R}$  be a subnormalised density matrix. Define  $\hat{A}_1 := A_1 \otimes \mathbb{C}^2$ . Similarly define  $\hat{A}'_1, \hat{A}_2, \hat{A}'_2$ . Define  $\rho^{\hat{A}_1 \hat{A}_2 R} := \rho^{A_1 A_2 R} \otimes$*

$|0\rangle\langle 0|^{\mathbb{C}^2} \otimes |0\rangle\langle 0|^{\mathbb{C}^2}$ ,  $\tau^{\hat{A}_1\hat{A}_2E} := \tau^{A'_1A'_2E} \otimes |0\rangle\langle 0|^{\mathbb{C}^2} \otimes |0\rangle\langle 0|^{\mathbb{C}^2}$ . Let  $\mathcal{T}^{\hat{A}_1\hat{A}_2 \rightarrow E}$  be the completely positive superoperator corresponding to the inverse Choi image of  $\tau^{\hat{A}_1\hat{A}_2E}$ . Then  $\mathcal{T}^{\hat{A}_1\hat{A}_2 \rightarrow E}$  is the completely positive superoperator corresponding to projecting from  $\hat{A}_1 \otimes \hat{A}_2$  onto  $(A_1 \otimes |0\rangle\langle 0|^{\mathbb{C}^2}) \otimes (A_2 \otimes |0\rangle\langle 0|^{\mathbb{C}^2})$  followed by applying  $4\mathcal{T}^{A_1A_2 \rightarrow E}$ .

Let  $U^{\hat{A}_1}$ ,  $U^{\hat{A}_2}$  be unitary matrices on their respective Hilbert spaces. Let  $0 < \epsilon < 1$ . Then,

$$\mathbb{E}_{U^{\hat{A}_1}U^{\hat{A}_2}} [\|(\mathcal{T}^{\hat{A}_1\hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \rho^{\hat{A}_1\hat{A}_2R}) - \tau^E \otimes \rho^R\|_1] \leq 18\epsilon^{1/128}$$

if

$$\begin{aligned} H_{\min}^\epsilon(A_1|R)_\rho + H_{\min}^\epsilon(A'_1|E)_\tau &> \log \epsilon^{-1}, \\ H_{\min}^\epsilon(A_2|R)_\rho + H_{\min}^\epsilon(A'_2|E)_\tau &> \log \epsilon^{-1}, \\ H_{\min}^\epsilon(A_1A_2|R)_\rho + H_{\min}^\epsilon(A'_1A'_2|E)_\tau &> \log \epsilon^{-1}, \\ |A_1|^2 &> \epsilon^{-1}, \quad |A_2|^2 > \epsilon^{-1}, \end{aligned}$$

where the expectation on the left is taken over independent choices of unitaries  $U^{\hat{A}_1}$ ,  $U^{\hat{A}_2}$  from their respective Haar measures.

We now proceed with the constructions required to prove Theorem 3. Observe without loss of generality that  $\rho^{A_1A_2R}$  can be assumed to be a normalised quantum state. Let  $L_R$  and  $L_E$  be the ancilla Hilbert spaces required for flattening  $\rho^R$  and  $\tau^E$  respectively. By flattening using the CPTP superoperators  $\mathcal{F}_\rho^{R \rightarrow L_R R}$ ,  $\mathcal{F}_\tau^{E \rightarrow L_E E}$ , and by applying Proposition 3 we observe that it suffices to show

$$\begin{aligned} \mathbb{E}_{U^{\hat{A}_1}U^{\hat{A}_2}} [\|((\mathcal{F}_\tau^{E \rightarrow L_E E} \cdot \mathcal{T}^{\hat{A}_1\hat{A}_2 \rightarrow E}) \otimes \mathcal{F}_\rho^{R \rightarrow L_R R})((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \rho^{\hat{A}_1\hat{A}_2R}) \\ - \mathcal{F}_\tau^{E \rightarrow L_E E}(\tau^E) \otimes \mathcal{F}_\rho^{R \rightarrow L_R R}(\rho^R)\|_1] \leq 81\epsilon^{1/64}. \end{aligned}$$

Henceforth, we shall redefine

$$\begin{aligned} L_R R &\equiv R, \\ L_E E &\equiv E, \\ \mathcal{F}_\tau^{E \rightarrow L_E E} \cdot \mathcal{T}^{\hat{A}_1\hat{A}_2 \rightarrow E} &\equiv \mathcal{T}^{\hat{A}_1\hat{A}_2 \rightarrow E}, \\ (\mathbb{I}^{\hat{A}_1} \otimes \mathbb{I}^{\hat{A}_2} \otimes \mathcal{F}_\rho^{R \rightarrow L_R R})(\rho^{\hat{A}_1\hat{A}_2R}) &\equiv \rho^{\hat{A}_1\hat{A}_2R}, \\ ((\mathcal{F}_\tau^{E \rightarrow L_E E} \cdot \mathcal{T}^{\hat{A}_1\hat{A}_2 \rightarrow E}) \otimes \mathbb{I}^{\hat{A}'_1\hat{A}'_2})(\Phi^{\hat{A}'_1\hat{A}_1} \otimes \Phi^{\hat{A}'_2\hat{A}_2}) &= (\mathcal{F}_\tau^{E \rightarrow L_E E} \otimes \mathbb{I}^{\hat{A}'_1\hat{A}'_2})(\tau^{\hat{A}'_1\hat{A}'_2E}) \equiv \tau^{\hat{A}'_1\hat{A}'_2E}. \end{aligned}$$

Thus with this changed notation we will show

$$\mathbb{E}_{U^{\hat{A}_1}U^{\hat{A}_2}} [\|(\mathcal{T}^{\hat{A}_1\hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \rho^{\hat{A}_1\hat{A}_2R}) - \tau^E \otimes \rho^R\|_1] \leq 81\epsilon^{1/64}. \quad (75)$$

Note that from its definition,  $\mathcal{T}^{\hat{A}_1\hat{A}_2 \rightarrow E}$  increases the trace by at most a multiplicative factor of 4.

Define the tilting map

$$T_{\epsilon^{1/4}}^{A_1 \rightarrow \hat{A}_1} : |a_1\rangle^{A_1} \mapsto \sqrt{1 - \epsilon^{1/4}}|a_1\rangle^{A_1}|0\rangle + \epsilon^{1/8}|a_1\rangle^{A_1}|1\rangle.$$

Define  $T_{\epsilon^{1/4}}^{A_2 \rightarrow \hat{A}_2}$ ,  $T_{\epsilon^{1/4}}^{A'_1 \rightarrow \hat{A}'_1}$ ,  $T_{\epsilon^{1/4}}^{A'_2 \rightarrow \hat{A}'_2}$  similarly. Let  $\Pi(1)^{A_1 R}$ ,  $\Pi(2)^{A_2 R}$ ,  $\Pi(3)^{A_1 A_2 R}$ ,  $\Pi(1)^{A'_1 E}$ ,  $\Pi(2)^{A'_2 E}$ ,  $\Pi(3)^{A'_1 A'_2 E}$  be the projectors promised by Proposition 2 achieving the respective smooth conditional min entropy quantities for the flattened states. More precisely, Propositions 3, 2 guarantee, when the flattening parameter  $\delta$  is small enough, that

$$\begin{aligned}
\Pi(1)^{A_1 R} \circ \rho^{A_1 R} &\leq \frac{(1+3\sqrt{\epsilon})2^{-H_{\min}^{\epsilon}(A_1|R)_{\rho}}}{|F'_{\rho}|} \mathbb{1}^{A_1 F'_{\rho}}, & \Pi(2)^{A_2 R} \circ \rho^{A_2 R} &\leq \frac{(1+3\sqrt{\epsilon})2^{-H_{\min}^{\epsilon}(A_2|R)_{\rho}}}{|F'_{\rho}|} \mathbb{1}^{A_2 F'_{\rho}}, \\
\Pi(3)^{A_1 A_2 R} \circ \rho^{A_1 A_2 R} &\leq \frac{(1+3\sqrt{\epsilon})2^{-H_{\min}^{\epsilon}(A_1 A_2|R)_{\rho}}}{|F'_{\rho}|} \mathbb{1}^{A_1 A_2 F'_{\rho}}, \\
\Pi(1)^{A'_1 E} \circ \rho^{A'_1 E} &\leq \frac{(1+3\sqrt{\epsilon})2^{-H_{\min}^{\epsilon}(A'_1|E)_{\rho}}}{|F'_{\rho}|} \mathbb{1}^{A'_1 F'_{\rho}}, & \Pi(2)^{A'_2 E} \circ \rho^{A'_2 E} &\leq \frac{(1+3\sqrt{\epsilon})2^{-H_{\min}^{\epsilon}(A'_2|E)_{\rho}}}{|F'_{\rho}|} \mathbb{1}^{A'_2 F'_{\rho}}, \\
\Pi(3)^{A'_1 A'_2 E} \circ \rho^{A'_1 A'_2 E} &\leq \frac{(1+3\sqrt{\epsilon})2^{-H_{\min}^{\epsilon}(A'_1 A'_2|E)_{\rho}}}{|F'_{\rho}|} \mathbb{1}^{A'_1 A'_2 F'_{\rho}}.
\end{aligned} \tag{76}$$

Define

$$\begin{aligned}
\hat{\rho}^{\hat{A}_1 \hat{A}_2 R} &:= (T_{\epsilon^{1/4}}^{A_1 \rightarrow \hat{A}_1} \otimes T_{\epsilon^{1/4}}^{A_2 \rightarrow \hat{A}_2} \otimes \mathbb{I}^R)(\rho^{A_1 A_2 R}), \\
\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E} &:= (T_{\epsilon^{1/4}}^{A'_1 \rightarrow \hat{A}'_1} \otimes T_{\epsilon^{1/4}}^{A'_2 \rightarrow \hat{A}'_2} \otimes \mathbb{I}^E)(\tau^{A'_1 A'_2 E}), \\
\hat{\Pi}^{\hat{A}_1 \hat{A}_2 R} &:= \mathbb{1}^{\hat{A}_1 \hat{A}_2 R} \\
&\quad - \text{span} \{((T_{\epsilon^{1/4}}^{A_1 \rightarrow \hat{A}_1} \otimes \mathbb{I}^R)(\mathbb{1}^{A_1 R} - \Pi(1)^{A_1 R})) \otimes \mathbb{1}^{\hat{A}_2}, \\
&\quad ((T_{\epsilon^{1/4}}^{A_2 \rightarrow \hat{A}_2} \otimes \mathbb{I}^R)(\mathbb{1}^{A_2 R} - \Pi(2)^{A_2 R})) \otimes \mathbb{1}^{\hat{A}_1}, \\
&\quad (T_{\epsilon^{1/4}}^{A_1 \rightarrow \hat{A}_1} \otimes T_{\epsilon^{1/4}}^{A_2 \rightarrow \hat{A}_2} \otimes \mathbb{I}^R)(\mathbb{1}^{A_1 A_2 R} - \Pi(3)^{A_1 A_2 R})\}, \\
\hat{\Pi}^{\hat{A}'_1 \hat{A}'_2 E} &:= \mathbb{1}^{\hat{A}'_1 \hat{A}'_2 E} \\
&\quad - \text{span} \{((T_{\epsilon^{1/4}}^{A'_1 \rightarrow \hat{A}'_1} \otimes \mathbb{I}^E)(\mathbb{1}^{A'_1 E} - \Pi(1)^{A'_1 E})) \otimes \mathbb{1}^{\hat{A}'_2}, \\
&\quad ((T_{\epsilon^{1/4}}^{A'_2 \rightarrow \hat{A}'_2} \otimes \mathbb{I}^E)(\mathbb{1}^{A'_2 E} - \Pi(2)^{A'_2 E})) \otimes \mathbb{1}^{\hat{A}'_1}, \\
&\quad (T_{\epsilon^{1/4}}^{A'_1 \rightarrow \hat{A}'_1} \otimes T_{\epsilon^{1/4}}^{A'_2 \rightarrow \hat{A}'_2} \otimes \mathbb{I}^E)(\mathbb{1}^{A'_1 A'_2 E} - \Pi(3)^{A'_1 A'_2 E})\}, \\
\hat{\rho}^{\hat{A}_1 \hat{A}_2 R} &:= \hat{\Pi}^{\hat{A}_1 \hat{A}_2 R} \circ \rho^{\hat{A}_1 \hat{A}_2 R}, & \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E} &:= \hat{\Pi}^{\hat{A}'_1 \hat{A}'_2 E} \circ \tau^{\hat{A}'_1 \hat{A}'_2 E}, \\
\sigma(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} &:= (\mathcal{T}^{\hat{A}_1 \hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \rho^{\hat{A}_1 \hat{A}_2 R}), \\
\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} &:= (\hat{\mathcal{T}}^{\hat{A}_1 \hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \hat{\rho}^{\hat{A}_1 \hat{A}_2 R}), \\
\hat{\hat{\sigma}}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} &:= (\hat{\hat{\mathcal{T}}}^{\hat{A}_1 \hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \hat{\hat{\rho}}^{\hat{A}_1 \hat{A}_2 R}),
\end{aligned} \tag{77}$$

where  $\hat{\mathcal{T}}^{\hat{A}_1 \hat{A}_2 \rightarrow E}$ ,  $\hat{\hat{\mathcal{T}}}^{\hat{A}_1 \hat{A}_2 \rightarrow E}$  are the completely positive superoperators arising as the inverse Choi images of  $\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}$ ,  $\hat{\hat{\tau}}^{\hat{A}'_1 \hat{A}'_2 E}$ . Note that  $\rho^{A_1 A_2 R}$ ,  $\rho^{\hat{A}_1 \hat{A}_2 R}$ ,  $\hat{\rho}^{\hat{A}_1 \hat{A}_2 R}$  are normalised density matrices, and  $\hat{\rho}^{\hat{A}_1 \hat{A}_2 R}$ ,  $\tau^{A'_1 A'_2 E}$ ,  $\tau^{\hat{A}'_1 \hat{A}'_2 E}$ ,  $\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}$ ,  $\hat{\hat{\tau}}^{\hat{A}'_1 \hat{A}'_2 E}$  are subnormalised density matrices. Even though  $\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}$ ,  $\hat{\hat{\tau}}^{\hat{A}'_1 \hat{A}'_2 E}$  are subnormalised, their inverse Choi images  $\hat{\mathcal{T}}^{\hat{A}_1 \hat{A}_2 \rightarrow E}$ ,  $\hat{\hat{\mathcal{T}}}^{\hat{A}_1 \hat{A}_2 \rightarrow E}$  can increase trace but by at most a multiplicative factor of 4.

The following properties can be proved easily as in Sections 4, 7.

$$\begin{aligned}
\frac{(1-\sqrt{\epsilon})(\text{Tr } \tau)}{|F'_\tau||F'_\rho|} \mathbb{1}^{F'_\tau F'_\rho} &\leq \tau^E \otimes \rho^R \leq \frac{(1+\sqrt{\epsilon})(\text{Tr } \tau)}{|F'_\tau||F'_\rho|} \mathbb{1}^{F'_\tau F'_\rho}, \\
\hat{\rho}^{\hat{A}_1 R} &= (T_{\epsilon^{1/4}}^{A_1 \rightarrow \hat{A}_1} \otimes \mathbb{I}^R)(\rho^{A_1 R}), \quad \hat{\rho}^{\hat{A}_2 R} = (T_{\epsilon^{1/4}}^{A_2 \rightarrow \hat{A}_2} \otimes \mathbb{I}^R)(\rho^{A_2 R}), \\
\hat{\tau}^{\hat{A}'_1 R} &= (T_{\epsilon^{1/4}}^{A'_1 \rightarrow \hat{A}'_1} \otimes \mathbb{I}^R)(\tau^{A'_1 R}), \quad \hat{\tau}^{\hat{A}'_2 R} = (T_{\epsilon^{1/4}}^{A'_2 \rightarrow \hat{A}'_2} \otimes \mathbb{I}^R)(\tau^{A'_2 R}), \\
\|\hat{\rho}^{\hat{A}_1 \hat{A}_2 R} - \rho^{\hat{A}_1 \hat{A}_2 R}\|_1 &\leq 4\sqrt{2}\epsilon^{1/8}, \quad \|\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 R} - \tau^{\hat{A}'_1 \hat{A}'_2 R}\|_1 \leq 4\sqrt{2}\epsilon^{1/8}(\text{Tr } \tau), \\
\text{Tr } [\hat{\rho}^{\hat{A}_1 \hat{A}_2 R}] &\geq 1 - 121\epsilon^{1/4}, \quad \text{Tr } [\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 R}] \geq (1 - 121\epsilon^{1/4})(\text{Tr } \tau), \\
\Rightarrow \|\hat{\rho}^{\hat{A}_1 \hat{A}_2 R} - \rho^{\hat{A}_1 \hat{A}_2 R}\|_1 &\leq 22\epsilon^{1/8}, \quad \|\hat{\rho}^{\hat{A}_1 \hat{A}_2 R} - \hat{\rho}^{\hat{A}_1 \hat{A}_2 R}\|_1 \leq 28\epsilon^{1/8}, \\
\Rightarrow \|\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 R} - \tau^{\hat{A}'_1 \hat{A}'_2 R}\|_1 &\leq 22\epsilon^{1/8}(\text{Tr } \tau), \quad \|\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 R} - \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 R}\|_1 \leq 28\epsilon^{1/8}(\text{Tr } \tau).
\end{aligned} \tag{78}$$

Following an argument in the proof of Proposition 3 in [Sen24], which in turn originated from a method in [SDTR13], and using Equation 78 and the fact that  $\hat{\mathcal{T}}^{\hat{A}_1 \hat{A}_2 \rightarrow E}$  can increase the trace by at most a factor of 4, we get

$$\begin{aligned}
&\mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|\sigma(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} - \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}\|_1] \\
&\leq \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\mathcal{T}^{\hat{A}_1 \hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \rho^{\hat{A}_1 \hat{A}_2 R}) \\
&\quad - (\hat{\mathcal{T}}^{\hat{A}_1 \hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \hat{\rho}^{\hat{A}_1 \hat{A}_2 R})\|_1] \\
&\leq \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\mathcal{T}^{\hat{A}_1 \hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \rho^{\hat{A}_1 \hat{A}_2 R}) \\
&\quad - (\mathcal{T}^{\hat{A}_1 \hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \hat{\rho}^{\hat{A}_1 \hat{A}_2 R})\|_1] \\
&\quad + \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\mathcal{T}^{\hat{A}_1 \hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \hat{\rho}^{\hat{A}_1 \hat{A}_2 R}) \\
&\quad - (\hat{\mathcal{T}}^{\hat{A}_1 \hat{A}_2 \rightarrow E} \otimes \mathbb{I}^R)((U^{\hat{A}_1} \otimes U^{\hat{A}_2} \otimes \mathbb{1}^R) \circ \hat{\rho}^{\hat{A}_1 \hat{A}_2 R})\|_1] \\
&\leq 4\|\rho^{\hat{A}_1 \hat{A}_2 R} - \hat{\rho}^{\hat{A}_1 \hat{A}_2 R}\|_1 + \|\tau^{\hat{A}'_1 \hat{A}'_2 E} - \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}\|_1 \leq 20\sqrt{2}\epsilon^{1/8}, \\
\text{Similarly, } &\mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} - \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}\|_1] \\
&\leq 4\|\hat{\rho}^{\hat{A}_1 \hat{A}_2 R} - \hat{\rho}^{\hat{A}_1 \hat{A}_2 R}\|_1 + \|\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E} - \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}\|_1 \leq 140\epsilon^{1/8}.
\end{aligned} \tag{79}$$

Define the set  $\text{Good} := \{(U^{\hat{A}_1}, U^{\hat{A}_2}) : \|\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} - \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}\|_1 < 140\epsilon^{1/16}\}$ . Applying Markov's inequality to the last inequality in Equation 79, we get

$$\Pr_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\text{Good}] > 1 - \epsilon^{1/16}.$$

Applying Lemma 1, we get

$$\begin{aligned}
&\forall (U^{\hat{A}_1}, U^{\hat{A}_2}) \in \text{Good} : \exists (\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} : \\
&\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} - \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}\|_1 < 12\epsilon^{1/32}, \\
&(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} \leq \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}, \\
&\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_2^2 \leq (1 + 24\epsilon^{1/32}) \text{Tr } [\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}], \\
&\forall (U^{\hat{A}_1}, U^{\hat{A}_2}) \notin \text{Good} : (\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} := \mathbf{0}^{ER}, \\
&\Rightarrow \forall (U^{\hat{A}_1}, U^{\hat{A}_2}) : (\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} \leq \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}, \\
&\|(\sigma'((U^{\hat{A}_1}, U^{\hat{A}_2})))^{ER}\|_2^2 \leq (1 + 24\epsilon^{1/32}) \text{Tr } [\hat{\sigma}((U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} \hat{\sigma}((U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}].
\end{aligned} \tag{80}$$

Using Equations 79 and 80 and the property that  $\mathcal{T}^{\hat{A}_1\hat{A}_2 \rightarrow E}$  increases the trace by at most a factor of 4, we get

$$\begin{aligned}
& \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|\sigma(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} - (\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_1] \\
&= \mathbb{E}_{(U^{\hat{A}_1}, U^{\hat{A}_2}) \in \text{Good}} [\|\sigma(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} - (\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_1] \\
&\quad + \mathbb{E}_{(U^{\hat{A}_1}, U^{\hat{A}_2}) \notin \text{Good}} [\|\sigma(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}\|_1] \\
&\leq \mathbb{E}_{(U^{\hat{A}_1}, U^{\hat{A}_2}) \in \text{Good}} [\|\sigma(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} - \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}\|_1] \\
&\quad + \mathbb{E}_{(U^{\hat{A}_1}, U^{\hat{A}_2}) \in \text{Good}} [\|\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} - (\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_1] + 4(1 - \Pr_{U^{\hat{A}_1}, U^{\hat{A}_2}}[\text{Good}]) \\
&\leq \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|\sigma(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} - \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}\|_1] + 12\epsilon^{1/32} + 4\epsilon^{1/16} \\
&\leq 20\sqrt{2}\epsilon^{1/8} + 12\epsilon^{1/32} + 4\epsilon^{1/16} < 45\epsilon^{1/32}.
\end{aligned}$$

Thus in order to prove the present proposition, from Equation 75, it suffices to show that

$$\mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} - \tau^E \otimes \rho^R\|_1] < 36\epsilon^{1/64}. \quad (81)$$

Define  $\Pi^{ER} := \text{supp}(\tau^E \otimes \rho^R) = F'_\tau \otimes F'_\rho$ . Observe that for all  $(U^{\hat{A}_1}, U^{\hat{A}_2})$ ,

$$\text{supp}((\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}) \leq \text{supp}(\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}) \leq \Pi^{ER}, \quad \text{supp}(\sigma(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}) \leq \Pi^{ER},$$

where the first operator inequality follows from Equation 80 and the second operator inequality follows from the fact that  $\hat{\mathcal{T}}^{\hat{A}_1\hat{A}_2 \rightarrow E}$  is the completely positive superoperator that projects from  $\hat{A}_1 \otimes \hat{A}_2$  onto the subspace  $T_{\epsilon^{1/4}}^{A_1 \rightarrow \hat{A}_1} \otimes T_{\epsilon^{1/4}}^{A_2 \rightarrow \hat{A}_2}$ , followed by rotating the subspace onto  $(A_1 \otimes |0\rangle\langle 0|^{\mathbb{C}^2}) \otimes (A_2 \otimes |0\rangle\langle 0|^{\mathbb{C}^2})$ , followed by applying  $4\mathcal{T}^{A_1A_2 \rightarrow E}$ . By Proposition 1, Equation 81 and the convexity of the square function, it suffices to show

$$\begin{aligned}
& \sqrt{\text{Tr } \Pi^{ER}} \cdot \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} - \tau^E \otimes \rho^R\|_1] < 36\epsilon^{1/64}, \\
& \implies \text{it suffices to show } \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} - \tau^E \otimes \rho^R\|_2^2] < \frac{1296\epsilon^{1/32}}{F'_\tau F'_\rho}. \quad (82)
\end{aligned}$$

Using Equations 78, 80, the method of the proof of Proposition 3 of [Sen24] and the fact that  $\hat{\mathcal{T}}^{\hat{A}_1\hat{A}_2 \rightarrow E}$  is a completely positive superoperator that increases the trace by at most a factor of 4, we have,

$$\begin{aligned}
& \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} - \tau^E \otimes \rho^R\|_2^2] \\
&= \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_2^2 + \|\tau^E \otimes \rho^R\|_2^2 - 2 \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\text{Tr} [(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} (\tau^E \otimes \rho^R)]]] \\
&\leq \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_2^2] + \frac{(1 + \sqrt{\epsilon})^2 (\text{Tr } \tau)^2}{|F'_\tau|^2 |F'_\rho|^2} \text{Tr } [\mathbb{1}^{F'_\tau F'_\rho}]
\end{aligned}$$

$$\begin{aligned}
& - \frac{2(1-\sqrt{\epsilon})(\text{Tr } \tau)}{|F'_\tau||F'_\rho|} \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\text{Tr} [(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} \Pi^{ER}]] \\
\leq & \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_2^2] + \frac{(1+\sqrt{\epsilon})^2(\text{Tr } \tau)^2}{|F'_\tau||F'_\rho|} \\
& - \frac{2(1-\sqrt{\epsilon})(\text{Tr } \tau)}{|F'_\tau||F'_\rho|} \mathbb{E}_{(U^{\hat{A}_1}, U^{\hat{A}_2}) \in \text{Good}} [\text{Tr} [(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER} \Pi^{ER}]] \\
\leq & \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_2^2] + \frac{(1+\sqrt{\epsilon})^2(\text{Tr } \tau)^2}{|F'_\tau||F'_\rho|} \\
& - \frac{2(1-\sqrt{\epsilon})(\text{Tr } \tau)}{|F'_\tau||F'_\rho|} \mathbb{E}_{(U^{\hat{A}_1}, U^{\hat{A}_2}) \in \text{Good}} [\text{Tr} [\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} \Pi^{ER}]] \\
& + \frac{2(1-\sqrt{\epsilon})(\text{Tr } \tau)}{|F'_\tau||F'_\rho|} \mathbb{E}_{(U^{\hat{A}_1}, U^{\hat{A}_2}) \in \text{Good}} [\|\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} - (\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_1] \\
\leq & \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_2^2] + \frac{(1+\sqrt{\epsilon})^2(\text{Tr } \tau)^2}{|F'_\tau||F'_\rho|} \\
& - \frac{2(1-\sqrt{\epsilon})(\text{Tr } \tau)}{|F'_\tau||F'_\rho|} \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\text{Tr} [\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} \Pi^{ER}]] \\
& + \frac{8(1-\sqrt{\epsilon})(\text{Tr } \tau)(1 - \Pr_{U^{\hat{A}_1}, U^{\hat{A}_2}}[\text{Good}])}{|F'_\tau||F'_\rho|} + \frac{24\epsilon^{1/32}(1-\sqrt{\epsilon})(\text{Tr } \tau)}{|F'_\tau||F'_\rho|} \\
\leq & \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_2^2] + \frac{(1+\sqrt{\epsilon})^2(\text{Tr } \tau)^2}{|F'_\tau||F'_\rho|} + \frac{32\epsilon^{1/32}}{|F'_\tau||F'_\rho|} \\
& - \frac{2(1-\sqrt{\epsilon})(\text{Tr } \tau)}{|F'_\tau||F'_\rho|} \text{Tr} [(\tau^E \otimes \tau^R) \Pi^{ER}] \\
= & \frac{32\epsilon^{1/32}}{|F'_\tau||F'_\rho|} + \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_2^2] + \frac{(1+\sqrt{\epsilon})^2(\text{Tr } \tau)^2}{|F'_\tau||F'_\rho|} - \frac{2(1-\sqrt{\epsilon})(\text{Tr } \tau)^2}{|F'_\tau||F'_\rho|} \\
< & \frac{32\epsilon^{1/32}}{|F'_\tau||F'_\rho|} + \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_2^2] - \frac{(1-5\sqrt{\epsilon})(\text{Tr } \tau)^2}{|F'_\tau||F'_\rho|}.
\end{aligned}$$

Thus from Equations 82, 80, it suffices to show that

$$\begin{aligned}
& \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\|(\sigma'(U^{\hat{A}_1}, U^{\hat{A}_2}))^{ER}\|_2^2] \leq \frac{(1+1259\epsilon^{1/32})(\text{Tr } \tau)^2}{|F'_\tau||F'_\rho|}, \\
\implies & \text{suffices to show } \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\text{Tr} [\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}]] \leq \frac{(1+30\epsilon^{1/32})(\text{Tr } \tau)^2}{|F'_\tau||F'_\rho|}. \quad (83)
\end{aligned}$$

Following the arguments in [CNS21] involving the use of the *swap trick* and the *twirling operator* over the Haar measure, we get

$$\begin{aligned}
& \mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\text{Tr} [\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}]] \\
= & \alpha_0 \text{Tr} [\hat{\rho}^R \hat{\rho}^R] + \alpha_1 \text{Tr} [\hat{\rho}^{\hat{A}_1 R} \hat{\rho}^{\hat{A}_1 R}] + \alpha_2 \text{Tr} [\hat{\rho}^{\hat{A}_2 R} \hat{\rho}^{\hat{A}_2 R}] + \alpha_{12} \text{Tr} [\hat{\rho}^{\hat{A}_1 \hat{A}_2 R} \hat{\rho}^{\hat{A}_1 \hat{A}_2 R}], \quad (84)
\end{aligned}$$

where the 4-tuple  $(\alpha_0, \alpha_1, \alpha_2, \alpha_{12})$  is defined to be the solution of the following linear equation:

$$\begin{bmatrix} \alpha_0 \\ \alpha_1 \\ \alpha_2 \\ \alpha_{12} \end{bmatrix} = \frac{|A_1||A_2|}{(|A_1|^2-1)(|A_2|^2-1)} \cdot \begin{bmatrix} |A_1||A_2| & -|A_2| & -|A_1| & 1 \\ -|A_2| & |A_1||A_2| & 1 & -|A_1| \\ -|A_1| & 1 & |A_1||A_2| & -|A_2| \\ 1 & -|A_1| & -|A_2| & |A_1||A_2| \end{bmatrix} \begin{bmatrix} \text{Tr} [\hat{\tau}^E \hat{\tau}^E] \\ \text{Tr} [\hat{\tau}^{\hat{A}'_1 E} \hat{\tau}^{\hat{A}'_1 E}] \\ \text{Tr} [\hat{\tau}^{\hat{A}'_2 E} \hat{\tau}^{\hat{A}'_2 E}] \\ \text{Tr} [\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E} \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}] \end{bmatrix}.$$

We bound the terms  $\alpha_0, \alpha_1, \alpha_2, \alpha_{12}$  as follows, using Fact 3.

$$\begin{aligned} \alpha_0 &= \frac{|A_1||A_2|}{(|A_1|^2-1)(|A_2|^2-1)} \cdot (|A_1||A_2| \cdot \text{Tr} [\hat{\tau}^E \hat{\tau}^E] - |A_2| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 E} \hat{\tau}^{\hat{A}'_1 E}] \\ &\quad - |A_1| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_2 E} \hat{\tau}^{\hat{A}'_2 E}] + \text{Tr} [\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E} \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}]) \\ &\leq \frac{|A_1||A_2|}{(|A_1|^2-1)(|A_2|^2-1)} \cdot (|A_1||A_2| \cdot \text{Tr} [\hat{\tau}^E \hat{\tau}^E] \\ &\quad - |A_1| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_2 E} \hat{\tau}^{\hat{A}'_2 E}] + |A_1| \text{Tr} [(\mathbb{1}^{\hat{A}'_1} \otimes \hat{\tau}^{\hat{A}'_2 E}) \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}]) \\ &= \frac{|A_1||A_2|}{(|A_1|^2-1)(|A_2|^2-1)} \cdot (|A_1||A_2| \cdot \text{Tr} [\hat{\tau}^E \hat{\tau}^E] \\ &\quad - |A_1| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_2 E} \hat{\tau}^{\hat{A}'_2 E}] + |A_1| \text{Tr} [\hat{\tau}^{\hat{A}'_2 E} \hat{\tau}^{\hat{A}'_2 E}]) \\ &= \frac{|A_1|^2|A_2|^2}{(|A_1|^2-1)(|A_2|^2-1)} \cdot \text{Tr} [\hat{\tau}^E \hat{\tau}^E], \\ \alpha_1 &= \frac{|A_1||A_2|}{(|A_1|^2-1)(|A_2|^2-1)} \cdot (-|A_2| \cdot \text{Tr} [\hat{\tau}^E \hat{\tau}^E] + |A_1||A_2| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 E} \hat{\tau}^{\hat{A}'_1 E}] \\ &\quad + \text{Tr} [\hat{\tau}^{\hat{A}'_2 E} \hat{\tau}^{\hat{A}'_2 E}] - |A_1| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E} \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}]) \\ &\leq \frac{|A_1||A_2|}{(|A_1|^2-1)(|A_2|^2-1)} \cdot (-|A_2| \cdot \text{Tr} [\hat{\tau}^E \hat{\tau}^E] + |A_1||A_2| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 E} \hat{\tau}^{\hat{A}'_1 E}] \\ &\quad + |A_2| \cdot \text{Tr} [(\mathbb{1}^{\hat{A}'_2} \otimes \hat{\tau}^E) \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}]) \\ &= \frac{|A_1||A_2|}{(|A_1|^2-1)(|A_2|^2-1)} \cdot (-|A_2| \cdot \text{Tr} [\hat{\tau}^E \hat{\tau}^E] + |A_1||A_2| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 E} \hat{\tau}^{\hat{A}'_1 E}] \\ &\quad + |A_2| \cdot \text{Tr} [\hat{\tau}^E \hat{\tau}^E] +) \\ &= \frac{|A_1|^2|A_2|^2}{(|A_1|^2-1)(|A_2|^2-1)} \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 E} \hat{\tau}^{\hat{A}'_1 E}], \\ \alpha_2 &\leq \frac{|A_1|^2|A_2|^2}{(|A_1|^2-1)(|A_2|^2-1)} \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_2 E} \hat{\tau}^{\hat{A}'_2 E}] \text{ similarly,} \\ \alpha_{12} &= \frac{|A_1||A_2|}{(|A_1|^2-1)(|A_2|^2-1)} \cdot (\text{Tr} [\hat{\tau}^E \hat{\tau}^E] - |A_1| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 E} \hat{\tau}^{\hat{A}'_1 E}] \\ &\quad - |A_2| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_2 E} \hat{\tau}^{\hat{A}'_2 E}] + |A_1||A_2| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E} \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}]) \\ &\leq \frac{|A_1||A_2|}{(|A_1|^2-1)(|A_2|^2-1)} \cdot (\text{Tr} [\hat{\tau}^E \hat{\tau}^E] + |A_1||A_2| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E} \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}]). \end{aligned} \tag{85}$$

From Equations 84, 85 we get,

$$\begin{aligned} &\mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\text{Tr} [\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}]] \\ &\leq \frac{|A_1|^2|A_2|^2}{(|A_1|^2-1)(|A_2|^2-1)} \cdot \text{Tr} [\hat{\tau}^E \hat{\tau}^E] \cdot \text{Tr} [\hat{\rho}^R \hat{\rho}^R] \\ &\quad + \frac{|A_1|^2|A_2|^2}{(|A_1|^2-1)(|A_2|^2-1)} \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_2 E} \hat{\tau}^{\hat{A}'_2 E}] \cdot \text{Tr} [\hat{\rho}^{\hat{A}_1 R} \hat{\rho}^{\hat{A}_1 R}] \\ &\quad + \frac{|A_1|^2|A_2|^2}{(|A_1|^2-1)(|A_2|^2-1)} \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 E} \hat{\tau}^{\hat{A}'_1 E}] \cdot \text{Tr} [\hat{\rho}^{\hat{A}_2 R} \hat{\rho}^{\hat{A}_2 R}] \\ &\quad + \frac{|A_1||A_2|}{(|A_1|^2-1)(|A_2|^2-1)} \cdot (\text{Tr} [\hat{\tau}^E \hat{\tau}^E] + |A_1||A_2| \cdot \text{Tr} [\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E} \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}]) \cdot \text{Tr} [\hat{\rho}^{\hat{A}_1 \hat{A}_2 R} \hat{\rho}^{\hat{A}_1 \hat{A}_2 R}]. \end{aligned} \tag{86}$$

As in Section 7, we can prove the following lemma.



**Lemma 25**

$$\begin{aligned}
\text{Tr} [\hat{\tau}^E \hat{\tau}^E] &\leq \frac{(1 + \sqrt{\epsilon})(\text{Tr } \tau)^2}{|F'_\tau|}, \quad \text{Tr} [\hat{\rho}^R \hat{\rho}^R] \leq \frac{1 + \sqrt{\epsilon}}{|F'_\rho|}, \\
\text{Tr} [\hat{\tau}^{\hat{A}'_1 E} \hat{\tau}^{\hat{A}'_1 E}] &\leq \frac{(1 + 3\sqrt{\epsilon})(\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^\epsilon(A'_1|E)_\tau}}{|F'_\tau|}, \\
\text{Tr} [\hat{\rho}^{\hat{A}_1 R} \hat{\rho}^{\hat{A}_1 R}] &\leq \frac{(1 + 3\sqrt{\epsilon}) \cdot 2^{-H_{\min}^\epsilon(A_1|R)_\rho}}{|F'_\rho|}, \\
\text{Tr} [\hat{\tau}^{\hat{A}'_2 E} \hat{\tau}^{\hat{A}'_2 E}] &\leq \frac{(1 + 3\sqrt{\epsilon})(\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^\epsilon(A'_2|E)_\tau}}{|F'_\tau|}, \\
\text{Tr} [\hat{\rho}^{\hat{A}_2 R} \hat{\rho}^{\hat{A}_2 R}] &\leq \frac{(1 + 3\sqrt{\epsilon}) \cdot 2^{-H_{\min}^\epsilon(A_2|R)_\rho}}{|F'_\rho|}, \\
\text{Tr} [\hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E} \hat{\tau}^{\hat{A}'_1 \hat{A}'_2 E}] &\leq \frac{(1 + 3\sqrt{\epsilon})(\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^\epsilon(A'_1 A'_2|E)_\tau}}{|F'_\tau|}, \\
\text{Tr} [\hat{\rho}^{\hat{A}_1 \hat{A}_2 R} \hat{\rho}^{\hat{A}_1 \hat{A}_2 R}] &\leq \frac{(1 + 3\sqrt{\epsilon}) \cdot 2^{-H_{\min}^\epsilon(A_1 A_2|R)_\rho}}{|F'_\rho|}.
\end{aligned}$$

Plugging Lemma 25 into Equation 86 and using the constraints on the entropic terms and Hilbert space dimensions given in the assumption of the theorem, we get

$$\begin{aligned}
&\mathbb{E}_{U^{\hat{A}_1}, U^{\hat{A}_2}} [\text{Tr} [\hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER} \hat{\sigma}(U^{\hat{A}_1}, U^{\hat{A}_2})^{ER}]] \\
&\leq \frac{|A_1|^2 |A_2|^2}{(|A_1|^2 - 1)(|A_2|^2 - 1)} \cdot \frac{(1 + \sqrt{\epsilon})^2 (\text{Tr } \tau)^2}{|F'_\tau| |F'_\rho|} \\
&\quad + \frac{|A_1|^2 |A_2|^2}{(|A_1|^2 - 1)(|A_2|^2 - 1)} \cdot \frac{(1 + 3\sqrt{\epsilon})^2 (\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^\epsilon(A'_1|E)_\tau - H_{\min}^\epsilon(A_1|R)_\rho}}{|F'_\tau| |F'_\rho|} \\
&\quad + \frac{|A_1|^2 |A_2|^2}{(|A_1|^2 - 1)(|A_2|^2 - 1)} \cdot \frac{(1 + 3\sqrt{\epsilon})^2 (\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^\epsilon(A'_2|E)_\tau - H_{\min}^\epsilon(A_2|R)_\rho}}{|F'_\tau| |F'_\rho|} \\
&\quad + \frac{|A_1| |A_2|}{(|A_1|^2 - 1)(|A_2|^2 - 1)} \cdot \frac{(1 + \sqrt{\epsilon})(1 + 3\sqrt{\epsilon})(\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^\epsilon(A_1 A_2|R)_\rho}}{|F'_\tau| |F'_\rho|} \\
&\quad + \frac{|A_1|^2 |A_2|^2}{(|A_1|^2 - 1)(|A_2|^2 - 1)} \cdot \frac{(1 + 3\sqrt{\epsilon})^2 (\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^\epsilon(A'_1 A'_2|E)_\tau - H_{\min}^\epsilon(A_1 A_2|R)_\rho}}{|F'_\tau| |F'_\rho|} \\
&\leq \frac{|A_1|^2 |A_2|^2}{(|A_1|^2 - 1)(|A_2|^2 - 1)} \cdot \frac{(1 + 3\sqrt{\epsilon})(\text{Tr } \tau)^2}{|F'_\tau| |F'_\rho|} \\
&\quad + \frac{|A_1|^2 |A_2|^2}{(|A_1|^2 - 1)(|A_2|^2 - 1)} \cdot \frac{(1 + 9\sqrt{\epsilon})(\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^\epsilon(A'_1|E)_\tau - H_{\min}^\epsilon(A_1|R)_\rho}}{|F'_\tau| |F'_\rho|} \\
&\quad + \frac{|A_1|^2 |A_2|^2}{(|A_1|^2 - 1)(|A_2|^2 - 1)} \cdot \frac{(1 + 9\sqrt{\epsilon})(\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^\epsilon(A'_2|E)_\tau - H_{\min}^\epsilon(A_2|R)_\rho}}{|F'_\tau| |F'_\rho|} \\
&\quad + \frac{|A_1|^2 |A_2|^2}{(|A_1|^2 - 1)(|A_2|^2 - 1)} \cdot \frac{(1 + 7\sqrt{\epsilon})(\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^\epsilon(A'_1 A'_2|E)_\tau - H_{\min}^\epsilon(A_1 A_2|R)_\rho}}{|F'_\tau| |F'_\rho|}
\end{aligned}$$

$$\begin{aligned}
& + \frac{|A_1|^2 |A_2|^2}{(|A_1|^2 - 1)(|A_2|^2 - 1)} \cdot \frac{(1 + 9\sqrt{\epsilon})(\text{Tr } \tau)^2 \cdot 2^{-H_{\min}^{\epsilon}(A'_1 A'_2 | E)_{\tau} - H_{\min}^{\epsilon}(A_1 A_2 | R)_{\rho}}}{|F'_{\tau}| |F'_{\rho}|} \\
& \leq (1 - \epsilon)^{-2} \cdot \frac{(1 + 3\sqrt{\epsilon})(\text{Tr } \tau)^2}{|F'_{\tau}| |F'_{\rho}|} + (1 - \epsilon)^{-2} \cdot \frac{\epsilon(1 + 9\sqrt{\epsilon})(\text{Tr } \tau)^2}{|F'_{\tau}| |F'_{\rho}|} \\
& \quad + (1 - \epsilon)^{-2} \cdot \frac{\epsilon(1 + 9\sqrt{\epsilon})(\text{Tr } \tau)^2}{|F'_{\tau}| |F'_{\rho}|} + (1 - \epsilon)^{-2} \cdot \frac{\epsilon(2 + 16\sqrt{\epsilon})(\text{Tr } \tau)^2}{|F'_{\tau}| |F'_{\rho}|} \\
& \leq \frac{(1 + 9\sqrt{\epsilon})(\text{Tr } \tau)^2}{|F'_{\tau}| |F'_{\rho}|} + \frac{14\epsilon(\text{Tr } \tau)^2}{|F'_{\tau}| |F'_{\rho}|} < \frac{(1 + 23\sqrt{\epsilon})(\text{Tr } \tau)^2}{|F'_{\tau}| |F'_{\rho}|},
\end{aligned}$$

where we used Fact 4 in the second inequality above. Combined with Equation 83, Theorem 3 is finally proved.

## 9 Conclusion

In this paper, we have noted how the telescoping technique to prove multipartite soft covering results fails when pairwise independence is lost amongst certain probability distributions involved in soft covering. We have then developed a sophisticated machinery that recovers the desired smooth inner bounds for multipartite soft covering under a slight loss of pairwise independence described precisely in Definition 3. To develop this machinery, we had to combine existing results on tilting and augmentation smoothing of quantum states from [Sen21b] together with a novel observation that the naive classical strategy for flattening probability distributions in fact preserves the fidelity of quantum states. We have then seen how this machinery allows us to prove a new inner bound for sending private classical information over a wiretap quantum interference channel by combining existing inner bounds in the non wiretap case from [Sen21a] together with our smooth soft covering results. Such an inner bound in the non pairwise independent case was unknown earlier even in the classical asymptotic iid setting.

Our work leads to natural avenues for further research. One direction to pursue is to further extend our tilting, augmentation smoothing and flattening based machinery, and find more applications to quantum Shannon theory. In spirit, our machinery is arguably much closer to simultaneous smoothing than telescoping. It gives a unified treatment of one shot multiparty smooth packing type questions as in [Sen21b, Sen21a, DGHW20], as well as one one shot smooth multiparty convex split and soft covering questions described in this paper. ‘ However, one important application where our machinery fails is in proving a fully smooth multipartite expander matrix Chernoff bound. The paper [Sen25] proves a fully smooth multipartite classical quantum soft covering lemma in concentration, aka fully smooth multipartite matrix Chernoff bound, but for independent choices of samples by each classical party. That paper also proves a smooth unipartite soft covering lemma in concentration when the samples are taken by the single classical party from a random walk on an expander graph, which can be called a unipartite expander matrix Chernoff bound. However, the techniques of that paper fail to prove a multipartite expander matrix Chernoff bound. Expander walks are a key example where pairwise independence amongst the classical samples of a party is lost. Unfortunately the loss of pairwise independence does not fit the framework behind Definition 3. Hence even fully smooth multipartite classical quantum soft covering in expectation under expander walks cannot be handled by the techniques of the present paper. Proving fully smooth multipartite soft covering results in expectation and concentration for expander walks is thus a very important open problem.

It is also important to find limitations of our tilting and augmentation smoothing based machinery. One immediate limitation that comes to mind is that augmenting quantum states requires tensoring with fresh completely mixed ancilla states. This is required so that tracing out a set of registers essentially removes the tilts along the traced out directions when quantified in terms of the Schatten- $\ell_\infty$  norm. Augmentation can be done without loss of generality for the packing type questions studied in earlier works, as well as in convex split and soft covering problems. However, high dimensional augmentation becomes a bottleneck for decoupling problems. This is why we gave the simpler proof for fully smooth convex split in Section 7 which did not need augmentation for smoothing purposes. The proof in Section 7 only had to enlarge the dimensions of the Hilbert spaces  $X$  and  $Y$  by factor of 2 in order to achieve the correct tilting. It led to the proof of a fully smooth decoupling theorem in Section 8. Though the decoupling theorem in Section 8 suffices for all the applications of decoupling in quantum information theory that we know of, the factor of 2 blowup is still mathematically unsatisfactory. In contrast, the telescoping based proof of fully smooth decoupling in [Sen24] has no such blowup. Removing this factor of 2 for decoupling for our machinery is an interesting open problem.

## Acknowledgements

I sincerely thank Rahul Jain for many stimulating discussions on smooth convex split and flattening quantum states. Many ideas in this work were slowly developed over several visits hosted by him during the last few years at the Centre for Quantum Technologies, National University of Singapore. I also thank the staff at the Centre for their efficient hospitality. This research is supported in part by the National Research Foundation, Singapore and A\*STAR under the CQT Bridging Grant and the Quantum Engineering Programme Award number NRF2021-QEP2-02-P05. I acknowledge support of the Department of Atomic Energy, Government of India, under project no. RTI4001.

## References

- [ABJT19] A. Anshu, M. Berta, R. Jain, and M. Tomamichel. A minimax approach to one-shot entropy inequalities. *Journal of Mathematical Physics*, 60:122201:1–122201:9, 2019.
- [AJ22] A. Anshu and R. Jain. Efficient methods for one-shot quantum communication. *npj Quantum Information*, 8:97:1–97:46, 2022.
- [AJW18] A. Anshu, R. Jain, and N. Warsi. A generalized quantum Slepian–Wolf. *IEEE Transactions on Information Theory*, 64:1436–1453, 2018.
- [CGB23] H-C. Cheng, L. Gao, and M. Berta. Quantum broadcast channel simulation via multipartite convex splitting. arXiv:2304.12056, 2023.
- [CMGE08] Chong, H., Motani, M., Garg, H., and El Gamal, H. On the Han-Kobayashi region for the interference channel. *IEEE Transactions on Information Theory*, 54:3188–3195, 2008.
- [CNS21] S. Chakraborty, A. Nema, and P. Sen. A multi-sender decoupling theorem and simultaneous decoding for the quantum mac. In *IEEE International Symposium on Information Theory (ISIT)*, 2021. Full version in arXiv::2102.02187.

- [CW23] P. Colomer and A. Winter. Decoupling by local random unitaries without simultaneous smoothing, and applications to multi-user quantum information tasks. *arXiv:2304.12114*, 2023.
- [DF13] L. Drescher and O. Fawzi. On simultaneous min-entropy smoothing. In *IEEE International Symposium on Information Theory (ISIT)*, pages 161–165, 2013.
- [DGHW20] Ding, D., Gharibyan, H., Hayden, P., and Walter, M. A quantum multiparty packing lemma and the relay channel. *IEEE Transactions on Information Theory*, 66:3500–3519, 2020.
- [RSW17] J. Radhakrishnan, P. Sen, and N. Warsi. One-shot private classical capacity of quantum wiretap channel: Based on one-shot quantum covering lemma. Proceedings of QCRYPT workshop. Also *arXiv:1703.01932*, 2017.
- [SDTR13] O. Szehr, F. Dupuis, M. Tomamichel, and R. Renner. Decoupling with unitary approximate two-designs. *New Journal of Physics*, 15:053022:1–053022:20, 2013.
- [Sen12] P. Sen. Achieving the Han-Kobayashi inner bound for the quantum interference channel. In *IEEE International Symposium on Information Theory (ISIT)*, pages 736–740, 2012. Full version at *arXiv:1109.0802*.
- [Sen21a] P. Sen. Inner bounds via simultaneous decoding in quantum network information theory. *Sadhana*, 46:0018:1–0018:20, 2021. Also available at *arXiv:1806.07276*.
- [Sen21b] P. Sen. Unions, intersections and a one shot quantum joint typicality lemma. *Sadhana*, 46:0057:1–0057:44, 2021. Also available at *arXiv:1806.07278*.
- [Sen24] P. Sen. Fully smooth one shot multipartite covering and decoupling of quantum states via telescoping. *arXiv:2410.17893*, 2024.
- [Sen25] P. Sen. Matrix Chernoff concentration bounds for multipartite soft covering and expander walks. *arXiv:2504.04067*, 2025.
- [Wil17] M. Wilde. Position-based coding and convex splitting for private communication over quantum channels. *Quantum Information Processing*, 16:264:1–264:31, 2017.