

AN EIGENRING CONSTRUCTION FOR FREE GROUPS AND FREE MODULES

MINKYU KIM

ABSTRACT. Let \mathbb{k} be a commutative unital ring. For a \mathbb{k} -algebra A with a multiplicative submodule J , we have a canonical subquotient algebra $E_A(J)$ called the *eigenring*. In the case where J is a left ideal, an adjunction exists between the category of A -modules and that of $E_A(J)$ -modules. The derived functors associated with this adjunction recover several well-known homological invariants, including group (co)homology and Hochschild (co)homology.

The purpose of this paper is to present an extension of these constructions, where \mathbb{k} -algebras are replaced with monads within a specified bicategory. This approach may be regarded as a categorification of eigenrings, given that monads in the bicategory are equivalent with \mathbb{k} -linear categories. Furthermore, we generalize the aforementioned adjunction, demonstrating how the associated derived functors provide examples of functor homology.

For a category C , a module over C is defined as a functor from C to the category of \mathbb{k} -modules and \mathbb{k} -linear maps. In the case where C is either the opposite category of *free groups* or that of *free modules* over a ring R , specific types of modules over C naturally emerge within the context of algebraic topology. These include polynomial modules, analytic modules, outer modules and primitively generated modules. The majority of this paper is devoted to applying the general framework to the study of these module types. The categorification of eigenrings leads to a unified framework in which these types are characterized by certain internal data of the monad L_C associated with C . In applications, we obtain calculations of particular functor homologies and uncover new aspects of the above module types.

CONTENTS

1. Introduction	2
1.1. A bicategory of matrices $\text{Mat}_{\mathbb{k}}$	3
1.2. Categorification of eigenrings	4
1.3. Internalization problem for classes of modules	5
1.4. Solutions for specific classes of modules	6
1.5. Further results	11
1.6. Organization	12
Acknowledgements	13
2. Notation	13
Part 1. General eigenring construction	13
3. The bicategory of matrices $\text{Mat}_{\mathbb{k}}$	13
4. Monad theory for bi-indexed modules	15
4.1. Review of monads	15
4.2. Monads in $\text{Mat}_{\mathbb{k}}$	16
4.3. Modules in $\text{Mat}_{\mathbb{k}}$	17
4.4. Hom and Tensor	18
5. Eigenmonads	20
5.1. Definition of eigenmonads	20
5.2. Construction of eigenmonad adjunctions	21
5.3. Vanishing modules	22
5.4. Example: algebras	23
6. Ideals and classes of modules	24

6.1. Refined eigenmonad adjunctions	24
6.2. Internalizers of classes of modules	26
7. Example: polynomial functors	28
7.1. Polynomial degrees	28
7.2. Polynomial ideals	29
7.3. Core internalizer for polynomial modules	30
7.4. Internalizers for analytic modules	32
7.5. Comparison with augmentation ideals	33
Part 2. Application to free groups	40
8. Notation	40
9. Polynomial ideals for free groups	41
9.1. Eigenmonads by polynomial ideals	41
9.2. Internalizer for analytic modules	43
10. Outer modules over free groups	43
10.1. Core internalizer for outer modules	43
10.2. Eigenmonad and group homology	44
11. Primitively generated modules over free groups	45
11.1. Definition of primitivity ideal	45
11.2. Canonical bimodule associated with primitivity ideal	45
11.3. Eigenmonad by primitivity ideal	47
11.4. Primitively generated modules	49
12. Comparison of primitivity and analyticity	49
Part 3. Application to free modules	50
13. Notation	50
14. Abelianization	51
15. Polynomial ideals for free modules	53
15.1. Eigenmonads for polynomial ideals	53
15.2. Internalizers for analytic modules	53
15.3. Comparison of polynomial ideals for gr^0 and $\text{fr}_{\mathbb{Z}}^0$	54
16. Primitively generated modules over free modules	55
16.1. Eigenmonad associated with primitivity ideal	56
16.2. Concrete description of the canonical bimodule	58
16.3. Proof of (1) of Theorem 16.4	59
16.4. Proof of (2) of Theorem 16.4	61
16.5. Refined eigenmonad adjunctions	62
17. Comparison of primitivity and analyticity	63
Appendix A. Enough injectives and projectives in τMod	65
Appendix B. Graded coaugmented coalgebras	65
Appendix C. A grading on free Lie algebras	67
References	69

1. INTRODUCTION

Let \mathbb{k} be a commutative unital ring. For a \mathbb{k} -algebra A with a submodule $J \subset A$ closed under the multiplication, there exists a canonical subquotient algebra of A by J called the *eigenring* [17, 3], which we denote by $E_A(J)$. This is defined as the quotient algebra of the *idealizer* that is the maximal subalgebra of A containing J as a two-sided ideal. The notion of

an eigenring generalizes that of a quotient algebra, since if J is a two-sided ideal, then $E_A(J)$ coincides with the quotient algebra A/J .

If $J \subset A$ is a left ideal, then the eigenring can be described in equivalent ways using homological invariants. Indeed, the opposite of the eigenring $E_A(J)^\circ$ is isomorphic to the algebra of endomorphisms on the left A -module A/J , so that it is embedded into the Ext ring on A/J :

$$(1) \quad \text{Ext}_A^0(A/J, A/J) \cong E_A(J)^\circ.$$

These observations give rise to the following tensor-hom adjunction between the category ${}_A\text{Mod}$ of left A -modules and the category ${}_{E_A(J)}\text{Mod}$ of left $E_A(J)$ -modules:

$$(2) \quad A/J \otimes_{E_A(J)} (-) : {}_{E_A(J)}\text{Mod} \xrightleftharpoons[\tau]{\tau} {}_A\text{Mod} : \text{Hom}_A(A/J, -).$$

We also have an analog of (2) for right A -modules and right $E_A(J)$ -modules. The derived functors associated these adjunctions contain basic homological invariants of fundamental importance. Group (co)homology and Hochschild (co)homology are illustrative examples, as demonstrated in section 5.4.

This paper presents a categorification of the aforementioned constructions, in which algebras are replaced with \mathbb{k} -linear categories (or, equivalently, *monads* in a bicategory of matrices, which will be explained later). We give a generalization of (2) whose associated derived functors are endowed with a canonical action of the categorified eigenring, which we refer to as an *eigenmonad*. The derived functors are examples of functor homology (see, for example, [7]).

The results presented herein provide a unified framework that encompasses a specific type of functors from a category C to the category of \mathbb{k} -modules ${}_{\mathbb{k}}\text{Mod}$, which naturally emerges within the context of algebraic topology:

- *Polynomial* functors and *analytic* functors for an abelian category C [5, 6, 4]; for the opposite category $C = \text{gr}^\circ$ of finitely generated free groups and group homomorphisms [21, 11, 29]; and for general C [9, section 2].
- *Outer* functors for $C = \text{gr}^\circ$ which arise from Hochschild-Pirashvili homology for wedge of circles [19, 24] and in quantum topology [11, 29].
- *Primitively generated* functors for $C = \text{gr}^\circ$ [12].

Based on our framework, we characterize these particular types of functors by certain *internal data* of C . The reader is referred to Problem 1.8 for further details. In consequence, we present new aspects of the functors described above (see Theorems 1.11, 1.14, 1.15 and 1.16) and their relations (see Theorem 1.18 and section 1.5.3).

In section 1.1, we introduce a bicategory $\text{Mat}_{\mathbb{k}}$, which is used to develop a categorification of eigenrings in section 1.2. In section 1.3, we present the central problem of this paper by using the results of section 1.2. This is connected to the preceding examples, and solved for them in section 1.4. In section 1.5, we present further results by applying the solutions.

1.1. A bicategory of matrices $\text{Mat}_{\mathbb{k}}$. A bicategory is a concept that extends the notion of category by allowing the composition to be associative up to an isomorphism [1]. The concept of a monad in a bicategory serves to generalise the formal aspect of a monoid or an algebra. In this paper, we develop a categorification of eigenrings by treating *monads* in a specific bicategory $\text{Mat}_{\mathbb{k}}$ called the *bicategory of matrices* [27].

Definition Sketch 1.1 (see Definition 3.1). The bicategory $\text{Mat}_{\mathbb{k}}$ consists of the following data:

- (1) Objects are sets $\mathcal{X}, \mathcal{Y}, \mathcal{Z} \dots$.

- (2) A morphism F from \mathcal{X} to \mathcal{Y} is given by a *matrix* whose rows (columns, resp.) are parametrized by \mathcal{Y} (\mathcal{X} , resp.) and whose entries $F(Y, X)$ are \mathbb{k} -modules for $Y \in \mathcal{Y}, X \in \mathcal{X}$:

$$F = \begin{pmatrix} & \vdots & \\ \cdots & F(Y, X) & \cdots \\ & \vdots & \end{pmatrix}.$$

The composition of morphisms is defined by the *matrix product* formula using the direct sum and tensor product of \mathbb{k} -modules. For composable morphisms F, G , we denote by $F \otimes G$ their composition. The identity morphism $\mathbb{1}_{\mathcal{X}}$ on \mathcal{X} is given by the diagonal matrix whose nontrivial components are \mathbb{k} .

- (3) A 2-morphism between morphisms is given by a \mathbb{k} -linear map entry-wise.

Definition Sketch 1.2 (see section 4.1). For an object \mathcal{X} , a *monad* in $\text{Mat}_{\mathbb{k}}$ on \mathcal{X} is defined to be a morphism T from \mathcal{X} to itself, which is endowed with 2-morphisms $\nabla : T \otimes T \rightarrow T$ and $\eta : \mathbb{1}_{\mathcal{X}} \rightarrow T$ satisfying *algebra-like* axioms. Furthermore, for monads T and S in $\text{Mat}_{\mathbb{k}}$ (on, possibly, distinct objects), the notions of a left (or right) T -module and a (T, S) -bimodule are defined in a manner analogous to those in ring-module theory.

There are two reasons why it is *advantageous* to deal with $\text{Mat}_{\mathbb{k}}$. Firstly, we may study \mathbb{k} -linear categories through monad theory in $\text{Mat}_{\mathbb{k}}$ since there is one-to-one correspondence between monads in $\text{Mat}_{\mathbb{k}}$ and \mathbb{k} -linear categories. The reader is referred to section 4 for further details.

Secondly, the bicategory $\text{Mat}_{\mathbb{k}}$ allows us general constructions and notions beyond \mathbb{k} -linear categories. We can extend the usual concepts for \mathbb{k} -modules such as a quotient module, a submodule and a (left or right) ideal to indexed \mathbb{k} -modules, i.e. morphisms of $\text{Mat}_{\mathbb{k}}$. These are essential ingredients for our categorification of eigenrings. Along with these, we can still do homological algebra on T -modules.

1.2. Categorification of eigenrings. Let T be a monad in $\text{Mat}_{\mathbb{k}}$ on an object $\mathcal{X} \in \text{Mat}_{\mathbb{k}}$.

Definition Sketch 1.3 (see Definition 5.3). A submodule $J \subset T$ (as an indexed module) is *multiplicative* if it is closed under the monad operation of T . For a multiplicative submodule $J \subset T$, we define the *eigenmonad* $E_T(J)$ of T by J as a canonical subquotient monad of T by mimicking the concept of an eigenring.

The formal resemblance of monads to algebras lead to the following fundamental proposition:

Proposition 1.1. *Let T be a monad in $\text{Mat}_{\mathbb{k}}$ on $\mathcal{X} \in \text{Mat}_{\mathbb{k}}$, and $J \subset T$ be a left ideal.*

- (1) *(Proof in Proposition 5.13) We have a natural $(T, E_T(J))$ -bimodule structure on the quotient T/J , which induces the following tensor-hom adjunction called the eigenmonad adjunction:*

$$(3) \quad T/J \otimes_{E_T(J)} (-) :_{E_T(J)} \text{Mod} \xrightleftharpoons{T} {}_T \text{Mod} : \text{Hom}_T(T/J, -).$$

- (2) *(Proof in Proposition 5.20) Let $\text{Ext}_T^\bullet(T/J, -)$ be the derived functors of the right adjoint in (3). Then $\text{Ext}_T^\bullet(T/J, T/J)$ is a graded monad in $\text{Mat}_{\mathbb{k}}$ on \mathcal{X} , and we have a monad isomorphism:*

$$\text{Ext}_T^0(T/J, T/J) \cong E_T(J)^\circ.$$

Example 1.2. The classical results (1) and (2) are special cases of (2) and (1) of the theorem respectively, since a monad in $\text{Mat}_{\mathbb{k}}$ on a one-point set is nothing but a \mathbb{k} -algebra.

There are classical examples where the adjunction (3) yields an equivalence of categories, as demonstrated in sections 5.4.2 and 5.4.3. However, in general, this does not necessarily lead to an equivalence. In this paper, we go further by giving a canonical refinement of the eigenmonad adjunction (see Proposition 6.7), which contains a number of equivalences of categories (see Corollaries 6.8, 7.15, 10.9, 12.3 and 17.6).

Remark 1.3. We also have a similar adjunction for right modules, as shown in (2) of Proposition 5.13. Analogously, we have a Tor functor $\mathrm{Tor}_{\bullet}^{\mathbb{T}}(-, \mathbb{T}/J)$ on right \mathbb{T} -modules.

1.3. Internalization problem for classes of modules. As mentioned earlier, this work investigates certain adjunctions that appear in the context of algebraic topology. One objective of this paper is to derive those from eigenmonad adjunctions in Proposition 1.1. In order to formulate a more precise problem, we introduce the following notion:

Definition 1.4. Let \mathbb{M} be a left \mathbb{T} -module. We say that \mathbb{M} is *J-vanishingly generated* if the counit of the eigenmonad adjunction is an epimorphism:

$$\mathbb{T}/J \otimes_{E_{\mathbb{T}}(J)} \mathrm{Hom}_{\mathbb{T}}(\mathbb{T}/J, \mathbb{M}) \twoheadrightarrow \mathbb{M}.$$

This terminology is motivated by the equivalent condition that \mathbb{M} is generated by the maximal submodule of \mathbb{M} on which J acts trivially (see Definition 6.1).

Definition 1.5. Denote by ${}_{\mathbb{T}}\mathrm{Mod}$ ¹ the class of isomorphism classes of left \mathbb{T} -modules. For a left ideal $J \subset \mathbb{T}$, we define $\mathcal{V}(J) \subset {}_{\mathbb{T}}\mathrm{Mod}$ to be the subclass of isomorphism classes of J -vanishingly generated left \mathbb{T} -modules. This gives rise to a map:

$$(4) \quad \mathcal{V} : \{\text{Left ideals of } \mathbb{T}\} \longrightarrow \{\text{Subclasses of } {}_{\mathbb{T}}\mathrm{Mod}\}.$$

Definition 1.6. Let \mathcal{S} be a subclass of ${}_{\mathbb{T}}\mathrm{Mod}$. A left ideal J of \mathbb{T} such that $\mathcal{S} = \mathcal{V}(J)$ is called a *T-internalizer of \mathcal{S}* . If it is a two-sided ideal, then it is called a *core T-internalizer of \mathcal{S}* .

These terminologies are predicated on our expectation that \mathcal{S} is essentially characterised by the internal data of \mathbb{T} , specifically the ideal J .

Example 1.7. We give examples of core internalizers in Theorems 1.11 and 1.14. In contrast, Theorems 1.13 and 1.15 provide examples of non-core internalizers.

Problem 1.8. In the present paper, we discuss the following problems with respect to particular \mathbb{T} and $\mathcal{S} \subset {}_{\mathbb{T}}\mathrm{Mod}$:

- (A) Does there exist a (core) \mathbb{T} -internalizer of \mathcal{S} ?
- (B) For a \mathbb{T} -internalizer J of \mathcal{S} , investigate the structure of the quotient \mathbb{T}/J and the eigenmonad $E_{\mathbb{T}}(J)$ (or, equivalently, $\mathrm{Ext}_{\mathbb{T}}^0(\mathbb{T}/J, \mathbb{T}/J)$).

If we have affirmative answers to Problem 1.8, then we obtain the eigenmonad adjunction associated with J . It is anticipated that the adjunction will serve as an ‘‘approximation’’ of the category of \mathbb{T} -modules by the category of $E_{\mathbb{T}}(J)$ -modules, since the notion of an adjunction is defined as a weak form of an equivalence of categories. Indeed, this contains several well-known examples such as a polynomial approximation (see Corollary 7.14) and an outer approximation (see Corollary 10.9). This is sometimes refined to an equivalence of categories. This is the underlying motivation behind Problem 1.8, which is illustrated in the

¹Note that its initial letter is curly, whereas the symbol used to represent a category of modules is written as Mod

following schematic diagram:

$$(5) \quad \begin{array}{ccc} & \text{Internalizers} & \\ & \curvearrowright & \\ \{\text{Left ideals of } T\} & \xrightarrow{\mathcal{V}} & \{\text{Subclasses of } {}_T\text{Mod}\} \\ \text{Eigenmonad adj.} \downarrow & & \\ \{\text{“Approximations” of } {}_T\text{Mod}\} & & \end{array}$$

Here, we notify that the dashed arrow “Internalizers” is not a map. In fact, there exists a subclass \mathcal{S} of ${}_T\text{Mod}$ having numerous T -internalizers.

One of the main results is the following theorem which demonstrates that a *core* internalizer has a distinctive nature:

Theorem 1.9. *Let $\mathcal{S} \subset {}_T\text{Mod}$.*

- (1) *A core T -internalizer of \mathcal{S} , if it exists, is contained in any T -internalizer of \mathcal{S} . In particular, there exists at most one core T -internalizer of \mathcal{S} .*
- (2) *A left T -module whose isomorphism class lies in \mathcal{S} is canonically a left $E_T(\mathcal{J})$ -module:*

$$\mathcal{S} \cong_{E_T(\mathcal{J})} \text{Mod}.$$

The part (1) follows from Propositions 6.15. The part (2) is proved by applying Example 5.9 to Corollary 6.8.

It should be noted that, in general, a (non-core) T -internalizer is not unique. There might be uncountably many T -internalizers (see (A-1) of Theorem 1.13).

Remark 1.10. In this paper, we go further by proving that a core T -internalizer of \mathcal{S} , if it exists, is equal to the intersection of the annihilators of all the T -modules in \mathcal{S} (see Theorem 6.20).

1.4. Solutions for specific classes of modules. This section is dedicated to an investigation of Problem 1.8 with respect to a particular T induced by a category \mathcal{C} . There are two categories of our main interest. The first is the opposite category gr° of finitely generated free groups and group homomorphisms. The second one is the opposite category fr_R° of free R -modules and R -homomorphisms for a unital (possibly, noncommutative) ring R . In this paper, we identify the object sets of gr° (fr_R° , resp.) with $\mathbb{N} = \{0, 1, \dots\}$ based on the number of generators of free groups (free modules, resp.). For such a category \mathcal{C} , we define $L_{\mathcal{C}}$ as the monad in $\text{Mat}_{\mathbb{k}}$ on \mathbb{N} induced by the \mathbb{k} -linearization of the morphism sets of \mathcal{C} :

$$L_{\mathcal{C}}(Y, X) := \mathbb{k}C(X, Y), \quad X, Y \in \text{Obj}(\mathcal{C}).$$

The monad structure is induced by the category structure of \mathcal{C} .

For a category \mathcal{C} , a *module over \mathcal{C}* is a functor from \mathcal{C} to the category ${}_{\mathbb{k}}\text{Mod}$. A module over \mathcal{C} is nothing but a left $L_{\mathcal{C}}$ -module in the sense of monad theory in $\text{Mat}_{\mathbb{k}}$, as we mentioned in section 1.1 (or, see Example 4.16).

In section 1.4.1, we give our answers to Problem 1.8 concerning polynomial modules. In section 1.4.2, we similarly present the answers on analytic modules. In section 1.4.3, we consider outer modules on gr° . In section 1.4.4, we study a certain type of modules called primitively generated modules, which generalizes the notion of primitively generated Hopf algebras.

1.4.1. Polynomial modules. If \mathcal{C} is a symmetric monoidal category whose unit object is a zero object, then the *polynomial degree* of a module over \mathcal{C} is defined by using certain conditions that are analogous to those satisfied by polynomial functions [9, section 2]. The origin

of this notion goes back to Eilenberg-MacLane [5] in the case that C is the category of abelian groups. For an abelian category C , the study of polynomial modules over C has been applied to algebraic topology (see, for example, [6]). Recently, the study of polynomial modules over \mathfrak{gr}^0 was spotlighted in the context of quantum topology [8, 11, 29]. The present paper is partly motivated by the algebraic aspects of the literature.

For a group G , we denote by $I_{\mathbb{k}}(G)$ the augmentation ideal of G over \mathbb{k} . For $d \in \mathbb{N}$, we define $\mathcal{P}^d(G)$ to be the quotient $\mathbb{k}[G]/I_{\mathbb{k}}(G)^{d+1}$. This is called the d -th *Passi functor* (see, for example, [22]) which was introduced to study polynomial maps on a monoid [18]. For $C \in \{\mathfrak{gr}^0, \mathfrak{fr}_R^0\}$, we define an $(\mathbb{N} \times \mathbb{N})$ -indexed module \mathcal{P}_C^d to be

$$\mathcal{P}_C^d(m, n) := \mathcal{P}^d(C_n^{\times m}), \quad n, m \in \mathbb{N}$$

where C_n is the free group F_n generated by n elements if $C = \mathfrak{gr}^0$; or C_n is the underlying group of the free R -module R^n generated by n elements if $C = \mathfrak{fr}_R^0$.

In what follows, we give the answers to (A) and (B) of Problem 1.8:

Theorem 1.11. *Suppose that C is a category with finite products and a zero object. For $d \in \mathbb{N}$, let \mathcal{S}_C^d be the class of isomorphism classes of left L_C -modules whose corresponding modules over C have the polynomial degree $\leq d$.*

(A) *The subclass $\mathcal{S}_C^d \subset {}_{L_C}\mathbf{Mod}$ has a unique core L_C -internalizer $\mathbb{I}_C^{(d+1)}$. In particular, by Theorem 1.9, we have*

$$\mathcal{S}_C^d \cong_{E_{L_C}(\mathbb{I}_C^{(d+1)})} \mathbf{Mod}.$$

(B) *For $C \in \{\mathfrak{gr}^0, \mathfrak{fr}_R^0\}$, we have an isomorphism of indexed \mathbb{k} -modules:*

$$E_{L_C}(\mathbb{I}_C^{(d+1)}) \cong \mathcal{P}_C^d.$$

The part (A) is proved in Theorem 7.13. Note that the uniqueness follows from Theorem 1.9. The part (B) implies that \mathcal{P}_C^d is a monad in $\mathbf{Mat}_{\mathbb{k}}$ on $\mathbf{Obj}(C) = \mathbb{N}$. The part (B) is proved from more general context in Theorem 7.30.

A key distinction between polynomial *functions* and general ones is that the former are described by finite data, typically the coefficients of monomials. This contrasts with the definition of polynomial *functors*, which does not explicitly indicate that they can be described by “finite” data. Is it possible to describe polynomial functors by utilizing “finite” data in any sense?

Corollary 1.12. *Let $C \in \{\mathfrak{gr}^0, \mathfrak{fr}_R^0\}$. For a left L_C -module \mathbf{M} , we have $\deg \mathbf{M} \leq d$ if and only if, for $n, m \in \mathbb{N}$, the L_C -action of \mathbf{M} restricted to the monoid $C_n^{\times m} \cong C(n, m)$ is a polynomial map of degree $\leq d$ [18, Definition 1.1, Chapter V]:*

$$C_n^{\times m} \hookrightarrow L_C(m, n) \xrightarrow{\mathbf{M}} \mathbf{hom}_{\mathbb{k}}(\mathbf{M}(n), \mathbf{M}(m)).$$

This is proved by (B) of Theorem 1.11, which indicates that the eigenmonad factors through the Passi functor. The corollary gives an answer to the above question, since, for $C \in \{\mathfrak{gr}^0, \mathfrak{fr}_R^0\}$, the \mathbb{k} -module $\mathcal{P}^d(C_n^{\times m})$ has a finite rank (see Theorems 9.1 and 15.4). The reader is referred to Remark 7.33 for further details.

1.4.2. *Analytic modules.* This section is dedicated to the discussion on analytic modules over C . We shall continue to consider the hypothesis on C of Theorem 1.11. It is known that every module over C possesses a filtration indexed by polynomial degrees (see, for instance, [12]). If the filtration converges to itself, the module over C is called *analytic*.

We now solve Problem 1.8 for analytic modules. The answer to (A) of Problem 1.8 is contingent upon whether an internalizer is considered to be core or not. We give a partial

answer to (B) of Problem 1.8. Let $\Gamma(C)$ be the set of a map from $\text{Obj}(C)$ to \mathbb{N} whose image is unbounded.

Theorem 1.13. *Suppose that C is a category with finite products and a zero object. Let \mathcal{S}_C^ω be the class of isomorphism classes of L_C -modules whose corresponding modules over C are analytic.*

- (A-1) *The subclass $\mathcal{S}_C^\omega \subset {}_{L_C}\text{Mod}$ has an L_C -internalizer. Furthermore, we have an assignment of an L_C -internalizer \mathbb{I}_C^ν to $\nu \in \Gamma(C)$, which is injective for $C \in \{\text{gr}^\circ, \text{fr}_\mathbb{Z}^\circ\}$.*
 (A-2) *For $C \in \{\text{gr}^\circ, \text{fr}_\mathbb{Z}^\circ\}$, the subclass $\mathcal{S}_C^\omega \subset {}_{L_C}\text{Mod}$ does not allow a core L_C -internalizer.*
 (B) *Let $C \in \{\text{gr}^\circ, \text{fr}_\mathbb{R}^\circ\}$ and $\nu \in \Gamma(C)$. For $X, Y \in \text{Obj}(C)$ such that $\nu(Y) \geq \nu(X)$, we have an isomorphism*

$$(E_{L_C}(\mathbb{I}_C^\nu))(Y, X) \cong \mathcal{P}_C^{\nu(X)-1}(Y, X).$$

In Definition 7.21, we give a concrete construction of \mathbb{I}_C^ν and, in Theorem 7.25, we prove (A-1). In particular, the injectiveness follows from Propositions 9.4 and 15.5. The part (A-2) is proved in Theorems 9.5 and 15.6 respectively. The part (B) follows from (B) of Theorem 1.11 and Proposition 7.27.

1.4.3. *Outer modules.* A module over gr° is *outer* if it assigns the identity to any inner automorphisms on each group [24, 11, 29].

Let $H_0(F_n; M)$ be the zeroth group homology of F_n with coefficients in a F_n -module M . We define an $(\mathbb{N} \times \mathbb{N})$ -indexed \mathbb{k} -module H_0 to be

$$H_0(m, n) := H_0(F_n; \mathbb{k}[F_n^{\times m}]), n, m \in \mathbb{N}$$

where the group F_n acts on $\mathbb{k}[F_n^{\times m}]$ conjugately.

The answers to (A) and (B) of Problem 1.8 for outer modules over gr° is provided as follows:

Theorem 1.14. *Let $\mathcal{S}^{\text{out}} \subset {}_{L_{\text{gr}^\circ}}\text{Mod}$ be the subclass of isomorphism classes of L_{gr° -modules whose corresponding module over gr° is outer.*

- (A) *The subclass $\mathcal{S}^{\text{out}} \subset {}_{L_{\text{gr}^\circ}}\text{Mod}$ has a unique core L_{gr° -internalizer \mathbb{I}^{out} . In particular, by Theorem 1.9, we have*

$$\mathcal{S}^{\text{out}} \cong_{E_{L_{\text{gr}^\circ}}(\mathbb{I}^{\text{out}})} \text{Mod}.$$

- (B) *We have an isomorphism of indexed \mathbb{k} -modules:*

$$E_{L_{\text{gr}^\circ}}(\mathbb{I}^{\text{out}}) \cong H_0,$$

The part (B) implies that H_0 is a monad in $\text{Mat}_{\mathbb{k}}$ on \mathbb{N} . The parts (A) and (B) follow from Theorems 10.4 and 10.8 respectively.

1.4.4. *Primitively generated modules.* Let \mathcal{O} be a (symmetric) \mathbb{k} -linear operad [13]. A \mathbb{k} -linear operad induces a \mathbb{k} -linear category by collecting all abstract operations lying in \mathcal{O} . We define $\mathbf{A}_{\mathcal{O}}$ to be the monad in $\text{Mat}_{\mathbb{k}}$ corresponding to the category (see section 1.1). To be precise, for $n, m \in \mathbb{N}$, we define $\mathbf{A}_{\mathcal{O}}$ to be the monad in $\text{Mat}_{\mathbb{k}}$ on \mathbb{N} such that:

$$\mathbf{A}_{\mathcal{O}}(m, n) := \bigoplus_f \bigotimes_{i=1}^m \mathcal{O}(f^{-1}(i)),$$

where f runs over maps from $\underline{n} := \{1, 2, \dots, n\}$ to \underline{m} . Let $\mathfrak{A}_{\text{ss}^u}$ be the operad for unital associative algebras and \mathfrak{A}_{ie} be the operad for Lie algebras. In [21, 12], the following adjunction is provided:

$$(6) \quad \tilde{\mathfrak{A}}_{\mathfrak{A}_{\text{ss}^u}} \otimes_{\mathfrak{A}_{\text{ie}}} (-) : \mathfrak{A}_{\text{ie}} \text{Mod} \begin{array}{c} \xleftarrow{\quad \top \quad} \\ \xrightarrow{\quad \text{L}_{\text{gr}^\circ} \quad} \end{array} \text{L}_{\text{gr}^\circ} \text{Mod} : \text{Hom}_{\text{L}_{\text{gr}^\circ}}(\tilde{\mathfrak{A}}_{\mathfrak{A}_{\text{ss}^u}}, -).$$

Here, $\tilde{\mathbf{A}}_{\mathfrak{L}(S^u)}$ is a $(L_{\text{gr}^0}, \mathbf{A}_{\mathfrak{Qie}})$ -bimodule whose underlying indexed module coincides with that of $\mathbf{A}_{\mathfrak{L}(S^u)}$. This adjunction generalizes the classical relationship between cocommutative Hopf algebras and Lie algebras; and captures the relationship between open Jacobi diagrams and Jacobi diagrams in handlebodies [29].

A left L_{gr^0} -module \mathbf{M} is *primitively generated* if the counit $\tilde{\mathbf{A}}_{\mathfrak{L}(S^u)} \otimes_{\mathbf{A}_{\mathfrak{Qie}}} \text{Hom}_{L_{\text{gr}^0}}(\tilde{\mathbf{A}}_{\mathfrak{L}(S^u)}, \mathbf{M}) \rightarrow \mathbf{M}$ of the adjunction (6) is an epimorphism². This terminology is inspired by the notion of a primitively generated Hopf algebra [16] (see Remark 11.18 for details).

Theorem 1.15. *Let $\mathcal{S}_{\text{gr}^0}^{\text{pr}}$ be the class of isomorphism classes of primitively generated L_{gr^0} -modules.*

- (A) *The subclass $\mathcal{S}_{\text{gr}^0}^{\text{pr}} \subset L_{\text{gr}^0} \text{Mod}$ has a canonical L_{gr^0} -internalizer $\mathbf{I}_{\text{gr}^0}^{\text{pr}}$.*
- (B) *For the left ideal \mathbf{I}^{pr} , we have a monad isomorphism:*

$$E_{L_{\text{gr}^0}}(\mathbf{I}_{\text{gr}^0}^{\text{pr}}) \cong \mathbf{A}_{\mathfrak{Qie}}.$$

Based on this, the $(L_{\text{gr}^0}, E_{L_{\text{gr}^0}}(\mathbf{I}_{\text{gr}^0}^{\text{pr}}))$ -bimodule $L_{\text{gr}^0}/\mathbf{I}^{\text{pr}}$ given by (1) of Proposition 1.1 is isomorphic to the $(L_{\text{gr}^0}, \mathbf{A}_{\mathfrak{Qie}})$ -bimodule $\tilde{\mathbf{A}}_{\mathfrak{L}(S^u)}$.

The part (A) follows from Proposition 11.17 where $\mathbf{I}_{\text{gr}^0}^{\text{pr}}$ is defined in Definition 11.1 and Notation 11.2. The proof of part (B) is presented in Theorems 11.9 and 11.13.

It is notable that the application of (B) to the adjunction (3) reproduces the adjunction (6).

If \mathbb{k} is a field of characteristic zero, (6) induces an equivalence between left $\mathbf{A}_{\mathfrak{Qie}}$ -modules and analytic left L_{gr^0} -modules [21]. By applying this to Theorem 1.15, we also prove that a natural refinement of the eigenmonad adjunction (3) yields an equivalence of categories (see Corollary 12.3).

There is an *abelian* version of the adjunction (6) (see, for instance, [29, section 2.4]).

$$(7) \quad \tilde{L}_{\text{Fin}} \otimes_{L_{\mathfrak{E}}} (-) : L_{\mathfrak{E}} \text{Mod} \xleftarrow[\text{L}_{\mathfrak{Z}}]{\text{L}_{\mathfrak{Z}}} \text{Mod} : \text{Hom}_{L_{\mathfrak{Z}}}(\tilde{L}_{\text{Fin}}, -).$$

Here, Fin is the category of finite sets and maps; and \tilde{L}_{Fin} is a $(L_{\text{fr}_{\mathbb{Z}}^0}, L_{\mathfrak{E}})$ -bimodule defined by deforming the monad L_{Fin} . In this paper, we prove analogous statements of Theorem 1.15 by replacing gr^0 with $\text{fr}_{\mathbb{Z}}^0$. We go further by giving a generalization by considering general unital (not necessarily commutative) ring R instead of \mathbb{Z} . Let $\hat{R} := \mathbb{k} \otimes_{\mathbb{Z}} R$ be the coefficient extension of R to \mathbb{k} . For a subcategory \mathcal{D} of Fin , we define an $(\mathbb{N} \times \mathbb{N})$ -indexed \mathbb{k} -module $L_{\mathcal{D}}^{\hat{R}}$ by tensoring \hat{R} 's (see Definition 16.1 for details). Furthermore, if \mathcal{D} is \mathfrak{S} the category of bijections, then we can define a monad structure on $L_{\mathfrak{S}}^{\hat{R}}$ in $\text{Mat}_{\mathbb{k}}$ on \mathbb{N} .

Theorem 1.16. *Let $\mathcal{S}_{\text{fr}_{\mathbb{Z}}^0}^{\text{pr}}$ be the class of isomorphism classes of primitively generated $L_{\text{fr}_{\mathbb{Z}}^0}$ -modules.*

- (A) *The subclass $\mathcal{S}_{\text{fr}_{\mathbb{Z}}^0}^{\text{pr}} \subset L_{\text{fr}_{\mathbb{Z}}^0} \text{Mod}$ has a canonical $L_{\text{fr}_{\mathbb{Z}}^0}$ -internalizer $\mathbf{I}_{\text{fr}_{\mathbb{Z}}^0}^{\text{pr}}$.*
- (B) *For the left ideal $\mathbf{I}_{\text{fr}_{\mathbb{Z}}^0}^{\text{pr}}$, we have a monad isomorphism:*


$$E_{L_{\text{fr}_{\mathbb{Z}}^0}}(\mathbf{I}_{\text{fr}_{\mathbb{Z}}^0}^{\text{pr}}) \cong L_{\mathfrak{S}}^{\hat{R}}.$$

Based on this, the $(L_{\text{fr}_{\mathbb{Z}}^0}, E_{L_{\text{fr}_{\mathbb{Z}}^0}}(\mathbf{I}_{\text{fr}_{\mathbb{Z}}^0}^{\text{pr}}))$ -bimodule $L_{\text{fr}_{\mathbb{Z}}^0}/\mathbf{I}_{\text{fr}_{\mathbb{Z}}^0}^{\text{pr}}$ given by (1) of Proposition 1.1 is isomorphic to the $(L_{\text{fr}_{\mathbb{Z}}^0}, L_{\mathfrak{S}}^{\hat{R}})$ -bimodule $\tilde{L}_{\text{Fin}}^{\hat{R}}$.

The part (A) follows from Corollary 16.8 where $\mathbf{I}_{\text{fr}_{\mathbb{Z}}^0}^{\text{pr}}$ is defined in Definition 11.1. The part (B) is proved in Theorem 16.4.

²The definition is presented in a different form, as given in Definition 11.1, but it can be shown to be equivalent to the original by Proposition 11.17.

1.4.5. *Summary of the solutions.* In this section, we summarize the results in the preceding sections. We first show how Problem 1.8 is solved. Let T be a monad in $\text{Mat}_{\mathbb{k}}$. The following schematic diagram presents the main results in accordance with (5):



Let \mathcal{S} be a class of isomorphism classes of left T -modules. In the following tables 1 and 2, we present our findings regarding a T -internalizer J of \mathcal{S} , and computations of the quotient T/J and the eigenmonad $E_T(J)$.

\mathcal{S}	$\mathcal{S}_{\text{gr}^0}^d$	$\mathcal{S}_{\text{gr}^0}^\omega$	$\mathcal{S}_{\text{gr}^0}^{\text{pr}}$	\mathcal{S}^{out}	\mathcal{S}^{ab}
J	$\mathbb{I}_{\text{gr}^0}^{(d+1)}$	$\mathbb{I}_{\text{gr}^0}^\vee$	\mathbb{I}^{pr}	\mathbb{I}^{out}	\mathbb{I}^{ab}
T/J	$\mathcal{P}_{\text{gr}^0}^d$	$L_{\text{gr}^0}/\mathbb{I}_{\text{gr}^0}^\vee$	$\tilde{\mathbb{A}}_{\text{LSS}^t}$	H_0	$L_{\text{fr}_Z}^{\text{gr}^0}$
$E(J)$	$\mathcal{P}_{\text{gr}^0}^d$	$E(\mathbb{I}_{\text{gr}^0}^\vee)$	\mathbb{A}_{Eie}	H_0	$L_{\text{fr}_Z}^{\text{gr}^0}$

TABLE 1. Classes of modules, ideals and eigenmonads for $T = L_{\text{gr}^0}$

\mathcal{S}	$\mathcal{S}_{\text{fr}_R^0}^d$	$\mathcal{S}_{\text{fr}_R^0}^\omega$	$\mathcal{S}_{\text{fr}_R^0}^{\text{pr}}$
J	$\mathbb{I}_{\text{fr}_R^0}^{(d+1)}$	$\mathbb{I}_{\text{fr}_R^0}^\vee$	$\mathbb{I}_{\text{fr}_R^0}^{\text{pr}}$
T/J	$\mathcal{P}_{\text{fr}_R^0}^d$	$L_{\text{fr}_R^0}/\mathbb{I}_{\text{fr}_R^0}^\vee$	$\tilde{L}_{\text{Fin}}^{\hat{R}}$
$E(J)$	$\mathcal{P}_{\text{fr}_R^0}^d$	$E(\mathbb{I}_{\text{fr}_R^0}^\vee)$	$L_{\Xi}^{\hat{R}}$

TABLE 2. Classes of modules, ideals and eigenmonads for $T = L_{\text{fr}_R^0}$

The first column of each table is obtained from Theorem 1.11. The second column of each table follows from Theorem 1.13. The third column of table 1 is obtained from Theorem 1.15. The fourth column of table 2 is obtained from Theorem 1.16. The fourth column of table 1 appears in Theorem 1.14. The fifth column of table 1 is related to the abelianization

of groups and not presented in the preceding sections, since it is easy to deduce. This is described in section 14.

1.5. Further results. This section presents further results derived primarily from the solutions given in section 1.4.

1.5.1. Calculations of functor homology. This section presents an interpretation of the preceding results as the computation of the simplest homological invariants, which are analogous to the zeroth Ext rings.

We notify that (B) of Theorem 1.11 implies two structures on \mathcal{P}_C^d . One such structure is the monad structure in $\text{Mat}_{\mathbb{k}}$ on \mathbb{N} . The other one is L_C -module structure induced by the quotient homomorphism $L_C \rightarrow E_{L_C}(\mathbb{I}_C^{(d+1)})$, which arises since $\mathbb{I}_C^{(d+1)}$ is a two-sided ideal. We have similar structures on H_0 by Theorem 1.14.

Theorem 1.17. *We have the following calculations of the zeroth Ext monads:*

- (1) $\text{Ext}_{L_C}^0(\mathcal{P}_C^d, \mathcal{P}_C^d) \cong (\mathcal{P}_C^d)^{\circ}$ where C is a category with finite products and a zero object..
- (2) $\text{Ext}_{L_{\text{gr}^{\circ}}}^0(H_0, H_0) \cong H_0^{\circ}$.
- (3) $\text{Ext}_{L_{\text{gr}^{\circ}}}^0(\tilde{A}_{\mathbb{I}_{\text{ss}}^{\circ}}, \tilde{A}_{\mathbb{I}_{\text{ss}}^{\circ}}) \cong A_{\mathbb{I}_{\text{ss}}^{\circ}}^{\circ}$.
- (4) $\text{Ext}_{L_{\text{fr}_R^{\circ}}}^0(\tilde{L}_{\text{Fin}}^{\hat{R}}, \tilde{L}_{\text{Fin}}^{\hat{R}}) \cong (L_{\mathbb{E}}^{\hat{R}})^{\circ}$.

These are proved by applying (2) of Proposition 1.1 to (B) of Theorems 1.11, 1.14, 1.15 and 1.16 respectively.

1.5.2. Comparison of classes. The preceding sections present the respective studies of specific classes of modules. This paper also demonstrates a methodology for comparing those classes based on the associated *internalizers*, as shown in Lemma 6.11. For instance, let C be a category with finite products and a zero object. We derive a well-known result of polynomial degrees (see, for instance, [12]) that $\mathcal{S}_C^d \subset \mathcal{S}_C^{d+1}$ by comparing the internalizers in Theorem 1.11. Likewise, by using the internalizers $\mathbb{I}_C^v, \mathbb{I}_C^{\text{pr}}$ in Theorems 1.13, 1.15 and 1.16, we obtain the following:

Theorem 1.18. *Let $C \in \{\text{gr}^{\circ}, \text{fr}_R^{\circ}\}$. A primitively generated L_C -module is analytic:*

$$(8) \quad \mathcal{S}_C^{\text{pr}} \subset \mathcal{S}_C^{\omega}.$$

In fact, we have a stronger result that there exists $v_0 \in \Gamma(C)$ such that $\mathbb{I}_C^{v_0} \subset \mathbb{I}_C^{\text{pr}}$.

The first statement is proved in Corollaries 12.2 and 17.4 respectively. The last statement appears in Theorems 12.1 and 17.3.


Remark 1.19. In the case that $C = \text{gr}^{\circ}$, the first statement was already proved in [12].

By combining the results in the literature [21], we can go further: if \mathbb{k} is a field of characteristic zero, then we have $\mathcal{S}_C^{\text{pr}} = \mathcal{S}_C^{\omega}$ (see Corollary 12.3 and Proposition 17.5).

Remark 1.20. Note that Theorem 1.18 extends the classical fact that primitively generated cocommutative Hopf algebras are conilpotent under the equivalence between exponential modules over gr° (modules over $\text{fr}_{\mathbb{Z}}^{\circ}$, resp.) and cocommutative (bicocommutative, resp.) Hopf algebras [20, 12]. In fact, an exponential module over C is analytic (primitively generated, resp.) if and only if the corresponding Hopf algebra is conilpotent (primitively generated, resp.) [12].

1.5.3. *Subquotient preorder.* This section illustrates a consequence of section 1.4. In fact, the calculations of eigenmonads lead to a new relation between the monads of interest. This should be considered as a relation between the \mathbb{k} -linear categories corresponding to the monads, as we mentioned in section 1.1. Let $\mathcal{X} \in \text{Mat}_{\mathbb{k}}$ be an object. For monads T, S in $\text{Mat}_{\mathbb{k}}$ on \mathcal{X} , we define a preorder relation \gg by $T \gg S$ if S is isomorphic to a quotient monad of a submonad of T . By the definition, for a multiplicative submodule $J \subset T$, we have $T \gg E_T(J)$. In (9), for $\mathcal{X} = \mathbb{N}$, we present a *genealogy* of monads, which implies the complexity of the monads:

Notation 1.21. If $T \gg S$, then the former is positioned higher than the latter, with the two entities being linked by a line.



The part (B) of Theorem 1.11 implies the relation $\mathcal{P}_C^{d+1} \gg \mathcal{P}_C^d$ by Example 5.9 and Proposition 7.10; and the relation $\mathcal{P}_{gr^o}^d \gg \mathcal{P}_{fr^o_Z}^d$ with the isomorphisms for $d = 0, 1$ by Theorem 15.10. The part (B) of Theorem 1.14 implies the relation $L_{gr^o} \gg H_0$; and the relation $H_0 \gg L_{fr^o_Z}$ by Corollary 14.10. The part (B) of Theorem 1.15 implies the relation $L_{gr^o} \gg A_{Qie}$. The part (B) of Theorem 1.16 implies the relation $L_{fr^o_Z} \gg L_{\emptyset}$; and the relation $A_{Qie} \gg L_{\emptyset}$, as described in Example 16.6.

1.6. **Organization.** This paper is organized as follows.

In part 1, we present a categorification of eigenrings and related constructions. In section 3, we introduce the bicategory $\text{Mat}_{\mathbb{k}}$ in which we formulate the categorification. In section 4, we study monad theory in the bicategory $\text{Mat}_{\mathbb{k}}$. In section 5, we introduce the notion of an eigenmonad. In section 6, we construct a correspondence between left ideals and classes of modules, which is closely related to Problem 1.8. In section 7, we study polynomial functors based on our framework.

In part 2, we present applications of the general theory of part 1 to the opposite category of free groups. In section 8, we explain the notation for part 2. In section 9, we give an application of the results presented in section 7 to free groups. In section 10, we study outer

functors on free groups based on our framework. In section 11, we introduce and study primitively generated L_{gr° -modules. In section 12, we present a comparison of analytic L_{gr° -modules and primitively generated L_{gr° -modules.

In part 3, we present applications of the general theory of part 1 to the opposite category of free modules over a ring. In section 13, we explain the notation for part 3. In section 14, we study the abelianization of free groups based on our setting. In section 15, we investigate polynomial $L_{\text{fr}_R^\circ}$ -modules by applying the results presented in section 7. In section 16, we study primitively generated $L_{\text{fr}_R^\circ}$ -modules. In section 17, we present a comparison of analytic $L_{\text{fr}_R^\circ}$ -modules and primitively generated $L_{\text{fr}_R^\circ}$ -modules.

In appendix A, we prove that the category of T-modules has enough projectives and injectives. In appendix B, we investigate fundamental properties of graded coaugmented coalgebras. In appendix C, we study a grading on free Lie algebras, which is applied to compute an eigenmonad in section 11.3.

ACKNOWLEDGEMENTS

The author is supported by a KIAS Individual Grant MG093701 at Korea Institute for Advanced Study. The author would like to express his sincere gratitude to Geoffrey Powell and Christine Vespa for carefully reading the earlier versions of this paper and providing him constructive comments. The author also thanks many people he encountered at the International Category Theory Conference CT2024. In particular, the author appreciates Rui Prezado and Sori Lee who demonstrated remarkable patience in listening to his presentation. He acknowledges that the reference for the bicategory of matrices [27] is due to Rui Prezado. The author appreciates Sori Lee for his advice on the formulation of Problem 1.8. Finally, the author wish to thank Habiro Kazuo, Katada Mai for the valuable discussion on this subject.

2. NOTATION

Throughout this paper, we make use of the following notation.

- \mathbb{k} : a commutative unital ring with the unit $1_{\mathbb{k}} \in \mathbb{k}$.
- \otimes : the tensor product over \mathbb{k} , or the composition of morphisms in $\text{Mat}_{\mathbb{k}}$.
- \mathbb{N} : the set of non-negative integers.
- \mathfrak{S}_n : the n -th symmetric group.
- \underline{n} : the set $\{1, 2, \dots, n\}$.

Part 1. General eigenring construction

In this part, we give a categorification of eigenrings and related constructions by replacing algebras with linear categories. For the convenience, we consider a bicategory $\text{Mat}_{\mathbb{k}}$ whose monads are equivalent with \mathbb{k} -linear categories. The categorified eigenring is called the eigenmonad which is induced by a monad T with a multiplicative submodule $J \subset T$. An adjunction of T -modules and modules over the eigenmonad naturally arises. Based on the adjunction, we give a correspondence (see (4)) between left ideals of T and classes of T -modules, which makes Problem 1.8 clear. Furthermore, we revisit the polynomial functor theory in section 7 where we encode the polynomial property into a left ideal.

3. THE BICATEGORY OF MATRICES $\text{Mat}_{\mathbb{k}}$

In this section, we introduce a bicategory in which we develop a categorification of eigenrings.

First, we review some concepts related to indexed \mathbb{k} -modules. For a set \mathcal{Z} , a \mathcal{Z} -indexed (\mathbb{k} -)module is a family of \mathbb{k} -modules $F = \{F(Z)\}_{Z \in \mathcal{Z}}$ indexed by the set \mathcal{Z} . A \mathcal{Z} -homomorphism

$\xi : F' \rightarrow F$ is a family of \mathbb{k} -linear maps $\xi_Z : F'(Z) \rightarrow F(Z)$ parametrized by $Z \in \mathcal{Z}$. We sometimes write this as simply $\xi : F' \rightarrow F$. It is obvious that the category of \mathcal{Z} -indexed modules and their homomorphisms constitutes an abelian category, with the (co)kernel objects calculated indexwise. Furthermore, most concepts associated with \mathbb{k} -modules can be readily extended to indexed modules:

- A *submodule* of F is an \mathcal{Z} -indexed \mathbb{k} -module G such that $G(Z)$ is a submodule of $F(Z)$ for $Z \in \mathcal{Z}$. we denote by $G \subset F$ when so it is.
- A *quotient module* F/G is defined to be a \mathcal{Z} -indexed module such that $(F/G)(Z) := F(Z)/G(Z)$.
- For a homomorphism $\xi : F' \rightarrow F$ and a submodule $G \subset F$, we define the *preimage* $\xi^{-1}(G)$ to be the \mathcal{Z} -indexed module such that $(\xi^{-1}(G))(Z) := \xi_Z^{-1}(G(Z))$.
- A homomorphism $\xi : F' \rightarrow F$ is an *epimorphism* (a *monomorphism*, resp.) if each map $\xi_Z : F'(Z) \rightarrow F(Z)$ is surjective (injective, resp.).

In this paper, we freely make use of these concepts and symbols.

Definition 3.1. We define a bicategory $\text{Mat}_{\mathbb{k}}$ by the following data:

- Its *objects* are sets. We use symbols $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$ to denote its objects.
- For objects \mathcal{X}, \mathcal{Y} , a *morphism* from \mathcal{X} to \mathcal{Y} is given by an $(\mathcal{Y} \times \mathcal{X})$ -indexed \mathbb{k} -module. For such a morphism $F = \{F(Y, X)\}_{(Y, X) \in \mathcal{Y} \times \mathcal{X}}$, we denote this by $F : \mathcal{X} \rightsquigarrow \mathcal{Y}$. For morphisms $F : \mathcal{X} \rightsquigarrow \mathcal{Y}$ and $G : \mathcal{Y} \rightsquigarrow \mathcal{Z}$, their *composition* $G \otimes F : \mathcal{X} \rightsquigarrow \mathcal{Z}$ is defined by

$$(G \otimes F)(Z, X) := \bigoplus_{Y \in \mathcal{Y}} G(Z, Y) \otimes F(Y, X), \quad (Z, X) \in \mathcal{Z} \times \mathcal{X}.$$

- For each object \mathcal{X} , the *identity morphism* $\mathbb{1}_{\mathcal{X}} : \mathcal{X} \rightsquigarrow \mathcal{X}$ on \mathcal{X} is defined to be

$$\mathbb{1}_{\mathcal{X}}(Y, X) := \begin{cases} \mathbb{k}, & (Y = X), \\ 0, & (Y \neq X). \end{cases}$$

- For morphisms F and G from \mathcal{X} to \mathcal{Y} , a *2-morphism* Φ from F to G , denoted as $\Phi : F \rightarrow G$, is defined to be a $(\mathcal{Y} \times \mathcal{X})$ -homomorphism between $(\mathcal{Y} \times \mathcal{X})$ -indexed \mathbb{k} -modules, i.e. a family of \mathbb{k} -linear maps $\Phi_{(Y, X)} : F(Y, X) \rightarrow G(Y, X)$ indexed by $(Y, X) \in \mathcal{Y} \times \mathcal{X}$. Denote by id_F the *identity 2-morphism* on F . Denote by $\text{hom}_{\mathcal{Y}, \mathcal{X}}(F, G)$ the set of 2-morphisms from F to G .

Remark 3.2. It is recommended that the reader thinks of a morphism $F : \mathcal{X} \rightsquigarrow \mathcal{Y}$ as a matrix whose entries consist of \mathbb{k} -modules; and whose columns and rows are parameterized by \mathcal{X} and \mathcal{Y} respectively. From this perspective, the composition of morphisms is consistent with the matrix product, and the identity morphism $\mathbb{1}_{\mathcal{X}}$ is the diagonal matrix. In fact, $\text{Mat}_{\mathbb{k}}$ is a special case of the *bicategory of \mathbb{W} -matrices* [27, section 1] where \mathbb{W} is the bicategory with one object, \mathbb{k} -modules as morphisms and \mathbb{k} -linear maps as 2-morphisms.

In this paper, we mainly use the terminology *homomorphism* (*isomorphism*, resp.) of indexed modules instead of 2-morphisms (2-isomorphism, resp.) in $\text{Mat}_{\mathbb{k}}$.

Notation 3.3. For a morphism $F : \mathcal{X} \rightsquigarrow \mathcal{Y}$, the *opposite* of F is denoted by $F^{\circ} : \mathcal{Y} \rightsquigarrow \mathcal{X}$ such that $F^{\circ}(X, Y) := F(Y, X)$ for $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$.

This can be interpreted as the transposition of matrices. We have $\mathbb{1}_{\mathcal{X}}^{\circ} = \mathbb{1}_{\mathcal{X}}$, and an obvious natural isomorphism $(G \otimes F)^{\circ} \cong F^{\circ} \otimes G^{\circ}$.

Definition 3.4 (Inner Hom). Let $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$ be objects of $\text{Mat}_{\mathbb{k}}$. For a morphism $E : \mathcal{Y} \rightsquigarrow \mathcal{Z}$, we define $\text{Hom}_{\mathcal{Z}}^{\text{L}}(E, -)$ to be a right adjoint to $E \otimes (-)$: for an object \mathcal{X} , morphisms

$G : \mathcal{X} \rightsquigarrow \mathcal{Z}$ and $F : \mathcal{X} \rightsquigarrow \mathcal{Y}$, the morphism $\text{Hom}^L_{\mathcal{Z}}(E, G) : \mathcal{X} \rightsquigarrow \mathcal{Y}$ satisfies a natural isomorphism:

$$\text{hom}_{\mathcal{Z}, \mathcal{X}}(E \otimes F, G) \cong \text{hom}_{\mathcal{Y}, \mathcal{X}}(F, \text{Hom}^L_{\mathcal{Z}}(E, G)).$$

In fact, we have an explicit description that $(\text{Hom}^L_{\mathcal{Z}}(E, G))(Y, X)$ consists of \mathcal{Z} -homomorphisms $E(-, Y) \rightarrow G(-, X)$ for $Y \in \mathcal{Y}, X \in \mathcal{X}$. Analogously, we introduce $\text{Hom}^R_{\mathcal{X}}(F, -)$ as a right adjoint to $(-) \otimes F$:

$$\text{hom}_{\mathcal{Z}, \mathcal{X}}(E \otimes F, G) \cong \text{hom}_{\mathcal{Z}, \mathcal{Y}}(E, \text{Hom}^R_{\mathcal{X}}(F, G)).$$

In this case, $(\text{Hom}^R_{\mathcal{X}}(F, G))(Z, Y)$ is a \mathbb{k} -module generated by \mathcal{X} -homomorphisms from $F(Y, -)$ to $G(Z, -)$.

The reader can observe that $\text{Hom}^L_{\mathcal{X}}$ and $\text{Hom}^R_{\mathcal{X}}$ serve as a realization of the assignment $\text{hom}_{\mathcal{Y} \times \mathcal{X}}$ (see Definition 3.1) as indexed modules, so-called an *internal hom*. Let $\mathcal{X}, \mathcal{Y}_1, \mathcal{Y}_2, \mathcal{Y}_3$ be objects. For morphisms $F_i : \mathcal{Y}_i \rightsquigarrow \mathcal{X}$, $i \in \{1, 2, 3\}$, the composition of homomorphisms induce the following associative pairing:

$$(10) \quad \text{Hom}^L_{\mathcal{X}}(F_1, F_2) \otimes \text{Hom}^L_{\mathcal{X}}(F_2, F_3) \rightarrow \text{Hom}^L_{\mathcal{X}}(F_1, F_3).$$

By a careful treatment of the order of components, we also have an associative pairing for $\text{Hom}^R_{\mathcal{X}}$. Let us consider $G_i : \mathcal{X} \rightsquigarrow \mathcal{Y}_i$, $i \in \{1, 2, 3\}$:

$$\text{Hom}^R_{\mathcal{X}}(G_2, G_3) \otimes \text{Hom}^R_{\mathcal{X}}(G_1, G_2) \rightarrow \text{Hom}^R_{\mathcal{X}}(G_1, G_3).$$

4. MONAD THEORY FOR BI-INDEXED MODULES

The objective of this section is to present a toolkit for the description of our results. We investigate monad theory within the bicategory $\text{Mat}_{\mathbb{k}}$. It will be shown that monads in that bicategory are equivalent with \mathbb{k} -linear categories; and modules over a monad are equivalent with modules over the corresponding \mathbb{k} -linear category (for example, in the context of [21, 29, 12]). This allows us to treat \mathbb{k} -linear categories and modules over them in a manner consistent with in the classical ring-module theory. Moreover, we show that the category of modules over a monad allows homological algebra.

4.1. Review of monads. The notion of a monad in a bicategory is a generalization of that of an algebra. It is useful to treat monads when we categorify eigenrings and the related constructions. In this section, we give a brief review of some notions in monad theory [1].

Fix a bicategory \mathbb{B} . For consistency with the preceding section, we denote by $f : x \rightsquigarrow y$ a morphism f from x to y . Let $g : y \rightsquigarrow z$ be another morphism. We also make use of the symbol \otimes for the composition $g \otimes f : x \rightsquigarrow z$ of the morphisms f, g . For an object x , let $\mathbb{1}_x$ denote the identity morphism on x . For a morphism f , we denote by id_f the identity 2-morphism on f .

A monad T in \mathbb{B} on an object x consists of 2-morphisms $\nabla : T \otimes T \rightarrow T$ and $\eta : \mathbb{1}_x \rightarrow T$, for which the following diagrams are commutative:

$$\begin{array}{ccc} T \otimes T \otimes T & \xrightarrow{\text{id}_T \otimes \nabla} & T \otimes T \\ \downarrow \nabla \otimes \text{id}_T & & \downarrow \nabla \\ T \otimes T & \xrightarrow{\nabla} & T \end{array} \quad \begin{array}{ccc} T \otimes \mathbb{1}_x & \xrightarrow{\cong} & T \longleftarrow \cong \mathbb{1}_x \otimes T \\ \downarrow \text{id}_T \otimes \eta & & \parallel \quad \downarrow \eta \otimes \text{id}_T \\ T \otimes T & \xrightarrow{\nabla} & T \longleftarrow \nabla T \otimes T \end{array}$$

Remark 4.1. Note that one should read these diagrams by properly inserting the associator of \mathbb{B} , i.e. the natural isomorphism for associativity.

Let T be a monad in \mathbb{B} on an object x . For an object y of \mathbb{B} , a *left T -module* from y is a morphism $M : y \rightsquigarrow x$ with a 2-morphism $\triangleright : T \otimes M \rightarrow M$ for which the diagrams below commutes:

$$\begin{array}{ccc} T \otimes T \otimes M & \xrightarrow{\text{id}_T \otimes \triangleright} & T \otimes M \\ \downarrow \nabla \otimes \text{id}_M & & \downarrow \triangleright \\ T \otimes M & \xrightarrow{\triangleright} & M \end{array} \quad \begin{array}{ccc} M & \xleftarrow{\cong} & \mathbb{1}_x \otimes M \\ \parallel & & \downarrow \eta \otimes \text{id}_M \\ M & \xleftarrow{\triangleright} & T \otimes M \end{array}$$

A *right T -module* to y is defined to be a morphism $M : x \rightsquigarrow y$ with a 2-morphism $\triangleleft : M \otimes T \rightarrow M$ satisfying similar axioms. For left (or right) T -modules M, M' , a *T -homomorphism* $\xi : M \rightarrow M'$ is defined to be a 2-morphism compatible with the T -actions.

Consider monads T, S in \mathbb{B} on x and y respectively. A *(T, S) -bimodule* is a morphism $M : y \rightsquigarrow x$ endowed with 2-morphisms $\triangleright : T \otimes M \rightarrow M$ and $\triangleleft : M \otimes S \rightarrow M$. It is required that \triangleright gives a left T -module, \triangleleft gives a right S -module and the following diagram commutes:

$$\begin{array}{ccc} T \otimes M \otimes S & \xrightarrow{\triangleright \otimes \text{id}_S} & M \otimes S \\ \downarrow \text{id}_T \otimes \triangleleft & & \downarrow \triangleleft \\ T \otimes M & \xrightarrow{\triangleright} & M \end{array}$$

For (T, S) -bimodules M, M' , a *(T, S) -homomorphism* $\xi : M \rightarrow M'$ is a 2-morphism which is compatible with the actions of the monads T and S .

4.2. Monads in $\text{Mat}_{\mathbb{k}}$. This section presents preliminaries for the study of monads in $\text{Mat}_{\mathbb{k}}$. We first give an equivalent description of monads in the bicategory $\text{Mat}_{\mathbb{k}}$. Let \mathcal{X} be an object of $\text{Mat}_{\mathbb{k}}$ and T be a monad in $\text{Mat}_{\mathbb{k}}$ on \mathcal{X} .

Notation 4.2. We often use the *composition* notation for the monad operation:

$$f \circ g := \nabla(f \otimes g),$$

for $X, Y, Z \in \mathcal{X}$, $f \in T(Z, Y)$ and $g \in T(Y, X)$. Here, we regard $f \otimes g \in T(Z, Y) \otimes T(Y, X) \subset (T \otimes T)(Z, X)$. For $X \in \mathcal{X}$, we also denote by $1_X \in T(X, X)$ the element corresponding to the unit $1_{\mathbb{k}} \in \mathbb{k}$ by the monad unit $\eta : 1_{\mathcal{X}} \rightarrow T$.

One can realize that a monad on \mathcal{X} gives rise to a \mathbb{k} -linear category whose composition is given by the operation \circ in Notation 4.2; and whose object set is \mathcal{X} . This assignment is invertible in the following sense. Denote by $\text{Mnd}^{\mathcal{X}}$ the category of monads in $\text{Mat}_{\mathbb{k}}$ on \mathcal{X} and monad homomorphisms. Denote by $\text{Lin}_{\mathbb{k}}^{\mathcal{X}}$ the category of \mathbb{k} -linear categories with object set \mathcal{X} and \mathbb{k} -linear functors. Then we have an equivalence of categories [27, section 2]:

$$(11) \quad \text{Lin}_{\mathbb{k}}^{\mathcal{X}} \rightarrow \text{Mnd}^{\mathcal{X}}.$$

In this paper, we identify \mathbb{k} -linear categories with monads in $\text{Mat}_{\mathbb{k}}$ based on the equivalence (11).

In the followings, we present examples of monads in $\text{Mat}_{\mathbb{k}}$.

Example 4.3. Let $\mathcal{X} = *$ be the one-point set. The category $\text{Lin}_{\mathbb{k}}^{\mathcal{X}}$ is nothing but the category of \mathbb{k} -algebras and algebra homomorphisms.

Example 4.4. For objects \mathcal{X}, \mathcal{Y} and a morphism $F : \mathcal{Y} \rightsquigarrow \mathcal{X}$, the pairing (10) induces a monad structure on $\text{Hom}_{\mathcal{X}}^l(F, F) : \mathcal{Y} \rightsquigarrow \mathcal{Y}$. Similarly, $\text{Hom}_{\mathcal{Y}}^r(F, F)$ is a monad on \mathcal{X} .

Example 4.5. Let \mathcal{X} be a set and \mathcal{C} be a category with objects \mathcal{X} . The \mathbb{k} -linearization of categories gives a linear category $\mathbb{k}\mathcal{C}$ with objects \mathcal{X} . Denote by $L_{\mathcal{C}}$ the monad in $\text{Mat}_{\mathbb{k}}$ on \mathcal{X} induced by the equivalence (11):

$$L_{\mathcal{C}}(Y, X) := \mathbb{k}\mathcal{C}(X, Y), \quad X, Y \in \mathcal{X}.$$

Here, $C(X, Y)$ is the set of morphisms from X to Y . By definitions, we have an isomorphism of monads $L_{C^\circ} \cong L_C^\circ$.

Example 4.6. Let \mathcal{O} be a \mathbb{k} -linear (symmetric) operad. The assignment of $\mathcal{O}(n)$ to $n \in \mathbb{N}$ naturally extends to finite sets: for a finite set X with order n , we define

$$\mathcal{O}(X) := \mathbb{k}\text{Bij}(\underline{n}, X) \otimes_{\mathfrak{S}_n} \mathcal{O}(n).$$

Here, $\text{Bij}(\underline{n}, X)$ is the set of bijections from $\underline{n} := \{1, 2, \dots, n\}$ to X ; and the \mathfrak{S}_n -action on it is defined by the precomposition. Denote by $\text{Cat}_{\mathcal{O}}$ the linear category associated with \mathcal{O} . It is a \mathbb{k} -linear category with objects \mathbb{N} ; and the \mathbb{k} -module formed by morphisms is given by

$$\text{Cat}_{\mathcal{O}}(n, m) := \bigoplus_{f \in \text{Fin}(n, m)} \bigotimes_{i=1}^m \mathcal{O}(f^{-1}(i)).$$

We denote by $\mathbf{A}_{\mathcal{O}}$ the monad in $\text{Mat}_{\mathbb{k}}$ on \mathbb{N} induced by the equivalence (11). Especially, we have $\mathbf{A}_{\mathcal{O}}(m, n) = \text{Cat}_{\mathcal{O}}(n, m)$.

Before we close this section, we define a basic relation of monads in $\text{Mat}_{\mathbb{k}}$. These allow us to illustrate our computation of eigenmonads, as shown in section 1.5.3.

Definition 4.7 (Subquotient preorder). Let \mathcal{X} be an object in $\text{Mat}_{\mathbb{k}}$. For monads T, S in $\text{Mat}_{\mathbb{k}}$ on \mathcal{X} , we say that S is a *submonad* (a *quotient monad*, resp.) of T if there exists a monad monomorphism (monad epimorphism, resp.) $S \rightarrow T$. Moreover, S is a *subquotient monad* of T if S is a quotient monad of a submonad of T . We define a relation $T \gg S$ if S is a subquotient monad of T . In Proposition 4.9, we prove that the relation \gg is a preorder.

Remark 4.8. The relation \gg obviously depends on \mathcal{X} , which we omit for the sake of simplicity.

In the following, the notation of *preimage* ξ^{-1} is explained in section 4.2.

Proposition 4.9. *The relation \gg gives a preorder of monads in $\text{Mat}_{\mathbb{k}}$ on \mathcal{X} .*

Proof. The reflexivity is validated since T is a subquotient of itself. Hence, it suffices to prove the transitivity. Suppose that $T \gg S \gg R$ are monads on \mathcal{X} . Let T' (S' , resp.) be a submonad of T (S , resp.) with a monad epimorphism $\xi : T' \rightarrow S$ ($\pi : S' \rightarrow R$, reps.). Here, we regard T' and S' as a submodule of T and S by using the images of the associated monomorphisms. Then $\xi^{-1}(S')$ is a submonad of T , and $\xi^{-1}(S') \xrightarrow{\xi} S' \xrightarrow{\pi} R$ is a monad epimorphism. Thus, we obtain $T \gg R$. \square

4.3. Modules in $\text{Mat}_{\mathbb{k}}$. In this section, we study modules over monads in the bicategory $\text{Mat}_{\mathbb{k}}$. In particular, we present a fact that T -modules in $\text{Mat}_{\mathbb{k}}$ are equivalent with linear functors from the category corresponding to T to the category of \mathbb{k} -modules.

Definition 4.10. Denote by ${}_T\text{Mod}_S$ the category of (T, S) -bimodules and (T, S) -homomorphisms. For objects M, N , denote by $\text{hom}_{T, S}(M, N)$ the \mathbb{k} -module generated by (T, S) -homomorphisms from M to N .

The category ${}_T\text{Mod}_S$ is an abelian category. In particular, (co)kernel objects of (T, S) -homomorphisms consist of (co)kernel modules index-wise.

Note that the opposite of morphisms in $\text{Mat}_{\mathbb{k}}$ induces an equivalence between the categories ${}_T\text{Mod}_S$ and ${}_S\text{Mod}_T$.

Let \mathcal{A}, \mathcal{B} be \mathbb{k} -linear categories with objects \mathcal{Y} and \mathcal{X} respectively. We set T and S to be the corresponding monads on \mathcal{Y} and \mathcal{X} respectively by the equivalence (11). Denote by ${}_{\mathcal{A}}\mathcal{F}_{\mathcal{B}}$

the category of \mathbb{k} -bilinear functors $\mathcal{B}^0 \times \mathcal{A} \rightarrow \mathbb{k}\text{Mod}$ and natural transformations between them. Then we have an equivalence of categories [27, section 3]:

$$(12) \quad \mathcal{A}\mathcal{F}_{\mathcal{B}} \rightarrow {}_T\text{Mod}_S.$$

The equivalence (12) assigns a (T, S) -bimodule $\mathbb{M} : \mathcal{X} \rightsquigarrow \mathcal{Y}$ such that $\mathbb{M}(Y, X) := L(X, Y)$ for $X \in \mathcal{X}, Y \in \mathcal{Y}$ to a \mathbb{k} -bilinear functor $L : \mathcal{B}^0 \times \mathcal{A} \rightarrow \mathbb{k}\text{Mod}$. Under this point of view, we identify \mathbb{k} -bilinear functors to $\mathbb{k}\text{Mod}$ with bimodules in $\text{Mat}_{\mathbb{k}}$. It is convenient to consider monads and modules over them when we develop the categorification in the next section.

Definition 4.11. Let $*$ be a one-point set. Consider an object X of $\text{Mat}_{\mathbb{k}}$ and a monad T on \mathcal{X} . In this paper, a *left T -module* is defined to be a left T -module from $*$ in the sense of section 4.1. Denote by ${}_T\text{Mod}$ the category of left T -modules and T -homomorphisms. Similarly, a *right T -module* is defined to be a right T -module to $*$. Denote by Mod_T the category of right T -modules and T -homomorphisms.

Notation 4.12. By the definition, a left T -module \mathbb{M} consists of a \mathbb{k} -module $T(X, *)$ parametrized by $X \in \mathcal{X}$. In this paper, we use the following simplified notation:

$$\mathbb{M}(X) := \mathbb{M}(X, *).$$

Remark 4.13. Recall the endomorphism monads in Example 4.4. A left T -module from \mathcal{Y} is nothing but a morphism $\mathbb{M} : \mathcal{Y} \rightsquigarrow \mathcal{X}$ endowed with a monad homomorphism $T \rightarrow \text{Hom}_{\mathcal{Y}}^R(\mathbb{M}, \mathbb{M})$. Hence, a left T -module \mathbb{M} is a family $\mathbb{M}(X)$, $X \in \mathcal{X}$ with a monad homomorphism $T \rightarrow \text{Hom}_{*}^R(\mathbb{M}, \mathbb{M})$.

Example 4.14. The assignment of the opposites (see Notation 3.3) gives an equivalence of categories ${}_T\text{Mod} \simeq \text{Mod}_{T^{\circ}}$.

Example 4.15. Let T be a monad in $\text{Mat}_{\mathbb{k}}$ and \mathcal{A} be the \mathbb{k} -linear category corresponding to T by (11). By the equivalence (12), one may see that the category ${}_T\text{Mod}$ is equivalent with the category of \mathbb{k} -linear functors $\mathcal{A} \rightarrow \mathbb{k}\text{Mod}$.

Example 4.16. Recall Example 4.5. By the above observation, left L_C -modules are equivalent to \mathbb{k} -linear functors $\mathbb{k}C \rightarrow \mathbb{k}\text{Mod}$. The \mathbb{k} -linearization gives a one-to-one correspondence between \mathbb{k} -linear functors $\mathbb{k}C \rightarrow \mathbb{k}\text{Mod}$ and functors $C \rightarrow \mathbb{k}\text{Mod}$. Let $\mathcal{F}(C)$ be the category of functors from C to $\mathbb{k}\text{Mod}$ and natural transformations. The previous correspondences induce an isomorphism of categories:

$$\mathcal{F}(C) \cong {}_{L_C}\text{Mod}.$$

4.4. Hom and Tensor. This section gives some formal constructions of modules which generalize hom functor and tensor product of modules. The former induces a monad generated by T -endomorphisms on a T -module for monad T . Moreover, we introduce the associated derived functors, which generalize Ext and Tor functors.

Definition 4.17 (Balanced Tensor product). Let $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$ be objects and S, E, T be monads in $\text{Mat}_{\mathbb{k}}$ on each object. Consider a (T, E) -bimodule $\mathbb{M} : \mathcal{Y} \rightsquigarrow \mathcal{Z}$ and an (E, S) -bimodule $\mathbb{N} : \mathcal{X} \rightsquigarrow \mathcal{Y}$. We define the morphism $\mathbb{M} \otimes_E \mathbb{N} : \mathcal{X} \rightsquigarrow \mathcal{Z}$ in $\text{Mat}_{\mathbb{k}}$ to be the coequalizer (as an $(\mathcal{X} \times \mathcal{Z})$ -indexed module) of

$$\mathbb{M} \otimes E \otimes \mathbb{N} \begin{array}{c} \xrightarrow{\triangleright \otimes \text{id}_{\mathbb{N}}} \\ \xrightarrow{\text{id}_{\mathbb{M}} \otimes \triangleleft} \end{array} \mathbb{M} \otimes \mathbb{N}$$

Here, \triangleleft is the right E -action on \mathbb{M} and \triangleright is the left E -action on \mathbb{N} . We endow $\mathbb{M} \otimes_E \mathbb{N}$ with the (T, S) -bimodule structure inherited from \mathbb{M} and \mathbb{N} .

Definition 4.18 (Inner Hom). Let \mathcal{X}, \mathcal{Y} be objects of $\text{Mat}_{\mathbb{k}}$ and T, S be monads on each object. Recall the notation in Definition 4.10. For a (T, S) -bimodule M , we define $\text{Hom}^L_T(M, -)$ to be a right adjoint to $M \otimes_S (-)$: for an object Z with a monad U on itself, a (S, U) -bimodule N and a (T, U) -bimodule P , the (S, U) -bimodule $\text{Hom}^L_T(M, P)$ satisfies a natural isomorphism:

$$(13) \quad \text{hom}_{T,U}(M \otimes_S N, P) \cong \text{hom}_{S,U}(N, \text{Hom}^L_T(M, P)).$$

The reader is referred to Notation 4.20 for further notation. Analogously, we define the assignment $\text{Hom}^R_U(N, -)$ to be characterized by

$$\text{hom}_{T,U}(M \otimes_S N, P) \cong \text{hom}_{T,S}(M, \text{Hom}^R_U(N, P)).$$

These have a concrete description. For instance, for $Y \in \mathcal{Y}, Z \in \mathcal{Z}$, the \mathbb{k} -module $(\text{Hom}^L_T(M, P))(Y, Z)$ is generated by T -homomorphisms from $M(-, Y)$ to $P(-, Z)$. Equivalently, by using $\text{Hom}^L_{\mathcal{X}}$ in Definition 3.4, $\text{Hom}^L_T(M, P)$ is a submodule of $\text{Hom}^L_{\mathcal{X}}(M, P)$ consisting of \mathcal{X} -homomorphisms commuting with the T -actions.

Example 4.19. We have $\text{Hom}^{\star}_{\mathbb{1}_{\mathcal{X}}} = \text{Hom}^{\star}_{\mathcal{X}}$ for $\star \in \{L, R\}$ where the latter is explained in Definition 3.4.

Notation 4.20. In this paper, we are mainly interested in *left* T -modules, so that we use the following notation, unless specified otherwise:

$$\text{Hom}_T(M, P) := \text{Hom}^L_T(M, P).$$

By using Notation 3.3, it is clear that $\text{Hom}^R_U(N, P) = \text{Hom}_{U^{\circ}}(N^{\circ}, P^{\circ})^{\circ}$.

Let T be a monad in $\text{Mat}_{\mathbb{k}}$ on an object \mathcal{X} . For objects $\mathcal{Y}_1, \mathcal{Y}_2, \mathcal{Y}_3$ of $\text{Mat}_{\mathbb{k}}$, consider a left T -module M_i from \mathcal{Y}_i for $i \in \{1, 2, 3\}$. Then the pairing (10) gives the following homomorphism:

$$(14) \quad \text{Hom}_T(M_1, M_2) \otimes \text{Hom}_T(M_2, M_3) \rightarrow \text{Hom}_T(M_1, M_3).$$

It is merely a collection of the compositions of T -homomorphisms. This gives a monad consisting of T -endomorphisms:

Definition 4.21. Let M be a left T -module from an object \mathcal{Y} of $\text{Mat}_{\mathbb{k}}$. In this paper, we regard $\text{Hom}_T(M, M)$ as a monad in $\text{Mat}_{\mathbb{k}}$ on \mathcal{Y} whose monad operation is induced by the pairing (14). The unit $\eta : \mathbb{1}_{\mathcal{Y}} \rightarrow \text{Hom}_T(M, M)$ is given by the assignment of the identities on $M(Y)$ for each $Y \in \mathcal{Y}$. We call this the *T-endomorphism monad on M*.

Before we close this section, we show that a T -endomorphism monad is a part of a homological invariant. It is well-known that the category of modules over a ring has enough injectives and projectives (see, for example, [30]). More generally, for a monad T in $\text{Mat}_{\mathbb{k}}$, the category ${}_T\text{Mod}$ is an abelian category with enough injectives and projectives (see appendix A for the proof). In consequence, we obtain the following homological invariants:

Definition 4.22. Let T, S be monads on objects $\mathcal{X}, \mathcal{Y} \in \text{Mat}_{\mathbb{k}}$ respectively, and M be a (T, S) -module. We introduce Ext modules $\text{Ext}^{\bullet}_T(M, -)$ as the right derived functors of $\text{Hom}_T(M, -) : {}_T\text{Mod} \rightarrow {}_S\text{Mod}$ (see Notation 4.20).

Remark 4.23. In a symmetric fashion, we can define Tor modules $\text{Tor}^T_{\bullet}(-, M)$ to be the left derived functors of $(-) \otimes_T M : \text{Mod}_T \rightarrow \text{Mod}_S$, which are beyond our interest of this paper.

Remark 4.24. These can be regarded as a functor homology (see, for example, [28]) under the identification (12).

Example 4.25. By the definitions, we have $\text{Ext}^0_T(M, -) \cong \text{Hom}_T(M, -)$.

There exists a rich structure on Ext functors, which is called the composition, pairing or Yoneda product. This is derived from the composition of Hom functors by using formal properties of the category of modules that it is an abelian category with enough injectives and projectives (see [15, Chapter III] or [31]). By extending this construction, we obtain the following pairing for left T-modules M_i from an object $\mathcal{Y}_i \in \text{Mat}_{\mathbb{k}}$:

$$\text{Ext}_{\mathbb{T}}^p(M_1, M_2) \otimes \text{Ext}_{\mathbb{T}}^q(M_2, M_3) \rightarrow \text{Ext}_{\mathbb{T}}^{p+q}(M_1, M_3).$$

If $p = q = 0$, then this coincides with (14) under the isomorphism in Example 4.25.

Definition 4.26. For a left T-module M from an object \mathcal{Y} , the *Ext monad* on M , denoted by $\text{Ext}_{\mathbb{T}}^*(M, M)$, is defined to be a graded monad $\bigoplus_{p \in \mathbb{Z}} \text{Ext}_{\mathbb{T}}^p(M, M)$ whose monad operation is induced by the previous pairing.

5. EIGENMONADS

In this section, we present a generalization of idealizers and eigenrings for monad theory in $\text{Mat}_{\mathbb{k}}$. Furthermore, we construct several adjunctions of categories of modules associated with eigenmonads, which extends those for eigenrings in (2). One noteworthy outcome of this section is the presentation of eigenrings in various forms (see Proposition 5.20). In section 5.4, we also give some examples that arise from algebras.

Throughout this section, we fix an object $\mathcal{X} \in \text{Mat}_{\mathbb{k}}$, a monad T in $\text{Mat}_{\mathbb{k}}$ on \mathcal{X} .

5.1. Definition of eigenmonads. In this section, we extend the notion of an eigenring [17, 3] to monads in $\text{Mat}_{\mathbb{k}}$.

Definition 5.1. Let \mathcal{Y} be an object of $\text{Mat}_{\mathbb{k}}$ and M be a left T-bimodule from \mathcal{Y} (see section 4.1). For submodules $T' \subset T$ and $M' \subset M$, we define a submodule $T' \triangleright M' \subset M$ by

$$(T' \triangleright M')(Y, X) := \{f \triangleright g \in M(Y, X) \mid Z \in \mathcal{X}, f \in T'(Y, Z), g \in M'(Z, X)\}, \forall X, Y \in \mathcal{X}.$$

If $M = T$ is the regular left T-module, then we write $T' \circ M' := T' \triangleright M'$.

For later use, we also introduce an analogue by replacing the submodule M' with a distinguished element $h \in M(Y, X)$ for some $X, Y \in \mathcal{X}$. Let $M(-, X)$ be the \mathcal{X} -indexed module such that $(M(-, X))(Z) := M(Z, X)$ for $Z \in \mathcal{Y}$. We define a morphism $T' \triangleright h : \{X\} \rightsquigarrow \mathcal{X}$ to be an $(\mathcal{X} \times \{X\})$ -indexed submodule $T' \triangleright h \subset M(-, X)$ by

$$(T' \triangleright h)(Z, X) := \{f \circ h \in M(Z, X) \mid f \in T'(Z, Y)\}, \forall Z \in \mathcal{X}.$$

The following definitions are made using this notation.

Definition 5.2. A submodule $J \subset T$ is *multiplicative* if $J \circ J \subset J$.

In the following, we give a generalization of the notion of an idealizer.

Definition 5.3 (Eigenmonad). Let J be a multiplicative submodule of T . We define the *idealizer monad* of J to be the submonad $I_{\mathbb{T}}(J)$ of T such that

$$(15) \quad (I_{\mathbb{T}}(J))(Y, X) := \{f \in T(Y, X) \mid J \circ f \subset J(-, X), f \circ J \subset J(Y, -)\}.$$

It is clear that J is a submodule of $I_{\mathbb{T}}(J)$, since J is assumed to be multiplicative. Hence, $I_{\mathbb{T}}(J)$ is the maximal submonad of T which contains J as a two-sided ideal. The *eigenmonad* of T by J is defined to be the quotient

$$E_{\mathbb{T}}(J) := I_{\mathbb{T}}(J)/J.$$

This inherits a monad structure from $I_{\mathbb{T}}(J)$ since J is its two-sided ideal of $I_{\mathbb{T}}(J)$.

Example 5.4. By Example 4.3, the above definitions implies the classical notions of idealizer and eigenring if $\mathcal{X} = *$.

Example 5.5. Recall the subquotient preorder in Definition 4.7. For a multiplicative submodule $J \subset T$, we have $T \gg E_T(J)$.

Proposition 5.6. *Let $J \subset T$ be a multiplicative submodule. For the opposite monad T° , J is a multiplicative submodule of T° , and we have $E_{T^\circ}(J) \cong E_T(J)^\circ$.*

Proof. By the definition of idealizer monads, we have an isomorphism $I_{T^\circ}(J) \cong I_T(J)^\circ$ of monads. Hence, we obtain the statement. \square

5.2. Construction of eigenmonad adjunctions. In this paper, we mainly study eigenmonads by a particular multiplicative submodule, called a *left ideal*. This section presents a definition of a left ideal of a monad in $\text{Mat}_{\mathbb{k}}$. Moreover, for a left ideal $J \subset T$, we give a construction of an adjunction between T -modules and $E_T(J)$ -modules which we refer to as the *eigenmonad adjunction*.

Definition 5.7. A submodule $J \subset T$ is a *left ideal* of T if we have $T \circ J \subset J$. In a symmetric fashion, A submodule $J \subset T$ is a *right ideal* of T if we have $J \circ T \subset J$. A *two-sided ideal* of T is a left ideal which is also a right ideal of T .

Remark 5.8. By Example 4.3, it is obvious that our notions generalize the classical notion of an ideal of an algebra.

In what follows, we study eigenmonads by left ideals.

Example 5.9. If J is a two-sided ideal, then the idealizer is equal to T , so that the eigenmonad becomes the quotient monad T/J .

Remark 5.10. If J is a left (right, resp.) ideal of T , the definition of the eigenmonad $E_T(J)$ is simplified. Indeed, the second (first, resp.) condition of (15) is redundant.

Proposition 5.11. *Let T be a monad in $\text{Mat}_{\mathbb{k}}$ on \mathcal{X} and J be a left ideal of T . The monad structure on T induces a $(T, E_T(J))$ -bimodule structure on T/J .*

Proof. The left T -action on T itself induces that on T/J . Moreover, the right T -action on T itself induces the right $E_T(J)$ -action on T/J . In fact, for $f \in (I_T(J))(Y, X)$, we have $J(Z, Y) \circ f = 0$ so that it gives a well-defined map

$$(T/J)(Z, Y) \otimes (I_T(J))(Y, X) \rightarrow (T/J)(Z, X).$$

It induces the right $E_T(J)$ -action

$$(T/J)(Z, Y) \otimes (E_T(J))(Y, X) \rightarrow (T/J)(Z, X),$$

since J is a left ideal. \square

Definition 5.12. The quotient T/J equipped with the $(T, E_T(J))$ -bimodule structure in Proposition 5.11 is referred to as the *canonical bimodule associated with (T, J)* , or, if T is specified, the *canonical bimodule associated with J* .

This bimodule structure of T/J is of fundamental importance for our main results. By applying the constructions in Definitions 4.17 and 4.18, we obtain the following adjunctions where Hom^R and Hom^L are defined in Definition 4.18:

Proposition 5.13 (Eigenmonad adjunctions). *Let J be a left ideal of a monad T in $\text{Mat}_{\mathbb{k}}$.*

(1) *We have an adjunction of the category of left $E_T(J)$ -modules and that of left T -modules:*

$$T/J \otimes_{E_T(J)} (-) : {}_{E_T(J)}\text{Mod} \begin{array}{c} \xleftarrow{T} \\ \xrightarrow{T} \end{array} {}_T\text{Mod} : \text{Hom}_{T}^L(T/J, -).$$

(2) We have an adjunction of the category of right $E_T(\mathbf{J})$ -modules and that of right T -modules:

$$(-) \otimes_T T/\mathbf{J} : \mathbf{Mod}_T \xrightleftharpoons{T} \mathbf{Mod}_{E_T(\mathbf{J})} : \mathrm{Hom}_{E_T(\mathbf{J})}^R(T/\mathbf{J}, -).$$

Proof. The statements follow from the tensor-hom adjunction (13). \square

Example 5.14. In this paper, we are mainly interested in (1) of Proposition 5.13 which are closely related to several well known adjunctions:

- In section 5.4, we present some examples arising from algebras.
- The adjunction for polynomial approximation of modules over C is refined in Corollary 7.14.
- The adjunction for outer approximation of modules over gr^0 is refined in Corollary 10.9.
- The adjunction related to Lie-algebraic structures and free groups [21, 12] is reproduced from Corollary 11.14.
- In Corollary 16.7, we give an example related to symmetric groups and free modules.

On the one hand, the adjunction of (2) also naturally arises in the context of algebraic topology. The readers are referred to Remark 7.16 for an example.

Definition 5.15. In this paper, the adjunction in (1) of Proposition 5.13 is called the *eigenmonad adjunctions associated with (T, \mathbf{J})* .

Remark 5.16. It is important to note that each left ideal of T naturally induces homological invariants of T -modules. Indeed, it is immediate from the eigenmonad adjunction that we have an assignment of a left $E_T(\mathbf{J})$ -module $\mathrm{Ext}_T^*(T/\mathbf{J}, \mathbf{M})$ to a left T -module \mathbf{M} (see Definition 4.22). In a parallel way, the Tor modules give an assignment of a right $E_T(\mathbf{J})$ -module $\mathrm{Tor}_T^*(\mathbf{N}, T/\mathbf{J})$ to a right T -module \mathbf{N} . Group (co)homology and Hochschild (co)homology are of these types, as demonstrated in section 5.4.

5.3. Vanishing modules. In this section, we give a concise description of the functor $\mathrm{Hom}_T(T/\mathbf{J}, -)$ in Proposition 5.13. Subsequently, we give four distinct descriptions of an eigenmonad. The results presented here are applied concrete examples later.

Definition 5.17. For a set \mathcal{Y} , let $\mathbf{M} : \mathcal{Y} \rightsquigarrow \mathcal{X}$ be a left T -module from \mathcal{Y} . For a left ideal \mathbf{J} of T , we define the *\mathbf{J} -vanishing module $\mathbf{V}_{\mathbf{J}}(\mathbf{M}) : \mathcal{Y} \rightsquigarrow \mathcal{X}$* to be a submodule of \mathbf{M} such that

$$(\mathbf{V}_{\mathbf{J}}(\mathbf{M}))(X, Y) := \{f \in \mathbf{M}(X, Y) \mid \mathbf{J} \triangleright f \cong 0\}, \quad \forall X \in \mathcal{X}, Y \in \mathcal{Y}.$$

We endow the \mathbf{J} -vanishing module with a left $E_T(\mathbf{J})$ -module structure as we see below.

Proposition 5.18. *Let \mathbf{M} be a left T -module. The left T -module structure on \mathbf{M} induces a left $E_T(\mathbf{J})$ -module structure on $\mathbf{V}_{\mathbf{J}}(\mathbf{M})$.*

Proof. It is sufficient to show that the assignment of $f \triangleright x \in (\mathbf{V}_{\mathbf{J}}(\mathbf{M}))(Z, X)$ to $f \bmod \mathbf{J} \in (\mathbf{I}_T(\mathbf{J}))(Z, Y)/\mathbf{J}(Z, Y)$ and $x \in (\mathbf{V}_{\mathbf{J}}(\mathbf{M}))(Y, X)$ is well-defined for any $Y, Z \in \mathcal{X}$ and $X \in \mathcal{Y}$. Note that $g \circ f$ lives in \mathbf{J} by the assumption that f lies in the idealizer monad. Hence, we have $f \triangleright x \in (\mathbf{V}_{\mathbf{J}}(\mathbf{M}))(Z, X)$ since $g \triangleright (f \triangleright x) = (g \circ f) \triangleright x = 0$ for any g in \mathbf{J} . \square

Recall that the hom module in Definition 4.18 inherits a natural bimodule structure from the components. For a left T -module \mathbf{M} , the left T -module $\mathrm{Hom}_T(T, \mathbf{M})$ is naturally isomorphic to \mathbf{M} . By the Yoneda lemma, we have

$$\mathrm{Hom}_T(T, \mathbf{M}) \xrightarrow{\cong} \mathbf{M}.$$

This observation leads to another description of the vanishing module by using homomorphism sets:

Proposition 5.19. *The evaluation at the unit of \mathbb{T} induces a natural isomorphism of left $E_{\mathbb{T}}(\mathbb{J})$ -modules.*

$$\mathrm{Hom}_{\mathbb{T}}(\mathbb{T}/\mathbb{J}, \mathbb{M}) \xrightarrow{\cong} V_{\mathbb{J}}(\mathbb{M}).$$

Proof. Note that the precomposition gives a submodule $\iota : \mathrm{Hom}_{\mathbb{T}}(\mathbb{T}/\mathbb{J}, \mathbb{M}) \hookrightarrow \mathrm{Hom}_{\mathbb{T}}(\mathbb{T}, \mathbb{M})$. It is easy to see that the composition of ι with the above Yoneda isomorphism factors via $V_{\mathbb{J}}(\mathbb{M}) \subset \mathbb{M}$. It suffices to prove that the factorization $\mathrm{Hom}_{\mathbb{T}}(\mathbb{T}/\mathbb{J}, \mathbb{M}) \rightarrow V_{\mathbb{J}}(\mathbb{M})$ is surjective pointwisely. For $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$, let $x \in (V_{\mathbb{J}}(\mathbb{M}))(X, Y)$. We define the \mathbb{T} -homomorphism $\xi : (\mathbb{T}/\mathbb{J})(-, X) \rightarrow \mathbb{M}(-, Y)$. For each $X' \in \mathcal{X}$, we set $\xi_{X'}(u) := u' \triangleright x$ where $u' \in \mathbb{T}(X', X)$ is a representative of $u \in (\mathbb{T}/\mathbb{J})(X', X)$. The collection of $\xi_{X'}$ for $X' \in \mathcal{X}$ gives a \mathbb{T} -homomorphism $\xi : (\mathbb{T}/\mathbb{J})(-, X) \rightarrow \mathbb{M}(-, Y)$. By definitions, the evaluation of ξ at the unit of \mathbb{T} (modular \mathbb{J}) coincides with x . It completes the proof. \square

Proposition 5.20. *The eigenmonad has equivalent descriptions as follows:*

- (1) *The quotient monad $I_{\mathbb{T}}(\mathbb{J})/\mathbb{J}$ of the idealizer monad.*
- (2) *The vanishing module $V_{\mathbb{J}}(\mathbb{T}/\mathbb{J})$.*
- (3) *The opposite of the endomorphism monad $\mathrm{Hom}_{\mathbb{T}}(\mathbb{T}/\mathbb{J}, \mathbb{T}/\mathbb{J})^{\circ}$ (see Definition 4.21).*
- (4) *The opposite of the zeroth Ext monad $\mathrm{Ext}_{\mathbb{T}}^0(\mathbb{T}/\mathbb{J}, \mathbb{T}/\mathbb{J})^{\circ}$ (see Definition 4.26).*

Proof. The inclusion $I_{\mathbb{T}}(\mathbb{J}) \rightarrow \mathbb{T}$ induces a monomorphism $E_{\mathbb{T}}(\mathbb{J}) \rightarrow \mathbb{T}/\mathbb{J}$. It is obvious that the image is contained in $V_{\mathbb{J}}(\mathbb{T}/\mathbb{J})$, so (2) follows. (3) follows from the first one and the isomorphism in Proposition 5.19. One can directly check that it is an anti-homomorphism between monads. (3) and (4) are the same since the zeroth Ext functor is isomorphic to hom functor. \square

Remark 5.21. In practice, it is useful to apply the second form of Proposition 5.20 when we compute the eigenmonad. See Theorem 11.13 and (2) of Theorem 16.4.

By Proposition 5.18, we obtain a functor $V_{\mathbb{J}} : {}_{\mathbb{T}}\mathrm{Mod} \rightarrow E_{\mathbb{T}}(\mathbb{J})\mathrm{Mod}$. This is a right adjoint:

Proposition 5.22. *We have an adjunction of categories:*

$$\mathbb{T}/\mathbb{J} \otimes_{E_{\mathbb{T}}(\mathbb{J})} (-) : E_{\mathbb{T}}(\mathbb{J})\mathrm{Mod} \xleftarrow{\quad \tau \quad} {}_{\mathbb{T}}\mathrm{Mod} : V_{\mathbb{J}}.$$

Proof. By Proposition 5.19, it suffices to prove that the functor $\mathbb{T}/\mathbb{J} \otimes_{E_{\mathbb{T}}(\mathbb{J})} (-)$ is a left adjoint to $\mathrm{Hom}_{\mathbb{T}}(\mathbb{T}/\mathbb{J}, -) : {}_{\mathbb{T}}\mathrm{Mod} \rightarrow E_{\mathbb{T}}(\mathbb{J})\mathrm{Mod}$. It follows from Proposition 5.13. \square

5.4. Example: algebras. For the sake of convenience, this section presents some familiar examples of the eigenmonad adjunction (see Definition 5.15) by considering a monad in $\mathrm{Mat}_{\mathbb{k}}$ on $\mathcal{X} = *$. As illustrated in Example 4.3, such a monad is nothing but a \mathbb{k} -algebra. Consequently, the examples of the eigenmonad adjunction presented here appear in the classical algebra theory.

Note that this section is *not* directly related to the main theorems of this paper. Readers interested in the main results are directed to sections 6, 7, parts 2 and 3.

5.4.1. Two-sided ideals. Let A be a \mathbb{k} -algebra and $J \subset A$ be a two-sided ideal. The eigenring $E_A(J)$ is isomorphic to the quotient algebra A/J . The associated eigenmonad adjunction yields the following one where $q : A \rightarrow A/J$ is the canonical surjection:

$$(16) \quad q^* : A/J\mathrm{Mod} \xleftarrow{\quad \tau \quad} A\mathrm{Mod} : V_J.$$

For instance, let us consider the group algebra $\mathbb{k}[G]$ for a group G . The augmentation ideal $I_{\mathbb{k}}(G)$ is defined to be the kernel of $\mathbb{k}[G] \rightarrow \mathbb{k}; \sum_{g \in G} a_g g \mapsto \sum_{g \in G} a_g$, which is a two-sided ideal. By the isomorphism $\mathbb{k}[G]/I_{\mathbb{k}}(G) \cong \mathbb{k}$, (16) yields an adjunction between the category of \mathbb{k} -modules and that of $\mathbb{k}[G]$ -modules. It can be observed that the vanishing

module $V_{\mathbb{I}_k(G)}(M)$ of a $\mathbb{k}[G]$ -module M is the same as the invariant submodule M^G by the definition. Furthermore, the homological invariants in Remark 5.16 with respect to $T = \mathbb{k}[G]$ and $J = \mathbb{I}_k(G)$ are nothing but group (co)homology by the definitions [30].

5.4.2. Enveloping algebras. Consider a \mathbb{k} -algebra B . We denote by $B^e := B \otimes B^o$ the *enveloping algebra* of B where B^o is the opposite algebra of B . The multiplication of B gives a B^e -homomorphism $\nabla : B^e \rightarrow B$. Let J_B be the kernel of ∇ which is a left ideal of B^e . Under the usual identification of left B^e -modules with (B, B) -bimodules, the J_B -vanishing module $V_{J_B}(M)$ of a B^e -module M coincides with the center $Z(M)$ of the corresponding (B, B) -bimodule. The adjunction in Proposition 5.22 is equivalent with the following one:

$$(17) \quad B \otimes_{Z(B)} (-) : {}_{Z(B)}\text{Mod} \xrightleftharpoons[T]{\quad} {}_B\text{Mod}_B : Z.$$

By the definitions, the Ext and Tor in Remark 5.16 with respect to $T = B^e$ and $J = J_B$ are equal to Hochschild cohomology and Hochschild homology respectively [30].

The algebra B is an *Azumaya algebra* if it is a central \mathbb{k} -algebra which is separable, i.e. a projective \mathbb{k} -module over its enveloping algebra B^e . For a central \mathbb{k} -algebra B , the adjunction (17) induces an equivalence of categories if and only if B is an Azumaya algebra (for example, see [2, section 1]).

5.4.3. Matrix algebras. Let $n \in \mathbb{N}$ and $M_n(\mathbb{k})$ be the matrix algebra. For I an ideal of \mathbb{k} and $u \in \mathbb{k}^n$, we define a left ideal $D(I, u) \subset M_n(\mathbb{k})$ by

$$D(I, u) := \{K \in M_n(\mathbb{k}) \mid Ku \in I^n\}.$$

The eigenmonad adjunction associated with $(M_n(\mathbb{k}), D(I, u))$ contains the following one:

$$\mathbb{k}^n \otimes (-) : {}_{\mathbb{k}}\text{Mod} \xrightleftharpoons[T]{\quad} {}_{M_n(\mathbb{k})}\text{Mod} : \text{Hom}_{M_n(\mathbb{k})}(\mathbb{k}^n, -).$$

For instance, by setting $I = 0$ and $u_0 = (0, 1, 1, \dots, 1) \in \mathbb{k}^n$, one may directly prove an isomorphism $M_n(\mathbb{k})/D(0, u_0) \cong \mathbb{k}^n$ of left $M_n(\mathbb{k})$ -modules; and $E_{M_n(\mathbb{k})}(D(0, u_0)) \cong \mathbb{k}$ of \mathbb{k} -algebras. It is well known that the above adjunction gives an equivalence of categories. In particular, the right adjoint is an exact functor, so that $\text{Ext}_{M_n(\mathbb{k})}^q(\mathbb{k}^n, -) \cong 0$ for $q > 0$.

6. IDEALS AND CLASSES OF MODULES

This section presents a correspondence between a left ideal of a monad T and a class of (isomorphism classes of) T -modules by applying the previous constructions. This leads to a problem of finding out a left ideal corresponding to a given class of T -modules, which is solved for concrete cases in the following parts.

6.1. Refined eigenmonad adjunctions. In this section, we introduce some properties of modules which depend on a left ideal of the base monad. These naturally yield a refinement of eigenmonad adjunctions (see Definition 5.15). The refined adjunction is designed to construct an equivalence between T -modules and $E_T(J)$ -modules. Indeed, it often gives an equivalence of categories (see Corollaries 6.8, 12.3 and 17.6). In section 6.2, we explain another motivation of specifying these properties.

Definition 6.1. Let T be a monad in $\text{Mat}_{\mathbb{k}}$ on an object \mathcal{X} and $J \subset T$ be a left ideal.

- A left T -module M is *J-vanishingly generated* if the counit of the eigenmonad adjunction is an epimorphism:

$$T/J \otimes_{E_T(J)} V_J(M) \twoheadrightarrow M.$$

Denote by ${}_T\text{Mod}^{J\text{-v.g.}}$ the full subcategory of left T -modules which are J -vanishingly generated.

- A left $E_T(J)$ -module N is *J-vanishingly extensible* if the unit of the eigenmonad adjunction is a monomorphism:

$$N \rightarrow V_J (T/J \otimes_{E_T(J)} N).$$

Denote by ${}_{E_T(J)}\text{Mod}^{J\text{-v.e.}}$ the full subcategory of left $E_T(J)$ -modules which are *J-vanishingly extensible*.

Remark 6.2. It is clear from definitions that a left T -module M is *J-vanishingly generated* if and only if the morphism induced by the action of T gives a epimorphism $T \otimes V_J(M) \twoheadrightarrow M$. Similarly, for a left $E_T(J)$ -module N , it is *J-vanishingly extensible* if and only if the tensor product with the unit of T induces a monomorphism $N \rightarrow T/J \otimes_{E_T(J)} N$.

Example 6.3. By (2) of Proposition 5.20, the regular left $E_T(J)$ -module $E_T(J)$ itself is *J-vanishingly extensible*.

Let $F : C \rightarrow D$ be a functor between categories, and G be a right adjoint to F . An elementary argument on adjunctions shows that, for $C \in C$ and $D = F(C) \in D$, the counit $F(G(D)) \rightarrow D$ is a (split) epimorphism. There is an analogous statement for the unit. This observation leads to the following:

Example 6.4. The T -module $T/J \otimes_{E_T(J)} N$ is *J-vanishingly generated* for a left $E_T(J)$ -module N . Similarly, for a left T -module M , $V_J(M)$ is *J-vanishingly extensible*.

Example 6.5. The left T -module T/J is *J-vanishingly generated*; but the regular left T -module T itself might not be *J-vanishingly generated* (for instance, $(T, J) = (L_{\text{gr}^o}, \mathbb{I}^{\text{pr}})$). Hence, the functor $T/J \otimes_{E_T(J)} (-)$ in Proposition 5.22 is not essentially surjective in general. In fact, its image is contained in ${}_{T}\text{Mod}^{J\text{-v.g.}}$ by Example 6.4.

Proposition 6.6. *Let $J \subset T$ be a two-sided ideal.*

- (1) *A left T -module M is *J-vanishingly generated* if and only if $V_J(M) = M$.*
- (2) *Every left $E_T(J)$ -module is *J-vanishingly extensible*.*

Proof. The eigenmonad $E_T(J)$ by J is isomorphic to T/J since J is two-sided. Hence, (1) follows from the definition. Moreover, every left $E_T(J)$ -module N is *J-vanishingly extensible* since we have $T/J \otimes_{E_T(J)} N \cong N$. \square

Proposition 6.7 (Refined eigenmonad adjunction). *The restriction of the eigenmonad adjunction associated with (T, J) induces the following adjunction:*

$$T/J \otimes_{E_T(J)} (-) : {}_{E_T(J)}\text{Mod}^{J\text{-v.e.}} \xrightleftharpoons[T]{\quad} {}_T\text{Mod}^{J\text{-v.g.}} : V_J.$$

Furthermore, both of these restrictions are faithful.

Proof. Proposition 5.22 implies the first assertion by Example 6.4. The last assertion follows from more formal statement, but let us sketch the proof here. We prove that the restriction $V_J : {}_T\text{Mod}^{J\text{-v.g.}} \rightarrow {}_{E_T(J)}\text{Mod}^{J\text{-v.e.}}$ is faithful. Let M, M' be *J-vanishingly generated* left T -modules. The evaluation for M induces the following map:

$$(18) \quad \text{Hom}_T(M, M') \rightarrow \text{Hom}_T(T \otimes_{E_T(J)} V_J(M), M') \cong \text{Hom}_{E_T(J)}(V_J(M), V_J(M')).$$

This map is injective since M is *J-vanishingly generated*. Analogously, the functor $T/J \otimes_{E_T(J)} (-)$ is faithful. \square

Corollary 6.8. *Let $J \subset T$ be a two-sided ideal. The refined eigenmonad adjunction associated with (T, J) gives an adjoint equivalence of categories:*

$$q^* : {}_{T/J}\text{Mod} \xrightleftharpoons[T]{\quad} {}_T\text{Mod}^{J\text{-v.g.}} : V_J.$$

Here, q^ is the functor induced by the quotient homomorphism $q : T \rightarrow T/J$.*

Proof. Proposition 6.7 gives the adjunction in the statement by (2) of Proposition 6.6. The functor V_J clearly induces an equivalence by (1) of Proposition 6.6. \square

6.2. Internalizers of classes of modules. Let ${}_{\mathsf{T}}\mathcal{M}\text{od}$ be the class of isomorphism classes of left T -modules. For a subclass $\mathcal{S} \subset {}_{\mathsf{T}}\mathcal{M}\text{od}$, we say that a left ideal $J \subset \mathsf{T}$ is a T -*internalizer* of \mathcal{S} when a left T -module is J -vanishingly generated if and only if its isomorphism class lies in \mathcal{S} . The existence of a T -internalizer, as we name it, makes us to understand the subclass \mathcal{S} by using internal data of T , namely J . In particular, the full subcategory of ${}_{\mathsf{T}}\mathcal{M}\text{od}$ generated by representatives of isomorphism classes lying in \mathcal{S} naturally induces an adjunction of categories of modules as Proposition 6.7. In sections 7, 9.2, and 10.1, we prove or disprove the existence of a T -internalizer of \mathcal{S} with respect to concrete T and \mathcal{S} .

Definition 6.9. A left ideal $J \subset \mathsf{T}$ is a T -*internalizer* of $\mathcal{S} \subset {}_{\mathsf{T}}\mathcal{M}\text{od}$ if $\mathcal{S} = \mathcal{V}(J)$. If J is a two-sided ideal, then it is called a *core* T -*internalizer*.

Example 6.10. The zero ideal of T is a core T -internalizer of ${}_{\mathsf{T}}\mathcal{M}\text{od}$ since $\mathcal{V}(0) = {}_{\mathsf{T}}\mathcal{M}\text{od}$.

Lemma 6.11. *Let J, J' be left ideals of T . Consider the following conditions:*

- (1) $J \subset J'$.
- (2) $\mathcal{V}(J') \subset \mathcal{V}(J)$.

The condition (1) implies (2). Furthermore, if J is a two-sided ideal, then (2) implies (1).

Proof. We prove the first assertion. Let M be a left T -module. By definition, we have $V_{J'}(\mathsf{M}) \subset V_J(\mathsf{M})$. Hence, if M is J' -vanishingly generated, then it is J -vanishingly generated.

We now prove the second statement. Note that T/J' is J' -vanishingly generated. By the assumption, this is J -vanishingly generated. By Proposition 6.6, we have $V_J(\mathsf{T}/J') = \mathsf{T}/J'$ which implies $J \subset J'$. \square

In the following two examples, we give illustrative observations on algebras. Note that these imply that an internalizer of a given class is not unique.

Example 6.12. Recall the discussion in section 5.4.3. It implies that every $M_n(\mathbb{k})$ -module is $D(0, u_0)$ -vanishingly generated; and every \mathbb{k} -module is $D(0, u_0)$ -vanishingly extensible. Hence, the left ideal $D(0, u_0)$ is a $M_n(\mathbb{k})$ -internalizer of ${}_{M_n(\mathbb{k})}\mathcal{M}\text{od}$.

Corollary 6.13. *Suppose that \mathbb{k} is a field. All the left ideals of $M_n(\mathbb{k})$ but itself are $M_n(\mathbb{k})$ -internalizers of ${}_{M_n(\mathbb{k})}\mathcal{M}\text{od}$.*

Proof. For a commutative unital ring \mathbb{k} , all the maximal left ideals of $M_n(\mathbb{k})$ are given by $D(I, u)$ for a maximal ideal $I \subset \mathbb{k}$ and $u \in \mathbb{k}^n \setminus I^n$ [26, Theorem 1.2]. Hence, all the left ideals of $M_n(\mathbb{k})$ are contained in some $D(I, u)$. For a field \mathbb{k} , every maximal ideal of \mathbb{k} is zero so that, by the above observation and Lemma 6.11, we obtain the statement. \square

Example 6.14. Recall the discussion in section 5.4.2. If B is an Azuyama algebra, then the left ideal $J \subset B^e$ is a B^e -internalizer of ${}_{B^e}\mathcal{M}\text{od}$. The zero ideal is also a B^e -internalizer of ${}_{B^e}\mathcal{M}\text{od}$ by Example 6.10.

We now return to a general monad T .

Proposition 6.15. *Let $\mathcal{S} \subset {}_{\mathsf{T}}\mathcal{M}\text{od}$. A core T -internalizer of \mathcal{S} , if it exists, is contained in every T -internalizer of \mathcal{S} . In particular, there exists at most one core T -internalizer.*

Proof. Let J be a core T -internalizer of \mathcal{S} . For a T -internalizer J' of \mathcal{S} , we have $\mathcal{V}(J') = \mathcal{S} = \mathcal{V}(J)$, so that we obtain $J \subset J'$ by the last statement of Lemma 6.11. The last assertion is immediate from the previous result. \square

We now introduce a method to construct a left ideal from a given subclass $\mathcal{S} \subset {}_{\mathsf{T}}\mathcal{M}\text{od}$, which provides a candidate for a T -internalizer.

Definition 6.16. Let \mathbf{M} be a left T -module. The *annihilator* of \mathbf{M} , denoted by $\text{Ann}(\mathbf{M})$, is defined to be a left ideal $\text{Ann}(\mathbf{M}) \subset T$ such that

$$(\text{Ann}(\mathbf{M})) (Y, X) := \{f \in T(Y, X) \mid f \triangleright \mathbf{M}(X) \cong 0\}, \forall X, Y \in \mathcal{X}.$$

Definition 6.17. For a subclass $\mathcal{S} \subset {}_T\text{Mod}$, we define a left ideal $\text{Ann}(\mathcal{S})$ of T to be the intersection of the annihilator of \mathbf{M} for any T -module \mathbf{M} whose isomorphism class lies in \mathcal{S} :

$$\text{Ann}(\mathcal{S}) := \bigcap_{[\mathbf{M}] \in \mathcal{S}} \text{Ann}(\mathbf{M}).$$

Proposition 6.18. (1) For a subclass $\mathcal{S} \subset {}_T\text{Mod}$, we have $\mathcal{S} \subset \mathcal{V}(\text{Ann}(\mathcal{S}))$.

(2) If $\mathcal{S} \subset \mathcal{S}'$, then $\text{Ann}(\mathcal{S}') \subset \text{Ann}(\mathcal{S})$.

Proof. We first prove (1). Let \mathbf{M} be a left T -module such that $[\mathbf{M}] \in \mathcal{S}$. Let $\mathbf{J} = \text{Ann}(\mathbf{M})$. Then, by the definition, we see that $V_{\mathbf{J}}(\mathbf{M}) = \mathbf{M}$. Especially, \mathbf{M} is \mathbf{J} -vanishingly generated, i.e. $[\mathbf{M}] \in \mathcal{V}(\mathbf{J})$. The definition of $\text{Ann}(\mathcal{S})$ implies $\text{Ann}(\mathcal{S}) \subset \mathbf{J}$, so, by Lemma 6.11, we obtain $\mathcal{V}(\mathbf{J}) \subset \mathcal{V}(\text{Ann}(\mathcal{S}))$. Hence, we obtain $[\mathbf{M}] \in \mathcal{V}(\text{Ann}(\mathcal{S}))$. The part (2) is tautological. \square

Lemma 6.19. For a left ideal $\mathbf{J} \subset T$, we have $\text{Ann}(\mathcal{V}(\mathbf{J})) \subset \mathbf{J}$. Moreover, we have $\text{Ann}(\mathcal{V}(\mathbf{J})) = \mathbf{J}$ if and only if \mathbf{J} is a two-sided ideal.

Proof. Since T/\mathbf{J} is \mathbf{J} -vanishingly generated, we have

$$(19) \quad \text{Ann}(\mathcal{V}(\mathbf{J})) \subset \text{Ann}(T/\mathbf{J}) \subset \mathbf{J}.$$

Hence, we obtain $\text{Ann}(\mathcal{V}(\mathbf{J})) \subset \mathbf{J}$. Moreover, if $\text{Ann}(\mathcal{V}(\mathbf{J})) = \mathbf{J}$, then, by (19), we obtain $\text{Ann}(T/\mathbf{J}) = \mathbf{J}$ which implies that \mathbf{J} is a two-sided ideal. Conversely, if \mathbf{J} is a two-sided ideal, for a \mathbf{J} -vanishingly generated \mathbf{M} , we have $V_{\mathbf{J}}(\mathbf{M}) = \mathbf{M}$ by Proposition 6.6. This implies that $\text{Ann}(\mathbf{M})$ contains \mathbf{J} . Thus, we obtain $\mathbf{J} \subset \text{Ann}(\mathcal{V}(\mathbf{J}))$ which leads to our statement. \square

In the following, we give a characterization of a core internalizer.

Theorem 6.20. If a subclass $\mathcal{S} \subset {}_T\text{Mod}$ has a core T -internalizer, then it should be $\text{Ann}(\mathcal{S})$.

Proof. Let \mathbf{J} be a core internalizer of \mathcal{S} . By the definition, we have $\mathcal{S} = \mathcal{V}(\mathbf{J})$. By Lemma 6.19, we have $\text{Ann}(\mathcal{S}) = \text{Ann}(\mathcal{V}(\mathbf{J})) = \mathbf{J}$. \square

Remark 6.21. Let \mathcal{L} be the poset of left ideals of T and \mathcal{P} be the poset of subclasses of ${}_T\text{Mod}$. A poset induces a category by regarding each inequality as a morphism. In spite of Proposition 6.18 and Lemma 6.19, note that Ann and \mathcal{V} do not induce an adjunction between \mathcal{L} and \mathcal{P} since Ann, \mathcal{V} are contravariant. Nevertheless, for any \mathcal{S} and \mathbf{J} , we have a map $\mathcal{L}^{\circ}(\text{Ann}(\mathcal{S}), \mathbf{J}) \rightarrow \mathcal{P}(\mathcal{S}, \mathcal{V}(\mathbf{J}))$ between morphism sets. This is automatically a bijection if and only if the left hand side is not empty.

Before we close this section, we show a basic statement for internalizers related to a filtration of two-sided ideals. This is applied to prove that the category of analytic modules do not have a core internalizer (see Proposition 7.26 and Theorem 9.5).

Proposition 6.22. Consider a descending filtration of T consisting of two-sided ideals:

$$\dots \subset \mathbf{J}^{(d+1)} \subset \mathbf{J}^{(d)} \subset \dots \subset T.$$

Denote by \mathcal{S}_{ω} the class of isomorphism classes of a T -module \mathbf{M} such that $\varinjlim_d V_{\mathbf{J}^{(d)}}(\mathbf{M}) \cong \mathbf{M}$. If \mathbf{J} is a core T -internalizer of \mathcal{S}_{ω} , then, for every $X \in \mathcal{X}$, there exists $d_0 \in \mathbb{N}$ such that the filtration below converges to $\mathbf{J}(-, X)$ from the degree d_0 :

$$\mathbf{J}(-, X) = \dots = \mathbf{J}^{(d_0+1)}(-, X) = \mathbf{J}^{(d_0)}(-, X) \subset T(-, X).$$

Proof. We first prove that $\mathcal{J} \subset \bigcap_{d \in \mathbb{N}} \mathcal{J}^{(d)}$. Note that, for $k \in \mathbb{N}$, a $\mathcal{J}^{(k)}$ -vanishingly generated left \mathcal{T} -module \mathbb{M} satisfies $\varinjlim_d V_{\mathcal{J}^{(d)}}(\mathbb{M}) \cong V_{\mathcal{J}^{(k)}}(\mathbb{M}) \cong \mathbb{M}$. Hence, we obtain $[\mathbb{M}] \in \mathcal{S}_\omega = \mathcal{V}(\mathcal{J})$ so that $\mathcal{V}(\mathcal{J}^{(k)}) \subset \mathcal{V}(\mathcal{J})$. By Lemma 6.11, we see that $\mathcal{J} \subset \mathcal{J}^{(k)}$ since we assume that \mathcal{J} is a two-sided ideal. This holds for any k so that we have $\mathcal{J} \subset \bigcap_{d \in \mathbb{N}} \mathcal{J}^{(d)}$.

Let $X \in \mathcal{X}$. We now prove that there exists $d_0 \in \mathbb{N}$ such that $\mathcal{J}^{(d_0)}(-, X) \subset \mathcal{J}(-, X)$. Note that \mathcal{T}/\mathcal{J} is \mathcal{J} -vanishingly generated. By the hypothesis, we have $\varinjlim_d V_{\mathcal{J}^{(d)}}(\mathcal{T}/\mathcal{J}) \cong \mathcal{T}/\mathcal{J}$. Hence, there exists $d_0 \in \mathbb{N}$ such that $(1_X \bmod \mathcal{J}(X, X)) \in (V_{\mathcal{J}^{(d_0)}}(\mathcal{T}/\mathcal{J}))(X, X)$ where $1_X \in \mathcal{T}(X, X)$ is the unit. This implies that the left action of any $f \in \mathcal{J}^{(d_0)}(Y, X)$ to 1_X , i.e. f itself, lies in $\mathcal{J}(Y, X)$ for any $Y \in \mathcal{Y}$. The combination of the above two results lead to our statement. \square

7. EXAMPLE: POLYNOMIAL FUNCTORS

In this section, we revisit the polynomial functor theory based on the preceding results. Consider a category \mathcal{C} with a zero object $*$ and a symmetric monoidal structure whose unit is $*$ - for example, the (co)product if it exists in \mathcal{C} . A functor $F : \mathcal{C} \rightarrow \mathbb{k}\text{Mod}$ is a polynomial functor (of a given degree d) if it is subject to an analogue of equations (see the equation (20)) satisfied by polynomial functions [9, 12]. By the isomorphism in Example 4.16, all the notions in polynomial functor theory for \mathcal{C} are inherited to left $L_{\mathcal{C}}$ -modules. The main result of this section is that, if \mathcal{C} has finite products, then the polynomial property is encoded into an action of an eigenmonad of $L_{\mathcal{C}}$. We go further by showing that the associated eigenmonad adjunction is equivalent with the adjunction induced by the polynomial approximation.

Throughout this section, we assume that \mathcal{C} has arbitrary finite products and a zero object $*$. Let \mathcal{X} be the object set of \mathcal{C} .

Notation 7.1. We denote by \times the binary product on \mathcal{C} . We also denote the product of morphisms g_k 's in \mathcal{C} by $\times_{k=1}^m g_k := g_1 \times \cdots \times g_m$.

7.1. Polynomial degrees. We give a brief review of the polynomial degree of functors. For an object X , let $\eta_X : * \rightarrow X$ and $\varepsilon_X : X \rightarrow *$ be the unique morphisms in \mathcal{C} . Put $z_X = \eta_X \circ \varepsilon_X$. For an endomorphism $f : X \rightarrow X$ and $\epsilon \in \{0, 1\}$, we set $f^\epsilon := f$ if $\epsilon = 1$; and $f^\epsilon := \text{id}_X$ if $\epsilon = 0$. Let $F \in \mathcal{F}(\mathcal{C})$. The polynomial degree of F does not exceed d if, for any objects X_1, \dots, X_{d+1} of \mathcal{C} , we have

$$(20) \quad \sum (-1)^\epsilon \cdot F(z_{X_1}^{\epsilon_1} \times z_{X_2}^{\epsilon_2} \times \cdots \times z_{X_{d+1}}^{\epsilon_{d+1}}) = 0.$$

Here, the sum is taken over $\epsilon_1, \dots, \epsilon_{d+1} \in \{0, 1\}$ and $\epsilon = \sum_{j=1}^{d+1} \epsilon_j$. In that case, we merely write $\deg F \leq d$. For $d \in \mathbb{N}$, the d -th polynomial approximation $P_d(F) \in \mathcal{F}(\mathcal{C})$ is defined to be the maximal subfunctor of F such that $\deg F \leq d$. Let $\mathcal{F}_{\leq d}(\mathcal{C})$ be the full subcategory of $\mathcal{F}(\mathcal{C})$ generated by functors with $\deg \leq d$. It is characterized as a right adjoint to the inclusion functor $\mathcal{F}_{\leq d}(\mathcal{C}) \rightarrow \mathcal{F}(\mathcal{C})$ [12]. In [12], we give an explicit description of the polynomial approximation.

We now recall the monad $L_{\mathcal{C}}$ defined in Example 4.5.

Definition 7.2. By using the isomorphism in Example 4.16, we define the category $L_{\mathcal{C}}\text{Mod}^{\leq d}$ to be the full subcategory of $L_{\mathcal{C}}\text{Mod}$ corresponding to $\mathcal{F}_{\leq d}(\mathcal{C})$. For a left $L_{\mathcal{C}}$ -module \mathbb{M} , we say that the polynomial degree of F does not exceed d , and denote by $\deg \mathbb{M} \leq d$, if $\mathbb{M} \in L_{\mathcal{C}}\text{Mod}^{\leq d}$. We denote by $P_d : L_{\mathcal{C}}\text{Mod} \rightarrow L_{\mathcal{C}}\text{Mod}^{\leq d}$ the functor induced by $P_d : \mathcal{F}(\mathcal{C}) \rightarrow \mathcal{F}_{\leq d}(\mathcal{C})$ with a slight abuse of notation. For a left $L_{\mathcal{C}}$ -module \mathbb{M} , we call $P_d(\mathbb{M})$ the d -th polynomial approximation.

Definition 7.3. Let $\mathcal{S}_{\mathcal{C}}^d$ be the class of isomorphism classes of $L_{\mathcal{C}}$ -modules with polynomial degree $\leq d$.

The characterization of $P_d : \mathcal{F}(C) \rightarrow \mathcal{F}_{\leq d}(C)$ as a right adjoint gives the following adjunction of categories:

$$(21) \quad \iota : {}_{L_C} \text{Mod}^{\leq d} \xrightleftharpoons[\tau]{\leftarrow} {}_{L_C} \text{Mod} : P_d.$$

7.2. Polynomial ideals. In this section, we introduce a two-sided ideal of the monad L_C called the polynomial ideal, which conceptualizes the polynomial condition. We prove that the polynomial approximation is computed by using the associated vanishing module (see Theorem 7.9). It implies that an L_C -module is polynomial of degree $\leq d$ if and only if the module structure is reduced to a module over the associated eigenmonad. Moreover, the polynomial filtration is encoded into a filtration consisting of the polynomial ideals.

In the following, recall the notation in section 7.1.

Definition 7.4. Let d be a positive integer. For $X \in \mathcal{X}$, we define

$$\pi_X^{C,d} := (\text{id}_X - z_X)^{\times d} \circ \Delta_X^{(d)} \in L_C(X^{\times d}, X) := \mathbb{k}C(X, X^{\times d})$$

where $\Delta_X^{(d)} \in C(X, X^{\times d})$ is the d -fold diagonal map on X . If C is specified, then we omit $\pi_X^d := \pi_X^{C,d}$. We also give an equivalent definition of π_X^d in an expanded form. Let $p_k \in C(X^{\times d}, X)$ be the k -th projection for $k \in \underline{d}$. For $T \subset \underline{d}$, we define $D_X^T \in C(X, X^{\times d})$ characterized by

$$(22) \quad p_k \circ D_X^T = \begin{cases} \text{id}_X & (k \in T), \\ z_X & (\text{otherwise}). \end{cases}$$

Then π_X^d is equivalently defined by

$$(23) \quad \pi_X^{C,d} := \sum_{T \subset \underline{d}} (-1)^{d-|T|} \cdot D_X^T.$$

Remark 7.5. One may describe D_X^T explicitly. We set $\zeta_T(i)$ to be 0 if $i \in T$; and 1 otherwise. Then $D_X^T = \left(\times_{i=1}^d z_X^{\zeta_T(i)} \right) \circ \Delta_X^{(d)} \in C(X, X^{\times d})$.

Definition 7.6. For positive $d \in \mathbb{N}$, we define $\mathbb{I}_C^{(d)} \subset L_C$ as the left ideal generated by π_X^d for $X \in \mathcal{X}$. We also define $\mathbb{I}_C^{(0)} := L_C$. We call $\mathbb{I}_C^{(d)}$ as the d -th polynomial ideal for C .

Lemma 7.7. For $f \in C(X, Y) \subset L_C(Y, X)$, we have $\pi_Y^d \circ f = (f^{\times d}) \circ \pi_X^d$.

Proof. The universality of the product gives $\Delta_Y^{(d)} \circ f = (f^{\times d}) \circ \Delta_X^{(d)}$, and that of the zero object $*$ gives $z_Y \circ f = f \circ z_X$. Hence, the statement follows from the definitions. \square

Lemma 7.8. Let $d \in \mathbb{N}$. The left ideal $\mathbb{I}_C^{(d)}$ is a two-sided ideal of L_C .

Proof. Let $X, Y \in \mathcal{X}$. Any element in $L_C(Y, X)$ is given by a linear combination of some elements of $C(X, Y)$, say f 's. By Lemma 7.7, we obtain $\pi_Y^d \circ f = (f^{\times d}) \circ \pi_X^d \in \mathbb{I}_C^{(d)}(Y^{\times d}, X)$. It implies that $\mathbb{I}_C^{(d)}$ is a right ideal of L_C . \square

One simple consequence is that the eigenmonad by $\mathbb{I}_C^{(d+1)}$ is isomorphic to the quotient monad $L_C/\mathbb{I}_C^{(d+1)}$. Furthermore, by Proposition 5.18, $\mathbb{V}_{\mathbb{I}_C^{(d+1)}}(\mathbb{M})$ is a left $L_C/\mathbb{I}_C^{(d+1)}$ -module. The $\mathbb{I}_C^{(d+1)}$ -vanishing module turns out to be the polynomial approximation:

Theorem 7.9. Let $d \in \mathbb{N}$. For $\mathbb{M} \in {}_{L_C} \text{Mod}$, we have an isomorphism of left L_C -modules:

$$\mathbb{V}_{\mathbb{I}_C^{(d+1)}}(\mathbb{M}) \cong P_d(\mathbb{M}).$$

The module \mathbb{M} is $\mathbb{I}_C^{(d+1)}$ -vanishingly generated if and only if $\deg \mathbb{M} \leq d$.

Proof. Let $F \in \mathcal{F}(C)$ be the one induced by \mathbf{M} via the isomorphism in Example 4.16. Recall the notation in Definition 7.4. By [12], for each object X of C , the module $(P_d(F))(X)$ coincides with the kernel of

$$\sum_{T \subset \underline{d}} (-1)^{d-|T|} \cdot F(D_X^T).$$

This is isomorphic to $(V_{\mathbb{I}_C^{(d+1)}}(\mathbf{M}))(X)$ by the definition of π_X^{d+1} in (23). Hence, we obtain $V_{\mathbb{I}_C^{(d+1)}}(\mathbf{M}) \cong P_d(\mathbf{M})$.

We now see that $\deg \mathbf{M} \leq d$ if and only if $V_{\mathbb{I}_C^{(d+1)}}(\mathbf{M}) = \mathbf{M}$. By Proposition 6.6, the module \mathbf{M} is $\mathbb{I}_C^{(d+1)}$ -vanishingly generated if and only if $V_{\mathbb{I}_C^{(d+1)}}(\mathbf{M}) = \mathbf{M}$, since $\mathbb{I}_C^{(d+1)}$ is a two-sided ideal. \square

It is well-known that the polynomial approximation gives rise to a filtration of functors (for example, see [12]). This fact is conceptualized by using the polynomial ideals:

Proposition 7.10. *For $d \in \mathbb{N}$, we have $\mathbb{I}_C^{(d+1)} \subset \mathbb{I}_C^{(d)}$.*

Proof. It suffices to prove that $\pi_X^{d+1} \in \mathbb{I}_C^{(d)}(X^{\times(d+1)}, X)$ since $\mathbb{I}_C^{(d)}$ is a left ideal.

$$\begin{aligned} (\text{id}_{X^{\times(d-1)}} \times \pi_X^2) \circ \pi_X^d &= (\text{id}_{X^{\times(d-1)}} \times \pi_X^2) \circ (e_X^{\times(d-1)} \times e_X) \circ \Delta_X^{(d)}, \\ &= (e_X^{\times(d-1)} \times (\pi_X^2 \circ e_X)) \circ \Delta_X^{(d)}, \\ &= (e_X^{\times(d-1)} \times (\pi_X^2 \circ (\text{id}_X - \eta_X \circ \varepsilon_X))) \circ \Delta_X^{(d)}, \\ &= (e_X^{\times(d-1)} \times (\pi_X^2 \circ \text{id}_X)) \circ \Delta_X^{(d)} - (e_X^{\times(d-1)} \times (\pi_X^2 \circ \eta_X \circ \varepsilon_X)) \circ \Delta_X^{(d)}, \\ &= (e_X^{\times(d-1)} \times (\pi_X^2 \circ \text{id}_X)) \circ \Delta_X^{(d)}, \\ &= \pi_X^{d+1}. \end{aligned}$$

Hence, we see that $\pi_X^{d+1} = (\text{id}_{X^{\times(d-1)}} \times \pi_X^2) \circ \pi_X^d \in \mathbb{I}_C^{(d)}(X^{\times(d+1)}, X)$. \square

Remark 7.11. For $C \in \{\text{gr}^0, \text{fr}_R^0\}$, we have $\mathbb{I}_C^{(d+1)} \subsetneq \mathbb{I}_C^{(d)}$. It follows from the comparison of our polynomial ideals and certain augmentation ideals in (29) and (40). For instance, the rank of $L_{\text{gr}^0}(m, n)/\mathbb{I}_C^{(d)}(n, m) \cong \mathbb{k}[\mathbb{F}_n^{\times m}]/\mathbb{I}_{\mathbb{k}}(\mathbb{F}_n^{\times m})^d$ is strictly monotone with respect to d , as shown in Theorem 9.1.

By combining Lemma 6.11, this proposition leads to the following. The first statement gives the polynomial filtration of L_C -modules by Theorem 7.9.

Corollary 7.12. \bullet *For \mathbf{M} a left L_C -module, the vanishing modules give a filtration of \mathbf{M} :*

$$V_{\mathbb{I}_C^{(d)}}(\mathbf{M}) \subset V_{\mathbb{I}_C^{(d+1)}}(\mathbf{M}).$$

\bullet *We have $\mathcal{S}_C^d \subset \mathcal{S}_C^{d+1}$.*

7.3. Core internalizer for polynomial modules. Let $d \in \mathbb{N}$. In this section, we discuss an existence of a core L_C -internalizer of the class of L_C -modules with polynomial degree $\leq d$. One consequence is that we obtain a new viewpoint to describe polynomial L_C -modules by using an action of an eigenmonad. See the paragraph below Corollary 7.15 for further details.

Let us recall Definitions 6.17 and 7.3.

Theorem 7.13. *Let $d \in \mathbb{N}$. The two-sided ideal $\mathbb{I}_C^{(d+1)}$ of L_C is the unique core L_C -internalizer of \mathcal{S}_C^d . In particular, we have $\text{Ann}(\mathcal{S}_C^d) = \mathbb{I}_C^{(d+1)}$.*

Proof. To prove that $\mathbb{I}_C^{(d+1)}$ is a core L_C -internalizer of \mathcal{S}_C^d , it suffices to check that, for a left module \mathbb{M} , we have $\deg \mathbb{M} \leq d$ if and only if $\mathbb{V}_{\mathbb{I}_C^{(d+1)}}(\mathbb{M}) = \mathbb{M}$ by Proposition 6.6. In fact, it follows from Theorem 7.9. The uniqueness follows from Proposition 6.15. The last assertion is immediate from Theorem 6.20. \square

The theorem is rephrased without class-set theory language: there exists a unique two-sided ideal $\mathbb{I}_C^{(d+1)} \subset L_C$ such that a left L_C -module has the polynomial degree $\leq d$ if and only if it is $\mathbb{I}_C^{(d+1)}$ -vanishingly generated. Moreover, $\mathbb{I}_C^{(d+1)}$ could be computed by taking the intersection of all the annihilators of L_C -modules with polynomial degree $\leq d$.

Corollary 7.14. *The eigenmonad adjunction associated with $(L_C, \mathbb{I}_C^{(d+1)})$ coincides with (21).*

Proof. It is immediate from Theorem 7.9. \square

Corollary 7.15. *The eigenmonad adjunction associated with $(L_C, \mathbb{I}_C^{(d+1)})$ gives an adjoint equivalence of categories:*

$$\iota : {}_{L_C/\mathbb{I}_C^{(d+1)}}\mathbf{Mod} \xrightleftharpoons[\top]{{}^{\top}} L_C \mathbf{Mod}^{\leq d} : \mathbb{V}_{\mathbb{I}_C^{(d+1)}}.$$

Proof. By Theorem 7.13, we have $\mathcal{S}_C^d = \mathcal{V}(\mathbb{I}_C^{(d+1)})$ so that we obtain $L_C \mathbf{Mod}^{\leq d} \simeq L_C \mathbf{Mod}^{\mathbb{I}_C^{(d+1)-\text{v.g.}}}$. Then our statement follows from Corollary 6.8. \square

Remark 7.16. The above adjunctions are derived from (1) of Proposition 5.13. Note that the adjunction of (2) of Proposition 5.13 also naturally arises in polynomial modules over \mathfrak{gr} (not over \mathfrak{gr}^0) [4], which is not covered by the previous discussion since \mathfrak{gr} does not satisfy the assumptions on C in section 7. However, we can still apply (2) of Proposition 5.13 to $\mathbb{T} = L_{\mathfrak{gr}^0}$ and $\mathbb{J} = \mathbb{I}_{\mathfrak{gr}^0}^{(d+1)}$. By Proposition 5.6, we have $E_{L_{\mathfrak{gr}^0}}(\mathbb{I}_{\mathfrak{gr}^0}^{(d+1)}) \cong E_{L_{\mathfrak{gr}}}(\mathbb{I}_{\mathfrak{gr}^0}^{(d+1)})^{\circ}$. Hence, we obtain the following by the equivalence in Example 4.14 where $q : L_{\mathfrak{gr}} \rightarrow E_{L_{\mathfrak{gr}}}(\mathbb{I}_{\mathfrak{gr}^0}^{(d+1)})$ is the canonical quotient homomorphism:

$$L_{\mathfrak{gr}}/\mathbb{I}_{\mathfrak{gr}^0}^{(d+1)} \otimes_{L_{\mathfrak{gr}}} (-) : L_{\mathfrak{gr}} \mathbf{Mod} \xrightleftharpoons[\top]{{}^{\top}} E_{L_{\mathfrak{gr}}}(\mathbb{I}_{\mathfrak{gr}^0}^{(d+1)}) \mathbf{Mod} : q^*.$$

It is easy to see that, for a left $L_{\mathfrak{gr}}$ -module \mathbb{M} , $L_{\mathfrak{gr}}/\mathbb{I}_{\mathfrak{gr}^0}^{(d+1)} \otimes_{L_{\mathfrak{gr}}} \mathbb{M}$ is isomorphic to the quotient $\mathbb{M} / \left(\mathbb{I}_{\mathfrak{gr}^0}^{(d+1)} \triangleright \mathbb{M} \right)$ which should be called the $\mathbb{I}_{\mathfrak{gr}^0}^{(d+1)}$ -*covanishing* module. In consequence, the above adjunction is the d -th polynomial approximation of \mathfrak{gr} -modules considered in the literature.

Recall that polynomial degrees are characterized by the equation (20). By the above corollary, (for C having finite products) a polynomial L_C -module with polynomial degree $\leq d$ is nothing but a module over the monad $L_C/\mathbb{I}_C^{(d+1)}$. In other words, the polynomial degree condition is encoded into $L_C/\mathbb{I}_C^{(d+1)}$. We give an explicit description of $L_C/\mathbb{I}_C^{(d+1)}$ for specific C in Theorems 9.1 and 15.4.

Before we leave this section, we give a simple application of the previous results by studying the 1st polynomial ideal $\mathbb{I}_C^{(1)}$.

Definition 7.17. Let C_0 be the quotient category of C by identifying any morphisms from X to Y for every object X, Y . Denote by $q_C : C \rightarrow C_0$ the canonical functor.

Proposition 7.18. *The induced monad homomorphism $L_C \rightarrow L_{C_0}$ gives an isomorphism of monads:*

$$L_C/\mathbb{I}_C^{(1)} \rightarrow L_{C_0}.$$

Proof. Recall that $\pi_X^1 = \text{id}_X - \eta_X \circ \varepsilon_X$. Hence, for $f \in \mathcal{C}(X, Y) \subset \mathbf{L}_C(Y, X)$, we have $f = f \circ \text{id}_X = f \circ \eta_X \circ \varepsilon_X = \eta_Y \circ \varepsilon_X$ modulo $\mathbf{I}_{C_0}(Y, X)$. Hence, the quotient morphism $\mathbf{L}_C \rightarrow \mathbf{L}_{C_0}$ induces $\mathbf{L}_C/\mathbf{I}_C^{(1)} \rightarrow \mathbf{L}_{C_0}$. Based on this observation, one may also verify that the assignment of trivial morphisms in \mathcal{C} induces the inverse of $\mathbf{L}_C/\mathbf{I}_C^{(1)} \rightarrow \mathbf{L}_{C_0}$. \square

The application of this proposition to Corollary 7.15 leads to the following statement.

Corollary 7.19. *We have an equivalence of categories ${}_{\mathbf{L}_C}\mathbf{Mod}^{\leq 0} \simeq {}_{\mathbf{L}_{C_0}}\mathbf{Mod}$.*

7.4. Internalizers for analytic modules. Recall the filtration of a left \mathbf{L}_C -module \mathbf{M} presented by Corollary 7.12. A left \mathbf{L}_C -module is *analytic* if the filtration converges to itself. In this section, we prove that there exists an \mathbf{L}_C -internalizer of analytic \mathbf{L}_C -modules. In contrast to the case of polynomial modules, such a left ideal can not be a two-sided ideal in general.

Definition 7.20. For $\mathbf{M} \in {}_{\mathbf{L}_C}\mathbf{Mod}$, we define $\mathbf{M}^\omega := \varinjlim_d \mathbf{V}_{\mathbf{I}_C^{(d)}}(\mathbf{M})$. A left \mathbf{L}_C -module \mathbf{M} is *analytic* if $\mathbf{M} = \mathbf{M}^\omega$. Let $\mathcal{S}_C^\omega \subset {}_{\mathbf{L}_C}\mathcal{M}\text{od}$ be the subclass of isomorphism classes of analytic \mathbf{L}_C -modules.

Definition 7.21. Let $\nu : \mathcal{X} \rightarrow \mathbb{N}$ be a map. We define $\mathbf{I}_C^\nu \subset \mathbf{L}_C$ by

$$\mathbf{I}_C^\nu(Y, X) := \mathbf{I}_C^{(\nu(X))}(Y, X), \quad X, Y \in \mathcal{X}.$$

This is proved to be a left ideal below, so this is referred to as the ν -*analyticity ideal*.

Lemma 7.22. *The submodule $\mathbf{I}_C^\nu \subset \mathbf{L}_C$ is a left ideal.*

Proof. Fix $X \in \mathcal{X}$ and put $d = \nu(X)$. For $Y, Z \in \mathcal{X}$, if $f \in \mathbf{L}_C(Z, Y)$ and $g \in \mathbf{I}_C^\nu(Y, X)$, then we have $g \circ f \in \mathbf{I}_C^\nu(Z, X)$ since $\mathbf{I}_C^{(d)} \subset \mathbf{L}_C$ is a left ideal. \square

Definition 7.23. For a map $\nu : \mathcal{X} \rightarrow \mathbb{N}$, we define $\text{sup}(\nu)$ to be the supremum of the image of ν .

Lemma 7.24. *Let \mathbf{M} be a left \mathbf{L}_C -module. The image of the \mathbf{L}_C -action $\mathbf{L}_C \otimes \mathbf{V}_{\mathbf{I}_C^\nu}(\mathbf{M}) \rightarrow \mathbf{M}$ coincides with $\mathbf{V}_{\mathbf{I}_C^{(\text{sup}(\nu))}}(\mathbf{M})$ if $\text{sup}(\nu) < \infty$; and \mathbf{M}^ω otherwise.*

Proof. Let $X, Y \in \mathcal{X}$. We unpack the action index-wise: $\mathbf{L}_C(Y, X) \otimes (\mathbf{V}_{\mathbf{I}_C^\nu}(\mathbf{M}))(X) \rightarrow \mathbf{M}(Y)$. By definition, this map is equal to

$$\mathbf{L}_C(Y, X) \otimes (\mathbf{V}_{\mathbf{I}_C^{(\nu(X))}}(\mathbf{M}))(X) \rightarrow \mathbf{M}(Y).$$

The image of this map is contained in $(\mathbf{V}_{\mathbf{I}_C^{(\nu(X))}}(\mathbf{M}))(Y)$ since $\mathbf{I}_C^{(\nu(X))} \subset \mathbf{L}_C$ is a two-sided ideal by Lemma 7.8. Moreover, if $X = Y$, the image coincides with $(\mathbf{V}_{\mathbf{I}_C^{(\nu(X))}}(\mathbf{M}))(Y)$. Hence, the image of $(\mathbf{L}_C \otimes \mathbf{V}_{\mathbf{I}_C^\nu}(\mathbf{M}))(Y) \rightarrow \mathbf{M}(Y)$ is $(\mathbf{V}_{\mathbf{I}_C^{(\text{sup}(\nu))}}(\mathbf{M}))(Y)$ if $\text{sup}(\nu) < \infty$; and $\mathbf{M}^\omega(Y)$ otherwise. \square

Theorem 7.25. *Let $\nu : \mathcal{X} \rightarrow \mathbb{N}$ be a map such that $\text{sup}(\nu) = \infty$. The left ideal \mathbf{I}_C^ν is an \mathbf{L}_C -internalizer of \mathcal{S}_C^ω . In other words, a left \mathbf{L}_C -module \mathbf{M} is analytic if and only if it is \mathbf{I}_C^ν -vanishingly generated.*

Proof. It suffices to prove that $\mathbf{M} \in {}_{\mathbf{L}_C}\mathbf{Mod}$ is analytic if and only if it is \mathbf{I}_C^ν -vanishingly generated. Recall that, by definition, the \mathbf{L}_C -module \mathbf{M} is \mathbf{I}_C^ν -vanishingly generated if the image of the action in Lemma 7.24 is an epimorphism. It holds if and only if $\mathbf{M} = \mathbf{M}^\omega$. \square

The subclass $\mathcal{S}_C^\omega \subset {}_{\mathbf{L}_C}\mathcal{M}\text{od}$ does not have a core \mathbf{L}_C -internalizer in general:

Proposition 7.26. *If there exists $X \in \mathcal{X}$ such that $\mathbf{I}_C^{(\bullet)}(-, X)$ gives a strictly descending filtration of $\mathbf{L}_C(-, X)$, i.e. for each $d \in \mathbb{N}$, there exists $Y \in \mathcal{X}$*

$$\mathbf{I}_C^{(d+1)}(Y, X) \subsetneq \mathbf{I}_C^{(d)}(Y, X),$$

then \mathcal{S}_C^ω has no core L_C -internalizer.

Proof. The contraposition of Proposition 6.22 proves the statement. \square

In Theorems 9.5 and 15.6, we give examples satisfying this assumption.

In what follows, we present a partial computation of the eigenmonad by the ν -analyticity ideal.

Proposition 7.27. *For $X, Y \in \mathcal{X}$, if $\nu(Y) \geq \nu(X)$, then we have an isomorphism*

$$(E_{L_C}(\mathbb{I}_C^\nu))(Y, X) \cong (L_C/I^{(\nu(X))})(Y, X).$$

Proof. We first compute the idealizer monad of \mathbb{I}_C^ν . By Lemma 7.22, $(I_{L_C}(\mathbb{I}_C^\nu))(Y, X)$ consists of $f \in L_C(Y, X)$ such that $\mathbb{I}_C^\nu \circ f \subset \mathbb{I}_C^\nu(-, X)$ (see Remark 5.10). By the definition of \mathbb{I}_C^ν , this is equivalent with the condition that, for $Z \in \mathcal{X}$, $\mathbb{I}_C^{(\nu(Y))}(Z, Y) \circ f \subset \mathbb{I}_C^{(\nu(X))}(Z, X)$. On the one hand, by Lemma 7.8, we have $\mathbb{I}_C^{(\nu(Y))}(Z, Y) \circ f \subset \mathbb{I}_C^{(\nu(Y))}(Z, X)$. By the assumption that $\nu(Y) \geq \nu(X)$ and Proposition 7.10, we see that $\mathbb{I}_C^{(\nu(Y))}(Z, X) \subset \mathbb{I}_C^{(\nu(X))}(Z, X)$. Hence, we obtain $(I_{L_C}(\mathbb{I}_C^\nu))(Y, X) = L_C(Y, X)$. Therefore, we have

$$(E_{L_C}(\mathbb{I}_C^\nu))(Y, X) \cong L_C(Y, X)/I^\nu(Y, X) \cong L_C(Y, X)/I^{(\nu(X))}(Y, X).$$

\square

7.5. Comparison with augmentation ideals. In this section, we investigate polynomial ideals by requiring additional assumptions on the category C . We show that a polynomial ideal for C is identified with an *augmentation ideal* of a certain monoid algebra. We assume the following hypothesis on C :

- (P) C is a PROP whose monoidal structure is given by (categorical) products and $0 \in C$ is a zero object. Here, a PROP means a strict symmetric monoidal category, whose set of objects is \mathbb{N} and whose monoidal structure on objects is given by the addition of natural numbers [14].
- (M) The object $1 \in C$ is a monoid object. This hypothesis yields following structures. Let $\nabla : (1 + 1) \rightarrow 1$ be the multiplication on the monoid object 1 . The set $C_n := C(n, 1)$ has a monoid structure: for $f, g \in C_n$, we define

$$f \cdot g := \nabla \circ (f \times g) \circ \Delta_n$$

where $\Delta_n : n \rightarrow (n + n) = 2n$ is the diagonal map. The zero morphism $e : n \rightarrow 1$, i.e. the unique morphism which factors through 0 , gives the unit of the monoid of C_n .

- (G) Note that every morphism $\xi : n \rightarrow m$ in C induces a *monoid homomorphism* $\xi^* : C_m \rightarrow C_n$ by precomposition. Let $p_k^{m,n} : nm \rightarrow m$ be the k -th projection from $nm = \overbrace{m + \cdots + m}^n$ to m . We assume that $\bigcup_{k=1}^n (p_k^{1,n})^*(C_1)$ generates the monoid C_n where $(p_k^{1,n})^* : C_1 \rightarrow C_n$ is the precomposition.

The assumption (P) implies that C is subject to the conditions in section 7.

Example 7.28. The opposite category gr° of finitely generated free groups satisfies these assumptions. The monoid C_n is nothing but the underlying monoid of the free group F_n .

Example 7.29. The opposite category fr_R° of free R -modules gives another example. The monoid C_n is equal to the underlying (abelian) monoid of R^n .

A monoid G induces an algebra $\mathbb{k}[G]$ over \mathbb{k} which we call the *monoid algebra*. It is the \mathbb{k} -module generated by G , and its multiplication is induced by the monoid structure of G . The *augmentation ideal* $I_{\mathbb{k}}(G)$ is defined to be the two-sided ideal of the monoid algebra $\mathbb{k}[G]$ generated by $(g - h)$ for $g, h \in G$. Denote by $I_{\mathbb{k}}(G)^d$ the d -th power of the ideal $I_{\mathbb{k}}(G)$.

From now on, we fix $n, m, d \in \mathbb{N}$. The hypothesis (P) on C implies that we have $C(n, m) \cong C(n, 1)^{\times m} = C_n^{\times m}$. This induces an isomorphism $\vartheta : \mathbb{k}[C_n^{\times m}] \rightarrow L_C(m, n)$. Under this identification, the following statement holds:

Theorem 7.30. *The isomorphism $\vartheta : \mathbb{k}[C_n^{\times m}] \rightarrow L_C(m, n)$ gives*

$$\vartheta \left(\mathbb{I}_{\mathbb{k}} (C_n^{\times m})^d \right) = \mathbb{I}_C^{(d)}(m, n).$$

Definition 7.31. For C subject to the aforementioned hypothesis, we define an $(\mathbb{N} \times \mathbb{N})$ -indexed module \mathcal{P}_C^d to be

$$\mathcal{P}_C^d(m, n) := \mathbb{k}[C_n^{\times m}] / \mathbb{I}_{\mathbb{k}} (C_n^{\times m})^{d+1}, \quad n, m \in \mathbb{N}.$$

This is proved to be a monad in $\text{Mod}_{\mathbb{k}}$ on \mathbb{N} , as shown in Corollary 7.32, and this is called the *d-th Passi monad for C*.

Corollary 7.32. *The underlying indexed module of the eigenmonad $E_{L_C}(\mathbb{I}_C^{(d+1)})$ of L_C by $\mathbb{I}_C^{(d+1)}$ is isomorphic to \mathcal{P}_C^d .*

Proof. The polynomial ideal $\mathbb{I}_C^{(d+1)}$ is a two-sided ideal by Lemma 7.8, so that the associated eigenmonad is isomorphic to the quotient $L_C / \mathbb{I}_C^{(d+1)}$. By applying Theorem 7.30, we obtain

$$\left(L_C / \mathbb{I}_C^{(d+1)} \right) (m, n) = L_C(m, n) / \mathbb{I}_C^{(d+1)}(m, n) \cong \mathbb{k}[C_n^{\times m}] / \mathbb{I}_{\mathbb{k}} (C_n^{\times m})^{d+1} = \mathcal{P}_C^d(m, n).$$

□

Remark 7.33. For $C \in \{\text{gr}^0, \text{fr}_R^0\}$, polynomial L_C -modules are characterized by finite data indexwise. Indeed, we will prove that \mathcal{P}_C^d has a finite rank indexwise (see Theorems 9.1 and 15.4). In those cases, for $m, n \in \mathbb{N}$, we can choose finitely many elements $v_1, \dots, v_r \in L_C(m, n)$ which induce a basis of $\left(E_{L_C}(\mathbb{I}_C^{(d+1)}) \right) (m, n)$ (recall that we have $E_{L_C}(\mathbb{I}_C^{(d+1)}) \cong L_C / \mathbb{I}_C^{(d+1)}$). By Theorem 7.13 and Corollary 7.32, for a left L_C -module \mathbb{M} with $\text{deg} \leq d$, the map given by the L_C -action on \mathbb{M} is uniquely determined by $\mathbb{M}(v_1), \dots, \mathbb{M}(v_r) \in \text{hom}_{\mathbb{k}}(\mathbb{M}(n), \mathbb{M}(m))$:

$$L_C(m, n) = \mathbb{k}C(n, m) \xrightarrow{\mathbb{M}} \text{hom}_{\mathbb{k}}(\mathbb{M}(n), \mathbb{M}(m)).$$

In terms of function theory, v_1, \dots, v_r play a role of *monomials*, and $\mathbb{M}(v_1), \dots, \mathbb{M}(v_r)$ are treated as the associated *coefficients*. See also Corollary 1.12.

The remaining part of this section is dedicated to the proof of Theorem 7.30.

7.5.1. Key lemmas. In this section, we give another description of $\mathbb{I}_C^{(d)}(m, n)$. This is proved to be \tilde{J}_{W_1} in Definition 7.41.

We choose a generator set W_1 of the monoid C_1 . By the hypothesis (G) on C , the union

$$W_n := \bigcup_{k=1}^n (p_k^{1,n})^*(W_1)$$

generates the monoid C_n . We assume that

$$(24) \quad e \notin W_1.$$

One may consider a typical choice $W_1 = C_1 \setminus \{e\}$.

Notation 7.34. We denote by $t_k^{n,d} : n \rightarrow nd = \overbrace{(n + \dots + n)}^d$ the k -th insertion.

Note that $(p_k^{1,n})^* : C_1 \rightarrow C_n$ is a split homomorphism with a retract $(\iota_k^{n,d})^*$. Hence, (24) implies

$$(25) \quad e \notin W_n.$$

Proposition 7.35. *For $x \in W_n$, there exists a unique $1 \leq k \leq n$ such that $x \in (p_k^{1,n})^*(W_1)$. For such k , we have*

$$(\iota_l^{1,n})^*(x) \begin{cases} \neq e & (l = k), \\ = e & (l \neq k). \end{cases}$$

Proof. By the definition of W_n , there exist some $1 \leq k \leq n$ and $y \in W_1$ such that $x = (p_k^{1,n})^*(y)$. The projection $p_k^{1,n}$ splits by the insertion $\iota_k^{1,n}$; and $p_k^{1,n} \circ \iota_k^{1,n}$ is the zero morphism if $l \neq k$. Hence, we obtain $(\iota_l^{1,n})^*(x) = (p_k^{1,n} \circ \iota_l^{1,n})^*(y) = e$ if $k \neq l$; and $(\iota_k^{1,n})^*(x) = y$. By the assumption (24), we have $(\iota_k^{1,n})^*(x) \neq e$.

If there exists $k' \neq k$ such that $x \in (p_{k'}^{1,n})^*(W_1)$, then $(\iota_{k'}^{1,n})^*(x) \neq e$ gives a contradiction by the above result. \square

Notation 7.36. In Proposition 7.35, we denote by $\partial(x) := k$. This gives a map $\partial : W_n \rightarrow \underline{n}$.

Definition 7.37. Let S be a finite toset (totally ordered set). For a set X , let X^S be the set of maps from S to X . We define a map

$$\varphi : (W_n \times \underline{d})^S \rightarrow W_{nd}^S$$

For a map $f = (f_1, f_2) : S \rightarrow W_n \times \underline{d}$, we define a map $\varphi(f) : S \rightarrow W_{nd}$ as

$$(\varphi(f))(s) := (p_{f_2(s)}^{n,d})^*(f_1(s)).$$

The inverse of φ (see the lemma below) is given as follows. Let $\lceil r \rceil$ be the least integer greater than or equal to $r \in \mathbb{R}$. For $g \in W_{nd}^S$, we define $\varphi^{-1}(g) = (f_1, f_2) : S \rightarrow W_n \times \underline{d}$ as follows:

$$f_2(s) := \lceil \frac{\partial(g(s))}{n} \rceil,$$

$$f_1(s) := (\iota_{f_2(s)}^{n,d})^*(g(s)).$$

Lemma 7.38. *For $1 \leq a \leq d$, we have the following:*

$$\bigcup_{l=1}^n (p_{l+(a-1)n}^{nd,1})^*(W_1) = (p_a^{n,d})^*(W_n).$$

Proof.

$$\begin{aligned} (p_a^{n,d})^*(W_n) &= (p_a^{n,d})^* \left(\bigcup_{k=1}^n (p_k^{1,n})^*(W_1) \right), \\ &= \bigcup_{k=1}^n (p_k^{1,n} \circ p_a^{n,d})^*(W_1), \\ &= \bigcup_{l=1}^n (p_{l+(a-1)n}^{nd,1})^*(W_1). \end{aligned}$$

\square

Lemma 7.39. *The maps φ, φ^{-1} are inverses of each other.*

Proof. Let $f = (f_1, f_2) : S \rightarrow W_n \times \underline{d}$ be a map and put $g = \varphi(f)$. By definitions, we have

$$\begin{aligned} (\iota_{f_2(s)}^{n,d})^*(g(s)) &= (\iota_{f_2(s)}^{n,d})^* \left((p_{f_2(s)}^{n,d})^*(f_1(s)) \right), \\ &= (p_{f_2(s)}^{n,d} \circ \iota_{f_2(s)}^{n,d})^*(f_1(s)) = f_1(s). \end{aligned}$$

In particular, (25) implies $(\iota_{f_2(s)}^{n,d})^*(g(s)) \neq e$, so that $(f_2(s) - 1)n < \partial(g(s)) \leq f_2(s)n$ by a slight refinement of Proposition 7.35. Hence, we obtain the following:

$$\left\lceil \frac{\partial(g(s))}{n} \right\rceil = f_2(s).$$

Thus, we have $\varphi^{-1}(\varphi(f)) = f$.

We now prove that $\varphi(\varphi^{-1}(g)) = g$ for $g \in W_{nd}^S$. Put $f = \varphi^{-1}(g) = (f_1, f_2)$. Then $(p_{f_2(s)}^{n,d})^*(f_1(s)) = (p_{f_2(s)}^{n,d})^* \left((\iota_{f_2(s)}^{n,d})^*(g(s)) \right) = \left(\iota_{f_2(s)}^{n,d} \circ p_{f_2(s)}^{n,d} \right)^*(g(s))$. By the definition of f_2 , we have $(f_2(s) - 1)n < \partial(g(s)) \leq f_2(s)n$. Thus, $g(s) \in \bigcup_{l=1}^n (p_{l+(f_2(s)-1)n}^{nd,1})^*(W_1)$. Hence, $g(s) \in (p_{f_2(s)}^{n,d})^*(W_n)$ by Lemma 7.38. Hence, we obtain

$$\left(\iota_{f_2(s)}^{n,d} \circ p_{f_2(s)}^{n,d} \right)^*(g(s)) = g(s).$$

□

Notation 7.40. Let G be a monoid. For a finite toset S and a map $f : S \rightarrow G$, we define

$$\mu(f) := \begin{cases} 1_G & \text{if } S = \emptyset, \\ a_1 a_2 \cdots a_k & \text{if } S = \{a_1 \leq a_2 \leq \cdots \leq a_k\}. \end{cases}$$

Definition 7.41. Let S be a finite toset and $f = (f_1, f_2, f_3) : S \rightarrow W_n \times \underline{d} \times \underline{m}$ be a map. We define

$$\tilde{\kappa}_{S,f} := \sum_{T \subset \underline{d}} (-1)^{|T|} \cdot \left(\bigotimes_{k=1}^m \mu \left(f_1|_{f_2^{-1}(T) \cap f_3^{-1}(k)} \right) \right) \circ \Delta_n^{(m)} \in L_C(m, n).$$

Here, $\Delta_n^{(m)} \in C(n, nm) \subset L_C(nm, n)$ is the m -fold diagonal morphism $n \rightarrow \overbrace{(n + \cdots + n)}^m$. By using this, we define \tilde{J}_{W_1} to be the submodule of $L_C(m, n)$ generated by any $\tilde{\kappa}_{S,f}$.

Let us apply Remark 7.5 to C . The characterization of D_X^T leads to the following:

Lemma 7.42. Let $f = (f_1, f_2) \in (W_n \times \underline{d})^S$. For $T \subset \underline{d}$, we have

$$\mu(f_1|_{f_2^{-1}(T)}) = \mu(\varphi(f)) \circ D_n^T$$

Proof. Let $g := \varphi(f)$. For any morphism $\xi : s \rightarrow t$, the induced map $\xi^* : C_t \rightarrow C_s$ is a monoid homomorphism so that $\xi^*(\mu(z)) = \mu(\xi^* \circ z)$ for $z \in C_t^S$. To prove our assertion, by the definition of φ , it suffices to check

$$g(s) \circ D_n^T = \begin{cases} f_1(s) & (s \in f_2^{-1}(T)), \\ e & (\text{otherwise}), \end{cases}$$

for each $s \in S$. In fact, it follows from the definition $g(s) := f_1(s) \circ p_{f_2(s)}^{n,d}$ and (22). □

Lemma 7.43. Let $f = (f_1, f_2, f_3) : S \rightarrow W_n \times \underline{d} \times \underline{m}$ be a map and $g = \varphi(f_1, f_2) \in W_{nd}^S$. For $T \subset \underline{d}$, we have

$$\left(\bigotimes_{k=1}^m \mu \left(f_1|_{f_2^{-1}(T) \cap f_3^{-1}(k)} \right) \right) \circ \Delta_n^{(m)} = \left(\bigotimes_{k=1}^m \mu \left(g|_{f_3^{-1}(k)} \right) \right) \circ \Delta_{nd}^{(m)} \circ D_n^T$$

Proof. By the assumption that the PROP structure on C is given by (categorical) products, it suffices to verify the equality after composing the projection $p_k^{1,m} : m \rightarrow 1$ for each $k \in \underline{m}$:

$$\mu \left(f_1|_{f_2^{-1}(T) \cap f_3^{-1}(k)} \right) = \mu \left(g|_{f_3^{-1}(k)} \right) \circ D_n^T.$$

This is proved by applying Lemma 7.42 to $f|_{f_3^{-1}(k)} \in (W_n \times \underline{d})^{f_3^{-1}(k)}$ by replacing S with $f_3^{-1}(k)$. \square

Lemma 7.44. *We have*

$$(26) \quad \tilde{J}_{W_1} = \mathbf{I}_C^{(d)}(m, n).$$

Proof. We first show that $\mathbf{I}_C^{(d)}(m, n) \subset \tilde{J}_{W_1}$. It suffices to prove that $\rho \circ \pi_n^d \in \tilde{J}_{W_1}$ for an arbitrary $\rho \in C(nd, m) \subset L_C(m, nd)$. By $C(nd, m) \cong C(nd, 1)^{\times m}$, we have $\rho = (w_1 \times \cdots \times w_m) \circ \Delta_{nd}^{(m)}$ for some $w_j \in C(nd, 1)$. For $N \in \mathbb{N}$, any element of C_N is equal to $\mu(h)$ for a finite toset S and a map $h : S \rightarrow W_N$ by the hypothesis (G) on C and the assumption that W_1 is a generator set of C_1 . Hence, for each $1 \leq i \leq m$, we may choose a finite toset S_i and a map $g_i : S_i \rightarrow W_{nd}$ such that $\mu(g_i) = w_i$. Let $S = \sqcup_{i=1}^m S_i$ (with the total ordering such that $x \leq y$ if $x \in S_i$ and $y \in S_{i+1}$) and $g := \sqcup_{i=1}^m g_i : S \rightarrow W_{nd}$. Let $f_3 : S \rightarrow \underline{m}$ be the grading, i.e. $f_3(s) = k$ if $s \in S_k$. Let $f = (f_1, f_2) := \varphi^{-1}(g)$. By the definitions, we have $w_k = \mu \left(g|_{f_3^{-1}(k)} \right)$. Recall the equivalent definition of π_n^d in Definition 7.4. By Lemma 7.39, we have $g = \varphi(f)$. We now apply Lemma 7.43:

$$\begin{aligned} \rho \circ \pi_n^d &= \left(\bigotimes_{k=1}^m \mu \left(g|_{f_3^{-1}(k)} \right) \right) \circ \Delta_{nd}^{(m)} \circ \sum_{T \subset \underline{d}} (-1)^{d-|T|} \cdot D_n^T, \\ &= \sum_{T \subset \underline{d}} (-1)^{d-|T|} \cdot \left(\bigotimes_{k=1}^m \mu \left(f_1|_{f_2^{-1}(T) \cap f_3^{-1}(k)} \right) \right) \circ \Delta_n^{(m)}. \end{aligned}$$

Therefore, we obtain $\rho \circ \pi_n^d = \tilde{\kappa}_{S,f} \in \tilde{J}_{W_1}$.

Conversely, we prove that $\tilde{J}_{W_1} \subset \mathbf{I}_C^{(d)}(m, n)$. Let S be a finite toset and $f = (f_1, f_2, f_3) : S \rightarrow W_n \times \underline{d} \times \underline{m}$ be a map. It is sufficient to prove an existence of $\rho \in C(nd, m)$ such that $\rho \circ \pi_n^d = \tilde{\kappa}_{S,f}$. Let $g = \varphi(f_1, f_2) \in W_{nd}^S$. We set ρ to be $\left(\bigotimes_{k=1}^m \mu \left(g|_{f_3^{-1}(k)} \right) \right) \circ \Delta_{nd}^{(m)} : nd \rightarrow m$. Then, by Lemma 7.43 again, we obtain $\rho \circ \pi_n^d = \tilde{\kappa}_{S,f}$. \square

7.5.2. Proof of Theorem 7.30. This section aims to give formal results on a generalization of augmentation ideals (see Theorem 7.51). These are applied to prove Theorem 7.30.

We fix an algebra A over \mathbb{k} with unit 1_A . The monoid algebra $\mathbb{k}[C_n^{\times m}]$ is a typical example of A . We extend Notation 7.40 to the algebra A . For instance, if $a_1, a_2, \dots, a_n \in A$, we define

$$\prod_{i=1}^n a_i := a_1 a_2 \cdots a_{n-1} a_n \in A.$$

Definition 7.45. Let $Y \subset A$ be a subset and $d \in \mathbb{N}$.

- We define $I_{Y,d}$ to be the *two-sided ideal* generated by $\prod_{i=1}^d (y_i - 1_A)$ where all the y_i 's run over Y .
- We define $\tilde{I}_{Y,d}$ to be the *submodule* generated by $\prod_{i=1}^k (y_i - 1_A)$ where $k \geq d$ and all the y_i 's run over Y .

Example 7.46. If G is a monoid and $A = \mathbb{k}[G]$, then we have $I_{G,d} = \tilde{I}_{G,d} = \mathbf{I}_{\mathbb{k}}(G)^d$.

Lemma 7.47. *If the algebra A is generated by $Y \subset A$, then we have $I_{Y,d} = \tilde{I}_{Y,d}$.*

Proof. It is obvious $\tilde{I}_{Y,d} \subset I_{Y,d}$. On the one hand, every element of $I_{Y,d}$ is a linear combination of some products

$$\prod_{i=1}^k z_i \cdot \prod_{i=1}^d (y_i - 1_A) \cdot \prod_{i=1}^l z'_i$$

where $z_i, y_i, z'_i \in Y$. We iteratively apply $z = (z - 1_A) + 1_A$ to expand both sides $\prod_{i=1}^k z_i$ and $\prod_{i=1}^l z'_i$. This leads to $I_{Y,d} \subset \tilde{I}_{Y,d}$. \square

Definition 7.48. For a subset $Y \subset A$, a d -family in Y is a triple of a finite toset S , a map $f = (f_1, f_2) : S \rightarrow Y \times \underline{d}$. For such a d -family, we define

$$\kappa_{S,f} := \sum_{T \subset \underline{d}} (-1)^{|T|} \mu(f_1|_{f_2^{-1}(T)}) \in A.$$

Here, we regard $f_2^{-1}(T)$ as an ordered subset of S .

Proposition 7.49. Suppose that $f_2 : S \rightarrow \underline{d}$ preserves the order. We have

$$\kappa_{S,f} = \prod_{i=1}^d \left(\mu(f_1|_{f_2^{-1}(i)}) - 1_A \right).$$

Proof. For a subset $T = \{t_1 \leq \dots \leq t_k\} \subset S$, we have $\mu(f_1|_{f_2^{-1}(T)}) = \prod_{i=1}^k \mu(f_1|_{f_2^{-1}(i)})$ since f_2 preserves the order. It leads to our statement. \square

Definition 7.50. Let $Y \subset A$ be a subset and $d \in \mathbb{N}$.

- We define $J_{Y,d}$ to be the *two-sided ideal* generated by $\kappa_{S,f}$ for (S, f) a d -family in Y such that $f_2 : S \rightarrow \underline{d}$ is an order-preserving map.
- We define $\tilde{J}_{Y,d}$ to be the *submodule* generated by $\kappa_{S,f}$ for (S, f) a d -family in Y .

Theorem 7.51. Let $Y \subset A$ be a subset and $d \in \mathbb{N}$. If Y generates the algebra A , then we have

$$I_{Y,d} = \tilde{I}_{Y,d} = J_{Y,d} = \tilde{J}_{Y,d}.$$

Proof. We defer the proof to the following lemmas. \square

Lemma 7.52. We have $I_{Y,d} = J_{Y,d}$ for $d \in \mathbb{N}$.

Proof. We first prove that $I_{Y,d} \subset J_{Y,d}$, or equivalently $\prod_{i=1}^d (y_i - 1_A) \in J_{Y,d}$ for $y_i \in Y$. Let $S = \underline{d}$ be the ordered set and $f_2 : S \rightarrow \underline{d}$ be the identity which is an order-preserving map. We define $f_1 : S = \underline{d} \rightarrow Y$ to be $f_1(i) = y_i$ for any i . Put $f := (f_1, f_2) : S \rightarrow Y \times \underline{d}$. By definitions, we have $\prod_{i=1}^d (y_i - 1_A) = \kappa_{S,f} \in J_{Y,d}$.

We now prove that $J_{Y,d} \subset I_{Y,d}$. Let (S, f) be a d -family in Y such that $f_2 : S \rightarrow \underline{d}$ is an order-preserving map. As we observed in Proposition 7.49, $\kappa_{S,f} \in A$ is given by a d -fold product of the differences between a product of elements in Y and the unit 1_A . It turns out that such a difference between a *product of elements* in Y and 1_A is equal to linear combination of some differences between *one element* in Y and 1_A . It is deduced by iteratively applying the formula below:

$$xy - 1_A = (x - 1_A)(y - 1_A) + (x - 1_A) + (y - 1_A), \quad x, y \in A.$$

\square

Lemma 7.53. If the algebra A is generated by Y , then we have $\tilde{J}_{Y,d} = J_{Y,d}$ for $d \in \mathbb{N}$.

Proof. We prove that $J_{Y,d} \subset \tilde{J}_{Y,d}$. The set Y generates the algebra A if and only if A is a module generated by $\mu(f)$ where f is a map from a finite toset S to Y . Hence, by Proposition 7.49, $J_{Y,d}$ is the submodule generated by $\mu(g) \cdot \prod_{i=1}^d \tilde{\mu}(f_i) \cdot \mu(h)$ where f_i, g, h run over all

the maps from some finite toset to Y where we set $\tilde{\mu}(f) := \mu(f) - 1_A$. It suffices to show that $\mu(g) \cdot \prod_{i=1}^d \tilde{\mu}(f_i) \cdot \mu(h)$ lies in $\tilde{J}_{Y,d}$. For maps $f : S \rightarrow Y$ and $g : S' \rightarrow Y$ with finite tosets S, S' , note that we have $\mu(f \amalg g) = \mu(f) \cdot \mu(g)$ where $f \amalg g : S \amalg S' \rightarrow Y$. By using this fact, we compute as follows:

$$\begin{aligned} \mu(g) \cdot \prod_{i=1}^d \tilde{\mu}(f_i) \cdot \mu(h) &= (\tilde{\mu}(g \amalg f_1) - \tilde{\mu}(g)) \cdot \prod_{i=2}^{d-1} \tilde{\mu}(f_i) \cdot (\tilde{\mu}(f_d \amalg h) - \tilde{\mu}(h)), \\ &= \tilde{\mu}(g \amalg f_1) \cdot \prod_{i=2}^{d-1} \tilde{\mu}(f_i) \cdot \tilde{\mu}(f_d \amalg h) - \tilde{\mu}(g) \cdot \prod_{i=2}^{d-1} \tilde{\mu}(f_i) \cdot \tilde{\mu}(f_d \amalg h) \\ &\quad - \tilde{\mu}(g \amalg f_1) \cdot \prod_{i=2}^{d-1} (\mu(f_i) - 1_A) \cdot \tilde{\mu}(g) + \tilde{\mu}(g) \cdot \prod_{i=2}^{d-1} \tilde{\mu}(f_i) \cdot \tilde{\mu}(h). \end{aligned}$$

By Proposition 7.49 again, this sum lies in $\tilde{J}_{Y,d}$.

We now prove $\tilde{J}_{Y,d} \subset J_{Y,d}$ for all $d \in \mathbb{N}$. If $d = 0$, then $\tilde{J}_{Y,d} = 0 = J_{Y,d}$. It suffices to prove $\tilde{J}_{Y,d} \subset J_{Y,d}$ under the assumption $\tilde{J}_{Y,d-1} \subset J_{Y,d-1}$. Let (S, f) be a d -family in Y and $f = (f_1, f_2) : S \rightarrow Y \times \underline{d}$ as before. We prove $\kappa_{S,f} \in J_{Y,d}$ by using the induction on the order of S . It is clear that it holds when $|S| = 0$. We assume that $|S| > 0$. Let $M_0 \in S$ be the maximum and suppose that $f_2(M_0) = j_0$ for a certain $1 \leq j_0 \leq d$. We set $S' = S \setminus \{M_0\}$. Put $f' := f|_{S'} = (f'_1, f'_2)$. Note that we have

$$\sum_{j_0 \in T \subset \underline{d}} (-1)^{|T|} \mu(f_1|_{f_2^{-1}(T)}) = \sum_{j_0 \in T \subset \underline{d}} (-1)^{|T|} \mu(f'_1|_{(f'_2)^{-1}(T)}) \cdot f_1(M_0),$$

since M_0 is the maximum. Hence, we obtain

$$\begin{aligned} \kappa_{S',f'} \cdot f(M_0) - \kappa_{S,f} &= \sum_{j_0 \notin T \subset \underline{d}} (-1)^{|T|} \left(\mu(f'_1|_{(f'_2)^{-1}(T)}) f_1(M_0) - \mu(f_1|_{f_2^{-1}(T)}) \right), \\ &= \sum_{j_0 \notin T \subset \underline{d}} (-1)^{|T|} \left(\mu(f_1|_{f_2^{-1}(T)}) f_1(M_0) - \mu(f_1|_{f_2^{-1}(T)}) \right), \\ &= \left(\sum_{j_0 \notin T \subset \underline{d}} (-1)^{|T|} \mu(f_1|_{f_2^{-1}(T)}) \right) \cdot (f_1(M_0) - 1_A). \end{aligned}$$

Here, the second equality follows from $j_0 \notin T$. We investigate the left component of this result. Let $\tilde{S} = S \setminus f_2^{-1}(j_0)$ be the ordered subset of S . By choosing a bijection $\underline{d} \setminus \{j_0\} \cong \underline{(d-1)}$, we define $\tilde{f}_2 : \tilde{S} \rightarrow \underline{(d-1)}$ to be the composition $\tilde{S} \xrightarrow{f_2|_{\tilde{S}}} \underline{d} \setminus \{j_0\} \cong \underline{(d-1)}$, and put $\tilde{f}_1 = f_1|_{\tilde{S}}$. These give a $(d-1)$ -family in Y , $\tilde{f} = (\tilde{f}_1, \tilde{f}_2) : \tilde{S} \rightarrow Y \times \underline{(d-1)}$. Then we have

$$\begin{aligned} \sum_{j_0 \notin T \subset \underline{d}} (-1)^{|T|} \mu(f_1|_{f_2^{-1}(T)}) &= \sum_{\tilde{T} \subset \underline{(d-1)}} (-1)^{|\tilde{T}|} \mu(\tilde{f}_1|_{\tilde{f}_2^{-1}(\tilde{T})}), \\ &= \kappa_{\tilde{S}, \tilde{f}}. \end{aligned}$$

By the assumption $\tilde{J}_{Y,d-1} \subset J_{Y,d-1}$, we have $\kappa_{\tilde{S}, \tilde{f}} \in J_{Y,d-1}$. By Lemma 7.52 and 7.47, we see that $\kappa_{\tilde{S}, \tilde{f}} \in \tilde{I}_{Y,d-1}$ since Y generates A . It implies that

$$\kappa_{\tilde{S}, \tilde{f}} \cdot (f_1(M_0) - 1_A) \in \tilde{I}_{Y,d}.$$

By Lemma 7.52 and 7.47 again, $\kappa_{\tilde{S}, \tilde{f}} \cdot (f_1(M_0) - 1_A) \in J_{Y,d}$ so that the previous result leads to $(\kappa_{S',f'} \cdot f(M_0) - \kappa_{S,f}) \in J_{Y,d}$. The hypothesis of the induction (on the order of S) implies

that $\kappa_{S',f'} \in J_{Y,d}$ due to $|S'| < |S|$. Hence, $\kappa_{S',f'} \cdot f(M_0) \in J_{Y,d}$ since $J_{Y,d}$ is a two-sided ideal. Therefore, we obtain $\kappa_{S,f} \in J_{Y,d}$. \square

We now complete the proof of Theorem 7.30 by applying the preceding results. Recall the setting in section 7.5.1. We introduce a set which plays the role of $Y \subset A$ in this section where $A = \mathbb{k}[C_n^{\times m}]$ is the monoid algebra for fixed n, m .

$$Y := \{(\overbrace{e, \dots, e}^{(i-1)}, x, \overbrace{e, \dots, e}^{m-i}) \in C_n^{\times m} \mid x \in W_n, i \in \underline{m}\} \subset A.$$

By the hypothesis (24), we have an obvious bijection $W_n \times \underline{m} \rightarrow Y$. Denote by the induced bijection $h : W_n \times \underline{d} \times \underline{m} \rightarrow Y \times \underline{d}$. Let S be a finite toset and $f : S \rightarrow W_n \times \underline{d} \times \underline{m}$ be a map. Then the composition with h induces a d -family in Y :

$$h_*(f) : S \xrightarrow{f} W_n \times \underline{d} \times \underline{m} \xrightarrow{h} Y \times \underline{d}.$$

The following is immediate from the monoid structure on $C_n^{\times m}$ where $\kappa_{S,h_*(f)}$ and $\tilde{\kappa}_{S,f}$ are defined in Definition 7.48 and Definition 7.41 respectively:

$$(27) \quad \vartheta(\kappa_{S,h_*(f)}) = \tilde{\kappa}_{S,f}.$$

Indeed, if we put $h_*(f) = (g_1, g_2) : S \rightarrow Y \times \underline{d}$, then we have $g_1(s) = (e, \dots, e, f_1(s), e, \dots, e)$ where the $f_3(s)$ -th component is $f_1(s)$; and $f_2(s) = g_2(s)$ for $s \in S$. Hence, we obtain

$$\mu(g_1|_{g_2^{-1}(T)}) = \left(\bigotimes_{k=1}^m \mu(f_1|_{f_2^{-1}(T) \cap f_3^{-1}(k)}) \right) \circ \Delta_n^{(m)}$$

for $T \subset \underline{d}$. By the definitions, this leads to the equation (27), which implies $\tilde{J}_{W_1} = \vartheta(\tilde{J}_{Y,d})$. Therefore, by Theorem 7.51 and Lemma 7.44, the claim leads to

$$\mathbb{I}_C^{(d)}(m, n) = \tilde{J}_{W_1} = \vartheta(I_{Y,d}) = \vartheta\left(\mathbb{I}_{\mathbb{k}}(C_n^{\times m})^d\right).$$

Part 2. Application to free groups

Let us denote the opposite category of finitely generated free groups and group homomorphisms by gr° . There naturally arise modules over gr° in the context of quantum topology [8, 11, 29]. In light of these examples, algebraic properties of modules over gr° were studied well, for instance, in [21, 23, 12]. In this part, we give applications of part 1 to modules over the category gr° . Indeed, the monad of primary interest is L_{gr° defined in Example 4.5. By introducing specific left ideals of this monad L_{gr° , we study the classes of polynomial modules, analytic modules, outer modules and primitively generated modules respectively.

8. NOTATION

We explain our convention for the category gr° . For a nonnegative integer n , we identify the free group generated n elements with n . We explain our notation of morphisms in gr° . Let F_n be the free group generated by n letters x_1, \dots, x_n . For a group homomorphism $\rho : F_m \rightarrow F_n$, we denote the corresponding morphism $n \rightarrow m$ in gr° by

$$(28) \quad [\rho(x_1)|\rho(x_2)|\dots|\rho(x_m)]_n \in \text{gr}^\circ(n, m).$$

This works well since a group homomorphism $\rho : F_m \rightarrow F_n$ is determined by the m -tuple of $(\rho(x_1), \rho(x_2), \dots, \rho(x_m)) \in F_n^{\times m}$.

Our convention for the operad $\mathbf{A}_{\text{ass}^u}$ for associative algebras is given as follows. Recall that we fix letters x_1, x_2, \dots, x_n which are generators of our free group F_n . Let $X_n := \{x_1, x_2, \dots, x_n\}$.

Denote by $\mathbb{k}\langle X_n \rangle$ the noncommutative polynomial algebra generated by X_n over \mathbb{k} . The operad for associative algebras \mathfrak{Ass}^u consists of a submodule $\mathfrak{Ass}^u(n) \subset \mathbb{k}\langle X_n \rangle$, $n \in \mathbb{N}$ which is generated by products of letters in X_n appearing exactly once:

$$\mathfrak{Ass}^u(n) := \bigoplus_{\sigma \in \mathfrak{S}_n} \mathbb{k}x_{\sigma^{-1}(1)}x_{\sigma^{-1}(2)} \cdots x_{\sigma^{-1}(n)}.$$

The \mathfrak{S}_n -action is given by the permutation of letters. The operad operation is induced by substitutions of words.

We elucidate our convention for the Lie operad \mathfrak{Lie} . By using the Lie bracket associated with the algebra $\mathbb{k}\langle X_n \rangle$, we set $\mathfrak{Lie}(n)$ to be the submodule of $\mathbb{k}\langle X_n \rangle$ generated by Lie polynomials [25] which contain each letter in X_n precisely once. Then the Lie operad \mathfrak{Lie} consists of the sequence of $\mathfrak{Lie}(n)$, $n \in \mathbb{N}$ with the operad operation described by the substitution of Lie brackets.

9. POLYNOMIAL IDEALS FOR FREE GROUPS

In this section, we study polynomial ideals for the category \mathfrak{gr}^0 by employing the general results of section 7. There are two principal applications. The first concerns a finiteness of the components of the associated eigenmonad indexwise. This provides a more concrete description of polynomial functors, as elucidated in Remark 7.33. The second application is the study of the class of analytic $L_{\mathfrak{gr}^0}$ -modules.

9.1. Eigenmonads by polynomial ideals. It is readily checked that the category \mathfrak{gr}^0 satisfies the assumptions on C in section 7.5. We derive a familiar aspect of polynomial ideals for \mathfrak{gr}^0 by relating them to augmentation ideals of groups. For $C = \mathfrak{gr}^0$, we have an obvious isomorphism $C_n \cong F_n$ as monoids. Hence, Theorem 7.30 implies the following isomorphism:

$$(29) \quad \mathbb{I}_{\mathbb{k}}(F_n^{\times m})^d \cong \mathbb{I}_{\mathfrak{gr}^0}^{(d)}(m, n).$$

In what follows, we give a computation of the eigenmonad by polynomial ideals. By Corollary 7.32, it suffices to compute the Passi monad $\mathcal{P}_{\mathfrak{gr}^0}^d$.

Theorem 9.1. *For $n, m \in \mathbb{N}$, the module $\mathcal{P}_{\mathfrak{gr}^0}^d(m, n)$ is a free \mathbb{k} -module with rank of*

$$\sum_{k=0}^d n^k \cdot \frac{(m+k-1)!}{k!(m-1)!}.$$

Proof. We defer the proof to later in this section. □

For a group G , let $\text{Gr}(G) := \bigoplus_{k \in \mathbb{N}} \mathbb{I}_{\mathbb{Z}}(G)^k / \mathbb{I}_{\mathbb{Z}}(G)^{k+1}$ be the graded algebra over \mathbb{Z} .

Definition 9.2. Let $m \in \mathbb{N}$. An *ordered m -partition* of d is an m -tuple $\vec{p} = (p_1, p_2, \dots, p_m)$ of natural numbers such that $p_1 + p_2 + \dots + p_m = d$. For such an m -partition, we define the map $Z_{\vec{p}} : (\mathbb{k}^n)^{\otimes d} \rightarrow \mathbb{k}[F_n]^{\otimes m}$ by

$$Z_{\vec{p}} \left(\bigotimes_{j=1}^d e_{k_j} \right) := \prod_{j=1}^{p_1} (x_{k_j} - e) \otimes \prod_{j=1}^{p_2} (x_{k_{p_1+j}} - e) \otimes \cdots \otimes \prod_{j=1}^{p_m} (x_{k_{p_1+\dots+p_{m-1}+j}} - e).$$

By identifying $\mathbb{k}[F_n^{\times m}]$ with $\mathbb{k}[F_n]^{\otimes m}$, we regard $Z_{\vec{p}}$ as a map $(\mathbb{k}^n)^{\otimes d} \rightarrow \mathbb{k}[F_n^{\times m}]$.

Let G, H be groups. Suppose that $\text{Gr}(G), \text{Gr}(H)$ are torsion-free. Then we have $\text{Gr}(G \times H) \cong \text{Gr}(G) \otimes \text{Gr}(H)$, and $\text{Gr}(G \times H)$ is torsion-free [4, Proposition 1.6]. In the application, one may see that $Z_{\vec{p}}$ factors via $\mathbb{I}_{\mathbb{k}}(F_n^{\times m})^d$. Based on this observation, we regard $Z_{\vec{p}}$ as the map from $(\mathbb{k}^n)^{\otimes d}$ to $\mathbb{I}_{\mathbb{k}}(F_n^{\times m})^d$. In the following, we consider the sum of $Z_{\vec{p}}$'s:

Lemma 9.3. *The sum of $Z_{\vec{p}}$'s induces an isomorphism where \vec{p} runs over m -partitions of d :*

$$\bigoplus_{\vec{p}} \mathbb{k}^{n^d} \cong \mathbb{I}_{\mathbb{k}}(\mathbb{F}_n^{\times m})^d / \mathbb{I}_{\mathbb{k}}(\mathbb{F}_n^{\times m})^{d+1}.$$

In particular, this has a basis with order $n^d \cdot \frac{(m+d-1)!}{d!(m-1)!}$.

Proof. Consider $\mathbb{k} = \mathbb{Z}$. By [18, Theorem 6.2, Chapter VIII], we have an isomorphism

$$\begin{aligned} (\mathbb{Z}^n)^{\otimes d} &\cong \mathbb{I}_{\mathbb{Z}}(\mathbb{F}_n)^d / \mathbb{I}_{\mathbb{Z}}(\mathbb{F}_n)^{d+1}, \\ e_{i_1} \otimes e_{i_2} \otimes \cdots \otimes e_{i_d} &\mapsto (x_{i_1} - e)(x_{i_2} - e) \cdots (x_{i_d} - e). \end{aligned}$$

where e_1, e_2, \dots, e_n is the standard basis for \mathbb{Z}^n . This implies that $\text{Gr}(\mathbb{F}_n)$ is free, in particular, torsion-free. By applying [4, Proposition 1.6] recursively, we obtain

$$\text{Gr}_d(\mathbb{F}_n^{\times m}) \cong \bigoplus_{\vec{p}} \bigotimes_{k=1}^m \text{Gr}_{p_k}(\mathbb{F}_n) \cong \bigoplus_{\vec{p}} \bigotimes_{k=1}^m (\mathbb{Z}^n)^{\otimes p_k},$$

where \otimes is the tensor product over \mathbb{Z} . We see that this isomorphism coincides with the sum of $Z_{\vec{p}}$'s in our statement since we have $\bigotimes_{k=1}^m (\mathbb{Z}^n)^{\otimes p_k} \cong \bigotimes_{k=1}^m \mathbb{Z}^{n^{p_k}} \cong \mathbb{Z}^{n^d}$.

Let us consider a general \mathbb{k} . By the previous results, it suffices to prove that the canonical map $\mathbb{k} \otimes_{\mathbb{Z}} (\mathbb{I}_{\mathbb{Z}}(G)^d / \mathbb{I}_{\mathbb{Z}}(G)^{d+1}) \rightarrow \mathbb{I}_{\mathbb{k}}(G)^d / \mathbb{I}_{\mathbb{k}}(G)^{d+1}$ is an isomorphism for $G = \mathbb{F}_n^{\times m}$. This map is already surjective, so that we should show that its kernel, denoted by Q_d , is trivial. The above results imply that the short exact sequence below splits:

$$0 \rightarrow \mathbb{I}_{\mathbb{Z}}(G)^{d+1} \rightarrow \mathbb{I}_{\mathbb{Z}}(G)^d \rightarrow \mathbb{I}_{\mathbb{Z}}(G)^d / \mathbb{I}_{\mathbb{Z}}(G)^{d+1} \rightarrow 0.$$

It implies that the application of the tensor product $\mathbb{k} \otimes_{\mathbb{Z}} (-)$ to this sequence gives a short exact sequence of \mathbb{k} -modules. Let K_d be the kernel of the canonical map $\mathbb{k} \otimes_{\mathbb{Z}} \mathbb{I}_{\mathbb{Z}}(G)^d \rightarrow \mathbb{I}_{\mathbb{k}}(G)^d$. By the definitions, we obtain the following commutative diagram whose columns and the bottom two rows are short exact sequences:

$$\begin{array}{ccccc} K_{d+1} & \longrightarrow & K_d & \longrightarrow & Q_d \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{k} \otimes_{\mathbb{Z}} \mathbb{I}_{\mathbb{Z}}(G)^{d+1} & \longrightarrow & \mathbb{k} \otimes_{\mathbb{Z}} \mathbb{I}_{\mathbb{Z}}(G)^d & \longrightarrow & \mathbb{k} \otimes_{\mathbb{Z}} (\mathbb{I}_{\mathbb{Z}}(G)^d / \mathbb{I}_{\mathbb{Z}}(G)^{d+1}) \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{I}_{\mathbb{k}}(G)^{d+1} & \longrightarrow & \mathbb{I}_{\mathbb{k}}(G)^d & \longrightarrow & \mathbb{I}_{\mathbb{k}}(G) / \mathbb{I}_{\mathbb{k}}(G)^{d+1} \end{array}$$

By the nine lemma, the first row is a short exact sequence for any d . Therefore, it is inductively proved that $K_d \cong 0$ and $Q_d \cong 0$ since we have $K_0 \cong 0$. \square

We now prove Theorem 9.1 by freely using (29). The assertion in the case of $d = 0$ is immediate from Proposition 7.18. Consider the short exact sequence below:

$$0 \rightarrow \mathbb{I}_{\text{gr}^0}^{(k)} / \mathbb{I}_{\text{gr}^0}^{(k+1)} \rightarrow \mathcal{P}_{\text{gr}^0}^k \rightarrow \mathcal{P}_{\text{gr}^0}^{k-1} \rightarrow 0$$

The exactness inductively implies that $\mathcal{P}_{\text{gr}^0}^k$ is (non-canonically) isomorphic to the direct sum $\bigoplus_{k=0}^d \mathbb{I}_{\text{gr}^0}^{(k)} / \mathbb{I}_{\text{gr}^0}^{(k+1)}$. Thus, our assertion follows from (29) and Lemma 9.3. In particular, there exists an isomorphism

$$\mathcal{P}_{\text{gr}^0}^d(m, n) \cong \bigoplus_{0 \leq k \leq d} \bigoplus_{\vec{p}} \mathbb{k}^{n^k}$$

where \vec{p} runs over m -partitions of k .

9.2. Internalizer for analytic modules. By Theorem 7.25, we know that the class of analytic L_{gr° -modules has an L_{gr° -internalizer. In this section, we prove that there exist uncountably many L_{gr° -internalizers and no core L_{gr° -internalizer for analytic modules.

Proposition 9.4. *The assignment of $\mathbf{I}_{\text{gr}^\circ}^\nu$ to a map $\nu : \mathbb{N} \rightarrow \mathbb{N}$ is injective.*

Proof. It basically follows from Remark 7.11. If $\nu \neq \nu'$, then there exists $n \in \mathbb{N}$ such that $\nu(n) \neq \nu'(n)$. Hence, we have $\mathbf{I}_{\text{gr}^\circ}^\nu(-, n) = \mathbf{I}_{\text{gr}^\circ}^{(\nu(n))}(-, n) \neq \mathbf{I}_{\text{gr}^\circ}^{(\nu'(n))}(-, n) = \mathbf{I}_{\text{gr}^\circ}^{\nu'}(-, n)$. \square

The class of polynomial modules has a core internalizer by Theorem 7.13. However, it is not true for analytic L_{gr° -modules:

Theorem 9.5. *The subclass $\mathcal{S}_{\text{gr}^\circ, \omega} \subset L_{\text{gr}^\circ} \text{Mod}$ does not allow a core L_{gr° -internalizer.*

Proof. By using (29), we identify the filtration $\mathbf{I}_{\text{gr}^\circ}^\bullet(1, 1)$ with $\mathbf{I}_{\mathbb{k}}(\mathbf{F}_1)^\bullet$. If we regard the group algebra of \mathbf{F}_1 as the polynomial algebra $\mathbb{k}[t]$, then the d -th power of the augmentation ideal becomes the ideal generated by $(t - 1)^d$. Hence, the filtration $\mathbf{I}_{\mathbb{k}}(\mathbf{F}_1)^\bullet$ is strictly descending, and so is $\mathbf{I}_{\text{gr}^\circ}^\bullet(1, 1)$. We apply Proposition 7.26 to the polynomial ideal filtration of L_{gr° . \square

10. OUTER MODULES OVER FREE GROUPS

A functor $F : \text{gr}^\circ \rightarrow \mathbb{k} \text{Mod}$ is an outer functor if inner automorphisms act trivially on $F(n)$ for all $n \in \mathbb{N}$ [24]. A left L_{gr° -module is *outer* if the corresponding functor from gr° to $\mathbb{k} \text{Mod}$ by the isomorphism in Example 4.16 is an outer functor. In this section, we prove that the class of outer modules has a core internalizer and compute the eigenmonad.

10.1. Core internalizer for outer modules. Let $n, m \in \mathbb{N}$. Consider the *conjugate action* of \mathbf{F}_n on $L_{\text{gr}^\circ}(m, n)$:

$$\text{Ad}_g([w_1|w_2|\cdots|w_m]_n) := [gw_1g^{-1}|gw_2g^{-1}|\cdots|gw_mg^{-1}]_n, \quad \forall g \in \mathbf{F}_n,$$

for $[w_1|w_2|\cdots|w_m]_n \in L_{\text{gr}^\circ}(m, n)$.

Definition 10.1. We define \mathbf{I}^{out} to be the left ideal of the monad L_{gr° generated by $\text{Aut}_g(\text{id}_n) - \text{id}_n$ for $n \in \mathbb{N}$ and $g \in \mathbf{F}_n$.

Proposition 10.2. *The left ideal \mathbf{I}^{out} is a two-sided ideal of the monad L_{gr° .*

Proof. It suffices to prove that $\text{Ad}_g(\text{id}_m) \circ \rho = \rho \pmod{\mathbf{I}^{\text{out}}(m, n)}$ for any $m, n \in \mathbb{N}$, $g \in \mathbf{F}_m$ and $\rho \in \text{gr}^\circ(n, m) \subset L_{\text{gr}^\circ}(m, n)$. Let $h \in \mathbf{F}_n$ be the element such that $[h]_n = [g]_m \circ \rho$. If $\rho = [w_1|w_2|\cdots|w_m]_n$ with $w_j \in \mathbf{F}_m$, then, by the definitions, we have

$$\begin{aligned} \text{Ad}_g(\text{id}_m) \circ \rho &= \text{Ad}_h([w_1|w_2|\cdots|w_m]_n), \\ &= [w_1|w_2|\cdots|w_m]_n \circ \text{Ad}_h([x_1|x_2|\cdots|x_n]_n), \\ &= [w_1|w_2|\cdots|w_m]_n \pmod{\mathbf{I}^{\text{out}}(m, n)}, \\ &= \rho. \end{aligned}$$

\square

Definition 10.3. Let $\mathcal{S}^{\text{out}} \subset L_{\text{gr}^\circ} \text{Mod}$ be the subclass of isomorphism classes of outer left L_{gr° -modules.

In the following, recall Definition 6.17.

Theorem 10.4. *The two-sided ideal \mathbf{I}^{out} of L_{gr° is the unique core L_{gr° -internalizer of \mathcal{S}^{out} . In particular, we have $\text{Ann}(\mathcal{S}^{\text{out}}) = \mathbf{I}^{\text{out}}$.*

Proof. By Proposition 6.6, we see that a left L_{gr° -module \mathbf{M} is \mathbf{I}^{out} -vanishingly generated if and only if we have $\mathbf{V}_{\mathbf{I}^{\text{out}}}(\mathbf{M}) = \mathbf{M}$, which is equivalent with the definition of an outer L_{gr° -module. The last assertion follows from Theorem 6.20. \square

10.2. Eigenmonad and group homology. By Theorem 10.4, the outer property of L_{gr^0} -modules is encoded into the eigenmonad of L_{gr^0} by \mathbf{I}^{out} . In this section, we define an indexed \mathbb{k} -module arising from group homology of free groups. We prove that the eigenmonad by the ideal \mathbf{I}^{out} is identified with that induced by the zeroth group homology.

We give a brief review of group homology. Let G be a group and M be a left G -module over \mathbb{k} . The *group homology* $H_*(G; M)$ of G with coefficients in M is defined to be the left derived functors of the coinvariant assignment [30, section 6]. Here, we use the following concrete model. In fact, the G -module M induces a simplicial \mathbb{k} -module whose r -th component is $\mathbb{k}G^{\otimes r} \otimes M$. The associated homology is the same as $H_*(G; M)$.

In this section, we regard $\mathbb{k}[\mathbf{F}_n^{\times m}]$ as a left \mathbf{F}_n -module by the conjugate action.

Definition 10.5. Let $q \in \mathbb{N}$. We introduce a morphism $H_q : \mathbb{N} \rightsquigarrow \mathbb{N}$ in $\text{Mat}_{\mathbb{k}}$, i.e. an $(\mathbb{N} \times \mathbb{N})$ -indexed \mathbb{k} -module. For $n, m \in \mathbb{N}$, we define

$$H_q(m, n) := H_q(\mathbf{F}_n; \mathbb{k}[\mathbf{F}_n^{\times m}])$$

where the coefficient module is induced by the conjugate action.

Note that the group homology is a functor with respect to the pair (G, M) . A morphism from (G, M) to (G', M') is a pair of group homomorphism $\rho : G \rightarrow G'$ and a linear map $f : M \rightarrow M'$ such that $f(g \cdot x) = \rho(g) \cdot f(x)$ for $g \in G$ and $x \in M$. Such a morphism induces a linear map $H_*(G; M) \rightarrow H_*(G'; M')$ in covariant way.

Proposition 10.6. Let $q \in \mathbb{N}$. The functoriality of group homology induces an $(L_{\text{gr}^0}, L_{\text{gr}^0})$ -bimodule structure on H_q .

Proof. By using the functoriality of group homology, we define the $(L_{\text{gr}^0}, L_{\text{gr}^0})$ -bimodule H_q by extending this assignment. The assignment of $H_q(m, n)$ to $(m, n) \in \mathbb{N} \times \mathbb{N}$ extends to a bilinear functor $H_q : (\mathbb{k}\text{gr}^0)^\circ \times \mathbb{k}\text{gr}^0 \rightarrow \mathbb{k}\text{Mod}$ by the functoriality of group homology. In fact, a morphism $m \rightarrow l$ in gr^0 induces a morphism $(\mathbf{F}_m, \mathbb{k}[\mathbf{F}_m^{\times m}]) \rightarrow (\mathbf{F}_l, \mathbb{k}[\mathbf{F}_l^{\times l}])$ in covariant way; and a morphism $k \rightarrow n$ in gr^0 induces a morphism $(\mathbf{F}_n, \mathbb{k}[\mathbf{F}_n^{\times m}]) \rightarrow (\mathbf{F}_k, \mathbb{k}[\mathbf{F}_k^{\times m}])$ in contravariant way. By the equivalence (12), this bilinear functor induces the $(L_{\text{gr}^0}, L_{\text{gr}^0})$ -bimodule H_q . \square

Definition 10.7. Denote by $L_{\text{gr}^0}\text{Mod}^{\text{out}}$ the full subcategory of outer L_{gr^0} -modules.

Theorem 10.8. We have an isomorphism of $(\mathbb{N} \times \mathbb{N})$ -indexed \mathbb{k} -modules:

$$E_{L_{\text{gr}^0}}(\mathbf{I}^{\text{out}}) \cong H_0.$$

In particular, we have $L_{\text{gr}^0}\text{Mod}^{\text{out}} \simeq_{H_0} \text{Mod}$.

Proof. The zeroth homology $H_0(m, n) = H_0(\mathbf{F}_n; \mathbb{k}[\mathbf{F}_n^{\times m}])$ is the quotient of $\mathbb{k}[\mathbf{F}_n^{\times m}]$ by the relation:

$$[gw_1g^{-1}|gw_2g^{-1}|\cdots|gw_ng^{-1}]_n \equiv [w_1|w_2|\cdots|w_n]_n.$$

On the one hand, by Proposition 10.2, we have $E_{L_{\text{gr}^0}}(\mathbf{I}^{\text{out}}) = L_{\text{gr}^0}/\mathbf{I}^{\text{out}}$ so that $E_{L_{\text{gr}^0}}(\mathbf{I}^{\text{out}})$ is isomorphic to the quotient of $L_{\text{gr}^0}(m, n)$ by $\mathbf{I}^{\text{out}}(m, n)$. Hence, the isomorphism $\mathbb{k}[\mathbf{F}_n^{\times m}] \cong L_{\text{gr}^0}(m, n)$ induces $(E_{L_{\text{gr}^0}}(\mathbf{I}^{\text{out}}))(m, n) \cong H_0(m, n)$. \square

Corollary 10.9. The eigenmonad adjunction yields an adjunction where ι is the inclusion functor:

$$\iota : {}_{H_0}\text{Mod} \xleftarrow{\quad \tau \quad} L_{\text{gr}^0}\text{Mod} : V_{\mathbf{I}^{\text{out}}}.$$

The refined eigenmonad adjunction gives an adjoint equivalence:

$$\iota : {}_{H_0}\text{Mod} \xleftarrow{\quad \tau \quad} L_{\text{gr}^0}\text{Mod}^{\text{out}} : V_{\mathbf{I}^{\text{out}}}.$$

Proposition 5.19 implies analogous statements based on $\text{Hom}_{L_{\text{gr}^0}}(H_0, -)$ instead of $V_{\mathbf{I}^{\text{out}}}$.

Proof. The first assertion follows from Proposition 5.22 and Theorem 10.8. The second one follows from Corollary 6.8 and Theorem 10.8. \square

11. PRIMITIVELY GENERATED MODULES OVER FREE GROUPS

In this section, we introduce a left ideal \mathbb{I}^{Pr} of $L_{\text{gr}^{\circ}}$, for which the eigenmonad is isomorphic to the moand $\mathbf{A}_{\mathcal{U}ie}$ associated with the Lie operad; and the quotient module is the module $\mathbf{A}_{\mathcal{U}ss''}$ associated with the unital associative operad. This implies that a functor cohomology related to $\mathbf{A}_{\mathcal{U}ss''}$ is isomorphic to $\mathbf{A}_{\mathcal{U}ie}$. In consequence, the eigenmonad adjunction reproduces the adjunction of $L_{\text{gr}^{\circ}}$ -modules and $\mathbf{A}_{\mathcal{U}ie}$ -modules [21, 12].

11.1. Definition of primitivity ideal. In this section, we introduce the primitivity ideal for a PROP C satisfying certain assumptions. We then examine the case of $C = \text{gr}^{\circ}$.

Definition 11.1. Let C be a PROP whose monoidal structure is the binary product \times and whose object $0 \in C$ is a zero object. Let $\Delta : 1 \rightarrow (1 \times 1) = 2$ be the diagonal morphism and $\eta : 0 \rightarrow 1$ be the unique morphism. We define $\theta_C \in \mathbb{k}C(1, 2) = L_C(2, 1)$ as

$$\theta_C := \Delta - (\text{id}_1 \times \eta) - (\eta \times \text{id}_1).$$

We define the *primitivity ideal*³ for C to be the left ideal of L_C generated by $\text{id}_k \times \theta_C \times \text{id}_l$ for $k, l \in \mathbb{N}$. It is denoted by \mathbb{I}_C^{Pr} . In other words, for each $n, m \in \mathbb{N}$, the module $\mathbb{I}_C^{\text{Pr}}(m, n) \subset L_C(m, n)$ is generated by $f \circ (\text{id}_{i-1} \times \theta \times \text{id}_{n-i})$ for $1 \leq i \leq n$ and $f \in L_C(m, n+1)$.

Notation 11.2. In this paper, we simply denote by $\mathbb{I}^{\text{Pr}} := \mathbb{I}_{\text{gr}^{\circ}}^{\text{Pr}}$.

Remark 11.3. The aforementioned definition is sufficient for our applications to gr° and fr_R° . It is possible to extend this to a PROP C whose object $1 \in C$ is a coaugmented coalgebra object. By using the comultiplication $\Delta : 1 \rightarrow (1 + 1) = 2$ and the coaugmentation $\eta : 0 \rightarrow 1$, we can define a similar notion. We wish to give an application of such a general construction somewhere.

Remark 11.4. The ideal \mathbb{I}^{Pr} implicitly appears in [12] based on different settings.

The category gr° satisfies the above conditions on C , and, by using the notation (28), we have

$$(30) \quad \theta_{\text{gr}^{\circ}} = [x_1 | x_1]_1 - [e | x_1]_1 - [x_1 | e]_1.$$

Hence, for $1 \leq i \leq n$, one may observe

$$\begin{aligned} \text{id}_{i-1} * \theta_{\text{gr}^{\circ}} * \text{id}_{n-i} &= [x_1 | \cdots | x_{i-1} | x_i | x_i | x_{i+1} | \cdots | x_n]_n \\ &\quad - ([x_1 | \cdots | x_{i-1} | x_i | e | x_{i+1} | \cdots | x_n]_n + [x_1 | \cdots | x_{i-1} | e | x_i | x_{i+1} | \cdots | x_n]_n). \end{aligned}$$

11.2. Canonical bimodule associated with primitivity ideal. In this section, we relate the canonical bimodule associated with the ideal \mathbb{I}^{Pr} to the operad for unital associative algebras. We prove an isomorphism between the quotient module $L_{\text{gr}^{\circ}}/\mathbb{I}^{\text{Pr}}$ and $\mathbf{A}_{\mathcal{U}ss''}$. As a consequence, $\mathbf{A}_{\mathcal{U}ss''}$ inherits a $(L_{\text{gr}^{\circ}}, E_{L_{\text{gr}^{\circ}}}(\mathbb{I}^{\text{Pr}}))$ -bimodule structure by Definition 5.12.

Recall our notation for the operad $\mathcal{A}ss''$ in section 8. Under our convention, we regard $\mathbf{A}_{\mathcal{U}ss''}(m, n)$ as a submodule of $\mathbb{k}\langle X_n \rangle^{\otimes m}$ via the following map ι :

$$(31) \quad \iota : \mathbf{A}_{\mathcal{U}ss''}(m, n) = \bigoplus_{f \in \text{Fin}(n, m)} \bigotimes_{i=1}^m \mathbf{A}_{\mathcal{U}ss''}(f^{-1}(i)) \rightarrow \mathbb{k}\langle X_n \rangle^{\otimes m}.$$

³Note that this is not related to *primitive ideals* in ring theory. Our terminology is motivated by that of primitively generated Hopf algebra. See Remark 11.18.

To be concrete, the \mathbb{k} -module $\mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(m, n)$ is a submodule of $\mathbb{k}\langle X_n \rangle^{\otimes m}$ generated by $w_1 \otimes \cdots \otimes w_m$ where w_1, \dots, w_m are words written by letters in X_n ; such that each letter in the concatenation of words $w_1 \cdots w_m$ appears exactly once. Especially,

$$(x_1 \otimes \cdots \otimes x_n) \in \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(n, n)$$

is the identity on n .

Definition 11.5. For $n, m \in \mathbb{N}$, we define a \mathbb{k} -linear map $\mathcal{E} : \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(m, n) \rightarrow \mathbf{L}_{\text{gr}^0}(m, n)$ by

$$\mathcal{E}(w_1 \otimes w_2 \otimes \cdots \otimes w_m) := [w_1 | w_2 | \cdots | w_m]_n,$$

where w_j 's are words based on X_n such that the concatenation of words $w_1 \cdots w_m$ is a permutation of the word $x_1 x_2 \cdots x_n$. The right hand side follows the notation in section 11.1. Its collection for all n, m defines a monad homomorphism $\mathcal{E} : \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u} \rightarrow \mathbf{L}_{\text{gr}^0}$.

Let $n \in \mathbb{N}$. The algebra $\mathbb{k}\langle X_n \rangle$ has a Hopf algebra structure whose comultiplication Δ is characterized by

$$(32) \quad \Delta(x_i) = x_i \otimes 1 + 1 \otimes x_i, \quad x_i \in X_n.$$

This structure leads to a (symmetric monoidal) functor $\text{gr}^0 \rightarrow \mathbb{k}\text{Mod}$ which assigns $\mathbb{k}\langle X_n \rangle^{\otimes m}$ to $m \in \mathbb{N}$ [20]. By the equivalence (12), we regard this functor as a left \mathbf{L}_{gr^0} -module \mathbf{M}_n . For instance, the action of $g = \text{id}_{j-1} * [x_1 | x_1]_1 * \text{id}_{m-j} \in \text{gr}^0(m, m+1)$ on \mathbf{M}_n is described by using a commutative diagram:

$$(33) \quad \begin{array}{ccc} \mathbf{M}_n(m) & \xrightarrow{g \triangleright (-)} & \mathbf{M}_n(m+1) \\ \parallel & & \parallel \\ \mathbb{k}\langle X_n \rangle^{\otimes m} & \xrightarrow{\text{id}^{\otimes (j-1)} \otimes \Delta \otimes \text{id}^{\otimes (m-j)}} & \mathbb{k}\langle X_n \rangle^{\otimes (m+1)} \end{array}$$

Lemma 11.6. Let $m, l \in \mathbb{N}$. For $f \in \text{gr}^0(m, l) \subset \mathbf{L}_{\text{gr}^0}(l, m)$, the restriction of the action $f \triangleright (-) : \mathbf{M}_n(m) \rightarrow \mathbf{M}_n(l)$ to $\mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(m, n)$ factors through $\mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(l, n)$:

$$\begin{array}{ccc} \mathbf{M}_n(m) & \xrightarrow{f \triangleright (-)} & \mathbf{M}_n(l) \\ \uparrow \iota & & \uparrow \iota \\ \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(m, n) & \xrightarrow{\exists} & \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(l, n) \end{array}$$

Proof. The PROP gr^0 is generated by the morphisms $[x_1 | x_1]_1 : 1 \rightarrow 2$, $[x_1 x_2]_2 : 2 \rightarrow 1$, $[x_1^{-1}]_1 : 1 \rightarrow 1$ and the trivial morphisms $0 \rightarrow 1$ and $1 \rightarrow 0$. Hence, it is sufficient to prove our assertion when f is given by taking free products of one of these generators and some identity morphisms. This is immediate from definitions. For instance, we consider $f = \text{id}_{j-1} * [x_1 | x_1]_1 * \text{id}_{m-j} \in \text{gr}^0(m, m+1)$ for $1 \leq j \leq m$. The commutative diagram (33) implies that, by (32), this maps $\mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(m, n)$ to $\mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(m+1, n)$. Similarly, it is verified for other types of morphisms. \square

Definition 11.7. We define a homomorphism $\mathcal{R} : \mathbf{L}_{\text{gr}^0} \rightarrow \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}$ between $(\mathbb{N} \times \mathbb{N})$ -indexed \mathbb{k} -modules. For $n, m \in \mathbb{N}$, the homomorphism $\mathcal{R} : \mathbf{L}_{\text{gr}^0}(m, n) \rightarrow \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(m, n)$ is defined as follows. For $f \in \mathbf{L}_{\text{gr}^0}(m, n)$, by using Lemma 11.6, we define

$$\mathcal{R}(f) := f \triangleright (x_1 \otimes x_2 \otimes \cdots \otimes x_n) \in \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(m, n)$$

where $x_1 \otimes x_2 \otimes \cdots \otimes x_n \in \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(n, n)$.

Lemma 11.8. The homomorphism $\mathcal{R} : \mathbf{L}_{\text{gr}^0} \rightarrow \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}$ induces a homomorphism $\bar{\mathcal{R}} : \mathbf{L}_{\text{gr}^0} / \mathbf{I}^{\text{Pr}} \rightarrow \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}$.

Proof. We first prove that $\mathcal{R}(\text{id}_{j-1} * \theta_{\text{gr}^0} * \text{id}_{n-j}) = 0$ for any $n \in \mathbb{N}$ and $1 \leq j \leq n$. By definition of the Hopf algebra structure on $\mathbb{k}\langle X_n \rangle$, we have

$$\begin{aligned} & (\text{id}_{j-1} * \theta_{\text{gr}^0} * \text{id}_{n-j}) \triangleright (x_1 \otimes x_2 \otimes \cdots \otimes x_n), \\ &= (\text{id}_{j-1} * ([x_1 | x_1]_1 - [e | x_1]_1 - [x_1 | e]_1) * \text{id}_{n-j}) \triangleright (x_1 \otimes x_2 \otimes \cdots \otimes x_n), \\ &= (x_1 \otimes \cdots \otimes \Delta(x_j) \otimes \cdots \otimes x_n) \\ &\quad - (x_1 \otimes \cdots \otimes x_j \otimes 1 \otimes \cdots \otimes x_n) - (x_1 \otimes \cdots \otimes 1 \otimes x_j \otimes \cdots \otimes x_n), \\ &= 0. \end{aligned}$$

Recall that \mathbf{I}^{pr} is the left ideal generated by $(\text{id}_{j-1} * \theta * \text{id}_{n-j})$'s. Hence, for $n, m \in \mathbb{N}$, every $f \in \mathbf{I}^{\text{pr}}(m, n)$ is written by $f' \circ (\text{id}_{j-1} * \theta * \text{id}_{n-j})$ for some f' , so that we have

$$\begin{aligned} \mathcal{R}(f) &= (f' \circ (\text{id}_{j-1} * \theta_{\text{gr}^0} * \text{id}_{n-j})) \triangleright (x_1 \otimes x_2 \otimes \cdots \otimes x_n), \\ &= f' \triangleright ((\text{id}_{j-1} * \theta_{\text{gr}^0} * \text{id}_{n-j}) \triangleright (x_1 \otimes x_2 \otimes \cdots \otimes x_n)) = 0. \end{aligned}$$

□

Theorem 11.9. *Let $p : \mathbf{L}_{\text{gr}^0} \rightarrow \mathbf{L}_{\text{gr}^0}/\mathbf{I}^{\text{pr}}$ be the canonical projection. The homomorphisms $p \circ \mathcal{E}$ and $\tilde{\mathcal{R}}$ are inverses to each other. In particular, we have an isomorphism of $(\mathbb{N} \times \mathbb{N})$ -indexed \mathbb{k} -modules:*

$$\mathbf{A}_{\mathfrak{U}_{\text{SS}}^u} \cong \mathbf{L}_{\text{gr}^0}/\mathbf{I}^{\text{pr}}.$$

Proof. In [12], the composition $p \circ \mathcal{E} : \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(m, n) \rightarrow (\mathbf{L}_{\text{gr}^0}/\mathbf{I}^{\text{pr}})(m, n)$ is proved to be surjective for any n, m . To prove our statement, it suffices to show that $\tilde{\mathcal{R}} \circ p \circ \mathcal{E} = \mathcal{R} \circ \mathcal{E}$ is the identity. Consider a generator $f = w_1 \otimes \cdots \otimes w_m \in \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}(m, n)$ for some words w_1, \dots, w_m consisting of x_1, \dots, x_n . Then, by definition of the left \mathbf{L}_{gr^0} -module \mathbf{M}_n , we have

$$\mathcal{R}(\mathcal{E}(f)) = [w_1 | \cdots | w_m]_n \triangleright (x_1 \otimes \cdots \otimes x_n) = w_1 \otimes \cdots \otimes w_m = f.$$

□

Remark 11.10. Note that the composition $(p \circ \mathcal{E})^{-1} \circ p : \mathbf{L}_{\text{gr}^0} \rightarrow \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}$ is not a monad homomorphism. In other words, \mathbf{I}^{pr} is not a two-sided ideal.

The underlying module of $\mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}$ has a canonical $(\mathbf{L}_{\text{gr}^0}, \mathbf{E}_{\mathbf{L}_{\text{gr}^0}}(\mathbf{I}^{\text{pr}}))$ -bimodule structure by Proposition 5.11 and 11.9.

Notation 11.11. We denote this by $\tilde{\mathbf{A}}_{\mathfrak{U}_{\text{SS}}^u}$, which turns out to be a $(\mathbf{L}_{\text{gr}^0}, \mathbf{A}_{\mathfrak{Q}_{\text{ie}}})$ -bimodule by Theorem 11.13.

Notation 11.12. Let $\beta : \mathbf{A}_{\mathfrak{Q}_{\text{ie}}} \rightarrow \mathbf{A}_{\mathfrak{U}_{\text{SS}}^u}$ be the monad homomorphism induced by the canonical operad homomorphism $\mathfrak{Q}_{\text{ie}} \rightarrow \mathfrak{U}_{\text{SS}}^u$. We regard this as the homomorphism between $\beta : \mathbf{A}_{\mathfrak{Q}_{\text{ie}}} \rightarrow \tilde{\mathbf{A}}_{\mathfrak{U}_{\text{SS}}^u}$ right $\mathbf{A}_{\mathfrak{Q}_{\text{ie}}}$ -modules.

11.3. Eigenmonad by primitivity ideal. In this section, we prove a monad isomorphism between the eigenmonad $\mathbf{E}_{\mathbf{L}_{\text{gr}^0}}(\mathbf{I}^{\text{pr}})$ and the monad $\mathbf{A}_{\mathfrak{Q}_{\text{ie}}}$ by refining the results of section 11.2. To this end, we show that β (see Notation 11.12) induces an isomorphism $\mathbf{A}_{\mathfrak{Q}_{\text{ie}}} \rightarrow \mathbf{V}_{\mathbf{I}^{\text{pr}}}(\tilde{\mathbf{A}}_{\mathfrak{U}_{\text{SS}}^u})$ of right $\mathbf{A}_{\mathfrak{Q}_{\text{ie}}}$ -modules.

We investigate the relationship between free algebras and free Lie algebras, since the operads $\mathfrak{U}_{\text{SS}}^u$ and \mathfrak{Q}_{ie} are closely related to them. For a finite set X , we denote by $\mathbb{k}\langle X \rangle$ the free algebra generated by X ; and by $L(X) \subset \mathbb{k}\langle X \rangle$ the free Lie algebra generated by X . Then $\mathbb{k}\langle X \rangle$ has a grading by the number of letters in X . To be precise, for a map $\delta : X \rightarrow \mathbb{N}$, if we define $\mathbb{k}\langle X \rangle_{\delta}$ to be the submodule generated by words consisting of $\delta(x)$ copies of $x \in X$, then we have $\mathbb{k}\langle X \rangle = \bigoplus_{\delta \in V_X} \mathbb{k}\langle X \rangle_{\delta}$ where V_X is the set of maps $X \rightarrow \mathbb{N}$. Then $L(X)$ inherits the V_X -grading of $\mathbb{k}\langle X \rangle$ (see Theorem C.5).

For $n, m \in \mathbb{N}$, the modules $\mathbf{A}_{\mathfrak{A}_{\text{ss}}^u}(m, n)$ and $\mathbf{A}_{\mathfrak{A}_{\text{ie}}}(m, n)$ can be described by using these graded modules. Let $X_n := \{x_1, \dots, x_n\}$ and $1 \equiv \delta_0 \in V_{X_n}$. By the definitions, the module $\mathbf{A}_{\mathfrak{A}_{\text{ss}}^u}(m, n)$ is identified with the δ_0 -component of $\mathbb{k}\langle X_n \rangle^{\otimes m}$. Here, we consider the tensor product of graded modules (see section B for further details). Similarly, $\mathbf{A}_{\mathfrak{A}_{\text{ie}}}(m, n)$ is the δ_0 -component of $L(X_n)^{\otimes m}$.

Theorem 11.13. (1) *The homomorphism β induces an isomorphism $\beta_0 : \mathbf{A}_{\mathfrak{A}_{\text{ie}}} \rightarrow \mathbf{V}_{\text{I}^{\text{pr}}}(\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u})$ of right $\mathbf{A}_{\mathfrak{A}_{\text{ie}}}$ -modules.*

(2) *We have an isomorphism $\beta_1 : \mathbf{A}_{\mathfrak{A}_{\text{ie}}} \rightarrow \mathbf{E}_{\mathbf{L}_{\text{gr}}^{\text{o}}}(\text{I}^{\text{pr}})$ of monads which fits into the following commutative diagram:*

$$(34) \quad \begin{array}{ccc} \tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u} & \xrightarrow{p \circ \mathcal{E}} & \mathbf{L}_{\text{gr}}^{\text{o}} / \text{I}^{\text{pr}} \\ \beta \uparrow & & \uparrow \\ \mathbf{A}_{\mathfrak{A}_{\text{ie}}} & \xrightarrow{\beta_1} & \mathbf{E}_{\mathbf{L}_{\text{gr}}^{\text{o}}}(\text{I}^{\text{pr}}) \end{array}$$

Proof. We use the aforementioned observations to prove (1). By the definition of vanishing module, we have

$$(\mathbf{V}_{\text{I}^{\text{pr}}}(\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u}))(m, n) = \bigcap_{j=1}^m \{f \in \tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u}(m, n) \mid (\text{id}_{j-1} * \theta_{\text{gr}^{\text{o}}} * \text{id}_{m-j}) \triangleright f = 0\}.$$

Theorem 11.9 implies that the left $\mathbf{L}_{\text{gr}}^{\text{o}}$ -module $\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u}(-, n)$ is a submodule of \mathbf{M}_n in Lemma 11.6. Hence, by applying the identification $\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u}(m, n) = (\mathbb{k}\langle X_n \rangle^{\otimes m})_{\delta_0}$, we obtain

$$(\mathbf{V}_{\text{I}^{\text{pr}}}(\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u}))(m, n) = \bigcap_{j=1}^m \{f \in (\mathbb{k}\langle X_n \rangle^{\otimes m})_{\delta_0} \mid (\text{id}_{j-1} * \bar{\Delta} * \text{id}_{m-j}) \triangleright f = 0\}$$

where $\bar{\Delta}$ is the reduced comultiplication on the Hopf algebra $\mathbb{k}\langle X_n \rangle$. By a careful modification of the basis theorem on free Lie algebras [25], we see that the right hand side is isomorphic to $(L(X_n)^{\otimes m})_{\delta_0}$. The reader is referred to Corollary C.6 for further details. By applying the identification $(L(X_n)^{\otimes m})_{\delta_0} \cong \mathbf{A}_{\mathfrak{A}_{\text{ie}}}(m, n)$, we obtain an isomorphism $\beta_0 : \mathbf{A}_{\mathfrak{A}_{\text{ie}}}(m, n) \rightarrow (\mathbf{V}_{\text{I}^{\text{pr}}}(\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u}))(m, n)$, which is, by the definitions, compatible with β . Furthermore, the induced isomorphism $\beta_0 : \mathbf{A}_{\mathfrak{A}_{\text{ie}}} \rightarrow \mathbf{V}_{\text{I}^{\text{pr}}}(\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u})$ automatically preserves right $\mathbf{A}_{\mathfrak{A}_{\text{ie}}}$ -module structure, since the vanishing module is taken with respect to the *left* $\mathbf{L}_{\text{gr}}^{\text{o}}$ -action.

By Proposition 5.20 and Theorem 11.9, the composition $p \circ \mathcal{E}$ induces an isomorphism $\mathbf{V}_{\text{I}^{\text{pr}}}(\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u}) \cong \mathbf{E}_{\mathbf{L}_{\text{gr}}^{\text{o}}}(\text{I}^{\text{pr}})$. By taking a composition with the isomorphism β_0 , we obtain an isomorphism $\beta_1 : \mathbf{A}_{\mathfrak{A}_{\text{ie}}} \rightarrow \mathbf{E}_{\mathbf{L}_{\text{gr}}^{\text{o}}}(\text{I}^{\text{pr}})$. By the definitions, for $n, m \in \mathbb{N}$ and $f \in \mathbf{A}_{\mathfrak{A}_{\text{ie}}}(m, n)$, we have $\beta_1(f) = \mathcal{E}(\beta(f)) \text{ mod } \text{I}^{\text{pr}}(m, n)$. Hence, β_1 is a monad homomorphism, since \mathcal{E}, β are monad homomorphisms. \square

Corollary 11.14. *The eigenmonad adjunction induces an adjunction:*

$$\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u} \otimes_{\mathbf{A}_{\mathfrak{A}_{\text{ie}}}} (-) : \mathbf{A}_{\mathfrak{A}_{\text{ie}}} \text{Mod} \xrightleftharpoons[\tau]{\epsilon} \mathbf{L}_{\text{gr}}^{\text{o}} \text{Mod} : \mathbf{V}_{\text{I}^{\text{pr}}}.$$

Proposition 5.19 implies analogous statements based on $\text{Hom}_{\mathbf{L}_{\text{gr}}^{\text{o}}}(\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u}, -)$ instead of $\mathbf{V}_{\text{I}^{\text{pr}}}$.

Remark 11.15. One may realize that, through the equivalence (12), the $(\mathbf{L}_{\text{gr}}^{\text{o}}, \mathbf{A}_{\mathfrak{A}_{\text{ie}}})$ -bimodule $\tilde{\mathbf{A}}_{\mathfrak{A}_{\text{ss}}^u}$ corresponds to the \mathbb{k} -bilinear functor $\Delta \text{Cat}_{\mathfrak{A}_{\text{ss}}^u} : \mathbb{k}\text{gr}^{\text{o}} \times \text{Cat}_{\mathfrak{A}_{\text{ie}}} \rightarrow \mathbb{k}\text{Mod}$ in [21, 12]. The above adjunction is directly constructed in the literature while we derive it from a general construction.

11.4. Primitively generated modules. In this section, we introduce the notion of a primitively generated L_{gr^0} -module. We also give its equivalent condition by using the previous results.

Definition 11.16. Let C be a PROP satisfying the assumptions in Definition 11.1. We say that a left L_C -module is *primitively generated* if it is \mathbb{I}_C^{Pr} -vanishingly generated (see Definition 6.1). Denote by $L_C \text{Mod}^{\text{prim}}$ the full subcategory of primitively generated left L_C -modules.

Proposition 11.17. *A left L_{gr^0} -module \mathbb{M} is primitively generated if and only if the following evaluation map is an epimorphism:*

$$(35) \quad \tilde{\mathbb{A}}_{\mathbb{I}_{\text{SS}}^u} \otimes_{\mathbb{A}_{\text{Vic}}} \text{Hom}_{L_{\text{gr}^0}}(\tilde{\mathbb{A}}_{\mathbb{I}_{\text{SS}}^u}, \mathbb{M}) \rightarrow \mathbb{M}.$$

Proof. The homomorphism in (35) is the same as the unit of the adjunction in Corollary 11.14. By Theorems 11.9 and 11.13, this is an epimorphism if and only if \mathbb{M} is $\mathbb{I}_{\text{gr}^0}^{\text{Pr}}$ -vanishingly generated, i.e. primitively generated. \square

Remark 11.18. By Example 4.16, a left L_{gr^0} -module is nothing but a functor from gr^0 to ${}_{\mathbb{k}}\text{Mod}$. Under this identification, the notion of a primitively generated L_{gr^0} -module coincides with that of a primitive gr^0 -module introduced in [12]. In that paper, we prove that the category of *exponential* functors from gr^0 to ${}_{\mathbb{k}}\text{Mod}$ is equivalent with that of primitively generated Hopf algebras.

12. COMPARISON OF PRIMITIVITY AND ANALYTICITY

This section presents a proof that primitively generated L_{gr^0} -modules are analytic. Furthermore, a more robust assertion is made with respect to the associated left ideals, i.e. the primitivity ideal \mathbb{I}^{Pr} and the left ideal \mathbb{I}_C^v in Definition 7.21.

Theorem 12.1. *Let $\nu_0 : \mathbb{N} \rightarrow \mathbb{N}$ such that $\nu_0(n) = n + 1$. The primitivity ideal contains the ν_0 -analyticity ideal:*

$$\mathbb{I}_{\text{gr}^0}^{\nu_0} \subset \mathbb{I}^{\text{Pr}}.$$

Proof. Let $d \in \mathbb{N}$. The polynomial degree of the left L_{gr^0} -module $\tilde{\mathbb{A}}_{\mathbb{I}_{\text{SS}}^u}(-, d)$ is d [12], and so does the left L_{gr^0} -module $(L_{\text{gr}^0}/\mathbb{I}^{\text{Pr}})(-, d)$ by Theorem 11.9. By Theorem 7.9, we have

$$V_{\mathbb{I}_{\text{gr}^0}^{(d+1)}}((L_{\text{gr}^0}/\mathbb{I}^{\text{Pr}})(-, d)) = (L_{\text{gr}^0}/\mathbb{I}^{\text{Pr}})(-, d).$$

In particular, for $n, m \in \mathbb{N}$ and $g \in \mathbb{I}_{\text{gr}^0}^{(d+1)}(m, n)$, every $f \in L_{\text{gr}^0}(n, d)$ satisfies $g \circ f \in \mathbb{I}^{\text{Pr}}(m, d)$. If we choose $n = d$ and f to be the identity on d , then we obtain $g \in \mathbb{I}^{\text{Pr}}(m, d)$. This proves

$$\mathbb{I}_{\text{gr}^0}^{\nu_0}(m, d) = \mathbb{I}_{\text{gr}^0}^{(d+1)}(m, d) \subset \mathbb{I}^{\text{Pr}}(m, d).$$

\square

Corollary 12.2. *Let \mathbb{M} be a left L_{gr^0} -module. If \mathbb{M} is primitively generated, then it is analytic.*

Proof. By Lemma 6.11 and Theorem 12.1, if a left L_{gr^0} -module \mathbb{M} is primitively generated, i.e. \mathbb{I}^{Pr} -vanishingly generated, then it is $\mathbb{I}_{\text{gr}^0}^{\nu_0}$ -vanishingly generated. Hence, \mathbb{M} is analytic by Theorem 7.25. \square

By applying the results in [21], we obtain the following:

Corollary 12.3. *Suppose that the ground ring \mathbb{k} is a field of characteristic zero. Then we have the followings.*

- A left L_{gr^0} -module is primitively generated if and only if it is analytic.
- Every left $E_{L_{\text{gr}^0}}(\mathbb{I}^{\text{Pr}})$ -module is \mathbb{I}^{Pr} -vanishingly extensible.

In particular, the refined eigenmonad adjunction (see Proposition 6.7) leads to an adjoint equivalence:

$$(36) \quad \tilde{\mathbf{A}}_{\mathbb{L}_{\text{gr}^0}} \otimes_{\mathbf{A}_{\text{vic}}} (-) : \mathbf{A}_{\text{vic}} \mathbf{Mod} \begin{array}{c} \xleftarrow{\quad \tau \quad} \\ \xrightarrow{\quad \tau \quad} \end{array} \mathbf{L}_{\text{gr}^0} \mathbf{Mod}^{\text{prim}} : \mathbf{V}_{\mathbb{I}^{\text{Pr}}}.$$

Proposition 5.19 implies analogous statements based on $\text{Hom}_{\mathbf{L}_{\text{gr}^0}}(\tilde{\mathbf{A}}_{\mathbb{L}_{\text{gr}^0}}, -)$ instead of $\mathbf{V}_{\mathbb{I}^{\text{Pr}}}$.

Proof. For an analytic left \mathbf{L}_{gr^0} -module \mathbf{M} , we have $\mathbf{M} \cong \tilde{\mathbf{A}}_{\mathbb{L}_{\text{gr}^0}} \otimes_{\mathbf{E}_{\mathbf{L}_{\text{gr}^0}}(\mathbb{I}^{\text{Pr}})} \mathbf{V}_{\mathbb{I}^{\text{Pr}}}(\mathbf{M})$ by [21]. Hence, \mathbf{M} is primitively generated since $\tilde{\mathbf{A}}_{\mathbb{L}_{\text{gr}^0}} \otimes_{\mathbf{A}_{\text{vic}}} \mathbf{N} \cong (\mathbf{L}_{\text{gr}^0}/\mathbb{I}^{\text{Pr}}) \otimes_{\mathbf{E}_{\mathbf{L}_{\text{gr}^0}}} \mathbf{N}$ by Theorem 11.9 and Theorem 11.13; and $(\mathbf{L}_{\text{gr}^0}/\mathbb{I}^{\text{Pr}}) \otimes_{\mathbf{E}_{\mathbf{L}_{\text{gr}^0}}} \mathbf{N}$ is primitively generated for any $\mathbf{N} \in \mathbf{A}_{\text{vic}} \mathbf{Mod}$. Furthermore, by Corollary 12.2, every primitively generated module is analytic.

We now prove the second part. Let \mathbf{N} be a left $\mathbf{E}_{\mathbf{L}_{\text{gr}^0}}(\mathbb{I}^{\text{Pr}})$ -module. By [21] again, we have $\mathbf{N} \cong \mathbf{V}_{\mathbb{I}^{\text{Pr}}}((\mathbf{L}_{\text{gr}^0}/\mathbb{I}^{\text{Pr}}) \otimes_{\mathbf{E}_{\mathbf{L}_{\text{gr}^0}}} \mathbf{N})$. Hence, the canonical homomorphism $\mathbf{N} \rightarrow (\mathbf{L}_{\text{gr}^0}/\mathbb{I}^{\text{Pr}}) \otimes_{\mathbf{E}_{\mathbf{L}_{\text{gr}^0}}} \mathbf{N}$ is a monomorphism.

To prove the equivalence in (36), we apply the previous results together with Theorem 11.9 and Theorem 11.13 to the adjunction in Proposition 6.7. \square

Part 3. Application to free modules

The study of modules over an *additive* category has been applied to the field of algebraic topology [10, 6, 7], since polynomial \mathbf{Ab} -modules were introduced in [5] to study the homology of Eilenberg-MacLane spaces where \mathbf{Ab} is the category of abelian groups. In this part, we study an additive category which is partly related to those literature. For the purposes of this discussion, we fix a unital ring R which is not necessarily commutative. It is hoped that readers will not confuse R with the commutative ring \mathbb{k} which is freely used throughout this paper. Let us denote by fr_R° the opposite category of finitely generated free right R -modules and R -homomorphisms. The purpose of this part is to study polynomial modules, primitively generated modules over the monad $\mathbf{L}_{\text{fr}_R^{\circ}}$ based on results of part 1. Additionally, when $R = \mathbb{Z}$, we provide a comparison with the results in part 2 via the abelianization.

13. NOTATION

The symbols \mathbf{Fin} , \mathfrak{S} stand for the following categories whose objects are natural numbers.

– $\mathbf{Fin}(n, m)$: the set of maps from $\underline{n} = \{1, 2, \dots, n\}$ to \underline{m} .

–

$$\mathfrak{S}(n, m) := \begin{cases} \mathfrak{S}_n & (n = m), \\ \emptyset & (n \neq m). \end{cases}$$

We explain our notation for morphisms in fr° which is freely used in this part. For $n \in \mathbb{N}$, we identify the object $R^n \in \text{fr}_R^{\circ}$ freely generated by n elements with n , so that we regard fr_R° as a category with objects \mathbb{N} . By definition, $\text{fr}_R^{\circ}(n, m)$ consists of R -homomorphisms from R^m to R^n . It is clear that, for $n, m \in \mathbb{N}$, we have a bijection:

$$\mathbf{M}_{n,m}(R) \xrightarrow{\cong} \text{fr}_R^{\circ}(n, m).$$

Here, $\mathbf{M}_{n,m}(R)$ is the set of R -matrices with n rows and m columns. In spite of the familiarity, we attempt to avoid the matrix expression in order that we distinguish sums in the \mathbb{k} -module $\mathbf{L}_{\text{fr}^{\circ}}(m, n) = \mathbb{k}\text{fr}^{\circ}(n, m)$ from that of matrices. Instead, we introduce a notation $\tau^X \in \text{fr}_R^{\circ}(n, m)$ for the morphism assigning to $X \in \mathbf{M}_{n,m}(R)$ where τ is a fixed formal letter.

Based on our notation, the abelian group structure on $\mathbf{M}_{n,m}(R)$ is inherited to $\text{fr}_R^{\circ}(n, m)$ as follows:

$$(37) \quad \tau^X \cdot \tau^Y = \tau^{X+Y}, \quad \forall X, Y \in \mathbf{M}_{n,m}(R).$$

As a special case, let $A_n(R) := \text{fr}_R^0(n, 1)$ be the abelian group with unit $e := \tau^0$. Then $C_n = A_n(R)$ for $C = \text{fr}^0$ in section 7.5.

By using our notation, the composition of fr_R^0 (hence, the monad operation of $L_{\text{fr}_R^0}$) is described by

$$(38) \quad \tau^Y \circ \tau^X = \tau^{XY}, \quad \forall X \in M_{n,m}(R), \forall Y \in M_{m,l}(R).$$

Formally speaking, the composition behaves like a substitution of the right morphism into the left τ , i.e. $\tau^Y \circ \tau^X = (\tau^X)^Y$.

The category fr_R^0 has product given by direct sum of R -modules. It assigns the following to morphisms τ^X, τ^Y :

$$\tau^X \oplus \tau^Y = \tau^{X \oplus Y}$$

$$\text{where } X \oplus Y := \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}.$$

14. ABELIANIZATION

It is necessary to recall the monads L_{gr^0} and $L_{\text{fr}_{\mathbb{Z}}^0}$ on \mathbb{N} described in Example 4.5. This section examines the relationship between $L_{\text{fr}_{\mathbb{Z}}^0}$ -modules and L_{gr^0} -modules by using the *abelianization* functor $\text{gr} \rightarrow \text{fr}_{\mathbb{Z}}$. As a consequence, we obtain a two-sided ideal of L_{gr^0} whose eigenmonad is given by $L_{\text{fr}_{\mathbb{Z}}^0}$. This is closely related to *abelianizable* L_{gr^0} -modules, which is introduced below. The results of this section are useful to understand the comparisons of L_{gr^0} -modules and $L_{\text{fr}_{\mathbb{Z}}^0}$ -modules in sections 15.3 and 17.

Definition 14.1. Let $\alpha_n : F_n \rightarrow \mathbb{Z}^n$ be the quotient map (associated with the abelianization). It assigns $e_i \in \mathbb{Z}^n$ to $x_i \in F_n$ where e_i has 1 in the i -th component and 0 in other components. We define a monad homomorphism $\alpha : L_{\text{gr}^0} \rightarrow L_{\text{fr}_{\mathbb{Z}}^0}$ to be

$$\alpha([w_1 | w_2 | \cdots | w_m]_n) := \tau^{[\alpha_n(w_1), \alpha_n(w_2), \dots, \alpha_n(w_m)]}$$

where $w_j \in F_n$ and we regard $[\alpha_n(w_1), \alpha_n(w_2), \dots, \alpha_n(w_m)]$ as a matrix in $M_{n,m}(\mathbb{Z})$.

Definition 14.2. For a left $L_{\text{fr}_{\mathbb{Z}}^0}$ -module \mathbb{M} , we define $\alpha^*(\mathbb{M})$ to be the left L_{gr^0} -module induced by α . We also denote by $\alpha^* : L_{\text{fr}_{\mathbb{Z}}^0} \text{Mod} \rightarrow L_{\text{gr}^0} \text{Mod}$ the induced functor.

Definition 14.3. A left L_{gr^0} -module \mathbb{N} is *abelianizable* if there exists a left $L_{\text{fr}_{\mathbb{Z}}^0}$ -module \mathbb{M} with $\alpha^*(\mathbb{M}) \cong \mathbb{N}$. We define a subclass $\mathcal{S}^{\text{ab}} \subset L_{\text{gr}^0} \text{Mod}$ to be the class of isomorphism classes of an abelianizable left L_{gr^0} -module.

In the following, recall $\text{Ann}(\mathcal{S})$ in Definition 6.17.

Theorem 14.4. *The subclass $\mathcal{S}^{\text{ab}} \subset L_{\text{gr}^0} \text{Mod}$ has a unique core L_{gr^0} -internalizer \mathbb{I}^{ab} . In particular, we have $\text{Ann}(\mathcal{S}^{\text{ab}}) = \mathbb{I}^{\text{ab}}$.*

Proof. Note that the kernel of α in Definition 14.1 is a two-sided ideal of L_{gr^0} , since α is a monad homomorphism. We denote by \mathbb{I}^{ab} the ideal. It is immediate from the definition that we have

$$(39) \quad L_{\text{gr}^0} / \mathbb{I}^{\text{ab}} \cong L_{\text{fr}_{\mathbb{Z}}^0}.$$

By Corollary 6.8, we see that a left L_{gr^0} -module is \mathbb{I}^{ab} -vanishingly generated if and only if it is in the image of α^* , i.e. abelianizable. Hence, $\mathcal{S}^{\text{ab}} = \mathcal{V}(\mathbb{I}^{\text{ab}})$. The last statement follows from Theorem 6.20. \square

Corollary 14.5. *The functor α^* is a left adjoint:*

$$\alpha^* : {}_{L_{\text{fr}}^{\circ}}\text{Mod} \begin{array}{c} \longleftarrow \\ \xrightarrow{\tau} \end{array} {}_{L_{\text{gr}}^{\circ}}\text{Mod} : \text{Hom}_{\mathbb{T}}(L_{\text{fr}}^{\circ}, -) (\cong V_{\mathbb{I}^{\text{ab}}}).$$

This induces an adjoint equivalence where ${}_{L_{\text{gr}}^{\circ}}\text{Mod}^{\text{ab}}$ is the full subcategory of ${}_{L_{\text{gr}}^{\circ}}\text{Mod}$ generated by abelianizable modules:

$$\alpha^* : {}_{L_{\text{fr}}^{\circ}}\text{Mod} \begin{array}{c} \longleftarrow \\ \xrightarrow{\tau} \end{array} {}_{L_{\text{gr}}^{\circ}}\text{Mod}^{\text{ab}} : \text{Hom}_{\mathbb{T}}(L_{\text{fr}}^{\circ}, -) (\cong V_{\mathbb{I}^{\text{ab}}}).$$

Proof. The first is proved by the eigenmonad adjunction in (1) of Proposition 5.13 and Proposition 5.19. The second claim follows from Corollary 6.8 and Theorem 14.4. \square

Remark 14.6. We have similar results for the *non-opposite* ones L_{fr_Z} and L_{gr} , but our discussion here aims to give some preliminaries for section 15.3.

We now give a presentation of \mathbb{I}^{ab} given in the proof of Theorem 14.4. In fact, the homomorphism α has a section homomorphism which is given by *rearranging the letters*:

Definition 14.7. Let $\gamma_n : \mathbb{Z}^n \rightarrow \mathbb{F}_n$ be a map defined as

$$\begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} \in \mathbb{Z}^n \mapsto x_1^{a_1} \cdots x_n^{a_n} \in \mathbb{F}_n$$

By extending this, we define a homomorphism $\gamma : L_{\text{fr}_Z}^{\circ} \rightarrow L_{\text{gr}}^{\circ}$ between $(\mathbb{N} \times \mathbb{N})$ -indexed \mathbb{k} -modules as follows:

$$L_{\text{fr}_Z}^{\circ}(m, n) \rightarrow L_{\text{gr}}^{\circ}(m, n); \tau^{[u_1, u_2, \dots, u_m]} \mapsto [\gamma_n(u_1) | \gamma_n(u_2) | \cdots | \gamma_n(u_m)]_n$$

where $u_j \in \mathbb{Z}^n$.

Lemma 14.8. *The homomorphism γ is a section of α , i.e. $\alpha \circ \gamma = \text{id}_{L_{\text{fr}_Z}^{\circ}}$.*

Proof. It follows from the fact that each map γ_n is a section of α_n , i.e. $\alpha_n \circ \gamma_n = \text{id}_{\mathbb{Z}^n}$. \square

Proposition 14.9. *The two-sided ideal \mathbb{I}^{ab} is generated by such elements as*

$$[w_1 | w_2 | \cdots | w_m]_n - [\gamma_n(\alpha_n(w_1)) | \gamma_n(\alpha_n(w_2)) | \cdots | \gamma_n(\alpha_n(w_m))]_n, \forall w_j \in \mathbb{F}_n.$$

Proof. By Lemma 14.8, we see that the kernel of α , i.e. \mathbb{I}^{ab} , coincides with the image of the endomorphism $(\text{id}_{L_{\text{gr}}^{\circ}} - \gamma \circ \alpha) : L_{\text{gr}}^{\circ} \rightarrow L_{\text{gr}}^{\circ}$. \square

Recall the ideal \mathbb{I}^{out} of L_{gr}° in Definition 10.1.

Corollary 14.10. *We have the followings:*

- (1) $\mathbb{I}^{\text{out}} \subset \mathbb{I}^{\text{ab}}$.
- (2) *The monad $L_{\text{fr}_Z}^{\circ}$ is a quotient monad of \mathbb{H}_0 .*
- (3) $\mathcal{S}^{\text{ab}} \subset \mathcal{S}^{\text{out}}$.

Proof. We prove the part (1). Let $g \in \mathbb{F}_n$. We put $[w_1 | w_2 | \cdots | w_n]_n := \text{Ad}_g([x_1 | x_2 | \cdots | x_n]_n)$. By the definition of γ_n , we have

$$[\gamma_n(\alpha_n(w_1)) | \gamma_n(\alpha_n(w_2)) | \cdots | \gamma_n(\alpha_n(w_m))]_n = [x_1 | x_2 | \cdots | x_n]_n.$$

Hence, we obtain $\text{Ad}_g(\text{id}_n) - \text{id}_n$ lies in \mathbb{I}^{ab} . This yields $\mathbb{I}^{\text{out}} \subset \mathbb{I}^{\text{ab}}$.

By (1), we have a monad epimorphism $\mathbb{H}_0 \rightarrow L_{\text{fr}_Z}^{\circ}$ by Theorem 10.8 and (39). It proves the parts (2) and (3). \square

15. POLYNOMIAL IDEALS FOR FREE MODULES

This section employs the same methodology as that used in section 9 to study polynomial ideals for fr_R^0 . Moreover, we give a comparison between polynomial ideals for $\text{fr}_\mathbb{Z}^0$ and those for gr^0 where the latter are studied in section 9.

15.1. Eigenmonads for polynomial ideals. In this section, we give a computation of the eigenmonad by polynomial ideals by following the strategy of section 7.5. The category fr_R^0 satisfies the assumptions on \mathcal{C} in that section. By Corollary 7.32, it suffices to compute the Passi monad $\mathcal{P}_{\text{fr}_R^0}^d$. It is readily verified that the monoid $C_n = \text{fr}_R^0(n, 1)$ is isomorphic to the underlying monoid of $\mathbf{A}_1(R)$ with the monoid operation given by (37). The augmentation ideal of $C_n^{\times m}$ is identified with that of $(\mathbf{A}_n(R))^{\times m} \cong \mathbf{A}_{nm}(R)$. Hence, Theorem 7.30 implies the following isomorphism:

$$(40) \quad \mathbb{I}_{\mathbb{k}}(\mathbf{A}_{nm}(R))^d \cong \mathbb{I}_{\text{fr}_R^0}^{(d)}(m, n), \quad n, m \in \mathbb{N}.$$

Remark 15.1. Note that fr_R itself also satisfies the assumptions on \mathcal{C} . Indeed, we have $\text{fr}_R \cong \text{fr}_R^0$ given by the dual functor $\text{Hom}_R(-, R)$. The reason that we consider the opposite category is to compare the results with those of gr^0 later.

Notation 15.2. $\mathbf{A}_n := \mathbf{A}_n(\mathbb{Z})$.

Lemma 15.3. *For $N \in \mathbb{N}$, we have an isomorphism where \vec{p} runs over all the ordered N -partitions of d :*

$$\bigoplus_{\vec{p}} \mathbb{k} \cong \mathbb{I}_{\mathbb{k}}(\mathbf{A}_N)^d / \mathbb{I}_{\mathbb{k}}(\mathbf{A}_N)^{d+1}.$$

In particular, its rank is $\frac{(N+d-1)!}{d!(N-1)!}$.

Proof. The algebra $\mathbb{k}[\mathbf{A}_N]$ is the (commutative) polynomial algebra with N variables t_i 's. The right hand side has a basis consisting of $\prod_{i=1}^N (t_i - 1)^{p_i} \in \mathbb{k}[\mathbf{A}_N]$ modulo $\mathbb{I}_{\mathbb{k}}(\mathbf{A}_N)^{d+1}$ for $p = (p_1, \dots, p_N)$ is an ordered N -partition of d . This argument leads to the statement. \square

Theorem 15.4. *For $n, m \in \mathbb{N}$, the module $\mathcal{P}_{\text{fr}_\mathbb{Z}^0}^d(m, n)$ is a free \mathbb{k} -module with rank of*

$$\sum_{k=0}^d \frac{(nm + k - 1)!}{k!(nm - 1)!}.$$

Proof. The strategy is the same as that of Theorem 9.1. We apply (40) and Lemma 15.3 with some short exact sequences. \square

15.2. Internalizers for analytic modules. By Theorem 7.25, we know that the class of analytic $\text{L}_{\text{fr}_R^0}$ -modules has an $\text{L}_{\text{fr}_R^0}$ -internalizer. For $R = \mathbb{Z}$, we can go further. In fact, there are uncountably many $\text{L}_{\text{fr}_\mathbb{Z}^0}$ -internalizers as follows:

Proposition 15.5. *The assignment of $\mathbb{I}_{\text{fr}_\mathbb{Z}^0}^v$ to a map $v : \mathbb{N} \rightarrow \mathbb{N}$ is injective.*

Proof. It basically follows from Remark 7.11. The proof is parallel with that of Proposition 9.4. \square

By Theorem 7.13, the class $\mathcal{S}_{\text{fr}_R^0}^d$ of modules of polynomial degree $\leq d$ has a core $\text{L}_{\text{fr}_R^0}$ -internalizer; but it is not true for the class $\mathcal{S}_{\text{fr}_R^0}^\omega$ of analytic modules when $R = \mathbb{Z}$:

Theorem 15.6. *The subclass $\mathcal{S}_{\text{fr}_\mathbb{Z}^0}^\omega \subset \text{L}_{\text{fr}_\mathbb{Z}^0} \text{Mod}$ does not allow a core $\text{L}_{\text{fr}_\mathbb{Z}^0}$ -internalizer.*

Proof. The proof is parallel with Theorem 9.5 since $\mathbf{F}_1 \cong \mathbf{A}_1$. \square

15.3. Comparison of polynomial ideals for gr^0 and $\text{fr}_{\mathbb{Z}}^0$. Recall the concepts related to the abelianization in section 14. It is easy to see that, under the functor α^* , the categories $\text{L}_{\text{fr}_{\mathbb{Z}}^0} \text{Mod}^{\leq 1}$ and $\text{L}_{\text{gr}^0} \text{Mod}^{\leq 1}$ are equivalent. For instance, the degree zero part follows from Proposition 7.18 (or see Corollary 15.11). It is natural to ask the following question: how about the cases of polynomial degrees greater than one? In this section, we give a negative answer to this question by comparing the eigenmonads by the polynomial ideals for $\text{fr}_{\mathbb{Z}}^0$ and gr^0 .

Lemma 15.7. *For $d, n, m \in \mathbb{N}$, we have $\alpha(\mathbb{I}_{\text{gr}^0}^{(d)}(m, n)) = \mathbb{I}_{\text{fr}_{\mathbb{Z}}^0}^{(d)}(m, n)$. Hence, α induces a homomorphism between $(\mathbb{N} \times \mathbb{N})$ -indexed \mathbb{k} -modules:*

$$\bar{\alpha} : \mathbb{I}_{\text{gr}^0}^{(d)}/\mathbb{I}_{\text{gr}^0}^{(d+1)} \rightarrow \mathbb{I}_{\text{fr}_{\mathbb{Z}}^0}^{(d)}/\mathbb{I}_{\text{fr}_{\mathbb{Z}}^0}^{(d+1)}.$$

Proof. We recall some definitions in section 7.2. By definitions, we have $\alpha(\pi_n^d) = \pi_{\alpha(n)}^d$. The functoriality of α implies $\alpha(f \circ \pi_n^d) = \alpha(f) \circ \pi_{\alpha(n)}^d$ for any $f \in \text{L}_{\text{gr}^0}(m, nd)$. Hence, $\alpha(\mathbb{I}_{\text{gr}^0}^{(d)}(m, n)) \subset \mathbb{I}_{\text{fr}_{\mathbb{Z}}^0}^{(d)}(m, n)$. Moreover, for any element of $\mathbb{I}_{\text{fr}_{\mathbb{Z}}^0}^{(d)}(m, n)$, say $g \circ \pi_{\alpha(n)}^d$ for some $g \in \text{L}_{\text{fr}_{\mathbb{Z}}^0}(m, nd)$, Lemma 14.8 implies $g \circ \pi_{\alpha(n)}^d = \alpha(\gamma(g)) \circ \alpha(\pi_n^d) = \alpha(\gamma(g) \circ \pi_n^d) \in \alpha(\mathbb{I}_{\text{gr}^0}^{(d)}(m, n))$ since α is a monad homomorphism. \square

We now recall the section γ of α in Definition 14.7.

Lemma 15.8. *For $n, m \in \mathbb{N}$, we have $\gamma(\mathbb{I}_{\text{fr}_{\mathbb{Z}}^0}^{(d)}(m, n)) \subset \mathbb{I}_{\text{gr}^0}^{(d)}(m, n)$.*

Proof. It suffices to prove that the map $\gamma : \mathbb{k}[\mathbf{A}_n^{\times m}] \rightarrow \mathbb{k}[\mathbf{F}_n^{\times m}]$ induced by the identifications (29) and (40) satisfies $\gamma(\mathbb{I}(\mathbf{A}_n^{\times m})^d) \subset \mathbb{I}_{\mathbb{k}}(\mathbf{F}_n^{\times m})^d$. To do that, we introduce a generator set Z of $\mathbb{k}[\mathbf{A}_n^{\times m}]$ with a total ordering \leq as follows:

- Let $z_{i,j}^{\pm 1} := (\overbrace{e, \dots, e}^{(i-1)}, \tau^{\pm e_j}, \overbrace{e, \dots, e}^{m-i}) \in Z$. If $i_0 < i_1$, then $z_{i_0, j}^{\epsilon} \leq z_{i_1, j'}^{\epsilon'}$ for any ϵ, ϵ' and j, j' .
- If $j_0 < j_1$, then $z_{i, j_0}^{\epsilon} \leq z_{i, j_1}^{\epsilon'}$ for any ϵ, ϵ' .
- $z_{i, j}^{-1} \leq z_{i, j}^{+1}$.

Note that such an ordering \leq is determined uniquely. Denote by $g_1 \leq g_2 \leq \dots \leq g_{2nm}$ all the elements of Z by considering the ordering. Then, by definition of γ , we have

$$(41) \quad \gamma\left(\prod_{k=1}^r g_{j_k}\right) = \prod_{k=1}^r \gamma(g_{j_k}),$$

for any sequence $1 \leq j_1 \leq j_2 \leq \dots \leq j_r \leq 2nm$ and $r \in \mathbb{N}$.

By definitions, $\mathbb{I}_{\mathbb{k}}(\mathbf{A}_n^{\times m})^d \subset \mathbb{k}[\mathbf{A}_n^{\times m}]$ is the submodule generated by $\prod_{k=1}^N (g_{i_k} - 1)$ for $N \geq d$ and $1 \leq i_1 \leq i_2 \leq \dots \leq i_N \leq 2nm$ since $\mathbf{A}_n^{\times m}$ is an abelian group. Hence, it is sufficient to show that $\gamma\left(\prod_{k=1}^N (g_{i_k} - 1)\right) \in \mathbb{I}_{\mathbb{k}}(\mathbf{F}_n^{\times m})^d$. If one expands $\prod_{k=1}^N (g_{i_k} - 1)$, then it is a sum of $\prod_{k=1}^r g_{j_k}$ for some sequence $1 \leq j_1 \leq j_2 \leq \dots \leq j_r \leq 2nm$ and $r \in \mathbb{N}$. By applying the equation (41) to such summands, we can factorize the sum to conclude

$$\gamma\left(\prod_{k=1}^N (g_{i_k} - 1)\right) = \prod_{k=1}^N (\gamma(g_{i_k}) - 1) \in \mathbb{I}_{\mathbb{k}}(\mathbf{F}_n^{\times m})^d.$$

\square

Lemma 15.9. *The homomorphism $\bar{\alpha} : \mathbb{I}_{\text{gr}^0}^{(d)}/\mathbb{I}_{\text{gr}^0}^{(d+1)} \rightarrow \mathbb{I}_{\text{fr}_{\mathbb{Z}}^0}^{(d)}/\mathbb{I}_{\text{fr}_{\mathbb{Z}}^0}^{(d+1)}$ is a split epimorphism. It is an isomorphism if and only if $d \in \{0, 1\}$.*

Proof. By Lemma 15.8, the homomorphism $\gamma : L_{\text{fr}_Z^0} \rightarrow L_{\text{gr}^0}$ induces $\bar{\gamma} : (L_{\text{fr}_Z^0}/\mathbb{I}_{\text{fr}_Z^0}^{(d)}) \rightarrow (L_{\text{gr}^0}/\mathbb{I}_{\text{gr}^0}^{(d)})$ which is a section of $\bar{\alpha}$ by Lemma 14.8.

Recall the notation in section 7.3. For $d = 0$, the statement follows from Proposition 7.18 since the category $(\text{gr}^0)_0$ is equivalent with $(\text{fr}_Z^0)_0$. Let $d = 1$. By the isomorphisms (29) and (40), it suffices to show that the following map gives an isomorphism:

$$\bar{\alpha} : \mathbb{I}_k(\mathbb{F}_n^{\times m})/\mathbb{I}_k(\mathbb{F}_n^{\times m})^2 \rightarrow \mathbb{I}_k(\mathbb{A}_n^{\times m})/\mathbb{I}_k(\mathbb{A}_n^{\times m})^2$$

This follows from Lemmas 9.3 and 15.3.

We now prove that, for $d \geq 2$, the homomorphism $\bar{\alpha} : \mathbb{I}_{\text{gr}^0}^{(d)}/\mathbb{I}_{\text{gr}^0}^{(d+1)} \rightarrow \mathbb{I}_{\text{fr}_Z^0}^{(d)}/\mathbb{I}_{\text{fr}_Z^0}^{(d+1)}$ is not an isomorphism. Recall the ranks of the domain and the codomain (indexwise) in Theorems 9.1 and 15.4. Note that we have an inequality for $nm \neq 0$:

$$\frac{(nm + d - 1)!}{d!(nm - 1)!} \leq n^d \cdot \frac{(m + d - 1)!}{d!(m - 1)!}.$$

It is easy to check that the equality holds if and only if either $d \in \{0, 1\}$ or $n = 1$. Therefore, if $d \geq 2$, then $\bar{\alpha}$ is not an isomorphism. \square

Theorem 15.10. *The epimorphism $\bar{\alpha} : E_{L_{\text{gr}^0}}(\mathbb{I}_{\text{gr}^0}^{(d+1)}) \rightarrow E_{L_{\text{fr}_Z^0}}(\mathbb{I}_{\text{fr}_Z^0}^{(d+1)})$ gives an isomorphism if and only if $d \in \{0, 1\}$.*

Proof. One may verify the first assertion directly by using γ . The second assertion for $d = 1$ is immediate from Lemma 15.9. To prove the case $d = 2$, we use the following commutative diagram such that the sequence in each row is exact:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{I}_{\text{gr}^0}^{(1)}/\mathbb{I}_{\text{gr}^0}^{(2)} & \longrightarrow & L_{\text{gr}^0}/\mathbb{I}_{\text{gr}^0}^{(2)} & \longrightarrow & L_{\text{gr}^0}/\mathbb{I}_{\text{gr}^0}^{(1)} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \mathbb{I}_{\text{fr}_Z^0}^{(1)}/\mathbb{I}_{\text{fr}_Z^0}^{(2)} & \longrightarrow & L_{\text{fr}_Z^0}/\mathbb{I}_{\text{fr}_Z^0}^{(2)} & \longrightarrow & L_{\text{fr}_Z^0}/\mathbb{I}_{\text{fr}_Z^0}^{(1)} & \longrightarrow & 0 \end{array}$$

By Lemma 15.9, the homomorphisms in the second and fourth columns (from the left side) are isomorphisms. We complete the proof by applying the 5-lemma. \square

Corollary 15.11. *The functor $\alpha^* : L_{\text{fr}_Z^0} \text{Mod}^{\leq d} \rightarrow L_{\text{gr}^0} \text{Mod}^{\leq d}$ gives an equivalence if and only if $d \in \{0, 1\}$.*

Proof. Let $T_1 = E_{L_{\text{gr}^0}}(\mathbb{I}_{\text{gr}^0}^{(d+1)})$ and $T_2 = E_{L_{\text{fr}_Z^0}}(\mathbb{I}_{\text{fr}_Z^0}^{(d+1)})$. By Corollary 7.15, it suffices to consider the functor $\bar{\alpha}^* : T_2 \text{Mod} \rightarrow T_1 \text{Mod}$ instead of α^* . By Theorem 15.10, it is clear that $\bar{\alpha}^*$ induces an equivalence of categories if $d \in \{0, 1\}$. This is what we mention in the beginning of this section. For $d > 1$, we claim that the functor $\bar{\alpha}^*$ is not essentially surjective. In fact, the homomorphism $\bar{\alpha} : T_1 \rightarrow T_2$ is an epimorphism with nontrivial kernel by Theorem 15.10. Hence, there exists $n, m \in \mathbb{N}$ such that $\bar{\alpha} : T_1(m, n) \rightarrow T_2(m, n)$ is surjective but not injective. We choose a nonzero $f \in T_1(m, n)$ lying in the kernel. If the left T_1 -module $T_1(-, n)$ lies in the essential image of $\bar{\alpha}^*$, then f should act on $T_1(-, n)$ trivially. Hence, for the unit $1_n \in T_1(n, n)$, we have $f \triangleright 1_n = 0$ which implies $f = 0$. There is a contradiction in the assumption that f is nonzero. \square

16. PRIMITIVELY GENERATED MODULES OVER FREE MODULES

In this section, we study the primitivity ideal $\mathbb{I}_{\text{fr}_R^0}^{\text{pr}}$ for fr_R^0 , which is introduced in Definition 11.1. Note that the category fr_R^0 fulfills the conditions on C in Definition 11.1. We investigate the structure of the associated eigenmonad based on a discussion analogous to section 11.

16.1. Eigenmonad associated with primitivity ideal. This section presents a proof that the canonical bimodule associated with the primitivity ideal for the category $\text{fr}_{\mathbb{Z}}^0$ is isomorphic to the \mathbb{k} -linearization of the category Fin of finite sets and maps. Furthermore, a generalization of this result based on a general unital ring R is provided. The main theorem of this section is then presented, preceded by a preliminary definition:

Definition 16.1. Let B be a \mathbb{k} -module. For a subcategory $\mathcal{D} \subset \text{Fin}$, we define a morphism $L_{\mathcal{D}}^B : \mathbb{N} \rightsquigarrow \mathbb{N}$ in $\text{Mat}_{\mathbb{k}}$, i.e. an $(\mathbb{N} \times \mathbb{N})$ -indexed module, to be

$$L_{\mathcal{D}}^B(m, n) := \bigoplus_{f \in \mathcal{D}(n, m)} B^{\otimes n}, \quad n, m \in \mathbb{N}.$$

For $f \in \mathcal{D}(n, m)$ and $b_j \in B$, we denote by $(b_1 \otimes \cdots \otimes b_n)_f$ the element of $L_{\mathcal{D}}^B(m, n)$ whose only nonzero component is the f -component given by $b_1 \otimes \cdots \otimes b_n \in B^{\otimes n}$.

Let B be a \mathbb{k} -algebra with unit 1_B . For the subcategory $\mathfrak{S} \subset \text{Fin}$ of bijections, we endow a monad structure on $L_{\mathfrak{S}}^B$ as follows:

- The unit $\eta : \mathbb{1}_{\mathbb{N}} \rightarrow L_{\mathfrak{S}}^B$ is characterized by $\eta(\text{id}_n) := (1_B^{\otimes n})_{\text{id}_n} \in L_{\mathfrak{S}}^B(n, n)$ for $n \in \mathbb{N}$.
- The monad operation is characterized by

$$(42) \quad (b'_1 \otimes \cdots \otimes b'_n)_g \circ (b_1 \otimes \cdots \otimes b_n)_f := (b_1 b'_{f(1)} \otimes \cdots \otimes b_n b'_{f(n)})_{g \circ f},$$

for $n \in \mathbb{N}$, $f, g \in \mathfrak{S}(n, n) = \mathfrak{S}_n$ and $b_j, b'_i \in B$.

Lemma 16.2. *The monad $L_{\mathfrak{S}}^B$ in $\text{Mat}_{\mathbb{k}}$ on \mathbb{N} is well-defined.*

Proof. Since any morphism in \mathfrak{S} is an bijective map, the formula (42) gives a well-defined \mathbb{k} -linear map

$$L_{\mathfrak{S}}^B(n, n) \otimes L_{\mathfrak{S}}^B(n, n) \rightarrow L_{\mathfrak{S}}^B(n, n),$$

which extends to a homomorphism $\nabla : L_{\mathfrak{S}}^B \otimes L_{\mathfrak{S}}^B \rightarrow L_{\mathfrak{S}}^B$. This satisfies the associativity:

$$\begin{aligned} \left(\left(\bigotimes_k b''_k \right)_h \circ \left(\bigotimes_j b'_j \right)_g \right) \circ \left(\bigotimes_i b_i \right)_f &= \left(\bigotimes_j b'_j b''_{g(j)} \right)_{h \circ g} \circ \left(\bigotimes_i b_i \right)_f, \\ &= \left(\bigotimes_i b_i b'_{f(i)} b''_{g(f(i))} \right)_{h \circ g \circ f}, \\ &= \left(\bigotimes_k b''_k \right)_h \circ \left(\bigotimes_i b_i b'_{f(i)} \right)_{g \circ f}, \\ &= \left(\bigotimes_k b''_k \right)_h \circ \left(\left(\bigotimes_j b'_j \right)_g \circ \left(\bigotimes_i b_i \right)_f \right). \end{aligned}$$

It is easy to check that $\eta : \mathbb{1}_{\mathbb{N}} \rightarrow L_{\mathfrak{S}}^B$ serves as the monad unit. □

Example 16.3. The above one is a generalization of the monad in Example 4.5. Indeed, there is an obvious isomorphism $L_{\mathfrak{S}}^{\mathbb{k}} \cong L_{\mathfrak{S}}$ of monads.

We are interested in the following algebra as B : let \hat{R} be the coefficient extension of R by \mathbb{k} , i.e.

$$\hat{R} := \mathbb{k} \otimes_{\mathbb{Z}} R.$$

We now recall the notation in Definition 11.1. The following sections present the proof of the following theorem.

Theorem 16.4. *We have a homomorphism $\mathcal{E}_R : L_{\text{Fin}}^{\hat{R}} \rightarrow L_{\text{fr}_R^0} / \mathbb{I}_{\text{fr}_R^0}^{\text{pr}}$ (see Definition 16.15) which gives the following isomorphisms:*

(1) We have an isomorphism of $(\mathbb{N} \times \mathbb{N})$ -indexed \mathbb{k} -modules:

$$\mathbf{L}_{\text{fr}_R^{\circ}} / \mathbf{I}_{\text{fr}_R^{\circ}}^{\text{pr}} \cong \mathbf{L}_{\text{Fin}}^{\hat{R}}.$$

(2) We have an isomorphism of monads in $\text{Mat}_{\mathbb{k}}$ on \mathbb{N} :

$$\mathbf{E}_{\mathbf{L}_{\text{fr}_R^{\circ}}}(\mathbf{I}_{\text{fr}_R^{\circ}}^{\text{pr}}) \cong \mathbf{L}_{\mathfrak{S}}^{\hat{R}}.$$

Proof. Let us defer the proof to later. Each statement is proved in sections 16.4 and 16.3 respectively. \square

Definition 16.5. Recall that $\mathbf{L}_{\text{fr}_R^{\circ}} / \mathbf{I}_{\text{fr}_R^{\circ}}^{\text{pr}}$ has a canonical bimodule structure by Definition 5.12. By Theorem 16.4, $\mathbf{L}_{\text{Fin}}^{\hat{R}}$ has a $(\mathbf{L}_{\text{fr}_R^{\circ}}, \mathbf{L}_{\mathfrak{S}}^{\hat{R}})$ -bimodule structure. We denote by $\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{R}}$ the bimodule.

Example 16.6. Let $R = \mathbb{Z}$. It is clear that $\hat{R} \cong \mathbb{k}$. The results in Theorem 16.4 lead to the isomorphisms

$$\begin{aligned} \mathbf{L}_{\text{fr}_{\mathbb{Z}}^{\circ}} / \mathbf{I}_{\text{fr}_{\mathbb{Z}}^{\circ}}^{\text{pr}} &\cong \mathbf{L}_{\text{Fin}}, \\ \mathbf{E}_{\mathbf{L}_{\text{fr}_{\mathbb{Z}}^{\circ}}}(\mathbf{I}_{\text{fr}_{\mathbb{Z}}^{\circ}}^{\text{pr}}) &\cong \mathbf{L}_{\mathfrak{S}}. \end{aligned}$$

By combining the results in section 11, we obtain the following well-known homomorphisms. Recall the abelianization $\alpha : \mathbf{L}_{\text{gr}^{\circ}} \rightarrow \mathbf{L}_{\text{fr}_{\mathbb{Z}}^{\circ}}$ in Definition 14.1. This induces a homomorphism

$$\tilde{\mathbf{A}}_{\mathfrak{S}^u} \cong \mathbf{L}_{\text{gr}^{\circ}} / \mathbf{I}^{\text{pr}} \rightarrow \mathbf{L}_{\text{fr}_{\mathbb{Z}}^{\circ}} / \mathbf{I}_{\text{fr}_{\mathbb{Z}}^{\circ}}^{\text{pr}} \cong \tilde{\mathbf{L}}_{\text{Fin}}$$

between the underlying indexed modules. The induced homomorphism is nothing but the one induced by $\mathfrak{S}^u \rightarrow \mathfrak{C}om^u$ under the isomorphism $\mathbb{k}\text{Fin} \cong \text{Cat}_{\mathfrak{C}om^u}$ where $\mathfrak{C}om^u$ is the operad for unital commutative algebras. Furthermore, we have $\alpha^{-1} \left(\mathbf{I}_{\text{fr}_{\mathbb{Z}}^{\circ}}^{\text{pr}} \right) = \mathbf{I}^{\text{pr}}$ (see Definition 11.1), so that we obtain a monad epimorphism

$$\mathbf{A}_{\mathfrak{S}^u} \cong \mathbf{E}_{\mathbf{L}_{\text{gr}^{\circ}}}(\mathbf{I}^{\text{pr}}) \cong \mathbf{V}_{\mathbf{I}^{\text{pr}}}(\mathbf{L}_{\text{gr}^{\circ}} / \mathbf{I}^{\text{pr}}) \rightarrow \mathbf{V}_{\mathbf{I}_{\text{fr}_{\mathbb{Z}}^{\circ}}^{\text{pr}}}(\mathbf{L}_{\text{fr}_{\mathbb{Z}}^{\circ}} / \mathbf{I}_{\text{fr}_{\mathbb{Z}}^{\circ}}^{\text{pr}}) \cong \mathbf{E}_{\mathbf{L}_{\text{gr}^{\circ}}}(\mathbf{I}^{\text{pr}}) \cong \mathbf{L}_{\mathfrak{S}}.$$

Corollary 16.7. *The eigenmonad adjunction gives an adjunction:*

$$\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{R}} \otimes_{\mathbf{L}_{\mathfrak{S}}^{\hat{R}}} (-) : \mathbf{L}_{\mathfrak{S}}^{\hat{R}} \text{Mod} \xrightleftharpoons[\mathbb{T}]{\mathbb{T}} \mathbf{L}_{\text{fr}_R^{\circ}} \text{Mod} : \mathbf{V}_{\mathbf{I}_{\text{fr}_R^{\circ}}^{\text{pr}}}.$$

Proposition 5.19 implies analogous statements based on $\text{Hom}_{\mathbf{L}_{\text{fr}_R^{\circ}}}(\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{R}}, -)$ instead of $\mathbf{V}_{\mathbf{I}_{\text{fr}_R^{\circ}}^{\text{pr}}}$.

Proof. It follows from Propositions 5.13, 5.22 and Theorem 16.4. \square

In the following, recall Definition 11.16.

Corollary 16.8. *A left $\mathbf{L}_{\text{fr}_R^{\circ}}$ -module \mathbf{M} is primitively generated if and only if the following evaluation map is an epimorphism:*

$$\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{R}} \otimes_{\mathbf{L}_{\mathfrak{S}}^{\hat{R}}} \text{Hom}_{\mathbf{L}_{\text{fr}_R^{\circ}}}(\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{R}}, \mathbf{M}) \rightarrow \mathbf{M}.$$

Proof. It is immediate from Corollary 16.7. The proof is parallel with that of Proposition 11.17. \square

16.2. Concrete description of the canonical bimodule. In advance of further discussion, we present a concrete description of the bimodule structure of $\tilde{\mathcal{L}}_{\text{Fin}}^{\hat{R}}$. Let us introduce the following notation.

Definition 16.9. For $1 \leq m \in \mathbb{N}$ and $1 \leq j \leq (m-1)$, we define $c_{m,j} \in \text{Fin}(m, m-1)$ to be the map such that

$$c_{m,j}(k) := \begin{cases} k & (k \leq j), \\ (k-1) & (k > j). \end{cases}$$

For $m \in \mathbb{N}$ and $0 \leq j \leq m$, we define $h_{m,j} \in \text{Fin}(m, m+1)$ to be

$$h_{m,j}(k) := \begin{cases} k & (k \leq j), \\ k+1 & (k > j). \end{cases}$$

Firstly, we explain *the left $\mathcal{L}_{\text{fr}_R^{\circ}}$ -module structure* of $\tilde{\mathcal{L}}_{\text{Fin}}^{\hat{R}}$. All the results here are derived from the isomorphism \mathcal{E}_R (see Definition 16.15) which yields Theorem 16.4. Let $(r_1 \otimes \cdots \otimes r_n)_f \in \tilde{\mathcal{L}}_{\text{Fin}}^{\hat{R}}(n, m)$ for $n, m \in \mathbb{N}$, $r_j \in R \subset \hat{R}$ and $f \in \text{Fin}(m, n)$.

- For $1 \leq j \leq m$, we have

$$(43) \quad \tau^{I_{j-1} \oplus [1 \ 1] \oplus I_{m-j}} \triangleright (r_1 \otimes \cdots \otimes r_n)_f = \sum_{\substack{f' \in \text{Fin}(n, m+1) \\ c_{m+1, j} \circ f' = f}} (r_1 \otimes \cdots \otimes r_n)_{f'}.$$

- Let X^t be the transposition of a matrix X . For $1 \leq j \leq (m-1)$, we have

$$\tau^{I_{j-1} \oplus [1 \ 1]^t \oplus I_{m-j-1}} \triangleright (r_1 \otimes \cdots \otimes r_n)_f = (r_1 \otimes \cdots \otimes r_n)_{c_{m, j} \circ f}.$$

- Let $\varrho \in \mathcal{M}_{0,1}(R)$ be the unique element. For $0 \leq j \leq m$, we have

$$(44) \quad \tau^{I_j \oplus \varrho \oplus I_{m-j}} \triangleright (r_1 \otimes \cdots \otimes r_n)_f = (r_1 \otimes \cdots \otimes r_n)_{h_{m, j} \circ f}.$$

- For $1 \leq j \leq m$, we have

$$(45) \quad \tau^{I_{j-1} \oplus \varrho^t \oplus I_{m-j}} \triangleright (r_1 \otimes \cdots \otimes r_n)_f = \sum_{\substack{f' \in \text{Fin}(n, m-1) \\ h_{m-1, j-1} \circ f' = f}} (r_1 \otimes \cdots \otimes r_n)_{f'}.$$

Note that, if such f' exists, then it is unique since $h_{m-1, j-1}$ is injective.

- For $1 \leq j \leq m$, we have

$$\tau^{I_{j-1} \oplus [a] \oplus I_{m-j}} \triangleright (r_1 \otimes \cdots \otimes r_n)_f = (r'_1 \otimes \cdots \otimes r'_n)_f.$$

where we set r'_i to be

$$r'_i := \begin{cases} r_i a & (f(i) = j), \\ r_i & (\text{otherwise}). \end{cases}$$

Note that the PROP fr_R° is generated by the morphisms $\tau^{[1 \ 1]} : 1 \rightarrow 2$, $\tau^{[1 \ 1]^t} : 2 \rightarrow 1$; the trivial morphisms $\tau^{\varrho} : 0 \rightarrow 1$, $\tau^{\varrho^t} : 1 \rightarrow 0$; and $\tau^{[a]} : 1 \rightarrow 1$ for $a \in R$. Thus, the left fr_R° -action on $\tilde{\mathcal{L}}_{\text{Fin}}^{\hat{R}}$ is described based on the above results in principle.

Secondly, we explain *the right $\mathcal{L}_{\mathfrak{S}}^{\hat{R}}$ -module structure* of $\tilde{\mathcal{L}}_{\text{Fin}}^{\hat{R}}$. For $n, m, f \in \text{Fin}(n, m)$ and $\sigma \in \mathfrak{S}(n, n) = \mathfrak{S}_n$, let $(b_1 \otimes \cdots \otimes b_n)_f \in \tilde{\mathcal{L}}_{\text{Fin}}^{\hat{R}}(m, n)$ and $(b'_1 \otimes \cdots \otimes b'_n)_{\sigma} \in \mathcal{L}_{\mathfrak{S}}^{\hat{R}}(n, n)$. Then we have

$$(b_1 \otimes \cdots \otimes b_n)_f \triangleleft (b'_1 \otimes \cdots \otimes b'_n)_{\sigma} = (b'_1 b_{\sigma(1)} \otimes \cdots \otimes b'_n b_{\sigma(n)})_{f \circ \sigma}.$$

16.3. Proof of (1) of Theorem 16.4. We now give an explicit construction of the isomorphism in Theorem 16.4, which is denoted by \mathcal{E}_R . To do that, it is useful to get used to the computation modulo $\mathbb{I}_{\text{fr}_R}^{\text{pr}}$.

Notation 16.10. For $v, w \in \mathbb{L}_{\text{fr}_R}^{\text{pr}}(m, n)$, we write $v \equiv_R w$ if $v - w \in \mathbb{I}_{\text{fr}_R}^{\text{pr}}(m, n)$.

Proposition 16.11. For $X_1, X_2, X_3 \in \mathbb{M}_{n,m}(R)$ such that

$$X_1 = \begin{bmatrix} r_{1,1} & \cdots & r_{1,m} \\ & \vdots & \\ r_{i,1} + r'_{i,1} & \cdots & r_{i,m} + r'_{i,m} \\ & \vdots & \\ r_{n,1} & \cdots & r_{n,m} \end{bmatrix}, X_2 = \begin{bmatrix} r_{1,1} & \cdots & r_{1,m} \\ & \vdots & \\ r_{i,1} & \cdots & r_{i,m} \\ & \vdots & \\ r_{n,1} & \cdots & r_{n,m} \end{bmatrix}, X_3 = \begin{bmatrix} r_{1,1} & \cdots & r_{1,m} \\ & \vdots & \\ r'_{i,1} & \cdots & r'_{i,m} \\ & \vdots & \\ r_{n,1} & \cdots & r_{n,m} \end{bmatrix},$$

we have

$$\tau^{X_1} \equiv_R \tau^{X_2} + \tau^{X_3}.$$

Proof. Let $X \in \mathbb{M}_{n+1,m}(R)$ be

$$X = \begin{bmatrix} r_{1,1} & \cdots & r_{1,m} \\ & \vdots & \\ r_{i,1} & \cdots & r_{i,m} \\ r'_{i,1} & \cdots & r'_{i,m} \\ r_{i+1,1} & \cdots & r_{i+1,m} \\ & \vdots & \\ r_{n,1} & \cdots & r_{n,m} \end{bmatrix}.$$

By Definition 11.1, we have

$$\theta_{\text{fr}_R}^{\text{pr}} = \tau^{[1 \ 1]} - \tau^{[1 \ 0]} - \tau^{[0 \ 1]}$$

so that

$$\begin{aligned} \tau^X \circ (\text{id}_{i-1} \oplus \theta_{\text{fr}_R}^{\text{pr}} \oplus \text{id}_{n-i}) &= \tau^X \circ (\tau^{I_{i-1}} \oplus \theta_{\text{fr}_R}^{\text{pr}} \oplus \tau^{I_{n-i}}), \\ &= \tau^X \circ (\tau^{I_{i-1} \oplus [1 \ 1] \oplus I_{n-i}} - \tau^{I_{i-1} \oplus [1 \ 0] \oplus I_{n-i}} - \tau^{I_{i-1} \oplus [0 \ 1] \oplus I_{n-i}}), \\ &= \tau^{X_1} - \tau^{X_2} - \tau^{X_3}. \end{aligned}$$

□

Definition 16.12. For a map $f \in \text{Fin}(n, m)$, we define a map $\tilde{v}_f : R^{\times n} \rightarrow \mathbb{M}_{n,m}(R)$ as follows. Let $r_1, \dots, r_n \in R$. The (i, j) -entry of the matrix $\tilde{v}_f(r_1, \dots, r_n)$ is defined to be r_i if $f(i) = j$; and 0 otherwise.

Lemma 16.13. Let $n, m \in \mathbb{N}$. The \mathbb{k} -module $(\mathbb{L}_{\text{fr}_R}^{\text{pr}} / \mathbb{I}_{\text{fr}_R}^{\text{pr}})(m, n)$ is generated by $\tau^{\tilde{v}_f(r_1, \dots, r_n)}$ mod $\mathbb{I}_{\text{fr}_R}^{\text{pr}}(m, n)$ for $r_j \in R$ and $f \in \text{Fin}(n, m)$.

Proof. The module $\mathbb{L}_{\text{fr}_R}^{\text{pr}}(m, n)$ is generated by τ^Y for $Y \in \mathbb{M}_{n,m}(R)$. Applying Proposition 16.11 to the i -th row, one may prove that $\tau^Y \pmod{\mathbb{I}_{\text{fr}_R}^{\text{pr}}(m, n)}$ is a linear combination of τ^Z 's where the i -th row of Z has at most one nonzero entry. We repeat this for every i . Then it turns out

that $\tau^Y \pmod{\mathbb{I}_{\text{tr}_R^{\text{pr}}}^{\text{pr}}(m, n)}$ is a linear combination of $\tau^{Z'}$'s where, for each i , the i -th row of Z' has at most one nonzero entry. Such $\tau^{Z'}$ is expressed as $\tilde{v}_f(r_1, \dots, r_n)$ for some $f \in \text{Fin}(n, m)$ and $r_j \in R$. \square

Note that \tilde{v}_f is not bilinear; but if we embed matrices into $\mathbb{L}_{\text{tr}_R^{\text{pr}}}$ by using $\tau^{(-)}$, then it becomes bilinear modulo \mathbb{I}^{pr} :

Lemma 16.14. *Let $n, m \in \mathbb{N}$ and $f \in \text{Fin}(n, m)$. There exists a well-defined \mathbb{k} -linear map $v_f : \hat{R}^{\otimes n} \rightarrow (\mathbb{L}_{\text{tr}_R^{\text{pr}}}/\mathbb{I}_{\text{tr}_R^{\text{pr}}}^{\text{pr}})(m, n)$ characterized by*

$$v_f(r_1 \otimes \cdots \otimes r_n) \equiv_R \tau^{\tilde{v}_f(r_1, \dots, r_n)},$$

for $r_j \in R \subset \hat{R}$.

Proof. It is sufficient to show that v_f gives a well-defined map $\overbrace{R \otimes_{\mathbb{Z}} \cdots \otimes_{\mathbb{Z}} R}^n = R^{\otimes_{\mathbb{Z}} n} \rightarrow (\mathbb{L}_{\text{tr}_R^{\text{pr}}}/\mathbb{I}_{\text{tr}_R^{\text{pr}}}^{\text{pr}})(m, n)$, since this always extends to $\hat{R}^{\otimes n} \cong \mathbb{k} \otimes_{\mathbb{Z}} R^{\otimes_{\mathbb{Z}} n}$ by the \mathbb{k} -linearity. By the universality of tensor product, it suffices to check

$$(46) \quad \tau^{\tilde{v}_f(r_1, \dots, (r_i + r'_i), \dots, r_n)} \equiv_R \tau^{\tilde{v}_f(r_1, \dots, r_i, \dots, r_n)} + \tau^{\tilde{v}_f(r_1, \dots, r'_i, \dots, r_n)}$$

for $1 \leq i \leq n$. Let us recall Definition 16.9. Then the equation (46) is proved by applying Proposition 16.11 to $g := f \circ c_{n+1, i}$ and $X := \tilde{v}_g(r_1, \dots, r_i, r'_i, \dots, r_n) \in \mathbb{M}_{n+1, m}(R)$. \square

Definition 16.15. We define a homomorphism $\mathcal{E}_R : \mathbb{L}_{\text{Fin}}^{\hat{R}} \rightarrow \mathbb{L}_{\text{tr}_R^{\text{pr}}}/\mathbb{I}_{\text{tr}_R^{\text{pr}}}^{\text{pr}}$ by using Lemma 16.14. For $n, m \in \mathbb{N}$, the map $\mathcal{E}_R : \mathbb{L}_{\text{Fin}}^{\hat{R}}(m, n) = \bigoplus_{f \in \text{Fin}(n, m)} \hat{R}^{\otimes n} \rightarrow (\mathbb{L}_{\text{tr}_R^{\text{pr}}}/\mathbb{I}_{\text{tr}_R^{\text{pr}}}^{\text{pr}})(m, n)$ is defined by

$$\mathcal{E}_R \left((b_1 \otimes \cdots \otimes b_n)_f \right) := v_f(b_1 \otimes \cdots \otimes b_n),$$

for $f \in \text{Fin}(n, m)$ and $b_j \in \hat{R}$.

Definition 16.16. We define a homomorphism $\mathcal{E}_R^{-1} : (\mathbb{L}_{\text{tr}_R^{\text{pr}}}/\mathbb{I}_{\text{tr}_R^{\text{pr}}}^{\text{pr}}) \rightarrow \mathbb{L}_{\text{Fin}}^{\hat{R}}$ which is proved to be the inverse of \mathcal{E}_R . For $n, m \in \mathbb{N}$, we define $\mathcal{E}_R^{-1} : (\mathbb{L}_{\text{tr}_R^{\text{pr}}}/\mathbb{I}_{\text{tr}_R^{\text{pr}}}^{\text{pr}})(m, n) \rightarrow \mathbb{L}_{\text{Fin}}^{\hat{R}}(m, n)$ by

$$\mathcal{E}_R^{-1}(\tau^X) := \sum_{f \in \text{Fin}(n, m)} (r_{1f(1)} \otimes r_{2f(2)} \otimes \cdots \otimes r_{nf(n)})_f.$$

where $X = [r_{ij}]_{i \in \underline{n}, j \in \underline{m}} \in \mathbb{M}_{n, m}(R)$.

Lemma 16.17. *The homomorphism \mathcal{E}_R^{-1} is well-defined, and it gives the inverse of \mathcal{E}_R .*

Proof. To prove the well-definedness, it suffices to show that $\mathcal{E}_R^{-1}(\tau^X \circ (\text{id}_{i-1} \oplus \theta_{\text{tr}_R^{\text{pr}}} \oplus \text{id}_{n-i})) = 0$ for $1 \leq i \leq n$ and $X \in \mathbb{M}_{n+1, m}(R)$. By applying (38), it is equivalent with

$$\mathcal{E}_R^{-1}(\tau^{(I_{i-1} \oplus [1 \ 1] \oplus I_{n-i})X} - \tau^{(I_{i-1} \oplus [0 \ 1] \oplus I_{n-i})X} + \tau^{(I_{i-1} \oplus [1 \ 0] \oplus I_{n-i})X}) = 0.$$

This is immediate from the definition of \mathcal{E}_R^{-1} since $\tau^{(I_{i-1} \oplus [1 \ 1] \oplus I_{n-i})X} \in \mathbb{M}_{n, m}(R)$ is the matrix whose i -th row is the sum of i -th and $(i+1)$ -th rows of X ; and all other rows are the same with those of X .

We now prove that $\mathcal{E}_R^{-1}(\mathcal{E}_R((r_1 \otimes \cdots \otimes r_n)_g)) = (r_1 \otimes \cdots \otimes r_n)_g$ for $g \in \text{Fin}(n, m)$ and $r_j \in R \subset \hat{R}$.

$$\begin{aligned} \mathcal{E}_R^{-1}(\mathcal{E}_R((r_1 \otimes \cdots \otimes r_n)_g)) &= \mathcal{E}_R^{-1}(v_g(r_1 \otimes \cdots \otimes r_n)), \\ &= \mathcal{E}_R^{-1}(\tau^{\tilde{v}_g(r_1, \dots, r_n)}). \end{aligned}$$

By Definition 16.12, we have $\mathcal{E}_R^{-1}(\tau^{\tilde{v}_g(r_1, \dots, r_n)}) = (r_1 \otimes \cdots \otimes r_n)_g$.

The above result implies that \mathcal{E}_R is a monomorphism. By Lemma 16.13, \mathcal{E}_R is an epimorphism. Therefore, \mathcal{E}_R is an isomorphism. \square

16.4. Proof of (2) of Theorem 16.4. In this section, we prove (2) of Theorem 16.4 by computing the eigenmonad associated with the primitivity ideal for fr_R^0 . To do that, we freely use the concrete description of the left $L_{\text{fr}_R^0}$ -action on $\tilde{L}_{\text{Fin}}^{\hat{R}}$ in section 16.2. In the following, let us recall Definition 16.5.

Lemma 16.18. *For $m, n \in \mathbb{N}$, the $\mathbb{I}_{\text{fr}_R^0}^{\text{pr}}$ -vanishing module $(V_{\mathbb{I}_{\text{fr}_R^0}^{\text{pr}}}(\tilde{L}_{\text{Fin}}^{\hat{R}}))(m, n)$ is isomorphic to $L_{\mathfrak{E}}^{\hat{R}}(m, n)$.*

Proof. To prove our statement, we compute the intersection of the kernels of the left action of $\text{id}_{l-1} \oplus \theta_{\text{fr}_R^0} \oplus \text{id}_{m-l}$ on $\tilde{L}_{\text{Fin}}^{\hat{R}}(m, n)$ for $1 \leq l \leq m$. Here, we consider the case of $l = 1$; and other cases are shown similarly. Let $f \in \text{Fin}(n, m)$ and $(r_1 \otimes \cdots \otimes r_n)_f \in L_{\text{Fin}}^{\hat{R}}(m, n)$. By (44), we have

$$\begin{aligned} (\tau^{[1 \ 0]} \oplus \text{id}_{m-1}) \triangleright (r_1 \otimes \cdots \otimes r_n)_f &= \tau^{l_1 \oplus 0 \oplus l_{n-1}} \triangleright (r_1 \otimes \cdots \otimes r_n)_f, \\ &= (r_1 \otimes \cdots \otimes r_n)_{h_{m,1} \circ f}. \end{aligned}$$

Note that $h_{m,1} \circ f$ is the unique map $g \in \text{Fin}(n, m+1)$ such that $c_{m+1,1} \circ g = f$, $g^{-1}(2) = \emptyset$. Similarly, we obtain

$$(\tau^{[0 \ 1]} \oplus \text{id}_{m-1}) \triangleright (r_1 \otimes \cdots \otimes r_n)_f = (r_1 \otimes \cdots \otimes r_n)_{h_{m,0} \circ f},$$

and $h_{m,0} \circ f$ is the unique map $g \in \text{Fin}(n, m+1)$ such that $c_{m+1,1} \circ g = f$, $g^{-1}(1) = \emptyset$. Thus, by (43) and the definition of $\theta_{\text{fr}_R^0}$, we have

$$(\theta_{\text{fr}_R^0} \oplus \text{id}_{m-1}) \triangleright (r_1 \otimes \cdots \otimes r_n)_f = \sum_{\substack{c_{m+1,1} \circ g = f \\ g^{-1}(1) \neq \emptyset \neq g^{-1}(2)}} (r_1 \otimes \cdots \otimes r_n)_g.$$

Thus, the kernel of $(\theta_{\text{fr}_R^0} \oplus \text{id}_{m-1}) \triangleright (-) : L_{\text{Fin}}^{\hat{R}}(m, n) \rightarrow L_{\text{Fin}}^{\hat{R}}(m+1, n)$ is generated by $(r_1 \otimes \cdots \otimes r_n)_f$ for $f \in \text{Fin}(n, m)$ such that there does not exist $g \in \text{Fin}(n, m+1)$ satisfying $c_{m+1,1} \circ g = f$ and $g^{-1}(1) \neq \emptyset \neq g^{-1}(2)$. The latter condition is equivalent with $|f^{-1}(1)| = 1$. Similarly, the kernel of the left action of $\text{id}_l \oplus \theta_{\text{fr}_R^0} \oplus \text{id}_{m-l-1}$ is generated by $(r_1 \otimes \cdots \otimes r_n)_f$ for $f \in \text{Fin}(n, m)$ such that $|f^{-1}(l)| = 1$. Therefore, we obtain

$$\bigoplus_{\substack{f \in \text{Fin}(n, m) \\ |f^{-1}(l)| = 1, \forall l \in \underline{m}}} \hat{R}^{\otimes n} \cong V_{\mathbb{I}_{\text{fr}_R^0}^{\text{pr}}}(\tilde{L}_{\text{Fin}}^{\hat{R}})(m, n).$$

The left hand side is $L_{\mathfrak{E}}^{\hat{R}}(m, n)$ by the definition. \square

We now present a proof of (2) of Theorem 16.4. We first give an isomorphism as $(\mathbb{N} \times \mathbb{N})$ -indexed \mathbb{k} -modules. By Proposition 5.20 and Theorem 16.4, it suffices to construct an isomorphism between the $\mathbb{I}_{\text{fr}_R^0}^{\text{pr}}$ -vanishing module of $\tilde{L}_{\text{Fin}}^{\hat{R}}$ and $L_{\mathfrak{E}}^{\hat{R}}$. This is done in Lemma 16.18. We now have to verify that this isomorphism preserves the monad structure. For $n \in \mathbb{N}$, let $\sigma \in \mathfrak{S}(n, n)$. Then, for $r_1, \dots, r_n \in R$, $(r_1 \otimes \cdots \otimes r_n)_{\sigma} \in L_{\mathfrak{E}}^{\hat{R}}(n, n)$ corresponds to $\mathcal{E}_R((r_1 \otimes \cdots \otimes r_n)_{\sigma}) = \tau^{\tilde{\nu}_{\sigma}(r_1, \dots, r_n)} \text{ mod } \mathbb{I}_{\text{fr}_R^0}^{\text{pr}} \in \left(E_{L_{\text{fr}_R^0}^{\text{pr}}}(\mathbb{I}_{\text{fr}_R^0}^{\text{pr}}) \right)(n, n)$ via the above isomorphism. We now consider $\sigma_1, \sigma_2 \in \mathfrak{S}(n, n)$; and $r_j, r'_j \in R$ for $j = 1, \dots, n$. It is elementary to validate that

$$\tilde{\nu}_{\sigma_1}(r_1, \dots, r_n) \cdot \tilde{\nu}_{\sigma_2}(r'_1, \dots, r'_n) = \tilde{\nu}_{\sigma_2 \circ \sigma_1}(r_1 r'_{\sigma_1(1)}, \dots, r_n r'_{\sigma_1(n)}).$$

On the one hand, we have $(r'_1 \otimes \cdots \otimes r'_n)_{\sigma_2} \circ (r_1 \otimes \cdots \otimes r_n)_{\sigma_1} = (r_1 r'_{\sigma_1(1)} \otimes \cdots \otimes r_n r'_{\sigma_1(n)})_{\sigma_2 \circ \sigma_1}$ in the monad $L_{\mathfrak{E}}^{\hat{R}}$ by definitions.

16.5. Refined eigenmonad adjunctions. Recall that, for a monad T in $\text{Mat}_{\mathbb{k}}$ and its left ideal J , a left $E_T(J)$ -module N is $\mathbb{I}_{\text{fr}_R}^{\text{pr}}$ -vanishingly extensible if the counit $N \rightarrow T/J \otimes_{E_T(J)} N$ is a monomorphism. The goal of this section is to prove that every left $E_{L_{\text{fr}_R}^{\text{pr}}}(\mathbb{I}_{\text{fr}_R}^{\text{pr}})$ -module is $\mathbb{I}_{\text{fr}_R}^{\text{pr}}$ -vanishingly extensible.

In the following, let us recall the notation in Definition 16.1.

Definition 16.19. Let B be a \mathbb{k} -algebra. We define a homomorphism $r : \tilde{L}_{\text{Fin}}^B \rightarrow L_{\mathfrak{S}}^B$. For each $n, m \in \mathbb{N}$, the map $r : \tilde{L}_{\text{Fin}}^B(m, n) \rightarrow L_{\mathfrak{S}}^B(m, n)$ is defined by

$$r \left((b_1 \otimes \cdots \otimes b_n)_f \right) := \begin{cases} (b_1 \otimes \cdots \otimes b_n)_f & (n = m, \text{ and } f \in \mathfrak{S}(n, n)), \\ 0 & (\text{otherwise}), \end{cases}$$

where $b_1, \dots, b_n \in B$ and $f \in \text{Fin}(n, m)$.

Lemma 16.20. *The homomorphism r preserves the right $L_{\mathfrak{S}}^B$ -action, and the diagram below commutes:*

$$\begin{array}{ccc} L_{\mathfrak{S}}^B & \xrightarrow{\quad} & \tilde{L}_{\text{Fin}}^B \\ & \searrow & \downarrow r \\ & & L_{\mathfrak{S}}^B \end{array}$$

Proof. It follows from definition that the diagram commutes. We prove that r preserves the right $L_{\mathfrak{S}}^B$ -action. Let $r_j \in R$, $f \in \text{Fin}(n, m)$ and $\sigma \in \mathfrak{S}(n, n) = \mathfrak{S}_n$.

$$\begin{aligned} & r \left((b_1 \otimes \cdots \otimes b_n)_f \triangleleft (b'_1 \otimes \cdots \otimes b'_n)_{\sigma} \right), \\ &= r \left((b'_1 b_{\sigma(1)} \otimes \cdots \otimes b'_n b_{\sigma(n)})_{f \circ \sigma} \right), \\ &= \begin{cases} (b'_1 b_{\sigma(1)} \otimes \cdots \otimes b'_n b_{\sigma(n)})_{f \circ \sigma}, & (n = m, \text{ and } f \in \mathfrak{S}(n, n)) \\ 0, & (\text{otherwise}), \end{cases} \\ &= r \left((b_1 \otimes \cdots \otimes b_n)_f \triangleleft (b'_1 \otimes \cdots \otimes b'_n)_{\sigma} \right). \end{aligned}$$

□

Theorem 16.21. *Every left $E_{L_{\text{fr}_R}^{\text{pr}}}(\mathbb{I}_{\text{fr}_R}^{\text{pr}})$ -module is $\mathbb{I}_{\text{fr}_R}^{\text{pr}}$ -vanishingly extensible.*

Proof. Let N be a left $E_{L_{\text{fr}_R}^{\text{pr}}}(\mathbb{I}_{\text{fr}_R}^{\text{pr}})$ -module, equivalently a left $L_{\mathfrak{S}}^{\hat{R}}$ -module by (2) of Theorem 16.4. It suffices to prove that the counit $N \rightarrow V_{\mathbb{I}_{\text{fr}_R}^{\text{pr}}} \left((L_{\text{fr}_R}^{\text{pr}}/\mathbb{I}_{\text{fr}_R}^{\text{pr}}) \otimes_{L_{\mathfrak{S}}^{\hat{R}}} N \right)$ is a monomorphism. By Lemma 16.20, r extends to a retract of the counit $N \rightarrow (L_{\text{fr}_R}^{\text{pr}}/\mathbb{I}_{\text{fr}_R}^{\text{pr}}) \otimes_{L_{\mathfrak{S}}^{\hat{R}}} N \cong \tilde{L}_{\text{Fin}}^{\hat{R}} \otimes_{L_{\mathfrak{S}}^{\hat{R}}} N$. □

Let ${}_{L_{\text{fr}_R}^{\text{pr}}} \text{Mod}^{\text{prim}}$ be the full subcategory of ${}_{L_{\text{fr}_R}^{\text{pr}}} \text{Mod}$ generated by primitively generated modules.

Corollary 16.22. *The refined eigenmonad adjunction (see Proposition 6.7) gives an adjunction:*

$$\tilde{L}_{\text{Fin}}^{\hat{R}} \otimes_{L_{\mathfrak{S}}^{\hat{R}}} (-) : {}_{L_{\mathfrak{S}}^{\hat{R}}} \text{Mod} \xleftarrow{\quad} \xrightarrow{\quad} {}_{L_{\text{fr}_R}^{\text{pr}}} \text{Mod}^{\text{prim}} : V_{\mathbb{I}_{\text{fr}_R}^{\text{pr}}}.$$

Proof. It follows from Propositions 16.4 and 16.21. □

Example 16.23. For general \mathbb{k} , the adjunction in Corollary 16.22 does not give an equivalence of categories. We give an example such that the counit is not a natural isomorphism. Suppose that \mathbb{k} is a ring such that $2 \cdot 1_{\mathbb{k}} = 0$; and $R = \mathbb{Z}$. For a \mathbb{k} -module W , we define $\bigwedge W \in {}_{L_{\mathfrak{S}}} \text{Mod}$ to

be $(\bigwedge W)(n) := \bigwedge^n W$, $\forall n \in \mathbb{N}$ where the symmetric group action is induced by the signature. For the unique map $f_n \in \text{Fin}(n, 1)$, the map $\bigwedge^n W \rightarrow \tilde{\mathbf{L}}_{\text{Fin}}^{\hat{Z}}(1, n) \otimes_{\mathcal{E}_n} \bigwedge^n W$; $z \mapsto f_n \otimes z$ gives an isomorphism by the hypothesis $2 \cdot 1_{\mathbb{k}} = 0$ where we identify $\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{Z}}$ with \mathbf{L}_{Fin} (see Example 16.3). Hence, we have an isomorphism

$$\bigoplus_{n \in \mathbb{N}} \bigwedge^n W \cong (\tilde{\mathbf{L}}^{\hat{Z}} \otimes_{\mathcal{L}_{\mathcal{E}}} \bigwedge W)(1).$$

We now compute the action of $\theta_{\text{fr}_Z^0} \in \mathbf{L}_{\text{fr}_Z^0}(2, 1)$ on $f_2 \otimes (w_1 \wedge w_2) \in \mathbf{L}_{\text{Fin}}(1, 2) \otimes_{\mathcal{E}_2} \bigwedge^2 W \subset (\tilde{\mathbf{L}}^{\hat{Z}} \otimes_{\mathcal{L}_{\mathcal{E}}} \bigwedge W)(1)$. By using the concrete description of the bimodule $\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{Z}}$ in section 16.2, we have

$$\begin{aligned} \theta_{\text{fr}_Z^0} \triangleright (f_2 \otimes (w_1 \wedge w_2)) &= \text{id}_2 \otimes (w_1 \wedge w_2) + \sigma \otimes (w_1 \wedge w_2), \\ &= \text{id}_2 \otimes (w_1 \wedge w_2) + \text{id}_2 \otimes \sigma_*(w_1 \wedge w_2), \\ &= \text{id}_2 \otimes (w_1 \wedge w_2) - \text{id}_2 \otimes (w_1 \wedge w_2) = 0. \end{aligned}$$

where $\sigma \in \text{Fin}(2, 2)$ is the nontrivial permutation. This implies that $\bigwedge^2 W \subset \mathbf{V}_{\text{fr}_Z^0}^{\text{pr}}(\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{Z}} \otimes_{\mathcal{L}_{\mathcal{E}}} \bigwedge W)(1) \cong \bigoplus_{n \in \mathbb{N}} \bigwedge^n W$. Hence, we obtain

$$(\bigwedge W)(1) \subsetneq \mathbf{V}_{\text{fr}_Z^0}^{\text{pr}}(\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{Z}} \otimes_{\mathcal{L}_{\mathcal{E}}} \bigwedge W)(1).$$

17. COMPARISON OF PRIMITIVITY AND ANALYTICITY

In this section, we prove that primitively generated $\mathbf{L}_{\text{fr}_R^0}$ -modules are analytic. This result is derived from a more robust argument with respect to the associated left ideals.

Lemma 17.1. *Let $n \in \mathbb{N}$. The polynomial degree of the left $\mathbf{L}_{\text{fr}_R^0}$ -module $\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{R}}(-, n)$ does not exceed n .*

Proof. One may prove the statement by calculating the associated cross-effects [12]; but here, we give a proof based on Theorem 7.9. We prove that $\mathbf{V}_{\text{fr}_R^0}^{(n+1)}(\mathbf{F}) = \mathbf{F}$ for $\mathbf{F} := \tilde{\mathbf{L}}_{\text{Fin}}^{\hat{R}}(-, n)$. For the embedding $\iota = R \otimes_{\mathbb{Z}} (-) : \text{fr}_{\mathbb{Z}} \hookrightarrow \text{fr}_R$, we have $\pi_m^{\text{fr}_R^0, d+1} = \iota(\pi_m^{\text{fr}_{\mathbb{Z}}^0, d+1})$ for any object m of $\text{fr}_{\mathbb{Z}}$ (see Definition 7.4). Hence, if we write $\iota^* \mathbf{M}$ the induced $\mathbf{L}_{\text{fr}_Z^0}$ -module for a left $\mathbf{L}_{\text{fr}_R^0}$ -module \mathbf{M} , we have $\mathbf{V}_{\text{fr}_R^0}^{(d+1)}(\mathbf{M}) = \mathbf{V}_{\text{fr}_Z^0}^{(d+1)}(\iota^* \mathbf{M})$. It implies that we need to focus on the $\mathbf{L}_{\text{fr}_Z^0}$ -module $\iota^* \mathbf{F}$ for the left $\mathbf{L}_{\text{fr}_R^0}$ -module \mathbf{F} to discuss the polynomial degree of \mathbf{F} . Note that, by Definition 16.1, we have an isomorphism $\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{R}}(-, n) \cong \hat{R}^{\otimes n} \otimes \tilde{\mathbf{L}}_{\text{Fin}}^{\hat{Z}}(-, n)$ indexwise. By the concrete description of the $\mathbf{L}_{\text{fr}_R^0}$ -module in section 16.2, this gives an isomorphism of $\mathbf{L}_{\text{fr}_Z^0}$ -modules, and leads to the following:

$$\mathbf{V}_{\text{fr}_Z^0}^{(d+1)}(\iota^* \mathbf{F}) \cong \hat{R}^{\otimes n} \otimes \mathbf{V}_{\text{fr}_Z^0}^{(d+1)}(\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{Z}}(-, n)).$$

We now investigate the vanishing module $\mathbf{V}_{\text{fr}_Z^0}^{(d+1)}(\tilde{\mathbf{L}}_{\text{Fin}}^{\hat{Z}}(-, n))$. To do that, we introduce an embedding $u : \tilde{\mathbf{L}}_{\text{Fin}}^{\hat{Z}}(-, n) \rightarrow \mathbf{G}$. Here, \mathbf{G} is a left $\mathbf{L}_{\text{fr}_Z^0}$ -module determined as follows. Let $\mathbb{k}[t_1, \dots, t_n]$ be the bicommutative Hopf algebra induced by polynomials with variables t_1, \dots, t_n and coefficients in \mathbb{k} . Then the assignment of $\mathbb{k}[t_1, \dots, t_n]^{\otimes m}$ to $m \in \mathbb{N}$ yields a (symmetric monoidal) functor from $\text{fr}_{\mathbb{Z}}$ to the tensor category ${}_{\mathbb{k}}\text{Mod}$. We define \mathbf{G} to be the corresponding left $\mathbf{L}_{\text{fr}_Z^0}$ -module [12]. We assign

$$(47) \quad \prod_{i_1 \in f^{-1}(1)} t_{i_1} \otimes \prod_{i_2 \in f^{-1}(2)} t_{i_2} \otimes \cdots \otimes \prod_{i_m \in f^{-1}(m)} t_{i_m} \in \mathbf{G}(m)$$

to $f \in \text{Fin}(n, m) \subset \tilde{\mathbb{L}}_{\text{Fin}}^{\hat{\mathbb{Z}}}(m, n)$. This induces a monomorphic $L_{\text{fr}_{\mathbb{Z}}^{\circ}}$ -homomorphism $u : \tilde{\mathbb{L}}_{\text{Fin}}^{\hat{\mathbb{Z}}}(-, n) \rightarrow \mathbb{G}$. Note that $\mathbb{G}(m) = \mathbb{k}[t_1, \dots, t_n]^{\otimes m}$ is the m -fold tensor product of the Hopf algebra $\mathbb{k}[t_1, \dots, t_n]$. Denote by $\bar{\Delta}_{\mathbb{G}(m)}$ the reduced comultiplication. For an object $m \in \text{fr}_{\mathbb{Z}}^{\circ}$, the action of $\pi_m^{\text{fr}_{\mathbb{Z}}^{\circ}, d+1}$ on \mathbb{G} coincides with the $(d+1)$ -fold reduced comultiplication $\bar{\Delta}_{H_m}^{(d+1)} : \mathbb{G}(m) \rightarrow \mathbb{G}(m)^{\otimes (d+1)}$. It is easy to see that the element of the form (47) vanishes under $\bar{\Delta}_{\mathbb{G}(m)}^{(n+1)}$. In other words, the kernel of $\bar{\Delta}_{\mathbb{G}(m)}^{(n+1)}$ contains the image of $u : \tilde{\mathbb{L}}_{\text{Fin}}^{\hat{\mathbb{Z}}}(n+1, n) \rightarrow \mathbb{G}(n+1)$. Thus, we obtain $V_{\mathbb{I}_{\text{fr}_{\mathbb{Z}}^{\circ}}^{(n+1)}}(\tilde{\mathbb{L}}_{\text{Fin}}^{\hat{\mathbb{Z}}}(-, n)) = \tilde{\mathbb{L}}_{\text{Fin}}^{\hat{\mathbb{Z}}}(-, n)$ since u is a monomorphic $L_{\text{fr}_{\mathbb{Z}}^{\circ}}$ -homomorphism $u : \tilde{\mathbb{L}}_{\text{Fin}}^{\hat{\mathbb{Z}}}(-, n) \rightarrow \mathbb{G}$. \square

Remark 17.2. In fact, the polynomial degree of $\tilde{\mathbb{L}}_{\text{Fin}}^{\hat{R}}(-, n)$ is explicitly n . In other words, there exists an element of $\tilde{\mathbb{L}}_{\text{Fin}}^{\hat{R}}(n, n)$ which does not vanish under the action of $\pi_n^{\text{fr}_R^{\circ}, n}$. For $u_n : \tilde{\mathbb{L}}_{\text{Fin}}^{\hat{\mathbb{Z}}}(n, n) \rightarrow \mathbb{G}(n)$ (in the previous proof), we have $u_n(\text{id}_n) = (t_1 \otimes t_2 \otimes \dots \otimes t_n) \in \mathbb{G}(n)$ which does not lie in the kernel of $\bar{\Delta}_{\mathbb{G}(n)}^{(n)}$. Hence, the polynomial degree of $\tilde{\mathbb{L}}_{\text{Fin}}^{\hat{R}}(-, n)$ exceed $(n-1)$.

In the following, we recall the left ideal \mathbb{I}_C^{ν} in Definition 7.21.

Theorem 17.3. *Let $\nu_0 : \mathbb{N} \rightarrow \mathbb{N}$ such that $\nu_0(n) = n+1$. The primitivity ideal contains the ν_0 -analyticity ideal:*

$$\mathbb{I}_{\text{fr}_R^{\circ}}^{\nu_0} \subset \mathbb{I}_{\text{fr}_R^{\circ}}^{\text{pr}}.$$

Proof. The proof is parallel with that of Theorem 12.1. Instead of using the fact in [12], we apply Lemma 17.1. \square

Corollary 17.4. *Let \mathbb{M} be a left $L_{\text{fr}_R^{\circ}}$ -module. If \mathbb{M} is primitively generated, then it is analytic.*

Proof. By Theorem 17.3, if a left $L_{\text{fr}_R^{\circ}}$ -module \mathbb{M} is primitive, then it is $\mathbb{I}_{\text{fr}_R^{\circ}}^{\nu_0}$ -vanishingly generated. Thus, our assertion follows from Theorem 7.25. \square

Before we close this section, we explain a special case that the analyticity and the primitivity become equivalent to each other.

Proposition 17.5. *Suppose that the ground ring \mathbb{k} is a field of characteristic zero. A left $L_{\text{fr}_{\mathbb{Z}}^{\circ}}$ -module is analytic if and only if it is primitively generated.*

Proof. Let $\alpha : L_{\text{gr}^{\circ}} \rightarrow L_{\text{fr}_{\mathbb{Z}}^{\circ}}$ be the abelianization. We have $\alpha(\mathbb{I}^{\text{pr}}) = \mathbb{I}_{\text{fr}_{\mathbb{Z}}^{\circ}}^{\text{pr}}$ since α is an epimorphism where \mathbb{I}^{pr} is the primitivity ideal for free groups (see (30)). Hence, a left $\text{fr}_{\mathbb{Z}}^{\circ}$ -module \mathbb{M} is primitively generated if and only if the left $L_{\text{gr}^{\circ}}$ -module $\alpha^*(\mathbb{M})$ is primitively generated. Similarly, we have $\alpha(\mathbb{I}_{\text{gr}^{\circ}}^{\nu}) = \mathbb{I}_{\text{fr}_{\mathbb{Z}}^{\circ}}^{\nu}$ for any map $\nu : \mathbb{N} \rightarrow \mathbb{N}$ where \mathbb{I}_C^{ν} is the ν -analyticity ideal (see Definition 7.21). Thus, \mathbb{M} is analytic as an $L_{\text{fr}_{\mathbb{Z}}^{\circ}}$ -module if and only if $\alpha^*(\mathbb{M})$ is analytic as an $L_{\text{gr}^{\circ}}$ -module. Therefore, by Corollary 12.3, a left $L_{\text{fr}_{\mathbb{Z}}^{\circ}}$ -module \mathbb{M} is analytic if and only if it is primitively generated. \square

Corollary 17.6. *Suppose that the ground ring \mathbb{k} is a field of characteristic zero. The refined eigenmonad adjunction gives an adjoint equivalence:*

$$\tilde{\mathbb{L}}_{\text{Fin}}^{\hat{\mathbb{Z}}} \otimes_{L_{\varepsilon}} (-) : L_{\varepsilon} \mathbf{Mod} \xrightleftharpoons[\text{pr}]{\text{pr}} L_{\text{fr}_{\mathbb{Z}}^{\circ}} \mathbf{Mod}^{\text{prim}} : V_{\mathbb{I}_{\text{fr}_{\mathbb{Z}}^{\circ}}^{\text{pr}}}.$$

Proof. The adjunction follows from Corollary 16.22 and the observation in Example 16.6. The functor $\tilde{\mathbb{L}}_{\text{Fin}}^{\hat{\mathbb{Z}}} \otimes_{L_{\varepsilon}} (-) : L_{\varepsilon} \mathbf{Mod} \rightarrow L_{\text{fr}_{\mathbb{Z}}^{\circ}} \mathbf{Mod}$ yields an equivalence between the categories of L_{ε} -modules and analytic $L_{\text{fr}_{\mathbb{Z}}^{\circ}}$ -modules [29, section 2.4]. By combining this fact with Corollary 17.5, it turns out that the refined eigenmonad adjunction gives an adjoint equivalence. \square

APPENDIX A. ENOUGH INJECTIVES AND PROJECTIVES IN ${}_{\mathbb{T}}\text{Mod}$

Let \mathbb{T} be a monad in $\text{Mat}_{\mathbb{k}}$ on an object \mathcal{X} . In this section, we prove that the category ${}_{\mathbb{T}}\text{Mod}$ of left \mathbb{T} -modules and \mathbb{T} -homomorphisms has enough injectives and projectives. The proof is based on a slight modification of the classical one. This allows Definition 4.22 to be well-defined.

Lemma A.1. *The category ${}_{\mathbb{T}}\text{Mod}$ has enough injectives.*

Proof. This is proved by carefully refining [30, Example 2.3.7]. We would like to apply the fact that the category ${}_{\mathbb{k}}\text{Mod}$ has enough injectives, and thus we prepare a preliminary. For $X \in \mathcal{X}$, consider the evaluation functor $\text{ev}_X : {}_{\mathbb{T}}\text{Mod} \rightarrow {}_{\mathbb{k}}\text{Mod}; \mathbb{M} \mapsto \mathbb{M}(X)$. It is easy to see that this is a left adjoint:

$$\text{ev}_X : {}_{\mathbb{T}}\text{Mod} \xleftarrow[{}_{\mathbb{T}}]{\quad} {}_{\mathbb{k}}\text{Mod} : \text{ev}_X^{\mathbb{R}}.$$

In fact, the right adjoint $\text{ev}_X^{\mathbb{R}}$ is given by

$$(\text{ev}_X^{\mathbb{R}}(V))(Y) := \text{hom}_{\mathbb{k}}(\mathbb{T}(X, Y), V), \quad V \in {}_{\mathbb{k}}\text{Mod}.$$

The regular right \mathbb{T} -action on \mathbb{T} gives a left \mathbb{T} -action on the assignment $\text{ev}_X^{\mathbb{R}}(V)$.

Let $\mathbb{M} \in {}_{\mathbb{T}}\text{Mod}$. For $X \in \mathcal{X}$, we can choose an injective \mathbb{k} -module V_X which contains $\mathbb{M}(X)$ as a submodule since the category ${}_{\mathbb{k}}\text{Mod}$ has enough injectives. By the previous adjunction, this inclusion induces a \mathbb{T} -homomorphism $\mathbb{M} \rightarrow \text{ev}_X^{\mathbb{R}}(V_X)$. Hence, we obtain a \mathbb{T} -homomorphism $\mathbb{M} \rightarrow \prod_{X \in \mathcal{X}} \text{ev}_X^{\mathbb{R}}(V_X)$ where a \mathbb{T} -module $\prod_{X \in \mathcal{X}} \mathbb{N}_X$ is defined as $(\prod_{X \in \mathcal{X}} \mathbb{N}_X)(Y) := \prod_{X \in \mathcal{X}} \mathbb{N}_X(Y)$ for \mathbb{T} -modules \mathbb{N}_X . Note that $\text{ev}_X^{\mathbb{R}}(V_X)$ is an injective, since $\text{ev}_X^{\mathbb{R}}$ is a right adjoint. Hence, the \mathbb{T} -module $\prod_{X \in \mathcal{X}} \text{ev}_X^{\mathbb{R}}(V_X)$ is injective, since the product of injectives is an injective. It is readily checked that the \mathbb{T} -homomorphism $\mathbb{M} \rightarrow \prod_{X \in \mathcal{X}} \text{ev}_X^{\mathbb{R}}(V_X)$ is a monomorphism. Hence, \mathbb{M} is embedded into an injective \mathbb{T} -module. \square

Lemma A.2. *The category ${}_{\mathbb{T}}\text{Mod}$ has enough projectives.*

Proof. It is notable that the functor ev_X in the proof of Lemma A.1 is representable, and the representing object is the regular left \mathbb{T} -module $\mathbb{T}(-, X)$:

$$\text{hom}_{\mathbb{T}}(\mathbb{T}(-, X), \mathbb{M}) \cong \text{ev}_X(\mathbb{M}) (= \mathbb{M}(X)).$$

Hence, $\mathbb{T}(-, X)$ is a projective object in ${}_{\mathbb{T}}\text{Mod}$ since the left adjoint ev_X preserves epimorphisms.

We now prove that an arbitrary \mathbb{T} -module \mathbb{M} is a quotient of a projective \mathbb{T} -module. For $v \in \mathbb{M}(X)$, we define a \mathbb{T} -homomorphism $g_{X,v} : \mathbb{T}(-, X) \rightarrow \mathbb{M}$ to be

$$\mathbb{T}(Y, X) \rightarrow \mathbb{M}(Y); f \mapsto f \triangleright v.$$

Then it is easy to check that $\bigoplus_{X \in \mathcal{X}} \bigoplus_{v \in \mathbb{M}(X)} g_{X,v} : \bigoplus_{X \in \mathcal{X}} \bigoplus_{v \in \mathbb{M}(X)} \mathbb{T}(-, X) \rightarrow \mathbb{M}$ is an epimorphism. The domain is projective since the direct sum of projectives is projective. \square

APPENDIX B. GRADED COAUGMENTED COALGEBRAS

Let V be an abelian monoid with the monoid operation $+$ and the unit $0 \in V$. A V -graded $(\mathbb{k}-)$ module is a \mathbb{k} -module C endowed with a decomposition $C = \bigoplus_{\delta \in V} C_{\delta}$. A homomorphism between V -graded modules is a \mathbb{k} -linear map which respects the grading. The tensor product of V -graded modules is given by $(C \otimes D)_{\delta} := \bigoplus_{\substack{\delta_1, \delta_2 \in V \\ \delta = \delta_1 + \delta_2}} C_{\delta_1} \otimes C_{\delta_2}$. Denote by ${}_{\mathbb{k}}\text{Mod}[V]$ the monoidal category of V -graded \mathbb{k} -modules. The unit object is given by \mathbb{k} which we regard as a V -graded \mathbb{k} -module concentrated on the grade $0 \in V$.

We are interested in coaugmented coalgebra objects in the monoidal category ${}_{\mathbb{k}}\text{Mod}[V]$: an object $C \in {}_{\mathbb{k}}\text{Mod}[V]$ with morphisms $\Delta : C \rightarrow C \otimes C$, $\varepsilon_C : C \rightarrow \mathbb{k}$ and $\eta_C : \mathbb{k} \rightarrow C$ in ${}_{\mathbb{k}}\text{Mod}[V]$ which fulfill the conditions of coaugmented coalgebras. This is also called a

V -graded coaugmented coalgebra. Let $1_C := \eta_C(1_{\mathbb{k}})$ and $\bar{\Delta} := \Delta - \text{id}_C \otimes 1_C - 1_C \otimes \text{id}_C$ be the reduced comultiplication. For $m \in \mathbb{N}$, we define a V -graded module $\text{Pr}^m(C)$ as

$$\text{Pr}^m(C) := \{v \in C \mid (\text{id}_C^{\otimes(j-1)} \otimes \bar{\Delta} \otimes \text{id}_C^{\otimes(m-j)})(v) = 0, 1 \leq j \leq m\}.$$

Now, we consider specific coaugmented coalgebras. Let S be a set. It induces a noncommutative polynomial algebra $\mathbb{k}\langle S \rangle$. It has a Hopf algebra structure whose comultiplication Δ is characterized by $\Delta(x) = x \otimes 1 + 1 \otimes x$ for $x \in S$. We are interested in the underlying coaugmented coalgebra of $\mathbb{k}\langle S \rangle$. There are some submodules determined by a total order on S :

- If S is endowed with a total order \leq , then we define $\mathbb{k}\langle S, \leq \rangle$ to be the submodule of $\mathbb{k}\langle S \rangle$ generated by the products $s_1 s_2 \cdots s_r$ for $r \in \mathbb{N}$ and $s_1 \geq s_2 \geq \cdots \geq s_r$. Then $\mathbb{k}\langle S, \leq \rangle$ inherits a coaugmented coalgebra structure from $\mathbb{k}\langle S \rangle$.
- Denote by $a > b$ when $a \geq b$ but $a \neq b$. Let $\mathbb{k}\langle S, < \rangle$ be the submodule generated by the products $s_1 s_2 \cdots s_r$ for $r \in \mathbb{N}$ and $s_1 > s_2 > \cdots > s_r$. This submodule also inherits the coaugmented coalgebra structure from $\mathbb{k}\langle S, \leq \rangle$.

Let $W(S)$ be the set of words generated by letters in S . We regard it as a monoid by the concatenation of words. We say that the monoid $W(S)$ is V -graded if $W(S) = \coprod_{\delta \in V} W_\delta(S)$; the empty word lies in $W_0(S)$; and $w \in W_\delta(S), w' \in W_{\delta'}(S)$ imply $ww' \in W_{\delta+\delta'}(S)$. It is easy to show that, if the monoid $W(S)$ is V -graded, then $\mathbb{k}\langle S \rangle = \mathbb{k}W(S)$ is a V -graded coaugmented coalgebra.

Proposition B.1. *Let (S, \leq) be a toset with a V -grading on the monoid $W(S)$. The inclusion $(\mathbb{k}S)^{\otimes m} \rightarrow \mathbb{k}\langle S, < \rangle^{\otimes m}$ induces an isomorphism of V -graded modules:*

$$(\mathbb{k}S)^{\otimes m} \cong \text{Pr}^m(\mathbb{k}\langle S, < \rangle).$$

Proof. The inclusion $(\mathbb{k}S)^{\otimes m} \rightarrow \mathbb{k}\langle S, < \rangle^{\otimes m}$ factors through $\text{Pr}^m(\mathbb{k}\langle S, < \rangle)$ since $\bar{\Delta}(s) = 0$ in $\mathbb{k}\langle S, < \rangle$ for $s \in S$. It suffices to prove that the factorization $(\mathbb{k}S)^{\otimes m} \rightarrow \text{Pr}^m(\mathbb{k}\langle S, < \rangle)$ is surjective. Suppose that $z \in \text{Pr}^m(\mathbb{k}\langle S, < \rangle)$. Let $1 \leq j \leq m$. Since $z \in \mathbb{k}\langle S, < \rangle^{\otimes m}$, we have

$$z = \sum_{S' \subset S} \sum_{i=1}^{N_{S'}} a_{S',i} \otimes v(S') \otimes b_{S',i}$$

for some $N_{S'} \in \mathbb{N}$, $a_{S',i} \in \mathbb{k}\langle S, < \rangle^{\otimes(j-1)}$, $b_{S',i} \in \mathbb{k}\langle S, < \rangle^{\otimes(m-j)}$ where $v(S') := s_1 s_2 \cdots s_k$ if $S' = \{s_1 < \cdots < s_k\}$. Of course, there are finite S' such that $\sum_{i=1}^{N_{S'}} a_{S',i} \otimes b_{S',i}$ is nonzero. The assumption that $z \in \text{Pr}^m(\mathbb{k}\langle S, < \rangle)$ implies

$$\begin{aligned} 0 &= \left(\text{id}^{\otimes(j-1)} \otimes \bar{\Delta} \otimes \text{id}^{\otimes(m-j)} \right) (z), \\ &= \sum_{S' \subset S} \sum_{i=1}^{N_{S'}} a_{S',i} \otimes \left(\sum_{\substack{S'_1 \cup S'_2 \\ S'_1 \neq \emptyset \neq S'_2}} v(S'_1) \otimes v(S'_2) \right) \otimes b_{S',i}. \end{aligned}$$

Note that $v(S'_1) \otimes v(S'_2)$, $S'_1, S'_2 \subset S$ are linearly independent in $\mathbb{k}\langle S, < \rangle^{\otimes 2}$. Hence, $\sum_{i=1}^{N_{S'}} a_{S',i} \otimes b_{S',i} = 0$ unless $|S'| = 1$. It implies that

$$z \in \mathbb{k}\langle S, < \rangle^{\otimes(j-1)} \otimes \mathbb{k}S \otimes \mathbb{k}\langle S, < \rangle^{\otimes(m-j)}.$$

By applying this to every $1 \leq j \leq m$, we obtain $z \in (\mathbb{k}S)^{\otimes m}$. □

Remark B.2. Note that $(\mathbb{k}S)^{\otimes m} \not\cong \text{Pr}^m(\mathbb{k}\langle S, \leq \rangle)$ in general.

APPENDIX C. A GRADING ON FREE LIE ALGEBRAS

This section presents a refinement of the basis theorem for free Lie algebras in [25]. We introduce a grading on free Lie algebras and Hall sets, and show that the results in that paper holds in the graded sense. The results of this section are applied to compute the eigenmonad by the primitivity ideal in section 11.3.

For a set X , we introduce the following notation:

- Let $M(X)$ be the free magma generated by X . We use the parenthesis $(-, -)$ to denote the magma operation. In accordance with [25], an element of $M(X)$ is called a *tree*. We fix a total order \leq on $M(X)$ such that, for any tree $t = (t', t'') \in M(X)$, we have $t < t''$.
- Let $H(X)$ be the Hall set induced by the toset $(M(X), \leq)$: we set $X \subset H(X)$; for $t \in M(X) \setminus X$, we have $t \in H(X)$ if and only if $t', t'' \in H(X)$, $t' < t''$ and either $t' \in X$ or $(t')'' \geq t''$.

Let V_X be the set of maps δ from X to \mathbb{N} such that $\delta^{-1}(\mathbb{N} \setminus \{0\})$ is finite. We regard V_X as an abelian monoid by the addition. For $\delta \in V_X$, we define

$$|\delta| := \sum_{x \in X} \delta(x) \in \mathbb{N}.$$

We now give particular V_X -gradings on $W(X)$, $H(X)$ and $W(H(X))$ as follows:

- For $\delta \in V_X$, we denote by $W_\delta(X) \subset W(X)$ the subset consisting of words formed by $\delta(x)$ copies of $x \in X$.
- For $\delta \in V_X$, the subset $H_\delta(X) \subset H(X)$ consists of Hall trees which have $\delta(x)$ copies of x as leaves for $x \in X$.
- By using this, we define $W_\delta(H(X)) \subset W(H(X))$ as the subset consisting of words $t_1 t_2 \cdots t_k$ with $t_j \in H_{\delta_j}(X)$, $\delta_j \in V_X$ and $\delta = \sum_{j=1}^k \delta_j$.

The monoids $W(X)$ and $W(H(X))$ are V_X -graded in the sense of appendix B.

Following [25], we define a map $g : M(X) \rightarrow \mathbb{k}\langle X \rangle$ by $g(x) := x$ for $x \in X$; and $g(t) = [g(t'), g(t'')]$ if $t = (t', t'')$. It induces a bialgebra homomorphism $\tilde{g} : \mathbb{k}\langle M(X) \rangle \rightarrow \mathbb{k}\langle X \rangle$ which assigns $g(t_1)g(t_2) \cdots g(t_k)$ to $t_1 t_2 \cdots t_k$ for $t_j \in M(X)$. Hence, the restriction of \tilde{g} to $\mathbb{k}\langle H(X), \leq \rangle \subset \mathbb{k}\langle M(X) \rangle$ gives a coaugmented coalgebra homomorphism $\tilde{g} : \mathbb{k}\langle H(X), \leq \rangle \rightarrow \mathbb{k}\langle X \rangle$. By [25, Theorem 3], the map $\tilde{g} : \mathbb{k}\langle H(X), \leq \rangle \rightarrow \mathbb{k}\langle X \rangle$ is an isomorphism of coaugmented coalgebras.

We refine the aforementioned fact to a graded sense. Note that the decomposition $W(X) = \coprod_{\delta \in V_X} W_\delta(X)$ yields a V_X -grading on the module $\mathbb{k}\langle X \rangle$ by $\mathbb{k}\langle X \rangle \cong \mathbb{k}W(X)$. Likewise, the aforementioned grading on $W(H(X))$ also gives a V_X -grading on $\mathbb{k}\langle H(X), \leq \rangle \subset \mathbb{k}W(H(X))$. Then it is immediate from definitions that \tilde{g} preserves the V_X -grading. Hence, we obtain the following statement:

Proposition C.1. *The map $\tilde{g} : \mathbb{k}\langle H(X), \leq \rangle \rightarrow \mathbb{k}\langle X \rangle$ gives an isomorphism of V_X -graded coaugmented coalgebras.*

In what follows, we give a similar version of the proposition for free Lie algebras. We first derive a lemma by refining the proof of [25, Theorem 4]:

Lemma C.2. *Let $\delta_1, \delta_2 \in V_X \setminus \{0\}$. For $t_1 \in H_{\delta_1}(X)$ and $t_2 \in H_{\delta_2}(X)$, the Lie bracket $[g(t_1), g(t_2)]$ is a linear combination over \mathbb{Z} of $g(t)$ for $t = (t', t'') \in H_{\delta_1 + \delta_2}(X)$ with $t'' \leq \max(t_1, t_2)$.*

Proof. We carefully follow the strategy in the proof of [25, Theorem 4]. We may suppose that X is finite. We prove the statement by induction on the couple $(|\delta_1| + |\delta_2|, \max(t_1, t_2))$ where these couples are ordered by $(d, u) < (e, v)$ if and only if either $d < e$; or $d = e$ and $u < v$. The first step of the induction is obvious.

By the antisymmetry of the Lie bracket, it suffices to consider the case that $t_1 < t_2$. Suppose that $(t_1, t_2) \in H_{\delta_1 + \delta_2}(X)$, i.e. either $t_1 \in X$ or $t_1 = (t'_1, t''_1)$ with $t''_1 \geq t_2$. Then, we set $t = (t_1, t_2)$, which implies $t'' = t_2 \leq \max(t_1, t_2)$. Hence, we obtain $[g(t_1), g(t_2)] = g(t)$ with the desirable property.

We now assume that $(t_1, t_2) \notin H_{\delta_1 + \delta_2}(X)$, i.e. $t_1 = (t'_1, t''_1)$ (derived from $t_1 \notin X$) with $t''_1 < t_2$. By the property of the order, we have $t_1 < t''_1$, hence $t_1 < t''_1 < t_2$. Moreover, $t'_1 < t''_1$ (deduced from $t_1 \in H(X)$) implies $t'_1 < t''_1 < t_2$.

Choose $\delta'_1, \delta'_2 \in V_X$ with $t'_1 \in H_{\delta'_1}(X)$ and $t''_1 \in H_{\delta''_1}(X)$. It is clear that $\delta_1 = \delta'_1 + \delta'_2$ and $|\delta'_1| \neq 0 \neq |\delta''_1|$. By the Jacobi identity, we have

$$[g(t_1), g(t_2)] = [[g(t'_1), g(t_2)], g(t''_1)] + [g(t'_1), [g(t''_1), g(t_2)]].$$

Since $|\delta'_1| + |\delta_2|$ is strictly smaller than $|\delta_1| + |\delta_2|$, the induction hypothesis shows that

$$[g(t'_1), g(t_2)] = \sum n_i g(u_i)$$

where $n_i \in \mathbb{Z}$ and $u_i \in H_{\delta'_1 + \delta_2}(X)$ with $u''_i \leq \max(t'_1, t_2) = t_2$. Similarly, we have $[g(t''_1), g(t_2)] = \sum m_j g(v_j)$ with $v_j \in H_{\delta''_1 + \delta_2}$ and $v''_j \leq \max(t''_1, t_2) = t_2$. Hence, we obtain

$$[g(t_1), g(t_2)] = \sum n_i [g(u_i), g(t''_1)] + \sum m_j [g(t'_1), g(v_j)].$$

We have $|\delta'_1 + \delta_2| + |\delta''_1| = |\delta_1| + |\delta_2|$. We have $\max(u_i, t''_1) < t_2 = \max(t_1, t_2)$ since $u_i < u''_i \leq t_2$. By the induction hypothesis, we deduce that $[g(u_i), g(t''_1)]$ is a linear combination over \mathbb{Z} of $g(t)$ with $t \in H_{\delta_1 + \delta_2}(X)$ and $t'' \leq \max(u_i, t''_1) < \max(t_1, t_2)$. Similarly, we have $|\delta'_1| + |\delta''_1 + \delta_2| = |\delta_1| + |\delta_2|$ and $\max(t'_1, v_j) < t_2 = \max(t_2, t_2)$. Thus the induction implies that $[g(t'_1), g(v_j)]$ is a linear combination of $g(t)$ with $t \in H_{\delta_1 + \delta_2}(X)$ and $t'' \leq \max(t'_1, v_j) < \max(t_1, t_2)$. \square

Definition C.3. Let $L(X)$ be Lie subalgebra of $\mathbb{k}\langle X \rangle$ generated by X . Let $L_\delta(X)$ be the submodule of $L(X)$ generated by Lie brackets consisting of $\delta(x)$ copies of $x \in X$. In other words, it is recursively defined by

$$L_\delta(X) := \sum_{\substack{\delta_1 + \delta_2 = \delta \\ |\delta_1| \neq 0 \neq |\delta_2|}} [L_{\delta_1}(X), L_{\delta_2}(X)]$$

By the definitions, the map $g : M(X) \rightarrow \mathbb{k}\langle X \rangle$ induces a map $g : H_\delta(X) \rightarrow L_\delta(X)$. It is not obvious whether $L_\delta(X)$'s give a grading on $L(X)$; but it is shown as follows.

Lemma C.4. *The map g induces $\mathbb{k}H_\delta(X) \cong L_\delta(X)$.*

Proof. We prove that the elements $g(t), t \in H_\delta(X)$, give a basis of $L_\delta(X)$. Note that $g(t), t \in H_\delta(X)$, are linearly independent [25, Theorem 3]. It is sufficient to prove that they span $L_\delta(X)$. We apply the induction on $|\delta|$. If $|\delta| = 1$, then it is clear since $H_{\delta_x}(X) = \{x\}$ and $L_{\delta_x}(X) = \mathbb{k}x$ where δ_x is the delta function at $x \in X$. We now suppose that $|\delta| > 1$. By definition, the module $L_\delta(X)$ is generated by $[v_1, v_2]$ where $v_i \in L_{\delta_i}(X)$ and $\delta_i \in V_X$ are nonzero maps such that $\delta = \delta_1 + \delta_2$. By the induction hypothesis, we see that each v_i is a linear combination of $g(t), t \in H_{\delta_i}(X)$. By Lemma C.2, $[v_1, v_2]$ is a linear combination of $g(t), t \in H_{\delta_1 + \delta_2}(X) = H_\delta(X)$. \square

Theorem C.5. *We have a V_X -grading on $L(X)$ such that $L(X) \cong \bigoplus_{\delta \in V_X} L_\delta(X)$. Moreover, the map \tilde{g} induces an isomorphism of V_X -graded modules $\mathbb{k}H(X)$ and $L(X)$.*

Proof. By [25, Theorem 4], we have $L(X) \cong \bigoplus_{t \in H(X)} \mathbb{k}g(t)$. By Lemma C.4, we obtain $\bigoplus_{t \in H(X)} \mathbb{k}g(t) = \bigoplus_{\delta \in V_X} \bigoplus_{t \in H_\delta(X)} \mathbb{k}g(t) \cong \bigoplus_{\delta \in V_X} L_\delta(X)$. This implies that $\tilde{g} : \mathbb{k}H(X) \rightarrow L(X)$ is an isomorphism of V_X -graded modules. \square

Corollary C.6. Consider $\delta \in V_X$ such that $\delta(X) \subset \{0, 1\}$. The inclusion $L(X) \rightarrow \mathbb{k}\langle X \rangle$ induces the following isomorphism of δ -components:

$$(L(X)^{\otimes m})_{\delta} \cong \text{Pr}^m(\mathbb{k}\langle X \rangle)_{\delta}.$$

Proof. By Proposition B.1, we have

$$(48) \quad (\mathbb{k}H(X)^{\otimes m})_{\delta} \cong \text{Pr}^m(\mathbb{k}\langle H(X), \langle \rangle \rangle)_{\delta}.$$

The hypothesis that $\delta(X) \subset \{0, 1\}$ implies that $(\mathbb{k}\langle (H(X), \langle \rangle)^{\otimes m} \rangle)_{\delta} = (\mathbb{k}\langle (H(X), \leq \rangle)^{\otimes m} \rangle)_{\delta}$. Hence, Proposition C.1 leads to $\text{Pr}^m(\mathbb{k}\langle X \rangle)_{\delta} \cong \text{Pr}^m(\mathbb{k}\langle (H(X), \leq \rangle)_{\delta}) = \text{Pr}^m(\mathbb{k}\langle (H(X), \langle \rangle)_{\delta})$. On the one hand, by Theorem C.5, the left hand side of (48) is isomorphic to $(L(X)^{\otimes m})_{\delta}$. \square

REFERENCES

- [1] Jean Bénabou, R Davis, A Dold, J Isbell, S MacLane, U Oberst, J E Roos, and Jean Bénabou. Introduction to bicategories. In *Reports of the midwest category seminar*, pages 1–77. Springer, 1967.
- [2] Francis Borceux and Enrico Vitale. Azumaya categories. *Applied Categorical Structures*, 10:449–467, 2002.
- [3] P. M. Cohn. *Free rings and their relations*, volume 19 of *London Mathematical Society Monographs*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London, second edition, 1985.
- [4] Aurélien Djament, Teimuraz Pirashvili, and Christine Vespa. Cohomologie des foncteurs polynomiaux sur les groupes libres. *Documenta Mathematica*, 21:205–222, 2016.
- [5] Samuel Eilenberg and Saunders Mac Lane. On the groups $H(\Pi, n)$. II. Methods of computation. *Ann. of Math. (2)*, 60:49–139, 1954.
- [6] Vincent Franjou, Eric M Friedlander, Teimuraz Pirashvili, and Lionel Schwartz. *Rational representations, the Steenrod algebra and functor homology*, volume 16. Société mathématique de France Paris, 2003.
- [7] Vincent Franjou and Antoine Touzé. *Lectures on functor homology*. Springer, 2015.
- [8] Kazuo Habiro and Gwénaél Massuyeau. The Kontsevich integral for bottom tangles in handlebodies. *Quantum Topology*, 12(4):593–703, 2021.
- [9] Manfred Hartl, Teimuraz Pirashvili, and Christine Vespa. Polynomial functors from algebras over a set-operad and nonlinear Mackey functors. *International Mathematics Research Notices*, 2015(6):1461–1554, 2015.
- [10] Hans-Werner Henn, Jean Lannes, and Lionel Schwartz. The categories of unstable modules and unstable algebras over the Steenrod algebra modulo nilpotent objects. *American Journal of Mathematics*, 115(5):1053–1106, 1993.
- [11] Mai Katada. Actions of automorphism groups of free groups on spaces of Jacobi diagrams. I. In *Annales de l’Institut Fourier*, pages 1–44, 2023.
- [12] Minkyu Kim and Christine Vespa. On analytic exponential functors on free groups. *arXiv preprint arXiv:2401.09151*, 2024.
- [13] Jean-Louis Loday and Bruno Vallette. *Algebraic operads*, volume 346 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, 2012.
- [14] Saunders Mac Lane. Categorical algebra. *Bull. Amer. Math. Soc.*, 71:40–106, 1965.
- [15] Saunders MacLane. *Homology*. Springer Science & Business Media, 2012.
- [16] John W. Milnor and John C. Moore. On the structure of Hopf algebras. *Ann. of Math. (2)*, 81:211–264, 1965.
- [17] Oystein Ore. Formale theorie der linearen Differentialgleichungen. (zweiter teil). *Journal für die reine und angewandte Mathematik*, 1932(168):233–252, 1932.
- [18] Inder Bir S. Passi. *Group rings and their augmentation ideals*, volume 715 of *Lecture Notes in Mathematics*. Springer, Berlin, 1979.
- [19] Teimuraz Pirashvili. Hodge decomposition for higher order Hochschild homology. In *Annales Scientifiques de l’École Normale Supérieure*, volume 33, pages 151–179. Elsevier, 2000.
- [20] Teimuraz Pirashvili. On the PROP corresponding to bialgebras. *Cah. Topol. Géom. Différ. Catég.*, 43(3):221–239, 2002.
- [21] Geoffrey Powell. On analytic contravariant functors on free groups. *arXiv preprint arXiv:2110.01934*, 2021.
- [22] Geoffrey Powell. On the Passi and the Mal’cev functors. *arXiv preprint arXiv:2309.07605*, 2023.
- [23] Geoffrey Powell. Outer functors and a general operadic framework. *Journal of Algebra*, 644:526–562, 2024.

- [24] Geoffrey Powell and Christine Vespa. Higher hochschild homology and exponential functors. *arXiv preprint arXiv:1802.07574*, 2018.
- [25] Christophe Reutenauer. Free Lie algebras. In *Handbook of algebra*, volume 3, pages 887–903. Elsevier, 2003.
- [26] David R Stone. Maximal left ideals and idealizers in matrix rings. *Canadian Journal of Mathematics*, 32(6):1397–1410, 1980.
- [27] Ross Street. Variation through enrichment. *Journal of Pure and Applied Algebra*, 29:109–127, 1983.
- [28] Antoine Touzé. Prerequisites of homological algebra. *Lectures on Functor Homology*, pages 99–149, 2015.
- [29] Christine Vespa. On the functors associated with beaded open Jacobi diagrams. *arXiv preprint arXiv:2202.10907*, 2022.
- [30] Charles A Weibel. *An introduction to homological algebra*. Number 38. Cambridge university press, 1994.
- [31] Nobuo Yoneda. On the homology theory of modules. *J. Fac. Sci. Univ. Tokyo. Sect. I*, 7:193–227, 1954.